

Entanglement detection via entropies in spinor Bose gases

Oliver Stockdale

Kirchhoff Institut für Physik, Universität Heidelberg



UNIVERSITÄT
HEIDELBERG
ZUKUNFT
SEIT 1386

Entanglement in Quantum Fields Workshop

29 June 2021



@GaerrtnerGroup



mbqd.de

Acknowledgements

Theory



Martin
Gärttner



Stefan
Floerchinger



Tobi
Haas

Funding




Experiments




Philipp
Kunkel



Maximilian
Prüfer



Stefan
Lannig



Robin
Strohmaier



Helmut
Strobel




Markus
Oberthaler


How to measure many-body entanglement?

Partition system, make measurements

With access to full state ρ



$$|\psi\rangle = |\psi\rangle_A \otimes |\psi\rangle_B$$



$$|\psi\rangle \neq |\psi\rangle_A \otimes |\psi\rangle_B$$




R. Islam, *et al.*, Nature 528, 77 (2015).

How to measure many-body entanglement?

Partition system, make measurements

With access to full state ρ ... not likely




R. Islam, *et al.*, Nature 528, 77 (2015).


How to measure many-body entanglement?

Partition system, make measurements

With access to full state ρ ... not likely



$$|\psi\rangle = |\psi\rangle_A \otimes |\psi\rangle_B$$



$$|\psi\rangle \neq |\psi\rangle_A \otimes |\psi\rangle_B$$



With access to uncertainties

$$\langle(\Delta\hat{u})^2\rangle_\rho + \langle(\Delta\hat{v})^2\rangle_\rho \geq a^2 + \frac{1}{a^2}$$

Violation flags entanglement


R. Islam, *et al.*, Nature 528, 77 (2015).

L. M. Duan, *et al.*, PRL 84, 2722 (2000).


How to measure many-body entanglement?

Partition system, make measurements

With access to full state ρ ... not likely



$$|\psi\rangle = |\psi\rangle_A \otimes |\psi\rangle_B$$



$$|\psi\rangle \neq |\psi\rangle_A \otimes |\psi\rangle_B$$



With access to uncertainties

$$\langle(\Delta\hat{u})^2\rangle_\rho + \langle(\Delta\hat{v})^2\rangle_\rho \geq a^2 + \frac{1}{a^2}$$

Violation flags entanglement

With access to entropies

$$H(X_A | X_B) + H(Z_A | Z_B) \geq -\log c + H(A | B)$$

R. Islam, *et al.*, Nature **528**, 77 (2015).


L. M. Duan, *et al.*, PRL **84**, 2722 (2000).

M. Berta, *et al.*, Nature Physics **6**, 659 (2012)


How to measure many-body entanglement?

Partition system, make measurements

With access to full state ρ ... not likely



$$|\psi\rangle = |\psi\rangle_A \otimes |\psi\rangle_B$$



$$|\psi\rangle \neq |\psi\rangle_A \otimes |\psi\rangle_B$$



With access to uncertainties

$$\langle(\Delta\hat{u})^2\rangle_\rho + \langle(\Delta\hat{v})^2\rangle_\rho \geq a^2 + \frac{1}{a^2}$$

Violation flags entanglement

With access to entropies

$$H(X_A | X_B) + H(Z_A | Z_B) \geq -\log c + H(A | B)$$

R. Islam, *et al.*, Nature 528, 77 (2015).

L. M. Duan, *et al.*, PRL 84, 2722 (2000).

M. Berta, *et al.*, Nature Physics 6, 659 (2012)

Outline

- Introduce the system: ^{87}Rb spin-1 gas
- Experimental readout: how do we measure entropy?
- An entropic entanglement measurement
- Numerical modelling
- Experimental data



The system

The system

Rubidium-87 BEC in $F = 1$
hyperfine manifold

Characterised by

$$|N_{-1}, N_0, N_{+1}\rangle, \hat{a}_{-1}, \hat{a}_0, \hat{a}_{+1}$$



q represents the
quadratic Zeeman shift



$$H = g[(2\hat{N}_0 - 1)(\hat{N}_{+1} + \hat{N}_{-1}) + 2\hat{a}_0^\dagger \hat{a}_0^\dagger \hat{a}_1 \hat{a}_{-1} + 2\hat{a}_1^\dagger \hat{a}_{-1}^\dagger \hat{a}_0 \hat{a}_0] + q(\hat{N}_{+1} + \hat{N}_{-1})$$

The system

Rubidium-87 BEC in $F = 1$
hyperfine manifold

Characterised by

$$|N_{-1}, N_0, N_{+1}\rangle, \hat{a}_{-1}, \hat{a}_0, \hat{a}_{+1}$$



$$H = g[(2\hat{N}_0 - 1)(\hat{N}_{+1} + \hat{N}_{-1}) + 2\hat{a}_0^\dagger\hat{a}_0^\dagger\hat{a}_1\hat{a}_{-1} + 2\hat{a}_1^\dagger\hat{a}_{-1}^\dagger\hat{a}_0\hat{a}_0] + q(\hat{N}_{+1} + \hat{N}_{-1})$$



Consider only a single spatial mode (for now)

The system

Rubidium-87 BEC in $F = 1$
hyperfine manifold

Characterised by

$$|N_{-1}, N_0, N_{+1}\rangle, \hat{a}_{-1}, \hat{a}_0, \hat{a}_{+1}$$




$$H = g[(2\hat{N}_0 - 1)(\hat{N}_{+1} + \hat{N}_{-1}) + 2\hat{a}_0^\dagger \hat{a}_0^\dagger \hat{a}_1 \hat{a}_{-1} + 2\hat{a}_1^\dagger \hat{a}_{-1}^\dagger \hat{a}_0 \hat{a}_0] + q(\hat{N}_{+1} + \hat{N}_{-1})$$

Consider only a single spatial mode (for now)


Spin-changing collision dynamics

$$H = g[(2\hat{N}_0 - 1)(\hat{N}_{+1} + \hat{N}_{-1}) + 2\hat{a}_0^\dagger \hat{a}_0^\dagger \hat{a}_1 \hat{a}_{-1} + 2\hat{a}_1^\dagger \hat{a}_{-1}^\dagger \hat{a}_0 \hat{a}_0] + q(\hat{N}_{+1} + \hat{N}_{-1})$$




$$|\psi(t=0)\rangle = |0, N, 0\rangle$$

Propagate in time via exact diagonalisation




Measuring entropy: Experimental readout

Rubidium-87 has two hyperfine manifolds





No need for state tomography!
fewer measurements

Transfer ~50% of atoms to $F = 2$,
simultaneously measure spin-1 observables




Measuring entropy: Experimental readout

Rubidium-87 has two hyperfine manifolds



Transfer ~50% of atoms to $F=2$, simultaneously measure spin-1 observables




Quasiprobability distribution

Our spin observables:

$$\hat{S}_x = \hat{a}_0^\dagger(\hat{a}_{+1} + \hat{a}_{-1})/\sqrt{2} + \text{h.c.}$$

$$\hat{Q}_{yz} = i\hat{a}_0^\dagger(\hat{a}_{+1} - \hat{a}_{-1})/\sqrt{2} + \text{h.c.}$$




Quasiprobability distribution

Our spin observables:

$$\hat{S}_x = \hat{a}_0^\dagger(\hat{a}_{+1} + \hat{a}_{-1})/\sqrt{2} + \text{h.c.}$$

$$\hat{Q}_{yz} = i\hat{a}_0^\dagger(\hat{a}_{+1} - \hat{a}_{-1})/\sqrt{2} + \text{h.c.}$$



What **information** can we extract from this distribution?

Distribution is:


- Non-negative
- Normalisable

Quasiprobability distribution

Our spin observables:

$$\hat{S}_x = \hat{a}_0^\dagger(\hat{a}_{+1} + \hat{a}_{-1})/\sqrt{2} + \text{h.c.}$$

$$\hat{Q}_{yz} = i\hat{a}_0^\dagger(\hat{a}_{+1} - \hat{a}_{-1})/\sqrt{2} + \text{h.c.}$$



What **information** can we extract from this distribution?

Distribution is:

- Non-negative
- Normalisable

We associate an **entropy** S_W (called Wehrl entropy) to the distribution Q_ρ via


$$S_W(\rho) = - \int \frac{dS_x dQ_{yz}}{2\pi} Q_\rho(S_x, Q_{yz}) \ln Q_\rho(S_x, Q_{yz})$$

Quasiprobability distribution

Our spin observables:

$$\hat{S}_x = \hat{a}_0^\dagger(\hat{a}_{+1} + \hat{a}_{-1})/\sqrt{2} + \text{h.c.}$$

$$\hat{Q}_{yz} = i\hat{a}_0^\dagger(\hat{a}_{+1} - \hat{a}_{-1})/\sqrt{2} + \text{h.c.}$$



What **information** can we extract from this distribution?

Distribution is:

- Non-negative
- Normalisable

DISCLAIMER

We associate an **entropy** S_W (called Wehrl entropy) to the distribution Q_ρ via


$$S_W(\rho) = - \int \frac{dS_x dQ_{yz}}{2\pi} Q_\rho(S_x, Q_{yz}) \ln Q_\rho(S_x, Q_{yz})$$

Quasiprobability distribution

Our spin observables:

$$\hat{S}_x = \hat{a}_0^\dagger(\hat{a}_{+1} + \hat{a}_{-1})/\sqrt{2} + \text{h.c.}$$

$$\hat{Q}_{yz} = i\hat{a}_0^\dagger(\hat{a}_{+1} - \hat{a}_{-1})/\sqrt{2} + \text{h.c.}$$



Witnessing correlations

$$S_W = - \int \frac{dS_x dQ_{yz}}{2\pi N} Q_\rho(S_x, Q_{yz}) \log Q_\rho(S_x, Q_{yz})$$

Introduce **Wehrl mutual information**

$$I_W(\rho_A : \rho_B) = S_W(\rho_A) - S_W(\rho_A | \rho_B)$$

Measures *total* correlations, inc. classical!

Wehrl mutual information is a
perfect witness for pure states

Witnessing correlations

$$S_W = - \int \frac{dS_x dQ_{yz}}{2\pi N} Q_\rho(S_x, Q_{yz}) \log Q_\rho(S_x, Q_{yz})$$

Introduce **Wehrl mutual information**

$$I_W(\rho_A : \rho_B) = S_W(\rho_A) - S_W(\rho_A | \rho_B)$$

Measures *total* correlations, inc. classical!

Wehrl mutual information is a
perfect witness for pure states

Lower bound on quantum MI

$$I_W(\rho_A : \rho_B) \leq I(\rho_A : \rho_B)$$


Regardless of squeezing parameter, POVM
measurement **bounds quantum MI**

$$I_{\text{POVM}}(\rho_A : \rho_B) \leq I_Q(\rho_A : \rho_B)$$

Theoretical modelling

Time dynamics of correlations


Truncated Wigner result



Numerical simulation of
experimental readout


Time dynamics of correlations

Truncated Wigner result



Numerical simulation of
experimental readout

Measuring variances



$$I_W(\rho_A : \rho_B) = \frac{1}{2} \frac{\det C_A \det C_B}{\det C}$$


$C = \text{inverse cov. matrix}$

Recall, entanglement
witnessed if

$$I_W > 0$$


Time dynamics of correlations

Truncated Wigner result




Numerical simulation of experimental readout

Measuring variances



Gaussian non-Gaussian

Density estimation
How to accurately
measure probability
density function?



$$I_W(\rho_A : \rho_B) = \frac{1}{2} \frac{\det C_A \det C_B}{\det C}$$

$C = \text{inverse cov. matrix}$


Recall, entanglement witnessed if

$$I_W > 0$$

Experimental analysis

Experimental results


Two spatial modes are excited



Different behaviour with
spatially excited modes?


Experimental results

Two spatial modes are excited




Different behaviour with
spatially excited modes?


Total signal, i.e., both modes together




Experimental results





Exact calculation of mutual information



Experimental results





Exact calculation of mutual information





Error via jackknife resampling then
naive error propagation

Experimental results



Error via jackknife resampling then
naive error propagation

Experimental results



Error via jackknife resampling then
naive error propagation

Conclusions & Future Direction

Entropies offer benefits to witnessing entanglement


$$S_W = - \int \frac{dS_x dQ_{yz}}{2\pi N} Q_\rho(S_x, Q_{yz}) \log Q_\rho(S_x, Q_{yz})$$

Conclusions & Future Direction

Entropies offer benefits to witnessing entanglement

$$S_W = - \int \frac{dS_x dQ_{yz}}{2\pi N} Q_\rho(S_x, Q_{yz}) \log Q_\rho(S_x, Q_{yz})$$

Benchmarked the method in Gaussian regime




Conclusions & Future Direction



Entropies offer benefits to witnessing entanglement

$$S_W = - \int \frac{dS_x dQ_{yz}}{2\pi N} Q_\rho(S_x, Q_{yz}) \log Q_\rho(S_x, Q_{yz})$$

Benchmarked the method in **Gaussian** regime



To reap benefits, we need to approximate pdf




Conclusions & Future Direction



Entropies offer benefits to witnessing entanglement

$$S_W = - \int \frac{dS_x dQ_{yz}}{2\pi N} Q_\rho(S_x, Q_{yz}) \log Q_\rho(S_x, Q_{yz})$$

Benchmarked the method in **Gaussian** regime



To reap benefits, we need to approximate pdf



Test the *true* entanglement witness

$$S_M(Q_\pm) \geq 1 + \ln 2$$


Floerchinger, *et al.* arXiv:2106.08788 (2021).

Conclusions & Future Direction

Entropies offer benefits to witnessing entanglement

$$S_W = - \int \frac{dS_x dQ_{yz}}{2\pi N} Q_\rho(S_x, Q_{yz}) \log Q_\rho(S_x, Q_{yz})$$

Benchmarked the method in **Gaussian** regime





More details!



@GaerrtnerGroup mbqd.de

To reap benefits, we need to **approximate pdf**



Test the **true** entanglement witness

$$S_M(Q_\pm) \geq 1 + \ln 2$$

Floerchinger, et al. arXiv:2106.08788 (2021).



Thank you! Questions?

Entropic uncertainty relations

Single system:



$$H(X) + H(Z) \geq -\log_2 c + S(\rho)$$




Not state dependent!

Measure two non-commuting observables

$$c = \max_{x,z} \left(|\langle X_x^A | Z_z^A \rangle|^2 \right)$$

Bipartite system:



$$H(X_A | X_B) + H(Z_A | Z_B) \geq -\log_2 c + S(\rho_A | \rho_B)$$

For a separable state (not entangled)

$$S(\rho_A | \rho_B) \geq 0$$

Lower bound on distillable entanglement: $-\log_2 c - H(X_A | X_B) - H(Z_A | Z_B)$

Theoretical modelling


When state is **Gaussian**, covariance matrix holds all information: $I_W(\rho_A : \rho_B) = \frac{1}{2} \frac{\det C_A \det C_B}{\det C}$

Analytic description

Two-mode squeezed state

$$|\psi\rangle = \sqrt{1 - \lambda^2} \sum_n (-\lambda)^n |n\rangle$$

Approximation for **very short t** when state is **Gaussian**




Calculate covariance matrix elements

Numerical description

Truncated Wigner: evolve operators semi-classically

$$i\hbar \frac{\partial a_i}{\partial t} = \frac{\partial H_W}{\partial a_i^*}$$

Approximate for short t



Simulating the experiment

Truncated Wigner approximation


Each mode is evolved via the equation

$$i\hbar \frac{\partial a_i}{\partial t} = \frac{\partial H_W}{\partial a_i^*}$$

$$\begin{pmatrix} a_{-1} \\ a_0 \\ a_{+1} \end{pmatrix} = \begin{pmatrix} X_1 \\ \sqrt{N} \\ X_2 \end{pmatrix}$$

and beam split via


$$\begin{pmatrix} a_i^A \\ a_i^B \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} a_i^{\text{sys}} \\ a_i^{\text{vac}} \end{pmatrix}$$



where X_i are complex random numbers with $\langle X_i \rangle = 0$ and $\Delta^2 X_i = 1/2$

Error calculation

Jackknife resampling



Calculate variance of jackknife set i : $\text{Var}(X)_i$

Repeat over i data points

Error in variance:

$$\sqrt{n-1} \text{std}(\{\text{Var}(X)_i\})$$

Variance error propagation

$$I_W(\rho_A : \rho_B) = \frac{1}{2} \frac{\det C_A \det C_B}{\det C}$$