

Reasons and means to model preferences as incomplete^{*}

Olivier Cailloux¹ and Sébastien Destercke²

¹ Université Paris-Dauphine, PSL Research University, CNRS, LAMSADE, 75016
PARIS, FRANCE

`olivier.cailloux@dauphine.fr`

² Sorbonne Université, UMR CNRS 7253 Heudiasyc,
Université de Technologie de Compiègne CS 60319 - 60203 Compiègne cedex, France

Abstract. Literature involving preferences of artificial agents or human beings often assume their preferences can be represented using a complete transitive binary relation. Much has been written however on different models of preferences. We review some of the reasons that have been put forward to justify more complex modeling, and review some of the techniques that have been proposed to obtain models of such preferences.

1 Introduction

Preferences of agents are usually assumed to be representable with a weak order (a complete and transitive binary relation). We are interested in discussing the completeness assumption.

Preference models are especially important in two fields: choosing an alternative when it is evaluated according to different aspects (multi-criteria decision making, or MCDM), and picking an alternative whose quality depends on states of the world that are uncertainly known (decision making under uncertainty, or DMU). In MCDM, the common assumption is that the alternatives, i.e., the state of the world, is known without ambiguity, and the difficulty is to determine the structure of the user's preferences over these well-defined alternatives. In DMU, the alternatives are usually not described over several criteria, but the problem is to recommend an alternative given our uncertainty about the world.

In this paper, we review some reasons to relax preference completeness and modeling approaches (either in MCDM or DMU) that support this relaxation. We discuss in particular reasons to consider that the assumption of completeness is empirically falsified. Although these reasons are not new, we think it is interesting to discuss this question here and now because of (as we perceive it) a relative ignorance of these discussions in research fields that use preference models but are not specialized in preference modeling per se, and because of recent and ongoing advances in analysis of incomplete preferences. We try to cover a wide scope by

^{*} This work has received support under the program “LABEX MS2T” launched by the French Government and implemented by ANR with the references ANR-11-IDEX-0004-02.

discussing some of the goals, assumptions and basic definitions related to preference modeling and reviewing a wide range of techniques for obtaining such models. In counterpart, this review does not claim to be comprehensive and does not provide technical details. To further simplify the discussion, we pretend that the MCDM and DMU contexts are sharply separated. (In reality, it is often possible to cover MCDM contexts while taking uncertainty into account [Keeney and Raiffa, 1993].) We also do not discuss transitivity.

We briefly present the MCDM and DMU settings considering completeness in Section 2. Section 3 discusses completeness in descriptive and normative approaches (recalling their difference at the same time). Finally, we review models that depart from completeness in Section 4.

2 Completeness in classical and less classical settings

In this section, we are going to recall the main models that consider completeness and transitivity of preferences as a consequence of natural requirements, if not as pre-requisite of any preference modeling. We will also recall normative views and descriptive views of these concepts.

2.1 MCDM

We consider a simple and classical setting in MCDM. We assume that the alternatives are evaluated using a set of criteria \mathcal{G} , each having an evaluation scale X_g . The set of all possible alternatives is $\mathcal{X} = \prod_{g \in \mathcal{G}} X_g$, that is, every combination of evaluations are considered possible. We are interested in a preference relation \succeq defined as a binary relation over \mathcal{X} .

Example 1. Say the Decision Maker (DM) must choose what to plant in her garden. The set of alternatives \mathcal{X} are all possible vegetables, the criteria $\mathcal{G} = \{g_1, g_2, g_3\}$ measure the taste, quantity, and price of each vegetable, and $A \subseteq \mathcal{X}$ are the vegetables that are available this year for planting. $X_{g_1} = \{A, B, C, D\}$, a set of labels, with x_1 representing the taste of the vegetable $x \in \mathcal{X}$ as considered by the DM (A is the worst taste, D the best), $X_{g_2} = [0, 100]$, with x_2 representing the number of meals that the DM would enjoy if deciding to plant x , and $X_{g_3} = \mathbb{R}$, where x_3 indicates the price to pay for planting x .

Typical approaches in MCDM assume that there is some real-valued function $v : \mathcal{X} \rightarrow \mathbb{R}$ mapping alternatives to their values, and that $x \succeq y$ iff $v(x) \geq v(y)$.

2.2 DMU

In the simplest form of DMU considered here (SDMR, for Simple Decision Making under Risk), we consider a set \mathcal{S} of possible states of the world, a finite set of consequences \mathcal{C} , and each act $x : \mathcal{S} \rightarrow \mathcal{C}$ is modeled

as a function where $x(s)$ is the consequence of performing x when s is the actual state of the world. Define \mathcal{X} in this case as the set of possible acts (thus $\mathcal{X} = C^S$). In SDMR, uncertainty is modeled by a probability measure p over the power set of S , $\mathcal{P}(S)$, thus with $p(s) \in [0, 1]$ indicating the probability of occurrence of s (with $s \subseteq S$), and $p(S) = 1$. We consider a preference relation \succeq defined as a binary relation over \mathcal{X} . Given an act x and a probability measure p , it is usually convenient to view x as p_x , a probability mass over the consequences: define $p_x : C \rightarrow [0, 1]$ as $p_x(c) = p(x^{-1}(c))$, where $x^{-1}(c)$ designate the set of states in which x leads to the consequence c . Such a p_x is usually called a lottery.

Example 2. Assume you want to go out and wonder about taking or leaving your umbrella. You consider relevant weather state to be going from A = “shiny” to D = “stormy”, with $A, B, C, D \subseteq S$. Two simple actions are x_1 : “leave the umbrella home” and x_2 : “take the umbrella”, and the consequences are c_1 : “encumbered” (when taking your umbrella, irrelevant of the weather), c_2 : “free” (when leaving your umbrella and weather is A), and c_3, c_4, c_5 being “slightly wet”, “wet” and “very wet”, when leaving your umbrella and weather being respectively B, C , or D . Assume the probabilities of the states A, B, C, D are respectively 0.2, 0.3, 0.4 and 0.1. Then the act x_1 can also be described as p_{x_1} defined as $p_{x_1}(c_1) = 0$, $p_{x_1}(c_2) = 0.2$, $p_{x_1}(c_3) = 0.3$, $p_{x_1}(c_4) = 0.4$, $p_{x_1}(c_5) = 0.1$, and similarly the constant act x_2 can be associated to p_{x_2} , with $p_{x_2}(c_1) = 1$.

In most DMU frameworks, consequences can be mapped to a real-valued reward or utility through a function $u_1 : C \rightarrow \mathbb{R}$, in which case $u_1(x(s))$ denotes the utility of performing x in state s , and acts can be evaluated using a utility function $u : \mathcal{X} \rightarrow \mathbb{R}$ defined as $u(x) = \sum_{s \in S} p(s)u_1(x(s))$, such that $u(x) \geq u(y)$ iff $x \succeq y$. It follows from this definition that u and u_1 are coherent, in the following sense: given an act x that brings a consequence c with probability one, $u(x) = u_1(c)$.

Expected utility has been justified axiomatically by different authors, the main ones being Savage [1972], De Finetti [2017] and von Neumann and Morgenstern [2004] (hereafter, vNM). In the De Finetti setting, utilities are given as random variables, and a precise price can be associated to each random variable. That reasoning should be probabilistic and choices made according to expected utility follow from two axioms: linearity and boundedness of those prices. vNM postulate conditions on \succeq ensuring that utility functions u and u_1 satisfying the above conditions exist. The axioms assume completeness of the preferences, and the probabilities are assumed to be given. In the Savage setting, both probabilities and expected utility follow from axioms about preferences between acts. In particular, his first axiom (P1) is that any pair of act should be comparable. Completeness is therefore postulated in the axioms, and expected utility and probabilistic reasoning follow from the axioms.

While these theoretical constructs have set very strong foundations for the use of probabilities, in practice experiments such as the Ellsberg [1961] urn (contradicting Savage sure-thing principle) suggest that people do not always act according to expected utility.

Since then, many different extensions have been proposed [Wakker, 2010, Quiggin, 2012]. Others propose to relax the probabilistic assump-

tion, for instance by considering a possibilistic setting (e.g., Dubois et al. [2003] discusses Savage-like axioms), by considering sets of probabilities such as in decision under ambiguity [Gajdos et al., 2008], or by simply considering completely missing information, such as Wald’s 1992 celebrated maximin criterion.

Such extensions address some of the observed behaviours deviating from expected utility, but keep assuming that all acts are comparable, i.e., that preferences are complete. Yet, if one considers that probabilities or utilities may be incompletely known, the next step is to consider that preferences as well could be.

All models presented or advocated thus far assume that it is possible to order alternatives according to their value, as given by a real function. Thus, they require \succeq to be complete and transitive (by which we mean that if \succeq is incomplete or not transitive, then no suitable function exists).

The two contexts we have introduced, multiple criteria and risk, use as a basis the notion of a preference relation. We need to say a word about what those preferences really represent and what the goal of modeling those may be. This is required to give meaning to discussion about which conditions are reasonable to postulate about \succeq , and how to check whether they hold.

3 Completeness in descriptive and normative approaches

3.1 Descriptive approach

In the descriptive approach to preferences, the goal of the model is to reflect the observed behavior of a DM. Typically, a set of sample choices of the DM is first collected, say, of choices of food products in his favorite store, and we would then try to obtain the model that best reflects his choice attitude. Or, we would query an individual’s preference about pairs of objects, and then try to build a predictive model on the whole set of possible pairs of alternatives (a method called active learning in the machine learning community). Such a model may be used to predict his behavior, e.g. for marketing or regulation purposes. In those cases where its purpose is to replace the DM by automatic choices, it is debatable whether the normative approach would not be better suited.

Here is what von Neumann and Morgenstern (vNM) say about the preference relation: “It is clear that every measurement — or rather every claim of measurability — must ultimately be based on some immediate sensation...In the case of utility the immediate sensation of preference...provides this basis...Let us for the moment accept the picture of an individual whose system of preferences is all-embracing and complete...we expect him, for any two alternative events which are put before him as possibilities, to be able to tell which of the two he prefers.”

As the preference relation is considered basic under a purely descriptive approach, the preference itself should be easily observable, and uncontroversial. This permits to empirically test conditions postulated on \succeq . Either by asking the individual what she prefers (in which case we

“observe” her answer), or by presenting a choice set and observing what she picks from that set (in real life or in a laboratory experiment). In the first case, we have however to be clear about what is meant by “prefers”, which is a reason why many experimenters prefer to go for the latter strategy.

3.2 Normative approach

Under the normative approach, the goal is to model on how the DM ought to choose rationally. Rationality may corresponds to accepted external norms (axioms), or to rules accepted by the DM after careful thinking. Hence, the decision outcome using such approach may differ from empirically observed decisions. Consider as an example a recruiter in an enterprise who wants to model the recruitment procedure. After having collected data, it may appear that for some (possibly unconscious) reason, the recruitment is biased against some particular socio-economic category. The DM may then want to find a recruitment strategy that avoids such biases, therefore actively trying to build a model contradicting empirical observations.

When the focus is on providing decision help to a DM by letting him think about the norms he prefers, rather than considering them as external rationality norms, the approach is often termed prescriptive, or constructive. Carefully and formally defining such terms is important, especially as different authors use them somewhat differently [Roy, 1993, Tsoukiàs, 2007]. As we do not want to focus on such issues, we will use the term normative as an umbrella.

McClennen [1990], Guala [2000] discuss philosophical grounds for accepting a normative model. Anand [1987], Mandler [2001] discuss normative grounds for usual axioms about preferences, including completeness.

What is the best approach to adopt is not always obvious. For instance recommender systems often adopt a descriptive approach, mimicking choice behaviour to increase profit, for example. Yet, such an approach reflect our cognitive limitations, and it is not clear that a more thoughtful or normative-oriented (at least partially) process would not lead to more satisfying, diverse or surprising recommendations that may moreover be explainable.

3.3 Defining and testing incompleteness

A first step to define and test incompleteness in preferences is to define what is meant by preference and by the relation \succeq , as its everyday use can be ambiguous: Frankfurt [1971] gives seven interpretations of “to want to”, and this exercise transposes, *mutatis mutandis*, to the notion of preference.

Expanding on vNM, we define that the DM *prefers* a to b when expressing an intuitive attraction towards a when presented with a and b , or an equal attraction towards a and b ; and this attraction does not change along a reasonable time span and as well as when irrelevant changes in the context happen. Here, we assume that a, b are alternatives in \mathcal{X} described by their evaluations on the criteria (in MCDM) or by the relevant

probability distributions and consequences (in SDMR), and consider as irrelevant changes anything that does not change those descriptions. Under this definition, postulating completeness of \geq amounts to say that choices of the DM will not change along time or when irrelevant changes happen. While this is not the only possible definition (others will be mentioned), it appears reasonable and sufficiently formal to make the condition empirically testable.

A first, immediate argument against completeness is that preferences are not stable over time, a well-accepted fact in experimental psychology. Quoting Tversky [1969], individuals “are not perfectly consistent in their choices...people often choose x in some instances and y in others. Furthermore, such inconsistencies are observed even in the absence of systematic changes in the decision maker’s taste...inconsistencies reflect inherent variability...in the evaluative process.” This argument alone would not be strong enough, at least from a normative standpoint, as we may consider that underlying preferences are complete, but that choices are noisy observations of such preferences.

However, a second argument also inherited from empirical psychology is that preferences may change in a non-random, consistent way according to changes in the presentation of the alternatives or the context that should have no impact from a normative point of view.

Let us also note that von Neumann and Morgenstern [2004, p. 630] themselves considered completeness as a strong condition: “it is very dubious, whether the idealization of reality which treats this postulate as a valid one, is appropriate or even convenient”.

3.4 Empirical evidence of incompleteness

In multicriteria contexts, psychologists have shown systematic differences between the so-called choice and matching elicitation procedures [Tversky et al., 1988]. Assume you want to know which of two alternatives x, y the DM prefers, in a problem involving two criteria. You can present both and directly ask for a choice. Alternatively, the matching procedure consists in presenting x with its two evaluations $g_1(x), g_2(x)$, and present y' with only $g_1(y') = g_1(y)$, and ask the DM for which value $g_2(y')$ y' would be indifferent to x . Assuming \geq satisfies dominance and transitivity, you then know that $x \geq y$ iff $g_2(y') \geq g_2(y)$. The authors confirm the prominence hypothesis stating that the more prominent criterion has more importance in choice than in matching. One of their study confront the subject to a hypothetical choice between two programs for control of a polluted beach. Program x completely cleans up the beach at a yearly cost of \$750 000; program y partially cleans it up for a yearly cost of \$250 000. They assume that pollution is the more prominent criterion here, hence expect that x will be chosen more often in choice than in matching. Indeed, 48% out of the 104 subjects confronted with a choice procedure selected x , whereas only 12% out of the 170 subjects selected it in a matching procedure. Note that similar studies hold for lotteries in SDMR [Luce, 2000].

This phenomenon is known as preference reversal due to a breach of procedure invariance. Another reversal is the one due to description

invariance (or framing effect), showing that preferences can change by changing alternative descriptions. In Tversky and Kahneman [1981], two groups have to choose a program to prepare against an epidemic outbreak that would result otherwise in 600 deaths. The two groups are presented with the same numeric alternatives x and y , but on the first group the alternatives is presented in terms of numbers of life saved, while in the second they are presented in terms of death counts. The experiment shows that preferences are consistently different in the two groups. It is indeed well-known that losses and gains, even if identical, are perceived differently Thaler [1980].

Numerous other studies exist that show and discuss preference reversal effects [Deparis, 2012, Ch. 2], [Lichtenstein and Slovic, 2006a, Tversky et al., 1990, Kahneman et al., 1981, Kahneman and Tversky, 2000]. How to best account for and predict preference reversals is still debated, but their existence is consensual [Wakker, 2010, Birnbaum, 2017]. Some skeptic did try to show that preference reversals could be attributed to deficiencies in the design of the studies or lack of realism, but finally came around Slovic and Lichtenstein [1983].

This shows that the \succeq relation cannot be expected to be complete given our definition. For some alternatives, individuals may be led to declare different preferences, denoting an absence of a clear, intuitive preference for each pairs of alternatives. This has been studied empirically [Slovic and Tversky, 1974, MacCrimmon and Larsson, 1979, Lichtenstein and Slovic, 2006b], and Savage [1972, pp. 101 – 103] famously reported that it happened to him. Luckily, preference reversals are not universal [Lichtenstein and Slovic, 2006a, p. xvi]: individuals preferences are not entirely manipulable.

One may of course want to preserve completeness of preferences, for example to preserve mathematical and computational simplicity. One way to do so, common in experimental psychology, is to restrict further the frame in which preferences are considered. For instance, Luce [2000] indicates clearly that he studies preferences in terms of choice, not judgment; MacCrimmon et al. [1980] excludes some kind loteries from the scope of the model. In such cases, completeness may well be justified, at the price of losing generality. In other settings, such as normative approaches or recommender systems, it is unclear that such reductions should be enforced, as they may lead to constraints or axioms that may appear unnatural to the user, or hard to impose in practice.

Note that in such experiments, incomparability is actually not observed *per se*, as experimental protocol forces to make a choice. Some other studies Deparis et al. [2012] are interested in the behavior of individuals when they are allowed to make explicit statements of incomparability. Others such Danan and Ziegelmeyer [2006] propose to consider that an incomparability is observed whenever the DM is ready to pay a small price to postpone the decision.

Above experiments suggest that incomparability may be, to some extent, an intrinsic property of individuals preferences. Yet, even when assuming completeness of preferences, for example in normative view, it may well be that provided information is insufficient to obtain a fully precise models, in which case incompleteness can be due to an incompletely

specified model.

4 Dropping completeness

4.1 Incompleteness in MCDM

Some approaches in MCDM in the family of outranking methods [Roy, 1996, Greco et al., 2016, Bouyssou et al., 2000, 2006], which are in the process of being axiomatized [Bouyssou and Pirlot, 2015], can represent incomparabilities. A much used idea is to take into account two points of view leading to weak-orders \succeq^1 and \succeq^2 , then define $\succeq = \succeq^1 \cap \succeq^2$. Thus, when the two weak-orders strongly disagree about some pair of objects, the result can declare them incomparable. As an example, consider (a simplification of) the ELECTRE III method (our much simplified description only consider the aspects sufficient to obtain incomparabilities). It builds a concordance relation C that determines whether alternative x is sufficiently better than y , by accounting only for the criteria in favor of x . For example, a model could declare that xCy iff x is better than or equal to y for at least two criteria. It also defines a discordance relation D . In ELECTRE III, the performance of x on a given criterion g may be considered as “unacceptably low” compared to the performance of y on g . For example, a model could consider that x is unacceptably low compared to y on g iff $y_g - x_g \geq 2$, where X_g would be numerical. Then define $x \succeq y$ iff xCy and $\neg(xDy)$.

Consider an example with $\mathcal{X} = \mathbb{R}^3$, each criteria to be maximized, and the following two alternatives: $x = (0, 0, 2), y = (1, 1, 0)$. Using the relations C and D defined above, we obtain xCy and xDy , thus x and y are incomparable in the resulting model. Such approaches tend to consider incomparabilities as intrinsic to the system, since even a completely specified model could lead to incomparabilities.

Robust methods in MCDM exist that distinguish conclusions (i.e., preferences) that hold for sure, given limited preferential information from the DM, from conclusions that possibly hold. Such methods typically start from a classe \mathbf{M} of possible models (equivalent to version or hypothesis spaces in machine learning) assumed to represent the DM preferences. A robust method, given a class \mathbf{M} and a set of constraints \mathbf{C} reducing the set of possibles models (typically preference statements of the DM), will consider that a is necessarily preferred to b , $a \succeq^N b$, whenever $a \succeq b$ for all relations \succeq in \mathbf{M} that satisfy \mathbf{C} [Greco et al., 2008].

Example 3. Assume that the only thing you know about the DM is that she prefers $x = (0, 0, 2)$ to $y = (0, 4, 0)$, and you assume that \succeq satisfies preferential independence, meaning that the way two alternatives compare does not change when changing equal values on a given criterion. Thus, \mathbf{M} contains all relations that satisfy preferential independence, and \mathbf{C} is the constraint $x \succ y$. You may then conclude that $a = (3, 0, 2)$ is preferred to $b = (3, 4, 0)$, thus, $a \succ^N b$, but you ignore whether $c = (1, 1, 1)$ is preferred to $d = (0, 2, 2)$, thus, $\neg(c \succeq^N d)$ and $\neg(d \succeq^N c)$.

Such approaches consider that incomparabilities follow from a lack of knowledge, and are not intrinsic to the modeled preference relation, as in principle one could collect enough constraints C about M to identify a unique model (or a set of model leading to only one possible relation $>$ on a set of alternatives).

4.2 Incompleteness in DMU

As recalled in Section 1, probability theory and expected utility are the most widely used tools when having to decide under uncertainty, and naturally induce completeness of preferences. It should however be noted early scholars were critical about the fact that completeness could hold in practice, as recalled previously. Many attempts to relax the completeness axioms does so by considering axioms leading to deal with sets of utilities and sets of probabilities Aumann [1962], entangling together aspects about decision and about information modeling.

Keeping precise probabilities but not expected utility Even when having precise probabilities, there are alternatives to expected utility that induce incomplete preferences. One of them that is particularly interesting is the notion of stochastic dominance Levy [1992]. Assuming that the set of consequences is completely ordered by preference, which we denote by $C = \{c_1, \dots, c_n\}$ where c_i is preferred to c_{i-1} , then a lottery p_x is said to stochastically dominate p_y iff

$$P_x(\{c_1, \dots, c_i\}) = \sum_{j=1}^i p_x(c_j) \leq P_y(\{c_1, \dots, c_i\}) = \sum_{j=1}^i p_y(c_j). \quad (1)$$

Since Inequality (1) can be satisfied for some consequence c_i and not for others, possible incomparabilities immediately follow.

Example 4. Consider again the space $C = \{c_1, \dots, c_3\}$ of consequences and the following probability masses (induced by different acts x_1, x_2, x_3) defined over them, given in vectorial forms: $p_1 = (0.2, 0.3, 0.5)$, $p_2 = (0.1, 0.3, 0.6)$ and $p_3 = (0.3, 0, 0.7)$. From those, it can be checked that that x_2 stochastically dominates x_1 , while x_3 is incomparable to both x_1 and x_2 , according to stochastic dominance.

The notion of stochastic dominance has some very attractive properties, as:

1. it does not necessitate to define utilities over consequences, and merely requires them to be linearly ordered;
2. it can be perceived as a criterion allowing for utilities to be ill-defined, as p_x stochastically dominates p_y if and only if x has a higher expected utility than y for any increasing utility function u defined over C .

Incompleteness from non-precise probabilities In the past few decades, different scholars have challenged the need for precise probabilities associated to classical axiomatics, advocating the use of imprecisely defined prices (expected values) or of imprecisely defined probabilities. To mention but a few:

- Levi Levi [1983] advocates the uses of sets of probabilities within a logical interpretation of probabilities;
- Walley Walley [1991] extends de Finetti axioms by assuming that an agent would give different buying and selling prices for an act, therefore allowing indecision if the given price is between these two bounds;
- Shafer and Vovk Shafer and Vovk [2005] explores a probabilistic setting centered on the notion of Martingale.

Such theories can most of the time be associated to the use convex sets of probabilities, and give rise to decision rules that extend expected utility but do allow incomparabilities. Once we accept that a convex set \mathcal{P} of probabilities (or a formally equivalent representation) can represent our knowledge, incompleteness ensues.

A prototypical way to induce incompleteness between acts from incompleteness in probabilities is to adapt expected utility criterion, and among rules doing so, maximality is a popular one (it is championed by Walley, but is considered as early as the 60's Aumann [1962]). Given acts x_1, x_2 , maximality says that

$$x_1 \succ x_2 \text{ iff } u(x_1) \geq u(x_2) \text{ for all } p \in \mathcal{P},$$

still with $u(x) = \sum_{s \in S} p(s)u_1(x(s))$. It is clear that this reduces to expected utility when \mathcal{P} is a singleton.

Example 5. Going back to Example 2, we have that

$$u(x_1) = p(A) - p(B) \text{ and } u(x_2) = p(B) - p(A)$$

which means that the two acts will be incomparable according to maximality as soon as \mathcal{P} contains a probability where $p(A) = p(B)$ that is not on the border of \mathcal{P} (i.e., it contains at least one mass where $p(A) < p(B)$, and another where $p(B) > p(A)$).

It should be noted that other authors have proposed different rules: for instance Levi Levi [1983] recommends to use a decision rule, often called E-admissibility, that does not give rise to an incomplete order between acts, but rather selects all the acts that are Bayes optimal according to at least one probability $p \in \mathcal{P}$. In terms of order, this comes down to consider a set of possible linear ordering, and to retain only those elements that are maximal for at least one of them.

Working with sets of probabilities and utilities Sets of probabilities are helpful to represent incomplete beliefs or lack of information, yet it is natural to also consider cases where the DM cannot provide a fully accurate estimation of utilities associated to consequences, or even

to completely order them. In some sense, stochastic dominance is an extreme view of such a case, where consequences are ordered but the utility function is left totally unspecified.

Other works have dealt with partially specified utilities:

- Dubra et al. [2004] represents preferences over lotteries by a set of utility functions. Preference holds whenever the expected utility for the preferred alternative is higher for all utility functions. This idea has been applied in other contexts [Ok, 2002].
- Dubra and Ok [2002] propose to view the preference relation as a completion of an intuitive partial preference relation: the DM knows intuitively the result of some comparisons, and compute the other ones by applying some reasoning process. They also obtain a preference relation that is representable using a set of utility functions. This approach directly tackles some of the shortcomings described in Section 3.3.
- Manzini and Mariotti [2008] use a utility function and a vagueness function, representing the preference using intervals of utilities rather than real valued utilities.

There exist a few works where both requirements of precise probabilities and utilities are relaxed. This can be traced back at least to Aumann [1962] whose axioms do not require uniqueness of utilities: $x > y \Rightarrow u(x) > u(y)$, without requiring the reverse. More recently, Galaabaatar and Karni [2013] are interested in the Savage-like context where probabilities are unknown and represent an incomplete preference relation in uncertainty using a set of pairs of probabilities and utilities.

5 Incompleteness: absence of knowledge or knowledge of absence?

We have tried to browse a general picture of reasons why preference modeling should accommodate for incompleteness, and how it can do so in multi-criteria problems and uncertainty modeling.

One issue that transpired in most of the paper is whether incompleteness should be considered as an intrinsic, or ontic property of the preferences, in which case incomparability express a knowledge of absence of relation, or if incompleteness should be considered as an incomplete, epistemic description of a complete order, in which case it expresses an absence of knowledge. This mirrors different views about probability sets (Walley’s consider that they model belief, without assuming an existing precise unknown distribution, while robust Bayesians consider the opposite).

Our opinion is that both views can be legitimate in different settings, and also that beyond the philosophical interest of distinguishing the two, this can have an important practical impact: knowing that incomparabilities are observable facts may influence strongly our information collection protocol; also, a same piece of information will be interpreted differently. If a DM pick act a among three acts $\{a, b, c\}$, in the epistemic interpretation, we would deduce $a > \{b, c\}$, but in the ontic one we could only deduce that a is a maximal element ($\neg(b > a)$ and $\neg(c > a)$)

Bibliography

- P. Anand. Are the preference axioms really rational? *Theory and Decision*, 23(2):189–214, Sept. 1987. ISSN 0040-5833, 1573-7187. doi:10.1007/BF00126305.
- R. J. Aumann. Utility Theory without the Completeness Axiom. *Econometrica*, 30(3):445–462, 1962. ISSN 0012-9682. doi:10.2307/1909888.
- M. H. Birnbaum. Empirical evaluation of third-generation prospect theory. *Theory and Decision*, pages 1–17, June 2017. ISSN 0040-5833, 1573-7187. doi:10.1007/s11238-017-9607-y.
- D. Bouyssou and M. Pirlot. A consolidated approach to the axiomatization of outranking relations: a survey and new results. *Annals of Operations Research*, 229(1):159–212, June 2015. ISSN 0254-5330, 1572-9338. doi:10.1007/s10479-015-1803-y.
- D. Bouyssou, T. Marchant, M. Pirlot, P. Perny, A. Tsoukiàs, and P. Vincke. *Evaluation and decision models: a critical perspective*. Number 32 in International Series in Operations Research & Management Science. Kluwer Academic, Dordrecht, 2000. URL <http://doi.org/10.1007/978-1-4615-1593-7>.
- D. Bouyssou, T. Marchant, M. Pirlot, A. Tsoukiàs, and P. Vincke. *Evaluation and decision models with multiple criteria: Stepping stones for the analyst*. International Series in Operations Research and Management Science, Volume 86. Springer, Boston, 1st edition, 2006. ISBN 0-387-31098-3. URL <http://doi.org/10.1007/0-387-31099-1>.
- E. Danan and A. Ziegelmeyer. Are preferences complete? An experimental measurement of indecisiveness under risk, 2006. URL <http://www.ericdanan.net/research>. Working paper.
- B. De Finetti. *Theory of probability: A critical introductory treatment*, volume 6. John Wiley & Sons, 2017.
- S. Deparis. *Etude de l'Effet du Conflit Multicritère sur l'Expression des Préférences : une Approche Empirique*. PhD thesis, École Centrale Paris, Châtenay-Malabry, France, June 2012.
- S. Deparis, V. Mousseau, M. Öztürk, C. Pallier, and C. Huron. When conflict induces the expression of incomplete preferences. *European Journal of Operational Research*, 221(3):593–602, Sept. 2012. ISSN 03772217. doi:10.1016/j.ejor.2012.03.041.
- D. Dubois, H. Fargier, and P. Perny. Qualitative decision theory with preference relations and comparative uncertainty: An axiomatic approach. *Artificial Intelligence*, 148(1-2):219–260, 2003.
- J. Dubra and E. A. Ok. A Model of Procedural Decision Making in the Presence of Risk*. *International Economic Review*, 43(4):1053–1080, Nov. 2002. ISSN 1468-2354. doi:10.1111/1468-2354.t01-1-00048.
- J. Dubra, F. Maccheroni, and E. A. Ok. Expected utility theory without the completeness axiom. *Journal of Economic Theory*, 115(1):118–133, Mar. 2004. ISSN 0022-0531. doi:10.1016/S0022-0531(03)00166-2.
- D. Ellsberg. Risk, Ambiguity, and the Savage Axioms. *Quarterly Journal of Economics*, 75:643–669, 1961.

- H. G. Frankfurt. Freedom of the Will and the Concept of a Person. *The Journal of Philosophy*, 68(1):5–20, Jan. 1971. doi:10.2307/2024717.
- T. Gajdos, T. Hayashi, J.-M. Tallon, and J.-C. Vergnaud. Attitude toward imprecise information. *Journal of Economic Theory*, 140(1): 27–65, 2008.
- T. Galaabaatar and E. Karni. Subjective Expected Utility With Incomplete Preferences. *Econometrica*, 81(1):255–284, Jan. 2013. ISSN 1468-0262. doi:10.3982/ECTA9621.
- S. Greco, V. Mousseau, and R. Słowiński. Ordinal regression revisited: multiple criteria ranking using a set of additive value functions. *European Journal of Operational Research*, 191(2):415–435, Dec. 2008. doi:10.1016/j.ejor.2007.08.013.
- S. Greco, M. Ehrgott, and J. R. Figueira, editors. *Multiple Criteria Decision Analysis - State of the Art Surveys*. Springer, 2nd edition, 2016. URL <http://doi.org/10.1007/978-1-4939-3094-4>.
- F. Guala. The logic of normative falsification: rationality and experiments in decision theory. *Journal of Economic Methodology*, 7(1): 59–93, Jan. 2000. ISSN 1350-178X. doi:10.1080/135017800362248.
- D. Kahneman and A. Tversky, editors. *Choices, Values, and Frames*. Cambridge University Press, 2000. ISBN 978-0-521-62749-8.
- D. Kahneman, P. Slovic, and A. Tversky. *Judgement under uncertainty - Heuristics and biases*. Cambridge University Press, Cambridge, 1981.
- R. L. Keeney and H. Raiffa. *Decisions with Multiple Objectives: Preferences and Value Tradeoffs*. Cambridge University Press, 2nd edition, Aug. 1993. ISBN 978-0-521-43883-4.
- I. Levi. *The enterprise of knowledge: An essay on knowledge, credal probability, and chance*. MIT press, 1983.
- H. Levy. Stochastic dominance and expected utility: survey and analysis. *Management science*, 38(4):555–593, 1992.
- S. Lichtenstein and P. Slovic, editors. *The Construction of Preference*. Cambridge University Press, 2006a. ISBN 0-521-83428-7.
- S. Lichtenstein and P. Slovic. Reversals of preferences between bids and choices gambling decisions. In S. Lichtenstein and P. Slovic, editors, *The Construction of Preference*, pages 52–68. Cambridge University Press, 2006b. ISBN 0-521-83428-7.
- R. D. Luce. *Utility of Gains and Losses: Measurement-theoretical and Experimental Approaches*. Lawrence Erlbaum Publishers, Mahwah, New Jersey, 2000.
- MacCrimmon, Stanbury, and Wehrung. Real money lotteries: A study of ideal risk, context effects, and simple processes. In T. S. Wallsten, editor, *Cognitive processes in choice and decision behavior*. L. Erlbaum Associates, Hillsdale, N.J., 1980.
- K. R. MacCrimmon and S. Larsson. Utility Theory: Axioms Versus ‘Paradoxes’. In M. Allais and O. Hagen, editors, *Expected Utility Hypotheses and the Allais Paradox*, number 21 in Theory and Decision Library, pages 333–409. Springer Netherlands, 1979. ISBN 978-90-481-8354-8 978-94-015-7629-1. URL http://doi.org/10.1007/978-94-015-7629-1_15.
- M. Mandler. A difficult choice in preference theory: rationality implies completeness or transitivity but not both. In E. Millgram, editor, *Varieties of practical reasoning*. MIT Press, Cambridge, Mass., 2001.

- P. Manzini and M. Mariotti. On the Representation of Incomplete Preferences Over Risky Alternatives. *Theory and Decision*, 65(4):303–323, Dec. 2008. ISSN 0040-5833, 1573-7187. doi:10.1007/s11238-007-9086-7.
- E. McClennen. *Rationality and dynamic choice: Foundational explorations*. Cambridge University Press, Cambridge, 1990.
- E. A. Ok. Utility Representation of an Incomplete Preference Relation. *Journal of Economic Theory*, 104(2):429–449, June 2002. ISSN 0022-0531. doi:10.1006/jeth.2001.2814.
- J. Quiggin. *Generalized expected utility theory: The rank-dependent model*. Springer Science & Business Media, 2012.
- B. Roy. Decision science or decision-aid science? *European Journal of Operational Research*, 66:184–203, 1993. doi:10.1016/0377-2217(93)90312-B.
- B. Roy. *Multicriteria Methodology for Decision Aiding*. Kluwer Academic, Dordrecht, 1996.
- L. J. Savage. *The foundations of statistics*. Dover Publications, New York, second revised edition, 1972. ISBN 978-0-486-62349-8.
- G. Shafer and V. Vovk. *Probability and finance: it's only a game!*, volume 491. John Wiley & Sons, 2005.
- P. Slovic and S. Lichtenstein. Preference reversals: A broader perspective. *American Economic Review*, 73:596–605, 1983.
- P. Slovic and A. Tversky. Who accepts Savage's axiom? *Behavioral Science*, 19:368–373, 1974. doi:10.1002/bs.3830190603.
- R. Thaler. Toward a positive theory of consumer choice. *Journal of Economic Behavior and Organization*, 1:39–60, 1980.
- A. Tsoukiàs. On the concept of decision aiding process: an operational perspective. *Annals of Operations Research*, 154(1):3–27, Oct. 2007. ISSN 0254-5330. doi:10.1007/s10479-007-0187-z.
- A. Tversky. Intransitivity of preferences. *Psychological Review*, 76:31–48, 1969.
- A. Tversky and D. Kahneman. The framing of decisions and the psychology of choice. *Science*, 211:453–458, 1981.
- A. Tversky, S. Sattath, and P. Slovic. Contingent weighting in judgment and choice. *Psychological Review*, 95(3):371–384, 1988. ISSN 0033-295X. doi:10.1037/0033-295X.95.3.371.
- A. Tversky, P. Slovic, and D. Kahneman. The causes of preference reversals. *American Economic Review*, 80(1):204–217, Mar. 1990.
- J. von Neumann and O. Morgenstern. *Theory of games and economic behavior*. Princeton classic editions. Princeton University Press, Princeton, N.J. ; Woodstock, 60th anniversary edition, 2004. ISBN 978-0-691-13061-3.
- P. P. Wakker. *Prospect Theory: For Risk and Ambiguity*. Cambridge University Press, July 2010. ISBN 978-1-139-48910-2.
- A. Wald. Statistical decision functions. In *Breakthroughs in Statistics*, pages 342–357. Springer, 1992.
- P. Walley. *Statistical Reasoning with Imprecise Probabilities*. Chapman & Hall/CRC Monographs on Statistics & Applied Probability. Taylor & Francis, 1991. ISBN 9780412286605. URL <https://books.google.fr/books?id=-hbvAAAAMAAJ>.