

		$\mathcal{B}_1$	$\mathcal{B}_2$	$\mathcal{B}_3$	$\mathcal{B}_4$
		1 2 5 9 3 5 4 0	1 1 1 1 3 9 0 8	1 1 1 1 1 1 7 6 2 4 2 1 6 8	2 1 7
$\mathcal{B}_1$	5	-00001	1111	111011111	11
	9	0-0000	0011	111111111	11
	3	01-011	0011	111111101	11
	15	111-11	1111	111111111	11
	4	0000-1	0011	111101111	10
	20	00010-	0001	111111111	10
$\mathcal{B}_2$	13	110000	-001	000011100	00
	19	101101	0-11	100011100	01
	10	101111	11-1	110001111	00
	18	111111	111-	100011111	11
$\mathcal{B}_3$	11	000000	0000	-00001100	01
	17	000010	0000	0-0001100	11
	6	000000	0000	00-000000	10
	12	000000	0000	000-00000	11
	14	000000	0001	0000-1000	11
	2	000000	0000	00100-000	11
	1	000010	0001	000001-11	10
	16	000000	0011	0000011-0	00
$\mathcal{B}_4$	8	000010	0011	10100100-	11
	21	001011	0001	0111111001	- 1
	7	000000	0001	1111111000	1 -

Fig. 9.10. Advice sociomatrix for Krackhardt's high-tech managers permuted according to positions from hierarchical clustering of correlations

Density Tables. A useful way to summarize the ties between positions is in a *density table*, or *density matrix*. A density table is a matrix that has positions rather than individual actors as its rows and columns, and the values in the matrix are the proportion of ties that present from the actors in the row position to the actors in column position. That is, the entries in the density table are Δ 's. For the densities in positions, diagonal elements (x_{ii}) are excluded from calculations if ties are undefined.

We will continue to use the example of the positions identified by hierarchical clustering of the correlations on the advice relation for Krackhardt's high-tech managers. Consider the ties from members of $\mathcal{B}_1$ to members of $\mathcal{B}_2$. Since there are six people in $\mathcal{B}_1$, and four people in $\mathcal{B}_2$, there are $6 \times 4 = 24$ possible ties that could be present from members