


Complex numbers

MOPSS



Suggested readings

- Evan Chen's advice [On reading solutions](https://blog.evanchen.cc/2017/03/06/on-reading-solutions/), available at <https://blog.evanchen.cc/2017/03/06/on-reading-solutions/>.
- Evan Chen's [Advice for writing proofs/Remarks on English](https://web.evanchen.cc/handouts/english/english.pdf), available at <https://web.evanchen.cc/handouts/english/english.pdf>.
- [Notes on proofs](#) by Evan Chen from [OTIS Excerpts](#) [[Che25](#), Chapter 1].
- [Tips for writing up solutions](https://www.math.utoronto.ca/barbeau/writingup.pdf) by Edward Barbeau, available at <https://www.math.utoronto.ca/barbeau/writingup.pdf>.
- Evan Chen discusses why [math olympiads](#) are a valuable experience for [high schoolers](#) in the post on [Lessons from math olympiads](#), available at <https://blog.evanchen.cc/2018/01/05/lessons-from-math-olympiads/>.

List of problems and examples

1.1	Exercise (Moldova National Olympiad 2001 Grade 11 Day 2 P6, AoPS)	2
-----	---	---

§1 Complex numbers

Exercise 1.1 (Moldova National Olympiad 2001 Grade 11 Day 2 P6, AoPS).

For a positive integer n , denote $A_n = \{(x, y) \in \mathbb{Z}^2 \mid x^2 + xy + y^2 = n\}$.

- Prove that the set A_n is finite.
- Prove that the number of elements of A_n is divisible by 6 for all $n \geq 1$.
- For which n is the number of elements of A_n divisible by 12?

Remark. The following walkthrough may be skipped if one does not have an exposure to the notion of **groups**.

Walkthrough —

(a) Note that $x^2 + xy + y^2 \geq (x + \frac{y}{2})^2 + \frac{3}{4}y^2$ holds.

(b) For part (b), consider the transformations

$$(x, y) \mapsto (y, -x - y), \quad (x, y) \mapsto (y, x).$$

Show that these transformations^a map A_n to itself and generate a group of order 6 acting on A_n without fixed points.

(c) For part (c), consider the transformations

$$(x, y) \mapsto (x, y), \quad (x, y) \mapsto (-x, -y).$$

Show that these transformations map A_n to itself, and provides an action of the group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ on A_n . Determine the fixed points of this action.

^aNote that

$$(x - y\omega)\omega = x\omega - y\omega^2 = y - (x + y)\omega$$

holds.

Solution 1. Note that any $(x, y) \in A_n$ satisfies

$$n = x^2 + xy + y^2 = \frac{1}{2}((x + y)^2 + x^2 + y^2),$$

which implies that $|x|, |y| \leq \sqrt{2n}$. Thus, there are only finitely many pairs (x, y) of integers satisfying $x^2 + xy + y^2 = n$. This proves part (a).

To establish part (b), assume that n is a positive integer. For any $(x, y) \in A_n$, consider the pairs

$$(x, y), (-x - y, x), (y, -x - y),$$

and note that they all belong to A_n . Indeed,

$$\begin{aligned} (-x - y)^2 + (-x - y)x + x^2 &= x^2 + xy + y^2, \\ y^2 + y(-x - y) + (-x - y)^2 &= x^2 + xy + y^2. \end{aligned}$$

Further, if any two of these three pairs were equal, then we would have $x = y = 0$, which is impossible since $n \geq 1$. Thus, the three pairs above are distinct. Hence, the elements of A_n can be grouped into sets of three, proving that $|A_n|$ is divisible by 3. Similarly, for any $(x, y) \in A_n$, the pair (y, x) also belongs to A_n , and is distinct from (x, y) unless $n/3$ is a perfect square. Moreover, for any $(x, y) \in A_n$, the pair $(-y, -x)$ also belongs to A_n , and is distinct from (x, y) unless n is a perfect square. Thus, the elements of A_n can be grouped into sets of two, proving that $|A_n|$ is even. Combining these results, we conclude that $|A_n|$ is divisible by 6.

Finally, to determine when $|A_n|$ is divisible by 12, it suffices to determine when $|A_n|$ is divisible by 4. For any $(x, y) \in A_n$, consider the pairs

$$(x, y), (y, x), (-x, -y), (-y, -x),$$

and note that they all belong to A_n . Moreover, if any two of these four pairs were equal, then we would have either $x = y$ with $n = 3x^2$, or $x = -y$ with $n = x^2$. Therefore, if $n/3$ is not a perfect square and n is not a perfect square, then the four pairs above are distinct, and consequently $|A_n|$ is divisible by 4. Moreover, if $n = 3k^2$ holds for some integer k , then the pairs $(k, k), (-k, -k)$ belongs to A_n , and the remaining elements of A_n can be grouped into sets of four, showing that $|A_n| \equiv 2 \pmod{4}$. Similarly, if $n = k^2$ holds for some integer k , then the pairs $(k, -k), (-k, k)$ belongs to A_n , and the remaining elements of A_n can be grouped into sets of four, showing that $|A_n| \equiv 2 \pmod{4}$. Thus, $|A_n|$ is divisible by 4 if and only if $n/3$ is not a perfect square and n is not a perfect square. ■

Exercise 1.2 (Putnam 2018 B2, AoPS). Let n be a positive integer, and let $f_n(z) = n + (n-1)z + (n-2)z^2 + \cdots + 2z^{n-2} + z^{n-1}$. Show that all the roots of $f_n(z)$ lie outside the closed unit disk $\{z \in \mathbb{C} : |z| \leq 1\}$.

Walkthrough —

(a)

Solution 2. Let us assume that there exists a root z of f_n such that $|z| \leq 1$. Note that

$$(1 - z)f_n(z) = n - z - z^2 - \cdots - z^{n-1} - z^n.$$

Using the triangle inequality, we have

$$\begin{aligned}
 0 &= |(1-z)f_n(z)| &= |n - z - z^2 - \dots - z^{n-1} - z^n| \\
 &\geq n - |z + z^2 + \dots + z^{n-1} + z^n| \\
 &\geq n - |z| - |z^2| - \dots - |z^{n-1}| - |z^n| \\
 &> n - 1 - 1 - \dots - 1 - 1 \\
 &= 0.
 \end{aligned}$$

This shows that $z, z^2, \dots, z^n$ have absolute value 1, and have the same argument, and hence $z = 1$. However, $f_n(1) = n + (n-1) + (n-2) + \dots + 2 + 1$, which is nonzero. This is a contradiction. Hence, all the roots of f_n lie outside the closed unit disk. ■

Solution 3. Let $g_n(z)$ denote the polynomial $z^n + z^{n-1} + \dots + z + 1$. Note that

$$g'_n(z) = nz^{n-1} + (n-1)z^{n-2} + \dots + 2z + 1 = z^{n-1}f_n\left(\frac{1}{z}\right).$$

Observe that the roots of the polynomials g'_n, f_n are nonzero, and a nonzero complex number z is a root of f_n if and only if $\frac{1}{z}$ is a root of g'_n . By the Gauss–Lucas theorem (Theorem ??), all the roots of g'_n lie in the convex hull of the set of roots of g_n . Note that the roots of g_n are precisely the n -th roots of unity other than 1. Since g_n has no multiple roots, it follows that g'_n does not vanish at any of the roots of g_n . Thus, all the roots of g'_n lie in the interior of the convex hull of the set of roots of g_n , and consequently, the roots of g'_n lie in the open unit disk. Therefore, all the roots of f_n lie outside the closed unit disk. ■

References

- [Che25] EVAN CHEN. *The OTIS Excerpts*. Available at <https://web.evanchen.cc/excerpts.html>. 2025, pp. vi+289 (cited p. 1)