

# Metastability for the Contact Process on $\mathbb{Z}$ , Part 1.

Following [Sch85]

O. Lynch<sup>1</sup>

<sup>1</sup>Department of Mathematics  
Universiteit Utrecht

December 6, 2020

# Outline

1. Introduction
  - 1.1 Metastability
  - 1.2 Contact Process
2. Crash Course in Contact Process
  - 2.1 Toolbox
  - 2.2 Review
3. Theorem 1
  - 3.1 Overview
  - 3.2 An Excerpt from Schonman Op. 1
  - 3.3 Conclusion

# Introduction

## 1. Introduction

### 1.1 Metastability

### 1.2 Contact Process

## 2. Crash Course in Contact Process

### 2.1 Toolbox

### 2.2 Review

## 3. Theorem 1

### 3.1 Overview

### 3.2 An Excerpt from Schonman Op. 1

### 3.3 Conclusion

# Informal Metastability

- ▶ Markov process  $X_N(t)$

# Informal Metastability

- ▶ Markov process  $X_N(t)$
- ▶ As  $N$  goes to  $\infty$ , looks like figure 1.




Figure: Coarse Graining as  $N \rightarrow \infty$

# Formal Metastability

## 1. Exponential hitting time

- ▶ There is a “trap state”, with hitting time  $T_N$
- ▶ Asymptotically as  $N \rightarrow \infty$ ,  $T_N$  has an exponential distribution.

# Formal Metastability

1. Exponential hitting time
  - ▶ There is a “trap state”, with hitting time  $T_N$
  - ▶ Asymptotically as  $N \rightarrow \infty$ ,  $T_N$  has an exponential distribution.
2. Quasi-stationary distribution before hitting time
  - ▶ There is a “approximate invariant distribution”  $\mu$
  - ▶ Up until  $T_N$ , temporal means of  $X_N(t)$  approximate  $\mu$

# Contact Process, Generator Definition

- ▶  $\xi(t)$  is a Markov process taking values in  $2^{\mathbb{Z}} = \mathcal{P}(\mathbb{Z})$
- ▶ Characterized by

$$Lf(\eta) = \sum_x c(x, \eta)(f(\eta^x) - f(\eta)) \quad (1)$$

- ▶ With rates

$$c(x, \eta) = \begin{cases} 1 & \text{if } \eta(x) = 1 \\ \lambda(\eta(x-1) + \eta(x+1)) & \text{otherwise} \end{cases} \quad (2)$$



# Contact Process, Percolation Structure Definition

▶ (show animation 1)

# Contact Process, Percolation Structure Definition

- ▶ (show animation 1)
- ▶ 3 processes for each site  $x \in \mathbb{Z}$ :  $P_x$  (rate 1),  $P_{x \rightarrow x+1}$  and  $P_{x \rightarrow x-1}$  (both rate  $\lambda$ ).

# Contact Process, Percolation Structure Definition

- ▶ (show animation 1)
- ▶ 3 processes for each site  $x \in \mathbb{Z}$ :  $P_x$  (rate 1),  $P_{x \rightarrow x+1}$  and  $P_{x \rightarrow x-1}$  (both rate  $\lambda$ ).
- ▶  $\xi^A(t)$  is the set of  $x \in \mathbb{Z}$  such that there is a path  $(y, 0) \rightarrow (x, t)$  for  $y \in A$ .

# Contact Process, Percolation Structure Definition

- ▶ (show animation 1)
- ▶ 3 processes for each site  $x \in \mathbb{Z}$ :  $P_x$  (rate 1),  $P_{x \rightarrow x+1}$  and  $P_{x \rightarrow x-1}$  (both rate  $\lambda$ ).
- ▶  $\xi^A(t)$  is the set of  $x \in \mathbb{Z}$  such that there is a path  $(y, 0) \rightarrow (x, t)$  for  $y \in A$ .
- ▶ A path is...

# Contact Process, Percolation Structure Definition

- ▶ (show animation 1)
- ▶ 3 processes for each site  $x \in \mathbb{Z}$ :  $P_x$  (rate 1),  $P_{x \rightarrow x+1}$  and  $P_{x \rightarrow x-1}$  (both rate  $\lambda$ ).
- ▶  $\xi^A(t)$  is the set of  $x \in \mathbb{Z}$  such that there is a path  $(y, 0) \rightarrow (x, t)$  for  $y \in A$ .
- ▶ A path is... easier to see intuitively in than write down formally

# Contact Process, Percolation Structure Definition

- ▶ (show animation 1)
- ▶ 3 processes for each site  $x \in \mathbb{Z}$ :  $P_x$  (rate 1),  $P_{x \rightarrow x+1}$  and  $P_{x \rightarrow x-1}$  (both rate  $\lambda$ ).
- ▶  $\xi^A(t)$  is the set of  $x \in \mathbb{Z}$  such that there is a path  $(y, 0) \rightarrow (x, t)$  for  $y \in A$ .
- ▶ A path is... easier to see intuitively in than write down formally
- ▶ Percolation structure is very useful for proofs

# Smaller Processes

- ▶  $\xi_B^A(t)$  is the set of  $x \in B$  such that there is a path  $(y, 0) \rightarrow (x, t)$  with  $y \in A$  and the path not leaving  $B$

# Smaller Processes

- ▶  $\xi_B^A(t)$  is the set of  $x \in B$  such that there is a path  $(y, 0) \rightarrow (x, t)$  with  $y \in A$  and the path not leaving  $B$
- ▶ Special cases
  - ▶  $\xi_N(t) = \xi_{[-N, N]}(t)$
  - ▶  $\xi_{[-N, \infty)}(t)$
  - ▶  $\xi_{(-\infty, N]}(t)$



# Crash Course in Contact Process

## 1. Introduction

### 1.1 Metastability

### 1.2 Contact Process

## 2. Crash Course in Contact Process

### 2.1 Toolbox

### 2.2 Review

## 3. Theorem 1

### 3.1 Overview

### 3.2 An Excerpt from Schonman Op. 1

### 3.3 Conclusion

# Critical $\lambda$

## Proposition

*There exists  $\lambda_c$  such that for  $\lambda < \lambda_c$ ,  $\xi(t)$  has only one invariant measure, which is concentrated at  $\emptyset$ . For  $\lambda > \lambda_c$ , there is also another extremal invariant measure, which we call  $\mu$ , which is obtained by time-averaging  $\xi(t)$ . This also holds for  $\xi_{[-N, \infty)}$  and  $\xi_{(-\infty, N]}$*

# Monotone Convergence

- ▶  $\mathbb{P}(\xi^A(t) \cap B \neq \emptyset)$  is a nonincreasing function of  $t$ .

# Monotone Convergence

- ▶  $\mathbb{P}(\xi^A(t) \cap B \neq \emptyset)$  is a nonincreasing function of  $t$ .
- ▶ Converges to  $\mu_B(\eta \mid \eta \cap A \neq \emptyset)$ , where  $\mu_B$  is the invariant measure of  $\xi$  started in state  $B$ .

# Self-duality

As long as  $A$  or  $B$  is finite, then for all  $t$ .

$$\mathbb{P}(\xi^A(t) \cap B \neq \emptyset) = \mathbb{P}(\xi^B(t) \cap A \neq \emptyset) \quad (3)$$

# Self-duality

As long as  $A$  or  $B$  is finite, then for all  $t$ .

$$\mathbb{P}(\xi^A(t) \cap B \neq \emptyset) = \mathbb{P}(\xi^B(t) \cap A \neq \emptyset) \quad (3)$$

“The probability of reaching somewhere in  $B$  starting from  $A$  is the same as the probability of reaching somewhere in  $A$  starting from  $B$ ”

# Self-duality

As long as  $A$  or  $B$  is finite, then for all  $t$ .

$$\mathbb{P}(\xi^A(t) \cap B \neq \emptyset) = \mathbb{P}(\xi^B(t) \cap A \neq \emptyset) \quad (3)$$

“The probability of reaching somewhere in  $B$  starting from  $A$  is the same as the probability of reaching somewhere in  $A$  starting from  $B$ ”

Specifically, for  $A$  finite

$$\mathbb{P}(\xi^A(t) \neq \emptyset) = \mathbb{P}(\xi(t) \cap A \neq \emptyset) \quad (4)$$

# Consequence of Monotone Convergence + Self-duality

$$\begin{aligned}\mathbb{P}(\xi^A(t) \neq \emptyset, \forall t > 0) &= \lim_{t \rightarrow \infty} \mathbb{P}(\xi^A(t) \neq \emptyset) \\ &= \lim_{t \rightarrow \infty} \mathbb{P}(\xi(t) \cap A \neq \emptyset) \\ &= \mu(\eta \mid \eta \cap A \neq \emptyset)\end{aligned}$$



# Characterizing $\mu$

- ▶  $\mu$  is translation invariant

# Characterizing $\mu$

- ▶  $\mu$  is translation invariant
- ▶ For  $\lambda > \lambda_c$ , given  $\varepsilon$ , we can find  $n(\varepsilon)$  such that  $\mu(\eta|_{\eta \cap [1, n(\varepsilon)]} \neq \emptyset) > 1 - \varepsilon$

# Characterizing $\mu$

- ▶  $\mu$  is translation invariant
- ▶ For  $\lambda > \lambda_c$ , given  $\varepsilon$ , we can find  $n(\varepsilon)$  such that  $\mu(\eta|_{\eta \cap [1, n(\varepsilon)]} \neq \emptyset) > 1 - \varepsilon$
- ▶ Therefore, we can find  $n(\varepsilon)$  so that if you start out in  $[1, n(\varepsilon)]$ , you are almost sure to survive.

# Characterizing $\mu$

- ▶  $\mu$  is translation invariant
- ▶ For  $\lambda > \lambda_c$ , given  $\varepsilon$ , we can find  $n(\varepsilon)$  such that  $\mu(\eta | \eta \cap [1, n(\varepsilon)] \neq \emptyset) > 1 - \varepsilon$
- ▶ Therefore, we can find  $n(\varepsilon)$  so that if you start out in  $[1, n(\varepsilon)]$ , you are almost sure to survive.
- ▶ It turns out that you can generalize this: if you start out *with at least*  $n(\varepsilon)$  elements, you are almost sure to survive.

What have we learned so far?

- ▶ An intuition for metastability — two key properties

What have we learned so far?

- ▶ An intuition for metastability — two key properties
- ▶ How to use the percolation structure to construct many related contact processes

What have we learned so far?

- ▶ An intuition for metastability — two key properties
- ▶ How to use the percolation structure to construct many related contact processes
- ▶ Some miscellaneous useful facts about the contact process

What have we learned so far?

- ▶ An intuition for metastability — two key properties
- ▶ How to use the percolation structure to construct many related contact processes
- ▶ Some miscellaneous useful facts about the contact process
  - ▶ Critical  $\lambda$



What have we learned so far?

- ▶ An intuition for metastability — two key properties
- ▶ How to use the percolation structure to construct many related contact processes
- ▶ Some miscellaneous useful facts about the contact process
  - ▶ Critical  $\lambda$
  - ▶ Monotone convergence

What have we learned so far?

- ▶ An intuition for metastability — two key properties
- ▶ How to use the percolation structure to construct many related contact processes
- ▶ Some miscellaneous useful facts about the contact process
  - ▶ Critical  $\lambda$
  - ▶ Monotone convergence
  - ▶ Self-duality

What have we learned so far?

- ▶ An intuition for metastability — two key properties
- ▶ How to use the percolation structure to construct many related contact processes
- ▶ Some miscellaneous useful facts about the contact process
  - ▶ Critical  $\lambda$
  - ▶ Monotone convergence
  - ▶ Self-duality
  - ▶ How to stay alive in the cruel, hard world of the contact process: be big!

# Theorem 1

## 1. Introduction

### 1.1 Metastability

### 1.2 Contact Process

## 2. Crash Course in Contact Process

### 2.1 Toolbox

### 2.2 Review

## 3. Theorem 1

### 3.1 Overview

### 3.2 An Excerpt from Schonman Op. 1

### 3.3 Conclusion

# Goals

## Theorem

If  $T_N$  is hitting time of  $\emptyset$ , then

$$\frac{T_N}{\mathbb{E} T_N} \xrightarrow{w} \text{Exp}(1)$$

That is, it converges to  $\text{Exp}(1)$  in distribution.

# Goals

## Theorem

If  $T_N$  is hitting time of  $\emptyset$ , then

$$\frac{T_N}{\mathbb{E} T_N} \xrightarrow{w} \text{Exp}(1)$$

That is, it converges to  $\text{Exp}(1)$  in distribution.

- ▶ Not feasible to give whole proof

# Goals

## Theorem

If  $T_N$  is hitting time of  $\emptyset$ , then

$$\frac{T_N}{\mathbb{E} T_N} \xrightarrow{w} \text{Exp}(1)$$

That is, it converges to  $\text{Exp}(1)$  in distribution.

- ▶ Not feasible to give whole proof
- ▶ Instead, will give strategy and then give detailed proof of just one part

# Strategy Part 1

- ▶ Use  $\beta_N$  instead of  $\mathbb{E} T_N$ , where  $\beta_N$  is unique number such that

$$\mathbb{P}(T_N > \beta_N) = e^{-1}$$



# Strategy Part 1

- ▶ Use  $\beta_N$  instead of  $\mathbb{E} T_N$ , where  $\beta_N$  is unique number such that

$$\mathbb{P}(T_N > \beta_N) = e^{-1}$$

- ▶ Let  $G_N(t) = \mathbb{P}\left[\frac{T_N}{\beta_N} > t\right]$  be CDF for  $T_N/\beta_N$

# Strategy Part 1

- ▶ Use  $\beta_N$  instead of  $\mathbb{E} T_N$ , where  $\beta_N$  is unique number such that

$$\mathbb{P}(T_N > \beta_N) = e^{-1}$$

- ▶ Let  $G_N(t) = \mathbb{P}\left[\frac{T_N}{\beta_N} > t\right]$  be CDF for  $T_N/\beta_N$
- ▶ (note that  $\{\frac{T_N}{\beta_N} > t\} = \{\xi_N(\beta_N t) \neq \emptyset\}$ )

# Strategy Part 1

- ▶ Use  $\beta_N$  instead of  $\mathbb{E} T_N$ , where  $\beta_N$  is unique number such that

$$\mathbb{P}(T_N > \beta_N) = e^{-1}$$

- ▶ Let  $G_N(t) = \mathbb{P}\left[\frac{T_N}{\beta_N} > t\right]$  be CDF for  $T_N/\beta_N$
- ▶ (note that  $\{\frac{T_N}{\beta_N} > t\} = \{\xi_N(\beta_N t) \neq \emptyset\}$ )
- ▶ Prove that

$$\lim_{N \rightarrow \infty} |G_N(t)G_N(s) - G_N(t+s)| = 0$$

- ▶ The only function with  $f(t+s) = f(t)f(s)$  and  $\int_0^\infty f(t) dt = 1$  is  $f(t) = e^{-t}$ .

## Strategy Part 2

- ▶ Use a set  $F_b \subseteq \mathcal{P}([-N, N])$  of “sufficiently dense” initial conditions

## Strategy Part 2

- ▶ Use a set  $F_b \subseteq \mathcal{P}([-N, N])$  of “sufficiently dense” initial conditions
- ▶ If we are not in  $F_b$ , then we are most likely already dead. Therefore, the probability that we end up in  $F_b$  at time  $\beta_N t$  is approximately  $G_N(t)$

## Strategy Part 2

- ▶ Use a set  $F_b \subseteq \mathcal{P}([-N, N])$  of “sufficiently dense” initial conditions
- ▶ If we are not in  $F_b$ , then we are most likely already dead. Therefore, the probability that we end up in  $F_b$  at time  $\beta_N t$  is approximately  $G_N(t)$
- ▶ Starting from  $A \in F_b$  is “just as good” as starting from  $[-N, N]$ , so if we end up in  $A \in F_b$  at time  $\beta_N t$ , then we have approximately  $G_N(s)$  chance of still being alive at time  $\beta_N(s + t)$ .

## Strategy Part 2

- ▶ Use a set  $F_b \subseteq \mathcal{P}([-N, N])$  of “sufficiently dense” initial conditions
- ▶ If we are not in  $F_b$ , then we are most likely already dead. Therefore, the probability that we end up in  $F_b$  at time  $\beta_N t$  is approximately  $G_N(t)$
- ▶ Starting from  $A \in F_b$  is “just as good” as starting from  $[-N, N]$ , so if we end up in  $A \in F_b$  at time  $\beta_N t$ , then we have approximately  $G_N(s)$  chance of still being alive at time  $\beta_N(s + t)$ .
- ▶ Put together, these two imply that  $G_N(t)G_N(s) \sim G_N(t + s)$

# Into the Unknown

- ▶ Show that  $\mathbb{P}[T_N = T_N^A] > 1 - \varepsilon$  when  $A \in F_b$ , for sufficiently large  $N$  and  $b$ . (i.e.,  $A \in F_b$  is “just as good” as  $[-N, N]$ )

- ▶ Let

$$F_b = \{A \in \mathbb{Z} \mid \frac{|A \cap [-b, -1]|}{b} \geq \frac{\rho}{2}, \frac{|A \cap [1, b]|}{b} \geq \frac{\rho}{2}\}$$

- ▶ “sufficiently dense on both sides”
- ▶  $\rho = \mathbb{P}[\xi^{\{0\}}(t) \neq \emptyset, \forall t] = \mu(\eta \mid \eta(0) = 1).$



- ▶ Let  $n = n(\varepsilon)$  large enough so that starting out with  $n$  elements will let you survive forever with probability  $1 - \varepsilon/2$

- ▶ Let  $n = n(\varepsilon)$  large enough so that starting out with  $n$  elements will let you survive forever with probability  $1 - \varepsilon/2$
- ▶ Then if  $b\rho/2 > n(\varepsilon)$ , for  $A$  in  $F_b$  starting in  $[1, b] \cap A$  or  $[-b, -1] \cap A$  will let you survive forever with probability  $1 - \varepsilon/2$

- ▶ Let  $n = n(\varepsilon)$  large enough so that starting out with  $n$  elements will let you survive forever with probability  $1 - \varepsilon/2$
- ▶ Then if  $b\rho/2 > n(\varepsilon)$ , for  $A$  in  $F_b$  starting in  $[1, b] \cap A$  or  $[-b, -1] \cap A$  will let you survive forever with probability  $1 - \varepsilon/2$
- ▶ Let  $E$  be the event that  $\xi_{[-N, \infty)}^{A \cap [-b, -1]}(t)$  and  $\xi_{(-\infty, N]}^{A \cap [1, b]}(t)$  survive forever (the invariant distribution for  $[-N, \infty)$  is the same as that for  $\mathbb{Z}$ )

- ▶ Let  $n = n(\varepsilon)$  large enough so that starting out with  $n$  elements will let you survive forever with probability  $1 - \varepsilon/2$
- ▶ Then if  $b\rho/2 > n(\varepsilon)$ , for  $A$  in  $F_b$  starting in  $[1, b] \cap A$  or  $[-b, -1] \cap A$  will let you survive forever with probability  $1 - \varepsilon/2$
- ▶ Let  $E$  be the event that  $\xi_{[-N, \infty)}^{A \cap [-b, -1]}(t)$  and  $\xi_{(-\infty, N]}^{A \cap [1, b]}(t)$  survive forever (the invariant distribution for  $[-N, \infty)$  is the same as that for  $\mathbb{Z}$ )
- ▶  $\mathbb{P}(E) > 1 - \varepsilon$

- ▶ Let  $n = n(\varepsilon)$  large enough so that starting out with  $n$  elements will let you survive forever with probability  $1 - \varepsilon/2$
- ▶ Then if  $b\rho/2 > n(\varepsilon)$ , for  $A$  in  $F_b$  starting in  $[1, b] \cap A$  or  $[-b, -1] \cap A$  will let you survive forever with probability  $1 - \varepsilon/2$
- ▶ Let  $E$  be the event that  $\xi_{[-N, \infty)}^{A \cap [-b, -1]}(t)$  and  $\xi_{(-\infty, N]}^{A \cap [1, b]}(t)$  survive forever (the invariant distribution for  $[-N, \infty)$  is the same as that for  $\mathbb{Z}$ )
- ▶  $\mathbb{P}(E) > 1 - \varepsilon$
- ▶ Therefore, we are done if in  $E$ ,  $T_N = T_N^A$ .

- Define two stopping times


$$U = \inf\{t \mid N \in \xi_{[-N, \infty)}^{A \cap [-b, -1]}(t)\}$$

$$V = \inf\{t \mid -N \in \xi_{(-\infty, N]}^{A \cap [1, b]}(t)\}$$

► Define two stopping times

$$U = \inf\{t \mid N \in \xi_{[-N, \infty)}^{A \cap [-b, -1]}(t)\}$$


$$V = \inf\{t \mid -N \in \xi_{(-\infty, N]}^{A \cap [1, b]}(t)\}$$



- ▶  $U$  and  $V$  are almost surely finite on  $E$ .
- ▶ At time  $U$ ,  $\xi_N^A$  is alive, because  $\xi_N^A(U) \supset \xi_{[-N, \infty)}^{A \cap [-b, -1]}(t)$ , and similarly for  $V$ , so  $T_N \geq T_N^A > \max(U, V)$



- After  $U$  and  $V$ , there is a path from  $[-b, -1] \cap A$  to  $N$ , and a path from  $[1, b] \cap A$  to  $-N$ . Intuitively, one of those paths intersects any path from  $x \in [-N, N]$  to  $y \in \xi_N(t)$ .



- ▶ For  $t > \max(U, V)$ ,  $\xi_N(t) = \xi_N^A(t)$
- ▶ Therefore,  $T_N = T_N^A$  on  $E$ , and we have shown that  $\mathbb{P}(T_N = T_N^A) > 1 - \varepsilon$ .

# Summary

We have learned

# Summary

We have learned

- ▶ What metastability is

# Summary

We have learned

- ▶ What metastability is
- ▶ Basic properties of the contact process

# Summary

We have learned

- ▶ What metastability is
- ▶ Basic properties of the contact process
- ▶ A rough sketch of how the contact process is metastable (part 1)

# Summary

We have learned

- ▶ What metastability is
- ▶ Basic properties of the contact process
- ▶ A rough sketch of how the contact process is metastable (part 1)
- ▶ What working with the contact process in practice looks like

# Citations

- [Sch85] Roberto H. Schonmann. “Metastability for the contact process.” In: *Journal of Statistical Physics* 41.3 (Nov. 1, 1985), pp. 445–464. ISSN: 1572-9613. DOI: 10.1007/BF01009017. URL: <https://doi.org/10.1007/BF01009017> (visited on 11/12/2020).