

Metastability for the Contact Process on \mathbb{Z} , Part 1.

Following [Sch85]

O. Lynch¹

¹Department of Mathematics
Universiteit Utrecht

December 8, 2020

Outline

1. Introduction
 - 1.1 Metastability
 - 1.2 Contact Process
2. Crash Course in Contact Process
 - 2.1 Toolbox
 - 2.2 Review
3. Theorem 1
 - 3.1 Overview
 - 3.2 An Excerpt from Schonman Op. 1
 - 3.3 Conclusion

Introduction

1. Introduction

1.1 Metastability

1.2 Contact Process

2. Crash Course in Contact Process

2.1 Toolbox

2.2 Review

3. Theorem 1

3.1 Overview

3.2 An Excerpt from Schonman Op. 1

3.3 Conclusion

Informal Metastability

- ▶ Markov process $\xi_N(t) \subseteq \mathbb{Z}$, “infected states at time t ”

Informal Metastability

- ▶ Markov process $\xi_N(t) \subseteq \mathbb{Z}$, “infected states at time t ”
- ▶ As N goes to ∞ , looks like figure 1.




Figure: Coarse Graining as $N \rightarrow \infty$

Formal Metastability

1. Exponential hitting time

- ▶ There is a “trap state”, with hitting time T_N
- ▶ Asymptotically as $N \rightarrow \infty$, T_N has an exponential distribution.

Formal Metastability

1. Exponential hitting time
 - ▶ There is a “trap state”, with hitting time T_N
 - ▶ Asymptotically as $N \rightarrow \infty$, T_N has an exponential distribution.
2. Quasi-stationary distribution before hitting time
 - ▶ There is a “approximate invariant distribution” μ
 - ▶ Up until T_N , temporal means of $X_N(t)$ approximate μ

Contact Process, Generator Definition

- ▶ $\xi(t)$ is a Markov process taking values in $2^{\mathbb{Z}} = \mathcal{P}(\mathbb{Z})$
- ▶ Characterized by

$$Lf(\eta) = \sum_x c(x, \eta)(f(\eta^x) - f(\eta)) \quad (1)$$

- ▶ With rates

$$c(x, \eta) = \begin{cases} 1 & \text{if } \eta(x) = 1 \\ \lambda(\eta(x-1) + \eta(x+1)) & \text{otherwise} \end{cases} \quad (2)$$

Contact Process, Percolation Structure Definition

▶ (show animation 1)

Contact Process, Percolation Structure Definition

- ▶ (show animation 1)
- ▶ 3 processes for each site $x \in \mathbb{Z}$: P_x (rate 1), $P_{x \rightarrow x+1}$ and $P_{x \rightarrow x-1}$ (both rate λ).

Contact Process, Percolation Structure Definition

- ▶ (show animation 1)
- ▶ 3 processes for each site $x \in \mathbb{Z}$: P_x (rate 1), $P_{x \rightarrow x+1}$ and $P_{x \rightarrow x-1}$ (both rate λ).
- ▶ $\xi^A(t)$ is the set of $x \in \mathbb{Z}$ such that there is a path $(y, 0) \rightarrow (x, t)$ for $y \in A$.

Contact Process, Percolation Structure Definition

- ▶ (show animation 1)
- ▶ 3 processes for each site $x \in \mathbb{Z}$: P_x (rate 1), $P_{x \rightarrow x+1}$ and $P_{x \rightarrow x-1}$ (both rate λ).
- ▶ $\xi^A(t)$ is the set of $x \in \mathbb{Z}$ such that there is a path $(y, 0) \rightarrow (x, t)$ for $y \in A$.
- ▶ A path is...

Contact Process, Percolation Structure Definition

- ▶ (show animation 1)
- ▶ 3 processes for each site $x \in \mathbb{Z}$: P_x (rate 1), $P_{x \rightarrow x+1}$ and $P_{x \rightarrow x-1}$ (both rate λ).
- ▶ $\xi^A(t)$ is the set of $x \in \mathbb{Z}$ such that there is a path $(y, 0) \rightarrow (x, t)$ for $y \in A$.
- ▶ A path is... easier to see intuitively in than write down formally

Contact Process, Percolation Structure Definition

- ▶ (show animation 1)
- ▶ 3 processes for each site $x \in \mathbb{Z}$: P_x (rate 1), $P_{x \rightarrow x+1}$ and $P_{x \rightarrow x-1}$ (both rate λ).
- ▶ $\xi^A(t)$ is the set of $x \in \mathbb{Z}$ such that there is a path $(y, 0) \rightarrow (x, t)$ for $y \in A$.
- ▶ A path is... easier to see intuitively in than write down formally
- ▶ Percolation structure is very useful for proofs

Smaller Processes

- ▶ $\xi_B^A(t)$ is the set of $x \in B$ such that there is a path $(y, 0) \rightarrow (x, t)$ with $y \in A$ and the path not leaving B

Smaller Processes

- ▶ $\xi_B^A(t)$ is the set of $x \in B$ such that there is a path $(y, 0) \rightarrow (x, t)$ with $y \in A$ and the path not leaving B
- ▶ Superscript: “starts in”, subscript: “contained in”

Smaller Processes

- ▶ $\xi_B^A(t)$ is the set of $x \in B$ such that there is a path $(y, 0) \rightarrow (x, t)$ with $y \in A$ and the path not leaving B
- ▶ Superscript: “starts in”, subscript: “contained in”
- ▶ If B is omitted, assumed to be \mathbb{Z} . If A is omitted, assumed to be B .

Smaller Processes

- ▶ $\xi_B^A(t)$ is the set of $x \in B$ such that there is a path $(y, 0) \rightarrow (x, t)$ with $y \in A$ and the path not leaving B
- ▶ Superscript: “starts in”, subscript: “contained in”
- ▶ If B is omitted, assumed to be \mathbb{Z} . If A is omitted, assumed to be B .
- ▶ Special cases
 - ▶ $\xi_N(t) = \xi_{[-N, N]}(t)$
 - ▶ $\xi_{[-N, \infty)}(t)$
 - ▶ $\xi_{(-\infty, N]}(t)$

Smaller Processes

- ▶ $\xi_B^A(t)$ is the set of $x \in B$ such that there is a path $(y, 0) \rightarrow (x, t)$ with $y \in A$ and the path not leaving B
- ▶ Superscript: “starts in”, subscript: “contained in”
- ▶ If B is omitted, assumed to be \mathbb{Z} . If A is omitted, assumed to be B .
- ▶ Special cases
 - ▶ $\xi_N(t) = \xi_{[-N, N]}(t)$
 - ▶ $\xi_{[-N, \infty)}(t)$
 - ▶ $\xi_{(-\infty, N]}(t)$
- ▶ We are interested in metastability for $\xi_N(t)$, with trap state \emptyset .

Crash Course in Contact Process

1. Introduction
 - 1.1 Metastability
 - 1.2 Contact Process
2. Crash Course in Contact Process
 - 2.1 Toolbox
 - 2.2 Review
3. Theorem 1
 - 3.1 Overview
 - 3.2 An Excerpt from Schonman Op. 1
 - 3.3 Conclusion

Critical λ

- Survival/extinction of $\xi(t)$ depends on infection rate λ

Critical λ

- ▶ Survival/extinction of $\xi(t)$ depends on infection rate λ
- ▶ For $\lambda < \lambda_c$, goes extinct in finite time a.s. (unique ergodic measure δ_\emptyset)

Critical λ

- ▶ Survival/extinction of $\xi(t)$ depends on infection rate λ
- ▶ For $\lambda < \lambda_c$, goes extinct in finite time a.s. (unique ergodic measure δ_\emptyset)
- ▶ For $\lambda > \lambda_c$, two extremal invariant measures: δ_\emptyset and μ

Critical λ

- ▶ Survival/extinction of $\xi(t)$ depends on infection rate λ
- ▶ For $\lambda < \lambda_c$, goes extinct in finite time a.s. (unique ergodic measure δ_\emptyset)
- ▶ For $\lambda > \lambda_c$, two extremal invariant measures: δ_\emptyset and μ
- ▶ This also holds for $\xi_{[-N, \infty)}(t)$ and $\xi_{(-\infty, N]}(t)$

Monotone Convergence

- ▶ $\mathbb{P}(\xi^A(t) \cap B \neq \emptyset)$ is a nonincreasing function of t .

Monotone Convergence

- ▶ $\mathbb{P}(\xi^A(t) \cap B \neq \emptyset)$ is a nonincreasing function of t .
- ▶ Converges to $\mu_A(\{\eta \mid \eta \cap B \neq \emptyset\})$, where μ_B is the invariant measure of ξ started in state B .

Self-duality

As long as A or B is finite, then for all t .

$$\mathbb{P}(\xi^A(t) \cap B \neq \emptyset) = \mathbb{P}(\xi^B(t) \cap A \neq \emptyset) \quad (3)$$

Self-duality

As long as A or B is finite, then for all t .

$$\mathbb{P}(\xi^A(t) \cap B \neq \emptyset) = \mathbb{P}(\xi^B(t) \cap A \neq \emptyset) \quad (3)$$

“The probability of reaching somewhere in B starting from A is the same as the probability of reaching somewhere in A starting from B ”

Self-duality

As long as A or B is finite, then for all t .

$$\mathbb{P}(\xi^A(t) \cap B \neq \emptyset) = \mathbb{P}(\xi^B(t) \cap A \neq \emptyset) \quad (3)$$

“The probability of reaching somewhere in B starting from A is the same as the probability of reaching somewhere in A starting from B ”

Specifically, for A finite

$$\mathbb{P}(\xi^A(t) \neq \emptyset) = \mathbb{P}(\xi(t) \cap A \neq \emptyset) \quad (4)$$

Consequence of Monotone Convergence + Self-duality

$$\begin{aligned}\mathbb{P}(\xi^A(t) \neq \emptyset, \forall t > 0) &= \lim_{s \rightarrow \infty} \mathbb{P}(\xi^A(t) \neq \emptyset, \forall s \geq t > 0) \\ &= \lim_{s \rightarrow \infty} \mathbb{P}(\xi^A(s) \neq \emptyset) \\ &= \lim_{s \rightarrow \infty} \mathbb{P}(\xi(s) \cap A \neq \emptyset) \\ &= \mu(\{\eta \mid \eta \cap A \neq \emptyset\})\end{aligned}$$

Characterizing μ

- ▶ μ is translation invariant

Characterizing μ

- ▶ μ is translation invariant
- ▶ For $\lambda > \lambda_c$, given ε , we can find $n(\varepsilon)$ such that $\mu(\eta|_{\eta \cap [1, n(\varepsilon)]} \neq \emptyset) > 1 - \varepsilon$

Characterizing μ

- ▶ μ is translation invariant
- ▶ For $\lambda > \lambda_c$, given ε , we can find $n(\varepsilon)$ such that $\mu(\eta|_{\eta \cap [1, n(\varepsilon)]} \neq \emptyset) > 1 - \varepsilon$
- ▶ Therefore, we can find $n(\varepsilon)$ so that if you start out in $[1, n(\varepsilon)]$, you are almost sure to survive.

Characterizing μ

- ▶ μ is translation invariant
- ▶ For $\lambda > \lambda_c$, given ε , we can find $n(\varepsilon)$ such that $\mu(\eta | \eta \cap [1, n(\varepsilon)] \neq \emptyset) > 1 - \varepsilon$
- ▶ Therefore, we can find $n(\varepsilon)$ so that if you start out in $[1, n(\varepsilon)]$, you are almost sure to survive.
- ▶ It turns out that you can generalize this: if you start out *with at least* $n(\varepsilon)$ elements, you are almost sure to survive.

What have we learned so far?

- ▶ An intuition for metastability — two key properties

What have we learned so far?

- ▶ An intuition for metastability — two key properties
- ▶ How to use the percolation structure to construct many related contact processes

What have we learned so far?

- ▶ An intuition for metastability — two key properties
- ▶ How to use the percolation structure to construct many related contact processes
- ▶ Some miscellaneous useful facts about the contact process

What have we learned so far?

- ▶ An intuition for metastability — two key properties
- ▶ How to use the percolation structure to construct many related contact processes
- ▶ Some miscellaneous useful facts about the contact process
 - ▶ Critical λ

What have we learned so far?

- ▶ An intuition for metastability — two key properties
- ▶ How to use the percolation structure to construct many related contact processes
- ▶ Some miscellaneous useful facts about the contact process
 - ▶ Critical λ
 - ▶ Monotone convergence

What have we learned so far?

- ▶ An intuition for metastability — two key properties
- ▶ How to use the percolation structure to construct many related contact processes
- ▶ Some miscellaneous useful facts about the contact process
 - ▶ Critical λ
 - ▶ Monotone convergence
 - ▶ Self-duality

What have we learned so far?

- ▶ An intuition for metastability — two key properties
- ▶ How to use the percolation structure to construct many related contact processes
- ▶ Some miscellaneous useful facts about the contact process
 - ▶ Critical λ
 - ▶ Monotone convergence
 - ▶ Self-duality
 - ▶ How an infection can persist forever with high probability: it has to start out with enough sites infected

Theorem 1

1. Introduction

1.1 Metastability

1.2 Contact Process

2. Crash Course in Contact Process

2.1 Toolbox

2.2 Review

3. Theorem 1

3.1 Overview

3.2 An Excerpt from Schonman Op. 1

3.3 Conclusion

Goals

Theorem

Let

$$T_N = \inf\{t \mid \xi_N(t) = \emptyset\}$$

Then for $\lambda > \lambda_c$

$$\frac{T_N}{\mathbb{E} T_N} \xrightarrow{w} \text{Exp}(1)$$

That is, it converges to $\text{Exp}(1)$ in distribution.

Goals

Theorem

Let

$$T_N = \inf\{t \mid \xi_N(t) = \emptyset\}$$

Then for $\lambda > \lambda_c$

$$\frac{T_N}{\mathbb{E} T_N} \xrightarrow{w} \text{Exp}(1)$$

That is, it converges to $\text{Exp}(1)$ in distribution.

- Not feasible to give whole proof

Goals

Theorem

Let

$$T_N = \inf\{t \mid \xi_N(t) = \emptyset\}$$

Then for $\lambda > \lambda_c$

$$\frac{T_N}{\mathbb{E} T_N} \xrightarrow{w} \text{Exp}(1)$$

That is, it converges to $\text{Exp}(1)$ in distribution.

- ▶ Not feasible to give whole proof
- ▶ Instead, will give strategy and then give detailed proof of just one part

Strategy Part 1

- ▶ Use β_N instead of $\mathbb{E} T_N$, where β_N is unique number such that

$$\mathbb{P}(T_N > \beta_N) = e^{-1}$$

Strategy Part 1

- ▶ Use β_N instead of $\mathbb{E} T_N$, where β_N is unique number such that

$$\mathbb{P}(T_N > \beta_N) = e^{-1}$$

- ▶ Let $G_N(t) = \mathbb{P}\left[\frac{T_N}{\beta_N} > t\right]$ be the tail distribution for T_N/β_N

Strategy Part 1

- ▶ Use β_N instead of $\mathbb{E} T_N$, where β_N is unique number such that

$$\mathbb{P}(T_N > \beta_N) = e^{-1}$$

- ▶ Let $G_N(t) = \mathbb{P}\left[\frac{T_N}{\beta_N} > t\right]$ be the tail distribution for T_N/β_N
- ▶ (note that $\{\frac{T_N}{\beta_N} > t\} = \{\xi_N(\beta_N t) \neq \emptyset\}$)

Strategy Part 1

- ▶ Use β_N instead of $\mathbb{E} T_N$, where β_N is unique number such that

$$\mathbb{P}(T_N > \beta_N) = e^{-1}$$

- ▶ Let $G_N(t) = \mathbb{P}\left[\frac{T_N}{\beta_N} > t\right]$ be the tail distribution for T_N/β_N
- ▶ (note that $\{\frac{T_N}{\beta_N} > t\} = \{\xi_N(\beta_N t) \neq \emptyset\}$)
- ▶ Prove that

$$\lim_{N \rightarrow \infty} |G_N(t)G_N(s) - G_N(t+s)| = 0$$

- ▶ The only function with $f(t+s) = f(t)f(s)$ and $\int_0^\infty f(t) dt = 1$ is $f(t) = e^{-t}$.

Strategy Part 2

- ▶ Use a set $F_b \subseteq \mathcal{P}([-N, N])$ of “sufficiently dense” initial conditions

Strategy Part 2

- ▶ Use a set $F_b \subseteq \mathcal{P}([-N, N])$ of “sufficiently dense” initial conditions
- ▶ If we are not in F_b , then we are most likely already extinct. Therefore, the probability that we end up in F_b at time $\beta_N t$ is approximately $G_N(t)$

Strategy Part 2

- ▶ Use a set $F_b \subseteq \mathcal{P}([-N, N])$ of “sufficiently dense” initial conditions
- ▶ If we are not in F_b , then we are most likely already extinct. Therefore, the probability that we end up in F_b at time $\beta_N t$ is approximately $G_N(t)$
- ▶ Starting from $A \in F_b$ is “just as good” as starting from $[-N, N]$, so if we end up in $A \in F_b$ at time $\beta_N t$, then we have approximately $G_N(s)$ chance of still being alive at time $\beta_N(s + t)$.

Strategy Part 2

- ▶ Use a set $F_b \subseteq \mathcal{P}([-N, N])$ of “sufficiently dense” initial conditions
- ▶ If we are not in F_b , then we are most likely already extinct. Therefore, the probability that we end up in F_b at time $\beta_N t$ is approximately $G_N(t)$
- ▶ Starting from $A \in F_b$ is “just as good” as starting from $[-N, N]$, so if we end up in $A \in F_b$ at time $\beta_N t$, then we have approximately $G_N(s)$ chance of still being alive at time $\beta_N(s + t)$.
- ▶ Put together, these two imply that $G_N(t)G_N(s) \sim G_N(t + s)$

Into the Unknown

- ▶ Show that $\mathbb{P}[T_N = T_N^A] > 1 - \varepsilon$ when $A \in F_b$, for sufficiently large N and b . (i.e., $A \in F_b$ is “just as good” as $[-N, N]$)

- ▶ Let

$$F_b = \{A \in \mathbb{Z} \mid \frac{|A \cap [-b, -1]|}{b} \geq \frac{\rho}{2}, \frac{|A \cap [1, b]|}{b} \geq \frac{\rho}{2}\}$$

- ▶ “sufficiently dense on both sides”
- ▶ $\rho = \mathbb{P}[\xi^{\{0\}}(t) \neq \emptyset, \forall t] = \mu(\eta \mid \eta(0) = 1).$

- ▶ Let $n = n(\varepsilon)$ large enough so that starting out with n infected sites will let the epidemic persist forever with probability $1 - \varepsilon/2$

- ▶ Let $n = n(\varepsilon)$ large enough so that starting out with n infected sites will let the epidemic persist forever with probability $1 - \varepsilon/2$
- ▶ Then if $b\rho/2 > n(\varepsilon)$, for A in F_b starting in $[1, b] \cap A$ or $[-b, -1] \cap A$ will persist forever with probability $1 - \varepsilon/2$

- ▶ Let $n = n(\varepsilon)$ large enough so that starting out with n infected sites will let the epidemic persist forever with probability $1 - \varepsilon/2$
- ▶ Then if $b\rho/2 > n(\varepsilon)$, for A in F_b starting in $[1, b] \cap A$ or $[-b, -1] \cap A$ will persist forever with probability $1 - \varepsilon/2$
- ▶ Let E be the event that $\xi_{[-N, \infty)}^{A \cap [-b, -1]}(t)$ and $\xi_{(-\infty, N]}^{A \cap [1, b]}(t)$ persist forever (the invariant distribution for $[-N, \infty)$ is the same as that for \mathbb{Z})

- ▶ Let $n = n(\varepsilon)$ large enough so that starting out with n infected sites will let the epidemic persist forever with probability $1 - \varepsilon/2$
- ▶ Then if $b\rho/2 > n(\varepsilon)$, for A in F_b starting in $[1, b] \cap A$ or $[-b, -1] \cap A$ will persist forever with probability $1 - \varepsilon/2$
- ▶ Let E be the event that $\xi_{[-N, \infty)}^{A \cap [-b, -1]}(t)$ and $\xi_{(-\infty, N]}^{A \cap [1, b]}(t)$ persist forever (the invariant distribution for $[-N, \infty)$ is the same as that for \mathbb{Z})
- ▶ $\mathbb{P}(E) > 1 - \varepsilon$

- ▶ Let $n = n(\varepsilon)$ large enough so that starting out with n infected sites will let the epidemic persist forever with probability $1 - \varepsilon/2$
- ▶ Then if $b\rho/2 > n(\varepsilon)$, for A in F_b starting in $[1, b] \cap A$ or $[-b, -1] \cap A$ will persist forever with probability $1 - \varepsilon/2$
- ▶ Let E be the event that $\xi_{[-N, \infty)}^{A \cap [-b, -1]}(t)$ and $\xi_{(-\infty, N]}^{A \cap [1, b]}(t)$ persist forever (the invariant distribution for $[-N, \infty)$ is the same as that for \mathbb{Z})
- ▶ $\mathbb{P}(E) > 1 - \varepsilon$
- ▶ Therefore, we are done if in E , $T_N = T_N^A$.

- Define two stopping times


$$U = \inf\{t \mid N \in \xi_{[-N, \infty)}^{A \cap [-b, -1]}(t)\}$$

$$V = \inf\{t \mid -N \in \xi_{(-\infty, N]}^{A \cap [1, b]}(t)\}$$

► Define two stopping times


$$U = \inf\{t \mid N \in \xi_{[-N, \infty)}^{A \cap [-b, -1]}(t)\}$$

$$V = \inf\{t \mid -N \in \xi_{(-\infty, N]}^{A \cap [1, b]}(t)\}$$



- ▶ U and V are almost surely finite on E .
- ▶ At time U , ξ_N^A is alive, because $\xi_N^A(U) \supset \xi_{[-N, \infty)}^{A \cap [-b, -1]}(t)$, and similarly for V , so $T_N \geq T_N^A > \max(U, V)$

- After U and V , there is a path from $[-b, -1] \cap A$ to N , and a path from $[1, b] \cap A$ to $-N$. Intuitively, one of those paths intersects any path from $x \in [-N, N]$ to $y \in \xi_N(t)$.



- ▶ For $t > \max(U, V)$, $\xi_N(t) = \xi_N^A(t)$
- ▶ Therefore, $T_N = T_N^A$ on E , and we have shown that $\mathbb{P}(T_N = T_N^A) > 1 - \varepsilon$.

Summary

We have learned

Summary

We have learned

- ▶ What metastability is

Summary

We have learned

- ▶ What metastability is
- ▶ Basic properties of the contact process

Summary

We have learned

- ▶ What metastability is
- ▶ Basic properties of the contact process
- ▶ A rough sketch of how the contact process is metastable (part 1)

Summary

We have learned

- ▶ What metastability is
- ▶ Basic properties of the contact process
- ▶ A rough sketch of how the contact process is metastable (part 1)
- ▶ What working with the contact process in practice looks like

Citations

- [Sch85] Roberto H. Schonmann. “Metastability for the contact process.” In: *Journal of Statistical Physics* 41.3 (Nov. 1, 1985), pp. 445–464. ISSN: 1572-9613. DOI: 10.1007/BF01009017. URL: <https://doi.org/10.1007/BF01009017> (visited on 11/12/2020).