

# Doob's Proof of MLE Consistency

Omar Hijab

Department of Mathematics, Temple University

2026

## Abstract

We present a concise exposition of Doob's proof of MLE consistency using surprisal and relative information. We clarify historical entropy terminology and explicitly verify all assumptions for the multivariate normal distribution using elementary matrix algebra.

*Keywords:* maximum likelihood estimator, true parameter, surprisal, relative information

# 1 Introduction

The consistency of the [Fisher \(1922\)](#) maximum likelihood estimator (MLE) is a basic result discovered more than a century ago. This method is typically treated as a special case of  $M$ -estimators, and derived via standard techniques dating back to [Cramér \(1946\)](#) and [Wald \(1949\)](#). A modern treatment is [van der Vaart \(1998\)](#). Because of its intuitive appeal and broad applicability, the method is a dominant tool of statistical inference.

In teaching this material, I found the [Doob \(1934\)](#) consistency proof of MLE convergence more motivated and accessible to students, particularly for students already familiar with information and entropy as part of a data science or machine learning course. The fact that the proof covers general parameter spaces and general state spaces with no extra effort only adds to its appeal.

The contribution of this note is expository: it gives a formulation of Doob's consistency argument in surprisal/information terminology, emphasizing the role of properness and separability, and works out the multivariate normal example in detail. The proofs are designed to be self-contained throughout: in particular, the consistency argument and the matrix lemmas are developed directly, without relying on determinant multiplicativity, eigenvalue continuity, or other standard results that can obscure the elementary structure of the argument.

We start with the setup and assumptions, then state and prove consistency. After that, we give a historical account of entropy terminology, which is an interesting story in itself, then we end by showing the assumptions hold for the multivariate normal case.

## 2 Surprisal and Information

A random variable  $X$  is sampled repeatedly, resulting in independent and identically distributed copies  $X_1, X_2, \dots$  of  $X$ . We assume the density  $p(X, \theta^*)$  of  $X$  is one of a parametric family  $p(X, \theta)$  of positive functions, and we wish to recover  $\theta^*$  from the samples  $X_1, X_2, \dots$ .

Stated in this manner, one may interpret the family  $p(X, \theta)$  as a probe of the underlying expectation  $E_*$ . While  $p(X, \theta)$  may be chosen arbitrarily, based on prior knowledge, we have no control over  $E_*$ . Given this, how do we use the samples  $X_1, X_2, \dots$  to recover  $\theta^*$ ?

Define the *surprisal* [Samson \(1951\)](#), [Tribus \(1961\)](#)

$$S(X, \theta) = \log(1/p(X, \theta)).$$

This terminology is intuitive: low probability equals high surprisal. Assume  $S(X, \theta)$  is integrable for all  $\theta$ . Then it is natural to call a minimizer  $\theta^*$  of the *expected surprisal*  $E_*(S(X, \theta))$  a *true parameter*, and the corresponding density  $p(X, \theta^*)$  a *true density*.

Equivalently, given  $\theta^*$ , the *relative information* is

$$I(\theta^*, \theta) = E_*(S(X, \theta)) - E_*(S(X, \theta^*)). \quad (1)$$

The relative information is the excess expected surprisal of  $\theta$  over  $\theta^*$ . Then  $\theta^*$  is a true parameter iff  $I(\theta^*, \theta) \geq 0$  for all  $\theta$ . The relative information  $I$  appears in many fields, in different forms. Because of widely differing terminology, in §4 we give an account of its history.

While  $p(X, \theta)$  have so far been arbitrary positive functions, we now specialize to the case where  $p(X, \theta)$  are densities in the usual sense. We say  $\theta^*$  is a *reference parameter* if

$$E_* \left( \frac{p(X, \theta)}{p(X, \theta^*)} \right) = 1 \quad \text{for all } \theta. \quad (2)$$

Then the corresponding density  $p(X, \theta^*)$  is a *reference density*.

Given  $\theta^*$ , define a background expectation  $E$  by  $E(f(X)) = E_*(f(X)/p(X, \theta^*))$ . Then, if  $\theta^*$  is a reference parameter,

$$E_*(f(X)) = E(p(X, \theta^*)f(X)) \quad \text{and} \quad E_\theta(f(X)) = E(p(X, \theta)f(X)) \quad (3)$$

hold, with  $E_\theta(1) = 1$  for all  $\theta$ . When  $\theta^*$  is a reference parameter, the relative information takes the familiar form [Donsker & Varadhan \(1975\)](#), [Dembo & Zeitouni \(1998\)](#) (see §4 for terminology)

$$I(\theta^*, \theta) = E \left( p(X, \theta^*) \log \left( \frac{p(X, \theta^*)}{p(X, \theta)} \right) \right). \quad (4)$$

Since  $j(p) = \log(1/p)$  is convex, by Jensen's inequality,

$$I(\theta^*, \theta) = E_* \left( j \left( \frac{p(X, \theta)}{p(X, \theta^*)} \right) \right) \geq j \left( E_* \left( \frac{p(X, \theta)}{p(X, \theta^*)} \right) \right) = j(1) = 0, \quad \text{for all } \theta.$$

It follows that a reference density is a true density.

If  $\theta$  is any other true parameter, then  $I(\theta^*, \theta) = 0$ , hence, by strict convexity of  $j(p)$ , there is a scalar  $t$  such that  $p(X, \theta) = t \cdot p(X, \theta^*)$ . Since  $\theta^*$  is a reference parameter,  $t = 1$ , hence  $p(X, \theta) = p(X, \theta^*)$ : every true density equals the reference density.

It follows there is at most one reference density; when there is a reference density, it is the unique true density. We say parameters  $\theta, \theta'$  *share a density* if  $p(X, \theta) = p(X, \theta')$ . When parameters do not share densities, there is at most one reference parameter; when there is a reference parameter, it is the unique true parameter.

The simplest example is as follows. A vector  $\theta = (\theta_1, \theta_2, \dots, \theta_d)$  is a probability vector if its components are nonnegative and sum to one. Suppose  $\theta^* = (\theta_1^*, \theta_2^*, \dots, \theta_d^*)$  is a probability vector and a random variable  $X$  satisfies  $P_*(X = i) = \theta_i^*, i = 1, 2, \dots, d$ . Then the mass function of  $X$  is  $p(X, \theta^*) = \theta_X^*$ . If  $\theta$  is any other probability vector, the surprisal is  $S(X, \theta) = \log(1/\theta_X)$ , and

$$I(\theta^*, \theta) = \sum_{i=1}^d \theta_i^* \log(\theta_i^*/\theta_i). \quad (5)$$

Turning back to the general setting, the *likelihood* and (empirical) *mean surprisal* are

$$L_n(\theta) = \prod_{k=1}^n p(X_k, \theta), \quad \bar{S}_n(\theta) = \frac{1}{n} \log(1/L_n(\theta)) = \frac{1}{n} \sum_{k=1}^n S(X_k, \theta).$$

The mean surprisal is also called the negative log-likelihood.

By definition,  $\theta^*$  minimizes  $E_*(S(X, \theta))$ . By the (strong) law of large numbers (LLN),  $\bar{S}_n(\theta)$  converges to  $E_*(S(X, \theta))$  as  $n \rightarrow \infty$ , with probability one, for each  $\theta$ . Hence it is natural to estimate  $\theta^*$  by a minimizer of  $\bar{S}_n(\theta)$  or, equivalently, a maximizer of  $L_n(\theta)$ : For each  $n \geq 1$ , a *maximum likelihood estimator (MLE)*, if it exists, is a parameter  $\hat{\theta}_n$ , depending on the samples, maximizing  $L_n(\theta)$  over all  $\theta$ . In what follows, only  $E_*$ ,  $p(X, \theta)$ , and true parameters enter;  $E$ ,  $E_\theta$ , and reference parameters play no role.

### 3 Consistency

Our first assumption is

**A. (Identifiability):** There is at most one true parameter.

As we saw above, **A** follows if there is a reference parameter and parameters do not share densities. A function  $F(\theta)$  is *proper* if its sublevel sets  $F(\theta) \leq c$  are compact subsets of the parameter space, for every level  $c$ . Since  $\hat{\theta}_n$  should minimize  $\bar{S}_n(\theta)$ , this leads us to the second assumption

**B. (Properness):** There is an  $N \geq 1$ , depending on the samples, and a proper  $F(\theta)$ , not depending on the samples, such that, with probability one,

$$F(\theta) \leq \bar{S}_n(\theta), \quad \text{for all } \theta \text{ and } n \geq N. \quad (6)$$

When the parameter space is compact, every continuous function  $F(\theta)$  is proper, and the proof below shows assumption **B** is superfluous. We will need the LLN convergence  $\bar{S}_n(\theta) \rightarrow E_*(S(X, \theta))$  across  $\theta$ , in the sense

$$\bar{S}_n(\theta) \rightarrow E_*(S(X, \theta)), \quad \text{as } n \rightarrow \infty, \quad \text{for all } \theta, \quad (7)$$

with probability one. For this we need the third assumption

**C. (Continuity):** The parameter space is separable and there is a continuous  $\delta(\theta, \theta')$  and an integrable nonnegative  $g(X)$  satisfying

$$|S(X, \theta) - S(X, \theta')| \leq \delta(\theta, \theta')g(X) \quad \text{and} \quad \delta(\theta, \theta) = 0. \quad (8)$$

Under **C**, a standard separability argument, given in the appendix, establishes (7) with probability one. Under **C**, the expected surprisal  $E_*(S(X, \theta))$  is continuous in  $\theta$ . If moreover **B** holds, then by sending  $n \rightarrow \infty$  in (6),  $F(\theta) \leq E_*(S(X, \theta))$  for all  $\theta$ , hence the expected surprisal is proper. Since a proper continuous function has a minimizer, under **A**, **B**, and **C**, there is a unique true parameter. Similarly, under **B** and **C**, the mean surprisal is continuous in  $\theta$  and proper, hence an MLE  $\hat{\theta}_n$  exists for  $n \geq N$ , with probability one.

These assumptions are particularly tractable for *exponential families* Cover & Thomas (2006), Murphy (2022), Schervish (1995), van der Vaart (1998), the densities  $p(X, \theta)$  where  $S(X, \theta)$  is a sum of products of the form  $f(\theta)g(X)$ .

In (7), there is no presumption of uniform convergence. This is a key feature of Doob’s proof: no uniform convergence of the mean surprisal is required. The argument instead combines properness, the LLN, and the continuity estimate in (8) to control limit points of the MLE sequence.

**Theorem** (Consistency). *Let  $X_1, X_2, \dots$  be independent and identically distributed, and let  $p(X, \theta)$  be a parametric family of positive functions. Under the above assumptions,*

1. *the expected surprisal is proper and there is a unique true parameter  $\theta^*$ ,*
2. *with probability one, an MLE sequence  $\hat{\theta}_n$  exists for sufficiently large  $n$ ,*
3. *with probability one, any MLE sequence converges to the true parameter,  $\hat{\theta}_n \rightarrow \theta^*$ .*

*Proof.* The first two items were derived above. For the third item, by the LLN,

$$\frac{1}{n} \sum_{k=1}^n g(X_k) \rightarrow E_*(g(X)), \quad \text{as } n \rightarrow \infty, \quad (9)$$

with probability one. Fix samples for which (6), (7) and (9) hold. We show  $\hat{\theta}_n \rightarrow \theta^*$  for these samples. By (6),

$$F(\hat{\theta}_n) \leq \bar{S}_n(\hat{\theta}_n) \leq \bar{S}_n(\theta^*), \quad \text{for } n \geq N.$$

Since  $\bar{S}_n(\theta^*) \rightarrow E_*(S(X, \theta^*))$ , the right side is bounded. Since  $F(\theta)$  is proper, the sequence  $\hat{\theta}_n$  lies in a compact set in the parameter space, and thus has limit points.

Suppose  $\theta$  is a limit point of  $\hat{\theta}_n$ . Then  $\hat{\theta}_n \rightarrow \theta$  along a subsequence. By (8),

$$\bar{S}_n(\theta) - \bar{S}_n(\theta^*) \leq \bar{S}_n(\theta) - \bar{S}_n(\hat{\theta}_n) \leq \delta(\theta, \hat{\theta}_n) \cdot \frac{1}{n} \sum_{k=1}^n g(X_k). \quad (10)$$

Since the left side of (9) is bounded as  $n \rightarrow \infty$ , and  $\delta(\theta, \hat{\theta}_n) \rightarrow 0$  along a subsequence, the right side of (10) goes to zero along a subsequence. By (1) and (7), the left side of (10) converges to  $I(\theta^*, \theta)$ , hence  $I(\theta^*, \theta) \leq 0$ . But  $I(\theta^*, \theta) \geq 0$ , hence  $I(\theta^*, \theta) = 0$ , showing  $\theta = \theta^*$ . Thus  $\hat{\theta}_n \rightarrow \theta^*$  as  $n \rightarrow \infty$ .  $\square$

## 4 Terminology History

The relative information  $I(\theta^*, \theta)$  is often called “relative entropy” Cover & Thomas (2006), Murphy (2022). The term “entropy,” originating in thermodynamics, was coined by Clausius (1865). Shortly afterward, Boltzmann (1872) derived his  $H$  theorem, using a quantity  $H$  proportional to the negative of entropy. Gibbs (1902), building on the combinatorial interpretation of entropy Boltzmann (1877) and Planck (1901), reversed Boltzmann’s sign convention, writing entropy with the sign convention used today.

Shannon (1948) independently defined

$$H(p) = - \sum_{i=1}^d p_i \log p_i.$$

von Neumann (1932) had earlier defined quantum entropy. According to a widely repeated account attributed to Shannon and reported in Tribus & McIrvine (1971), von Neumann suggested the name “entropy” when Shannon was casting about for terminology after developing his communication theory in the mid-1940s; later Shannon (1982) expressed doubt that the conversation took place. Whatever the origin of the name, in his 1948 paper, Shannon explicitly links his  $H$  to Boltzmann’s: “ $H$  is then, for example, the  $H$  in Boltzmann’s  $H$  theorem,” connecting his terminology to the earlier usage despite the difference in sign conventions.

The relative information  $I(\theta^*, \theta)$  (4) is jointly convex in the densities  $p(X, \theta^*)$  and  $p(X, \theta)$ , as is apparent in the special case (5), whereas the entropy  $H(p)$  above is strictly concave in  $p$ . From this perspective, the terminology “relative entropy” can seem counterintuitive. This terminology extends into the Python `scipy.stats` package: `entropy(p)` returns the concave  $H$ , whereas `entropy(p, q)` returns the convex  $I$ . The issue is the conceptual opposition between entropy and information: entropy is concave, whereas information is convex.

Indeed, based on curvature, one might argue that the *negative* of (5) is the more natural candidate for “relative entropy.” Similar remarks apply to the “cross-entropy” loss  $J$  of machine learning. Since  $J$  differs from  $I$  by a constant, one might analogously call  $J$  the *cross-information*, and its negative “cross-entropy.”

While many authors use terminology different from ours, some align partially with our usage. Schervish (1995), uses “Kullback-Leibler information” for  $I$ , which is more descriptive than the widely used “Kullback-Leibler divergence” Kullback & Leibler (1951), Bishop (2006). In particular, Kullback (1959) refers to  $I$  as an “information function.” Donsker & Varadhan (1975) denote their rate function by  $I$  rather than  $H$ ; the relative information is the i.i.d. special case of their  $I$ .

Doob’s 1934 proof appeared at the dawn of modern probability theory, within the milieu of Daniell, Fréchet, Hopf, Khintchine, Kolmogorov, Lévy, and Wiener, and within a year of the Kolmogorov extension theorem Kolmogorov (1933).

## 5 Multivariate Normal

Let  $\mu$  be in  $\mathbf{R}^d$ ,  $Q$  a positive symmetric  $d \times d$  matrix,  $\theta = (\mu, Q)$ , and let  $v \cdot w$  be the dot product in  $\mathbf{R}^d$ . The determinant  $\det Q$  is the product of the eigenvalues of  $Q$ . With  $X$  in  $\mathbf{R}^d$ , the *normal density with mean  $\mu$  and variance  $Q$*  is  $p(X, \theta) = e^{-S(X, \theta)}$  where

$$S(X, \theta) = \frac{1}{2} \log \det(2\pi Q) + \frac{1}{2} (X - \mu) \cdot Q^{-1} (X - \mu). \quad (11)$$

This defines the parametric family.

A random variable  $X$  in  $\mathbf{R}^d$  is *normally distributed with parameter  $\theta^* = (\mu^*, Q^*)$*  if its moment-generating function is

$$E_*(e^{v \cdot X}) = \exp \left( v \cdot \mu^* + \frac{1}{2} v \cdot Q^* v \right). \quad (12)$$

Suppose  $X$  is normally distributed with parameter  $\theta^* = (\mu^*, Q^*)$  as in (12), and let  $X_1, X_2, \dots$  be independent and identically distributed copies of  $X$ . We show  $\theta^*$  is the true parameter, and we verify A, B, and C. We review the necessary matrix algebra.

We say a symmetric matrix  $Q$  is *positive*, and we write  $Q > 0$ , if  $v \cdot Qv > 0$  for  $v \neq 0$ . We say a symmetric matrix  $Q$  is *nonnegative*, and we write  $Q \geq 0$ , if  $v \cdot Qv \geq 0$  for all  $v$ . Given symmetric matrices  $Q, R$  we write  $Q \geq R$  if  $Q - R \geq 0$ . If  $Q > 0$ , then  $Q \geq \delta I$  for some  $\delta > 0$ . If  $Q \geq 0$  and  $R \geq S$ , then  $\text{trace}(QR) \geq \text{trace}(QS)$ . Let  $\|Q\| = \sqrt{\text{trace}(Q^2)}$ . Then  $\text{trace}(QR) = \text{trace}(RQ)$  and we have the Cauchy-Schwarz inequality

$$\text{trace}(QR) \leq \|Q\| \|R\|, \quad Q, R \text{ symmetric.}$$

Given vectors  $v, w$  in  $\mathbf{R}^d$ , the *tensor product*  $v \otimes w$  is the matrix whose  $ij$ -th entry is  $v_i w_j$ . Then  $v \otimes v$  is nonnegative and

$$v \cdot Qv = \text{trace}(Q(v \otimes v)).$$

Let  $R = v \otimes v$ . Since  $R^2 = |v|^2 R$  and  $\text{trace}(R) = |v|^2$ ,  $\|R\| = |v|^2$ . Using this and Cauchy-Schwarz,

$$\delta = \|Q\| \implies Q \leq \delta I. \quad (13)$$

If  $u = \sqrt{\epsilon}v - w/\sqrt{\epsilon}$ , then  $u \otimes u \geq 0$ , hence

$$v \otimes w + w \otimes v \leq \epsilon v \otimes v + \frac{1}{\epsilon} w \otimes w. \quad (14)$$

If  $Q_n$  is a sequence of symmetric matrices with  $Q_n \rightarrow 0$ , then  $\|Q_n\| \rightarrow 0$ . By (13), given  $Q > 0$ , there is  $N \geq 1$  such that  $Q_n \leq Q$  for  $n \geq N$ .

Let  $\theta' = (\mu', Q')$ ,  $g(X) = 1 + |X|^2$  and let  $2\delta(\theta, \theta')$  equal

$$|\log \det Q - \log \det Q'| + |Q^{-1}\mu - (Q')^{-1}\mu'| + \|Q^{-1} - (Q')^{-1}\| + |\mu \cdot Q^{-1}\mu - \mu' \cdot (Q')^{-1}\mu'|.$$

Since<sup>1</sup> the determinant is continuous,  $\delta(\theta, \theta')$  is continuous. By the triangle inequality and Cauchy-Schwarz, (8) follows from (11), establishing **C**.

The proof of the following is in the appendix.

**Proposition.** For  $Q^* > 0$  and  $c$  scalar, let  $\epsilon = e^{-2c-1} \det(2\pi e Q^*)$  and

$$F_*(Q, Q^*) = \frac{1}{2} \log \det(2\pi Q) + \frac{1}{2} \text{trace}(Q^{-1} Q^*), \quad Q > 0. \quad (15)$$

Then  $F_*(Q, Q^*)$  is minimized iff  $Q = Q^*$ , and  $F_*(Q, Q^*) \leq c$  implies  $\epsilon Q^* \leq Q \leq Q^*/\epsilon$ .

Let  $f(\mu) = Q^* + (\mu^* - \mu) \otimes (\mu^* - \mu)$ . Given  $\theta = (\mu, Q)$ , let

$$F(\mu, Q) = F_*(Q, f(\mu)) = F_*(Q, Q^*) + \frac{1}{2}(\mu^* - \mu) \cdot Q^{-1}(\mu^* - \mu).$$

By the proposition,  $(\mu^*, Q^*)$  is the unique minimizer of  $F(\mu, Q)$ .

In (12), multiply both sides by  $e^{-v \cdot \mu}$ , then replace  $v$  by  $tv$  and differentiate twice at  $t = 0$ . This yields

$$E_*(X) = \mu^*, \quad E_*((X - \mu) \otimes (X - \mu)) = f(\mu).$$

It follows  $E_*(S(X, \theta)) = F(\mu, Q)$ . It follows  $\theta^* = (\mu^*, Q^*)$  is the unique true parameter, establishing **A**.

The second consequence of the proposition is properness of  $F(\mu, Q)$ . If  $F(\mu, Q) \leq c$ , then  $F_*(Q, Q^*) \leq c$ , hence  $\epsilon Q^* \leq Q \leq Q^*/\epsilon$ , confining  $Q$  to a compact subset of the space of positive definite matrices. From this,

$$\begin{aligned} c &\geq F_*(Q, Q^*) + \frac{1}{2}(\mu^* - \mu) \cdot Q^{-1}(\mu^* - \mu) \\ &\geq F_*(Q^*, Q^*) + \frac{\epsilon}{2}(\mu^* - \mu) \cdot (Q^*)^{-1}(\mu^* - \mu), \end{aligned}$$

confining  $\mu$  to a compact subset of  $\mathbf{R}^d$ . Thus  $F(\mu, Q)$  is proper.

Let

$$f_\epsilon(\mu) = (1 - \epsilon)f(\mu) - \epsilon Q^*, \quad F_\epsilon(\theta) = F_\epsilon(\mu, Q) = F_*(Q, f_\epsilon(\mu)).$$

Since  $f_\epsilon(\mu) \geq (1 - 2\epsilon)f(\mu)$ ,

$$F_\epsilon(\mu, (1 - 2\epsilon)Q) \geq \frac{1}{2} \log \det((1 - 2\epsilon)I) + F(\mu, Q),$$

hence  $F_\epsilon(\theta)$  is proper.

With

$$Q_n(\mu) = \frac{1}{n} \sum_{k=1}^n (X_k - \mu) \otimes (X_k - \mu), \quad \hat{\mu}_n = \frac{1}{n} \sum_{k=1}^n X_k, \quad \hat{Q}_n = Q_n(\hat{\mu}_n),$$

---

<sup>1</sup>See Lemma 2 in the appendix.

let  $\hat{\theta}_n = (\hat{\mu}_n, \hat{Q}_n)$ , and let  $0 < \epsilon < 1/2$ .

By the LLN, with probability one,

$$(\hat{\mu}_n - \mu^*) \otimes (\hat{\mu}_n - \mu^*) \rightarrow 0, \quad Q^* - Q_n(\mu^*) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Thus, from above, there is  $N \geq 1$ , depending on the samples, such that, with probability one,

$$(\hat{\mu}_n - \mu^*) \otimes (\hat{\mu}_n - \mu^*) \leq \epsilon^2 Q^*, \quad Q_n(\mu^*) \geq (1 - \epsilon)Q^*, \quad \text{for } n \geq N. \quad (16)$$

Fix samples for which (16) holds. We establish (6) for these samples and  $F(\theta) = F_\epsilon(\theta)$ .

With  $w = \hat{\mu}_n - \mu^*$  and  $v = \mu - \mu^*$ , using (14),

$$v \otimes w + w \otimes v \leq \epsilon(\mu^* - \mu) \otimes (\mu^* - \mu) + \frac{1}{\epsilon}(\hat{\mu}_n - \mu^*) \otimes (\hat{\mu}_n - \mu^*) \leq \epsilon f(\mu)$$

for all  $\mu$  and  $n \geq N$ . Writing  $X_k - \mu = (X_k - \mu^*) - v$ ,

$$Q_n(\mu) = Q_n(\mu^*) - (v \otimes w + w \otimes v) + v \otimes v.$$

Combined with the above inequalities, this leads to

$$Q_n(\mu) \geq f_\epsilon(\mu), \quad \text{for all } \mu \text{ and } n \geq N.$$

In particular, since  $f_\epsilon(\mu) \geq (1 - 2\epsilon)Q^*$ , we have shown  $\hat{Q}_n > 0$  for  $n \geq N$ .

Since  $\bar{S}_n(\theta) = F_*(Q, Q_n(\mu))$  and  $Q_n(\mu) \geq \hat{Q}_n$  by least squares, by the proposition,

$$\bar{S}_n(\theta) = F_*(Q, Q_n(\mu)) \geq F_*(Q, \hat{Q}_n) \geq F_*(\hat{Q}_n, \hat{Q}_n) = \bar{S}_n(\hat{\theta}_n),$$

for all  $\theta$  and  $n \geq N$ . Thus  $\hat{\theta}_n$  is the MLE for  $n \geq N$ , and

$$\bar{S}_n(\theta) = F_*(Q, Q_n(\mu)) \geq F_*(Q, f_\epsilon(\mu)) = F_\epsilon(\theta), \quad \text{for all } \theta \text{ and } n \geq N.$$

This establishes (6) and completes the derivation of B.

## 6 Disclosures

The author reports no funding was obtained for this work and there are no competing interests to declare. The author used a generative AI tool to assist with language editing and bibliographic searches. All AI-assisted content was reviewed, verified, and edited by the author, who takes full responsibility for the content, accuracy, and originality of this work.

## A Auxiliary Proofs

*Proof of (7) with probability one.* By separability of the parameter space, there is a countable dense set  $D$  of parameters. Since  $D$  is countable, with probability one,

$$\bar{S}_n(\theta') \rightarrow E_*(S(X, \theta')), \quad \text{as } n \rightarrow \infty, \quad \text{for all } \theta' \text{ in } D. \quad (17)$$

Fix samples for which (9) and (17) hold. We establish (7) for these samples. If  $\theta, \theta'$  are any parameters, by (8) and the triangle inequality,

$$\begin{aligned} |\bar{S}_n(\theta) - E_*(S(X, \theta))| &\leq |\bar{S}_n(\theta') - E_*(S(X, \theta'))| \\ &\quad + \delta(\theta, \theta') \left( E_*(g(X)) + \frac{1}{n} \sum_{k=1}^n g(X_k) \right). \end{aligned} \quad (18)$$

Now take the  $\limsup_{n \rightarrow \infty}$  in (18). By (9) and (17),

$$\limsup_{n \rightarrow \infty} |\bar{S}_n(\theta) - E_*(S(X, \theta))| \leq 2\delta(\theta, \theta') E_*(g(X)), \quad \text{for all } \theta' \text{ in } D \text{ and all } \theta.$$

Since  $D$  is dense,  $\delta(\theta, \theta')$  is continuous, and  $\delta(\theta, \theta) = 0$ , for fixed  $\theta$  and varying  $\theta'$ , the right side is arbitrarily small, while the left side does not depend on  $\theta'$ . This establishes (7).  $\square$

To derive the proposition, we need three Lemmas. The proofs below are included to keep the paper self-contained: the point is not merely to verify the required facts, but to expose the elementary mechanism behind them.

By the eigenvalue decomposition, every symmetric matrix is a linear combination of matrices of the form  $v \otimes v$ , and every positive matrix is a sum of  $\delta I$  and matrices of the form  $v \otimes v$ .

**Lemma 1.** *With  $Q$  invertible symmetric and  $t$  a scalar,  $\det(Q + tv \otimes v) = \det(Q) (1 + tv \cdot Q^{-1}v)$ .*

*Proof.* Assume  $t \neq 0$ . If  $v$  is orthogonal to an eigenvector  $x$  of  $Q$ , then we reduce to a lower-dimensional case by restricting to  $x^\perp$ . Hence we may assume  $v \cdot x \neq 0$  for all eigenvectors  $x$  of  $Q$ . If  $v$  is orthogonal to an eigenvector  $x'$  of  $Q' = Q + tv \otimes v$ , then  $x'$  is an eigenvector of  $Q$ , contradicting our assumption. Hence we may assume  $v \cdot x' \neq 0$  for all eigenvectors  $x'$  of  $Q'$ . If  $x'_1$  and  $x'_2$  are linearly independent eigenvectors of  $Q'$  corresponding to the same eigenvalue, then, for suitable  $r_1, r_2$ ,  $x' = r_1 x'_1 + r_2 x'_2$  is an eigenvector of  $Q'$  satisfying  $v \cdot x' = 0$ , contradicting our assumptions. Hence we may assume the eigenvalues of  $Q'$  are simple. If  $x$  and  $x'$  are eigenvectors for  $Q$  and  $Q'$  corresponding to the same eigenvalue, then

$$t(v \cdot x')(v \cdot x) = x' \cdot (tv \otimes v)x = x' \cdot Q'x - x' \cdot Qx = 0,$$

contradicting our assumptions. Hence we may assume the eigenvalues of  $Q$  and the eigenvalues of  $Q'$  do not overlap. Let  $\lambda_1, \lambda_2, \dots, \lambda_d$  be the eigenvalues of  $Q$  and  $\lambda'_1, \lambda'_2, \dots, \lambda'_d$  the eigenvalues of  $Q'$ . Let  $v_1, v_2, \dots, v_d$  be the corresponding orthonormal basis of eigenvectors of  $Q$ . Then the secular equation is

$$\sigma(z) = 1 + tv \cdot (Q - zI)^{-1}v = 1 + t \sum_{i=1}^d \frac{(v \cdot v_i)^2}{\lambda_i - z} = 0.$$

If  $\lambda'$  is an eigenvalue of  $Q'$  with eigenvector  $x'$ , then  $(Q' - \lambda'I)x' = 0$ , hence  $(Q - \lambda'I)x' + t(v \cdot x')v = 0$ . Taking the dot product with  $(Q - \lambda'I)^{-1}v$  yields  $v \cdot x' + t(v \cdot x')(v \cdot (Q - \lambda'I)^{-1}v)$ , hence  $\sigma(\lambda') = 0$ . If we let

$$\det(Q - zI) = \prod_{i=1}^d (\lambda_i - z), \quad p(z) = \det(Q - zI)\sigma(z),$$

then  $p(z)$  is a polynomial and  $p(\lambda') = 0$ . Since  $\lambda'_1, \lambda'_2, \dots, \lambda'_d$  are simple,  $p(z) = \prod_{i=1}^d (\lambda'_i - z)$ . Since the product of the roots is  $p(0)$ , the identity follows.  $\square$

**Lemma 2.** *If  $Q > 0$ ,  $S > 0$ , and  $Z > 0$ , then  $\det Q$  is continuous, and  $\det(SZS) = \det(Z)\det(S)^2$ .*

*Proof.* By Lemma 1,  $\det(Q + \lambda v \otimes v)$  is a continuous function of  $\lambda$  and  $v$ . By Lemma 1 again,  $\det(Q + \lambda_1 v_1 \otimes v_1 + \lambda_2 v_2 \otimes v_2)$  is a continuous function of  $\lambda_1, \lambda_2$  and  $v_1, v_2$ . Continuing in this manner, with  $Z = \lambda_1 v_1 \otimes v_1 + \lambda_2 v_2 \otimes v_2 + \dots + \lambda_d v_d \otimes v_d$ ,  $\det(Q + Z)$  is a continuous function of  $\lambda_1, \lambda_2, \dots, \lambda_d$  and  $v_1, v_2, \dots, v_d$ . If  $Z_n$  is a sequence of symmetric matrices converging to zero, then  $\|Z_n\| \rightarrow 0$ , hence the eigenvalues of  $Z_n$  converge to zero, hence we may write  $Z_n = t_n Z'_n$  with  $t_n \rightarrow 0$  and  $\|Z'_n\| \leq 1$ . By passing to a subsequence, we may assume the eigenvalues and eigenvectors of  $Z'_n$  converge. Since  $\det(Q + t_n Z'_n)$  is a continuous function of the scalars  $t_n$ , these eigenvalues, and eigenvectors,  $\det(Q + Z_n) \rightarrow \det Q$ . This establishes continuity.

Assume the second result holds for  $Z$ . By Lemma 1,

$$\begin{aligned} \det(S(Z + v \otimes v)S) &= \det(SZS + Sv \otimes Sv) \\ &= \det(SZS)(1 + Sv \cdot (SZS)^{-1}Sv) \\ &= \det(Z)\det(S)^2(1 + v \cdot Z^{-1}v) = \det(Z + v \otimes v)\det(S)^2, \end{aligned}$$

hence the second result holds for  $Z + v \otimes v$ . If  $Z = \delta I$ , the second result holds. Since every  $Z > 0$  is a sum of  $\delta I$  and a sum of matrices of the form  $v \otimes v$ , the second result follows by induction over the number of terms in the sum.  $\square$



**Lemma 3.** For  $c$  scalar, let  $\epsilon = e^{-c}$  and  $h(\lambda) = \log \lambda + 1/\lambda$ ,  $\lambda > 0$ . Then  $h(\lambda)$  is minimized iff  $\lambda = 1$ , and  $h(\lambda) \leq c$  implies  $\epsilon \leq \lambda \leq 1/\epsilon$ .

*Proof.* The first part is checked by differentiation. For the second part, clearly  $h(\lambda) \leq c$  implies  $\lambda \leq 1/\epsilon$ . Since  $\lambda \log \lambda \geq -1/e$  and  $(e-1)e^c \geq ec$  for  $c \geq 1$ ,  $h(\lambda) \leq c$  and  $c \geq 1$  implies  $\lambda \geq (1-1/e)/c \geq \epsilon$ .  $\square$

*Proof of the proposition.* Since  $Q^* > 0$ , there is  $S > 0$  with  $Q^* = S^2$ . Let  $Z = S^{-1}QS^{-1}$  and let  $\lambda_1, \lambda_2, \dots, \lambda_d$  be the eigenvalues of  $Z$ . Then  $\text{trace}(Q^{-1}Q^*) = \text{trace}(SQ^{-1}S) = \text{trace}(Z^{-1})$ . By Lemmas 2 and 3,

$$\begin{aligned} 2F_*(Q, Q^*) - 2F_*(Q^*, Q^*) &= \log \det Q - \log \det Q^* + \text{trace}(Q^{-1}Q^* - I) \\ &= \log \det(SZS) - \log \det(S)^2 + \text{trace}(Z^{-1} - I) \\ &= \log \det Z + \text{trace}(Z^{-1} - I) \\ &= \sum_{i=1}^d (h(\lambda_i) - 1) \geq 0, \end{aligned} \tag{19}$$

establishing  $F_*(Q, Q^*) \geq F_*(Q^*, Q^*)$ . If (19) is an equality,  $Z = I$  hence  $Q = Q^*$ . Now suppose  $F_*(Q, Q^*) \leq c$ . Then

$$2F_*(Q, Q^*) - 2F_*(Q^*, Q^*) \leq 2c - \log \det(2\pi Q^*) - \text{trace}(I) = \log(1/\epsilon) - 1.$$

Since the terms in the sum in (19) are nonnegative,  $h(\lambda_i) - 1 \leq \log(1/\epsilon) - 1$  for  $i = 1, 2, \dots, d$ . By Lemma 3,  $\epsilon \leq \lambda_i \leq 1/\epsilon$  for  $i = 1, 2, \dots, d$ . This implies  $\epsilon I \leq Z \leq I/\epsilon$ , hence  $\epsilon Q^* \leq Q \leq Q^*/\epsilon$ .  $\square$

## References

- Bishop, C. M. (2006), *Pattern Recognition and Machine Learning*, Information Science and Statistics, Springer, Cham.
- Boltzmann, L. (1872), ‘Weitere Studien über das Wärmegleichgewicht unter Gas-molekülen’, *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe. Abteilung II* **66**, 275–370.
- Boltzmann, L. (1877), ‘Über die beziehung zwischen dem zweiten hauptsatze der mechanischen wärmetheorie und der wahrscheinlichkeitsrechnung respektive den sätzen über das wärmegleichgewicht’, *Wiener Berichte* **76**, 373–435.
- Clausius, R. (1865), ‘Über verschiedene für die anwendung bequeme formen der hauptgleichungen der mechanischen wärmetheorie’, *Annalen der Physik und Chemie* **125**, 353–400.
- Cover, T. M. & Thomas, J. A. (2006), *Elements of Information Theory*, 2nd edn, Wiley-Interscience.
- Cramér, H. (1946), *Mathematical Methods of Statistics*, Princeton University Press.
- Dembo, A. & Zeitouni, O. (1998), *Large Deviations Techniques and Applications*, Vol. 38 of *Stochastic Modelling and Applied Probability*, 2nd edn, Springer, New York.
- Donsker, M. D. & Varadhan, S. R. S. (1975), ‘Asymptotic evaluation of certain markov process expectations for large time, I’, *Communications on Pure and Applied Mathematics* **28**(1), 1–47.
- Doob, J. L. (1934), ‘Probability and statistics’, *Transactions of the American Mathematical Society* **36**, 759–775.
- Fisher, R. A. (1922), ‘On the mathematical foundations of theoretical statistics’, *Philosophical Transactions of the Royal Society of London, Series A* **222**, 309–368.

- Gibbs, J. W. (1902), *Elementary Principles in Statistical Mechanics*, Yale University Press, New Haven.
- Kolmogorov, A. N. (1933), *Grundbegriffe der Wahrscheinlichkeitsrechnung*, Springer, Berlin.
- Kullback, S. (1959), *Information Theory and Statistics*, Wiley, New York.
- Kullback, S. & Leibler, R. A. (1951), ‘On information and sufficiency’, *Annals of Mathematical Statistics* **22**(1), 79–86.
- Murphy, K. P. (2022), *Probabilistic Machine Learning: An Introduction*, MIT Press.
- Planck, M. (1901), ‘Ueber das gesetz der energieverteilung im normalspectrum’, *Annalen der Physik* **309**(3), 553–563.
- Samson, E. W. (1951), Fundamental natural concepts of information theory, Report E5079, Communications Laboratory, Electronics Research Division, Air Force Cambridge Research Center, Cambridge, MA.
- Schervish, M. J. (1995), *Theory of Statistics*, Springer, New York.
- Shannon, C. E. (1948), ‘A mathematical theory of communication’, *Bell System Technical Journal* **27**(3), 379–423.
- Shannon, C. E. (1982), ‘Oral-History: Claude E. Shannon’, Engineering and Technology History Wiki (ETHW), interview by Robert Price. Interview conducted 28 July 1982.  
**URL:** [https://ethw.org/Oral-History:Claude\\_E.\\_Shannon](https://ethw.org/Oral-History:Claude_E._Shannon)
- Tribus, M. (1961), *Thermostatistics and Thermodynamics: An Introduction to Energy, Information and States of Matter, With Engineering Applications*, D. Van Nostrand Company, Princeton, NJ.
- Tribus, M. & McIrvine, E. C. (1971), ‘Energy and information’, *Scientific American* **225**(3), 179–190.
- van der Vaart, A. W. (1998), *Asymptotic Statistics*, Vol. 3 of *Cambridge Series in Statistical and Probabilistic Mathematics*, Cambridge University Press, Cambridge.
- von Neumann, J. (1932), *Mathematische Grundlagen der Quantenmechanik*, Springer, Berlin.
- Wald, A. (1949), ‘Note on the consistency of the maximum likelihood estimate’, *Annals of Mathematical Statistics* **20**(4), 595–601.