

Doob's Proof of MLE Strong Consistency: A Surprisal and Information Perspective

Omar Hijab

Department of Mathematics, Temple University

2026

Abstract

We give a concise, self-contained exposition of Doob's proof of strong maximum likelihood consistency, formulated in terms of surprisal and relative information. This formulation clarifies the roles of identifiability, properness, and continuity, and highlights that consistency holds without uniform convergence of the log-likelihood. We also verify these conditions for the multivariate normal distribution, recovering the sample mean and biased sample covariance as the maximum likelihood estimators.

Keywords: maximum likelihood estimator, true parameter, relative information

1 Introduction

The consistency of the [Fisher \(1922\)](#) maximum likelihood estimator (MLE) is a basic result discovered more than a century ago. This method is typically treated as a special case of M -estimators, and derived via standard techniques dating back to [Cramér \(1946\)](#) and [Wald \(1949\)](#). A modern treatment is [van der Vaart \(1998\)](#). Because of its intuitive appeal and broad applicability, the method is a dominant tool of statistical inference.

In teaching this material, I found the [Doob \(1934\)](#) strong consistency proof of MLE convergence more motivated and accessible to students, particularly for students already familiar with information and entropy as part of a data science or machine learning course. The fact that the proof covers general parameter spaces and general state spaces with essentially no extra effort only adds to its appeal.

The contribution of this note is expository and pedagogical: We give a formulation of Doob's consistency argument in surprisal/information terminology, emphasizing that strong consistency holds without assuming uniform convergence, and we work out the multivariate normal example.

2 Surprisal and Information

A random variable X is sampled repeatedly, resulting in independent and identically distributed copies $X_1, X_2, \dots$ of X . We assume the density $p(X, \theta^*)$ of X is one of a parametric family $p(X, \theta)$ of positive functions, and we wish to recover θ^* from the samples $X_1, X_2, \dots$. While $p(X, \theta)$ may be chosen arbitrarily, based on prior knowledge, we have no control over the underlying expectation E_* . Given this, how do we use the samples $X_1, X_2, \dots$ to recover θ^* ?

Define the *surprisal* [Samson \(1951\)](#), [Tribus \(1961\)](#)

$$S(X, \theta) = \log(1/p(X, \theta)).$$

This terminology is intuitive: low probability equals high surprisal. Assume $S(X, \theta)$ is integrable for all θ . Then it is natural to call a minimizer θ^* of the *expected surprisal* $E_*(S(X, \theta))$ a *true parameter*, and the corresponding density $p(X, \theta^*)$ a *true density*. Equivalently, given θ^* and θ , the *relative information* is

$$I(\theta^*, \theta) = E_*(S(X, \theta)) - E_*(S(X, \theta^*)). \quad (1)$$

The relative information is the excess expected surprisal of θ over θ^* . Then θ^* is a true parameter iff $I(\theta^*, \theta) \geq 0$ for all θ .

While $p(X, \theta)$ have so far been arbitrary positive functions, we now specialize to the case where $p(X, \theta)$ are densities in the usual sense. We say θ^* is a *reference parameter* if

$$E_* \left(\frac{p(X, \theta)}{p(X, \theta^*)} \right) = 1 \quad \text{for all } \theta.$$

Then the corresponding density $p(X, \theta^*)$ is a *reference density*.

Let θ^* be a reference parameter, and define E and E_θ by

$$E_*(f(X)) = E(p(X, \theta^*)f(X)) \quad \text{and} \quad E_\theta(f(X)) = E(p(X, \theta)f(X)).$$

Then the relative information takes the familiar form [Donsker & Varadhan \(1975\)](#), [Dembo & Zeitouni \(1998\)](#)

$$I(\theta^*, \theta) = E \left(p(X, \theta^*) \log \left(\frac{p(X, \theta^*)}{p(X, \theta)} \right) \right), \quad (2)$$

and $E_\theta(1) = 1$ for all θ . Since $j(p) = \log(1/p)$ is convex, by Jensen's inequality,

$$I(\theta^*, \theta) = E_* \left(j \left(\frac{p(X, \theta)}{p(X, \theta^*)} \right) \right) \geq j \left(E_* \left(\frac{p(X, \theta)}{p(X, \theta^*)} \right) \right) = j(E_\theta(1)) = j(1) = 0, \quad \text{for all } \theta.$$

It follows that a reference density is a true density.

If θ is any other true parameter, then $I(\theta^*, \theta) = 0$, hence, by strict convexity of $j(p)$, there is a scalar t such that $p(X, \theta) = t \cdot p(X, \theta^*)$. Since θ^* is a reference parameter, $t = 1$, hence $p(X, \theta) = p(X, \theta^*)$: every true density equals the reference density.

It follows that there is at most one reference density; when there is a reference density, it is the unique true density. We say parameters θ, θ' *share a density* if $p(X, \theta) = p(X, \theta')$. When parameters do not share densities, there is at most one reference parameter; when there is a reference parameter, it is the unique true parameter.

The simplest example is as follows. A vector $\theta = (\theta_1, \theta_2, \dots, \theta_d)$ is a probability vector if its components are nonnegative and sum to one. Suppose $\theta^* = (\theta_1^*, \theta_2^*, \dots, \theta_d^*)$ is a probability vector and a random variable X satisfies $P_*(X = i) = \theta_i^*$, $i = 1, 2, \dots, d$. Then the mass function of X is $p(X, \theta^*) = \theta_X^*$. If θ is any other probability vector, the surprisal is $S(X, \theta) = \log(1/\theta_X)$, and

$$I(\theta^*, \theta) = \sum_{i=1}^d \theta_i^* \log(\theta_i^*/\theta_i). \quad (3)$$

In the form (2), $I(\theta^*, \theta)$ is more commonly known as Kullback-Leibler divergence [van der Vaart \(1998\)](#) or relative entropy [Cover & Thomas \(2006\)](#), [Dembo & Zeitouni \(1998\)](#) or both [Bishop \(2006\)](#), [Murphy \(2022\)](#), and appears as the rate function in Sanov's theorem [Sanov \(1957\)](#). We regard it as information, consistent with some authors [Kullback \(1959\)](#), [Schervish \(1995\)](#), since information and entropy are conceptual opposites: the [Shannon \(1948\)](#) entropy $-\sum_{i=1}^d p_i \log p_i$ is concave, while (3) is convex.

Returning to the general setting, the *likelihood* and (empirical) *mean surprisal* are

$$L_n(\theta) = \prod_{k=1}^n p(X_k, \theta), \quad S_n(\theta) = \frac{1}{n} \log(1/L_n(\theta)) = \frac{1}{n} \sum_{k=1}^n S(X_k, \theta).$$

The mean surprisal is also called the negative log-likelihood.

By definition, θ^* minimizes $E_*(S(X, \theta))$. By the strong law of large numbers (LLN), $S_n(\theta)$ converges to $E_*(S(X, \theta))$ as $n \rightarrow \infty$, with probability one, for each θ . Hence it is natural to estimate θ^* by a minimizer of $S_n(\theta)$ or, equivalently, a maximizer of $L_n(\theta)$: For each $n \geq 1$, a *maximum likelihood estimator (MLE)*, if it exists, is a parameter $\hat{\theta}_n$, depending on the samples, maximizing $L_n(\theta)$ over all θ . In what follows, only E_* , $p(X, \theta)$, and true parameters enter; E , E_θ , and reference parameters play no role.

3 Consistency

In addition to integrability of $S(X, \theta)$, we make three assumptions. Our first assumption is

A (Identifiability). *There is at most one true parameter.*

A follows if the expected surprisal is strictly convex. Also, as we saw above, **A** follows if there is a reference parameter and parameters do not share densities.

A sequence of parameters *subconverges* to θ if a subsequence converges to θ . If a sequence subconverges to θ , θ is a *limit point of the sequence*. A sequence has exactly one limit point θ iff the sequence converges to θ . A subset K of the parameter space is *compact* if every sequence in K has at least one limit point in K .

A function $F(\theta)$ is *proper* if its sublevel sets $F(\theta) \leq c$ are compact subsets of the parameter space, for every level c . Since a proper continuous function has a minimizer and $\hat{\theta}_n$ should minimize $S_n(\theta)$, this leads us to the second assumption

B (Properness). *There is an $N \geq 1$, depending on the samples, and a proper $F(\theta)$, not depending on the samples, such that, with probability one,*

$$F(\theta) \leq S_n(\theta), \quad \text{for all } \theta \text{ and } n \geq N. \quad (4)$$

When the parameter space is compact, every continuous function $F(\theta)$ is proper, and the proof below shows assumption **B** is superfluous. We will need the pointwise LLN convergence $S_n(\theta) \rightarrow E_*(S(X, \theta))$ simultaneously for all parameters, in the sense

$$S_n(\theta) \rightarrow E_*(S(X, \theta)), \quad \text{as } n \rightarrow \infty, \quad \text{for all } \theta, \quad (5)$$

with probability one. We emphasize this is not uniform convergence, since the rate of convergence may vary across parameters θ . For this we need the third assumption

C (Continuity). *The parameter space is separable and there is a continuous $\delta(\theta, \theta')$ and an integrable non-negative $g(X)$ satisfying*

$$|S(X, \theta) - S(X, \theta')| \leq \delta(\theta, \theta')g(X) \quad \text{and} \quad \delta(\theta, \theta) = 0. \quad (6)$$

Under **C**, a standard separability argument establishes (5) with probability one. Under **C**, the expected surprisal $E_*(S(X, \theta))$ is continuous in θ . If moreover **B** holds, then by sending $n \rightarrow \infty$ in (4), $F(\theta) \leq E_*(S(X, \theta))$ for all θ , hence the expected surprisal is proper. Under **A**, **B**, and **C**, there is a unique true parameter. Similarly, under **B** and **C**, the mean surprisal is continuous in θ and proper, hence an MLE $\hat{\theta}_n$ exists for $n \geq N$, with probability one.

These assumptions are particularly tractable for *exponential families* Cover & Thomas (2006), Murphy (2022), Schervish (1995), van der Vaart (1998), in which $S(X, \theta)$ is essentially a sum of products of the form $f(\theta)g(X)$. In particular, we verify these assumptions for the multivariate normal in §4.

In (5), there is no presumption of uniform convergence. This is a key feature of Doob's proof. The argument instead combines properness, the LLN, and the continuity estimate in (6) to control limit points of the MLE sequence.

Theorem (Consistency). *Let $X_1, X_2, \dots$ be independent and identically distributed, and let $p(X, \theta)$ be a parametric family of positive functions. Under the above assumptions,*

1. *there is a unique true parameter θ^* ,*
2. *with probability one, an MLE sequence $\hat{\theta}_n$ exists for sufficiently large n ,*
3. *with probability one, any MLE sequence converges to the true parameter, $\hat{\theta}_n \rightarrow \theta^*$.*

Proof. The first two items were derived above. For the third item, by the LLN,

$$\frac{1}{n} \sum_{k=1}^n g(X_k) \rightarrow E_*(g(X)), \quad \text{as } n \rightarrow \infty, \quad (7)$$

with probability one. Fix samples $X_1, X_2, \dots$ for which (4), (5) and (7) hold. We show $\hat{\theta}_n \rightarrow \theta^*$ for these samples. By (4),

$$F(\hat{\theta}_n) \leq S_n(\hat{\theta}_n) \leq S_n(\theta^*), \quad \text{for } n \geq N.$$

By (5), the right side of this last display is bounded as $n \rightarrow \infty$. Since $F(\theta)$ is proper, the sequence $\hat{\theta}_n$ lies in a compact set in the parameter space, and thus has limit points.

Suppose θ is a limit point of $\hat{\theta}_n$. Then $\hat{\theta}_n$ subconverges to θ . By (6),

$$S_n(\theta) - S_n(\theta^*) \leq S_n(\theta) - S_n(\hat{\theta}_n) \leq \delta(\theta, \hat{\theta}_n) \cdot \frac{1}{n} \sum_{k=1}^n g(X_k). \quad (8)$$

Since the left side of (7) is bounded as $n \rightarrow \infty$, and $\delta(\theta, \hat{\theta}_n)$ subconverges to zero, the right side of (8) subconverges to zero. By (1) and (5), the left side of (8) converges to $I(\theta^*, \theta)$, hence $I(\theta^*, \theta) \leq 0$. But $I(\theta^*, \theta) \geq 0$ since θ^* is a true parameter, hence $I(\theta^*, \theta) = 0$. Since there is only one true parameter, $\theta = \theta^*$. Thus θ^* is the only limit point, hence $\hat{\theta}_n \rightarrow \theta^*$ as $n \rightarrow \infty$. $\square$

4 Multivariate Normal

Let μ be in $\mathbf{R}^d$, Q a positive definite¹ $d \times d$ matrix, $\theta = (\mu, Q)$, and let $v \cdot w$ be the dot product in $\mathbf{R}^d$. With X in $\mathbf{R}^d$, the *normal density with mean μ and covariance Q* is $p(X, \theta) = e^{-S(X, \theta)}$ where

$$S(X, \theta) = \frac{1}{2} \log \det(2\pi Q) + \frac{1}{2} (X - \mu) \cdot Q^{-1} (X - \mu).$$

¹Background linear algebra is summarized in the appendix.

This defines the parametric family.

A random variable X in $\mathbf{R}^d$ is *normally distributed with parameter* $\theta^* = (\mu^*, Q^*)$ if its moment-generating function is

$$E_*(e^{v \cdot X}) = \exp \left(v \cdot \mu^* + \frac{1}{2} v \cdot Q^* v \right).$$

Let $X_1, X_2, \dots$ be independent and identically distributed copies of X . We show $S(X, \theta)$ is integrable and θ^* is the true parameter, and we verify [A](#), [B](#), and [C](#).

Let $\theta' = (\mu', Q')$, $g(X) = 1 + |X|^2$ and let $\delta(\theta, \theta')$ equal half of

$$|\log \det Q - \log \det Q'| + |Q^{-1}\mu - (Q')^{-1}\mu'| + \|Q^{-1} - (Q')^{-1}\| + |\mu \cdot Q^{-1}\mu - \mu' \cdot (Q')^{-1}\mu'|.$$

Then $\delta(\theta, \theta')$ is continuous. By standard manipulations ([6](#)) follows, establishing [C](#).

Proposition. For $Q^* > 0$ and c scalar, let $\alpha = e^{-2c-1} \det(2\pi e Q^*)$ and

$$F_*(Q, Q^*) = \frac{1}{2} \log \det(2\pi Q) + \frac{1}{2} \text{trace}(Q^{-1} Q^*), \quad Q > 0.$$

Then $F_*(Q, Q^*)$ is minimized iff $Q = Q^*$, and $F_*(Q, Q^*) \leq c$ implies $\alpha Q^* \leq Q \leq Q^*/\alpha$.

Proof. Let $h(\lambda) = \log \lambda + 1/\lambda$, $\lambda > 0$. Then $h(\lambda)$ is minimized iff $\lambda = 1$ and $h(\lambda)$ is proper in the sense $h(\lambda) \leq \log(1/\alpha)$ implies $\alpha \leq \lambda \leq 1/\alpha$. Write $Q^* = S^2$ with $S > 0$, and define Z by $Q = SZS$. Suppose $F_*(Q, Q^*) \leq c$ and let $\lambda_1, \lambda_2, \dots, \lambda_d$ be the eigenvalues of Z . Then the proposition follows from

$$\log(1/\alpha) - 1 \geq 2F_*(Q, Q^*) - 2F_*(Q^*, Q^*) = \log \det Z + \text{trace}(Z^{-1} - I) = \sum_{i=1}^d (h(\lambda_i) - 1). \quad \square$$

Using the moment generating function,

$$E_*(S(X, \theta)) = F_*(Q, Q^*) + \frac{1}{2}(\mu^* - \mu) \cdot Q^{-1}(\mu^* - \mu).$$

Since $S(X, \theta)$ is bounded below for each θ , this shows $S(X, \theta)$ is integrable. By the proposition, (μ^*, Q^*) is the unique minimizer of the expected surprisal, establishing [A](#).

Now let $0 < \epsilon < 1/2$ and

$$F_\epsilon(\theta) = F_*(Q, (1 - 2\epsilon)Q^*) + \frac{1}{2}(1 - \epsilon)(\mu^* - \mu) \cdot Q^{-1}(\mu^* - \mu), \quad \theta = (\mu, Q).$$

By the proposition with $(1 - 2\epsilon)Q^*$ replacing Q^* , if $F_\epsilon(\theta) \leq c$, then Q is confined to a compact subset of positive definite matrices and, subsequently, μ is confined to a compact subset of $\mathbf{R}^d$. Hence $F_\epsilon(\theta)$ is proper.

With (see the appendix for the tensor product $\otimes$)

$$Q_n(\mu) = \frac{1}{n} \sum_{k=1}^n (X_k - \mu) \otimes (X_k - \mu), \quad \hat{\mu}_n = \frac{1}{n} \sum_{k=1}^n X_k, \quad \hat{Q}_n = Q_n(\hat{\mu}_n),$$

let $\hat{\theta}_n = (\hat{\mu}_n, \hat{Q}_n)$. Then $\hat{\mu}_n$ and $\hat{Q}_n$ are the usual sample mean and the biased sample covariance.

By the LLN applied to their entries, with probability one, the matrix sequences

$$(\hat{\mu}_n - \mu^*) \otimes (\hat{\mu}_n - \mu^*) \rightarrow 0, \quad Q^* - Q_n(\mu^*) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

If Q_n is any sequence of symmetric matrices with $Q_n \rightarrow 0$, then $\|Q_n\| \rightarrow 0$. Since $Q_n \leq \|Q_n\| I$, given any $Q > 0$, there is $N \geq 1$ such that $Q_n \leq Q$ for $n \geq N$. By choosing $Q = \epsilon^2 Q^*$ and $Q = \epsilon Q^*$, it follows there is $N \geq 1$, depending on the samples, such that, with probability one,

$$(\hat{\mu}_n - \mu^*) \otimes (\hat{\mu}_n - \mu^*) \leq \epsilon^2 Q^*, \quad Q_n(\mu^*) \geq (1 - \epsilon)Q^*, \quad \text{for } n \geq N. \quad (9)$$

Fix samples $X_1, X_2, \dots$ for which (9) holds. We establish (4) for these samples and $F(\theta) = F_\epsilon(\theta)$.

Let

$$f_\epsilon(\mu) = (1 - 2\epsilon)Q^* + (1 - \epsilon)(\mu - \mu^*) \otimes (\mu - \mu^*)$$

and let $f_0(\mu)$ be obtained by inserting $\epsilon = 0$. Then $F_\epsilon(\theta) = F_*(Q, f_\epsilon(\mu))$. With $w = \hat{\mu}_n - \mu^*$ and $v = \mu - \mu^*$, using (10) in the appendix,

$$v \otimes w + w \otimes v \leq \epsilon(\mu^* - \mu) \otimes (\mu^* - \mu) + \frac{1}{\epsilon}(\hat{\mu}_n - \mu^*) \otimes (\hat{\mu}_n - \mu^*) \leq \epsilon f_0(\mu)$$

for all μ and $n \geq N$. Writing $X_k - \mu = (X_k - \mu^*) - v$,

$$Q_n(\mu) = Q_n(\mu^*) - (v \otimes w + w \otimes v) + v \otimes v.$$

Combined with the above inequalities, this leads to

$$Q_n(\mu) \geq f_\epsilon(\mu), \quad \text{for all } \mu \text{ and } n \geq N.$$

In particular, since $f_\epsilon(\mu) \geq (1 - 2\epsilon)Q^*$, we have shown $\hat{Q}_n > 0$ for $n \geq N$.

Since $S_n(\theta) = F_*(Q, Q_n(\mu))$ and $Q_n(\mu) \geq \hat{Q}_n$ by least squares, by the proposition,

$$S_n(\theta) = F_*(Q, Q_n(\mu)) \geq F_*(Q, \hat{Q}_n) \geq F_*(\hat{Q}_n, \hat{Q}_n) = S_n(\hat{\theta}_n),$$

for all θ and $n \geq N$. Thus $\hat{\theta}_n$ is the MLE for $n \geq N$, and

$$S_n(\theta) = F_*(Q, Q_n(\mu)) \geq F_*(Q, f_\epsilon(\mu)) = F_\epsilon(\theta), \quad \text{for all } \theta \text{ and } n \geq N.$$

This establishes (4) and completes the derivation of B.

5 Disclosures

The author reports no funding was obtained for this work and there are no competing interests to declare. The author used a generative AI tool (Anthropic's *Claude Sonnet*) to assist with language editing and bibliographic searches. All AI-assisted content was reviewed, verified, and edited by the author, who takes full responsibility for the content, accuracy, and originality of this work.

Appendix

This is a review of the linear algebra used above. Let Q and Z be symmetric matrices. We say Q is *positive definite*, and write $Q > 0$, if $v \cdot Qv > 0$ for all vectors $v \neq 0$. We say Q is *nonnegative definite*, and write $Q \geq 0$, if $v \cdot Qv \geq 0$ for all vectors v . We write $Q \geq Z$ if $Q - Z \geq 0$. Then $aI \leq Q \leq bI$ iff the eigenvalues λ of Q satisfy $a \leq \lambda \leq b$. In particular, if $Q \geq 0$, the eigenvalues satisfy $\lambda \geq 0$, and, if $Q > 0$, the eigenvalues satisfy $\lambda > 0$. From this, if $Q \geq 0$, there is $S \geq 0$ satisfying $Q = S^2$, and $S > 0$ if $Q > 0$.

Given vectors $v = (s_1, s_2, \dots, s_d)$ and $w = (t_1, t_2, \dots, t_d)$, the *tensor product* $v \otimes w$ is the matrix whose ij -th entry is $s_i t_j$. Then $u \otimes u \geq 0$. In particular, taking $u = \sqrt{\epsilon}v - w/\sqrt{\epsilon}$, $u \otimes u \geq 0$ implies

$$v \otimes w + w \otimes v \leq \epsilon v \otimes v + \frac{1}{\epsilon} w \otimes w. \quad (10)$$

If $u_1, u_2, \dots, u_d$ is any orthonormal basis and $e_1, e_2, \dots, e_d$ is the standard one-hot encoded basis, then $\sum_{i=1}^d e_i \otimes e_i$ and $\sum_{i=1}^d u_i \otimes u_i$ both equal I .

For any square matrix A , the *trace* of A is the sum $\text{trace}(A) = \sum_{i=1}^d e_i \cdot A e_i$ of the diagonal entries of A . Then $\text{trace}(QZ) = \text{trace}(ZQ)$ and $\text{trace}(Q(u \otimes u)) = u \cdot Qu$ follow immediately. If $u_1, u_2, \dots, u_d$ is an orthonormal basis of eigenvectors of Q , $\text{trace}(Q)$ is the sum $\sum_{i=1}^d u_i \cdot Qu_i$ of the eigenvalues of Q , hence $\text{trace}(Q) \geq 0$ if $Q \geq 0$. Since $Z \geq 0$ and S symmetric imply $SZS \geq 0$, $\text{trace}(QZ) = \text{trace}(S^2Z) = \text{trace}(SZS) \geq 0$. It follows the trace is *monotone*,

$$Q \geq S \text{ and } Z \geq 0 \quad \implies \quad \text{trace}(QZ) \geq \text{trace}(SZ).$$

The *norm squared* $\|Q\|^2 = \text{trace}(Q^2)$ is the sum of the squares of the eigenvalues of Q , hence $Q \leq \|Q\| I$. Since $\|Q\|^2$ is the sum of the diagonal entries of Q^2 , $\|Q\|^2$ is the sum of the squares of the entries of Q , and we have the *Cauchy-Schwarz inequality* $|\text{trace}(QZ)| \leq \|Q\| \|Z\|$.

The *determinant* of Q is the product $\det Q$ of the eigenvalues of Q . Then $Q > 0$ implies $\det Q > 0$ and the determinant is *multiplicative*,

$$Z > 0 \text{ and } S > 0 \quad \implies \quad \det(SZS) = \det S \cdot \det Z \cdot \det S.$$

For this restricted setting, multiplicativity is a consequence of the *matrix determinant lemma* (Hijab 2026, Exercise 3.2.18). By reducing to eigenvalues, for Z symmetric, one has

$$1 + |\det(I + Z) - 1| \leq \exp\left(\sqrt{d} \|Z\|\right).$$

This implies $\det Q$ is continuous at $Q = I$. Writing $Q = S^2$ with $S > 0$, multiplicativity extends this to continuity at all $Q > 0$.

References

- Bishop, C. M. (2006), *Pattern Recognition and Machine Learning*, Information Science and Statistics, Springer, New York.
- Cover, T. M. & Thomas, J. A. (2006), *Elements of Information Theory*, 2nd edn, Wiley-Interscience.
- Cramér, H. (1946), *Mathematical Methods of Statistics*, Princeton University Press.
- Dembo, A. & Zeitouni, O. (1998), *Large Deviations Techniques and Applications*, Vol. 38 of *Stochastic Modelling and Applied Probability*, 2nd edn, Springer, New York.
- Donsker, M. D. & Varadhan, S. R. S. (1975), ‘Asymptotic evaluation of certain markov process expectations for large time, I’, *Communications on Pure and Applied Mathematics* **28**(1), 1–47.
- Doob, J. L. (1934), ‘Probability and statistics’, *Transactions of the American Mathematical Society* **36**, 759–775.
- Fisher, R. A. (1922), ‘On the mathematical foundations of theoretical statistics’, *Philosophical Transactions of the Royal Society of London, Series A* **222**, 309–368.
- Hijab, O. (2026), *Math for Data Science*, Springer, Cham. Corrected printing.
- Kullback, S. (1959), *Information Theory and Statistics*, Wiley, New York.
- Murphy, K. P. (2022), *Probabilistic Machine Learning: An Introduction*, MIT Press.
- Samson, E. W. (1951), Fundamental natural concepts of information theory, Report E5079, Communications Laboratory, Electronics Research Division, Air Force Cambridge Research Center, Cambridge, MA.
- Sanov, I. N. (1957), ‘On the probability of large deviations of random variables’, *Matematicheskii Sbornik* **42**, 11–44. (In Russian). English translation in *Selected Translations in Mathematical Statistics and Probability*, Vol. 1 (1961), pp. 213–244.
- Schervish, M. J. (1995), *Theory of Statistics*, Springer, New York.
- Shannon, C. E. (1948), ‘A mathematical theory of communication’, *Bell System Technical Journal* **27**(3), 379–423.
- Tribus, M. (1961), *Thermostatistics and Thermodynamics: An Introduction to Energy, Information and States of Matter, With Engineering Applications*, D. Van Nostrand Company, Princeton, NJ.
- van der Vaart, A. W. (1998), *Asymptotic Statistics*, Vol. 3 of *Cambridge Series in Statistical and Probabilistic Mathematics*, Cambridge University Press, Cambridge.
- Wald, A. (1949), ‘Note on the consistency of the maximum likelihood estimate’, *Annals of Mathematical Statistics* **20**(4), 595–601.