

Doob's Proof of MLE Strong Consistency: A Surprisal and Information Perspective

Omar Hijab

Department of Mathematics, Temple University

2026

Abstract

We give a concise, self-contained exposition of Doob's proof of strong maximum likelihood consistency, formulated in terms of surprisal and relative information. This formulation clarifies the roles of identifiability, properness, convergence, and continuity, and highlights that consistency holds without uniform convergence of the log-likelihood. We then apply the result to two special cases, i.i.d. sequences and stationary ergodic Markov processes, illustrated by an i.i.d. Gaussian sequence and an autoregressive Gaussian process.

Keywords: maximum likelihood estimator, true parameter, relative information, autoregressive, markov process, multivariate normal

1 Introduction

The consistency of the [Fisher \(1922\)](#) maximum likelihood estimator (MLE) is a classical result discovered more than a century ago. This method is typically treated as a special case of M -estimators, and derived via techniques dating back to [Cramér \(1946\)](#) and [Wald \(1949\)](#). A modern treatment is [van der Vaart \(1998\)](#). Because of its intuitive appeal and broad applicability, the method is a dominant tool of statistical inference.

In teaching this material, I found the [Doob \(1934\)](#) strong consistency proof of MLE convergence more motivated and accessible to students, particularly for students already familiar with information and entropy as part of a data science or machine learning course. The fact that the proof covers general parameter spaces and general state spaces with essentially no extra effort only adds to its appeal.

The contribution of this note is expository and pedagogical: We give a formulation of Doob's consistency argument in surprisal/information terminology, in sufficient generality to encompass i.i.d. sequences and Markov processes as special cases. This is done without assuming uniform convergence. To highlight the applicability of our assumptions, we also work out two examples, i.i.d. Gaussian sequences and autoregressive Gaussian processes.

2 Surprisal and Information

A random variable x is sampled repeatedly, resulting in samples¹ $x_1, x_2, \dots$. We assume the density $p(x, \theta^*)$ of x is one of a parametric family $p(x, \theta)$ of positive functions, and we wish to recover θ^* from $x_1, x_2, \dots$. While $p(x, \theta)$ may be chosen arbitrarily, based on prior knowledge, we have no control over the underlying expectation E_* . Given this, how do we recover θ^* from the process $x_1, x_2, \dots$?

Define the *surprisal* [Samson \(1951\)](#), [Tribus \(1961\)](#)

$$S(x, \theta) = \log(1/p(x, \theta)).$$

This terminology is intuitive: low probability equals high surprisal.

Assume $S(x, \theta)$ is integrable for all θ . Then it is natural to call a minimizer θ^* of the *expected surprisal* $E_*(S(x, \theta))$ a *true parameter*, and the corresponding density $p(x, \theta^*)$ a *true density*. Equivalently, given θ^* and θ , the *relative information* is

$$I(\theta^*, \theta) = E_*(S(x, \theta)) - E_*(S(x, \theta^*)). \quad (1)$$

The relative information is the excess expected surprisal of θ over θ^* . Then θ^* is a true parameter iff $I(\theta^*, \theta) \geq 0$ for all θ .

While $p(x, \theta)$ have so far been arbitrary positive functions, we now specialize to the case where $p(x, \theta)$ are densities in the usual sense. We say θ^* is a *reference parameter* if

$$E_* \left(\frac{p(x, \theta)}{p(x, \theta^*)} \right) = 1 \quad \text{for all } \theta.$$

Then the corresponding density $p(x, \theta^*)$ is a *reference density*.

Let θ^* be a reference parameter, and define E and E_θ by

$$E_*(f(x)) = E(p(x, \theta^*)f(x)) \quad \text{and} \quad E_\theta(f(x)) = E(p(x, \theta)f(x)).$$

Then the relative information takes the familiar form [Kullback & Leibler \(1951\)](#), [Sanov \(1957\)](#), [Donsker & Varadhan \(1975\)](#), [Dembo & Zeitouni \(1998\)](#)

$$I(\theta^*, \theta) = E \left(p(x, \theta^*) \log \left(\frac{p(x, \theta^*)}{p(x, \theta)} \right) \right), \quad (2)$$

and $E_\theta(1) = 1$ for all θ . Since $j(p) = \log(1/p)$ is convex, by Jensen's inequality,

$$I(\theta^*, \theta) = E_* \left(j \left(\frac{p(x, \theta)}{p(x, \theta^*)} \right) \right) \geq j \left(E_* \left(\frac{p(x, \theta)}{p(x, \theta^*)} \right) \right) = j(E_\theta(1)) = j(1) = 0, \quad \text{for all } \theta.$$

¹We use the same letter (e.g. x) to denote both a random variable and a sample of it.

It follows that a reference density is a true density.

If θ is any other true parameter, then $I(\theta^*, \theta) = 0$, hence, by strict convexity of $j(p)$, there is a scalar t such that $p(x, \theta) = t \cdot p(x, \theta^*)$. Since θ^* is a reference parameter, $t = 1$, hence $p(x, \theta) = p(x, \theta^*)$: every true density equals the reference density. It follows that there is at most one reference density; when there is a reference density, it is the unique true density.

We say parameters θ, θ' *share a density* if $p(x, \theta) = p(x, \theta')$. If parameters do not share densities, there is at most one reference parameter; moreover, when there is a reference parameter, it is the unique true parameter.

The simplest example is as follows. A vector $\theta = (\theta_1, \theta_2, \dots, \theta_d)$ is a probability vector if its components are nonnegative and sum to one. Suppose $\theta^* = (\theta_1^*, \theta_2^*, \dots, \theta_d^*)$ is a probability vector and a random variable x satisfies $P_*(x = i) = \theta_i^*$, $i = 1, 2, \dots, d$. Then the mass function of x is $p(x, \theta^*) = \theta_x^*$. If θ is any other probability vector, the surprisal is $S(x, \theta) = \log(1/\theta_x)$, and

$$I(\theta^*, \theta) = \sum_{i=1}^d \theta_i^* \log(\theta_i^* / \theta_i). \quad (3)$$

In the form (2), $I(\theta^*, \theta)$ is more commonly known as Kullback-Leibler divergence [van der Vaart \(1998\)](#) or relative entropy [Cover & Thomas \(2006\)](#), [Dembo & Zeitouni \(1998\)](#) or both [Bishop \(2006\)](#), [Murphy \(2022\)](#). It appears as the rate function in Sanov's theorem [Sanov \(1957\)](#). It is natural to regard $I(\theta^*, \theta)$ as information, following some authors [Kullback \(1959\)](#), [Schervish \(1995\)](#), since information and entropy are conceptual opposites: the [Shannon \(1948\)](#) entropy $-\sum_{i=1}^d p_i \log p_i$ is concave, while (3) is convex.

Returning to the general setting, the *likelihood* and (empirical) *mean surprisal* are

$$L_n(\theta) = \prod_{k=1}^n p(x_k, \theta), \quad S_n(\theta) = \frac{1}{n} \log(1/L_n(\theta)) = \frac{1}{n} \sum_{k=1}^n S(x_k, \theta).$$

The mean surprisal is also called the negative log-likelihood.

By definition, θ^* minimizes the expected surprisal. The MLE method rests on the convergence of the mean surprisal to the expected surprisal for large samples. While this is not the case in general, this convergence is guaranteed in two classical cases: an i.i.d. sequence [Fisher \(1922\)](#), and a stationary ergodic Markov process [Anderson & Goodman \(1957\)](#), [Billingsley \(1961\)](#). In the former case, the convergence is a consequence of the law of large numbers; in the latter case, of the ergodic theorem.

Because of this, it is natural to estimate θ^* by a minimizer of $S_n(\theta)$ or, equivalently, a maximizer of $L_n(\theta)$: For each $n \geq 1$, a *maximum likelihood estimator (MLE)*, if it exists, is a parameter $\hat{\theta}_n$, depending on the samples, maximizing $L_n(\theta)$ over all θ .

In what follows, only E_* , $p(x, \theta)$, and true parameters enter; E , E_θ , and reference parameters play no role.

3 Consistency

In addition to integrability of $S(x, \theta)$, we make four assumptions. The first is

A (Identifiability). *There is at most one true parameter.*

A follows if the expected surprisal is strictly convex. Also, as we saw above, **A** follows if there is a reference parameter and parameters do not share densities.

A sequence of parameters *subconverges* to θ if a subsequence converges to θ . If a sequence subconverges to θ , θ is a *limit point of the sequence*. A sequence has exactly one limit point θ iff the sequence converges to θ . A subset K of the parameter space is *compact* if every sequence in K has at least one limit point in K .

A function $F(\theta)$ is *proper* if its sublevel sets $F(\theta) \leq c$ are compact subsets of the parameter space, for every level c . Since a proper continuous function has a minimizer and $\hat{\theta}_n$ should minimize $S_n(\theta)$, this leads to the second

B (Properness). *There is an $N \geq 1$, depending on the samples, and a proper $F(\theta)$, not depending on the samples, such that, with probability one,*

$$F(\theta) \leq S_n(\theta), \quad \text{for all } \theta \text{ and } n \geq N. \quad (4)$$

When the parameter space is compact, the proof below shows assumption **B** is superfluous. The assumption crucial to the method is the third

C (Convergence). *For every integrable $g(x)$*

$$\frac{1}{n} \sum_{k=1}^n g(x_k) \rightarrow E_*(g(x)), \quad \text{as } n \rightarrow \infty, \quad \text{with probability one.} \quad (5)$$

We will need the convergence (5) with $g(x) = S(x, \theta)$ simultaneously for all θ , in the sense

$$S_n(\theta) \rightarrow E_*(S(x, \theta)), \quad \text{as } n \rightarrow \infty, \quad \text{for all } \theta, \text{ with probability one.} \quad (6)$$

We emphasize this is not uniform convergence, since the rate of convergence may vary with parameters θ . For this we need the fourth

D (Continuity). *The parameter space is separable and there is a continuous $\delta(\theta, \theta')$ and an integrable non-negative $g(x)$ satisfying*

$$|S(x, \theta) - S(x, \theta')| \leq \delta(\theta, \theta')g(x) \quad \text{and} \quad \delta(\theta, \theta) = 0. \quad (7)$$

Under **C**, **D**, a standard separability argument establishes (6). Under **D**, the expected surprisal $E_*(S(x, \theta))$ is continuous in θ . If moreover **B** holds, then by sending $n \rightarrow \infty$ in (4), $F(\theta) \leq E_*(S(x, \theta))$ for all θ , hence the expected surprisal is proper. Hence there is a unique true parameter under **A**, **B**, **C**, and **D**. Similarly, under **B**, **D**, the mean surprisal is continuous in θ and proper, hence an MLE $\hat{\theta}_n$ exists for $n \geq N$, with probability one.

These assumptions are particularly tractable for *exponential families* Cover & Thomas (2006), Murphy (2022), Schervish (1995), van der Vaart (1998), in which $S(x, \theta)$ is essentially a sum of products of the form $f(\theta)g(x)$. In §4, we verify these assumptions for a multivariate normal i.i.d process and an autoregressive Gaussian process.

In (6), there is no presumption of uniform convergence. This is a key feature of Doob's proof. The argument instead combines properness, convergence (6), and the continuity estimate in (7) to control limit points of the MLE sequence.

Theorem (Consistency). *Let $x_1, x_2, \dots$ be a process, and let $p(x, \theta)$ be a parametric family of positive functions. Under the above assumptions,*

1. *there is a unique true parameter θ^* ,*
2. *with probability one, an MLE sequence $\hat{\theta}_n$ exists for sufficiently large n ,*
3. *with probability one, any MLE sequence converges to the true parameter, $\hat{\theta}_n \rightarrow \theta^*$.*

Proof. The first two items were derived above. For the third item, with $g(x)$ as in (7), fix samples $x_1, x_2, \dots$ for which (4), (5), and (6) hold. We show $\hat{\theta}_n \rightarrow \theta^*$ for these samples. By (4),

$$F(\hat{\theta}_n) \leq S_n(\hat{\theta}_n) \leq S_n(\theta^*), \quad \text{for } n \geq N.$$

By (6), the right side of this last display is bounded as $n \rightarrow \infty$. Since $F(\theta)$ is proper, the sequence $\hat{\theta}_n$ lies in a compact set in the parameter space, and thus has limit points.

Suppose θ is a limit point of $\hat{\theta}_n$. Then $\hat{\theta}_n$ subconverges to θ . By (7),

$$S_n(\theta) - S_n(\theta^*) \leq S_n(\theta) - S_n(\hat{\theta}_n) \leq \delta(\theta, \hat{\theta}_n) \cdot \frac{1}{n} \sum_{k=1}^n g(x_k). \quad (8)$$

Since the left side of (5) is bounded as $n \rightarrow \infty$, and $\delta(\theta, \hat{\theta}_n)$ subconverges to zero, the right side of (8) subconverges to zero. By (1) and (6), the left side of (8) converges to $I(\theta^*, \theta)$, hence $I(\theta^*, \theta) \leq 0$. But $I(\theta^*, \theta) \geq 0$ since θ^* is a true parameter, hence $I(\theta^*, \theta) = 0$. Since there is only one true parameter, $\theta = \theta^*$. Thus θ^* is the only limit point, hence $\hat{\theta}_n \rightarrow \theta^*$ as $n \rightarrow \infty$. $\square$

4 Examples

The first example is MLE for an i.i.d. Gaussian sequence, and the second example is MLE for an autoregressive Gaussian process. In the first example, the sequence is denoted $z_1, z_2, \dots$ —anticipating its role as the innovations driving the autoregressive process of the second example—reserving $x_1, x_2, \dots$ for that Markov process.

4.1 I.I.D. Gaussian Sequence

Let μ be in $\mathbf{R}^d$, Q a positive definite² $d \times d$ matrix, and let $v \cdot w$ be the dot product in $\mathbf{R}^d$. With z in $\mathbf{R}^d$, the *normal density with parameter* $\theta = (\mu, Q)$ is $p(z, \theta) = e^{-S(z, \theta)}$ where

$$S(z, \theta) = \frac{1}{2} \log \det(2\pi Q) + \frac{1}{2} (z - \mu) \cdot Q^{-1} (z - \mu).$$

This defines the parametric family. A random variable z in $\mathbf{R}^d$ is *normally distributed with parameter* $\theta^* = (\mu^*, Q^*)$ if its moment-generating function is

$$E_*(e^{v \cdot z}) = \exp \left(v \cdot \mu^* + \frac{1}{2} v \cdot Q^* v \right). \quad (9)$$

Let $z_1, z_2, \dots$ be independent and identically distributed copies of z . We show $S(z, \theta) = \log(1/p(z, \theta))$ is integrable and θ^* is the true parameter, and we verify [A](#), [B](#), [C](#), and [D](#). Assumption [C](#) is verified, as it is the law of large numbers (LLN).

Let $\theta' = (\mu', Q')$, $g(z) = 1 + |z|^2$ and let $\delta(\theta, \theta')$ equal half of

$$|\log \det Q - \log \det Q'| + |Q^{-1} \mu - (Q')^{-1} \mu'| + \|Q^{-1} - (Q')^{-1}\| + |\mu \cdot Q^{-1} \mu - \mu' \cdot (Q')^{-1} \mu'|.$$

Then $\delta(\theta, \theta')$ is continuous. By standard manipulations ([7](#)) follows, establishing [D](#).

Proposition 1. *For $Q^* > 0$ and c scalar, let $\alpha = e^{-2c-1} \det(2\pi e Q^*)$ and*

$$F_*(Q, Q^*) = \frac{1}{2} \log \det(2\pi Q) + \frac{1}{2} \text{trace}(Q^{-1} Q^*), \quad Q > 0.$$

Then $F_(Q, Q^*)$ is minimized iff $Q = Q^*$, and $F_*(Q, Q^*) \leq c$ implies $\alpha Q^* \leq Q \leq Q^*/\alpha$.*

Proof. Let $h(\lambda) = \log \lambda + 1/\lambda$, $\lambda > 0$. Then $h(\lambda)$ is minimized iff $\lambda = 1$ and $h(\lambda)$ is proper in the sense $h(\lambda) \leq \log(1/\alpha)$ implies $\alpha \leq \lambda \leq 1/\alpha$. Write $Q^* = S^2$ with $S > 0$, and define Z by $Q = SZS$. Suppose $F_*(Q, Q^*) \leq c$ and let $\lambda_1, \lambda_2, \dots, \lambda_d$ be the eigenvalues of Z . Then the proposition follows from

$$\log(1/\alpha) - 1 \geq 2F_*(Q, Q^*) - 2F_*(Q^*, Q^*) = \log \det Z + \text{trace}(Z^{-1} - I) = \sum_{i=1}^d (h(\lambda_i) - 1). \quad \square$$

Using the moment generating function,

$$E_*(S(z, \theta)) = F_*(Q, Q^*) + \frac{1}{2} (\mu^* - \mu) \cdot Q^{-1} (\mu^* - \mu).$$

Since $S(z, \theta)$ is bounded below for each θ , $S(z, \theta)$ is integrable. By the proposition, (μ^*, Q^*) is the unique minimizer of the expected surprisal, establishing [A](#).

Now let $0 < \epsilon < 1/2$ and

$$F_\epsilon(\theta) = F_*(Q, (1 - 2\epsilon)Q^*) + \frac{1}{2} (1 - \epsilon) (\mu^* - \mu) \cdot Q^{-1} (\mu^* - \mu), \quad \theta = (\mu, Q).$$

By the proposition with $(1 - 2\epsilon)Q^*$ replacing Q^* , if $F_\epsilon(\theta) \leq c$, then Q is confined to a compact subset of positive definite matrices and, subsequently, μ is confined to a compact subset of $\mathbf{R}^d$. Hence $F_\epsilon(\theta)$ is proper.

²Linear algebra is reviewed in [Appendix A](#).

With

$$Q_n(\mu) = \frac{1}{n} \sum_{k=1}^n (z_k - \mu) \otimes (z_k - \mu), \quad \hat{\mu}_n = \frac{1}{n} \sum_{k=1}^n z_k, \quad \hat{Q}_n = Q_n(\hat{\mu}_n),$$

let $\hat{\theta}_n = (\hat{\mu}_n, \hat{Q}_n)$. Then $\hat{\mu}_n$ and $\hat{Q}_n$ are the usual sample mean and the biased sample covariance.

By the LLN, with probability one, the matrix sequences

$$(\mu^* - \hat{\mu}_n) \otimes (\mu^* - \hat{\mu}_n) \rightarrow 0, \quad Q^* - Q_n(\mu^*) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

If Q_n is any sequence of symmetric matrices with $Q_n \rightarrow 0$, then $\|Q_n\| \rightarrow 0$. Since $Q_n \leq \|Q_n\| I$, given any $Q > 0$, there is $N \geq 1$ such that $Q_n \leq Q$ for $n \geq N$. By choosing $Q = \epsilon^2 Q^*$ and $Q = \epsilon Q^*$, it follows there is $N^* \geq 1$, depending on the samples, such that, with probability one,

$$(\mu^* - \hat{\mu}_n) \otimes (\mu^* - \hat{\mu}_n) \leq \epsilon^2 Q^*, \quad Q_n(\mu^*) \geq (1 - \epsilon) Q^*, \quad \text{for } n \geq N^*. \quad (10)$$

Let

$$f_0(\mu) = Q^* + (\mu^* - \mu) \otimes (\mu^* - \mu), \quad f_\epsilon(\mu) = (1 - \epsilon) f_0(\mu) - \epsilon Q^*.$$

Then $F_\epsilon(\theta) = F_*(Q, f_\epsilon(\mu))$. With $w = \hat{\mu}_n - \mu^*$ and $v = \mu - \mu^*$, using (17) in the appendix,

$$v \otimes w + w \otimes v \leq \epsilon(\mu^* - \mu) \otimes (\mu^* - \mu) + \frac{1}{\epsilon}(\hat{\mu}_n - \mu^*) \otimes (\hat{\mu}_n - \mu^*) \leq \epsilon f_0(\mu), \quad \text{for all } \mu \text{ and } n \geq N^*.$$

Writing $z_k - \mu = (z_k - \mu^*) - v$, we have $Q_n(\mu) = Q_n(\mu^*) - (v \otimes w + w \otimes v) + v \otimes v$. Combined with the above inequalities, this leads to

$$Q_n(\mu) \geq f_\epsilon(\mu), \quad \text{for all } \mu \text{ and } n \geq N^*.$$

In particular, since $f_\epsilon(\mu) \geq (1 - 2\epsilon) Q^*$, we have shown $\hat{Q}_n > 0$ for $n \geq N^*$.

Since $S_n(\theta) = F_*(Q, Q_n(\mu))$ and $Q_n(\mu) \geq \hat{Q}_n$ by least squares, by the proposition,

$$S_n(\theta) = F_*(Q, Q_n(\mu)) \geq F_*(Q, \hat{Q}_n) \geq F_*(\hat{Q}_n, \hat{Q}_n) = S_n(\hat{\theta}_n), \quad \text{for all } \theta \text{ and } n \geq N^*.$$

Thus $\hat{\theta}_n$ is the MLE for $n \geq N^*$, and

$$S_n(\theta) = F_*(Q, Q_n(\mu)) \geq F_*(Q, f_\epsilon(\mu)) = F_\epsilon(\theta), \quad \text{for all } \theta \text{ and } n \geq N^*.$$

This establishes (4) and completes the derivation of B.

4.2 Autoregressive Gaussian Process

Let A be a $d \times d$ matrix. With x_0, x_1 in $\mathbf{R}^d$, the *transition density with parameter* $\theta = (\mu, A, Q)$ is $p(x_0, x_1, \theta) = e^{-S(x_0, x_1, \theta)}$ where

$$S(x_0, x_1, \theta) = \frac{1}{2} \log \det(2\pi Q) + \frac{1}{2} (x_1 - Ax_0 - \mu) \cdot Q^{-1} (x_1 - Ax_0 - \mu).$$

This defines the parametric family.

Let $\theta^* = (\mu^*, A^*, Q^*)$, and assume A^* is stable (Appendix A). Define

$$c^* = (I - A^*)^{-1} \mu^*, \quad L^* = \sum_{n=0}^{\infty} (A^*)^n Q^* ((A^*)^t)^n, \quad (11)$$

both well-defined by stability of A^* ; as shown below, these are the values of the mean and covariance of x_0 needed to make $x_1, x_2, \dots$ (defined via the recurrence (12) below) stationary.

Let z be normally distributed with parameter (μ^*, Q^*) , and let x_0 be normally distributed with parameter (c^*, L^*) . The *innovations process* $z_1, z_2, \dots$ consists of i.i.d. copies of z , assumed independent of x_0 . The

autoregressive Gaussian process (of order one), with parameter $\theta^* = (\mu^*, A^*, Q^*)$, is the process $x_1, x_2, \dots$ generated by the recurrence

$$x_n = A^* x_{n-1} + z_n, \quad n \geq 1. \quad (12)$$

Then z_n is independent of $x_1, \dots, x_{n-1}$, and $x_1, x_2, \dots$ is a Gaussian Markov process.

Let $(x_0, x_1), (x_1, x_2), \dots$, be the *pair process*. We show $S(x_0, x_1, \theta)$ is integrable and θ^* is the true parameter, and we verify [A](#), [B](#), [C](#), and [D](#) for the pair process, assuming stability of A^* .

For the pair process, the likelihood and mean surprisal are

$$L_n(\theta) = \prod_{k=1}^n p(x_{k-1}, x_k, \theta), \quad S_n(\theta) = \frac{1}{n} \log(1/L_n(\theta)) = \frac{1}{n} \sum_{k=1}^n S(x_{k-1}, x_k, \theta).$$

We verify [C](#) first. Because the series in (11) converges, we can go backward in time and present x_0 more explicitly. Let $z_0, z_{-1}, z_{-2}, \dots$ be a Gaussian i.i.d. process, with parameter (μ^*, Q^*) , independent of $z_1, z_2, \dots$. Then $z_0, z_{\pm 1}, z_{\pm 2}, \dots$ is the *extended innovations* process. If we set

$$x_0 \equiv \sum_{k=0}^{\infty} (A^*)^k z_{-k},$$

then by (11) x_0 is normally distributed with parameter (c^*, L^*) and independent of $z_1, z_2, \dots$. Moreover iterating (12) and combining,

$$x_n = (A^*)^n x_0 + \sum_{k=1}^n (A^*)^{n-k} z_k = \sum_{k=-\infty}^n (A^*)^{n-k} z_k, \quad n \geq 1. \quad (13)$$

Thus $x_1, x_2, \dots$ is the image of the extended innovations process under a shift-invariant map (see Appendix [B](#) for definitions). Since the extended innovations process is i.i.d., it is stationary and ergodic, hence $x_1, x_2, \dots$ is stationary and ergodic. Since $x_1, x_2, \dots$ is stationary, by the ergodic theorem (Appendix [B](#)), [Billingsley \(1961\)](#),

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n g(x_{k-1}, x_k) \quad (14)$$

exists and is an invariant function of $x_1, x_2, \dots$, with probability one. Since $x_1, x_2, \dots$ is ergodic, this limit is constant with probability one. Taking expectations of (14), the constant equals $E_*(g(x_0, x_1))$. This establishes [C](#) for the pair process.

For [D](#), with $\theta' = (\mu', A', Q')$, replace the previous $g(z)$ by $g(x_0, x_1) = 1 + |x_0|^2 + |x_1|^2$, and augment the previous $\delta(\theta, \theta')$ by three additional terms, equal to half of

$$\|Q^{-1}A - (Q')^{-1}A'\| + |A^t Q^{-1} \mu - (A')^t (Q')^{-1} \mu'| + \|A^t Q^{-1} A - (A')^t (Q')^{-1} (A')\|.$$

This establishes [D](#).

For [A](#), the mean and variance of $x_1 - Ax_0 - \mu = z_1 + (A^* - A)x_0 - \mu$ are

$$c^* - Ac^* - \mu = (A^* - A)c^* + (\mu^* - \mu) \quad \text{and} \quad Q^* + (A^* - A)L^*(A^* - A)^t.$$

Here we used $c^* - A^*c^* = \mu^*$ ([11](#)). If we let $v = (\mu^* - \mu) + (A^* - A)c^*$ and

$$f_0(\mu, A) = Q^* + (A^* - A)L^*(A^* - A)^t + v \otimes v,$$

we conclude

$$E_*(S(x_0, x_1, \theta)) = F_*(Q, f_0(\mu, A)) \equiv F_0(\mu, A, Q) \quad (15)$$

is the sum of three terms. Since the first term is $F_*(Q, Q^*)$, $Q = Q^*$ is its unique minimizer. Since $L^* > 0$, $A = A^*$ is the unique minimizer of the second term, hence $(\mu, A) = (\mu^*, A^*)$ is the unique minimizer of the sum of the second and third terms. This establishes [A](#). Moreover, since the second and third terms are nonnegative, inspecting $F_0(\mu, A, Q) \leq c$ implies Q , then A , then μ are confined to compact subsets of their domains, hence $F_0(\mu, A, Q)$ is proper.

Our last item is establishing B. With the previously defined $\hat{\mu}_n$, here we also need

$$\hat{c}_n = \frac{1}{n} \sum_{k=1}^n x_{k-1}, \quad \hat{L}_n = \frac{1}{n} \sum_{k=1}^n (x_{k-1} - c^*) \otimes (x_{k-1} - c^*), \quad \hat{C}_n = \frac{1}{n} \sum_{k=1}^n (z_k - \mu^*) \otimes (x_{k-1} - c^*).$$

By the ergodic theorem, with probability one, $\hat{c}_n$ and $\hat{L}_n$ converge to c^* and L^* , and $\hat{C}_n$ converges to $E_*((z_1 - \mu^*) \otimes (x_0 - c^*))$ which vanishes by independence of z_1 and x_0 . Let

$$Q_n(\mu, A) = \frac{1}{n} \sum_{k=1}^n (x_k - Ax_{k-1} - \mu) \otimes (x_k - Ax_{k-1} - \mu).$$

Then $S_n(\theta) = F_*(Q, Q_n(\mu, A))$, and

$$x_k - Ax_{k-1} - \mu = z_k + (A^* - A)x_{k-1} - \mu = (z_k - \mu^*) + (A^* - A)(x_{k-1} - c^*) + v.$$

The trinomial expansion of $Q_n(\mu, A)$ is the sum of three square terms and three cross terms. The sums of the square terms are $Q_n(\mu^*, A^*)$, $v \otimes v$, and $(A^* - A)\hat{L}_n(A^* - A)^t$. With $w = \hat{\mu}_n - \mu^*$ and $u = (A^* - A)(\hat{c}_n - c^*)$, the sums of the cross terms are $u \otimes v + v \otimes u$, $w \otimes v + v \otimes w$, and $(A^* - A)\hat{C}_n^t + \hat{C}_n(A^* - A)^t$.

For $0 < \epsilon < 1/3$, set

$$f_\epsilon(\mu, A) = (1 - 3\epsilon)Q^* + (1 - 3\epsilon)(A^* - A)L^*(A^* - A)^t + (1 - 2\epsilon)v \otimes v,$$

and $F_\epsilon(\theta) = F_*(Q, f_\epsilon(\mu, A))$. Then, as before, $F_\epsilon(\theta)$ is proper.

By the ergodic theorem, there is $N^* \geq 1$, depending on the samples, such that

$$\begin{aligned} Q_n(\mu^*, A^*) &\geq (1 - \epsilon)Q^*, & \hat{L}_n &\geq (1 - \epsilon)L^*, & (\hat{c}_n - c^*) \otimes (\hat{c}_n - c^*) &\leq \epsilon^2 L^*, \\ \hat{C}_n(L^*)^{-1}\hat{C}_n^t &\leq \epsilon^2 Q^*, & (\hat{\mu}_n - \mu^*) \otimes (\hat{\mu}_n - \mu^*) &\leq \epsilon^2 Q^*. \end{aligned}$$

for $n \geq N^*$, with probability one. Fix samples $x_1, x_2, \dots$ for which these hold. We establish (4) for these samples and $F(\theta) = F_\epsilon(\theta)$.

Using (16) and (17), this implies the sum of the sums of the cross terms is bounded by $2\epsilon f_0(\mu, A)$ for $n \geq N^*$, hence

$$Q_n(\mu, A) \geq f_\epsilon(\mu, A), \quad \text{for all } \mu, A \text{ and } n \geq N^*.$$

Since $S_n(\theta) = F_*(Q, Q_n(\mu, A))$, (4) follows.

The MLE $\hat{\theta}_n$ is obtained by doing least squares of $Q_n(\mu, A)$ over (μ, A) . This returns the (multivariate) Yule–Walker estimator Yule (1927), Walker (1931), which for the Gaussian case coincides with the MLE.

5 Disclosures

The author reports no funding was obtained for this work and there are no competing interests to declare. The author used a generative AI tool (Anthropic’s *Claude Sonnet*) to assist with language editing and bibliographic searches. All AI-assisted content was reviewed, verified, and edited by the author, who takes full responsibility for the content, accuracy, and originality of this work.

A Linear Algebra

This is a review of the linear algebra used above. Let Q and Z be symmetric matrices. We say Q is *positive definite*, and write $Q > 0$, if $v \cdot Qv > 0$ for all vectors $v \neq 0$. We say Q is *nonnegative definite*, and write $Q \geq 0$, if $v \cdot Qv \geq 0$ for all vectors v . We write $Q \geq Z$ if $Q - Z \geq 0$. Then $aI \leq Q \leq bI$ iff the eigenvalues λ of Q satisfy $a \leq \lambda \leq b$. In particular, if $Q \geq 0$, the eigenvalues satisfy $\lambda \geq 0$, and, if $Q > 0$, the eigenvalues satisfy $\lambda > 0$. From this, if $Q \geq 0$, there is $S \geq 0$ satisfying $Q = S^2$, and $S > 0$ if $Q > 0$.

Given matrices V, W , let $A = \sqrt{\epsilon}V + W/\sqrt{\epsilon}N$. Then $AA^t \geq 0$ implies

$$VW^t + WV^t \leq \epsilon VV^t + \frac{1}{\epsilon}WW^t. \quad (16)$$

Given vectors $v = (s_1, s_2, \dots, s_d)$ and $w = (t_1, t_2, \dots, t_d)$, the *tensor product* $v \otimes w$ is the matrix whose ij -th entry is $s_i t_j$. Taking $V = v$ and $W = w$ in (16) implies

$$v \otimes w + w \otimes v \leq \epsilon v \otimes v + \frac{1}{\epsilon} w \otimes w. \quad (17)$$

If $u_1, u_2, \dots, u_d$ is any orthonormal basis and $e_1, e_2, \dots, e_d$ is the standard one-hot encoded basis, then $\sum_{i=1}^d e_i \otimes e_i$ and $\sum_{i=1}^d u_i \otimes u_i$ both equal I .

For any square matrix A , the *trace* of A is the sum $\text{trace}(A) = \sum_{i=1}^d e_i \cdot A e_i$ of the diagonal entries of A . Then $\text{trace}(QZ) = \text{trace}(ZQ)$ and $\text{trace}(Q(u \otimes u)) = u \cdot Qu$ follow immediately. If $u_1, u_2, \dots, u_d$ is an orthonormal basis of eigenvectors of Q , $\text{trace}(Q)$ is the sum $\sum_{i=1}^d u_i \cdot Qu_i$ of the eigenvalues of Q , hence $\text{trace}(Q) \geq 0$ if $Q \geq 0$. Since $Z \geq 0$ and S symmetric imply $SZS \geq 0$, $\text{trace}(QZ) = \text{trace}(S^2 Z) = \text{trace}(SZS) \geq 0$. It follows the trace is *monotone*,

$$Q \geq S \text{ and } Z \geq 0 \quad \implies \quad \text{trace}(QZ) \geq \text{trace}(SZ).$$

The *norm squared* $\|Q\|^2 = \text{trace}(Q^2)$ is the sum of the squares of the eigenvalues of Q , hence $Q \leq \|Q\| I$. Since $\|Q\|^2$ is the sum of the diagonal entries of Q^2 , $\|Q\|^2$ is the sum of the squares of the entries of Q , and we have $|Ax| \leq \|A\| |x|$ and the *Cauchy-Schwarz inequality* $|\text{trace}(QZ)| \leq \|Q\| \|Z\|$.

The *determinant* of Q is the product $\det Q$ of the eigenvalues of Q . Then $Q > 0$ implies $\det Q > 0$ and the determinant is *multiplicative*,

$$Z > 0 \text{ and } S > 0 \quad \implies \quad \det(SZS) = \det S \cdot \det Z \cdot \det S.$$

For this restricted setting, multiplicativity is a consequence of the *matrix determinant lemma* (Hijab 2026, Exercise 3.2.18). By reducing to eigenvalues, for Z symmetric, one has

$$1 + |\det(I + Z) - 1| \leq \exp(\sqrt{d} \|Z\|).$$

This implies $\det Q$ is continuous at $Q = I$. Writing $Q = S^2$ with $S > 0$, multiplicativity extends this to continuity at all $Q > 0$.

A matrix A is *stable* if its (possibly complex) eigenvalues satisfy $|\lambda| < 1$. If $\rho(A) = \max |\lambda|$ is the radius of the smallest disk containing the eigenvalues, Gelfand's formula states $\rho(A) = \lim_{n \rightarrow \infty} \|A^n\|^{1/n}$. If $\rho(A) < r < 1$, then $\|A^n\| \leq r^n$ eventually, so the geometric series $\sum A^n$ converges, and hence $(I - A)^{-1}$ exists. Similarly, the Lyapunov series in (11) also converges.

B Ergodicity

For

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n g(x_k) = E_*(g(x_1)), \quad \text{with probability one,} \quad (5)$$

we are interested in two cases, when the process is an i.i.d. sequence, and when it is a Markov process. In the former case, (5) is the strong law of large numbers (LLN). In the latter case, it is a consequence of the ergodic theorem.

The first step in deconstructing (5) is to take the expectation of its left side. For this to equal the right side, it is simplest to assume the marginals $E_*(g(x_n))$ are all equal: A process $x_1, x_2, \dots$ is stationary if expectations are unchanged under the shift $x_n \rightarrow x_{n+1}$. More precisely, the process is *stationary* if, for any function $G(x_1, x_2, \dots)$ of the process,

$$E_*(G(x_1, x_2, \dots)) = E_*(G(x_{n+1}, x_{n+2}, \dots)), \quad \text{for every } n \geq 1. \quad (18)$$

Here the function G may be a function of a single sample $G(x_1)$, or two samples $G(x_1, x_2)$, or all the samples $G(x_1, x_2, \dots)$. For example, because of the limit, the left side of (5) is a function of all the samples. In practice the process generating the samples may not be stationary, and assuming stationarity is considered a simplifying first step.

Let $\bar{g}$ denote the left side of (5), when the limit exists. Since shifting the index by one $x_n \rightarrow x_{n+1}$ changes only the first and last terms of a partial average, these discrepancies vanish in the limit, and $G = \bar{g}$ satisfies (18) with E_* deleted:

$$G(x_1, x_2, \dots) = G(x_{n+1}, x_{n+2}, \dots), \quad \text{for every } n \geq 1. \quad (19)$$

A function G satisfying this is called *invariant*. Thus, if the limit in (5) exists, it is an invariant function.

The ergodic theorem Birkhoff (1931), Durrett (2019) guarantees the limit in (5) exists, with probability one and in the mean (L^1), for every stationary process. A process is *ergodic* if every invariant function is a constant, with probability one. The second step is to assume ergodicity. When this happens, the limit in (5) is a constant; since L^1 convergence implies the expectation of the limit in (5) equals the limit of the expectations, this constant equals, by the first step, the expectation $E_*(g(x_1))$. We conclude (5) holds when $x_1, x_2, \dots$ is stationary and ergodic.

Suppose $x_1, x_2, \dots$ are independent and identically distributed. Then (18) follows immediately, yielding stationarity. For such a sequence, ergodicity is particularly simple to verify, and follows from the Kolmogorov zero-one law Kolmogorov (1933, 1950), Durrett (2019). Indeed, since an invariant function does not depend on $x_1, x_2, \dots, x_n$ for any n , it is a tail event, and Kolmogorov's zero-one law states every tail event has probability zero or one — hence every invariant function is constant with probability one, i.e., the process is ergodic.

A doubly-infinite process is a process $\dots, z_{-2}, z_{-1}, z_0, z_1, z_2, \dots$. Suppose x_0 is a function of a doubly-infinite process

$$x_0 = F(\dots, z_{-2}, z_{-1}, z_0, z_1, z_2, \dots).$$

Let σ be the shift, sending $z_n \rightarrow z_{n+1}$. Then we may define a process $x_1, x_2, \dots$ by

$$x_n = F(\sigma^n(\dots, z_{-2}, z_{-1}, z_0, z_1, z_2, \dots)) = F(\dots, z_{n-2}, z_{n-1}, z_n, z_{n+1}, z_{n+2}, \dots), \quad n \geq 1. \quad (20)$$

Now assume the doubly-infinite process is stationary, so shifting leaves its law unchanged. Since shifting $x_n \rightarrow x_{n+1}$ corresponds to shifting $z_n \rightarrow z_{n+1}$ in (20), $x_1, x_2, \dots$ is stationary. Moreover, if $G(x_1, x_2, \dots)$ is an invariant function of $x_1, x_2, \dots$, then, from (20), it is an invariant function of the doubly-infinite process. We have derived

Proposition 2. *If $\dots, z_{-2}, z_{-1}, z_0, z_1, z_2, \dots$ is a stationary and ergodic process and (20) holds, then $x_1, x_2, \dots$ is stationary and ergodic.*

References

- Anderson, T. W. & Goodman, L. A. (1957), ‘Statistical inference about markov chains’, *Annals of Mathematical Statistics* **28**(1), 89–110.
- Billingsley, P. (1961), *Statistical Inference for Markov Processes*, University of Chicago Press, Chicago.
- Birkhoff, G. D. (1931), ‘Proof of the ergodic theorem’, *Proceedings of the National Academy of Sciences* **17**(12), 656–660.
- Bishop, C. M. (2006), *Pattern Recognition and Machine Learning*, Information Science and Statistics, Springer, New York.
- Cover, T. M. & Thomas, J. A. (2006), *Elements of Information Theory*, 2nd edn, Wiley-Interscience.
- Cramér, H. (1946), *Mathematical Methods of Statistics*, Princeton University Press.
- Dembo, A. & Zeitouni, O. (1998), *Large Deviations Techniques and Applications*, Vol. 38 of *Stochastic Modelling and Applied Probability*, 2nd edn, Springer, New York.
- Donsker, M. D. & Varadhan, S. R. S. (1975), ‘Asymptotic evaluation of certain markov process expectations for large time, I’, *Communications on Pure and Applied Mathematics* **28**(1), 1–47.

- Doob, J. L. (1934), ‘Probability and statistics’, *Transactions of the American Mathematical Society* **36**, 759–775.
- Durrett, R. (2019), *Probability: Theory and Examples*, 5th edn, Cambridge University Press, Cambridge.
- Fisher, R. A. (1922), ‘On the mathematical foundations of theoretical statistics’, *Philosophical Transactions of the Royal Society of London, Series A* **222**, 309–368.
- Hijab, O. (2026), *Math for Data Science*, Springer, Cham. Corrected printing.
- Kolmogorov, A. N. (1933), *Grundbegriffe der Wahrscheinlichkeitsrechnung*, Springer, Berlin.
- Kolmogorov, A. N. (1950), *Foundations of the Theory of Probability*, Chelsea Publishing Company, New York. Translation edited by Nathan Morrison.
- Kullback, S. (1959), *Information Theory and Statistics*, Wiley, New York.
- Kullback, S. & Leibler, R. A. (1951), ‘On information and sufficiency’, *Annals of Mathematical Statistics* **22**(1), 79–86.
- Murphy, K. P. (2022), *Probabilistic Machine Learning: An Introduction*, MIT Press.
- Samson, E. W. (1951), Fundamental natural concepts of information theory, Report E5079, Communications Laboratory, Electronics Research Division, Air Force Cambridge Research Center, Cambridge, MA.
- Sanov, I. N. (1957), ‘On the probability of large deviations of random variables’, *Matematicheskii Sbornik* **42**, 11–44. (In Russian). English translation in *Selected Translations in Mathematical Statistics and Probability*, Vol. 1 (1961), pp. 213–244.
- Schervish, M. J. (1995), *Theory of Statistics*, Springer, New York.
- Shannon, C. E. (1948), ‘A mathematical theory of communication’, *Bell System Technical Journal* **27**(3), 379–423.
- Tribus, M. (1961), *Thermostatistics and Thermodynamics: An Introduction to Energy, Information and States of Matter, With Engineering Applications*, D. Van Nostrand Company, Princeton, NJ.
- van der Vaart, A. W. (1998), *Asymptotic Statistics*, Vol. 3 of *Cambridge Series in Statistical and Probabilistic Mathematics*, Cambridge University Press, Cambridge.
- Wald, A. (1949), ‘Note on the consistency of the maximum likelihood estimate’, *Annals of Mathematical Statistics* **20**(4), 595–601.
- Walker, G. (1931), ‘On periodicity in series of related terms’, *Proceedings of the Royal Society of London, Series A* **131**(818), 518–532.
- Yule, G. U. (1927), ‘On a method of investigating periodicities in disturbed series, with special reference to wolfer’s sunspot numbers’, *Philosophical Transactions of the Royal Society of London, Series A* **226**, 267–298.