1. The probability of a leap year selected at random contain 53					
Sunday is:					
(a) 53/ 366	(b) 1/7	(c) 2/7	(d) 53/365		
2. A bag contain	s 3 red and 2 b	lue marbles. A n	narble is drawn at		
random. The prol	bability of drav	ving a black ball	is:		
• •	` '	(c) $0/5$			
3. The probabilit	y that it will rai	in tomorrow is 0	.85. What is the		
probability that it					
* *	• •		(d) none of these		
•	•		ed from the numbers		
(1, 2, 3,,15	•		(1)		
	* *	(c) 2/15	• •		
5. What are the					
* *	* *	(c) 8	* *		
-	-	number selecte	d at random from the		
numbers (1,2,3, .		( ) 40 (05	(1)		
• •		1 7	(d) none of these		
7. The sum of th					
		0 (d) none			
_	-	are given; cnoos	e the correct answer		
for that which is		(a) 7/F	(d) none of these		
			(d) none of these.		
		iuitaneousiy, tha	n the probability of		
getting at least ty		(0) 1/	(4) 1 /0		
10. A letter is ch		(C) ½			
(a) 6/12	(h) 7/12	(a) 1	letter chosen has: (d) none of these.		
(a) 0/13	(b) // 13	(C) I	(d) Holle of these.		
11 A dica is thro	wn Find the n	rohability of gett	ing an even number.		
(A) 2/3	-	· · · · · · · · · · · · · · · · · ·	D) 1/2		
(A) 2/3	(6) 1	(0) 3/0	0) 1/2		
12. Two coins are thrown at the same time. Find the probability of getting both heads.					
(A) 3/4 (B) 1/4		(D) 0			
13. Two dice are thrown simultaneously. The probability of getting a					

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4	/9	
	ds are numbere ime number.	ed from 1 to 10	00. Find the	probability of	
(A) 3/4	(B) 27/50	(C) 1/4	(D)	29/100	
of drawing a blue balls in	a blue ball is do a bag is:	ouble that of a	red ball, the	f the probability en the number of	
(A) 5	(B) 10 (	C) 15	(D) 20		
	f 600 bulbs con random from t ve bulb is:				
(A) 143/150	(B) 147/	<b>150</b> (C)	1/25	(D) 1/50	
mixed thoro	earked with numbughly. One car lity that the num (B) 1/10	d is drawn fro	m this box r s a perfect s	andomly, then square.	
<b>18. What is</b> (A) 1/7	the probability (B) 53/366	•	Mondays in (D) 7	• •	
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit.  (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13					
equally likel 1,2,312 ,	of chance con y to come to re then the proba B) 1/12	est pointing to	one of the i	number n odd number is:	
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game.  (A) 3/4 (B) 1/2 (C) 1 (D) 1/4					

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:							
(A) 364/365	(B) 31/365	(C) 1/365	(D) 1/133225				
2. Then the pr	r <i>x</i> is chosen at ra robability that x <sup>2</sup> < 2/5 (C) 3/5	<pre>2 is?</pre>	umbers -2, -1, 0 , 1,				
a marble is dr red is 2/3, the	24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is:  (A) 10 (B) 6 (C) 8 (D) 7						
Then the prob		multiple of 3 and	0 natural numbers.   4 is:				
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?							
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) 4/21				
	_		nes are 50, 70, 82,				
	ne standard devia b) 25.49	c) 25.29	d) 25.69				
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.							
a) 13, 15	b) 13, 18	c) 18, 15	d) 13, 16				
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is							
a) $^{1}/_{2}$	b) $^{1}/_{3}$		d) $^{1}/_{6}$				
	ate between 0 and o) 7	d 3. The value of c) 27 d	E(X²) is <mark>) 9</mark>				
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?							

32.Out of t probability?	~	alues, whic	h one is not poss	sible in
a) $P(x) = 1$	b) ∑ x d) P(x	P(x) = 3 x = -0.5		
33.If E(x) =	<b>2 and E(z) = 4</b> b) 6	c) 0	•	sufficient data
34.The cov	ariance of two	independe	nt random variab	ole is
a) 1	b) 0	c) - 1	d) Ur	ndefined
<b>35.If Σ P(x</b> ) a) 0	<b>) = k<sup>2</sup> – 8 then,</b> b) 1	the value o		sufficient data
, ,	<b>0.5 and x = 4,</b> b) 0.5	, ,	? d) 2	
is always?	-		on, the sum of al	-
a) 0	b) Infinite	c) 1	d) Uno	defined
38.If the provenue of the second seco	obability of hi	tting the tai	get is 0.4, find m	nean and
	b) 0.6,	0.24	c) 0.4, 0.16	d) 0.6, 0.16
-	% and if 10 bo	mbs are dr	pped from a plac opped, find mear 0.4, 0.16	
a) 2	e mean of toss b) 4 s the mean and	c) 8	d) 1 <b>or standard norm</b>	nal distribution?

c) 5

d) 7

a) 3

b) 4

a) Mean is 0 and variance is 1 b) Mean is 1 and variance is 0								
c) Mean is 0 and variance is $\infty$ d) Mean is $\infty$ and variance is 0								
	Variance (X)	e of a rand b) E(X	om varia (2)		•		. d) (E(X))2	
	<b>43.Mean of a random variable X is given by a)</b> E(X)							
44.N a) 0	44.Mean of a constant 'a' is a) 0							
	45. Variance of a constant 'a' is a) 0							
46.Find the mean and variance of X?								
,	Х	0	1	2	3	4		
	f(x)	1/9	2/9	3/9	2/9	1/9		
a) 2, 4/3 b) 3, 4/3 c) 2, 2/3 d)						d) 3, 2/3		
47 Find the expectation of a random variable X2								

	Х	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) (	).5		b) 1.5		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by



c) np2q

d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a)  $\sqrt{np}$  b)  $\sqrt{pq}$  c) (np)2