

BWH

BridgeSplat: Bidirectionally Coupled CT and Non-Rigid Gaussian Splatting for Deformable Intraoperative Surgical Navigation


Maximilian Fehrentz^{1,2}, Alexander Winkler², Thomas Heiliger², Nazim Haouchine³, Christian Heiliger²,

Department of General, Visceral, and Transplantation Surgery, Hospital of the LMU Munich, Munich, Germany



Motivation


- Currently, there are no (deformable) navigation systems for
- Problem: Rigid registration is not sufficient for navigation in
- Goal: Using initial rigid registration and monocular laparoscopic video to deform preoperative CT during surgery



Preop: Coupling CT & 4DGS

• We parametrize 3D Gaussians w.r.t. a mesh of the abdominal cavity extracted from preoperative CT

Initialization




Intraop: Deforming CT via 4DGS

• Initial rigid registration & tracked imaging (laparoscope)



 Satisying the photometric loss drives optimization of deformation field that can be propagated to CT




Results

Simulated Data




Example sequence of a simulated scene: textured mesh, tool, and a realistic background, underneath the 3D model with stress



Exemplary results on a simulated sequence. Input image, rendered image from Gaussian Splatting, and deformed mesh

Clinical Data



Examples from a tool-tissue interaction sequence. Rows from top to bottom: input images acquired by the laparoscope, images rendered with our method, deforming

Evaluation on Simulated Data

Quant. Evaluation on Cartage SimCircular SimU	in
Quant. Evaluation on SimLeft SimCircular SimU	P
SimLeft SimChes	124
SimDown Shitter 210+0.19 0.1210	
Simin 5. 2.84	-
SimIn SimDown SimLeft SimCircuit SimIn SimIn SimDown SimLeft SimCircuit O.11±0.19 O.14±0.27 O.09±0.16 O.10±0.19 O.12±0.19 O.11±0.19 O.14±0.27 O.09±0.16 O.10±0.19 O.12±0.19 O.11±0.19 O.14±0.27 O.09±0.16 O.10±0.19 O.12±0.19 O.11±0.19 O.11±0.19 O.14±0.27 O.09±0.16 O.10±0.19 O.12±0.19 O.11±0.19	
Metric Second Point Contract Contract	m
Metric Euclidean Distance (mm) Euclidean Distance (mm) Sevelidean Distance (mm)	
Euclidean Distance (mm) 3.17 Euclidean Distance (mm) 3.17 Max Euclidean Distance (mm) istance dataset with Euclidean distance and max experiences interaction sequences	
Euclidean Distance	SEATTINE.
Max Euchuce ha simulated uate thissue interaction	
tion on the sill lated tool-uses	

Quantitative evaluation on the simulated datas error (in mm) on five simulated tool-tissue interact.