CLINICAL · EAR

Facial nerve palsy

A face that will not move is frightening for the patient and a test of method for the doctor. First decide whether the forehead is spared; then remember that Bell's palsy is a diagnosis of exclusion, examine the ear, parotid, neck and other cranial nerves, and protect the eye today.

AT A GLANCE

- Forehead spared means central.
- Bell's is a diagnosis of exclusion; check ear, parotid and neck.
- Protect the eye and give early steroids.
- ENoG and EMG guide decompression and repair.
- Slow or recurrent palsy needs imaging.

1 CHAPTER ONE Overview

One nerve, a long course, many causes.

1.1 Definition

Facial nerve palsy is weakness of the muscles of facial expression from a lesion anywhere between the motor cortex and the muscles. The approach depends on where the lesion is, how complete it is, and whether the nerve is likely to recover.

1.2 Why it matters

The eye is at risk within hours, the face carries a person's expression and identity, and a palsy may be the first sign of a tumour, a cholesteatoma or a fracture.

2 CHAPTER TWO Applied anatomy

A long and narrow course.

2.1 Central connections

The part of the facial nucleus that supplies the forehead receives fibres from both cerebral hemispheres; the part that supplies the lower face is controlled only from the opposite side. An upper motor neurone (*supranuclear*) lesion therefore spares the forehead. Emotional movement travels by separate pathways, so a smile may survive a stroke that abolishes voluntary movement.




FIGURE 1 The forehead receives bilateral cortical innervation and is spared in supranuclear paralysis; emotional movements travel by separate pathways.
Dhingra, fig. 14.1

2.2 The course of the nerve

2.2.1 Intracranial

From the pons across the cerebellopontine angle with the eighth nerve.

2.2.2 Intratemporal

- **Meatal segment:** in the internal acoustic meatus, anterosuperior.
- **Labyrinthine segment:** about 4 mm long, the narrowest part of the canal, ending at the geniculate ganglion. Swelling here is thought to cause Bell's palsy.
- **Tympanic segment:** along the medial wall of the middle ear above the oval window; the commonest site of bony dehiscence.

- **Mastoid segment:** vertically down to the stylomastoid foramen.

2.2.3 Extratemporal

Through the parotid gland, dividing into temporal, zygomatic, buccal, marginal mandibular and cervical branches.




FIGURE 2 Course of the facial nerve: meatal, labyrinthine, tympanic and mastoid segments, and its branches.
Dhingra, fig. 14.2

2.3 Branches inside the temporal bone

- **Greater superficial petrosal nerve:** lacrimation.
- **Nerve to stapedius:** the stapedial reflex.
- **Chorda tympani:** taste from the anterior two-thirds of the tongue and submandibular and sublingual secretion.

These allow a lesion to be localised (see 4.3).




Fig. 25: Nerves in relation with the middle ear. Note secretomotor pathway of salivary, lacrimal and nasal glands

FIGURE 3 The course of the facial nerve through the temporal bone, and the petrosal nerves and tympanic plexus of the middle ear.

Anatomy atlas · Doss p. 149, ENT Steps p. 53

2.4 Surgical landmarks




FIGURE 4 First, landmarks for finding the nerve in parotid surgery. Second, variations in its course, including bony dehiscence.

Dhingra, figs 14.4 and 14.5

2.5 Nerve injury

SUNDERLAND CLASSIFICATION

DEGREE	WHAT IS DAMAGED	OUTCOME
1	Conduction block (neurapraxia)	Full recovery
2	Axons, tubes intact (axonotmesis)	Full recovery, slower
3	Endoneurium	Incomplete, synkinesis likely
4	Perineurium	Poor without repair
5	Whole nerve (neurotmesis)	No recovery without repair

3 CHAPTER THREE Causes

Think beyond Bell's palsy.

CAUSES OF FACIAL PALSY

- CENTRAL **Stroke, tumour** forehead spared
- INFECTION **Bell's palsy, Ramsay Hunt** otitis media, cholesteatoma, necrotising otitis externa, Lyme disease
- TRAUMA **Temporal bone fracture** surgery, birth (forceps), facial wounds
- TUMOUR **Parotid cancer** facial schwannoma, haemangioma, glomus, vestibular schwannoma, temporal bone cancer
- SYSTEMIC **Diabetes** sarcoid (Heerfordt), Guillain-Barré, leukaemia, granulomatosis
- CONGENITAL **Möbius syndrome** birth injury

3.1 Bell's palsy

3.1.1 What it is

An acute idiopathic lower motor neurone palsy, 60 to 75% of all facial palsies. Reactivation of herpes simplex virus is the likely cause; the nerve swells inside the tight labyrinthine segment. It is commoner in diabetes and pregnancy.

3.1.2 Course

Weakness comes on over hours and peaks within 72 hours, often after a day of pain behind the ear. About 85 to 90% recover fully, and 95% if the palsy never becomes complete.




FIGURE 5 First, the features of a left-sided facial palsy. Second, Bell's palsy on the left in (A) an adult and (B) a child.

Dhingra, fig. 14.7

3.2 Ramsay Hunt syndrome

Herpes zoster oticus: vesicles in the concha, canal or palate, severe ear pain, facial palsy, and often sensorineural hearing loss and vertigo. The outlook for the face is worse than in Bell's palsy.



FIGURE 6 Ramsay Hunt syndrome: facial palsy with vesicles in the concha.
Dhingra, fig. 14.9

3.3 Otitis media and cholesteatoma

Acute otitis media can inflame a dehiscent nerve; cholesteatoma erodes the canal, usually in the tympanic segment. Both need urgent otological care.

3.4 Trauma

TEMPORAL BONE FRACTURES AND THE FACIAL NERVE

	LONGITUDINAL	TRANSVERSE
Frequency	About 80%	About 20%
Blow	Side of the head	Back of the head
Bleeding from the ear	Common	Usually absent
Hearing	Conductive	Sensorineural
Vertigo	Less often	Severe
Facial palsy	About 20%, often delayed	About 50%, often immediate




FIGURE 7 Longitudinal and transverse fractures of the temporal bone.
Dhingra, fig. 14.10

3.5 Tumours

A slowly progressive palsy, a recurrent palsy on the same side, or palsy with twitching suggests a tumour. A facial palsy with a parotid lump is almost always malignant.

3.6 Bilateral or recurrent palsy

Bilateral: Lyme disease, Guillain-Barré, sarcoidosis, leukaemia, Möbius syndrome, HIV, neurofibromatosis 2 and bilateral fractures. Recurrent on the same side: a tumour in about a third; *Melkersson-Rosenthal syndrome* combines recurrent palsy, lip swelling and a fissured tongue.

4 CHAPTER FOUR Assessment

Where, how bad, and why.

4.1 History

Speed of onset, pain, vesicles, hearing loss and vertigo, ear discharge, trauma or surgery, previous episodes, and systemic illness.

4.2 Examination

4.2.1 The face

Ask the patient to raise the eyebrows, close the eyes tightly, smile and puff the cheeks. On attempted eye closure the eye rolls up and out (*Bell's phenomenon*). Note drooling and tearing.

4.2.2 Grading

HOUSE-BRACKMANN GRADING

GRADE	DESCRIPTION
I	Normal
II	Slight weakness; complete eye closure with minimal effort
III	Obvious weakness, complete eye closure with effort, synkinesis may appear
IV	Obvious disfiguring weakness, incomplete eye closure
V	Barely perceptible movement
VI	No movement

4.2.3 Everything else

Otoscopy, the parotid and neck, the other cranial nerves, and hearing. A sixth nerve palsy with it points to the brainstem; fifth, eighth or lower nerve involvement points to the cerebellopontine angle; a motor-only palsy points to the parotid.

4.3 Topodiagnosis

LOCALISING THE LESION

LOST FUNCTION	LESION LIES
Lacrimation (Schirmer test)	At or above the geniculate ganglion

LOST FUNCTION	LESION LIES
Stapedial reflex	Above the nerve to stapedius
Taste and submandibular flow	Above the chorda tympani
Motor only	Below the chorda, in the mastoid or parotid




FIGURE 8 Topographical localisation of facial nerve lesions.
Dhingra, fig. 14.12

4.4 Electrical tests

4.4.1 Electroneuronography (ENoG)

Compares the compound muscle response on the two sides. Degeneration of 90% or more within the first three weeks predicts a poor recovery.

4.4.2 Electromyography (EMG)

Fibrillation potentials at 14 to 21 days mean denervation; polyphasic potentials appear 6 to 12 weeks before visible recovery; electrical silence means the motor end plates are lost.

WHERE THE BOOKS DIFFER · THE ENOG WINDOW

Dhingra	Lee


From day 4 to day 21.	Days 3 to 12 when deciding on decompression.

4.5 Imaging


Contrast MRI of the whole course, and CT of the temporal bone for fractures and cholesteatoma. Indicated for an atypical, slow, recurrent or bilateral palsy, or no recovery by three to four months.

IMAGING IN THE APP 2 cases

Facial nerve schwannoma and geniculate haemangioma, the tumours behind a slow or recurrent palsy.



Schwannoma of the Facial Nerve
Tumours · 3 images



Hemangioma of the Geniculate Ganglion
Tumours · 3 images

5 CHAPTER FIVE Management

Eye first, then the nerve.

5.1 Eye protection

Lubricating drops by day, ointment and taping at night, sunglasses or a moisture chamber. If closure stays poor, a gold or platinum weight in the upper lid, a lateral tarsal strip or tarsorrhaphy.

5.2 Bell's palsy

Oral steroids started within 72 hours, for example prednisolone 60 mg daily for five days followed by a five-day taper. An antiviral may add a small benefit in severe palsy.

WHERE THE BOOKS DIFFER · STEROIDS IN BELL'S PALSY

Dhingra	Lee
Prednisolone 1 mg/kg/day, extended if the palsy stays complete; notes that benefit is not proved beyond doubt. Current practice gives early steroids.	Recommends steroids within 48 to 72 hours.

Be cautious with steroids in diabetes, hypertension, peptic ulcer, tuberculosis, glaucoma and pregnancy.

5.3 Ramsay Hunt syndrome

High-dose steroids with an antiviral such as aciclovir or valaciclovir, and analgesia.

5.4 Otitis media and cholesteatoma

Acute otitis media: antibiotics with myringotomy. Chronic disease: urgent exploration, clearing disease from the nerve; granulation that has invaded the nerve is left alone.

5.5 Trauma

An immediate, complete palsy with a fracture through the canal is explored and decompressed or repaired. A delayed palsy is managed conservatively, with serial ENoG.

WHERE THE BOOKS DIFFER · DELAYED TRAUMATIC PALSY

Dhingra	Lee
Conservative treatment.	Serial ENoG; decompress if the response falls below 10% of normal.

5.6 Palsy after ear surgery

Remove the packing and any tight dressing first. If the nerve was not seen and the palsy is complete, explore unless continuity is certain.

5.7 Decompression

Opening the bony canal, and sometimes the sheath, at the site of compression. For Bell's palsy it is controversial and, if done, must include the labyrinthine segment within two weeks.

5.8 Repair and reanimation

5.8.1 Nerve repair

Tension-free primary anastomosis, or a cable graft from the great auricular (not when there are neck metastases), sural or medial antebrachial cutaneous nerve.

5.8.2 Nerve transfer

When the distal nerve and muscle survive but the proximal stump is unavailable, ideally within 18 months: the masseteric nerve is most used; hypoglossal transfer costs some tongue function; a cross-face graft can give a spontaneous smile.

5.8.3 Muscle transfer

For long-standing palsy with atrophied muscle: temporalis transfer, or a free gracilis transfer powered by the masseteric nerve (one stage, about 92% success, not spontaneous) or by a cross-face graft (two stages, about 80%, spontaneous).

5.8.4 Static procedures

Brow lift, fascia lata sling and lid procedures.

OPERATIVE GUIDES IN THEATRE 7 operations

Facial Nerve Decompression
Decompression
Tumors of the Facial Nerve
Facial nerve tumours
Temporal Bone Trauma
Temporal bone trauma
Nerve Graft Techniques
Nerve grafts
Eyelid Reanimation in Patients With Facial Paralysis
Eyelid reanimation
Facial Reanimation
Facial reanimation
Mastoid Surgery
Mastoid surgery when ear disease is the cause

CHAPTER SIX

6 Complications

When recovery is incomplete.

- **Exposure keratitis.**
- **Synkinesis:** the eye closes when the patient smiles, from aberrant regeneration; treated with physiotherapy and botulinum toxin.
- **Contracture and hemifacial spasm.**
- **Crocodile tears (Bogorad syndrome):** tearing while eating, as salivary fibres regrow into the lacrimal pathway.
- **Psychological distress.**



FIGURE 9 Hemifacial spasm: all the facial muscles and platysma contract together.
Dhingra, fig. 14.13

6.1 Hemifacial spasm

Usually from a vascular loop compressing the nerve at the brainstem. Treated with botulinum toxin, or microvascular decompression.

7 CHAPTER SEVEN For the examination

What the examiner wants to hear.

7.1 On call

ACUTE FACIAL WEAKNESS

1. Forehead spared with emotional movement kept: central; follow the stroke pathway if acute.
2. Protect the eye now: drops by day, ointment and tape at night.
3. Look in the ear: vesicles mean Ramsay Hunt, so add an antiviral; a foul ear or cholesteatoma needs urgent otology review.

4 Examine the parotid, neck and other cranial nerves.

5 After trauma, record whether the palsy was immediate or delayed.

6 Start steroids within 72 hours for Bell's palsy.

7.2 Viva essentials

WHAT THE EXAMINER IS LOOKING FOR

- **Upper versus lower motor neurone** palsy, and why the forehead is spared.
- **Segments of the nerve,** and the narrow labyrinthine segment.
- **Topodiagnosis:** Schirmer, stapedial reflex, taste.
- **House-Brackmann grading.**
- **ENoG and EMG:** timing and meaning.
- **Longitudinal versus transverse fractures.**
- **Bell's palsy is a diagnosis of exclusion.**

7.3 Common pitfalls

COMMON PITFALLS

- Calling it Bell's palsy without looking in the ear or feeling the parotid.
- Forgetting eye care.
- Missing a slowly progressive or recurrent palsy that needs imaging.
- Not recording the timing of a post-traumatic palsy.

Written for this app from Oxford Textbook of Otolaryngology ch. 10, Oxford Specialist Handbook ch. 17, Dhingra ch. 14 and K. J. Lee ch. 21. Figures reproduced from the books for private study. Study reference only: follow local protocols and senior advice for patients.

IMAGING


On imaging

The imaging of this topic: key findings, differentials and representative scans.

FACIAL NERVE · TUMOURS

Schwannoma of the Facial Nerve

- CT and MRI are complementary in the evaluation of the extension of the schwannoma to the middle or posterior cranial fossa as well as to the inner ear structures. Compared with hemangiomas, schwannomas of the facial nerve can be quite large before symptoms occur.
- Surgical removal with facial nerve grafting is only considered in (near) total paralysis of the facial nerve or when expansion of the lesion would lead to the damage of surrounding critical structures.




CT, coronal. CT demonstrates an opacification (1) in the medial and anterior epitympanum, suggestive of an expanding mass with destruction …

FACIAL NERVE · TUMOURS

Hemangioma of the Geniculate Ganglion

- Signs of infiltration or expansion. An MRI may be necessary to exclude or evaluate intracranial involvement.
- Most tumors, such as schwannoma, demonstrate more sharply defined enhancement.
- Outlines are often poorly defined in hemangiomas. A characteristic salt and pepper configuration is due to accumulation of methemoglobin and flow voids from the increased vascularization. Ossifying hemangiomas might demonstrate bone spicules.
- Hemangiomas of the facial nerve become symptomatic at an earlier stage compared with schwannomas, which might be quite large before symptoms occur.



CT, axial. There is a spiculated area of bone destruction or a mass with calcifications in the region of the geniculate ganglion (1), just distal to…