CLINICAL · LARYNX & AIRWAY

Congenital lesions of the larynx

A newborn's airway is narrow, soft and quick to swell, and most of its congenital problems announce themselves as noise: a high-pitched crowing that is worse on feeding and crying and better lying prone. Most of these babies have laryngomalacia and will outgrow it. The art is to recognise the few who will not, the stenosis, the paralysed folds, the haemangioma, the cleft, and to look at the whole airway before deciding.

AT A GLANCE

- Laryngomalacia is the commonest; normal cry, better prone.
- Subglottic stenosis: under 4 mm in a term neonate.
- Subglottic haemangioma: propranolol.
- Palpate the interarytenoid area for a cleft.

1 CHAPTER ONE Overview

The congenital larynx.

1.1 Lesions

- Laryngomalacia (the commonest).
- Vocal fold paralysis (the second commonest).
- Congenital subglottic stenosis.
- Laryngeal web and atresia.
- Subglottic haemangioma.
- Laryngeal cysts and laryngocele.
- Laryngo-oesophageal cleft.
- Tracheomalacia and vascular rings (in the trachea, outside the larynx).

1.2 Clues from the cry

A *normal cry* with stridor points away from the glottis (laryngomalacia, subglottic stenosis, haemangioma); a *weak, hoarse or absent cry* points to the vocal folds (paralysis, web). Stridor eased by lying prone suggests laryngomalacia, micrognathia or macroglossia. Choking and cyanosis with feeds suggest a cleft, paralysis or a tracheo-oesophageal fistula.

CHAPTER TWO

2 Laryngomalacia

Floppy supraglottis.

2.1 Features

Excessive flaccidity of the supraglottis, which is drawn into the airway on inspiration. Stridor appears at birth or within the first weeks, is inspiratory and fluttering, worse with crying, feeding, excitement and lying supine, and better prone; the cry is normal. Symptoms usually peak at six to eight months and resolve by 12 to 24 months. Reflux often coexists and aggravates it.



FIGURE 1 Laryngomalacia: the epiglottis folded longitudinally into an omega.
Dhingra, fig. 59.1

2.2 Appearances

On flexible endoscopy in the awake infant: an *omega-shaped*, elongated epiglottis curled on itself; short, tight aryepiglottic folds tethering it; and redundant mucosa over large arytenoids that prolapses into the glottis on inspiration. Vocal fold movement is normal. Lee classifies the collapse as type 1 (short aryepiglottic folds), type 2 (redundant arytenoid tissue) and type 3 (posterior collapse of the epiglottis, often with neuromuscular disease). It cannot be judged in a paralysed, anaesthetised child.

2.3 Treatment

- **Most:** reassurance and observation; positioning; anti-reflux treatment.
- **Severe disease** (failure to thrive, feeding difficulty, apnoeas, cyanosis, pulmonary hypertension): microlaryngoscopy and bronchoscopy to exclude a second lesion, then endoscopic *supraglottoplasty*: division of the tight aryepiglottic folds and trimming of redundant arytenoid mucosa with cold steel or laser.
- **Tracheostomy** rarely.

OPERATIVE GUIDES IN THEATRE 1 operation

Supraglottoplasty for laryngomalacia.

Laryngomalacia
Supraglottoplasty

3 CHAPTER THREE Vocal fold paralysis

The second commonest.

3.1 Causes

Birth trauma (stretching of the recurrent nerve in breech or forceps delivery), central nervous system anomalies (Arnold–Chiari malformation and hydrocephalus, usually bilateral), cardiac surgery and ligation of the ductus arteriosus, cardiomegaly and mediastinal lesions (usually left-sided), and idiopathic.

3.2 Presentation

- **Unilateral:** a weak, breathy cry, mild stridor, and choking or aspiration with feeds.
- **Bilateral:** a good cry but marked inspiratory stridor and respiratory distress, because both folds lie near the midline.

3.3 Assessment

Flexible endoscopy in the awake child; imaging of the brain, neck and chest; and a swallowing assessment.

3.4 Treatment

Observation for unilateral paralysis, with feeding support. Bilateral paralysis often needs a tracheostomy; definitive airway surgery is deferred, because many recover within the first year or two, and the larynx grows. Shunting for Arnold–Chiari malformation may restore movement.

4 CHAPTER FOUR Subglottic stenosis

The narrow ring.

4.1 Congenital

Thickening of the cricoid cartilage (cartilaginous) or of the submucosa (soft tissue) below the vocal folds. Diagnosed when the subglottic lumen is under *4 mm in a term neonate* (normal 4.5 to 5.5 mm) or under 3 mm in a premature baby. The child may be well until a cold produces stridor, recurrent or prolonged croup; the cry is normal.

4.2 Acquired

Far commoner now: from prolonged or repeated *intubation* in premature and sick neonates, especially with oversized tubes and movement; presenting as failure to extubate or stridor after extubation.

4.3 Grading

Cotton–Myer: grade I up to 50% obstruction, II 51 to 70%, III 71 to 99%, IV no lumen; assessed by the largest endotracheal tube that allows a leak.

4.4 Treatment

Many mild congenital stenoses improve as the larynx grows. Options: observation; endoscopic balloon dilatation and radial incision; anterior cricoid split (in neonates failing extubation); laryngotracheal reconstruction with costal cartilage grafts; and cricotracheal resection for severe stenosis; tracheostomy where needed.

OPERATIVE GUIDES IN THEATRE 1 operation

Subglottic stenosis surgery.

Subglottic Stenosis
Subglottic stenosis

5 CHAPTER FIVE Webs, cysts and haemangioma

Space-occupying lesions.

5.1 Laryngeal web

Incomplete recanalisation of the larynx in the tenth week. Usually an anterior glottic web between the vocal folds with a concave posterior edge; associated with 22q11 deletion (velocardiofacial syndrome), which should be tested for. A weak or absent cry from birth, and airway obstruction when the web is large. Thin webs are divided endoscopically with knife or CO2 laser; thick webs often have subglottic extension and need excision through a laryngofissure with a silicone keel and later dilatations. *Laryngeal atresia* causes complete obstruction at birth and death unless an emergency airway is made (or a tracheo-oesophageal fistula provides one).




FIGURE 2 A laryngeal web.
Dhingra, fig. 59.2

5.2 Subglottic haemangioma

Asymptomatic at birth; stridor appears at six weeks to six months as the haemangioma proliferates, often presenting as recurrent croup, with a normal cry; worse with crying because of venous engorgement. About half have a cutaneous haemangioma, especially in the beard distribution; some have mediastinal lesions. Endoscopy shows a soft, compressible, reddish-blue swelling, usually

posterolateral in the subglottis; biopsy is not needed and may bleed. MRI or CT shows the extent.

- **Propranolol** is now first-line and often produces dramatic involution.
- Systemic steroids (Dhingra describes dexamethasone, then prednisolone for many months) and intralesional steroid.
- CO2 or KTP laser or open excision for resistant lesions.
- Tracheostomy while awaiting involution, which is the natural history.

5.3 Laryngeal cysts

Saccular cysts (a mucus-filled saccule) and *ductal cysts* in the aryepiglottic fold, vallecula or subglottis appear as smooth, bluish, fluid-filled swellings that obstruct the airway. Needle aspiration can provide an emergency airway; endoscopic deroofing or excision with laser is definitive. Subglottic cysts follow intubation in premature infants.

5.4 Laryngocele

An air-filled dilatation of the saccule rising between the false fold and the thyroid lamina: *internal* (within the larynx, a supraglottic bulge), *external* (through the thyrohyoid membrane, a neck swelling that enlarges with Valsalva), or *combined*. Rare in infants; in adults linked to wind instrument players and glass blowers, and sometimes to a ventricular carcinoma blocking the saccule, which must be excluded. Infection forms a *laryngopyocele*. Endoscopic excision for internal ones, external excision for external and combined.




FIGURE 3 First, a laryngocele of the mixed type with internal and external components. Second, a left laryngocele as seen on Valsalva (arrow).

Dhingra, figs 61.8 and 61.9





FIGURE 4 Laryngocele.

Anatomy atlas · ENT Steps p. 240

IMAGING IN THE APP 1 case

Laryngocele on imaging.



Laryngocele
Infection & inflammation · 10 images

6 CHAPTER SIX Laryngeal cleft

A gap at the back.

Incomplete fusion of the posterior cricoid lamina and tracheo-oesophageal septum, leaving a communication between larynx and oesophagus.

BENJAMIN–INGLIS CLASSIFICATION

TYPE	EXTENT
I	Interarytenoid, above the vocal folds
II	Partly into the cricoid lamina
III	Through the whole cricoid, into the cervical trachea
IV	Into the thoracic trachea

Choking, coughing and cyanosis with feeds, recurrent aspiration pneumonia, a hoarse cry and stridor. Diagnosed at microlaryngoscopy by *palpating* the interarytenoid area with a probe, since small clefts are hidden by redundant mucosa. Type I clefts may be managed by thickened feeds or endoscopic injection or repair; deeper clefts need endoscopic or open repair.

OPERATIVE GUIDES IN THEATRE 1 operation

Laryngeal cleft repair.

Surgery for Aspiration: Laryngeal Cleft Repair and Laryngotracheal Separation
Laryngeal cleft repair

7 CHAPTER SEVEN
The trachea and beyond

Not always the larynx.

- **Tracheomalacia:** a soft, collapsing trachea, primary or after compression or tracheo-oesophageal fistula repair; expiratory stridor and a barking cough; most improve with growth; aortopexy, stenting or tracheostomy for severe cases.
- **Vascular rings and slings:** a double aortic arch, right arch with ligamentum, anomalous innominate artery or pulmonary artery sling compress the trachea and oesophagus, causing stridor and dysphagia. Contrast CT or MRI and a barium swallow; surgical division.
- **Tracheo-oesophageal fistula and oesophageal atresia.**
- **Cri-du-chat syndrome** (deletion of the short arm of chromosome 5): a cat-like cry with a narrow, diamond-shaped glottis, learning disability and hypotonia.

8 CHAPTER EIGHT
For the examination

What the examiner wants to hear.

8.1 Viva essentials

WHAT THE EXAMINER IS LOOKING FOR

- **Laryngomalacia:** presentation, endoscopic appearance, natural history, supraglottoplasty.
- **Normal versus hoarse cry** as a localising sign.
- **Subglottic stenosis:** the 4 mm rule, causes, grading and treatment.
- **Subglottic haemangioma:** age of onset, beard haemangioma, propranolol.
- **Laryngeal web** and its association with 22q11 deletion.
- **Laryngeal cleft:** palpate the interarytenoid area.

8.2 Common pitfalls

COMMON PITFALLS

- Labelling every noisy baby laryngomalacia without looking at the whole airway when symptoms are severe.
- Biopsying a subglottic haemangioma.
- Missing a cleft because the interarytenoid area was not palpated.

Written for this app from Oxford Textbook of Otolaryngology ch. 74, 79, Oxford Specialist Handbook ch. 23, Oxford Handbook of ENT ch. 17, Dhingra ch. 59 and K. J. Lee ch. 50, 53. Figures reproduced from the books for private study. Study reference only: follow local protocols and senior advice for patients.

IMAGING

On imaging

The imaging of this topic: key findings, differentials and representative scans.

LARYNX & TRACHEA · INFECTION & INFLAMMATION

Laryngocele

IMAGING FINDINGS


- Air or fluid-filled sac from the laryngeal ventricle
- Internal (paraglottic) or external through the thyrohyoid membrane
- Rim enhancement if infected (pyolaryngocele)

TOP DIFFERENTIALS


- Thyroglossal duct cyst
- Second branchial cleft cyst
- Saccular cyst

CHECKLIST

- Exclude a carcinoma at the ventricle in adults
- Airway compromise



CT with intravenous contrast enhancement, axial. The axial slice at the lower border of the hyoid (1) reveals a homogeneous, expanding…



Coronal CECT reformat shows a similar lesion. This is also known as a "mixed" laryngocele because it contains internal…

Key facts after Koch, Hamilton, Hudgins and Harnsberger, Diagnostic Imaging: Head and Neck, 3rd ed., p. 292.