CLINICAL · MOUTH & THROAT

Dysphagia and pharyngeal pouch

Swallowing is a reflex so well rehearsed that we notice it only when it fails. Dysphagia is a symptom, never a diagnosis, and in an adult it is cancer until proven otherwise. The elderly patient who gurgles when he drinks, brings back yesterday's food and coughs at night has a pharyngeal pouch, the classic ENT cause, through the weak spot between the two parts of the inferior constrictor.

AT A GLANCE

- Regurgitation and aspiration in the elderly suggest a pouch.
- Barium before scopes.
- Button batteries are an emergency.
- Fever or crepitus means perforation.
- Cricopharyngeal bars are seen in a third of normal people; stapling is less effective for small pouches.

1 CHAPTER ONE Dysphagia

Where is the hold-up?

1.1 Definitions

Dysphagia is difficulty in swallowing; *odynophagia* is pain on swallowing; *globus* is a lump feeling in the throat between meals without difficulty swallowing food; *aspiration* is passage of material below the vocal folds.

1.2 Types by level

- **Oral:** poor lip seal, chewing or bolus control; drooling and food left in the mouth (stroke, facial palsy, tongue surgery, neuromuscular disease).
- **Oropharyngeal:** difficulty starting a swallow, coughing or choking as soon as one swallows, nasal regurgitation, food sticking in the throat (stroke, motor neurone disease, myasthenia, Parkinson disease, head and neck cancer and its treatment, cricopharyngeal dysfunction, pouch).

- **Oesophageal:** food sticking behind the sternum seconds after swallowing (strictures, rings and webs, achalasia and spasm, tumours, extrinsic compression).

CAUSES OF DYSPHAGIA

GROUP	EXAMPLES
Lumen	Foreign body, food bolus
Wall: inflammation	Pharyngitis, tonsillitis, quinsy, oesophagitis (reflux, candida, herpes, eosinophilic, pill, caustic)
Wall: tumour	Carcinoma of the oropharynx, hypopharynx or oesophagus
Wall: structural	Pouch, webs (Plummer–Vinson), Schatzki ring, peptic and caustic strictures
Wall: motility	Achalasia, diffuse spasm, presbyoesophagus, scleroderma, polymyositis
Outside the wall	Goitre, cervical osteophytes, mediastinal nodes or tumour, aortic aneurysm, aberrant right subclavian artery (dysphagia lusoria)
Neurological	Stroke, motor neurone disease, multiple sclerosis, Parkinson disease, myasthenia, bulbar and pseudobulbar palsy, vagal palsy

RED FLAGS

Progressive dysphagia for solids then liquids, weight loss, odynophagia, haematemesis or haemoptysis, voice change, referred otalgia, a neck mass, or age over fifty with new dysphagia: urgent endoscopy for malignancy.

1.3 Evaluation

- **History:** level, solids or liquids, progression, regurgitation, cough with eating, weight, voice, heartburn, neurological symptoms.
- **Examination:** mouth, oropharynx, flexible laryngoscopy (pooling, vocal fold movement, sensation), neck, cranial nerves.
- **Videofluoroscopy (modified barium swallow):** the oral and pharyngeal phases, penetration and aspiration, and the effect of postures and textures.
- **Fibreoptic endoscopic evaluation of swallowing (FEES):** swallowing coloured food under endoscopic view, with or without sensory testing.
- **Barium swallow:** pouches, webs, rings, strictures, motility.

- **Endoscopy:** rigid or flexible oesophagoscopy, or unsedated transnasal oesophagoscopy, with biopsy.
- **Manometry:** motility disorders (achalasia, spasm).
- **CT** for tumours and extrinsic compression.




FIGURE 1 The constrictions of the oesophagus and their distance from the incisors.
Anatomy atlas · ENT Steps p. 226

1.4 Oropharyngeal dysfunction and its treatment

- **Tongue base weakness:** residue in the valleculae; chin tuck, tongue base exercises.
- **Pharyngeal sensory loss:** silent aspiration, worse with thin liquids; thickened fluids, head turned to the weak side.
- **Vocal fold palsy:** aspiration of thin liquids and a breathy voice; medialisation.
- **Poor laryngeal elevation:** residue and failure of the sphincter to open; the Mendelsohn manoeuvre, exercises and stimulation.
- **Failure of upper oesophageal sphincter opening:** a cricopharyngeal bar (present in many asymptomatic elderly people) or a pouch; dilatation, botulinum toxin or cricopharyngeal myotomy.

Speech and language therapists guide postures, textures and exercises; a gastrostomy feeds those who cannot swallow safely.

CHAPTER TWO

2 Pharyngeal pouch

Zenker diverticulum.

2.1 Definition

A *pulsion diverticulum* in which pharyngeal mucosa herniates through *Killian's dehiscence*, the weak triangle in the posterior pharyngeal wall between the oblique *thyropharyngeus* above and the transverse *cricopharyngeus* below, the two parts of the inferior constrictor. Also called a hypopharyngeal diverticulum.




FIGURE 2 First, a hypopharyngeal (Zenker) diverticulum herniating through Killian's dehiscence. Second, the potential sites of diverticula: Killian's dehiscence, the Killian–Jamieson area and Laimer's dehiscence.

Dhingra, figs 54.1 and 54.2




FIGURE 3 Killian's dehiscence and the pharyngeal pouch, and Killian's and Laimer's triangles.
Anatomy atlas · ENT Steps p. 198, ENT Steps p. 225

2.2 Cause

Not fully known. Incoordination between pharyngeal contraction and cricopharyngeal relaxation, failure of the sphincter to open (fibrosis of cricopharyngeus, poor laryngeal elevation) and raised pressure above it force mucosa out through the weak area. Reflux may contribute. It is a disease of the elderly, usually over sixty, men more than women.

2.3 Pathology

The pouch begins in the midline, lies first behind the oesophagus, then enlarges downwards and comes to lie to its left. As it grows, its mouth comes to lie in line with the pharynx while the oesophageal opening is pushed forward, so food enters the sac preferentially.

2.4 Clinical features

- **Dysphagia:** often appearing after a few swallows, once the pouch fills and presses on the oesophagus; progressive.
- **Gurgling** in the neck on swallowing (*borborygmi*); a swelling in the left side of the neck that empties with pressure.
- **Regurgitation** of undigested food, sometimes hours later and at night when lying down, with *nocturnal cough* and *aspiration pneumonia*.

- **Halitosis**, weight loss and malnutrition.
- Associated hiatus hernia; rarely carcinoma within a long-standing pouch.

2.5 Diagnosis

A *barium swallow* shows the sac and its size, best in the lateral view. Endoscopy carries a risk of entering and perforating the sac, but is needed if carcinoma is suspected; nasogastric tubes are passed with care.



FIGURE 4 Barium swallow showing an early Zenker's diverticulum (arrow) at the level of the sixth cervical vertebra in a 31-year-old woman.

Dhingra, fig. 54.3


IMAGING IN THE APP 3 cases

Zenker diverticulum, oesophageal passage disorders and aspiration.




Zenker Diverticulum (Pharyngeal Pouch)

Infection & inflammation · 9 images



Esophageal Passage Disorders

Infection & inflammation · 2 images



Aspiration

Infection & inflammation · 3 images

2.6 Treatment

Small asymptomatic pouches can be watched. For symptoms:

2.6.1 Endoscopic

- **Endoscopic stapling diverticulotomy:** the party wall (the *diverticular bar*, containing cricopharyngeus) between the pouch and the oesophagus is divided and sealed with a linear stapler through a Weerda diverticuloscope; the pouch becomes part of the oesophagus and the muscle is divided at the same time. Now the commonest operation.
- **Dohlman's procedure / CO2 laser diverticulotomy:** the bar is divided by diathermy or laser through the diverticuloscope; suited to poor-risk and elderly patients.

Endoscopic methods are quick, with rapid return to eating, but are less effective for very small pouches (too little bar to staple) and may be impossible when the neck cannot be extended or the mouth opened.

2.6.2 External (cervical)

Through an incision along the anterior border of sternomastoid, usually on the left, the pouch is exposed, and either excised, inverted or suspended (diverticulopexy), *always with cricopharyngeal myotomy* to prevent recurrence.

MYOTOMY IS THE KEY

Whatever is done to the sac, dividing cricopharyngeus treats the cause. Excision without myotomy invites recurrence and fistula.

2.6.3 Complications

> Mediastinitis from perforation, pharyngocutaneous fistula, recurrent laryngeal nerve palsy, recurrence, and, with endoscopic methods, failure to complete the division.

OPERATIVE GUIDES IN THEATRE 4 operations

Upper oesophageal sphincter surgery and oesophagoscopy.

Dysfunction of the Upper Esophageal Sphincter
Upper oesophageal sphincter
Esophagoscopy and Hypopharyngoscopy
Oesophagoscopy
Partial Pharyngectomy
Partial pharyngectomy
Total Laryngopharyngectomy
Total laryngopharyngectomy

3 CHAPTER THREE Other causes

Webs, rings and motility.

3.1 Plummer–Vinson syndrome

Iron deficiency anaemia, dysphagia from *postcricoid webs*, glossitis, angular cheilitis and koilonychia, mostly in women; a predisposition to postcricoid carcinoma. Barium swallow shows the web; iron replacement and dilatation of the web; surveillance.

3.2 Rings and strictures

A *Schatzki ring* at the gastro-oesophageal junction causes intermittent dysphagia for solids and food impaction; dilatation. Strictures from reflux, pills, caustics, radiotherapy or surgery are dilated and the cause treated.

3.3 Motility disorders

- **Achalasia:** loss of the myenteric plexus; progressive dysphagia for solids and liquids, regurgitation and chest pain; a dilated oesophagus tapering to a *bird's beak*; aperistalsis with failure of the lower sphincter to relax on manometry. Pneumatic

dilatation, surgical or peroral endoscopic myotomy, or botulinum toxin; exclude a tumour at the cardia (pseudoachalasia).

- **Diffuse oesophageal spasm:** chest pain and dysphagia, a "corkscrew" oesophagus; nitrates or calcium channel blockers.
- **Scleroderma, polymyositis, diabetes and hypothyroidism** also impair peristalsis.

3.4 Eosinophilic oesophagitis

Young atopic adults with dysphagia and food bolus obstruction; rings on endoscopy and eosinophils on biopsy; elimination diets, swallowed steroids and dilatation.

3.5 Extrinsic compression

Goitre, cervical osteophytes, mediastinal masses, and *dysphagia lusoria* from an aberrant right subclavian artery passing behind the oesophagus (often with a non-recurrent right laryngeal nerve).

CRICOPHARYNGEAL DYSFUNCTION AND NEWER OPTIONS (OXFORD TEXTBOOK)

The upper oesophageal sphincter is formed by the lower inferior constrictor, the cricopharyngeus and the upper oesophagus. In **cricopharyngeal dysfunction** the sphincter opens too little or relaxes poorly; videofluoroscopy may show a *cricopharyngeal bar* at C5–C6, but a bar is seen in about a third of people without symptoms, so it must be matched to the complaint. **High-resolution pharyngeal manometry** with videofluoroscopy shows whether the pharynx contracts normally against a high residual sphincter pressure, which predicts benefit from dilatation, botulinum toxin or myotomy. **Transnasal oesophagoscopy** under local anaesthetic examines the oesophagus in clinic and allows balloon dilatation of strictures and placement of feeding tubes. For pouches, endoscopic stapling remains first line for medium and large pouches in fit patients; it may not divide enough of the septum in small pouches (under about 2 to 3 cm), where laser or open myotomy with diverticulectomy or inversion is preferred. Flexible endoscopic septotomy, including the submucosal tunnelling technique (Z-POEM), is increasingly offered by gastroenterologists, especially to frail patients who cannot extend the neck for rigid access.

4 CHAPTER FOUR For the examination

What the examiner wants to hear.

4.1 Viva essentials

WHAT THE EXAMINER IS LOOKING FOR

- **Causes of dysphagia** by lumen, wall and outside, and the red flags.
- **Investigations:** videofluoroscopy, FEES, barium swallow, endoscopy, manometry.
- **Killian's dehiscence** and the anatomy of a pouch.
- **Clinical features** of a pharyngeal pouch.
- **Treatment:** endoscopic stapling, Dohlman, excision with cricopharyngeal myotomy.
- **Plummer–Vinson syndrome.**

4.2 Common pitfalls

COMMON PITFALLS

- Passing a rigid oesophagoscope blindly in a patient with a pouch.
- Excising a pouch without cricopharyngeal myotomy.
- Attributing progressive dysphagia in an older smoker to "age" without endoscopy.

Written for this app from Oxford Textbook of Otolaryngology ch. 48, Oxford Specialist Handbook ch. 10, Oxford Handbook of ENT ch. 13, Dhingra ch. 54 and K. J. Lee ch. 36. Figures reproduced from the books for private study. Study reference only: follow local protocols and senior advice for patients.

IMAGING

On imaging

The imaging of this topic: key findings, differentials and representative scans.

PHARYNX, TONSILS & OESOPHAGUS · INFECTION & INFLAMMATION

Zenker Diverticulum (Pharyngeal Pouch)

IMAGING FINDINGS

- Outpouching from the posterior hypopharynx above the cricopharyngeus on barium swallow
- Air–fluid level behind the oesophagus on CT
- Residual barium after the swallow

TOP DIFFERENTIALS


- Lateral pharyngeal or Killian–Jamieson diverticulum
- Paratracheal air cyst

CHECKLIST

- Size and neck of the pouch
- Aspiration on the swallow



Barium swallow, anteroposterior view. After swallowing contrast, some contrast has remained in the hypopharynx (1) due to physiologic…




Barium esophagram demonstrates a large diverticulum with retained layering contrast from the posterior lateral junction of the hypopharynx…

Key facts after Koch, Hamilton, Hudgins and Harnsberger, Diagnostic Imaging: Head and Neck, 3rd ed., p. 258.

PHARYNX, TONSILS & OESOPHAGUS · INFECTION & INFLAMMATION

Esophageal Passage Disorders

- Patients with removable dentures may lack oral sensory input regarding the size of their food. This may also happen with sharp foreign bodies, which may cause perforations of the esophagus.
- Anatomic structures such as the aortic arch or an aberrant right esophageal artery are demonstrated as persistent impressions without stasis of contrast.




Barium swallow, lateral view. Problems with esophageal passage can be a result of a variety of etiologies occurring either singly or in…

LARYNX & TRACHEA · INFECTION & INFLAMMATION

Aspiration

- Occasionally aspiration is caused by mechanical failure of the epiglottis to close because of the presence of tumor, but most cases are due to neurologic (sensory) disorders without any visible pathology apart from the aspiration. Beware of aspiration pneumonia.
- The above-mentioned pathologies are partly demonstrated in other sections.



First stage: the barium contrast (1) just reaches the esophagus. For orientation note the mandible (2), hyoid (3), and vallecula (4)....