

The Margin That Matters: Pathology, Disease Mechanism, and Community Access in Surgical Oncology

Introduction

In the operating room, the surgeon's job often ends where the pathologist's begins. A tumor is excised, the specimen is inked and oriented, and whether that patient goes home cured or returns for a second surgery frequently comes down to a few millimeters of tissue read under a microscope. Surgical margin assessment is one of the most consequential steps in oncologic care, yet it is also one of the least understood outside of pathology and surgery themselves. Margin status is not a single number; it is the endpoint of several distinct biological processes, including how a tumor grows within tissue, how it acquires the ability to spread, and how the body's own wound-healing response behaves once the tumor has been removed.

This paper takes a case-anchored approach to that biology. Rather than reviewing margin statistics in the abstract, it works outward from a single published clinical case, using each pathologic finding in that case as a starting point for explaining the underlying disease mechanism. It then connects that mechanism-level understanding to a broader, community-facing problem: whether patients outside major cancer centers have equitable access to the standardized pathology reporting and timely surgical care that this biology requires. Having spent time in both a surgical oncology theatre and an emergency department, I wanted to examine this topic the way a pathologist and a treatment team actually would, moving from a specific specimen to the general principles it illustrates.

Literature Review

1. A Case to Ground the Discussion

A case published in the journal *Oncology* illustrates how easily margin status can be misjudged even by an experienced team (Brown et al., 2017). A 46-year-old woman presenting after a screening mammogram showing new calcifications underwent biopsy of three sites within the same breast. One site showed high-grade ductal carcinoma in situ (DCIS) with comedonecrosis, strongly positive for estrogen and progesterone receptors. She proceeded to lumpectomy, and the final pathology report described a small invasive tumor, under one centimeter, with a positive margin for invasive carcinoma and lymphovascular space invasion present. Given the extent and grade of disease, she opted for bilateral mastectomy rather than a second, more limited excision. The mastectomy specimen revealed two separate foci of high-grade DCIS, one of them 20 millimeters across, along with two positive sentinel lymph nodes containing metastatic deposits over a centimeter in size (Brown et al., 2017).

Nothing about this case involved a technical failure. The surgeon removed what imaging showed, and the pathologist read the specimen correctly. What the case demonstrates instead is how much biological information a single margin report can carry, and how each finding within it points to a distinct physiological process that helps explain why the disease behaved the way it did.

2. DCIS Growth Pattern and Margin Positivity

The mastectomy in this case uncovered a second, previously undetected 20 millimeter focus of high-grade DCIS. This is a direct consequence of how DCIS grows. Ductal carcinoma in situ does not expand as a single solid mass; it proliferates within the existing branching architecture of the breast's ductal system, filling and distending individual ducts as it spreads along their length (Hanna et al., 2019). Because a mammogram, an ultrasound, or even a single pathology slide can only sample a two-dimensional cross-section of a three-dimensional branching structure, disease can appear to stop in one duct and then reappear, unconnected on that slide, in a neighboring one. This is part of why consensus guidelines require a full 2 millimeter margin of normal tissue around DCIS rather than the simpler "no ink on tumor" standard used for invasive cancer: a narrower buffer leaves too much room for unsampled, branching disease to be left behind (Chauhan et al., 2024). Reported rates of positive or close margins in DCIS surgery span a wide range, from roughly 5 to 57 percent, in large part because the disease's own growth pattern makes it inherently difficult to fully visualize before the tissue is removed (Chauhan et al., 2024).

3. Comedonecrosis as a Marker of Aggressive Biology

The DCIS in this case was described as high grade with comedonecrosis, meaning the centers of the affected ducts contained dead tumor cells. This is not incidental detail. High-grade DCIS cells proliferate faster than the duct's existing blood supply can support, and the cells furthest from the duct wall, and therefore furthest from any nearby capillary, become starved of oxygen and nutrients and die, leaving a necrotic core surrounded by a rim of viable, dividing tumor cells (Hanna et al., 2019). Comedonecrosis is one of the most consistently reported predictors of local recurrence in the DCIS literature, and it is associated with a shorter interval to recurrence compared to low-grade disease without necrosis, often under five years rather than more than fifteen (Hanna et al., 2019; Pathology Outlines, n.d.). The mechanistic reason this matters clinically is that a lesion aggressive enough to outgrow its own blood supply is also a lesion whose true margins are harder to predict, which is part of why surgeons are typically advised to pursue wider excision margins whenever comedonecrosis is present.

4. Lymphovascular Invasion and the Route to Metastasis

The invasive component of the tumor in this case showed lymphovascular space invasion, meaning tumor cells were found inside small blood or lymphatic channels adjacent to the main lesion. This finding reflects a specific step in tumor biology. For a cancer cell to appear inside a vessel, it must first detach from the primary tumor mass, a process that typically involves loss of cell adhesion proteins such as E-cadherin, and then physically breach the basement membrane and endothelial wall of a nearby vessel using matrix-degrading enzymes. Once inside, that cell has access to the body's circulatory or lymphatic transport network, the same route metastatic disease uses to reach lymph nodes and distant organs. This is precisely what happened in the case described: both sentinel lymph nodes removed during the mastectomy contained metastatic deposits, at 1.3 and 1.6 centimeters respectively, confirming that disease had already used this route before the mastectomy was performed (Brown et al., 2017). Lymphovascular invasion is one of the strongest independent predictors of both margin positivity and recurrence risk in the breast cancer literature.

5. The Wound Microenvironment and Residual Tumor Survival

Had this patient chosen re-excision instead of mastectomy, the tissue left behind after that second surgery would not simply sit inert while waiting for follow-up imaging. Surgical wounds are sites

of active biological signaling. Wound healing involves the release of cytokines and growth factors, including cyclooxygenase-2 (COX-2), a prostaglandin-producing enzyme, and transforming growth factor-beta (TGF-beta), both of which support normal tissue repair but have also been shown to promote the survival and proliferation of nearby residual tumor cells (Predina et al., 2012). In animal models, tumor cells injected next to a fresh surgical wound grow measurably more aggressively than identical cells injected into unwounded tissue, and this effect has been traced in head and neck cancer models to activation of the STAT3 signaling pathway inside the tumor cells themselves, driven largely by interleukin-6 released at the wound site (Predina et al., 2012; Ekblad et al., 2013). This is one reason aggressive breast cancers show a disproportionately high rate of local recurrence in the first two to three years after surgery, a window that tracks with the timeline of postoperative tissue remodeling rather than random chance, and it is a central reason adjuvant radiation and systemic therapy are used after re-excision rather than treating a clean margin alone as sufficient.

6. Minimal Residual Disease Versus Field Cancerization

Not every local recurrence traces back to cells left behind at the surgical edge. In head and neck cancer, where 10 to 30 percent of patients with histologically negative margins still develop local recurrence, molecular studies have identified two distinct underlying mechanisms (Pierik et al., 2021). The first is minimal residual disease: small numbers of tumor cells present in the tissue but too sparse to be caught on a standard slide. The second is field cancerization, in which a broad area of tissue surrounding the original tumor has already accumulated genetic changes, typically from shared carcinogen exposure, sufficient to give rise to an entirely new tumor later from tissue that looked histologically normal at the time of surgery (Pierik et al., 2021). These are functionally different diseases that can produce the same clinical picture, and a margin report reading "negative" cannot distinguish between them, which is part of why long-term surveillance remains essential even after a technically successful surgery.

7. Standardizing Pathology Communication

None of this biology is useful to a treatment team unless it is communicated clearly. For decades, pathology reports were written as narrative free text, a format prone to omission and inconsistent terminology. The College of American Pathologists developed standardized, checklist-style synoptic reporting protocols to address this directly, and synoptic reports have been shown to improve completeness and reduce ambiguity between pathologists and treating clinicians (Torous et al., 2021). Yet adoption remains uneven: a 2024 study across the U.S. Department of Veterans Affairs health system found that only 48 percent of reviewed pathology reports for four common cancers used a synoptic format, with specific required elements consistently missing from the rest (Ould Ismail et al., 2024). From what I saw shadowing in surgical oncology, a pathology report is often read quickly, sometimes mid-case, and its structure directly affects how fast and how accurately a surgeon can act on findings like the ones in the case above.

8. Access, Distance, and Who Benefits from the Biology

The reliability of margin assessment means little if a patient cannot reach the surgical and pathology infrastructure required to act on it. Cancer disparities affecting rural populations, racial and ethnic minorities, and patients in persistent poverty are well documented and described in the surgical oncology literature as largely systemic and preventable (Johnson et al., 2024). Rural regions have far fewer oncologists, pathologists, and surgeons per capita, along with fewer

dedicated cancer centers and diagnostic laboratories (ASCO, 2020). Greater travel distance to a treatment center has been independently associated with a lower likelihood of receiving guideline-recommended chemotherapy, regardless of insurance status. In lung cancer care, rural residence has been shown to independently predict a longer interval between diagnostic testing and surgical treatment, even after adjusting for comorbidities (Minerva et al., 2022). Every mechanism described above unfolds on a biological clock. A patient who needs a second surgery after a positive margin but has to wait weeks for a slot at a distant regional center is facing a delay layered directly on top of a wound-healing environment that pathology research suggests is actively working against them.

Synthesis: What the Case Demonstrates

Read together, the findings in this single case trace a chain of biological reasoning that a margin report is meant to compress into a few lines:

- Growth pattern: DCIS follows the branching architecture of the ductal system, which is why a second, undetected 20 mm focus was found only after mastectomy.
- Necrosis: comedonecrosis signals a lesion outgrowing its blood supply, correlating with higher-grade, less predictable disease.
- Vascular breach: lymphovascular invasion marks the point at which a tumor gains access to the body's transport systems, consistent with the two positive sentinel nodes found in this case.
- Wound biology: had re-excision been chosen, COX-2, TGF-beta, and STAT3-driven signaling in the healing wound could have actively favored survival of any residual tumor cells.
- Reporting and access: none of this information changes outcomes unless it reaches the surgical team in a usable format, and unless the patient can act on it within a clinically meaningful window.

This is why the treatment team's decision to pursue mastectomy, rather than a second limited excision, reflects an integration of several independent physiological signals rather than a response to any single finding in isolation.

Theoretical and Practical Implications

- Margin width guidelines (e.g., the 2 mm DCIS standard) are not arbitrary; they are a direct response to the branching, under-sampled growth pattern of in situ disease.
- Adjuvant radiation and systemic therapy after re-excision are not simply "extra insurance" but a targeted countermeasure against a wound-healing environment shown to favor residual tumor cell survival.

- A negative margin cannot, on its own, rule out either minimal residual disease or field cancerization, which is part of the rationale for continued long-term surveillance.
- Standardized synoptic reporting is a patient-safety issue, not an administrative preference, given documented gaps in adoption even within well-resourced health systems.
- Because every mechanism above unfolds on a biological timeline, delays introduced by geographic distance from surgical and pathology services are not neutral; they compound the underlying disease process.

Limitations and Future Research Directions

- This paper is anchored to a single published case; while the underlying mechanisms are supported by broader literature, case-based reasoning cannot establish population-level frequencies or effect sizes on its own.
- Much of the wound-microenvironment literature (COX-2, TGF-beta, STAT3, NETosis) is derived from animal models or non-breast cancer types; direct human breast-tissue confirmation of each pathway remains an active area of research.
- Minimal residual disease and field cancerization are best characterized in head and neck cancer; the degree to which the same two-mechanism framework applies to breast DCIS recurrence after negative margins warrants further study.
- Rural and access-related findings are drawn largely from lung cancer and general surgical oncology cohorts; breast-specific, margin-specific access data across varied health systems would strengthen the community-access argument made here.
- Longitudinal studies tracking synoptic reporting adoption alongside re-excision and recurrence rates would help clarify how much of the outcome gap between well-resourced and under-resourced settings is attributable to reporting quality versus access to timely surgery.

References

- American Society of Clinical Oncology. (2020). Closing the rural cancer care gap: Three institutional approaches. JCO Oncology Practice.
- Brown, R. J., Sams, S., Wolverton, D. E., Chong, T., Borges, V. F., Rabinovitch, R. A., & Murphy, C. D. (2017). Don't be fooled by DCIS. *Oncology (Williston Park)*, 31(7), 549–551, 560–561.
- Chauhan, H., Jiwa, N., Nagarajan, V. R., Thiruchelvam, P., Hogben, K., Al-Mufti, R., Hadjiminis, D., Shousha, S., Cutress, R., Ashrafian, H., Takats, Z., & Leff, D. R. (2024). Clinicopathological predictors of positive resection margins in breast-conserving surgery. *Annals of Surgical Oncology*, 31(6), 3939–3947. <https://doi.org/10.1245/s10434-024-15153-8>

- Ekblad, L., Lindgren, G., Persson, E., Kjellén, E., & Wennerberg, J. (2013). Cell-line-specific stimulation of tumor cell aggressiveness by wound healing factors — a central role for STAT3. *BMC Cancer*, 13, 33. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3599402/>
- Hanna, W. M., Parra-Herran, C., Lu, F. I., Slodkowska, E., Nofech-Mozes, S., & Rakovitch, E. (2019). Ductal carcinoma in situ of the breast: An update for the pathologist in the era of individualized risk assessment and tailored therapies. *Modern Pathology*, 32(7), 896–915. <https://www.nature.com/articles/s41379-019-0204-1>
- Johnson, S. M., Miller-Hammond, K., McKenzie-Johnson, T., & Stewart, J. (2024). Disparities in health outcomes in surgical oncology. *Journal of Surgical Oncology*. <https://doi.org/10.1002/jso.27891>
- Minerva, E. M., Tessitore, A., Cafarotti, S., & Patella, M. (2022). Urban–rural disparities in the lung cancer surgical treatment pathway: The paradox of a rich, small region. *Frontiers in Surgery*, 9, 884048. <https://doi.org/10.3389/fsurg.2022.884048>
- Ould Ismail, A. A., Kale, S., McGonagle, K., Hill, B., Pettus, J. R., DuVall, S. L., Ferraro, J. P., & Schroeck, F. R. (2024). Adherence to synoptic cancer pathology reporting among pathologists in the national Department of Veterans Affairs health care system. *Archives of Pathology & Laboratory Medicine*. <https://doi.org/10.5858/arpa.2024-0229-OA>
- Pathology Outlines. (n.d.). Breast malignant — DCIS. Retrieved August 11, 2026, from <https://www.pathologyoutlines.com/topic/breastmalignantdcis.html>
- Pierik, A. S., Leemans, C. R., & Brakenhoff, R. H. (2021). Resection margins in head and neck cancer surgery: An update of residual disease and field cancerization. *Cancers*, 13(11), 2635. <https://doi.org/10.3390/cancers13112635>
- Predina, J. D., Judy, B., Fridlender, Z. G., Aliperti, L. A., Madajewski, B., Kapoor, V., Cheng, G., Quatromoni, J., Okusanya, O., & Singhal, S. (2012). A positive-margin resection model recreates the postsurgical tumor microenvironment and is a reliable model for adjuvant therapy evaluation. *Cancer Biology & Therapy*, 13(9), 745–755. <https://doi.org/10.4161/cbt.20557>
- Torous, V. F., Simpson, R. W., Balani, J. P., Baras, A. S., Berman, M. A., Birdsong, G. G., Giannico, G. A., Paner, G. P., Pettus, J. R., Sessions, Z., Sirintrapun, S. J., Srigley, J. R., & Spencer, S. (2021). College of American Pathologists cancer protocols: From optimizing cancer patient care to facilitating interoperable reporting and downstream data use. *JCO Clinical Cancer Informatics*, 5, 47–55. <https://doi.org/10.1200/CCI.20.00104>

AI statement: AI was consulted in the editing process of this paper.