


3. Find the point  $(x_1, x_2)$  that lies on the line  $x_1 + 2x_2 = 4$  and on the line  $x_1 - x_2 = 1$ . See the figure.



4. Find the point of intersection of the lines  $x_1 + 2x_2 = -13$  and  $3x_1 - 2x_2 = 1$

Determine if the systems in Exercises 15 and 16 are consistent. Do not completely solve the systems.

15. 
$$\begin{aligned} x_1 - 6x_2 &= 5 \\ x_2 - 4x_3 + x_4 &= 0 \\ -x_1 + 6x_2 + x_3 + 5x_4 &= 3 \\ -x_2 + 5x_3 + 4x_4 &= 0 \end{aligned}$$

17. Do the three lines  $2x_1 + 3x_2 = -1$ ,  $6x_1 + 5x_2 = 0$ , and  $2x_1 - 5x_2 = 7$  have a common point of intersection? Explain.

In Exercises 19–22, determine the value(s) of  $h$  such that the matrix is the augmented matrix of a consistent linear system.

19. 
$$\begin{bmatrix} 1 & h & 4 \\ 3 & 6 & 8 \end{bmatrix}$$

20. 
$$\begin{bmatrix} 1 & h & -5 \\ 2 & -8 & 6 \end{bmatrix}$$

21. 
$$\begin{bmatrix} 1 & 4 & -2 \\ 3 & h & -6 \end{bmatrix}$$

22. 
$$\begin{bmatrix} -4 & 12 & h \\ 2 & -6 & -3 \end{bmatrix}$$

25. Find an equation involving  $g$ ,  $h$ , and  $k$  that makes this augmented matrix correspond to a consistent system:

$$\left[ \begin{array}{ccc|c} 1 & -4 & 7 & g \\ 0 & 3 & -5 & h \\ -2 & 5 & -9 & k \end{array} \right]$$

26. Suppose the system below is consistent for all possible values of  $f$  and  $g$ . What can you say about the coefficients  $c$  and  $d$ ? Justify your answer.

$$2x_1 + 4x_2 = f$$

$$cx_1 + dx_2 = g$$

In Exercises 17 and 18, determine the value(s) of  $h$  such that the matrix is the augmented matrix of a consistent linear system.

17.  $\left[ \begin{array}{ccc|c} 1 & -1 & 4 & \\ -2 & 3 & h & \end{array} \right]$

18.  $\left[ \begin{array}{ccc|c} 1 & -3 & 1 & \\ h & 6 & -2 & \end{array} \right]$

In Exercises 1 and 2, compute  $\mathbf{u} + \mathbf{v}$  and  $\mathbf{u} - 2\mathbf{v}$ .

1.  $\mathbf{u} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} -3 \\ -1 \end{bmatrix}$       2.  $\mathbf{u} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$

In Exercises 5 and 6, write a system of equations that is equivalent to the given vector equation.

5.  $x_1 \begin{bmatrix} 3 \\ -2 \\ 8 \end{bmatrix} + x_2 \begin{bmatrix} 5 \\ 0 \\ -9 \end{bmatrix} = \begin{bmatrix} 2 \\ -3 \\ 8 \end{bmatrix}$

6.  $x_1 \begin{bmatrix} 3 \\ -2 \end{bmatrix} + x_2 \begin{bmatrix} 7 \\ 3 \end{bmatrix} + x_3 \begin{bmatrix} -2 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$

In Exercises 9 and 10, write a vector equation that is equivalent to the given system of equations.

9.  $x_2 + 5x_3 = 0$       10.  $3x_1 - 2x_2 + 4x_3 = 3$   
 $4x_1 + 6x_2 - x_3 = 0$        $-2x_1 - 7x_2 + 5x_3 = 1$   
 $-x_1 + 3x_2 - 8x_3 = 0$        $5x_1 + 4x_2 - 3x_3 = 2$

In Exercises 11 and 12, determine if  $\mathbf{b}$  is a linear combination of  $\mathbf{a}_1, \mathbf{a}_2$ , and  $\mathbf{a}_3$ .

11.  $\mathbf{a}_1 = \begin{bmatrix} 1 \\ -2 \\ 0 \end{bmatrix}, \mathbf{a}_2 = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix}, \mathbf{a}_3 = \begin{bmatrix} 5 \\ -6 \\ 8 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 2 \\ -1 \\ 6 \end{bmatrix}$

12.  $\mathbf{a}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \mathbf{a}_2 = \begin{bmatrix} -2 \\ 3 \\ -2 \end{bmatrix}, \mathbf{a}_3 = \begin{bmatrix} -6 \\ 7 \\ 5 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 11 \\ -5 \\ 9 \end{bmatrix}$

In Exercises 13 and 14, determine if  $\mathbf{b}$  is a linear combination of the vectors formed from the columns of the matrix  $A$ .

13.  $A = \begin{bmatrix} 1 & -4 & 2 \\ 0 & 3 & 5 \\ -2 & 8 & -4 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 3 \\ -7 \\ -3 \end{bmatrix}$

14.  $A = \begin{bmatrix} 1 & 0 & 5 \\ -2 & 1 & -6 \\ 0 & 2 & 8 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 2 \\ -1 \\ 6 \end{bmatrix}$

15. Let  $\mathbf{a}_1 = \begin{bmatrix} 1 \\ 3 \\ -1 \end{bmatrix}$ ,  $\mathbf{a}_2 = \begin{bmatrix} -5 \\ -8 \\ 2 \end{bmatrix}$ , and  $\mathbf{b} = \begin{bmatrix} 3 \\ -5 \\ h \end{bmatrix}$ . For what value(s) of  $h$  is  $\mathbf{b}$  in the plane spanned by  $\mathbf{a}_1$  and  $\mathbf{a}_2$ ?

16. Let  $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 0 \\ -2 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} -2 \\ 1 \\ 7 \end{bmatrix}$ , and  $\mathbf{y} = \begin{bmatrix} h \\ -3 \\ -5 \end{bmatrix}$ . For what value(s) of  $h$  is  $\mathbf{y}$  in the plane generated by  $\mathbf{v}_1$  and  $\mathbf{v}_2$ ?


In Exercises 17 and 18, list five vectors in  $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2\}$ . For each vector, show the weights on  $\mathbf{v}_1$  and  $\mathbf{v}_2$  used to generate the vector and list the three entries of the vector. Do not make a sketch.

17.  $\mathbf{v}_1 = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} -4 \\ 0 \\ 1 \end{bmatrix}$

21. Let  $\mathbf{u} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$  and  $\mathbf{v} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ . Show that  $\begin{bmatrix} h \\ k \end{bmatrix}$  is in  $\text{Span}\{\mathbf{u}, \mathbf{v}\}$  for all  $h$  and  $k$ .

32. Consider the vectors  $\mathbf{v}_1$ ,  $\mathbf{v}_2$ ,  $\mathbf{v}_3$ , and  $\mathbf{b}$  in  $\mathbb{R}^2$ , shown in the figure. Does the equation  $x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + x_3\mathbf{v}_3 = \mathbf{b}$  have a

solution? Is the solution unique? Use the figure to explain your answers.





## 1.4 EXERCISES

1.  $\begin{bmatrix} -4 & 2 \\ 1 & 6 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ -2 \\ 7 \end{bmatrix}$

multiply  
←

Write as a vector Eq<sup>n</sup>s

5.  $\begin{bmatrix} 1 & 2 & -3 & 1 \\ -2 & -3 & 1 & -1 \end{bmatrix} \begin{bmatrix} 2 \\ -1 \\ 1 \\ -1 \end{bmatrix} = \begin{bmatrix} -4 \\ 1 \end{bmatrix}$

write As matrix Equation and vector Equation

9.  $5x_1 + x_2 - 3x_3 = 8$

$2x_2 + 4x_3 = 0$

10.  $4x_1 - x_2 = 8$

$5x_1 + 3x_2 = 2$

$3x_1 - x_2 = 1$

Given  $A$  and  $\mathbf{b}$  in Exercises 11 and 12, write the augmented matrix for the linear system that corresponds to the matrix equation  $A\mathbf{x} = \mathbf{b}$ . Then solve the system and write the solution as a vector.

11.  $A = \begin{bmatrix} 1 & 3 & -4 \\ 1 & 5 & 2 \\ -3 & -7 & 6 \end{bmatrix}$ ,  $\mathbf{b} = \begin{bmatrix} -2 \\ 4 \\ 12 \end{bmatrix}$

13. Let  $\mathbf{u} = \begin{bmatrix} 0 \\ 4 \\ 4 \end{bmatrix}$  and  $A = \begin{bmatrix} 3 & -5 \\ -2 & 6 \\ 1 & 1 \end{bmatrix}$ . Is  $\mathbf{u}$  in the plane in

$\mathbb{R}^3$  spanned by the columns of  $A$ ? (See the figure.) Why or why not?

•  $\mathbf{u}$ ?

15. Let  $A = \begin{bmatrix} 3 & -1 \\ -9 & 3 \end{bmatrix}$  and  $\mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$ . Show that the equation  $A\mathbf{x} = \mathbf{b}$  does not have a solution for all possible  $\mathbf{b}$ , and describe the set of all  $\mathbf{b}$  for which  $A\mathbf{x} = \mathbf{b}$  does have a solution.

22. Let  $\mathbf{v}_1 = \begin{bmatrix} 0 \\ 0 \\ -3 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} 0 \\ -3 \\ 9 \end{bmatrix}$ ,  $\mathbf{v}_3 = \begin{bmatrix} 4 \\ -2 \\ -6 \end{bmatrix}$ . Does

$\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  span  $\mathbb{R}^3$ ? Why or why not?

25. Note that  $\begin{bmatrix} 4 & -3 & 1 \\ 5 & -2 & 5 \\ -6 & 2 & -3 \end{bmatrix} \begin{bmatrix} -3 \\ -1 \\ 2 \end{bmatrix} = \begin{bmatrix} -7 \\ -3 \\ 10 \end{bmatrix}$ . Use this

fact (and no row operations) to find scalars  $c_1, c_2, c_3$  such

that  $\begin{bmatrix} -7 \\ -3 \\ 10 \end{bmatrix} = c_1 \begin{bmatrix} 4 \\ 5 \\ -6 \end{bmatrix} + c_2 \begin{bmatrix} -3 \\ -2 \\ 2 \end{bmatrix} + c_3 \begin{bmatrix} 1 \\ 5 \\ -3 \end{bmatrix}$ .

26. Let  $\mathbf{u} = \begin{bmatrix} 7 \\ 2 \\ 5 \end{bmatrix}$ ,  $\mathbf{v} = \begin{bmatrix} 3 \\ 1 \\ 3 \end{bmatrix}$ , and  $\mathbf{w} = \begin{bmatrix} 5 \\ 1 \\ 1 \end{bmatrix}$ . It can be

shown that  $2\mathbf{u} - 3\mathbf{v} - \mathbf{w} = \mathbf{0}$ . Use this fact (and no row operations) to find  $x_1$  and  $x_2$  that satisfy the equation

$$\begin{bmatrix} 7 & 3 \\ 2 & 1 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 5 \\ 1 \\ 1 \end{bmatrix}.$$

## 1.5 EXERCISES

In Exercises 1–4, determine if the system has a nontrivial solution. Try to use as few row operations as possible.

$$\begin{array}{ll} \mathbf{1.} & \begin{array}{l} 2x_1 - 5x_2 + 8x_3 = 0 \\ -2x_1 - 7x_2 + x_3 = 0 \\ 4x_1 + 2x_2 + 7x_3 = 0 \end{array} \\ \mathbf{2.} & \begin{array}{l} x_1 - 2x_2 + 3x_3 = 0 \\ -2x_1 - 3x_2 - 4x_3 = 0 \\ 2x_1 - 4x_2 + 9x_3 = 0 \end{array} \end{array}$$

**18.** As in Exercise 17, describe the solutions of the following system in parametric vector form, and provide a geometric comparison with the solution set in Exercise 6.

$$\begin{array}{rcl} x_1 + 2x_2 - 3x_3 & = & 5 \\ 2x_1 + x_2 - 3x_3 & = & 13 \\ -x_1 + x_2 & = & -8 \end{array}$$

In Exercises 19 and 20, find the parametric equation of the line through **a** parallel to **b**.


$$\mathbf{19.} \quad \mathbf{a} = \begin{bmatrix} -2 \\ 0 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} -5 \\ 3 \end{bmatrix} \quad \mathbf{20.} \quad \mathbf{a} = \begin{bmatrix} 3 \\ -2 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} -7 \\ 6 \end{bmatrix}$$



In Exercises 21 and 22, find a parametric equation of the line  $M$  through  $\mathbf{p}$  and  $\mathbf{q}$ . [Hint:  $M$  is parallel to the vector  $\mathbf{q} - \mathbf{p}$ . See the figure below.]

21.  $\mathbf{p} = \begin{bmatrix} 3 \\ -3 \end{bmatrix}, \mathbf{q} = \begin{bmatrix} 4 \\ 1 \end{bmatrix}$

22.  $\mathbf{p} = \begin{bmatrix} -3 \\ 2 \end{bmatrix}, \mathbf{q} = \begin{bmatrix} 0 \\ -3 \end{bmatrix}$



check pages  
44, 45, 46

## 1.7 EXERCISES

In Exercises 1–4, determine if the vectors are linearly independent. Justify each answer.

1.  $\begin{bmatrix} 5 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 7 \\ 2 \\ -6 \end{bmatrix}, \begin{bmatrix} 9 \\ 4 \\ -8 \end{bmatrix}$       2.  $\begin{bmatrix} 0 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ -8 \end{bmatrix}, \begin{bmatrix} -1 \\ 3 \\ 1 \end{bmatrix}$

3.  $\begin{bmatrix} 2 \\ -3 \end{bmatrix}, \begin{bmatrix} -4 \\ 6 \end{bmatrix}$       4.  $\begin{bmatrix} -1 \\ 3 \end{bmatrix}, \begin{bmatrix} -3 \\ -9 \end{bmatrix}$

In Exercises 9 and 10, (a) for what values of  $h$  is  $\mathbf{v}_3$  in  $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2\}$ , and (b) for what values of  $h$  is  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  linearly dependent? Justify each answer.

9.  $\mathbf{v}_1 = \begin{bmatrix} 1 \\ -3 \\ 2 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} -3 \\ 9 \\ -6 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} 5 \\ -7 \\ h \end{bmatrix}$

10.  $\mathbf{v}_1 = \begin{bmatrix} 1 \\ -3 \\ -5 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} -3 \\ 9 \\ 15 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} 2 \\ -5 \\ h \end{bmatrix}$