

STATISTICAL ANALYSIS - LECTURE 6

Dr. Mahmoud Mounir

mahmoud.mounir@cis.asu.edu.eg

The Normal Distribution


Note: This PowerPoint is only a summary and your main source should be the book.

Introduction

- Normal Distribution.
- Applications of the Normal Distribution.


The Normal Distribution

Note: This PowerPoint is only a summary and your main source should be the book.




Mean Median Mode

(b) Negatively skewed




Mode Median Mean

(c) Positively skewed



→
**(a) Normal
Mean =Median=Mode**

□ A normal distribution is a continuous , symmetric , bell shaped distribution of a variable.



□ A normal distribution curve depend on two parameters .

μ → Position parameter

σ → shape parameter

The mathematical equation for the normal distribution:

$$y = \frac{e^{-(x-\mu)^2/2\sigma^2}}{\sigma\sqrt{2\pi}}$$

where


$e \approx 2.718$

$\pi \approx 3.14$

$\mu \approx$ population mean

$\sigma \approx$ population standard deviation


1



Normal curves
with $\mu_1 \neq \mu_2$ and
 $\sigma_1 = \sigma_2$

(1) Different
means but same
standard
deviations.


2



Normal curves
with $\mu_1 = \mu_2$ and
 $\sigma_1 < \sigma_2$

(2) Same means
but different
standard
deviations .


3



Normal curves
with $\mu_1 < \mu_2$ and
 $\sigma_1 < \sigma_2$

(3) Different
means and
different standard
deviations .

Properties of the Normal Distribution




- The normal distribution curve is **bell-shaped**.
- The mean, median, and mode are **equal** and located at the center of the distribution.
- The normal distribution curve is **unimodal** (single mode).
- The curve is **symmetrical** about the mean.
- The curve is **continuous**.
- The curve **never touches the x -axis**.
- The total area under the normal distribution curve is **equal to 1 or 100%**.

- The area under **the normal distribution curve** that lies within
 - one standard deviation of the mean is approximately 0.68 (68%). The interval between $(\bar{x} - s, \bar{x} + s)$
 - two standard deviations of the mean is approximately 0.95 (95%). The interval between $(\bar{x} - 2s, \bar{x} + 2s)$
 - three standard deviations of the mean is approximately 0.997 (99.7%). The interval between $(\bar{x} - 3s, \bar{x} + 3s)$

Empirical Rule: Normal Distribution

(Areas Under the Normal Curve)



Note: This PowerPoint is only a summary and your main source should be the book.

The Standard Normal Distribution

- The standard normal distribution is a normal distribution with a mean of 0 and a standard deviation of 1.
- The formula for the standard normal distribution is

$$y = \frac{e^{\frac{-z^2}{2}}}{\sqrt{2\pi}}$$

The diagram illustrates the formula for the standard normal distribution probability density function. The formula is $y = \frac{e^{\frac{-z^2}{2}}}{\sqrt{2\pi}}$. Two arrows point from the formula to two red-bordered boxes. The top arrow points to the term $\mu = 0$, and the bottom arrow points to the term $\sigma = 1$.


- All Normal Distribution can be transformed into standard Distribution.

$$z = \frac{\text{value} - \text{mean}}{\text{standard deviation}}$$


or

$$z = \frac{x - \mu}{\sigma}$$

Empirical Rule: Standard Normal Distribution



$$\begin{array}{l} \mu = 0 \\ \sigma = 1 \end{array}$$



Normal Distribution Curve


Standard Normal Distribution Curve

- The area under the **standard normal distribution curve** that lies within
 - one standard deviation of the mean is approximately 0.68 (68%). The interval between (-1,1).
 - two standard deviations of the mean is approximately 0.95 (95%). The interval between (-2,2).
 - three standard deviations of the mean is approximately 0.997 (99.7%). The interval between (-3,3).


Finding Areas Under the Standard Normal Distribution Curve:

1. To the left of any Z value

(a)




=



$$P(z < -a) = P(z > a) = Q(a)$$


2. To the right of any Z value

(b)




$$P(z > -a) = P(z > a) = 1 - Q(a)$$


3.Between any two Z values




$$P(-b < z < -a) = P(a < z < b) = P(z > a) - P(z > b)$$

$$= Q(a) - Q(b)$$






<i>z</i>	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
-3.4	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0002
-3.3	0.0005	0.0005	0.0005	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0003
-3.2	0.0007	0.0007	0.0006	0.0006	0.0006	0.0006	0.0006	0.0005	0.0005	0.0005
-3.1	0.0010	0.0009	0.0009	0.0009	0.0008	0.0008	0.0008	0.0008	0.0007	0.0007
-3.0	0.0013	0.0013	0.0013	0.0012	0.0012	0.0011	0.0011	0.0011	0.0010	0.0010
-2.9	0.0019	0.0018	0.0018	0.0017	0.0016	0.0016	0.0015	0.0015	0.0014	0.0014
-2.8	0.0026	0.0025	0.0024	0.0023	0.0023	0.0022	0.0021	0.0021	0.0020	0.0019
-2.7	0.0035	0.0034	0.0033	0.0032	0.0031	0.0030	0.0029	0.0028	0.0027	0.0026
-2.6	0.0047	0.0045	0.0044	0.0043	0.0041	0.0040	0.0039	0.0038	0.0037	0.0036
-2.5	0.0062	0.0060	0.0059	0.0057	0.0055	0.0054	0.0052	0.0051	0.0049	0.0048
-2.4	0.0082	0.0080	0.0078	0.0075	0.0073	0.0071	0.0069	0.0068	0.0066	0.0064
-2.3	0.0107	0.0104	0.0102	0.0099	0.0096	0.0094	0.0091	0.0089	0.0087	0.0084
-2.2	0.0139	0.0136	0.0132	0.0129	0.0125	0.0122	0.0119	0.0116	0.0113	0.0110
-2.1	0.0179	0.0174	0.0170	0.0166	0.0162	0.0158	0.0154	0.0150	0.0146	0.0143
-2.0	0.0228	0.0222	0.0217	0.0212	0.0207	0.0202	0.0197	0.0192	0.0188	0.0183
-1.9	0.0287	0.0281	0.0274	0.0268	0.0262	0.0256	0.0250	0.0244	0.0239	0.0233
-1.8	0.0359	0.0351	0.0344	0.0336	0.0329	0.0322	0.0314	0.0307	0.0301	0.0294
-1.7	0.0446	0.0436	0.0427	0.0418	0.0409	0.0401	0.0392	0.0384	0.0375	0.0367
-1.6	0.0548	0.0537	0.0526	0.0516	0.0505	0.0495	0.0485	0.0475	0.0465	0.0455
-1.5	0.0668	0.0655	0.0643	0.0630	0.0618	0.0606	0.0594	0.0582	0.0571	0.0559
-1.4	0.0808	0.0793	0.0778	0.0764	0.0749	0.0735	0.0721	0.0708	0.0694	0.0681
-1.3	0.0968	0.0951	0.0934	0.0918	0.0901	0.0885	0.0869	0.0853	0.0838	0.0823
-1.2	0.1151	0.1131	0.1112	0.1093	0.1075	0.1056	0.1038	0.1020	0.1003	0.0985
-1.1	0.1357	0.1335	0.1314	0.1292	0.1271	0.1251	0.1230	0.1210	0.1190	0.1170
-1.0	0.1587	0.1562	0.1539	0.1515	0.1492	0.1469	0.1446	0.1423	0.1401	0.1379
-0.9	0.1841	0.1814	0.1788	0.1762	0.1736	0.1711	0.1685	0.1660	0.1635	0.1611
-0.8	0.2119	0.2090	0.2061	0.2033	0.2005	0.1977	0.1949	0.1922	0.1894	0.1867
-0.7	0.2420	0.2389	0.2358	0.2327	0.2296	0.2266	0.2236	0.2206	0.2177	0.2148
-0.6	0.2743	0.2709	0.2676	0.2643	0.2611	0.2578	0.2546	0.2514	0.2483	0.2451
-0.5	0.3085	0.3050	0.3015	0.2981	0.2946	0.2912	0.2877	0.2843	0.2810	0.2776
-0.4	0.3446	0.3409	0.3372	0.3336	0.3300	0.3264	0.3228	0.3192	0.3156	0.3121
-0.3	0.3821	0.3783	0.3745	0.3707	0.3669	0.3632	0.3594	0.3557	0.3520	0.3483
-0.2	0.4207	0.4168	0.4129	0.4090	0.4052	0.4013	0.3974	0.3936	0.3897	0.3859
-0.1	0.4602	0.4562	0.4522	0.4483	0.4443	0.4404	0.4364	0.4325	0.4286	0.4247
-0.0	0.5000	0.4960	0.4920	0.4880	0.4840	0.4801	0.4761	0.4721	0.4681	0.4641



z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319

Examples


a. Area to the left of $z = 1.36$: $P(z < 1.36)$



$$= 0.9131$$

Examples

b. Area to the left of $z = -0.60$: $P(z < -0.60)$




= 0.2743

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
-3.4	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0002
-3.3	0.0005	0.0005	0.0005	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0003
-3.2	0.0007	0.0007	0.0006	0.0006	0.0006	0.0006	0.0006	0.0005	0.0005	0.0005
-3.1	0.0010	0.0009	0.0009	0.0009	0.0008	0.0008	0.0008	0.0008	0.0007	0.0007
-3.0	0.0013	0.0013	0.0013	0.0012	0.0012	0.0011	0.0011	0.0011	0.0010	0.0010
-2.9	0.0019	0.0018	0.0018	0.0017	0.0016	0.0016	0.0015	0.0015	0.0014	0.0014
-2.8	0.0026	0.0025	0.0024	0.0023	0.0023	0.0022	0.0021	0.0021	0.0020	0.0019
-2.7	0.0035	0.0034	0.0033	0.0032	0.0031	0.0030	0.0029	0.0028	0.0027	0.0026
-2.6	0.0047	0.0045	0.0044	0.0043	0.0041	0.0040	0.0039	0.0038	0.0037	0.0036
-2.5	0.0062	0.0060	0.0059	0.0057	0.0055	0.0054	0.0052	0.0051	0.0049	0.0048
-2.4	0.0082	0.0080	0.0078	0.0075	0.0073	0.0071	0.0069	0.0068	0.0066	0.0064
-2.3	0.0107	0.0104	0.0102	0.0099	0.0096	0.0094	0.0091	0.0089	0.0087	0.0084
-2.2	0.0139	0.0136	0.0132	0.0129	0.0125	0.0122	0.0119	0.0116	0.0113	0.0110
-2.1	0.0179	0.0174	0.0170	0.0166	0.0162	0.0158	0.0154	0.0150	0.0146	0.0143
-2.0	0.0228	0.0222	0.0217	0.0212	0.0207	0.0202	0.0197	0.0192	0.0188	0.0183
-1.9	0.0287	0.0281	0.0274	0.0268	0.0262	0.0256	0.0250	0.0244	0.0239	0.0233
-1.8	0.0359	0.0351	0.0344	0.0336	0.0329	0.0322	0.0314	0.0307	0.0301	0.0294
-1.7	0.0446	0.0436	0.0427	0.0418	0.0409	0.0401	0.0392	0.0384	0.0375	0.0367
-1.6	0.0548	0.0537	0.0526	0.0516	0.0505	0.0495	0.0485	0.0475	0.0465	0.0455
-1.5	0.0668	0.0655	0.0643	0.0630	0.0618	0.0606	0.0594	0.0582	0.0571	0.0559
-1.4	0.0808	0.0793	0.0778	0.0764	0.0749	0.0735	0.0721	0.0708	0.0694	0.0681
-1.3	0.0968	0.0951	0.0934	0.0918	0.0901	0.0885	0.0869	0.0853	0.0838	0.0823
-1.2	0.1151	0.1131	0.1112	0.1093	0.1075	0.1056	0.1038	0.1020	0.1003	0.0985
-1.1	0.1357	0.1335	0.1314	0.1292	0.1271	0.1251	0.1230	0.1210	0.1190	0.1170
-1.0	0.1587	0.1562	0.1539	0.1515	0.1492	0.1469	0.1446	0.1423	0.1401	0.1379
-0.9	0.1841	0.1814	0.1788	0.1762	0.1736	0.1711	0.1685	0.1660	0.1635	0.1611
-0.8	0.2119	0.2090	0.2061	0.2033	0.2005	0.1977	0.1949	0.1922	0.1894	0.1867
-0.7	0.2420	0.2389	0.2358	0.2327	0.2296	0.2266	0.2236	0.2206	0.2177	0.2148
-0.6	0.2743	0.2709	0.2676	0.2643	0.2611	0.2578	0.2546	0.2514	0.2483	0.2451
-0.5	0.3085	0.3050	0.3015	0.2981	0.2946	0.2912	0.2877	0.2843	0.2810	0.2776
-0.4	0.3446	0.3409	0.3372	0.3336	0.3300	0.3264	0.3228	0.3192	0.3156	0.3121
-0.3	0.3821	0.3783	0.3745	0.3707	0.3669	0.3632	0.3594	0.3557	0.3520	0.3483
-0.2	0.4207	0.4168	0.4129	0.4090	0.4052	0.4013	0.3974	0.3936	0.3897	0.3859
-0.1	0.4602	0.4562	0.4522	0.4483	0.4443	0.4404	0.4364	0.4325	0.4286	0.4247
-0.0	0.5000	0.4960	0.4920	0.4880	0.4840	0.4801	0.4761	0.4721	0.4681	0.4641

Examples


c. Area to the right of $z = 1.47$: $P(z > 1.47)$



$$\begin{aligned}
 &= 1 - P(z < 1.47) \\
 &= 1 - 0.9292 \\
 &= 0.0708
 \end{aligned}$$

Examples

d. Area to the right of $z = -0.33$: $P(z > -0.33)$




$$\begin{aligned}
 &= 1 - P(z < -0.33) \\
 &= 1 - 0.3707 \\
 &= 0.6293
 \end{aligned}$$

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
-3.4	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0002
-3.3	0.0005	0.0005	0.0005	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0003
-3.2	0.0007	0.0007	0.0006	0.0006	0.0006	0.0006	0.0006	0.0005	0.0005	0.0005
-3.1	0.0010	0.0009	0.0009	0.0009	0.0008	0.0008	0.0008	0.0008	0.0007	0.0007
-3.0	0.0013	0.0013	0.0013	0.0012	0.0012	0.0011	0.0011	0.0011	0.0010	0.0010
-2.9	0.0019	0.0018	0.0018	0.0017	0.0016	0.0016	0.0015	0.0015	0.0014	0.0014
-2.8	0.0026	0.0025	0.0024	0.0023	0.0023	0.0022	0.0021	0.0021	0.0020	0.0019
-2.7	0.0035	0.0034	0.0033	0.0032	0.0031	0.0030	0.0029	0.0028	0.0027	0.0026
-2.6	0.0047	0.0045	0.0044	0.0043	0.0041	0.0040	0.0039	0.0038	0.0037	0.0036
-2.5	0.0062	0.0060	0.0059	0.0057	0.0055	0.0054	0.0052	0.0051	0.0049	0.0048
-2.4	0.0082	0.0080	0.0078	0.0075	0.0073	0.0071	0.0069	0.0068	0.0066	0.0064
-2.3	0.0107	0.0104	0.0102	0.0099	0.0096	0.0094	0.0091	0.0089	0.0087	0.0084
-2.2	0.0139	0.0136	0.0132	0.0129	0.0125	0.0122	0.0119	0.0116	0.0113	0.0110
-2.1	0.0179	0.0174	0.0170	0.0166	0.0162	0.0158	0.0154	0.0150	0.0146	0.0143
-2.0	0.0228	0.0222	0.0217	0.0212	0.0207	0.0202	0.0197	0.0192	0.0188	0.0183
-1.9	0.0287	0.0281	0.0274	0.0268	0.0262	0.0256	0.0250	0.0244	0.0239	0.0233
-1.8	0.0359	0.0351	0.0344	0.0336	0.0329	0.0322	0.0314	0.0307	0.0301	0.0294
-1.7	0.0446	0.0436	0.0427	0.0418	0.0409	0.0401	0.0392	0.0384	0.0375	0.0367
-1.6	0.0548	0.0537	0.0526	0.0516	0.0505	0.0495	0.0485	0.0475	0.0465	0.0455
-1.5	0.0668	0.0655	0.0643	0.0630	0.0618	0.0606	0.0594	0.0582	0.0571	0.0559
-1.4	0.0808	0.0793	0.0778	0.0764	0.0749	0.0735	0.0721	0.0708	0.0694	0.0681
-1.3	0.0968	0.0951	0.0934	0.0918	0.0901	0.0885	0.0869	0.0853	0.0838	0.0823
-1.2	0.1151	0.1131	0.1112	0.1093	0.1075	0.1056	0.1038	0.1020	0.1003	0.0985
-1.1	0.1357	0.1335	0.1314	0.1292	0.1271	0.1251	0.1230	0.1210	0.1190	0.1170
-1.0	0.1587	0.1562	0.1539	0.1515	0.1492	0.1469	0.1446	0.1423	0.1401	0.1379
-0.9	0.1841	0.1814	0.1788	0.1762	0.1736	0.1711	0.1685	0.1660	0.1635	0.1611
-0.8	0.2119	0.2090	0.2061	0.2033	0.2005	0.1977	0.1949	0.1922	0.1894	0.1867
-0.7	0.2420	0.2389	0.2358	0.2327	0.2296	0.2266	0.2236	0.2206	0.2177	0.2148
-0.6	0.2743	0.2709	0.2676	0.2643	0.2611	0.2578	0.2546	0.2514	0.2483	0.2451
-0.5	0.3085	0.3050	0.3015	0.2981	0.2946	0.2912	0.2877	0.2843	0.2810	0.2776
-0.4	0.3446	0.3409	0.3372	0.3336	0.3300	0.3264	0.3228	0.3192	0.3156	0.3121
-0.3	0.3821	0.3783	0.3745	0.3707	0.3669	0.3632	0.3594	0.3557	0.3520	0.3483
-0.2	0.4207	0.4168	0.4129	0.4090	0.4052	0.4013	0.3974	0.3936	0.3897	0.3859
-0.1	0.4602	0.4562	0.4522	0.4483	0.4443	0.4404	0.4364	0.4325	0.4286	0.4247
-0.0	0.5000	0.4960	0.4920	0.4880	0.4840	0.4801	0.4761	0.4721	0.4681	0.4641

Examples

e. Area between $z = -2.16$ and $z = 0.45$: $P(-2.16 < z < 0.45)$




$$\begin{aligned}
 &= P(z < 0.45) - P(z < -2.16) \\
 &= 0.6736 - 0.0154 \\
 &= 0.6582
 \end{aligned}$$

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
-3.4	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0002
-3.3	0.0005	0.0005	0.0005	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0003
-3.2	0.0007	0.0007	0.0006	0.0006	0.0006	0.0006	0.0006	0.0006	0.0005	0.0005
-3.1	0.0010	0.0009	0.0009	0.0009	0.0008	0.0008	0.0008	0.0008	0.0007	0.0007
-3.0	0.0013	0.0013	0.0013	0.0012	0.0012	0.0011	0.0011	0.0011	0.0010	0.0010
-2.9	0.0019	0.0018	0.0018	0.0017	0.0016	0.0016	0.0015	0.0015	0.0014	0.0014
-2.8	0.0026	0.0025	0.0024	0.0023	0.0023	0.0022	0.0021	0.0021	0.0020	0.0019
-2.7	0.0035	0.0034	0.0033	0.0032	0.0031	0.0030	0.0029	0.0028	0.0027	0.0026
-2.6	0.0047	0.0045	0.0044	0.0043	0.0041	0.0040	0.0039	0.0038	0.0037	0.0036
-2.5	0.0062	0.0060	0.0059	0.0057	0.0055	0.0054	0.0052	0.0051	0.0049	0.0048
-2.4	0.0082	0.0080	0.0078	0.0075	0.0073	0.0071	0.0069	0.0068	0.0066	0.0064
-2.3	0.0107	0.0104	0.0102	0.0099	0.0096	0.0094	0.0091	0.0089	0.0087	0.0084
-2.2	0.0139	0.0136	0.0132	0.0129	0.0125	0.0122	0.0119	0.0116	0.0113	0.0110
-2.1	0.0179	0.0174	0.0170	0.0166	0.0162	0.0158	0.0154	0.0150	0.0146	0.0143
-2.0	0.0228	0.0222	0.0217	0.0212	0.0207	0.0202	0.0197	0.0192	0.0188	0.0183
-1.9	0.0287	0.0281	0.0274	0.0268	0.0262	0.0256	0.0250	0.0244	0.0239	0.0233
-1.8	0.0359	0.0351	0.0344	0.0336	0.0329	0.0322	0.0314	0.0307	0.0301	0.0294
-1.7	0.0446	0.0436	0.0427	0.0418	0.0409	0.0401	0.0392	0.0384	0.0375	0.0367
-1.6	0.0548	0.0537	0.0526	0.0516	0.0505	0.0495	0.0485	0.0475	0.0465	0.0455
-1.5	0.0668	0.0655	0.0643	0.0630	0.0618	0.0606	0.0594	0.0582	0.0571	0.0559
-1.4	0.0808	0.0793	0.0778	0.0764	0.0749	0.0735	0.0721	0.0708	0.0694	0.0681
-1.3	0.0968	0.0951	0.0934	0.0918	0.0901	0.0885	0.0869	0.0853	0.0838	0.0823
-1.2	0.1151	0.1131	0.1112	0.1093	0.1075	0.1056	0.1038	0.1020	0.1003	0.0985
-1.1	0.1357	0.1335	0.1314	0.1292	0.1271	0.1251	0.1230	0.1210	0.1190	0.1170
-1.0	0.1587	0.1562	0.1539	0.1515	0.1492	0.1469	0.1446	0.1423	0.1401	0.1379
-0.9	0.1841	0.1814	0.1788	0.1762	0.1736	0.1711	0.1685	0.1660	0.1635	0.1611
-0.8	0.2119	0.2090	0.2061	0.2033	0.2005	0.1977	0.1949	0.1922	0.1894	0.1867
-0.7	0.2420	0.2389	0.2358	0.2327	0.2296	0.2266	0.2236	0.2206	0.2177	0.2148
-0.6	0.2743	0.2709	0.2676	0.2643	0.2611	0.2578	0.2546	0.2514	0.2483	0.2451
-0.5	0.3085	0.3050	0.3015	0.2981	0.2946	0.2912	0.2877	0.2843	0.2810	0.2776
-0.4	0.3446	0.3409	0.3372	0.3336	0.3300	0.3264	0.3228	0.3192	0.3156	0.3121
-0.3	0.3821	0.3783	0.3745	0.3707	0.3669	0.3632	0.3594	0.3557	0.3520	0.3483
-0.2	0.4207	0.4168	0.4129	0.4090	0.4052	0.4013	0.3974	0.3936	0.3897	0.3859
-0.1	0.4602	0.4562	0.4522	0.4483	0.4443	0.4404	0.4364	0.4325	0.4286	0.4247
-0.0	0.5000	0.4960	0.4920	0.4880	0.4840	0.4801	0.4761	0.4721	0.4681	0.4641

Examples


f. Area between $z = 1.13$ and $z = 3.10$: $P(1.13 < z < 3.10)$



$$\begin{aligned}
 &= P(z < 3.1) - P(z < 1.13) \\
 &= 0.9990 - 0.8708 \\
 &= 0.1282
 \end{aligned}$$

Example (1):

Find the area to the left of $z=1.99$



$$P(Z < 1.99) = 0.9767$$

Column 1.9 row

Table E (continued)


Cumulative Standard Normal Distribution

<i>z</i>	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817

Note: This PowerPoint is only a summary and your main source should be the book.

Example (2):


Find the area to the right of $z=-1.16$



$$\begin{aligned} P(Z > -1.16) &= 1 - P(Z < -1.16) \\ &= 1 - 0.1230 \\ &= 0.8770 \end{aligned}$$

Example (3):

Find the area between $z=1.68$ and $z=-1.37$



$$\begin{aligned} P(-1.37 < Z < 1.68) &= P(Z < 1.68) - P(Z < -1.37) \\ &= 0.9535 - 0.0853 \\ &= 0.8682 \end{aligned}$$


A Normal Distribution Curve as a Probability Distribution Curve:

- The area under the standard normal distribution curve can also be thought of as a probability .


Example (4):

Find probability for each


- a) $P(0 < z < 2.23)$
- b) $P(z < 1.65)$
- c) $P(z > 1.91)$



$$\begin{aligned} \text{a) } P(0 < Z < 2.23) &= P(Z < 2.23) - P(Z < 0) \\ &= 0.9898 - 0.5000 \\ &= 0.4898 \end{aligned}$$




b) $P(Z < 1.65) = 0.9505$



c) $P(Z > 1.91) = 1 - P(Z < 1.91)$
 $= 1 - 0.9719$
 $= 0.0281$


Find the Z value that corresponds to given area

1




we cannot find the area in the table

2




Look up the area in table E to
find the z value

3



we cannot find the area in the table


4



Look up the area in table E to
find the z value


For examples 5 to 12:

5




$$z = -2.08$$

6



$$z = 1.84$$


7



$$1 - 0.0239 = 0.9761$$


$$z = 1.98$$

8




$$1 - 0.8962 = 0.1038$$

$$z = -1.26$$

5


$$0.5000 + a = \text{----}$$

z = look in the table E

10


$$0.5000 - a = \text{----}$$

z = look in the table E

11

$$0.5000 + 0.4066 = 0.9066$$

$z = 1.32$


12

$$0.5000 - 0.4175 = 0.0825$$

$z = -1.39$

Example (13):

Find the z value such that the area under the standard normal distribution curve between 0 and the z value is 0.2123



$$0.2123 + 0.5000 = 0.7123$$

$$Z=0.56$$

Table E *(continued)*

Cumulative Standard Normal Distribution

<i>z</i>	.00	.01	.02	.03	.04	.05	.06	.07
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577

Note: This PowerPoint is only a summary and your main source should be the book.

Questions ???



- 1) Find the z value to the left of the mean so that 98.87% of the area under the distribution curve lies to the right of it.
- 2) Find two z values so that 48% of the middle area is bounded by them.
- 3) Find two z values, one positive and one negative, that are equidistant from the mean so that the area in the two tails total the following values (5%).

Example (1):

A survey by the National Retail Federation found that women spend on average \$146.21 for the Christmas holidays. Assume the standard deviation is \$29.44. Find the **percentage** of women who spend less than \$160. Assume the variable is **normally distributed**.


Solution:

FIND $P(X < 160)$

Step 1: Find the z value .

$$z = \frac{X - \mu}{\sigma} = \frac{160 - 146.21}{29.44} = 0.47$$

Step 2: Draw the figure



Step 3: Find the area using Z-table

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879

$$P(Z < 0.47) = \underline{\underline{0.6808}}$$

68.08% of the women spend less than 160\$ at Christmas time.

Example (2):

Each month, an American household generates an average of 28 pounds of newspaper for garbage or recycling. Assume the standard deviation is 2 pounds. If a household is selected at random, Find the probability of its generating.

- a) Between 27 and 31 pounds per month
- b) More than 30.2 pounds per month

Assume the variable is approximately **normally distributed**.

Solution (a) :


FIND $P(27 < X < 31)$

Step 1: Find the two z values

$$Z_1 = \frac{X - \mu}{\sigma} = \frac{27 - 28}{2} = -0.5$$

$$Z_2 = \frac{X - \mu}{\sigma} = \frac{31 - 28}{2} = 1.5$$

Step 2: Draw the figure



Note: This PowerPoint is only a summary and your main source should be the book.

Step 3: Find the area using Z-table

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
-0.5	0.3085	0.3050	0.3015	0.2981	0.2946	0.2912	0.2877	0.2843	0.2810	0.2776

$$\begin{aligned}P(-0.5 < Z < 1.5) &= P(Z < 1.5) - P(Z < -0.5) \\&= 0.9332 - 0.3085 = \underline{\underline{0.6247}}\end{aligned}$$

The probability that household generates between 27 and 31 pounds of newspapers per month is 62.47%


Solution (b) :

FIND $P(X > 30.2)$

Step 1: Find the z value

$$Z = \frac{X - \mu}{\sigma} = \frac{30.2 - 28}{2} = 1.1$$

Step 2: Draw the figure



Step 3: Find the area using Z-table

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830

$$P(Z > 1.1) = 1 - P(Z < 1.1) = 1 - 0.8643 = \underline{0.1375}$$

The probability that household generates more than 30.2 pounds of newspapers is 0.1375 or 13.75%

Example (3):

The American Automobile Association reports that the average time it takes to respond to an emergency call is 25 minutes. Assume the variable is approximately normally distributed and the standard deviation is 4.5 minutes. If 80 calls are randomly selected, approximately how many will be responded to in less than 15 minutes?


Solution:

FIND $P(X < 15)$

Step 1: Find the z value

$$Z = \frac{X - \mu}{\sigma} = \frac{15 - 25}{4.5} = -2.22$$

Step 2: Draw the figure



Step 3: Find the area using Z-table

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
-2.2	0.0139	0.0136	0.0132	0.0129	0.0125	0.0122	0.0119	0.0116	0.0113	0.0110

$$P(Z < -2.22) = 0.0132$$

Step 4: to find how many calls

$$0.0132 \times 80 = 1.056 \approx 1$$

Finding Data Values Given Specific Probabilities

$$Z = \frac{X - \mu}{\sigma} \longrightarrow Z \sigma = X - \mu$$

Formula for Finding X:

$$X = z \cdot \sigma + \mu$$

we must use that **three** steps.

First Step: draw the figure and Shading the area.

Second Step: find the z , using Z-table.


Third Step : find the X ,using the formula .

Example (4):

To qualify for a police academy, candidates must score in the top **10%** on a general abilities test. The test has a mean of 200 and standard deviation of 20. Find the lowest possible score to qualify. Assume the test scores is **normally distributed**.

Solution:

Step 1: Draw the figure



Step 2: Find the z value

$$P(Z > z) = 0.1$$

$$P(Z > z) = 1 - 0.1 = 0.9$$

$$\underline{Z=1.28}$$

Note: This PowerPoint is only a summary and your main source should be the book.

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830
1.2	0.8849	0.8860	0.8882	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974
2.8	0.9974	0.9975	0.9976	0.9977	0.9977	0.9978	0.9979	0.9979	0.9980	0.9981
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986
3.0	0.9987	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9989	0.9990	0.9990
3.1	0.9990	0.9991	0.9991	0.9991	0.9992	0.9992	0.9992	0.9992	0.9993	0.9993
3.2	0.9993	0.9993	0.9994	0.9994	0.9994	0.9994	0.9994	0.9995	0.9995	0.9995
3.3	0.9995	0.9995	0.9995	0.9996	0.9996	0.9996	0.9996	0.9996	0.9996	0.9997
3.4	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9998

Step 3: Find the value of x

$$X = (1.28)(20) + 200 = 226$$


Any body scoring 226 or higher will qualify

Example (5):

For a medical study, a researcher wishes to select people in the middle **60%** of the population based on blood pressure. If the mean systolic blood pressure is 120 and the standard deviation is 8, find the upper and lower readings that would qualify people to participate in the study. Assume that blood pressure readings is **normally distributed**.

Solution:

Step 1: Draw the figure



Step 2 : Find the two z values . $X = z \cdot \sigma + \mu$

$$P(Z > Z_1) = 0.2$$

$$P(Z < Z_1) = 1 - 0.2 = 0.8$$

$$Z_1 = 0.84$$

$$P(Z < Z_2) = 0.2$$

$$Z_2 = -0.84$$

Step 3 :Find the two values of x.

$$X_1 = Z_1 \cdot \sigma + \mu$$

$$X_1 = (0.84)(8) + 120 = 126.72$$

$$X_2 = Z_2 \cdot \sigma + \mu$$

$$X_2 = (-0.84)(8) + 120 = 113.28$$

*The middle 60% will have blood pressure reading of
113.28 < X < 126.72*

Example (6):

Given a normal distribution with a mean of 25, what is the standard deviation if 18% of the values are above 29?

Solution:

$$P(X > 29) = 0.18$$

$$P(X < 29) = 1 - 0.18 = 0.82$$

$$P\left(Z < \frac{x - \mu}{\sigma}\right) = 0.82$$

$$P\left(Z < \frac{29 - 25}{\sigma}\right) = 0.82$$

$$P\left(Z < \frac{4}{\sigma}\right) = 0.82$$

From the Z-Table

$$\frac{4}{\sigma} = 0.92$$

$$0.92 \sigma = 4$$

$$\underline{\sigma = 4.35}$$

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133
0.9	0.8159	0.8180	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974
2.8	0.9974	0.9975	0.9976	0.9977	0.9977	0.9978	0.9979	0.9979	0.9980	0.9981
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986
3.0	0.9987	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9989	0.9990	0.9990
3.1	0.9990	0.9991	0.9991	0.9991	0.9992	0.9992	0.9992	0.9992	0.9993	0.9993
3.2	0.9993	0.9993	0.9994	0.9994	0.9994	0.9994	0.9994	0.9995	0.9995	0.9995
3.3	0.9995	0.9995	0.9995	0.9996	0.9996	0.9996	0.9996	0.9996	0.9996	0.9997
3.4	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9998

Note: This PowerPoint is only a summary and your main source should be the book.

Example (7):

Given a normal distribution with a standard deviation of 10, what is the mean if 21% of the values are below 50?

Solution:

$$P(X < 50) = 0.21$$

$$P\left(Z < \frac{x - \mu}{\sigma}\right) = 0.21$$

$$P\left(Z < \frac{50 - \mu}{10}\right) = 0.21$$

From the Z-Table

$$\frac{50 - \mu}{10} = -0.81$$

$$50 - \mu = -8.1$$

$\mu = 58.1$

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
-3.4	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0002
-3.3	0.0005	0.0005	0.0005	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0003
-3.2	0.0007	0.0007	0.0006	0.0006	0.0006	0.0006	0.0006	0.0005	0.0005	0.0005
-3.1	0.0010	0.0009	0.0009	0.0009	0.0008	0.0008	0.0008	0.0008	0.0007	0.0007
-3.0	0.0013	0.0013	0.0013	0.0012	0.0012	0.0011	0.0011	0.0011	0.0010	0.0010
-2.9	0.0019	0.0018	0.0018	0.0017	0.0016	0.0016	0.0015	0.0015	0.0014	0.0014
-2.8	0.0026	0.0025	0.0024	0.0023	0.0023	0.0022	0.0021	0.0021	0.0020	0.0019
-2.7	0.0035	0.0034	0.0033	0.0032	0.0031	0.0030	0.0029	0.0028	0.0027	0.0026
-2.6	0.0047	0.0045	0.0044	0.0043	0.0041	0.0040	0.0039	0.0038	0.0037	0.0036
-2.5	0.0062	0.0060	0.0059	0.0057	0.0055	0.0054	0.0052	0.0051	0.0049	0.0048
-2.4	0.0082	0.0080	0.0078	0.0075	0.0073	0.0071	0.0069	0.0068	0.0066	0.0064
-2.3	0.0107	0.0104	0.0102	0.0099	0.0096	0.0094	0.0091	0.0089	0.0087	0.0084
-2.2	0.0139	0.0136	0.0132	0.0129	0.0125	0.0122	0.0119	0.0116	0.0113	0.0110
-2.1	0.0179	0.0174	0.0170	0.0166	0.0162	0.0158	0.0154	0.0150	0.0146	0.0143
-2.0	0.0228	0.0222	0.0217	0.0212	0.0207	0.0202	0.0197	0.0192	0.0188	0.0183
-1.9	0.0287	0.0281	0.0274	0.0268	0.0262	0.0256	0.0250	0.0244	0.0239	0.0233
-1.8	0.0359	0.0351	0.0344	0.0336	0.0329	0.0322	0.0314	0.0307	0.0301	0.0294
-1.7	0.0446	0.0436	0.0427	0.0418	0.0409	0.0401	0.0392	0.0384	0.0375	0.0367
-1.6	0.0548	0.0537	0.0526	0.0516	0.0505	0.0495	0.0485	0.0475	0.0465	0.0455
-1.5	0.0668	0.0655	0.0643	0.0630	0.0618	0.0606	0.0594	0.0582	0.0571	0.0559
-1.4	0.0808	0.0793	0.0778	0.0764	0.0749	0.0735	0.0721	0.0708	0.0694	0.0681
-1.3	0.0968	0.0951	0.0934	0.0918	0.0901	0.0885	0.0869	0.0853	0.0838	0.0823
-1.2	0.1151	0.1131	0.1112	0.1093	0.1075	0.1056	0.1038	0.1020	0.1003	0.0985
-1.1	0.1357	0.1335	0.1314	0.1292	0.1271	0.1251	0.1230	0.1210	0.1190	0.1170
-1.0	0.1587	0.1562	0.1539	0.1515	0.1492	0.1469	0.1446	0.1423	0.1401	0.1379
-0.9	0.1841	0.1814	0.1788	0.1762	0.1736	0.1711	0.1685	0.1660	0.1635	0.1611
-0.8	0.2119	0.2090	0.2061	0.2033	0.2005	0.1977	0.1949	0.1922	0.1894	0.1867
-0.7	0.2420	0.2389	0.2358	0.2327	0.2296	0.2266	0.2236	0.2206	0.2177	0.2148
-0.6	0.2743	0.2709	0.2676	0.2643	0.2611	0.2578	0.2546	0.2514	0.2483	0.2451
-0.5	0.3085	0.3050	0.3015	0.2981	0.2946	0.2912	0.2877	0.2843	0.2810	0.2776
-0.4	0.3446	0.3409	0.3372	0.3336	0.3300	0.3264	0.3228	0.3192	0.3156	0.3121
-0.3	0.3821	0.3783	0.3745	0.3707	0.3669	0.3632	0.3594	0.3557	0.3520	0.3483
-0.2	0.4207	0.4168	0.4129	0.4090	0.4052	0.4013	0.3974	0.3936	0.3897	0.3859
-0.1	0.4602	0.4562	0.4522	0.4483	0.4443	0.4404	0.4364	0.4325	0.4286	0.4247
-0.0	0.5000	0.4960	0.4920	0.4880	0.4840	0.4801	0.4761	0.4721	0.4681	0.4641

Note: This PowerPoint is only a summary and your main source should be the book.

Example (8):

Given a normal distribution with 80% of the values are above 125 and 90% of the values are above 110, what are the mean and standard deviation of this distribution?

Solution:

$$P(X > 125) = 0.8$$

$$P(X < 125) = 1 - 0.8 = 0.2$$

$$P\left(Z < \frac{x - \mu}{\sigma}\right) = 0.2$$

$$P\left(Z < \frac{125 - \mu}{\sigma}\right) = 0.2$$

From the Z- Table

$$\frac{125 - \mu}{\sigma} = -0.84 \rightarrow (1)$$

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
-3.4	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0002
-3.3	0.0005	0.0005	0.0005	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0003
-3.2	0.0007	0.0007	0.0006	0.0006	0.0006	0.0006	0.0006	0.0005	0.0005	0.0005
-3.1	0.0010	0.0009	0.0009	0.0009	0.0008	0.0008	0.0008	0.0008	0.0007	0.0007
-3.0	0.0013	0.0013	0.0013	0.0012	0.0012	0.0011	0.0011	0.0011	0.0010	0.0010
-2.9	0.0019	0.0018	0.0018	0.0017	0.0016	0.0016	0.0015	0.0015	0.0014	0.0014
-2.8	0.0026	0.0025	0.0024	0.0023	0.0023	0.0022	0.0021	0.0021	0.0020	0.0019
-2.7	0.0035	0.0034	0.0033	0.0032	0.0031	0.0030	0.0029	0.0028	0.0027	0.0026
-2.6	0.0047	0.0045	0.0044	0.0043	0.0041	0.0040	0.0039	0.0038	0.0037	0.0036
-2.5	0.0062	0.0060	0.0059	0.0057	0.0055	0.0054	0.0052	0.0051	0.0049	0.0048
-2.4	0.0082	0.0080	0.0078	0.0075	0.0073	0.0071	0.0069	0.0068	0.0066	0.0064
-2.3	0.0107	0.0104	0.0102	0.0099	0.0096	0.0094	0.0091	0.0089	0.0087	0.0084
-2.2	0.0139	0.0136	0.0132	0.0129	0.0125	0.0122	0.0119	0.0116	0.0113	0.0110
-2.1	0.0179	0.0174	0.0170	0.0166	0.0162	0.0158	0.0154	0.0150	0.0146	0.0143
-2.0	0.0228	0.0222	0.0217	0.0212	0.0207	0.0202	0.0197	0.0192	0.0188	0.0183
-1.9	0.0287	0.0281	0.0274	0.0268	0.0262	0.0256	0.0250	0.0244	0.0239	0.0233
-1.8	0.0359	0.0351	0.0344	0.0336	0.0329	0.0322	0.0314	0.0307	0.0301	0.0294
-1.7	0.0446	0.0436	0.0427	0.0418	0.0409	0.0401	0.0392	0.0384	0.0375	0.0367
-1.6	0.0548	0.0537	0.0526	0.0516	0.0505	0.0495	0.0485	0.0475	0.0465	0.0455
-1.5	0.0668	0.0655	0.0643	0.0630	0.0618	0.0606	0.0594	0.0582	0.0571	0.0559
-1.4	0.0808	0.0793	0.0778	0.0764	0.0749	0.0735	0.0721	0.0708	0.0694	0.0681
-1.3	0.0968	0.0951	0.0934	0.0918	0.0901	0.0885	0.0869	0.0853	0.0838	0.0823
-1.2	0.1151	0.1131	0.1112	0.1093	0.1075	0.1056	0.1038	0.1020	0.1003	0.0985
-1.1	0.1357	0.1335	0.1314	0.1292	0.1271	0.1251	0.1230	0.1210	0.1190	0.1170
-1.0	0.1587	0.1562	0.1539	0.1515	0.1492	0.1469	0.1446	0.1423	0.1401	0.1379
-0.9	0.1841	0.1814	0.1788	0.1762	0.1736	0.1711	0.1685	0.1660	0.1635	0.1611
-0.8	0.2113	0.2090	0.2061	0.2030	0.2005	0.1977	0.1949	0.1922	0.1894	0.1867
-0.7	0.2420	0.2389	0.2358	0.2327	0.2296	0.2266	0.2236	0.2206	0.2177	0.2148
-0.6	0.2743	0.2709	0.2676	0.2643	0.2611	0.2578	0.2546	0.2514	0.2483	0.2451
-0.5	0.3085	0.3050	0.3015	0.2981	0.2946	0.2912	0.2877	0.2843	0.2810	0.2776
-0.4	0.3446	0.3409	0.3372	0.3336	0.3300	0.3264	0.3228	0.3192	0.3156	0.3121
-0.3	0.3821	0.3783	0.3745	0.3707	0.3669	0.3632	0.3594	0.3557	0.3520	0.3483
-0.2	0.4207	0.4168	0.4129	0.4090	0.4052	0.4013	0.3974	0.3936	0.3897	0.3859
-0.1	0.4602	0.4562	0.4522	0.4483	0.4443	0.4404	0.4364	0.4325	0.4286	0.4247
-0.0	0.5000	0.4960	0.4920	0.4880	0.4840	0.4801	0.4761	0.4721	0.4681	0.4641

Note: This PowerPoint is only a summary and your main source should be the book.

$$P(X > 110) = 0.9$$

$$P(X < 110) = 1 - 0.9 = 0.1$$

$$P\left(Z < \frac{x - \mu}{\sigma}\right) = 0.1$$

$$P\left(Z < \frac{110 - \mu}{\sigma}\right) = 0.1$$

From the Z- Table

$$\frac{110 - \mu}{\sigma} = -1.28 \rightarrow (2)$$

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
-3.4	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0002
-3.3	0.0005	0.0005	0.0005	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0003
-3.2	0.0007	0.0007	0.0006	0.0006	0.0006	0.0006	0.0006	0.0005	0.0005	0.0005
-3.1	0.0010	0.0009	0.0009	0.0009	0.0008	0.0008	0.0008	0.0008	0.0007	0.0007
-3.0	0.0013	0.0013	0.0013	0.0012	0.0012	0.0011	0.0011	0.0011	0.0010	0.0010
-2.9	0.0019	0.0018	0.0018	0.0017	0.0016	0.0016	0.0015	0.0015	0.0014	0.0014
-2.8	0.0026	0.0025	0.0024	0.0023	0.0023	0.0022	0.0021	0.0021	0.0020	0.0019
-2.7	0.0035	0.0034	0.0033	0.0032	0.0031	0.0030	0.0029	0.0028	0.0027	0.0026
-2.6	0.0047	0.0045	0.0044	0.0043	0.0041	0.0040	0.0039	0.0038	0.0037	0.0036
-2.5	0.0062	0.0060	0.0059	0.0057	0.0055	0.0054	0.0052	0.0051	0.0049	0.0048
-2.4	0.0082	0.0080	0.0078	0.0075	0.0073	0.0071	0.0069	0.0068	0.0066	0.0064
-2.3	0.0107	0.0104	0.0102	0.0099	0.0096	0.0094	0.0091	0.0089	0.0087	0.0084
-2.2	0.0139	0.0136	0.0132	0.0129	0.0125	0.0122	0.0119	0.0116	0.0113	0.0110
-2.1	0.0179	0.0174	0.0170	0.0166	0.0162	0.0158	0.0154	0.0150	0.0146	0.0143
-2.0	0.0228	0.0222	0.0217	0.0212	0.0207	0.0202	0.0197	0.0192	0.0188	0.0183
-1.9	0.0287	0.0281	0.0274	0.0268	0.0262	0.0256	0.0250	0.0244	0.0239	0.0233
-1.8	0.0359	0.0351	0.0344	0.0336	0.0329	0.0322	0.0314	0.0307	0.0301	0.0294
-1.7	0.0446	0.0436	0.0427	0.0418	0.0409	0.0401	0.0392	0.0384	0.0375	0.0367
-1.6	0.0548	0.0537	0.0526	0.0516	0.0505	0.0495	0.0485	0.0475	0.0465	0.0455
-1.5	0.0668	0.0655	0.0643	0.0630	0.0618	0.0606	0.0594	0.0582	0.0571	0.0559
-1.4	0.0808	0.0793	0.0778	0.0764	0.0749	0.0735	0.0721	0.0708	0.0694	0.0681
-1.3	0.0968	0.0951	0.0934	0.0918	0.0901	0.0885	0.0869	0.0853	0.0838	0.0823
-1.2	0.1151	0.1151	0.1112	0.1093	0.1075	0.1056	0.1036	0.1020	0.1003	0.0985
-1.1	0.1357	0.1335	0.1314	0.1292	0.1271	0.1251	0.1230	0.1210	0.1190	0.1170
-1.0	0.1587	0.1562	0.1539	0.1515	0.1492	0.1469	0.1446	0.1423	0.1401	0.1379
-0.9	0.1841	0.1814	0.1788	0.1762	0.1736	0.1711	0.1685	0.1660	0.1635	0.1611
-0.8	0.2119	0.2090	0.2061	0.2033	0.2005	0.1977	0.1949	0.1922	0.1894	0.1867
-0.7	0.2420	0.2389	0.2358	0.2327	0.2296	0.2266	0.2236	0.2206	0.2177	0.2148
-0.6	0.2743	0.2709	0.2676	0.2643	0.2611	0.2578	0.2546	0.2514	0.2483	0.2451
-0.5	0.3085	0.3050	0.3015	0.2981	0.2946	0.2912	0.2877	0.2843	0.2810	0.2776
-0.4	0.3446	0.3409	0.3372	0.3336	0.3300	0.3264	0.3228	0.3192	0.3156	0.3121
-0.3	0.3821	0.3783	0.3745	0.3707	0.3669	0.3632	0.3594	0.3557	0.3520	0.3483
-0.2	0.4207	0.4168	0.4129	0.4090	0.4052	0.4013	0.3974	0.3936	0.3897	0.3859
-0.1	0.4602	0.4562	0.4522	0.4483	0.4443	0.4404	0.4364	0.4325	0.4286	0.4247
-0.0	0.5000	0.4960	0.4920	0.4880	0.4840	0.4801	0.4761	0.4721	0.4681	0.4641

Note: This PowerPoint is only a summary and your main source should be the book.

From Equations (1) and (2)

$$\frac{125 - \mu}{\sigma} = -0.84$$

$$125 - \mu = -0.84\sigma \rightarrow (1)$$

$$\frac{110 - \mu}{\sigma} = -1.28$$

$$110 - \mu = -1.28\sigma \rightarrow (2)$$

Subtract (2) from (1)

$$-15 = -0.442\sigma$$

$$\underline{\sigma = 33.9}$$

Substitute in (2)







$$110 - \mu = -1.28(33.9)$$

$$\underline{\mu = 153.5}$$

Questions ???



1-If X is normally distributed random variable with $\mu = 5$,
 $\sigma = 4$, find the $P(x > -1.4)$??



Discrete Probability Distributions

Note: This PowerPoint is only a summary and your main source should be the book.

The Binomial Distribution

Mean, Variance and Standard deviation for The Binomial Distribution

- Many types of probability problems have only two possible outcomes or they can be reduced to two outcomes.

- Examples include: when a coin is tossed it can land on heads or tails, when a baby is born it is either a boy or girl, etc.

The **binomial experiment** is a probability experiment that satisfies these requirements:

1. Each trial can have only **two** possible outcomes—success or failure.
2. There must be a **fixed number** of trials.
3. The outcomes of each trial must be **independent** of each other.
4. The probability of success must remain the **same** for each trial.

Notation for the Binomial Distribution

$P(S)$:The symbol for the probability of success

$P(F)$:The symbol for the probability of failure

p :The numerical probability of success

q :The numerical probability of failure

$$P(S) = p \text{ and } P(F) = 1 - p = q$$

n :The number of trials

X :The number of successes

Note that $X = 0, 1, 2, 3, \dots, n$

In a binomial experiment, the probability of exactly X successes in n trials is

$$P(X) = {}_n C_x \cdot \underbrace{p^X \cdot q^{n-X}}_{\text{probability of a desired outcome}}$$

number of possible desired outcomes


$$P(X) = \frac{n!}{(n - X)! X!} \cdot p^X \cdot q^{n-X}$$

Example (1): Tossing Coins

A coin is tossed 3 times. Find the probability of getting exactly 2 heads.

Solution :

$$p(2 \text{ heads}) = \frac{n!}{(n - x)! x!} \cdot p^x \cdot q^{n-x}$$



$$p(2 \text{ heads}) = \frac{3!}{(3 - 2)! 2!} \cdot \left(\frac{1}{2}\right)^2 \cdot \left(\frac{1}{2}\right)^2 = \frac{3}{8} = 0.375$$

Example (2): Survey on Doctor Visits

A survey found that one out of five Americans say he or she has visited a doctor in any given month. If 10 people are selected at random, find the probability that exactly 3 will have visited a doctor last month.

Solution :

$$P(X) = \frac{n!}{(n-X)!X!} \cdot p^X \cdot q^{n-X}$$

$$P(3) = \frac{10!}{7!3!} \cdot \left(\frac{1}{5}\right)^3 \cdot \left(\frac{4}{5}\right)^7 = \boxed{0.201}$$

$$\begin{aligned} n &= 10 \\ x &= 3 \\ p &= \frac{1}{5} \\ q &= 1 - \frac{1}{5} = \frac{4}{5} \end{aligned}$$

Example (3): Survey on Employment

A survey from Teenage Research Unlimited (Northbrook, Illinois) found that 30% of teenage consumers receive their spending money from part-time jobs. If 5 teenagers are selected at random, find the probability that at least 3 of them will have part-time jobs.


Solution :

$$P(3) = \frac{5!}{2!3!} \cdot (0.30)^3 \cdot (0.70)^2 = 0.132$$

$$P(4) = \frac{5!}{1!4!} \cdot (0.30)^4 \cdot (0.70)^1 = 0.028$$

$$P(5) = \frac{5!}{0!5!} \cdot (0.30)^5 \cdot (0.70)^0 = 0.002$$

$$\begin{aligned} P(X \geq 3) &= 0.132 \\ &\quad + 0.028 \\ &\quad + 0.002 \\ &= \boxed{0.162} \end{aligned}$$



Mean, Variance and Standard deviation for the binomial

The mean , variance and SD of a variable that the binomial distribution can be found by using the following formulas:

$$\text{Mean: } \mu = np$$

$$\text{Variance: } \sigma^2 = npq$$

$$\text{Standard Deviation: } \sigma = \sqrt{npq}$$

Example (4): Tossing A Coin

A coin is tossed 4 times. Find the mean, variance and standard deviation of number of heads that will be obtained.

Solution :

$$\mu = n \cdot p = 4 \cdot \frac{1}{2} = 2$$

$$n = 4$$
$$p = \frac{1}{2}$$
$$q = 1 - \frac{1}{2} = \frac{1}{2}$$

$$\sigma^2 = n \cdot p \cdot q = 4 \cdot \frac{1}{2} \cdot \frac{1}{2} = 1$$

$$\sigma = \sqrt{\sigma^2} = \sqrt{n \cdot p \cdot q} = \sqrt{1} = 1$$

Example (5): Rolling a die

A die is rolled 360 times , find the mean , variance and standard deviation of the number of 4s that will be rolled .

Solution :

$$\mu = n \cdot p = 360 \cdot \frac{1}{6} = 60$$

$$n = 360$$
$$p = \frac{1}{6}$$
$$q = 1 - \frac{1}{6} = \frac{5}{6}$$

$$\sigma^2 = n \cdot p \cdot q = 360 \cdot \frac{1}{6} \cdot \frac{1}{6} = 50$$

$$\sigma = \sqrt{\sigma^2} = \sqrt{n \cdot p \cdot q} = \sqrt{50} = 7.07$$

Example (6):

The probability that a driver must stop at any one traffic light coming to Lincoln University is 0.2. There are 15 sets of traffic lights on the journey.

- a) What is the probability that a student must stop at exactly 2 of the 15 sets of traffic lights?
- b) What is the probability that a student will be stopped at 1 or more of the 15 sets of traffic lights?

Solution (6.a) :

$$P(X) = \frac{n!}{(n-X)!X!} \cdot p^X \cdot q^{n-X}$$

n = 15
x = 2
p = 0.2
q = 1 - 0.2 = 0.8

$$P(x = 2) = \frac{15!}{(15-2)!2!} (0.2)^2 (0.8)^{(15-2)}$$

$$P(x = 2) = \frac{15!}{(13!)(2!)} (0.2)^2 (0.8)^{13} = 0.2309$$

$$= 23.09 \%$$

Solution (6.b) :

$$P(X) = \frac{n!}{(n-X)!X!} \cdot p^X \cdot q^{n-X}$$

$$P(X \geq 1) = 1 - P(X < 1) = 1 - P(X = 0)$$

$$P(x = 0) = \frac{15!}{(15-0)!0!} (0.2)^0 (0.8)^{(15-0)}$$

$$P(x = 0) = \frac{15!}{(15!)(0!)} (0.2)^0 (0.8)^{15} = 0.0352$$

$$P(X \geq 1) = 1 - P(X = 0) = 1 - 0.0352 = 0.9648 = \mathbf{96.48\%}$$

n = 15
x = 1, 2, ..., 15
p = 0.2
q = 1 - 0.2 = 0.8

A coin is tossed 72 times. The standard deviation for the number of heads that will be tossed is

- A) 18
- B) 4.24
- C) 6
- D) 36

A student takes a 6 question multiple choice quiz with 4 choices for each question. If the student guesses at random on each question, what is the probability that the student gets exactly 3 questions correct?

- A) 0.088
- B) 0.0512
- C) 0.132
- D) 0.022

