

Nonparametric Bayes

Omar Gutiérrez

@omargz

June 15, 2015

Mostly based on **A Tutorial on Bayesian Nonparametric Models** by Samuel J. Gershman.

Outline

Introduction

Example: clustering

Bayesian nonparametric models

Chinese Restaurant Process

Introduction

- ▶ Before to start: Statistics was already there, even before than Machine Learning (ML)
- ▶ What we do in ML is fitting a model to the data
- ▶ That is, we adjust the values of certain parameters

Linear Regression




Figure 1: Linear Regression

Neural Networks




Figure 2: Perceptron

Hidden Markov Models




Figure 3: Hidden Markov Models

Bayesian Learning

$$P(h|D) = \frac{P(D|h)P(h)}{P(D)} \quad (1)$$

Maximum Likelihood Estimation

$$\begin{aligned}h_{MAP} &\equiv \arg \max_{h \in H} P(h|D) \\&= \arg \max_{h \in H} \frac{P(D|h)P(h)}{P(D)} \\&= \arg \max_{h \in H} P(D|h)P(h) \\h_{MLE} &= \arg \max_{h \in H} P(D|h)\end{aligned}\tag{2}$$

Data is a mess

- ▶ The articles in Wikipedia
- ▶ The species in the planet
- ▶ The hashtags on Twitter

How the problem is *sometimes* addressed

- ▶ Let's start with the classic approach
- ▶ Let's do clustering
- ▶ Let's use Gaussian Mixture Models (GMM)
- ▶ We can fit several models and then compare them with some metric.

How the problem is *sometimes* addressed



Figure 4: Points classified with GMM

How we can *alternatively* approach the problem?




Figure 5: Points classified with Infinite GMM

How do we approach the problem?

- ▶ Another interesting approach is to use Bayesian Nonparametric (BNP) models
- ▶ BNP models will build a model that can adapt its complexity to the data

Bayesian nonparametric models



Recap: Bayesian parametric vs nonparametric models

- ▶ Traditional approach (finite)
 - ▶ The number of parameters θ (e.g. clusters) is prespecified
 - ▶ We have a prior distribution over parameters $P(\theta)$
 - ▶ For example, in the Gaussian mixture model, each cluster will be modelled using a parametric model (e. g. Gaussian)
- ▶ Bayesian nonparametric models
 - ▶ We assume that there is an **infinite** number of latent clusters
 - ▶ A finite number of clusters is *inferred* from data
 - ▶ The number of clusters grow as new data points are observed

Chinese Restaurant Process



- ▶ Imagine a restaurant with an infinite number of tables,
- ▶ and a sequence of customers entering the restaurant and sitting down.
- ▶ The first customer enters and sits at the first table
- ▶ The second customer enters and sits...
 - ▶ at the first table with probability $\frac{1}{1+\alpha}$
 - ▶ at the second table with probability $\frac{\alpha}{1+\alpha}$
- ▶ We realize that CRP is a form of clustering: K groups and each group with size N_k

What else?

- ▶ Dirichlet distribution is a generalization of β distribution.
- ▶ Dirichlet process

Thank you
Questions?