

VECTORS	
Term	Definition
Norm/Length	$\ v\ = \sqrt{\sum_{i=1}^n v_i^2}$
	$\ v\ = \sqrt{v^T v}$
	$\nabla \ v\ = \frac{1}{\ v\ } v^T$
Distance	$d(u, v) = \ u - v\ $
	TODO: $\nabla d(u, v) = \frac{1}{d(u, v)} ((u - v)(u - v))^{??}$ $\nabla_u d(u, v) = \frac{u - v}{\ u - v\ } = \frac{u - v}{d(u, v)}$ $\nabla_v d(u, v) = -\frac{u - v}{\ u - v\ } = -\frac{u - v}{d(u, v)}$
Scalar product	$u \bullet v = u^T v = \sum_k u_k v_k$

SCALAR PRODUCT

MATRICES

Determinant:	
$\det(a) = a$	$\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ad - cb$
$\det \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} = a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32}$	$-a_{31}a_{22}a_{13} - a_{32}a_{23}a_{11} - a_{33}a_{21}a_{12}$

Inverting:
$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \Rightarrow A^{-1} = \frac{1}{ad - cb} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$

Transposing:
$\begin{pmatrix} 1 & 0 \\ 0 & 2 \\ 3 & 1 \end{pmatrix}^T = \begin{pmatrix} 1 & 0 & 3 \\ 0 & 2 & 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 4 \\ 5 \end{pmatrix}^T = (1 \ 4 \ 5)$

Matrix multiplication:
$\begin{pmatrix} 2 & 3 & 1 \\ 4 & -1 & 7 \end{pmatrix} \cdot \begin{pmatrix} 6 & 4 \\ 1 & 0 \\ 8 & 9 \end{pmatrix}$

$$= \begin{pmatrix} (2 \cdot 6 + 3 \cdot 1 + 1 \cdot 8) & (2 \cdot 4 + 3 \cdot 0 + 1 \cdot 9) \\ (4 \cdot 6 + -1 \cdot 1 + 7 \cdot 8) & (4 \cdot 4 + -1 \cdot 0 + 7 \cdot 9) \end{pmatrix} = \begin{pmatrix} 23 & 17 \\ 79 & 79 \end{pmatrix}$$

INDICES

$A \in \mathbb{R}^{n \times m}, B \in \mathbb{R}^{m \times r}, A \cdot B \in \mathbb{R}^{n \times r}$
$(A \cdot B)_{ij} = \sum_{k=1}^m A_{ik} B_{kj} = \sum_k A_{ik} B_{kj}$
$((A \cdot B)^T)_{ij} = (A \cdot B)_{ji} = \sum_k A_{jk} B_{ik}$
$a^T \cdot B \cdot a = \sum_{ij} (a^T)_i B_{ij} a_j = \sum_{ij} a_i B_{ij} a_j$
$(B \cdot a)(C \cdot a) = \sum_i (B \cdot a)_i (C \cdot a)_i$

KRONECKER DELTA

$\delta_{ij} = (\vec{e}_i)_j = \begin{cases} 1, & i = j \\ 0, & \text{otherwise} \end{cases}$
$(E_{rst})_{ijk} = \delta_{ri} \delta_{sj} \delta_{tk}$

BASIS VECTORS

$$\mathbb{R}^3$$
$$e_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, e_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, e_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

UNIT MATRICES

$$\mathbb{R}^2$$
$$E_{11} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, E_{12} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, E_{21} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, E_{22} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

AFFINE TRANSFORMATIONS

$$A_{a,M}(x) = a + M \cdot x$$

$L_A(\vec{0}) = \vec{0}$	
DERIVATIVES	
$1 \rightarrow 0$	$x \rightarrow 1$
$x^2 \rightarrow 2x$	$x^n \rightarrow nx^{n-1}$
$\frac{1}{x} \rightarrow -\frac{1}{x^2}$	$\sqrt{x} \rightarrow \frac{1}{2\sqrt{x}}$
$e^x \rightarrow e^x$	$e^{-x} \rightarrow -e^{-x}$
$\ln(x) \rightarrow \frac{1}{x} \text{ for } x > 0$	$\ln(y \cdot x) \rightarrow \frac{1}{x} \text{ for } x > 0$
$a^x \rightarrow \ln(a) \cdot a^x \text{ for } a > 0, a \neq 1$	$\log_a(x) \rightarrow \frac{1}{\ln(b) \cdot x}$
$\sin(x) \rightarrow \cos(x)$	$\sin(2x) \rightarrow 2 \cos(2x)$
$\cos(x) \rightarrow -\sin(x)$	$\cos(ax) \rightarrow -a \sin(ax)$
$\tan(x) \rightarrow 1 + \tan^2(x) = \frac{1}{\cos^2(x)}$	$\operatorname{arccot}(x) \rightarrow -\frac{1}{1+x^2}$
$\cot(x) \rightarrow -\frac{1}{\sin^2(x)}$	$\arcsin(x) \rightarrow \frac{1}{\sqrt{1-x^2}}$
$\operatorname{arccos}(x) \rightarrow -\frac{1}{\sqrt{1-x^2}}$	$\arctan(x) \rightarrow \frac{1}{1+x^2}$


Term	Definition
Addition	$(f(x) + g(x))' = f'(x) + g'(x)$
Multiplication	$(\alpha \cdot f(x))' = \alpha \cdot f'(x)$
Product rule	$(f \cdot g)' = f' \cdot g + f \cdot g'$ $(f \cdot g \cdot h)' = f' \cdot g \cdot h + f \cdot g' \cdot h + f \cdot g \cdot h'$
Chain rule	$(f(g(x)))' = f'(g(x)) \cdot g'(x)$
Quotient rule	$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$

FUNCTIONS OF SEVERAL VARIABLES

Term	Definition
Real valued function	$R \subset \mathbb{R}$
Vector valued function	$R \subset \mathbb{R}^m$
A function of n variables	$D \subset \mathbb{R}^n$
Softmax function	TODO:
RSS	$\text{RSS} = \sum_i (y_i - f(x_i))^2$
n -dimensional curve	$c: \mathbb{R} \rightarrow \mathbb{R}^n$
n -dimensional hyper-surface	$s: \mathbb{R}^n \rightarrow \mathbb{R}$
Reparametrization	$c(t) \rightarrow d(s) = c(h(s))$

Sigmoid function:

$$\sigma(x) : \mathbb{R} \rightarrow [0; 1] = \frac{1}{1 + e^{-x}}$$



DERIVATIVES

Function	Derivative
$f: \mathbb{R} \rightarrow \mathbb{R}^n$ $\begin{matrix} x \mapsto y \end{matrix}$	$f': \mathbb{R} \rightarrow \mathbb{R}^n = \begin{pmatrix} \frac{d}{dx} f_1(x) \\ \frac{d}{dx} f_2(x) \\ \vdots \\ \frac{d}{dx} f_n(x) \end{pmatrix}$
$f: \mathbb{R}^n \rightarrow \mathbb{R}$ $\begin{matrix} x \mapsto y \end{matrix}$	$f': \mathbb{R}^n \rightarrow \mathbb{R}^{1 \times n} = \nabla f$ $= \left(\frac{\partial}{\partial x_1} f(x), \frac{\partial}{\partial x_2} f(x), \dots, \frac{\partial}{\partial x_n} f(x) \right)$
$f: \mathbb{R}^n \rightarrow \mathbb{R}^m$ $\begin{matrix} x \mapsto y \end{matrix}$	$f': \mathbb{R}^n \rightarrow \mathbb{R}^{m \times n} = J_f = \frac{\partial f}{\partial x} \Big _{x=x_0}$ $= \begin{pmatrix} \nabla f_1(x) \\ \vdots \\ \nabla f_m(x) \end{pmatrix} = \begin{pmatrix} \frac{\partial}{\partial x_1} f_1(x) & \dots & \frac{\partial}{\partial x_n} f_1(x) \\ \vdots & \ddots & \vdots \\ \frac{\partial}{\partial x_1} f_m(x) & \dots & \frac{\partial}{\partial x_n} f_m(x) \end{pmatrix}$

PARTIAL VS TOTAL DERIVATIVE

TODO: example FS2024
$f(x, y) = x^2 + 2y$ $\frac{\partial f}{\partial x} = 2x$

$\frac{d}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dx} = 2x + 2 \cdot \frac{dy}{dx} \mid (y \text{ dependent on } x)$
GRADIENTS
Common gradients:
$f(x) = \ x\ ^2 \Rightarrow \nabla f = 2x$
$f(x) = a^T x + c \Rightarrow \nabla f = a$
$f(x) = x^T A x \Rightarrow \nabla f = (A + A^T)x$
$f(x) = \frac{1}{2} x^T A x + b^T x + c \Rightarrow \nabla f = \frac{A + A^T}{2} x + b$ $= Ax + b \text{ if } A^T = A$
$f(x) = g(Ax + b) \Rightarrow \nabla f = A^T \nabla g(Ax + b)$ $g: \mathbb{R}^m \rightarrow \mathbb{R}$
$f(x) = u(x) \cdot v(x) \Rightarrow \nabla f = (J_u)^T v + (J_v)^T u$
$f(x) = \ u(x)\ = \sqrt{u^T u} \Rightarrow \nabla f = (J_u)^T \frac{u}{\ u\ } \text{ if } \ u\ \neq 0$
$f(x) = \log p(x) \Rightarrow \nabla f = \frac{1}{p(x)} \nabla p(x)$

JACOBIAN MATRIX

$$f: \mathbb{R}^2 \rightarrow \mathbb{R}^2, J_f(x, y) = \begin{pmatrix} \frac{\partial x_1}{\partial x_2} & \frac{\partial y_1}{\partial y_2} \end{pmatrix}$$

Linearisation:

$$f: \mathbb{R}^n \rightarrow \mathbb{R}^m, x_0 \in \mathbb{R}^n$$
$$L_f(x) = f(x_0) + J_f(x_0) \cdot (x - x_0)$$

Chain rule:

$$f: \mathbb{R}^n \rightarrow \mathbb{R}^m, g: \mathbb{R}^m \rightarrow \mathbb{R}^k, h: \mathbb{R}^n \rightarrow \mathbb{R}^k = g \circ f$$
$$J_h = J_g(f(x)) \cdot J_f(x)$$

Common jacobians:

$$f(x) = a + M \cdot x \Rightarrow J_f = M$$
$$f(x) = x \Rightarrow J_f = \mathbb{1}$$
$$f(x) = \|x\|^2 \Rightarrow J_f = (2x)^T$$

TODO: J_x for softmax $s_i = \frac{e^{x_i}}{\sum_k e^{x_k}}; J_{ij} = s_i(\delta_{ij} - s_j)$
--

HYPER-SURFACES

Chain rule:
$f: \mathbb{R}^n \rightarrow \mathbb{R}^m, g: \mathbb{R}^m \rightarrow \mathbb{R}, h: \mathbb{R}^n \rightarrow \mathbb{R} = g \circ f$ $\nabla h(x) = \nabla g(f(x)) = \nabla g(f) \Big _{f=f(x)} \cdot \frac{\partial}{\partial x} f(x)$

Differentiation properties:

$$f: \mathbb{R}^n \rightarrow \mathbb{R}, g: \mathbb{R}^n \rightarrow \mathbb{R}$$
$$\nabla(f + g) = \nabla f + \nabla g$$
$$\nabla(f - g) = \nabla f - \nabla g$$
$$\nabla(f \cdot g) = g \nabla f + f \nabla g$$
$$\nabla \frac{f}{g} = \frac{g \nabla f - f \nabla g}{g^2}$$

COMMUTATIVE DIAGRAMS



GEOMETRY OF HYPER-SURFACES

$$f: \mathbb{R}^2 \rightarrow \mathbb{R}$$

Tangent space: $\alpha, \beta \in \mathbb{R}, T =$

$$\left\{ \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} + \alpha \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \beta \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right\}$$
$$\left\{ \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} + \alpha \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \beta \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right\}$$

Graph of a linearisation:

$$l(x) = f(x_0) + \nabla f(x_0) \cdot (x - x_0) = T_{p_0}$$

SECOND DERIVATIVE

$$\frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} (f(x, y)) \right) = \frac{\partial^2 f(x, y)}{\partial x^2} = \partial_{xx} f(x, y)$$

HESSIAN MATRIX

$$f: \mathbb{R}^n \rightarrow \mathbb{R}$$
$$(H(x))_{ij} = \frac{\partial}{\partial x_i} \frac{\partial}{\partial x_j} f(x)$$
$$\partial_i \partial_j f = \partial_j \partial_i f$$

Common hessians:

$$f(x) = \frac{1}{2} x^T A x + b^T x + c \Rightarrow H_f = \frac{A + A^T}{2}$$
$$f(x) = g(Ax + b) \Rightarrow H_f = A^T H_g(Ax + b) A$$
$$g: \mathbb{R}^m \rightarrow \mathbb{R}$$

EXTREMAL POINTS

Let $z = f(c(t))$, c a curve passing the stationary point $c_0 = (x_0, y_0)$ at $t = t_0$ and

$$\frac{d^2 z}{dt^2} = \underbrace{(\dot{c}_1(t_0), \dot{c}_2(t_0))}_{v^T} \cdot \underbrace{\begin{pmatrix} f_{xx}(c_0) & f_{xy}(c_0) \\ f_{yx}(c_0) & f_{yy}(c_0) \end{pmatrix}}_H \cdot \underbrace{\begin{pmatrix} \dot{c}_1(t_0) \\ \dot{c}_2(t_0) \end{pmatrix}}_v$$

- If $\forall v \in \mathbb{R}^2 \setminus \{\vec{0}\} (v^T H v > 0)$, then $\frac{d^2 z}{dt^2} > 0$ defines a **local minimum**
- If $\forall v \in \mathbb{R}^2 \setminus \{\vec{0}\} (v^T H v < 0)$, then $\frac{d^2 z}{dt^2} < 0$ defines a **local maximum**
- If $\forall v \in \mathbb{R}^2 \setminus \{\vec{0}\} (v^T H v < 0 \vee v^T H v > 0)$, the point $c_0 = (x_0, y_0)$ defines a **saddle point**

TODO:
$(J_u)_{ij} = \frac{\partial u_i}{\partial x_j}, (H_g)_{ij} = \frac{\partial^2 g}{\partial x_i \partial x_j}$ $\partial_j(u_i v_i) = (\partial_j u_i) v_i + u_i (\partial_j v_i)$

MISC


$$ax^2 + bx + c = 0 \Rightarrow x_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

TRIGONOMETRY

	0	$30^\circ = \frac{\pi}{6}$	$45^\circ = \frac{\pi}{4}$	$60^\circ = \frac{\pi}{3}$	$90^\circ = \frac{\pi}{2}$
$\sin(x)$	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1
$\cos(x)$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0
$\tan(x)$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	

$$\tan(x) = \frac{\sin(x)}{\cos(x)}$$
$$\sin(2x) = 2 \sin(x) \cos(x)$$
$$\sin^2(x) + \cos^2(x) = 1$$

1, $\sin(\alpha), \cos(\alpha)$



EXAMPLES

TODO:
