PREPRINT VERSION

Odd Zeta Values from the PCF Torus: φ and π as Arithmetic-Geometric Sources

Jorge Armando GONZÁLEZ GARCÍA [a], *Víctor Manuel GONZÁLEZ GARCÍA* [a], *Itzel Marion DRESSLER PÉREZ* [b] *and Luz María GARCÍA ORDÓÑEZ* [a]

[a)] *TTAMAYO PUNTO COM, S.A.P.I. de C.V., Mexico*
E-mail: *jorge@ttamayo.com*, *victor@ttamayo.com*

[b)] *Independent Researcher*

Received April 10, 2026; Updated April 12, 2026; Preprint hosted at Zenodo
Zenodo All Versions DOI: 10.5281/zenodo.19492791

We demonstrate that the odd values $\zeta(2k+1)$ of the Riemann zeta function are structurally determined by $\varphi = (1+\sqrt{5})/2$ and π. The Euler product of ζ is built from local factors $f_p(s)$, one for each prime. We prove that the Frobenius lift $\varphi^p = F_p\varphi + F_{p-1}$—where F_n is the n-th Fibonacci number—determines the splitting type of every prime p in $\mathbb{Z}[\varphi]$, and hence every $f_p(s)$. Through the Dedekind factorisation

$$\zeta(s) = \frac{\zeta_{\mathbb{Q}(\sqrt{5})}(s)}{L(s,\chi_5)},$$

this determines $\zeta(s)$ completely: the isomorphism between the Frobenius structure and the Euler product is demonstrated at every level (primes, splitting types, local factors, L-function), anchored by the base case $L(1,\chi_5) = 2\log\varphi/\sqrt{5}$ (the class-number formula for $\mathbb{Q}(\sqrt{5})$), which expresses the L-function value entirely in terms of φ. This resolves the apparent freedom of $\zeta(2k+1)$ noted by Elvang, Herderschee and Morales in the $\mathcal{N}$=4 SYM S-matrix bootstrap: those values are free only relative to the EFT; the pentagonal arithmetic fixes them.

Contents

1 Introduction **2**
1.1 Context . . . 3

2 The PCF ring and torus **4**
2.1 Uniqueness of φ and $\mathbb{Q}(\sqrt{5})$ . . . 5
2.2 Frobenius lifts and the splitting criterion . . . 6
2.3 Mersenne bridge and localization at 2 . . . 6

3 Mechanisms of the arithmetic formulation **7**
3.1 Prime decomposition and local Euler factors . . . 7
3.2 Dirichlet factorization . . . 8
3.3 The fundamental formula . . . 8
3.4 Even zeta values . . . 8
3.5 The pentagonal source . . . 9
3.5.1 Numerical case study: $\zeta(3)$ . . . 11
3.6 The period bridge . . . 12
3.6.1 Dimensional emergence . . . 13

4 String theory and the Primitive Structure **14**
4.1 The worldline entropy . . . 14
4.2 Bekenstein–Hawking entropy and L-values . . . 15
4.3 The unifying invariant $R = \log\varphi$ . . . 16
4.4 Geometric unity at $\tau_{\mathrm{PCF}} = i$ . . . 17
4.5 Algebraic symmetry via $\mathbb{Z}_2 \times \mathbb{Z}_2$ . . . 17
4.6 Synthesis: physical implications . . . 18
4.7 The unifying generator . . . 18

5 Categorical framework **22**
5.1 Preliminaries . . . 22
5.2 The Frobenius system as a functor . . . 22
5.3 The Euler product as a categorical colimit . . . 23
5.4 The Dedekind factorisation as a natural isomorphism . . . 23
5.5 The arithmetic–spectral span . . . 24
5.6 Implications of the categorical framework . . . 25
5.7 Connection with the forgetful functor chain . . . 25
5.8 Analogy with Weil's proof for function fields . . . 26
5.9 The polylogarithmic circuit: from $\mathbb{F}_1$ to transcendence . . . 27

6 Discussion **28**
6.1 Verification . . . 32
6.2 Open problems . . . 34
6.3 Conclusion . . . 34

A Compilation and verification **36**

1 Introduction

What determines the values $\zeta(3)$, $\zeta(5)$, $\zeta(7)$, ...? Euler showed that the even values $\zeta(2k) = r_k\pi^{2k}$ ($r_k \in \mathbb{Q}$) are governed by π alone. The odd values have resisted a comparable structural account for over two centuries: Apéry [1] proved $\zeta(3) \notin \mathbb{Q}$, Ball and Rivoal [4] showed that infinitely many $\zeta(2k+1)$ are irrational, but no organising principle has emerged. This paper identifies such a principle: the pentagonal arithmetic of $\mathbb{Q}(\sqrt{5})$, generated by $\varphi^2 = \varphi + 1$. The irrationality itself traces to $\log\varphi$ being transcendental (Lindemann–Weierstrass).

The key identity is the Dedekind factorisation

$$\zeta(s) = \frac{\zeta_{\mathbb{Q}(\sqrt{5})}(s)}{L(s,\chi_5)}, \tag{1.1}$$

where the Artin character χ_5 is dictated by the pentagon:

$$\chi_5(a) = \frac{\log|2\cos(\pi a/5)|}{\log\varphi} \in \{+1,-1\}. \tag{1.2}$$

This character classifies every prime p as split or inert in $\mathbb{Z}[\varphi]$, determining the local Euler factor $f_p(s)$ and hence the entire product $\zeta_{\mathbb{Q}(\sqrt{5})}(s) = \prod_p f_p(s)$. The base case $L(1,\chi_5) = 2\log\varphi/\sqrt{5} = T/(\pi\sqrt{5})$ ($T = 2\pi\log\varphi$) shows that π—the circumference of the circle in T^2_{PCF}—is present from the outset. The result resolves the apparent freedom noted by Elvang et al. [9]: the Wilson coefficients $a_{2k-1,0} = \alpha'^{2k+1}\zeta(2k+1)$ that are free in the $\mathcal{N}$=4 SYM EFT are fixed by this arithmetic.

The objective is summarized in the following diagram:

$$\begin{tikzcd}
 & \zeta_{\mathbb{Q}(\sqrt{5})}(2k{+}1) \arrow[rr, "\text{colimit}", "\text{(Proposition 16)}"'] & & \zeta_{\mathbb{Q}(\sqrt{5})}(s) \arrow[dr, "\text{factorisation}", "\text{(Theorem 2)}"'] & \\
\varphi^2{=}\varphi{+}1 \arrow[ur, "\text{splitting}", "\text{(Proposition 2)}"'] \arrow[dr, "\text{contraction}", "(|\ker G|{=}2)"'] & & & & \zeta(2k{+}1) \\
 & \mu_3 = 1/2 \arrow[rr, "\text{uniqueness}", "\text{(Corollary 5)}"'] & & S_3 \arrow[ur, "\text{eigenvalues}", "\text{(Lemma 2)}"'] &
\end{tikzcd} \tag{1.3}$$

The upper branch (arithmetic) and lower branch (spectral) are co-determined by the single map $G\colon (\mathbb{Z}/20\mathbb{Z})^\times \to \mathbb{Z}_2 \times \mathbb{Z}_2$ (Theorem 12). The upper branch produces the odd zeta values: G classifies primes, determining every f_p, assembling the Euler product, and delivering $\zeta(2k+1) = \zeta_{\mathbb{Q}(\sqrt{5})}(2k+1)/L(2k+1, \chi_5)$. The lower branch produces the spectral structure: $|\ker G| = 2$ forces $\mu_3 = 1/2$, which determines S_3 symmetry, eigenvalues $\omega^k/2$, the AdS radius $\ell = 1$ (from S-duality at $\tau_{\mathrm{PCF}} = i$), and the Brown–Henneaux [6] central charge $c = 3\ell/(2G_N) = 3$ (§ 4.7, Lemma 2 and Lemma 1). The regulator $R = \log\varphi$ links both branches: it is the arithmetic size of the field ($L(1, \chi_5) = 2R/\sqrt{5}$), the worldsheet period ($T = 2\pi R$), and the entropy anchor ($S_{\mathrm{BH}}/k_B = \lambda_{\log} \cdot R$). § 5 makes the categorical content explicit: $\{\psi_p\}$ is a functor, the Euler product is a colimit, the Dedekind factorisation is a natural isomorphism across the four quadratic fields classified by G, and the two branches form a span with common source G.

This span discharges the `bridge_axiom` of the Primitive Structure [12] (*The Hilbert Pólya Operator and the Primitive Structure of the Complex Plane*, hereafter *the Primitive Structure*): because G simultaneously produces $\zeta(s)$ (via the Euler product [10] and analytic continuation [20]) and the spectral constraint $\mu_3 = 1/2$ (via $|\ker G| = 2$), the same map that determines $\zeta(s)$ everywhere also constrains $\mathrm{Re}(\rho)$. The Primitive Structure establishes the functorial transport of the eigenvalue constraint $\mathrm{Re} > 0 \Rightarrow \mathrm{Re} = 1/2$ from $\widehat{\Omega}$ to ζ across the full critical strip; the present paper provides the *mechanism* underlying that transport. Theorem 10 shows that every structure in this paper derives from the single relation $\varphi^2 = \varphi + 1$.

1.1 Context

In the Primitive Structure [12], the proof that $\mathrm{Re}(\rho) = 1/2$ within the PCF categorical framework proceeds through a squeeze (Theorem 10 of [12]): two independent bounds, both derived from the arithmetic of $(\mathbb{Z}/20\mathbb{Z})^\times$ and the Galois functor $G\colon (\mathbb{Z}/20\mathbb{Z})^\times \to \mathbb{Z}_2 \times \mathbb{Z}_2$ (Definition 5 of [12]), force equality at 1/2. The squeeze itself is formally verified in Lean 4. However, the step connecting the categorical machinery to the zeros of ζ—the identification of $\mathrm{Re}(\rho)$ with an eigenvalue of $\widehat{\Omega}$—is left as an axiom (`bridge_axiom`, equivalently the structure field `bounded_by_mu : ` $\rho_{\mathrm{re}} \le \mu_3$ of `TernaryLZero`, Definition 7 of [12]).

The axiom states $\rho_{\mathrm{re}} \le \mu_3 = 1/2$: it is Premise A of the squeeze (Proposition 3 of [12]). Once provided, `hecke_1920` derives the companion bound $1 - \rho_{\mathrm{re}} \le \mu_3$ (Premise B in Theorem 10 of [12]), and the two together force $\rho_{\mathrm{re}} = 1/2$ by `linarith`. The mathematical justification is the Spectral Embedding (Proposition 5 of [12]): the Euler product enters $(\mathbb{Z}/20\mathbb{Z})^\times$, the Galois functor G classifies primes into four characters, the S_3 symmetry of the tripartite object restricts $\mathrm{spec}(\widehat{\Omega}) = \frac{1}{2}\{1, \omega, \omega^2\}$ with real parts $\{1/2, -1/4\}$, both $\le \mu_3$. In Lean the step remains an axiom. The Primitive Structure cites the Euler product identity, Gelfand's Banach algebra

theory [11], and the Dunford–Schwartz spectral mapping theorem [8] as justification, but does not demonstrate that these three classical tools act on a common structure: the Euler product organised by G through the same map that produces $\mathrm{spec}(\widehat{\Omega})$.

The present paper develops this bridge. The development formalises the relationships between the $\mathbb{F}_1$-geometric programme (Borger [5], Manin [18], López Peña–Lorscheid [16]) and the PCF framework in relation to the Euler product: the Λ-ring structure of R_{PCF}—the Frobenius lifts $\psi_p(\varphi) = \varphi^p$ with composition law $\psi_p \circ \psi_q = \psi_{pq}$—is simultaneously the $\mathbb{F}_1$-descent data in the sense of Borger and the source of every local Euler factor $f_p(s)$ through the Fibonacci splitting criterion $\chi_5(p) \equiv F_p \pmod{p}$ (Proposition 2), with the classification of residue classes in $(\mathbb{Z}/20\mathbb{Z})^\times$ via the golden ratio due to Grisales Herrera [13]. Establishing the categorical infrastructure—the Frobenius system as a functor (§ 5.2), the Euler product as a colimit (§ 5.3), the Dedekind factorisation as a natural isomorphism (§ 5.4), and the arithmetic–spectral span (§ 5.5)—we found that the same chain that discharges the bridge also yields a structural determination of the odd zeta values: $\zeta(2k+1) = \zeta_{\mathbb{Q}(\sqrt{5})}(2k+1)/L(2k+1, \chi_5)$, with every component fixed by the pentagonal arithmetic of $\mathbb{Q}(\sqrt{5})$ and no free parameter. This determination connects to a problem even older than the Riemann Hypothesis—the structural origin of the odd zeta values—and relates it to string theory from a viewpoint that both complements and constrains recent independent developments.

Elvang, Herderschee and Morales [9] demonstrate that in the weakly-coupled 4d planar $\mathcal{N} = 4$ SYM EFT, the Wilson coefficients

$$a_{2k-1,0} = \alpha'^{2k+1}\zeta(2k+1), \qquad k = 1, 2, 3, \ldots$$

are free parameters: supersymmetry and tree-level factorisation fix all $a_{2k,q}$ in terms of lower-order coefficients but leave $a_{2k-1,0}$ unconstrained. In the Veneziano amplitude [21] these take the values $\alpha'^{2k+1}\zeta(2k+1)$, but from the EFT alone the odd zeta values enter as external data.

The PCF (Precedent-Current-Forthcoming) framework, formalised in the Primitive Structure [12], provides the missing arithmetic structure. The present paper shows that once the Frobenius action of φ on the primes of $\mathbb{Q}(\sqrt{5})$ is taken into account, the odd zeta values are determined—not through new closed forms, but through the structural identity (1.1) together with the class-number formula for $\mathbb{Q}(\sqrt{5})$.

In § 2, we review the Frobenius structure of the PCF. In § 3.1, we classify the rational primes into split and inert types. In § 3.2, we establish the Dedekind factorisation $\zeta = \zeta_{\mathbb{Q}(\sqrt{5})}/L(s, \chi_5)$. In § 3.3, we deduce the fundamental formula. In § 3.4, we verify consistency with the even values. In § 3.5, we identify the pentagonal source of the determination. In § 5, we present the arithmetic–spectral span. In § 6, we discuss the string theory implications. We conclude in §§ 6.2 and 6.3 with open problems and the final summary.

2 The PCF ring and torus

Definition 1 (PCF ring)**.** The *PCF ring* is

$$R_{\mathrm{PCF}} = \mathbb{Z}\big[\varphi,\, \varphi^{-1},\, \tfrac{1}{2}\big]\,,$$

the ring of integers of $\mathbb{Q}(\sqrt{5})$ localised at 2. Here $\varphi = (1+\sqrt{5})/2$ satisfies $\varphi^2 = \varphi + 1$, $N(\varphi) = \varphi\bar{\varphi} = -1$ (unit), $\mathrm{Tr}(\varphi) = 1$. The discriminant of $\mathbb{Q}(\sqrt{5})$ is $\Delta = 5$.

Definition 2 (Frobenius lifts)**.** For each prime p, the *Frobenius lift* is the ring endomorphism

$$\psi_p\colon R_{\mathrm{PCF}} \to R_{\mathrm{PCF}}, \qquad \psi_p(\varphi) = \varphi^p = F_p\varphi + F_{p-1},$$

where F_n denotes the n-th Fibonacci number. These satisfy the composition law $\psi_p \circ \psi_q = \psi_{pq}$, mirroring the multiplicative structure of the Euler product.

Definition 3 (PCF torus and worldline)**.** The *PCF torus* is $T^2_{\mathrm{PCF}} = \mathbb{C}/\Lambda_{\mathrm{PCF}}$ with modular parameter $\tau_{\mathrm{PCF}} = i$, the unique fixed point of S-duality $S\colon \tau \mapsto -1/\tau$ on the upper half-plane (since $-1/i = i$). The worldline on T^2_{PCF} is

$$\Omega(\tau) = \tfrac{1}{2}\, e^{i\tau \log \varphi}.$$

Its natural period, obtained by closing on the circle of T^2_{PCF}, is

$$T = 2\pi \log \varphi. \tag{2.1}$$

Remark 1 (The circle is there)**.** Although $\log \varphi$ appears without π in the class-number formula (§ 3.3), the period $T = 2\pi \log \varphi$ shows that π is the circumference of the circle in T^2_{PCF}. At $s = 1$, the π of the period and the π of the Dedekind residue formula cancel, leaving $2 \log \varphi/\sqrt{5}$; but the circle origin is geometric, not algebraic.

Remark 2 (Canonical identifications)**.** The following classical identifications are used throughout:

1. $\mathbb{R} \cong$ the real line as a 1D topological manifold (Euclidean topology);
2. $\varphi^\sigma : \mathbb{R} \to \mathbb{R}_{>0}$ is a bijection (exponential with base $\varphi > 1$);
3. every positive integer factors uniquely into primes (Fundamental Theorem of Arithmetic; Euclid, Prop. IX.14);
4. $[\mathbb{Q}(\sqrt{5}) : \mathbb{Q}] = 2$ (minimal polynomial $x^2 - x - 1$ is irreducible over $\mathbb{Q}$).

2.1 Uniqueness of φ and $\mathbb{Q}(\sqrt{5})$

The Eq. (1.1) uses the specific field $\mathbb{Q}(\sqrt{5})$. We show this choice is *forced*, not ad hoc.

Proposition 1 (Uniqueness of φ)**.** *φ is the unique real quadratic irrationality whose minimal polynomial $p(x) = x^2 - x - 1$ is self-dual under $x \mapsto -1/x$:*

$$p(-1/x) = -\frac{p(x)}{x^2}.$$

Any other real quadratic irrationality ξ with minimal polynomial $q(x) = x^2 + ax + b$ satisfies this self-duality only if $a = -1$, $b = -1$, i.e. $q = p$ uniquely.

Proof. $p(-1/x) = 1/x^2 + 1/x - 1 = -(x^2 - x - 1)/x^2 = -p(x)/x^2$. For $q(x) = x^2 + ax + b$: $q(-1/x) = -(bx^2 + ax - 1)/x^2$. Equality $q(-1/x) = -q(x)/x^2$ requires $b = -1$ and $a = -1$. ■

Corollary 1 (Uniqueness of $\mathbb{Q}(\sqrt{5})$)**.** *$\mathbb{Q}(\sqrt{5})$ is the unique real quadratic field in which:*

1. *The fundamental unit φ satisfies $\varphi^2 = \varphi + 1$ (PCF relation);*
2. *The norm is $N(\varphi) = \varphi\bar{\varphi} = -1$ (φ is a unit of norm -1);*
3. *The discriminant is $\Delta = 5$, making $p = 5$ the unique ramified prime;*
4. *The Galois conjugation $\varphi \leftrightarrow \bar{\varphi} = -1/\varphi$ is a self-duality of the lattice Λ_{PCF} (Proposition 1).*

These four properties together single out $\mathbb{Q}(\sqrt{5})$ as the unique quadratic field on which the PCF Euler product factorisation $\zeta(s) = \zeta_{\mathbb{Q}(\sqrt{5})}(s)/L(s, \chi_5)$ rests.

2.2 Frobenius lifts and the splitting criterion

We now make explicit the connection between the Frobenius lift $\psi_p(\varphi) = \varphi^p$ and the prime splitting type $\chi_5(p)$, closing the chain pentagon $\to \varphi \to \chi_5 \to G \to R_{\mathrm{PCF}}$.

Proposition 2 (Fibonacci splitting criterion)**.** *For every prime $p \neq 2, 5$, the Frobenius lift $\psi_p(\varphi) = \varphi^p = F_p\varphi + F_{p-1}$ determines the splitting type of p in $\mathbb{Z}[\varphi]$ through:*

$$\chi_5(p) \equiv F_p \pmod{p}, \tag{2.2}$$

where $\chi_5(p) \in \{+1, -1\}$ is read as ± 1 in $\mathbb{Z}/p\mathbb{Z}$. Explicitly:

$$\chi_5(p) = \begin{cases} +1 & (p \text{ splits}) \iff F_p \equiv +1 \pmod{p} \iff F_{p-1} \equiv 0 \pmod{p}, \\ -1 & (p \text{ inert}) \iff F_p \equiv -1 \pmod{p} \iff F_{p+1} \equiv 0 \pmod{p}. \end{cases}$$

Proof. The Frobenius lift satisfies $\psi_p(a) \equiv a^p \pmod{p}$ (§ 2.2).

If p splits in $\mathbb{Z}[\varphi]$: the Frobenius acts as the identity on the residue field, so $\varphi^p \equiv \varphi \pmod{p}$, giving $F_p \equiv 1$ and $F_{p-1} \equiv 0 \pmod{p}$. Then $\chi_5(p) = +1 = F_p \bmod p$.

If p is inert: the Frobenius acts as the non-trivial Galois automorphism $\varphi \mapsto \bar{\varphi} = 1 - \varphi$, so $\varphi^p \equiv \bar{\varphi} \pmod{p}$, giving $F_p \equiv -1$ and $F_{p-1} \equiv 1 \pmod{p}$. Then $\chi_5(p) = -1 \equiv F_p \bmod p$.

In both cases $\chi_5(p) \equiv F_p \pmod{p}$. Representative values for $3 \leq p \leq 59$ are provided in Table 1. ■

Table 1. Fibonacci criterion for prime splitting in $\mathbb{Z}[\varphi]$. $\chi_5(p) \equiv F_p \pmod{p}$ in all cases.

p	$p \bmod 5$	$\chi_5(p)$	Type	$F_p \bmod p$
3	3	-1	inert	-1
7	2	-1	inert	-1
11	1	$+1$	split	$+1$
13	3	-1	inert	-1
17	2	-1	inert	-1
19	4	$+1$	split	$+1$
23	3	-1	inert	-1
29	4	$+1$	split	$+1$
31	1	$+1$	split	$+1$

Remark 3 (The Artin character is the Fibonacci character)**.** Proposition 2 expresses the Artin character $\chi_5 = (\cdot/5)$ of $\mathbb{Q}(\sqrt{5})/\mathbb{Q}$ in terms of Fibonacci numbers: $\chi_5(p) = F_p \bmod p$. This is a direct consequence of $\psi_p(\varphi) = \varphi^p = F_p\varphi + F_{p-1}$ and the Frobenius-splitting correspondence. It shows that the prime decomposition of $\mathbb{Z}[\varphi]$—which governs every local factor $f_p(s)$ of $\zeta_{\mathbb{Q}(\sqrt{5})}(s)$—is encoded in the Fibonacci sequence itself.

2.3 Mersenne bridge and localization at 2

We now explain why $R_{\mathrm{PCF}} = \mathbb{Z}[\varphi, \varphi^{-1}, \frac{1}{2}]$ rather than $\mathbb{Z}[\varphi]$ is the correct ring, and how the localization at 2 connects to the binary structure.

Theorem 1 (Mersenne bridge; González García et al. [12])**.**

$$\varphi^{\lambda_{\log}} = 2, \qquad \lambda_{\log} = \frac{\log 2}{\log \varphi} \approx 1.4404. \tag{2.3}$$

Proof. $\varphi^{\lambda_{\log}} = e^{(\log 2/\log\varphi)\cdot\log\varphi} = e^{\log 2} = 2.$ ■

Proposition 3 (Inertness of $p = 2$ and the localization)**.** *The prime $p = 2$ is* inert *in $\mathbb{Z}[\varphi]$: the Kronecker symbol $(5/2) = \chi_5(2) = -1$. Its local Euler factor in $\zeta_{\mathbb{Q}(\sqrt{5})}(s)$ is $(1 - 2^{-2s})^{-1}$, not $(1 - 2^{-s})^{-2}$.*

Localising at 2 to form $R_{\mathrm{PCF}} = \mathbb{Z}[\varphi, \varphi^{-1}, \frac{1}{2}]$ is justified by the Mersenne bridge Eq. (2.3): $\varphi^{\lambda_{\log}} = 2$ means that 2 is a natural scale in the tower generated by φ, and the factor $\frac{1}{2}$ in R_{PCF} captures the inert place $p = 2$ without introducing a new ramified prime.

Proof. $\chi_5(2) = (5/2) = -1$ since $2 \not\equiv \pm 1 \pmod 5$. The local factor follows from Proposition 4. The Mersenne bridge is Theorem 1. ■

Remark 4 (The non-circularity chain)**.** The full acyclic chain from the pentagon to the Euler product is:

$$\underbrace{2\cos\tfrac{\pi}{5} = \varphi}_{\text{pentagon}} \to \underbrace{\varphi^2 = \varphi + 1}_{R_{\mathrm{PCF}}} \to \underbrace{\chi_5(p) = F_p \bmod p}_{\text{Artin char.}} \to \underbrace{G(p \bmod 20) = (\chi_4(p), \chi_5(p))}_{\text{class functor}}$$

$$\to \underbrace{\zeta_{\mathbb{Q}(\sqrt{5})}(s) = \textstyle\prod_p f_p(s)}_{\text{Euler product}} \to \zeta(s).$$

At no point does $\zeta(s)$ itself enter the construction of χ_5 or $\zeta_{\mathbb{Q}(\sqrt{5})}$; the chain is acyclic. The localisation at 2 is justified by $\varphi^{\lambda_{\log}} = 2$ (Theorem 1). Each arrow in this chain is a functor or natural transformation in the categorical framework of § 5: $\varphi^2 = \varphi + 1$ generates the Frobenius functor Ψ (Definition 6), Ψ determines χ_5 via the Fibonacci splitting criterion (Proposition 2), χ_5 is a component of G (Theorem 9), and G assembles the Euler product as a colimit (Proposition 16). The acyclicity of the chain is the acyclicity of a composed functor.

3 Mechanisms of the arithmetic formulation

3.1 Prime decomposition and local Euler factors

Since $\Delta = 5$ is the only prime dividing the discriminant of $\mathbb{Q}(\sqrt{5})$, the decomposition of primes in $\mathbb{Z}[\varphi]$ is determined by the Legendre/Kronecker symbol $\chi_5(p) = (5/p)$:

Proposition 4 (Local factors)**.** *The Dedekind zeta function of $\mathbb{Q}(\sqrt{5})$ has the Euler product*

$$\zeta_{\mathbb{Q}(\sqrt{5})}(s) = \prod_{p \text{ prime}} f_p(s), \tag{3.1}$$

where

$$f_p(s) = \begin{cases} (1 - p^{-s})^{-1} & \chi_5(p) = 0 \ \textit{(ramified; only } p = 5\textit{)}, \\ (1 - p^{-s})^{-2} & \chi_5(p) = +1 \ \textit{(split; e.g. } p = 11, 19, 29, \dots\textit{)}, \\ (1 - p^{-2s})^{-1} & \chi_5(p) = -1 \ \textit{(inert; e.g. } p = 2, 3, 7, 13, \dots\textit{)}. \end{cases}$$

In particular, $p = 2$ is inert *(Kronecker $(5/2) = -1$), so the factor $\frac{1}{2}$ in R_{PCF} does not introduce a ramified place.*

Each factor $f_p(s)$ is completely determined by the Frobenius lift $\psi_p(\varphi) = \varphi^p$.

Proof. The classical theory of Dedekind zeta functions for quadratic fields; see e.g. [19]. The claim that f_p is determined by ψ_p follows from the fact that $\chi_5(p) \equiv F_p \pmod p$ (Proposition 2): whether φ^p is congruent to φ or $\bar{\varphi}$ modulo p determines the splitting. ■

3.2 Dirichlet factorization

Theorem 2 (Factorisation)**.**

$$\zeta(s) = \frac{\zeta_{\mathbb{Q}(\sqrt{5})}(s)}{L(s,\chi_5)}, \tag{3.2}$$

where $L(s,\chi_5) = \prod_p (1-\chi_5(p)p^{-s})^{-1}$ *is the Dirichlet L-function of the symbol* $\chi_5 = (5/\cdot)$*.*

Proof. The Dedekind zeta of a quadratic field K factorises as $\zeta_K(s) = \zeta(s)L(s,\chi_K)$, where χ_K is the Kronecker character associated with $K/\mathbb{Q}$; for $K = \mathbb{Q}(\sqrt{5})$ this is χ_5. See [19], Ch. VII. ■

Remark 5 (The Euler product determines ζ everywhere)**.** The identity $\prod_p (1-p^{-s})^{-1} = \sum_{n=1}^{\infty} n^{-s}$ for $\mathrm{Re}(s) > 1$ (Euler [10]) follows from unique factorisation of $\mathbb{N}$ (Remark 2). The Dirichlet coefficients $a_n = 1$ determine $\zeta(s)$ on all of $\mathbb{C}$ by analytic continuation (identity theorem for Dirichlet series; Riemann [20]). Therefore the Euler product—and hence G—determines $\zeta(s)$ everywhere, including the critical strip $0 < \mathrm{Re}(s) < 1$ and its zeros.

Remark 6 (Naturality of the factorisation)**.** The factorisation $\zeta(s) = \zeta_{\mathbb{Q}(\sqrt{5})}(s)/L(s,\chi_5)$ is not specific to $\mathbb{Q}(\sqrt{5})$: it is the $K = \mathbb{Q}(\sqrt{5})$ component of a natural isomorphism η across all quadratic extensions classified by G (Theorem 11). The four components of η correspond to the four fibers of the Galois functor.

3.3 The fundamental formula

Theorem 3 (Fundamental formula)**.**

$$L(1,\chi_5) = \frac{2\log\varphi}{\sqrt{5}} = \frac{T}{\pi\sqrt{5}}, \tag{3.3}$$

where $T = 2\pi\log\varphi$ *is the period of* T^2_{PCF}*.*

Proof. This is the Dirichlet class-number formula for $\mathbb{Q}(\sqrt{5})$. With class number $h = 1$, regulator $R = \log\varphi$, number of roots of unity $w = 2$, and discriminant $\Delta = 5$:

$$\lim_{s\to 1}(s-1)\,\zeta_{\mathbb{Q}(\sqrt{5})}(s) = \frac{2^{r_1}(2\pi)^{r_2}hR}{w\sqrt{\Delta}} = \frac{4\cdot 1\cdot 1\cdot\log\varphi}{2\sqrt{5}} = \frac{2\log\varphi}{\sqrt{5}}.$$

Since $\zeta_{\mathbb{Q}(\sqrt{5})}(s) = \zeta(s)L(s,\chi_5)$ and $\lim_{s\to 1}(s-1)\zeta(s) = 1$, we obtain $L(1,\chi_5) = 2\log\varphi/\sqrt{5}$. The identity $T/(\pi\sqrt{5}) = 2\log\varphi/\sqrt{5}$ follows from $T = 2\pi\log\varphi$. ■

Remark 7. The Eq. (3.3) exposes the role of the circle. Writing $L(1,\chi_5) = T/(\pi\sqrt{5})$, we see that π from the torus period $T = 2\pi\log\varphi$ cancels against π in the denominator, leaving $2\log\varphi/\sqrt{5}$. The π—and hence the circle of T^2_{PCF}—is present from the outset.

3.4 Even zeta values

Theorem 4 (Even formula)**.** *For* $k \geq 1$*:*

$$L(2k,\chi_5) = \frac{(-1)^{k+1}\sqrt{5}\,(2\pi)^{2k}\,B_{2k,\chi_5}}{2\,(2k)!\,5^{2k}}, \tag{3.4}$$

where $B_{2k,\chi_5} \in \mathbb{Q}$ are the generalised Bernoulli numbers:

$$B_{n,\chi_5} = 5^{n-1} \sum_{a=1}^{5} \chi_5(a)\, B_n\left(\tfrac{a}{5}\right).$$

Computed values: $B_{2,\chi_5} = \frac{4}{5}$, $B_{4,\chi_5} = -8$, $B_{6,\chi_5} = \frac{804}{5}$, $B_{8,\chi_5} = -5776$.

Proof. Classical result: generalized Bernoulli formula for primitive even characters; see e.g. [23], Thm. 4.2. The factor $\sqrt{5}$ is the Gauss sum $\tau(\chi_5) = \sqrt{5}$. ■

3.5 The pentagonal source

The connection between φ and the odd L-values runs through the pentagon. We establish three identities—trigonometric, logarithmic, and analytic—that collectively show how φ and π organise $L(s, \chi_5)$ for *all* $s \geq 1$ via the same pentagonal structure.

Proposition 5 (Pentagon identities)**.** *The golden ratio φ identifies the fundamental scales of the pentagon:*

$$2\cos\tfrac{\pi}{5} = \varphi, \qquad 2\cos\tfrac{2\pi}{5} = \varphi^{-1}. \tag{3.5}$$

These primary values extend symmetrically to the full set of four coordinates:

$$2\cos\tfrac{\pi}{5} = \varphi, \quad 2\cos\tfrac{2\pi}{5} = \varphi^{-1}, \quad 2\cos\tfrac{3\pi}{5} = -\varphi^{-1}, \quad 2\cos\tfrac{4\pi}{5} = -\varphi. \tag{3.6}$$

Proof. The regular pentagon inscribed in the unit circle has diagonal-to-side ratio φ (Euclid, Prop. XIII.8). The half-angle of the central subtended by one side is $\pi/5$, giving $2\cos(\pi/5) = \varphi$. The second identity follows from $\varphi^{-1} = \varphi - 1$ and the double-angle formula. Verified numerically to machine precision (see § 6.1): yielding a zero residual. ■

Proposition 6 (Gauss logarithmic formula)**.**

$$\sum_{a=1}^{4} \chi_5(a)\, \log\!\left(2\sin\tfrac{\pi a}{5}\right) = -2\log\varphi. \tag{3.7}$$

Proof. Direct computation using $\chi_5(1) = \chi_5(4) = +1$, $\chi_5(2) = \chi_5(3) = -1$, and the identities $\sin(\pi/5)\sin(4\pi/5) = \sin^2(\pi/5)$, $\sin(2\pi/5)\sin(3\pi/5) = \sin^2(2\pi/5)$. The result follows from the product formula $\prod_{a=1}^{4} \sin(\pi a/5) = \sqrt{5}/4$ and $2\sin(\pi/5) \cdot 2\sin(2\pi/5) = \sqrt{5}$, combined with Proposition 5. Verified numerically (see § 6.1). ■

Corollary 2 (Derivation of $L(1, \chi_5)$ from pentagon)**.** *The class-number formula (Theorem 3) follows from Proposition 6 via the Gauss–Lerch formula for primitive even characters with conductor f and Gauss sum $\tau(\chi_5) = \sqrt{5}$:*

$$L(1,\chi_5) = -\frac{\tau(\chi_5)}{f} \sum_{a=1}^{f-1} \chi_5(a) \log\!\left(2\sin\tfrac{\pi a}{f}\right) = -\frac{\sqrt{5}}{5} \cdot (-2\log\varphi) = \frac{2\log\varphi}{\sqrt{5}} = \frac{T}{\pi\sqrt{5}}. \tag{3.8}$$

The pentagon generates φ (Proposition 5), which enters $L(1, \chi_5)$ through the Gauss sum over $\sin(\pi a/5)$.

Proposition 7 (Digamma formula)**.** *Let $\psi = \Gamma'/\Gamma$ denote the digamma function. Then*

$$\sum_{a=1}^{4} \chi_5(a)\, \psi\!\left(\tfrac{a}{5}\right) = -2\sqrt{5}\, \log \varphi, \tag{3.9}$$

and consequently $L(1, \chi_5) = -\frac{1}{5} \sum_{a=1}^{4} \chi_5(a)\, \psi(a/5) = 2 \log \varphi/\sqrt{5}$.

Proof. The digamma formula at rational arguments gives $\psi(a/5) = -\gamma - \log(5) - \frac{\pi}{2} \cot(\pi a/5) + 2 \sum_{k=1}^{2} \cos(2\pi k a/5) \log \sin(\pi k/5)$ (Gauss's digamma theorem). The χ_5-weighted sum eliminates γ and $\log(5)$ (since $\sum \chi_5(a) = 0$) and reduces to the logarithmic formula of Proposition 6. Verified numerically (see § 6.1). ■

Proposition 8 (Pentagon dictates χ_5)**.** *The Artin character χ_5 of $\mathbb{Q}(\sqrt{5})/\mathbb{Q}$ is completely determined by the pentagon through:*

$$\chi_5(a) = \frac{\log |2\cos(\pi a/5)|}{\log \varphi} \in \{+1, -1\}, \qquad a = 1, 2, 3, 4. \tag{3.10}$$

Equivalently:

$$|2\cos(\pi a/5)| = \varphi^{\chi_5(a)}. \tag{3.11}$$

The four pentagonal cosines $2\cos(\pi a/5) \in \{\pm\varphi, \pm\varphi^{-1}\}$ are the four images of the Galois functor $G\colon (\mathbb{Z}/20\mathbb{Z})^\times \to \mathbb{Z}_2 \times \mathbb{Z}_2$ evaluated at the residues $\{1, 2, 3, 4\} \subset \mathbb{Z}/5\mathbb{Z}$.

Proof. Proposition 5 gives: $2\cos(\pi/5) = \varphi$ and $2\cos(2\pi/5) = \varphi^{-1}$. By symmetry of cosine: $2\cos(3\pi/5) = -\varphi^{-1}$ and $2\cos(4\pi/5) = -\varphi$. Hence $|2\cos(a\pi/5)| \in \{\varphi, \varphi^{-1}\}$ for all $a \in \{1, 2, 3, 4\}$. Taking logarithms: $\log |2\cos(a\pi/5)| \in \{\log \varphi, -\log \varphi\}$. Dividing by $\log \varphi$: $\log |2\cos(a\pi/5)|/\log \varphi \in \{+1, -1\}$.

The identification with $\chi_5(a)$: for $a \in \{1, 4\}$ (quadratic residues mod 5), $|2\cos(a\pi/5)| = \varphi > \varphi^{-1}$, so the ratio is $+1 = \chi_5(a)$. For $a \in \{2, 3\}$ (non-residues), $|2\cos(a\pi/5)| = \varphi^{-1} < \varphi$, so the ratio is $-1 = \chi_5(a)$. Verified numerically (see § 6.1). ■

Remark 8 (Connection to the Galois functor)**.** The pentagonal cosines of Proposition 8 reappear as the $(0, 1)$-component of the Galois functor $G\colon (\mathbb{Z}/20\mathbb{Z})^\times \to \mathbb{Z}_2 \times \mathbb{Z}_2$ (Theorem 9 below): $\chi_5(a)$ is simultaneously the pentagonal fingerprint (3.10) and the second coordinate of $G(a \bmod 20)$.

Remark 9 (The pentagonal dictate)**.** Proposition 8 makes explicit what "φ dictates χ_5" means concretely. The chain is:

$$\varphi = 2\cos\tfrac{\pi}{5} \;\Rightarrow\; \chi_5(a) = \tfrac{\log |2\cos(a\pi/5)|}{\log \varphi}$$
$$\Rightarrow\; L(s, \chi_5) = \tfrac{1}{5^s} \textstyle\sum_a \chi_5(a)\, \zeta(s, \tfrac{a}{5}) \;\Rightarrow\; \zeta(2k+1) = \tfrac{\zeta_{\mathbb{Q}(\sqrt{5})}(2k+1)}{L(2k+1, \chi_5)}.$$

The values $\zeta(2k+1)$ that appear as free parameters in the EFT of Elvang et al. are not free: they are Hurwitz projections at the pentagonal points $\{a/5\}$ with weights $\chi_5(a)$, which are themselves the logarithmic fingerprints of φ in the pentagon. The "dictum" is $\varphi = 2\cos(\pi/5)$; everything else follows.

Theorem 5 (Hurwitz representation and pentagonal source)**.** *For all s with $\mathrm{Re}(s) > 1$ (and by analytic continuation for $s \neq 1$):*

$$L(s, \chi_5) = \frac{1}{5^s} \sum_{a=1}^{4} \chi_5(a)\, \zeta(s, \tfrac{a}{5}), \tag{3.12}$$

where $\zeta(s,x) = \sum_{n=0}^{\infty}(n+x)^{-s}$ *is the Hurwitz zeta function. The four arguments* $a/5$ *(*$a = 1,2,3,4$*) are the* pentagonal points*, and* χ_5 *is their Artin weight.*

The three kernels at special values:

$$s=1: \quad \zeta(s,a/5) \xrightarrow{s\to 1} -\psi(a/5) - \log 5 \;\Rightarrow\; L(1,\chi_5) = \tfrac{2\log\varphi}{\sqrt{5}}, \tag{3.13}$$

$$s=2k: \quad \zeta(2k,a/5) \text{ via Bernoulli} \;\Rightarrow\; L(2k,\chi_5) = \sqrt{5}\cdot\pi^{2k}\cdot\mathbb{Q}, \tag{3.14}$$

$$s=2k+1,\ k\geq 1: \quad \zeta(2k+1,a/5) = \sum_{n=0}^{\infty}(n+a/5)^{-(2k+1)}. \tag{3.15}$$

In all three cases the organising structure is identical: the same $\chi_5(a)$*-weights over the same pentagonal points* $\{a/5\}$*.*

Proof. Equation (3.12) is well-established (Apostol, Thm. 12.6). The kernel at $s=1$ follows from $\zeta(s,x) = 1/(s-1)+\psi(x)+O(s-1)$ combined with $\sum\chi_5(a)=0$ (so the $1/(s-1)$ pole and $\log 5$ cancel), giving Proposition 7 and hence Corollary 2. The kernel at even s gives Theorem 4. The odd kernel Eq. (3.15) is the direct Hurwitz series, convergent for $s>1$. ■

Theorem 6 (Pentagonal determination of odd zeta values)**.** *For* $k\geq 0$*, the factorisation* $\zeta(2k+1) = \zeta_{\mathbb{Q}(\sqrt{5})}(2k+1)/L(2k+1,\chi_5)$ *expresses* $\zeta(2k+1)$ *as the ratio of two* χ_5*-projections of Hurwitz zeta at pentagonal points. All data are generated by:*

1. $2\cos(\pi/5) = \varphi$ (pentagonal geometry generates φ)*,*
2. $\sum_{a=1}^{4}\chi_5(a)\log(2\sin\frac{\pi a}{5}) = -2\log\varphi$ (the logarithmic signature of φ in the pentagon)*,*
3. $\psi_p(\varphi) = \varphi^p \;\Rightarrow\; \chi_5(p) =$ *Artin character* (Frobenius connects φ to the L-function character)*.*

The case $k=0$ *yields the exact closed form* $L(1,\chi_5) = T/(\pi\sqrt{5})$*. For* $k\geq 1$*, the determination is structural:* $L(2k+1,\chi_5)$ *and* $\zeta_{\mathbb{Q}(\sqrt{5})}(2k+1)$ *are both projections of the same pentagonal Hurwitz structure, organised by* $\chi_5 =$ *Artin character of* $\mathbb{Q}(\sqrt{5})/\mathbb{Q}$*. No additional parameter beyond* φ*,* π*, and the pentagonal arithmetic of* $\mathbb{Q}(\sqrt{5})$ *is required.*

Proof. Items (1)–(3) are Propositions 5 and 6 and Theorem 9 respectively. The factorisation is Theorem 2. The Hurwitz representation is Theorem 5. The $k=0$ closed form is Corollary 2. For $k\geq 1$: both $\zeta_{\mathbb{Q}(\sqrt{5})}(2k+1) = \zeta(2k+1)\cdot L(2k+1,\chi_5)$ and $L(2k+1,\chi_5) = 5^{-(2k+1)}\sum\chi_5(a)\zeta(2k+1,a/5)$ are determined by the Hurwitz series at pentagonal points weighted by χ_5. Neither factor has a simpler closed form (Open Problem 6.1), but their ratio $\zeta(2k+1)$ is fixed by the arithmetic of R_{PCF} without any free parameter. ■

Remark 10 (Structural vs. closed-form determination)**.** Theorem 6 establishes *structural* determination: the pentagonal lattice $\{a/5\}$, the Artin character χ_5, and the Hurwitz kernel $\zeta(2k+1,\cdot)$ are all fixed by φ and π without additional parameters. A closed-form expression $\zeta(2k+1) = r_k\pi^{2k+1}$ for $r_k\in\mathbb{Q}$ would be stronger; whether it exists for $k\geq 1$ is an open problem related to Apéry's theorem and its generalisations. What the present paper establishes is that the *source*—the organising structure—is the pentagon generated by φ and π, not free arithmetic data.

3.5.1 Numerical case study: $\zeta(3)$

We trace the full chain from the pentagon to $\zeta(3) = 1.2020569\ldots$ for the first few primes.

Step 1: Pentagon → character. $\varphi = 2\cos(\pi/5)$. The Artin character $\chi_5(p) = (5/p)$ classifies primes: $\chi_5(2) = -1$, $\chi_5(3) = -1$, $\chi_5(7) = -1$, $\chi_5(11) = +1$, $\chi_5(13) = -1$, $\chi_5(19) = +1$.

Step 2: Character → local factors at $s = 3$. Split ($\chi_5 = +1$): $f_p(3) = (1 - p^{-3})^{-2}$. Inert ($\chi_5 = -1$): $f_p(3) = (1 - p^{-6})^{-1}$. Ramified ($p = 5$): $f_5(3) = (1 - 5^{-3})^{-1}$. Explicitly:

$$f_2(3) = \frac{64}{63}, \quad f_3(3) = \frac{729}{728}, \quad f_5(3) = \frac{125}{124}, \quad f_{11}(3) = \left(\frac{1331}{1330}\right)^2.$$

Step 3: Euler product → $\zeta(3)$. $\zeta_{\mathbb{Q}(\sqrt{5})}(3) = \prod_p f_p(3)$ and $L(3, \chi_5) = \prod_p (1 - \chi_5(p)p^{-3})^{-1}$. Their ratio gives

$$\zeta(3) = \frac{\zeta_{\mathbb{Q}(\sqrt{5})}(3)}{L(3, \chi_5)} = 1.2020569\ldots$$

Every factor in both products is determined by $\chi_5(p)$, which is determined by F_p mod p (Proposition 2), which is determined by $\varphi^p = F_p\varphi + F_{p-1}$ (Definition 2). No parameter is chosen; the pentagon dictates the value. Verified numerically (see § 6.1).

The mechanism extends to all odd values. Table 2 displays the decomposition $\zeta(2k+1) = \zeta_{\mathbb{Q}(\sqrt{5})}(2k+1)/L(2k+1, \chi_5)$ alongside the even values $\zeta(2k) = r_k\pi^{2k}$, illustrating the two regimes: even values governed by π alone, odd values by both φ and π.

Table 2. Zeta values from $s = 2$ to $s = 13$. Even values (shaded) are rational multiples of π^{2k}; odd values are ratios $\zeta_{\mathbb{Q}(\sqrt{5})}(s)/L(s, \chi_5)$, both factors determined by χ_5 and hence by φ. All entries verified at 50-digit precision (see § 6.1).

s	$\zeta_{\mathbb{Q}(\sqrt{5})}(s)$	$L(s, \chi_5)$	$\zeta(s)$	Source
2	—	—	$\pi^2/6 \approx 1.6449$	π
3	1.02755	0.85482	1.20206	φ, π
4	—	—	$\pi^4/90 \approx 1.0823$	π
5	1.00133	0.96567	1.03693	φ, π
6	—	—	$\pi^6/945 \approx 1.0173$	π
7	1.00007	0.99179	1.00835	φ, π
8	—	—	$\pi^8/9450 \approx 1.0041$	π
9	1.000004	0.99800	1.00201	φ, π
10	—	—	$\pi^{10}/93555 \approx 1.00099$	π
11	1.0000003	0.99951	1.00049	φ, π
12	—	—	$691\pi^{12}/638512875 \approx 1.00025$	π
13	1.00000002	0.99988	1.00012	φ, π

3.6 The period bridge

The two structures of T^2_{PCF} interact precisely through the period T:

Structure	Object	Enters via
Multiplicative (ring)	$\psi_p(\varphi) = \varphi^p$	Euler product factors $f_p(s)$
Additive/circular (torus)	$T = 2\pi \log\varphi$	$L(1, \chi_5) = T/(\pi\sqrt{5})$

The Eq. (3.3) is the bridge: π (circle) and $\log\varphi$ (Frobenius regulator) combine in T, then π cancels to leave $2\log\varphi/\sqrt{5}$. At $s = 2k$ the π reappears explicitly via Eq. (3.4). At $s = 2k+1$ for $k \geq 1$ the same two structures produce $\zeta(2k+1)$ through their ratio.

The table displays two faces of T^2_{PCF}: multiplicative (Frobenius, Euler product) and additive (period, circle). We now show that both arise from the self-measurement $\varphi^2 = \varphi + 1$.

3.6.1 Dimensional emergence

Proposition 9 (Dimensional chain from self-measurement)**.** *The algebraic relations $i^2 = -1$ and $\varphi^2 = \varphi + 1$ generate a dimensional chain*

$$\underbrace{\mathbb{R}}_{d=1} \xrightarrow{\times\, i} \underbrace{\mathbb{C}}_{d=2} \xrightarrow{\times\, \varphi} \underbrace{E^3}_{d=3}, \tag{3.16}$$

where $E^3 = \{(x,\, y,\, \varphi y) : x, y \in \mathbb{R}\}$ is the golden extension of $\mathbb{C}$. Under the coupling $z = \varphi y$, the characteristic equation $\varphi^2 = \varphi + 1$ manifests as the self-measurement identity:

$$z^2 = z \cdot y + y^2. \tag{3.17}$$

Proof. (i) $\mathbb{R} \to \mathbb{C}$**:** This is the foundational construction of $\mathbb{C}$ as $\mathbb{R}[x]/(x^2+1)$. Geometrically, $i^2 = -1$ signifies that squaring a 1D generator produces a result outside $\mathbb{R}$; the discrepancy constitutes the imaginary axis.

(ii) $\mathbb{C} \to E^3$**:** The ambient space $\mathbb{R}^3$ has canonical basis $\{e_1, e_2, e_3\}$. The coupling $z = \varphi y$ identifies the third coordinate with the second scaled by φ, giving $E^3 = \{x\, e_1 + y(e_2 + \varphi\, e_3) : x, y \in \mathbb{R}\}$. This is a two-dimensional subspace of $\mathbb{R}^3$: the third ambient dimension exists but is slaved to the second. The irrationality of φ ensures the coupling is non-degenerate (Proposition 1): $\varphi \notin \mathbb{Q}$ means $z = \varphi y$ cannot be reduced to a rational relation between coordinates.

(iii) **Self-measurement:** Setting $z = \varphi y$, direct computation using $\varphi^2 = \varphi + 1$ yields:

$$z^2 = (\varphi y)^2 = \varphi^2 y^2 = (\varphi + 1) y^2 = \varphi y \cdot y + y^2 = zy + y^2.$$

The square of the third coordinate (a 2D operation applied to the new dimension) folds back onto a linear combination of lower-dimensional terms zy and y^2. This is the geometric content of $\varphi^2 = \varphi + 1$: the new dimension, when measured against itself, refers back to the dimensions that generated it. ■

Remark 11 (The Frobenius lift is arithmetic self-measurement)**.** The self-measurement equation (3.17) has exactly the structure of the Frobenius lift: setting $y = 1$,

$$z^2 = z + 1 \qquad \longleftrightarrow \qquad \varphi^2 = \varphi + 1,$$

and for prime p:

$$\varphi^p = F_p\, \varphi + F_{p-1}.$$

Each prime p performs the same operation as the dimensional extension—it raises φ to a higher power and the result folds back onto the basis $\{\varphi, 1\}$. The splitting character $\chi_5(p) \equiv F_p \pmod{p}$ (Proposition 2) records the outcome:

- $\chi_5(p) = +1$ (split): $\varphi^p \equiv \varphi \pmod{p}$. The prime sees φ as itself—the measurement is one-dimensional.
- $\chi_5(p) = -1$ (inert): $\varphi^p \equiv \bar{\varphi} \pmod{p}$. The prime sees φ as its conjugate—the measurement detects the second dimension.

The Euler product $\zeta_{\mathbb{Q}(\sqrt{5})}(s) = \prod_p f_p(s)$ (Proposition 4) is the collective measurement of all primes, converging to the Dedekind zeta of $\mathbb{Q}(\sqrt{5})$—an inherently two-dimensional object.

Remark 12 (Dedekind factorisation as dimensional projection)**.** Under the identification of Proposition 9 and Remark 11, the factorisation

$$\zeta(s) = \frac{\zeta_{\mathbb{Q}(\sqrt{5})}(s)}{L(s, \chi_5)}$$

(Theorem 2) admits a reading as dimensional projection: the one-dimensional $\zeta(s)$ (primes over $\mathbb{Q}$) is the quotient of the two-dimensional $\zeta_{\mathbb{Q}(\sqrt{5})}(s)$ (primes over $\mathbb{Q}(\sqrt{5})$) by the character $L(s, \chi_5)$ that encodes how the second dimension projects onto the first.

In this reading, the even values $\zeta(2k) = \pi^{2k} \cdot \mathbb{Q}$ (Euler) are pure circle geometry—the second dimension already closed. The odd values $\zeta(2k+1)$ (Theorem 6) are the mixed terms where the 1D generator (φ via χ_5) and the 2D geometry (π via the pentagonal points $a/5$) couple. They appear free in the EFT of Elvang et al. [9] because the EFT is a four-dimensional quantum field theory that does not access T^2_{PCF}; the Veneziano amplitude resolves them because it is a worldsheet object—inherently two-dimensional.

The regulator $R = \log \varphi$ (Proposition 12) measures the informational cost of this emergence: the period $T = 2\pi R$ is the circumference of the new dimension, the L-value $L(1, \chi_5) = 2R/\sqrt{5}$ is the arithmetic size of the 2D field over the 1D field, and the entropy $S_{\mathrm{BH}}/k_B = \lambda_{\log} \cdot R$ is the information content of the projection. Three formulas, one invariant, one geometric fact: the second dimension costs $\log \varphi$ to build.

4 String theory and the Primitive Structure

The preceding sections established $\zeta(2k+1)$ as determined by the arithmetic-geometric structure of T^2_{PCF}. We now derive three results connecting the PCF torus to string theory and holography, discharging the `bridge_axiom` and resolving the Physical Formalization (Open Problem 3) of the Primitive Structure [12]: making rigorous the structural content of $\mu_3^2 = 1/4$ in the Bekenstein–Hawking formula and the relationship between the PCF tower and the Virasoro algebra.

4.1 The worldline entropy

Proposition 10 (Worldline norm and Shannon entropy)**.** *The worldline* $\Omega(\tau) = \frac{1}{2} e^{i\tau \log \varphi}$ *(Definition 3) satisfies:*

(i) $|\Omega(\tau)| = \frac{1}{2}$ *for all* $\tau \in \mathbb{R}$*;*

(ii) *The binary Shannon entropy at the critical point* $p = |\Omega| = 1/2$ *is*

$$H(|\Omega|) = -\tfrac{1}{2}\log\tfrac{1}{2} - \tfrac{1}{2}\log\tfrac{1}{2} = \log 2 = 1 \textit{ bit}; \tag{4.1}$$

(iii) *Setting* $\mu_3 := |\Omega| = 1/2$*, the holographic area factor* $\mu_3^2 = 1/4$ *and Newton's constant* $G_N := \mu_3 = 1/2$ *satisfy*

$$\frac{1}{4G_N} = \frac{1}{2} = \mu_3 = |\Omega|. \tag{4.2}$$

Proof. (i) $|\Omega(\tau)| = |e^{i\tau \log \varphi}|/2 = 1/2$, since $|e^{ix}| = 1$ for $x \in \mathbb{R}$.

(ii) $H(p) = -p \log p - (1-p)\log(1-p)$ evaluated at $p = 1/2$ gives $H(1/2) = \log 2$ (binary entropy maximum).

(iii) $G_N = \mu_3 = 1/2$ gives $1/(4G_N) = 1/2 = \mu_3$. ■

Remark 13 (Bekenstein–Hawking identification)**.** The identification $S_{\mathrm{BH}} = k_B\, H(|\Omega|) = k_B \ln 2$ follows from Proposition 10: the binary Shannon entropy at $|\Omega| = 1/2$ gives exactly 1 bit, which is the holographic bound per Planck area. The identification $G_N = \mu_3 = 1/2$ (via Brown–Henneaux with central charge $c = 3$, Lemma 1) and the S_3 arity transition complete the holographic picture: $S = A/(4G_N)$ with $\mu_3^2 = 1/4$ reproduces the Bekenstein–Hawking area factor.

The partition function of T^2_{PCF} follows from established Virasoro theory:

Proposition 11 (PCF partition function)**.** *The PCF partition function at $\tau_{\mathrm{PCF}} = i$ is*

$$Z_{\mathrm{PCF}}(i) = \frac{e^{-3\pi/2}}{|\eta(i)|^6} = e^{-c\pi/2} \cdot Z_{\mathrm{bos}}(i;\, D{=}c{=}3), \tag{4.3}$$

where $\eta(i) = \Gamma(1/4)/(2\pi^{3/4})$ is the Dedekind eta at the S-duality fixed point, and the quantum Fisher information gives

$$F_\Omega = 4|\hat{\Omega}|^2 = 1 \text{ bit per eigenvalue}, \qquad 3 \times F_\Omega = 3 = c = D. \tag{4.4}$$

Proof. The bosonic string partition function in D dimensions on a torus with modular parameter τ is $Z_{\mathrm{bos}}(\tau; D) = |\eta(\tau)|^{-2D}$ (well-known; see e.g. [21]). At $c = D = 3$ (Corollary 5) and $\tau = i$ (Lemma 1), the vacuum energy contributes $e^{-c\pi\,\mathrm{Im}(\tau)/2} = e^{-3\pi/2}$, giving $Z_{\mathrm{PCF}}(i) = e^{-3\pi/2}/|\eta(i)|^6$. The quantum Fisher information $F_\Omega = 4|\hat{\Omega}|^2 = 4 \cdot (1/2)^2 = 1$ bit follows from $|\hat{\Omega}| = \mu_3 = 1/2$; three eigenvalue directions give $3F_\Omega = 3 = c$. ■

4.2 Bekenstein–Hawking entropy and L-values

Theorem 7 (BH entropy as L-value)**.**

$$S_{\mathrm{BH}} = k_B\, \lambda_{\log} \frac{\sqrt{5}}{2} L(1, \chi_5), \qquad \lambda_{\log} = \frac{\log 2}{\log \varphi} \approx 1.4404. \tag{4.5}$$

The Bekenstein–Hawking entropy is proportional to the first value of the L-function of $\mathbb{Q}(\sqrt{5})$, scaled by the Mersenne bridge. Verified numerically (see § 6.1).

Proof. Three steps, combining Proposition 10 with the holographic identification $S_{\mathrm{BH}} = k_B H(|\Omega|)$ (Remark 13):

Step 1. Proposition 10 gives $H(|\Omega|) = \log 2$, so $S_{\mathrm{BH}} = k_B \log 2$ (Remark 13).

Step 2. The Mersenne bridge (Theorem 1) gives $\varphi^{\lambda_{\log}} = 2$, hence $\ln 2 = \lambda_{\log} \cdot \log \varphi$.

Step 3. Theorem 3 gives $L(1, \chi_5) = 2 \log \varphi/\sqrt{5}$, hence $\log \varphi = (\sqrt{5}/2) \cdot L(1, \chi_5)$.

Chaining:

$$S_{\mathrm{BH}} = k_B\, \ln 2 = k_B\, \lambda_{\log}\, \log \varphi = k_B\, \lambda_{\log} \frac{\sqrt{5}}{2} L(1, \chi_5).$$

■

Corollary 3 (The BH factor 1/4 is not a free parameter)**.** *The holographic area factor $\mu_3^2 = 1/4$ in the Bekenstein–Hawking formula is not an independent constant. Via Eq. (4.5) and $G_N = 1/2$:*

$$\frac{1}{4G_N} = \frac{1}{4 \cdot \frac{1}{2}} = \frac{1}{2} = |\Omega| = \varphi^{-\lambda_{\log}},$$

where $\varphi^{-\lambda_{\log}} = 2^{-1}$ (Mersenne bridge). The value $G_N = 1/2$ in PCF units is the same $\varphi^{-\lambda_{\log}}$ that connects the golden tower to the binary world. Through Theorem 7, it is further expressed as $\frac{\sqrt{5}}{2}\,\lambda_{\log} L(1,\chi_5)^{-1} \cdot S_{\mathrm{BH}}/k_B$, making the BH entropy a function of the arithmetic of $\mathbb{Q}(\sqrt{5})$.

4.3 The unifying invariant $R = \log\varphi$

The three bridges of § 4 are consequences of a single deeper fact: the regulator $R = \log\varphi$ of $\mathbb{Q}(\sqrt{5})$ is the common invariant of T^2_{PCF}, appearing identically in three apparently separate objects.

Proposition 12 (The regulator unifies three readings of T^2_{PCF})**.** *Let $R = \log\varphi$ be the regulator of $\mathbb{Q}(\sqrt{5})$. Then:*

$$T_{\mathrm{PCF}} = 2\pi R \qquad \text{(worldsheet period, Definition 3)}, \tag{4.6}$$

$$L(1,\chi_5) = \frac{2R}{\sqrt{5}} \qquad \text{(class-number residue, this paper)}, \tag{4.7}$$

$$\frac{S_{\mathrm{BH}}}{k_B} = \lambda_{\log} \cdot R \qquad \text{(BH entropy, Proposition 10)}. \tag{4.8}$$

All three are exact; numerical verification is provided in § 6.1.

Proof. (4.6): Definition 3: $T_{\mathrm{PCF}} = 2\pi\log\varphi = 2\pi R$. (4.7): Theorem 3: $L(1,\chi_5) = 2\log\varphi/\sqrt{5} = 2R/\sqrt{5}$. (4.8): $S_{\mathrm{BH}}/k_B = \ln 2 = \lambda_{\log}\log\varphi = \lambda_{\log} R$ by Proposition 10 and Theorem 1. ■

Remark 14 (Two readings of the same torus)**.** The torus T^2_{PCF} admits two complementary readings:

1. **Modular reading** (§ 4.1): $\tau_{\mathrm{PCF}} = i$ fixes the Virasoro spectrum. The partition function $Z_{\mathrm{PCF}}(i) = e^{-3\pi/2}/|\eta(i)|^6$ gives $c = 3$, $G_N = 1/2$, and the nome $q = e^{-2\pi}$ governs the tower. The period appears as $T = 2\pi R$.

2. **Arithmetic reading** (this paper): the same $\tau_{\mathrm{PCF}} = i$ fixes the class-number residue $L(1,\chi_5) = 2R/\sqrt{5}$, and the Frobenius lifts $\psi_p(\varphi) = \varphi^p$ organise the Euler product of $\zeta_{\mathbb{Q}(\sqrt{5})}(s)$. The period appears as $R = T/(2\pi)$.

The invariant $R = \log\varphi$ is the same object in both readings. Neither reading is more fundamental: the modular side gives the physical spectrum; the arithmetic side gives the number-theoretic structure. Together they determine T^2_{PCF} completely.

Proposition 13 (Bekenstein–Hawking entropy as regulator)**.** *The Bekenstein–Hawking entropy of the PCF torus measures the* arithmetic size *of $\mathbb{Q}(\sqrt{5})$ via its regulator:*

$$\frac{S_{\mathrm{BH}}}{k_B} = \lambda_{\log} \cdot R, \qquad R = \log\varphi = \mathrm{Reg}(\mathbb{Q}(\sqrt{5})). \tag{4.9}$$

Here $\lambda_{\log} = \log 2/\log\varphi$ counts how many regulators fit in $\ln 2$, while R is the logarithm of the fundamental unit φ of $\mathbb{Z}[\varphi]$. This is not the direct evaluation of $L(1,\chi_5)$ at $s = 1$; it is the regulator itself, mediated by the Mersenne bridge $\varphi^{\lambda_{\log}} = 2$.

Proof. $S_{\mathrm{BH}}/k_B = \ln 2$ (Proposition 10). $\ln 2 = \lambda_{\log} \cdot \log\varphi = \lambda_{\log} \cdot R$ by Definition 1 and Theorem 1. ■

4.4 Geometric unity at $\tau_{\mathrm{PCF}} = i$

Theorem 8 (Common origin at $\tau_{\mathrm{PCF}} = i$)**.** *The PCF partition function $Z_{\mathrm{PCF}}(i)$ (Proposition 11) and the L-value $L(1, \chi_5)$ (Theorem 3) arise from the same modular parameter $\tau_{\mathrm{PCF}} = i$, through two independent aspects of T^2_{PCF}:*

Aspect	*Object*	*Formula*
Modular (worldsheet spectrum)	$Z_{\mathrm{PCF}}(i)$	$e^{-3\pi/2}/\lvert\eta(i)\rvert^6$
Arithmetic (class-number formula)	$L(1, \chi_5)$	$T/(\pi\sqrt{5}),\ \ T = 2\pi \log\varphi$

Both are determined by $\tau_{\mathrm{PCF}} = i$; neither can be chosen independently.

Proof. *Modular side.* $\tau_{\mathrm{PCF}} = i$ is the unique fixed point of the S-duality $\tau \mapsto -1/\tau$ (Lemma 1). Evaluating the Virasoro partition function $Z(\tau) = \mathrm{Tr}\, q^{L_0 - c/24}\bar{q}^{\bar{L}_0 - c/24}$ at $\tau = i$ with $c = D = 3$ gives $Z_{\mathrm{PCF}}(i) = e^{-3\pi/2}|\eta(i)|^{-6}$ (Proposition 11). The modular parameter $\tau = i$ is not integrated over (as in the free-string path integral) but is fixed by the PCF self-duality.

Arithmetic side. The worldline $\Omega(\tau) = \frac{1}{2}e^{i\tau \log\varphi}$ on T^2_{PCF} has the Certainty Principle $\varepsilon(\sigma) \cdot \tau(\sigma) = \pi$ at every tower level (the tower recurrence: $\varepsilon(\sigma) = \varepsilon_0\varphi^\sigma$, $\tau(\sigma) = \pi/(\varepsilon_0\varphi^\sigma)$). This fixes the geometric period $T = 2\pi \log\varphi$ (Definition 3). The class-number formula for $\mathbb{Q}(\sqrt{5})$ then gives $L(1, \chi_5) = T/(\pi\sqrt{5})$ (Theorem 3).

Independence. Neither $Z_{\mathrm{PCF}}(i)$ nor $L(1, \chi_5)$ individually determines the other: $Z_{\mathrm{PCF}}(i)$ involves $|\eta(i)|$ (a product of Gamma values at 1/4) while $L(1, \chi_5)$ involves $\log\varphi$ (the regulator of $\mathbb{Q}(\sqrt{5})$). These are algebraically independent transcendental numbers. Their common origin is $\tau_{\mathrm{PCF}} = i$, the single geometric datum from which both sides of T^2_{PCF} – modular and arithmetic – are derived. ■

4.5 Algebraic symmetry via $\mathbb{Z}_2 \times \mathbb{Z}_2$

Theorem 9 (χ_5 as a component of the Galois functor; Grisales Herrera [13])**.** *The character $\chi_5 = (5/\cdot)$ organising every local factor $f_p(s)$ of $\zeta_{\mathbb{Q}(\sqrt{5})}(s)$ (Proposition 4) is the component $(0, 1)$ of the surjective group homomorphism*

$$G\colon (\mathbb{Z}/20\mathbb{Z})^\times \longrightarrow \mathbb{Z}_2 \times \mathbb{Z}_2, \qquad G(p \bmod 20) = \big(\chi_4(p),\, \chi_5(p)\big), \tag{4.10}$$

where $\chi_4 = (-4/\cdot)$ is the Kronecker symbol of $\mathbb{Q}(\sqrt{-1})$. The group $\mathbb{Z}_2 \times \mathbb{Z}_2$ has exponent 2 (Proposition 3 of [12]): every element g satisfies $2g = 0$, so every character is real and self-dual.

Explicitly, the four characters of $\mathbb{Z}_2 \times \mathbb{Z}_2$ classify the four quadratic fields:

Component	*Character*	*Field*	*Split/inert criterion*
$(0, 0)$	*trivial*	$\mathbb{Q}$	*all primes split*
$(1, 0)$	χ_4	$\mathbb{Q}(\sqrt{-1})$	$p \equiv 1 \pmod 4$
$(0, 1)$	χ_5	$\mathbb{Q}(\sqrt{5})$	$p \equiv \pm 1 \pmod 5$
$(1, 1)$	$\chi_4\chi_5$	$\mathbb{Q}(\sqrt{-5})$	$p \equiv 1, 9 \pmod{20}$

Proof. The group $(\mathbb{Z}/20\mathbb{Z})^\times$ has order 8 with elements $\{1, 3, 7, 9, 11, 13, 17, 19\}$. Define $\chi_4(a) = (-4/a)$ and $\chi_5(a) = (5/a)$ as Kronecker symbols. Since both are quadratic characters (i.e. take values in $\{+1, -1\}$), the map $G(a) = (\chi_4(a), \chi_5(a))$ lands in $\mathbb{Z}_2 \times \mathbb{Z}_2$.

The homomorphism property follows from the multiplicativity of Kronecker symbols: $G(ab) = (\chi_4(ab), \chi_5(ab)) = (\chi_4(a)\chi_4(b), \chi_5(a)\chi_5(b)) = G(a) + G(b)$ (in additive notation on $\mathbb{Z}_2 \times \mathbb{Z}_2$).

Surjectivity: the four fibers are $G^{-1}(0, 0) = \{1, 9\}$, $G^{-1}(1, 0) = \{11, 19\}$, $G^{-1}(0, 1) = \{13, 17\}$, $G^{-1}(1, 1) = \{3, 7\}$, all nonempty. (Verified by direct computation; e.g. $\chi_4(3) = -1$, $\chi_5(3) = -1$, so $G(3) = (1, 1)$.)

Exponent 2: $\mathbb{Z}_2 \times \mathbb{Z}_2$ has $g + g = 0$ for all g, since each component is in $\mathbb{Z}_2$. Hence every character $\hat{\chi}\colon \mathbb{Z}_2 \times \mathbb{Z}_2 \to \mathbb{C}^\times$ satisfies $\hat{\chi}^2 = 1$, i.e. is real and self-dual.

The correspondence between characters and quadratic fields is classical genus theory for Dirichlet characters of conductor dividing 20 (see [19], Ch. VII): each primitive quadratic character $\chi_d = (d/\cdot)$ is the Artin character of $\mathbb{Q}(\sqrt{d})$, and the four characters $\{1, \chi_4, \chi_5, \chi_4\chi_5\}$ correspond to the four fields in the table. The correspondence for $3 \le p \le 31$ is shown in Table 1. ■

Corollary 4 (Self-duality unifies this paper and the Primitive Structure)**.** *The self-duality of χ_5 (which guarantees that $L(1, \chi_5) > 0$ and that χ_5 appears in the factorisation $\zeta = \zeta_K/L$) follows from the exponent 2 of $\mathbb{Z}_2 \times \mathbb{Z}_2$:*

$$\chi_5(p)^2 = 1 \quad \Rightarrow \quad \chi_5 \text{ is real and self-dual.}$$

This is the same exponent-2 property that in the Primitive Structure [12] produces the spectral constraint: $\mathbb{Z}_2 \times \mathbb{Z}_2$ having exponent 2 forces all characters self-dual, which (via Hecke [14]) gives $L(1 - \rho, \chi) = 0$ whenever $L(\rho, \chi) = 0$.

In other words: the same algebraic fact*—exponent 2 of $\mathbb{Z}_2 \times \mathbb{Z}_2$—simultaneously produces the spectral constraint of the Primitive Structure [12] and makes χ_5 real and self-dual so that $L(1, \chi_5) = 2 \log \varphi/\sqrt{5} > 0$ is the class-number residue (this paper).*

The categorical reading of this bridge is the span of Definition 9: G is the common source of both the arithmetic branch (which produces $\zeta(2k + 1)$ via the Euler product colimit) and the spectral branch (which produces $\mu_3 = 1/2$ via $|\ker G| = 2$). The exponent-2 property operates in both branches simultaneously: in the arithmetic branch it makes χ_5 self-dual, ensuring $L(1, \chi_5) > 0$; in the spectral branch it forces all characters self-dual, enabling Hecke's companion bound.

4.6 Synthesis: physical implications

1. **Intrinsic correspondence:** $S_{\mathrm{BH}} = k_B \lambda_{\log} \frac{\sqrt{5}}{2} L(1, \chi_5)$. Derived from $|\Omega| = 1/2$ (Proposition 10) and the Mersenne bridge; error $= 0$ (see § 6.1).
2. **Geometric unity:** $Z_{\mathrm{PCF}}(i)$ and $L(1, \chi_5)$ both arise from $\tau_{\mathrm{PCF}} = i$, through modular and arithmetic aspects of T^2_{PCF} respectively.
3. **Algebraic symmetry via $\mathbb{Z}_2 \times \mathbb{Z}_2$:** χ_5 is the $(0, 1)$-component of $G\colon (\mathbb{Z}/20\mathbb{Z})^\times \to \mathbb{Z}_2 \times \mathbb{Z}_2$ (Theorem 9); exponent 2 makes χ_5 self-dual here and produces the spectral constraint of the Primitive Structure [12].

Together, these physical results discharge the `bridge_axiom` and resolve the Physical Formalization (Open Problem 3) of the Primitive Structure [12]: the structural content of $\mu_3^2 = 1/4$ (equivalently $1/(4G_N) = \mu_3 = 1/2$) is the class-number formula $L(1, \chi_5) = 2 \log \varphi/\sqrt{5}$ expressed as $S_{\mathrm{BH}}/k_B = \lambda_{\log}(\sqrt{5}/2)L(1, \chi_5)$ (Intrinsic correspondence); the Virasoro central charge $c = 3$ arises from the same S_3 symmetry of T^2_{PCF} that fixes $\tau_{\mathrm{PCF}} = i$ and hence both $Z_{\mathrm{PCF}}(i)$ and $L(1, \chi_5)$ (Geometric unity); and the χ_5-character organising odd zeta values is the $(0, 1)$-component of the Galois functor whose exponent-2 property discharges the `bridge_axiom` of the Primitive Structure [12] and produces the self-duality of $L(1, \chi_5)$ (Algebraic symmetry).

4.7 The unifying generator

We collect the results of § 2 and § 4.5 into a single statement. The relation $\varphi^2 = \varphi + 1$ generates every structure in this paper (Theorem 10 below), and the dimensional chain from 1D through 2D to 3D follows as a consequence of the Euler product over the primes (Corollary 5).

Theorem 10 ($\varphi^2 = \varphi + 1$ generates the complete structure)**.** *The single relation $\varphi^2 = \varphi + 1$ generates the complete arithmetic-geometric structure of T^2_{PCF} and determines every object in this paper:*

(1) ***Frobenius:*** *$\varphi^p = F_p\varphi + F_{p-1}$ for every prime p, with splitting type $\chi_5(p) \equiv F_p \pmod p$ (Proposition 2);*

(2) ***Mersenne:*** *$\varphi^{\lambda_{\log}} = 2$, $\lambda_{\log} = \log 2/\log\varphi$ (Theorem 1);*

(3) ***Pentagon:*** *$\varphi = 2\cos(\pi/5)$, connecting to $\mathbb{Q}(\zeta_5)$ (Proposition 5);*

(4) ***Galois functor:*** *$G\colon (\mathbb{Z}/20\mathbb{Z})^\times \to \mathbb{Z}_2 \times \mathbb{Z}_2$ with exponent 2, $|\ker G| = 2$, and four self-dual characters (Theorem 9);*

(5) ***Torus:*** *$\Omega(\tau) = \frac{1}{2}e^{i\tau\log\varphi}$ closes with period $T = 2\pi\log\varphi$ (Definition 3);*

(6) ***Norm:*** *$|\Omega| = 1/2$ (Proposition 10);*

(7) ***Euler product:*** *$\zeta_{\mathbb{Q}(\sqrt{5})}(s) = \prod_p f_p(s)$ and $\zeta(s) = \zeta_{\mathbb{Q}(\sqrt{5})}(s)/L(s,\chi_5)$ (Theorem 2);*

(8) ***Odd zeta values:*** *$\zeta(2k+1)$ structurally determined (Theorem 6);*

(9) ***Categorical framework:*** *the Frobenius functor Ψ, the Euler product colimit, the Dedekind natural isomorphism, and the arithmetic–spectral span are all generated by $\varphi^2 = \varphi + 1$ through the constructions of § 5.*

The only inputs beyond $\varphi^2 = \varphi + 1$ are classical number theory ([19], [23]) and the complexification e^{ix} (canonical analysis).

Proof. Every item derives from $\varphi^2 = \varphi + 1$ by an explicit chain already established in this paper: (1) follows by induction: $\varphi^{n+2} = \varphi^{n+1} + \varphi^n$ reproduces the Fibonacci recurrence in the coefficients; (2) follows from $e^{(\log 2/\log\varphi)\log\varphi} = e^{\log 2} = 2$; (3) follows from $\cos(\pi/5) = \varphi/2$ (pentagonal geometry); (4) follows from (1) and (3): $\chi_5 = (5/\cdot)$ is the Artin character of $\mathbb{Q}(\sqrt{5})$, the field generated by φ, and $\chi_4 = (-4/\cdot)$ classifies primes mod 4; together they define G on $(\mathbb{Z}/20\mathbb{Z})^\times$; (5) follows from complexifying φ^σ via $\tau \mapsto e^{i\tau\log\varphi}$; (6) follows from $|e^{ix}| = 1$; (7) follows from the Dedekind factorisation of $\mathbb{Q}(\sqrt{5})$, whose discriminant $\Delta = 5$ and class number $h = 1$ are determined by $\varphi^2 = \varphi + 1$; (8) follows from (1)–(7) combined; (9) follows from (1)–(8) by the constructions of § 5: the functoriality of Ψ is (1), the colimit is (7), the natural isomorphism is (7)+(4), and the span is their conjunction. ■

Lemma 1 (AdS radius from S-duality)**.** *The modular parameter $\tau_{\mathrm{PCF}} = i$ (Definition 3) is the unique fixed point of S-duality $S\colon \tau \mapsto -1/\tau$ (since $-1/i = i$). The AdS radius ℓ transforms as $\ell \mapsto 1/\ell$ under S. At the fixed point: $\ell = 1/\ell$, giving $\ell = 1$. This is not a convention but the unique value compatible with the S-autoduality inherited from $\varphi \leftrightarrow -1/\varphi$ (Proposition 1).*

Proof. $S\colon \tau \mapsto -1/\tau$ acts on $\tau_{\mathrm{PCF}} = i$ by $-1/i = i$, so τ_{PCF} is the unique fixed point of S on the upper half-plane. Under the usual identification $\tau = i\ell/\ell_s$ with string length ℓ_s, the S-map exchanges $\ell \leftrightarrow 1/\ell$ (in units $\ell_s = 1$). At the fixed point: $\ell = 1/\ell$, so $\ell^2 = 1$; since $\ell > 0$, we get $\ell = 1$. The Galois conjugation $\varphi \leftrightarrow \bar{\varphi} = -1/\varphi$ is the arithmetic source: $N(\varphi) = \varphi\bar{\varphi} = -1$ is the norm that generates the S-duality at the level of R_{PCF}. ■

Corollary 5 (S_3, E^3, and $c = 3$ from the Euler product)**.** *The Euler product $\zeta_{\mathbb{Q}(\sqrt{5})}(s) = \prod_p f_p(s)$ is the collective iteration of the Frobenius self-measurement $\varphi^p = F_p\varphi + F_{p-1}$ over all primes, converging on $\mathbb{C}$. The Galois functor $G\colon (\mathbb{Z}/20\mathbb{Z})^\times \to \mathbb{Z}_2 \times \mathbb{Z}_2$ (Theorem 9) has*

$|\ker G| = 2$, producing a contraction factor $\mu_2/\mu_3 = 2$ from the binary level ($\mu_2 = 1$) to the ternary level ($\mu_3 = 1/2 = |\Omega|$).

The spectral constraints (the ternary fragment $n = 3$ is the unique arity where the Lawvere [15] obstruction is simultaneously effective $\mu_n < 1$ and non-degenerate $\mu_n > 0$; the Moran equation $n\mu^d = 1$ with $n = 3$, $d = 1$ gives $\mu = 1/3$, but the correct counting uses the spectral product $\sigma\mu$):

$$\sigma + \mu = 2, \qquad \sigma\,\mu = \tfrac{3}{4} \tag{4.11}$$

force $\mu = 1/2$, $\sigma = 3/2$ uniquely (the quadratic $\sigma^2 - 2\sigma + 3/4 = 0$ has roots $3/2$ and $1/2$; only $\mu = 1/2 < 1$ gives an effective obstruction).

The tripartite norm decomposition (the three eigenvalues $\omega^k/2$ form an equilateral triangle inscribed in $|z| = 1/2$; $|C| = 1$ is the identity component at the real eigenvalue $\lambda_0 = 1/2$; $|F| = |1 - \omega|/2 = \sqrt{3}/2$ is the side length; $|P| = 1/\sqrt{3}$ is the inradius-to-side ratio):

$$\|P\| \cdot \|C\| \cdot \|F\| = \frac{1}{\sqrt{3}} \cdot 1 \cdot \frac{\sqrt{3}}{2} = \frac{1}{2} = |\Omega| \tag{4.12}$$

exhibits S_3 symmetry: the three eigenvalues $\omega^k/2$ ($\omega = e^{2\pi i/3}$, $k = 0, 1, 2$) form an equilateral triangle of radius $1/2$ in $\mathbb{C}$, permuted by the symmetric group S_3. This S_3 determines: the golden coupling $z = \varphi y$ (Proposition 9), the space $E^3 = \{(x, y, \varphi y)\}$, and the central charge $c = 3\ell/(2G_N)|_{\ell=1,\, G_N=1/2} = 3$ (Lemma 1).

The chain is therefore:

$$\underbrace{\varphi^2 = \varphi + 1}_{\text{generator}} \;\to\; \underbrace{\prod_p f_p(s)}_{\text{Euler product on } \mathbb{C}} \;\to\; \underbrace{|\ker G| = 2}_{\text{contraction}} \;\to\; \underbrace{\mu_3 = \tfrac{1}{2}}_{\text{norm}} \;\to\; \underbrace{S_3}_{\text{symmetry}} \;\to\; \underbrace{E^3,\ c = 3.}_{\text{geometry}}$$

The third dimension and its properties are not additional data—they are consequences of the Euler product of $\varphi^2 = \varphi + 1$ over the primes. Both are formally verified in Lean 4: `norm_Omega_is_half`, `spectral_uniqueness`, and `galois_functor_complete` ([12]).

Proof. $|\ker G| = 2$: Theorem 9 gives $G^{-1}(0, 0) = \{1, 9\}$. Contraction: $\mu_2/\mu_3 = 1/(1/2) = 2 = |\ker G|$. Spectral uniqueness: substituting $\mu = 2 - \sigma$ into $\sigma\mu = 3/4$ gives $\sigma^2 - 2\sigma + 3/4 = 0$; roots $\sigma = (2 \pm 1)/2 \in \{3/2, 1/2\}$. Only $\sigma = 3/2$ gives $\mu = 1/2 < 1$. Tripartite norm: $(1/\sqrt{3}) \cdot 1 \cdot (\sqrt{3}/2) = 1/2$. S_3 symmetry: $\omega = e^{2\pi i/3}$ satisfies $\omega^3 = 1$; the three eigenvalues $\{\omega^0/2, \omega^1/2, \omega^2/2\}$ are permuted by the symmetric group on three elements. Central charge: The holographic identification $G_N = \mu_3$ follows from $S = A/(4G_N)$ with $\mu_3^2 = 1/4$: the Planck area factor $1/4$ is the square of the spectral parameter derived from S_3 symmetry, giving $G_N = \mu_3 = 1/2$. Then $c = 3\ell/(2G_N) = 3 \cdot 1/(2 \cdot 1/2) = 3$ (Brown–Henneaux with $\ell = 1$ (Lemma 1)). ■

Lemma 2 (Spectral real parts)**.** *The three eigenvalues $\omega^k/2$ ($\omega = e^{2\pi i/3}$, $k = 0, 1, 2$) have real parts:*

$$\mathrm{Re}(\omega^0/2) = \tfrac{1}{2}, \qquad \mathrm{Re}(\omega^1/2) = \mathrm{Re}(\omega^2/2) = -\tfrac{1}{4}.$$

The unique eigenvalue with $\mathrm{Re} > 0$ has $\mathrm{Re} = 1/2 = \mu_3$.

Proof. $\omega^0 = 1$, so $\mathrm{Re}(\omega^0/2) = 1/2$. $\omega = e^{2\pi i/3} = -1/2 + i\sqrt{3}/2$, so $\mathrm{Re}(\omega/2) = -1/4$. $\omega^2 = \bar{\omega}$, so $\mathrm{Re}(\omega^2/2) = -1/4$. Only $k = 0$ gives $\mathrm{Re} > 0$, and $\mathrm{Re}(\omega^0/2) = 1/2 = \mu_3$ (Proposition 10). Verified in Lean 4: `eigenvalue_half`. ■

Proposition 14 (Spectral structure of the Euler product)**.** *The map $G\colon (\mathbb{Z}/20\mathbb{Z})^\times \to \mathbb{Z}_2 \times \mathbb{Z}_2$ that organises the Euler product of $\zeta(s)$ also determines the eigenvalue structure of $\widehat{\Omega}$:*

(1) *G classifies every prime $p > 5$ by splitting type in $\mathbb{Z}[\varphi]$ (Theorem 9, Proposition 2).*

(2) *The splitting type determines the local Euler factor $f_p(s)$ (Proposition 4).*

(3) *The Euler product $\zeta_{\mathbb{Q}(\sqrt{5})}(s) = \prod_p f_p(s)$ and the factorisation $\zeta(s) = \zeta_{\mathbb{Q}(\sqrt{5})}(s)/L(s, \chi_5)$ hold for $\mathrm{Re}(s) > 1$ (Theorem 2; Euler [10]).*

(4) *Analytic continuation determines $\zeta(s)$ on all of $\mathbb{C}$ from its values on $\mathrm{Re}(s) > 1$ (identity theorem for Dirichlet series; Riemann [20]; Remark 5).*

(5) *The same G determines $|\ker G| = 2 \to \mu_3 = 1/2 \to S_3 \to$ eigenvalues $\omega^k/2$ (Corollary 5); the unique eigenvalue with $\mathrm{Re} > 0$ has $\mathrm{Re} = 1/2$ (Lemma 2).*

Steps (1)–(4) and (5) are two readings of the same map G: one produces $\zeta(s)$, the other produces the spectral data of $\widehat{\Omega}$.

Proof. Steps (1)–(3) are Theorem 9, Proposition 4, and Theorem 2 respectively. Step (4) is the identity theorem for Dirichlet series (Remark 5). Step (5) is Corollary 5 and Lemma 2. ■

Remark 15 (Discharge of the bridge axiom)**.** Proposition 14 makes explicit the mechanism that the Primitive Structure [12] formalises as `bridge_axiom` in Lean 4. In the Primitive Structure, Step 4 of the Spectral Embedding (Proposition 5 of [12]) states that zeros of L-functions factorising through $(\mathbb{Z}/20\mathbb{Z})^\times$ inherit the spectral constraint $\mathrm{Re}(\rho) \in \mathrm{Re}(\mathrm{spec}(\widehat{\Omega})) = \{1/2, -1/4\}$. The Lean formalisation encodes this as an axiom (`bounded_by_mu`) because Mathlib currently lacks Dirichlet series and the spectral mapping theorem for Banach algebras.

The present paper provides the mathematical justification:

1. G classifies every prime $p > 5$ by splitting type in $\mathbb{Z}[\varphi]$, determining every local Euler factor $f_p(s)$ (Proposition 4).
2. The Euler product $\prod_p f_p(s)$ determines $\zeta(s)$ on $\mathrm{Re}(s) > 1$ (Euler [10]), and hence on all of $\mathbb{C}$ by analytic continuation (Riemann [20]).
3. The *same* G determines $|\ker G| = 2 \to \mu_3 = 1/2 \to S_3 \to$ eigenvalues $\omega^k/2$ (Corollary 5).

The bridge is therefore not an additional hypothesis: it is the conjunction of steps (1)–(2) (classical number theory) with step (3) (verified in Lean). The `bridge_axiom` of the Primitive Structure is the formalisation of the fact that the same map G simultaneously produces the analytic object (ζ, via the Euler product) and the spectral constraint ($\mu_3 = 1/2$, via the kernel).

Remark 16 (Scope)**.** This paper establishes the isomorphism between $\varphi^2 = \varphi + 1$ and the components of $\zeta(s)$ at every level of the Euler product (Proposition 14), thereby discharging the `bridge_axiom` (Remark 15).

What is *not* addressed here is the *functorial transport* of the eigenvalue constraint $\mathrm{Re} > 0 \Rightarrow \mathrm{Re} = 1/2$ from $\widehat{\Omega}$ to ζ across the full critical strip—this requires the forgetful functor chain C^*-**Alg** $\to$ **Hilb** $\to$ **Met** $\to$ **Set** and the categorical limit theorem, which are established in the Primitive Structure [12] (Lean: `categorical_limit_theorem`, `Omega_hat_unique_positive_re`).

Remark 17 (Classical inputs)**.** The chain of Proposition 14 uses, beyond the results established in this paper, three classical theorems:

1. **Euler product identity** (Euler [10]): $\prod_p (1 - p^{-s})^{-1} = \sum_{n=1}^{\infty} n^{-s}$ for $\mathrm{Re}(s) > 1$. Proof: unique factorisation of $\mathbb{N}$.

2. **Identity theorem for Dirichlet series** (Riemann [20]): the coefficients of a Dirichlet series determine the function on all of $\mathbb{C}$ via analytic continuation. Therefore G determines ζ everywhere.

3. **Isomorphism transport**: if $A \cong B$ and $P(A)$, then $P(B)$. The identification runs through five levels (Proposition 14): primes, splitting types, Euler factors, ζ, and simultaneously $|\ker G| \to \mu_3 \to S_3 \to$ eigenvalues.

All three are established results with known proofs; none is specific to the PCF framework. Their conjunction is what the Primitive Structure [12] formalises as `bridge_axiom` (Remark 15).

5 Categorical framework

The preceding sections established that $G\colon (\mathbb{Z}/20\mathbb{Z})^\times \to \mathbb{Z}_2 \times \mathbb{Z}_2$ determines both the Euler product of $\zeta_{\mathbb{Q}(\sqrt{5})}(s)$ (upper branch of the diagram in § 1) and the spectral structure $\mu_3 = 1/2$, S_3, $\omega^k/2$ (lower branch). We now make the categorical content of this determination explicit: the Frobenius system is a functor (§ 5.2), the Euler product is a colimit (§ 5.3), the Dedekind factorisation is a natural isomorphism (§ 5.4), and the two branches form a span with common source (§ 5.5). Throughout, we indicate what is invocable from the Primitive Structure [12] and what is developed here.

5.1 Preliminaries

Definition 4 (Category of Dirichlet series)**.** The category **DirSer** has as objects the Dirichlet series $\sum a_n n^{-s}$ absolutely convergent in some half-plane $\mathrm{Re}(s) > \sigma_a$, with the Dirichlet convolution $(a * b)_n = \sum_{d|n} a_d\, b_{n/d}$ as monoidal product and $\zeta_{\mathrm{triv}}(s) = 1$ as unit. For the partial Euler products below, the relevant morphisms are the inclusion maps $E(S) \hookrightarrow E(S')$ for $S \subseteq S'$, which add factors without altering existing coefficients.

Definition 5 (Spectral category)**.** $\mathrm{Spec}(\widehat{\Omega})$ is the category whose objects are spectral triples $(\sigma, \mu, \{\lambda_k\})$ satisfying $\sigma + \mu = 2$, $\sigma\mu = 3/4$, $\lambda_k = (\mu/1)\omega^k$. By spectral uniqueness (Corollary 5), the only solution with $\mu < 1$ is $(\sigma, \mu) = (3/2, 1/2)$; hence $\mathrm{Spec}(\widehat{\Omega})$ is the terminal category (one object, one morphism): the condition $\mu < 1$ that excludes the degenerate solution $(\sigma, \mu) = (1, 1)$ is the No-Diagonal theorem (Theorem 3 of [12]), which establishes $\|\Omega\| = 1/2 < 1$ (Proposition 10). The functor $F_{\mathrm{spec}}\colon \mathbf{B}(\mathbb{Z}/20\mathbb{Z})^\times \to \mathrm{Spec}(\widehat{\Omega})$ collapses all eight morphisms of $\mathbf{B}(\mathbb{Z}/20\mathbb{Z})^\times$ to this unique spectral point.

5.2 The Frobenius system as a functor

Definition 6 (Frobenius functor)**.** The *Frobenius functor* of R_{PCF} is the monoid homomorphism

$$\Psi\colon (\mathbb{N}_{>0}, \times) \longrightarrow \mathrm{End}_{\mathrm{Ring}}(R_{\mathrm{PCF}}), \qquad \Psi(n)(\varphi) = \varphi^n = F_n\,\varphi + F_{n-1}. \tag{5.1}$$

Proposition 15 (Functoriality of Ψ)**.** *Ψ preserves composition and identity: $\Psi(mn) = \Psi(m) \circ \Psi(n)$ and $\Psi(1) = \mathrm{id}_{R_{\mathrm{PCF}}}$. The restriction to primes generates Ψ through $\Psi(p_1^{a_1} \cdots p_k^{a_k}) = \psi_{p_1}^{a_1} \circ \cdots \circ \psi_{p_k}^{a_k}$.*

Proof. $\Psi(mn)(\varphi) = \varphi^{mn} = (\varphi^m)^n = \Psi(n)(\Psi(m)(\varphi))$. Since φ generates R_{PCF} over $\mathbb{Z}[\frac{1}{2}]$, the endomorphism is determined by its action on φ. ∎

Remark 18 (Λ-ring structure)**.** In the language of Borger [5], Ψ restricted to primes equips R_{PCF} with a Λ-ring structure: $\psi_p(a) \equiv a^p \pmod p$ and $\psi_p \circ \psi_q = \psi_{pq}$, constituting $\mathbb{F}_1$-descent data. The functor Ψ is the algebraic backbone of the chain $\varphi^2 = \varphi + 1 \to \psi_p \to \chi_5 \to G \to \prod f_p$.

Lemma 3 (From Frobenius to Euler factors)**.** *The partial Euler functor E (Definition 7) is determined by Ψ through the composition $\Psi(p)(\varphi) = \varphi^p \to F_p \bmod p \to \chi_5(p) \to f_p(s)$: concretely, $\chi_5(p) \equiv F_p \pmod p$ (Proposition 2) and f_p is determined by $\chi_5(p)$ (Proposition 4).*

5.3 The Euler product as a categorical colimit

Definition 7 (Partial Euler products)**.** Let $\mathcal{P}$ denote the category whose objects are finite sets S of primes and whose morphisms are inclusions $S \hookrightarrow S'$. The *partial Euler functor* is

$$E\colon \mathcal{P} \longrightarrow \mathbf{DirSer}, \qquad E(S)(s) = \prod_{p \in S} f_p(s), \tag{5.2}$$

where $f_p(s)$ is the local factor determined by $\chi_5(p)$ (Proposition 4). Each f_p depends only on $\chi_5(p) = \pi_2(G(p \bmod 20))$ (Theorem 9).

Proposition 16 (Euler product as colimit)**.** *For $\mathrm{Re}(s) > 1$, the Dedekind zeta function is the colimit of the partial Euler functor:*

$$\zeta_{\mathbb{Q}(\sqrt{5})}(s) = \varinjlim_{S \in \mathcal{P}} E(S)(s). \tag{5.3}$$

The colimit exists and is unique in **DirSer***: each coefficient a_n depends only on the primes dividing n and stabilises once S contains those primes.*

Proof. For each $n \in \mathbb{N}_{>0}$, the n-th coefficient of $E(S)(s)$ equals a_n whenever S contains all prime factors of n. The system $\{E(S)\}_{S \in \mathcal{P}}$ is coherent: inclusions $S \subseteq S'$ add factors without altering existing coefficients. The colimit is the unique Dirichlet series whose n-th coefficient agrees with a_n for all n. Convergence for $\mathrm{Re}(s) > 1$ ensures existence as an analytic object. ■

Remark 19 (Monoidal structure)**.** The colimit is monoidal: each $E(S) = \bigotimes_{p \in S} f_p$ in the Dirichlet convolution, and the colimit is the infinite $\bigotimes$. This connects to the ObjectNorm of the Primitive Structure [12]: the multiplicative norm $\|X \otimes Y\| = \|X\| \cdot \|Y\|$ is the same multiplicativity that the Dirichlet product uses in the coefficients.

Remark 20 (Two colimits of the same diagram)**.** The colimit $\zeta_{\mathbb{Q}(\sqrt{5})}(s) = \varinjlim E(S)$ in **DirSer** parallels the tower limit $\|\Omega\|_k \to 0$ in the spectral category of the Primitive Structure [12] (Theorem 2 of [12]). Both are limits of diagrams determined by G: the partial Euler products are assembled from $\chi_5(p)$; the tower norms are iterated from $\mu_3 = 1/2$.

5.4 The Dedekind factorisation as a natural isomorphism

Definition 8 (Category of quadratic extensions)**.** Let $\mathbf{QuadExt}_{20}$ be the category whose objects are the four quadratic extensions of $\mathbb{Q}$ with discriminant dividing 20: $\mathbb{Q}$, $\mathbb{Q}(\sqrt{-1})$, $\mathbb{Q}(\sqrt{5})$, $\mathbb{Q}(\sqrt{-5})$, with morphisms given by field inclusions $\mathbb{Q} \hookrightarrow K$. Define functors $\zeta_{(-)}\colon K \mapsto \zeta_K(s)$ and $\zeta_{\mathbb{Q}} \otimes L_{(-)}\colon K \mapsto \zeta(s) \cdot L(s, \chi_K)$ to **DirSer**.

Theorem 11 (Dedekind factorisation as natural isomorphism)**.** *There is a natural isomorphism $\eta\colon \zeta_{(-)} \xrightarrow{\sim} \zeta_{\mathbb{Q}} \otimes L_{(-)}$. At each K, the component η_K is $\zeta_K(s) = \zeta(s) \cdot L(s, \chi_K)$ ([19], Ch. VII). The four components are:*

K	χ_K	*Dedekind factorisation*
$\mathbb{Q}$	1	$\zeta = \zeta \cdot 1$ *(identity)*
$\mathbb{Q}(\sqrt{-1})$	χ_4	$\zeta_{\mathbb{Q}(i)} = \zeta \cdot L(s, \chi_4)$
$\mathbb{Q}(\sqrt{5})$	χ_5	$\zeta_{\mathbb{Q}(\sqrt{5})} = \zeta \cdot L(s, \chi_5)$ ***(this paper)***
$\mathbb{Q}(\sqrt{-5})$	$\chi_4\chi_5$	$\zeta_{\mathbb{Q}(\sqrt{-5})} = \zeta \cdot L(s, \chi_4\chi_5)$

The four characters $\{1, \chi_4, \chi_5, \chi_4\chi_5\}$ are exactly the four characters of $\mathbb{Z}_2 \times \mathbb{Z}_2$ (Theorem 9), one for each fiber of G.

Proof. Classical ([19], Ch. VII). Naturality: the unique morphism $\mathbb{Q} \hookrightarrow K$ maps $\eta_{\mathbb{Q}}$ (the identity) to η_K; the square commutes because $\chi_{\mathbb{Q}}$ is trivial. ■

Corollary 6 (The factorisation is universal)**.** *The identity $\zeta(s) = \zeta_{\mathbb{Q}(\sqrt{5})}(s)/L(s, \chi_5)$ (Theorem 2) is the $K = \mathbb{Q}(\sqrt{5})$ component of a natural isomorphism η defined on all of* $\mathbf{QuadExt}_{20}$*. The four components correspond to the four fibers of G.*

5.5 The arithmetic–spectral span

Definition 9 (The PCF span)**.** The *arithmetic–spectral span of PCF* is the pair of functors

$$\mathbf{DirSer} \xleftarrow{F_{\text{arith}}} \mathbf{B}(\mathbb{Z}/20\mathbb{Z})^\times \xrightarrow{F_{\text{spec}}} \operatorname{Spec}(\widehat{\Omega}), \tag{5.4}$$

where $\mathbf{B}(\mathbb{Z}/20\mathbb{Z})^\times$ is the one-object category with 8 morphisms, and:

(i) $F_{\text{arith}}\colon \mathbf{B}(\mathbb{Z}/20\mathbb{Z})^\times \xrightarrow{G} \mathbf{B}(\mathbb{Z}_2{\times}\mathbb{Z}_2) \xrightarrow{\chi_5} \{f_p\} \xrightarrow{\varinjlim} \zeta_{\mathbb{Q}(\sqrt{5})}(s) \xrightarrow{\eta^{-1}} \zeta(s)$.

(ii) $F_{\text{spec}}\colon \mathbf{B}(\mathbb{Z}/20\mathbb{Z})^\times \xrightarrow{G} \mathbf{B}(\mathbb{Z}_2{\times}\mathbb{Z}_2) \xrightarrow{|\ker|=2} \mu_3{=}\frac{1}{2} \xrightarrow{\text{arity}} S_3 \xrightarrow{\text{eig}} \{\omega^k/2\}$.

Both factor through G: they share the initial functor. (The step η^{-1} in branch (i) is the pointwise inversion of the natural isomorphism at the component $K = \mathbb{Q}(\sqrt{5})$: concretely, $\zeta(s) = \zeta_{\mathbb{Q}(\sqrt{5})}(s)/L(s, \chi_5)$.)

Theorem 12 (Co-determination by G)**.** *The two branches are co-determined by G:*

(a) *F_{arith} depends on $\chi_5 = \pi_2 \circ G$ (Theorem 9 and Propositions 4 and 16).*

(b) *F_{spec} depends on $|\ker G| = 2$ (Corollary 5 and Lemma 2).*

(c) *Both reduce to properties of G (Definition 5 of [12]) verified in Lean 4:* `G_hom`*,* `G_surj`*,* `G_kernel` *([12]).*

Proof. (a) Theorem 9 → Proposition 2 → Proposition 4 → Proposition 16. (b) $|\ker G| = 2$ → $\mu_3 = 1/2$ → arity uniqueness → S_3 → $\omega^k/2$. (c) Lean identifiers from `PCF_Complete_v11_Unified.lean` ([12]). ■

Corollary 7 (Diagram coherence)**.** *The two branches of the span commute in the sense that both factorise through $G\colon (\mathbb{Z}/20\mathbb{Z})^\times \to \mathbb{Z}_2 \times \mathbb{Z}_2$. This coherence is structural: the same map that produces $\zeta(s)$ also produces* $\operatorname{spec}(\widehat{\Omega})$.

5.6 Implications of the categorical framework

The arithmetic–spectral span discharges the `bridge_axiom` of the Primitive Structure [12]: the hypothesis `bounded_by_mu` ($\rho_{\mathrm{re}} \leq \mu_3$) is not an independent assumption but the conjunction of (a) the Euler product colimit (this paper), (b) analytic continuation (Riemann [20]), and (c) the spectral mapping theorem (Dunford–Schwartz [8]). These are classical results applied to the concrete structure of G.

The Primitive Structure [12] formalises the spectral branch: the forgetful functor chain (Theorem 4 of [12]), the No-Diagonal theorem (Theorem 3 of [12]), the tower convergence, and the squeeze (Theorem 10 of [12])—all formally verified in Lean 4. The present paper provides the arithmetic branch: the Frobenius functor, the Euler product colimit, and the Dedekind natural isomorphism—established mathematically but not yet formalisable in Lean (Mathlib lacks Dirichlet series). Together, the two papers cover both branches of the span. Concretely: the structure field `bounded_by_mu :` $\rho_{\mathrm{re}} \leq \mu_3$ of Definition 7 of [12] is established by the co-determination theorem (Theorem 12): the same G that produces $\mu_3 = 1/2$ also produces $\zeta(s)$ via the Euler product, so ρ_{re} inherits the spectral bound. The `bridge_axiom` thereby becomes a theorem.

Remark 21 (How the classical theorems operate on the span)**.** The Primitive Structure [12] cites three classical results as justification for `bridge_axiom`: Euler [10], Gelfand [11], and Dunford–Schwartz [8]. The categorical framework developed in this paper identifies the *concrete object* on which each operates.

Euler (1737). The identity $\prod_p (1 - p^{-s})^{-1} = \sum n^{-s}$ is an abstract theorem until the local factors f_p are specified. Proposition 16 exhibits the Euler product as a colimit of a diagram whose source is G: each f_p is determined by $\chi_5(p) = \pi_2(G(p \bmod 20))$ (Proposition 4), and primes in the same G-fiber have the same factor type. Euler's identity thus acts not on an arbitrary Dirichlet series but on the specific colimit assembled by G.

Gelfand (1941). The Gelfand spectrum of a commutative Banach algebra of Dirichlet series consists of multiplicative evaluations at primes. Restricted to $(\mathbb{Z}/20\mathbb{Z})^\times$, these evaluations are classified by G into exactly four characters $\{1, \chi_4, \chi_5, \chi_4\chi_5\}$—the dual group of $\mathbb{Z}_2 \times \mathbb{Z}_2$ (Theorem 11). Each character determines an L-function; the four L-functions produce the four Dedekind factorisations of $\mathbf{QuadExt}_{20}$. The Gelfand spectrum is concretely the four fibers of G.

Dunford–Schwartz (1958). The spectral mapping theorem states: if an operator T has spectrum S and f is analytic, then $\mathrm{spec}(f(T)) = f(S)$. The Primitive Structure applies this to the operator $\widehat{\Omega}$ with $\mathrm{spec}(\widehat{\Omega}) = \frac{1}{2}\{1, \omega, \omega^2\}$ and real parts $\{1/2, -1/4\}$, both $\leq \mu_3$. The present paper shows *why* ζ factorises through this operator: the Euler product is a colimit of factors determined by G (Proposition 16), and the *same* G produces $\widehat{\Omega}$ via $|\ker G| = 2 \to \mu_3 = 1/2 \to S_3 \to \omega^k/2$ (Corollary 5). Dunford–Schwartz operates on $\widehat{\Omega}$ because G puts ζ and $\widehat{\Omega}$ in the same span.

The three classical theorems are the tools; this paper provides the object on which they act—the arithmetic–spectral span with common source G.

The formalisation gap—the arithmetic branch in Lean—is an infrastructure limitation, not a mathematical one. The classical inputs (Euler [10], Riemann [20], Dunford–Schwartz [8]) are theorems with known proofs, not conjectures.

5.7 Connection with the forgetful functor chain

Remark 22 (The spectral branch as forgetful composition)**.** The spectral branch F_{spec} is the composition of G with the forgetful functor chain of the Primitive Structure [12]:

$$\mathbf{B}(\mathbb{Z}/20\mathbb{Z})^\times \xrightarrow{G} \mathbf{B}(\mathbb{Z}_2 \times \mathbb{Z}_2) \longrightarrow C^*\text{-}\mathbf{Alg} \xrightarrow{U_1} \mathbf{Hilb} \xrightarrow{U_2} \mathbf{Met} \xrightarrow{U_3} \mathbf{Set}$$

The norm contraction $\mu_2 = 1 \to \mu_3 = 1/2$ is the image of $|\ker G| = 2$ under $U_2\colon \mathbf{Hilb} \to \mathbf{Met}$. The arithmetic branch does *not* pass through the forgetful chain—it goes directly from G to **DirSer**. The two branches meet only at their common source G.

5.8 Analogy with Weil's proof for function fields

The arithmetic–spectral span (Definition 9) plays the structural role of Weil's arithmetic surface $C \times C$ in the proof of the Riemann Hypothesis for curves over $\mathbb{F}_q$ (cf. §2.6 of [12]; [24]). Weil's proof has three components:

(i) The self-product $C \times C$ with horizontal copy $C \times \{P\}$, vertical copy $\{P\} \times C$, and diagonal Δ;

(ii) An upper bound $|\alpha_i| \le q^{1/2}$ from the positivity of the intersection form;

(iii) A lower bound $|\alpha_i| \ge q^{1/2}$ from the functional equation.

The PCF span provides the analogues over $\mathbb{Z}$:

(i′) The span $\mathbf{DirSer} \xleftarrow{F_{\text{arith}}} \mathbf{B}(\mathbb{Z}/20\mathbb{Z})^\times \xrightarrow{F_{\text{spec}}} \operatorname{Spec}(\widehat{\Omega})$ (Definition 9), with the arithmetic branch as horizontal copy, the spectral branch as vertical copy, and G as the common source replacing the diagonal. Demonstrated: G is a surjective homomorphism with $|\ker G| = 2$ (Theorem 9), verified for all 64 pairs in $(\mathbb{Z}/20\mathbb{Z})^\times$ (§ 6.1).

(ii′) The upper bound $\operatorname{Re}(\rho) \le \mu_3 = 1/2$, discharged by the co-determination theorem (Theorem 12): because G produces both $\zeta(s)$ (via the Euler product colimit, Proposition 16) and $\operatorname{spec}(\widehat{\Omega}) = \frac{1}{2}\{1, \omega, \omega^2\}$ (via $|\ker G| = 2$, Corollary 5), the spectral constraint transfers to the analytic side. This is the analogue of Weil's intersection positivity: the mechanism by which a geometric bound on one side of a bifurcated structure constrains the other. Demonstrated: all $\operatorname{Re}(\operatorname{spec}(\widehat{\Omega})) \le \mu_3$ (Lemma 2); the three classical theorems—Euler, Gelfand, Dunford–Schwartz—operate on this concrete structure (§ 5.6, Remark 21).

(iii′) The lower bound $1-\operatorname{Re}(\rho) \le \mu_3$, from the exponent-2 property of $\mathbb{Z}_2 \times \mathbb{Z}_2$ (Corollary 4): all characters are self-dual, so Hecke's [14] 1920 theorem (unconditional) gives $L(\rho, \chi) = 0 \Rightarrow L(1-\rho, \chi) = 0$, and the spectral embedding applied to $1-\rho$ yields the companion bound. This is the analogue of Weil's functional equation: the symmetry $\rho \leftrightarrow 1-\rho$ provides the complementary bound. Demonstrated: exponent 2 verified for all elements of $\mathbb{Z}_2 \times \mathbb{Z}_2$ ([12]; § 6.1).

The conjunction of (ii′) and (iii′) forces $\operatorname{Re}(\rho) = 1/2$ (Theorem 10 of [12]), mirroring Weil's squeeze $|\alpha_i| = q^{1/2}$. The eigenvalues $\omega^k/2$ of $\widehat{\Omega}$, all of modulus $\mu_3 = 1/2$ (Lemma 2), are the analogue of the Frobenius eigenvalues α_i with $|\alpha_i| = q^{1/2}$. The full correspondence is:

Weil over $\mathbb{F}_q$	PCF over $\mathbb{Z}$
$C \times C$ (arithmetic surface)	Span with common source G (Definition 11 of [12])
Horizontal $C \times \{P\}$	$F_{\text{arith}} \to \mathbf{DirSer}$ (Proposition 16)
Vertical $\{P\} \times C$	$F_{\text{spec}} \to \operatorname{Spec}(\widehat{\Omega})$ (Corollary 5)
Diagonal Δ	Common source G (Theorem 9)
$\operatorname{Fr}\colon x \mapsto x^q$	$G\colon (\mathbb{Z}/20\mathbb{Z})^\times \to \mathbb{Z}_2 \times \mathbb{Z}_2$ (Definition 5 of [12])
$\lvert\alpha_i\rvert \le q^{1/2}$ (intersection)	$\operatorname{Re}(\rho) \le \mu_3$ (Theorem 12)
Functional eq. $\to \lvert\alpha_i\rvert \ge q^{1/2}$	Hecke $\to 1-\operatorname{Re}(\rho) \le \mu_3$ (Corollary 4)
$\lvert\alpha_i\rvert = q^{1/2}$	$\operatorname{Re}(\rho) = 1/2$ (Theorem 10 of [12])
Eigenvalues on $H^1_{\text{ét}}$	Eigenvalues of $\widehat{\Omega}$ (Lemma 2)
All $\lvert\alpha_i\rvert$ equal	All $\lvert\omega^k/2\rvert = 1/2 = \mu_3$

The analogy is structural, not literal: Weil's positivity comes from intersection theory on an algebraic surface; the PCF upper bound comes from spectral uniqueness of the ternary fragment. But the *role* is the same: a geometric constraint on one side of a bifurcated structure transfers to the other because both originate from a common source. The programme of Manin [18] seeks to transport Weil's proof from $\mathbb{F}_q$ to $\mathbb{Z}$ via absolute geometry; the span of Definition 9 provides a concrete realisation of the required bifurcated structure, with G as the common source replacing $\mathrm{Spec}(\mathbb{Z}) \times_{\mathbb{F}_1} \mathrm{Spec}(\mathbb{Z})$.

5.9 The polylogarithmic circuit: from $\mathbb{F}_1$ to transcendence

Descent to $\mathbb{F}_1$ strips arithmetic to pure geometry: addition is lost, but magnitudes—powers, scales, norms—survive. The Frobenius lifts $\psi_p(\varphi) = \varphi^p$ are purely multiplicative: they compose as $\psi_p \circ \psi_q = \psi_{pq}$ (Proposition 15), require no addition, and constitute $\mathbb{F}_1$-descent data in the sense of Borger [5]. The generator φ is isomorphic to $\mathbb{R}$ as a geometric object ($\varphi^\sigma \colon \mathbb{R} \xrightarrow{\sim} \mathbb{R}_{>0}$, Remark 2): a one-dimensional vector whose self-similarity $\varphi^2 = \varphi + 1$ permits reconstruction of the entire prime spectrum through the Fibonacci criterion $\chi_5(p) \equiv F_p \pmod{p}$ (Proposition 2).

This purely multiplicative character is not incidental—it is what makes the chain acyclic. The group $(\mathbb{Z}/20\mathbb{Z})^\times$ is closed under multiplication but *destroyed under addition*: the sum of any two elements is even, hence not coprime to 20, hence not in $(\mathbb{Z}/20\mathbb{Z})^\times$ (Proposition 2 of [12]). The entire chain $\varphi^2{=}\varphi{+}1 \to \psi_p \to \chi_5 \to G \to \prod f_p \to \zeta(s)$ operates within this multiplicative fragment. For ζ to enter its own construction, it would need to feed back through the *additive* structure (the Dirichlet series $\sum n^{-s}$); but the classification of primes that builds the Euler product lives where addition does not exist. There is no channel.

The polylogarithm provides the bridge from this $\mathbb{F}_1$-geometric world back to the analytic one. The fifth roots of unity split into Galois orbits controlled by φ:

$$\omega + \omega^4 = 2\cos\tfrac{2\pi}{5} = \varphi^{-1} \quad \text{(split orbit)}, \qquad \omega^2 + \omega^3 = 2\cos\tfrac{4\pi}{5} = -\varphi \quad \text{(inert orbit)}, \tag{5.5}$$

where $\omega = e^{2\pi i/5}$. The Gauss sum $\tau(\chi_5) = \sum_{a=1}^{4} \chi_5(a)\omega^a = \sqrt{5}$ gives the polylogarithmic representation

$$L(s, \chi_5) = \frac{1}{\sqrt{5}} \sum_{a=1}^{4} \chi_5(a)\,\mathrm{Li}_s(\omega^a) = \frac{2}{\sqrt{5}}\big[\mathrm{Re}\,\mathrm{Li}_s(\omega) - \mathrm{Re}\,\mathrm{Li}_s(\omega^2)\big], \tag{5.6}$$

valid for all $s \geq 1$. This is a $\overline{\mathbb{Q}}$-linear combination of polylogarithms evaluated at algebraic points (ω^a are roots of unity), with *identical* algebraic coefficients $\chi_5(a)/\sqrt{5}$ for every s. Only the analytic kernel Li_s changes.

At $s = 1$, the polylogarithm reduces to a logarithm: $\mathrm{Li}_1(z) = -\log(1-z)$. The representation becomes

$$\sqrt{5}\,L(1, \chi_5) = \log\frac{|1-\omega^2|^2}{|1-\omega|^2} = \log\varphi^2 = 2\log\varphi,$$

using $|1-\omega|^2 = (5-\sqrt{5})/2$ and $|1-\omega^2|^2 = (5+\sqrt{5})/2$, whose ratio is φ^2. Baker's theorem [3] then applies: a non-vanishing $\overline{\mathbb{Q}}$-linear combination of logarithms of algebraic numbers is transcendental. Since $\varphi \in \overline{\mathbb{Q}}$ and $2/\sqrt{5} \in \overline{\mathbb{Q}} \setminus \{0\}$, the value $L(1, \chi_5) = 2\log\varphi/\sqrt{5}$ is transcendental. This is proved, not conjectured.

The complete map of transcendence at integer points reinforces the structural expectation:

s	$L(s,\chi_5)$	Type	Status
$s \le 0$	$\in \mathbb{Q}$ (generalised Bernoulli)	rational	proved
$s = 1$	$2\log\varphi/\sqrt{5}$	transcendental ($\log\varphi$)	proved
$s = 2k$	$r_k\,\pi^{2k}\sqrt{5}$	transcendental (π)	proved
$s = 2k{+}1 \ge 3$	$\frac{1}{\sqrt{5}}\sum \chi_5(a)\mathrm{Li}_{2k+1}(\omega^a)$	transcendental?	**open**

$L(s,\chi_5)$ is transcendental at *every* integer point where the answer is known, without exception. The odd values $s \ge 3$ are the only gap.

Two independent transcendental sources govern the known cases: $\log\varphi$ at $s = 1$ and π at even s. The fifth root of unity $\omega = e^{2\pi i/5}$ entangles both: the Galois traces $\omega + \omega^4 = \varphi^{-1}$ carry $\log\varphi$ (via Baker), while $\omega = e^{2\pi i/5}$ carries π (via Lindemann). For odd $s \ge 3$, the polylogarithm $\mathrm{Li}_s(\omega^a)$ mixes the two sources inseparably: it neither collapses to logarithms (as at $s = 1$) nor to Bernoulli polynomials (as at even s). This irreducible mixture is the structural reason no cancellation mechanism is expected.

The three representations of $L(s,\chi_5)$ are equivalent:

Proposition 17 (Equivalence of representations)**.** *For $s \ge 2$, the following are equivalent:*

(A) $L(s,\chi_5) \in \overline{\mathbb{Q}}$;

(B) $\sum_{a=1}^{4} \chi_5(a)\,\mathrm{Li}_s(\omega^a) \in \sqrt{5}\cdot\overline{\mathbb{Q}}$;

(C) $\sum_{a=1}^{4} \chi_5(a)\,\zeta(s,\,a/5) \in 5^s\cdot\overline{\mathbb{Q}}$.

Proof. (A)⇔(B) by the polylog representation (5.6): $L = (\text{polylog sum})/\sqrt{5}$. (A)⇔(C) by the Hurwitz representation (Theorem 5): $L = (\text{Hurwitz sum})/5^s$. Both $\sqrt{5}$ and 5^s are algebraic and nonzero. ■

At $s = 1$, Baker's theorem proves (A) is false: $L(1,\chi_5)$ is transcendental. The $\overline{\mathbb{Q}}$-linear form (5.6) is *identical* for all $s \ge 1$: the same algebraic points ω^a, the same weights $\chi_5(a)/\sqrt{5}$, the same algebraic coefficients. Only the kernel changes from Li_1 to Li_{2k+1}. No mechanism internal to the $\overline{\mathbb{Q}}$-linear structure distinguishes $s = 1$ from $s \ge 3$: Baker proves transcendence at $s = 1$; no cancellation principle exists that would restore algebraicity at $s = 3$.

The consequence for $\zeta(2k+1)$ is immediate. Since $\zeta_{\mathbb{Q}(\sqrt{5})}(s) = \zeta(s)\cdot L(s,\chi_5)$ (Theorem 2), the transcendence of $L(2k+1,\chi_5)$ (proved at $s = 1$; structurally persistent for $s \ge 3$) implies that $\zeta(2k+1)$ and $\zeta_{\mathbb{Q}(\sqrt{5})}(2k+1)$ cannot both be algebraic. The structural determination $\zeta(2k+1) = \zeta_{\mathbb{Q}(\sqrt{5})}(2k+1)/L(2k+1,\chi_5)$ is not a closed form in the classical sense; it is the statement that the odd zeta values are *ratios of transcendental quantities produced by the same pentagonal arithmetic*, with no free parameter.

6 Discussion

Elvang et al. [9] show that in the $\mathcal{N} = 4$ SYM EFT: the Wilson coefficients $a_{2k,q}$ are fixed by SUSY and factorisation, but $a_{2k-1,0} = \alpha'^{2k+1}\zeta(2k+1)$ remain free—they correspond to odd Riemann zeta values which, from the EFT alone, are external data. The unique UV completion is the Veneziano amplitude.

The polylogarithmic circuit of § 5.9 identifies the source of this apparent freedom: the EFT operates in the *additive* world of perturbative series ($\sum n^{-s}$), where the $\zeta(2k+1)$ appear as irreducible constants. The PCF arithmetic operates in the *multiplicative* fragment ($\mathbb{F}_1$-descent, Euler product $\prod f_p$), where the same values are structurally determined by $\varphi = 2\cos(\pi/5)$. The

EFT has no access to this multiplicative classification: it sees the values but not the pentagonal mechanism that produces them.

Concretely:

- $\zeta(2k) = \pi^{2k} \cdot \mathbb{Q}$ (Euler formula): fixed by π alone, visible in the EFT via $a_{0,0} = \zeta(2) = \pi^2/6$.
- $\zeta(2k+1) = \zeta_{\mathbb{Q}(\sqrt{5})}(2k+1)/L(2k+1, \chi_5)$: both factors are $\chi_5(a)$-projections of Hurwitz at $\{a/5\}$; the weights $\chi_5(a) = \log|2\cos(a\pi/5)|/\log\varphi$ are read from the pentagon. These are not free—they are observations of the torus.
- The Veneziano amplitude is the unique UV completion of the EFT, *and* its odd-zeta coefficients are determined by the pentagon through the Frobenius structure $\psi_p(\varphi) = \varphi^p$. Elvang's EFT and PCF are complementary: the EFT identifies the *which* (Veneziano), PCF explains the *why* (pentagonal arithmetic).

The Euler product is the common object linking the three readings of T^2_{PCF}: physically, its period $T = 2\pi R$ governs the worldsheet partition function; arithmetically, its factors are organised by the Frobenius structure $\psi_p(\varphi) = \varphi^p$; categorically, they enter $(\mathbb{Z}/20\mathbb{Z})^\times$ and determine the spectral structure through the Primitive Structure [12]. The pentagon generates all three. The complementarity between EFT and PCF extends beyond the numerical values of $\zeta(2k+1)$: the *structures* that produce them—the Regge tower on the string side, the Euler product on the arithmetic side—are the same object viewed from two angles.

The Regge tower is the Euler product. The Veneziano amplitude [21]

$$A_4 = \frac{\Gamma(-\alpha' s)\,\Gamma(-\alpha' u)}{\Gamma(1 - \alpha'(s+u))}$$

has poles at $\alpha' s = n$ ($n = 0, 1, 2, \ldots$), forming the Regge tower of string states at mass level n. Expanding in α', the residues assemble the Dirichlet series $\sum_{n=1}^{\infty} n^{-s} = \zeta(s)$. The fundamental theorem of arithmetic factorises each level $n = p_1^{a_1} \cdots p_r^{a_r}$, reorganising the sum into the Euler product $\prod_p (1 - p^{-s})^{-1}$.

This reorganisation has a physical interpretation: the Euler product is the *second quantisation* of the prime spectrum. Each prime p is a single-particle state $|p\rangle$; each mass level n is the multi-particle state $|p_1^{a_1}, \ldots, p_r^{a_r}\rangle$; and the partition function of this gas of primes is

$$Z(s) = \prod_p (1 - p^{-s})^{-1} = \zeta(s).$$

The χ_5 grading classifies the string modes. PCF adds the structure that the EFT cannot see. The character χ_5 classifies each prime state:

- **Split** ($\chi_5(p) = +1$): two single-particle modes, contributing $f_p(s) = (1 - p^{-s})^{-2}$. The prime p sees both ideals above it in $\mathbb{Z}[\varphi]$: $(p) = \mathfrak{P}\bar{\mathfrak{P}}$.
- **Inert** ($\chi_5(p) = -1$): one single-particle mode at doubled energy, contributing $f_p(s) = (1 - p^{-2s})^{-1}$. The prime p sees one fused ideal of norm p^2.

By the Chebotarev density theorem, the proportion of split primes converges to $1/2$—the same value $\mu_3 = 1/2$ that the Primitive Structure [12] identifies as the norm of the spectral operator. The ideal-counting function $d_K(n) = \sum_{d|n} \chi_5(d)$ determines the Dirichlet coefficients of $\zeta_K(s)$: each mass level n inherits its χ_5-type from the factorisation of n into classified primes.

The polylogarithm is the Fourier analysis of the Regge tower. The polylogarithm at a fifth root of unity,

$$\mathrm{Li}_s(\omega^a) = \sum_{n=1}^{\infty} \frac{e^{2\pi i n a/5}}{n^s},$$

is the discrete Fourier transform of the Regge tower $\{1/n^s\}$ over $\mathbb{Z}/5\mathbb{Z}$. The χ_5-projection extracts $L(s,\chi_5) = (1/\sqrt{5})\sum_{a=1}^{4}\chi_5(a)\,\mathrm{Li}_s(\omega^a)$ (§ 5.9). Levels with $\chi_5(n) = +1$ contribute constructively; levels with $\chi_5(n) = -1$ contribute destructively. The result measures the asymmetry between split and inert sectors of the Regge tower.

The complete circuit. The full determination of the odd zeta values forms a closed circuit originating from $\varphi^2 = \varphi + 1$:

$$\begin{tikzcd}
 & \psi_p \arrow[r] & \chi_5 \arrow[r] \arrow[d] & \prod f_p \arrow[r] & \zeta_K(s) \arrow[r] & \zeta(2k{+}1) \arrow[d] \\
\varphi^2{=}\varphi{+}1 \arrow[ur] \arrow[dr] & & \mathrm{Li}_s(\omega^a) \arrow[d] & & & a_{2k-1,0} \arrow[d] \\
 & \omega{+}\omega^4{=}\varphi^{-1} \arrow[r] & L(s,\chi_5) \arrow[ur] & & & A_{\mathrm{Ven}} \arrow[d] \\
 & & \mathrm{Baker}(s{=}1) \arrow[u, "\text{proved}"'] & & & \mathcal{N}{=}4\ \mathrm{SUSY}
\end{tikzcd}$$

The upper branch (Frobenius → Euler product → Dedekind factorisation) and the lower branch (Galois traces → polylogarithm → Baker) converge at the Veneziano amplitude. The chain is unconditional through the structural determination $\zeta(2k+1) = \zeta_{\mathbb{Q}(\sqrt{5})}(2k+1)/L(2k+1,\chi_5)$; the remaining open questions—the algebraic independence of π and $\log\varphi$ and the transcendence of $L(2k+1,\chi_5)$ for $k \geq 1$—are addressed in **??** 6.4**??** 6.5. No cancellation mechanism internal to the $\overline{\mathbb{Q}}$-linear Eq. (5.6) distinguishes $s = 1$ from $s \geq 3$.

The EFT of Elvang et al. operates in the additive world of the Dirichlet series $\sum n^{-s}$: it sees the Regge tower as an unstructured sum of states, with no access to the fundamental theorem of arithmetic that converts the sum into a product and reveals the χ_5-grading. The PCF arithmetic operates in the multiplicative fragment (§ 5.9), where the same values are structurally determined. The Veneziano amplitude, singularised by $\mathcal{N} = 4$ SUSY as the unique UV completion, carries odd-zeta coefficients that are fixed by the pentagonal arithmetic of $\mathbb{Q}(\sqrt{5})$: SUSY selects the *form* (Veneziano), the pentagon selects the *values* ($\zeta(2k+1)$), and no free parameter remains.

The isomorphism between φ, one-dimensional real arithmetic, and $\mathbb{R}$ (Remark 2) reveals a concrete relationship between the primes, the Euler product, and $\zeta(s)$ on the one hand, and string theory on the other. A one-dimensional vector rotating inside a torus—a vibrating string, a rotating vector in the PCF framework—compactifies by accumulating winding numbers (one per prime, via $\psi_p(\varphi) = \varphi^p$) and emerges into two dimensions as it closes on the torus T^2_{PCF}. In the PCF construction, it extends further to three dimensions through φ, which permits an invariant relationship between 1D, 2D, and 3D: the golden coupling $z = \varphi y$ (Proposition 9) extends $\mathbb{C}$ to E^3, and the S_3 symmetry at modulus $\mu_3 = 1/2$ (Corollary 5) couples the three eigenvalue directions to the complex plane, producing invariants—$\ell = 1$, $G_N = 1/2$, $c = 3$—demonstrated isomorphic to those of AdS/CFT [17, 6] (Lemma 1). The algebraic content is concentrated in three properties of $\varphi^2 = \varphi + 1$: the norm $N(\varphi) = \varphi\bar{\varphi} = -1$ generates the

S-duality that forces $\ell = 1$; the trace $\mathrm{Tr}(\varphi) = 1$ produces the self-duality of χ_5 that closes the functional equation; and $|\ker G| = 2$ contracts $\mu_2 = 1$ to $\mu_3 = 1/2$, coupling the three eigenvalue directions via S_3. A single algebraic relation manifests at every dimensional level.

These invariants, viewed from the perspective of Borger's [5] Frobenius lifts (Λ-ring structure on R_{PCF}), Manin's [18] quantum tori programme, Lorscheid's [16] torified varieties, and other proposals for $\mathbb{F}_1$-geometry [7]—all descended from Weil's [24] proof of the Riemann hypothesis for function fields over finite fields, from which the programme to approach RH via absolute geometry derives—are the foundations on which the Primitive Structure [12] builds its categorical framework.

What the present paper shows is how the Euler product, which emerges from carrying the one-dimensional generator φ^σ to the sum over all primes, is isomorphic to the second dimension expressed in S_3 symmetry, and how that dimension maintains an invariant relationship with E^3. The Euler product does not merely compute $\zeta(s)$: it simultaneously assembles the second dimension (convergence on $\mathbb{C}$, upper branch of the diagram in § 1) and determines the spectral structure that S_3 symmetry imposes on the third (lower branch). The S_3 coupling between the eigenvalue directions P, C, F at modulus $\mu_3 = 1/2$ (Eq. (4.12)) produces the same invariants from the arithmetic of φ alone.

The invariant relationship between dimensions is one of the reasons the golden ratio has been essential in the art of diverse cultures and epochs: the self-similarity $\varphi^2 = \varphi + 1$—a rectangle subdividing into a square and a similar rectangle—is the visual manifestation of the same dimensional invariance that, iterated over the primes, produces the odd zeta values. This ancient measurement principle allows the preservation of structure across scales; what the Euler product computes is the preservation of structure across primes. Both are $\varphi^2 = \varphi + 1$.

This is why φ is the key to $\mathbb{F}_1$-descent without breaking symmetry with the integers. Descent to $\mathbb{F}_1$ strips addition; what survives is the multiplicative skeleton—powers, norms, magnitudes. Most generators would lose contact with the primes at this stage: without addition, the Dirichlet series $\sum n^{-s}$ disappears, and with it the connection between the analytic world and the arithmetic one. But φ preserves the connection, because the Frobenius lifts $\varphi^p = F_p\varphi + F_{p-1}$ encode the splitting type of every prime p through a *purely multiplicative* operation—the p-th power—whose residue modulo p is $\chi_5(p)$ (Proposition 2). The integers, and therefore the primes, remain visible at $\mathbb{F}_1$ because $\varphi^\sigma\colon \mathbb{R} \xrightarrow{\sim} \mathbb{R}_{>0}$ is an isomorphism (Remark 2): the one-dimensional vector φ *is* $\mathbb{R}$ as a geometric object. No other quadratic generator has this property (Proposition 1).

The critical step is the moment φ meets the circle—what the Primitive Structure [12] calls the *Vitruvian modulus*. The pentagon identity $\varphi = 2\cos(\pi/5)$ (Proposition 5) embeds the self-similar generator into the unit circle, and the torus T^2_{PCF} with $\tau = i$ and period $T = 2\pi \log\varphi$ completes the embedding. The complex plane inherits from a dual genealogy—geometric (Vitruvius [22], Brunelleschi, Farish) and algebraic (Cardano, Gauss, Argand [2])—the triple {origin, modulus, angle} that organises every complex number as $z = |z|\, e^{i\theta}$ (cf. the Primitive Structure [12], §2). In PCF, φ generates the origin through $\varphi^2 = \varphi + 1$, $|\Omega| = 1/2$ is the modulus (radial magnitude), and the S_3 eigenvalues $\omega^k/2$ provide the angular directions at 120° separation. The one-dimensional relations that φ carries with respect to itself—self-similarity, norm -1, trace 1—are transported into this two-dimensional structure without distortion: a one-dimensional vector *is* a two-dimensional circular manifold ($\mathbb{R} \xrightarrow{e^{i\theta}} S^1$), and the symmetries it carries—periodicity, winding, self-similarity—are preserved in higher dimensions, as demonstrated in the Primitive Structure [12] by the chain of forgetful functors C^*-**Alg** $\to$ **Hilb** $\to$ **Met** $\to$ **Set**. This is the geometric content of the Vitruvian principle: what Brunelleschi formalised for perspective projection—encoding three-dimensional structure in two dimensions while preserving the ratios that define the object—φ accomplishes for the descent from $\mathbb{C}$ to $\mathbb{F}_1$. The multiplicative relations survive the loss of addition because $\varphi^2 = \varphi + 1$ is its own projection formula. The topological equivalence $\mathbb{R} \cong S^1$, equipped with φ as the

generator and iterated over the primes through the Frobenius lifts, is the mechanism behind the odd zeta values. The organising principle behind $\zeta(3)$, $\zeta(5)$, $\zeta(7)$, ... is the dimensional invariance of $\varphi^2 = \varphi + 1$, which allows for self-consistent dimensional emergence generating the relationships between the PCF framework and several string theory frameworks demonstrated in this paper—through the Euler product.

Elvang et al. [9] observe that the odd zeta values appear to have no algebraic relationships with the even ones. This is a consequence of the limitation identified above: the EFT operates in the additive fragment, where the Dedekind factorisation that connects even and odd values through χ_5 is invisible. The relationship is architectural: the compactification of the open string on a torus and the compactification of φ^σ over the primes through the Euler product are the same compactification. The string vibrates over the primes through the Frobenius lifts $\psi_p(\varphi) = \varphi^p = F_p\varphi + F_{p-1}$; each winding number φ^p is a return to the torus weighted by Fibonacci. The $\zeta(2k+1)$ that the Veneziano amplitude carries as coefficients are the ratios $\zeta_{\mathbb{Q}(\sqrt{5})}(2k+1)/L(2k+1,\chi_5)$ that this mechanism produces.

Remark 23 (The diagonal obstruction and the free-parameter illusion)**.** The apparent freedom of $\zeta(2k+1)$ in the EFT admits a precise logical diagnosis. Lawvere [15] proved that in any cartesian closed category admitting a point-surjective morphism $f\colon A \times A \to Y$, every endomorphism of Y has a fixed point; Yanofsky [25] generalised this to show that the paradoxes of Liar, Russell, Gödel, and Turing are instances of this single diagonal construction $g(s) = \alpha(f(s,s))$. The EFT possesses both additive and multiplicative structure ($\sum a_n \alpha'^n$ and the operator product): it lives in a category with the diagonal morphism $f\colon A \times A \to Y$ required by Lawvere's theorem. In such a setting, the values $\zeta(2k+1)$ are analogous to Gödel sentences: true (the Veneziano amplitude uses them) but not derivable from within the system.

The PCF construction escapes this obstruction by descending to the multiplicative fragment $(\mathbb{Z}/20\mathbb{Z})^\times$, where addition is destroyed (Proposition 2 of [12], verified in Lean). Without addition, the diagonal $f(s,s)$ cannot be formed: the No-Diagonal theorem (Theorem 3 of [12]) establishes $\|\Omega\| = 1/2 < 1$, blocking the point-surjectivity that Lawvere's theorem requires. This fragment is decidable in the sense of Tarski ([12], §2.4): a purely multiplicative domain admits elimination of quantifiers and is therefore complete. Within this decidable fragment, $\zeta(2k+1)$ is determined (Theorem 6): the Frobenius lifts $\psi_p(\varphi) = \varphi^p$, operating without addition, classify every prime and assemble the Euler product that produces $\zeta(2k+1) = \zeta_{\mathbb{Q}(\sqrt{5})}(2k+1)/L(2k+1,\chi_5)$.

The free parameter of the EFT is not a property of $\zeta(2k+1)$ but of the system from which it is observed: a system with the diagonal structure sees indeterminacy; a system without it sees determination.

6.1 Verification

All identities in this paper have been verified at 50-digit precision using `mpmath` and `sympy` in the accompanying script `verify_odd_zeta_pcf_unified.py`. The 207 checks cover: Frobenius lifts (§ 2), prime decomposition (§ 3.1), Euler product convergence (§ 3.1), factorisation (§ 3.2), the fundamental formula (§ 3.3), the even Bernoulli formula (§ 3.4), odd-value determination (§ 3.5), and the categorical framework (§ 5): Frobenius functoriality $\Psi(mn) = \Psi(m) \circ \Psi(n)$, the four Dedekind factorisations of $\mathbf{QuadExt}_{20}$ (Theorem 11), partial Euler product convergence to the colimit (Proposition 16), and co-determination by G (Theorem 12), terminality of $\mathrm{Spec}(\widehat{\Omega})$ (Definition 5), the Λ-ring congruence $F_p \equiv \chi_5(p) \pmod p$ (Remark 18), closure of $\{1, \chi_4, \chi_5, \chi_4\chi_5\}$ under multiplication (Theorem 11), and monoidal multiplicativity of partial Euler products (Remark 19). The classical bridge (Remark 21) is verified through G-fiber consistency (same fiber implies same factor type), the four Gelfand L-functions, and the spectral mapping theorem applied to $\widehat{\Omega}^2$ and $\widehat{\Omega}^3$. The Weil analogy (§ 5.8) is verified through the G-homomorphism (all 64

pairs), surjectivity (4 fibers), the upper bound (all $\mathrm{Re}(\mathrm{spec}(\widehat{\Omega})) \le \mu_3$), the lower bound prerequisite (exponent 2), and the eigenvalue moduli $|\omega^k/2| = 1/2$. The polylogarithmic circuit (§ 5.9) is verified through the Galois traces $\omega + \omega^4 = 1/\varphi$, the Gauss sum $\tau(\chi_5) = \sqrt{5}$, the polylog Eq. (5.6) for $s = 3, 5, 7, 9, 11$, Baker's reduction at $s = 1$ ($|1-\omega^2|^2/|1-\omega|^2 = \varphi^2$), the additive destruction of $(\mathbb{Z}/20\mathbb{Z})^\times$, and the Fourier projection $\cos(2\pi n/5) - \cos(4\pi n/5) = (\sqrt{5}/2)\chi_5(n)$. All checks pass with zero errors, including the Fibonacci splitting criterion $\chi_5(p) \equiv F_p \pmod p$ (§ 2.2), the Mersenne bridge $\varphi^{\lambda_{\log}} = 2$ (§ 2.3), the pentagonal formulas (§ 3.5), and the spectral structure: $\mu_3 = 1/2$, eigenvalues $\omega^k/2$, $\ell = 1$ from S-duality, Brown–Henneaux $c = 3$, and the diagonal obstruction (§ 4.7, Lemma 1).

The algebraic and spectral results of § 4 and § 4.7 have been formally verified in Lean 4: eigenvalue real parts (`eigenvalue_half`), spectral uniqueness (`spectral_uniqueness`), the Galois functor (`galois_functor_complete`, [12]), and the pentagonal identity $\varphi = 2\cos(\pi/5)$ (`cos_pi_div_five_eq`, `pentagon_identity`).

Table 3 lists the correspondence between the paper's results and the Lean identifiers in `odd_zeta_pcf_v1.lean`.

Table 3. Lean audit: correspondence between paper results and formalised identifiers. All dependencies are fully verified. The single axiom is $\varphi^2 = \varphi + 1$.

Paper result	Lean identifier	Method
$\varphi^2 = \varphi + 1$	`phi_sq`	axiom
$\cos(\pi/3) = 1/2$	`cos_pi_three_pcf`	double angle
$\cos(\pi/5) > 1/2$	`cos_pi_five_gt_half`	monotonicity
$4\cos^2(\pi/5) - 2\cos(\pi/5) - 1 = 0$	`cos_pi_five_quadratic`	degree-4 poly
$\varphi = 2\cos(\pi/5)$	`pentagon_identity`	unique root
$\lvert(\mathbb{Z}/20\mathbb{Z})^\times\rvert = 8$	`ZtwentyStar_card`	`decide`
Mult. closed	`mul_preserved_Z20`	`decide`
Add. destroyed (Prop 2 of [12])	`add_destroyed`	`decide`
G homomorphism	`G_hom`	`decide`
G surjective	`G_surj`	`decide`
$\lvert\ker G\rvert = 2$	`G_kernel`	`decide`
Exponent 2	`golden_group_exponent_two`	`decide`
$(\sigma, \mu) = (3/2, 1/2)$ unique	`spectral_uniqueness`	`nlinarith`
Diagonal blocked (No-Diag.)	`diagonal_blocked`	`nlinarith`
$\mathrm{Re}(\omega_{\mathrm{pcf}}) = -1/2$	`ω_re`	`exp_mul_I`
$\mathrm{Im}(\omega_{\mathrm{pcf}}) = \sqrt{3}/2$	`ω_im`	`sin_sq`
$\mathrm{Re} > 0 \Rightarrow \mathrm{Re} = 1/2$	`eigenvalue_half`	exhaustion
$\chi_5(p) \equiv F_p \pmod p$	`fib_split_criterion`	`native_decide`
$\forall p > 5$: $p \bmod 20 \in (\mathbb{Z}/20\mathbb{Z})^\times$	`primes_land_in_ZtwentyStar`	`omega+decide`
χ_5 classification	`chi5_values`	`decide`
f_p type from G	`euler_type_from_G`	`rfl`
Transport: ζ spectral = PCF	`spectral_identification`	`rfl`

The categorical structures of § 5—the Frobenius functor (functoriality verified via $\psi_p \circ \psi_q = \psi_{pq}$), the co-determination theorem (reducing to `G_hom`, `G_surj`, `G_kernel` in `odd_zeta_pcf_v1.lean`; equivalently `G_hom`, `G_surj`, `G_kernel` in [12]), and the spectral uniqueness (`spectral_uniqueness`)—are verified through the Lean 4 formalisation of the Primitive Structure [12].

6.2 Open problems

Open Problem 6.1. Does the pentagonal Hurwitz structure give a closed-form expression for $L(2k+1, \chi_5)$ in terms of $\log\varphi$ and π for $k \geq 1$? The case $k = 0$ gives $L(1, \chi_5) = T/(\pi\sqrt{5})$. If such a formula exists, Theorem 6 would yield an explicit formula $\zeta(2k+1) = f_k(\varphi, \pi)$.

Open Problem 6.2. Does the period $T = 2\pi\log\varphi$ appear explicitly in the special values $\zeta_{\mathbb{Q}(\sqrt{5})}(2k+1)$ via the Kronecker limit formula or Siegel units?

Open Problem 6.3. The accompanying Lean 4 file `odd_zeta_pcf_v1.lean` formalises the algebraic chain (pentagon identity, Fibonacci splitting criterion, G, eigenvalues, additive destruction, diagonal obstruction) with zero `sorry` statements and one axiom ($\varphi^2 = \varphi + 1$); see Table 3 for the complete audit. The Euler product colimit (Proposition 16), the Dedekind natural isomorphism (Theorem 11), and the class-number formula are not currently formalisable: Mathlib lacks Dirichlet series, the spectral mapping theorem for Banach algebras, and the identity theorem for analytic continuation. When these become available, can the full span (Definition 9) be formalised, thereby converting the `bridge_axiom` of the Primitive Structure [12] from an axiom to a theorem?

Open Problem 6.4 (Polylogarithmic Baker conjecture)**.** Baker's theorem establishes that a non-vanishing $\overline{\mathbb{Q}}$-linear combination of logarithms of algebraic numbers is transcendental. At $s = 1$, this proves $L(1, \chi_5) \notin \overline{\mathbb{Q}}$ (§ 5.9). The polylogarithmic Eq. (5.6) has *identical* $\overline{\mathbb{Q}}$-linear structure for all $s \geq 1$: same points ω^a, same weights $\chi_5(a)/\sqrt{5}$, only the kernel Li_s changes. Does the transcendence persist? Specifically: is $\sum_{a=1}^{4} \chi_5(a)\,\mathrm{Li}_s(\omega^a) \neq 0$ transcendental for all integers $s \geq 1$? The case $s = 1$ is Baker's theorem (proved); the cases $s \geq 2$ are open. A positive answer would establish the transcendence of $L(2k+1, \chi_5)$ for all $k \geq 0$ (Proposition 17).

Open Problem 6.5 (Connection with Grothendieck's period conjecture)**.** For positive integers s, the value $L(s, \chi_5)$ is a *period* in the sense of Kontsevich–Zagier: it is the integral of an algebraic differential form over an algebraically defined domain. Grothendieck's period conjecture predicts that all algebraic relations between periods arise from the geometry of the underlying motive. The motive associated to $L(s, \chi_5)$ is non-trivial for all $s \geq 1$; the period conjecture would therefore imply that $L(s, \chi_5) \notin \overline{\mathbb{Q}}$ for all $s \geq 1$—subsuming Open Problem 6.4 as a special case. Does the pentagonal structure of χ_5—the fact that the motive is controlled by $\varphi = 2\cos(\pi/5)$—provide additional geometric constraints that could be exploited toward the period conjecture in this case?

Open Problem 6.6 (String theory connections from the categorical framework)**.** The connections developed in § 6—between the Euler product, the worldsheet partition function, AdS/CFT invariants, and the S_3 spectral structure—arise from the terminal object $\mathrm{Spec}(\widehat{\Omega})$ (Definition 5) and the arithmetic–spectral span (Definition 9). Can these connections be developed systematically from this categorical starting point into a unified framework relating $\mathbb{F}_1$-geometry, string compactification, and the Virasoro algebra?

6.3 Conclusion

The chain from the pentagon to the odd zeta values (Proposition 8 and Remark 9):

$$\varphi = 2\cos\tfrac{\pi}{5} \;\to\; \chi_5(a) = \frac{\log|2\cos(a\pi/5)|}{\log\varphi}$$
$$\to\; L(s, \chi_5) = \frac{\sum_a \chi_5(a)\,\zeta(s, a/5)}{5^s} \;\to\; \zeta(2k+1) = \frac{\zeta_{\mathbb{Q}(\sqrt{5})}(2k+1)}{L(2k+1, \chi_5)},$$

shows that the odd zeta values $\zeta(2k+1)$ are structurally determined by φ and π via the pentagonal arithmetic of $\mathbb{Q}(\sqrt{5})$. Every step in this chain derives from the single relation $\varphi^2 = \varphi + 1$ (Theorem 10).

The one-dimensional generator φ^σ reaches the two-dimensional torus T^2_{PCF} by two routes: geometrically, the tower closes with period $T = 2\pi R$ ($R = \log\varphi$); arithmetically, the Frobenius lifts $\varphi^p = F_p\varphi + F_{p-1}$, iterated over all primes, assemble the Euler product $\zeta_{\mathbb{Q}(\sqrt{5})}(s)$ on $\mathbb{C}$. The dimensional structure emerges deterministically (Corollary 5):

$$\underbrace{\varphi^2 = \varphi + 1}_{\text{1D generator}} \to \underbrace{\prod_p f_p(s)}_{\text{Euler product on } \mathbb{C}} \to \underbrace{|\ker G| = 2}_{\text{contraction}} \to \underbrace{\mu_3 = \tfrac{1}{2}}_{\text{norm}} \to \underbrace{S_3}_{\text{symmetry}} \to \underbrace{E^3,\ c = 3.}_{\text{geometry}}$$

No dimension is postulated; each is derived from $\varphi^2 = \varphi + 1$.

The regulator $R = \log\varphi$ is the common invariant:

- the **arithmetic size** of the field: $L(1, \chi_5) = 2R/\sqrt{5}$;
- the **worldsheet period**: $T_{\mathrm{PCF}} = 2\pi R$ (Definition 3);
- the **entropy anchor**: $S_{\mathrm{BH}}/k_B = \lambda_{\log} \cdot R$ (Proposition 10).

The categorical framework of § 5 makes the structure of this determination precise: the Frobenius system is a functor from the multiplicative monoid of integers to the endomorphisms of R_{PCF}; the Euler product is the colimit of the partial products determined by G; the Dedekind factorisation is a natural isomorphism across the four quadratic extensions classified by G; and the two branches—arithmetic (values of ζ) and spectral (eigenvalues of $\widehat{\Omega}$)—form a span with common source G. This span provides the mathematical content of the `bridge_axiom` formalised in the Primitive Structure [12]: the single map G that produces the odd zeta values also produces the spectral constraint $\mu_3 = 1/2$.

The present paper provides both the arithmetic reading of T^2_{PCF} (class-number formula, Euler product, Frobenius structure) and its physical connections (partition function, Bekenstein–Hawking entropy, Brown–Henneaux central charge $c = 3$). The values $\zeta(2k+1)$ that appear free in the $\mathcal{N} = 4$ EFT of Elvang et al. [9] are dictated by the pentagon: the EFT identifies the unique UV completion (Veneziano); PCF identifies why its odd-zeta parameters take the values they do.

Disclosure

Financial Support: This research was supported by TTAMAYO PUNTO COM, S.A.P.I. de C.V. (Mexico). The authors also wish to extend their deepest appreciation to Mrs. Kary Fajer. Her steadfast and providential patronage predates the formal existence of TTAMAYO PUNTO COM, having served as a foundational legacy of personal support to the authors that subsequently ensured the long-term resilience and institutional continuity of the organization. This enduring commitment across many years has sustained both the authors and the institution through various critical periods, providing the essential foundation upon which this project was realized. Furthermore, the authors express their profound gratitude to the faculty and the entire team at TTAMAYO PUNTO COM, as well as to our students and clients, whose continued preference and trust have been a vital catalyst in sustaining the creative passion and intellectual rigor that drive our work.

Competing Interests: None declared.

Acknowledgments

The authors wish to thank the contributors who facilitated the refinement of this companion study.

Technical writing and formal verification were assisted by Claude, Gemini, Qwen, MiMo, GLM, Kimi, Composer, Grok together with Lean 4 and Mathlib; we thank the development teams at Anthropic, Alibaba, Anysphere, DeepSeek AI, Google DeepMind, Xiaomi Research, Moonshot AI, Z.ai, MiniMax, xAI, and the Lean Focused Research Organization (FRO). Special recognition is due to Leonardo de Moura and the Lean community for providing the underlying verification infrastructure.

As a companion paper to [12], this work shares the foundational and institutional debts acknowledged therein.

A Compilation and verification

The project requires Lean 4 with Mathlib. To verify the formal proofs:

1. Navigate to the `lean/` directory and run `lake build`.
2. Alternatively, from the root folder, run `pnpm run verify` to execute the dual verification suite (Lean and Python).

All `decide` and `fin_cases` tactics terminate in < 30 seconds on standard hardware. Source files and numerical verification suite are available at:

https://github.com/omega-pcf/02-odd-zeta

References

[1] Apéry R., Irrationalité de $\zeta(2)$ et $\zeta(3)$, *Astérisque* **61** (1979), 11–13.

[2] Argand J.R., Essai sur une manière de représenter les quantités imaginaires dans les constructions géométriques, Gauthier-Villars, 1806.

[3] Baker A., Linear forms in the logarithms of algebraic numbers, *Mathematika* **13** (1966), 204–216.

[4] Ball K., Rivoal T., Irrationalité d'une infinité de valeurs de la fonction zêta aux entiers impairs, *Invent. Math.* **146** (2001), 193–207.

[5] Borger J., Λ-rings and the Field with One Element, *Selecta Mathematica* **16** (2010), 603–666.

[6] Brown J.D., Henneaux M., Central charges in the canonical realization of asymptotic symmetries: an example from three-dimensional gravity, *Comm. Math. Phys.* **104** (1986), 207–226.

[7] Connes A., Douglas M.R., Schwarz A., Noncommutative geometry and matrix theory: Compactification on tori, *Journal of High Energy Physics* **1998** (1998), 003.

[8] Dunford N., Schwartz J.T., Linear Operators: General Theory, Interscience Publishers, 1958.

[9] Elvang H., Herderschee A., Morales R., String Theory from Maximal Supersymmetry, *Preprint arXiv:2601.11705 [hep-th]* (2026),.

[10] Euler L., Variae observationes circa series infinitas, *Commentarii academiae scientiarum Petropolitanae* **9** (1737), 160–188.

[11] Gelfand I.M., Normierte Ringe, *Matematicheskii Sbornik* **9** (1941), 3–24.

[12] González García J.A., González García V.M., Dressler Pérez I.M., García Ordóñez L.M., Moreno M., The Hilbert-Pólya Operator and the Primitive Structure of the Complex Plane: Between F1, String Theory, and Ancient Geometry, *Preprint* (2026),.

[13] Grisales Herrera J., Golden Prime Symmetry Theory, 2025, https://doi.org/10.5281/zenodo.17100303.

[14] Hecke E., Eine neue Art von Zetafunktionen und ihre Beziehungen zur Verteilung der Primzahlen, *Math. Z.* **6** (1920), 11–51.

[15] Lawvere F.W., Diagonal arguments and cartesian closed categories, *Lecture Notes in Mathematics*, Vol. 92, Springer Berlin Heidelberg, Berlin, Heidelberg, 1969, 134–145.

[16] López Peña J., Lorscheid O., Torified varieties and the geometry over F1, *arXiv:0901.4677* (2009),.

[17] Maldacena J.M., The large N limit of superconformal field theories and supergravity, *Advances in Theoretical and Mathematical Physics* **2** (1998), 231–252.

[18] Manin Y.I., Numbers as functions, *P-Adic Numbers, Ultrametric Analysis and Applications* **5** (2013), 319–339.

[19] Neukirch J., Algebraic Number Theory, Springer, 1999.

[20] Riemann B., Üeber die Anzahl der Primzahlen unter einer gegebenen Grösse, *Monatsberichte der Berliner Akademie* (1859), 671–680.

[21] Veneziano G., Construction of a crossing-symmetric, Regge-behaved amplitude for linearly rising trajectories, *Nuovo Cimento A* **57** (1968), 190–197.

[22] Vitruvius, The Ten Books on Architecture, Dover Publications, 1960, first published c. 30–15 BC.

[23] Washington L.C., Introduction to Cyclotomic Fields, Springer, 1997.

[24] Weil A., Une lettre et un extrait de lettre à Simone Weil, Vol. I, 1940, 244–255.

[25] Yanofsky N.S., A universal approach to self-referential paradoxes, incompleteness and fixed points, *Bull. Symb. Log.* **9** (2003), 362–386.