

Reductions for the finite implication $E677 \models_{\text{fin}} E255$

Anonymous note

May 2026

Abstract

The Equational Theories Project leaves open, up to duality, the finite implication $E677 \models_{\text{fin}} E255$ for magmas, while the corresponding unrestricted implication is known to be false. This note records a checked package of reductions for that remaining finite problem. The main output is a route graph: the target law at a point is equivalent to a unique right fixed point, to a left-division identity called Lemma L, and to several constrained collision and counterexample conditions. The note also isolates false trails that should not be reused, including a previously suspected quotient family and a full-shift law. No proof or finite counterexample is claimed.

2020 Mathematics Subject Classification. Primary 08A05; Secondary 08B05, 03C13, 68V15.

Keywords. Finite magma, equational law, finite implication, left division, equational theories.

1 Introduction

A *magma* is a set M equipped with a binary operation $\diamond$. The two laws studied here are

$$\begin{aligned} E677 : \quad & x = y \diamond (x \diamond ((y \diamond x) \diamond y)), \\ E255 : \quad & x = ((x \diamond x) \diamond x) \diamond x. \end{aligned}$$

The problem is the finite implication

$$E677 \models_{\text{fin}} E255,$$

that is, whether every finite magma satisfying $E677$ also satisfies $E255$. The unrestricted implication $E677 \models E255$ is false. The Equational Theories Project (ETP) paper records the finite problem as the only unresolved finite implication up to duality, equivalently together with the dual implication $E2910 \models_{\text{fin}} E47$, and tentatively conjectures the implication to be false [1, Problem 8.1]. The ETP blueprint chapter on Equation 677 contains the local structure theory and the known no-go statements for standard counterexample families [4]; the dashboard records the graph status [3].

This note is not a computational search report. Its purpose is to make the remaining mathematical problem sharper. The reductions below are organized around a fixed element x of a finite $E677$ -magma. They show that the target law at x is the same as a single right fixed point, that any such fixed-point witness is unique, and that the same target can be expressed as a left-division identity (Lemma L). The complementary sections describe what a collision proof or a finite counterexample would have to supply.

2 Preliminaries and external input

For $a \in M$, write

$$L_a(t) = a \diamond t, \quad R_a(t) = t \diamond a, \quad S(t) = t \diamond t.$$

If L_a is bijective, write $a \setminus b$ for the unique element u satisfying $a \diamond u = b$. All products in this note are non-associative and are parenthesized when there is any possible ambiguity.

The finite hypothesis first enters through the following standard ETP fact.

Proposition 1 (Left division in finite E677-magmas). *Let M be a finite magma satisfying E677. Then every left translation L_a is bijective. Moreover*

$$a \setminus b = b \diamond ((a \diamond b) \diamond a)$$

for all $a, b \in M$.

Proof. Law E677 gives

$$b = a \diamond (b \diamond ((a \diamond b) \diamond a)).$$

Thus L_a is surjective. Since M is finite, L_a is bijective, and the displayed preimage formula is exactly the formula for $a \setminus b$. $\square$

The ETP chapter also records several strategic facts that are used here only as external input. The unrestricted implication $\text{E677} \not\models \text{E255}$ is witnessed by the free E677-magma [4, Section 13.2]. Linear models satisfying E677 satisfy E255, and affine-fiber extensions over such models do not produce counterexamples [4, Lemmas 13.3–13.4]. The same chapter gives finite E677-magmas that are not right-cancellative [4, Section 13.1]. Hence no argument below may use right cancellation, associativity, an identity element, or group or quasigroup intuition unless such structure has first been derived. The finite-map lemmas in the ETP finite chapter are a natural source of future tools [5, Lemmas 5.5–5.7].

Lemma 2 (Transformed identity). *Let M be a finite magma satisfying E677. Then*

$$z = (y \diamond z) \diamond ((y \diamond (y \diamond z)) \diamond y)$$

for all $y, z \in M$.

Proof. Substitute $y \diamond z$ for x in E677. This gives

$$y \diamond z = y \diamond ((y \diamond z) \diamond ((y \diamond (y \diamond z)) \diamond y)).$$

Cancel the common left factor y using Proposition 1. $\square$

3 The orbit package

Fix a finite E677-magma M and an element $x \in M$. Define

$$c_0 = x, \quad c_{k+1} = x \diamond c_k,$$

and let d be the minimal period of this L_x -orbit, so that $c_d = x$ and $c_0, \dots, c_{d-1}$ are pairwise distinct. Indices on the c_k are read modulo d . Put

$$d_k = c_k \diamond x.$$

Lemma 3 (Orbit recurrence). *For every k ,*

$$c_{k-1} = c_k \diamond d_{k+1}.$$

Consequently $d_1 = c_{d-2}$.

Proof. Apply E677 with $y = x$ and the law variable x replaced by c_k :

$$c_k = x \diamond (c_k \diamond ((x \diamond c_k) \diamond x)) = x \diamond (c_k \diamond d_{k+1}).$$

Since also $c_k = x \diamond c_{k-1}$, left cancellation gives the recurrence. Taking $k = 0$ gives $c_{d-1} = x \diamond d_1$, so $d_1 = x \backslash c_{d-1} = c_{d-2}$. $\square$

Lemma 4 (Basic left-division identities). *For every k ,*

$$x \backslash c_k = c_{k-1}, \quad c_k \backslash c_{k-1} = d_{k+1}.$$

If

$$p = x \backslash x = c_{d-1}, \quad q = x \backslash p,$$

then

$$q = d_1 = c_{d-2} = (x \diamond x) \diamond x,$$

and

$$p \diamond (x \diamond x) = d_1, \quad p \backslash d_1 = x \diamond x.$$

Proof. The first identity follows from $c_k = x \diamond c_{k-1}$. The second is precisely the recurrence in Lemma 3. Since $x \diamond c_{d-1} = c_d = x$, one has $p = c_{d-1}$, and then

$$q = x \backslash p = x \backslash c_{d-1} = c_{d-2} = d_1.$$

Also $d_1 = c_1 \diamond x = (x \diamond x) \diamond x$. Finally, applying the general identity $a \backslash (a \backslash b) = (a \backslash b) \diamond (b \diamond a)$, which follows immediately from Proposition 1, with $a = b = x$, gives

$$x \backslash p = p \diamond (x \diamond x).$$

The left side is $q = d_1$, and hence $p \diamond (x \diamond x) = d_1$. The last displayed identity is the definition of left division by p . $\square$

Lemma 5 (A fixed point has the forced index). *If $d_j = x$, then $j \equiv d - 2 \pmod{d}$.*

Proof. The condition $d_j = x$ says $c_j \diamond x = x$. Lemma 2 with $y = c_j, z = x$ gives

$$x = x \diamond (x \diamond c_j).$$

On the other hand, E677 with both variables specialized to x gives

$$x = x \diamond (x \diamond d_1).$$

Two left cancellations yield $c_j = d_1 = c_{d-2}$, so the minimality of the orbit period gives the claimed congruence. $\square$

4 The route graph

The following display is the dependency graph used in the rest of the note. The first row contains equivalent fixed-point targets. The two lower branches describe the two ways an unfinished proof can still move: collision lifting or explicit counterexample construction.

$$\begin{array}{ccccc}
 \text{E255 at } x & \iff & q \diamond x = x & \iff & \exists u (u \diamond x = x) & \iff & q \backslash x = x \\
 & & \Downarrow & & & & \Downarrow \\
 & & \text{unique witness } q & & & & ((x \diamond x) \diamond ((x \diamond x) \diamond x)) \diamond (x \diamond x) = x \\
 & & & & & & \Downarrow \\
 & & & & & & (q \diamond x) \diamond q = p.
 \end{array}$$

If the fixed point is not established, the two honest remaining branches are

$$\begin{array}{lcl}
 d_i = d_j = e & \implies & (c_i \diamond e) \diamond c_i = (c_j \diamond e) \diamond c_j \quad \dashrightarrow \quad c_i \diamond e = c_j \diamond e \text{ or a fixed point,} \\
 x \notin \text{im } R_x & \implies & y \diamond x = z \diamond x = r \quad \implies \quad (y \diamond r) \diamond y = r \backslash x.
 \end{array}$$

The dashed arrow is a missing lemma, not a theorem.

5 The fixed-point frontier

The fixed-point route is sharper than the existential formulation suggests.

Proposition 6 (Unique right fixed-point witness). *Let M be a finite E677-magma and fix $x \in M$. If $u \diamond x = x$, then*

$$u = q = (x \diamond x) \diamond x.$$

Consequently E255 at x is equivalent to the existence of u with $u \diamond x = x$.

Proof. The first half of the proof is the argument already used in Lemma 5. If $u \diamond x = x$, then Lemma 2 gives

$$x = x \diamond (x \diamond u).$$

The specialization $y = x$, $x = x$ in E677 gives

$$x = x \diamond (x \diamond q).$$

Two left cancellations give $u = q$. Since $q = (x \diamond x) \diamond x$, the identity $q \diamond x = x$ is exactly E255 at x . Thus E255 at x is equivalent to the existence of a right fixed point over x , and the witness is forced to be q . $\square$

Proposition 7 (Equivalent local targets). *With the notation above, the following are equivalent:*

1. E255 at x , equivalently $q \diamond x = x$;
2. $q \backslash x = x$;
3. $((x \diamond x) \diamond ((x \diamond x) \diamond x)) \diamond (x \diamond x) = x$;
4. $(q \diamond x) \diamond q = p$.

Proof. The equivalence of (1) and (2) is just uniqueness of left division by q . To identify the left quotient, apply Lemma 2 with $y = x \diamond x$ and $z = x$:

$$x = q \diamond ((x \diamond x) \diamond q) \diamond (x \diamond x).$$

Therefore

$$q \backslash x = ((x \diamond x) \diamond q) \diamond (x \diamond x),$$

which proves the equivalence of (2) and (3), because $q = (x \diamond x) \diamond x$.

For (4), use Proposition 1:

$$q \backslash x = x \diamond ((q \diamond x) \diamond q).$$

Since $p = x \backslash x$, injectivity of L_x gives

$$x \diamond ((q \diamond x) \diamond q) = x \iff (q \diamond x) \diamond q = p.$$

Together with (1) $\iff$ (2), this proves the claim. $\square$

Definition 8 (Lemma L). For the fixed element x , write $L(x)$ for the identity

$$((x \diamond x) \diamond ((x \diamond x) \diamond x)) \diamond (x \diamond x) = x.$$

By Proposition 7, $L(x)$ is equivalent, in the present finite E677 setting, to E255 at x .

Remark 9. Lemma L is a target, not an established lemma. Because the unrestricted implication $E677 \models E255$ is false [4, Section 13.2], any successful proof of the fixed-point route must use finiteness in a visible way: for example through left bijection, orbit periodicity, a finite-map lemma, or a collision argument. A derivation that simply rewrites E677 into Lemma L without using finite structure should be treated as suspect.

6 Collision frontier

The orbit recurrence suggests a collision strategy, but the exact missing step is narrow.

Proposition 10 (Collision lift). *If $d_i = d_j = e$, then*

$$(c_i \diamond e) \diamond c_i = (c_j \diamond e) \diamond c_j. \tag{C}$$

If in addition $i \not\equiv j \pmod{d}$, then

$$c_i \diamond e \neq c_j \diamond e.$$

Proof. Lemma 2 with $y = c_k$ and $z = x$ gives

$$x = d_k \diamond ((c_k \diamond d_k) \diamond c_k).$$

If $d_i = d_j = e$, left cancellation by e yields (C). If also $c_i \diamond e = c_j \diamond e = s$, then (C) becomes $s \diamond c_i = s \diamond c_j$. Left cancellation by s gives $c_i = c_j$, contradicting the minimal period unless $i \equiv j \pmod{d}$. $\square$

Problem 11 (Collision-lift target). Prove or refute the following implication inside finite E677-magmas: if $d_i = d_j = e$, then either $q \diamond x = x$ already, or

$$c_i \diamond e = c_j \diamond e.$$

Together with Proposition 10, this would rule out nontrivial collisions unless the desired fixed point has already been found.

The common shortcut “prove $k \mapsto d_k$ injective” is not by itself a proof of the main theorem. It becomes useful only if the argument also connects the resulting injectivity to the forced fixed point $q \diamond x = x$, or if it supplies the missing alternative in Problem 11.

7 Counterexample frontier

Suppose a finite counterexample exists and choose $x \in M$ with $q \diamond x \neq x$. Proposition 6 says that no element u has $u \diamond x = x$. Thus $x \notin \text{im } R_x$. Since M is finite, R_x is not injective, and there exist $y \neq z$ and $r \neq x$ such that

$$y \diamond x = z \diamond x = r.$$

The following proposition records the forced shape of such a right collision.

Proposition 12 (Right-collision rectangles). *Let $y \diamond x = r$. Then*

$$(y \diamond r) \diamond y = r \setminus x. \tag{R}$$

Consequently, if $y \diamond x = z \diamond x = r$ and $y \diamond r = z \diamond r$, then $y = z$.

Proof. Lemma 2 with $y = y$ and $z = x$ gives

$$x = r \diamond ((y \diamond r) \diamond y),$$

which is exactly (R). If $y \diamond r = z \diamond r = s$, then (R) gives $s \diamond y = s \diamond z$. Left cancellation by s gives $y = z$. $\square$

This proposition is a counterexample design constraint. A candidate finite counterexample must contain right-collision fibers for every bad point x , but those fibers must split under the second probe $t \mapsto t \diamond r$, while all elements in the fiber preserve the same twisted diagonal value $(t \diamond r) \diamond t = r \setminus x$.

There are two plausible construction languages that respect this constraint. First, because all left translations are permutations, one may write

$$a \diamond b = \sigma(a)(b)$$

with $\sigma(a) \in \text{Sym}(M)$. Law E677 is then the labeled- permutation condition

$$\sigma(y)(\sigma(x)(\sigma(\sigma(y)(x))(y))) = x.$$

A counterexample at x_0 would additionally require $\sigma(a)(x_0) \neq x_0$ for every label a . This is not merely a group-theoretic condition on a subgroup of $\text{Sym}(M)$; the labeling map $a \mapsto \sigma(a)$ is part of the data.

Second, one may seek nonlinear fiber extensions over positive E677-models. The ETP affine-extension obstruction [4, Lemma 13.4] rules out the obvious affine version, so a successful extension would have to be genuinely nonlinear and would still need to realize the rectangles in Proposition 12.

8 Obstructions and false routes

Several attractive shortcuts are invalid or unsupported.

1. The family

$$d_k \setminus x = c_{k-2} \diamond c_{k-1}$$

should not be used. The old derivation confused $c_{k-1} \diamond x$ with d_k ; by definition it is d_{k-1} . The active repository does not contain a publication-ready finite table witness for this failure, so the present paper records only the derivation error and quarantines the identity.

2. The full-shift law

$$p \diamond d_k = d_{k+1}$$

is also quarantined. It is not a consequence of the trusted local package.

3. A proof of universal idempotence, $d = 1$, or any universal small-period conclusion would contradict known nontrivial positive models from the ETP finite theory. Such a claim should therefore be audited before it is used.
4. A nontrivial collision $d_i = d_j$ does not automatically propagate backwards to $d_{i-1} = d_{j-1}$. Any cascade proof must derive the propagation or avoid it.

9 Open frontiers

The work above leaves three precise frontiers.

Problem 13 (Fixed-point route). Prove $q \diamond x = x$, equivalently Lemma L from Definition 8, by an argument that uses finite structure explicitly.

Problem 14 (Collision route). Resolve Problem 11: from $d_i = d_j = e$, derive either the fixed point $q \diamond x = x$ or the equality $c_i \diamond e = c_j \diamond e$.

Problem 15 (Counterexample route). Construct a finite labeled-permutation model or a genuinely nonlinear fiber extension satisfying E677, violating E255, and satisfying the right-collision rectangle constraints of Proposition 12. A table-free parameter search is not a construction; any proposed example should include a complete finite table or a checkable structural verification.

A Proof audit checklist

The reductions in this note use left cancellation only after Proposition 1, where finiteness has made each L_a bijective. They do not use right cancellation, associativity, an identity element, or group/quasigroup structure. Future work on this problem should state which of the three frontiers it attacks and end, if unfinished, with one missing lemma rather than a broad search prescription.

Every new proof attempt should check the following list before being added to the mathematical record.

1. All products are parenthesized.
2. Every cancellation is a left cancellation by a known bijective left translation.
3. Every use of finiteness is named: left bijectivity, orbit periodicity, a finite-map lemma, or pigeonhole reasoning for a self-map.
4. The false quotient family and false full-shift law are not used.
5. Any counterexample claim contains either a complete finite operation or a structural verification of E677 and failure of E255.

References

- [1] Matthew Bolan, Joachim Breitner, Jose Brox, Nicholas Carlini, Mario Carneiro, Floris van Doorn, Martin Dvorak, Andres Goens, Aaron Hill, Harald Husum, Hernan Ibarra Mejia, Zoltan A. Kocsis, Bruno Le Floch, Amir Livne Bar-on, Lorenzo Luccioli, Douglas McNeil, Alex Meiburg, Pietro Monticone, Pace P. Nielsen, Emmanuel Osalotioman Osazuwa, Giovanni Paolini, Marco Petracci, Bernhard Reinke, David Renshaw, Marcus Rossel, Cody Roux, Jeremy Scanvic, Shreyas Srinivas, Anand Rao Tadipatri, Terence Tao, Vlad Tsyrklevich, Fernando Vaquerizo-Villar, Daniel Weber, and Fan Zheng. The Equational Theories Project: Advancing collaborative mathematical research at scale. arXiv:2512.07087, 2025. <https://arxiv.org/abs/2512.07087>; DOI: <https://doi.org/10.48550/arXiv.2512.07087>.
- [2] Equational Theories Project contributors. *The Equational Theories Project*. GitHub repository, 2024–2026. https://github.com/teorth/equational_theories.
- [3] Equational Theories Project. Dashboard. https://teorth.github.io/equational_theories/dashboard/.
- [4] Equational Theories Project. Blueprint, Chapter 13: Equation 677. https://teorth.github.io/equational_theories/blueprint/677-chapter.html.
- [5] Equational Theories Project. Blueprint, Chapter 5: Finite implications and infinite models. https://teorth.github.io/equational_theories/blueprint/infinite-model-chapter.html.
- [6] Equational Theories Project. Blueprint, Chapter 27: Hard implications. https://teorth.github.io/equational_theories/blueprint/hard-implications-chapter.html.
- [7] A. K. Austin. A note on models of identities. *Proceedings of the American Mathematical Society*, 16:522–523, 1965. DOI: <https://doi.org/10.1090/S0002-9939-1965-0174649-7>; JSTOR: <https://doi.org/10.2307/2034688>.
- [8] A. K. Austin. Finite models for laws in two variables. *Proceedings of the American Mathematical Society*, 17:1410–1412, 1966. DOI: <https://doi.org/10.1090/S0002-9939-1966-0199134-9>; JSTOR: <https://doi.org/10.2307/2035753>.
- [9] Andrzej Kisielwicz. Varieties of algebras with no nontrivial finite members. In *Lattices, Semigroups, and Universal Algebra*, pages 129–136. Plenum, New York, 1990. DOI: https://doi.org/10.1007/978-1-4899-2608-1_14.
- [10] A. Kisielwicz. Austin identities. *Algebra Universalis*, 38(3):324–328, 1997. DOI: <https://doi.org/10.1007/s000120050057>.
- [11] Joel Berman and Stanley Burris. A computer study of 3-element groupoids. In *Logic and Algebra*, Lecture Notes in Pure and Applied Mathematics 180, pages 379–429. Marcel Dekker, 1996. Reprint DOI: <https://doi.org/10.1201/9780203748671-18>.
- [12] Stanley Burris and H. P. Sankappanavar. *A Course in Universal Algebra*. Graduate Texts in Mathematics 78. Springer, New York, 1981. ISBN 978-0-387-90578-5.
- [13] Donald E. Knuth. Notes on central groupoids. *Journal of Combinatorial Theory*, 8:376–390, 1970. DOI: [https://doi.org/10.1016/S0021-9800\(70\)80032-1](https://doi.org/10.1016/S0021-9800(70)80032-1).

- [14] Donald E. Knuth and Peter B. Bendix. Simple word problems in universal algebras. In *Computational Problems in Abstract Algebra*, pages 263–297. Pergamon, 1970. DOI: <https://doi.org/10.1016/B978-0-08-012975-4.50028-X>; reprint DOI: https://doi.org/10.1007/978-3-642-81955-1_23.
- [15] Franz Baader and Tobias Nipkow. *Term Rewriting and All That*. Cambridge University Press, 1998. DOI: <https://doi.org/10.1017/CB09781139172752>.
- [16] Max Willsey, Chandrakana Nandi, Yisu Remy Wang, Oliver Flatt, Zachary Tatlock, and Pavel Panchekha. egg: Fast and extensible equality saturation. *Proceedings of the ACM on Programming Languages*, 5(POPL):1–29, 2021. DOI: <https://doi.org/10.1145/3434304>.
- [17] Thomas Koehler, Andres Goens, Siddharth Bhat, Tobias Grosser, Phil Trinder, and Michel Steuwer. Guided equality saturation. *Proceedings of the ACM on Programming Languages*, 8(POPL):1727–1758, 2024. DOI: <https://doi.org/10.1145/3632900>.