

# The Finite Free Stam Inequality

## 1 Setup and statement

Let  $p(x) = \sum_{k=0}^n a_k x^{n-k}$  and  $q(x) = \sum_{k=0}^n b_k x^{n-k}$  be monic ( $a_0 = b_0 = 1$ ) real-rooted polynomials of degree  $n$ . Their *symmetric additive convolution* is

$$(p \boxplus_n q)(x) = \sum_{k=0}^n c_k x^{n-k}, \quad c_k = \sum_{i+j=k} \frac{(n-i)!(n-j)!}{n!(n-k)!} a_i b_j.$$

For  $p(x) = \prod_{i=1}^n (x - \lambda_i)$  with distinct roots define the *scores* and *finite free Fisher information*:

$$V_i := \sum_{j \neq i} \frac{1}{\lambda_i - \lambda_j}, \quad \Phi_n(p) := \sum_{i=1}^n V_i^2,$$

with  $\Phi_n(p) := \infty$  when  $p$  has a repeated root.

**Definition 1.1** (Variance).  $\sigma^2(p) := \frac{1}{n} \sum_{i=1}^n (\lambda_i - \bar{\lambda})^2$  where  $\bar{\lambda} = \frac{1}{n} \sum_i \lambda_i$ .

**Definition 1.2** (Score-gap form).  $\mathcal{S}(p) := \sum_{i < j} \frac{(V_i - V_j)^2}{(\lambda_i - \lambda_j)^2}$ .

**Theorem 1.1** (Finite Free Stam Inequality). *For  $p, q \in \mathcal{P}_n^{\mathbb{R}}$  with distinct roots,*

$$\frac{1}{\Phi_n(p \boxplus_n q)} \geq \frac{1}{\Phi_n(p)} + \frac{1}{\Phi_n(q)}. \quad (1)$$

We prove (1) in full for  $n = 2$  (Section 4) and  $n = 3$  (Section 5), and reduce the general case to a single pointwise inequality (Section 7).

## 2 Preliminary identities

All polynomials below are monic of degree  $n$  with distinct roots.

**Lemma 2.1** (Score-root identity). *For centered  $p$  (i.e.  $\bar{\lambda} = 0$ ),  $\sum_{i=1}^n \lambda_i V_i = \frac{n(n-1)}{2}$ .*

*Proof.*  $\sum_i \lambda_i V_i = \sum_i \sum_{j \neq i} \frac{\lambda_i}{\lambda_i - \lambda_j} = \sum_{i < j} \frac{\lambda_i - \lambda_j}{\lambda_i - \lambda_j} = \binom{n}{2}$ . □

**Lemma 2.2** (Score sum).  $\sum_{i=1}^n V_i = 0$ .

*Proof.*  $\sum_i V_i = \sum_i \sum_{j \neq i} \frac{1}{\lambda_i - \lambda_j} = \sum_{i < j} \left( \frac{1}{\lambda_i - \lambda_j} + \frac{1}{\lambda_j - \lambda_i} \right) = 0$ . □

**Lemma 2.3** (Score via derivatives).  $V_i = \frac{r''(\lambda_i)}{2r'(\lambda_i)}$ , where  $r = p$ .

*Proof.* Since  $r'(\lambda_i) = \prod_{j \neq i} (\lambda_i - \lambda_j)$ , differentiating  $r'(x) = \sum_{i=1}^n \prod_{j \neq i} (x - \lambda_j)$  yields  $r''(\lambda_i) = 2 \sum_{k \neq i} \prod_{j \neq i, j \neq k} (\lambda_i - \lambda_j) = 2r'(\lambda_i) \sum_{k \neq i} \frac{1}{\lambda_i - \lambda_k} = 2r'(\lambda_i) V_i$ .  $\square$

**Lemma 2.4** (Fisher–variance inequality).  $\Phi_n(p) \sigma^2(p) \geq \frac{n(n-1)^2}{4}$ , with equality iff  $n = 2$ .

*Proof.* By Cauchy–Schwarz,  $(\sum_i \lambda_i V_i)^2 \leq (\sum_i \lambda_i^2) (\sum_i V_i^2) = n \sigma^2(p) \Phi_n(p)$ . By Lemma 2.1 the left side is  $n^2(n-1)^2/4$ .  $\square$

**Lemma 2.5** (Variance additivity).  $\sigma^2(p \boxplus_n q) = \sigma^2(p) + \sigma^2(q)$ .

*Proof.* The coefficient formula gives  $c_1 = a_1 + b_1$  and  $c_2 = a_2 + b_2$ , so the variance (a function of  $c_1, c_2$  alone) is additive.  $\square$

### 3 Critical-value formula for $\Phi_n$

**Theorem 3.1** (Critical-value formula). *Let  $r$  be a monic polynomial of degree  $n$  with distinct roots, and let  $\zeta_1, \dots, \zeta_{n-1}$  be the zeros of  $r'$  (assumed simple). Then*

$$\Phi_n(r) = -\frac{1}{4} \sum_{j=1}^{n-1} \frac{r''(\zeta_j)}{r(\zeta_j)}. \quad (2)$$

*Proof.* By Lemma 2.3,  $\Phi_n = \frac{1}{4} \sum_{i=1}^n \frac{r''(\lambda_i)^2}{r'(\lambda_i)^2}$ . Consider the meromorphic function

$$F(x) = \frac{r''(x)^2}{r'(x)r(x)}.$$

*Residues at the roots  $\lambda_i$ .* Since  $r$  has a simple zero at  $\lambda_i$ ,

$$\text{Res}_{x=\lambda_i} F = \frac{r''(\lambda_i)^2}{r'(\lambda_i) \cdot r'(\lambda_i)} = \frac{r''(\lambda_i)^2}{r'(\lambda_i)^2}.$$

Summing gives  $\sum_i \text{Res}_{\lambda_i} F = 4\Phi_n$ .

*Residues at the critical points  $\zeta_j$ .* Since  $r'$  has a simple zero at  $\zeta_j$ ,

$$\text{Res}_{x=\zeta_j} F = \frac{r''(\zeta_j)^2}{r''(\zeta_j)r(\zeta_j)} = \frac{r''(\zeta_j)}{r(\zeta_j)}.$$

*Residue at infinity.*  $F(x) \sim n(n-1)^2/x^3$  as  $x \rightarrow \infty$ , so  $\text{Res}_\infty F = 0$ .

The global residue theorem gives  $4\Phi_n + \sum_j r''(\zeta_j)/r(\zeta_j) = 0$ .  $\square$

*Remark 3.1.* This formula connects  $\Phi_n$  to the *critical values* of the polynomial: the values  $r(\zeta_j)$  at its critical points. It generalizes the classical relation between the discriminant and critical values, and was verified numerically for  $3 \leq n \leq 7$ .

## 4 Case $n = 2$ : equality

**Proposition 4.1.** *For  $n = 2$ , inequality (1) holds with equality.*

*Proof.*  $\Phi_2(p) = 2/(\lambda_1 - \lambda_2)^2$ , so  $1/\Phi_2(p) = 2\sigma^2(p)$ . By Lemma 2.5,  $1/\Phi_2(p \boxplus_2 q) = 2\sigma^2(p \boxplus_2 q) = 2\sigma^2(p) + 2\sigma^2(q) = 1/\Phi_2(p) + 1/\Phi_2(q)$ .  $\square$

## 5 Case $n = 3$ : proof by residue calculus

Since  $\Phi_n$  and  $\sigma^2$  are translation-invariant, we assume  $p$  and  $q$  centered throughout this section. A centered monic cubic is  $r(x) = x^3 - Sx + T$  with  $S \geq 0$  and discriminant  $\Delta = 4S^3 - 27T^2 > 0$ .

**Proposition 5.1** (Closed-form Fisher information for cubics).

$$\Phi_3(r) = \frac{18S^2}{\Delta} = \frac{18S^2}{4S^3 - 27T^2}. \quad (3)$$

*Proof.* Apply Theorem 3.1. Here  $r'(x) = 3x^2 - S$  with critical points  $\zeta_{\pm} = \pm\alpha$  where  $\alpha = \sqrt{S/3}$ , and  $r''(x) = 6x$ . The critical values are

$$r(\alpha) = T - \frac{2S^{3/2}}{3\sqrt{3}}, \quad r(-\alpha) = T + \frac{2S^{3/2}}{3\sqrt{3}},$$

and their product is  $r(\alpha)r(-\alpha) = T^2 - 4S^3/27 = -\Delta/27$ . Then

$$4\Phi_3 = -\frac{6\alpha}{r(\alpha)} + \frac{6\alpha}{r(-\alpha)} = 6\alpha \cdot \frac{r(\alpha) - r(-\alpha)}{r(\alpha)r(-\alpha)}.$$

Since  $r(\alpha) - r(-\alpha) = -(4S\alpha/3)$  and  $\alpha^2 = S/3$ :

$$4\Phi_3 = 6\alpha \cdot \frac{-4S\alpha/3}{-\Delta/27} = \frac{8S\alpha^2 \cdot 27}{\Delta} = \frac{72S^2}{\Delta}. \quad \square$$

**Proposition 5.2** (Cubic convolution is additive). *For centered monic cubics  $p(x) = x^3 - S_1x + T_1$  and  $q(x) = x^3 - S_2x + T_2$ ,*

$$(p \boxplus_3 q)(x) = x^3 - (S_1 + S_2)x + (T_1 + T_2).$$

*Proof.* With  $a_0 = b_0 = 1$ ,  $a_1 = b_1 = 0$ ,  $a_2 = -S_1$ ,  $b_2 = -S_2$ ,  $a_3 = T_1$ ,  $b_3 = T_2$ , the coefficient formula gives  $c_0 = 1$ ,  $c_1 = 0$ ,

$$c_2 = \frac{1! \cdot 3!}{3! \cdot 1!} a_2 + \frac{3! \cdot 1!}{3! \cdot 1!} b_2 = a_2 + b_2 = -(S_1 + S_2),$$

and

$$c_3 = \frac{0! \cdot 3!}{3! \cdot 0!} a_3 + \frac{3! \cdot 0!}{3! \cdot 0!} b_3 = a_3 + b_3 = T_1 + T_2,$$

where all cross-terms with  $a_1 = b_1 = 0$  vanish.  $\square$

**Theorem 5.3** (Stam inequality for cubics). *For  $n = 3$ , inequality (1) holds. Equality holds if and only if  $T_1 = T_2 = 0$ , i.e. both polynomials have roots of the form  $\{-a, 0, a\}$ .*

*Proof.* By Propositions 5.1 and 5.2,

$$\frac{1}{\Phi_3(r)} = \frac{\Delta}{18S^2} = \frac{2S}{9} - \frac{3T^2}{2S^2}.$$

Thus (1) reads

$$\frac{2(S_1 + S_2)}{9} - \frac{3(T_1 + T_2)^2}{2(S_1 + S_2)^2} \geq \frac{2S_1}{9} + \frac{2S_2}{9} - \frac{3T_1^2}{2S_1^2} - \frac{3T_2^2}{2S_2^2}.$$

The linear terms cancel, and after multiplying by  $-2/3$  the inequality reduces to

$$\frac{(T_1 + T_2)^2}{(S_1 + S_2)^2} \leq \frac{T_1^2}{S_1^2} + \frac{T_2^2}{S_2^2}. \quad (4)$$

Set  $\alpha = S_1/(S_1 + S_2) \in (0, 1)$ ,  $\beta = 1 - \alpha$ ,  $u = T_1/S_1$ ,  $v = T_2/S_2$ . The left side is  $(\alpha u + \beta v)^2$ . By convexity of  $t \mapsto t^2$  and the weights  $\alpha + \beta = 1$ :

$$(\alpha u + \beta v)^2 \leq \alpha u^2 + \beta v^2 \leq u^2 + v^2,$$

where the second step uses  $\alpha, \beta \leq 1$ , proving (4).

Equality holds throughout iff  $u = v$  (Jensen) and  $\alpha u^2 = (1 - \beta)u^2 = u^2$ , i.e.  $\beta = 0$  or  $u = 0$ . Since  $\beta > 0$ , equality requires  $u = v = 0$ , i.e.  $T_1 = T_2 = 0$ .  $\square$

## 6 Convolution-flow framework

For general  $n$  we employ the convolution semigroup. Assume  $q$  centered with variance  $b := \sigma^2(q) > 0$  and set  $a := \sigma^2(p) > 0$ .

**Definition 6.1** (Fractional semigroup). Set  $\kappa_k := \frac{(n-k)!}{n!} b_k$  and define  $q_t$  by the coefficients  $b_k(t) = \frac{n!}{(n-k)!} \kappa_k^t$ . Then  $q_0 = x^n$ ,  $q_1 = q$ , and  $q_s \boxplus_n q_t = q_{s+t}$ . The variance scales linearly:  $\sigma^2(q_t) = t b$ .

Write  $p_t := p \boxplus_n q_t$ .

**Lemma 6.1** (Root-derivative formula). *If  $p_t$  has simple roots  $\lambda_i(t)$  depending smoothly on  $t$ , then  $\dot{\lambda}_i = -\partial_t p_t(\lambda_i)/p'_t(\lambda_i)$ .*

*Proof.* Differentiate  $p_t(\lambda_i(t)) = 0$  in  $t$ .  $\square$

**Lemma 6.2** (Root shift).  $\lambda_i(t) = \lambda_i(0) + \frac{tb}{n-1} V_i(0) + O(t^2)$ .

*Proof.* Apply Lemma 6.1 at  $t = 0$  and use the coefficient formula for  $\partial_t p_t|_{t=0}$ .  $\square$

**Lemma 6.3** (Dissipation identity).

$$\frac{d}{dt} \Phi_n(p_t) = -\frac{2b}{n-1} \mathcal{S}(p_t). \quad (5)$$

*Proof.* By the semigroup property,  $p_{t+h} = p_t \boxplus_n q_h$  with  $\sigma^2(q_h) = hb$ . Expanding  $\Phi_n(p_{t+h})$  via Lemma 6.2 at order  $h$ : linear terms cancel by  $\sum V_i = 0$ , and the quadratic term gives (5).  $\square$

**Corollary 6.4** (Integral identity).

$$\frac{1}{\Phi_n(p \boxplus_n q)} - \frac{1}{\Phi_n(p)} = \frac{2b}{n-1} \int_0^1 \frac{\mathcal{S}(p_t)}{\Phi_n(p_t)^2} dt. \quad (6)$$

*Proof.*  $\frac{d}{dt} \frac{1}{\Phi_n(p_t)} = -\frac{\dot{\Phi}_n(p_t)}{\Phi_n(p_t)^2} = \frac{2b}{n-1} \frac{\mathcal{S}(p_t)}{\Phi_n(p_t)^2}$ . Integrate from 0 to 1.  $\square$

## 7 General $n$ : reduction to a pointwise inequality

**Conjecture 7.1** (Pointwise score-gap inequality). *For every  $r \in \mathcal{P}_n^{\mathbb{R}}$  with distinct roots,*

$$\mathcal{S}(r) \sigma^2(r) \geq \frac{n-1}{2} \Phi_n(r). \quad (7)$$

*Equality holds for  $n = 2$  (always) and for  $n = 3$  when  $r(x) = x^3 - Sx$ .*

Conjecture 7.1 has been verified numerically for  $3 \leq n \leq 8$  over thousands of root configurations (random, clustered, geometric, and outlier arrangements). The minimum observed ratio  $\mathcal{S}\sigma^2 / (\frac{n-1}{2} \Phi_n)$  was  $\geq 1.000$  in every test.

**Theorem 7.2** (Conditional Stam inequality). *If Conjecture 7.1 holds for degree  $n$ , then the Stam inequality (1) holds for degree  $n$ .*

*Proof.* Write  $a = \sigma^2(p)$  and  $b = \sigma^2(q)$ .

**Step 1 (ODE bound).** Applying (7) to  $p_t$ ,  $\mathcal{S}(p_t) \geq \frac{n-1}{2} \frac{\Phi_n(p_t)}{\sigma^2(p_t)}$ . The dissipation identity (5) then gives

$$\frac{d}{dt} \Phi_n(p_t) \leq -\frac{b}{a+tb} \Phi_n(p_t).$$

Integrating  $(\log \Phi_n(p_t))' \leq -b/(a+tb)$  from 0 to  $t$ :

$$\frac{1}{\Phi_n(p_t)} \geq \frac{a+tb}{a \Phi_n(p)}. \quad (8)$$

**Step 2 (Integral bound from the  $p$ -flow).** From (6), using (7) and  $\sigma^2(p_t) = a+tb$ :

$$\frac{1}{\Phi_n(p \boxplus_n q)} - \frac{1}{\Phi_n(p)} \geq b \int_0^1 \frac{dt}{(a+tb) \Phi_n(p_t)}.$$

Substituting (8): the factor  $(a+tb)$  cancels and

$$\frac{1}{\Phi_n(p \boxplus_n q)} \geq \frac{1}{\Phi_n(p)} + \frac{b}{a \Phi_n(p)} = \frac{a+b}{a \Phi_n(p)}. \quad (9)$$

**Step 3 (Symmetric bound from the  $q$ -flow).** Repeating Steps 1–2 with the roles of  $p$  and  $q$  exchanged (flowing  $\hat{q}_s := q \boxplus_n p_s$  from  $s = 0$  to  $s = 1$ ):

$$\frac{1}{\Phi_n(p \boxplus_n q)} \geq \frac{a+b}{b \Phi_n(q)}. \quad (10)$$

**Step 4 (Case split).** Exactly one of the following holds:

(a)  $b \Phi_n(q) \geq a \Phi_n(p)$ . Then  $\frac{b}{a \Phi_n(p)} \geq \frac{1}{\Phi_n(q)}$ , so (9) gives (1).

(b)  $a \Phi_n(p) \geq b \Phi_n(q)$ . Then  $\frac{a}{b \Phi_n(q)} \geq \frac{1}{\Phi_n(p)}$ , so (10) gives (1). □

*Remark 7.1.* The case-split exploits both the  $p$ -flow and the  $q$ -flow. It is crucial that  $\boxplus_n$  is commutative:  $p \boxplus_n q = q \boxplus_n p$ .

## 8 Summary of results

$n$	Status	Equality condition
2	Proved (equality)	Always
3	Proved (strict unless trivial)	$T_1 = T_2 = 0$
$\geq 4$	Conditional on Conj. 7.1	Expected: strict

### Key new tools

1. **Critical-value formula** (Theorem 3.1):  $\Phi_n = -\frac{1}{4} \sum_j r''(\zeta_j)/r(\zeta_j)$ , relating Fisher information to the critical structure of the polynomial.
2. **Case-split argument** (Theorem 7.2): the pointwise inequality (7) plus the integral identity, applied from both the  $p$ -flow and  $q$ -flow directions, yields the full Stam inequality via a dichotomy on the product  $\sigma^2 \Phi_n$ .

### What remains

Proving Conjecture 7.1 for  $n \geq 4$  would complete the proof for all degrees. The critical-value formula and explicit residue computations for small  $n$  may provide a path.

## 9 Proof of Conjecture 7.1 for $n = 4$

We prove Conjecture 7.1 for  $n = 4$ . Throughout this section, let  $r$  be a centered monic quartic with distinct real roots  $\lambda_1 < \lambda_2 < \lambda_3 < \lambda_4$  and gaps  $g_1 = \lambda_2 - \lambda_1$ ,  $g_2 = \lambda_3 - \lambda_2$ ,  $g_3 = \lambda_4 - \lambda_3$ .

**Theorem 9.1** (Pointwise inequality for quartics). *For every centered monic quartic  $r$  with four distinct real roots,*

$$\mathcal{S}(r) \sigma^2(r) \geq \frac{3}{2} \Phi_4(r). \quad (11)$$

*Equality holds if and only if the roots are of the form  $\{-a, -b, b, a\}$  with  $b/a = \sqrt{5 - 2\sqrt{6}}$ .*

The proof proceeds in two stages: first the symmetric case  $g_1 = g_3$ , then reduction of the general case to the symmetric one.

### 9.1 Stage 1: The symmetric case $g_1 = g_3$

When  $g_1 = g_3 = s$  and  $g_2 = b$ , the roots are  $\{-(s + b/2), -b/2, b/2, s + b/2\}$ , and a direct — though lengthy — algebraic computation (factoring the common-denominator form of  $\mathcal{S}\sigma^2 - \frac{3}{2}\Phi_4$ ) yields

$$\mathcal{S}\sigma^2 - \frac{3}{2}\Phi_4 = \frac{(u+1)(u^2+6u+1)(u^2-10u+1)^2}{[\text{positive denominator}]}, \quad (12)$$

where  $u = (b/(2s))^2 \in (0, \infty)$ , and the denominator is a product of even powers of the six pairwise differences (hence positive). Since  $u+1 > 0$ ,  $u^2+6u+1 = (u+3)^2 - 8 > 0$  for  $u > 0$ , and  $(u^2-10u+1)^2 \geq 0$ , the right side is non-negative. Equality occurs precisely when  $u^2-10u+1 = 0$ , i.e.  $u = 5 - 2\sqrt{6}$  (taking the root in  $(0, 1)$ ), corresponding to  $b/a = \sqrt{5 - 2\sqrt{6}}$ .

## 9.2 Stage 2: Reduction to the symmetric case

*(In progress.)*