

Worksheet 4: Local illumination – Gouraud and Phong shading

Reading	Angel: Sections 5.8 and 6-6.10.
Purpose	The purpose of this set of exercises is to get acquainted with lighting and shading in WebGL and GLSL. We compute the local illumination of an object based on ambient, diffuse and specular material properties, and a light source.
Part 1	<p>Draw a sphere in perspective view.</p> <ul style="list-style-type: none"> Start from Part 2 of Worksheet 3, which draws three wireframe cubes in perspective view. Simplify such that you draw just one cube in the image center and switch to drawing triangles instead of wireframe. [Angel 2.4.2, 4.6, 5.3, 5.6] Draw a unit sphere instead of a unit cube using recursive subdivision of a tetrahedron. [Angel 6.6] Insert two buttons: one which increments the subdivision level and one which decrements the subdivision level. [Angel 3.6.2]
Part 2	<p>Use depth buffer and back face culling to remove hidden surfaces.</p> <ul style="list-style-type: none"> Draw the vertex positions as colors ($\mathbf{c} = 0.5 \cdot \mathbf{p} + 0.5$). [Part 3 of Worksheet 1, Angel 2.10] Use the depth buffer to ensure that you are looking at the nearest part of the surface of the sphere. [Angel 2.10.4, 5.8] Enable back face culling to improve efficiency. [Angel 5.8] <p>If your result is different from the previous step, you should flip the winding order of your triangles. With the subdivided tetrahedron, you can flip the winding order by flipping the sign of the z-coordinate in each of the four initial vertices.</p>
Part 3	<p>Use Gouraud shading (with true normals) to draw a diffuse sphere lit by a distant, white, directional light with direction $(0, 0, -1)$.</p> <ul style="list-style-type: none"> Obtain the surface normal in the vertex shader. [Angel 6.9] Think of the color of the sphere as its diffuse reflection coefficient k_d. Introduce a distant light with direction $\mathbf{l}_e = (0, 0, -1)$ and light emission $L_d = L_e = (1, 1, 1)$, no distance attenuation. Compute the diffusely reflected light in the vertex shader and set the vertex color to this result (note that $\mathbf{l} = -\mathbf{l}_e$). [Angel 6.3.2, 6.7.1] Let the camera orbit the sphere over time. [Angel 3.1]

Worksheet 4: Local illumination – Gouraud and Phong shading

Part 4	<p>Implement the full Phong reflection model in the vertex shader and create sliders for material parameters (k_a, k_d, k_s, α) and light emission (L_e). [Angel 3.6.5, 6.3, 6.7-6.8]</p> <p>Use the “input” event instead of the onchange function if you prefer immediate response while using a slider (instead of only getting response when the slider is released).</p> <p>Please use just a single slider for each parameter and the same light emission for all lighting terms ($L_a = L_d = L_s = L_e$). This means that your solution can be significantly simpler than the example in the textbook.</p> <p>[Section 6.8 of the textbook has many minor glitches. Please consult the Wiki page with textbook clarifications before using the code example in Section 6.8.]</p> <p>If your final hand-in is not a webpage, create a 5x5 table of sphere images where the value of only one material parameter changes in each row. Let the type of material parameter change from row to row. Start from the upper row and work your way downwards successively changing one of the parameters k_a, k_d, k_s, α, and L_e while keeping the others constant.</p>
Part 5	<p>Use Phong shading by moving your implementation of the Phong reflection model to the fragment shader and varying positions and normals across triangles instead of colors. [Angel 6.5.3, 6.10]</p> <p>Is Gouraud or Phong shading the best method for simulating highlight? Explain.</p>
Part 6	<p>Answer the following questions: [Angel 5.2, 6.2, 6.3, 6.5, 6.8]</p> <ol style="list-style-type: none"> What is the difference between Phong shading and Phong lighting (the Phong reflection model)? What is the difference between flat shading, Gouraud shading, and Phong shading? List pros and cons of each. What is the difference between a directional light and a point light? Does the eye position influence the shading of an object in any way? What is the effect of setting the specular term to $(0, 0, 0)$? What is the effect of increasing the shininess exponent (α)? In what coordinate space did you compute the lighting?