

Liver Lesion Comparison Report

High-resolution 3D segmentation and
lesion-by-lesion CT comparison

BASELINE CT

25 Dec 2025

PET/CT diagnostic CT

FOLLOW-UP CT

19 Jan 2026

Diagnostic contrast CT

CANDIDATES

16 + 1


Matched plus uncertain

LATEST BURDEN

10.22%

170.34 mL tumor volume

Important: This document is generated from automated segmentation. It is not a radiologist-signed diagnosis, formal RECIST report, or surgical-planning document.



Overall disease burden appears broadly stable

TUMOR VOLUME

+2.1%

166.82 → 170.34 mL

TUMOR BURDEN

+0.44 pp

9.78% → 10.22%

LARGEST DIAMETER

+0.4%

64.3 → 64.5 mm

MATCHED LESIONS

16

One additional uncertain focus

Key interpretation

01

Dominant segment 3 mass

Stable axial diameter; automated volume increased 10.2%. The official report described minimal growth from approximately 62 to 64 mm.

02

Segment 8 mass

Stable by diameter, consistent with the official radiology report.

03


Remaining lesions

Predominantly stable. Volume percentages for small lesions are highly contour-sensitive.

04

Low attenuation

No reliable evidence of extensive macroscopic necrosis. Contrast timing differs between scans.



Low attenuation is not synonymous with necrosis. Normal-liver median attenuation differed substantially between scans (137 HU versus 117 HU), demonstrating non-equivalent contrast conditions.

L01


Liver segment 3

BROADLY STABLE

Lesion L01 — automated CT comparison Volume change +10.2% | Long-axis change +0.4%


December 25, 2025

S3 | Inst 425
64.3 × 49.6 × 60.0 mm
Volume 96.36 mL
Median 82 HU | <40 HU 1.4%



January 19, 2026

S3 | Inst 185
64.5 × 51.3 × 66.0 mm
Volume 106.21 mL
Median 68 HU | <40 HU 5.4%



Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

64.3 → 64.5 mm

Change +0.4%

3D VOLUME

96.36 → 106.21 mL

Change +10.2%

CRANIOCAUDAL

60.0 → 66.0 mm

Third spatial dimension

CORE BELOW 40 HU

1.4% → 5.4%

Low-attenuation proxy only



Assessment: Axial diameter is broadly stable.

L02

Liver segment 8

BROADLY STABLE

Lesion L02 — automated CT comparison Volume change -6.7% | Long-axis change -2.0%



Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

46.0 → 45.0 mm

Change -2.0%

3D VOLUME

44.81 → 41.81 mL

Change -6.7%

CRANIOCAUDAL

56.2 → 55.5 mm

Third spatial dimension

CORE BELOW 40 HU

0.5% → 0.4%

Low-attenuation proxy only

Assessment: Axial diameter is broadly stable.


L03

Liver segment 7


BROADLY STABLE

Lesion L03 — automated CT comparison Volume change -1.9% | Long-axis change -3.1%

December 25, 2025



January 19, 2026



Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

29.4 → 28.5 mm

Change -3.1%

3D VOLUME

10.03 → 9.83 mL

Change -1.9%

CRANIOCAUDAL

30.0 → 30.0 mm

Third spatial dimension

CORE BELOW 40 HU

0.1% → 0.0%

Low-attenuation proxy only



Assessment: Axial diameter is broadly stable.

L04

Liver segment 4

BROADLY STABLE

Lesion L04 — automated CT comparison Volume change +0.4% | Long-axis change -2.3%



Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

23.5 → 22.9 mm

Change -2.3%

3D VOLUME

2.36 → 2.37 mL

Change +0.4%

CRANIOCAUDAL

20.0 → 21.0 mm

Third spatial dimension

CORE BELOW 40 HU

0.0% → 0.3%

Low-attenuation proxy only



Assessment: Axial diameter is broadly stable.

L05

Liver segment 8

BROADLY STABLE

Lesion L05 — automated CT comparison Volume change -17.0% | Long-axis change +3.5%



Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

17.7 → 18.3 mm

Change +3.5%

3D VOLUME

2.28 → 1.89 mL

Change -17.0%

CRANIOCAUDAL

15.0 → 15.0 mm

Third spatial dimension

CORE BELOW 40 HU

0.0% → 0.0%

Low-attenuation proxy only

Assessment: Axial diameter is broadly stable. The larger volume percentage is likely contour-sensitive.


L06

Liver segment 7


BROADLY STABLE

Lesion L06 — automated CT comparison Volume change +5.7% | Long-axis change +1.4%

December 25, 2025



January 19, 2026



Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

15.9 → 16.1 mm

Change +1.4%

3D VOLUME

1.76 → 1.85 mL

Change +5.7%

CRANIOCAUDAL

16.2 → 16.5 mm

Third spatial dimension

CORE BELOW 40 HU

1.2% → 0.0%

Low-attenuation proxy only

Assessment: Axial diameter is broadly stable.

L07


Liver segment 6

BROADLY STABLE

Lesion L07 — automated CT comparison
Volume change +28.2% | Long-axis change +1.7%


December 25, 2025

S6 | Inst 494
14.9 × 12.0 × 13.8 mm
Volume 1.08 mL
Median 104 HU | <40 HU 0.0%



January 19, 2026

S6 | Inst 248
15.1 × 12.5 × 15.0 mm
Volume 1.38 mL
Median 86 HU | <40 HU 0.0%



Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

14.9 → 15.1 mm

Change +1.7%

3D VOLUME

1.07 → 1.38 mL

Change +28.2%

CRANIOCAUDAL

13.8 → 15.0 mm

Third spatial dimension

CORE BELOW 40 HU

0.0% → 0.0%

Low-attenuation proxy only

Assessment: Axial diameter is broadly stable. The larger volume percentage is likely contour-sensitive.

L08


Liver segment 5

BROADLY STABLE

Lesion L08 — automated CT comparison Volume change -0.6% | Long-axis change -4.8%


December 25, 2025

S5 | Inst 441
12.9 × 12.1 × 12.5 mm
Volume 0.96 mL
Median 95 HU | <40 HU 0.0%



January 19, 2026

S5 | Inst 204
12.3 × 12.0 × 13.5 mm
Volume 0.96 mL
Median 70 HU | <40 HU 0.0%



Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

12.9 → 12.3 mm

Change -4.8%

3D VOLUME

0.96 → 0.96 mL

Change -0.6%

CRANIOCAUDAL

12.5 → 13.5 mm

Third spatial dimension

CORE BELOW 40 HU

0.0% → 0.0%

Low-attenuation proxy only



Assessment: Axial diameter is broadly stable.

L09

Liver segment 5

BROADLY STABLE

Lesion L09 — automated CT comparison Volume change -2.8% | Long-axis change -2.3%



Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

13.2 → 12.9 mm

Change -2.3%

3D VOLUME

0.90 → 0.88 mL

Change -2.8%

CRANIOCAUDAL

12.5 → 13.5 mm

Third spatial dimension

CORE BELOW 40 HU

0.0% → 0.0%

Low-attenuation proxy only

Assessment: Axial diameter is broadly stable.

L10

Liver segment 7

BROADLY STABLE

Lesion L10 — automated CT comparison Volume change +13.5% | Long-axis change +0.8%

December 25, 2025

S7 | Inst 405
11.9 × 9.7 × 11.2 mm
Volume 0.62 mL
Median 93 HU | <40 HU 0.0%

R

L

January 19, 2026

S7 | Inst 181
12.0 × 11.0 × 12.0 mm
Volume 0.71 mL
Median 76 HU | <40 HU 0.0%

R

L

Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

11.9 → 12.0 mm

Change +0.8%

3D VOLUME

0.62 → 0.71 mL

Change +13.5%

CRANIOCAUDAL

11.2 → 12.0 mm

Third spatial dimension

CORE BELOW 40 HU

0.0% → 0.0%

Low-attenuation proxy only

Assessment: Axial diameter is broadly stable.

L11

Liver segment 6

BROADLY STABLE

Lesion L11 — automated CT comparison Volume change +10.5% | Long-axis change +4.0%

December 25, 2025

S6 | Inst 451
11.2 × 9.0 × 12.5 mm
Volume 0.62 mL
Median 90 HU | <40 HU 0.0%

R

January 19, 2026

S6 | Inst 215
11.6 × 10.1 × 12.0 mm
Volume 0.68 mL
Median 74 HU | <40 HU 0.0%

R

L

Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

11.2 → 11.6 mm

Change +4.0%

3D VOLUME

0.62 → 0.68 mL

Change +10.5%

CRANIOCAUDAL

12.5 → 12.0 mm

Third spatial dimension

CORE BELOW 40 HU

0.0% → 0.0%

Low-attenuation proxy only


Assessment: Axial diameter is broadly stable.

L12

Liver segment 8

MINOR MEASURED DECREASE

Lesion L12 — automated CT comparison Volume change -6.7% | Long-axis change -8.3%



Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

12.4 → 11.4 mm

Change -8.3%

3D VOLUME

0.67 → 0.62 mL

Change -6.7%

CRANIOCAUDAL

11.2 → 9.0 mm

Third spatial dimension

CORE BELOW 40 HU

0.7% → 0.0%

Low-attenuation proxy only

Assessment: A measured decrease is present, but protocol and contour variability limit certainty.

L13


Liver segment 4/5

UNRELIABLE COMPARISON

Lesion L13 — automated CT comparison Volume change -86.1% | Long-axis change -41.7%


December 25, 2025

S4 | Inst 463
20.2 × 16.8 × 23.8 mm
Volume 3.46 mL
Median 88 HU | <40 HU 2.0%



January 19, 2026

S5 | Inst 215
11.8 × 11.0 × 9.0 mm
Volume 0.48 mL
Median 71 HU | <40 HU 2.4%



Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

20.2 → 11.8 mm

Change -41.7%

3D VOLUME

3.46 → 0.48 mL

Change -86.1%

CRANIOCAUDAL

23.8 → 9.0 mm

Third spatial dimension

CORE BELOW 40 HU

2.0% → 2.4%

Low-attenuation proxy only

Assessment: The apparent reduction conflicts with the overall imaging report and is most likely caused by segmentation or matching inconsistency.

L14


Liver segment 6

SUB-CENTIMETER

Lesion L14 — automated CT comparison Volume change +25.4% | Long-axis change -1.5%


December 25, 2025

S6 | Inst 446
8.0 × 8.0 × 5.0 mm
Volume 0.19 mL
Median 111 HU | <40 HU 0.0%



January 19, 2026

S6 | Inst 211
7.9 × 7.3 × 7.5 mm
Volume 0.23 mL
Median 84 HU | <40 HU 0.0%



Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

8.0 → 7.9 mm

Change -1.5%

3D VOLUME

0.19 → 0.23 mL

Change +25.4%

CRANIOCAUDAL

5.0 → 7.5 mm

Third spatial dimension

CORE BELOW 40 HU

0.0% → 0.0%

Low-attenuation proxy only


Assessment: Small boundary shifts strongly affect volume. Diameter is more dependable than percentage volume change.

L15

Liver segment 2

SUB-CENTIMETER

Lesion L15 — automated CT comparison Volume change -21.9% | Long-axis change -11.9%



Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

9.4 → 8.3 mm

Change -11.9%

3D VOLUME

0.30 → 0.23 mL

Change -21.9%

CRANIOCAUDAL

8.8 → 7.5 mm

Third spatial dimension

CORE BELOW 40 HU

9.4% → 0.3%

Low-attenuation proxy only

Assessment: Small boundary shifts strongly affect volume. Diameter is more dependable than percentage volume change.

L16


Liver segment 7

SUB-CENTIMETER

Lesion L16 — automated CT comparison Volume change -2.9% | Long-axis change +2.2%


December 25, 2025

S7 | Inst 365
7.5 × 7.1 × 7.5 mm
Volume 0.18 mL
Median 82 HU | <40 HU 0.6%



January 19, 2026

S7 | Inst 149
7.7 × 6.7 × 7.5 mm
Volume 0.18 mL
Median 82 HU | <40 HU 0.0%



Yellow: automated lesion boundary | Red: axial long axis | Cyan: perpendicular axis | Review required

AXIAL LONG AXIS

7.5 → 7.7 mm

Change +2.2%

3D VOLUME

0.18 → 0.18 mL

Change -2.9%

CRANIOCAUDAL

7.5 → 7.5 mm

Third spatial dimension

CORE BELOW 40 HU

0.6% → 0.0%

Low-attenuation proxy only

Assessment: Small boundary shifts strongly affect volume. Diameter is more dependable than percentage volume change.

U01


Liver segment 8

UNMATCHED CANDIDATE

Old-only candidate U01 — uncertain sub-centimeter focus

December 25, 2025

S8 | Inst 452
8.4 × 6.2 × 10.0 mm
Volume 0.24 mL
Median 109 HU | <40 HU 0.0%



No distinct corresponding automated component on January CT

AXIAL LONG AXIS

8.4 → — mm

Change —

3D VOLUME

0.24 → — mL

Change —

CRANIOCAUDAL

10.0 → — mm

Third spatial dimension

CORE BELOW 40 HU

0.0% → —%

Low-attenuation proxy only

Assessment: No separate January component was identified. This may reflect merging, non-detection or artifact.

Consolidated lesion measurements

ID	SEG	DEC LONG	JAN LONG	LONG Δ	DEC VOL	JAN VOL	VOL Δ	JAN <40 HU
L01	3	64.3	64.5	+0.4%	96.36	106.21	+10.2%	5.4%
L02	8	46.0	45.0	-2.0%	44.81	41.81	-6.7%	0.4%
L03	7	29.4	28.5	-3.1%	10.03	9.83	-1.9%	0.0%
L04	4	23.5	22.9	-2.3%	2.36	2.37	+0.4%	0.3%
L05	8	17.7	18.3	+3.5%	2.28	1.89	-17.0%	0.0%
L06	7	15.9	16.1	+1.4%	1.76	1.85	+5.7%	0.0%
L07	6	14.9	15.1	+1.7%	1.07	1.38	+28.2%	0.0%
L08	5	12.9	12.3	-4.8%	0.96	0.96	-0.6%	0.0%
L09	5	13.2	12.9	-2.3%	0.90	0.88	-2.8%	0.0%
L10	7	11.9	12.0	+0.8%	0.62	0.71	+13.5%	0.0%
L11	6	11.2	11.6	+4.0%	0.62	0.68	+10.5%	0.0%
L12	8	12.4	11.4	-8.3%	0.67	0.62	-6.7%	0.0%
L13	4/5	20.2	11.8	-41.7%	3.46	0.48	-86.1%	2.4%
L14	6	8.0	7.9	-1.5%	0.19	0.23	+25.4%	0.0%
L15	2	9.4	8.3	-11.9%	0.30	0.23	-21.9%	0.3%
L16	7	7.5	7.7	+2.2%	0.18	0.18	-2.9%	0.0%
U01	8	8.4	—	—	0.24	—	—	—%

Volume change is substantially less reproducible than long-axis change for small lesions. U01 was identified only as a separate component on the older scan.

How to interpret this report

Source examinations

25 Dec 2025 PET/CT diagnostic contrast CT, series 3 (approximately $0.98 \times 0.98 \times 1.25$ mm), compared with 19 Jan 2026 contrast CT, series 2 (approximately $0.70 \times 0.70 \times 1.5$ mm).

Segmentation

TotalSegmentator liver-lesion and liver-segment models were used. Candidate components of at least 0.05 mL were measured and matched by segment and normalized intrahepatic position.

Dimensions and volume

Long and perpendicular axes were calculated on the largest axial cross-section. Craniocaudal extent was measured across slices. Volumes depend directly on contour accuracy.

Necrosis limitation

The percentage below 40 HU is a low-attenuation proxy only. True viable-versus-necrotic assessment requires suitable multiphase imaging, consistent acquisition, expert review and sometimes pathology.

Response assessment

This is not a formal RECIST, mRECIST or volumetric response classification. Target selection and non-target disease assessment must be performed by the treating team.

Clinical safety

Do not use this automated document alone to start, stop or alter treatment. Provide the original DICOM studies and this package to a qualified radiologist for correction and sign-off.