

You are an expert neuroimaging researcher and academic writer. I am writing my Master's thesis in Neuroscience investigating the effects of ketamine-augmented Prolonged Exposure (PE) therapy for PTSD.

My project uses resting-state fMRI (rs-fMRI) across 3 timepoints: Baseline (T1), Day 3 post-ketamine (T2), and Month 1 post-treatment (T3), utilizing different dosing cohorts.

I need to integrate two distinct but complementary analytical dimensions into the thesis introduction, methods, and discussion. Please use the following detailed summary of my framework to help draft and refine the manuscript:

DIMENSION 1: Baseline Prognostic Biomarker & The "Neuroplastic Rescue" (Core Thesis)

- Focus Axis: Dynamic functional connectivity (dFC) variance between the Salience Network (SN) and Default Mode Network (DMN), calculated via a 60-second sliding window.
- Clinical Framing: Shifting from a diagnostic framing to a prognostic framing. Baseline SN-DMN dFC-variance serves as a pre-treatment quantitative predictor of PE treatment outcomes (measured by change in CAPS-5 scores).
- The Decoupling Effect: In the placebo group, high baseline network instability predicts PE non-response (treatment failure). Crucially, ketamine "uncouples" this risk, rescuing individuals with high-risk neural phenotypes so they can successfully benefit from exposure therapy.
- Methodological Rigor: The pipeline explicitly identifies and corrects for Oldham-type artifacts (mathematical coupling inherent when correlating baseline values with subsequent longitudinal changes).

DIMENSION 2: Transient Entropic/Flexibility States (Additions based on PI Feedback)

- Rationale: Ketamine induces a transient window of heightened neural plasticity/flexibility that peaks around Day 3 (coinciding with our T2 scan), which chemically optimizes the brain for behavioral extinction learning during PE.
- Signal-Level Complexity: Incorporating Sample Entropy (SampEn) / Approximate Entropy (ApEn) and Lempel-Ziv Complexity (LZC) of the BOLD signal to quantify the transition from a "rigid/trauma-locked" state to a highly complex, "entropic" state.
- Network-Level Flexibility: Utilizing multilayer modularity (graph theory community detection over time) to calculate "Network Flexibility"—specifically looking at how frequently regional nodes switch network allegiances (e.g., transitioning fluently between DMN and SN).
- dFC State Metrics: Extracting Fractional Occupancy, Dwell Time, and State Transition Entropy via K-means clustering of sliding windows to track fluid transitions vs. "stuck" clinical states.

Statistical Framework to Implement:

1. Linear Mixed-Effects Models (LMM): To model the longitudinal data (T1, T2, T3) across dosing groups, handling missing data cleanly, with fixed effects for Time, Dosing, and the Time x Dosing interaction.
2. Mediation/Prediction Analysis: Testing the ultimate mechanistic hypothesis: whether the transient spike in brain entropy/flexibility at Day 3 (T2) mathematically mediates the relationship between Ketamine Dosing and the long-term clinical reduction of PTSD symptoms at Month 1 (T3).

Please keep this framework in mind for all subsequent drafting tasks.

Ketamine-augmented PE, dynamic connectivity, and entropy/flexibility in PTSD

Your framework aligns well with emerging work on dynamic connectivity, signal complexity, and ketamine-related plasticity. Below is an integrated, literature-grounded scaffold you can adapt for your Introduction/Methods/Discussion, keeping Dimension 1 as the core thesis and Dimension 2 as a mechanistic extension.

Mechanisms of Action: Ketamine, Plasticity Windows, and Network Dynamics

Ketamine and related antidepressants induce rapid and sustained neuroplastic changes, partly via BDNF/TrkB and p75NTR signaling and glutamate-dependent synaptogenesis in prefrontal–hippocampal circuits, supporting extinction-memory and cortical reorganization (■ Diniz et al. 2024; ■ Pardossi et al. 2024; Brown et al. 2026). Human 7T rs-fMRI shows ketamine-related BDNF increases are coupled to alterations in prefrontal–DMN connectivity, interpreted as plasticity-related RSFC reconfiguration (■ Woelfer et al. 2020, 696 - 710).

In PTSD, dynamic FC abnormalities are repeatedly observed within and between DMN, salience-related, and cerebellar/visual systems, often interpreted as disrupted emotional modulation and self-referential processing (Dai et al. 2023, 294-305; Dwulit et al. 2025). Large consortia show that state dwell time and transitions can be quantified via sliding windows and k-means clustering at scale, even if case–control differences are modest at the diagnostic level (■ Tomas et al. 2025).

Signal Entropy and Network Flexibility (Dimension 2)

BOLD-signal entropy (e.g., SampEn/MSE) reliably distinguishes regions and networks, with intrinsic networks like DMN and frontoparietal control showing higher complexity than sensory networks, supporting its interpretation as an intra-voxel marker of information-processing capacity (Nezafati et al. 2020; Trevino et al. 2024). Methodologically, minimizing bias in SampEn/MSE requires careful choice of pattern length and tolerance, especially for relatively short rs-fMRI runs (Yang et al. 2018).

Across psychiatric disorders, meta-analytic work suggests shared increases in rs-fMRI entropy in frontal and posterior cortical regions, implicating complexity changes as a transdiagnostic feature (■ Ji et al. 2022, 336 - 343). This supports your conceptualization of a ketamine-induced shift from “rigid/trauma-locked” to more entropic brain states as a candidate mediator of treatment response.

Dynamic multilayer modularity approaches quantify **network flexibility** (frequency of community switches) and related metrics, summarizing temporal reconfiguration of large-scale systems (■ Al-Sharoa et al. 2019, 695-709; Ting et al. 2020, 468-480). Flexibility is sensitive to experience-dependent plasticity (e.g., long-term motor training) and psychopathology, with group differences at both network and nodal levels (Cao et al. 2025; ■ Gifford et al. 2020, S285 - S285).

dFC States, Fractional Occupancy, and PTSD (Dimension 1 & 2 Bridge)

Sliding-window dFC with k-means clustering routinely yields a small set of recurrent connectivity “states,” allowing estimation of fractional occupancy, dwell time, and transition profiles in PTSD and related conditions (Dai et al. 2023, 294-305; ■ Tomas et al. 2025; 2021). In comorbid PTSD+mTBI, greater SN-related VFC and altered dwell time in externally focused and transition states link dynamic instability to hypervigilance and reduced network segregation (Dwulit et al. 2025).

These methods operationalize exactly the metrics you propose (variance, fractional occupancy, dwell time, transition dynamics) and can be embedded in linear mixed-effects and mediation models to relate baseline instability and T2 entropy/flexibility shifts to CAPS-5 change.

Example Applications to Your Thesis (Methods/Discussion Phrasing)

Methods hooks you can adapt

- “Dynamic FC between SN and DMN will be estimated using sliding-window correlations and summarized as variance and state metrics (fractional occupancy, dwell time, transition entropy), paralleling established rs-fMRI dFC workflows in PTSD and large consortia samples.” (Dai et al. 2023, 294-305; ■ Tomas et al. 2025)- “Signal-level complexity (SampEn/LZC) will be computed within SN, DMN, and cerebellar/limbic ROIs as an intra-voxel marker of temporal complexity, using parameter choices that minimize known SampEn bias in short rs-fMRI series.” (Nezafati et al. 2020; Yang et al. 2018)- “Multilayer modularity will be used to estimate network flexibility (community switching over time) across timepoints, extending multilayer community detection approaches shown to capture temporal brain dynamics and disease-related reconfiguration.” (■ Al-Sharoa et al. 2019, 695-709; ■ Canal-Garcia et al. 2024)### Conceptual mediation framing

You can explicitly frame T2 entropy/flexibility measures as candidate mediators of the dosing → T3 symptom-change pathway, grounded in ketamine–BDNF–plasticity models and their coupling to rs-fMRI connectivity changes (■ Woelfer et al. 2020, 696 - 710; ■ Pardossi et al. 2024).

Summary

The literature supports your two-dimensional thesis: (1) baseline SN–DMN dFC instability as a prognostic biomarker in PTSD, and (2) a ketamine-induced, time-sensitive plasticity window indexed by increases in BOLD entropy and network flexibility. Established rs-fMRI tools—entropy measures, multilayer modularity, and dFC state metrics—provide a clear methodological backbone for your LMM and mediation analyses, and link naturally to ketamine’s BDNF-dependent neuroplastic mechanisms and extinction-learning enhancement.

These search results were found and analyzed using Consensus, an AI-powered search engine for research. Try it at <https://consensus.app>. © 2026 Consensus NLP, Inc. Personal, non-commercial use only; redistribution requires copyright holders’ consent.

References

"Using deep clustering to improve fMRI dynamic functional connectivity analysis." *Neuroimage* 257 (2021). <https://doi.org/10.1016/j.neuroimage.2022.119288>.

Al-Sharoa, Esraa, M. Al-khassaweneh, and Selin Aviyente. "Tensor Based Temporal and Multilayer Community Detection for Studying Brain Dynamics During Resting State fMRI." *IEEE Transactions on Biomedical Engineering* 66 (2019): 695-709. <https://doi.org/10.1109/tbme.2018.2854676>.

Brown, Kyle A., Patrick J. Morris, Craig J. Thomas, and Todd D. Gould. "A Time-Sensitive Plasticity Distinguishes the Rapid and Sustained Synaptic Actions of Ketamine from Its (2R,6R)-Hydroxynorketamine Metabolite." *The Journal of Neuroscience* 46 (2026). <https://doi.org/10.1523/jneurosci.0958-25.2026>.

Canal-Garcia, Anna, Dániel Veréb, Mite Mijalkov, E. Westman, Giovanni Volpe, and J. Pereira. "Dynamic multilayer functional connectivity detects preclinical and clinical Alzheimer’s disease." *Cerebral Cortex (New York, NY)* 34 (2024). <https://doi.org/10.1093/cercor/bhad542>.

Cao, Bolin, Yu Guo, Fengguang Xia, Lunxiong Li, Zhanbing Ren, Min-ju Lu, Jun Wang, and Ruiwang Huang. "Dynamic reconfiguration of brain functional networks in world class gymnasts: a resting-state functional MRI study." *Brain Communications* 7 (2025). <https://doi.org/10.1093/braincomms/fcaf083>.

Dai, Yingliang, Zhou Zhou, F. Chen, Li Zhang, J. Ke, R. Qi, G. Lu, and Y. Zhong. "Altered dynamic functional connectivity associates with post-traumatic stress disorder." *Brain Imaging and Behavior* 17 (2023): 294-305. <https://doi.org/10.1007/s11682-023-00760-y>.

Diniz, C., A. P. Crestani, P. Casarotto, C. Biojone, C. Cannarozzo, Frederike Winkel, Mikhail A. Prozorov, et al. "Fluoxetine and ketamine trigger the p75NTR proteolytic pathway and enhance extinction memory and brain plasticity through p75NTR.." *Biological psychiatry* (2024). <https://doi.org/10.1016/j.biopsych.2024.06.021>.

Dwulit, Alexandra K., Delin Sun, Courtney C. Haswell, Ahmed Hussain, Emily L. Dennis, Elisabeth A Wilde, Mary Newsome, et al. "Dynamic and Static Resting-State Functional Connectivity of Canonical Networks in Military and Civilian Populations with Posttraumatic Stress Disorder and/or Mild Traumatic Brain Injury.." *Biological psychiatry. Cognitive neuroscience and neuroimaging* (2025). <https://doi.org/10.1016/j.bpsc.2025.08.002>.

Gifford, G., N. Crossley, K. Matthew, Morgan Sarah, P. Dazzan, Jonathan Young, and P. McGuire. "T143. RESTING STATE FMRI BASED MULTILAYER NETWORK CONFIGURATION IN PATIENTS WITH SCHIZOPHRENIA." *Schizophrenia Bulletin* 46 (2020): S285 - S285. <https://doi.org/10.1093/schbul/sbaa029.703>.

Ji, Shanling, Yinghui Zhang, Nan Chen, Xia Liu, Yongchao Li, Xuexiao Shao, Zhengwu Yang, Zhijun Yao, and Bin Hu. "Shared increased entropy of brain signals across patients with different mental illnesses: A coordinate-based activation likelihood estimation meta-analysis." *Brain Imaging and Behavior* 16 (2022): 336 - 343. <https://doi.org/10.1007/s11682-021-00507-7>.

Nezafati, M., Hisham Temmar, and S. Keilholz. "Functional MRI Signal Complexity Analysis Using Sample Entropy." *Frontiers in Neuroscience* 14 (2020). <https://doi.org/10.3389/fnins.2020.00700>.

Pardossi, S., A. Fagiolini, and A. Cuomo. "Variations in BDNF and Their Role in the Neurotrophic Antidepressant Mechanisms of Ketamine and Esketamine: A Review." *International Journal of Molecular Sciences* 25 (2024). <https://doi.org/10.3390/ijms252313098>.

Ting, C., S. Samdin, Meini Tang, H. Ombao, and C. Ting. "Detecting Dynamic Community Structure in Functional Brain Networks Across Individuals: A Multilayer Approach." *IEEE Transactions on Medical Imaging* 40 (2020): 468-480. <https://doi.org/10.1109/tmi.2020.3030047>.

Tomas, C., Jacklynn M. Fitzgerald, C. Baird, C. Haswell, C. Abdallah, Mike Angstadt, Justin Baker, et al. "Data-Driven Approach to Dynamic Resting State Functional Connectivity in Post-Traumatic Stress Disorder: An ENIGMA-PGC PTSD Study." *Human Brain Mapping* 46 (2025). <https://doi.org/10.1002/hbm.70116>.

Trevino, Gabriel, John J Lee, Joshua S. Shimony, Patrick H. Luekett, and Eric C. Leuthardt. "Complexity organization of resting-state functional-MRI networks." *Human Brain Mapping* 45 (2024). <https://doi.org/10.1002/hbm.26809>.

Woelfer, Marie, Meng Li, L. Colic, T. Liebe, Xin Di, B. Biswal, J. Murrough, V. Lessmann, T. Brigadski, and M. Walter. "Ketamine-induced changes in plasma brain-derived neurotrophic factor (BDNF) levels are associated with the resting-state functional connectivity of the prefrontal cortex." *The World Journal of Biological Psychiatry* 21 (2020): 696 - 710. <https://doi.org/10.1080/15622975.2019.1679391>.

Yang, A., S. Tsai, Ching-Po Lin, and Chung-Kang Peng. "A Strategy to Reduce Bias of Entropy Estimates in Resting-State fMRI Signals." *Frontiers in Neuroscience* 12 (2018). <https://doi.org/10.3389/fnins.2018.00398>.