

Preparing and labelling probe from a complex oligo library

Omid Gholamalamdari

Updated: 8/9/2018

Introduction

This protocol employs emulsion PCR (ePCR) to enrich sub libraries from a complex oligo library. I have prepared this protocol based on “*Amplification of complex gene libraries by emulsion PCR*” R. William, *et.al*, *Nature Methods*¹ and “*Preparing Single-Stranded Labelled Probes from MYtags Immortal Library*”², Arbor Biosciences. Labelling protocol is based terminal transferase protocol from Roche.

Materials

- Oligo nucleotide library
- Enrichment primer sets
- Nuclease-free water
- Span 80 (Sigma S6760)
- Tween 80 (Sigma P5188)
- Triton X-100 (Sigma T8787)
- Minral Oil (Sigma M5904)
- Bovine Serum Albumin (BSA) (Sigma B6917)
- diethyl ether

Equipments

- BRAND magnetic stirring bar, PTFE, cylindrical with pivot ring (Sigma Z329061). pre-cleaned (See “Cleaning stir bars”).
 - Magnetic stirrer with speed control (Burrell Scientific)
 - PCR cleanup kit
-

Before you begin

Prepare library

1. Prepare a stock solution of your library.
2. Resuspend at 1 ng/ μ l in 10 mM Tris-HCl pH 7.5 or nuclease-free water.
 - Keep the stock at -80° C.
3. Prepare working stock aliquots (0.07 ng/ μ l) by diluting 2 μ l of immortal library in 28 μ l nuclease-free water.

¹<https://www.ncbi.nlm.nih.gov/pubmed/16791213>

²Version 1.4

Prepare primers

Prepare forward and reverse primers working stocks($10\mu\text{M}$) by diluting the $100\mu\text{M}$ stock in nuclease-free water.

Prepare water saturated diethyl ether.

Mix 100 ml of diethyl ether with 100 ml of nuclease-free water in a glass bottle by shaking for 30s. Allow the phases to settle before use (the top phase is diethyl ether). The water saturated solvent can be stored at RT for 3 months.

Procedure

Prepare ePCR mix.

1. Prepare the oil-surfactant mixture by thoroughly mixing the following components in a 50ml conical tube.

Component	Amount	Final concentration
Span 80	2.25ml	4.5% (vol/vol)
Tween 80	200 μl	0.4% (vol/vol)
Triton X-100	25 μl	0.05% (vol/vol)
Mineral oil	to 50ml	

2. Transfer 400 μl of the oil-surfactant mixture to a CryoTube vial, and add a 3x8 mm stir bar(pre-cleaned)
3. Begin stirring the mixture at 1,000 r.p.m. on the magnetic stirrer.
4. Prepare the aqueous phase for the emulsion by mixing the following:

Component	Amount
10X PCR buffer	26 μl
BSA (100 mg/ml stock)	26 μl
Forward primer (10 μM stock)	7.8 μl
Reverse primer (10 μM stock)	7.8 μl
dNTPs (5mM stock)	10.4 μl
DNA polymerase	5.2 μl
Template DNA	$\leq 10^9$ molecules (1.66 fmol)
Nuclease-free water	to 260 μl

5. Add 200 μl of the aqueous phase to the oil-surfactant mixture in dropwise manner over a period of 1.5 min.
 - After the addition is complete, continue stirring for another 5 minutes.
 - Keep the aqueous phase leftovers as the without emulsion control.

▲ CRITICAL STEP

6. Pipet the emulsion into the wells of a PCR plate as 10 aliquots of 50 μl .
 - Pipet 50 μl of the aqueous phase left over to a well as nonemulsified control.
 - Overlay the emulsified and nonemulsified reactions with mineral oil.

PCR

7. Subject the PCR plate to the following program.

Cycle number	Denaturation	Annealing	Polymerization
1	2 min at 95°C		
2-26	30 sec at 95°C	30 sec at A ¹	B ¹ min at 72°C
27			10 min at 72°C

¹ $A = C - 5^{\circ}\text{C}$ where C is the T_m of the primer with lowest T_m . B is equal to length of amplification in kbp.

8. Place the nonemulsified reaction aside.

■ **PAUSE POINT** You can keep both the w and w/o emulsion reaction at 4°C overnight!

9. Pool the ePCR reactions in a 1.7 ml eppendorf tube.

- Centrifuge at 13,000g for 5 minutes.
- Discard the upper (oil) phase.

10. To remove the remaining oil from the emulsion, perform the following extraction twice:

- Add 1 ml of water-saturated diethyl ether, vortex the tube and let it settle.
- Discard the upper (solvent) phase.

▲ **CRITICAL STEP**

11. Remove the residual solvent from the broken emulsion by centrifuging under vacuum for 5 min at 25°C.

12. Clean the emulsified and nonemulsified PCR products using a PCR clean-up kit.

13. Analyze Aliquots of the emulsified and nonemulsified PCR products by electrophoresis.

- 2% agarose gel in 1X TBE.