

# 3D-FISH in HCT116

*Omid Gholamalamdari*

*Updated on 10/22/2018*

## Cell Preparation

1. Seed the 50k cells per 12mm, #1.5 coverslips (0.5 mL of 100k/mL). Let the cells grow for 2 days. Gorde88
2. Rinse the cells briefly in two to three changes of warm PBS.
3. Fix the cells in 3% paraformaldehyde (freshly made) for 10 minutes at room temperature (RT).
4. Rinse the cells three times with PBS.
5. Permeabilize the cells in PBS-0.5% Triton X-100 for 10 minutes at RT.
6. Rinse the cells three times with PBS.
7. Incubate the cells in 20% glycerol for a minimum of 60 minutes (preferably overnight) at RT.
8. Freeze the cells by dipping the coverslips into liquid nitrogen (for ~30 sec) and thaw on a paper towel. As soon as the frozen layer disappears, put the coverslips back into 20% glycerol. Repeat four times.
9. Wash the cells three times for 10 minutes each in PBS.
10. Rinse the cells briefly with 0.1 N HCl and incubate with a fresh portion of 0.1 N HCl for 10 minutes at RT.
  - The time may be in the range of 5-15 minutes for slides/coverslips with densely grown cells and/or with nuclei embedded in a voluminous cytoplasm. The concentration of HCl should not be varied.
11. Rinse the cells three times with 2X SSC.
12. Incubate the cells in the 50% formamide/2X SSC solution for at least 1 hour at RT (preferably overnight at 4°C) before proceeding with probe hybridization.
  - At this stage, cell can be stored for at least 3-4 months in 50% formamide/2X SSC. Longer storage may result in deterioration of the nuclear morphology after the denaturation step of 3D-FISH.

## Hybridization

1. Prepare the hybridization mixture as following to the total volume of 4 $\mu$ L.

Component	Amount
Probe(s)	80ng each
human COT1 DNA 1mg/ml	1 $\mu$ l
Hybridization buffer	to 4 $\mu$ l

2. Take a coverslip with cells out of the formamide-SSC and quickly drain off the excess fluid. Do not let cells dry out!
3. Place the coverslip onto the drop of hybridization mix (with cells facing the drop).
4. Seal the coverslip with rubber cement. Let the rubber cement dry completely.
5. Place slides on a heating block at 77°C for 3 minutes to denature cellular and probe DNA.
6. Perform hybridization at 37°C for 2-3 days in a humidified chamber.

## Washing and Detection

1. Before you begin warm-up a suitable volume of 2X SSC to 37°C and 0.1X SSC to 60°C.
2. After hybridization, peel off the rubber cement, gently remove the coverslip, and transfer the cells to 2X SSC in a 24 well plate.
  - You can add a small volume 2X SSC on top of the coverslips. This will both soften the rubber cement and coverslips will float easily.
  - Make sure not to move the coverslips with force as this will damage cellular and nuclear structures.
3. Wash in 2X SSC at 37°C, shaking 3x 5min.
  - This is for removing excess probe.
4. Wash in 0.1X SSC at 60°C (stringent washes), shaking 3x 5min.
5. If only fluorochrome-labeled probes are used directly, immediately proceed to DNA counterstaining.
6. Rinse the cells briefly in SSC/Tween (0.2% [v/v] Tween 20 in 4X SSC). Transfer coverslips to a humid chamber.
7. Wash once in 4X SSC at RT, 5 min.
8. Block with 4X SSC/0.1% Triton X-100/4% BSA (Blocking buffer) at RT, for 30 min in a humid chamber.
  - Make sure that you filter the blocking buffer with .2 $\mu$ m syringe filters.
9. Dilute the required antibody or avidin-conjugates to appropriate working concentration in blocking buffer.
  - 50-100 $\mu$ L per coverslip is enough.
10. Incubate cells with antibody for 2 hours in a dark humid chamber at RT.
11. Wash with 4X SSC/0.1% Triton X-100 at RT, shaking 3x 5min.
12. Mount in DAPI containing mounting media.
  - Alternatively, you can counterstain with DAPI (0.05 $\mu$ g/ml in 4X SSC) at RT for 5 min. Wash briefly in 4X SSC/0.1% Triton X-100.
13. If using hardening mounting media, wait overnight for curing.
14. Seal coverslips with colorless nail polish and store at 4°C.
  - FISH signals should be stable for at least 1 year.

HAPPY FISHing!!