Supporting Information for OmicsTopology

OmicsTopology: calibrated topology and phase diagnostics for transcriptomic state spaces

Xiaofeng Song[1]†, Pengyu Chen[1]† and Fei Liang[2]*

[1] North Henan Medical University, Xinxiang, Henan, China

[2] Key Laboratory of Tropical Fruit Biology of Ministry of Agriculture, South Subtropical Crops Research Institute, Chinese Academy of Tropical Agricultural Sciences, Zhanjiang, Guangdong, China

† These authors contributed equally. * Corresponding author: fyliangfei@163.com.

Evidence organization

This supplement contains eleven figures and four numerical tables, plus a visual crosswalk. Figures S1 and S2 are regenerated visual diagnostics, Figures S3–S8 redraw the historical v1 records, S9 summarizes the frozen v2 feature sensitivity, and S10 documents numerical provenance. Historical panels retain their original protocol identity and are not included in current v2 performance estimates.

README image coverage

README visual topic / asset	Publication location	Treatment in current manuscript
Main gallery: software logic architecture	Figure 1d-e	Workflow and evidence layers represented in the current concept-to-inference figure
Main gallery: calibrated detection and abstention	Figure 2	Recomputed from frozen v2 calibration and recovery records
Main gallery: axis-aware phase diagnostics	Figure 5	Updated categorical-axis and circular-phase results
Main gallery: numerical provenance	Figure S10	Provenance diagram aligned to source hashes, settings, and outputs
Main gallery: external ordering comparison	Figure 4; Table S2	Recomputed on common inputs; full and common-support scores retained
Main gallery: cross-experiment heat-response association	Figure 6; Table S1	Recomputed contrasts, contingency counts, odds ratios, and adjusted q

		values
Software overview: state-space topology compared	Figure 1a-c	Conceptual states redrawn; coordinates are illustrative only
Software overview: current v2 analysis workflow	Figure 1d-e	Current inference and interpretation workflow redrawn
v2 figure 2: frozen synthetic validation	Figure 2a,c,d	Current simulation outcomes and threshold sensitivity
v2 figure 3: axis simulations	Figure 5b	Current 30-scenario classification results
v2 figure 4: GSE3431 persistence and stability	Figure 3d-e; Figure S9	Current held-out analysis and feature-selection sensitivity
v2 figure 5: GSE158444 phase diagnostics	Figure 5c-d; Results 2.5	Current circular-dispersion summaries and wrapped phase differences
Historical gallery: circadian loop	Figure 3a; Figure S2	Trajectory display regenerated from public source matrices
Historical gallery: circularity and replicates	Figure S1	Recomputed using the exact v2-selected matrices
Historical gallery: benchmark trajectories	Figure S2	Public-source PCA trajectories regenerated; chronology is not forced closed
Historical gallery: H1 versus null	Figure 2b; Figure S3	Current calibration and historical v1 diagnostic kept separate
Historical gallery: circular versus linear Mantel	Figure S4	Historical v1 correlations retained with explicit legacy labels
Historical gallery: labels held out	Figure 4; Figure S5	Current common-input comparison and historical v1 audit are distinguished
Historical gallery: cycle-counting power	Figure 2d; Figure S6	Current cutoff sensitivity and historical signal-density sweep separated
Historical gallery: incremental value	Figure S7	Historical 22-method baseline benchmark retained as legacy evidence
Historical gallery: increment power	Figure S8	Historical expanded ordering benchmark retained as legacy evidence

Additional methods

The trajectory atlas follows the public registry preprocessing, retaining up to 3,000 variable features after expression filtering and log transformation. PCA is computed separately within each condition, and ordered centroids are joined without forcing the last time point back to the first. This visual analysis is distinct from the 1,000-feature v2 inference matrices. S1 uses those exact v2 matrices; pairwise medians are plotted as separate quantities. Source-file hashes and plotted coordinates accompany the package.

For gene-set enrichment, wild-type heat response is defined identically in both contrast workbooks, while mutant-versus-wild-type at 37 °C is a separate exploratory question. The

background consists of reliably phased genes with an exact identifier match and numeric fold-change/FDR entries for the chosen contrast. This explicit contrast definition supersedes the less specific heat-response summary in the earlier draft. No differential-expression model is refitted, and no inference of genotype-by-temperature interaction is made.

For the historical figures, all numerical values originate in archived JSON records. Their labels describe the original analysis targets; no archived operating characteristic is relabeled as a v2 estimate. The broader historical baseline benchmark has 22 recorded methods, and the expanded synthetic comparison has nine. Reference citations in this supplement refer to the main manuscript bibliography.

S1 Table. Recomputed enrichment contingency counts

a: candidate and externally positive; b: candidate and externally negative; c: background noncandidate and externally positive; d: background noncandidate and externally negative. OR is the sample odds ratio. The heat and genotype test families are adjusted separately.

RNA	Workbook	Test	a	b	c	d	OR	BH q
TOT	cca1lhy	heat	95	21	247	212	3.883	1.3e-08
TOT	cca1lhy	genotype	9	107	49	410	0.704	0.867
TOT	prr7prr9	heat	90	26	236	223	3.271	1.57e-07
TOT	prr7prr9	genotype	3	113	7	452	1.714	0.867
TR	cca1lhy	heat	111	26	302	248	3.506	1.22e-08
TR	cca1lhy	genotype	10	127	52	498	0.754	0.867
TR	prr7prr9	heat	105	32	272	278	3.354	1.22e-08
TR	prr7prr9	genotype	2	135	7	543	1.149	0.867

S2 Table. External ordering and common support

Succession is evaluated on each method’s visited samples. Common-support succession restricts every ordering to the H1-walk vertex set without refitting. Coverage counts unique samples. All scores are descriptive; fit times and complete index orders are retained in the accompanying JSON record.

Dataset subset	Method	Succession	Common support	Coverage

Mouse liver	H1 walk	0.545	0.545	22.9%
Mouse liver	PCA angle	0.271	0.182	100.0%
Mouse liver	Circle fit	0.354	0.091	100.0%
Mouse liver	MDS	0.458	0.364	100.0%
Mouse liver	Isomap	0.396	0.455	100.0%
Mouse liver	Laplacian	0.438	0.364	100.0%
Mouse liver	Tour	0.562	0.545	100.0%
LD whole leaf	H1 walk	0.900	0.900	83.3%
LD whole leaf	PCA angle	1.000	1.000	100.0%
LD whole leaf	Circle fit	1.000	1.000	100.0%
LD whole leaf	MDS	1.000	1.000	100.0%
LD whole leaf	Isomap	1.000	1.000	100.0%
LD whole leaf	Laplacian	0.833	0.800	100.0%
LD whole leaf	Tour	1.000	1.000	100.0%
LD mesophyll	H1 walk	1.000	1.000	66.7%
LD mesophyll	PCA angle	1.000	1.000	100.0%
LD mesophyll	Circle fit	1.000	1.000	100.0%
LD mesophyll	MDS	1.000	1.000	100.0%
LD mesophyll	Isomap	1.000	1.000	100.0%
LD mesophyll	Laplacian	1.000	1.000	100.0%
LD mesophyll	Tour	1.000	1.000	100.0%
LD vasculature	H1 walk	0.778	0.778	66.7%
LD vasculature	PCA angle	1.000	1.000	100.0%
LD vasculature	Circle fit	1.000	1.000	100.0%
LD vasculature	MDS	1.000	1.000	100.0%
LD vasculature	Isomap	1.000	1.000	100.0%
LD vasculature	Laplacian	0.917	0.750	100.0%
LD vasculature	Tour	0.667	0.500	100.0%
SD whole leaf	H1 walk	1.000	1.000	66.7%
SD whole leaf	PCA angle	1.000	1.000	100.0%
SD whole leaf	Circle fit	1.000	1.000	100.0%
SD whole leaf	MDS	1.000	1.000	100.0%
SD whole leaf	Isomap	1.000	1.000	100.0%
SD whole leaf	Laplacian	0.917	1.000	100.0%
SD whole leaf	Tour	1.000	1.000	100.0%
SD mesophyll	H1 walk	1.000	1.000	58.3%
SD mesophyll	PCA angle	0.917	1.000	100.0%
SD mesophyll	Circle fit	1.000	1.000	100.0%
SD mesophyll	MDS	0.917	1.000	100.0%
SD mesophyll	Isomap	1.000	1.000	100.0%
SD mesophyll	Laplacian	1.000	1.000	100.0%
SD mesophyll	Tour	0.917	1.000	100.0%
SD vasculature	H1 walk	1.000	1.000	58.3%
SD vasculature	PCA angle	1.000	1.000	100.0%
SD vasculature	Circle fit	1.000	1.000	100.0%
SD vasculature	MDS	1.000	1.000	100.0%
SD vasculature	Isomap	1.000	1.000	100.0%
SD vasculature	Laplacian	1.000	1.000	100.0%
SD vasculature	Tour	1.000	1.000	100.0%

S3 Table. Public source-file provenance and SHA-256 checksums

Hashes identify the exact downloaded bytes before parsing. The source matrices themselves are not redistributed in this package; the reproduction scripts retrieve the public files and verify these hashes before analysis. GSE116004_index.html is the retrieval index used to locate the published contrast workbooks.

Downloaded source file	Use in analysis	SHA-256
GSE3431_series_matrix.txt.gz	Held-out yeast analysis	6d7c9b65f710d588604d731e2d177264f06eaee36919aa6e3a8d89b56b5dda5b
GSE158444_Counts_Time.txt.gz	Main Arabidopsis v2 analysis	04c6b7e53b058fda37410cd7d9e78ac8be3a88b064a3f8f42f941bd939a0aeeb
GSE158444_Counts_Recovery.txt.gz	Trajectory atlas	fa376c7f095e0b4eaaf0d9e200189befdfb1e7cfc17089e2d2e4e2ac2ead173f
GSE11923_series_matrix.txt.gz	External ordering: mouse liver	2dd9f79ce8562155e003e43984e49b1ed8d190f4ac197f6d612274d21eab39ab
GSE50438_series_matrix.txt.gz	External ordering: Arabidopsis subsets	055a95e054e76bd0b24b4bc1faac99f6316ea4efd67b7d5a3c7bbf8b020a156a
GSE116004_cca1lhyData.xls.gz	Gene-set contrast: cca1lhy	b9b8e006d006cbdd9cdf81b44bda27a3e4aee39b15ca86ba34811e43b5a81cdd
GSE116004_prr7prr9Data.xls.gz	Gene-set contrast: prr7prr9	2e44fca65f324774037f6dee6235d83e8735c681bb5721c88c476d95b7503705
GSE116004_index.html	GEO retrieval index for published contrast workbooks	5b1c73aa8358db4cc1f0b0b5dbf858342a59f27bc7f66e5cf98b04ae31d6942e
GSE138193_AllSamples_FPKM.txt.gz	Trajectory atlas: Arabidopsis	20189037b108d2ecebdf149a7a7aae30c7f4befbb7bc66760018a38ecfba1898
GSE137732_LD_ZT2_47_transcript_expr.txt.gz	Trajectory atlas: long-day series	c7ffe960258227e70305577b7964dc1451b1b4c46e1d70745eb7934dc69adeff
GSE137732_LL_CT2_47_transcript_expr.txt.gz	Trajectory atlas: constant-light series	89b3b95768c4f9e500c11eea318bbe684ad2da38f8b381cfac7b73609b805a1e
GSE128739_devTom_tpm.xlsx	Trajectory atlas: tomato development	8cd24397e599532ba52ac1e7b686f866f3f755fd1a939f600943ebaab5987b9c


S4 Table. Sensitivity of external enrichment to the candidate cutoff

The primary analysis uses the top 20% of reliably fitted genes by absolute wrapped phase difference. The 10% and 30% cutoffs are sensitivity checks. For each row, a/b/c/d are candidate-positive, candidate-negative, noncandidate-positive, and noncandidate-negative genes. BH q values are adjusted across all 12 threshold-by-contrast tests within each outcome family; the heat-response and genotype-under-heat families are adjusted separately.


Outcome	Top fraction	RNA	Contrast	Cand./	a/b/c/d	OR	BH q (12

				mapped			**tests)**
Genotype under heat	10%	TOT	cca1lhy	59/575	4/55/54/462	0.62	1.00
Genotype under heat	10%	TOT	prr7prr9	59/575	0/59/10/506	0.00	1.00
Genotype under heat	10%	TR	cca1lhy	69/687	5/64/57/561	0.77	1.00
Genotype under heat	10%	TR	prr7prr9	69/687	1/68/8/610	1.12	1.00
Genotype under heat	20%	TOT	cca1lhy	116/575	9/107/49/410	0.70	1.00
Genotype under heat	20%	TOT	prr7prr9	116/575	3/113/7/452	1.71	1.00
Genotype under heat	20%	TR	cca1lhy	137/687	10/127/52/498	0.75	1.00
Genotype under heat	20%	TR	prr7prr9	137/687	2/135/7/543	1.15	1.00
Genotype under heat	30%	TOT	cca1lhy	174/575	12/162/46/355	0.57	1.00
Genotype under heat	30%	TOT	prr7prr9	174/575	4/170/6/395	1.55	1.00
Genotype under heat	30%	TR	cca1lhy	206/687	12/194/50/431	0.53	1.00
Genotype under heat	30%	TR	prr7prr9	206/687	2/204/7/474	0.66	1.00
Heat response	10%	TOT	cca1lhy	59/575	55/4/287/229	10.97	4.34e-09
Heat response	10%	TOT	prr7prr9	59/575	48/11/278/238	3.74	2.54e-05
Heat response	10%	TR	cca1lhy	69/687	63/6/350/268	8.04	4.34e-09
Heat response	10%	TR	prr7prr9	69/687	55/14/322/296	3.61	6.21e-06
Heat response	20%	TOT	cca1lhy	116/575	95/21/247/212	3.88	1.67e-08
Heat response	20%	TOT	prr7prr9	116/575	90/26/236/223	3.27	1.88e-07
Heat response	20%	TR	cca1lhy	137/687	111/26/302/248	3.51	1.22e-08
Heat response	20%	TR	prr7prr9	137/687	105/32/272/278	3.35	9.69e-09
Heat response	30%	TOT	cca1lhy	174/575	141/33/201/200	4.25	9.72e-12
Heat response	30%	TOT	prr7prr9	174/575	128/46/198/203	2.85	5.17e-08
Heat response	30%	TR	cca1lhy	206/687	162/44/251/230	3.37	1.47e-10
Heat response	30%	TR	prr7prr9	206/687	146/60/231/250	2.63	2.85e-08


S1 Fig. Updated circularity and replicate diagnostics in GSE158444 (TOT, total RNA; TR, translating RNA). (a,b,d,e) Pairwise correlation dissimilarity versus linear time separation for control and heat libraries in total and translating RNA. Points are sample pairs and lines join the median at each observed separation. (c,f) Medians for pairs at the same versus different sampling hours are displayed side by side, rather than added as if they were components of a total. This calculation uses the first 24 hours and the exact v2-selected matrices. Pairwise entries share samples and are descriptive, not independent replicates. This replaces the README circularity-and-replicates illustration with a calculation consistent with the current selection.




S2 Fig. Regenerated multi-study trajectory atlas corresponding to the README benchmark gallery. PCA is fitted separately to each dataset-condition matrix after standardization of the 3,000 features selected by the registry pipeline. Panels show Arabidopsis diurnal control and heat, long-day and constant-light series, recovery control and heat, three rice genotypes, and tomato fruit development. Color progresses from earlier to later time within each panel; gray lines join chronological centroids without forced closure. The registry includes the 24-hour endpoint in the diurnal control (26 libraries), whereas the current main analysis uses hours below 24 (23 control libraries). These displays do not inherit historical detection verdicts or define the geometry used in Figures 2-6.


a
Arabidopsis diurnal control
b
Arabidopsis diurnal heat
c
Arabidopsis long day (LD)
d
Arabidopsis constant light (LL)
e
Arabidopsis recovery control
f
Arabidopsis recovery heat
g
Rice diurnal LY2186
h
Rice diurnal MH86
i
Rice diurnal SE21s
j
Tomato fruit development
PC1
PC2


S3 Fig. Historical v1 H1-reference comparison. Bars show the normalized longest H1 lifetime and red marks the recorded 95th percentile of the v1 reference for ten dataset-condition clouds. Values are redrawn directly from benchmark/results.json. They are retained as a historical diagnostic and are not recalculated v2 tail scores. Experimental expectations of recurrence are not biological ground-truth H1 labels.




S4 Fig. Historical v1 time-association diagnostics. Gray and blue points show correlations with linear time and the declared circular period; orange marks show the largest correlation over the archived period scan. The optimized values are descriptive. Historical permutation p-values are omitted, and this figure is not used to infer an independent cyclic coordinate. Source: benchmark/results.json.




S5 Fig. Historical label-withholding and sample-order diagnostics. (a) Circular Mantel correlations with recorded time labels and the median after label permutation, from the v1 side-information audit. (b) H1-walk cyclic succession and unique-sample coverage for the archived ordering experiments. Time labels were withheld from construction of the representative walk and used to evaluate its order. High succession on restricted support does not imply full-sample reconstruction. Numerical records are retained from benchmark/topology_without_labels.json; v1 detection labels and permutation significance claims are not transferred to v2.




S6 Fig. Historical signal-density and sampling-density sweep. The number of signal-carrying features is varied among 3,000 total features under six or sixteen distinct phases, using eight seeds per setting. (a) Median longest-to-second-longest H1-bar ratio. (b) Fraction of simulations reported as one cycle by the historical procedure. Values are redrawn from benchmark/cycle_counting_power.json. The bar-ratio criterion belongs to the earlier protocol and is not the v2 decision rule. This figure illustrates why relative signal strength and phase coverage must be reported.




S7 Fig. Complete historical baseline comparison. Sensitivity, specificity, and mean fitting time are redrawn for every method in benchmark/incremental_value.json on the archived corpus of 12 designated positive and 23 negative clouds. Error bars retain the recorded intervals; timings describe that original environment. Design-based positive labels in real data are not independently established topological ground truth. These historical ordering results do not estimate the operating characteristics of the current v2 H1 calibration.




S8 Fig. Historical expanded ordering benchmark. (a) False declarations among 240 simulated negative clouds, with recorded Wilson intervals and counts. (b) Proportions by negative generator family, each containing 60 clouds. (c) The fraction producing no order, producing an order below its score threshold, or making a false declaration. Forty positive ring clouds were also evaluated; all nine methods detected 40 of 40 in the archived benchmark. The record evaluates ordering against a phase-score reference and is separate from the v2 maximum-H1 reference test. Source: benchmark/increment_power.json.



S9 Fig. Feature-selection sensitivity in the GSE3431 v2 analysis. (a) Full-cloud candidate counts at 500, 1,000, and all 9,335 varying features, annotated with the final status. (b) Agreement of reduced-cloud counts with the full-cloud count; the dashed line is the fixed 0.75 threshold. The two selected-feature analyses remain unresolved, while the all-feature analysis returns stable zero detection. Source: benchmark/v2/real_validation.json.



S10 Fig. Numerical provenance and figure generation. Public accessions and source hashes connect to declared preprocessing, seeds, algorithm version, and numerical environment. Scored records preserve spectra, reference maxima, reduced-cloud counts, phase fits, and external gene matches. Figure-generation scripts consume these records. The supplement distinguishes regenerated source-matrix analyses from historical records redrawn without rerunning their inference.



S11 Fig. Sensitivity of external heat-response enrichment to the candidate cutoff. Odds ratios and 95% log-odds Wald intervals are shown for all four combinations of RNA fraction and external contrast at top 10%, 20%, and 30% absolute wrapped phase-difference cutoffs. The 20% threshold remains the primary analysis; alternatives are sensitivity checks, not a threshold-selection procedure. Benjamini-Hochberg q values across all 12 threshold-by-contrast tests in the heat-response family are reported in S4 Table.

