

DYNAMICAL SYSTEMS AND OPERATOR ALGEBRAS ASSOCIATED TO ARTIN'S REPRESENTATION OF BRAID GROUPS

TRON OMLAND

ABSTRACT. Artin's representation is an injective homomorphism from the braid group B_n on n strands into $\text{Aut } \mathbb{F}_n$, the automorphism group of the free group \mathbb{F}_n on n generators. The representation induces maps $B_n \rightarrow \text{Aut } C_r^*(\mathbb{F}_n)$ and $B_n \rightarrow \text{Aut } C^*(\mathbb{F}_n)$ into the automorphism groups of the corresponding group C^* -algebras of \mathbb{F}_n . These maps also have natural restrictions to the pure braid group P_n . In this paper, we consider twisted versions of the actions by cocycles with values in the circle, and discuss the ideal structure of the associated crossed products. Additionally, we make use of Artin's representation to show that the braid groups B_∞ and P_∞ on infinitely many strands are both C^* -simple.

INTRODUCTION

The theory of braids, developed by Artin [1], has been of interest to mathematicians from a wide variety of fields, and provides an intriguing mixture of algebra, analysis, and geometry. Originally, the motivation for studying braid groups came from knot theory, and we refer to the book by Birman [8] for an introduction to braids in connection with knots, links, and mapping class groups. Our approach, on the other hand, will be almost purely algebraic.

The braid groups B_n on n strands can be formed from the standard presentation of the symmetric group S_n , by removing all order two relations. While S_n is the group of permutations of an n -point set, the braid group can also be viewed as being of dynamical nature, acting by permutations twisted by inner automorphisms, where the most natural example to consider is the action of B_n on the free group \mathbb{F}_n . This action is defined by an injective homomorphism $B_n \rightarrow \text{Aut } \mathbb{F}_n$, which is often called “Artin's representation”. In this paper, we study Artin's representation from an operator algebraic viewpoint, letting instead B_n act on the various group C^* -algebras of \mathbb{F}_n , and also allowing the permutations to be twisted by a scalar-valued cocycle.

After defining the C^* -dynamical systems, we investigate the corresponding crossed product C^* -algebras. For the reduced versions, we find conditions for the crossed products to be simple and have a unique tracial state, while for the full versions, we find conditions that ensure primitivity, that is, existence of a faithful irreducible representation.

We organize the paper as follows. First, in Section 1, we recall the definitions and some properties of braid groups and Artin's representation that we will need later, and include a short Section 2 with some preliminaries on group C^* -algebras and semidirect products. Then, in Section 3 we introduce the twisted versions of Artin's representations for operator algebras, and state our main theorems, before computing all the possible deformations, that is, the cocycles up to cohomology class. To prove our results, we apply techniques developed in [6] for reduced twisted group C^* -algebras, especially the so-called “relative Kleppner condition”. This is described in Section 4, which also contains the proofs of our main results, relying on several combinatorially involved arguments. In Section 5 we investigate the second cohomology groups and the twisted group C^* -algebras associated to different braid related

Date: April 24, 2019.

2010 Mathematics Subject Classification. 46L05 (Primary) 20F36, 22D25, 46L55 (Secondary).

groups in more detail. Finally, in Section 6 we prove that the braid groups B_∞ and P_∞ on infinitely many strands are both C^* -simple, once again by using Artin's representation and the relative Kleppner condition.

1. BRAID GROUPS AND ARTIN'S REPRESENTATION

For every $2 \leq n \leq \infty$, the *braid group* B_n is defined by generators s_1, s_2, \dots, s_{n-1} subject to relations

$$\begin{aligned} s_i s_{i+1} s_i &= s_{i+1} s_i s_{i+1} \text{ for all } 1 \leq i \leq n-2, \\ s_i s_j &= s_j s_i \text{ when } |i-j| \geq 2. \end{aligned}$$

Moreover, for every n there is a natural surjection from B_n onto the symmetric group S_n , sending the generator s_i to the permutation $(i, i+1)$, and thus s_i^2 to the identity of S_n for all i . The *pure braid group* P_n is defined as the kernel of the map $B_n \rightarrow S_n$.

There is an action α of B_n on the free group $\mathbb{F}_n = \langle x_1, \dots, x_n \rangle$ given by

$$(1.1) \quad \alpha(s_i)(x_j) = \begin{cases} x_{i+1} & \text{if } j = i, \\ x_{i+1}^{-1} x_i x_{i+1} & \text{if } j = i+1, \\ x_j & \text{else.} \end{cases}$$

This induces an injective homomorphism $B_n \rightarrow \text{Aut } \mathbb{F}_n$ called *Artin's representation of B_n* (cf. [8, Corollary 1.8.3]; note that we use the inverse of their representation, to ensure compatibility with our definition of the semidirect product in Section 2.2, see [18, Section 1.5.1]). Remark in particular that $\alpha(s_i)(x_1 x_2 \cdots x_n) = x_1 x_2 \cdots x_n$ for all i . We also use α to denote the restriction of this action to P_n . The pure braid groups can now be described as iterative semidirect products by free groups, that is,

$$P_{n+1} \simeq \mathbb{F}_n \rtimes_\alpha P_n \simeq \mathbb{F}_n \rtimes \mathbb{F}_{n-1} \rtimes \cdots \rtimes \mathbb{F}_2 \rtimes \mathbb{Z}.$$

It is possible to describe P_n by generators

$$a_{i,j} = s_{j-1} \cdots s_{i+1} s_i^2 s_{i+1}^{-1} \cdots s_{j-1}^{-1}$$

for $1 \leq i < j \leq n$ and certain relations (see [8, Lemma 1.8.2] or [18, Corollary 1.19]). The isomorphism between this version of P_{n+1} and the semidirect product described above identifies the free group generated by $a_{1,n+1}, a_{2,n+1}, \dots, a_{n,n+1}$ with the normal subgroup \mathbb{F}_n .

Moreover, define the *annular braid group* A_n as the (non-normal) subgroup of braids of B_n that fix the endpoint of the last string (note that P_n is the subgroup of braids of B_n that fix the endpoint of *all* strings). One can compute that

$$A_{n+1} = \langle s_1, \dots, s_{n-1}, s_n^2 \rangle \subseteq B_{n+1}.$$

The annular braid group was first studied by Chow [12]. The group does not seem to have a standardized notation, e.g. it is denoted by D_n in [8, p. 22] and by B_n^1 in [13, 3.8, 3.9, 4.6]. It is observed in several papers, see e.g. [14, Proposition 2.1], that

$$A_{n+1} \simeq \mathbb{F}_n \rtimes_\alpha B_n,$$

where $a_{1,n+1}, \dots, a_{n,n+1} \in A_{n+1}$ generate \mathbb{F}_n and $s_1, \dots, s_{n-1} \in A_{n+1}$ generate B_n . In fact, the surjections $A_{n+1} \rightarrow B_n$ and $P_{n+1} \rightarrow P_n$ considered in this section, both with kernel \mathbb{F}_n , can be viewed as forgetting the last string.

Clearly, P_n is also a normal subgroup of A_n for all $n \geq 2$, and $A_n/P_n \simeq S_{n-1}$. For completeness, we set $A_1 = P_1 = B_1 = \{e\}$. Thus $[B_n : P_n] = n!$, $[A_n : P_n] = (n-1)!$, and $[B_n : A_n] = n$ for all $n \geq 1$.

Define $\Delta := s_1(s_2s_1) \cdots (s_{n-1} \cdots s_2s_1)$, sometimes called the fundamental element of B_n . For $n \geq 3$, it is well-known (see e.g. [8, Corollary 1.8.4] or [18, Theorem 1.24, Exercise 1.3.2]) that the element $z := \Delta^2$ generates the center of B_n , and that

$$(1.2) \quad z = \Delta^2 = (s_1 \cdots s_{n-1})^n = a_{12}(a_{13}a_{23}) \cdots (a_{1n}a_{2n} \cdots a_{n-1,n}).$$

This element is a member of both A_n and P_n and generates the centers of those groups too. It is not hard to see that

$$(1.3) \quad \alpha(z)(y) = (x_1 \cdots x_n)^{-1} y (x_1 \cdots x_n)$$

for all $y \in \mathbb{F}_n$. Indeed, one checks that this holds when $y = x_i$, by applying (1.2) and an iterative process using that for all $1 \leq i \leq n-1$ we have

$$(1.4) \quad \alpha(s_1 \cdots s_{n-1})(x_i) = x_{i+1} \quad \text{and} \quad \alpha(s_1 \cdots s_{n-1})(x_n) = (x_2 \cdots x_n)^{-1} x_1 (x_2 \cdots x_n).$$

Thus, in the semidirect product description, the generator of the center of $\mathbb{F}_n \rtimes_\alpha B_n$ is

$$(1.5) \quad x_1 \cdots x_n z = z x_1 \cdots x_n,$$

where $z \in B_n$ is the element described in (1.2), and $\mathbb{F}_n = \langle x_1, \dots, x_n \rangle$.

Remark 1.1. The group A_{n+1} is isomorphic to the Artin group associated with the irreducible spherical (finite) Coxeter group with matrix entries $m_{n-1,n} = 4$, $m_{i,i+1} = 3$ when $1 \leq i \leq n-2$, and $m_{ij} = 2$ else. In the classification system for Coxeter groups, these groups are sometimes classified as type B , while the braid groups are of type A (i.e., opposite of the symbols used above for the corresponding Artin groups). In particular, we have

$$(1.6) \quad A_3 \simeq \langle t_1, t_2 : (t_1 t_2)^2 = (t_2 t_1)^2 \rangle.$$

Moreover, the map $P_3 \rightarrow \mathbb{F}_2 \times \mathbb{Z} = \langle v_1, v_2 \rangle \times \langle u \rangle$ given by

$$(1.7) \quad s_2 s_1^2 s_2^{-1} \mapsto v_1, \quad s_2^2 \mapsto v_2, \quad s_1^2 \mapsto (v_1 v_2)^{-1} u \quad (\text{and } z \mapsto u)$$

is an isomorphism. In fact, for all $n \geq 1$, we have $P_n \simeq (P_n/Z(P_n)) \times Z(P_n)$.

2. PRELIMINARIES

2.1. Group C^* -algebras. Here we give some brief preliminaries on group C^* -algebras in the slightly more general twisted setting, that we will need later.

Let G be any discrete group with identity e . A normalized 2-cocycle on G is a map $\sigma: G \times G \rightarrow \mathbb{T}$ satisfying

$$\begin{aligned} \sigma(a, b)\sigma(ab, c) &= \sigma(a, bc)\sigma(b, c) \\ \sigma(a, e) &= \sigma(e, a) = 1 \end{aligned}$$

for all $a, b, c \in G$.

The left regular σ -projective representation $\lambda_\sigma: G \rightarrow B(\ell^2(G))$ is defined by

$$\lambda_\sigma(a)f(b) = \sigma(a, a^{-1}b)f(a^{-1}b).$$

The reduced twisted group C^* -algebra $C_r^*(G, \sigma)$ is the C^* -subalgebra of $B(\ell^2(G))$ generated by $\lambda_\sigma(G)$, and the twisted group von Neumann algebra $W^*(G, \sigma)$ is the von Neumann subalgebra of $B(\ell^2(G))$ generated by $\lambda_\sigma(G)$.

The full twisted group C^* -algebra $C^*(G, \sigma)$ is the universal C^* -algebra generated by a set of unitaries $\{U_\sigma(a)\}_{a \in G}$ subject to the relations $U_\sigma(a)U_\sigma(b) = \sigma(a, b)U_\sigma(ab)$.

Let $Z^2(G, \mathbb{T})$ denote the group of normalized 2-cocycles on G , and let $B^2(G, \mathbb{T})$ denote its subgroup consisting of all $\sigma \in Z^2(G, \mathbb{T})$ for which there exists a function $\beta: G \rightarrow \mathbb{T}$ such that

$$\sigma(a, b) = \beta(a)\beta(b)\overline{\beta(ab)} \quad \text{for all } a, b \in G.$$

The second cohomology group of G with values in \mathbb{T} is then

$$H^2(G, \mathbb{T}) = Z^2(G, \mathbb{T})/B^2(G, \mathbb{T}).$$

We say that σ_1 and σ_2 in $Z^2(G, \mathbb{T})$ are *similar* and write $\sigma_1 \sim \sigma_2$ if their image coincide in $H^2(G, \mathbb{T})$. If this is the case, there is an isomorphism between $C_r^*(G, \sigma_1)$ and $C_r^*(G, \sigma_2)$ given by $\lambda_{\sigma_1}(a) \mapsto \beta(a)\lambda_{\sigma_2}(a)$, where β is the function implementing the similarity. The same also holds for the group von Neumann algebra and for the full group C^* -algebra, with the $\lambda_{\sigma_i}(a)$'s replaced by the universal generators $U_{\sigma_i}(a)$ in the latter case.

2.2. Semidirect products. An action α of a group K on another group H gives rise to a semidirect product G consisting of elements $(x, a) \in H \times K$ and with multiplication given by $(x, a)(y, b) = (x\alpha_a(y), ab)$. As is common, we use the notation $G = H \rtimes_{\alpha} K$ for a semidirect product. One often identifies H with a normal subgroup of G and K with a subgroup of G , and therefore drops the parentheses and just writes xa for an element of G , and the action is then given by $\alpha_a(x) = axa^{-1}$.

Moreover, recall that $Z^1(K, \text{Hom}(H, \mathbb{T}))$ consists of all functions $\varphi: K \times H \rightarrow \mathbb{T}$ satisfying

$$(2.1) \quad \begin{aligned} \varphi(ab, x) &= \varphi(a, \alpha_b(x))\varphi(b, x), \\ \varphi(a, xy) &= \varphi(a, x)\varphi(a, y). \end{aligned}$$

for $a, b \in K$ and $x, y \in H$. Such functions φ are often called 1-cocycles for α with values in \mathbb{T} . The subgroup $B^1(K, \text{Hom}(H, \mathbb{T}))$ of $Z^1(K, \text{Hom}(H, \mathbb{T}))$ consists of all functions $h: K \times H \rightarrow \mathbb{T}$ for which there exists $f \in \text{Hom}(H, \mathbb{T})$ such that

$$h(a, x) = f(\alpha_a(x))\overline{f(x)}$$

for all $a \in K$ and $x \in H$. The corresponding cohomology group is then defined as

$$H^1(K, \text{Hom}(H, \mathbb{T})) = Z^1(K, \text{Hom}(H, \mathbb{T}))/B^1(K, \text{Hom}(H, \mathbb{T})),$$

and we write $\varphi_1 \sim \varphi_2$ if $\varphi_1, \varphi_2 \in Z^1(K, \text{Hom}(H, \mathbb{T}))$ and $\varphi_1\overline{\varphi_2} \in B^1(K, \text{Hom}(H, \mathbb{T}))$.

Let $G = H \rtimes_{\alpha} K$ be any semidirect product and let $\varphi \in Z^1(K, \text{Hom}(H, \mathbb{T}))$. Define $\sigma^{\varphi} \in Z^2(G, \mathbb{T})$ by

$$\sigma^{\varphi}((x, a), (y, b)) = \varphi(a, y)$$

for $x, y \in H$ and $a, b \in K$. For the moment, write λ_{φ} for the left regular σ^{φ} -projective representation of G and λ for the left regular representation of H . Then

$$\lambda_{\varphi}(a)\lambda_{\varphi}(x)\lambda_{\varphi}(a)^* = \sigma^{\varphi}(a, x)\overline{\sigma^{\varphi}(\alpha_a(x), a)}\lambda_{\varphi}(\alpha_a(x)) = \varphi(a, x)\lambda(\alpha_a(x)),$$

so there are $*$ -automorphisms of $C_r^*(H)$ satisfying

$$\alpha_a^{\varphi}(\lambda(x)) = \varphi(a, x)\lambda(\alpha_a(x)).$$

It is now an easy exercise to check that $\alpha_a^{\varphi}\alpha_b^{\varphi} = \alpha_{ab}^{\varphi}$ using (2.1). Thus, α^{φ} defines an action of K on $C_r^*(H)$, giving rise to a C^* -dynamical system for which we can form the reduced crossed product $C_r^*(H) \rtimes_{\alpha^{\varphi}}^r K$.

Similarly, we get actions of K on $W^*(H)$ and $C^*(H)$, also denoted by α^{φ} , producing W^* - and full C^* -crossed products. It is well-known (see [3, Theorem 2.1], [2, Theorem 1], and [24, Theorem 4.1], respectively) that

$$C_r^*(G, \sigma^{\varphi}) \simeq C_r^*(H) \rtimes_{\alpha^{\varphi}}^r K, \quad W^*(G, \sigma^{\varphi}) \simeq W^*(H) \rtimes_{\alpha^{\varphi}} K,$$

and

$$C^*(G, \sigma^{\varphi}) \simeq C^*(H) \rtimes_{\alpha^{\varphi}} K.$$

If $\varphi_1 \sim \varphi_2$ via some f , define $\beta: G \rightarrow \mathbb{T}$ by $\beta(x, a) = f(x)$, and then $\alpha^{\varphi_1} \sim \alpha^{\varphi_2}$ via β .

2.3. Simplicity, primitivity, Kleppner's condition, and tracial states. Recall that a C^* -algebra is called *simple* if it has no nontrivial proper two-sided closed ideals, and *primitive* if it has a faithful irreducible representation. Simplicity clearly implies primitivity, which again implies triviality of the center. A von Neumann algebra is called a *factor* if it has trivial center (or equivalently, if it does not have any von Neumann algebra ideals).

A group G is called *icc* if every nontrivial conjugacy class in G is infinite, and this notion generalizes to the twisted setting as follows: an element $a \in G$ is σ -regular if $\sigma(a, b) = \sigma(b, a)$ whenever $ab = ba$, and σ -regularity is a property of conjugacy classes, so inspired by [19] we say that (G, σ) satisfies *Kleppner's condition* if there is no nontrivial finite σ -regular conjugacy class in G .

For a 2-cocycle σ on a countably infinite group G , it is explained in [22, Theorem 2.7] that $W^*(G, \sigma)$ is a II_1 factor if and only if $C_r^*(G, \sigma)$ is primitive, if and only if $C_r^*(G, \sigma)$ has trivial center, if and only if (G, σ) satisfies Kleppner's condition.

Moreover, a state on a C^* -algebra A is a linear functional $\phi: A \rightarrow \mathbb{C}$ that is positive, i.e., $\phi(S) \geq 0$ whenever $S \in A$ and $S \geq 0$, and unital, i.e., $\phi(1) = 1$. A state ϕ is called *tracial* if it satisfies the additional property that $\phi(ST) = \phi(TS)$ for all $S, T \in A$.

There is a canonical faithful tracial state τ on $C_r^*(G, \sigma)$, namely the vector state associated with δ_e , that is, $\tau(S) = \langle S\delta_e, \delta_e \rangle$ for all $S \in C_r^*(G, \sigma)$.

A group G is called C^* -*simple* if $C_r^*(G)$ is simple and is said to have *the unique trace property* if τ is the only tracial state on $C_r^*(G)$. It is shown in [9, Theorem 1.3], based on [17], that C^* -simplicity is stronger than the unique trace property (in fact, strictly stronger by [20, 16]). If G is C^* -simple (resp. has the unique trace property), then $C_r^*(G, \sigma)$ is simple (resp. has a unique tracial state) for every $\sigma \in Z^2(G, \mathbb{T})$, cf. [10, Corollaries 4.5 and 5.3]. In general, for a pair (G, σ) with $\sigma \not\sim 1$, it is not known whether simplicity of $C_r^*(G, \sigma)$ implies that τ is its only tracial state.

Note that in [5, 6] a pair (G, σ) is said to be C^* -simple (resp. have the unique trace property) when $C_r^*(G, \sigma)$ is simple (resp. has a unique tracial state).

Remark 2.1. The group $B_n/Z(B_n)$ is C^* -simple for all $n \geq 1$, according to [3, p. 536]. Thus its normal subgroup $P_n/Z(P_n)$ is also C^* -simple for all $n \geq 1$, by applying [9, Theorem 1.4]. Moreover, $A_n/Z(A_n)$ embeds into $B_n/Z(B_n)$ as a subgroup of finite index, and is therefore C^* -simple for all $n \geq 1$, by [7, p. 216].

As a consequence, all these quotient groups are icc, which by [5, Proposition 2.5] means that every conjugacy class in B_n , P_n , or A_n is either infinite or a one-element set.

Finally, we also note that for any of these groups, the center coincides with the amenable radical defined in [16, Section 6].

3. TWISTED VERSIONS OF ARTIN'S REPRESENTATION

We now apply Section 2.2 to Artin's representation, i.e., with $H = \mathbb{F}_n$ and $K = B_n$ or P_n .

Definition 3.1. Let $\varphi: B_n \times \mathbb{F}_n \rightarrow \mathbb{T}$ be an element of $Z^1(B_n, \text{Hom}(\mathbb{F}_n, \mathbb{T}))$, and let α be as in (1.1). Define the reduced φ -deformed version of Artin's representation of braid groups as the C^* -dynamical system $\alpha^\varphi: B_n \rightarrow \text{Aut } C_r^*(\mathbb{F}_n)$ given on generators by

$$\alpha_{s_j}^\varphi(\lambda(x_k)) = \varphi(s_j, x_k) \lambda(\alpha_{s_j}(x_k)).$$

The von Neumann algebra version $\alpha^\varphi: B_n \rightarrow \text{Aut } W^*(\mathbb{F}_n)$ is defined analogously.

Let $\mu^\varphi \in \mathbb{T}$ be the product of the values of φ on generators, i.e.,

$$\mu^\varphi = \prod_{\substack{1 \leq j \leq n-1 \\ 1 \leq k \leq n}} \varphi(s_j, x_k).$$

As usual, an element $w \in \mathbb{T}$ is called nontorsion if $w^m \neq 1$ for all $m \in \mathbb{Z} \setminus \{0\}$.

Theorem 3.2. *Let α , φ , α^φ , and μ^φ be as above. Then the following are equivalent:*

- (i) μ^φ is nontorsion,
- (ii) $C_r^*(\mathbb{F}_n) \rtimes_{\alpha^\varphi}^r B_n$ is simple,
- (iii) $C_r^*(\mathbb{F}_n) \rtimes_{\alpha^\varphi}^r B_n$ has a unique tracial state,
- (iv) $W^*(\mathbb{F}_n) \rtimes_{\alpha^\varphi} B_n$ is a factor.

The proof of Theorem 3.2 is given below in Section 4.

Definition 3.3. Let $\varphi: P_n \times \mathbb{F}_n \rightarrow \mathbb{T}$ be an element of $Z^1(P_n, \text{Hom}(\mathbb{F}_n, \mathbb{T}))$, and let α be as in (1.1). Define the reduced φ -deformed version of Artin's representation of pure braid groups as the C^* -dynamical system $\alpha^\varphi: P_n \rightarrow \text{Aut } C_r^*(\mathbb{F}_n)$ given by

$$\alpha_{a_{ij}}^\varphi(\lambda(x_k)) = \varphi(a_{ij}, x_k) \lambda(\alpha_{a_{ij}}(x_k)).$$

The von Neumann algebra version $\alpha^\varphi: P_n \rightarrow \text{Aut } W^*(\mathbb{F}_n)$ is defined analogously.

For $1 \leq k \leq n$, let $\nu_k^\varphi \in \mathbb{T}$ be the value

$$\nu_k^\varphi = \prod_{1 \leq i < j \leq n} \varphi(a_{ij}, x_k).$$

Theorem 3.4. *Let α , φ , α^φ , and ν_k^φ be as above. Consider the following conditions:*

- (i) ν_k^φ is nontorsion for at least one k ,
- (ii) $C_r^*(\mathbb{F}_n) \rtimes_{\alpha^\varphi}^r P_n$ is simple,
- (iii) $C_r^*(\mathbb{F}_n) \rtimes_{\alpha^\varphi}^r P_n$ has a unique tracial state,
- (iv) $W^*(\mathbb{F}_n) \rtimes_{\alpha^\varphi} P_n$ is a factor.

Then (i) \implies (ii) \implies (iv) and (i) \implies (iii) \implies (iv).

Moreover, if $n = 2$, then (iv) \implies (i).

The proof of Theorem 3.4 is given below in Section 4.

Example 3.5. Let us consider the case $n = 2$, where $B_2 \simeq \mathbb{Z}$ is generated by $s = s_1$ and $P_2 \simeq \mathbb{Z}$ is generated by $a = a_{12} = s^2$. First, for the braid group we have

$$\begin{aligned} \alpha_s^\varphi(\lambda(x_1)) &= \varphi(s, x_1) \lambda(x_2), \\ \alpha_s^\varphi(\lambda(x_2)) &= \varphi(s, x_2) \lambda(x_2^{-1} x_1 x_2). \end{aligned}$$

Moreover, $\varphi_1 \sim \varphi_2$ if and only if $\varphi_1(s, x_1) \varphi_1(s, x_2) = \mu^{\varphi_1} = \mu^{\varphi_2} = \varphi_2(s, x_1) \varphi_2(s, x_2)$, as explained in Proposition 3.10 below. For the pure braid group, we have

$$\begin{aligned} \alpha_a^\varphi(\lambda(x_1)) &= \varphi(a, x_1) \lambda(x_2^{-1} x_1 x_2), \\ \alpha_a^\varphi(\lambda(x_2)) &= \varphi(a, x_2) \lambda(x_2^{-1} x_1^{-1} x_2 x_1 x_2), \end{aligned}$$

In particular, we notice that $\alpha_a = \text{Ad } x_2^{-1} x_1^{-1}$ in this case. See Example 4.5 for more on this.

Question 3.6. In Theorem 3.4, does (iv) \implies (i), (ii), or (iii) for $n \geq 3$?

Definition 3.7. Let α be as in (1.1). First, for $\varphi \in Z^1(B_n, \text{Hom}(\mathbb{F}_n, \mathbb{T}))$, the full φ -deformed version of Artin's representation of braid groups is the map $\alpha^\varphi: B_n \rightarrow \text{Aut } C^*(\mathbb{F}_n)$, defined on generators similarly as in Definition 3.1.

Secondly, for $\varphi \in Z^1(P_n, \text{Hom}(\mathbb{F}_n, \mathbb{T}))$, the full φ -deformed version of Artin's representation of pure braid groups is the map $\alpha^\varphi: P_n \rightarrow \text{Aut } C^*(\mathbb{F}_n)$, defined on generators similarly as in Definition 3.3.

Theorem 3.8. *The full crossed product $C^*(\mathbb{F}_2) \rtimes_{\alpha^\varphi} B_2$ is primitive if and only if it has trivial center, if and only if $\mu^\varphi = \varphi(s, x_1)\varphi(s, x_2)$ is nontorsion.*

The full crossed product $C^(\mathbb{F}_2) \rtimes_{\alpha^\varphi} P_2$ is primitive if and only if it has trivial center, if and only if at least one of $\nu_1^\varphi = \varphi(a, x_1)$ and $\nu_2^\varphi = \varphi(a, x_2)$ is nontorsion.*

Proof. Assume $\mu = \mu^\varphi$ is nontorsion. We will first show that $C^*(\mathbb{F}_2) \rtimes_{\alpha^\varphi} B_2$ is primitive. According to work of Choi [11], there exists a faithful irreducible representation π of $C^*(\mathbb{F}_2)$ on some separable Hilbert space \mathcal{H} . Let U_1 and U_2 be the two generators of $C^*(\mathbb{F}_2)$ and set $V_1 = \pi(U_1)$ and $V_2 = \pi(U_2)$. We may assume that V_2 is diagonal with distinct diagonal entries (see [11, Proof of Theorem 6]). For every $z \in \mathbb{T}$ define an automorphism γ_z on $C^*(\mathbb{F}_2)$ by $\gamma_z(U_1) = U_1$ and $\gamma_z(U_2) = zU_2$, and set $\pi_z = \pi \circ \gamma_z$. Clearly, π_z is also faithful and irreducible for every choice of $z \in \mathbb{T}$, and we would like to show that $\pi_z \circ \alpha_{s^k}^\varphi \neq \pi_z$ for every $k \in \mathbb{Z} \setminus \{0\}$, and then apply [4, Theorem 2.1]. We compute that

$$\pi_z \circ \alpha_{s^{2k}}^\varphi(U_i) = z\mu^k \pi(U_1 U_2)^{-k} \pi(U_i) \pi(U_1 U_2)^k \simeq z\mu^k \pi(U_i)$$

for $i = 1, 2$ all $k \in \mathbb{Z}$, and moreover that

$$\pi_z \circ \alpha_{s^{2k+1}}^\varphi(U_1) = \pi_z \circ \alpha_{s^{2k}}^\varphi(\varphi(s, x_1)U_2) \simeq \varphi(s, x_1)z\mu^k \pi(U_2)$$

for all $k \in \mathbb{Z}$. Applying the same strategy as in [4, proof of Theorem 2.3], using that the point spectrum of V_2 is countable, it is not hard to see that

$$\Omega := \bigcup_{k \in \mathbb{Z}} (\{z : V_2 \not\simeq z\mu^k V_2\} \cup \{z : V_1 \not\simeq \varphi(s, x_1)z\mu^k V_2\})$$

is a countable set. Thus, by choosing $z \in \mathbb{T} \setminus \Omega$, we get that $\pi_z \circ \alpha_{s^k}^\varphi \neq \pi_z$ for every $k \in \mathbb{Z} \setminus \{0\}$.

Hence, it follows from [4, Theorem 2.1] that $C^*(\mathbb{F}_2) \rtimes_{\alpha^\varphi} B_2$ is primitive.

All the other implications follow from known results after identifying the crossed products with twisted group C^* -algebras as explained in Example 4.5 below. \square

Question 3.9. For $n \geq 3$, does condition (i) of Theorem 3.2 and 3.4 imply primitivity of $C^*(\mathbb{F}_n) \rtimes_{\alpha^\varphi} B_n$ and $C^*(\mathbb{F}_n) \rtimes_{\alpha^\varphi} P_n$, respectively?

Before we prove the above results in the next section, we compute the cohomology groups:

Proposition 3.10. *We have*

$$H^1(B_n, \text{Hom}(\mathbb{F}_n, \mathbb{T})) \simeq \begin{cases} \mathbb{T} & \text{for } n = 2, \\ \mathbb{T}^2 & \text{for } n \geq 3, \end{cases}$$

where the cohomology class $[\varphi]$ of $\varphi \in Z^1(B_n, \text{Hom}(\mathbb{F}_n, \mathbb{T}))$ is determined by the parameters (3.5) and (3.6) below.

Moreover, for all $n \geq 2$, we have

$$H^1(P_n, \text{Hom}(\mathbb{F}_n, \mathbb{T})) = Z^1(P_n, \text{Hom}(\mathbb{F}_n, \mathbb{T})) = \text{Hom}(P_n, \text{Hom}(\mathbb{F}_n, \mathbb{T})) \simeq \mathbb{T}^{\frac{1}{2}n(n-1) \cdot n}.$$

Proof. Let $n \geq 2$ and pick $\varphi \in Z^1(B_n, \text{Hom}(\mathbb{F}_n, \mathbb{T}))$. Computations using (2.1) and the fact that $\varphi(s_i s_{i+1} s_i, x_j) = \varphi(s_{i+1} s_i s_{i+1}, x_j)$ for all $1 \leq i \leq n-2$ and $1 \leq j \leq n$ show the following: By taking $j = i$ or $i+2$ we get that

$$(3.1) \quad \varphi(s_{i+1}, x_i) = \varphi(s_i, x_{i+2}) \text{ for all } 1 \leq i \leq n-2.$$

By letting $j = i+1$ and using (3.1) we get that

$$(3.2) \quad \varphi(s_i, x_i)\varphi(s_i, x_{i+1}) = \varphi(s_{i+1}, x_{i+1})\varphi(s_{i+1}, x_{i+2}) \text{ for all } 1 \leq i \leq n-2,$$

that is, $\varphi(s_i, x_i)\varphi(s_i, x_{i+1})$ takes the same value for all $1 \leq i \leq n-1$. Finally, we get that

$$(3.3) \quad \varphi(s_i, x_j) = \varphi(s_{i+1}, x_j) \text{ for all } 1 \leq i \leq n-2 \text{ and all } j \notin \{i, i+1, i+2\}.$$

If $n \geq 3$, then computations using (2.1) and the fact that $\varphi(s_i s_j, x_k) = \varphi(s_j s_i, x_k)$ whenever $|i - j| \geq 2$ and $1 \leq k \leq n$ show that

$$(3.4) \quad \varphi(s_i, x_k) = \varphi(s_i, x_\ell) \text{ for all } 1 \leq i \leq n-2 \text{ and all } k, \ell \notin \{i, i+1\}.$$

By combining (3.1), (3.2), (3.3), and (3.4), we see that there is $\mu_1 \in \mathbb{T}$ such that

$$(3.5) \quad \mu_1 = \varphi(s_i, x_i) \varphi(s_i, x_{i+1}) \text{ for all } 1 \leq i \leq n-1,$$

and if $n \geq 3$, there is $\mu_2 \in \mathbb{T}$ such that

$$(3.6) \quad \mu_2 = \varphi(s_i, x_j) \text{ for all } 1 \leq i \leq n-1 \text{ and all } j \notin \{i, i+1\}.$$

These two values, together with a choice of $\varphi(s_i, x_i)$ or $\varphi(s_i, x_{i+1})$ for every i , determine φ . Further computations show that $B^1(B_n, \text{Hom}(\mathbb{F}_n, \mathbb{T}))$ consists of all functions h satisfying

$$h(s_i, x_i) h(s_i, x_{i+1}) = h(s_i, x_j) = 1 \text{ for all } 1 \leq i \leq n-1 \text{ and all } j \notin \{i, i+1\}.$$

Indeed, if $h(a, x) = f(\alpha_a(x)) \overline{f(x)}$ for some $f \in \text{Hom}(\mathbb{F}_n, \mathbb{T})$, then

$$h(s_i, x_{i+1}) = f(x_i) \overline{f(x_{i+1})} = \overline{h(s_i, x_i)} \text{ for all } 1 \leq i \leq n-1,$$

and $h(s_i, x_j) = f(x_i) \overline{f(x_i)} = 1$ if $j \notin \{i, i+1\}$. Therefore, for $n \geq 3$ the values $\mu_1, \mu_2 \in \mathbb{T}$ determine the class of φ in $H^1(B_n, \text{Hom}(\mathbb{F}_n, \mathbb{T}))$, and two distinct such choices give rise to nonsimilar 1-cocycles. For $n = 2$, the class of φ is determined by the single parameter μ_1 .

The last statement follows from a fairly straightforward calculation, showing that for all $a \in P_n$, there is some $x \in \mathbb{F}_n$ such that $\alpha(a) = \text{Ad } x$. That is, the action α of P_n on $\text{Hom}(\mathbb{F}_n, \mathbb{T})$ is trivial. \square

Lemma 3.11. *Let $\varphi \in Z^1(B_n, \text{Hom}(\mathbb{F}_n, \mathbb{T}))$ and z is the generator of $Z(B_n)$ defined in (1.2). For every $1 \leq i \leq n$ and $1 \leq j \leq n-1$, we have*

$$(3.7) \quad \mu^\varphi = \varphi(z, x_i) = \varphi(s_j, x_1 \cdots x_n)^{n-1}.$$

Moreover, we note that z acts trivially on $\text{Hom}(\mathbb{F}_n, \mathbb{T})$, so that

$$(3.8) \quad \varphi(za, x) = \varphi(z, x) \varphi(a, x) \text{ and } \varphi(z, x) = \varphi(z, \alpha_a(x)) \text{ for all } a \in B_n \text{ and } x \in \mathbb{F}_n.$$

Proof. First, by combining (3.1) and (3.2) we have that

$$\varphi(s_i, x_i x_{i+1} x_{i+2}) = \varphi(s_{i+1}, x_i x_{i+1} x_{i+2}) \text{ for all } 1 \leq i \leq n-2.$$

Then, by using (3.3) we see that $\varphi(s_i, x_1 \cdots x_n)$ takes the same value for all $1 \leq i \leq n-1$, namely the value $\mu_1 \mu_2^{n-2}$, where μ_1, μ_2 are the values described in the proof of Proposition 3.10.

Calculations using (2.1) and (1.4) show that

$$\varphi((s_1, \dots, s_{n-1})^n, x_i) = \varphi(s_1, \dots, s_{n-1}, x_1 \cdots x_n),$$

for all i , so by (1.2) and the fact that all s_i act trivially on $x_1 \cdots x_n$, we get that

$$\varphi(z, x_i) = \prod_{\substack{1 \leq j \leq n-1 \\ 1 \leq k \leq n}} \varphi(s_j, x_k) = (\mu_1 \mu_2^{n-2})^{n-1},$$

and hence (3.7) holds. Showing (3.8) is straightforward from (2.1) and (1.3). \square

4. PROOFS VIA TWISTED GROUP C^* -ALGEBRAS

Let H be a normal subgroup of a discrete group G , and let σ be a 2-cocycle on G . Following [6, Definition 4.5], we say that $g \in G$ is σ -regular with respect to H if $\sigma(g, x) = \sigma(x, g)$ whenever $x \in H$ and $gx = xg$. Moreover, (G, H, σ) satisfies the relative Kleppner condition if $C_H(g) = \{xgx^{-1} : x \in H\}$ is infinite whenever $g \in G \setminus H$ is σ -regular with respect to H .

We refer to [6, Section 4] for a brief discussion on the connection between the relative Kleppner condition and freely acting (or equivalently, properly outer) automorphisms of the twisted group von Neumann algebra. In this paper we will repeatedly apply the following:

Proposition 4.1 ([6, Corollary 4.11]). *Let G be a discrete group with a normal subgroup H . Let σ be a 2-cocycle on G , and denote its restriction to H by σ' . Assume moreover that (G, H, σ) satisfies the relative Kleppner condition.*

If $C_r^(H, \sigma')$ is simple (resp. has a unique tracial state), then $C_r^*(G, \sigma)$ is simple (resp. has a unique tracial state).*

To prove Theorem 3.2 and 3.4 we will identify the crossed products with reduced twisted group C^* -algebras as explained in Section 2.2. We first need the following result:

Lemma 4.2. *Suppose that $g \in \mathbb{F}_n \rtimes_\alpha B_n$. Then $C_{\mathbb{F}_n}(g)$ is finite if and only if $g \in Z(\mathbb{F}_n \rtimes_\alpha B_n)$.*

Proof. Assume that $C_{\mathbb{F}_n}(g)$ is finite. It is explained in [6, Proof of Proposition 4.18] that $C_{\mathbb{F}_n}(g) = \{g\}$, but for the convenience of the reader we sketch the argument. Let $g' \in C_{\mathbb{F}_n}(g)$, so $g' = xgx^{-1}$ for some $x \in \mathbb{F}_n$. Then we have $g^{-1}g' = (g^{-1}xg)x^{-1} \in \mathbb{F}_n$. Moreover, $C_{\mathbb{F}_n}(g^{-1}g') \subseteq C_{\mathbb{F}_n}(g)^{-1}C_{\mathbb{F}_n}(g') = C_{\mathbb{F}_n}(g)^{-1}C_{\mathbb{F}_n}(g)$, so

$$|C_{\mathbb{F}_n}(g^{-1}g')| \leq |C_{\mathbb{F}_n}(g)^{-1}C_{\mathbb{F}_n}(g')| \leq |C_{\mathbb{F}_n}(g)|^2 < \infty.$$

Since \mathbb{F}_n is icc, we must have $g^{-1}g' = e$. Thus, $g' = g$, that is, $C_{\mathbb{F}_n}(g) = \{g\}$, and it follows that g belongs to the centralizer of \mathbb{F}_n in $\mathbb{F}_n \rtimes_\alpha B_n$.

Next, let $y \in \mathbb{F}_n$ and $a \in B_n$ so that $g = ya \in \mathbb{F}_n \rtimes_\alpha B_n$. Since g belongs to the centralizer, it commutes with x_i for all i , i.e., $ya x_i a^{-1} y^{-1} = x_i$, that is, $\alpha(a)(x_i) = y^{-1} x_i y$ for all i . Then

$$x_1 \cdots x_n = \alpha(a)(x_1 \cdots x_n) = y^{-1} x_1 \cdots x_n y,$$

meaning that y commutes with $x_1 \cdots x_n$, and thus $y = (x_1 \cdots x_n)^k$ for some $k \in \mathbb{Z}$ (recall that the centralizer of a single element in a free group is a cyclic subgroup). This again means by (1.3) that $\alpha(a) = \alpha(z^k)$, that is, $a = z^k$ because of injectivity of α . Hence, by (1.5),

$$g = ya = (x_1 \cdots x_n)^k z^k = (x_1 \cdots x_n z)^k \in Z(\mathbb{F}_n \rtimes_\alpha B_n). \quad \square$$

The analogous result and proof also holds for P_n in place of B_n .

Lemma 4.3. *Let $\varphi \in Z^1(B_n, \text{Hom}(\mathbb{F}_n, \mathbb{T}))$. The following are equivalent:*

- (i) $\varphi(z, x_j)$ is nontorsion for some $1 \leq j \leq n$,
- (ii) $(\mathbb{F}_n \rtimes_\alpha B_n, \mathbb{F}_n, \sigma^\varphi)$ satisfies the relative Kleppner condition,
- (iii) $(\mathbb{F}_n \rtimes_\alpha B_n, \sigma^\varphi)$ satisfies Kleppner's condition.

Proof. Assume that $\varphi(z, x_j)$ is nontorsion for some $1 \leq j \leq n$. Suppose that $g \notin \mathbb{F}_n$ and $C_{\mathbb{F}_n}(g)$ is finite. By Lemma 4.2 we have that $g \in Z(\mathbb{F}_n \rtimes_\alpha B_n)$, so (1.5) gives that $g = (x_1 \cdots x_n)^k z^k$ for some $k \in \mathbb{Z} \setminus \{0\}$. Then, using (3.8)

$$\sigma^\varphi(g, x_j) \overline{\sigma^\varphi(x_j, g)} = \varphi(z^k, x_j) = \varphi(z, x_j)^k \neq 1.$$

Hence, $(\mathbb{F}_n \rtimes_\alpha B_n, \mathbb{F}_n, \sigma^\varphi)$ satisfies the relative Kleppner condition.

Assume next that $\varphi(z, x_j)$ is torsion for all j , that is, there exists $k \in \mathbb{Z} \setminus \{0\}$ such that $\varphi(z^k, x_j) = \varphi(z, x_j)^k = 1$ for all j by (3.8). Set

$$g = (x_1 \cdots x_n)^{k(n-1)} z^{k(n-1)} \in Z(\mathbb{F}_n \rtimes_\alpha B_n).$$

We will show that g is σ^φ -regular, i.e., that $\sigma^\varphi(g, ya) = \sigma^\varphi(ya, g)$ for all $y \in \mathbb{F}_n$ and all $a \in B_n$. Note that

$$\sigma^\varphi(g, ya) \overline{\sigma^\varphi(ya, g)} = \varphi(z^{k(n-1)}, y) \overline{\varphi(a, (x_1 \cdots x_n)^{k(n-1)})}.$$

Since z is central, $\varphi(z^{k(n-1)}, y) = \varphi(z^k, y)^{n-1} = 1$. By applying (3.7) and (3.8), we get that

$$\varphi(s_i, x_1 \cdots x_n)^{k(n-1)} = \varphi(z^k, x_1 \cdots x_n) = 1$$

for all i , so $\varphi(a, x_1 \cdots x_n)^{k(n-1)} = 1$ for all $a \in B_n$ as well. Thus g is σ^φ -regular, so $(\mathbb{F}_n \rtimes_\alpha B_n, \sigma^\varphi)$ does not satisfy Kleppner's condition.

We skip the proof of (ii) \implies (iii) here, since it follows from the proof of Theorem 3.2. \square

Proof of Theorem 3.2. We identify the two algebras in question with $C_r^*(\mathbb{F}_n \rtimes_\alpha B_n, \sigma^\varphi)$ and $W^*(\mathbb{F}_n \rtimes_\alpha B_n, \sigma^\varphi)$, and recall that the latter is a factor if and only if $(\mathbb{F}_n \rtimes_\alpha B_n, \sigma^\varphi)$ satisfies Kleppner's condition (cf. [19]). Then (iv) \implies (i) immediately follows from Lemma 4.3.

Next, (i) \implies (ii) and (i) \implies (iii) follow from Lemma 4.3 and Proposition 4.1, together with the well-known fact that $C_r^*(\mathbb{F}_n)$ is simple and has a unique trace (cf. [26]).

Finally, (ii) \implies (iv) and (iii) \implies (iv) holds as Kleppner's condition is necessary both for simplicity and uniqueness of trace, see [6, Section 2.3] (this also takes care of the last implication from Lemma 4.3). \square

Lemma 4.4. *Let $\varphi \in Z^1(P_n, \text{Hom}(\mathbb{F}_n, \mathbb{T}))$. Consider the following conditions:*

- (i) $\varphi(z, x_k)$ is nontorsion for some $1 \leq k \leq n$,
- (ii) $(\mathbb{F}_n \rtimes_\alpha P_n, \mathbb{F}_n, \sigma^\varphi)$ satisfies the relative Kleppner condition,
- (iii) $(\mathbb{F}_n \rtimes_\alpha P_n, \sigma^\varphi)$ satisfies Kleppner's condition,
- (iv) $\varphi(a_{ij}, x_1 \cdots x_n)$ or $\varphi(z, x_k)$ is nontorsion for some $1 \leq i < j \leq n$ or $1 \leq k \leq n$.

Then (i) \iff (ii) \implies (iii) \iff (iv).

Proof. First, (i) \implies (ii) follows by the same argument as in Lemma 4.3, applying Lemma 4.2. For the converse, if $\varphi(z, x_i)$ is torsion for all i , we can find $\ell \in \mathbb{Z} \setminus \{0\}$ such that $\varphi(z, x_i)^\ell = 1$ for all i , and then choose $g = (x_1 \cdots x_n)^\ell z^\ell$. Then $C_{\mathbb{F}_n}(g) = \{g\}$ and $\sigma^\varphi(g, x) = \sigma^\varphi(x, g)$ for all $x \in \mathbb{F}_n$.

By Remark 2.1, to determine Kleppner's condition, it is enough to deal with central elements, i.e., of the form $g = (x_1 \cdots x_n)^\ell z^\ell$ for $\ell \in \mathbb{Z}$. We note that

$$\overline{\sigma^\varphi(g, a_{ij})} \sigma^\varphi(a_{ij}, g) = \varphi(a_{ij}, (x_1 \cdots x_n)^\ell) = \varphi(a_{ij}, x_1 \cdots x_n)^\ell$$

for all $1 \leq i < j \leq n$ and

$$\sigma^\varphi(g, x_k) \overline{\sigma^\varphi(x_k, g)} = \varphi(z^\ell, x_k) = \varphi(z, x_k)^\ell$$

for all $1 \leq k \leq n$. Hence, we conclude that (iii) \implies (iv), and then (i) \implies (iv) is obvious. \square

Proof of Theorem 3.4. This goes along similar lines as the proof of Theorem 3.2, except we do not have that Kleppner's condition implies the relative Kleppner condition for $n \geq 3$. For $n = 2$, see Example 4.5. \square

Example 4.5. When $n = 2$, then $(\mathbb{F}_2 \rtimes_\alpha B_2, \sigma^\varphi)$ satisfies Kleppner's condition precisely when $\mu = \mu^\varphi$ is nontorsion. This completes the proof of first statement of Theorem 3.8 since primitivity implies triviality of the center, which again implies Kleppner's condition by [22, Corollary 2.8].

For the second statement of Theorem 3.8, recall from (1.7) that $\mathbb{F}_2 \rtimes_\alpha P_2 \simeq \mathbb{F}_2 \times \mathbb{Z}$, so

$$C^*(\mathbb{F}_2) \rtimes_{\alpha^\varphi} P_2 \simeq C^*(\mathbb{F}_2 \times \mathbb{Z}, \sigma)$$

for some 2-cocycle σ on $\mathbb{F}_2 \times \mathbb{Z}$. This C^* -algebra is discussed in [22, Example 3.11], where it is shown to be primitive if and only if at least one of $\nu_1 = \varphi(a, x_1)$, $\nu_2 = \varphi(a, x_2)$ is nontorsion, which is again equivalent with Kleppner's condition.

Finally, the above also takes care of the very last statement of Theorem 3.2.

5. MORE ON TWISTED GROUP C^* -ALGEBRAS AND THE SECOND COHOMOLOGY GROUP

The nontrivial 2-cocycles on B_n are the ones lifted from S_n , that is,

$$H^2(B_n, \mathbb{T}) = H^2(S_n, \mathbb{T}) = \begin{cases} 1 & \text{if } n \leq 3, \\ \mathbb{Z}/2\mathbb{Z} & \text{if } n \geq 4. \end{cases}$$

Let $n \geq 4$ and σ denote a nontrivial 2-cocycle on B_n . The universal C^* -algebra generated by a set of unitaries $\{U_i\}_{i=1}^{n-1}$ subject to the relations

$$\begin{aligned} U_i U_{i+1} U_i &= U_{i+1} U_i U_{i+1} \text{ for all } 1 \leq i \leq n-2, \\ U_i U_j &= -U_j U_i \text{ when } |i-j| \geq 2. \end{aligned}$$

is isomorphic to $C^*(B_n, \sigma)$. Since the image of $Z(B_n)$ is trivial under the quotient map $B_n \rightarrow S_n$, every element of $Z(B_n)$ is σ -regular. It follows that (B_n, σ) never satisfies Kleppner's condition, so $C^*(B_n, \sigma)$ and $C_r^*(B_n, \sigma)$ both have nontrivial centers.

Lemma 5.1. *The second cohomology groups of A_{n+1} and P_{n+1} are*

$$\begin{aligned} H^2(A_{n+1}, \mathbb{T}) &\simeq H^2(B_n, \mathbb{T}) \times H^1(B_n, \text{Hom}(\mathbb{F}_n, \mathbb{T})), \\ H^2(P_{n+1}, \mathbb{T}) &\simeq H^2(P_n, \mathbb{T}) \times H^1(P_n, \text{Hom}(\mathbb{F}_n, \mathbb{T})), \end{aligned}$$

and every 2-cocycle on P_{n+1} is similar to one given by

$$(5.1) \quad \sigma(xa, yb) = \varphi(a, y)\omega(a, b) \text{ for } a, b \in P_n \text{ and } x, y \in \mathbb{F}_n,$$

where $\varphi \in Z^1(P_n, \text{Hom}(\mathbb{F}_n, \mathbb{T}))$ and ω is a 2-cocycle on P_n .

Moreover, if σ_1 and σ_2 are 2-cocycles on P_{n+1} of the form (5.1), then $\sigma_1 \sim \sigma_2$ if and only if $\varphi_1 \sim \varphi_2$ and $\omega_1 \sim \omega_2$.

The last two statements also hold when replacing P_{n+1} and P_n by A_{n+1} and B_n .

Proof. Since $H^2(\mathbb{F}_n, \mathbb{T})$ is trivial for all n , the first statement follows directly from the Lyndon-Hochschild-Serre spectral sequence (see e.g. [25, p. 715]). The second statement is deduced from [21, Theorem 9.4], and the third statement is explained in [23, 2.1-2.4] (see also [25, Appendix 2]). \square

It now follows from Lemma 5.1 that

$$H^2(A_3, \mathbb{T}) \simeq \mathbb{T}, \quad H^2(A_4, \mathbb{T}) \simeq \mathbb{T}^2, \quad H^2(A_n, \mathbb{T}) \simeq \mathbb{T}^2 \oplus \mathbb{Z}_2 \text{ for } n \geq 5,$$

and by applying some basic summation formulas, we get

$$H^2(P_n, \mathbb{T}) \simeq \mathbb{T}^{\frac{1}{24}n(n-1)(n-2)(3n-1)}.$$

Proposition 5.2. *Let $n \geq 3$ and $\sigma \in Z^2(A_n, \mathbb{T})$. Then the following are equivalent:*

- (i) (A_n, σ) is C^* -simple,
- (ii) (A_n, σ) has the unique trace property,
- (iii) (A_n, σ) satisfies Kleppner's condition.

In particular, A_n belongs to the class \mathcal{K} introduced in [6].

Proof. Let $n \geq 4$ and σ be a 2-cocycle on A_{n+1} of the form (5.1), coming from a pair (φ, ω) . We proceed as in the proof of Lemma 4.3, choosing g the same way, and compute that

$$\sigma(g, ya)\overline{\sigma(ya, g)} = \varphi(z^{k(n-1)}, y)\omega(z^{k(n-1)}, a)\overline{\varphi(a, (x_1 \cdots x_n)^{k(n-1)})\omega(a, z^{k(n-1)})}.$$

It is observed above that all elements of $Z(B_n)$ are ω -regular, so $\omega(z^{k(n-1)}, a) = \omega(a, z^{k(n-1)})$, meaning that the ω disappears, and the rest of the argument goes exactly as in Lemma 4.3 and the proof of Theorem 3.2. \square

Example 5.3. Let σ be a 2-cocycle on A_3 . By Proposition 5.1 and 3.10, we may, up to similarity, assume that $\sigma = \sigma^\varphi$, where $\varphi(s, x_1) = 1$ and $\sigma(s, x_2) = \mu$ for some $\mu \in \mathbb{T}$. By using (1.6), it is straightforward to check that $C^*(A_3, \sigma)$ is the universal C^* -algebra generated by unitaries U_1, U_2 subject to the relation

$$(U_1 U_2)^2 = \mu(U_2 U_1)^2,$$

and by Theorem 3.8, this algebra is primitive if and only if μ is nontorsion.

Proposition 5.4. *Let $\sigma \in Z^2(P_{n+1}, \mathbb{T})$ be of the type described in (5.1), coming from a pair (φ, ω) . Consider the following conditions:*

- (i) $\varphi(z, x_k)$ is nontorsion for some $1 \leq k \leq n$,
- (ii) (P_{n+1}, σ) is C^* -simple,
- (iii) (P_{n+1}, σ) has the unique trace property,
- (iv) at least one of the following hold:
 - $\varphi(a_{ij}, x_1 \cdots x_n)\omega(a_{ij}, z)\overline{\omega(z, a_{ij})}$ is nontorsion for some $1 \leq i < j \leq n$,
 - $\varphi(z, x_k)$ is nontorsion for some $1 \leq k \leq n$.

Then (i) \implies (ii) \implies (iv) and (i) \implies (iii) \implies (iv).

Moreover, if $n = 3$, then (iv) \implies (i).

Proof. A modification of Lemma 4.4 shows that (iv) is equivalent to Kleppner's condition, while (i) is equivalent to the relative Kleppner condition. The argument now goes along the same lines as the proof of Theorem 3.4. \square

6. C^* -SIMPLICITY OF BRAID GROUPS ON INFINITELY MANY STRANDS

The braid group on infinitely many strands is the group B_∞ generated by $\{s_i\}_{i=1}^\infty$ subject to the same relations as defining B_n . Moreover, there are embeddings $B_n \rightarrow B_{n+1}$, given by $s_i \mapsto s_i$ for $1 \leq i \leq n-1$, and it is easy to see that B_∞ coincides with the direct limit of this system. For $m < n \leq \infty$, we will consider B_m as the subgroup of B_n coming from the first m strands.

Recall from (1.2) that for $3 \leq n < \infty$, the center of B_n is generated by an element of length $n(n-1)$, and all generators s_i of B_n are needed to write word that is a central element. Thus, if $m < n$, then every nontrivial element of $Z(B_m)$ does not belong to $Z(B_n)$, that is,

$$(6.1) \quad Z(B_m) \cap Z(B_n) = \{e\}.$$

Therefore, B_∞ has trivial center. In fact, all of its nontrivial conjugacy classes are infinite, since if g has a finite conjugacy class in B_∞ , then there is some $N \geq 3$ such that $g \in B_n$ and g has a finite conjugacy class in B_n for all $n \geq N$. Then $g \in Z(B_n)$ for all $n \geq N$, so $g = e$. Thus $W^*(B_\infty)$ is a factor (see [15, Corollary 5.3] for a few more properties of $W^*(B_\infty)$).

Furthermore, let S_∞ denote the infinite symmetric group, and define the pure braid group on infinitely many strands, P_∞ , as the kernel of the surjection $B_\infty \rightarrow S_\infty$, sending s_i^2 to the identity for all i . Analogously to the above, it is not hard to see that P_∞ coincides with the direct limit of the embeddings $P_n \rightarrow P_{n+1}$ given by $a_{ij} \mapsto a_{ij}$ for $1 \leq i < j \leq n$.

Finally, we define Artin's representation $\alpha: B_\infty \rightarrow \text{Aut } \mathbb{F}_\infty$ similarly as in (1.1).

Lemma 6.1. *The semidirect products $\mathbb{F}_\infty \rtimes_\alpha B_\infty$ and $\mathbb{F}_\infty \rtimes_\alpha P_\infty$ are both C^* -simple.*

Proof. The triple $(\mathbb{F}_\infty \rtimes_\alpha B_\infty, \mathbb{F}_\infty, 1)$ always satisfies the relative Kleppner condition by modifying the argument of Lemma 4.2. Indeed, if there is $g \in \mathbb{F}_\infty \rtimes_\alpha B_\infty$ such that $C_{\mathbb{F}_\infty}(g)$ is finite, then $C_{\mathbb{F}_\infty}(g) = \{g\}$, that is, $g = ya$ for $y \in \mathbb{F}_\infty$, $a \in B_\infty$, and $gx_i = x_i g$ for all i , where $\{x_i\}_{i=1}^\infty$ are the generators of \mathbb{F}_∞ . There is some $N \geq 3$ such that $g \in \mathbb{F}_n \rtimes_\alpha B_n$ for all $n \geq N$, meaning that $C_{\mathbb{F}_n}(g) = \{g\}$ for all $n \geq N$, so $g \in Z(\mathbb{F}_n \rtimes_\alpha B_n)$ for all $n \geq N$ by Lemma 4.2. Then (1.5) says that we must have $a \in Z(B_n)$ for all $n \geq N$, so $a = e$ by (6.1), and hence $g \in \mathbb{F}_\infty$.

Therefore, it follows from Proposition 4.1 that $\mathbb{F}_\infty \rtimes_\alpha B_\infty$ is C^* -simple. Since $\mathbb{F}_\infty \rtimes_\alpha P_\infty$ is normal in $\mathbb{F}_\infty \rtimes_\alpha B_\infty$, it is also C^* -simple by [9, Theorem 1.4]. \square

Theorem 6.2. *The groups P_∞ and B_∞ are both C^* -simple.*

Proof. First, consider the surjection $P_\infty \rightarrow P_\infty$ given by $a_{1,i} \mapsto e$ and $a_{i,j} \mapsto a_{i-1,j-1}$ for all $2 \leq i < j$. This map restricts to surjections $P_{n+1} \rightarrow P_n$ with kernel being the free group \mathbb{F}_n generated by $\{a_{1,j}\}_{j=2}^{n+1}$ (see [18, Corollary 1.23]; this is just Artin's representation with the generators in reversed order). These restrictions give rise to short exact sequences $\mathbb{F}_n \rightarrow P_{n+1} \rightarrow P_n$ compatible with the embeddings $P_n \rightarrow P_{n+1}$, so $P_\infty \rightarrow P_\infty$ becomes its direct limit, and the kernel is the free group \mathbb{F}_∞ on the infinite number of generators $\{a_{1,j}\}_{j=2}^\infty$. Thus we have that $P_\infty \simeq \mathbb{F}_\infty \rtimes_\alpha P_\infty$ via the map $a_{1,i} \mapsto x_{i-1}$ and $a_{i,j} \mapsto a_{i-1,j-1}$ for all $2 \leq i < j$, and the latter group is shown to be C^* -simple in the previous lemma.

To show that B_∞ is C^* -simple, we will explain that $(B_\infty, P_\infty, 1)$ satisfies the relative Kleppner condition, and then use Proposition 4.1 once more to conclude.

So suppose that $g \in B_\infty$ is such that $C_{P_\infty}(g)$ is finite. Arguing as in Lemma 4.2, using that P_∞ is icc (since it is C^* -simple), we get that $C_{P_\infty}(g) = \{g\}$, i.e., $gx = xg$ for all $x \in P_\infty$. Moreover, there is some $N \geq 3$ such that $g \in B_n$ for all $n \geq N$, meaning that $C_{P_n}(g) = \{g\}$ for all $n \geq N$. Since P_n has finite index in B_n , for every $n \geq N$, there is some $k(n) \geq 1$ such that $g^{k(n)} \in P_n$, so if g commutes with all elements of P_n , then $g^{k(n)} \in Z(P_n)$. Therefore, if $N \leq m < n$, then $g^{k(m)k(n)} \in Z(P_m) \cap Z(P_n) = \{e\}$, using (6.1). Since B_n is torsion-free for all n (see e.g. [18, Corollary 1.29]), we must have $g = e$. This completes the proof. \square

ACKNOWLEDGEMENTS

The author is funded by the Research Council of Norway through FRINATEK, project no. 240913. Part of the work was done when attending the research program ‘‘Classification of operator algebras: complexity, rigidity, and dynamics’’ at the Mittag-Leffler Institute outside Stockholm, January–February 2016. Finally, the author would like to thank Fred Cohen, Alex Suciu, and Dan Cohen for helpful e-mail correspondence on braid groups.

REFERENCES

- [1] Emil Artin. Theory of braids. *Ann. of Math. (2)*, 48:101–126, 1947.
- [2] Erik Bédos. A decomposition theorem for regular extensions of von Neumann algebras. *Math. Scand.*, 68(1):108–114, 1991.
- [3] Erik Bédos. Discrete groups and simple C^* -algebras. *Math. Proc. Cambridge Philos. Soc.*, 109(3):521–537, 1991.
- [4] Erik Bédos and Tron Omland. The full group C^* -algebra of the modular group is primitive. *Proc. Amer. Math. Soc.*, 140(4):1403–1411, 2012.
- [5] Erik Bédos and Tron Omland. On twisted group C^* -algebras associated with FC-hypercentral groups and other related groups. *Ergodic Theory Dynam. Systems*, 36(6):1743–1756, 2016.
- [6] Erik Bédos and Tron Omland. On reduced twisted group C^* -algebras that are simple and/or have a unique trace. *J. Noncommut. Geom.*, 12(3):947–996, 2018.
- [7] M. Bachir Bekka and Pierre de la Harpe. Groups with simple reduced C^* -algebras. *Expo. Math.*, 18(3):215–230, 2000.

- [8] Joan S. Birman. *Braids, links, and mapping class groups*, vol. 82 of *Ann. of Math. Stud.* Princeton Univ. Press, Princeton, NJ, 1974.
- [9] Emmanuel Breuillard, Mehrdad Kalantar, Matthew Kennedy, and Narutaka Ozawa. C^* -simplicity and the unique trace property for discrete groups. *Publ. Math. Inst. Hautes Études Sci.*, 126:35–71, 2017.
- [10] Rasmus S. Bryder and Matthew Kennedy. Reduced twisted crossed products over C^* -simple groups. *Int. Math. Res. Not. IMRN*, 6:1638–1655, 2018.
- [11] Man Duen Choi. The full C^* -algebra of the free group on two generators. *Pacific J. Math.*, 87(1):41–48, 1980.
- [12] Wei-Liang Chow. On the algebraical braid group. *Ann. of Math. (2)*, 49:654–658, 1948.
- [13] Daniel C. Cohen and Alexander I. Suci. Homology of iterated semidirect products of free groups. *J. Pure Appl. Algebra*, 126(1-3):87–120, 1998.
- [14] John Crisp and Luis Paris. Artin groups of type B and D . *Adv. Geom.*, 5(4):607–636, 2005.
- [15] Rolf Gohm and Claus Köstler. Noncommutative independence from the braid group \mathbb{B}_∞ . *Comm. Math. Phys.*, 289(2):435–482, 2009.
- [16] Nikolay A. Ivanov and Tron Omland. C^* -simplicity of free products with amalgamation and radical classes of groups. *J. Funct. Anal.*, 272(9):3712–3741, 2017.
- [17] Mehrdad Kalantar and Matthew Kennedy. Boundaries of reduced C^* -algebras of discrete groups. *J. Reine Angew. Math.*, 727:247–267, 2017.
- [18] Christian Kassel and Vladimir Turaev. *Braid groups*, vol. 247 of *Grad. Texts in Math.* Springer, New York, 2008.
- [19] Adam Kleppner. The structure of some induced representations. *Duke Math. J.*, 29:555–572, 1962.
- [20] Adrien Le Boudec. C^* -simplicity and the amenable radical. *Invent. Math.*, 209(1):159–174, 2017.
- [21] George W. Mackey. Unitary representations of group extensions. I. *Acta Math.*, 99:265–311, 1958.
- [22] Tron Omland. Primeness and primitivity conditions for twisted group C^* -algebras. *Math. Scand.*, 114(2):299–319, 2014.
- [23] Tron Omland. C^* -algebras generated by projective representations of free nilpotent groups. *J. Operator Theory*, 73(1):3–25, 2015.
- [24] Judith A. Packer and Iain Raeburn. Twisted crossed products of C^* -algebras. *Math. Proc. Cambridge Philos. Soc.*, 106(2):293–311, 1989.
- [25] Judith A. Packer and Iain Raeburn. On the structure of twisted group C^* -algebras. *Trans. Amer. Math. Soc.*, 334(2):685–718, 1992.
- [26] Robert T. Powers. Simplicity of the C^* -algebra associated with the free group on two generators. *Duke Math. J.*, 42:151–156, 1975.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OSLO, NO-0316 OSLO, NORWAY AND DEPARTMENT OF
COMPUTER SCIENCE, OSLO METROPOLITAN UNIVERSITY, NO-0130 OSLO, NORWAY

Email address: trono@math.uio.no