ESCORWIN INNOVATIONS PVT. LTD.

MOLECULAR DOCKING & INTERACTION ANALYSIS REPORT

AutoDock Vina Engine v1.2.5 | PLIP Profiling | Job ID: amd_mw37pm8w

Date: Thu, 05 Nov 2026 07:16:39 GMT

SAMPLE DOCKING REPORT

SIMULATION INTEGRITY: VERIFIED & CONVERGED

Engine: AutoDock Vina v1.2.5 | Search: Iterated Local Search (16 Passes) | Target PDB: 1M17

1. Executive Computational Summary

Top Binding Affinity ("G)

-9.3 kcal/mol

Exhaustiveness: 16

Inhibition Constant (Ki)

152.1 nM

T = 298.15 K

Ligand Efficiency (LE)

0.3 kcal/mol/atom

"G / Heavy Atoms

2. Target Receptor & Ligand Profile

RECEPTOR: 1M17 (X-ray Diffraction)

Title: Epidermal Growth Factor Receptor (EGFR) Ki...

Organism: Homo sapiens

Resolution: 2.6 Å

Grid Center: [22.01, 0.25, 52.79] Å

Primary Activity: EGFR Inhibitor

LIGAND: Gefitinib (Iressa) (.cdx format)

Formula: C22H24ClFN4O3 | MW: 446.9 g/mol

Calculated LogP: 3.75 (Lipinski"d 5)

H-Bond Donors: 1 | H-Bond Acceptors: 7

Rotatable Bonds: 6 | Heavy Atoms: 31

Lipinski Rule of 5: Satisfied / Drug-like

3. AutoDock Vina Multi-Pose Binding Energetics

Mode / Pose	Affinity ("	RMSD l.b.	RMSD u.b.	Estimated Ki	Ligand Efficiency
Mode 1 (Best)	-9.3 kcal/mol	0.00 Å	0.00 Å	152.1 nM	0.300 kcal/mol/atom
Mode 2	-8.9 kcal/mol	1.55 Å	2.65 Å	298.8 nM	0.287 kcal/mol/atom
Mode 3	-8.5 kcal/mol	2.00 Å	3.25 Å	586.9 nM	0.274 kcal/mol/atom
Mode 4	-8.2 kcal/mol	2.45 Å	3.85 Å	973.9 nM	0.265 kcal/mol/atom
Mode 5	-7.8 kcal/mol	2.90 Å	4.45 Å	1.91 µM	0.252 kcal/mol/atom
Mode 6	-7.3 kcal/mol	3.35 Å	5.05 Å	4.45 µM	0.235 kcal/mol/atom
Mode 7	-6.8 kcal/mol	3.80 Å	5.65 Å	10.35 µM	0.219 kcal/mol/atom
Mode 8	-6.4 kcal/mol	4.25 Å	6.25 Å	20.33 µM	0.206 kcal/mol/atom
Mode 9	-6.0 kcal/mol	4.70 Å	6.85 Å	39.93 µM	0.194 kcal/mol/atom

4. Best Pose (Mode 1) Ligand-Protein Interaction Photograph




Figure 1: Best Pose (Mode 1, "G = -9.3 kcal/mol) Ligand-Protein Interaction Photograph showing active

5. Residue Interaction Profile (PLIP Standard Analysis)

Interaction	Residue	Ligand Atom	Receptor Atom	Distance	Est. Energy	Details
Hydrogen Bond	Leu718 (A)	N1 (Pyridine/Quinazoline)	NH (Main chain)	2.68 Å	-1.85 kcal/mol	Strong directional H-bond stabilizing the core scaffold inside the active pocket (2.68 Å).
Hydrogen Bond	Val726 (A)	O2 (Ether/Carbonyl)	OH (Side chain)	2.90 Å	-2.05 kcal/mol	Strong directional H-bond stabilizing the core scaffold inside the active pocket (2.9 Å).
Hydrogen Bond	Ala743 (A)	N1 (Pyridine/Quinazoline)	NH (Main chain)	3.09 Å	-2.25 kcal/mol	Strong directional H-bond stabilizing the core scaffold inside the active pocket (3.09 Å).
Hydrophobic Contact	Ala743 (A)	C4 (Aromatic ring)	C´/C² (Lipo pocket)	3.42 Å	-0.65 kcal/mol	Van der Waals lipophilic contact with hydrophobic sub-pocket lining (3.42 Å).
Hydrophobic Contact	Lys745 (A)	C7 (Aromatic ring)	C´/C² (Lipo pocket)	3.57 Å	-0.65 kcal/mol	Van der Waals lipophilic contact with hydrophobic sub-pocket lining (3.57 Å).
Hydrophobic Contact	Met793 (A)	C10 (Aromatic ring)	C´/C² (Lipo pocket)	3.72 Å	-0.65 kcal/mol	Van der Waals lipophilic contact with hydrophobic sub-pocket lining (3.72 Å).
Hydrophobic Contact	Pro794 (A)	C13 (Aromatic ring)	C´/C² (Lipo pocket)	3.87 Å	-0.65 kcal/mol	Van der Waals lipophilic contact with hydrophobic sub-pocket lining (3.87 Å).
Pi-Stacking	Lys745 (A)	Centroid 1 (Aromatic Core)	Centroid (Aromatic Ring)	3.82 Å	-1.25 kcal/mol	Face-to-edge / offset parallel Å-Å stacking sta orientation (3.82 Å).
Halogen Bond	Asp855 (A)	Cl1 (Aryl halide)	O (Backbone Carbonyl)	3.24 Å	-0.95 kcal/mol	Halogen bond σ-hole interaction with peptide carbonyl oxygen (3.24 Å).

6. Scientific Mechanistic Interpretation & Discussion

The molecular docking simulation executed via AutoDock Vina v1.2.5 successfully docked the ligand Gefitinib (Iressa) into the active binding pocket of Epidermal Growth Factor Receptor (EGFR) Kinase Domain with Erlotinib (1M17).

Thermodynamic Stability: A top binding affinity of -9.3 kcal/mol confirms strong non-covalent binding complementarity. The calculated inhibition constant (Ki = 152.1 nM) denotes potent sub-micromolar inhibition under standard biochemical conditions (T = 298.15 K).

Binding Mode & Interactions: As illustrated in the interaction photograph (Figure 1), the ligand forms multiple directional hydrogen bonds with conserved active residues (Leu718, Val726, Ala743). In addition, peripheral lipophilic substituents engage the hydrophobic envelope of the receptor, yielding high shape and electrostatic complementarity without steric clashes.

Drug-likeness & Pharmacological Alignment: The ligand satisfies Lipinski's Rule of Five with an estimated LogP of 3.75 and molecular weight of 446.9 g/mol. This scaffold shows favorable pharmacological alignment with the intended activity (EGFR Inhibitor).

Validation Guidance: Computational predictions provide strong structural rationale for subsequent in vitro assays (e.g. enzyme kinetics, surface plasmon resonance, or isothermal titration calorimetry).

OFFICIAL VERIFICATION & CERTIFICATION

Issued by: Escorwin Innovations Pvt. Ltd. (Computational Chemistry & Molecular Modeling Division)

Digital Verification Code: AMD-AMD_MW37PM8W-5275

Computational Protocol: Monte Carlo ILS with Lamarckian Genetic Algorithm energy minimization.

Quality Standard: Compliant with AutoDock Vina v1.2.5 scoring and PLIP v2.3.0 profiling protocol.