The auxiliary torus of 7+1 spacetime

What it is, how it is described, and what it does — a guide for students, with worked examples checked in sections 25 and 26 of the notebook 7+1.nb and by independent Python programs

Repository: github.com/once-ere/7plus1 — prepared for Nash
Notebook: 7+1.nb (Wolfram Language 15.0.1), evaluated Wed 28 Oct 2026 08:57:47
Document generated by tools/make_docs.py on 2026-10-28

Abstract

In the 7+1 model the four auxiliary space-like coordinates $x_4, \dots, x_7$ are periodic: they form a four-dimensional torus, $T^4 = \mathbb{R}^4/(L_c\mathbb{Z})^4$, the AUXILIARY TORUS. This document defines it precisely — as a quotient space, with its metric, its size b ($b^8 = \det(-g_{kl})$), its proper volume $V_\perp = b^4 L_c^4$ and its symmetries — and explains what it does: fields on it are Fourier series, and their momenta along the torus appear in ordinary spacetime as the masses of a Kaluza–Klein tower; the number of levels below a mass μ is $V_\perp \mu^4/(32\pi^2)$; the volume fixes the four-dimensional Newton constant, $8\pi G_4 = \kappa/V_\perp$; today ($M_8 = 10$ TeV) the torus has the proper size $\ell_\perp = 3.0792 \times 10^{-13}$ m. A central worked example shows how gravity changes from the eight-dimensional $1/r^5$ law to Newton's $1/r$ law: the sum over the images of a mass on the torus equals the sum over its Kaluza–Klein modes (Poisson summation), and the long-range force is $\frac{5}{3}G_4$ — the graviton plus the radion — until the stabilizing potential gives the radion its mass. The document follows the size of the torus through the three regimes of the model and its vacuum energy, and compares it with a 4-SPHERE of the same volume: curvature, levels, what a brane sees, Dirac zero modes, and how the size is held. Every equation used is written here. Section 25 of 7+1.nb contains 27 checks (27 True); the independent program tools/ref_torus_brane.py contains 38 (38 True); the 17 groups of numbers printed by both agree to a largest relative difference of 3.8×10^{-10}. The comparison with the sphere is checked in section 26 (33 checks, 33 True) and by tools/ref_radion.py (31, 31 True).

Contents

1. Results at a glance **2**

2. Conventions **3**

3. The definition of the auxiliary torus **3**
3.1 Periodic coordinates, and the torus as a quotient space . . . 3
3.2 Two examples to build intuition . . . 4
3.3 The spacetime of the model, and where the world brane is . . . 4
3.4 The metric, the size b and the proper volume $V_\perp$ . . . 4
3.5 Only the product bL_c is physical . . . 5
3.6 Symmetries . . . 5

4. Fields on the torus: Fourier modes and the Kaluza–Klein tower **6**
4.1 Fourier modes . . . 6
4.2 Example: orthonormality and Parseval's identity . . . 6
4.3 The Laplacian of the torus . . . 6
4.4 Kaluza–Klein masses: a scalar field . . . 7
4.5 Kaluza–Klein masses: dirac16complex . . . 7
4.6 Fermions on the torus: the boundary conditions . . . 8
4.7 The graviton on the torus . . . 8

5. Counting the levels **8**
5.1 Jacobi's four-square theorem . . . 8
5.2 Weyl's law: the number of levels grows like the volume of a ball . . . 8
5.3 Levels below a mass, and sums over levels . . . 9

6. The size of the torus today, and four-dimensional gravity **9**

6.1 From eight to four dimensions: $8\pi G_4 = \kappa/V_\perp$ . . . 9
6.2 The numbers today . . . 10
6.3 Newton's law on the torus: images and Kaluza–Klein modes (worked example) . . . 10
6.4 Graviton and radion: the factor $\frac{5}{3}$, and what the stabilizer changes . . . 11
6.5 If the size changes: the radion . . . 12
7. The life of the torus: its size in the three regimes of the model **12**
7.1 The primordial regime: the torus contracts . . . 12
7.2 Without a stabilizer the torus re-expands . . . 12
7.3 The stabilizer catches the torus . . . 13
7.4 Today: frozen . . . 13
7.5 The far future: frozen forever (or inflating without the stabilizer) . . . 13
8. The torus and the world brane: Kaluza–Klein momentum **13**
8.1 Conserved in the bulk . . . 13
8.2 Not conserved at the brane . . . 13
8.3 The delta function of the torus, and smeared sources . . . 14
9. The vacuum energy of the torus **14**
10. The torus compared with a 4-sphere **14**
10.1 Flat versus curved: the size as a field . . . 14
10.2 Levels and their counting . . . 15
10.3 What a brane sees . . . 15
10.4 Spinors: zero modes only on the torus . . . 15
10.5 Side by side . . . 15
11. Validity and limitations **16**
12. Verification **16**
13. How to reproduce **21**
14. References **22**

1. Results at a glance

question	answer	where
What is the auxiliary torus?	the space of the four auxiliary coordinates $\vec{y} = (x_4, \ldots, x_7)$ with $x_k \simeq x_k + L_c n_k$ ($n_k \in \mathbb{Z}$): $T^4 = \mathbb{R}^4/(L_c\mathbb{Z})^4$, compact, flat, without boundary	§3.1
How big is it?	size b with $b^8 = \det(-g_{kl})$; each circle has the proper circumference bL_c; proper volume $V_\perp = b^4 L_c^4$; only bL_c is physical	§3.4, §3.5
How big is it today?	$\ell_\perp = V_\perp^{1/4} = 1560.45$ GeV$^{-1} = 3.0792 \times 10^{-13}$ m $= 307.9$ fm: about 366 proton radii, or the Bohr radius divided by 172	§6.2
What are its symmetries?	the translations $\vec{y} \to \vec{y} + \vec{c}$ (their charge is the Kaluza–Klein momentum) and the 384 signed permutations of the axes, which fix the world brane at $\vec{y} = 0$	§3.6
What does a field on it look like?	a Fourier series in $\phi_{\vec{n}} = e^{i\vec{q}_{\vec{n}}\cdot\vec{y}}/\sqrt{V_\perp}$, $\vec{q}_{\vec{n}} = 2\pi\vec{n}/L_c$, $\vec{n} \in \mathbb{Z}^4$	§4.1
What does ordinary spacetime see?	a Kaluza–Klein tower of masses $\sqrt{m^2 + \mu_{\vec{n}}^2}$, $\mu_{\vec{n}} = 2\pi|\vec{n}|/\ell_\perp$; the lightest non-zero one is 4.027 MeV; for dirac16complex four brane Dirac fermions per $\vec{n}$	§4.4, §4.5
How many levels?	$r_4(k) = 8\sigma(k) - 32\sigma(k/4)$ vectors with $\|\vec{n}\|^2 = k$; $N(\mu) = V_\perp \mu^4/(32\pi^2)$ levels below μ: 1.17×10^{13} below 5 GeV, 1.88×10^{26} below M_8	§5
What does it do to gravity?	$8\pi G_4 = \kappa/V_\perp$; the force between two masses on the brane goes from $1/r^5$ ($r \ll \ell_\perp$) to $1/r$ ($r \gg \ell_\perp$); images = Kaluza–Klein modes; $\frac{5}{3} = 1 + \frac{2}{3}$ (graviton + radion)	§6
How does its size evolve?	it contracts in the primordial regime, would re-expand ($b \propto \tau^{2/5}$) without the stabilizing potential, is caught by it, and stays frozen today and in the far future	§7
How does the world brane see it?	the brane sits at the point $\vec{y} = 0$; it breaks the translations, so it couples to every Kaluza–Klein level with the same strength $1/\sqrt{V_\perp}$	§8
What is its vacuum energy?	a Casimir energy, for a massless dirac16complex $6.8 \times 10^{34}\,\rho_\Lambda$: one more constant term of the cosmological constant	§9
Why a torus and not a sphere?	a torus is flat: its size costs no curvature energy (a modulus until stabilized), it has Dirac zero modes, and a brane on it keeps 384 symmetries; a 4-sphere of the same volume (radius 1.359×10^{-13} m) is curved, needs $(\Lambda + \kappa U_0) r_0^2 = 6$ to be held, has its first level at 2.903 MeV and no Dirac zero modes	§10

2. Conventions

Units $\hbar = c = 1$ (energies, masses and inverse lengths in GeV; $\hbar c = 1.973269804 \times 10^{-16}$ GeV m). The coordinates of 7+1 spacetime are $X = (x_0, x_1, \ldots, x_7)$: x_0 is time (the direction of the octonion unit $e_0 = 1$), x_1, x_2, x_3 are ordinary space, and $x_4, \ldots, x_7$ are the four auxiliary space-like coordinates, collected in $\vec{y} = (x_4, x_5, x_6, x_7)$ (components $y_k = x_k$, $k = 4, \ldots, 7$); $\vec{x} = (x_1, x_2, x_3)$. Indices: $A, B = 0, \ldots, 7$ for all eight directions, $\mu, \nu = 0, \ldots, 3$ for the directions of the world brane, $k, l = 4, \ldots, 7$ for the directions of the torus. A prime is d/dx_0. The frame metric is $\eta = \mathrm{diag}(+1, -1, \ldots, -1)$, simply the squares of the octonion basis (explained in the next paragraph).

The signature: $\{7, 1, 0\}$ and $\mathrm{diag}(+1, -1, \ldots, -1)$. The 7+1 spacetime has one time-like dimension (x_0, the direction of the octonion unit $e_0 = 1$) and seven space-like dimensions (x_1, x_2, x_3 ordinary space, $x_4, \ldots, x_7$ auxiliary space). Two different objects describe this, and neither may be replaced by the other. (i) The tensor package xAct, used in section 12 of 7+1.nb exactly as in the author's notebook Einstein-Lovelock-7+1-Nash.nb, declares the metric with `DefMetric[{7, 1, 0}, g71[-a, -b], CD, …]`: the list $\{7, 1, 0\}$ declares seven space-like dimensions, one time-like dimension and no degenerate dimension — the numbers of entries $+1$ and -1 of $\eta_{71} = \mathrm{diag}(-1, +1, \ldots, +1)$ (`η71`), the same spacetime written in the sign convention of 4+4.nb (space-like $= +1$, time-like $= -1$). (ii) The matrix $\eta = -\eta_{71} = \mathrm{diag}(+1, -1, \ldots, -1)$ (`η17`) is simply the squares of the octonion basis: $e_0^2 = +1$ for the unit $e_0 = 1$ (time) and $e_i^2 = -1$ for the seven imaginary units $e_1, \ldots, e_7$ (space); more generally it is the real part of the product of two basis units, and it gives the quadratic form of the octonions:

$$\eta_{AB} = \mathrm{Re}(e_A e_B), \qquad e_0^2 = +1, \qquad e_i^2 = -1 \quad (i = 1, \ldots, 7),$$
$$\mathrm{Re}(x\,x) = x_0^2 - x_1^2 - \cdots - x_7^2 = \eta_{AB}\, x^A x^B \qquad (x = x^A e_A \in \mathbb{O}).$$

η gives the signs of the Clifford relation $\Gamma_A \Gamma_B + \Gamma_B \Gamma_A = 2\,\eta_{AB}\, 1_{16}$ of the real 16×16 Dirac matrices: time squares to $+1_{16}$ and each of the seven space directions to -1_{16}, exactly like the octonion units (in the notation $Cl(p, q)$, with p generators squaring to $+1$ and q to -1, the algebra $Cl(1, 7) \cong \mathbb{R}(16)$). It also gives the signs of the metric components: $g_{00} > 0$ along the time-like direction, negative squares along the space-like directions. Every component computed by 7+1.nb follows from the component matrix of the metric; the declared list fixes for xAct the sign of the determinant of the metric, `SignDetOfMetric[g71]` $= -1$, which is the same in both sign conventions, since $\det \eta = \det \eta_{71} = -1$. 7+1.nb checks `SignatureOfMetric[g71]` $= \{7, 1, 0\}$, `SignDetOfMetric[g71]` $= -1$, $\mathrm{Re}(e_A e_B) = \eta_{AB}$ and $\mathrm{Re}(x\,x) = \eta_{AB} x^A x^B$.

Physical constants (as in sections 17–24 of 7+1.nb): the reduced eight-dimensional Planck mass $M_8 = 10$ TeV, with $\kappa = 8\pi G_8 = M_8^{-6}$ (G_8 the Newton constant of eight dimensions); the reduced four-dimensional Planck mass $M_{\mathrm{Pl}} = 2.435 \times 10^{18}$ GeV, $8\pi G_4 = M_{\mathrm{Pl}}^{-2}$; the radion mass $m_r = 10$ GeV; the Higgs mass $m_h = 125.20$ GeV.

3. The definition of the auxiliary torus

3.1 Periodic coordinates, and the torus as a quotient space

Definition. The four auxiliary coordinates are periodic with the same coordinate period $L_c > 0$:

$$x_k \simeq x_k + L_c\, n_k \qquad (k = 4, \ldots, 7, \quad n_k \in \mathbb{Z}).$$

Two points $\vec{y}$ and $\vec{y}\,'$ of $\mathbb{R}^4$ describe the SAME point of the auxiliary space when their difference is a vector of the lattice $\Lambda = L_c \mathbb{Z}^4$. The auxiliary space is the set of these classes — the quotient space

$$T^4 = \mathbb{R}^4/\Lambda, \qquad \Lambda = L_c\mathbb{Z}^4 = \{L_c \vec{n} :\ \vec{n} \in \mathbb{Z}^4\}, \qquad \vec{y} \sim \vec{y}\,' \iff \vec{y} - \vec{y}\,' \in \Lambda,$$

the four-dimensional torus. Every class has exactly one representative in the FUNDAMENTAL DOMAIN

$$F = [0, L_c)^4 \qquad (\text{equivalently } [-\tfrac{L_c}{2}, \tfrac{L_c}{2})^4),$$

a hypercube whose opposite faces are glued together. The torus is the product of four circles, $T^4 = S^1 \times S^1 \times S^1 \times S^1$ with $S^1 = \mathbb{R}/L_c\mathbb{Z}$. A function on the torus is the same thing as a function on $\mathbb{R}^4$ that is periodic in each coordinate,

$$f(\vec{y} + L_c \vec{e}_k) = f(\vec{y}) \qquad (k = 4, \ldots, 7),$$

($\vec{e}_k$ the unit vector along y_k). Three properties matter throughout: the torus is **compact** (it has a finite volume), it is **flat** (its metric below has zero curvature), and it has **no boundary** and **no special point** — every point looks like every other, because the translations $\vec{y} \to \vec{y} + \vec{c}$ are symmetries. That is why the world brane can be put at the point $\vec{y} = 0$ without loss of generality.

3.2 Two examples to build intuition

Example 3.1 — the circle, and distances on the torus

The circle $S^1 = \mathbb{R}/L_c\mathbb{Z}$ is the interval $[0, L_c)$ with its two ends identified: walking past $y = L_c$ brings you back to $y = 0$. With the metric $b^2 dy^2$ (section 3.4) the proper distance between two points is the length of the SHORTER way around,

$$d(y, y') = b \min_{n\in\mathbb{Z}} |y - y' - nL_c|, \qquad d(\vec{y}, \vec{y}') = b \min_{\vec{n}\in\mathbb{Z}^4} |\vec{y} - \vec{y}' - L_c\vec{n}| \ \text{ on } T^4.$$

For $y = 0.9\,L_c$ and $y' = 0.1\,L_c$ the distance is $b \cdot |0.9L_c - 0.1L_c - L_c| = 0.2\,bL_c$, not $0.8\,bL_c$; no two points of the circle are farther apart than $bL_c/2$, and no two points of T^4 farther than $\sqrt{4}\,bL_c/2 = bL_c$.

Example 3.2 — the two-torus as a video-game screen

The two-torus $T^2 = \mathbb{R}^2/(L_c\mathbb{Z})^2$ is a square whose left and right edges are glued, and whose top and bottom edges are glued: an object leaving through the right edge re-enters at the left edge at the same height, as in the old arcade screens. The square is flat — straight lines and the angles of triangles are those of the plane — although the surface closes on itself. The auxiliary torus is the same construction with four directions: a hypercube with its four pairs of opposite faces glued.

3.3 The spacetime of the model, and where the world brane is

The spacetime of the model, the BULK, is the product

$$M = \mathbb{R}_{x_0} \times \mathbb{R}^3_{\vec{x}} \times T^4_{\vec{y}} \qquad \text{(time} \times \text{ordinary space} \times \text{auxiliary torus)}.$$

Gravity and the 16-component field dirac16complex live in the bulk. The Standard Model lives on the WORLD BRANE

$$W = \{X \in M :\ x_4 = x_5 = x_6 = x_7 = 0\} = \mathbb{R}_{x_0} \times \mathbb{R}^3_{\vec{x}} \times \{\vec{y} = 0\},$$

the 3+1-dimensional submanifold through the point $\vec{y} = 0$ of the torus. Its metric is the induced metric $h_{\mu\nu}(x) = g_{\mu\nu}(x, \vec{y} = 0)$ ($\mu, \nu = 0, \ldots, 3$). The brane is a point of the torus: everything said below about fields on the torus decides how the brane sees them (section 8).

3.4 The metric, the size b and the proper volume $V_\perp$

Write the metric of the bulk as $ds^2 = g_{AB}\, dX^A dX^B$ and its torus block as g_{kl} ($k, l = 4, \ldots, 7$; negative definite, since the four directions are space-like). **The size b of the torus is defined by**

$$b^8 = \det(-g_{kl}) \qquad (k, l = 4, \ldots, 7), \qquad \text{so that} \qquad dV_T = \sqrt{\det(-g_{kl})}\, d^4y = b^4\, d^4y$$

is the proper volume element of the torus. All metrics of 7+1.nb that describe the cosmology of the model have the form

$$ds^2 = n^2 dx_0^2 - a^2\,(dx_1^2 + dx_2^2 + dx_3^2) - b^2\,(dx_4^2 + dx_5^2 + dx_6^2 + dx_7^2), \qquad a = a(x_0), \quad b = b(x_0),$$

(n the lapse, $n = 1$ in the cosmic time x_0; a the scale factor of ordinary space), so $-g_{kl} = b^2\delta_{kl}$, and b is the scale factor of the auxiliary space. Then each of the four circles has the proper circumference, and the torus the proper volume,

$$C = \int_0^{L_c} b\, dx_k = b\, L_c, \qquad V_\perp = \int_F b^4\, d^4y = b^4 L_c^4, \qquad \ell_\perp = V_\perp^{1/4} = b\, L_c.$$

Today the size is frozen at $b = b_0$, so $V_\perp = b_0^4 L_c^4$ and $\ell_\perp = b_0 L_c$.

Example 3.3 — the volume of the torus for the cosmological metric

For the metric above the torus block is $g_{kl} = -b^2\delta_{kl}$, so $\det(-g_{kl}) = b^8$, and

$$V_\perp = \int_0^{L_c}\int_0^{L_c}\int_0^{L_c}\int_0^{L_c} \sqrt{\det(-g_{kl})}\, dx_4\, dx_5\, dx_6\, dx_7 = b^4 L_c^4,$$

as checked symbolically in section 25.1 of 7+1.nb (and by the Python program).

3.5 Only the product bL_c is physical

The coordinate $\vec{y}$ is a label; a rescaled label describes the same torus. Under $\vec{y} = \lambda\vec{y}'$ ($\lambda > 0$):

$$b^2\, d\vec{y}\cdot d\vec{y} = (\lambda b)^2\, d\vec{y}'\cdot d\vec{y}', \qquad L_c' = \frac{L_c}{\lambda}, \qquad b' = \lambda b, \qquad b'L_c' = bL_c, \qquad b'^4 L_c'^4 = b^4 L_c^4.$$

So the period L_c and the size b are not separately measurable — only the proper circumference bL_c and the volume $V_\perp$ are. A common choice is $b_0 = 1$, which makes $L_c = \ell_\perp$ the proper size today; 7+1.nb keeps both symbols. **Example 3.4:** the torus with $L_c = 1$ (in any unit of coordinate length) and $b_0 = 1560.45\ \text{GeV}^{-1}$ per unit, and the torus with $L_c = 1560.45\ \text{GeV}^{-1}$ and $b_0 = 1$, are the same torus (checked symbolically in section 25.1).

3.6 Symmetries

The isometries of the flat torus (the maps that preserve all distances) are of two kinds:

- the **translations** $\vec{y} \to \vec{y} + \vec{c}$ ($\vec{c} \in \mathbb{R}^4$, modulo Λ). They form the group $U(1)^4$; the conserved charge that belongs to them is the momentum along the torus, the Kaluza–Klein momentum $\vec{n}$ (section 8);
- the **signed permutations** of the four axes, $\vec{y} \to O\vec{y}$ with O a 4×4 matrix that has one entry ± 1 in each row and column. These are exactly the orthogonal matrices that map the lattice Λ onto itself (an orthogonal matrix with integer entries is a signed permutation). There are $2^4 \cdot 4! = 384$ of them (the hyperoctahedral group B_4); they fix the point $\vec{y} = 0$ and with it the world brane.

A rotation by a general angle is NOT a symmetry of the torus: it does not map the lattice to itself. The torus is flat and locally looks like $\mathbb{R}^4$, but globally it is not isotropic. The signed permutations permute the vectors $\vec{n} \in \mathbb{Z}^4$ of each shell $|\vec{n}|^2 = k$, and two vectors are in the same orbit exactly when they have the same multiset of $|n_i|$.

Example 3.5 — the shells of the lattice and their orbits

$k = \lvert\vec{n}\rvert^2$	$r_4(k)$	**orbit sizes**	**one vector of each orbit (in the same order)**
1	8	8	$(1,0,0,0)$
2	24	24	$(1,1,0,0)$
3	32	32	$(1,1,1,0)$
4	24	8 + 16	$(2,0,0,0)$, $(1,1,1,1)$
5	48	48	$(2,1,0,0)$
6	96	96	$(2,1,1,0)$
7	64	64	$(2,1,1,1)$
8	24	24	$(2,2,0,0)$
9	104	8 + 96	$(3,0,0,0)$, $(2,2,1,0)$
10	144	48 + 96	$(3,1,0,0)$, $(2,2,1,1)$
11	96	96	$(3,1,1,0)$
12	96	32 + 64	$(2,2,2,0)$, $(3,1,1,1)$

For example the shell $k = 4$ holds the 8 vectors $\pm 2\vec{e}_i$ and the 16 vectors $(\pm 1, \pm 1, \pm 1, \pm 1)$: they have the same length but are not related by a symmetry. The orbit sizes are printed by section 25.1 of 7+1.nb and agree with the Python program.

4. Fields on the torus: Fourier modes and the Kaluza–Klein tower

4.1 Fourier modes

A field on the torus is periodic in each y_k. A plane wave $e^{i\vec{q}\cdot\vec{y}}$ is periodic only if $e^{iq_kL_c}=1$, i.e. $q_k=2\pi n_k/L_c$ with $n_k\in\mathbb{Z}$: the allowed wave vectors form the DUAL LATTICE $(2\pi/L_c)\mathbb{Z}^4$. The normalized MODES are

$$\phi_{\vec{n}}(\vec{y})=\frac{e^{i\vec{q}_{\vec{n}}\cdot\vec{y}}}{\sqrt{V_\perp}},\qquad \vec{q}_{\vec{n}}=\frac{2\pi\vec{n}}{L_c},\qquad \vec{n}\in\mathbb{Z}^4.$$

($\vec{q}_{\vec{n}}$ is the wave vector in the coordinate $\vec{y}$; the physical momentum along the torus is $\vec{q}_{\vec{n}}/b$, because the proper distance is $b\,|d\vec{y}|$.) They are ORTHONORMAL for the proper volume element and COMPLETE:

$$\int_F b^4d^4y\;\phi^*_{\vec{n}}\,\phi_{\vec{m}}=\delta_{\vec{n}\vec{m}},$$

$$f(\vec{y})=\sum_{\vec{n}\in\mathbb{Z}^4}c_{\vec{n}}\,\phi_{\vec{n}}(\vec{y}),\qquad c_{\vec{n}}=\int_F b^4d^4y\;\phi^*_{\vec{n}}f,\qquad \int_F b^4d^4y\;|f|^2=\sum_{\vec{n}}|c_{\vec{n}}|^2\quad\text{(Parseval)}.$$

The proof of orthonormality is one line per direction: $\int_0^{L_c}e^{2\pi i(m_k-n_k)y_k/L_c}dy_k=L_c\,\delta_{m_kn_k}$, and $b^4L_c^4/V_\perp=1$.

4.2 Example: orthonormality and Parseval's identity

Example 4.1 — three exact integrals

Section 25.2 of 7+1.nb integrates exactly over the fundamental domain: $\int_F b^4\phi^*_{\vec{n}}\phi_{\vec{m}}=1$ for $\vec{n}=\vec{m}=(1,0,-2,3)$, $=0$ for $\vec{n}=(1,0,-2,3)$, $\vec{m}=(1,1,-2,3)$, and $=0$ for $\vec{n}=0$, $\vec{m}=(0,2,0,0)$.

Example 4.2 — a smooth bump and its 81 Fourier coefficients

Take $f(\vec{y})=\prod_{k=4}^{7}\cos^2(\pi y_k/L_c)$, a bump that is largest on the brane ($f=1$ at $\vec{y}=0$) and vanishes on the faces $y_k=\pm L_c/2$. Since $\cos^2\theta=\frac12+\frac14e^{2i\theta}+\frac14e^{-2i\theta}$,

$$f=\prod_{k=4}^{7}\Big(\frac12+\frac14e^{2\pi iy_k/L_c}+\frac14e^{-2\pi iy_k/L_c}\Big),\qquad c_{\vec{n}}=\sqrt{V_\perp}\prod_k w(n_k),\quad w(0)=\tfrac12,\ w(\pm1)=\tfrac14,$$

so exactly $3^4=81$ coefficients are non-zero ($n_k\in\{-1,0,1\}$). Parseval's identity holds:

$$\sum_{\vec{n}}|c_{\vec{n}}|^2=V_\perp\Big(\frac14+\frac{1}{16}+\frac{1}{16}\Big)^4=\Big(\frac38\Big)^4V_\perp=\int_F b^4d^4y\,f^2,$$

because $\int_0^{L_c}\cos^4(\pi y/L_c)\,dy=\frac38L_c$. Section 25.2 checks three coefficients by integration and both sides of the identity.

4.3 The Laplacian of the torus

The Laplacian of the flat torus with the metric $b^2\delta_{kl}$ is

$$\Delta_T=\frac{1}{b^2}\sum_{k=4}^{7}\frac{\partial^2}{\partial y_k^2},\qquad -\Delta_T\,\phi_{\vec{n}}=\frac{|\vec{q}_{\vec{n}}|^2}{b^2}\,\phi_{\vec{n}}=\Big(\frac{2\pi}{bL_c}\Big)^2|\vec{n}|^2\,\phi_{\vec{n}}:$$

the modes are its eigenfunctions, and the eigenvalue depends only on $k=|\vec{n}|^2$ (with the multiplicity $r_4(k)$ of section 5). **Example 4.3:** for $\vec{n}=(2,-1,0,3)$, $|\vec{n}|^2=14$ and $-\Delta_T\phi_{\vec{n}}=(2\pi/(bL_c))^2\cdot14\,\phi_{\vec{n}}$ (section 25.2).

4.4 Kaluza–Klein masses: a scalar field

Take a static torus ($b = b_0$) over flat ordinary spacetime, $g = \mathrm{diag}(1,-1,-1,-1,-b_0^2,-b_0^2,-b_0^2,-b_0^2)$, and a scalar field Φ of mass m in the bulk, with the Klein–Gordon equation

$$(\Box + m^2)\Phi = 0, \qquad \Box\Phi = \frac{1}{\sqrt{|g|}}\,\partial_A\big(\sqrt{|g|}\,g^{AB}\partial_B\Phi\big).$$

For $\Phi = e^{-iEx_0+i\vec{k}\cdot\vec{x}}\,\phi_{\vec{n}}(\vec{y})$ the operator gives a number times Φ:

$$(\Box + m^2)\Phi = \Big(-E^2 + |\vec{k}|^2 + \frac{|\vec{q}_{\vec{n}}|^2}{b_0^2} + m^2\Big)\Phi = 0 \quad\Longrightarrow\quad E^2 = |\vec{k}|^2 + M_{\vec{n}}^2,$$
$$M_{\vec{n}} = \sqrt{m^2 + \mu_{\vec{n}}^2}, \qquad \mu_{\vec{n}} = \frac{|\vec{q}_{\vec{n}}|}{b_0} = \frac{2\pi|\vec{n}|}{\ell_\perp}.$$

An observer who cannot resolve the torus sees, for every $\vec{n}$, a particle of mass $M_{\vec{n}}$: **momentum along the hidden directions looks like mass**. The set of these masses is the KALUZA–KLEIN TOWER; $\vec{n} = 0$ is the ZERO MODE (mass m, constant over the torus). Section 25.2 checks the dispersion relation symbolically for $\vec{n} = (1,-2,0,2)$.

4.5 Kaluza–Klein masses: dirac16complex

For the 16-component field the same happens with four 4-component spinors per mode. With the 4×4 Dirac matrices (Dirac representation) and four anti-Hermitian matrices ρ_k,

$$\gamma_0 = \mathrm{diag}(1,1,-1,-1), \quad \gamma_i = \begin{pmatrix} 0 & \sigma_i \\ -\sigma_i & 0 \end{pmatrix}\ (i=1,2,3), \quad \gamma_5 = i\gamma_0\gamma_1\gamma_2\gamma_3, \quad \gamma_\mu\gamma_\nu + \gamma_\nu\gamma_\mu = 2\eta_{\mu\nu}1_4,$$
$$\rho_4 = i\gamma_0, \quad \rho_5 = -\gamma_1, \quad \rho_6 = -\gamma_2, \quad \rho_7 = -\gamma_3, \qquad \rho_k\rho_l + \rho_l\rho_k = -2\delta_{kl}1_4,$$

(σ_i the Pauli matrices), the sixteen matrices $R_\mu = \gamma_\mu\otimes1_4$ ($\mu = 0,\dots,3$) and $R_k = \gamma_5\otimes\rho_k$ ($k = 4,\dots,7$) satisfy the same Clifford relation $R_AR_B + R_BR_A = 2\eta_{AB}1_{16}$ as the real 16×16 Dirac matrices Γ_A of dirac16complex, so there is a constant unitary 16×16 matrix U with $\Gamma_AU = UR_A$ for all A (both representations are irreducible). Writing the field as $\psi = U\zeta$, $\zeta\in\mathbb{C}^4\otimes\mathbb{C}^4$, separates a 4-component spinor index of the brane from a 4-valued index a of the transverse directions. For $\psi\propto e^{-iEx_0+i\vec{p}\cdot\vec{x}+i\vec{q}\cdot\vec{y}}$ at $b = b_0$ the Dirac equation $(i\Gamma^Ae_A{}^B\partial_B - m)\psi = 0$ of dirac16complex ($\Gamma^A = \eta^{AA}\Gamma_A$, frame $e_A{}^B = \mathrm{diag}(1,1,1,1,b_0^{-1},b_0^{-1},b_0^{-1},b_0^{-1})$) becomes

$$\big(\gamma\cdot p\otimes1_4 + X - m\big)\zeta = 0, \qquad \gamma\cdot p = E\gamma_0 - \sum_{i=1}^{3}p_i\gamma_i, \qquad X = \gamma_5\otimes\frac{\vec{\rho}\cdot\vec{q}}{b_0}, \qquad X^2 = -\frac{|\vec{q}|^2}{b_0^2}\,1_{16},$$

and X anticommutes with every $\gamma_\mu\otimes1_4$. Squaring, $E^2 = |\vec{p}|^2 + m^2 + |\vec{q}|^2/b_0^2$: each momentum vector $\vec{n}$ of the torus gives FOUR Dirac fermions of mass $M_{\vec{n}}$ in ordinary spacetime (section 18.1 of 7+1.nb).

Example 4.4 — the lowest levels today

$k = \vec{n}\,^2$	**momentum vectors** $r_4(k)$	**Dirac fermions** $4r_4(k)$	**mass** $2\pi\sqrt{k}/\ell_\perp$ **(MeV),** $m = 0$
1	8	32	4.02652
2	24	96	5.69437
3	32	128	6.97415
4	24	96	8.05305
5	48	192	9.00358
6	96	384	9.86293
7	64	256	10.6532
8	24	96	11.3887

With $m = 0$ there are 4 massless Dirac fermions (the zero mode), then 32 of mass 4.0265 MeV, 96 of mass 5.6944 MeV, and so on; the spacing of the masses shrinks like $1/\sqrt{k}$, and the number of states grows like k (section 5).

4.6 Fermions on the torus: the boundary conditions

A spinor need not return to itself when it goes once around a circle: it may return to itself or change sign. 7+1.nb uses PERIODIC conditions on all four circles (the modes $e^{i\vec{q}_{\vec{n}}\cdot\vec{y}}$ above; also in the Casimir energy of section 9). It is one of the $2^4 = 16$ possible choices (spin structures of T^4); with an antiperiodic circle k the momenta along it would be shifted to $q_k = 2\pi(n_k + \frac{1}{2})/L_c$ and there would be no zero mode.

4.7 The graviton on the torus

Gravity is also a bulk field. A massless graviton in D spacetime dimensions has $D(D-3)/2$ states: 20 for $D = 8$. Seen from ordinary spacetime they split as follows (counting the states of the little groups):

$$\text{zero mode (massless): } 20 = \underbrace{2}_{\text{graviton } g_{\mu\nu}} + \underbrace{4\times 2}_{\text{4 vectors } g_{\mu k}} + \underbrace{10}_{\text{scalars } g_{kl}},$$

$$\text{each massive level: } 20 = \underbrace{5}_{\text{massive spin 2}} + \underbrace{3\times 3}_{\text{3 massive vectors}} + \underbrace{6}_{\text{scalars}}.$$

The ten massless scalars g_{kl} are the MODULI of the torus: one is its volume (the RADION, section 6.5), nine describe its shape (the angles and ratios of the circles). The four vectors $g_{\mu k}$ are the graviphotons. The stabilizing potential $U(b)$ of 7+1.nb depends on the volume only; the shape moduli and the graviphotons are not given a potential or studied in 7+1.nb. Fields on the brane feel only the induced metric $h_{\mu\nu} = g_{\mu\nu}(x, 0)$, which contains neither of them (section 11).

5. Counting the levels

5.1 Jacobi's four-square theorem

The masses depend only on $k = |\vec{n}|^2$. The number of $\vec{n} \in \mathbb{Z}^4$ with $|\vec{n}|^2 = k$ is given by Jacobi's four-square theorem (1829):

$$r_4(k) = \#\{\vec{n} \in \mathbb{Z}^4 : |\vec{n}|^2 = k\} = 8\sigma(k) - 32\sigma(k/4) \quad (k \geq 1), \qquad r_4(0) = 1, \qquad \sigma(k) = \sum_{d|k} d,$$

with $\sigma(k/4) = 0$ unless 4 divides k. **Example 5.1:** $r_4(4) = 8\,\sigma(4) - 32\,\sigma(1) = 8 \cdot 7 - 32 \cdot 1 = 24 = 8 + 16$ (the two orbits of Example 3.5); $r_4(9) = 8 \cdot 13 = 104 = 8 + 96$. The first values:

k	**0**	**1**	**2**	**3**	**4**	**5**	**6**	**7**	**8**	**9**	**10**	**11**	**12**	**13**	**14**	**15**	**16**
$r_4(k)$	1	8	24	32	24	48	96	64	24	104	144	96	96	112	192	192	24

Section 25.3 counts all $\vec{n} \in \{-10, \ldots, 10\}^4$ directly and finds $r_4(k)$ for every $k \leq 100$.

5.2 Weyl's law: the number of levels grows like the volume of a ball

The number of lattice vectors in a ball of radius R approaches the volume $\frac{\pi^2}{2}R^4$ of the four-dimensional ball:

$$N(R) = \#\{\vec{n} \in \mathbb{Z}^4 : |\vec{n}| \leq R\} = \sum_{k=0}^{R^2} r_4(k) \;\simeq\; \frac{\pi^2}{2}R^4 \qquad (R \to \infty).$$

R	$N(R)$ **(lattice count)**	$\pi^2 R^4/2$	$N(R)/(\pi^2 R^4/2) - 1$
10	49689	49348.02201	0.00691
30	4000425	3.997189782×10^6	0.000809
100	493490641	4.934802201×10^8	2.11×10^{-5}
300	39972070929	$3.997189782 \times 10^{10}$	4.33×10^{-6}

R	$N(R)$ **(lattice count)**	$\pi^2R^4/2$	$N(R)/(\pi^2R^4/2)-1$
1000	4934805110729	4.934802201×10^{12}	5.90×10^{-7}

The relative deviation falls roughly like $1/R^2$ (the lattice points near the surface of the ball): Weyl's law (1911) for the eigenvalues of the Laplacian.

5.3 Levels below a mass, and sums over levels

A level with $\mu_{\vec n}\le\mu$ has $|\vec n|\le R=\mu\ell_\perp/(2\pi)$, so the number of levels below μ and their density are

$$N(\mu)\simeq\frac{\pi^2}{2}\Big(\frac{\mu\ell_\perp}{2\pi}\Big)^4=\frac{V_\perp\mu^4}{32\pi^2},\qquad dN=\frac{V_\perp\mu^3\,d\mu}{8\pi^2},\qquad \sum_{\vec n}F(\mu_{\vec n})\simeq\frac{V_\perp}{8\pi^2}\int_0^\infty\mu^3F(\mu)\,d\mu.$$

Example 5.2 — how many levels?

With $V_\perp=5.929\times10^{12}$ GeV^{-4} (section 6.2): $N(5\text{ GeV})=1.17335\times10^{13}$ (the lattice count, section 22.2 of 7+1.nb, is 11733511217161: they agree to 5×10^{-7}); $N(m_h)=4.61\times10^{18}$; $N(M_8)=M_{\rm Pl}^2/(32\pi^2M_8^2)=1.88\times10^{26}$, since $V_\perp M_8^4=M_{\rm Pl}^2/M_8^2$.

Example 5.3 — a weighted sum: the levels open to the decay of a Higgs boson

A Higgs boson can decay into a neutrino and any level lighter than m_h, with a phase-space weight $(1-\mu_{\vec n}^2/m_h^2)^2$ (section 24.4 of 7+1.nb). The weighted number of levels is

$$\sum_{\mu_{\vec n}<m_h}\Big(1-\frac{\mu_{\vec n}^2}{m_h^2}\Big)^2\simeq\frac{V_\perp}{8\pi^2}\int_0^{m_h}\mu^3\Big(1-\frac{\mu^2}{m_h^2}\Big)^2d\mu=\frac{V_\perp m_h^4}{8\pi^2}\int_0^1u^3(1-u^2)^2du=\frac{V_\perp m_h^4}{192\pi^2}=7.688\times10^{17},$$

since $\int_0^1u^3(1-u^2)^2du=\frac14-\frac13+\frac18=\frac{1}{24}$. It is smaller than the number $N(m_h)=4.61\times10^{18}$ of levels below m_h because the heavy levels have little phase space. (For a lattice radius $R=100$ the lattice sum agrees with the integral to 10^{-8}; section 24.4.)

6. The size of the torus today, and four-dimensional gravity

6.1 From eight to four dimensions: $8\pi G_4=\kappa/V_\perp$

The Einstein–Hilbert term of the bulk is $S_1=-\frac{1}{2\kappa}\int d^8X\sqrt{|g|}\,R$ ($\kappa=8\pi G_8$; the curvature conventions of 7+1.nb). For a static torus over a four-dimensional spacetime, $g=\mathrm{diag}\big(g^{(4)}_{\mu\nu}(x),-b_0^2\delta_{kl}\big)$, the volume element is $\sqrt{|g|}=b_0^4\sqrt{|g^{(4)}|}$ and, the torus being flat and static, the curvature of the product is that of the four-dimensional factor, $R=R^{(4)}$. Integrating over the torus,

$$S_1=-\frac{b_0^4L_c^4}{2\kappa}\int d^4x\sqrt{|g^{(4)}|}\,R^{(4)}=-\frac{1}{16\pi G_4}\int d^4x\sqrt{|g^{(4)}|}\,R^{(4)},$$
$$8\pi G_4=\frac{\kappa}{V_\perp},\qquad M_{\rm Pl}^2=\frac{1}{8\pi G_4}=\frac{V_\perp}{\kappa}=V_\perp M_8^6.$$

Example 6.1: for the cosmological metric 7+1.nb finds $R=-2(3H_a'+6H_a^2+4H_b'+10H_b^2+12H_aH_b)$ with $H_a=a'/a$, $H_b=b'/b$ (section 11.2 of 7+1.nb); for a static torus ($H_b=0$) this is $-6(H_a'+2H_a^2)$, the scalar curvature of the four-dimensional Friedmann–Robertson–Walker metric $\mathrm{diag}(1,-a^2,-a^2,-a^2)$ in the same conventions (checked by the Python program), and $\sqrt{|g|}=a^3b^4$. A bigger torus means weaker four-dimensional gravity: $G_4=G_8/V_\perp$.

6.2 The numbers today

With $M_8 = 10$ TeV and $M_{\rm Pl} = 2.435 \times 10^{18}$ GeV, $V_\perp = M_{\rm Pl}^2/M_8^6$. For comparison (CODATA 2018): the proton charge radius $r_p = 0.8414$ fm and the Bohr radius $a_0 = 5.29177210903 \times 10^{-11}$ m.

quantity	**value**	**formula**
proper volume	$V_\perp = 5.929225 \times 10^{12}$ GeV^{-4}	$M_{\rm Pl}^2/M_8^6$
proper size (circumference of each circle)	$\ell_\perp = 1560.4487$ GeV^{-1}	$V_\perp^{1/4} = b_0 L_c$
the same in metres	3.07919×10^{-13} m = 307.919 fm	$\ell_\perp \cdot \hbar c$
compared with the proton	$\ell_\perp / r_p = 365.96$	
compared with the hydrogen atom	$a_0/\ell_\perp = 171.86$	
lowest Kaluza–Klein mass	$2\pi/\ell_\perp = 4.02652$ MeV	$\mu_{\vec n}$ for $\lvert\vec n\rvert = 1$
radius of each circle (convention of the Particle Data Group, $V_\perp = (2\pi R_{\rm PDG})^4$)	$R_{\rm PDG} = 248.353$ GeV$^{-1} = 4.9007 \times 10^{-14}$ m	$\ell_\perp/(2\pi)$
fundamental scale in that convention	$M_D = 34.050$ TeV	$M_{\rm Pl}^2 = R_{\rm PDG}^4 M_D^6 \Rightarrow M_D = (2\pi)^{2/3} M_8$

So the auxiliary torus is about 366 times larger than a proton and 172 times smaller than a hydrogen atom. Searches for large extra dimensions at colliders exclude M_D below 5.9 to 11.2 TeV for 6 to 2 extra dimensions (Particle Data Group 2024); $M_D = 34.05$ TeV is above these bounds. The same numbers are printed by sections 17.5 and 22.2 of 7+1.nb.

6.3 Newton's law on the torus: images and Kaluza–Klein modes (worked example)

This example shows, in one formula, how a compact dimension hides itself. Work in LINEARIZED eight-dimensional gravity WITHOUT the stabilizing potential (section 6.4 adds it).

Step 1: Newton's law in eight dimensions. In D spacetime dimensions the Newtonian potential of a static mass density ρ obeys (the potential $\Phi_{\rm N}$ of a static weak field is defined by $g_{00} = 1 + 2\Phi_{\rm N}$; this follows from the linearized Einstein equation)

$$\nabla^2 \Phi_{\rm N} = 8\pi G_D \, \frac{D-3}{D-2} \, \rho, \qquad \nabla^2 \frac{1}{r^{d-2}} = -(d-2)\,\Omega_{d-1}\,\delta^d(\vec r), \qquad \Omega_{d-1} = \frac{2\pi^{d/2}}{\Gamma(d/2)} \quad (d = D-1),$$

(Ω_{d-1} the area of the unit sphere in d dimensions). For $D = 4$: $\nabla^2\Phi_{\rm N} = 4\pi G\rho$ and $\Phi_{\rm N} = -GM/r$. For $D = 8$, $d = 7$: $\Omega_6 = 16\pi^3/15$, $(D-3)/(D-2) = 5/6$, and a point mass M has

$$\Phi_8(r) = -\frac{5}{6}\,\frac{\kappa M}{5\,\Omega_6\, r^5} = -\frac{5\kappa M}{32\pi^3 r^5} \qquad (\kappa = 8\pi G_8).$$

Step 2: images. On the torus the mass at $\vec y = 0$ also sits, as far as the covering space $\mathbb{R}^4$ knows, at every lattice point $\ell_\perp \vec n$ (we set $b_0 = 1$, $L_c = \ell_\perp$). A second mass on the brane, at the distance r along the brane, feels all the images:

$$\Phi_{\rm N}(r) = -\frac{5\kappa M}{32\pi^3} \sum_{\vec n \in \mathbb{Z}^4} \frac{1}{(r^2 + \ell_\perp^2 |\vec n|^2)^{5/2}} = -\frac{5\kappa M}{32\pi^3 \ell_\perp^5}\, S\Big(\frac{r}{\ell_\perp}\Big), \qquad S(r) = \sum_{\vec n \in \mathbb{Z}^4} \frac{1}{(r^2 + |\vec n|^2)^{5/2}}.$$

Step 3: Poisson summation. For a smooth function f on $\mathbb{R}^4$ that decays fast enough, the sum over the lattice equals the sum of its Fourier transform over the dual lattice:

$$\sum_{\vec n \in \mathbb{Z}^4} f(\vec n) = \sum_{\vec m \in \mathbb{Z}^4} \hat f(2\pi \vec m), \qquad \hat f(\vec p) = \int d^4y \; e^{-i\vec p\cdot\vec y} f(\vec y).$$

The four-dimensional Fourier transform of $(r^2 + |\vec y|^2)^{-5/2}$ is

$$\int d^4y \, \frac{e^{-i\vec p\cdot\vec y}}{(r^2 + |\vec y|^2)^{5/2}} = \frac{4\pi^2}{p} \int_0^\infty \frac{y^2 J_1(py)}{(r^2 + y^2)^{5/2}}\, dy = \frac{4\pi^2}{3r}\, e^{-pr}$$

The first step is the Fourier transform of a radial function in four dimensions (J_1 the Bessel function); the second uses the standard integral

$$\int_0^\infty \frac{y^{\nu+1}J_\nu(py)}{(y^2+r^2)^{\mu+1}}\,dy = \frac{p^\mu r^{\nu-\mu}}{2^\mu\,\Gamma(\mu+1)}\,K_{\nu-\mu}(pr) \qquad (\nu=1,\ \mu=\tfrac{3}{2}), \qquad K_{-1/2}(z)=K_{1/2}(z)=\sqrt{\frac{\pi}{2z}}\,e^{-z},$$

(K the modified Bessel function), which gives $\frac{p}{3r}e^{-pr}$ for the radial integral; the Python program checks the whole transform numerically to 10^{-20} for three pairs (p,r). Hence the IDENTITY

$$\boxed{\sum_{\vec n\in\mathbb{Z}^4}\frac{1}{(r^2+|\vec n|^2)^{5/2}} = \frac{4\pi^2}{3r}\sum_{\vec m\in\mathbb{Z}^4} e^{-2\pi r|\vec m|}}$$

— a sum over IMAGES (left) equals a sum over KALUZA–KLEIN MODES (right). Section 25.4 of 7+1.nb evaluates both sides (the left with the shells $r_4(k)$ up to $k=1.5\times10^6$ and a continuum tail, the right with shells up to $k=20000$):

$r/\ell_\perp$	**images (left side)**	**Kaluza–Klein modes (right side)**	**relative difference**
1/4	1046.849375970	1046.849375970	-7.3×10^{-13}
1/2	51.12525179526	51.12525179603	-1.5×10^{-11}
1	13.40960454141	13.40960454217	-5.7×10^{-11}
2	6.579922931695	6.579922932459	-1.2×10^{-10}

Step 4: what it means. Restoring $\ell_\perp$ and using $8\pi G_4=\kappa/V_\perp$, $V_\perp=\ell_\perp^4$:

$$\Phi_{\rm N}(r) = -\frac{5\kappa M}{32\pi^3}\cdot\frac{4\pi^2}{3V_\perp r}\sum_{\vec m} e^{-\mu_{\vec m}r} = -\frac{5}{3}\,\frac{G_4M}{r}\sum_{\vec m\in\mathbb{Z}^4} e^{-\mu_{\vec m}r} = -\frac{5}{3}\,\frac{G_4M}{r}\Big(1+8e^{-2\pi r/\ell_\perp}+24e^{-2\sqrt{2}\pi r/\ell_\perp}+\cdots\Big),$$

with $\mu_{\vec m}=2\pi|\vec m|/\ell_\perp$ and $\frac{5}{32\pi^3}\cdot\frac{4\pi^2}{3}\cdot8\pi=\frac{5}{3}$. Every Kaluza–Klein level adds a YUKAWA potential $e^{-\mu r}/r$ of the same strength as the zero mode; the multiplicities are $r_4(k)$ (8, 24, …).

- **Far from the source** ($r\gg\ell_\perp$) only the zero mode survives: $\Phi_{\rm N}\to-\frac{5}{3}G_4M/r$, a four-dimensional $1/r$ law. Check: $\big(\frac{3r}{4\pi^2}S(r)-1\big)\big/\big(8e^{-2\pi r}\big) = 1.272,\ 1.017,\ 1.001$ at $r/\ell_\perp=1,\ 2,\ 3$ — approaching 1 as the eight lightest levels take over.
- **Near the source** ($r\ll\ell_\perp$) the nearest image dominates and the eight-dimensional law returns, $S(r)=r^{-5}+Z(5/2)+O(r^2)$ with the Epstein value $Z(5/2)=\sum_{\vec n\neq0}|\vec n|^{-5}=\sum_k r_4(k)k^{-5/2}=8(1-4^{-3/2})\zeta(\frac{5}{2})\zeta(\frac{3}{2})=7\zeta(\frac{5}{2})\zeta(\frac{3}{2})=24.531278$ (shell sum: 24.53127769). Check: $(r^5S(r)-1)/(r^5Z(5/2))=0.99703$ at $r/\ell_\perp=1/20$ and 0.98823 at 1/10: at a tenth of the torus size the force differs from the pure eight-dimensional one by $r^5Z(5/2)=0.00025$.

6.4 Graviton and radion: the factor $\frac{5}{3}$, and what the stabilizer changes

Why $\frac{5}{3}G_4$ and not G_4? Without the stabilizing potential the size of the torus is a massless field, the RADION, and it is exchanged between the two masses together with the four-dimensional graviton. In 7+1.nb (section 20.3) the radion $\beta=\ln(b/b_0)$ has the canonical form $\phi=\sqrt{12}\,M_{\rm Pl}\beta$ and couples to brane matter through the trace of its energy–momentum tensor,

$$\mathcal{L}_{\phi W} = \frac{\phi}{\sqrt{3}\,M_{\rm Pl}}\,T^\mu{}_\mu.$$

For two static masses $T^\mu{}_\mu=M\delta^3(\vec x)$; the static field equation $(-\nabla^2+m_r^2)\phi=M\delta^3/(\sqrt{3}M_{\rm Pl})$ gives $\phi=Me^{-m_rr}/(4\pi\sqrt{3}M_{\rm Pl}r)$, and the second mass feels the potential energy

$$V_\phi(r) = -\frac{\phi\,M'}{\sqrt{3}M_{\rm Pl}} = -\frac{MM'}{12\pi M_{\rm Pl}^2}\,\frac{e^{-m_rr}}{r} = -\frac{2}{3}\,\frac{G_4MM'}{r}\,e^{-m_rr}, \qquad \frac{1}{12\pi M_{\rm Pl}^2}=\frac{2}{3}\cdot\frac{1}{8\pi M_{\rm Pl}^2}=\frac{2}{3}G_4.$$

With $m_r = 0$: graviton (1) + radion $(\frac{2}{3}) = \frac{5}{3}$ — exactly the coefficient of the image sum. This checks two results of 7+1.nb at once: $8\pi G_4 = \kappa/V_\perp$ and the normalization $1/(\sqrt{3}M_{\rm Pl})$ of the radion coupling. **With the stabilizing potential** the radion has the mass m_r (10 GeV, a range $1/m_r = 0.1\ \text{GeV}^{-1} = 0.02$ fm, far below $\ell_\perp$), and at distances $r \gg \ell_\perp$, $1/m_r$ only the graviton remains: Newton's law $V = -G_4 MM'/r$ with $G_4 = G_8/V_\perp$. At $r \ll 1/m_r$ (where the potential is negligible) the eight-dimensional law of step 1 holds; between $1/m_r$ and $\ell_\perp$ the force is that of eight-dimensional gravity with a stabilized volume, which 7+1.nb does not compute.

6.5 If the size changes: the radion

Since $G_4 = \kappa/(8\pi b^4 L_c^4)$, a changing torus changes the strength of gravity on the brane: $\dot{G}_4/G_4 = -4H_b$ with $H_b = b'/b$. 7+1.nb holds the size with the STABILIZING POTENTIAL (an energy density in the bulk that depends on the size)

$$U(b) = U_0 + \frac{K_U}{2}\,\ln^2\frac{b}{b_0}, \qquad \beta = \ln\frac{b}{b_0}, \qquad m_r^2 = \frac{\kappa K_U}{12}, \qquad \phi = \sqrt{12}\,M_{\rm Pl}\,\beta,$$

whose minimum b_0 is the size today; m_r is the radion mass. In Einstein gravity the exact equation of the radion is

$$\beta'' + 3H_a\beta' + 4\beta'^2 = \kappa\Big(\frac{\varepsilon}{6} - \frac{p_a}{2} + \frac{p_b}{3}\Big) + \frac{\Lambda}{3},$$

with ε the energy density and p_a, p_b the pressures along ordinary and auxiliary space of everything in the bulk (the potential contributes $\varepsilon = U$, $p_a = -U$, $p_b = -U - \frac{b}{4}U'(b)$, which gives the restoring term $-m_r^2\beta$), and Λ the bulk cosmological constant. Brane fields see the metric $h = (b_0/b)^4 g_E$, with g_E the four-dimensional Einstein-frame metric; this is the origin of the coupling $\phi\,T^\mu{}_\mu/(\sqrt{3}M_{\rm Pl})$ of section 6.4.

7. The life of the torus: its size in the three regimes of the model

7+1.nb solves the coupled field equations of the model (Einstein–Lovelock gravity with the couplings α_2 (Gauss–Bonnet) and α_3 (cubic), dirac16complex, the stabilizing potential, the brane) in three regimes (sections 14, 21–23). Here is what the torus does in each; τ is the time since the singularity, $H_a = a'/a$, $H_b = b'/b$.

7.1 The primordial regime: the torus contracts

Near $x_0 = 0$ power laws $a \propto \tau^p$, $b \propto \tau^q$ solve the vacuum equations of each Lovelock order. In Einstein gravity $3p + 4q = 1$ and $3p^2 + 4q^2 = 1$, so

$$p_1 = \frac{1+2\sqrt{2}}{7} = 0.546918, \qquad q_1 = \frac{2-3\sqrt{2}}{14} = -0.160189.$$

So **ordinary space expands while the torus contracts** — a dynamical compactification. In the Gauss–Bonnet regime ($3p + 4q = 3$) $p_2 = 1.56115$, $q_2 = -0.420861$; closest to the singularity, in the cubic regime, ordinary space expands like $a \propto \tau$ while the torus contracts at the finite rate $H_b \to h_0 = -\sqrt{\alpha_2/(30\alpha_3)} = -182.574$ (for $\alpha_3/\alpha_2^2 = 10^{-6}$, in units $\alpha_2 = 1$); with a gas of dirac16complex, $H_b \to h_* = -\sqrt{\alpha_2/(54\alpha_3)} = -136.083$ (section 21.3 finds -136.083 numerically).

7.2 Without a stabilizer the torus re-expands

Once a gas of dirac16complex dominates, Einstein gravity with a fluid of pressures $p_a = w_a\varepsilon$, $p_b = w_b\varepsilon$ has power laws $a \propto \tau^P$, $b \propto \tau^Q$ with $3P(1+w_a) + 4Q(1+w_b) = 2$. The momenta along the torus are blue-shifted by its contraction, so the gas ends with $(w_a, w_b) = (0, \frac{1}{4})$ and

$$(P, Q) = \Big(0, \frac{2}{5}\Big): \qquad b \propto \tau^{2/5},$$

ordinary space stops expanding and the torus GROWS (section 21.3 finds $\tau H_b = 0.399995$ and $p_b/\varepsilon = 0.249999$ at $H_a = 10^{-14}$). Without the stabilizing potential the 7+1 universe would decompactify.

7.3 The stabilizer catches the torus

With $U(b)$ the radion is caught when the expansion rate falls below m_r and then oscillates about $\beta = 0$ like a four-dimensional massive field in a matter-dominated universe:

$$\beta \simeq \frac{A\,\sin(m_r\tau + \vartheta)}{m_r(\tau - \tau_E)}, \qquad H_E = H_a + 2H_b \simeq \frac{2}{3(\tau - \tau_E)}.$$

Section 22.1 integrates the coupled equations (units $\ell_2 = \sqrt{\alpha_2} = 1$, $m_r = 10^{-3}$, start at $\tau = 56.750$ with $\beta = 0.4596$) and fits $A = 0.4693$, $\vartheta = 2.136$, $\tau_E = -1564.36$ with a largest residual of 0.013 of the amplitude. After the capture the torus is static on average, and the gas of Kaluza–Klein modes (its frozen momenta now masses) is four-dimensional matter.

7.4 Today: frozen

With the torus static the brane obeys the four-dimensional Friedmann equations with $G_4 = G_8/V_\perp$. The radion sits at a displaced minimum,

$$\beta_{\rm eq} = \frac{\Lambda_4 + \kappa f_W/V_\perp + \frac{\kappa}{2}(\varepsilon - 3p_a) + \kappa p_b - 12\alpha_2 H_a^2(H_a' + H_a^2)}{3m_r^2},$$

(Λ_4 the four-dimensional cosmological constant, f_W the brane tension) — numerically $\beta_{\rm eq} = 1.7\times10^{-86}$ from Λ_4 and matter, 5.6×10^{-24} from a tension $f_W = M_8^4$ (section 22.4, $m_r = 10$ GeV): G_4 is constant to that accuracy, as experiments require. The Gauss–Bonnet and cubic corrections are $\alpha_2 H_0^2 = 2.1\times10^{-92}$ and $\alpha_3 H_0^4 = 4.3\times10^{-190}$.

7.5 The far future: frozen forever (or inflating without the stabilizer)

In the de Sitter future ($H_\infty = H_0\Omega_\Lambda^{1/2} = 1.808\times10^{-18}$ s^{-1}) the radion relaxes to $3m_r^2\beta_{\rm eq} = 3H^2 - 12\alpha_2 H^4$, of order H_∞^2/m_r^2 (section 23.2; a numerical test in units $\ell_2 = 1$ reaches $\beta_{\rm eq} = 0.0102041$). Without the stabilizer the far future would instead be eight-dimensional de Sitter space,

$$a = b = e^{Hx_0}, \qquad 21H^2 - 420\alpha_2 H^4 + 2520\alpha_3 H^6 = \Lambda,$$

in which the torus inflates and $G_4 \propto b^{-4}$ falls like e^{-4Hx_0} — excluded by the observed constancy of G.

8. The torus and the world brane: Kaluza–Klein momentum

8.1 Conserved in the bulk

The translations of the torus are symmetries of the bulk, so their charge, the Kaluza–Klein momentum $\vec{n}$, is conserved by every local interaction in the bulk. **Example 8.1:** the overlap of three modes, which is the amplitude of a cubic interaction,

$$\int_F b^4 d^4y\; \phi_{\vec{n}}\,\phi_{\vec{m}}\,\phi_{\vec{l}}^{\,*} = \frac{\delta_{\vec{n}+\vec{m},\vec{l}}}{\sqrt{V_\perp}}$$

is $1/\sqrt{V_\perp}$ for $\vec{n} = (1,0,0,2)$, $\vec{m} = (0,1,-1,0)$, $\vec{l} = (1,1,-1,2) = \vec{n} + \vec{m}$ and 0 for $\vec{l} = (1,1,-1,1)$ (section 25.5).

8.2 Not conserved at the brane

The world brane sits at the single point $\vec{y} = 0$ and is not invariant under the translations. At the brane every mode has the same value, $\phi_{\vec{n}}(0) = 1/\sqrt{V_\perp}$, so a bulk field seen from the brane is

$$\psi(x, \vec{y} = 0) = \frac{1}{\sqrt{V_\perp}} \sum_{\vec{n}\in\mathbb{Z}^4} \chi_{\vec{n}}(x), \qquad \big(\phi_{\vec{n}}\phi_{\vec{m}}\phi_{\vec{l}}^{\,*}\big)(0) = V_\perp^{-3/2} \;\text{ for all } \vec{n},\vec{m},\vec{l},$$

($\chi_{\vec{n}}$ the four-dimensional field of the level $\vec{n}$): an interaction on the brane couples EVERY level with the same strength and can change $\vec{n}$ by any amount. The brane couplings of 7+1.nb (the portals) therefore act on the whole tower at once — the reason why the Higgs portal of section 24 couples every pair of levels.

8.3 The delta function of the torus, and smeared sources

The delta function of the torus contains every mode with the same weight,

$$\delta^4(\vec{y}) = \frac{1}{L_c^4}\sum_{\vec{n}\in\mathbb{Z}^4} e^{i\vec{q}_{\vec{n}}\cdot\vec{y}}, \qquad \int_{-L_c/2}^{L_c/2} e^{-iq_n y}\,\delta(y)\,dy = 1,$$

and its average over the torus (its zero mode) is $1/L_c^4$. A source on the brane with the four-dimensional energy density ρ_4 has the eight-dimensional density $\rho_4\,\delta^4(\vec{y})/b^4$, whose average over the torus is

$$\varepsilon_8 = \frac{1}{V_\perp}\int_F b^4 d^4y\,\frac{\rho_4\,\delta^4(\vec{y})}{b^4} = \frac{\rho_4}{V_\perp}\,:$$

this SMEARING is how the homogeneous equations of 7+1.nb treat the brane (a brane tension f_W gives $\varepsilon = f_W/V_\perp$, $p_a = -f_W/V_\perp$, $p_b = 0$).

9. The vacuum energy of the torus

Quantum fields on the torus have a zero-point energy that depends on its size — a CASIMIR energy. For a field of mass m with periodic conditions, per real bosonic degree of freedom and per unit volume of the brane (section 24.5 of 7+1.nb),

$$\rho^{(1)}_{\rm Cas} = -\frac{2}{\ell_\perp^4}\sum_{\vec{n}\neq 0}\Big(\frac{m\ell_\perp}{2\pi|\vec{n}|}\Big)^4 K_4(m\ell_\perp|\vec{n}|) \;\xrightarrow{m\to 0}\; -\frac{\Gamma(4)}{\pi^4\ell_\perp^4}\,Z(4),$$

$$Z(4) = \sum_{\vec{n}\neq 0}\frac{1}{|\vec{n}|^8} = \sum_{k\geq 1}\frac{r_4(k)}{k^4} = 8(1-4^{-3})\,\zeta(4)\,\zeta(3),$$

(K_4 the modified Bessel function, ζ the Riemann zeta function); a fermion contributes with the opposite sign, and dirac16complex has 16 real degrees of freedom on shell. With $\ell_\perp$ of section 6.2 and the observed dark-energy density $\rho_\Lambda = 2.518\times 10^{-47}$ GeV^4, a massless dirac16complex gives $6.76\times 10^{34}\,\rho_\Lambda$ and the graviton (20 degrees of freedom) $-8.45\times 10^{34}\,\rho_\Lambda$; the dirac16complex term equals ρ_Λ only for $m\ell_\perp = 92.23$ ($m = 59.10$ MeV). While the torus is frozen these energies are constant ($w = -1$): they are terms of the four-dimensional cosmological constant, not a dynamical dark energy.

10. The torus compared with a 4-sphere

Why is the auxiliary space of 7+1 a torus? The clearest way to see what the choice implies is to replace it by the other simple compact four-dimensional space, the 4-SPHERE S^4 of radius r, and compare (section 26 of 7+1.nb). On S^4 the coordinates $x_4, x_5, x_6 \in [0,\pi]$, $x_7 \in [0, 2\pi)$ are angles, and

$$g_{kl} = -r^2\hat{\gamma}_{kl}, \qquad \hat{\gamma} = \mathrm{diag}\big(1, \sin^2 x_4, \sin^2 x_4 \sin^2 x_5, \sin^2 x_4 \sin^2 x_5 \sin^2 x_6\big), \qquad V = \frac{8\pi^2}{3}r^4, \qquad R_{kl} = 3\hat{\gamma}_{kl}.$$

10.1 Flat versus curved: the size as a field

Let the size depend on the point of ordinary spacetime, $b = b_0 e^{\beta(x)}$ (torus) or $r = r_0 e^{\beta(x)}$ (sphere), with the four-dimensional metric rescaled so that Newton's constant stays constant (the Einstein frame):

$$ds^2 = e^{-4\beta(x)} g_{E\mu\nu}dx^\mu dx^\nu - e^{2\beta(x)}\gamma_{kl}dy^k dy^l, \qquad \gamma = b_0^2\delta \;\;(T^4), \quad \gamma = r_0^2\hat{\gamma} \;\;(S^4).$$

Integrated over the extra space, the eight-dimensional action becomes $\sqrt{|g_E|}[-\frac{M_{\rm Pl}^2}{2}R_E + 6M_{\rm Pl}^2(\partial\beta)^2 - V(\beta)]$ with $M_{\rm Pl}^2 = V_0/\kappa$, so $\phi = \sqrt{12}M_{\rm Pl}\beta$ is the canonical radion on both spaces, and

$$V(\beta) = M_{\rm Pl}^2\Big[e^{-4\beta}\big(\Lambda + \kappa U(\beta)\big) - \frac{6\sigma}{r_0^2}e^{-6\beta}\Big], \qquad \sigma = 0\;(T^4), \quad \sigma = 1\;(S^4)$$

(Λ the bulk cosmological constant, U the stabilizing energy density). The flat torus is a MODULUS: for $\Lambda = U = 0$ every size is a solution. The curved sphere is not: its curvature energy $-6M_{\rm Pl}^2 e^{-6\beta}/r_0^2$ drives it to collapse. With $U = U_0 + U_1\beta + \frac{K_U}{2}\beta^2$ a flat four-dimensional vacuum ($V(0) = V'(0) = 0$) needs

$$T^4:\ \Lambda + \kappa U_0 = 0,\ U_1 = 0,\ m_r^2 = \frac{\kappa K_U}{12}; \qquad S^4:\ (\Lambda + \kappa U_0) r_0^2 = 6,\ \kappa U_1 r_0^2 = -12,\ m_r^2 = \frac{\kappa K_U}{12} - 2\,\frac{1}{r_0^2},$$

($m_r^2 = V''(0)/(12M_{\rm Pl}^2)$). A sphere held by Λ alone ($dS_4 \times S^4$, $\Lambda r_0^2 = 9$) is unstable, $m^2 = -6\,H^2$.

10.2 Levels and their counting

	T^4	S^4
eigenfunctions of $-\Delta$	$e^{2\pi i \vec{n}\cdot\vec{y}/\ell_\perp}/\sqrt{V}$	spherical harmonics of degree l
masses	$2\pi\lvert\vec{n}\rvert/\ell_\perp$	$\sqrt{l(l+3)}/r$
multiplicity	$r_4(k) = 8\sigma(k) - 32\sigma(k/4)$	$d_l = \frac{(2l+3)(l+1)(l+2)}{6}$
first multiplicities	$1, 8, 24, 32, 24, \ldots$ $(k = 0, 1, 2, 3, 4)$	$1, 5, 14, 30, 55, 91, \ldots$ $(l = 0, \ldots, 5)$
Weyl's law	$N(\mu) \simeq V\mu^4/(32\pi^2)$	the same; exactly $N = y^2(y^2-1)/12$, $y = L + 2$

Both obey Weyl's law, which depends only on the volume: the number of heavy levels cannot tell the two spaces apart. **Example 10.2:** with the same volume $V_\perp$ (the same G_4), the sphere has the radius $r_{S^4} = (3V_\perp/(8\pi^2))^{1/4} = 1.3595 \times 10^{-13}$ m; below 5 GeV the torus has 11733511217161 states (lattice count), the sphere 11737500461050 (exact), Weyl's law gives 1.17335×10^{13}. The LIGHTEST levels differ: the torus at $2\pi/\ell_\perp = 4.0265$ MeV (8 states), the sphere at $2/r_{S^4} = 2.9030$ MeV (5 states).

10.3 What a brane sees

A brane at $\vec{y} = 0$ of the torus sees every mode with the value $1/\sqrt{V}$: per level it couples to one combination of the $r_4(k)$ modes with the strength $\sqrt{r_4(k)/V}$. A brane at the north pole $x_4 = 0$ of the sphere sees only the ZONAL harmonic of each level,

$$Z_l = \sqrt{\frac{d_l}{V}}\,\frac{C_l^{(3/2)}(\cos x_4)}{C_l^{(3/2)}(1)}, \qquad Z_l(0) = \sqrt{\frac{d_l}{V}}$$

($C_l^{(3/2)}$ the Gegenbauer polynomial): one state per level with the strength $\sqrt{d_l/V}$. The brane keeps 384 symmetries of the torus and 6 of the 10 rotations of the sphere ($SO(4) \subset SO(5)$). On both spaces the brane sees the same induced metric $e^{-4\beta} g_E$, so the radion couples to brane matter in the same way, $\phi T^\mu{}_\mu/(\sqrt{3}M_{\rm Pl})$.

10.4 Spinors: zero modes only on the torus

With periodic spinors the torus has the constant spinors as zero modes of its Dirac operator: dirac16complex gives four massless brane Dirac fermions for $m = 0$ (section 4.5). The sphere has none: the eigenvalues of its Dirac operator are $\pm i(l+2)/r$ with the degeneracy $4\binom{l+3}{3}$ for each sign, the lowest belonging to the Killing spinors, which reach Friedrich's bound $\lvert\lambda\rvert^2 \geq nR/(4(n-1)) = 4/r^2$ ($R = 12/r^2$). On S^4 the lightest dirac16complex states are eight Dirac fermions of mass $\sqrt{m^2 + 4/r^2}$, at least 2.9030 MeV for $r = r_{S^4}$.

10.5 Side by side

property	T^4 **(7+1.nb)**	S^4 **(comparison)**
curvature	flat	$R_{kl} = 3\hat{\gamma}_{kl}$, $R = 12/r^2$
volume	$b^4 L_c^4$	$\frac{8\pi^2}{3} r^4$
size without stabilizer	modulus (any size)	collapses
stabilization	$\Lambda + \kappa U_0 = 0$, $U_1 = 0$	$(\Lambda + \kappa U_0) r_0^2 = 6$, $\kappa U_1 r_0^2 = -12$
radion mass	$\kappa K_U/12$	$\kappa K_U/12 - 2/r_0^2$

property	T^4 (7+1.nb)	S^4 (comparison)
size today ($V = V_\perp$)	$\ell_\perp = 3.079 \times 10^{-13}$ m	$r_{S^4} = 1.359 \times 10^{-13}$ m
lightest level	4.027 MeV, 8 states	2.903 MeV, 5 states
symmetry kept by a brane	384 signed permutations	$SO(4)$
Dirac zero modes	4	none

11. Validity and limitations

- The torus of 7+1.nb is the CUBIC torus: four circles of the same period, at right angles. A general flat torus $\mathbb{R}^4/\Lambda$ with any lattice Λ would have the masses $\mu^2 = \vec{q} \cdot G^{-1}\vec{q}$ (G the metric of the lattice, $\vec{q}$ in the dual lattice); its shape is not dynamical in 7+1.nb.
- Only the volume is stabilized ($U(b)$); the nine shape moduli and the four graviphotons (section 4.7) are not discussed in 7+1.nb.
- Fermions are periodic around all four circles (section 4.6).
- The 4-sphere of section 10 is a comparison worked out in section 26 of 7+1.nb; the model itself uses the torus.
- The Newton-law example (section 6.3) is linearized gravity of a point mass without the stabilizing potential; the stabilizer is added through the radion mass only (section 6.4). The gravitational field of the brane itself is not included.
- The numbers today assume $M_8 = 10$ TeV (and $M_{\rm Pl} = 2.435 \times 10^{18}$ GeV); $\ell_\perp \propto M_8^{-3/2}$, so a different M_8 changes them accordingly.
- Continuum formulas such as $N(\mu) = V_\perp \mu^4/(32\pi^2)$ hold for $\mu \gg 2\pi/\ell_\perp$; their accuracy is that of Weyl's law (section 5.2).

12. Verification

Every worked example of this document is a check of section 25 of 7+1.nb (Wolfram Language 15.0.1; evaluated in a fresh kernel, 0 message cells) and of the independent Python program tools/ref_torus_brane.py (sympy, numpy, mpmath; its own lattice sums, quadratures and Christoffel symbols). The checks of section 25:

#	check	result	section
1	auxiliary torus: for the cosmological metric det(-g_kl) == b^8 (k, l = 4..7), and the proper volume of the fundamental domain [0, Lc)^4, Integrate[Sqrt[det(-g_kl)], {y, 0, Lc}^4], is b^4 Lc^4 = V_perp	True	25.1
2	only the product b Lc of the auxiliary torus is physical: y = lambda y' maps the torus of period Lc and size b to the one of period Lc/lambda and size lambda b; the line element b^2 dy.dy, the proper period b Lc and V_perp = b^4 Lc^4 are unchanged	True	25.1
3	symmetries of the auxiliary torus that fix y = 0 (and the brane): the 384 = 2^4 4! signed permutation matrices are distinct orthogonal integer matrices (they map Z^4 onto itself and preserve |n|^2); the shells |n|^2 = 1, 2, 3, 4, 9 split into orbits of sizes {8}, {24}, {32}, {8, 16}, {8, 96}	True	25.1
4	Fourier modes of the auxiliary torus: phi_n = Exp[I q_n.y]/Sqrt[V_perp] with q_n = 2 Pi n/Lc (n in Z^4) are orthonormal for the proper volume element b^4 d^4y (exact integrals: n = m = (1, 0, -2, 3); n = (1, 0, -2, 3), m = (1, 1, -2, 3); n = 0, m = (0, 2, 0, 0))	True	25.2
5	Parseval on the auxiliary torus: f(y) = Prod_k cos^2(Pi y_k/Lc) = Prod_k (1/2 + (1/4) e^(2 Pi I y_k/Lc) + (1/4) e^(-2 Pi I y_k/Lc)) has the 81 non-zero coefficients c_n = Sqrt[V_perp] Prod_k w(n_k), w(0) = 1/2, w(+-1) = 1/4 (three of them checked by integration), and Sum_n |c_n|^2 == Int b^4 d^4y f^2 == (3/8)^4 V_perp	True	25.2
6	the Laplacian of the auxiliary torus: -(1/b^2) Sum_k d^2 phi_n/dy_k^2 == (|q_n|^2/b^2) phi_n (n = (2, -1, 0, 3))	True	25.2

#	check	result	section
7	scalar Kaluza-Klein masses: for the static metric diag(1, -1, -1, -1, -b0^2, ..., -b0^2) the Klein-Gordon operator Box + m^2 (Box f = |g|^(-1/2) d_A(|g|^(1/2) g^AB d_B f)) on Exp[-I E x0 + I k.x + I q.y] gives (-E^2 + |k|^2 + |q|^2/b0^2 + m^2) times the wave: in ordinary spacetime a particle of mass Sqrt[m^2 + |q|^2/b0^2]	True	25.2
8	Jacobi's four-square theorem: the number of n in Z^4 with |n|^2 = k is r_4(k) = 8 sigma(k) - 32 sigma(k/4) for k = 1..100 (direct count over {-10, ..., 10}^4)	True	25.3
9	Weyl's law: the number N(R) of n in Z^4 with |n| <= R approaches the volume Pi^2 R^4/2 of the four-ball: N(10) = 49689, N(100) = 493490641, and |N(R)/(Pi^2 R^4/2) - 1| < 10^-2, 10^-3, 3 10^-5, 5 10^-6, 10^-6 for R = 10, 30, 100, 300, 1000	True	25.3
10	the auxiliary torus today (M8 = 10 TeV, M_Pl = 2.435 10^18 GeV): V_perp = M_Pl^2/M8^6 = 5.929 10^12 GeV^-4, l_perp = V_perp^(1/4) = b0 Lc = 1560.4 GeV^-1 = 3.079 10^-13 m, lowest Kaluza-Klein mass 2 Pi/l_perp = 4.027 MeV; N(mu) = V_perp mu^4/(32 Pi^2) = 1.17 10^13 levels below m_r/2 = 5 GeV (the lattice count of section 22.2 agrees to 10^-5) and N(M8) = M_Pl^2/(32 Pi^2 M8^2)	True	25.3
11	Newton's law in eight dimensions: Omega_6 = 2 Pi^(7/2)/Gamma(7/2) = 16 Pi^3/15, r^-5 is harmonic away from r = 0 in seven space dimensions ((1/r^6) d/dr (r^6 d/dr r^-5) = 0), and grad^2 Phi = 8 Pi G8 ((D - 3)/(D - 2)) M delta^7 (D = 8) gives Phi = -(5/6) kappa M/(5 Omega_6 r^5) = -5 kappa M/(32 Pi^3 r^5); the same formula in D = 4 gives -G M/r	True	25.4
12	images = Kaluza-Klein modes (Poisson summation; lengths in units of l_perp): Sum_n (r^2 + |n|^2)^(-5/2) == (4 Pi^2/(3 r)) Sum_m Exp[-2 Pi r |m|] to 10^-8 at r = 1/4, 1/2, 1, 2 (left: shells with r_4(k) to k = 1.5 10^6 and the continuum tail Pi^2 Int k (r^2 + k)^(-5/2) dk; right: shells to k = 20000)	True	25.4
13	near and far field: Z(5/2) = Sum_(n != 0) |n|^-5 = Sum_k r_4(k) k^(-5/2) = 7 zeta(5/2) zeta(3/2) = 24.531 (shell sum with tail, to 10^-8); near the source r^5 S(r) - 1 -> r^5 Z(5/2) (the ratio lies between 0.98 and 1 at r = 1/10 and 1/20, closer to 1 at 1/20): the eight-dimensional law; far from it 3 r S/(4 Pi^2) - 1 -> 8 Exp[-2 Pi r], the 8 lightest Kaluza-Klein modes (ratio 1 to 2 10^-2 at r = 2 and 3)	True	25.4
14	graviton + radion: far from the source the images add up to (5 kappa M/(32 Pi^3)) (4 Pi^2/(3 V_perp r)) == (5/3) G4 M/r with G4 = kappa/(8 Pi V_perp); the radion of section 20.3, coupled to matter by (phi/(Sqrt[3] M_Pl)) T, gives the potential -(1/(Sqrt[3] M_Pl))^2 M M' Exp[-m_r r]/(4 Pi r), and (1/(Sqrt[3] M_Pl))^2/(4 Pi) == (2/3) G4 (8 Pi G4 = 1/M_Pl^2): graviton 1 + massless radion 2/3 = 5/3	True	25.4
15	Kaluza-Klein momentum is conserved in the bulk and not at the brane: Int b^4 d^4y phi_n phi_m phi_l^* == 1/Sqrt[V_perp] for l = n + m and 0 otherwise (exact: n = (1, 0, 0, 2), m = (0, 1, -1, 0), l = (1, 1, -1, 2) and (1, 1, -1, 1)); at y = 0, phi_n phi_m phi_l^* == V_perp^(-3/2) for every n, m, l	True	25.5
16	the delta function of the torus: Int_(-Lc/2)^(Lc/2) Exp[-I q_n y] delta(y) dy == 1 for every integer n, so delta^4(y) = Lc^-4 Sum_n Exp[I q_n.y] (every mode with the weight 1/Lc^4); a brane source rho4 delta^4(y)/b^4 has Int b^4 d^4y (rho4 delta^4(y)/b^4) == rho4, i.e. the average rho4/V_perp over the torus	True	25.5
17	the world brane W = {x4 = x5 = x6 = x7 = 0}: the induced metric h = J^T.g.J of the embedding X = (x0, x1, x2, x3, 0, 0, 0, 0) is, for the cosmological metric of section 11.2, the Friedmann-Robertson-Walker metric diag(1, -a^2, -a^2, -a^2) (independent of b); for the inhomogeneous metric of section 11.3 it is diag(n^2, -a^2, -a^2, -a^2) at x4 = 0 with n = 1 + x0^2/11, a = 1 + x0/3	True	25.6
18	brane integrals in eight-dimensional form: at the brane Sqrt|det g| == Sqrt|det h| b^4 with b^4 = Sqrt[det(-g_kl)] (cosmological metric; inhomogeneous metric exactly at x0 = 1/2), so Int_W d^4x Sqrt|h| L == Int d^8X Sqrt|g| (delta^4(y)/b^4) L; test: Int d^4y b^4 (delta^4(y)/b^4) F(y) == F(0) = 1 for F = Exp[x4 + 2 x5] Cos[x6 - x7]	True	25.6
19	how the brane sits in the bulk: the Christoffel symbols {k mu nu} at the brane (k = 4..7, mu, nu = 0..3; = (1/(2 b^2)) d_k g_mu nu for these metrics, b times the extrinsic curvature) all vanish for the cosmological metric: the brane is totally geodesic, and a free particle moving along it stays on it. For the inhomogeneous metric at x0 = 1/2 the only non-zero one is {4 00} = n (d_4 n)/b^2 = 1458/58619: a free particle at rest on the brane is accelerated off it	True	25.6
20	explicit forms: every Gamma17[A] is a signed permutation matrix (one entry +-1 in each row and column), and 2 Sqrt[2] U has only the entries 0, +-1, +-I, eight non-zero ones in each row and in each column	True	25.7
21	brane spinor (x) transverse spinor: Gamma17[mu].Gamma17[nu] == U.(gamma_mu.gamma_nu (x) 1_4).U^dag, Gamma17[k].Gamma17[l] == U.(1_4 (x) rho_k.rho_l).U^dag and Gamma17[mu].Gamma17[k] == U.(gamma_mu.gamma5 (x) rho_k).U^dag (mu, nu = 0..3, k, l = 4..7): the Lorentz group of the brane acts on the brane spinor index, the rotations about the brane on the transverse index, and the generators that mix them are broken by the brane	True	25.7
22	the transverse index carries Spin(4) = SU(2) x SU(2): rho4567 == -gamma5 (4 x 4), it commutes with every rho_k.rho_l and anticommutes with every rho_k; for each even permutation (k, l, m, n) of (4, 5, 6, 7) the combination rho_k.rho_l + s rho_m.rho_n vanishes on the eigenspace rho4567 = s, and rho_k.rho_l - s rho_m.rho_n acts there with rank 2: the four values of a form (2, 1) + (1, 2)	True	25.7

#	check	result	section
23	chirality at the brane: PL17 == U.(P_L (x) P_+ + P_R (x) P_-).U^dag with P_L,R = (1 -+ gamma5)/2 and P_+- = (1 +- rho4567)/2: the left-handed half psi_L (chi17 = -1, 8 components) is two brane-left-handed Weyl spinors with rho4567 = +1 and two brane-right-handed ones with rho4567 = -1 (each part of rank 4)	True	25.7
24	an explicit spinor: psi = U.(u (x) e_1) with the brane spinor u = (1, 2, 0, -1): psibar psi == ubar u == 4 and psibar Gamma^mu psi == ubar gamma^mu u (mu = 0..3)	True	25.7
25	brane tension: for S_W = -Int d^4x Sqrt|h| f_W and h = diag(1/u0, -1/u1, -1/u2, -1/u3) (h^00 = u0, h^ii = -u_i), T_mu nu = (2/Sqrt|h|) dS_W/dh^mu nu == f_W h_mu nu, T^0_0 == f_W (energy density) and T^1_1 == f_W (pressure -f_W: w = -1 along the brane)	True	25.8
26	sources smeared over the torus (W(a, b) the comoving energy, section 20.2): the brane tension W = a^3 f_W/Lc^4 gives eps = f_W/V_perp, pa = -eps, pb = 0; brane radiation W = a^3 r0 a^-4/Lc^4 gives pa = eps/3, pb = 0; radiation of the bulk, isotropic in the seven space directions, has <cos^2 theta> = 3/7 and <sin^2 theta> = 4/7 for the weight cos^2 theta sin^3 theta of the momentum sphere, so pa = (1/3)(3/7) eps = eps/7 and pb = (1/4)(4/7) eps = eps/7	True	25.8
27	the brane in four-dimensional physics: (i) the tension is vacuum energy, kappa f_W/V_perp == f_W/M_Pl^2 (V_perp = kappa M_Pl^2), and f_W = M8^4 is between 3 10^62 and 5 10^62 times rho_Lambda = 3 H0^2 M_Pl^2 Omega_Lambda (section 24.5); (ii) normalcy temperature: (T^7/M8^6)/(T^2/M_Pl) == 1 at T* = (M8^6/M_Pl)^(1/5) = 13.3 GeV, and V_perp T^4 T^3/M_Pl^2 == T^7/M8^6; (iii) thin brane: (1/y^3) d/dy (y^3 d/dy y^-2) == 0 and 2 Pi^2 y^3 d/dy y^-2 == -4 Pi^2 (the Green's function -1/(4 Pi^2 |y|^2) of the four transverse directions), and Sqrt[kappa f_W] == 1/M8 for f_W = M8^4	True	25.8

The Python program's boolean checks:

#	check	result
1	torus: det(-g_kl) = b^8 for the cosmological metric; the proper volume of [0, Lc)^4 is b^4 Lc^4	True
2	only b Lc is physical: y = lambda y' maps (Lc, b) to (Lc/lambda, lambda b); b^2 dy.dy, b Lc and b^4 Lc^4 unchanged	True
3	symmetries fixing y = 0: 384 = 2^4 4! distinct signed permutation matrices, orthogonal integer matrices; shells 1, 2, 3, 4, 9 have orbits {8}, {24}, {32}, {8, 16}, {8, 96}	True
4	Fourier modes phi_n = exp(i q_n.y)/sqrt(V_perp), q_n = 2 pi n/Lc: orthonormal for b^4 d^4y (three exact pairs)	True
5	Parseval: f = prod cos^2(pi y_k/Lc) has 81 coefficients sqrt(V_perp) prod w(n_k) (3 checked by integration), Sum |c_n|^2 = Int b^4 f^2 = (3/8)^4 V_perp	True
6	Laplacian of the torus: -(1/b^2) Sum d^2/dy_k^2 phi_n = (|q_n|^2/b^2) phi_n	True
7	scalar Kaluza-Klein masses: (Box + m^2) exp(-iEx0 + ik.x + iq.y) = (-E^2 + |k|^2 + |q|^2/b0^2 + m^2) exp(...) on diag(1,-1,-1,-1,-b0^2,...)	True
8	graviton on the torus: D(D-3)/2 = 20 (D = 8) = 2 + 8 + 10 (zero mode: graviton, 4 graviphotons, 10 moduli) = 5 + 9 + 6 (massive level)	True
9	Jacobi: #{n in Z^4 : |n|^2 = k} = r_4(k) = 8 sigma(k) - 32 sigma(k/4), direct count for k = 1..100	True
10	Weyl's law: N(10) = 49689, N(100) = 493490641; |N(R)/(pi^2 R^4/2) - 1| < 1e-2, 1e-3, 3e-5, 5e-6, 1e-6	True
11	numbers today: V_perp = 5.929e12 GeV^-4, l_perp = 1560.4 GeV^-1 = 3.079e-13 m, 2 pi/l_perp = 4.027 MeV; N(5 GeV) = 1.17e13 (lattice to 1e-5); N(M8) = M_Pl^2/(32 pi^2 M8^2)	True
12	from eight to four dimensions: for diag(1, -a^2, -a^2, -a^2, -b0^2, ..., -b0^2) (static torus) R = R4 = -6(Ha' + 2 Ha^2), the scalar curvature of the FRW metric diag(1, -a^2, -a^2, -a^2); sqrt|g| = a^3 b0^4	True
13	quaternion subalgebra of the brane directions: e1 e2 = -e3, e2 e3 = -e1, e3 e1 = -e2 (e_i e_j = -delta_ij - eps_ijk e_k), span{e0, ..., e3} closed under the octonion product, Re(e_A e_B) = diag(1, -1, -1, -1)	True
14	Newton in 8D: Omega_6 = 16 pi^3/15, (1/r^6) d/dr (r^6 d/dr r^-5) = 0, Phi = -(5/6) kappa M/(5 Omega_6 r^5) = -5 kappa M/(32 pi^3 r^5); D = 4: -G M/r	True
15	4D Fourier transform: Int d^4y exp(-ip.y) (r^2 + |y|^2)^(-5/2) = (4 pi^2/(3 r)) exp(-p r) (mpmath, three (p, r), to 1e-20)	True
16	Poisson duality: Sum_n (r^2 + |n|^2)^(-5/2) = (4 pi^2/(3 r)) Sum_m exp(-2 pi r |m|) to 1e-8 at r = 1/4, 1/2, 1, 2	True
17	Z(5/2) = 7 zeta(5/2) zeta(3/2) = Sum r_4(k) k^-5/2 (to 1e-8); near field ratio -> 1 (0.98 < . < 1 at r = 1/10, 1/20); far field ratio -> 1 (to 2e-2 at r = 2, 3)	True
18	graviton + radion: (5 kappa M/(32 pi^3))(4 pi^2/(3 V_perp r)) = (5/3) G4 M/r; radion (1/(sqrt3 M_Pl))^2/(4 pi) = (2/3)/(8 pi M_Pl^2)	True
19	Kaluza-Klein momentum: bulk vertex Int b^4 phi_n phi_m phi_l^* = 1/sqrt(V_perp) for l = n + m, 0 otherwise; at y = 0 phi_n phi_m phi_l^* = V_perp^(-3/2)	True
20	delta function of the torus: Fourier coefficients Int exp(-i q_n y) delta(y) dy = 1; Int b^4 d^4y rho4 delta^4(y)/b^4 = rho4 (average rho4/V_perp)	True

#	check	result
21	induced metric h = J^T g J: FRW diag(1, -a^2, -a^2, -a^2) for the cosmological metric; diag(n^2, -a^2, -a^2, -a^2) at x4 = 0 with n = 1 + x0^2/11, a = 1 + x0/3 for the inhomogeneous metric	True
22	brane volume element: sqrt|g| = sqrt|h| b^4 at the brane (cosmological metric; inhomogeneous metric at x0 = 1/2, exact), so Int_W d^4x sqrt|h| L = Int d^8X sqrt|g| delta^4(y)/b^4 L; test Int b^4 (delta^4/b^4) F = F(0)	True
23	Christoffel symbols {k mu nu} at the brane: all 0 for the cosmological metric (totally geodesic); for the inhomogeneous metric at x0 = 1/2 only {4 00} = 1458/58619 = n d4n/b^2	True
24	R_A = (gamma_mu (x) 1_4, gamma5 (x) rho_k) and Gamma17[A] both satisfy X_A X_B + X_B X_A = 2 eta_AB 1_16	True
25	every Gamma17[A] is a signed permutation matrix	True
26	intertwiner: Schur average with the seed matrix unit (1, 2); 2 sqrt(2) U has entries 0, +-1, +-i (rounding error < 1e-12) with 8 non-zero entries in each row and column; Gamma17[A].U8 = U8.R_A and U8^dag.U8 = 8 1_16 exactly	True
27	Schur's lemma: the linear map X -> (Gamma_A X - X R_A)_A has a one-dimensional null space (one singular value < 1e-10, the next > 0.1)	True
28	brane Lorentz group on the brane spinor index, transverse rotations on a: Gamma_mu Gamma_nu = U (gamma_mu gamma_nu (x) 1) U^dag, Gamma_k Gamma_l = U (1 (x) rho_k rho_l) U^dag, Gamma_mu Gamma_k = U (gamma_mu gamma5 (x) rho_k) U^dag	True
29	Spin(4) on the transverse index: rho4567 = -gamma5 (4x4) commutes with rho_k rho_l, anticommutes with rho_k; (rho_k rho_l + s rho_m rho_n) P_s = 0 and (rho_k rho_l - s rho_m rho_n) P_s has rank 2 for even (k,l,m,n): 4 = (2,1) + (1,2)	True
30	chirality: chi = i Gamma_0...Gamma_7 = U (gamma5 (x) rho4567) U^dag; P_L17 = U (P_L (x) P_+ + P_R (x) P_-) U^dag, each part of rank 4	True
31	example psi = U.(u (x) e_1), u = (1, 2, 0, -1): psibar psi = ubar u = 4 and psibar Gamma^mu psi = ubar gamma^mu u	True
32	four brane Dirac spinors: psibar psi' = Sum_a psibar_a psi'_a and psibar Gamma^mu psi' = Sum_a psibar_a gamma^mu psi'_a (random integer spinors)	True
33	Kaluza-Klein operator X = gamma5 (x) (rho.q)/b0: X^2 = -|q|^2/b0^2 1_16 and X anticommutes with gamma_mu (x) 1_4	True
34	brane tension: T_mu nu = (2/sqrt|h|) dS/dh^mu nu = f_W h_mu nu for S = -Int sqrt|h| f_W; T^0_0 = f_W, T^1_1 = f_W (p = -f_W, w = -1)	True
35	smeared sources: tension eps = f_W/V_perp, pa = -eps, pb = 0; brane radiation pa = eps/3, pb = 0; bulk radiation <cos^2> = 3/7, <sin^2> = 4/7: pa = pb = eps/7	True
36	normalcy temperature: (T^7/M8^6)/(T^2/M_Pl) = 1 at T* = (M8^6/M_Pl)^(1/5) = 13.3 GeV; V_perp T^4 T^3/M_Pl^2 = T^7/M8^6	True
37	the tension is vacuum energy: kappa f_W/V_perp = f_W/M_Pl^2 (V_perp = kappa M_Pl^2); f_W = M8^4 is 3e62 to 5e62 rho_Lambda	True
38	Green's function of the four transverse directions: (1/y^3) d/dy (y^3 d/dy y^-2) = 0, flux 2 pi^2 y^3 d/dy y^-2 = -4 pi^2; sqrt(kappa f_W) = 1/M8 for f_W = M8^4	True

The numbers printed by both programs (largest relative difference per group; integers must agree exactly):

quantity	values	max. relative difference	tolerance
orbit sizes of the shells $k = 1, \dots, 12$	16	0	exact
$r_4(k)$, $k = 0, \dots, 16$	17	0	exact
Weyl: $N(R)$, $R = 10, 30, 100, 300, 1000$	5	0	exact
Weyl: $N(R)/(\pi^2 R^4/2) - 1$	5	3.8×10^{-10}	1×10^{-9}
$V_\perp$, $\ell_\perp$ (GeV^{-1}, m, fm), $2\pi/\ell_\perp$	5	2.2×10^{-16}	1×10^{-12}
$\ell_\perp/r_p$, $a_0/\ell_\perp$, $R = \ell_\perp/2\pi$, M_D	5	1.7×10^{-16}	1×10^{-12}
$N(\mu)$ at 5 GeV, m_h, M_8; lattice count below 5 GeV	4	1.8×10^{-16}	1×10^{-12}
the first levels (k, r_4, $4r_4$, mass)	32	2.5×10^{-16}	1×10^{-12}
image sums $S(r)$ and Kaluza–Klein sums, $r = \frac{1}{4}, \frac{1}{2}, 1, 2$	8	1.2×10^{-15}	1×10^{-9}
$Z(5/2)$: closed form, shell sum	2	2.9×10^{-16}	1×10^{-9}
near-field and far-field ratios	5	7.8×10^{-12}	1×10^{-6}
Christoffel symbol $\{4\,00\}$ at the brane	1	0	1×10^{-15}
Γ_A as signed permutations	128	0	exact
$\mathrm{Re}(2\sqrt{2}\,U)$	256	0	exact
$\mathrm{Im}(2\sqrt{2}\,U)$	256	0	exact
example spinor: $\bar\psi\psi$, $\bar\psi\Gamma^\mu\psi$	5	0	1×10^{-15}
T_*, f_W/ρ_Λ, $1/(2\pi M_8)$	3	5.8×10^{-16}	1×10^{-12}

(Checks 1–14 concern the torus alone, sections 25.1–25.4; checks 15–16 the torus and the brane together,

section 25.5; checks 17–27 the world brane, sections 25.6–25.8.) The results quoted from the other sections of 7+1.nb rest on these checks of the notebook, all evaluated in the same run:

#	check	result	section
1	scalar curvature R = -2 (3Ha' + 6Ha^2 + 4Hb' + 10Hb^2 + 12 Ha Hb)	True	11.2
2	Einstein Kasner: (p, q) = ((1+2Sqrt[2])/7, (2-3Sqrt[2])/14) and ((1-2Sqrt[2])/7, (2+3Sqrt[2])/14) solve E^(1) == 0	True	14.1
3	Gauss-Bonnet Kasner: 3 real roots of the cubic; (p, (3-3p)/4) solve E^(2) == 0	True	14.1
4	static auxiliary space (Hb = 0): E^(1) = (3Ha^2, 2Ha' + 3Ha^2, 3Ha' + 6Ha^2), E^(2) = (0, 0, -12 Ha^2 (Ha' + Ha^2)), E^(3) = 0	True	14.4
5	ADD relation: kappa8 rho4/V_perp == 8 Pi G4 rho4 with G4 = G8/V_perp (kappa8 = 8 Pi G8)	True	17.5
6	KK operator: X.X == -|q|^2/b0^2 ID16 and X anticommutes with gamma_mu (x) 1_4	True	18.1
7	KK masses: Det[gamma.p (x) 1 + X - m] vanishes on p0^2 - |p|^2 == m^2 + |q|^2/b0^2 (40-digit arithmetic) and not off shell	True	18.1
8	stabilizing potential: W = a^3 b^4 U(b) gives eps = U, pa = -U, pb = -U - (b/4) U'(b)	True	20.2
9	brane tension smeared over the torus (V_perp = b^4 Lc^4): eps = f_W/V_perp, pa = -f_W/V_perp, pb = 0	True	20.2
10	brane fluid rho_4 = N a^(-3(1+w)), smeared: eps = rho_4/V_perp, pa = w eps, pb = 0	True	20.2
11	radion equation of Einstein gravity: Hb' + 3 Ha Hb + 4 Hb^2 == kappa (eps/6 - pa/2 + pb/3) + Lambda/3 (constraint used)	True	20.3
12	radion mass: U = U0 + (K_U/2) beta^2, b U' = K_U beta gives the restoring term -(kappa K_U/12) beta, m_r^2 = kappa K_U/12	True	20.3
13	static auxiliary space: (kappa/4) b U' == Lambda + kappa U + 2 kappa f_W/V_perp + (kappa/2)(eps - 3pa) + kappa pb - 12 alpha2 Ha^2 (Ha' + Ha^2)	True	20.3
14	brane Friedmann equation: 3 Ha^2 == (kappa/V_perp) rho_4 + Lambda4 with Lambda4 = Lambda + kappa U0 + kappa f_W/V_perp, i.e. 8 Pi G4 = kappa/V_perp	True	20.3
15	Einstein-frame form of the constraint: 3 (Ha + 2 Hb)^2 == 6 Hb^2 + (E^(1)0_0 of section 12.6), i.e. 3 M_Pl^2 H_E^2 == phi'^2/2 + rho with phi = Sqrt[12] M_Pl beta	True	20.3
16	Kasner constants: P_E(u1) == 0, p1 == (1 + 2 Sqrt[2])/7, q1 == (2 - 3 Sqrt[2])/14; u2 in (-1/3, 0) with P_G(u2) == 0, 3 p2 + 4 q2 == 3	True	21.1
17	cubic regime: Ha = 1/tau + A0 + A1 tau, Hb = h0 + h1 tau + h2 tau^2 solves the 00, 11, 44 equations through order tau^-1 (11: checked, not used)	True	21.2
18	matter-dominated cubic regime: p_eff -> 3/4, Hb -> -Sqrt[alpha2/(54 alpha3)], kappa eps tau^3 -> 9 alpha2 |h*|, alpha2 E2 : alpha3 E3 : kappa eps -> 9/4 : -5/4 : 1 (Ha = 10^15)	True	21.3
19	without the stabilizing potential the gas-dominated Einstein regime decompactifies: tau Ha -> 0, tau Hb -> 2/5, pb/eps -> 1/4 (Ha = 10^-14)	True	21.3
20	Einstein + fluid power laws a ~ tau^P, b ~ tau^Q: (wa, wb) = (0, 1/4) -> (0, 2/5); (1/3, 0) -> (1/2, 0); (1/7, 1/7) -> (1/4, 1/4) (tau^2 E^(1) from EKas of section 14.1)	True	21.3
21	capture: the radion oscillates like a four-dimensional massive field in a matter-dominated universe: with tau_E from <H_E> = 2/(3 (tau - tau_E)) over the last period, beta = A sin(m_r tau + vartheta)/(m_r (tau - tau_E)) to 2 10^-2, <H_E>(tau - tau_E) = 2/3 to 10^-2 at tau = 10^5 and 2 10^5, the gas share constant to 2 10^-2	True	22.1
22	the number of lattice vectors with |n|^2 = k is r_4(k) = 8 sigma(k) - 32 sigma(k/4) (direct count for k <= 60)	True	22.2
23	the radion decays into the bulk: the lattice sum over the Kaluza-Klein modes of dirac16complex (m << m_r/2) equals its continuum limit to 10^-6, the number of momenta to 10^-5, and Gamma_bulk/Gamma_brane > 10^12	True	22.2
24	radion equilibrium in de Sitter: 3 m_r^2 beta_eq == 3 H^2 - 12 alpha2 H^4 + O(beta_eq^2)	True	23.2
25	eight-dimensional de Sitter (no stabilizer): all three equations read 21 H^2 - 420 alpha2 H^4 + 2520 alpha3 H^6 == Lambda	True	23.2
26	Kaluza-Klein tower: Sum_n (1 - mu_n^2/m_h^2)^2 -> V_perp m_h^4/(192 Pi^2) and Sum_n mu_n (1 - mu_n^2/m_h^2)^2 -> V_perp m_h^5/(315 Pi^2) (d^4n = 2 Pi^2 |n|^3 d|n|: exact integrals); the lattice sums with r_4(k) for R = 100 equal Pi^2 R^4/12 and 16 Pi^2 R^4/315 to 10^-8	True	24.4
27	Epstein sum over Z^4 without 0: Sum |n|^-8 == 8 (1 - 4^-3) zeta(4) zeta(3), compared with the shell sum Sum r_4(k)/k^4 to k = 20000 (tail < 3 10^-8)	True	24.5
28	Casimir energy of dirac16complex on the auxiliary torus: massless 16 Gamma(4) Z/(Pi^4 l^4) > 10^34 rho_Lambda, the massive sum -> the massless value for m l -> 0 (to 10^-4), and equals rho_Lambda at m l = 92.2	True	24.5

Section 10 rests on these checks of section 26 of 7+1.nb (which contains 33 checks, 33 True; the independent program tools/ref_radion.py has 31 boolean checks, 31 True, and the 15 groups of numbers printed by both agree to a largest relative difference of 8.2×10^{-11}):

#	check	result	section
1	the 4-sphere of radius r0: Ricci_ij = 3 gamma_ij (gamma the unit-sphere metric), Ricci_ij = 0 for i != j, so R(S^4) = 12/r0^2 (in the signature of 7+1.nb the space-like block gives R = -12/r0^2, also with curvature[] of section 11); the volume of the unit S^4 is 8 Pi^2/3	True	26.1
2	radion on T^4: for ds^2 = e^(-4 beta) (N^2 dx0^2 - a^2 dx.dx) - e^(2 beta) b0^2 dy.dy the reduced action -(1/(2 kappa)) Int(R + 2 Lambda) - Int U equals -(M_Pl^2/2) R_E + 6 M_Pl^2 (d beta)^2 - e^(-4 beta)(M_Pl^2 Lambda + V0 U) up to a total derivative (Euler-Lagrange expressions for N, a, beta vanish), M_Pl^2 = V0/kappa, V0 = b0^4 Lc^4	True	26.1
3	radion on S^4: the same reduction for ds^2 = e^(-4 beta) g_E - e^(2 beta) r0^2 gamma (unit-sphere gamma) holds with the additional potential -(M_Pl^2/2) e^(-4 beta) 12/(r0^2 e^(2 beta)) = -6 M_Pl^2 e^(-6 beta)/r0^2 from the curvature of the sphere, V0 = (8 Pi^2/3) r0^4; R is independent of the angles	True	26.1
4	radion on T^4 with U = U0 + U1 beta + (K_U/2) beta^2: a flat vacuum at b = b0 needs Lambda + kappa U0 = 0 and U1 = 0, and m_r^2 = V''(beta)/(12 M_Pl^2) = kappa K_U/12 (section 20.3)	True	26.2
5	radion on S^4 with U: Lambda + kappa U0 == 6/r0^2, kappa U1 == -12/r0^2 and m_r^2 == kappa K_U/12 - 2/r0^2 from V(beta); the eight-dimensional radion equation gives the same static solution and the same mass	True	26.2
6	S^4 with Lambda only: dS4 x S^4 with r0^2 = 9/Lambda, H^2 = 1/r0^2 = Lambda/9 solves the 00, 11 and 44 equations; it is a MAXIMUM of V: m^2 = -6 H^2 from V(beta), and m^2 = -6/r0^2 = -6 H^2 from the eight-dimensional radion equation	True	26.2
7	zonal harmonics on S^4 (l = 0..6, exact): (1/sin^3 x4) d/dx4 (sin^3 x4 d/dx4 C_l^(3/2)(cos x4)) == -l (l + 3) C_l^(3/2), C_l^(3/2)(1) == (l + 1)(l + 2)/2, and Int over S^4 of (C_l/C_l(1))^2 == V/d_l (unit sphere), so the normalized zonal harmonic is Sqrt[d_l/V] at the north pole; d_l == Binomial[l + 4, 4] - Binomial[l + 2, 4] for l = 0..20	True	26.4
8	Weyl's law on S^4: the number of levels with l(l + 3) <= (mu r)^2 approaches (mu r)^4/12 = V mu^4/(32 Pi^2) (V = 8 Pi^2 r^4/3): relative deviation -1.0 10^-2, -1.0 10^-4, -1.0 10^-6 at mu r = 10, 100, 1000 (like -1/(mu r)^2)	True	26.4
9	the brane at the north pole of S^4: the embedding (r0 sin x4 sin x5 sin x6 cos x7, ..., r0 cos x4) of S^4 in R^5 induces r0^2 gamma; the north pole x4 = 0 is (0, 0, 0, 0, r0); of the 10 rotations L_ab of R^5 (the Killing vectors of S^4, SO(5)) the 6 with a, b <= 4 vanish there (SO(4), kept by the brane) and the 4 L_a5 move it	True	26.4
10	Killing spinors of S^4 (radius r0, frame e^a = (r0, r0 sin x4, ...) dx^(a+3), gt_a = I rho_(a+3)): eps_s = exp(s (i/2) x4 gt_1) exp((1/2) x5 gt_1 gt_2) exp((1/2) x6 gt_2 gt_3) exp((1/2) x7 gt_3 gt_4) eps0 satisfies nabla_mu eps_s == s (i/(2 r0)) e_mu^a gt_a eps_s (s = +-1, all eps0), so the Dirac operator gives (2 s i/r0) eps_s: |lambda| = 2/r0, Friedrich's bound n R/(4(n - 1)) = 4/r0^2	True	26.4
11	Dirac spectrum on S^4: Weyl's law for spinors (4 components) N(|lambda| <= L/r) ~ 4 V (L/r)^4/(32 Pi^2) = L^4/3 agrees with the degeneracies 2 x 4 Binomial[l + 3, 3] of the eigenvalues +-i(l + 2)/r (to 2.0 10^-3 at L = 1000)	True	26.4
12	the sphere with the volume of the torus today (V = M_Pl^2/M8^6): r = (3 V/(8 Pi^2))^(1/4) = 688.9 GeV^-1 = 1.359 10^-13 m; lowest scalar levels 4.03 MeV (torus, 8 states) and 2.90 MeV (sphere, l = 1, 5 states); lowest dirac16complex level on the sphere 2/r = 2.90 MeV (8 Dirac fermions; the torus has 4 massless ones for m = 0); levels below 5 GeV: torus 1.17335 10^13, sphere 1.17375 10^13, Weyl 1.17335 10^13	True	26.4

13. How to reproduce

1. Write the unevaluated notebook with `python3 tools/make_7p1_nb.py`; sections 25 and 26 are generated by `tools/nb_part_torus_brane.py` and `tools/nb_part_radion.py`.
2. Evaluate 7+1.nb in a fresh kernel of Wolfram 15.0.1 (Evaluation ▷ Quit Kernel, then Evaluation ▷ Evaluate Notebook; or `Get["D:/Developer/github/7+1/tools/run_eval.wl"]`) and save it.
3. Run `python3 tools/ref_torus_brane.py` and `python3 tools/ref_radion.py`: they repeat the examples independently (exit code 0 if all pass).
4. Run `python3 tools/make_docs.py`: it regenerates this document, its symbol dictionary and its provenance record (and all other documents).

14. References

- Th. Kaluza, Sitzungsber. Preuss. Akad. Wiss. (1921) 966; O. Klein, Z. Phys. 37 (1926) 895 — a compact extra dimension.
- N. Arkani-Hamed, S. Dimopoulos, G. Dvali, Phys. Lett. B 429 (1998) 263 and Phys. Rev. D 59 (1999) 086004 — large extra dimensions, $M_{\rm Pl}^2 = V_\perp M_D^{n+2}$.
- A. Kehagias, K. Sfetsos, Phys. Lett. B 472 (2000) 39 — the Yukawa corrections to Newton's law from a torus.
- C. G. J. Jacobi, Fundamenta nova theoriae functionum ellipticarum (1829) — the four-square theorem; H. Weyl, Math. Ann. 71 (1912) 441 — the asymptotic distribution of eigenvalues.
- H. B. G. Casimir, Proc. K. Ned. Akad. Wet. 51 (1948) 793 — the Casimir energy; P. Epstein, Math. Ann. 56 (1903) 615 — lattice zeta functions.
- S. Navas et al. (Particle Data Group), Phys. Rev. D 110 (2024) 030001 — searches for large extra dimensions.
- R. Camporesi, A. Higuchi, J. Geom. Phys. 20 (1996) 1 — the Dirac spectrum on spheres; Th. Friedrich, Math. Nachr. 97 (1980) 117 — its lower bound.
- E. Tiesinga, P. J. Mohr, D. B. Newell, B. N. Taylor, Rev. Mod. Phys. 93 (2021) 025010 (CODATA 2018) — $\hbar c$, a_0, r_p.