The generalized Kronecker delta and the Lovelock tensors of the author's primordial metric

GKD, the pure-Rust generalized Kronecker delta, and the exact Lovelock tensors of order k = 1, 2, 3 of Lovelock's equation (4.38) for the 4+4 metric with deflating extra times, with their independent verifications and an exact statement of what they establish

dirac16complex project (with Claude Opus 5.5)

September 2026

Contents

1 The task and the result **3**

2 Setting and conventions **4**

2.1 The author's metric and coordinates 4

2.2 Curvature conventions 4

3 The generalized Kronecker delta **5**

3.1 The author's definition 5

3.2 GKD and the proof that it equals the definition 5

3.3 The self-test of GKD 5

4 The Lovelock tensors **6**

4.1 Lovelock's equation (4.38) 6

4.2 Definitions and index conventions of this record 6

4.3 Normalisations 7

4.4 Why the series stops at k = 3 8

5 How the code computes them **8**

5.1 The crate 8

5.2 The enumeration 9

6 Results **10**

6.1 Christoffel symbols and curvature 10

6.2 The Lovelock scalars 11

6.3 The Lovelock tensors 11

6.4 The checks of the Rust program 16

7 Independent verifications **16**

7.1 The Wolfram check (29 checks) 16

7.2 The sympy checker (49 checks) 18

8 The comparison with the author's notebooks **19**

8.1 What was read for the computation 19

8.2 The comparison with the author's stored outputs (2026-10-08) 20

8.3 What the comparison does not establish 23

8.4 The metric with $b_4(x_4)$ 23

9 How the field equations for a_4 and b_4 use the Lovelock tensors **24**

10 Verification records **25**

11 Reproduction **26**

11.1 Commands 26

11.2 Expected output 27

11.3 Run times 28

11.4 The PDF 28

12 What is established and what is not **28**

12.1 Established 28

12.2 Not established 29

Abstract

This document records the computation of `Revision/gkd_lovelock`: the generalized Kronecker delta of the author's notebook, re-implemented in pure Rust as the function GKD, and the three nonzero Lovelock tensors $P_{(1)}$, $P_{(2)}$, $P_{(3)}$ of Lovelock's equation (4.38) for the author's primordial metric of signature (4,4), in which the three extra times x_5, x_6, x_7 deflate as $e^{-a_4(x_4)} b_4(x_4)^{1/2}$ while 3-space inflates as $e^{a_4(x_4)}$ (the metric function $b_4(x_4)$ of the extra-time components is never 1). Every component of $P_{(k)}$ is exact: a polynomial in H, a_4', a_4'', $u = b_4'/b_4$ and $v = u'$ (and, in the x_0-x_4 pair, $\cot z$); the diagonal components are free of the warp factor $\sin^{1/3} z$, of e^{a_4} and of x_0, with equal components in the three 3-space directions and in the three extra-time directions, and the only off-diagonal components are the pair x_0-x_4, x_4-x_0, proportional to b_4' (they would vanish only for a constant b_4); $P_{(4)} = 0$ identically in eight dimensions. The Rust program checks 19 identities (19 of 19 pass); GKD equals the author's determinant on all 16,777,216 pairs of index lists of length 4 and on 200,000 pseudo-random pairs of each length 5 to 9. Two independent verifications agree with every component: a Wolfram Language check that evaluates the author's definition verbatim (29 of 29 checks pass) and a sympy checker that shares no code with the Rust program (49 of 49 checks pass). The field equations for a_4 and b_4 use these tensors with the normalisation $E_{(k)} = -P_{(k)}/2^{k+1}$, $E_{(1)} = G$; with the author's Lovelock weights they give, on the history $a_4 = Hx_4$, $b_4 = b_4(0)e^{(4/3)Hx_4}$, exactly the constant source that this history needs. A later comparison with the stored outputs of the author's earlier notebook, which computes the withdrawn metric $b_4 = 1$ (`Revision/gkd_lovelock/comparison/`, 2026-10-08; 78 checks: 73 PASS, 0 FAIL, 5 NOT-AVAILABLE), finds the author's metric, Ricci scalar and all 64 components of the author's Einstein tensor exactly equal to this record under a function mapping stated before the comparison (the coordinates are the author's on both sides), so that $P_{(1)}$ and $L_{(1)}$ equal -4 times the author's Einstein tensor and 2 times the author's Ricci scalar; in the two notebooks of the author in the repository a keyword scan finds no stored output of the Lovelock tensors of order $k = 2$ and $k = 3$, so for those orders nothing could be compared. Not established here: any solution of the field equations, anything at or beyond the brane $z = \pi/2$, and agreement with any Lovelock tensor of order $k = 2$ or $k = 3$ computed by the author.

1 The task and the result

The author's test (2026-10-01, `Revision/README.md` and `Revision/SPEC.md` sections 5 and 10): refactor the author's Wolfram Language definition of the generalized Kronecker delta into a pure-Rust function GKD, use it to compute, with provenance, the three nonzero Lovelock tensors of Lovelock's equation (4.38) for the author's 8 x 8 metric, write every component, and do the work without reading the author's answers. The result, in one paragraph each:

1. GKD (section 3). For index lists of equal length, GKD returns the determinant of the author's 0/1 matrix without computing a determinant: it is the sign of the permutation that maps the lower list onto the upper list, or 0. The equality with the author's definition is a theorem (section 3.2), confirmed exhaustively and on random samples by three programs.

2. The Lovelock tensors (sections 4 to 6). ${P_{(k)}}^h{}_j$, ${A_{(k)}}^{lh}$ and $L_{(k)}$ for $k = 1, 2, 3$ are computed exactly from the metric alone, with every term of the sums weighted by an explicit call of GKD. Their nonzero components are listed in section 6.3.

3. The verifications (sections 7 and 10). The Rust program, the Wolfram check and the sympy checker agree exactly on every one of the 64 components of each tensor; every check of the three reports has the verdict PASS.

4. The use (section 9). The field equations for a_4 and b_4 of `Revision/field_equations_a4` recompute the three tensors in both of their verifiers and compare them with the file of this record.

5. The comparison with the author (sections 8.2 and 8.4). Made after the computation: under the mapping stated before the comparison, the stored metric, Ricci scalar and Einstein tensor of the author's earlier notebook (the withdrawn metric $b_4 = 1$) equal those of this record at $b_4 = 1$ exactly, component by component; for $k = 2$ and $k = 3$ that notebook holds no output to compare with. The author's new notebook, with b_4, is compared in `Revision/metric_b4`.

Every formula and number below is taken from the Revision files named with it; nothing is taken from the earlier stages of the repository. Interpretations are labelled as such.

2 Setting and conventions

2.1 The author's metric and coordinates

The coordinates are the author's, $x_0, x_1, \ldots, x_7$, in the author's order; arrays index them $0, \ldots, 7$ (index k is x_k). x_0 is the hidden space direction with $z = 6Hx_0 \in (0, \pi/2)$; x_1, x_2, x_3 are ordinary 3-space, which inflates with the scale factor $e^{a_4} \sin^{1/6} z$; x_4 is the time; x_5, x_6, x_7 are the three extra times, time-like, which DEFLATE exponentially with the scale factor $e^{-a_4} b_4^{1/2} \sin^{1/6} z$ as $a_4(x_4)$ increases (they are never treated as static); $H > 0$ is the author's constant and $a_4' = da_4/dx_4$. (Until 2026-10-14 this record named the coordinates $x_1, \ldots, x_8$, with the hidden direction x_8 listed last; the user ordered the author's coordinates on 2026-10-14.) The metric of `Revision/SPEC.md` section 1, as the Rust program stores it (`metricDiagonal` of `curvature.json`), is

$$\begin{aligned} g = \operatorname{diag}\big(&\cot^2 z,\ e^{2a_4} \sin^{1/3} z,\ e^{2a_4} \sin^{1/3} z,\ e^{2a_4} \sin^{1/3} z,\ -1, \\ &-e^{-2a_4} b_4 \sin^{1/3} z,\ -e^{-2a_4} b_4 \sin^{1/3} z,\ -e^{-2a_4} b_4 \sin^{1/3} z \big), \end{aligned}$$

with the metric function $b_4(x_4) > 0$ of the user's metric of 2026-10-14 multiplying the three extra-time components (b_4 is a function of the time and never 1; $b_4 = 1$ is the withdrawn metric used before 2026-10-14, section 8.4), of signature (4,4): space-like x_0, x_1, x_2, x_3, time-like x_4, x_5, x_6, x_7. Its determinant is $\det g = b_4^3 \cos^2 z$, so $\sqrt{|g|} = b_4^{3/2} \cos z$ on $0 < z < \pi/2$; the Rust program writes it as $\sin z \cot z\, b_4^{3/2}$ (`sqrtAbsDetG`). In the 7-volume the factors e^{3a_4} of the inflating 3-space and e^{-3a_4} of the extra times cancel, and $b_4^{3/2}$ remains. Below, $u = b_4'/b_4$ and $v = u'$ (so $b_4''/b_4 = v + u^2$); the log-derivative of the extra-time scale factor is $-(a_4' - u/2)$. The Revision's Kohn-Sham record uses the history $a_4 = Hx_4$, $b_4 = b_4(0) e^{(4/3)Hx_4}$, on which $a_4' = H$, $u = \frac{4}{3}H$ and $a_4' - u/2 = H/3$ (section 9).

2.2 Curvature conventions

Stated in `Revision/gkd_lovelock/code/src/geometry.rs` and used by all three programs:

- Christoffel symbols $\Gamma^a{}_{bc} = \frac{1}{2} g^{ad} (\partial_b g_{dc} + \partial_c g_{db} - \partial_d g_{bc})$.

- Riemann tensor (Misner-Thorne-Wheeler) $R^a{}_{bcd} = \partial_c \Gamma^a{}_{bd} - \partial_d \Gamma^a{}_{bc} + \Gamma^a{}_{ce}\Gamma^e{}_{bd} - \Gamma^a{}_{de}\Gamma^e{}_{bc}$; a round 2-sphere of radius r has $R^{12}{}_{12} = +1/r^2$.
- $R^{ab}{}_{cd} = g^{be} R^a{}_{ecd}$; Ricci $R_{bd} = R^a{}_{bad}$, so $R^h{}_j = R^{ha}{}_{ja}$; $R = R^{ab}{}_{ab}$; Einstein $G^h{}_j = R^h{}_j - \frac{1}{2}\delta^h_j R$.

3 The generalized Kronecker delta

3.1 The author's definition

The cell stored as `In[87]` in the author's notebook `Generalized _Kronecker_Delta_4+4.nb` (the author's `In[54]`) defines

```
kδ[lower_, upper_] /; Length[lower] == Length[upper] := Det[Outer[delta, lower, upper]]
```

For lists $l = (l_1, \ldots, l_p)$ and $u = (u_1, \ldots, u_p)$ the matrix `Outer[delta, lower, upper]` has the entries $M_{ij} = \delta(l_i, u_j)$ (1 if $l_i = u_j$, otherwise 0), and

$$\delta^{u_1 \cdots u_p}_{l_1 \cdots l_p} = \det\left[\delta(l_i, u_j)\right]_{i,j=1}^{p}.$$

For lists of different length the definition does not apply and the expression stays unevaluated. Because $\det M = \det M^T$, exchanging the two lists does not change the value.

3.2 GKD and the proof that it equals the definition

Theorem (GKD). For lists of equal length p, $\det M = 0$ if two lower indices are equal, if two upper indices are equal, or if some lower index is not among the upper indices; otherwise $\det M = \mathrm{sign}(\sigma)$, where σ is the permutation with $l_i = u_{\sigma(i)}$.

Proof (the comment of `Revision/gkd_lovelock/code/src/gkd.rs`). (a) Two equal lower indices give two equal rows of M, so $\det M = 0$. (b) Two equal upper indices give two equal columns, so $\det M = 0$. (c) A lower index l_i that is not among the upper indices gives a zero row i, so $\det M = 0$. (d) Otherwise both lists consist of p distinct indices and every l_i equals exactly one $u_{\sigma(i)}$; M has a single 1 in each row i, in column $\sigma(i)$, so it is the permutation matrix of σ, and in the Leibniz formula only the term of σ survives: $\det M = \mathrm{sign}(\sigma)$. QED.

`gkd.rs::GKD` implements cases (a) to (d) with a position table and counts the inversions of σ: $O(p^2)$ integer operations ($p \le 64$) instead of a $p \times p$ determinant. With more than eight indices over eight values two indices of a list are always equal (pigeonhole), so GKD is 0. `gkd.rs::kdelta_det` is a literal transcription of the definition (the Leibniz determinant of the 0/1 matrix, permutations by Heap's algorithm) used only to test GKD.

3.3 The self-test of GKD

`lovelock_gkd gkd-selftest -exhaustive-max 4` compares GKD with `gkd.rs::kdelta_det` on every pair of index lists of length 1 to 4 over the eight labels and on 200,000 pseudo-random pairs

(xorshift64*, half of them permutations of each other) of each length 5 to 9. The report `Revision/gkd_lovelock/results/gkd-selftest.json` has the verdict `SUCCESS`:

length p	**mode**	**pairs**	**nonzero values**	**mismatches**
1	exhaustive	64		0
2	exhaustive	4096		0
3	exhaustive	262144		0
4	exhaustive	16777216		0
5	random	200000	20538	0
6	random	200000	7812	0
7	random	200000	1949	0
8	random	200000	244	0
9	random	200000	0	0

At length 9 (nine indices over eight values) no pair has a nonzero value, as the pigeonhole argument requires. The Lovelock program records two values of GKD itself (`gkdSelfCheck` of `lovelock-report.json`): $+1$ for a 3-cycle (`length3Pair`) and -1 for a transposition (`transposition`).

4 The Lovelock tensors

4.1 Lovelock's equation (4.38)

The cell stored as `In[101]` in the author's notebook (the author's `In[68]`) is a picture of Lovelock's equation (4.38) (Lovelock and Rund), rendered to `Revision/gkd_lovelock/results/notebook-in68-image.png` and read from that picture. In n dimensions, with arbitrary constants $\alpha_{(k)}$ and λ and $m = n/2$ for even n,

$$A^{lh} = \sqrt{g} \sum_{k=1}^{m-1} \alpha_{(k)}\, g^{jl}\, \delta^{hh_1\cdots h_{2k}}_{jj_1\cdots j_{2k}}\, R^{j_1 j_2}{}_{h_1 h_2} \cdots R^{j_{2k-1} j_{2k}}{}_{h_{2k-1} h_{2k}} + \lambda \sqrt{g}\, g^{lh}.$$

For the author's metric $n = 8$ and $m = 4$, so the sum runs over $k = 1, 2, 3$; $\sqrt{g}$ means $\sqrt{|\det g|} = \cos z$ (section 2.1).

4.2 Definitions and index conventions of this record

For $k = 1, 2, 3$ (`lovelock-tensors.json`, key `definition`):

$$\begin{aligned}
P_{(k)}{}^{h}{}_{j} &= \delta^{hh_1\cdots h_{2k}}_{jj_1\cdots j_{2k}}\, R^{j_1 j_2}{}_{h_1 h_2} \cdots R^{j_{2k-1} j_{2k}}{}_{h_{2k-1} h_{2k}} \\
&= \sum \mathrm{GKD}([j, j_1, \ldots, j_{2k}], [h, h_1, \ldots, h_{2k}]) \prod_{i=1}^{k} R^{j_{2i-1} j_{2i}}{}_{h_{2i-1} h_{2i}}, \\
L_{(k)} &= \delta^{h_1\cdots h_{2k}}_{j_1\cdots j_{2k}}\, R^{j_1 j_2}{}_{h_1 h_2} \cdots R^{j_{2k-1} j_{2k}}{}_{h_{2k-1} h_{2k}}, \\
A_{(k)}{}^{lh} &= \sqrt{|g|}\, g^{jl}\, P_{(k)}{}^{h}{}_{j},
\end{aligned}$$

summed over all repeated indices. Conventions, stated once:

- The lower list of GKD is $[j, j_1, \ldots, j_{2k}]$ (the lower indices of δ, which carry the UPPER index pairs of the curvature factors) and the upper list is $[h, h_1, \ldots, h_{2k}]$; in the code (`lovelock.rs::dfs`) the upper pair a, b of a factor $R^{ab}{}_{cd}$ joins the lower row of δ and the lower pair c, d joins the upper row. The field-equations branch writes the lists in the other order; by section 3.1 the value is the same.
- $P_{(k)}$ is stored with its first index up and its second down (`P1_mixed_up_h_down_j`, `P2_mixed_up_h_down_j`, `P3_mixed_up_h_down_j`); $A_{(k)}$ with both indices up (`A1_contravariant_l_h`, `A2_contravariant_l_h`, `A3_contravariant_l_h`). For the diagonal metric $A_{(k)}{}^{lh} = \cos z \; g^{ll} \, P_{(k)}{}^{h}{}_{l}$.
- In this notation equation (4.38) reads $A^{lh} = \sum_{k=1}^{3} \alpha_{(k)} A_{(k)}{}^{lh} + \lambda \sqrt{|g|}\, g^{lh}$.

4.3 Normalisations

$P_{(k)}$ carries the normalisation of the δ sum. The usual normalised Lovelock tensors are (`lovelock-tensors.json`, key `normalisationNote`; `lovelock.rs::usual_normalisation`)

$$E_{(k)}{}^{h}{}_{j} = -\frac{P_{(k)}{}^{h}{}_{j}}{2^{k+1}}, \qquad E_{(1)} = G,$$

with the following exact relations, derived in the docstring of `Revision/gkd_lovelock/verification/check_lovelock_gkd.py` by the Laplace expansion of the determinant, $P_{(k)}{}^{h}{}_{j} = \delta^{h}_{j} L_{(k)} - 2k \, \delta^{h h_1 \cdots h_{2k}}_{j j_2 \cdots j_{2k}} R^{h j_2}{}_{h_1 h_2} \cdots$, and verified for this metric:

- $k = 1$: $P_{(1)} = 2R\,\delta - 4\,\mathrm{Ric} = -4G$ and $L_{(1)} = 2R$ (`k1_equals_minus_4_einstein`, `L1_equals_2R`).
- $k = 2$: $P_{(2)} = -8\mathcal{H}$, with the Gauss-Bonnet (Lanczos) tensor $\mathcal{H}^{h}{}_{j} = 2(RR^{h}{}_{j} - 2R^{h}{}_{a}R^{a}{}_{j} - 2R^{ha}{}_{jb}R^{b}{}_{a} + R^{habc}R_{jabc}) - \frac{1}{2}\delta^{h}_{j}(R^2 - 4R_{ab}R^{ab} + R_{abcd}R^{abcd})$, and $L_{(2)} = 4(R^2 - 4R^{a}{}_{b}R^{b}{}_{a} + R^{ab}{}_{cd}R^{cd}{}_{ab})$ (`k2_equals_minus_8_gauss_bonnet`, `L2_equals_4_gauss_bonnet`, `gauss_bonnet_index_placement`).
- $k = 3$: $L_{(3)} = 8(2T_1 + 8T_2 + 24T_3 + 3T_4 + 24T_5 + 16T_6 - 12T_7 + T_8)$ with the cubic invariants $T_1 = R^{ab}{}_{cd}R^{cd}{}_{ef}R^{ef}{}_{ab}$, $T_2 = R^{ab}{}_{cd}R^{ce}{}_{bf}R^{df}{}_{ae}$, $T_3 = R^{ab}{}_{cd}R^{cd}{}_{be}R^{e}{}_{a}$, $T_4 = R\,R^{ab}{}_{cd}R^{cd}{}_{ab}$, $T_5 = R^{ab}{}_{cd}R^{c}{}_{a}R^{d}{}_{b}$, $T_6 = R^{a}{}_{b}R^{b}{}_{c}R^{c}{}_{a}$, $T_7 = R\,R^{a}{}_{b}R^{b}{}_{a}$ and $T_8 = R^3$ (`L3_equals_8_cubic_lovelock_density`).
- The constants are also derived, not only assumed: from literal δ sums on random algebraic curvature tensors, $P_{(1)}/G$ takes the single value -4 in $d = 4$, $P_{(2)}/\mathcal{H}$ the single value -8 in $d = 5$, and in $d = 6$ the 9 x 8 system of nine random tensors has rank 8 with the unique solution c = [16, 64, 192, 24, 192, 128, -96, 8] = 8 x [2, 8, 24, 3, 24, 16, -12, 1] (`normalisation_P1_derived_minus_4`, `normalisation_P2_derived_minus_8`, `normalisation_L3_cubic_derived`).
- Trace identity: $\sum_h P_{(k)}{}^{h}{}_{h} = (n - 2k)L_{(k)} = (8 - 2k)L_{(k)}$.

4.4 Why the series stops at k = 3

$P_{(4)}$ would need δ with $2\cdot4+1 = 9$ upper and 9 lower indices over eight values: two indices of each list are equal, so every term vanishes and $P_{(4)} = 0$ identically (`k4_tensor_vanishes`, `k4_terms_vanish_literally`, `gkd_nine_indices_in_eight_dimensions_vanish`; `lovelock-tensors.json` key `k4`: "identically zero (GKD of 9 indices in 8 dimensions)"). Equation (4.38) stops at $k = m - 1 = 3$, so $k = 1, 2, 3$ is the complete series in eight dimensions.

5 How the code computes them

5.1 The crate

`Revision/gkd_lovelock/code` is the Rust crate `lovelock_gkd` (edition 2021, no dependencies, its own empty workspace). Its modules:

file	function or type	what it does
`gkd.rs`	`gkd.rs::GKD`	the generalized Kronecker delta by cases (a) to (d) of section 3.2
`gkd.rs`	`gkd.rs::kdelta_det`, `gkd.rs::compare_exhaustive`, `gkd.rs::compare_random`	the literal Leibniz determinant and the two self-test loops
`rational.rs`	`rational.rs::Rational`	exact rationals on `i128`; every operation that would overflow stops with a message
`poly.rs`	`poly.rs::Poly`, `poly.rs::d`	exact Laurent polynomials with rational coefficients in H, a_4' to a_4'''', $E = e^{a_4}$, $S = \sin^{1/3} z$, $C = \cot z$, with the exact derivatives in x_0 and x_4
`geometry.rs`	`geometry.rs::metric_diagonal`, `geometry.rs::Geometry`	the metric, its inverse, $\sqrt{\lvert g\rvert} = S^3C$, the Christoffel symbols, $R^a{}_{bcd}$, $R^{ab}{}_{cd}$, Ricci and R
`geometry.rs`	`geometry.rs::nonzero_mixed`, `geometry.rs::einstein_mixed`	the list of nonzero $R^{ab}{}_{cd}$; the Einstein tensor from the Ricci tensor
`lovelock.rs`	`lovelock.rs::lovelock_mixed`, `lovelock.rs::lovelock_scalar`, `lovelock.rs::dfs`	$P_{(k)}{}^h{}_j$ and $L_{(k)}$ by a depth-first search over the nonzero curvature entries
`lovelock.rs`	`lovelock.rs::lovelock_contravariant`, `lovelock.rs::divergence`	$A_{(k)}{}^{lh}$; the covariant divergence $\nabla_h P^h{}_j$

file	function or type	what it does
lovelock.rs	lovelock.rs::brute_force_numeric, lovelock.rs::usual_normalisation	the unpruned literal sum at a numerical point; the factor $-1/2^{k+1}$
output.rs	output.rs::tensor_json, output.rs::monomials_json, output.rs::tensor_markdown	deterministic JSON and Markdown writers
main.rs	main.rs::gkd_selftest, main.rs::run_lovelock, main.rs::print_config	the command line gkd-selftest, lovelock, print-config

The symbols S and C satisfy $S^6(1+C^2)=1$, so the representation of `poly.rs` is not canonical in general; S cancels identically in every mixed tensor the program prints, every zero test is done on mixed tensors, and the independent checkers use a canonical form (section 7.2).

5.2 The enumeration

For each pair (h,j), `lovelock.rs::dfs` chooses k ordered nonzero entries $R^{ab}{}_{cd}$ one after another and appends a,b to the lower list and c,d to the upper list. A branch is cut when an index would repeat within a list (GKD is 0 by cases (a) and (b)); at a leaf the two index SETS must be equal (otherwise GKD is 0 by case (c)), and only then is GKD called; a nonzero value multiplies the product of the chosen entries. A vanishing curvature factor gives a vanishing term, so only the nonzero entries are enumerated. The counters of `lovelock-report.json`:

k	**leaves**	**GKD calls**	**nonzero GKD values**	**GKD calls for $L_{(k)}$**	**nonzero components of $P_{(k)}$**
1	5616	696	696	108	10
2	176640	32640	32640	7200	10
3	1128960	495360	495360	213120	10
4	0	0	0	1105920	0

The pruning is checked, not assumed: `lovelock.rs::brute_force_numeric` evaluates the completely unpruned literal sum, every index list sent to GKD, at the point $H=0.23$, $a_4=0.17$, $a_4'=0.61$, $a_4''=-0.37$, $b_4=1.37$, $b_4'=0.31$, $b_4''=-0.23$, $x_0=0.41$: 262144 index lists for $k=1$ (maximal relative deviation from the exact $P_{(1)}$ 9.99e-16) and 1073741824 index lists for $k=2$ (maximal relative deviation from the exact $P_{(2)}$ 3.89e-14) (`k1_brute_force_numeric`, `k2_brute_force_numeric`). The two independent checkers repeat the unpruned sums exactly (section 7).

6 Results

6.1 Christoffel symbols and curvature

The formulas of this section are those of the metric with $b_4(x_4)$, read from `curvature.json` and written in a_4', a_4'', $u = b_4'/b_4$ and $v = u'$ (at constant b_4, the withdrawn metric, every term with u or v drops out).

`curvature.json` lists 25 nonzero Christoffel symbols $\Gamma^a{}_{bc}$ with $b \le c$. With $i = 1, 2, 3$ (3-space) and $t = 5, 6, 7$ (extra times):

$$
\begin{aligned}
&\Gamma^{x_i}{}_{x_i x_4} = a_4', \quad \Gamma^{x_t}{}_{x_4 x_t} = -(a_4' - u/2), \quad \Gamma^{x_i}{}_{x_i x_0} = \Gamma^{x_t}{}_{x_t x_0} = H \cot z,\\
&\Gamma^{x_4}{}_{x_i x_i} = a_4' e^{2a_4} \sin^{1/3} z, \quad \Gamma^{x_4}{}_{x_t x_t} = (a_4' - u/2)\, e^{-2a_4} b_4 \sin^{1/3} z,\\
&\Gamma^{x_0}{}_{x_i x_i} = -H e^{2a_4} \sin^{1/3} z \, \tan z, \quad \Gamma^{x_0}{}_{x_t x_t} = H e^{-2a_4} b_4 \sin^{1/3} z \, \tan z,\\
&\Gamma^{x_0}{}_{x_0 x_0} = -6H \tan z - 6H \cot z.
\end{aligned}
$$

The mixed Riemann tensor $R^{ab}{}_{cd}$ has 156 nonzero entries. Up to the antisymmetry in each index pair they are, with $i \neq i'$ in 3-space and $t \neq t'$ among the extra times:

entry	**value**
$R^{x_i x_{i'}}{}_{x_i x_{i'}}$	$(a_4')^2 - H^2$
$R^{x_t x_{t'}}{}_{x_t x_{t'}}$	$(a_4' - u/2)^2 - H^2$
$R^{x_i x_t}{}_{x_i x_t}$	$-a_4'(a_4' - u/2) - H^2$
$R^{x_i x_4}{}_{x_i x_4}$	$a_4'' + (a_4')^2$
$R^{x_4 x_t}{}_{x_4 x_t}$	$-a_4'' + v/2 + (a_4' - u/2)^2$
$R^{x_i x_0}{}_{x_i x_0}$ and $R^{x_t x_0}{}_{x_t x_0}$	$-H^2$
$R^{x_i x_4}{}_{x_i x_0}$	$H a_4' \cot z$
$R^{x_4 x_t}{}_{x_t x_0}$	$H(a_4' - u/2) \cot z$
$R^{x_i x_0}{}_{x_i x_4}$	$-H a_4' \tan z$
$R^{x_t x_0}{}_{x_4 x_t}$	$-H(a_4' - u/2) \tan z$

$R^{x_0 x_4}{}_{x_0 x_4} = 0$. Every $R^{ab}{}_{cd}$ is free of $\sin^{1/3} z$ and of e^{a_4} (`mixed_riemann_free_of_sin_third`, `mixed_riemann_free_of_warp_and_exponential`). The extra times enter through the rate $a_4' - u/2$ of their scale factor, with the opposite sign of the inflating directions in the x_4 entries: the sectional curvature of the plane (x_i, x_4) is $a_4'' + (a_4')^2$ and that of (x_4, x_t) is $-a_4'' + v/2 + (a_4' - u/2)^2$.

The mixed Ricci tensor is diagonal except for the pair x_4-x_0: $R^{x_i}{}_{x_i} = a_4'' - 6H^2 + \frac{3}{2} a_4' u$, $R^{x_4}{}_{x_4} = 6(a_4')^2 - 3a_4' u + \frac{3}{4} u^2 + \frac{3}{2} v$, $R^{x_t}{}_{x_t} = -a_4'' - 6H^2 - \frac{3}{2} a_4' u + \frac{3}{4} u^2 + \frac{1}{2} v$, $R^{x_0}{}_{x_0} = -6H^2$, $R^{x_4}{}_{x_0} = \frac{3}{2} H u \cot z$, $R^{x_0}{}_{x_4} = -\frac{3}{2} H u \tan z$; the Ricci scalar is $R = 6(a_4')^2 - 3a_4' u + 3u^2 + 3v - 42H^2$ (`ricciScalar`), and the Einstein tensor is

$$
\begin{aligned}
&G^{x_i}{}_{x_i} = a_4'' - 3(a_4')^2 + 3a_4' u + 15H^2 - \tfrac{3}{2} u^2 - \tfrac{3}{2} v, \quad G^{x_4}{}_{x_4} = 3(a_4')^2 - \tfrac{3}{2} a_4' u + 21H^2 - \tfrac{3}{4} u^2,\\
&G^{x_t}{}_{x_t} = -a_4'' - 3(a_4')^2 + 15H^2 - \tfrac{3}{4} u^2 - v, \quad G^{x_0}{}_{x_0} = -3(a_4')^2 + \tfrac{3}{2} a_4' u + 15H^2 - \tfrac{3}{2} u^2 - \tfrac{3}{2} v,\\
&G^{x_4}{}_{x_0} = \tfrac{3}{2} H u \cot z, \quad G^{x_0}{}_{x_4} = -\tfrac{3}{2} H u \tan z.
\end{aligned}
$$

6.2 The Lovelock scalars

From `lovelock-tensors.json` (keys `L1`, `L2`, `L3`), with $b_4''/b_4 = v + u^2$:

$$
\begin{aligned}
L_{(1)} &= 12\,(a_4')^2 - 6\,a_4'\,u + 6\,u^2 \\
&\quad + 6\,v - 84\,H^2, \\
L_{(2)} &= -96\,(a_4')^4 + 96\,(a_4')^3\,u - 144\,(a_4')^2\,u^2 \\
&\quad - 96\,(a_4')^2\,v - 2112\,(a_4')^2\,H^2 - 192\,a_4'\,a_4''\,u \\
&\quad + 60\,a_4'\,u^3 + 96\,a_4'\,u\,v + 1056\,a_4'\,u\,H^2 \\
&\quad + 48\,a_4''\,u^2 + 6\,u^4 + 12\,u^2\,v \\
&\quad - 552\,u^2\,H^2 - 720\,v\,H^2 + 3360\,H^4, \\
L_{(3)} &= 1152\,(a_4')^6 - 1728\,(a_4')^5\,u + 3456\,(a_4')^4\,u^2 \\
&\quad + 1728\,(a_4')^4\,v + 31104\,(a_4')^4\,H^2 + 6912\,(a_4')^3\,a_4''\,u \\
&\quad - 2736\,(a_4')^3\,u^3 - 3456\,(a_4')^3\,u\,v - 31104\,(a_4')^3\,u\,H^2 \\
&\quad - 5184\,(a_4')^2\,a_4''\,u^2 + 648\,(a_4')^2\,u^4 + 1296\,(a_4')^2\,u^2\,v \\
&\quad + 10368\,(a_4')^2\,u^2\,H^2 + 6912\,(a_4')^2\,v\,H^2 + 100224\,(a_4')^2\,H^4 \\
&\quad + 864\,a_4'\,a_4''\,u^3 + 13824\,a_4'\,a_4''\,u\,H^2 - 1296\,a_4'\,u^3\,H^2 \\
&\quad - 6912\,a_4'\,u\,v\,H^2 - 50112\,a_4'\,u\,H^4 - 3456\,a_4''\,u^2\,H^2 \\
&\quad - 432\,u^4\,H^2 - 864\,u^2\,v\,H^2 + 13824\,u^2\,H^4 \\
&\quad + 25920\,v\,H^4 - 40320\,H^6.
\end{aligned}
$$

6.3 The Lovelock tensors

Every ${P_{(k)}}^h{}_j$ has 10 nonzero components: the 8 diagonal ones and the pair ${P^{x_0}}_{x_4}$, ${P^{x_4}}_{x_0}$, which are proportional to b_4' (they would vanish only for a constant b_4, the withdrawn metric); the other 54 off-diagonal components vanish. The diagonal satisfies $x_1 = x_2 = x_3$ and $x_5 = x_6 = x_7$, so four components per order are independent. The formulas below are the LaTeX texts of `lovelock-tensors.json`, unchanged (only broken into lines at top-level signs).

$k = 1$:

$$
{P_{(1)}}^{x_0}{}_{x_0} = 6\,b_4^{-1}\,b_4'' - 6\,a_4'\,b_4^{-1}\,b_4' + 12\,(a_4')^2 - 60\,H^2
$$

$$
\begin{aligned}
{P_{(1)}}^{x_1}{}_{x_1} &= 6\,b_4^{-1}\,b_4'' - 4\,a_4'' - 12\,a_4'\,b_4^{-1}\,b_4' \\
&\quad + 12\,(a_4')^2 - 60\,H^2
\end{aligned}
$$

$$
{P_{(1)}}^{x_4}{}_{x_4} = 3\,b_4^{-2}\,(b_4')^2 + 6\,a_4'\,b_4^{-1}\,b_4' - 12\,(a_4')^2 - 84\,H^2
$$

$$
\begin{aligned}
{P_{(1)}}^{x_5}{}_{x_5} &= -b_4^{-2}\,(b_4')^2 + 4\,b_4^{-1}\,b_4'' + 4\,a_4'' + 12\,(a_4')^2 \\
&\quad - 60\,H^2
\end{aligned}
$$

$$P_{(1)}{}^{x_0}{}_{x_4} = 6\,H\,\cot^{-1}(6Hx_0)\,b_4^{-1}\,b_4'$$

$$P_{(1)}{}^{x_4}{}_{x_0} = -6\,H\,\cot(6Hx_0)\,b_4^{-1}\,b_4'$$

$k = 2$:

$$\begin{aligned}
P_{(2)}{}^{x_0}{}_{x_0} = &-6\,b_4^{-4}\,(b_4')^4 + 12\,b_4^{-3}\,(b_4')^2\,b_4'' \\
&+ 48\,a_4''\,b_4^{-2}\,(b_4')^2 - 36\,a_4'\,b_4^{-3}\,(b_4')^3 \\
&+ 96\,a_4'\,b_4^{-2}\,b_4'\,b_4'' - 192\,a_4'\,a_4''\,b_4^{-1}\,b_4' \\
&- 48\,(a_4')^2\,b_4^{-2}\,(b_4')^2 - 96\,(a_4')^2\,b_4^{-1}\,b_4'' \\
&+ 96\,(a_4')^3\,b_4^{-1}\,b_4' - 96\,(a_4')^4 \\
&+ 96\,H^2\,b_4^{-2}\,(b_4')^2 - 480\,H^2\,b_4^{-1}\,b_4'' \\
&+ 672\,H^2\,a_4'\,b_4^{-1}\,b_4' - 1344\,H^2\,(a_4')^2 \\
&+ 1440\,H^4
\end{aligned}$$

$$\begin{aligned}
P_{(2)}{}^{x_1}{}_{x_1} = &-6\,b_4^{-4}\,(b_4')^4 + 12\,b_4^{-3}\,(b_4')^2\,b_4'' \\
&+ 24\,a_4''\,b_4^{-2}\,(b_4')^2 - 24\,a_4'\,b_4^{-3}\,(b_4')^3 \\
&+ 48\,a_4'\,b_4^{-2}\,b_4'\,b_4'' - 192\,a_4'\,a_4''\,b_4^{-1}\,b_4' \\
&- 48\,(a_4')^2\,b_4^{-2}\,(b_4')^2 - 96\,(a_4')^2\,b_4^{-1}\,b_4'' \\
&+ 192\,(a_4')^2\,a_4'' + 192\,(a_4')^3\,b_4^{-1}\,b_4' \\
&- 96\,(a_4')^4 + 144\,H^2\,b_4^{-2}\,(b_4')^2 \\
&- 480\,H^2\,b_4^{-1}\,b_4'' + 320\,H^2\,a_4'' \\
&+ 960\,H^2\,a_4'\,b_4^{-1}\,b_4' - 1344\,H^2\,(a_4')^2 \\
&+ 1440\,H^4
\end{aligned}$$

$$\begin{aligned}
P_{(2)}{}^{x_4}{}_{x_4} = &\;36\,a_4'\,b_4^{-3}\,(b_4')^3 - 288\,(a_4')^3\,b_4^{-1}\,b_4' \\
&+ 288\,(a_4')^4 - 240\,H^2\,b_4^{-2}\,(b_4')^2 \\
&- 480\,H^2\,a_4'\,b_4^{-1}\,b_4' + 960\,H^2\,(a_4')^2 \\
&+ 3360\,H^4
\end{aligned}$$

$$\begin{aligned}
P_{(2)}{}^{x_5}{}_{x_5} = &\;24\,a_4''\,b_4^{-2}\,(b_4')^2 - 24\,a_4'\,b_4^{-3}\,(b_4')^3 \\
&+ 48\,a_4'\,b_4^{-2}\,b_4'\,b_4'' - 192\,(a_4')^2\,a_4'' \\
&- 96\,(a_4')^4 + 128\,H^2\,b_4^{-2}\,(b_4')^2 \\
&- 320\,H^2\,b_4^{-1}\,b_4'' - 320\,H^2\,a_4'' \\
&+ 384\,H^2\,a_4'\,b_4^{-1}\,b_4' - 1344\,H^2\,(a_4')^2 \\
&+ 1440\,H^4
\end{aligned}$$

$$
\begin{aligned}
P_{(2)}{}^{x_0}{}_{x_4} &= 12\,H\,\cot^{-1}(6Hx_0)\,b_4^{-3}\,(b_4')^3 \\
&\quad + 144\,H\,a_4'\,\cot^{-1}(6Hx_0)\,b_4^{-2}\,(b_4')^2 \\
&\quad - 288\,H\,(a_4')^2\,\cot^{-1}(6Hx_0)\,b_4^{-1}\,b_4' \\
&\quad - 480\,H^3\,\cot^{-1}(6Hx_0)\,b_4^{-1}\,b_4'
\end{aligned}
$$

$$
\begin{aligned}
P_{(2)}{}^{x_4}{}_{x_0} &= -12\,H\,\cot(6Hx_0)\,b_4^{-3}\,(b_4')^3 \\
&\quad - 144\,H\,a_4'\,\cot(6Hx_0)\,b_4^{-2}\,(b_4')^2 \\
&\quad + 288\,H\,(a_4')^2\,\cot(6Hx_0)\,b_4^{-1}\,b_4' \\
&\quad + 480\,H^3\,\cot(6Hx_0)\,b_4^{-1}\,b_4'
\end{aligned}
$$

$k = 3$:

$$
\begin{aligned}
P_{(3)}{}^{x_0}{}_{x_0} &= 864\,a_4'\,a_4''\,b_4^{-3}\,(b_4')^3 \\
&\quad - 648\,(a_4')^2\,b_4^{-4}\,(b_4')^4 \\
&\quad + 1296\,(a_4')^2\,b_4^{-3}\,(b_4')^2\,b_4'' \\
&\quad - 5184\,(a_4')^2\,a_4''\,b_4^{-2}\,(b_4')^2 \\
&\quad + 720\,(a_4')^3\,b_4^{-3}\,(b_4')^3 \\
&\quad - 3456\,(a_4')^3\,b_4^{-2}\,b_4'\,b_4'' \\
&\quad + 6912\,(a_4')^3\,a_4''\,b_4^{-1}\,b_4' \\
&\quad + 1728\,(a_4')^4\,b_4^{-2}\,(b_4')^2 \\
&\quad + 1728\,(a_4')^4\,b_4^{-1}\,b_4'' - 1728\,(a_4')^5\,b_4^{-1}\,b_4' \\
&\quad + 1152\,(a_4')^6 + 216\,H^2\,b_4^{-4}\,(b_4')^4 \\
&\quad - 432\,H^2\,b_4^{-3}\,(b_4')^2\,b_4'' \\
&\quad - 1728\,H^2\,a_4''\,b_4^{-2}\,(b_4')^2 \\
&\quad + 2160\,H^2\,a_4'\,b_4^{-3}\,(b_4')^3 \\
&\quad - 3456\,H^2\,a_4'\,b_4^{-2}\,b_4'\,b_4'' \\
&\quad + 6912\,H^2\,a_4'\,a_4''\,b_4^{-1}\,b_4' \\
&\quad + 1728\,H^2\,(a_4')^2\,b_4^{-2}\,(b_4')^2 \\
&\quad + 3456\,H^2\,(a_4')^2\,b_4^{-1}\,b_4'' \\
&\quad - 10368\,H^2\,(a_4')^3\,b_4^{-1}\,b_4' + 10368\,H^2\,(a_4')^4 \\
&\quad - 3456\,H^4\,b_4^{-2}\,(b_4')^2 + 8640\,H^4\,b_4^{-1}\,b_4'' \\
&\quad - 15552\,H^4\,a_4'\,b_4^{-1}\,b_4' + 31104\,H^4\,(a_4')^2 \\
&\quad - 5760\,H^6
\end{aligned}
$$

$$
\begin{aligned}
{P_{(3)}}^{x_1}{}_{x_1} = {} & 288\, a_4'\, a_4''\, b_4^{-3}\, (b_4')^3 \\
& - 216\, (a_4')^2\, b_4^{-4}\, (b_4')^4 \\
& + 432\, (a_4')^2\, b_4^{-3}\, (b_4')^2\, b_4'' \\
& - 2592\, (a_4')^2\, a_4''\, b_4^{-2}\, (b_4')^2 \\
& + 288\, (a_4')^3\, b_4^{-3}\, (b_4')^3 \\
& - 1728\, (a_4')^3\, b_4^{-2}\, b_4'\, b_4'' \\
& + 6912\, (a_4')^3\, a_4''\, b_4^{-1}\, b_4' \\
& + 1728\, (a_4')^4\, b_4^{-2}\, (b_4')^2 \\
& + 1728\, (a_4')^4\, b_4^{-1}\, b_4'' - 5760\, (a_4')^4\, a_4'' \\
& - 3456\, (a_4')^5\, b_4^{-1}\, b_4' + 1152\, (a_4')^6 \\
& + 216\, H^2\, b_4^{-4}\, (b_4')^4 \\
& - 432\, H^2\, b_4^{-3}\, (b_4')^2\, b_4'' \\
& - 864\, H^2\, a_4''\, b_4^{-2}\, (b_4')^2 \\
& + 2592\, H^2\, a_4'\, b_4^{-3}\, (b_4')^3 \\
& - 1728\, H^2\, a_4'\, b_4^{-2}\, b_4'\, b_4'' \\
& + 6912\, H^2\, a_4'\, a_4''\, b_4^{-1}\, b_4' \\
& - 3456\, H^2\, (a_4')^2\, b_4^{-2}\, (b_4')^2 \\
& + 3456\, H^2\, (a_4')^2\, b_4^{-1}\, b_4'' \\
& - 6912\, H^2\, (a_4')^2\, a_4'' \\
& - 6912\, H^2\, (a_4')^3\, b_4^{-1}\, b_4' + 10368\, H^2\, (a_4')^4 \\
& - 5184\, H^4\, b_4^{-2}\, (b_4')^2 + 8640\, H^4\, b_4^{-1}\, b_4'' \\
& - 5760\, H^4\, a_4'' - 17280\, H^4\, a_4'\, b_4^{-1}\, b_4' \\
& + 31104\, H^4\, (a_4')^2 - 5760\, H^6
\end{aligned}
$$

$$
\begin{aligned}
{P_{(3)}}^{x_4}{}_{x_4} = {} & 720\, (a_4')^3\, b_4^{-3}\, (b_4')^3 \\
& - 4320\, (a_4')^4\, b_4^{-2}\, (b_4')^2 \\
& + 8640\, (a_4')^5\, b_4^{-1}\, b_4' - 5760\, (a_4')^6 \\
& - 1296\, H^2\, a_4'\, b_4^{-3}\, (b_4')^3 \\
& + 10368\, H^2\, (a_4')^3\, b_4^{-1}\, b_4' - 10368\, H^2\, (a_4')^4 \\
& + 4320\, H^4\, b_4^{-2}\, (b_4')^2 + 8640\, H^4\, a_4'\, b_4^{-1}\, b_4' \\
& - 17280\, H^4\, (a_4')^2 - 40320\, H^6
\end{aligned}
$$

$$
\begin{aligned}
{P_{(3)}}^{x_5}{}_{x_5} = {} & 864\,(a_4')^2\,a_4''\,b_4^{-2}\,(b_4')^2 \\
& - 288\,(a_4')^3\,b_4^{-3}\,(b_4')^3 \\
& + 576\,(a_4')^3\,b_4^{-2}\,b_4'\,b_4'' \\
& - 4608\,(a_4')^3\,a_4''\,b_4^{-1}\,b_4' \\
& + 288\,(a_4')^4\,b_4^{-2}\,(b_4')^2 \\
& - 1152\,(a_4')^4\,b_4^{-1}\,b_4'' + 5760\,(a_4')^4\,a_4'' \\
& + 1152\,(a_4')^6 - 864\,H^2\,a_4''\,b_4^{-2}\,(b_4')^2 \\
& + 864\,H^2\,a_4'\,b_4^{-3}\,(b_4')^3 \\
& - 1728\,H^2\,a_4'\,b_4^{-2}\,b_4'\,b_4'' \\
& + 5184\,H^2\,(a_4')^2\,b_4^{-2}\,(b_4')^2 \\
& + 6912\,H^2\,(a_4')^2\,a_4'' \\
& - 13824\,H^2\,(a_4')^3\,b_4^{-1}\,b_4' + 10368\,H^2\,(a_4')^4 \\
& - 3168\,H^4\,b_4^{-2}\,(b_4')^2 + 5760\,H^4\,b_4^{-1}\,b_4'' \\
& + 5760\,H^4\,a_4'' - 13824\,H^4\,a_4'\,b_4^{-1}\,b_4' \\
& + 31104\,H^4\,(a_4')^2 - 5760\,H^6
\end{aligned}
$$

$$
\begin{aligned}
{P_{(3)}}^{x_0}{}_{x_4} = {} & 2160\,H\,(a_4')^2\,\cot^{-1}(6Hx_0)\,b_4^{-3}\,(b_4')^3 \\
& - 8640\,H\,(a_4')^3\,\cot^{-1}(6Hx_0)\,b_4^{-2}\,(b_4')^2 \\
& + 8640\,H\,(a_4')^4\,\cot^{-1}(6Hx_0)\,b_4^{-1}\,b_4' \\
& - 432\,H^3\,\cot^{-1}(6Hx_0)\,b_4^{-3}\,(b_4')^3 \\
& - 5184\,H^3\,a_4'\,\cot^{-1}(6Hx_0)\,b_4^{-2}\,(b_4')^2 \\
& + 10368\,H^3\,(a_4')^2\,\cot^{-1}(6Hx_0)\,b_4^{-1}\,b_4' \\
& + 8640\,H^5\,\cot^{-1}(6Hx_0)\,b_4^{-1}\,b_4'
\end{aligned}
$$

$$
\begin{aligned}
{P_{(3)}}^{x_4}{}_{x_0} = {} & -2160\,H\,(a_4')^2\,\cot(6Hx_0)\,b_4^{-3}\,(b_4')^3 \\
& + 8640\,H\,(a_4')^3\,\cot(6Hx_0)\,b_4^{-2}\,(b_4')^2 \\
& - 8640\,H\,(a_4')^4\,\cot(6Hx_0)\,b_4^{-1}\,b_4' \\
& + 432\,H^3\,\cot(6Hx_0)\,b_4^{-3}\,(b_4')^3 \\
& + 5184\,H^3\,a_4'\,\cot(6Hx_0)\,b_4^{-2}\,(b_4')^2 \\
& - 10368\,H^3\,(a_4')^2\,\cot(6Hx_0)\,b_4^{-1}\,b_4' \\
& - 8640\,H^5\,\cot(6Hx_0)\,b_4^{-1}\,b_4'
\end{aligned}
$$

Remarks, each an exact property of the listed components:

- $P_{(1)} = -4G$ component by component (compare section 6.1).
- Of the metric functions only derivatives occur: a_4' and a_4'', not a_4 itself, and b_4 only through $u = b_4'/b_4$ and $v = u'$ (a constant factor of b_4 is a rescaling of x_5, x_6, x_7; record `lovelock_depends_on_b4_through_u_and_v`). The x_4 components contain no a_4''.

- $P^{x_1}_{(k)x_1} + P^{x_5}_{(k)x_5} - 2P^{x_0}_{(k)x_0}$ contains u or v in every term, for $k = 1, 2, 3$: it vanishes for constant b_4, the identity $P^{x_1}_{(k)x_1} + P^{x_5}_{(k)x_5} = 2P^{x_0}_{(k)x_0}$ of the withdrawn metric, but not for the metric with b_4 (section 9).

- The contravariant densities are ${A_{(k)}}^{ll} = b_4^{3/2} \cos z \; g^{ll} \, {P_{(k)}}^{l}{}_{l}$ for every l ($\sqrt{|g|} = b_4^{3/2} \cos z$; the off-diagonal pair ${A_{(k)}}^{x_0x_4}$, ${A_{(k)}}^{x_4x_0}$ is built in the same way from ${P_{(k)}}^{x_0}{}_{x_4}$, ${P_{(k)}}^{x_4}{}_{x_0}$), with $g^{x_0x_0} = \tan^2 z$, $g^{x_ix_i} = e^{-2a_4} \sin^{-1/3} z$, $g^{x_4x_4} = -1$ and $g^{x_tx_t} = -e^{2a_4} b_4^{-1} \sin^{-1/3} z$; for example the file gives ${A_{(1)}}^{x_4x_4} = -3 \sin(6Hx_0) \cot(6Hx_0) \, b_4^{-1/2} \, (b_4')^2 - 6 \, a_4' \sin(6Hx_0) \cot(6Hx_0) \, b_4^{1/2} \, b_4' + 12 \, (a_4')^2 \sin(6Hx_0) \cot(6Hx_0) \, b_4^{3/2} + 84 \, H^2 \sin(6Hx_0) \cot(6Hx_0) \, b_4^{3/2}$. All 64 components of each $A_{(k)}$, each with its exact monomial list, are in `lovelock-tensors.json`; all 384 components of $P_{(k)}$ and $A_{(k)}$ are also written out in `lovelock-components.md`.

6.4 The checks of the Rust program

`lovelock_gkd lovelock -brute-force-k2` writes `curvature.json`, `lovelock-tensors.json`, `lovelock-components.md` and `lovelock-report.json`. The report gives `"checkCount": 19`, `"failedCheckCount": 0` and `"verdict": "SUCCESS"`; every one of its 19 checks has `"passed": true`:

check	check	check
riemann_antisymmetry	riemann_first_bianchi	mixed_riemann_free_of_sin_third
k1_equals_minus_4_einstein	k1_trace_identity	k1_divergence_free
k1_symmetric	k1_free_of_sin_third	k2_trace_identity
k2_divergence_free	k2_symmetric	k2_free_of_sin_third
k3_trace_identity	k3_divergence_free	k3_symmetric
k3_free_of_sin_third	k4_tensor_vanishes	k1_brute_force_numeric
k2_brute_force_numeric		

They are mathematical identities, not comparisons with the author's numbers: the antisymmetries and the first Bianchi identity of ${R^{ab}}_{cd}$ for all 4096 index lists, $P_{(1)} = -4G$ by an independent route, the trace identities, $\nabla_h {P_{(k)}}^{h}{}_{j} = 0$ for every j, the symmetry of $g_{hh} {P_{(k)}}^{h}{}_{j}$ (so ${A_{(k)}}^{lh} = {A_{(k)}}^{hl}$), the absence of the warp factor, $P_{(4)} = 0$ and the unpruned numerical sums. Two runs give byte-identical files (`Revision/gkd_lovelock/results/PROVENANCE_OF_THE_COMPUTATION.md`).

7 Independent verifications

7.1 The Wolfram check (29 checks)

`Revision/gkd_lovelock/verification/verify_lovelock_gkd.wls` with the package `Lovelock GKDCheck.wl` writes `wolfram-gkd-report.json`: `"checkCount": 29`, `"expectedCheckCount": 29`, `"failedCheckCount": 0`, `"verdict": "SUCCESS"`, producer Wolfram Language 15.0.1. What it does:

- The author's definition, re-typed verbatim, is the kernel's definition of `kδ` (compared with the text recorded in `PROVENANCE_OF_THE_COMPUTATION.md`), with `delta[a_Integer, b_Integer] := KroneckerDelta[a, b]`; unequal lengths stay unevaluated.
- GKD is NOT re-implemented in Wolfram Language: a small exporter compiled with cargo against the unchanged crate writes the values of the Rust function GKD for 346304 pairs (all 64, 4096 and 262144 pairs of lengths 1, 2, 3 and 20000 pseudo-random pairs of each length 4 to 7; values file of 3161984 bytes), and the verbatim `kδ` is evaluated on every pair: 0 mismatches. The negative control against −GKD reports a mismatch at exactly every pair where GKD is nonzero (8, 112, 2016, 10022, 10005, 10002, 10000 pairs for lengths 1 to 7).
- The curvature is recomputed from the metric with `D`, `Inverse` and `Simplify` (156 nonzero ${R^{ab}}_{cd}$, the same as `curvature.json`), and $\det g = \cos^2 z$.
- The Lovelock tensors are recomputed with the verbatim `kδ`: for $k = 1$ and $k = 2$ every k-tuple of the 156 nonzero entries for each of the 64 components (9984 and 1557504 calls), for $k = 3$ every list left after skipping lists with a repeated label in a row (1128960 calls). `FullSimplify` of the difference with the Rust tensors is 0 for all 64 components of each $P_{(k)}$ and $A_{(k)}$. The numbers of nonzero `kδ` terms, 696, 32640 and 495360, equal the Rust counters. A tampered $P_{(2)}$ component and a tampered $A_{(3)}$ component are flagged, and only those.

check	**check**
definition_is_the_authors_verbatim	definition_unequal_lengths_stay_unevaluated
gkd_values_file_layout	gkd_equals_kdelta_exhaustive_length_1
gkd_equals_kdelta_exhaustive_length_2	gkd_equals_kdelta_exhaustive_length_3
gkd_equals_kdelta_random_length_4	gkd_equals_kdelta_random_length_5
gkd_equals_kdelta_random_length_6	gkd_equals_kdelta_random_length_7
negative_control_gkd_comparison_detects_a_sign_flip	riemann_mixed_equals_rust_curvature
sqrt_det_g_is_cos	lovelock_kdelta_values_are_integers
k1_P_equals_rust_all_64_components	k1_A_equals_rust_all_64_components
k2_P_equals_rust_all_64_components	k2_A_equals_rust_all_64_components
k3_P_equals_rust_all_64_components	k3_A_equals_rust_all_64_components
negative_control_tensor_comparison_detects_a_change	k1_equals_minus_4_einstein
k1_trace_equals_6_L1	k2_trace_equals_4_L2
k3_trace_equals_2_L3	k1_nonzero_kdelta_terms_equal_rust_counter
k2_nonzero_kdelta_terms_equal_rust_counter	k3_nonzero_kdelta_terms_equal_rust_counter
rust_json_text_equals_its_monomials	

Scope. The Wolfram check is independent of the Rust code in its definition of δ (the author's verbatim text), its curvature and its Lovelock sums; it uses the Rust program for the GKD values it compares and for the tensors it compares with. Its default run skips, for $k = 3$, lists with a repeated label in a row (their determinant is 0 because two rows or columns are equal); an optional mode

recomputes the eight diagonal $k = 3$ components with no skip at all, and its single recorded run passed 30 of 30 checks (`Revision/gkd_lovelock/verification/WOLFRAMSCRIPT_PROVENANCE.md`, part 4.5); that report is not committed. Only Wolfram 15.0.1 was used.

7.2 The sympy checker (49 checks)

The sympy checker is the file `check_lovelock_gkd.py` in the folder `Revision/gkd_lovelock/verification/`. Its report `python-lovelock-report.json` gives `"checkCount": 49`, `"failedCheckCount": 0` and `"verdict": "SUCCESS"`. It shares no code with the Rust crate: the geometry is sympy's own differentiation, δ is the author's definition evaluated with sympy's determinant (memoised by the 0/1 matrix), and the sums are organised breadth-first (all k-fold products with pairwise different lower and pairwise different upper indices, then completed by every admissible h and j, one literal call of the determinant per term). Its counters:

k	**products**	**determinant calls**	**nonzero values**	**distinct multisets**
1	156	5772	804	156
2	11040	187680	39840	5520
3	282240	1411200	708480	47040

with 4096410 determinant calls in all and 29096 distinct 0/1 matrices determined by sympy (4096401 and 29030 before the relabel of 2026-10-14: the random samples of the checker are drawn over the labels in the new order). The skipped terms are checked as well: the completely unpruned literal sums for $k = 1$ (10140 calls) and $k = 2$ (1581840 calls) agree exactly with the pruned sums; 20000 random terms of the unpruned $k = 3$ sum, 19909 of them with a repeated index, give 0 whenever skipped; 300 random 9-index lists give 0. The comparison with the Rust output is exact three times over: by an exact canonical form (`canon`, which uses $s^6(1 + c^2) = 1$ to reach a unique form, so a zero difference is a proof of equality), by plain sympy simplification, and numerically with 60 digits at 5 random rational points.

check	**check**	**check**
`gkd_literal_equals_cofactor_expansion`	`gkd_examples`	`gkd_nine_indices_in_eight_dimensions_vanish`
`normalisation_P1_derived_minus_4`	`normalisation_P2_derived_minus_8`	`normalisation_L3_cubic_derived`
`metric_inverse`	`sqrt_abs_det_g`	`christoffel_symmetric`
`riemann_first_bianchi`	`mixed_riemann_free_of_warp_and_exponential`	`riemann_antisymmetry_and_pair_symmetry`
`k1_unpruned_literal_sum_agrees`	`k2_unpruned_literal_sum_agrees`	`k3_pruned_terms_vanish_literally`
`k4_terms_vanish_literally`	`k1_equals_minus_4_einstein`	`L1_equals_2R`
`k2_equals_minus_8_gauss_bonnet`	`L2_equals_4_gauss_bonnet`	`gauss_bonnet_index_placement`
`L3_equals_8_cubic_lovelock_density`	`k1_trace_identity`	`k2_trace_identity`

check	check	check
k3_trace_identity	k1_divergence_free	k1_symmetric
k1_free_of_warp	k2_divergence_free	k2_symmetric
k2_free_of_warp	k3_divergence_free	k3_symmetric
k3_free_of_warp	rust_k1_mixed_co mponents_agree	rust_k1_contrava riant_components_agree
rust_L1_agrees	rust_k2_mixed_co mponents_agree	rust_k2_contrava riant_components_agree
rust_L2_agrees	rust_k3_mixed_co mponents_agree	rust_k3_contrava riant_components_agree
rust_L3_agrees	negative_control s_detected	rust_metric_and_ sqrt_g_agree
rust_christoffels_agree	rust_riemann_agrees	rust_ricci_einst ein_scalar_agree
rust_text_matche s_monomials		

Scope. The sympy checker reads only the metric (typed in from the task text) and, for the comparison, `curvature.json` and `lovelock-tensors.json`; it never reads the author's notebook. Its literal δ is compared with an independent cofactor expansion on all 266304 pairs of lengths 1 to 3 and 7500 random pairs of lengths 4 to 8. The negative controls (a coefficient increased by 1, a wrong power of $\sin^{1/3} z$, a wrong power of e^{a_4}, an added $10^{-30}H^4$) are each detected by each of its three comparison methods.

8 The comparison with the author's notebooks

8.1 What was read for the computation

What was read is recorded in `PROVENANCE_OF_THE_COMPUTATION.md`, written before any comparison, and in the provenance file `WOLFRAMSCRIPT_PROVENANCE.md` of the folder `Revision/gkd_lovelock/notebook_reading/`: only the 58 INPUT cells of `Generalized _Kronecker_Delta_4+4.nb` (sha256 `23bb4e0c70943e766d9b081a3a399ef29664889aad088041329ce2e065b80afb`), converted to text without evaluation; the committed digest `notebook-input-cells.txt` gives, for each, its label, its length and its first 160 characters. The definition of kδ was taken from the cell `In[87]` and Lovelock's equation from the picture in the cell `In[101]` (1372 x 435 pixels). For the computation no output cell, no file `Generalized_Kronecker_Delta.txt` and no other Mathematica file of the author was opened; the metric was taken from the author's message.

What the notebook computes, as far as its input cells show (a reading of the digest, labelled as such): it sets up the xAct packages, an 8-dimensional manifold and a metric of signature (4,4) (`In[29]`); it builds the generalized delta in a second way, as contracted products of two Levi-Civita tensors divided by $(8-p)!$ (`delta11`, `delta22`, `delta33`, `delta55`); it defines kδ (`In[87]`) and evaluates it for 3, 5 and 7 index pairs (kδ33, kδ55, kδ77); it compares the two constructions (`In[102]`, `In[105]`, `In[108]`); and it contains the picture of (4.38). Its answers are in its output cells, which were not read for the computation; section 8.2 compares no output of this notebook either.

The checks against the author's definitions, made with the computation:

- The definition: `definition_is_the_authors_verbatim` shows that the definition evaluated by the Wolfram check is the author's text, character for character.
- The function: GKD equals the author's `k`δ on every pair of lists of length 1 to 3, on 20000 random pairs of each length 4 to 7 (Wolfram), and equals the literal determinant on every pair of length 1 to 4 and 200000 random pairs of each length 5 to 9 (Rust self-test), as the theorem of section 3.2 requires.
- The equation: the tensors are those of (4.38) as shown in the notebook's picture, with the conventions of section 4.2.

8.2 The comparison with the author's stored outputs (2026-10-08)

`PROVENANCE_OF_THE_COMPUTATION.md` announces that the comparison with the author's own answers is done afterwards, in a separate, later commit; that comparison is recorded in `Revision/gkd_lovelock/comparison/`: its `README.md`, the read-only extractor `extract_author_curvature_outputs.wls` with its output `author-curvature-outputs.json`, and the program `compare_with_author.py` with its report `author-comparison-report.json`. Its files were first committed on 2026-10-08, after the results of this record (commit 3e81eeb, 2026-10-01). By its README it changes no file of `Revision/gkd_lovelock/results/`, `Revision/gkd_lovelock/code/`, `Revision/gkd_lovelock/verification/` or `Revision/gkd_lovelock/notebook_reading/`.

This notebook of the author computes the earlier metric, without b_4 (its `MatrixMetric44` has $b_4 = 1$, the withdrawn metric); the comparison therefore evaluates this record at $b_4 = 1$, the only place where this document restricts it. The metric with b_4 is compared with the author's new notebook in `Revision/metric_b4` (section 8.4).

What was read. The author's notebook `Pair_Creation_of_Universes_WaveFunctionOfUniverse-4+4-Einstein-Lovelock-Nash.nb` in the repository root (sha256 `5ee5cb2a95146136ee65636a4aef174f303b6c41da130d2c57e4684f9c8ff69f`), read only: `Get` of the file gives the inert notebook expression, and the stored output boxes are converted with `ToExpression[boxes, StandardForm, HoldComplete]`, so nothing is evaluated and no input cell is run. Of its 2127 cells (996 Input, 805 Output) the comparison uses these:

cell	what it is
`In[79]`, `In[82]`	the definition of the author's metric `MatrixMetric44`, in which `a4` is a function of the argument Hx_4
`In[214]`	the definition of `rt[g]`: inverse metric, Christoffel symbols, Riemann tensor, Ricci tensor, Ricci scalar and Einstein tensor
`Out[235]`	`gtry = MatrixMetric44`, the stored 8 x 8 metric
`In[238]`	`rt[gtry]`, which assigns `RS` and `Einstein G`; its output is suppressed
`Out[245]`	`MatrixMetric44` printed again (a control: identical to `Out[235]`)
`Out[535]`	the stored Ricci scalar `RS`

cell	what it is
Out[536]	the stored Einstein tensor EinsteinG, both indices down (8 x 8)

That Out[535] and Out[536] are the values returned by rt[gtry] at In[238] is an inference from the file, labelled as such: among all input cells only In[238] assigns the global symbols RS and EinsteinG. The notebook was not re-run; its outputs are taken as stored.

The mapping (a definition stated before the comparison, not a fit). The author's array position 1 is the hidden direction x0, with $g_{00} = \cot^2(6Hx_0)$, and positions 2 to 8 are x1 to x7. This record uses the same coordinates in the same order, so the coordinate map is the identity and no component is permuted. (Until 2026-10-14 the record named the hidden direction x_8 and listed it last; the author's x0 was then mapped to x_8, with the same checks and verdicts.) The author's a4 is a function of the argument Hx_4, this record's of x_4: $a_4(x_4) := a_4^{\text{author}}(Hx_4)$, so by the chain rule $(a_4^{\text{author}})^{(n)}(Hx_4) = H^{-n} a_4^{(n)}(x_4)$ for $n = 0, 1, 2$; the program checks that no other argument of a4 occurs in the author's outputs. H is the same constant on both sides. The mapping is checked on the metric first; only then are the curvature outputs compared.

The conventions and normalisations (each stated and applied, none fitted). The author's rt[g] uses the Christoffel symbols, the Riemann tensor and the Ricci tensor of section 2.2, and $G = \text{Ric} - \frac{1}{2} g\, R$ with both indices down. The Ricci scalar is compared with no factor. The author's Einstein tensor is raised with the author's own (mapped) inverse metric, $G^{\mu}{}_{\nu} = g^{\mu s} G_{s\nu}$, with no other factor. For $k = 1$, section 4.3 gives $P_{(1)} = -4G$ and $L_{(1)} = 2R$, so P1_mixed_up_h_down_j is compared with -4 times the author's $G^{\mu}{}_{\nu}$ and L1 with 2 times the author's Ricci scalar.

The result. A check is PASS only when the difference, reduced exactly with sympy, is 0; NOT-AVAILABLE means that the author's files hold nothing to compare with (it is not a failure):

check	what is compared	verdict
author-json-matches-notebook	the notebook read by the extractor is the notebook of the working tree (sha256)	PASS
author-metric-outputs-agree	Out[235] and Out[245] are identical	PASS
revision-metric-spec-vs-curvature-json	the metric of Revision/SPEC.md section 1 equals metricDiagonal, 64 components	PASS
metric-mapping-author-Out235-vs-revision	the author's metric Out[235], mapped, equals this record's, 64 components	PASS
ricci-scalar-author-Out535-vs-curvature-json	the author's Ricci scalar Out[535] equals ricciScalar	PASS
einstein-mixed-<h>,<j>	each of the 64 components of Out[536], mapped and raised, equals einsteinMixed (64 checks)	PASS

check	what is compared	verdict
`lovelock-P1-vs-minus4-author-EinsteinG`	`P1_mixed_up_h_down_j` equals -4 times the author's $G^{\mu}{}_{\nu}$, 64 components	PASS
`lovelock-L1-vs-2-author-RS`	`L1` equals 2 times the author's Ricci scalar	PASS
`control-mapping-without-chain-rule-is-detected`	the mapping without the factor H^{-n} leaves a nonzero difference	PASS
`control-einstein-without-index-raising-is-detected`	the author's $G_{x_1x_1}$ compared without raising the index leaves a nonzero difference	PASS
`lovelock-k2-P2-and-L2`	$P_{(2)}$ and $L_{(2)}$	NOT-AVAILABLE
`lovelock-k3-P3-and-L3`	$P_{(3)}$ and $L_{(3)}$	NOT-AVAILABLE
`christoffel-components`	the Christoffel symbols	NOT-AVAILABLE
`riemann-components`	the Riemann tensor	NOT-AVAILABLE
`ricci-tensor-components`	the Ricci tensor	NOT-AVAILABLE

In all, the report `author-comparison-report.json` has 78 checks: 73 PASS, 0 FAIL, 5 NOT-AVAILABLE. Two examples of what was compared, as the report gives them: the author's stored Ricci scalar is `6*H^2*(-7 + Derivative[1][a4][H*x4]^2)`, which the mapping turns into $6(a_4')^2 - 42H^2$, the value R of section 6.1 at constant b_4; the author's stored $G_{x_4x_4}$ becomes $-3H^2(7 + (a_4')^2/H^2)$, and raised with $g^{x_4x_4} = -1$ it is $3(a_4')^2 + 21H^2 = G^{x_4}{}_{x_4}$. The controls show that the comparison is not vacuous: without the chain-rule factor the difference of the Ricci scalars is $6(H^2 - 1)(a_4')^2$, and without the index raising the difference of the x_1x_1 components is nonzero; both are detected.

Why five objects are NOT-AVAILABLE. The Christoffel symbols, the Riemann tensor and the Ricci tensor are computed by `rt[gtry]`, but `In[238]` ends with a semicolon, so no value of them is stored there, and no other stored output of their values was taken. For the Lovelock tensors the box strings of every cell of the main notebook and of `Generalized _Kronecker_Delta_4+4.nb` were scanned for the word Lovelock (in any case) and for box strings equal to one of the tokens `P2`, `P3`, `P4`, `LovelockP`, `kd` and `k`δ. In the main notebook 20 cells contain the word; the 4 Output cells among them hold only strings (file names), and none of its cells contains one of the tokens. In the Kronecker-delta notebook 3 cells contain the word, none of them an Output cell, and the cells with the token `k`δ are Input and Text cells. No stored output of $P_{(2)}$, $P_{(3)}$, $L_{(2)}$ or $L_{(3)}$ was found.

Re-run for this document (2026-10-08, into a scratch directory, the committed files untouched): the extractor reproduced `author-curvature-outputs.json` byte for byte (one run, 6 s of wall time including the start of the kernel), and `compare_with_author.py` reproduced `author-comparison-report.json` byte for byte in four runs (two of them timed: 0.85 s and 0.92 s). Later on 2026-10-08 the extractor was changed (it normalises the repository root, and it no longer writes the text of `$Version`, which names the operating system; only the version number 15.0.1 is written) and `author-curvature-outputs.json` was regenerated: the only change of that file is the removed line with the version text (its sha256 is now `f041955db36f67d5ec113ad65542f448af1de6eba159c85ecff73f83761039f0`), the only change of the report is the sha256 it records for that input, and the checks and verdicts are unchanged (78 checks: 73 PASS, 0 FAIL, 5 NOT-AVAILABLE). Re-run of both programs after that change, for this document (one run each into a scratch directory, the committed files untouched, on a loaded machine): both reproduced the

committed files byte for byte (the extractor 19.2 s, the comparison 5.3 s); the report then had the sha256 `6896fd78e327784d0c295e73eef4e45a127e88aa2d6990bf9855ac7b35796265`. These run times were measured while writing this document; no report records them. On 2026-10-14 the comparison was re-run for the metric with b_4 (at $b_4 = 1$) and for the author's coordinates (the identity mapping): the same 78 checks with the same verdicts; the Einstein components are now named `einstein-mixed-x0,x0` to `einstein-mixed-x7,x7`, and the report has the sha256 `1fc3c6fb2a73d7f361846b8971c669079e9f2d96e5094a512310ceab285fd545` (it records the sha256 of `Revision/SPEC.md`, which was reworded the same day; `Revision/gkd_lovelock/comparison/README.md`, section 7). On 2026-10-16, after section 1 of `Revision/SPEC.md` was changed for the history with b_4, the report was regenerated once more: the only change is the sha256 it records for `Revision/SPEC.md`, and the report has the sha256 `2a89ce981d4d2fd8e22875cf402de0f26eb1445a6f4824235ed7afd6f8bd7b55`.

8.3 What the comparison does not establish

- Nothing about $k = 2$ and $k = 3$: the tensors $P_{(2)}$ (Gauss-Bonnet) and $P_{(3)}$ and the scalars $L_{(2)}$ and $L_{(3)}$ of this record are compared with no output of the author, because the files read contain none. Agreement with the author's own Lovelock tensors of order $k = 2$ and $k = 3$ is therefore not established.
- The $k = 1$ comparison carries no information beyond the Einstein-tensor comparison together with this record's own identity $P_{(1)} = -4G$ (`k1_equals_minus_4_einstein`).
- The Christoffel symbols, the Riemann tensor and the Ricci tensor are not compared: no stored output of them was taken.
- The agreement is one between two computations of the same objects for this one metric on the patch $0 < z < \pi/2$; it is not an independent check of conventions beyond those stated above.
- The author's outputs are taken as stored; the notebook was not re-run, and that the stored values come from `In[238]` is the inference stated above.
- The keyword scan finds the Lovelock tensors by name; a value stored under an unrelated name would not be found by it. Only the two notebooks of the author in the repository were read; the main notebook's text says that the Einstein-Lovelock tensors are calculated in other notebooks of the author, and none of those was read (the repository holds none of them).
- Nothing about field equations, sources, solutions or physical interpretation.

8.4 The metric with $b_4(x_4)$

On 2026-10-14 the user changed the primordial metric: the three extra-time components gained the factor $b_4(x_4)$, $g_{x_5x_5} = g_{x_6x_6} = g_{x_7x_7} = -e^{-2a_4} b_4 \sin^{1/3} z$ (section 2.1), and on 2026-10-15 the user ordered that everything be moved to $b_4(x_4)$, b_4 never 1 ($b_4 = 1$ is only the withdrawn metric used before). Every file of `Revision/gkd_lovelock/results/` was recomputed with b_4 on 2026-10-14: the Rust program passes 19 of 19 checks (with the brute force of $k = 2$), the Wolfram check 29 of 29, the sympy checker 49 of 49; each $P_{(k)}$ has 10 nonzero components, the 8 diagonal ones and x_0-x_4, x_4-x_0, proportional to b_4' (section 6.3); the monomial vectors of `lovelock-tensors.json` have 13 exponents. Since 2026-10-16 this document quotes every formula for general b_4 (sections 6.1 to 6.3).

The author's new notebook `Dirac_Einstein-Lovelock_4+4_Nash.nb` stores the Lovelock tensors allLovelock2 and allLovelock3 with b_4, so the tensors of order $k = 2$ and $k = 3$, which were NOT-AVAILABLE in section 8.2, are compared exactly with the author's stored outputs in `Revision/metric_b4` (`Revision/metric_b4/METRIC_CHANGE_PROVENANCE.md`): the Wolfram verifier `metric_b4/wolfram/verify_metric_b4.wls` passes 27 of 27 checks and the independent sympy checker `metric_b4/python/check_metric_b4.py` 21 of 21, and the field-equations record checks that every component of $E_{(1)}$, $E_{(2)}$, $E_{(3)}$ for general b_4 equals that set (`lovelock_equals_metric_b4_set`). These are exact comparisons of the stored outputs of that notebook for this metric; they do not extend what section 8.3 says is not established about the earlier notebook.

9 How the field equations for a_4 and b_4 use the Lovelock tensors

`Revision/field_equations_a4` writes the Einstein-Lovelock equations of `Revision/SPEC.md` section 5 for the author's metric with $b_4(x_4)$,

$$\sum_{k=1}^{3} \alpha_k \, E_{(k)}{}^{\mu}{}_{\nu} + \Lambda \, \delta^{\mu}_{\nu} = \kappa \, T^{\mu}{}_{\nu}, \qquad E_{(k)} = -\frac{P_{(k)}}{2^{k+1}}, \qquad E_{(1)} = G,$$

with $\alpha_1 = 1$ for Einstein gravity ($\alpha_2 = \alpha_3 = 0$). Both of its verifiers recompute the three tensors with their own GKD (Wolfram: as the determinant of the delta matrix; sympy: as the permutation sign) and compare all 64 components of each with the exact monomial lists of `Revision/gkd_lovelock/results/lovelock-tensors.json`, for general b_4:

step	**Wolfram records (`wolfram-a4-report.json`)**	**sympy records (`python-a4-report.json`)**
$P_{(k)}$ equal to this record	`P1_direct_equals_gkd_branch_monomials`, `P2_direct_equals_gkd_branch_monomials`, `P3_direct_equals_gkd_branch_monomials`	`P1_equals_gkd_branch_monomials`, `P2_equals_gkd_branch_monomials`, `P3_equals_gkd_branch_monomials`
$L_{(k)}$ equal to this record	`L1_direct_equals_gkd_branch`, `L2_direct_equals_gkd_branch`, `L3_direct_equals_gkd_branch`	`L1_equals_gkd_branch`, `L2_equals_gkd_branch`, `L3_equals_gkd_branch`
$P_{(1)} = -4G$, traces	`P1_direct_equals_minus_4_Einstein`, `P1_trace_identity`, `P2_trace_identity`, `P3_trace_identity`	`P1_equals_minus_4_Einstein`, `P1_trace_identity`, `P2_trace_identity`, `P3_trace_identity`
structure, $P_{(4)} = 0$, b_4 through u and v	`P1_structure`, `P2_structure`, `P3_structure`, `independent_components`, `P4_vanishes_pigeonhole`, `lovelock_depends_on_b4_through_u_and_v`	`P1_structure`, `P2_structure`, `P3_structure`, `other_components_vanish`

step	Wolfram records (wolfram-a4-report.json)	sympy records (python-a4-report.json)
divergence of $E_{(k)}$; the $E_{(k)}$ of a4-equations.json	E1_divergence_free, E2_divergence_free, E3_divergence_free	E1_divergence_free, E2_divergence_free, E3_divergence_free, json_lovelock_components

What follows from the tensors of this record (taken from those reports, not derived here): the independent left-hand sides are x_0, $x_1 = x_2 = x_3$, x_4, $x_5 = x_6 = x_7$ and the pair x_4-x_0, x_0-x_4. The pair factorises, $\sum_k \alpha_k {E_{(k)}}^{x_4}{}_{x_0} = \cot z\, u\, Q(a_4', u)$ (offdiagonal_factorises), so a source without a ${T^{x_4}}_{x_0}$ component needs $Q = 0$ at every instant at which b_4 changes; in Einstein gravity $Q = \frac{3}{2}H$, which forces $b_4' = 0$ (einstein_offdiagonal_forces_b4_constant). The combination $\sum_k \alpha_k({E^{x_1}}_{x_1} + {E^{x_5}}_{x_5} - 2{E^{x_0}}_{x_0}) = I(a_4', a_4'', u, v)$ vanishes for constant b_4 (the source condition $p_3 + p_t = 2p_0$ of the withdrawn metric) but not in general (algebraic_identity_x1_plus_x5_minus_2x0, algebraic_identity). With the author's weights $\alpha_k = w_k/H^{2k}$, $w = (1, \frac{1}{32}, \frac{1}{1152})$ and $\Lambda = -10$ ($H = 1$), $Q = 0$ on $a_4 = Hx_4$ gives $d\ln b_4/da_4 \in \{0, \frac{4}{3}\}$ (author_weights_branch), and the history $a_4 = Hx_4$, $b_4 = b_4(0)e^{(4/3)Hx_4}$ needs the constant source $\kappa(\rho, p_3, p_t, p_0) = (-\frac{2}{27}, -\frac{2}{9}, -\frac{22}{27}, -\frac{14}{27})$ (author_weights_history_source); the Revision notebook lovelock_gkd.ipynb (section 8.4 of its text) reproduces this source from the tensors of this record. The solutions of these equations and their sources are the subject of the field-theory documents Revision/docs/DIRAC16COMPLEX_FIELD_THEORY.md and Revision/docs/DIRAC16COMPLEX00_FIELD_THEORY.md and of the Kohn-Sham record, not of this one.

10 Verification records

Every check has a name, a verdict and a detail. Counts at the time of writing, as the JSON files give them (the publication test re-reads them):

report	checks	PASS	FAIL
Revision/gkd_lovelock/results/lovelock-report.json	19	19	0
Revision/gkd_lovelock/results/wolfram-gkd-report.json	29	29	0
Revision/gkd_lovelock/results/python-lovelock-report.json	49	49	0
Revision/field_equations_a4/reports/wolfram-a4-report.json	59	59	0

report	checks	PASS	FAIL
Revision/field_equations_a4/reports/python-a4-report.json	65	65	0

The report of the comparison with the author (section 8.2) has a third verdict, NOT-AVAILABLE, for objects of which the author's files hold no stored output; it is not a failure:

report	checks	PASS	FAIL	NOT-AVAILABLE
Revision/gkd_lovelock/comparison/author-comparison-report.json	78	73	0	5

The Rust report stores each check with `"passed": true` instead of a verdict; its 19 checks are counted as PASS. The GKD self-test `Revision/gkd_lovelock/results/gkd-selftest.json` has no list of checks; its nine comparisons (section 3.3) all have 0 mismatches and its verdict is `SUCCESS`. The test file `Revision/tests/test_gkd_lovelock.py` pins the sha256 of the seven result files of the computation and its verifications (`curvature.json`, `lovelock-tensors.json`, `lovelock-components.md`, `lovelock-report.json`, `gkd-selftest.json`, `python-lovelock-report.json`, `wolfram-gkd-report.json`), requires every check to pass, and, in its slow tests, re-runs the Rust program, the sympy checker and the Wolfram check and requires byte-identical outputs.

Re-verification for this document (2026-10-08, on the shared machine of the provenance files; written to a scratch directory, the committed files untouched): `lovelock_gkd lovelock -brute-force-k2` gave 19 of 19 checks in 18.9 s and four files byte-identical to the committed ones; the sympy checker gave 49 of 49 checks in 40.3 s and a report byte-identical to the committed one. These two run times are measurements made while writing this document; they are not recorded in a report.

11 Reproduction

11.1 Commands

From the repository root (Rust 1.87 or newer, Python 3 with sympy and numpy, Wolfram Language with WolframScript). A backslash at the end of a line is the line continuation of a POSIX shell; in PowerShell write such a command on one line:

```
cargo build --release --manifest-path Revision/gkd_lovelock/code/Cargo.toml
Revision/gkd_lovelock/code/target/release/lovelock_gkd gkd-selftest \
    --exhaustive-max 4 --output Revision/gkd_lovelock/results
Revision/gkd_lovelock/code/target/release/lovelock_gkd lovelock \
```

```
    --output Revision/gkd_lovelock/results --brute-force-k2
python Revision/gkd_lovelock/verification/check_lovelock_gkd.py
wolframscript -file Revision/gkd_lovelock/verification/verify_lovelock_gkd.wls \
    Revision/gkd_lovelock/results/wolfram-gkd-report.json
wolframscript -file \
    Revision/gkd_lovelock/notebook_reading/lovelock_extract_nb_inputs.wls
python Revision/gkd_lovelock/notebook_reading/lovelock_digest_nb_inputs.py
wolframscript -file \
    Revision/gkd_lovelock/notebook_reading/lovelock_export_nb_image.wls
wolframscript -file \
    Revision/gkd_lovelock/comparison/extract_author_curvature_outputs.wls
python Revision/gkd_lovelock/comparison/compare_with_author.py
python -m unittest Revision/tests/test_gkd_lovelock.py -v
python scripts/build_provenance_pdf.py Revision/docs/LOVELOCK_GKD.md \
    --developer-layout --specifications Revision/pdf-specifications.json
python -m unittest Revision/tests/test_lovelock_gkd_publication.py -v
```

On Windows the binary is `lovelock_gkd.exe`. Each of the commands that write into `Revision/gkd_lovelock/results` overwrites a committed file with a byte-identical one; to leave the committed files untouched, give another output directory or report path and compare. The Wolfram check builds its exporter in `<$TemporaryDirectory>/revision_gkd_export` (or in `$LOVELOCK_GKD_EXPORT_DIR`); run only one copy at a time with the default directory. The notebook-reading commands need the author's notebook in the repository root and write the uncommitted text `build/lovelock_nb_inputs.txt`. The extractor needs both notebooks of the author in the repository root and the comparison the main one, `Pair_Creation_of_Universes_WaveFunctionOfUniverse-4+4-Einstein-Lovelock-Nash.nb`, whose sha256 it checks; the extractor writes `author-curvature-outputs.json` (or the file named by the environment variable `AUTHOR_CURVATURE_OUTPUTS`), and the comparison writes `author-comparison-report.json` (or the file given with `-output`). The gate `Revision/verify_revision.sh` and its PowerShell twin run them as the steps `gkd-author-extract` and `gkd-author-compare`.

11.2 Expected output

- `gkd-selftest`: nine lines starting with `PASS - GKD vs literal`, then `gkd-selftest: SUCCESS (...)`; the file `gkd-selftest.json` with the table of section 3.3.
- `lovelock`: the 19 check lines starting with `PASS - `, three lines starting with `info - ` (the scalar $L_{(4)}$ and the times of the two unpruned sums), the counter lines of section 5.2, then `check_count=19`, `failed_check_count=0` and `lovelock: SUCCESS (...)`; exit code 0.
- The sympy checker: 49 lines `[PASS] <name>` in six groups, then `total ... s; 49 checks, 0 failed; verdict SUCCESS`; exit code 0.
- The Wolfram check: ten `time_<step>=<seconds>` lines, the 29 lines `check_<name>=true`, then `check_count=29`, `failed_check_count=0`, `time_total=<seconds>` and `report=<path>`; exit code 0 (1 if a check fails, 2 on a load, build or input/output error).
- The notebook reading: `input cells written: 58`, `58 input cells`, `{1372, 435}` and `28229 bytes (117 bytes of text/time chunks removed)`, with both committed outputs reproduced byte for byte.

- The comparison with the author: the extractor prints `cells: 2127`, `inputs parsed: 8/8` and `outputs parsed: 4/4`, then the output dimensions and the keyword-scan counts; the comparison prints `checks: 78; PASS 73, FAIL 0, NOT-AVAILABLE 5`; both exit with code 0 and reproduce their committed outputs byte for byte.

11.3 Run times

From the provenance files (shared 24-core machine, Windows 11):

- The Wolfram check, from part 4.4 of the provenance file `WOLFRAMSCRIPT_PROVENANCE.md` in `Revision/gkd_lovelock/verification/`: `time_total` 68.4 s in the run of 2026-10-08 that regenerated the committed report, 68-135 s over seven runs; the $k = 2$ sum took 26-43 s and the $k = 3$ sum 35-77 s over eight runs; peak memory of the kernel about 486 MB working set. The optional no-skip mode for $k = 3$ took 1637.7 s (part 4.5).
- The notebook reading, from the provenance file `WOLFRAMSCRIPT_PROVENANCE.md` in `Revision/gkd_lovelock/notebook_reading/`: the whole set takes about 10 to 25 seconds, and up to about 40 seconds on a fully loaded machine.
- The GKD self-test with `-exhaustive-max 4`: several minutes (`Revision/tests/test_gkd_lovelock.py`, which runs it only when `LOVELOCK_GKD_SELFTEST=1`).
- The Rust `lovelock` run and the sympy checker: no run time is recorded in a provenance file; see the measurement of section 10 (18.9 s and 40.3 s).
- The comparison with the author, from section 7 of the README of `Revision/gkd_lovelock/comparison/`: the extractor 4.4 to 6.0 s (seven runs), the comparison 0.7 to 0.9 s; see also the measurement of section 8.2.

11.4 The PDF

The PDF build runs the Markdown-to-LaTeX builder twice (with different hash seeds), runs pdflatex three times into each of two fresh directories, requires warning-free logs and byte-identical PDFs, and checks the PDF against its entry `lovelock-gkd` in `Revision/pdf-specifications.json` (the Revision registry; the registry of the earlier stages is not touched).

12 What is established and what is not

12.1 Established

1. GKD equals the author's definition $\mathtt{k}\delta$ for all index lists of equal length (theorem of section 3.2, with its proof); confirmed exhaustively up to length 4 (Rust) and length 3 (Wolfram, sympy) and on pseudo-random samples up to length 9.
2. For the author's metric on $0 < z < \pi/2$, with the conventions of sections 2.2 and 4.2: the 25 nonzero Christoffel symbols, the 156 nonzero $R^{ab}{}_{cd}$, the Ricci and Einstein tensors and

the Lovelock tensors $P_{(k)}$, $A_{(k)}$ and scalars $L_{(k)}$, $k = 1, 2, 3$, exactly as listed in section 6 and in `lovelock-tensors.json`; three programs that share no code for the sums agree on every component.

3. $P_{(1)} = -4G$, $P_{(2)} = -8\mathcal{H}$ (Gauss-Bonnet), $L_{(1)} = 2R$, $L_{(2)} = 4\,GB$, $L_{(3)} = 8(2T_1 + \cdots + T_8)$; the trace identities; $\nabla_h {P_{(k)}}^h{}_j = 0$; the symmetry of $A_{(k)}$; $P_{(4)} = 0$ in eight dimensions.

4. The structure: diagonal except for the pair x_0-x_4, x_4-x_0, which is proportional to b_4'; the diagonal components free of $\sin^{1/3} z$, e^{a_4} and x_0, with $x_1 = x_2 = x_3$ and $x_5 = x_6 = x_7$; b_4 enters only through $u = b_4'/b_4$ and $v = u'$, and the x_4 components contain no a_4''.

5. Agreement with the author's stored outputs of the earlier notebook (section 8.2; the withdrawn metric $b_4 = 1$): under the mapping stated before the comparison, the author's metric `Out[235]`, Ricci scalar `Out[535]` and all 64 components of the Einstein tensor `Out[536]` (raised with the author's own metric) equal those of this record at $b_4 = 1$ exactly, and hence $P_{(1)}$ and $L_{(1)}$ equal -4 times the author's Einstein tensor and 2 times the author's Ricci scalar; of the 78 checks 73 pass, none fails and 5 are NOT-AVAILABLE.

12.2 Not established

1. No solution: this record gives the left-hand sides of the field equations only. Which a_4, which b_4 and which sources solve them is not part of it (section 9).

2. The domain: only the patch $0 < z < \pi/2$, where $\sqrt{|g|} = b_4^{3/2} \cos z$. Nothing is computed at the brane $z = \pi/2$, where $g_{00} = \cot^2 z = 0$ and $\det g = 0$, or on a mirror patch.

3. No comparison for $k = 2$ and $k = 3$: the author's files that were read hold no stored output of $P_{(2)}$, $P_{(3)}$, $L_{(2)}$ or $L_{(3)}$, nor of the Christoffel symbols, the Riemann tensor or the Ricci tensor (section 8.2). Agreement with the author's own Lovelock tensors of order $k = 2$ and $k = 3$ is therefore not established. The notebooks in which, by the main notebook's text, the author calculates the Einstein-Lovelock tensors were not read; the limits of the comparison that was made are listed in section 8.3.

4. For $k = 3$ the tensor $P_{(3)}$ is verified by three independent recomputations, the trace, the divergence and the symmetry, and the scalar $L_{(3)}$ against the classical cubic density; it is not compared with an independent closed formula for the third-order Lovelock tensor. $E_{(k)} = -P_{(k)}/2^{k+1}$ is a normalisation convention, which equals the Einstein tensor for $k = 1$ and the Gauss-Bonnet tensor for $k = 2$.

5. The unpruned sums are exact for $k = 1, 2$ (sympy) and numerical at one point (Rust); for $k = 3$ the skipped terms are confirmed by a random sample (sympy) and by the theorem of section 3.2; the Wolfram no-skip run of the diagonal $k = 3$ components is recorded in its provenance file only, and its report is not committed.

6. Only Wolfram Language 15.0.1 and the toolchains named in the provenance files were used.

7. No physical interpretation is attached to the components here; in particular nothing is said about the sign of an energy or about the stability of a solution.