UM-4 solution

The approximate solution of the coupled field equations of UM-4: gravity and the background, the volume of the extra times, the sources of the fold, the Higgs field in the ad hoc vacuum, the propagation in the bulk, the resonances and the Kohn-Sham states of dirac16complex, the residuals of every field equation, the neutrino flavours, the thermal history, the primordial era, and the CMB, the CNB and the BAO

Underlying theory: Patrick L. Nash Solved and verified by Claude (Anthropic)

Solution of UM-4, 2026-10-25 Repository once-ere/Dirac_claude, folder universe_model_4

Contents

1 The fold: definition and examples **2**

2 How to read this document **13**
2.1 What this document contains . . . 13
2.2 The status labels . . . 14
2.3 Every equation is followed by its symbols . . . 14
2.4 Coordinates, signature and units . . . 15
2.5 The solution at a glance . . . 15

3 Gravity **17**

4 The background **21**
4.1 The bulk metric of an expanding fold and the equations of the fold . . . 28
4.2 The volume of the extra times . . . 32

5 The sources of the fold **36**
5.1 The gravitational waves of the fold . . . 36
5.2 The gravity induced by the fields of the fold . . . 37
5.3 The fold fluid . . . 38

6 The Higgs field in the ad hoc vacuum **41**

7 Propagation in the bulk **45**
7.1 The modes of the fields of the bulk . . . 45
7.2 The propagators from one point of the fold to another . . . 50
7.3 The causal structure . . . 53

8 dirac16complex: the resonances at the fold **55**
8.1 The momentum along the extra times . . . 64

9 The Kohn-Sham states of dirac16complex **65**
9.1 The equations of the gas of particles χ . . . 68
9.2 The method . . . 73
9.3 The solution . . . 73
9.4 A weakly bound gas . . . 75
9.5 The Krein norm and the stability of the gas . . . 75

10 The residuals of every field equation **76**

11 The three neutrino flavours **86**
11.1 The Sun . . . 90
11.2 The envelope of a supernova . . . 91
11.3 The fast instability near the neutrinosphere . . . 92
11.4 The flavour equipartition and the heating behind the shock . . . 94
11.5 The newer global fit, the Dirac phase and the lightest mass . . . 95

12 The thermal history **97**
12.1 The relic density of the particles χ . . . 98
12.2 The portals of χ to the Standard Model . . . 101
12.3 The right-handed neutrinos . . . 107

13 The primordial era **109**

14 The CMB, the CNB and the BAO **112**
14.1 The background and the BAO ruler . . . 118
14.2 The CMB . . . 119
14.3 The BAO . . . 122
14.4 The CNB . . . 122

15 What the solution does not settle **123**

16 What was verified **124**

17 Further symbols **125**

1 The fold: definition and examples

Definition. The *fold* is the seven-dimensional surface of the author's eight-dimensional spacetime on which the cosine of the angle $z = 6Hx_0$ of the hidden direction vanishes:

$$\mathcal{F} = \big\{X = (x_0, x_1, \ldots, x_7) :\ \cos(6Hx_0) = 0\big\}, \qquad \text{in one period } 0 < z < \pi \text{ of the metric: } z = \frac{\pi}{2}, \quad x_0 = \frac{\pi}{12H} \tag{FD-1}$$

Symbols in (FD-1).

- $\mathcal{F}$: The fold: the seven-dimensional surface $\cos(6Hx_0) = 0$ of the eight-dimensional spacetime (FD-1), in one period $0 < z < \pi$ the surface $z = \pi/2$, $x_0 = \pi/(12H)$, where $\sigma = 0$; $|_{\mathcal{F}}$ means restricted to the fold. Status: DERIVED.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$, the fold is $z = \pi/2$. Unit: dimensionless. Status: AUTHOR.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

- Λ: The eight-dimensional cosmological constant of UM-4, $\Lambda = -21H^2$: the value for which the static block is a vacuum of Einstein gravity (its Einstein tensor is $21H^2$ in every direction). The author writes the pure number Λ_{N} in $0 = -\Lambda_{\mathrm{N}} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$, so that $\Lambda = -\Lambda_{\mathrm{N}} H^2$. Unit: GeV^2. Value: $-21H^2$; the author's original $-10H^2$; UM-3: $-7H^2$. Status: AUTHOR (the term), DERIVED (the value).
- Λ_{N}: The author's pure number in the vacuum equations of his notebook, $\Lambda = -\Lambda_{\mathrm{N}} H^2$: his original value 10; 21 in UM-4 (Einstein gravity), 7 in UM-3. Unit: dimensionless. Value: 21 (the author's original 10). Status: AUTHOR.
- $w_{(p)}$: The Lovelock weights of the author's Einstein-Lovelock action, $w_{(1)} = 1$ and $w_{(2)}$, $w_{(3)}$ of the Gauss-Bonnet and the cubic densities. UM-4 sets $w_{(2)} = w_{(3)} = 0$ (Einstein gravity): the only weights for which gravitons propagate on the light cone on every background, not only around the static block. The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) keep the gravitons on the light cone only around the static block (with $\Lambda = -161H^2/16$); UM-3 used $(1, 1/20, 1/360)$. Unit: dimensionless. Value: $(1, 0, 0)$ in UM-4; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), DERIVED (the values that the order of 2026-10-25 selects).

$X = \{x_0, \ldots, x_7\}$ are the author's coordinates: x_0 the hidden space-like direction, x_1, x_2, x_3 ordinary space, x_4 the time, x_5, x_6, x_7 the three extra times; H is the author's constant (an inverse length). The metric of UM-4 depends on x_0 only through $\cot^2 z$ and $(\sin^2 z)^{1/6}$, so it repeats with the period $\pi/(6H)$ of x_0, and every period $0 < z < \pi$ contains exactly one surface of $\mathcal{F}$, at $z = \pi/2$: $\mathcal{F}$ is the union of these copies, one per period (example 2), and "the fold" without a qualifier is the copy in the period that contains our world. A point of the fold is labelled by the seven coordinates $x_1, \ldots, x_7$, and the metric restricted to the fold ($\mathrm{d}x_0 = 0$, $z = \pi/2$) is

$$g_{MN}\,\mathrm{d}X^M \mathrm{d}X^N\big|_{\mathcal{F}} = \mathrm{e}^{2a_4}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{d}x_4^2 - \mathrm{e}^{-2a_4} b_4^2 \big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big) \tag{FD-2}$$

Symbols in (FD-2).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M \mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- $\mathcal{F}$: The fold: the seven-dimensional surface $\cos(6Hx_0) = 0$ of the eight-dimensional spacetime (FD-1), in one period $0 < z < \pi$ the surface $z = \pi/2$, $x_0 = \pi/(12H)$, where $\sigma = 0$; $|_{\mathcal{F}}$ means restricted to the fold. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.

- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.

with three space-like and four time-like directions; a_4 and b_4 are the author's two functions of the time x_4. Notation: ln is the natural logarithm, e Euler's number, $\exp(y) = \mathrm{e}^y$, sin, cos and cot = cos / sin the trigonometric functions, sinh, cosh, tanh = sinh / cosh and coth = cosh / sinh the hyperbolic functions, $|y|$ the absolute value, max and min the larger and the smaller of their arguments; det, tr and ^{T} are the determinant, the trace and the transpose of a square matrix, and $\mathbf{1}_4$, $\mathbf{1}_8$ and $\mathbf{1}_{16}$ the identity matrices of 4, 8 and 16 rows; $f(0^+)$ and $f(0^-)$ are the limits of $f(\sigma)$ as σ approaches 0 from above and from below.

Why this surface is called a fold. Measured by the proper distance along the hidden direction,

$$\sigma = \mathrm{sgn}\Big(\frac{\pi}{2} - z\Big)\frac{\ln\sin z}{6H}, \qquad \mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0, \qquad \sin z = \mathrm{e}^{-6H|\sigma|} \tag{FD-3}$$

Symbols in (FD-3).

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$, the fold is $z = \pi/2$. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.

(the fold is $\sigma = 0$; the half-period $0 < z < \pi/2$ is $\sigma < 0$, the half-period $\pi/2 < z < \pi$ is $\sigma > 0$; the two ends of the period, the tips $z \to 0$ and $z \to \pi$, lie at $\sigma \to -\infty$ and $\sigma \to +\infty$), the metric of UM-4 is

$$g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N = \mathrm{d}\sigma^2 + \mathrm{e}^{-2H|\sigma|}\Big[\mathrm{e}^{2a_4}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{d}x_4^2 - \mathrm{e}^{-2a_4}b_4^2\big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big)\Big] \tag{FD-4}$$

Symbols in (FD-4).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.

It depends on σ only through $|\sigma|$. The function $\mathcal{W}(\sigma) = \mathrm{e}^{-2H|\sigma|}$, which multiplies the squared steps of all seven directions of the fold (ordinary space, the time and the extra times), is the warp factor of the metric (its square root $\mathrm{e}^{-H|\sigma|}$ multiplies the lengths). The time x_4 is warped like the other directions (the author's order of 2026-10-25, 13:15 PDT; in UM-3 it was not), so that every surface $\sigma = \mathrm{const}$ has the light cone of the fold. Four properties make the surface $\sigma = 0$ a fold, and together they single it out within the period:

1. *It is a mirror.* The reflection $\sigma \to -\sigma$ (that is $z \to \pi - z$, or $x_0 \to \pi/(6H) - x_0$) maps the metric to itself, and the fold is the only surface of the period that the reflection leaves in place: the two halves of the period are mirror images of each other, joined along the fold, as the two halves of a folded sheet of paper are joined along the crease.
2. *The warp factor is largest on it.* The warp factor $(\sin^2 z)^{1/6} = \mathrm{e}^{-2H|\sigma|}$ equals 1 on the fold and decreases on both sides.
3. *The metric has a kink on it.* The metric is continuous across the fold, but its first derivative along σ jumps. The extrinsic curvature of the surfaces $\sigma = \mathrm{const}$ changes sign across the fold,

$$K^a{}_b = \frac{1}{2}\,g^{ac}\,\partial_\sigma g_{cb} = -H\,\mathrm{sgn}(\sigma)\,\delta^a{}_b \quad \text{(all seven directions of the fold)} \tag{FD-5}$$

Symbols in (FD-5).

- $K^a{}_b$: The extrinsic curvature of a surface of constant σ of the hidden direction, $K^a{}_b = \frac{1}{2}g^{ac}\partial_\sigma g_{cb}$ for a, b along the surface (FD-5): $-H\,\mathrm{sgn}(\sigma)$ along all seven directions of the fold for the author's form of the metric of UM-4; $K = K^c{}_c$ its trace. Unit: GeV (1/length). Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- c: A frame index (as a). Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+, +, +, +, -, -, -, -$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.

- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- η_{ab}: The flat metric of the frame, $\eta = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$ (eta4488 of 4+4.nb); η^{ab} is its inverse (the same matrix). Unit: dimensionless. Status: AUTHOR.
- $\hat{g}_{ab}$: The metric of the seven-dimensional fold (FD-2) in its coordinates $x_1, \ldots, x_7$ (or in its frame), $\hat{g}^{ab}$ its inverse; $\hat{h}_{\mu\nu}$ is its part along the four observable directions. Unit: dimensionless. Status: AUTHOR.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- x: The coordinates of the fold: x_1, x_2, x_3 of ordinary space, x_4 the time, x_5, x_6, x_7 the extra times; a field of the fold is a function of the four observable coordinates $x = (x_1, x_2, x_3, x_4)$, and $\mathrm{d}^4x = \mathrm{d}x_1\,\mathrm{d}x_2\,\mathrm{d}x_3\,\mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.

(a, b frame indices along the fold); it jumps by $-2H$ in every direction of the fold. The field equations of gravity of UM-4, $\mathcal{E}_{(1)}{}^M{}_N + \Lambda\delta^M_N = \kappa\mathcal{T}^M{}_N$ (Einstein gravity: the author's Einstein-Lovelock equations with the weights of the Gauss-Bonnet and cubic terms set to zero; $\mathcal{E}_{(1)}$ the Einstein tensor, $\Lambda = -21H^2$ the cosmological constant, $\kappa = 8\pi G_8$ the gravitational coupling of eight dimensions, $\mathcal{T}^M{}_N$ the energy-momentum tensor, δ^M_N the Kronecker delta), are linear in the second σ-derivatives of the metric, so the jump puts a Dirac delta function $\delta_\mathrm{D}(\sigma)$ into them: the metric solves them only if the fold carries a thin source $\mathcal{S}^a{}_b$, an energy-momentum tensor concentrated on $\sigma = 0$, given by the integral of the equations across the fold (the junction condition of Israel),

$$\kappa\,\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma\,\mathcal{E}_{(1)}{}^a{}_b = -\Big(\big(K^a{}_b\big|_{0^+} - K^a{}_b\big|_{0^-}\big) - \delta^a{}_b\big(K\big|_{0^+} - K\big|_{0^-}\big)\Big) \tag{FD-11}$$

Symbols in (FD-11).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma\,\mathcal{E}_{(1)}{}^a{}_b$ (the junction condition of Israel); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -12H\delta^a{}_b$ for every a_4, b_4 of the author's form. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\mathcal{E}_{(p)}{}^M{}_N$: The Lovelock tensor of order p, $\mathcal{E}_{(p)}{}^M{}_N = -2^{-(p+1)}\,\delta^{M P_1 \cdots P_{2p}}_{N Q_1 \cdots Q_{2p}}\,R^{Q_1 Q_2}{}_{P_1 P_2} \cdots R^{Q_{2p-1} Q_{2p}}{}_{P_{2p-1} P_{2p}}$ (with both indices down it is $1/\sqrt{|g|}$ times the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to g^{MN}; $\mathcal{E}_{(1)}$ the Einstein tensor,

$\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor, $\mathcal{E}_{(3)}$ the cubic Lovelock tensor); with frame indices its frame components. Unit: $1/\text{length}^{2p}$. Status: STANDARD.

- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number (UM-4 keeps only $p = 1$); in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- ${K^a}_b$: The extrinsic curvature of a surface of constant σ of the hidden direction, ${K^a}_b = \frac{1}{2} g^{ac} \partial_\sigma g_{cb}$ for a, b along the surface (FD-5): $-H \operatorname{sgn}(\sigma)$ along all seven directions of the fold for the author's form of the metric of UM-4; $K = {K^c}_c$ its trace. Unit: GeV (1/length). Status: DERIVED.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- G_8: The eight-dimensional Newton constant ($\kappa = 8\pi G_8$). Unit: GeV^{-6}. Status: AUTHOR.
- ${R^{MN}}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor ${R^M}_{NPQ}$ (in the convention in which the commutator of two covariant derivatives, acting on a vector, is the Riemann tensor, with its last two indices those of the two derivatives, contracted with the vector), with two indices the Ricci tensor $R_{MN} = {R^P}_{MPN}$, without indices the scalar curvature $R = g^{MN} R_{MN}$; with frame indices ${R^{(a)(b)}}_{(c)(d)}$ its frame components, ${R^{(a)(b)}}_{(a)(b)}$ (no sum) the sectional curvature of the frame plane (a, b). Unit: $1/\text{length}^2$. Status: STANDARD.
- $\mathcal{L}_{(p)}$: The Lovelock density of order p, $\mathcal{L}_{(p)} = 2^{-p}\, \delta^{P_1 \cdots P_{2p}}_{Q_1 \cdots Q_{2p}}\, {R^{Q_1 Q_2}}_{P_1 P_2} \cdots {R^{Q_{2p-1} Q_{2p}}}_{P_{2p-1} P_{2p}}$, a scalar built from p curvature tensors: $\mathcal{L}_{(1)} = R$ (Einstein-Hilbert), $\mathcal{L}_{(2)}$ the Gauss-Bonnet density, $\mathcal{L}_{(3)}$ the cubic Lovelock density. Unit: $1/\text{length}^{2p}$. Status: STANDARD.

($K = {K^c}_c$ the trace of the extrinsic curvature), with ${\mathcal{S}^4}_4 = -$(its energy density), ${\mathcal{S}^1}_1 = {\mathcal{S}^2}_2 = {\mathcal{S}^3}_3$ its pressure along ordinary space and ${\mathcal{S}^5}_5 = {\mathcal{S}^6}_6 = {\mathcal{S}^7}_7$ its pressure along the extra times; $\mathcal{S}$ is $\mathcal{T}$ integrated across the fold, an energy per unit volume of the six other directions of the fold (unit GeV^7). For the static block (a_4 and b_4 constant) and, to the first order in the curvature of the fold divided by H^2, for every fold

$$\begin{aligned} &\text{static block:} \quad \kappa {\mathcal{S}^a}_b = -12H\, {\delta^a}_b \quad \text{(a pure tension)}, \qquad \kappa\lambda_{\mathrm{F}} = 12H, \\ &\text{any fold:} \quad \kappa {\mathcal{S}^a}_b = -12H\, {\delta^a}_b + \frac{2}{5H}\, {\hat{G}^a}_b + \ldots \end{aligned} \tag{FD-12}$$

Symbols in (FD-12).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- ${\mathcal{S}^a}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa {\mathcal{S}^a}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma\, {\mathcal{E}_{(1)}}^a{}_b$ (the junction condition of Israel); ${\mathcal{S}^4}_4$ is minus its energy density, ${\mathcal{S}^1}_1$ and ${\mathcal{S}^5}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa {\mathcal{S}^a}_b = -12H {\delta^a}_b$ for every a_4, b_4 of the author's form. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \operatorname{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- λ_{F}: The tension that the static block requires of the fold, $\kappa\lambda_{\mathrm{F}} = 12H$: the total vacuum energy of the seven-dimensional fold, made of the tension of the fold fluid and the vacuum energy of the fields of the fold (both have the pressure minus their energy density along ordinary space and along the extra times), per

unit volume of the six other directions of the fold (eight-dimensional normalisation); in the four-dimensional normalisation it is $\lambda_F V_*$, and the excess of the total over $\lambda_F V_*$ is the dark energy ρ_Λ. Unit: GeV^7. Value: $12H/\kappa$; $\lambda_F V_* = 30H^2 M_{red}^2 = 1.1 \times 10^{45}$ GeV^4 for $H = 2534$ GeV. Status: DERIVED.

- $\hat{G}^\mu{}_\nu$: The Einstein tensor of the metric $\hat{g}_{ab}$ of the fold (FD-2), with frame indices along its seven directions; with four-dimensional indices, $\hat{G}^\mu{}_\nu$, its components along the four observable directions, which for frozen extra times are those of the Einstein tensor of the four-dimensional metric $\hat{h}_{\mu\nu}$. Unit: 1/length^2. Status: STANDARD.
- M_{red}: The reduced Planck mass, $M_{red} = (8\pi G_N)^{-1/2}$. Unit: GeV. Value: 2.435×10^{18} GeV. Status: STANDARD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_t(e^{-a_4}b_4)^3$ (with $V_t = \int dx_5\, dx_6\, dx_7$ the coordinate volume of the extra times) at a time when $e^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- $\hat{h}_{\mu\nu}$: The four-dimensional metric of the fold, $\hat{h}_{\mu\nu} = \mathrm{diag}(e^{2a_4}, e^{2a_4}, e^{2a_4}, -1)$ in the coordinates $x_1, \ldots, x_4$ (the induced metric at $\sigma = 0$ without the extra times); $\hat{h}^{\mu\nu}$ its inverse, $\hat{h}$ its determinant, $\sqrt{-\hat{h}} = e^{3a_4}$; $\hat{h}^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- ρ_Λ: The vacuum energy density of the fold above its static tension, the dark energy of UM-4: $\rho_\Lambda = \rho_{ff} + \rho_{vac} - \lambda_F V_*$ (four-dimensional normalisation). Unit: GeV^4. Value: $0.685\,\rho_c^{(0)} = 2.5 \times 10^{-47}$ GeV^4 (Planck 2018). Status: DERIVED.
- G_N: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_N = 1/M_{Pl}^2$ with the Planck mass $M_{Pl} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- V_t: The coordinate volume of the (compact) extra times, $V_t = \int dx_5\, dx_6\, dx_7$. Unit: length^3. Status: ASSUMED.
- ρ_c: The critical density, the energy density of a spatially flat fold at the expansion rate H_4; $\rho_c^{(0)}$ its value today. Unit: GeV^4. Value: $\rho_c^{(0)} = 1.0537 \times 10^{-5}\, h_{100}^2$ GeV/cm^3. Status: STANDARD.
- ρ_{ff}: The energy density of the fold fluid. Unit: GeV^4. Value: $\lambda_F V_* + \rho_\Lambda - \rho_{vac}$: together with the vacuum energy ρ_{vac} of the fields of the fold it makes up the tension $\lambda_F V_*$ that the static block requires plus the dark energy ρ_Λ. Status: ASSUMED (the fluid), DERIVED (the value).
- ρ_{vac}: The vacuum energy of the fields of the fold: the minimum of their potential (for the Standard Model the minimum of the Higgs potential with its loop corrections), a tension of the seven-dimensional fold. Unit: GeV^4. Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\, h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- M_{Pl}: The Planck mass, $M_{Pl} = G_N^{-1/2}$. Unit: GeV. Value: 1.221×10^{19} GeV. Status: STANDARD.
- h_{100}: The present expansion rate of the fold in units of 100 km/s/Mpc. Unit: dimensionless. Value: 0.674 (Planck 2018). Status: STANDARD.

($\hat{G}^a{}_b$ the Einstein tensor of the metric FD-2 of the fold): the tension plus matter that obeys the Einstein equations of seven dimensions, $\hat{G}^a{}_b = \kappa_7 \mathtt{t}^a{}_b$ with $\kappa_7 = 5\kappa H/2$ ($\mathtt{t}^a{}_b$ the matter of the fold), which for frozen extra times are the Friedmann equations of general relativity. The fold is therefore not only a place but a physical object: a thin layer that carries energy and momentum (a brane). 4. *The author's coordinate x_0 degenerates on it.* The metric component $g_{x_0x_0} = \cot^2(6Hx_0)$ vanishes on the fold and nowhere else in the period; near the fold the proper distance grows with the square of the coordinate distance,

$$|\sigma| = 3H\Big(x_0 - \frac{\pi}{12H}\Big)^2\Big[1 + 6H^2\Big(x_0 - \frac{\pi}{12H}\Big)^2 + \ldots\Big] \qquad \text{(FD-6)}$$

Symbols in (FD-6).

- σ: The proper coordinate of the hidden direction: $d\sigma = |\cot z|\, dx_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $e^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_F = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.

- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

so the two coordinate values on either side of $x_0 = \pi/(12H)$ at the same coordinate distance lie at the same proper distance from the fold: the coordinate line of x_0 is folded onto the line of proper distances (a fold map, example 5).

What lives on the fold. In UM-4 the Higgs doublet (with the author's ad hoc vacuum, a time-dependent choice of its vacuum value), three generations of quarks and leptons, three right-handed neutrinos and the fold fluid (its tension and the pressure along the extra times that keeps their volume fixed) live on the fold: they are fields at $\sigma = 0$, functions of $x_1, \dots, x_4$ (no field depends on the extra times). The gauge fields of $SU(3)_C \times SU(2)_L \times U(1)_Y$ live in the bulk; their massless modes are bound to the fold, so that the photon and the gluons of the four-dimensional world are waves of the bulk that the fold holds, and photons that leave the fold travel through the bulk at the speed of light. dirac16complex (the author's spinor field of eight dimensions) and gravity live in the bulk $\sigma \neq 0$ as well as on the fold; the states χ of dirac16complex that the fold holds (in UM-4 long-lived resonances) are the part of it near the fold, and the portals (couplings between dirac16complex and the fields of the fold) act at $\sigma = 0$. Integrated over the extra times, the fold is the four-dimensional universe that we observe, with the metric $\mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in $x_1, \dots, x_4$. In UM-4 files "the fold" means this surface; the examples below use the word for other surfaces with the same properties. The word is not used in its other senses: a set of equal energy levels is called degenerate, and a growth by the factor e is called an e-folding.

Examples of folds that are meaningful for UM-4.

1. *Our fold.* The fold $z = \pi/2$ of the period that contains our world (FD-1): the seven-dimensional world-volume of the quarks, leptons and the Higgs field, and after the integration over the extra times the four-dimensional universe that we observe.
2. *The copies of the fold.* Because the metric repeats with the period $\pi/(6H)$, every surface $x_0 = \pi/(12H) +$ (an integer multiple of $\pi/(6H)$) is a copy of the fold, one per period. Neighbouring copies are separated by a tip (z a multiple of π), where the warp factor $(\sin^2 z)^{1/6}$ vanishes and $g_{x_0x_0} = \cot^2 z$ diverges, and which lies at an infinite proper distance from both: in proper distance every copy has its own complete line $-\infty < \sigma < \infty$. A compact direction of length L would give every field a tower of particles with the masses $2\pi n/L$, $n = 1, 2, \dots$ (a Kaluza-Klein tower); because the copies are infinitely far apart, the period $\pi/(6H)$ of x_0 produces no such tower with the spacing $12H$: every field of the bulk has instead a continuum of modes that starts at zero mass, and the fold holds its massless mode.
3. *A folded sheet of paper.* The crease of a folded sheet is a fold in this sense: the two halves are mirror images (property 1) joined along the crease, the sheet is continuous there and its direction changes abruptly (property 3). In UM-4 the direction across the crease is the hidden direction σ, the seven directions along the crease are $x_1, \dots, x_7$, and the crease is the fold.
4. *A folded function.* The warp factor $\mathrm{e}^{-2H|\sigma|}$ is the simplest folded function: continuous, mirror symmetric, largest at $\sigma = 0$, with a jump of its derivative,

$$\frac{\mathrm{d}}{\mathrm{d}\sigma}\mathrm{e}^{-2H|\sigma|} = -2H\,\mathrm{sgn}(\sigma)\,\mathrm{e}^{-2H|\sigma|}, \qquad \frac{\mathrm{d}^2}{\mathrm{d}\sigma^2}\mathrm{e}^{-2H|\sigma|} = 4H^2\,\mathrm{e}^{-2H|\sigma|} - 4H\,\delta_{\mathrm{D}}(\sigma) \tag{FD-7}$$

Symbols in (FD-7).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.

- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- δ_{D}: The Dirac delta function of σ: zero for $\sigma \neq 0$, with the integral 1 over any range that contains $\sigma = 0$; the integral of $\delta_{\mathrm{D}}(\sigma)$ times a smooth function of σ is the value of that function at $\sigma = 0$. Unit: 1/length (for $\delta_{\mathrm{D}}(\sigma)$). Status: STANDARD.

whose delta function is the origin of the thin source of property 3. Every metric function of FD-4 depends on σ through $|\sigma|$ and is folded at $\sigma = 0$ in this sense. 5. *A fold map.* In the mathematics of smooth maps the fold (Whitney's fold singularity) is the map $u \mapsto u^2$ of a line onto a half-line: it folds the line at $u = 0$, two points $\pm u$ land on the same point, and the derivative of the map vanishes on the fold. Near the fold the map from the author's coordinate x_0 to the distance $|\sigma|$ from the fold is of this kind (FD-6), and $g_{x_0x_0} = \cot^2 z$, the square of the derivative $\mathrm{d}\sigma/\mathrm{d}x_0$, vanishes there. 6. *The brane of Randall and Sundrum.* The five-dimensional metric

$$\mathrm{d}y_{\mathrm{RS}}^2 + \mathrm{e}^{-2k_{\mathrm{RS}}|y_{\mathrm{RS}}|}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2 - \mathrm{d}x_4^2\big), \qquad \lambda_{\mathrm{RS}} = \frac{6k_{\mathrm{RS}}}{\kappa_5}, \qquad \Lambda_{\mathrm{RS}} = -6k_{\mathrm{RS}}^2 \tag{FD-8}$$

Symbols in (FD-8).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- y_{RS}: The proper coordinate across the brane of the five-dimensional model of Randall and Sundrum (FD-8, an example, not UM-4). Unit: length. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- k_{RS}: The inverse curvature length of the bulk of the model of Randall and Sundrum: the warp factor of the brane is $\mathrm{e}^{-2k_{\mathrm{RS}}|y_{\mathrm{RS}}|}$ (FD-8, an example, not UM-4). Unit: GeV (1/length). Status: STANDARD.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- λ_{RS}: The tension (energy per unit volume of ordinary space) of the brane of the model of Randall and Sundrum, $\lambda_{\mathrm{RS}} = 6k_{\mathrm{RS}}/\kappa_5$ (FD-8, an example, not UM-4). Unit: GeV^4. Status: STANDARD.
- κ_5: The gravitational coupling of five dimensions in the model of Randall and Sundrum, 8π times the Newton constant of five dimensions (FD-8, an example, not UM-4). Unit: GeV^{-3}. Status: STANDARD.
- Λ_{RS}: The cosmological constant of the bulk of the model of Randall and Sundrum, $\Lambda_{\mathrm{RS}} = -6k_{\mathrm{RS}}^2$, the vacuum energy density $\Lambda_{\mathrm{RS}}/\kappa_5$ (FD-8, an example, not UM-4). Unit: GeV^2. Status: STANDARD.

has a fold at $y_{\mathrm{RS}} = 0$ with the properties 1 to 3 (its coordinate y_{RS} is already the proper distance, so property 4 does not arise); Einstein's equations require there a brane of the tension λ_{RS} (Israel's junction condition) and in the bulk the cosmological constant Λ_{RS} (the vacuum energy $\Lambda_{\mathrm{RS}}/\kappa_5$), and the bulk is otherwise empty. The fold of UM-4 is of the same kind, with every direction of the fold warped by the same factor as in the model of Randall and Sundrum: its static block is a space of constant curvature $-H^2$ in all eight directions, a vacuum of Einstein gravity with $\Lambda = -21H^2$, with a fold of pure tension (FD-12), and the graviton is bound to the fold as there. It differs in that the fold has seven dimensions, three of them extra times, in that the space is not cut at the brane (Randall and Sundrum identify the two sides of their brane, one physical side; the fold of UM-4 has two distinct sides, example 7), and in that the gauge fields live in the bulk as well. 7. *A particle held by a thin layer.* A particle held by an attractive thin layer has a wave function that falls off exponentially with the distance on both sides, and the layer makes its derivative jump (for the first-order Dirac equation, the wave function itself). In UM-4 the states χ of dirac16complex near the fold are of this kind only near the fold: at rest and without their self-interaction (the field is $\Psi = \mathrm{e}^{(7/2)H|\sigma|}\chi(\sigma)\mathrm{e}^{-\mathrm{i}\omega x_4}$)

$$\begin{aligned} &\chi(\sigma) = \mathrm{e}^{-q_{\mathrm{b}}|\sigma|}\chi(0^+) \ \ (0 < \sigma \ll \sigma_{\mathrm{t}}), \qquad \chi(\sigma) = \mathrm{e}^{-q_{\mathrm{b}}|\sigma|}\chi(0^-) \ \ (0 < -\sigma \ll \sigma_{\mathrm{t}}), \\ &\chi(0^+) = \exp\Big(\gamma^{(0)}\big(\mu_{\mathrm{L}}P_{\mathrm{L}} + \mu_{\mathrm{R}}P_{\mathrm{R}}\big)\Big)\chi(0^-), \qquad \omega_0 = \omega_{\mathrm{r}} - \frac{\mathrm{i}}{2}\,\Gamma_\chi, \qquad \omega_{\mathrm{r}} \to \omega_{\mathrm{b}} = \sqrt{m_\Psi^2 - q_{\mathrm{b}}^2} \end{aligned} \tag{FD-9}$$

Symbols in (FD-9).

- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{\frac{7}{2}H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of the level of dirac16complex

at the fold (a long-lived resonance) and $\bar{\chi}$ its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- q_b: The inverse width of a state of dirac16complex at the fold: inside the turning point its profile falls as $\mathrm{e}^{-q_\mathrm{b}|\sigma|}$. Unit: GeV. Status: DERIVED.
- σ_t: The turning point of a state of dirac16complex of the frequency ω: the depth at which the warped frequency equals the bulk mass, $|\omega|\mathrm{e}^{H\sigma_\mathrm{t}} = m_\Psi$; beyond it every solution oscillates. Unit: length. Status: DERIVED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- μ_X, μ_L, μ_R: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_\mathrm{L}P_\mathrm{L} + \mu_\mathrm{R}P_\mathrm{R})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- P_R: The author's right-handed chiral projector, $P_\mathrm{R} = (\mathbf{1} + \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- ω_0: The energy at rest (zero momentum along ordinary space and along the extra times) of a state of dirac16complex at the fold; complex for a resonance, $\omega_0 = \omega_\mathrm{r} - \frac{\mathrm{i}}{2}\Gamma_\chi$. Unit: GeV. Status: DERIVED.
- ω_r: The mass of the particle χ: the real part of the energy at rest of the resonance of dirac16complex at the fold, $\omega_\mathrm{r} = \omega_\mathrm{b}(1 - H/(2q_\mathrm{b})) + \ldots$. Unit: GeV. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- Γ_χ: The width of the particle χ at rest: the rate of its decay into the waves of dirac16complex in the bulk. Unit: GeV. Value: below $1/(11.6\,t_0) = 1.3 \times 10^{-43}$ GeV for dark matter (a lifetime above 160 Gyr). Status: DERIVED.
- ω_b: The energy at zero momentum of the state of dirac16complex at the fold in the limit $H \to 0$ (the bound state of UM-3): the particle χ of UM-4 has the mass $\omega_\mathrm{r} = \omega_\mathrm{b}(1 - H/(2q_\mathrm{b})) + \ldots$, equal to ω_b up to the relative correction $H/(2q_\mathrm{b})$. Unit: GeV. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- Γ_8: The chirality matrix of eight dimensions, $\Gamma_8 = T_0T_1\cdots T_7$ (the author's T16^A[8]) $= \mathrm{diag}(-\mathbf{1}_8, \mathbf{1}_8)$; $\Gamma_8^2 = \mathbf{1}$; it anticommutes with every T_a. Unit: dimensionless. Status: AUTHOR.
- Ψ, $\bar{\Psi}$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar{\Psi} = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- $\mathbf{1}$, $\mathbf{1}_4$, $\mathbf{1}_8$, $\mathbf{1}_{16}$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_4$, $\mathbf{1}_8$ and $\mathbf{1}_{16}$ are the identity matrices of 4, 8 and 16 rows. Unit: dimensionless. Status: STANDARD.
- $\mathbf{e}_a{}^M$, $\mathbf{e}^a{}_M$: The vielbein (frame field) of the metric: $g_{MN} = \eta_{ab}\,\mathbf{e}^a{}_M\mathbf{e}^b{}_N$; $\mathbf{e}_a{}^M$ is its inverse; $\mathbf{e}_{bN} = \eta_{bc}\mathbf{e}^c{}_N$; $\mathbf{e}^{(a)}$ is the frame one-form $\mathbf{e}^a{}_M\,\mathrm{d}X^M$. Unit: dimensionless. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- $\vec{k}$: The (comoving) momentum along ordinary space, $\vec{k}^2$ its square. Unit: GeV. Status: STANDARD.
- $\vec{x}$: The position in ordinary space, $\vec{x} = (x_1, x_2, x_3)$. Unit: length. Status: AUTHOR.
- t_0: The age of the universe. Unit: time. Value: 13.8 Gyr. Status: STANDARD.
- σ_{16}: The matrix of the Dirac adjoint, $\sigma_{16} = T_0T_1T_2T_3$, real and symmetric; $\sigma_{16}T_a$ is real and antisymmetric. Unit: dimensionless. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.

where μ_L, μ_R are the strengths of the fold for the left- and right-handed halves of dirac16complex (set by the fields of the fold), $q_\mathrm{b} = -m_\Psi(\mu_\mathrm{L} + \mu_\mathrm{R})\tanh\varsigma/(2\varsigma)$, $\varsigma = \sqrt{\mu_\mathrm{L}\mu_\mathrm{R}}$ ($q_\mathrm{b} = -m_\Psi\mu_\mathrm{L}/2$ for $\mu_\mathrm{R} = 0$), and ω_r tends to the energy ω_b of the bound state of UM-3 when $H/q_\mathrm{b} \to 0$. Because the time is warped, the energy ω of the field enters its equation multiplied by $\mathrm{e}^{H|\sigma|}$: beyond the turning point σ_t ($\omega_\mathrm{r}\mathrm{e}^{H\sigma_\mathrm{t}} = m_\Psi$) the profile oscillates as a wave that leaves the fold, so the state is a resonance of the width $\Gamma_\chi = (2/\mathrm{e})(q_\mathrm{b}^2/m_\Psi)\,\mathrm{e}^{-2F(v_\mathrm{b})m_\Psi/H}$ (to the leading order in H/q_b),

with $F(v)=\ln((1+\sqrt{1-v^2})/v)-\sqrt{1-v^2}$ and $v_\mathrm{b}=\omega_\mathrm{b}/m_\Psi$ (the tunnelling through the region between the fold and σ_t, where the profile decays at the local rate $\sqrt{m_\Psi^2-\omega_\mathrm{r}^2\mathrm{e}^{2H|\sigma|}}$, which falls to 0 at σ_t). The profile is not a function of $|\sigma|$: it jumps at the fold, $\chi(0^+)\neq\chi(0^-)$. 8. *The fold as a mirror for dirac16complex.* The reflection across the fold acts on the profiles $\chi(\sigma)$ in two ways,

$$(\mathsf{R}_1\chi)(\sigma)=T_0T_1T_2T_3\,\chi(-\sigma),\qquad (\mathsf{R}_2\chi)(\sigma)=T_0T_5T_6T_7\,\chi(-\sigma) \tag{FD-10}$$

Symbols in (FD-10).

- R_1: The reflection across the fold acting on a profile of dirac16complex together with the reflection of ordinary space, $(\mathsf{R}_1\chi)(\sigma)=T_0T_1T_2T_3\,\chi(-\sigma)$ (FD-10): a symmetry of dirac16complex, of gravity and of the mean field of the gas of the particles χ; the states at the fold of Krein charge -1 are even, those of Krein charge $+1$ odd. Unit: dimensionless. Status: DERIVED.
- R_2: The reflection across the fold acting on a profile of dirac16complex together with the reflection of the extra times, $(\mathsf{R}_2\chi)(\sigma)=T_0T_5T_6T_7\,\chi(-\sigma)$ (FD-10): it maps the states at the fold of one Krein charge onto those of the other. Unit: dimensionless. Status: DERIVED.
- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi=\mathrm{e}^{\frac{7}{2}H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x}-\omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of the level of dirac16complex at the fold (a long-lived resonance) and $\bar{\chi}$ its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma=|\cot z|\,\mathrm{d}x_0$, $\sigma=\mathrm{sgn}(\pi/2-z)\ln(\sin z)/(6H)$ on one period $0<z<\pi$; $\sigma=0$ is the fold $z=\pi/2$, $\sigma\to\mp\infty$ the tips $z\to0,\pi$; one period of x_0 is the whole line $-\infty<\sigma<\infty$. Unit: length. Status: DERIVED.
- T_a, T^a: The author's eight real 16×16 Dirac matrices T16ˆA[a] of 4+4.nb ($a=0,\ldots,7$; $T_0,\ldots,T_7$), written out in FD-13: T_a has the block $\bar{\Xi}_a$ in its upper right and Ξ_a in its lower left corner; $\{T_a,T_b\}=2\eta_{ab}\mathbf{1}_{16}$; $T^a=\eta^{ab}T_b$. Unit: dimensionless. Status: AUTHOR.

(for states at rest R_1 is the reflection across the fold combined with the reflection of ordinary space, R_2 the reflection across the fold combined with the reflection of the extra times). T_a ($a=0,\ldots,7$) are the author's real 16×16 Dirac matrices `T16ˆA[a]` of `4+4.nb`, built there from six real 4×4 matrices:

$$\mathbf{s}_1=\begin{pmatrix}0&0&0&1\\0&0&1&0\\0&-1&0&0\\-1&0&0&0\end{pmatrix},\quad \mathbf{s}_2=\begin{pmatrix}0&0&-1&0\\0&0&0&1\\1&0&0&0\\0&-1&0&0\end{pmatrix},\quad \mathbf{s}_3=\begin{pmatrix}0&1&0&0\\-1&0&0&0\\0&0&0&1\\0&0&-1&0\end{pmatrix},$$
$$\mathbf{t}_1=\begin{pmatrix}0&0&0&-1\\0&0&1&0\\0&-1&0&0\\1&0&0&0\end{pmatrix},\quad \mathbf{t}_2=\begin{pmatrix}0&0&-1&0\\0&0&0&-1\\1&0&0&0\\0&1&0&0\end{pmatrix},\quad \mathbf{t}_3=\begin{pmatrix}0&1&0&0\\-1&0&0&0\\0&0&0&-1\\0&0&1&0\end{pmatrix},$$
$$\Xi_0=\mathbf{1}_8,\quad \Xi_1=\begin{pmatrix}0&\mathbf{s}_1\\\mathbf{s}_1&0\end{pmatrix},\quad \Xi_2=\begin{pmatrix}0&\mathbf{s}_2\\\mathbf{s}_2&0\end{pmatrix},\quad \Xi_3=\begin{pmatrix}0&\mathbf{s}_3\\\mathbf{s}_3&0\end{pmatrix},$$
$$\Xi_4=\begin{pmatrix}0&\mathbf{t}_3\\-\mathbf{t}_3&0\end{pmatrix},\quad \Xi_5=\begin{pmatrix}0&\mathbf{t}_2\\-\mathbf{t}_2&0\end{pmatrix},\quad \Xi_6=\begin{pmatrix}0&\mathbf{t}_1\\-\mathbf{t}_1&0\end{pmatrix},\quad \Xi_7=\Xi_1\Xi_2\Xi_3\Xi_4\Xi_5\Xi_6=\begin{pmatrix}-\mathbf{1}_4&0\\0&\mathbf{1}_4\end{pmatrix},$$
$$\bar{\Xi}_0=\mathbf{1}_8,\quad \bar{\Xi}_a=\sigma_8\,\Xi_a^T\,\sigma_8,\quad \sigma_8=\begin{pmatrix}0&\mathbf{1}_4\\\mathbf{1}_4&0\end{pmatrix},\qquad T_a=\begin{pmatrix}0&\bar{\Xi}_a\\\Xi_a&0\end{pmatrix} \tag{FD-13}$$

Symbols in (FD-13).

- $\mathbf{s}_1$, $\mathbf{s}_2$, $\mathbf{s}_3$: The author's three real antisymmetric 4×4 matrices s4by4[1], s4by4[2], s4by4[3] of 4+4.nb (written out in FD-13), the blocks of Ξ_1, Ξ_2, Ξ_3. Unit: dimensionless. Status: AUTHOR.
- $\mathbf{t}_1$, $\mathbf{t}_2$, $\mathbf{t}_3$: The author's three real antisymmetric 4×4 matrices t4by4[1], t4by4[2], t4by4[3] of 4+4.nb (written out in FD-13), the blocks of Ξ_6, Ξ_5, Ξ_4. Unit: dimensionless. Status: AUTHOR.

- Ξ_a: The author's eight real 8×8 matrices tau[a] of 4+4.nb ($a=0,\dots,7$; $\Xi_0=\mathbf{1}_8$, $\Xi_7=\Xi_1\Xi_2\cdots\Xi_6$), the lower left blocks of the Dirac matrices T_a (FD-13). Unit: dimensionless. Status: AUTHOR.
- $\bar\Xi_a$: The author's eight real 8×8 matrices taubar[a] of 4+4.nb, $\bar\Xi_0=\mathbf{1}_8$ and $\bar\Xi_a=\sigma_8\Xi_a^T\sigma_8$ for $a=1,\dots,7$, the upper right blocks of the Dirac matrices T_a (FD-13). Unit: dimensionless. Status: AUTHOR.
- $\mathbf{1}$, $\mathbf{1}_4$, $\mathbf{1}_8$, $\mathbf{1}_{16}$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_4$, $\mathbf{1}_8$ and $\mathbf{1}_{16}$ are the identity matrices of 4, 8 and 16 rows. Unit: dimensionless. Status: STANDARD.
- σ_8: The author's real symmetric 8×8 matrix sigma88 of 4+4.nb, with the blocks $\mathbf{1}_4$ off the diagonal and zeros on it (FD-13); $\sigma_8^2=\mathbf{1}_8$ and $\sigma_{16}=\mathrm{diag}(-\sigma_8,\sigma_8)$. Unit: dimensionless. Status: AUTHOR.
- ^{T}: The transpose of a matrix. Status: STANDARD.
- T_a, T^a: The author's eight real 16×16 Dirac matrices T16^A[a] of 4+4.nb ($a=0,\dots,7$; $T_0,\dots,T_7$), written out in FD-13: T_a has the block $\bar\Xi_a$ in its upper right and Ξ_a in its lower left corner; $\{T_a,T_b\}=2\eta_{ab}\mathbf{1}_{16}$; $T^a=\eta^{ab}T_b$. Unit: dimensionless. Status: AUTHOR.
- a: A frame (orthonormal) index running over $0,1,\dots,7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab}=\mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a,b\in\{1,\dots,7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat g_{ab}$ and the tensors built from it) a coordinate index of the fold, $a\in\{1,\dots,7\}$; a repeated index is summed. Status: STANDARD.
- η_{ab}: The flat metric of the frame, $\eta=\mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$ (eta4488 of 4+4.nb); η^{ab} is its inverse (the same matrix). Unit: dimensionless. Status: AUTHOR.
- σ_{16}: The matrix of the Dirac adjoint, $\sigma_{16}=T_0T_1T_2T_3$, real and symmetric; $\sigma_{16}T_a$ is real and antisymmetric. Unit: dimensionless. Status: AUTHOR.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0,x_1,\dots,x_7$ for a tensor of eight dimensions, $x_1,\dots,x_7$ for a tensor of the fold, $x_1,\dots,x_4$ for a four-dimensional tensor). Status: STANDARD.

($\bar\Xi_a=\sigma_8\Xi_a^T\sigma_8$ for $a=1,\dots,7$, and T_a for $a=0,\dots,7$; a matrix of four blocks, each block 4×4 or 8×8, 0 a block of zeros; $\mathbf{s}_1,\mathbf{s}_2,\mathbf{s}_3$ and $\mathbf{t}_1,\mathbf{t}_2,\mathbf{t}_3$ are `s4by4[1..3]` and `t4by4[1..3]`, Ξ_a and $\bar\Xi_a$ are `tau[a]` and `taubar[a]`, σ_8 is `sigma88` of the notebook). They obey $\{T_a,T_b\}=T_aT_b+T_bT_a=2\eta_{ab}\mathbf{1}_{16}$ with $\eta=\mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; $\gamma^{(a)}=\eta^{aa}T_a$ (no sum) are the Dirac matrices of the frame, $\gamma^{(0)}$ along σ; the chirality matrix is $\Gamma_8=T_0T_1\cdots T_7=\mathrm{diag}(-\mathbf{1}_8,\mathbf{1}_8)$, the chiral projectors are $P_\mathrm{L}=(\mathbf{1}_{16}-\Gamma_8)/2$ and $P_\mathrm{R}=(\mathbf{1}_{16}+\Gamma_8)/2$, and $\sigma_{16}=T_0T_1T_2T_3=\mathrm{diag}(-\sigma_8,\sigma_8)$ defines the Dirac adjoint $\bar\Psi=\Psi^\dagger\sigma_{16}$ ($\Psi^\dagger$ the complex conjugate transpose). The Krein charge of a state is its share of the conserved charge of dirac16complex, which is indefinite: a state can carry the charge -1 or $+1$. R_1 is a symmetry of dirac16complex, of gravity and of a mean field of a gas of the particles χ that is the same on both sides of the fold: the four resonances of Krein charge -1 (the particles χ) are even under it and the four of Krein charge $+1$ are odd, so such a mean field cannot mix the two signs of the indefinite charge. R_2 maps the states of one Krein charge onto those of the other and makes the two sets degenerate without the self-interaction. 9. *The planes of Horava and Witten.* In M-theory (the eleven-dimensional theory whose limits include the superstring theories) on an interval (a circle whose points are identified with their mirror images), the two ten-dimensional planes left in place by the reflection are mirror surfaces (property 1); each carries one E_8 gauge group while gravity fills the bulk. As in UM-4, gravity fills the bulk; unlike UM-4, the gauge fields live on the mirror planes (in UM-4 they live in the bulk), and the two sides of each plane are identified (one physical side).

What is not a fold.

- *The tips* $z=0$ and $z=\pi$ (mod π): the mirror exchanges them within a period, but there the warp factor vanishes, $g_{x_0x_0}$ diverges, and they lie at an infinite proper distance; they carry no thin source.
- *A smooth maximum of a warp factor*, such as $1/\cosh^2(H\sigma)$: mirror symmetric and largest at $\sigma=0$, but without a kink and without a thin source.

2 How to read this document

2.1 What this document contains

This document solves, approximately and step by step, the coupled field equations of the universe model UM-4: Einstein gravity in eight dimensions (the author's Einstein-Lovelock theory with the weights of its Gauss-Bonnet and cubic terms set to zero) with the metric of UM-4, the author's metric with the time warped like every other direction of the fold (the author's order of 2026-10-25, 13:15 PDT); the author's field dirac16complex (a 16-component complex

spinor field) and the gauge fields of the Standard Model in the bulk; and the Higgs doublet, the quarks and the leptons with three right-handed neutrinos on the fold. It is self-contained: every equation that is solved is written out here (with the tag it carries in every UM-4 document), every new equation of the solution carries a tag S-1, S-2, ..., and every symbol of every display equation is defined right after it. The sections are:

- **Gravity**: why UM-4 uses Einstein gravity with $\Lambda = -21H^2$ (the only choice of the author's Lovelock weights for which gravitons travel on the light cone on every background).
- **The background**: the exact bulk equations of the metric of UM-4, the pure tension of the fold, the static block, the Ricci-flat folds, the author's vacuum (not a solution), the bulk metric of an expanding fold and the equations of the fold, which to the first order in its curvature are the Einstein equations of seven dimensions.
- **The volume of the extra times**, which must be held fixed, and the pressure that holds it.
- **The sources of the fold**: the gravitational waves of the fold, the gravity that the fields of the fold may induce, the fold fluid and the tuning of its tension.
- **The Higgs field in the ad hoc vacuum**: the author's ad hoc vacuum written as a gauge rotation, the field of W^2 that the Gauss law requires, and the one residual that remains.
- **Propagation in the bulk**: the modes of every field of the bulk, the photon and the gravitational wave, the propagators between points of the fold, Newton's and Coulomb's laws, the W and Z bosons, the bounds on H, and the causal structure of the bulk.
- **dirac16complex**: its equations, the resonances that the fold holds (the particles χ, with their mass and their width), the lifetime that the dark matter requires, and the momentum along the extra times.
- **The Kohn-Sham states of dirac16complex**: the self-consistent states of a gas of the particles χ with the self-interaction, gravity and the mean fields of the Standard Model, with their residuals, and the Krein norm of the field.
- **The residuals of every field equation**: how well the approximate solution satisfies each equation.
- **The three neutrino flavours**: the Sun, the envelope of a supernova, the fast instability and the flavour equipartition, and what the newer global fit, the Dirac phase and the lightest mass change.
- **The thermal history**: the degrees of freedom, the relic density, the portals of χ to the Standard Model, when the particles χ of UM-4 could be a thermal relic (only if they are strongly bound) and why the portals cannot bind them to the fold, the right-handed neutrinos and the ad hoc vacuum before nucleosynthesis.
- **The primordial era**: the exact de Sitter fold, the inflation of the fold with frozen extra times and its scales.
- **The CMB, the CNB and the BAO** (the cosmic microwave background, the cosmic neutrino background and the baryon acoustic oscillations): the cosmological perturbations of the metric of the fold, against the measurements of Planck, of DESI DR2 (the second data release of the Dark Energy Spectroscopic Instrument) and of ACT DR6 (the sixth data release of the Atacama Cosmology Telescope).
- **What the solution does not settle**, and **what was verified**.

2.2 The status labels

Every equation and every result carries one of the labels

- **AUTHOR**: from the author (his notebooks `4+4.nb` and `Dirac_Einstein-Lovelock_4+4_Nash.nb`, and his orders of 2026-10-24 and 2026-10-25: the coordinates, the metric with its change of 2026-10-25 13:15 (the time warped), the Einstein-Lovelock action, the Dirac matrices, the chiral projectors, the list of fields, the Higgs potential and the ad hoc vacuum);
- **RECORD**: from the record of this repository (the Lagrangian of dirac16complex with its canonical spin connection and its contact self-interaction, the Kohn-Sham treatment);
- **STANDARD**: textbook physics, or a published measurement (the source is named);
- **ASSUMED**: a choice made in UM-4 without derivation;
- **INVENTED**: a new interaction of UM-4 (the portals of dirac16complex);
- **DERIVED**: computed and verified in UM-4 (the verification is stated);
- **ESTIMATE**: an order-of-magnitude evaluation;
- **OPEN**: not settled.

2.3 Every equation is followed by its symbols

Every display equation is followed by the list "Symbols in (S-n)" or "Symbols in (FE-n)" of EVERY symbol that appears in it: every quantity, field, coupling, constant and operator, every index letter with the values it takes, every mark (prime, dagger, transpose, star), the imaginary unit, Euler's number and the differential. Each item gives the

meaning, the unit (natural units, in which the reduced Planck constant $\hbar$ and the speed of light c are 1), the value where one exists, and the status. A symbol that one of these meanings uses is itself listed in this document. A field equation of UM-4 carries a tag FE-n and a term of the action a tag IN-n, the same in every UM-4 document; it is written here exactly as everywhere else. A new equation of this document carries a tag S-n.

2.4 Coordinates, signature and units

The author's coordinates are $X = \{x_0, x_1, \ldots, x_7\}$, always with the hidden direction first: x_0 is the hidden space-like coordinate, on which the metric depends periodically with the period $\pi/(6H)$ through the angle $z = 6Hx_0$; x_1, x_2, x_3 are ordinary space, x_4 is the time, and x_5, x_6, x_7 are three extra time-like coordinates. The signature is $(+,+,+,+,-,-,-,-)$ in this order; the four observable coordinates $x_1, \ldots, x_4$ have the signature $(+,+,+,-)$. H is the author's constant (an inverse length), a free parameter of UM-4, bounded from below by 2534 GeV (about 2.5 TeV). Units are natural: the reduced Planck constant $\hbar$ and the speed of light c are 1 ($1\ \mathrm{GeV}^{-1} = 1.973 \times 10^{-14}$ cm $= 6.582 \times 10^{-25}$ s; $1\ \mathrm{GeV}^{-2} = 3.894 \times 10^{-28}\ \mathrm{cm}^2$). A prime on a function of the time x_4 alone is its derivative $\mathrm{d}/\mathrm{d}x_4$, and on a profile (a function of σ alone) its derivative $\mathrm{d}/\mathrm{d}\sigma$. The proper coordinate of the hidden direction is σ (FE-27): $\sigma = 0$ is the fold $z = \pi/2$. The fields live as follows: gravity, dirac16complex and the gauge fields of the Standard Model in the bulk (their massless modes are bound to the fold); the Higgs doublet, the quarks and the leptons with three right-handed neutrinos, a thin fluid (the fold fluid) and, possibly, an Einstein-Hilbert term that the fields of the fold induce (with the coupling κ_{ind}) on the fold $\sigma = 0$.

2.5 The solution at a glance

part	result	status
gravity	Einstein gravity with $\Lambda = -21H^2$: the only choice of the author's Lovelock weights for which gravitons travel on the light cone on every background; the static block is its vacuum (a space of constant curvature $-H^2$; among the static warps of the author's form the only vacuum of Einstein gravity), and the fold of the metric of UM-4 is a pure tension $\kappa\lambda_{\mathrm{F}} = 12H$	DERIVED
background	the metric of the author's form is a vacuum of the bulk only for the Ricci-flat folds (the static fold and two Kasner folds); an expanding fold with matter has a bulk metric corrected at the order of its curvature over H^2, and its fold obeys the Einstein equations of seven dimensions with $\kappa_7 = 5\kappa H/2$, with frozen extra times the Friedmann equations of general relativity (corrections of the relative size $(a_4'/H)^2$, about 3×10^{-91} today); the author's vacuum is not a solution	DERIVED
volume of the extra times	a free volume would change Newton's constant 8500 times faster than lunar laser ranging allows; the fold fluid holds it fixed with the pressure $q_{\mathrm{hold}} = -0.842\rho_{\mathrm{c}}^{(0)}$ along the extra times today (with no pressure of the baryons along them)	DERIVED, ASSUMED (the holding pressure and the baryons), OPEN (its origin)
sources of the fold	the gravitational waves of the fold travel at the speed of light at every wavelength (GW170817 holds for every H); the induced gravity is negligible for a cutoff of the order of H; the tension $\lambda_{\mathrm{F}}V_* = 30H^2M_{\mathrm{red}}^2$ must be tuned to one part in 10^{92} against the observed dark energy	DERIVED, ESTIMATE, OPEN (the tuning)
Higgs field	the ad hoc vacuum is an $SU(2)$ rotation of the radial field $\sqrt{v^2 + \mathcal{A}^2}$ with a pure-gauge W^2 (the value at the fold of the field of the bulk); one radial residual remains, which vanishes as $\mathcal{A} \to 0$	DERIVED

part	result	status
propagation in the bulk	every massless field of the bulk travels at the speed of light at every depth and every wavelength; the photon is a transverse wave of helicity ±1, the gravitational wave a transverse traceless wave of helicity ±2 with the plus and cross polarisations; of the other polarisations of the graviton of the bulk the three vectors $h_{\mu r}$ are ghosts, which the matter of the fold does not emit at the linear order; dirac16complex is invariant under the boosts of the fold; nothing that stays at fixed extra times crosses the static bulk faster than light (along the extra times the light cone of signature (4, 4) allows any shortcut), and on an expanding fold the bulk is ahead by about 4×10^{-94} over the Hubble distance; the massive modes correct Newton's and Coulomb's laws by $0.34/(Hr)^5$ and $0.42/(Hr)^3$ and give the W and Z bosons widths into the bulk, which bound $H > 2534$ GeV	DERIVED, ESTIMATE (the bounds), OPEN (the ghost vectors beyond the linear order)
dirac16complex	the states at the fold are resonances (the particles χ): mass $\omega_\mathrm{r} = \omega_\mathrm{b}(1 - H/(2q_\mathrm{b}))$, width $\Gamma_\chi = (2/\mathrm{e})(q_\mathrm{b}^2/m_\Psi)\mathrm{e}^{-2F(v_\mathrm{b})m_\Psi/H}$ (derived, to the leading order in H/q_b); dark matter (a lifetime above 160 Gyr) needs $m_\Psi/H > 1135$ for $q_\mathrm{b} = m_\Psi/2$, so with $H > 2534$ GeV the particles χ are heavier than 2.5×10^6 GeV for that binding; the bound falls with the binding, to 1.2×10^4 GeV for $q_\mathrm{b} = 0.99\, m_\Psi$; they move slower than light ($\omega^2 = \omega_0^2 + \vec{k}^2$ exactly)	DERIVED, ESTIMATE (the bounds)
Kohn-Sham	self-consistent states of a homogeneous gas of the particles χ at $m_\Psi/H = 10^4$ ($q_\mathrm{b} = 3000H$) for $G_\mathrm{KS} = 10^{-6}$ to 0.5, with the four occupied orbitals in one multiplet (a stationary point of the first-order energy that is a saddle, not a minimum): first-order perturbation theory in the dilute limit, the gas confined to the fold within the box, a positive pressure that vanishes in the dilute limit (cold dark matter), a convergent Higgs-portal loop; the repulsive self-interaction lowers the barrier, and with H at its bound the gas lives longer than 160 Gyr only for $G_\mathrm{KS} < 0.139$; a weakly bound point ($q_\mathrm{b} = 0.03\, m_\Psi$)	DERIVED, ASSUMED (the symmetric occupation)
Krein norm	the vacuum cannot decay; whether the gas of χ is classically stable is not decided (the symmetric occupation is a saddle of the first-order energy, which has no lower bound on the occupations of negative norm); chiral couplings at the fold produce ghosts (the Higgs portal with exactly its elastic amplitude), so UM-4 with them is not unitary; the convective current of the current portals creates pairs only with an amplitude of the second order in the momentum	DERIVED, OPEN
residuals	every field equation is satisfied to the stated accuracy, except the radial residual of the ad hoc vacuum, the component of the equations of the fold along the extra times (carried by the fold fluid) and the projected Weyl tensor of the bulk that an expanding fold with dust requires at the order of the square of its density over the tension	DERIVED, ASSUMED, OPEN
neutrinos	the flavour conversion of UM-4 is that of the Standard Model with three Dirac neutrinos: solar survival 0.55 at low and 0.30 at high energy (Borexino 0.57, 0.53, 0.43, 0.37); adiabatic supernova resonances; fast instabilities within centimetres wherever the electron lepton number crosses; the equipartition raises or lowers the heating behind the shock depending on the spectra	DERIVED, ESTIMATE
neutrino parameters	NuFIT 6.0 changes the results by at most 1.5 per cent; the Dirac phase drops out of every result; DESI DR2 with NuFIT 6.0 bounds the lightest mass below 23 meV, which lowers the CNB capture rate from 6.9 to 4.1 per year at most	DERIVED

part	result	status
thermal history	a thermal relic would need $\langle\sigma_{\rm ann}\mathsf{v}\rangle = 2.2\times10^{-26}$ cm^3/s, which unitarity allows only up to 9.7×10^4 GeV (6.9×10^4 to 1.37×10^5 GeV with the bookkeeping of the channels): only particles χ bound more strongly than $q_{\rm b} = 0.906\,m_\Psi$ live long enough at such masses; the portals of IN-16 cannot bind them so: bound by the Higgs portal they would have to be bound so weakly (direct detection) that they live long enough only for $H < 2.5$ GeV, and a binding by the gluon condensate makes them scatter on nucleons 29 to 33 orders of magnitude above the limit of LZ; the current portals annihilate them only with the fourth power of the velocity, too weakly for a perturbatively coupled relic that LZ allows: the dark matter of UM-4 needs a binding to the fold that does not couple χ to nucleons; the right-handed neutrinos must not thermalise; the ad hoc vacuum must have decayed long before nucleosynthesis	DERIVED, ESTIMATE, OPEN
primordial era	the de Sitter fold is exact (the extra times inflate with ordinary space); an inflation with frozen extra times needs a Weyl tensor of the bulk and, in the regime of general relativity, a rate below $H/10$: below 253 GeV for $H = 2534$ GeV; Higgs inflation (an extension of the action) would need $H > 1.6\times10^{14}$ GeV	DERIVED, ESTIMATE, OPEN
CMB, CNB, BAO	with the Planck 2018 parameters UM-4 reproduces the CMB ($\chi^2 = 584$ for 613 high-l bins) and its derived quantities; the BAO of DESI DR2 show the same tension as the cosmological-constant model (2.3 standard deviations, DESI's number), since the dark energy of UM-4 has $w = -1$; the CNB of Dirac neutrinos has one helicity per state	DERIVED, STANDARD (the DESI tension)
extra times	every field with a momentum along the extra times larger than its mass and its momentum along ordinary space is tachyonic (a smaller one is a rotation of the state at rest); UM-4 keeps only fields without it and needs a gauge principle that removes it	DERIVED, OPEN

3 Gravity

Gravity in UM-4 is the author's Einstein-Lovelock theory in eight dimensions (his notebook `Dirac_Einstein-Lovelock_4+4_Nash.nb`, In[134]) with the weights of its Gauss-Bonnet and cubic terms set to zero: Einstein gravity with the cosmological constant $\Lambda = -21H^2$. The author's order of 2026-10-25, 13:15 PDT requires that gravitons propagate at the speed of light in the bulk; this section shows that this selects Einstein gravity. Every later section uses it.

$$S_{\rm EL} = \frac{1}{2\kappa}\int \mathrm{d}^8X\,\sqrt{|g|}\,\Big(\sum_{p=1}^{3}\alpha_{(p)}\,\mathcal{L}_{(p)} - 2\Lambda\Big) = \frac{1}{2\kappa}\int \mathrm{d}^8X\,\sqrt{|g|}\,\big(R - 2\Lambda\big), \qquad \kappa = 8\pi G_8,$$
$$\alpha_{(p)} = w_{(p)}\,H^{2-2p}, \qquad w_{(1)} = 1, \quad w_{(2)} = w_{(3)} = 0, \qquad \Lambda = -21H^2 \tag{IN-2}$$

Symbols in (IN-2).

- $S_{\rm EL}$: The Einstein-Lovelock action of gravity in eight dimensions; in UM-4 the Einstein-Hilbert action with the cosmological constant. Unit: dimensionless. Status: AUTHOR.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as dσ, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\dots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- p: The Lovelock order $p=1,2,3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number (UM-4 keeps only $p=1$); in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)}=w_{(p)}H^{2-2p}$: in UM-4 $\alpha_{(1)}=1$ and $\alpha_{(2)}=\alpha_{(3)}=0$ (Einstein gravity). Unit: GeV^{2-2p}. Status: AUTHOR.
- $\mathcal{L}_{(p)}$: The Lovelock density of order p, $\mathcal{L}_{(p)}=2^{-p}\,\delta^{P_1\cdots P_{2p}}_{Q_1\cdots Q_{2p}}\,R^{Q_1Q_2}{}_{P_1P_2}\cdots R^{Q_{2p-1}Q_{2p}}{}_{P_{2p-1}P_{2p}}$, a scalar built from p curvature tensors: $\mathcal{L}_{(1)}=R$ (Einstein-Hilbert), $\mathcal{L}_{(2)}$ the Gauss-Bonnet density, $\mathcal{L}_{(3)}$ the cubic Lovelock density. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant of UM-4, $\Lambda=-21H^2$: the value for which the static block is a vacuum of Einstein gravity (its Einstein tensor is $21H^2$ in every direction). The author writes the pure number Λ_{N} in $0=-\Lambda_{\mathrm{N}}+\sum_p H^{-2p}w_{(p)}\mathcal{E}_{(p)}$, so that $\Lambda=-\Lambda_{\mathrm{N}}H^2$. Unit: GeV^2. Value: $-21H^2$; the author's original $-10H^2$; UM-3: $-7H^2$. Status: AUTHOR (the term), DERIVED (the value).
- $R^{MN}{}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor $R^M{}_{NPQ}$ (in the convention in which the commutator of two covariant derivatives, acting on a vector, is the Riemann tensor, with its last two indices those of the two derivatives, contracted with the vector), with two indices the Ricci tensor $R_{MN}=R^P{}_{MPN}$, without indices the scalar curvature $R=g^{MN}R_{MN}$; with frame indices $R^{(a)(b)}{}_{(c)(d)}$ its frame components, $R^{(a)(b)}{}_{(a)(b)}$ (no sum) the sectional curvature of the frame plane (a,b). Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_8: The eight-dimensional Newton constant ($\kappa=8\pi G_8$). Unit: GeV^{-6}. Status: AUTHOR.
- $w_{(p)}$: The Lovelock weights of the author's Einstein-Lovelock action, $w_{(1)}=1$ and $w_{(2)}$, $w_{(3)}$ of the Gauss-Bonnet and the cubic densities. UM-4 sets $w_{(2)}=w_{(3)}=0$ (Einstein gravity): the only weights for which gravitons propagate on the light cone on every background, not only around the static block. The author's original weights $w_{(2)}=1/32$, $w_{(3)}=1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) keep the gravitons on the light cone only around the static block (with $\Lambda=-161H^2/16$); UM-3 used $(1,1/20,1/360)$. Unit: dimensionless. Value: $(1,0,0)$ in UM-4; the author's original $(1,1/32,1/1152)$. Status: AUTHOR (the form), DERIVED (the values that the order of 2026-10-25 selects).
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda=-21H^2$ and the tension $\kappa\lambda_{\mathrm{F}}=12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H>2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.

The gravitational action of UM-4. The weights multiply the Lovelock densities in units of the author's constant H ($\alpha_{(p)}=w_{(p)}H^{2-2p}$); with $w_{(2)}=w_{(3)}=0$ only the Einstein-Hilbert term and the cosmological constant remain. Status: AUTHOR (the action), DERIVED (the weights and Λ that the order selects, below).

$$
\begin{aligned}
&\sum_{p=1}^{3}\alpha_{(p)}\,\mathcal{E}_{(p)}{}^a{}_b+\Lambda\,\delta^a{}_b=\mathsf{P}(\mathsf{y})\,H^2\,\delta^a{}_b,\qquad \mathsf{P}(\mathsf{y})=21\mathsf{y}-420w_{(2)}\mathsf{y}^2+2520w_{(3)}\mathsf{y}^3+\frac{\Lambda}{H^2},\\
&\text{Einstein } (w_{(2)}=w_{(3)}=0,\ \Lambda=-21H^2):\quad \mathsf{P}=21(\mathsf{y}-1);\\
&\text{the author's original weights } \Big(\frac{1}{32},\frac{1}{1152}\Big),\ \Lambda=-\frac{161}{16}H^2:\quad \mathsf{P}=\frac{7}{16}(\mathsf{y}-1)\big(5\mathsf{y}^2-25\mathsf{y}+23\big);\\
&\text{UM-3's weights } \Big(\frac{1}{20},\frac{1}{360}\Big),\ \Lambda=-7H^2:\quad \mathsf{P}=7(\mathsf{y}-1)^3
\end{aligned}
\tag{FE-70}
$$

Symbols in (FE-70).

- p: The Lovelock order $p=1,2,3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number (UM-4 keeps only $p=1$); in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)}=w_{(p)}H^{2-2p}$: in UM-4 $\alpha_{(1)}=1$ and $\alpha_{(2)}=\alpha_{(3)}=0$ (Einstein gravity). Unit: GeV^{2-2p}. Status: AUTHOR.
- $\mathcal{E}_{(p)}{}^M{}_N$: The Lovelock tensor of order p, $\mathcal{E}_{(p)}{}^M{}_N=-2^{-(p+1)}\,\delta^{MP_1\cdots P_{2p}}_{NQ_1\cdots Q_{2p}}\,R^{Q_1Q_2}{}_{P_1P_2}\cdots R^{Q_{2p-1}Q_{2p}}{}_{P_{2p-1}P_{2p}}$ (with

both indices down it is $1/\sqrt{|g|}$ times the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to g^{MN}; $\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor, $\mathcal{E}_{(3)}$ the cubic Lovelock tensor); with frame indices its frame components. Unit: $1/\text{length}^{2p}$. Status: STANDARD.

- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \text{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- Λ: The eight-dimensional cosmological constant of UM-4, $\Lambda = -21H^2$: the value for which the static block is a vacuum of Einstein gravity (its Einstein tensor is $21H^2$ in every direction). The author writes the pure number Λ_N in $0 = -\Lambda_\text{N} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$, so that $\Lambda = -\Lambda_\text{N} H^2$. Unit: GeV^2. Value: $-21H^2$; the author's original $-10H^2$; UM-3: $-7H^2$. Status: AUTHOR (the term), DERIVED (the value).
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\text{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- $\mathsf{P}(\mathsf{y})$: The vacuum polynomial of the static blocks of curvature $-\mathsf{y}H^2$: the left-hand side of the bulk equations divided by H^2, the same in every direction, $\mathsf{P}(\mathsf{y}) = 21\mathsf{y} - 420 w_{(2)} \mathsf{y}^2 + 2520 w_{(3)} \mathsf{y}^3 + \Lambda/H^2$; $\mathsf{P}'(1)$ its derivative at $\mathsf{y} = 1$, which sets the strength of the graviton around the static block relative to Einstein gravity ($\mathsf{P}'(1)/21$). Unit: dimensionless. Value: $21(\mathsf{y} - 1)$ for Einstein gravity with $\Lambda = -21H^2$. Status: DERIVED.
- y: The curvature of a static block of constant curvature $-\mathsf{y}H^2$ in all eight directions, in units of $-H^2$ (its warp is $\text{e}^{-2\sqrt{\mathsf{y}} H|\sigma|}$ in all seven directions of the fold): $\mathsf{y} = 1$ is the static block of UM-4. Unit: dimensionless. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\text{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\text{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- $w_{(p)}$: The Lovelock weights of the author's Einstein-Lovelock action, $w_{(1)} = 1$ and $w_{(2)}$, $w_{(3)}$ of the Gauss-Bonnet and the cubic densities. UM-4 sets $w_{(2)} = w_{(3)} = 0$ (Einstein gravity): the only weights for which gravitons propagate on the light cone on every background, not only around the static block. The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) keep the gravitons on the light cone only around the static block (with $\Lambda = -161H^2/16$); UM-3 used $(1, 1/20, 1/360)$. Unit: dimensionless. Value: $(1, 0, 0)$ in UM-4; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), DERIVED (the values that the order of 2026-10-25 selects).

The solution, step 1: the vacua. On the static blocks of curvature $-\mathsf{y}H^2$ in all eight directions the left-hand side of the bulk equations is the same in every direction, $\mathsf{P}(\mathsf{y})H^2$. The static block of UM-4 ($\mathsf{y} = 1$) is a vacuum if $\mathsf{P}(1) = 0$ and a non-degenerate one if $\mathsf{P}'(1) \neq 0$. Einstein gravity has the single vacuum $\mathsf{y} = 1$ with $\mathsf{P}'(1) = 21$; the author's original weights would need $\Lambda = -161H^2/16$ (not his $-10H^2$), with three vacua ($\mathsf{y} = 1$, 1.2155 and 3.7845) and $\mathsf{P}'(1) = 21/16$; UM-3's weights make $\mathsf{y} = 1$ a triple root, a degenerate vacuum without linear gravitons. Status: DERIVED (`um4_geometry.py`, G4; `um4_gravity.py`, L1).

$$\frac{\mathsf{P}'(1)}{42}\Big[\text{e}^{2H|\sigma|}\Big(\frac{\partial^2 h_\text{gw}}{\partial x_4^2} - \frac{\partial^2 h_\text{gw}}{\partial x_3^2}\Big) - \frac{\partial^2 h_\text{gw}}{\partial \sigma^2} + 7H\,\text{sgn}(\sigma)\,\frac{\partial h_\text{gw}}{\partial \sigma}\Big] = 0, \qquad \mathsf{P}'(1) = 21 - 840 w_{(2)} + 7560 w_{(3)} \quad \text{(FE-68)}$$

Symbols in (FE-68).

- $\mathsf{P}(\mathsf{y})$: The vacuum polynomial of the static blocks of curvature $-\mathsf{y}H^2$: the left-hand side of the bulk equations divided by H^2, the same in every direction, $\mathsf{P}(\mathsf{y}) = 21\mathsf{y} - 420 w_{(2)} \mathsf{y}^2 + 2520 w_{(3)} \mathsf{y}^3 + \Lambda/H^2$; $\mathsf{P}'(1)$ its derivative at $\mathsf{y} = 1$, which sets the strength of the graviton around the static block relative to Einstein gravity ($\mathsf{P}'(1)/21$). Unit: dimensionless. Value: $21(\mathsf{y} - 1)$ for Einstein gravity with $\Lambda = -21H^2$. Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \text{d}a_4/\text{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- h_{gw}: The amplitude of a gravitational wave of the + polarisation travelling along x_3: the metric components along x_1 and x_2 multiplied by $\mathrm{e}^{h_{\mathrm{gw}}}$ and $\mathrm{e}^{-h_{\mathrm{gw}}}$; a function of σ, x_3 and x_4 in the bulk, and on the fold the gravitational wave of the four-dimensional world, to the first order its plus polarisation h_+. Unit: dimensionless. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- $w_{(p)}$: The Lovelock weights of the author's Einstein-Lovelock action, $w_{(1)} = 1$ and $w_{(2)}$, $w_{(3)}$ of the Gauss-Bonnet and the cubic densities. UM-4 sets $w_{(2)} = w_{(3)} = 0$ (Einstein gravity): the only weights for which gravitons propagate on the light cone on every background, not only around the static block. The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) keep the gravitons on the light cone only around the static block (with $\Lambda = -161H^2/16$); UM-3 used $(1, 1/20, 1/360)$. Unit: dimensionless. Value: $(1, 0, 0)$ in UM-4; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), DERIVED (the values that the order of 2026-10-25 selects).

The solution, step 2: the graviton around the static block, for any weights. A transverse traceless wave of the + polarisation along x_3 obeys the wave equation of the light cone of the fold at every depth, with the strength $\mathsf{P}'(1)/21$ relative to Einstein gravity: around the static block the order does not single out the weights. Status: DERIVED (`um4_gravity.py`, L2).

$$
\begin{aligned}
&\mathrm{d}\sigma^2 + \mathrm{e}^{-2\mathfrak{p}H\sigma}\,\mathrm{d}\vec{x}^2 - \mathrm{e}^{-2\mathfrak{q}H\sigma}\,\mathrm{d}x_4^2 - \mathrm{e}^{-2\mathfrak{r}H\sigma}\,\mathrm{d}\vec{x}_5^2 \quad (\sigma > 0):\\
&c_{\mathrm{x}} - c_{\mathrm{t}} = -4\,(\mathfrak{p} - \mathfrak{q})\big(18\mathfrak{r}^2 w_{(3)} - w_{(2)}\big)\big(\mathfrak{p} + \mathfrak{q} + 3\mathfrak{r}\big), \qquad \text{(FE-69)}\\
&c_\sigma - c_{\mathrm{t}} = -4\big[w_{(2)}\big(\mathfrak{p}\mathfrak{q} - \mathfrak{p}^2 + 3\mathfrak{q}\mathfrak{r} - 3\mathfrak{r}^2\big) + 18 w_{(3)}\mathfrak{r}^2\big(\mathfrak{p}^2 - 3\mathfrak{p}\mathfrak{q} + 2\mathfrak{p}\mathfrak{r} - \mathfrak{q}\mathfrak{r} + \mathfrak{r}^2\big)\big]
\end{aligned}
$$

Symbols in (FE-69).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- $\mathfrak{p}$, $\mathfrak{q}$, $\mathfrak{r}$: The rates of the warps of ordinary space ($\mathfrak{p}$), of the time ($\mathfrak{q}$) and of the extra times ($\mathfrak{r}$) of a static warped background, in units of H. Unit: dimensionless. Value: $\mathfrak{p} = \mathfrak{q} = \mathfrak{r} = 1$ for the static block of UM-4; $\mathfrak{p} = \mathfrak{r} = 1$, $\mathfrak{q} = 0$ for that of UM-3. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- $\vec{x}$: The position in ordinary space, $\vec{x} = (x_1, x_2, x_3)$. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\vec{x}_5$: The position along the extra times, $\vec{x}_5 = (x_5, x_6, x_7)$; $\mathrm{d}\vec{x}_5^2 = \mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2$. Unit: length. Status: AUTHOR.
- c_{x}: The coefficient of the second derivative along the proper length of x_3 in the same equation; the wave moves along x_3 with the local speed $\sqrt{c_{\mathrm{x}}/c_{\mathrm{t}}}$ in units of the local speed of light. Unit: dimensionless. Value: 1 for

Einstein gravity on every background; $1 - 24w_{(2)} + 72w_{(3)}$ at the static block of UM-3 ($c_\mathrm{x}/c_\mathrm{t} = 5$ with the author's original weights). Status: DERIVED.

- c_t: The coefficient of the second derivative along the proper time in the linearised equation of a gravitational wave of the + polarisation on a static warped background, normalised so that Einstein gravity gives 1. Unit: dimensionless. Value: 1 for Einstein gravity on every background; $1 - 40w_{(2)} + 360w_{(3)}$ at the static block of UM-3. Status: DERIVED.
- c_σ: The coefficient of the second derivative along σ in the linearised equation of a gravitational wave on a static warped background, normalised so that Einstein gravity gives 1; the wave moves along σ with the local speed $\sqrt{c_\sigma/c_\mathrm{t}}$ in units of the local speed of light. Unit: dimensionless. Value: 1 for Einstein gravity on every background. Status: DERIVED.
- $w_{(p)}$: The Lovelock weights of the author's Einstein-Lovelock action, $w_{(1)} = 1$ and $w_{(2)}$, $w_{(3)}$ of the Gauss-Bonnet and the cubic densities. UM-4 sets $w_{(2)} = w_{(3)} = 0$ (Einstein gravity): the only weights for which gravitons propagate on the light cone on every background, not only around the static block. The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) keep the gravitons on the light cone only around the static block (with $\Lambda = -161H^2/16$); UM-3 used $(1, 1/20, 1/360)$. Unit: dimensionless. Value: $(1, 0, 0)$ in UM-4; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), DERIVED (the values that the order of 2026-10-25 selects).

The solution, step 3: the gravitons on backgrounds that are not of constant curvature, the static warps with the rates $\mathfrak{p}$, $\mathfrak{q}$, $\mathfrak{r}$ of ordinary space, the time and the extra times. Einstein gravity gives the coefficients $c_\mathrm{t} = c_\mathrm{x} = c_\sigma = 1$ of the second derivatives along the proper time, the proper length of x_3 and σ on every such background, so its gravitons move on the local light cone; the Gauss-Bonnet and cubic terms give the deviations shown, which vanish for all rates only for $w_{(2)} = w_{(3)} = 0$. An expanding universe, a star or a gravitational wave of finite amplitude make the warps of the different directions differ, so any other choice of the weights would let gravitons leave the light cone there. Einstein gravity is therefore the gravity of UM-4. Status: DERIVED (`um4_gravity.py`, L3 and L4).

4 The background

The order 0 of the solution is the metric of UM-4 itself, with the author's two functions of the time, a_4 and b_4 (b_4 is a function of x_4 and never identically 1, by the author's order of 2026-10-24). Its exact curvature and its exact field equations in the bulk and at the fold are restated here; the solution asks which functions $a_4(x_4)$ and $b_4(x_4)$ they admit, what they require, and how the metric changes when the fold carries matter.

$$
\begin{aligned}
g_{MN} &= \mathrm{diag}\Big(\cot^2 z,\ \mathrm{e}^{2a_4}(\sin^2 z)^{1/6},\ \mathrm{e}^{2a_4}(\sin^2 z)^{1/6},\ \mathrm{e}^{2a_4}(\sin^2 z)^{1/6},\ -(\sin^2 z)^{1/6},\\
&\qquad -\mathrm{e}^{-2a_4}b_4^2(\sin^2 z)^{1/6},\ -\mathrm{e}^{-2a_4}b_4^2(\sin^2 z)^{1/6},\ -\mathrm{e}^{-2a_4}b_4^2(\sin^2 z)^{1/6}\Big), \qquad z = 6Hx_0,\\
\sigma &= \mathrm{sgn}\Big(\frac{\pi}{2} - z\Big)\frac{\ln\sin z}{6H}, \qquad \mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0,\\
g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N &= \mathrm{d}\sigma^2 + \mathrm{e}^{-2H|\sigma|}\Big[\mathrm{e}^{2a_4}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{d}x_4^2 - \mathrm{e}^{-2a_4}b_4^2\big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big)\Big]
\end{aligned}
\tag{FE-27}
$$

Symbols in (FE-27).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$, the fold is $z = \pi/2$. Unit: dimensionless. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\, \mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.

The metric of UM-4 (the author's metric with the time warped by the same factor $(\sin^2 z)^{1/6}$ as the other six directions of the fold) and the proper chart: one period of x_0 is the whole line of σ, the fold $z = \pi/2$ is $\sigma = 0$, and all seven directions of the fold are warped by $\mathrm{e}^{-H|\sigma|}$. Status: AUTHOR (the metric and the order), ASSUMED (the way the order is met), DERIVED (the chart; `um4_geometry.py`, G1).

$$
\begin{aligned}
&\mathbf{e}^{(0)} = \mathrm{d}\sigma, \quad \mathbf{e}^{(i)} = \mathrm{e}^{a_4 - H|\sigma|}\mathrm{d}x_i, \quad \mathbf{e}^{(4)} = \mathrm{e}^{-H|\sigma|}\mathrm{d}x_4, \quad \mathbf{e}^{(r)} = b_4\,\mathrm{e}^{-a_4 - H|\sigma|}\mathrm{d}x_r, \qquad \sqrt{|g|} = b_4^3\,\mathrm{e}^{-7H|\sigma|},\\
&R^{(a)(b)}{}_{(c)(d)} = \mathrm{e}^{2H|\sigma|}\,\hat{R}^{(a)(b)}{}_{(c)(d)} - H^2\,\delta^{ab}_{cd} \quad \text{(all four along the fold)}, \qquad R^{(0)(a)}{}_{(0)(b)} = -H^2\,\delta^a{}_b,\\
&\hat{R}^{(i)(j)}{}_{(i)(j)} = \mathsf{A}^2H^2, \quad \hat{R}^{(i)(4)}{}_{(i)(4)} = (\mathsf{A}^2 + \mathsf{A}_1)H^2, \quad \hat{R}^{(i)(r)}{}_{(i)(r)} = \mathsf{A}\mathsf{V}H^2,\\
&\hat{R}^{(r)(s)}{}_{(r)(s)} = \mathsf{V}^2H^2, \quad \hat{R}^{(4)(r)}{}_{(4)(r)} = (\mathsf{V}^2 + \mathsf{V}_1)H^2,\\
&\mathsf{A} = \frac{a_4'}{H}, \qquad \mathsf{U} = \frac{b_4'}{Hb_4}, \qquad \mathsf{V} = \mathsf{U} - \mathsf{A}, \qquad \mathsf{A}_1 = \frac{\mathsf{A}'}{H}, \qquad \mathsf{V}_1 = \frac{\mathsf{V}'}{H}
\end{aligned}
\tag{FE-28}
$$

Symbols in (FE-28).

- $\mathbf{e}_a{}^M$, $\mathbf{e}^a{}_M$: The vielbein (frame field) of the metric: $g_{MN} = \eta_{ab}\,\mathbf{e}^a{}_M\mathbf{e}^b{}_N$; $\mathbf{e}_a{}^M$ is its inverse; $\mathbf{e}_{bN} = \eta_{bc}\mathbf{e}^c{}_N$; $\mathbf{e}^{(a)}$ is the frame one-form $\mathbf{e}^a{}_M\,\mathrm{d}X^M$. Unit: dimensionless. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- x: The coordinates of the fold: x_1, x_2, x_3 of ordinary space, x_4 the time, x_5, x_6, x_7 the extra times; a field of the fold is a function of the four observable coordinates $x = (x_1, x_2, x_3, x_4)$, and $\mathrm{d}^4x = \mathrm{d}x_1\,\mathrm{d}x_2\,\mathrm{d}x_3\,\mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal), and in a tensor of ordinary space such as h_{ij} one of the directions x_1, x_2, x_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- $R^{MN}{}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor $R^M{}_{NPQ}$ (in the convention in which the commutator of two covariant derivatives, acting on a vector, is the Riemann tensor, with its last two indices those of the two derivatives, contracted with the vector), with two indices the Ricci tensor $R_{MN} = R^P{}_{MPN}$, without indices the scalar curvature $R = g^{MN}R_{MN}$; with frame indices $R^{(a)(b)}{}_{(c)(d)}$ its frame components, $R^{(a)(b)}{}_{(a)(b)}$ (no sum) the sectional curvature of the frame plane (a, b). Unit: 1/length^2. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- c: A frame index (as a). Status: STANDARD.
- d: A frame index (as a). Status: STANDARD.
- $\hat{R}$: The curvature of the metric $\hat{g}_{ab}$ of the fold (FD-2): with four frame indices its Riemann tensor, with two its Ricci tensor $\hat{R}_{ab}$, without indices its scalar curvature $\hat{R}$; for frozen extra times the scalar curvature equals that of the four-dimensional metric $\hat{h}_{\mu\nu}$. Unit: 1/length^2. Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1M_2\cdots}_{N_1N_2\cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- j: A second generation or mass-state index, or a second direction of ordinary space (as i). Status: STANDARD.
- s: A second extra-time direction of the fold (as r, different from it). Status: STANDARD.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); the first-order equations of the fold hold for $\mathsf{A}^2 \ll 1$; 1 at the author's vacuum, which is not a solution of UM-4. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- V: The rate of change of the scale factor $\mathrm{e}^{-a_4}b_4$ of the extra times on the fold in units of H, $\mathsf{V} = \mathsf{U} - \mathsf{A}$; zero with frozen extra times. Unit: dimensionless. Status: DERIVED.
- V_1: The time derivative of V in units of H, $\mathsf{V}_1 = \mathsf{V}'/H$. Unit: dimensionless. Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra

times. Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.

The orthonormal frame, the volume element, the exact frame curvature and the curvature of the metric of the fold, with the dimensionless rates of the background. The metric of UM-4 is a warped product: in every plane of two directions of the fold the curvature of the bulk is that of the fold enlarged by $\mathrm{e}^{2H|\sigma|}$, minus H^2; in every plane with the hidden direction it is $-H^2$. Status: DERIVED (`um4_geometry.py`, G2, exact, in the author's coordinates on both halves of the period).

$$\mathcal{E}_{(1)}{}^{(a)}{}_{(b)} + \Lambda\,\delta^a{}_b = \mathrm{e}^{2H|\sigma|}\,\hat{G}^a{}_b \quad \text{(both along the fold)}, \qquad \mathcal{E}_{(1)}{}^{(0)}{}_{(0)} + \Lambda = -\frac{1}{2}\,\mathrm{e}^{2H|\sigma|}\,\hat{R}, \qquad \mathcal{E}_{(1)}{}^{(0)}{}_{(a)} = 0 \quad \text{(FE-29)}$$

Symbols in (FE-29).

- $\mathcal{E}_{(p)}{}^{M}{}_{N}$: The Lovelock tensor of order p, $\mathcal{E}_{(p)}{}^{M}{}_{N} = -2^{-(p+1)}\,\delta^{MP_1\cdots P_{2p}}_{NQ_1\cdots Q_{2p}}\,R^{Q_1Q_2}{}_{P_1P_2}\cdots R^{Q_{2p-1}Q_{2p}}{}_{P_{2p-1}P_{2p}}$ (with both indices down it is $1/\sqrt{|g|}$ times the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to g^{MN}; $\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor, $\mathcal{E}_{(3)}$ the cubic Lovelock tensor); with frame indices its frame components. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number (UM-4 keeps only $p = 1$); in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- Λ: The eight-dimensional cosmological constant of UM-4, $\Lambda = -21H^2$: the value for which the static block is a vacuum of Einstein gravity (its Einstein tensor is $21H^2$ in every direction). The author writes the pure number Λ_N in $0 = -\Lambda_\mathrm{N} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$, so that $\Lambda = -\Lambda_\mathrm{N} H^2$. Unit: GeV^2. Value: $-21H^2$; the author's original $-10H^2$; UM-3: $-7H^2$. Status: AUTHOR (the term), DERIVED (the value).
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1M_2\cdots}_{N_1N_2\cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\hat{G}^\mu{}_\nu$: The Einstein tensor of the metric $\hat{g}_{ab}$ of the fold (FD-2), with frame indices along its seven directions; with four-dimensional indices, $\hat{G}^\mu{}_\nu$, its components along the four observable directions, which for frozen extra times are those of the Einstein tensor of the four-dimensional metric $\hat{h}_{\mu\nu}$. Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- $\hat{R}$: The curvature of the metric $\hat{g}_{ab}$ of the fold (FD-2): with four frame indices its Riemann tensor, with two its Ricci tensor $\hat{R}_{ab}$, without indices its scalar curvature $\hat{R}$; for frozen extra times the scalar curvature equals that of the four-dimensional metric $\hat{h}_{\mu\nu}$. Unit: $1/\mathrm{length}^2$. Status: STANDARD.

The bulk equations of the metric of UM-4 with Einstein gravity and $\Lambda = -21H^2$, exactly, for any $a_4(x_4)$ and $b_4(x_4)$: the metric of the author's form is a vacuum of the bulk exactly when the metric of the fold is Ricci flat. Status: DERIVED (`um4_geometry.py`, G5).

$$K^a{}_b = \frac{1}{2}\, g^{ac}\, \partial_\sigma g_{cb} = -H\, \mathrm{sgn}(\sigma)\, \delta^a{}_b, \qquad \kappa\, \mathcal{S}^a{}_b = -12H\, \delta^a{}_b \quad \big(\kappa\lambda_{\mathrm{F}} = 12H\big) \tag{FE-30}$$

Symbols in (FE-30).

- $K^a{}_b$: The extrinsic curvature of a surface of constant σ of the hidden direction, $K^a{}_b = \frac{1}{2} g^{ac} \partial_\sigma g_{cb}$ for a, b along the surface (FD-5): $-H\,\mathrm{sgn}(\sigma)$ along all seven directions of the fold for the author's form of the metric of UM-4; $K = K^c{}_c$ its trace. Unit: GeV (1/length). Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\, \mathrm{d}X^M \mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\, \mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- c: A frame index (as a). Status: STANDARD.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\, \mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma\, \mathcal{E}_{(1)}{}^a{}_b$ (the junction condition of Israel); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -12H\delta^a{}_b$ for every a_4, b_4 of the author's form. Status: DERIVED.
- λ_{F}: The tension that the static block requires of the fold, $\kappa\lambda_{\mathrm{F}} = 12H$: the total vacuum energy of the seven-dimensional fold, made of the tension of the fold fluid and the vacuum energy of the fields of the fold (both have the pressure minus their energy density along ordinary space and along the extra times), per unit volume of the six other directions of the fold (eight-dimensional normalisation); in the four-dimensional normalisation it is $\lambda_{\mathrm{F}} V_*$, and the excess of the total over $\lambda_{\mathrm{F}} V_*$ is the dark energy ρ_Λ. Unit: GeV^7. Value: $12H/\kappa$; $\lambda_{\mathrm{F}} V_* = 30H^2 M_{\mathrm{red}}^2 = 1.1 \times 10^{45}$ GeV^4 for $H = 2534$ GeV. Status: DERIVED.

The fold of the metric of the author's form: the extrinsic curvature is $-H\,\mathrm{sgn}(\sigma)$ in all seven directions, and Israel's junction condition makes the fold a pure tension $\kappa\lambda_{\mathrm{F}} = 12H$ for every a_4, b_4; for a fold of n dimensions in a bulk of $n+1$ the tension is $2(n-1)H$ with $\Lambda = -n(n-1)H^2/2$, which for $n = 4$ is the brane of Randall and Sundrum. Status: DERIVED (`um4_geometry.py`, G7; `um4_foldsources.py`, FS1).

$$a_4,\ b_4 \text{ constant}: \qquad R^{(a)(b)}{}_{(a)(b)} = -H^2 \text{ in all 28 frame planes}, \qquad \mathcal{E}_{(1)}{}^a{}_b = 21H^2\, \delta^a{}_b, \qquad \Lambda = -21H^2 \tag{FE-32}$$

Symbols in (FE-32).

- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.

- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\, \mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.
- $R^{MN}{}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor $R^M{}_{NPQ}$ (in the convention in which the commutator of two covariant derivatives, acting on a vector, is the Riemann tensor, with its last two indices those of the two derivatives, contracted with the vector), with two indices the Ricci tensor $R_{MN} = R^P{}_{MPN}$, without indices the scalar curvature $R = g^{MN} R_{MN}$; with frame indices $R^{(a)(b)}{}_{(c)(d)}$ its frame components, $R^{(a)(b)}{}_{(a)(b)}$ (no sum) the sectional curvature of the frame plane (a, b). Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- $\mathcal{E}_{(p)}{}^M{}_N$: The Lovelock tensor of order p, $\mathcal{E}_{(p)}{}^M{}_N = -2^{-(p+1)}\, \delta^{MP_1\cdots P_{2p}}_{NQ_1\cdots Q_{2p}}\, R^{Q_1Q_2}{}_{P_1P_2} \cdots R^{Q_{2p-1}Q_{2p}}{}_{P_{2p-1}P_{2p}}$ (with both indices down it is $1/\sqrt{|g|}$ times the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to g^{MN}; $\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor, $\mathcal{E}_{(3)}$ the cubic Lovelock tensor); with frame indices its frame components. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number (UM-4 keeps only $p = 1$); in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1M_2\cdots}_{N_1N_2\cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- Λ: The eight-dimensional cosmological constant of UM-4, $\Lambda = -21H^2$: the value for which the static block is a vacuum of Einstein gravity (its Einstein tensor is $21H^2$ in every direction). The author writes the pure number Λ_{N} in $0 = -\Lambda_{\mathrm{N}} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$, so that $\Lambda = -\Lambda_{\mathrm{N}} H^2$. Unit: GeV^2. Value: $-21H^2$; the author's original $-10H^2$; UM-3: $-7H^2$. Status: AUTHOR (the term), DERIVED (the value).

The static block (a_4 and b_4 constant): a space of constant curvature $-H^2$ in all eight directions, on each side of the fold a region of the space of constant negative curvature of eight dimensions, radius $1/H$ and signature (4, 4) (the analogue of anti-de Sitter space), a vacuum of Einstein gravity with $\Lambda = -21H^2$, with a fold of pure tension: the late universe on the largest scales, and the arrangement of Randall and Sundrum. Status: DERIVED (`um4_geometry.py`, G3, G4).

$$\begin{aligned} &\mathrm{e}^{a_4} = x_4^{\mathfrak{k}_1}, \quad \mathrm{e}^{-a_4} b_4 = x_4^{\mathfrak{k}_2} : \quad 3\mathfrak{k}_1 + 3\mathfrak{k}_2 = 1, \quad 3\mathfrak{k}_1^2 + 3\mathfrak{k}_2^2 = 1, \quad \{\mathfrak{k}_1, \mathfrak{k}_2\} = \Big\{\frac{1+\sqrt{5}}{6}, \frac{1-\sqrt{5}}{6}\Big\}; \\ &\mathsf{V} = 0 : \quad \hat{G}^{(4)}{}_{(4)} = -3a_4'^2 \end{aligned} \tag{FE-33}$$

Symbols in (FE-33).

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\, \mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.

- $\mathfrak{k}_1$, $\mathfrak{k}_2$: The exponents of the Kasner folds (the expanding folds without matter of the author's form of the metric): $\mathrm{e}^{a_4} = x_4^{\mathfrak{k}_1}$ and $\mathrm{e}^{-a_4} b_4 = x_4^{\mathfrak{k}_2}$. Unit: dimensionless. Value: $(1 \pm \sqrt{5})/6$. Status: DERIVED.
- V: The rate of change of the scale factor $\mathrm{e}^{-a_4} b_4$ of the extra times on the fold in units of H, $\mathsf{V} = \mathsf{U} - \mathsf{A}$; zero with frozen extra times. Unit: dimensionless. Status: DERIVED.
- $\hat{G}^{\mu}{}_{\nu}$: The Einstein tensor of the metric $\hat{g}_{ab}$ of the fold (FD-2), with frame indices along its seven directions; with four-dimensional indices, $\hat{G}^{\mu}{}_{\nu}$, its components along the four observable directions, which for frozen extra times are those of the Einstein tensor of the four-dimensional metric $\hat{h}_{\mu\nu}$. Unit: 1/length^2. Status: STANDARD.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.

The other vacua of the author's form: the Ricci-flat metrics of the fold are the static fold and two Kasner folds, in which ordinary space and the extra times expand and contract with powers of the time. With frozen extra times the only Ricci-flat fold is static: an expanding fold with frozen extra times needs matter, and with matter its bulk metric is not of the author's form (FE-34). Status: DERIVED (`um4_geometry.py`, G6; `um4_background.py`, BG1).

$$a_4 = Hx_4, \quad b_4 = \text{constant } (\mathsf{A} = 1,\ \mathsf{U} = 0): \qquad \hat{R}_{ab} \neq 0, \qquad \kappa_7\rho_7 = -3H^2, \quad \kappa_7 P_7 = \kappa_7 q_7 = -3H^2 \qquad \text{(FE-31)}$$

Symbols in (FE-31).

- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\, \mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); the first-order equations of the fold hold for $\mathsf{A}^2 \ll 1$; 1 at the author's vacuum, which is not a solution of UM-4. Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- $\hat{R}$: The curvature of the metric $\hat{g}_{ab}$ of the fold (FD-2): with four frame indices its Riemann tensor, with two its Ricci tensor $\hat{R}_{ab}$, without indices its scalar curvature $\hat{R}$; for frozen extra times the scalar curvature equals that of the four-dimensional metric $\hat{h}_{\mu\nu}$. Unit: 1/length^2. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- κ_7: The gravitational coupling of the fold: to the first order in its curvature over H^2 the fold obeys the Einstein equations of seven dimensions $\hat{G}^a{}_b = \kappa_7 \mathsf{t}^a{}_b$ with $\kappa_7 = 5\kappa H/2$, the coupling of the massless graviton of the bulk bound to the fold; four-dimensional gravity on the fold has $8\pi G_{\mathrm{N}} = \kappa_7/V_*$ (plus the induced term, which is negligible). Unit: GeV^{-5}. Value: $5\kappa H/2$. Status: DERIVED.
- ρ_7: The energy density of the matter of the fold above its tension in the seven-dimensional normalisation (minus the component $\mathsf{t}^{(4)}{}_{(4)}$: an energy per unit volume of the six other directions of the fold); $V_*\rho_7$ in the four-dimensional normalisation. Unit: GeV^7. Status: DERIVED.
- P_7: The pressure along ordinary space of the matter of the fold above its tension in the seven-dimensional normalisation (the component $\mathsf{t}^{(i)}{}_{(i)}$). Unit: GeV^7. Status: DERIVED.
- q_7: The pressure along the extra times of the matter of the fold above its tension in the seven-dimensional normalisation (the component $\mathsf{t}^{(r)}{}_{(r)}$). Unit: GeV^7. Status: DERIVED.

The author's vacuum ($a_4 = Hx_4$ with constant b_4), which the earlier versions of the model took as the primordial era, is not a solution of UM-4: the metric of its fold is not Ricci flat, and as a fold with matter it would need a negative energy density, at a rate outside the regime in which the equations of the fold hold. Status: DERIVED (`um4_primordial.py`, PR0).

$$\begin{aligned} \hat{G}^{(i)}{}_{(i)} &= -\big(2a_4'' + 3\mathsf{c}'' + 3a_4'^2 + 6a_4'\mathsf{c}' + 6\mathsf{c}'^2\big), \qquad \hat{G}^{(4)}{}_{(4)} = -\big(3a_4'^2 + 9a_4'\mathsf{c}' + 3\mathsf{c}'^2\big), \\ \hat{G}^{(r)}{}_{(r)} &= -\big(3a_4'' + 2\mathsf{c}'' + 6a_4'^2 + 6a_4'\mathsf{c}' + 3\mathsf{c}'^2\big), \qquad \mathsf{c} = \ln\big(\mathrm{e}^{-a_4} b_4\big) \end{aligned} \tag{FE-66}$$

Symbols in (FE-66).

- $\hat{G}^{\mu}{}_{\nu}$: The Einstein tensor of the metric $\hat{g}_{ab}$ of the fold (FD-2), with frame indices along its seven directions; with four-dimensional indices, $\hat{G}^{\mu}{}_{\nu}$, its components along the four observable directions, which for frozen extra times are those of the Einstein tensor of the four-dimensional metric $\hat{h}_{\mu\nu}$. Unit: 1/length^2. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal), and in a tensor of ordinary space such as h_{ij} one of the directions x_1, x_2, x_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- c: The logarithm of the scale factor of the extra times on the fold, $\mathsf{c} = \ln(\mathrm{e}^{-a_4} b_4)$; $\mathsf{c}' = \mathsf{V}H$; frozen extra times have $\mathsf{c}' = 0$. Unit: dimensionless. Status: DERIVED.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\, \mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

The Einstein tensor of the metric of the fold, written with the logarithm c of the scale factor of the extra times. Status: DERIVED (`um4_background.py`, BG2).

4.1 The bulk metric of an expanding fold and the equations of the fold

$$\begin{aligned} g_{MN}\, \mathrm{d}X^M \mathrm{d}X^N &= \mathrm{d}\sigma^2 + \mathrm{e}^{-2H|\sigma|}\, \tilde{g}_{ab}\, \mathrm{d}x^a \mathrm{d}x^b, \\ \tilde{g}_{ab} &= \hat{g}_{ab} - \frac{\mathrm{e}^{2H|\sigma|} - 1}{5H^2}\Big(\hat{R}_{ab} - \frac{\hat{R}}{12}\, \hat{g}_{ab}\Big) + \mathrm{e}^{7H|\sigma|}\, \mathfrak{c}_{ab} + \ldots \end{aligned} \tag{FE-34}$$

Symbols in (FE-34).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\, \mathrm{d}X^M \mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\, \mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\, \mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- $\tilde{g}_{ab}$: The metric of the slice at the depth σ divided by the warp factor $\mathrm{e}^{-2H|\sigma|}$: equal to the metric of the fold at $\sigma = 0$. Unit: dimensionless. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- x: The coordinates of the fold: x_1, x_2, x_3 of ordinary space, x_4 the time, x_5, x_6, x_7 the extra times; a field of the fold is a function of the four observable coordinates $x = (x_1, x_2, x_3, x_4)$, and $\mathrm{d}^4x = \mathrm{d}x_1\,\mathrm{d}x_2\,\mathrm{d}x_3\,\mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.
- $\hat{g}_{ab}$: The metric of the seven-dimensional fold (FD-2) in its coordinates $x_1, \ldots, x_7$ (or in its frame), $\hat{g}^{ab}$ its inverse; $\hat{h}_{\mu\nu}$ is its part along the four observable directions. Unit: dimensionless. Status: AUTHOR.
- $\hat{R}$: The curvature of the metric $\hat{g}_{ab}$ of the fold (FD-2): with four frame indices its Riemann tensor, with two its Ricci tensor $\hat{R}_{ab}$, without indices its scalar curvature $\hat{R}$; for frozen extra times the scalar curvature equals that of the four-dimensional metric $\hat{h}_{\mu\nu}$. Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- $\mathfrak{c}_{ab}$: A traceless conserved tensor of the fold that sets the homogeneous part of the bulk metric of an expanding fold: the Weyl tensor of the bulk that a field travelling through the bulk can leave (dark radiation); UM-4 takes $\mathfrak{c}_{ab} = 0$. Unit: dimensionless. Value: 0. Status: ASSUMED.

The solution of the bulk equations for a fold with any metric $\hat{g}_{ab}(x)$, expanded in the number of derivatives along the fold (the expansion parameter is the curvature of the fold divided by H^2, for a homogeneous fold A^2): the second term solves every component of the bulk equations to the first order and vanishes at the fold; the third, the bulk Weyl tensor that a field travelling through the bulk can leave (the dark radiation of the brane of Randall and Sundrum), is set to zero. Status: DERIVED (`um4_fold.py`, F1), ASSUMED ($\mathfrak{c}_{ab} = 0$).

$$\kappa\,\mathcal{S}^a{}_b = -12H\,\delta^a{}_b + \frac{2}{5H}\,\hat{G}^a{}_b + \ldots, \qquad \mathcal{S}^a{}_b = -\lambda_\mathrm{F}\,\delta^a{}_b + \mathfrak{t}^a{}_b : \qquad \hat{G}^a{}_b = \kappa_7\,\mathfrak{t}^a{}_b, \qquad \kappa_7 = \frac{5}{2}\,\kappa H \qquad \text{(FE-71)}$$

Symbols in (FE-71).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma\,\mathcal{E}_{(1)}{}^a{}_b$ (the junction condition of Israel); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -12H\delta^a{}_b$ for every a_4, b_4 of the author's form. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised

Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.

- $\hat{G}^{\mu}{}_{\nu}$: The Einstein tensor of the metric $\hat{g}_{ab}$ of the fold (FD-2), with frame indices along its seven directions; with four-dimensional indices, $\hat{G}^{\mu}{}_{\nu}$, its components along the four observable directions, which for frozen extra times are those of the Einstein tensor of the four-dimensional metric $\hat{h}_{\mu\nu}$. Unit: 1/length^2. Status: STANDARD.
- $\lambda_{\rm F}$: The tension that the static block requires of the fold, $\kappa\lambda_{\rm F} = 12H$: the total vacuum energy of the seven-dimensional fold, made of the tension of the fold fluid and the vacuum energy of the fields of the fold (both have the pressure minus their energy density along ordinary space and along the extra times), per unit volume of the six other directions of the fold (eight-dimensional normalisation); in the four-dimensional normalisation it is $\lambda_{\rm F} V_*$, and the excess of the total over $\lambda_{\rm F} V_*$ is the dark energy ρ_Λ. Unit: GeV^7. Value: $12H/\kappa$; $\lambda_{\rm F} V_* = 30 H^2 M_{\rm red}^2 = 1.1 \times 10^{45}$ GeV^4 for $H = 2534$ GeV. Status: DERIVED.
- $\mathtt{t}^a{}_b$, $\mathtt{t}$: The matter of the fold above its tension: the thin source minus the tension, $\mathcal{S}^a{}_b = -\lambda_{\rm F}\delta^a{}_b + \mathtt{t}^a{}_b$ (eight-dimensional normalisation); $\mathtt{t} = \mathtt{t}^c{}_c$ its trace; with frozen extra times $\mathtt{t}^a{}_b = \mathrm{diag}(P_7, P_7, P_7, -\rho_7, q_7, q_7, q_7)$. Unit: GeV^7. Status: DERIVED.
- κ_7: The gravitational coupling of the fold: to the first order in its curvature over H^2 the fold obeys the Einstein equations of seven dimensions $\hat{G}^a{}_b = \kappa_7 \mathtt{t}^a{}_b$ with $\kappa_7 = 5\kappa H/2$, the coupling of the massless graviton of the bulk bound to the fold; four-dimensional gravity on the fold has $8\pi G_{\rm N} = \kappa_7/V_*$ (plus the induced term, which is negligible). Unit: GeV^{-5}. Value: $5\kappa H/2$. Status: DERIVED.

The junction condition with the bulk metric FE-34: the thin source is the tension plus $2/(5H)$ times the Einstein tensor of the fold, so the matter of the fold obeys the Einstein equations of seven dimensions with $\kappa_7 = 5\kappa H/2$ and no cosmological constant of its own (for the critical tension). Integrated over the extra times (volume V_*) Newton's constant is $8\pi G_{\rm N} = \kappa_7/V_*$. Status: DERIVED (`um4_fold.py`, F2).

$$\begin{aligned} \hat{G}_{ab} &= \kappa_7\, \mathtt{t}_{ab} + \kappa^2\, \Pi_{ab} - \mathfrak{E}_{ab}, \\ \Pi_{ab} &= -\frac{1}{4}\, \mathtt{t}_{ac}\mathtt{t}^c{}_b + \frac{1}{24}\, \mathtt{t}\, \mathtt{t}_{ab} + \frac{1}{8}\, \hat{g}_{ab}\, \mathtt{t}_{cd}\mathtt{t}^{cd} - \frac{1}{48}\, \hat{g}_{ab}\, \mathtt{t}^2, \qquad \mathtt{t} = \mathtt{t}^c{}_c \end{aligned} \tag{FE-35}$$

Symbols in (FE-35).

- $\hat{G}^{\mu}{}_{\nu}$: The Einstein tensor of the metric $\hat{g}_{ab}$ of the fold (FD-2), with frame indices along its seven directions; with four-dimensional indices, $\hat{G}^{\mu}{}_{\nu}$, its components along the four observable directions, which for frozen extra times are those of the Einstein tensor of the four-dimensional metric $\hat{h}_{\mu\nu}$. Unit: 1/length^2. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- κ_7: The gravitational coupling of the fold: to the first order in its curvature over H^2 the fold obeys the Einstein equations of seven dimensions $\hat{G}^a{}_b = \kappa_7 \mathtt{t}^a{}_b$ with $\kappa_7 = 5\kappa H/2$, the coupling of the massless graviton of the bulk bound to the fold; four-dimensional gravity on the fold has $8\pi G_{\rm N} = \kappa_7/V_*$ (plus the induced term, which is negligible). Unit: GeV^{-5}. Value: $5\kappa H/2$. Status: DERIVED.
- $\mathtt{t}^a{}_b$, $\mathtt{t}$: The matter of the fold above its tension: the thin source minus the tension, $\mathcal{S}^a{}_b = -\lambda_{\rm F}\delta^a{}_b + \mathtt{t}^a{}_b$ (eight-dimensional normalisation); $\mathtt{t} = \mathtt{t}^c{}_c$ its trace; with frozen extra times $\mathtt{t}^a{}_b = \mathrm{diag}(P_7, P_7, P_7, -\rho_7, q_7, q_7, q_7)$. Unit: GeV^7. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- Π_{ab}: The correction of high energies in the exact equations of the fold, quadratic in the matter $\mathtt{t}_{ab}$ of the fold. Unit: GeV^{14}. Status: DERIVED.
- $\mathfrak{E}_{ab}$: The projection of the Weyl tensor of the bulk onto the fold (its electric part), which enters the exact equations of the fold; zero for a bulk of constant curvature, and not fixed by the fold alone. Unit: 1/length^2. Status: DERIVED.
- c: A frame index (as a). Status: STANDARD.
- d: A frame index (as a). Status: STANDARD.
- $\hat{g}_{ab}$: The metric of the seven-dimensional fold (FD-2) in its coordinates $x_1, \ldots, x_7$ (or in its frame), $\hat{g}^{ab}$ its inverse; $\hat{h}_{\mu\nu}$ is its part along the four observable directions. Unit: dimensionless. Status: AUTHOR.

The exact equations of the fold in a bulk of constant curvature (the method of Shiromizu, Maeda and Sasaki in eight dimensions): the term $\kappa^2\Pi_{ab}$, quadratic in the matter of the fold, is the correction of high energies, and $\mathfrak{E}_{ab}$ the projected Weyl tensor of the bulk. Status: DERIVED (`um4_fold.py`, F3).

$$\lambda_{\mathrm{F}}V_* = 30H^2M_{\mathrm{red}}^2, \qquad \frac{\kappa^2\rho_7^2}{\kappa_7\rho_7} = \frac{24}{5}\,\frac{\rho}{\lambda_{\mathrm{F}}V_*} = \frac{12}{25}\,\mathsf{A}^2 \qquad \left(\rho = 3M_{\mathrm{red}}^2 a_4'^2\right) \tag{S-1}$$

Symbols in (S-1).

- λ_{F}: The tension that the static block requires of the fold, $\kappa\lambda_{\mathrm{F}} = 12H$: the total vacuum energy of the seven-dimensional fold, made of the tension of the fold fluid and the vacuum energy of the fields of the fold (both have the pressure minus their energy density along ordinary space and along the extra times), per unit volume of the six other directions of the fold (eight-dimensional normalisation); in the four-dimensional normalisation it is $\lambda_{\mathrm{F}}V_*$, and the excess of the total over $\lambda_{\mathrm{F}}V_*$ is the dark energy ρ_Λ. Unit: GeV^7. Value: $12H/\kappa$; $\lambda_{\mathrm{F}}V_* = 30H^2M_{\mathrm{red}}^2 = 1.1\times10^{45}$ GeV^4 for $H = 2534$ GeV. Status: DERIVED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- M_{red}: The reduced Planck mass, $M_{\mathrm{red}} = (8\pi G_{\mathrm{N}})^{-1/2}$. Unit: GeV. Value: 2.435×10^{18} GeV. Status: STANDARD.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- ρ_7: The energy density of the matter of the fold above its tension in the seven-dimensional normalisation (minus the component $\mathsf{t}^{(4)}{}_{(4)}$: an energy per unit volume of the six other directions of the fold); $V_*\rho_7$ in the four-dimensional normalisation. Unit: GeV^7. Status: DERIVED.
- κ_7: The gravitational coupling of the fold: to the first order in its curvature over H^2 the fold obeys the Einstein equations of seven dimensions $\hat{G}^a{}_b = \kappa_7\mathsf{t}^a{}_b$ with $\kappa_7 = 5\kappa H/2$, the coupling of the massless graviton of the bulk bound to the fold; four-dimensional gravity on the fold has $8\pi G_{\mathrm{N}} = \kappa_7/V_*$ (plus the induced term, which is negligible). Unit: GeV^{-5}. Value: $5\kappa H/2$. Status: DERIVED.
- ρ: The total energy density of the matter of the fold (all species); $\Delta\rho$ its perturbation. Unit: GeV^4. Status: STANDARD.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); the first-order equations of the fold hold for $\mathsf{A}^2 \ll 1$; 1 at the author's vacuum, which is not a solution of UM-4. Status: DERIVED.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.

The size of the correction of high energies. The tension in the four-dimensional normalisation is $30H^2M_{\mathrm{red}}^2$ (M_{red} the reduced Planck mass; 1.1×10^{45} GeV^4 for $H = 2534$ GeV), and the quadratic term relative to the linear one is $12/25$ times the square of the expansion rate in units of H: today $(H_4^{(0)}/H)^2 = 3\times10^{-91}$ for $H = 2534$ GeV, and it stays below 0.005 ($\mathsf{A} < 0.1$) as long as the energy density of the fold is below $(3.3\times10^{10}\ \mathrm{GeV})^4$ for $H = 2534$ GeV (the primordial era, below). The late universe of UM-4 therefore expands as in general relativity. With frozen extra times the divergence of Π_{ab} does not vanish for matter other than a vacuum energy, radiation and stiff matter: for a fluid with the pressure $w\rho$ along ordinary space and the pressure along the extra times that the radion condition requires, it is $a_4'\rho^2$ times $-3(w-1)(w+1)(3w-1)/32$, so that for dust the projected Weyl tensor $\mathfrak{E}_{ab}$ must be non-zero at the order $\rho^2/(\lambda_{\mathrm{F}}V_*)$, 10^{-91} of the matter today; its value needs the bulk solution and is OPEN. Status: DERIVED (`um4_fold.py`, F5 and F6), OPEN ($\mathfrak{E}_{ab}$).

4.2 The volume of the extra times

$$\mathsf{c}' = 0: \quad 3a_4'^2 = \kappa_7\rho_7, \quad -\big(2a_4'' + 3a_4'^2\big) = \kappa_7 P_7, \quad -3\big(a_4'' + 2a_4'^2\big) = \kappa_7 q_7 \quad \Rightarrow \quad q_7 = -\frac{\rho_7 - 3P_7}{2};$$
$$q_7 = P_7 = 0,\ \mathsf{c}' = 0: \ \mathsf{c}'' = \frac{3}{5}\,a_4'^2 \tag{FE-72}$$

Symbols in (FE-72).

- c: The logarithm of the scale factor of the extra times on the fold, $\mathsf{c} = \ln(\mathrm{e}^{-a_4}b_4)$; $\mathsf{c}' = \mathsf{V}H$; frozen extra times have $\mathsf{c}' = 0$. Unit: dimensionless. Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- κ_7: The gravitational coupling of the fold: to the first order in its curvature over H^2 the fold obeys the Einstein equations of seven dimensions $\hat{G}^a{}_b = \kappa_7 \mathsf{t}^a{}_b$ with $\kappa_7 = 5\kappa H/2$, the coupling of the massless graviton of the bulk bound to the fold; four-dimensional gravity on the fold has $8\pi G_\mathrm{N} = \kappa_7/V_*$ (plus the induced term, which is negligible). Unit: GeV^{-5}. Value: $5\kappa H/2$. Status: DERIVED.
- ρ_7: The energy density of the matter of the fold above its tension in the seven-dimensional normalisation (minus the component $\mathsf{t}^{(4)}{}_{(4)}$: an energy per unit volume of the six other directions of the fold); $V_*\rho_7$ in the four-dimensional normalisation. Unit: GeV^7. Status: DERIVED.
- P_7: The pressure along ordinary space of the matter of the fold above its tension in the seven-dimensional normalisation (the component $\mathsf{t}^{(i)}{}_{(i)}$). Unit: GeV^7. Status: DERIVED.
- q_7: The pressure along the extra times of the matter of the fold above its tension in the seven-dimensional normalisation (the component $\mathsf{t}^{(r)}{}_{(r)}$). Unit: GeV^7. Status: DERIVED.

The fold with frozen extra times: the Friedmann equations of four-dimensional general relativity and the radion condition along the extra times. Radiation satisfies the radion condition by itself; dust and a vacuum energy do not, and without a pressure along the extra times they drive the volume of the extra times. Status: DERIVED (`um4_fold.py`, F4; `um4_background.py`, BG2).

$$g_1^2 = \frac{3H\mathsf{g}_1^2}{2V_*}, \qquad g_2^2 = \frac{3H\mathsf{g}_2^2}{2V_*}, \qquad g_3^2 = \frac{3H\mathsf{g}_3^2}{2V_*}, \qquad 8\pi G_\mathrm{N} = \frac{\kappa_7}{V_*},$$
$$g_{1,\mathrm{eff}} = \frac{g_1}{\sqrt{\mathcal{Z}}}, \quad g_{2,\mathrm{eff}} = \frac{g_2}{\sqrt{\mathcal{Z}}}, \quad g_{3,\mathrm{eff}} = \frac{g_3}{\sqrt{\mathcal{Z}}}, \qquad \lambda_\mathrm{eff} = \frac{\lambda}{\mathcal{Z}}, \qquad \mathcal{Y}_\mathrm{eff} = \frac{\mathcal{Y}}{\sqrt{\mathcal{Z}}}, \qquad G_{\mathrm{N,eff}} = \frac{G_\mathrm{N}}{\mathcal{Z}}, \tag{FE-44}$$
$$\mathcal{Z} = \frac{V_\mathrm{t}\big(\mathrm{e}^{-a_4}b_4\big)^3}{V_*}$$

Symbols in (FE-44).

- g_1: The $U(1)_Y$ gauge coupling of the four-dimensional world (usually g'; not normalised for grand unification): the coupling of the massless mode of the hypercharge field of the bulk, $g_1^2 = 3H\mathsf{g}_1^2/(2V_*)$. Unit: dimensionless. Value: $g_1(m_Z) = 0.3574$. Status: STANDARD.
- g_2: The $SU(2)_L$ gauge coupling of the four-dimensional world (usually g): the coupling of the massless mode of the $SU(2)_L$ field of the bulk, $g_2^2 = 3H\mathsf{g}_2^2/(2V_*)$. Unit: dimensionless. Value: $g_2(m_Z) = 0.6517$. Status: STANDARD.
- g_3: The $SU(3)_C$ gauge coupling of the four-dimensional world (the strong coupling g_s): the coupling of the massless mode of the gluon field of the bulk, $g_3^2 = 3H\mathsf{g}_3^2/(2V_*)$. Unit: dimensionless. Value: $g_3(m_Z) = 1.2177$ ($g_3^2/(4\pi) = 0.1180$). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- g_1, g_2, g_3: The gauge couplings of eight dimensions of $U(1)_Y$, $SU(2)_L$ and $SU(3)_C$: the coefficients of the kinetic terms of the gauge fields of the bulk in their geometric normalisation; the four-dimensional couplings are $g_k^2 = 3H\mathsf{g}_k^2/(2V_*)$. Unit: GeV^{-2} (g^2: length^4). Status: ASSUMED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the

eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.

- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_N: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_\mathrm{N} = 1/M_\mathrm{Pl}^2$ with the Planck mass $M_\mathrm{Pl} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- κ_7: The gravitational coupling of the fold: to the first order in its curvature over H^2 the fold obeys the Einstein equations of seven dimensions $\hat{G}^a{}_b = \kappa_7 \mathsf{t}^a{}_b$ with $\kappa_7 = 5\kappa H/2$, the coupling of the massless graviton of the bulk bound to the fold; four-dimensional gravity on the fold has $8\pi G_\mathrm{N} = \kappa_7/V_*$ (plus the induced term, which is negligible). Unit: GeV^{-5}. Value: $5\kappa H/2$. Status: DERIVED.
- $g_{k,\mathrm{eff}}$: The effective (canonically normalised) gauge couplings of the fold when the fold factor is $\mathcal{Z}$, $g_{k,\mathrm{eff}} = g_k/\sqrt{\mathcal{Z}}$ ($k = 1, 2, 3$): the kinetic terms of the massless modes of the gauge fields of the bulk carry the volume of the extra times as the terms of the fold do. Unit: dimensionless. Status: DERIVED.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4) = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold and the kinetic terms of the massless modes of the fields of the bulk; $\mathcal{Z} = 1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- λ_eff: The effective quartic Higgs coupling, $\lambda/\mathcal{Z}$. Unit: dimensionless. Status: DERIVED.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- $\mathcal{Y}_\mathrm{eff}$: The effective Yukawa matrices, $\mathcal{Y}/\sqrt{\mathcal{Z}}$ (for each of $\mathcal{Y}^u, \mathcal{Y}^d, \mathcal{Y}^e, \mathcal{Y}^\nu$). Unit: dimensionless. Status: DERIVED.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^\nu$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- $G_\mathrm{N,eff}$: The effective Newton constant of the fold, $G_\mathrm{N}/\mathcal{Z}$. Unit: GeV^{-2}. Status: DERIVED.
- V_t: The coordinate volume of the (compact) extra times, $V_\mathrm{t} = \int \mathrm{d}x_5\, \mathrm{d}x_6\, \mathrm{d}x_7$. Unit: length^3. Status: ASSUMED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.

The four-dimensional couplings: the gauge couplings of the massless modes of the bulk, Newton's constant of the massless graviton, and the factor $\mathcal{Z}$ that the volume of the extra times gives every coupling of the fold. Status: DERIVED.

$$\frac{\mathrm{d}\ln\mathcal{Z}}{\mathrm{d}x_4} = 3\mathsf{c}', \qquad q_7 = P_7 = 0,\ \mathsf{c}' = 0:\ \mathsf{c}'' = \frac{3}{5}\,a_4'^2, \qquad q_7 = P_7 = -\rho_7,\ \mathsf{c}' = 0:\ \mathsf{c}'' = \frac{6}{5}\,a_4'^2 \tag{S-2}$$

Symbols in (S-2).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4) = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold and the kinetic terms of the massless modes of the fields of the bulk; $\mathcal{Z} = 1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- c: The logarithm of the scale factor of the extra times on the fold, $\mathsf{c} = \ln(\mathrm{e}^{-a_4}b_4)$; $\mathsf{c}' = \mathsf{V}H$; frozen extra times have $\mathsf{c}' = 0$. Unit: dimensionless. Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- q_7: The pressure along the extra times of the matter of the fold above its tension in the seven-dimensional normalisation (the component $\mathsf{t}^{(r)}{}_{(r)}$). Unit: GeV^7. Status: DERIVED.
- P_7: The pressure along ordinary space of the matter of the fold above its tension in the seven-dimensional normalisation (the component $\mathsf{t}^{(i)}{}_{(i)}$). Unit: GeV^7. Status: DERIVED.

- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- ρ_7: The energy density of the matter of the fold above its tension in the seven-dimensional normalisation (minus the component $\mathsf{t}^{(4)}{}_{(4)}$: an energy per unit volume of the six other directions of the fold); $V_*\rho_7$ in the four-dimensional normalisation. Unit: GeV^7. Status: DERIVED.

A free volume is excluded. The factor $\mathcal{Z}$ changes at three times the rate of the scale factor of the extra times, so every coupling of the fold changes when the extra times do. Without a pressure along the extra times (dust: the pressure of a species along the extra times is its mean on-shell Lagrangian, FE-5, which vanishes for free particles, for a Dirac field because its Lagrangian vanishes on its equation of motion and for a free scalar because the averages of its kinetic and mass terms are equal; a vacuum energy: its pressure $-\rho_7$ along them, its on-shell Lagrangian) the volume accelerates (S-2). Integrated through a matter era from $\mathsf{c}' = 0$, the ratio c'/a_4' reaches 0.752, 0.984, 0.999 and 1.000 after 1, 2, 3 and 5 e-folds (the integration runs over more than 5 e-folds of ordinary space): the attractor is the seven-dimensional dust solution, in which the extra times expand as ordinary space. Newton's constant of the fold, $8\pi G_\mathrm{N} = \kappa_7/(V_\mathrm{t}\mathrm{e}^{3\mathsf{c}})$, then changes at the rate $-3\mathsf{c}' = -3.00H_4$; lunar laser ranging measures $(-5.0 \pm 9.6) \times 10^{-15}$ per year (L. Biskupek, J. Müller and J.-M. Torre, Universe 7 (2021) 34), which allows at most $3.5 \times 10^{-4} H_4^{(0)}$ (two standard deviations, $H_4^{(0)} = 67.4$ km/s/Mpc): a free volume is excluded by a factor of 8545 (for the baryons the pressure along the extra times is ASSUMED to vanish as well). Newton's constant, the squares of the gauge and Yukawa couplings and the quartic Higgs coupling would all fall like $1/\mathcal{Z}$. The volume of the extra times must be held fixed. Status: DERIVED (`um4_background.py`, BG3 and BG4), STANDARD (the measurement).

$$
\begin{aligned}
&H_4^2 = \frac{8\pi G_\mathrm{N}}{3}\big(\rho_\mathrm{SM} + \rho_\chi + \rho_\Lambda\big), \qquad 2H_4' + 3H_4^2 = -8\pi G_\mathrm{N}\,\mathcal{P}, \qquad \frac{1}{8\pi G_\mathrm{N}} = \frac{V_*}{\kappa_7} + \frac{1}{\kappa_\mathrm{ind}}, \qquad \kappa_7 = \frac{5}{2}\,\kappa H, \\
&\rho_\Lambda = \rho_\mathrm{ff} + \rho_\mathrm{vac} - \lambda_\mathrm{F} V_*, \qquad q_\mathrm{SM} + q_\chi + q_\mathrm{P} + q_\mathrm{ff} + \lambda_\mathrm{F} V_* + \frac{\rho - 3\mathcal{P}}{2} = 0, \\
&q_\mathrm{SM} = \big\langle \mathcal{L}_\mathrm{H} + \mathcal{L}_\mathrm{f} + \mathcal{L}_\mathrm{Y} \big\rangle, \qquad q_\mathrm{P} = \langle \mathcal{L}_\mathrm{P} \rangle
\end{aligned}
\tag{FE-59}
$$

Symbols in (FE-59).

- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_N: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_\mathrm{N} = 1/M_\mathrm{Pl}^2$ with the Planck mass $M_\mathrm{Pl} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- ρ_SM: The energy density of the Standard-Model particles. Unit: GeV^4. Status: STANDARD.
- ρ_χ: The energy density of the dark matter, the particles χ of dirac16complex; $\rho_\chi^{(0)}$ its value today. Unit: GeV^4. Status: DERIVED.
- ρ_Λ: The vacuum energy density of the fold above its static tension, the dark energy of UM-4: $\rho_\Lambda = \rho_\mathrm{ff} + \rho_\mathrm{vac} - \lambda_\mathrm{F} V_*$ (four-dimensional normalisation). Unit: GeV^4. Value: $0.685\,\rho_\mathrm{c}^{(0)} = 2.5 \times 10^{-47}$ GeV^4 (Planck 2018). Status: DERIVED.
- $\mathcal{P}$: The total pressure of the matter of the fold; $\Delta\mathcal{P}$ its perturbation. Unit: GeV^4. Status: STANDARD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- κ_7: The gravitational coupling of the fold: to the first order in its curvature over H^2 the fold obeys the Einstein equations of seven dimensions $\hat{G}^a{}_b = \kappa_7 \mathsf{t}^a{}_b$ with $\kappa_7 = 5\kappa H/2$, the coupling of the massless graviton of the bulk bound to the fold; four-dimensional gravity on the fold has $8\pi G_\mathrm{N} = \kappa_7/V_*$ (plus the induced term, which is negligible). Unit: GeV^{-5}. Value: $5\kappa H/2$. Status: DERIVED.
- κ_ind: The coupling of the Einstein-Hilbert term $\hat{R}/(2\kappa_\mathrm{ind})$ that the quantum fluctuations of the fields of the fold may induce, $\kappa_\mathrm{ind} = 8\pi G_\mathrm{ind}$; it adds to the gravity of the fold, $1/(8\pi G_\mathrm{N}) = V_*/\kappa_7 + 1/\kappa_\mathrm{ind}$. Unit: GeV^{-2}. Value: not fixed by UM-4; negligible for a cutoff of the order of H. Status: ESTIMATE.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.

- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- ρ_ff: The energy density of the fold fluid. Unit: GeV^4. Value: $\lambda_\mathrm{F}V_* + \rho_\Lambda - \rho_\mathrm{vac}$: together with the vacuum energy ρ_vac of the fields of the fold it makes up the tension $\lambda_\mathrm{F}V_*$ that the static block requires plus the dark energy ρ_Λ. Status: ASSUMED (the fluid), DERIVED (the value).
- ρ_vac: The vacuum energy of the fields of the fold: the minimum of their potential (for the Standard Model the minimum of the Higgs potential with its loop corrections), a tension of the seven-dimensional fold. Unit: GeV^4. Status: STANDARD.
- λ_F: The tension that the static block requires of the fold, $\kappa\lambda_\mathrm{F} = 12H$: the total vacuum energy of the seven-dimensional fold, made of the tension of the fold fluid and the vacuum energy of the fields of the fold (both have the pressure minus their energy density along ordinary space and along the extra times), per unit volume of the six other directions of the fold (eight-dimensional normalisation); in the four-dimensional normalisation it is $\lambda_\mathrm{F}V_*$, and the excess of the total over $\lambda_\mathrm{F}V_*$ is the dark energy ρ_Λ. Unit: GeV^7. Value: $12H/\kappa$; $\lambda_\mathrm{F}V_* = 30H^2M_\mathrm{red}^2 = 1.1 \times 10^{45}$ GeV^4 for $H = 2534$ GeV. Status: DERIVED.
- q_SM: The pressure of the Standard Model along the extra times at the fold: the mean on-shell Lagrangian density of the Standard-Model fields (the extra-time component of the energy-momentum tensor of every term of the fold is its Lagrangian, because the fields of the fold do not depend on the extra times). Unit: GeV^4. Status: DERIVED (for radiation and free fields), ASSUMED (for the baryons).
- q_χ: The pressure of the particles χ of dirac16complex along the extra times, integrated across the fold (their on-shell Lagrangian, the self-interaction energy per volume; zero for a dilute gas). Unit: GeV^4. Status: DERIVED.
- q_P: The pressure of the portals along the extra times: their mean on-shell Lagrangian density. Unit: GeV^4. Status: DERIVED.
- q_ff: The pressure of the fold fluid along the extra times: minus its energy density (its tension) plus the pressure q_hold that holds the volume of the extra times fixed. Unit: GeV^4. Value: $-\rho_\mathrm{ff} + q_\mathrm{hold}$. Status: ASSUMED (the fluid), DERIVED (the value).
- ρ: The total energy density of the matter of the fold (all species); $\Delta\rho$ its perturbation. Unit: GeV^4. Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- $\mathcal{L}_\mathrm{H}$: The Lagrangian of the Higgs doublet on the fold. Unit: GeV^4. Status: AUTHOR.
- $\mathcal{L}_\mathrm{f}$: The kinetic Lagrangian of the quarks and leptons on the fold. Unit: GeV^4. Status: STANDARD.
- $\mathcal{L}_\mathrm{Y}$: The Yukawa Lagrangian of the quarks and leptons with the Higgs doublet. Unit: GeV^4. Status: STANDARD.
- $\mathcal{L}_\mathrm{P}$: The Lagrangian of the portals of dirac16complex at the fold. Unit: GeV^4. Status: INVENTED.

The four-dimensional equations of the fold with the volume held fixed: the Friedmann equations of general relativity with Newton's constant $1/(8\pi G_\mathrm{N}) = V_*/\kappa_7 + 1/\kappa_\mathrm{ind}$ (the induced term is negligible, below), the dark energy ρ_Λ (the excess of the tension of the fold fluid and of the vacuum energy of the fields over the critical tension), and the radion condition along the extra times, which the fold fluid fulfils. Status: DERIVED.

$$q_\mathrm{hold} = -\Delta_\mathrm{r}, \qquad \Delta_\mathrm{r} = \frac{\rho_\chi}{2} + \frac{\rho_\mathrm{b}}{2} + \rho_\Lambda = (0.1325 + 0.0245 + 0.685)\,\rho_\mathrm{c}^{(0)} = 0.842\,\rho_\mathrm{c}^{(0)} \quad \text{today} \tag{S-3}$$

Symbols in (S-3).

- q_hold: The pressure along the extra times that the fold fluid adds to its tension to hold the volume of the extra times fixed (four-dimensional normalisation). Unit: GeV^4. Value: $-(\rho - 3\mathcal{P})/2 - q_\mathrm{SM} - q_\chi - q_\mathrm{P}$: 0 for radiation, $-\rho/2$ for dust, $-\rho$ for a vacuum energy; today $-0.842\,\rho_\mathrm{c}^{(0)}$ if the baryons have no pressure along the extra times (ASSUMED). Status: DERIVED (the value), ASSUMED (that the fold fluid supplies it).
- Δ_r: The mismatch of the radion condition: the sum over the matter of the fold of its pressure along the extra times and $(\rho - 3\mathcal{P})/2$ (zero for radiation). Unit: GeV^4. Value: today $0.842\,\rho_\mathrm{c}^{(0)}$ if the baryons have no pressure along the extra times (the assumption of UM-4), $0.793\,\rho_\mathrm{c}^{(0)}$ if they have the pressure $-\rho_\mathrm{b}$, $0.891\,\rho_\mathrm{c}^{(0)}$ for $+\rho_\mathrm{b}$. Status: DERIVED (the condition and the values for each choice), ASSUMED (the pressure of the baryons along the extra times).

- ρ_χ: The energy density of the dark matter, the particles χ of dirac16complex; $\rho_\chi^{(0)}$ its value today. Unit: GeV^4. Status: DERIVED.
- ρ_b: The (rest-mass) energy density of the baryons: in a supernova the local density of the matter (quoted in g/cm^3), in cosmology the mean density of the baryons ($\rho_\mathrm{b}^{(0)}$ today). Unit: GeV^4. Status: STANDARD.
- ρ_Λ: The vacuum energy density of the fold above its static tension, the dark energy of UM-4: $\rho_\Lambda = \rho_\mathrm{ff} + \rho_\mathrm{vac} - \lambda_\mathrm{F} V_*$ (four-dimensional normalisation). Unit: GeV^4. Value: $0.685\,\rho_\mathrm{c}^{(0)} = 2.5 \times 10^{-47}$ GeV^4 (Planck 2018). Status: DERIVED.
- ρ_c: The critical density, the energy density of a spatially flat fold at the expansion rate H_4; $\rho_\mathrm{c}^{(0)}$ its value today. Unit: GeV^4. Value: $\rho_\mathrm{c}^{(0)} = 1.0537 \times 10^{-5}\,h_{100}^2$ GeV/cm^3. Status: STANDARD.

What the fold fluid must supply today: the mismatch Δ_r of the radion condition, species by species the pressure along the extra times plus half of $\rho - 3\mathcal{P}$: $\rho_\chi/2$ for the dark matter (a dilute gas, no pressure along the extra times or along ordinary space), $\rho_\mathrm{b}/2$ for the baryons with no pressure along the extra times (ASSUMED: the on-shell Lagrangian of the quark and gluon fields inside a nucleon is not computed), $-\rho_\Lambda + 2\rho_\Lambda$ for the vacuum energy, and zero for radiation; with the densities of Planck 2018 (0.265, 0.049 and 0.685 of the critical density $\rho_\mathrm{c}^{(0)}$) it is $0.842\rho_\mathrm{c}^{(0)}$. If the baryons had the pressure $-\rho_\mathrm{b}$ along the extra times, $\Delta_\mathrm{r} = 0.793\rho_\mathrm{c}^{(0)}$; with $+\rho_\mathrm{b}$, $0.891\rho_\mathrm{c}^{(0)}$. Status: DERIVED (`um4_foldsources.py`, FS5), ASSUMED (the pressure of the baryons along the extra times).

5 The sources of the fold

The kink of the warp at the fold requires a thin source (FE-30), and the gravity of the fold must carry gravitational waves with the speed of light. This section solves both: the gravitational waves of the fold, the gravity that the fields of the fold may induce in addition to that of the bulk, and the fold fluid with the tuning of its tension.

5.1 The gravitational waves of the fold

$$\frac{1}{2}\Big(\frac{\partial^2 h_\mathrm{gw}}{\partial x_4^2} + 3a_4'\,\frac{\partial h_\mathrm{gw}}{\partial x_4} - \mathrm{e}^{-2a_4}\,\frac{\partial^2 h_\mathrm{gw}}{\partial x_3^2}\Big) = \kappa_7\,\Pi_\mathrm{gw}, \qquad \kappa_7 = \frac{5}{2}\,\kappa H, \qquad c_\mathrm{g} = 1 \tag{FE-63}$$

Symbols in (FE-63).

- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- h_gw: The amplitude of a gravitational wave of the + polarisation travelling along x_3: the metric components along x_1 and x_2 multiplied by $\mathrm{e}^{h_\mathrm{gw}}$ and $\mathrm{e}^{-h_\mathrm{gw}}$; a function of σ, x_3 and x_4 in the bulk, and on the fold the gravitational wave of the four-dimensional world, to the first order its plus polarisation h_+. Unit: dimensionless. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- κ_7: The gravitational coupling of the fold: to the first order in its curvature over H^2 the fold obeys the Einstein equations of seven dimensions $\hat{G}^a{}_b = \kappa_7 \mathsf{t}^a{}_b$ with $\kappa_7 = 5\kappa H/2$, the coupling of the massless graviton of the bulk bound to the fold; four-dimensional gravity on the fold has $8\pi G_\mathrm{N} = \kappa_7/V_*$ (plus the induced term, which is negligible). Unit: GeV^{-5}. Value: $5\kappa H/2$. Status: DERIVED.
- Π_gw: The transverse traceless stress of the matter of the fold that drives the gravitational wave h_gw (half the difference of its components along x_1 and x_2), integrated across the fold (eight-dimensional normalisation). Unit: GeV^7. Status: STANDARD.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free

parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.

- c_{g}: The speed of gravitational waves on the fold, in units of the speed of light. Unit: dimensionless. Value: 1 at every wavelength for the four-dimensional graviton on the static block and on every fold with frozen extra times, and for the massless graviton of the bulk at every depth; measured $-3 \times 10^{-15} \leq c_{\mathrm{g}} - 1 \leq 7 \times 10^{-16}$ (GW170817). Status: DERIVED.

The solution. The equations of the fold FE-71 were linearised for a gravitational wave of the + polarisation that travels along x_3 on an expanding fold with frozen extra times (any $a_4(x_4)$, the static block included): half the difference of the components of the Einstein tensor of the fold along x_1 and x_2 is exactly the wave operator of four-dimensional general relativity on the metric $\mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$, with the coupling κ_7 of the Friedmann equations and the transverse traceless stress Π_{gw} of the matter of the fold as its source. The wave is the zero mode of the graviton of the bulk (FE-76 below), which has the same amplitude at every depth: $c_{\mathrm{g}} = 1$ at every wavelength, and the Planck mass of the waves equals that of the expansion. The gravitational waves of the neutron-star merger GW170817 arrived from a distance of about 40 Mpc within 1.7 s of the gamma rays of its burst GRB 170817A, which bounds $-3 \times 10^{-15} \leq c_{\mathrm{g}} - 1 \leq 7 \times 10^{-16}$ (B. P. Abbott et al., Astrophys. J. Lett. 848 (2017) L13): UM-4 satisfies it for every value of H. The massive modes of the bulk graviton add to the propagator of the fold only terms analytic in the momentum (contact terms) and the correction of long range of Newton's law (FE-81 below), and they move at most at the speed of light. Status: DERIVED (`um4_foldsources.py`, FS2; `um4_bulk.py`, B4 and B5), STANDARD (the measurement).

5.2 The gravity induced by the fields of the fold

$$\mathcal{L}_{\mathrm{IG}} = \frac{\hat{R}}{2\kappa_{\mathrm{ind}}} \tag{IN-9}$$

Symbols in (IN-9).

- $\mathcal{L}_{\mathrm{IG}}$: The Lagrangian of the Einstein-Hilbert term of the four-dimensional metric of the fold that the quantum fluctuations of the fields of the fold induce; it adds to the Einstein-Hilbert term that the graviton of the bulk gives the fold and does not change the speed of gravitational waves. Unit: GeV^4. Status: ESTIMATE.
- $\hat{R}$: The curvature of the metric $\hat{g}_{ab}$ of the fold (FD-2): with four frame indices its Riemann tensor, with two its Ricci tensor $\hat{R}_{ab}$, without indices its scalar curvature $\hat{R}$; for frozen extra times the scalar curvature equals that of the four-dimensional metric $\hat{h}_{\mu\nu}$. Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- κ_{ind}: The coupling of the Einstein-Hilbert term $\hat{R}/(2\kappa_{\mathrm{ind}})$ that the quantum fluctuations of the fields of the fold may induce, $\kappa_{\mathrm{ind}} = 8\pi G_{\mathrm{ind}}$; it adds to the gravity of the fold, $1/(8\pi G_{\mathrm{N}}) = V_*/\kappa_7 + 1/\kappa_{\mathrm{ind}}$. Unit: GeV^{-2}. Value: not fixed by UM-4; negligible for a cutoff of the order of H. Status: ESTIMATE.

$$\frac{1}{16\pi G_{\mathrm{ind}}} = \frac{N_{\mathrm{ind}}\,\Lambda_{\mathrm{c}}^2}{192\pi^2}, \qquad N_{\mathrm{ind}} = \sum_{\text{real scalars}} \big(1 - 6\xi\big) + N_{\mathrm{Weyl}} - 4N_{\mathrm{gauge}} = 52 - 24\xi_{\mathrm{H}} \tag{FE-64}$$

Symbols in (FE-64).

- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_{ind}: The Newton constant of the induced Einstein-Hilbert term of the fold, $\kappa_{\mathrm{ind}} = 8\pi G_{\mathrm{ind}}$. Unit: GeV^{-2}. Status: ESTIMATE.
- N_{ind}: The weighted number of the fields of the fold that induce the Einstein-Hilbert term: each real scalar counts $1 - 6\xi$, each Weyl fermion 1, each gauge field -4 (with its ghost fields). Unit: dimensionless. Value: 52 for the fields of the fold of UM-4 (the four real scalars of the Higgs doublet and 48 Weyl fermions; the gauge fields live in the bulk) with $\xi_{\mathrm{H}} = 0$; $52 - 24\xi_{\mathrm{H}}$ with the curvature coupling ξ_{H} of an extension. Status: DERIVED.
- Λ_{c}: The cutoff of the proper-time integral of the one-loop effective action of the fields of the fold: the highest energy of the quantum fluctuations that induce gravity. Unit: GeV. Value: taken of the order of H (2534 GeV); the induced term alone would give the measured Newton constant for $\Lambda_{\mathrm{c}} = \pi M_{\mathrm{red}}\sqrt{96/52} = 1.0 \times 10^{19}$ GeV. Status: ESTIMATE.
- ξ: The coupling of a real scalar field to the scalar curvature $\hat{R}$ of the fold: the term $-\frac{1}{2}\xi\hat{R}$ times the square of the field in its Lagrangian ($\xi = 1/6$ is the conformal coupling). Unit: dimensionless. Status: STANDARD.

- N_{Weyl}: The number of Weyl (two-component) fermion fields of the fold. Unit: dimensionless. Value: 48 (three generations of quarks and leptons with three right-handed neutrinos). Status: STANDARD.
- N_{gauge}: The number of gauge fields of the fold. Unit: dimensionless. Value: 0 in UM-4 (the gauge fields live in the bulk); 12 (eight gluons, three W, one B) if they lived on the fold. Status: STANDARD.
- ξ_{H}: The coupling of the Higgs doublet to the scalar curvature of the fold, the term $-\xi_{\mathrm{H}}\hat{R}\,\Phi^{\dagger}\Phi$ in the Lagrangian of the fold: an extension of UM-4, whose Higgs Lagrangian has no such term ($\xi_{\mathrm{H}}=0$). Unit: dimensionless. Value: 0 in UM-4; about 1.6×10^{4} if the Higgs field drove inflation (ESTIMATE, an extension of the action). Status: ESTIMATE (an extension of the action).

The solution. The one-loop effective action of every field of the fold with a proper-time cutoff Λ_{c} contains an Einstein-Hilbert term; its coefficient follows from the second heat-kernel coefficient of each field: a real scalar with the curvature coupling ξ gives $1-6\xi$, a Weyl fermion 1, a gauge field with its ghost fields -4. In UM-4 the gauge fields live in the bulk, so the fields of the fold are the four real scalars of the Higgs doublet and the 48 Weyl fermions of the quarks and leptons with three right-handed neutrinos: $N_{\mathrm{ind}}=52-24\xi_{\mathrm{H}}$, and the Higgs Lagrangian of UM-4 has no coupling to the curvature of the fold, $\xi_{\mathrm{H}}=0$. With a cutoff of the order of H (the fields of the fold are not described beyond the scale of the bulk), at the lower bound $\Lambda_{\mathrm{c}}=H=2534$ GeV, $1/(8\pi G_{\mathrm{ind}})=5.9\times10^{-32}M_{\mathrm{red}}^{2}$: negligible against the gravity of the bulk graviton, V_{*}/κ_{7}. The same loops give a vacuum energy of about 6.0×10^{12} GeV^4, a contribution to the tension of the fold. The induced term alone would give Newton's constant only for $\Lambda_{\mathrm{c}}=\pi M_{\mathrm{red}}\sqrt{96/52}=1.0\times10^{19}$ GeV, near the Planck mass, where one loop cannot be trusted. Status: ESTIMATE (one loop, cutoff-dependent; `um4_foldsources.py`, FS3).

5.3 The fold fluid

$$\mathcal{T}_{\mathrm{ff}}{}^{a}{}_{b}=\mathrm{diag}\big(p_{\mathrm{ff}},p_{\mathrm{ff}},p_{\mathrm{ff}},-\rho_{\mathrm{ff}},q_{\mathrm{ff}},q_{\mathrm{ff}},q_{\mathrm{ff}}\big) \tag{IN-10}$$

Symbols in (IN-10).

- $\mathcal{T}_{\mathrm{ff}}{}^{a}{}_{b}$: The energy-momentum tensor of the fold fluid (frame components of the fold, four-dimensional normalisation). Unit: GeV^4. Status: ASSUMED.
- a: A frame (orthonormal) index running over $0,1,\ldots,7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab}=\mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a,b\in\{1,\ldots,7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^{a} (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a\in\{1,\ldots,7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0,x_1,\ldots,x_7$ for a tensor of eight dimensions, $x_1,\ldots,x_7$ for a tensor of the fold, $x_1,\ldots,x_4$ for a four-dimensional tensor). Status: STANDARD.
- p_{ff}: The pressure of the fold fluid along ordinary space. Unit: GeV^4. Value: $-\rho_{\mathrm{ff}}$ (a tension). Status: ASSUMED (the fluid), DERIVED (the value).
- ρ_{ff}: The energy density of the fold fluid. Unit: GeV^4. Value: $\lambda_{\mathrm{F}}V_{*}+\rho_{\Lambda}-\rho_{\mathrm{vac}}$: together with the vacuum energy ρ_{vac} of the fields of the fold it makes up the tension $\lambda_{\mathrm{F}}V_{*}$ that the static block requires plus the dark energy ρ_{Λ}. Status: ASSUMED (the fluid), DERIVED (the value).
- q_{ff}: The pressure of the fold fluid along the extra times: minus its energy density (its tension) plus the pressure q_{hold} that holds the volume of the extra times fixed. Unit: GeV^4. Value: $-\rho_{\mathrm{ff}}+q_{\mathrm{hold}}$. Status: ASSUMED (the fluid), DERIVED (the value).

$$\begin{aligned}&p_{\mathrm{ff}}=-\rho_{\mathrm{ff}},\qquad q_{\mathrm{ff}}=-\rho_{\mathrm{ff}}+q_{\mathrm{hold}},\qquad \kappa\lambda_{\mathrm{F}}=12H\quad\big(\lambda_{\mathrm{F}}V_{*}=30H^{2}M_{\mathrm{red}}^{2}\big),\\&q_{\mathrm{hold}}=-\frac{\rho-3\mathcal{P}}{2}-q_{\mathrm{SM}}-q_{\chi}-q_{\mathrm{P}}:\qquad \text{radiation }0,\quad \text{dust }-\frac{\rho}{2},\quad \text{a vacuum energy }-\rho\end{aligned} \tag{FE-65}$$

Symbols in (FE-65).

- p_{ff}: The pressure of the fold fluid along ordinary space. Unit: GeV^4. Value: $-\rho_{\mathrm{ff}}$ (a tension). Status: ASSUMED (the fluid), DERIVED (the value).
- ρ_{ff}: The energy density of the fold fluid. Unit: GeV^4. Value: $\lambda_{\mathrm{F}}V_{*}+\rho_{\Lambda}-\rho_{\mathrm{vac}}$: together with the vacuum energy ρ_{vac} of the fields of the fold it makes up the tension $\lambda_{\mathrm{F}}V_{*}$ that the static block requires plus the dark energy ρ_{Λ}. Status: ASSUMED (the fluid), DERIVED (the value).

- $q_{\rm ff}$: The pressure of the fold fluid along the extra times: minus its energy density (its tension) plus the pressure $q_{\rm hold}$ that holds the volume of the extra times fixed. Unit: GeV^4. Value: $-\rho_{\rm ff}+q_{\rm hold}$. Status: ASSUMED (the fluid), DERIVED (the value).
- $q_{\rm hold}$: The pressure along the extra times that the fold fluid adds to its tension to hold the volume of the extra times fixed (four-dimensional normalisation). Unit: GeV^4. Value: $-(\rho-3\mathcal{P})/2-q_{\rm SM}-q_\chi-q_{\rm P}$: 0 for radiation, $-\rho/2$ for dust, $-\rho$ for a vacuum energy; today $-0.842\,\rho_{\rm c}^{(0)}$ if the baryons have no pressure along the extra times (ASSUMED). Status: DERIVED (the value), ASSUMED (that the fold fluid supplies it).
- κ: The eight-dimensional gravitational coupling, $\kappa=8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\lambda_{\rm F}$: The tension that the static block requires of the fold, $\kappa\lambda_{\rm F}=12H$: the total vacuum energy of the seven-dimensional fold, made of the tension of the fold fluid and the vacuum energy of the fields of the fold (both have the pressure minus their energy density along ordinary space and along the extra times), per unit volume of the six other directions of the fold (eight-dimensional normalisation); in the four-dimensional normalisation it is $\lambda_{\rm F}V_*$, and the excess of the total over $\lambda_{\rm F}V_*$ is the dark energy ρ_Λ. Unit: GeV^7. Value: $12H/\kappa$; $\lambda_{\rm F}V_*=30H^2M_{\rm red}^2=1.1\times10^{45}$ GeV^4 for $H=2534$ GeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda=-21H^2$ and the tension $\kappa\lambda_{\rm F}=12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H>2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- V_*: The physical volume of the extra times x_5,x_6,x_7 on the fold in the late universe, $V_*=V_{\rm t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\rm t}=\int\mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- $M_{\rm red}$: The reduced Planck mass, $M_{\rm red}=(8\pi G_{\rm N})^{-1/2}$. Unit: GeV. Value: 2.435×10^{18} GeV. Status: STANDARD.
- ρ: The total energy density of the matter of the fold (all species); $\Delta\rho$ its perturbation. Unit: GeV^4. Status: STANDARD.
- $\mathcal{P}$: The total pressure of the matter of the fold; $\Delta\mathcal{P}$ its perturbation. Unit: GeV^4. Status: STANDARD.
- $q_{\rm SM}$: The pressure of the Standard Model along the extra times at the fold: the mean on-shell Lagrangian density of the Standard-Model fields (the extra-time component of the energy-momentum tensor of every term of the fold is its Lagrangian, because the fields of the fold do not depend on the extra times). Unit: GeV^4. Status: DERIVED (for radiation and free fields), ASSUMED (for the baryons).
- q_χ: The pressure of the particles χ of dirac16complex along the extra times, integrated across the fold (their on-shell Lagrangian, the self-interaction energy per volume; zero for a dilute gas). Unit: GeV^4. Status: DERIVED.
- $q_{\rm P}$: The pressure of the portals along the extra times: their mean on-shell Lagrangian density. Unit: GeV^4. Status: DERIVED.

The solution for the fold fluid. The source of the fold of the static block, $\kappa\mathcal{S}=-12H$ in all seven directions (FE-30), is a pure tension: its pressures are minus its energy density along ordinary space, the time and the extra times, like the tension of the brane of Randall and Sundrum. The vacuum energy of every field of the fold is a tension of this kind, because the fields do not depend on the extra times and their pressure along the extra times is their on-shell Lagrangian (FE-5), minus the potential at the minimum, the same as along ordinary space. The total tension of the fold fluid and of the vacuum energy of the fields must equal $\lambda_{\rm F}$, $\lambda_{\rm F}V_*=30H^2M_{\rm red}^2$ in the four-dimensional normalisation; its excess over $\lambda_{\rm F}$ is the dark energy (S-4). The component of the equations of the fold along the extra times is the second requirement: the fold fluid must add $q_{\rm hold}$ (S-3) to hold the volume of the extra times fixed (FE-72, S-2). Status: DERIVED (the values; `um4_foldsources.py`, FS1 and FS5), ASSUMED (that the fold fluid supplies $q_{\rm hold}$; the pressure of the baryons along the extra times), OPEN (its microscopic origin: it acts as a potential for the volume of the extra times, as the stabilisation of the sizes of compact extra dimensions does in other theories).

$$\mathcal{S}^a{}_b=-\big(\lambda_{\rm F}+\Delta\lambda_{\rm F}\big)\delta^a{}_b+\mathsf{t}^a{}_b:\qquad \hat{G}^a{}_b=\kappa_7\big(\mathsf{t}^a{}_b-\Delta\lambda_{\rm F}\,\delta^a{}_b\big),\qquad \rho_\Lambda=V_*\,\Delta\lambda_{\rm F},\qquad \frac{\Delta\lambda_{\rm F}}{\lambda_{\rm F}}=\frac{\rho_\Lambda}{30H^2M_{\rm red}^2}=2.2\times10^{-92} \tag{S-4}$$

Symbols in (S-4).

- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b=\int_{0^-}^{0^+}\mathrm{d}\sigma\,\mathcal{E}_{(1)}{}^a{}_b$ (the junction condition of Israel); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its

pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -12H\delta^a{}_b$ for every a_4, b_4 of the author's form. Status: DERIVED.

- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- λ_F: The tension that the static block requires of the fold, $\kappa\lambda_\mathrm{F} = 12H$: the total vacuum energy of the seven-dimensional fold, made of the tension of the fold fluid and the vacuum energy of the fields of the fold (both have the pressure minus their energy density along ordinary space and along the extra times), per unit volume of the six other directions of the fold (eight-dimensional normalisation); in the four-dimensional normalisation it is $\lambda_\mathrm{F}V_*$, and the excess of the total over $\lambda_\mathrm{F}V_*$ is the dark energy ρ_Λ. Unit: GeV^7. Value: $12H/\kappa$; $\lambda_\mathrm{F}V_* = 30H^2M_\mathrm{red}^2 = 1.1 \times 10^{45}$ GeV^4 for $H = 2534$ GeV. Status: DERIVED.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1M_2\cdots}_{N_1N_2\cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- $\mathtt{t}^a{}_b$, $\mathtt{t}$: The matter of the fold above its tension: the thin source minus the tension, $\mathcal{S}^a{}_b = -\lambda_\mathrm{F}\delta^a{}_b + \mathtt{t}^a{}_b$ (eight-dimensional normalisation); $\mathtt{t} = \mathtt{t}^c{}_c$ its trace; with frozen extra times $\mathtt{t}^a{}_b = \mathrm{diag}(P_7, P_7, P_7, -\rho_7, q_7, q_7, q_7)$. Unit: GeV^7. Status: DERIVED.
- $\hat{G}^\mu{}_\nu$: The Einstein tensor of the metric $\hat{g}_{ab}$ of the fold (FD-2), with frame indices along its seven directions; with four-dimensional indices, $\hat{G}^\mu{}_\nu$, its components along the four observable directions, which for frozen extra times are those of the Einstein tensor of the four-dimensional metric $\hat{h}_{\mu\nu}$. Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- κ_7: The gravitational coupling of the fold: to the first order in its curvature over H^2 the fold obeys the Einstein equations of seven dimensions $\hat{G}^a{}_b = \kappa_7\mathtt{t}^a{}_b$ with $\kappa_7 = 5\kappa H/2$, the coupling of the massless graviton of the bulk bound to the fold; four-dimensional gravity on the fold has $8\pi G_\mathrm{N} = \kappa_7/V_*$ (plus the induced term, which is negligible). Unit: GeV^{-5}. Value: $5\kappa H/2$. Status: DERIVED.
- ρ_Λ: The vacuum energy density of the fold above its static tension, the dark energy of UM-4: $\rho_\Lambda = \rho_\mathrm{ff} + \rho_\mathrm{vac} - \lambda_\mathrm{F}V_*$ (four-dimensional normalisation). Unit: GeV^4. Value: $0.685\,\rho_\mathrm{c}^{(0)} = 2.5 \times 10^{-47}$ GeV^4 (Planck 2018). Status: DERIVED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- M_red: The reduced Planck mass, $M_\mathrm{red} = (8\pi G_\mathrm{N})^{-1/2}$. Unit: GeV. Value: 2.435×10^{18} GeV. Status: STANDARD.

The dark energy and the tuning of the tension. A tension above the critical one by $\Delta\lambda_\mathrm{F}$ gives the fold, by FE-71, the cosmological constant $\kappa_7\Delta\lambda_\mathrm{F}$, which in the four-dimensional normalisation is the vacuum energy $\rho_\Lambda = V_*\Delta\lambda_\mathrm{F}$ with the equation of state $w = -1$ (its pressure minus its energy density, exactly). The observed dark energy, $\rho_\Lambda = 2.52 \times 10^{-47}$ $\mathrm{GeV}^4 = (2.24\ \mathrm{meV})^4$, needs the tension tuned to $\Delta\lambda_\mathrm{F}/\lambda_\mathrm{F} = 2.2 \times 10^{-92}$ for $H = 2534$ GeV (2.26×10^{-92} for $H = 2.5$ TeV), smaller still for a larger H: the cosmological-constant problem of the brane of Randall and Sundrum. UM-4 accommodates the value but does not explain it. Status: DERIVED (`um4_foldsources.py`, FS4), OPEN (the tuning).

6 The Higgs field in the ad hoc vacuum

$$\frac{1}{\sqrt{-\hat{h}}\,\mathcal{Z}}\,D_\mu\big(\sqrt{-\hat{h}}\,\mathcal{Z}\,\hat{h}^{\mu\nu}D_\nu\Phi\big)-\mu^2\Phi-2\lambda\big(\Phi^\dagger\Phi\big)\Phi-c_\Phi\,\bar{\Psi}P_{\mathrm{L}}\Psi\,\Phi=\Theta,$$
$$\Theta=\begin{pmatrix}\bar{d}_{\mathrm{R}}(\mathcal{Y}^d)^\dagger u_{\mathrm{L}}+\bar{e}_{\mathrm{R}}(\mathcal{Y}^e)^\dagger\nu_{\mathrm{L}}-\bar{d}_{\mathrm{L}}\mathcal{Y}^u u_{\mathrm{R}}-\bar{e}_{\mathrm{L}}\mathcal{Y}^\nu\nu_{\mathrm{R}}\\ \bar{d}_{\mathrm{R}}(\mathcal{Y}^d)^\dagger d_{\mathrm{L}}+\bar{e}_{\mathrm{R}}(\mathcal{Y}^e)^\dagger e_{\mathrm{L}}+\bar{u}_{\mathrm{L}}\mathcal{Y}^u u_{\mathrm{R}}+\bar{\nu}_{\mathrm{L}}\mathcal{Y}^\nu\nu_{\mathrm{R}}\end{pmatrix}\qquad\text{(FE-15)}$$

Symbols in (FE-15).

- $\hat{h}_{\mu\nu}$: The four-dimensional metric of the fold, $\hat{h}_{\mu\nu}=\mathrm{diag}(\mathrm{e}^{2a_4},\mathrm{e}^{2a_4},\mathrm{e}^{2a_4},-1)$ in the coordinates $x_1,\dots,x_4$ (the induced metric at $\sigma=0$ without the extra times); $\hat{h}^{\mu\nu}$ its inverse, $\hat{h}$ its determinant, $\sqrt{-\hat{h}}=\mathrm{e}^{3a_4}$; $\hat{h}^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4)=V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold and the kinetic terms of the massless modes of the fields of the bulk; $\mathcal{Z}=1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M=\partial_M+\Omega_M$ (with $D_M\bar{\Psi}=\partial_M\bar{\Psi}-\bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu=\hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A=\partial_\mu X^A+g_3f^{ABC}G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1,x_2,x_3,x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- μ^2: The mass parameter of the Higgs potential, $\mu^2<0$ (it enters only as μ^2). Unit: GeV^2. Value: $\mu^2=-\lambda v^2=-m_h^2/2=-(88.6\ \mathrm{GeV})^2$. Status: AUTHOR.
- λ: The quartic coupling of the Higgs potential, $\lambda>0$. Unit: dimensionless. Value: $\lambda=m_h^2/(2v^2)=0.129$. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- Ψ, $\bar{\Psi}$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar{\Psi}=\Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma=0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- P_{L}: The author's left-handed chiral projector, $P_{\mathrm{L}}=(\mathbf{1}-\Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- Θ: The Yukawa source of the Higgs equation, $\Theta=-\partial\mathcal{L}_{\mathrm{Y}}/\partial\Phi^\dagger$ (a doublet of fermion bilinears). Unit: GeV^3. Status: STANDARD.
- d_{R}: The right-handed down-type quarks (down, strange and bottom; hypercharge $-1/3$). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^\nu$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- u_{L}: The upper (up-type, left-handed) components of the quark doublets. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- e_{R}: The right-handed electron, muon and tau (singlets, hypercharge -1). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- ν_{L}: The upper (neutral, left-handed) components of the lepton doublets: the left-handed neutrinos. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- d_{L}: The lower (down-type, left-handed) components of the quark doublets. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- u_{R}: The right-handed up-type quarks (up, charm and top; singlets of $SU(2)_L$, hypercharge 2/3). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- ν_{R}: The three right-handed neutrinos (gauge singlets). Unit: $\mathrm{GeV}^{3/2}$. Status: ASSUMED.
- e_{L}: The lower (charged, left-handed) components of the lepton doublets. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.

$$\frac{1}{\sqrt{|g|}}\,D_M\big(\sqrt{|g|}\,W^{IMN}\big)=0 \qquad (\sigma\neq 0),$$
$$\frac{1}{\mathsf{g}_2^2}\Big(W^{I\sigma\nu}\big|_{0^+}-W^{I\sigma\nu}\big|_{0^-}\Big)=-\frac{1}{V_*}\Big(J_{\mathrm{w}}^{I\nu}+J_\Phi^{I\nu}+\frac{1}{g_2}\,D_\mu\big[(\mathcal{K}_W-1)W^{I\mu\nu}\big]\Big), \tag{FE-17}$$
$$\mathcal{K}_W=1+\sum_X c_X^W\,\bar\Psi P_X\Psi, \qquad J_{\mathrm{w}}^{I\nu}=\bar Q\,\hat\gamma^\nu\frac{\tau^I}{2}Q+\bar\ell\,\hat\gamma^\nu\frac{\tau^I}{2}\ell, \qquad J_\Phi^{I\nu}=\mathrm{i}\Big[(D^\nu\Phi)^\dagger\frac{\tau^I}{2}\Phi-\Phi^\dagger\frac{\tau^I}{2}D^\nu\Phi\Big]$$

Symbols in (FE-17).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\dots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M=\partial_M+\Omega_M$ (with $D_M\bar\Psi=\partial_M\bar\Psi-\bar\Psi\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu=\hat h^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A=\partial_\mu X^A+g_3 f^{ABC}G_\mu^B X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2,\dots$): it runs over $x_0, x_1,\dots,x_7$; a repeated index is summed. Status: STANDARD.
- W_M^I, W_μ^I, W_{MN}^I: The gauge field of $SU(2)_L$ ($I=1,2,3$): in the bulk W_M^I (geometric normalisation), on the fold the canonically normalised four-dimensional field $W_\mu^I(x)$ of its massless mode, $g_2W_\mu^I(x)=W_\mu^I(\sigma=0,x)$; W_{MN}^I, $W_{\mu\nu}^I$ their field strengths; $W_\mu^\pm=(W_\mu^1\mp\mathrm{i}W_\mu^2)/\sqrt{2}$. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- I: A weak-isospin (adjoint SU(2)) index running over $1,2,3$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2,\dots$), running over $x_0,\dots,x_7$; a repeated index is summed. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma=|\cot z|\,\mathrm{d}x_0$, $\sigma=\mathrm{sgn}(\pi/2-z)\ln(\sin z)/(6H)$ on one period $0<z<\pi$; $\sigma=0$ is the fold $z=\pi/2$, $\sigma\to\mp\infty$ the tips $z\to 0,\pi$; one period of x_0 is the whole line $-\infty<\sigma<\infty$. Unit: length. Status: DERIVED.
- g_1, g_2, g_3: The gauge couplings of eight dimensions of $U(1)_Y$, $SU(2)_L$ and $SU(3)_C$: the coefficients of the kinetic terms of the gauge fields of the bulk in their geometric normalisation; the four-dimensional couplings are $g_k^2=3H\mathsf{g}_k^2/(2V_*)$. Unit: GeV^{-2} (g^2: length^4). Status: ASSUMED.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_*=V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\mathrm{t}}=\int\mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- $J_{\mathrm{w}}^{I\nu}$: The weak-isospin current of the left-handed fermion doublets. Unit: GeV^3. Status: STANDARD.
- $J_\Phi^{I\nu}$: The weak-isospin current of the Higgs doublet (it contains the mass terms of the W fields); $J_\Phi^{2\,x_4}$ its isospin charge density with $I=2$. Unit: GeV^3. Status: STANDARD.
- g_2: The $SU(2)_L$ gauge coupling of the four-dimensional world (usually g): the coupling of the massless mode of the $SU(2)_L$ field of the bulk, $g_2^2=3H\mathsf{g}_2^2/(2V_*)$. Unit: dimensionless. Value: $g_2(m_Z)=0.6517$. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- $\mathcal{K}_G$, $\mathcal{K}_W$, $\mathcal{K}_B$: The portal factors of the gauge kinetic terms, $\mathcal{K}_G=1+\sum_X c_X^G\bar\Psi P_X\Psi$ (likewise for W and B): the gauge couplings at the fold are divided by the square root of the portal factor. Unit: dimensionless. Value: 1 where dirac16complex vanishes. Status: DERIVED.
- X: The label of a half of dirac16complex: $X\in\{\mathrm{L},\mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c_X^W: The W-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi=\Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma=0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- P_X: The chiral projector of the half X: $P_X=P_{\mathrm{L}}$ for $X=\mathrm{L}$, P_{R} for $X=\mathrm{R}$. Unit: dimensionless. Status: AUTHOR.
- Q, $\bar Q$: The left-handed quark doublets of the three generations, $Q=(u_{\mathrm{L}},d_{\mathrm{L}})$ (colour triplets, hypercharge 1/6). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.

- $\hat{\gamma}^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^\mu$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^\mu,\hat{\gamma}^\nu\}=-2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_\mu=\hat{h}_{\mu\nu}\hat{\gamma}^\nu$. Unit: dimensionless. Status: STANDARD.
- τ^I: The Pauli matrices acting on weak doublets (τ^1,τ^2,τ^3; τ^2 is the second one, not a square). Unit: dimensionless. Status: STANDARD.
- ℓ, $\bar{\ell}$: The left-handed lepton doublets of the three generations, $\ell=(\nu_\mathrm{L},e_\mathrm{L})$ (hypercharge $-1/2$). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge $1/2$), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.

The Higgs equation on the fold (with the factor $\mathcal{Z}$ of the extra times) and the equation of the $SU(2)$ gauge fields, which in UM-4 live in the bulk: away from the fold they obey the Yang-Mills equation of eight dimensions, and at the fold the jump of their field strength along σ is set by the currents of the fold, the isospin current of the Higgs doublet among them. For homogeneous fields the jump condition is the Gauss law of the fold. Status: DERIVED.

$$\begin{aligned}&\mathcal{R}_\mathcal{A}=\mathcal{A}''+3\frac{b_4'}{b_4}\mathcal{A}'+\lambda\mathcal{A}^3+c_\Phi\,\bar{\Psi}P_\mathrm{L}\Psi\,\mathcal{A}=\Big(4a_{40}^2-6a_{40}\frac{b_4'}{b_4}\Big)\mathcal{A}+\lambda\mathcal{A}^3+c_\Phi\,\bar{\Psi}P_\mathrm{L}\Psi\,\mathcal{A},\\&\mathcal{R}_v=\lambda\mathcal{A}^2v+c_\Phi\,\bar{\Psi}P_\mathrm{L}\Psi\,v,\qquad J_\Phi^{2\,x_4}=a_{40}\,v\,\mathcal{A},\qquad J_\Phi^{1\,x_4}=J_\Phi^{3\,x_4}=0\end{aligned}\tag{FE-26}$$

Symbols in (FE-26).

- $\mathcal{R}_\mathcal{A}$, $\mathcal{R}_v$: The residuals of the ad hoc Higgs vacuum in the upper ($\mathcal{R}_\mathcal{A}$) and lower ($\mathcal{R}_v$) components of the Higgs equation: what the ad hoc vacuum leaves unbalanced. Unit: GeV^3. Status: DERIVED.
- $\mathcal{A}$: The author's ad hoc upper component of the Higgs vacuum, almightyAdHoc, $\mathcal{A}(x_4)=v_\mathcal{A}\mathrm{e}^{-2a_{40}x_4}$. Unit: GeV. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4'=\mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4>0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- λ: The quartic coupling of the Higgs potential, $\lambda>0$. Unit: dimensionless. Value: $\lambda=m_h^2/(2v^2)=0.129$. Status: AUTHOR.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- Ψ, $\bar{\Psi}$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar{\Psi}=\Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma=0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L}=(\mathbf{1}-\Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- a_{40}: The decay constant of almightyAdHoc, $a_{40}>0$: $\mathcal{A}$ falls by e in the time $1/(2a_{40})$. Unit: GeV (1/length). Status: AUTHOR.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2=-\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $=(\sqrt{2}\,G_\mathrm{F})^{-1/2}$ with the Fermi constant $G_\mathrm{F}=1.1663788\times10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- $J_\Phi^{I\nu}$: The weak-isospin current of the Higgs doublet (it contains the mass terms of the W fields); $J_\Phi^{2\,x_4}$ its isospin charge density with $I=2$. Unit: GeV^3. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c=1$). Status: AUTHOR.

The author's ad hoc vacuum $\Phi=(\mathcal{A},v+h)/\sqrt{2}$ with $\mathcal{A}=v_\mathcal{A}\mathrm{e}^{-2a_{40}x_4}$, inserted literally into FE-15 and FE-17 (with $h=0$ and no gauge field): it leaves the residuals $\mathcal{R}_\mathcal{A}$ and $\mathcal{R}_v$ in the two components of the Higgs equation, and it carries a homogeneous charge density of the second isospin direction, which the Gauss law of FE-17 forbids for homogeneous gauge fields. Status: DERIVED.

$$\Phi = \frac{1}{\sqrt{2}}\begin{pmatrix}\mathcal{A}\\ v\end{pmatrix} = \mathcal{O}(\phi_{\rm rot})\,\frac{1}{\sqrt{2}}\begin{pmatrix}0\\ V_{\rm rad}\end{pmatrix}, \qquad V_{\rm rad} = \sqrt{v^2+\mathcal{A}^2}, \qquad \tan\frac{\phi_{\rm rot}}{2} = \frac{\mathcal{A}}{v},$$

$$\mathcal{O}(\phi_{\rm rot}) = \begin{pmatrix}\cos\frac{\phi_{\rm rot}}{2} & \sin\frac{\phi_{\rm rot}}{2}\\ -\sin\frac{\phi_{\rm rot}}{2} & \cos\frac{\phi_{\rm rot}}{2}\end{pmatrix}, \qquad W^2_{x_4} = \frac{\phi'_{\rm rot}}{g_2} = -\frac{4a_{40}\,v\,\mathcal{A}}{g_2\big(v^2+\mathcal{A}^2\big)}, \qquad J^{2\,x_4}_{\Phi} = 0, \tag{S-5}$$

$$\text{FE-15} \;\Leftrightarrow\; \mathcal{O}(\phi_{\rm rot})\,\frac{1}{\sqrt{2}}\begin{pmatrix}0\\ -\mathcal{R}_{\rm rad}\end{pmatrix} = 0, \qquad \mathcal{R}_{\rm rad} = V''_{\rm rad} + 3\,\frac{b'_4}{b_4}\,V'_{\rm rad} + \lambda\,\mathcal{A}^2\,V_{\rm rad} + c_\Phi\,\bar\Psi P_{\rm L}\Psi\,V_{\rm rad}$$

Symbols in (S-5).

- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- $\mathcal{A}$: The author's ad hoc upper component of the Higgs vacuum, almightyAdHoc, $\mathcal{A}(x_4) = v_{\mathcal{A}}\mathrm{e}^{-2a_{40}x_4}$. Unit: GeV. Status: AUTHOR.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_{\rm F})^{-1/2}$ with the Fermi constant $G_{\rm F} = 1.1663788\times10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- $\mathcal{O}(\phi_{\rm rot})$: The 2×2 isospin rotation by the angle $\phi_{\rm rot}$ about the second isospin axis (a real element of $SU(2)$). Unit: dimensionless. Status: DERIVED.
- $V_{\rm rad}$: The modulus of the Higgs doublet in the ad hoc vacuum, $V_{\rm rad} = \sqrt{v^2+\mathcal{A}^2}$ ($V^2_{\rm rad} = 2\Phi^\dagger\Phi$). Unit: GeV. Status: DERIVED.
- $\phi_{\rm rot}$: The angle of the isospin rotation that turns $(0, V_{\rm rad})/\sqrt{2}$ into the ad hoc vacuum: $\tan(\phi_{\rm rot}/2) = \mathcal{A}/v$. Unit: dimensionless. Status: DERIVED.
- W^I_M, W^I_μ, W^I_{MN}: The gauge field of $SU(2)_L$ ($I = 1, 2, 3$): in the bulk W^I_M (geometric normalisation), on the fold the canonically normalised four-dimensional field $W^I_\mu(x)$ of its massless mode, $g_2W^I_\mu(x) = W^I_\mu(\sigma = 0, x)$; W^I_{MN}, $W^I_{\mu\nu}$ their field strengths; $W^\pm_\mu = (W^1_\mu \mp \mathrm{i}W^2_\mu)/\sqrt{2}$. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a'_4 = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- g_2: The $SU(2)_L$ gauge coupling of the four-dimensional world (usually g): the coupling of the massless mode of the $SU(2)_L$ field of the bulk, $g_2^2 = 3H\mathsf{g}_2^2/(2V_*)$. Unit: dimensionless. Value: $g_2(m_Z) = 0.6517$. Status: STANDARD.
- a_{40}: The decay constant of almightyAdHoc, $a_{40} > 0$: $\mathcal{A}$ falls by e in the time $1/(2a_{40})$. Unit: GeV (1/length). Status: AUTHOR.
- $J^{I\nu}_\Phi$: The weak-isospin current of the Higgs doublet (it contains the mass terms of the W fields); $J^{2\,x_4}_\Phi$ its isospin charge density with $I = 2$. Unit: GeV^3. Status: STANDARD.
- $\mathcal{R}_{\rm rad}$: The radial residual of the Higgs equation in the ad hoc vacuum written as a gauge rotation: the amount by which $V_{\rm rad}$ fails to solve the radial equation; zero for a solution. Unit: GeV^3. Status: DERIVED.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: GeV$^{7/2}$. Status: AUTHOR.
- $P_{\rm L}$: The author's left-handed chiral projector, $P_{\rm L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.

The ad hoc vacuum is a rotation of the doublet $(0, V_{\rm rad})/\sqrt{2}$ by the angle $\phi_{\rm rot}(x_4)$ about the second isospin axis. Together with the pure-gauge field $W^2_{x_4} = \phi'_{\rm rot}/g_2$, which undoes the rotation in the covariant derivative, the isospin charge density of the Higgs field vanishes, so the Gauss law of FE-17 holds with homogeneous fields. In UM-4 this $W^2_{x_4}$ is the value at the fold of the field of the bulk: a homogeneous pure gauge, constant along σ (the profile of the zero mode), which is a solution of the bulk equation with no field strength and no jump at the fold. The

masses of the W and Z bosons and of the fermions depend on the doublet only through $\Phi^\dagger\Phi$ ($m_\mathrm{W}^2 = g_2^2(v^2+\mathcal{A}^2)/4$, $m_\mathrm{Z}^2 = (g_1^2+g_2^2)(v^2+\mathcal{A}^2)/4$, each fermion mass $\propto \sqrt{v^2+\mathcal{A}^2}$) and are unchanged by the rotation. The Higgs equation FE-15 then reduces exactly to one radial residual $\mathcal{R}_\mathrm{rad}$ (at 20 random points of the parameters the reduction holds exactly). With $\mu^2 = -\lambda v^2$ the ad hoc vacuum is a solution only in the limit $\mathcal{A} \to 0$, which it reaches exponentially. Status: DERIVED (`um4_background.py`, BG5).

a_{40}/v	$\mathcal{R}_\mathrm{rad}(x_4=0)/(\lambda v^3)$	$\rho_\mathrm{H}(x_4=0)/v^4$	time until $\mathcal{A} < 10^{-3}v$
0.1	1.74	0.042	$34.5/v$
1	34.2	1.03	$3.45/v$
10	3280	100	$0.345/v$

The radial residual at $x_4 = 0$ and the energy density of the Higgs field, $\rho_\mathrm{H} = \frac{1}{2}V_\mathrm{rad}'^2 + \frac{\lambda}{4}\mathcal{A}^4$, for $v_\mathcal{A} = v$, a constant b_4 and no dirac16complex at the fold, for three values of a_{40} (in units of $v = 246.22$ GeV); the ad hoc component is gone after a time of order $3.5/a_{40}$. It must be gone long before nucleosynthesis. Its potential energy $\lambda\mathcal{A}^4/4$ stays below the energy density $3M_\mathrm{Pl}^2/(32\pi x_4^2)$ of the radiation of an era that begins at $x_4 = 0$ only if $a_{40} > v_\mathcal{A}^2\sqrt{\lambda\pi/6}/(\mathrm{e}M_\mathrm{Pl}) = 4.8\times10^{-16}\,(v_\mathcal{A}/v)^2$ GeV $= 7.2\times10^8\,(v_\mathcal{A}/v)^2$ s^{-1} (the ratio of the two is largest at $x_4 = 1/(4a_{40})$; otherwise an epoch dominated by the energy of the prescribed vacuum would interrupt the radiation era, which the ad hoc vacuum does not describe). With such a rate the ad hoc vacuum has no effect on nucleosynthesis. The weak rates alone would allow far slower decays: $v_\mathrm{loc}^2 = v^2+\mathcal{A}^2$ lowers the Fermi constant, $G_\mathrm{F,loc} = 1/(\sqrt{2}v_\mathrm{loc}^2)$ (FE-49), and raises the electron mass and the part of the neutron-proton mass difference that comes from the quark masses (2.52 MeV of the 1.293 MeV; Sz. Borsanyi et al., Science 347 (2015) 1452), both in proportion to v_loc. An estimate of the helium abundance (the neutron fraction X_n from the Born rates of the conversion of neutrons into protons by the capture of neutrinos and positrons and by the decay of the neutron, and of the inverse processes, normalised to the neutron lifetime 878.4 s, with the transfer of the entropy of the electrons and positrons to the photons, from 10 MeV to the deuterium bottleneck at 0.078 MeV, and the helium mass fraction $Y_\mathrm{p} = 2X_\mathrm{n}$ there) gives $Y_\mathrm{p} = 0.2528$ without the ad hoc vacuum (observed 0.245 ± 0.003, Particle Data Group 2024); a constant $\mathcal{A}^2/v^2$ changes it by $+0.350$ (the Fermi constant) and -0.681 (the masses) per unit, and $|\Delta Y_\mathrm{p}| < 0.006$ (twice the error) requires $a_{40} > 1.4$ s^{-1} for $v_\mathcal{A} = v$ (4.4 s^{-1} for $10v$, 22.6 s^{-1} for $10^6 v$; none for $0.1v$), weaker than the bound from the energy by more than 10^8. Status: DERIVED (`um4_background.py`, BG6; `um4_thermal.py`, T6: the bound from the energy, with the numerical maximum of the ratio; the rates obey detailed balance and tend to the free decay), ESTIMATE (the helium abundance of a simple freeze-out; the radiation era from $x_4 = 0$).

7 Propagation in the bulk

In the metric of UM-4 the warp factor multiplies the whole metric of the fold, the time included, so every slice of constant σ of the static block is a copy of the flat fold scaled in all seven directions alike, with the light cone of the fold. For every field of the bulk the dependence on σ separates from that on the seven coordinates of the fold, and the field of the fold obeys the Lorentz-invariant equation of seven dimensions with a mass that its profile fixes. This section solves the profiles of every field of the bulk on the static block ($a_4 = 0$, $b_4 = 1$ by the choice of the units of $x_1, \ldots, x_7$), the photon and the gravitational wave as waves of four dimensions, the propagators of the bulk between points of the fold with Newton's and Coulomb's laws and the W and Z bosons, and the causal structure. (In UM-3, whose time was not warped, the zero modes of bulk scalars and photons moved $\sqrt{3/2} = 1.22$ and $\sqrt{2} = 1.41$ times as fast as light on the fold at long wavelengths; `um4_bulk.py`, B9, recomputes these numbers.)

7.1 The modes of the fields of the bulk

$$\phi_\mathrm{bulk} = u(\sigma)\,\mathrm{e}^{\mathrm{i}(kx_3-\omega x_4)}: \qquad u'' - 7H\,\mathrm{sgn}(\sigma)\,u' + \mathrm{e}^{2H|\sigma|}\,m^2\,u = 0, \qquad m^2 = \omega^2 - k^2,$$
$$u_0 = 1, \quad \int_{-\infty}^{\infty} \mathrm{d}\sigma\,\mathrm{e}^{-5H|\sigma|}\,u_0^2 = \frac{2}{5H} \tag{FE-73}$$

Symbols in (FE-73).

- ϕ_bulk: A massless scalar field of the bulk, $\nabla_M\nabla^M\phi_\mathrm{bulk} = 0$ (an illustration: each polarisation of the graviton obeys its equation). Unit: GeV^3. Status: STANDARD.

- $u(\sigma)$, u_0: The profile along the hidden direction of a mode of a field of the bulk (a scalar, a gauge field or a polarisation of the graviton): the field is $u(\sigma)$ times a field of the fold; $u_0 = 1$ is the profile of the massless mode (the zero mode); in a propagator the response of the field to a unit source on the fold. Unit: dimensionless (a relative amplitude); length in a propagator. Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- k: A wave number: the momentum of a plane wave along x_3, $\mathrm{e}^{\mathrm{i}(kx_3 - \omega x_4)}$; in the cosmological perturbations the comoving wave number of a Fourier mode, $k = |\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m^2_{ij} = m_i^2 - m_j^2$ the squared-mass difference of the states i and j (Δm^2_{21}, Δm^2_{31}, Δm^2_{32}); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by v^2_loc/v^2 where this differs from 1; m without a label the mass of a mode of a field of the bulk as the fold sees it, $m^2 = \omega^2 - k^2$; for the W and Z bosons their complex pole mass, $m^2 = m_0^2 - \mathrm{i}m_0\Gamma_\mathrm{bulk}$ to the leading order; in the relic abundance of the particles χ their mass. Unit: GeV. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.

A massless scalar of the bulk (each polarisation of the graviton obeys the same equation): ω and k enter only through $m^2 = \omega^2 - k^2$, so every mode is a field of the fold with the dispersion $\omega^2 = k^2 + m^2$. The zero mode has the profile 1 and the finite kinetic weight $2/(5H)$: a massless field of the fold, bound to it, at the speed of light at every wavelength; the other modes form a continuum from zero mass upwards and are slower. Status: DERIVED (`um4_bulk.py`, B1: the profile equation exactly on both halves; the lowest ω^2 of the operator in a box $|\sigma| < 6/H$ equals k^2 to 4×10^{-12} for $k = 0.1H$ to $2H$; B10: the continuum of massive modes from zero mass).

$$B_\sigma = 0, \quad B_a = u(\sigma)\,\hat{B}_a(x): \qquad u'' - 5H\,\mathrm{sgn}(\sigma)\,u' + \mathrm{e}^{2H|\sigma|}\,m^2\,u = 0, \qquad \eta^{ac}\,\partial_a \hat{B}_{cb} = m^2\,\hat{B}_b,$$
$$u_0 = 1, \quad \int_{-\infty}^{\infty} \mathrm{d}\sigma\,\mathrm{e}^{-3H|\sigma|}\,u_0^2 = \frac{2}{3H} \tag{FE-74}$$

Symbols in (FE-74).

- B_M, B_μ, B_{MN}: The hypercharge gauge field of $U(1)_Y$: in the bulk B_M (geometric normalisation), on the fold the canonically normalised four-dimensional field $B_\mu(x)$ of its massless mode, $g_1 B_\mu(x) = B_\mu(\sigma = 0, x)$; B_{MN}, $B_{\mu\nu}$ their field strengths. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- $u(\sigma)$, u_0: The profile along the hidden direction of a mode of a field of the bulk (a scalar, a gauge field or a polarisation of the graviton): the field is $u(\sigma)$ times a field of the fold; $u_0 = 1$ is the profile of the massless mode (the zero mode); in a propagator the response of the field to a unit source on the fold. Unit: dimensionless (a

relative amplitude); length in a propagator. Status: DERIVED.

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\hat{B}_a$, $\hat{B}_{ab}$: The field of the fold of a mode of the hypercharge field of the bulk (of any gauge field of the bulk to the first order in the field), $B_a = u(\sigma)\hat{B}_a(x)$; $\hat{B}_{ab} = \partial_a\hat{B}_b - \partial_b\hat{B}_a$ its field strength. Unit: GeV (field), GeV^2 (field strength). Status: STANDARD.
- x: The coordinates of the fold: x_1, x_2, x_3 of ordinary space, x_4 the time, x_5, x_6, x_7 the extra times; a field of the fold is a function of the four observable coordinates $x = (x_1, x_2, x_3, x_4)$, and $\mathrm{d}^4x = \mathrm{d}x_1\,\mathrm{d}x_2\,\mathrm{d}x_3\,\mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m^2_{ij} = m_i^2 - m_j^2$ the squared-mass difference of the states i and j (Δm^2_{21}, Δm^2_{31}, Δm^2_{32}); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by v_loc^2/v^2 where this differs from 1; m without a label the mass of a mode of a field of the bulk as the fold sees it, $m^2 = \omega^2 - k^2$; for the W and Z bosons their complex pole mass, $m^2 = m_0^2 - \mathrm{i}m_0\Gamma_\mathrm{bulk}$ to the leading order; in the relic abundance of the particles χ their mass. Unit: GeV. Status: STANDARD.
- η_{ab}: The flat metric of the frame, $\eta = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$ (eta4488 of 4+4.nb); η^{ab} is its inverse (the same matrix). Unit: dimensionless. Status: AUTHOR.
- c: A frame index (as a). Status: STANDARD.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.

The modes of a gauge field of the bulk in the gauge $B_\sigma = 0$: the field of the fold obeys the Proca equation with the mass m; the massless mode has the profile 1 and the kinetic weight $2/(3H)$, which gives the four-dimensional couplings $g_k^2 = 3H\mathtt{g}_k^2/(2V_*)$ (FE-44). A component along σ has no normalisable zero mode. Status: DERIVED (`um4_bulk.py`, B2).

$$
\begin{aligned}
&A_\mu = \mathfrak{e}_\mu\,\mathrm{e}^{\mathrm{i}k(x_3-x_4)}: \qquad \mathfrak{e}_3 + \mathfrak{e}_4 = 0, \qquad \mathfrak{e}_\mu \sim \mathfrak{e}_\mu + \beta_\mathrm{g}\,k\,(0,0,1,-1),\\
&\mathfrak{e}_\pm = \frac{1}{\sqrt{2}}\big(1, \pm\mathrm{i}, 0, 0\big) \to \mathrm{e}^{\mp\mathrm{i}\theta_\mathrm{r}}\,\mathfrak{e}_\pm, \qquad \vec{E}\cdot\vec{B} = 0, \quad \vec{E}^2 = \vec{B}^2
\end{aligned}
\tag{FE-75}
$$

Symbols in (FE-75).

- A_μ: The photon field of the fold (canonically normalised): the massless combination of the massless modes of W^1, W^2, W^3 and B that the Higgs vacuum leaves massless; a field of the bulk whose zero mode travels at the speed of light. Unit: GeV. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- $\mathfrak{e}_\mu$, $\mathfrak{e}_\pm$: The polarisation vector of a plane wave of the photon, with components along x_1, x_2, x_3, x_4; $\mathfrak{e}_\pm = (1, \pm\mathrm{i}, 0, 0)/\sqrt{2}$ the circular polarisations of a wave travelling along x_3. Unit: dimensionless. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- k: A wave number: the momentum of a plane wave along x_3, $\mathrm{e}^{\mathrm{i}(kx_3 - \omega x_4)}$; in the cosmological perturbations the comoving wave number of a Fourier mode, $k = |\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- β_g: Any constant: the size of a gauge transformation, which adds a multiple of the wave vector to the polarisation vector of the photon. Unit: dimensionless. Status: STANDARD.
- θ_r: The angle of a rotation about the direction of motion of a wave. Unit: dimensionless (radian). Status: STANDARD.
- $\vec{E}$: The electric field of the photon on the fold: its component along the direction x_i of ordinary space is $\partial A_4/\partial x_i - \partial A_i/\partial x_4$ ($i = 1, 2, 3$; A_4 and A_i the components of A_μ); $\vec{E}^2$ its square. Unit: GeV^2. Status: STANDARD.
- $\vec{B}$: The magnetic field of the photon on the fold, the curl of the components A_1, A_2, A_3 of A_μ along ordinary space (its component along x_1 is $\partial A_3/\partial x_2 - \partial A_2/\partial x_3$, and cyclically); $\vec{B}^2$ its square. Unit: GeV^2. Status: STANDARD.

The photon as a wave: a transverse wave of helicity ± 1 at the speed of light, with its electric and magnetic fields orthogonal to the direction of motion and to each other and equal in size; the longitudinal and the time-like polarisations are pure gauge. Status: DERIVED (`um4_bulk.py`, B3), ASSUMED (no components along the extra times).

$$\frac{1}{2}\Big[\mathrm{e}^{2H|\sigma|}\Big(\frac{\partial^2 h_\mathrm{gw}}{\partial x_4^2} - \frac{\partial^2 h_\mathrm{gw}}{\partial x_3^2}\Big) - \frac{\partial^2 h_\mathrm{gw}}{\partial \sigma^2} + 7H\,\mathrm{sgn}(\sigma)\,\frac{\partial h_\mathrm{gw}}{\partial \sigma}\Big] = 0: \qquad h_\mathrm{gw} = h_+(x_3 - x_4),$$
$$\mathsf{h}_{ij} = \begin{pmatrix} h_+ & h_\times & 0 \\ h_\times & -h_+ & 0 \\ 0 & 0 & 0 \end{pmatrix}, \qquad \frac{\Delta L}{L} = \frac{1}{2}\big(h_+ \cos 2\phi_\mathrm{r} + h_\times \sin 2\phi_\mathrm{r}\big), \qquad h_+ \pm \mathrm{i}h_\times \to \mathrm{e}^{\mp 2\mathrm{i}\theta_\mathrm{r}}\big(h_+ \pm \mathrm{i}h_\times\big) \tag{FE-76}$$

Symbols in (FE-76).

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- h_gw: The amplitude of a gravitational wave of the + polarisation travelling along x_3: the metric components along x_1 and x_2 multiplied by $\mathrm{e}^{h_\mathrm{gw}}$ and $\mathrm{e}^{-h_\mathrm{gw}}$; a function of σ, x_3 and x_4 in the bulk, and on the fold the gravitational wave of the four-dimensional world, to the first order its plus polarisation h_+. Unit: dimensionless. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- h_+: The plus polarisation of a gravitational wave travelling along x_3: $\mathsf{h}_{11} = -\mathsf{h}_{22} = h_+$, a function of $x_3 - x_4$; to the first order the zero mode of h_gw. Unit: dimensionless. Status: STANDARD.
- h_{ij}: The transverse traceless perturbation of the metric of ordinary space by a gravitational wave as the four-dimensional world sees it: $\hat{h}_{ij} = \mathrm{e}^{2a_4}(\delta_{ij} + \mathsf{h}_{ij})$ ($i, j = 1, 2, 3$). Unit: dimensionless. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal), and in a tensor of ordinary space such as h_{ij} one of the directions x_1, x_2, x_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- j: A second generation or mass-state index, or a second direction of ordinary space (as i). Status: STANDARD.

- $h_\times$: The cross polarisation of a gravitational wave travelling along x_3: $\mathsf{h}_{12} = \mathsf{h}_{21} = h_\times$, a function of $x_3 - x_4$. Unit: dimensionless. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- L: A distance between two points of the fold along ordinary space: for a gravitational wave the radius of a ring of free test masses (ΔL the change of the distance of one of them from the centre), for a signal the distance between its emitter and its receiver. Unit: length. Status: STANDARD.
- ϕ_r: The angle at which a test mass sits on a ring of test masses in the plane of x_1 and x_2, measured from the x_1 axis. Unit: dimensionless (radian). Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- θ_r: The angle of a rotation about the direction of motion of a wave. Unit: dimensionless (radian). Status: STANDARD.

The gravitational wave: in the bulk the + polarisation obeys the equation of the massless scalar, and its zero mode, any function of $x_3 - x_4$, solves the bulk equation and the junction condition; on the fold it is the transverse traceless wave with the plus and the cross polarisations, which stretch and squeeze a ring of test masses along perpendicular directions, and it has the helicities ± 2. Of the 14 polarisations of the massless graviton of seven dimensions the graviton (2) and the six scalars h_{rs} have kinetic terms of the right sign, the three vectors $h_{\mu r}$ (6) of the wrong sign: they are ghosts, like the components of a gauge field along the extra times that UM-4 excludes. Matter of the fold that does not depend on the extra times and is the same along the three of them emits at the linear order only the plus and the cross polarisations and the volume of the extra times, which the fold fluid holds fixed, and no ghost vectors. Status: DERIVED (`um4_bulk.py`, B4, B5 and B11: the signs of the kinetic terms from the Einstein-Hilbert action to the second order; `um4_foldsources.py`, FS2), ASSUMED (the volume held fixed for waves as well), OPEN (the ghost vectors beyond the linear order).

$$\begin{aligned} &\Psi_\mathrm{k} = \mathrm{e}^{\frac{7}{2}H|\sigma|}\,\mathrm{e}^{-M_\mathrm{k}|\sigma|}\,\chi_0(x), \qquad \gamma^{(0)}\chi_0 = -\chi_0, \qquad \gamma^{(a)}\,\partial_a\chi_0 = 0 \quad \text{(along the fold)}, \\ &\sqrt{|g|}\,\bar\Psi_\mathrm{k}\gamma^{x_4}\Psi_\mathrm{k} \propto \mathrm{e}^{(H-2M_\mathrm{k})|\sigma|} \end{aligned} \tag{FE-77}$$

Symbols in (FE-77).

- Ψ_k, $\bar\Psi_\mathrm{k}$: A fermion field of the bulk with the kink mass $M_\mathrm{k}\,\mathrm{sgn}(\sigma)$ (an illustration, not a field of UM-4). Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- M_k: The mass of a fermion of the bulk that changes sign at the fold (a kink mass $M_\mathrm{k}\,\mathrm{sgn}(\sigma)$), used to illustrate the speeds of bulk fermions. Unit: GeV. Status: ASSUMED.
- χ_0: The spinor of the fold of the massless mode of a fermion of the bulk: a solution of the massless Dirac equation of the fold of one chirality along σ. Unit: $\mathrm{GeV}^{7/2}$. Status: DERIVED.
- x: The coordinates of the fold: x_1, x_2, x_3 of ordinary space, x_4 the time, x_5, x_6, x_7 the extra times; a field of the fold is a function of the four observable coordinates $x = (x_1, x_2, x_3, x_4)$, and $\mathrm{d}^4x = \mathrm{d}x_1\,\mathrm{d}x_2\,\mathrm{d}x_3\,\mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat g_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.

- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\dots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c=1$). Status: AUTHOR.

A fermion of the bulk with a kink mass (an illustration: UM-4 has no such field): its zero mode is a massless fermion of the fold for $M_\mathrm{k} > H/2$, at the speed of light. Status: DERIVED (`um4_bulk.py`, B6).

$$\mathsf{S}=\frac{1}{2}\,\gamma^{(3)}\gamma^{(4)}: \qquad \big[\mathsf{S},\gamma^{(0)}\big]=\big[\mathsf{S},\Gamma_8\big]=0, \qquad \big[\mathsf{S},\gamma^{(3)}\big]=-\gamma^{(4)}, \qquad \big[\mathsf{S},\gamma^{(4)}\big]=-\gamma^{(3)} \qquad \Rightarrow \qquad \omega^2=\omega_0^2+k^2 \tag{FE-78}$$

Symbols in (FE-78).

- S: The generator of the boosts along x_3 on dirac16complex, $\mathsf{S}=\frac{1}{2}\gamma^{(3)}\gamma^{(4)}$. Unit: dimensionless. Status: DERIVED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M=\mathbf{e}_a{}^M T^a$, $\gamma_M=g_{MN}\gamma^N$; $\gamma^{(a)}=T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- Γ_8: The chirality matrix of eight dimensions, $\Gamma_8=T_0T_1\cdots T_7$ (the author's T16^A[8]) $=\mathrm{diag}(-\mathbf{1}_8,\mathbf{1}_8)$; $\Gamma_8^2=\mathbf{1}$; it anticommutes with every T_a. Unit: dimensionless. Status: AUTHOR.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- ω_0: The energy at rest (zero momentum along ordinary space and along the extra times) of a state of dirac16complex at the fold; complex for a resonance, $\omega_0=\omega_\mathrm{r}-\frac{\mathrm{i}}{2}\Gamma_\chi$. Unit: GeV. Status: DERIVED.
- k: A wave number: the momentum of a plane wave along x_3, $\mathrm{e}^{\mathrm{i}(kx_3-\omega x_4)}$; in the cosmological perturbations the comoving wave number of a Fourier mode, $k=|\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.

dirac16complex is invariant under the boosts of the fold, so every one of its states at the fold has $\omega^2=\omega_0^2+k^2$ exactly and moves slower than light. Status: DERIVED (`um4_bulk.py`, B7, with the author's matrices).

7.2 The propagators from one point of the fold to another

$$-\mathrm{e}^{nH|\sigma|}\,\partial_\sigma\big(\mathrm{e}^{-nH|\sigma|}\,\partial_\sigma u\big)+\mathrm{e}^{2H|\sigma|}\,\mathsf{p}^2\,u=\delta_\mathrm{D}(\sigma), \qquad \mathbb{G}_n(\mathsf{p})=u(0)=\frac{\mathrm{K}_{n/2}(\mathsf{p}/H)}{2\mathsf{p}\,\mathrm{K}_{n/2-1}(\mathsf{p}/H)} \tag{FE-79}$$

Symbols in (FE-79).

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- n: The label of a field of the bulk in its profile equation: the exponent of the weight $\mathrm{e}^{-nH|\sigma|}$ of its σ-derivatives, $n=7$ for a scalar and for each polarisation of the graviton, $n=5$ for a gauge field. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda=-21H^2$ and the tension $\kappa\lambda_\mathrm{F}=12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H>2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma=|\cot z|\,\mathrm{d}x_0$, $\sigma=\mathrm{sgn}(\pi/2-z)\ln(\sin z)/(6H)$ on one period $0<z<\pi$; $\sigma=0$ is the fold $z=\pi/2$, $\sigma\to\mp\infty$ the tips $z\to0,\pi$; one period of x_0 is the whole line $-\infty<\sigma<\infty$. Unit: length. Status: DERIVED.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- $u(\sigma)$, u_0: The profile along the hidden direction of a mode of a field of the bulk (a scalar, a gauge field or a polarisation of the graviton): the field is $u(\sigma)$ times a field of the fold; $u_0=1$ is the profile of the massless mode (the zero mode); in a propagator the response of the field to a unit source on the fold. Unit: dimensionless (a relative amplitude); length in a propagator. Status: DERIVED.

- p: The Euclidean momentum along the fold of a source on the fold: p^2 the square of its seven-dimensional momentum (for a static source spread uniformly over the extra times the square of its momentum along ordinary space); $\mathsf{p} = -im$ for a time-like momentum of the mass m with outgoing waves into the bulk. Unit: GeV. Status: STANDARD.
- δ_{D}: The Dirac delta function of σ: zero for $\sigma \neq 0$, with the integral 1 over any range that contains $\sigma = 0$; the integral of $\delta_{\mathrm{D}}(\sigma)$ times a smooth function of σ is the value of that function at $\sigma = 0$. Unit: 1/length (for $\delta_{\mathrm{D}}(\sigma)$). Status: STANDARD.
- $\mathbb{G}_n$, $\mathbb{G}_7$, $\mathbb{G}_5$: The propagator of a massless field of the bulk from one point of the fold to another, in the momentum space of the fold: the value at the fold of its response to a unit source on the fold; $\mathbb{G}_7$ for each polarisation of the graviton and for a scalar, $\mathbb{G}_5$ for a gauge field. Unit: GeV^{-1}. Status: DERIVED.
- K: The modified Bessel function of the second kind (of the order written as its subscript): the solution of the modified Bessel equation that decays exponentially for a large argument; for a half-integer order it is an exponential times a polynomial in the inverse of the argument, which makes the propagators of the bulk rational functions. Unit: dimensionless. Status: STANDARD.

$$\begin{aligned}\mathbb{G}_7 &= \frac{H}{2\mathsf{p}^2}\,\frac{x_{\mathrm{p}}^3 + 6x_{\mathrm{p}}^2 + 15x_{\mathrm{p}} + 15}{x_{\mathrm{p}}^2 + 3x_{\mathrm{p}} + 3} = \frac{5H}{2\mathsf{p}^2}\Big(1 + \frac{x_{\mathrm{p}}^2}{15} - \frac{x_{\mathrm{p}}^4}{45} + \frac{x_{\mathrm{p}}^5}{45} + \dots\Big),\\ \mathbb{G}_5 &= \frac{H}{2\mathsf{p}^2}\,\frac{x_{\mathrm{p}}^2 + 3x_{\mathrm{p}} + 3}{x_{\mathrm{p}} + 1} = \frac{3H}{2\mathsf{p}^2}\Big(1 + \frac{x_{\mathrm{p}}^2}{3} - \frac{x_{\mathrm{p}}^3}{3} + \dots\Big), \qquad x_{\mathrm{p}} = \frac{\mathsf{p}}{H}\end{aligned} \tag{FE-80}$$

Symbols in (FE-80).

- $\mathbb{G}_n$, $\mathbb{G}_7$, $\mathbb{G}_5$: The propagator of a massless field of the bulk from one point of the fold to another, in the momentum space of the fold: the value at the fold of its response to a unit source on the fold; $\mathbb{G}_7$ for each polarisation of the graviton and for a scalar, $\mathbb{G}_5$ for a gauge field. Unit: GeV^{-1}. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- p: The Euclidean momentum along the fold of a source on the fold: p^2 the square of its seven-dimensional momentum (for a static source spread uniformly over the extra times the square of its momentum along ordinary space); $\mathsf{p} = -im$ for a time-like momentum of the mass m with outgoing waves into the bulk. Unit: GeV. Status: STANDARD.
- x_{p}: The Euclidean momentum along the fold in units of H, $x_{\mathrm{p}} = \mathsf{p}/H$. Unit: dimensionless. Status: DERIVED.

The response of a massless field of the bulk to a unit source on the fold: the propagator of the bulk between points of the fold, rational in $x_{\mathrm{p}} = \mathsf{p}/H$ for the half-integer orders of the Bessel functions. For small momenta it is the massless propagator of the fold with the strength $\frac{5}{2}H$ (graviton) or $\frac{3}{2}H$ (gauge field); the even powers of x_{p} are contact terms, the first odd power gives a correction of long range. Status: DERIVED (`um4_propagators.py`, P1 and P2).

$$\mathcal{V}_7(r) = \frac{5H}{8\pi r}\Big[1 + \frac{16}{15\pi}\,\frac{1}{(Hr)^5} + \dots\Big], \qquad \mathcal{V}_5(r) = \frac{3H}{8\pi r}\Big[1 + \frac{4}{3\pi}\,\frac{1}{(Hr)^3} + \dots\Big] \tag{FE-81}$$

Symbols in (FE-81).

- $\mathcal{V}_7(r)$, $\mathcal{V}_5(r)$: The static potentials between two unit point sources on the fold at the distance r (spread uniformly over the extra times) from the exchange of the graviton ($\mathcal{V}_7$) and of a gauge field ($\mathcal{V}_5$) of the bulk: the three-dimensional Fourier transforms of $\mathbb{G}_7$ and $\mathbb{G}_5$. Unit: GeV^2. Status: DERIVED.
- r: A distance from a centre: the distance from the centre of the star along the path of a neutrino; in Newton's and Coulomb's laws the distance between two point sources on the fold. Unit: length. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

Newton's and Coulomb's laws on the fold: the massive modes of the bulk add the relative corrections $0.3395/(Hr)^5$ and $0.4244/(Hr)^3$. Status: DERIVED (`um4_propagators.py`, P3: the asymptotic coefficients exactly, and the numerical Fourier transforms at $Hr = 32$ within 1.2 per cent of them), ESTIMATE (the factor of order one of the tensor structure of the massive gravitons).

$$\Big(1+M_{\rm loc}^2\,\mathbb{G}_5(\mathsf{p})\Big)\Big|_{\mathsf{p}=-\mathrm{i}m}=0:\qquad m^2=m_0^2\Big(1-\frac{m_0^2}{3H^2}\Big)-\mathrm{i}\,\frac{m_0^5}{3H^3}+\ldots,\qquad m_0^2=\frac{3}{2}\,HM_{\rm loc}^2,\qquad \Gamma_{\rm bulk}=\frac{m_0^4}{3H^3}\ \text{(FE-82)}$$

Symbols in (FE-82).

- $M_{\rm loc}$: The mass term that the Higgs vacuum on the fold gives a gauge field of the bulk: $M_{\rm loc}^2\delta_{\rm D}(\sigma)$ in its profile equation; for the W, $M_{\rm loc}^2=\mathsf{g}_2^2(v^2+\mathcal{A}^2)/(4V_*)$. Unit: $\mathrm{GeV}^{1/2}$ ($M_{\rm loc}^2$ in GeV). Status: DERIVED.
- $\mathbb{G}_n$, $\mathbb{G}_7$, $\mathbb{G}_5$: The propagator of a massless field of the bulk from one point of the fold to another, in the momentum space of the fold: the value at the fold of its response to a unit source on the fold; $\mathbb{G}_7$ for each polarisation of the graviton and for a scalar, $\mathbb{G}_5$ for a gauge field. Unit: GeV^{-1}. Status: DERIVED.
- p: The Euclidean momentum along the fold of a source on the fold: p^2 the square of its seven-dimensional momentum (for a static source spread uniformly over the extra times the square of its momentum along ordinary space); $\mathsf{p}=-\mathrm{i}m$ for a time-like momentum of the mass m with outgoing waves into the bulk. Unit: GeV. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m_{ij}^2=m_i^2-m_j^2$ the squared-mass difference of the states i and j (Δm_{21}^2, Δm_{31}^2, Δm_{32}^2); Δm^2 without labels the difference of the two states of a resonance, $m_2^2-m_1^2$, multiplied by $v_{\rm loc}^2/v^2$ where this differs from 1; m without a label the mass of a mode of a field of the bulk as the fold sees it, $m^2=\omega^2-k^2$; for the W and Z bosons their complex pole mass, $m^2=m_0^2-\mathrm{i}m_0\Gamma_{\rm bulk}$ to the leading order; in the relic abundance of the particles χ their mass. Unit: GeV. Status: STANDARD.
- m_0: The mass of the W or the Z boson without the massive modes of the bulk (the mass of the massless mode with the Higgs vacuum), $m_0^2=\frac{3}{2}HM_{\rm loc}^2$; $m_0^2=g_2^2(v^2+\mathcal{A}^2)/4$ for the W. Unit: GeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda=-21H^2$ and the tension $\kappa\lambda_{\rm F}=12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H>2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- $\Gamma_{\rm bulk}$: The width of the W or the Z boson into the bulk (its decay into the massive modes of the gauge fields of the bulk), $\Gamma_{\rm bulk}=m_0(m_0/H)^3/3$. Unit: GeV. Value: 1.42 MeV for the Z and 0.85 MeV for the W at $H=2534$ GeV. Status: DERIVED.

The W and Z bosons are resonances of the bulk: the mass term that the Higgs vacuum gives the gauge fields at the fold moves the pole of the propagator to a complex mass, with the width $\Gamma_{\rm bulk}=m_0(m_0/H)^3/3$ into the bulk (m_0 the mass of the zero mode alone, $m_0^2=g_2^2(v^2+\mathcal{A}^2)/4$ for the W and $(g_1^2+g_2^2)(v^2+\mathcal{A}^2)/4$ for the Z); the squared masses are shifted by $-m_0^4/(3H^2)$ (the masses by $-m_0^3/(6H^2)$). Status: DERIVED (`um4_propagators.py`, P4: the exact pole of the rational propagator differs from the expansion by the relative amounts $1.83(m_0/H)^2$ in the imaginary part and $0.44(m_0/H)^4$ in the real part, the next orders, for $m_0/H=10^{-2}$ to 10^{-3}).

bound on H	from	value
invisible width of the Z boson	its decay into the bulk: at most 1.42 MeV of extra invisible width (the number of light neutrinos 2.9963 ± 0.0074 measured at the Z, P. Janot and S. Jadach, Phys. Lett. B 803 (2020) 135319, with 167.2 MeV per neutrino species; the exact pole gives 1.414 MeV, the formula 1.417 MeV)	$H>2534$ GeV

bound on H	from	value
contact interactions between quarks and leptons	the contact term $x_{\mathrm{p}}^2/3$ of the propagator of the photon (with the charge 2/3 of the up quark) against the limit $\Lambda_{\mathrm{CI}} > 26.0$ TeV of ATLAS for the destructive left-left structure in the spectra of electron and muon pairs (JHEP 11 (2020) 005; the photon exchange is a vector-vector interaction, so this is an estimate)	$H > 1083$ GeV (1327 GeV with unit charges)
the parameter rho of the electroweak fits	the shift $-m_0^4/(3H^2)$ of the squared masses of the W and Z, against 1.00031 ± 0.00019 (Particle Data Group 2024)	$H > 969$ GeV
Newton's law at 52 micrometres	the correction $0.3395/(Hr)^5$, taken to be below one per cent at 52 micrometres, the shortest separation of the torsion balances of J. G. Lee, E. G. Adelberger, T. S. Cook, S. M. Fleischer and B. R. Heckel, Phys. Rev. Lett. 124 (2020) 101101 (which test 52 micrometres to 3 mm and exclude Yukawa forces of gravitational strength with a range above 38.6 micrometres)	$H > 7.7 \times 10^{-3}$ eV

The bounds on the author's constant (Λ_{CI} the scale of a contact interaction, r the distance between the sources). The strongest is the invisible width of the Z boson: $H > 2534$ GeV (about 2.5 TeV), the value used everywhere in UM-4. At $H = 2534$ GeV the width of the W into the bulk is 0.85 MeV (its measured total width is 2.1 GeV), the correction of Newton's law at 52 micrometres is 2.6×10^{-75} (2.7×10^{-75} for $H = 2.5$ TeV) and that of Coulomb's law at the Bohr radius 1.4×10^{-27}. Status: ESTIMATE (`um4_propagators.py`, P5: the measured values are inputs).

7.3 The causal structure

$$\tilde{\sigma} = \mathrm{sgn}(\sigma)\,\frac{\mathrm{e}^{H|\sigma|} - 1}{H}: \qquad g_{MN}\,\mathrm{d}X^M \mathrm{d}X^N = \frac{\mathrm{d}\tilde{\sigma}^2 + \mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2 - \mathrm{d}x_4^2 - \mathrm{d}x_5^2 - \mathrm{d}x_6^2 - \mathrm{d}x_7^2}{\left(1 + H|\tilde{\sigma}|\right)^2}, \qquad \Delta x_4 \geq L \tag{FE-83}$$

Symbols in (FE-83).

- $\tilde{\sigma}$: The conformal coordinate of the hidden direction on the static block, $\tilde{\sigma} = \mathrm{sgn}(\sigma)(\mathrm{e}^{H|\sigma|} - 1)/H$, in which the static block is conformally flat. Unit: length. Status: DERIVED.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.

- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- L: A distance between two points of the fold along ordinary space: for a gravitational wave the radius of a ring of free test masses (ΔL the change of the distance of one of them from the centre), for a signal the distance between its emitter and its receiver. Unit: length. Status: STANDARD.

The static block is conformally flat: its null geodesics are the null straight lines of flat space of signature (4, 4), and a signal that leaves the fold and returns to it at fixed extra times arrives later than light along the fold (the triangle inequality of the section of fixed extra times). A null line that also moves by Y along an extra time reaches the distance L in the time $\sqrt{L^2 - Y^2}$, down to 0: the bound holds for the fields that do not depend on the extra times, not in the full spacetime with its four time-like directions. Status: DERIVED (`um4_causality.py`, C1: the geodesic equation; at fixed extra times 200 sampled paths through the bulk, the smallest delay 1.7×10^{-5} of the distance; the null line along an extra time).

$$c_{\mathrm{loc}}(\sigma) = 1 - \frac{a_4''}{5H^2}\left(\mathrm{e}^{2H|\sigma|} - 1\right), \qquad \sigma(x_3) = \frac{|a_4''|}{5H}\, x_3\,(L - x_3), \qquad \Delta x_4 = L - \frac{a_4''^2 L^3}{150H^2} \tag{FE-84}$$

Symbols in (FE-84).

- c_{loc}: The coordinate speed of light along x_3 at the depth σ (the rate of change of x_3 with x_4 along a null line at fixed σ), in units of that on the fold: $\mathrm{e}^{H|\sigma|}$ on the static block of UM-3, whose time was not warped, 1 on the static block of UM-4, and on an expanding fold of UM-4 different from 1 in proportion to the second derivative a_4'' of a_4 (the rate of change of the expansion rate a_4'). Unit: dimensionless. Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- L: A distance between two points of the fold along ordinary space: for a gravitational wave the radius of a ring of free test masses (ΔL the change of the distance of one of them from the centre), for a signal the distance

between its emitter and its receiver. Unit: length. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.

An expanding fold with frozen extra times: wherever the expansion rate falls ($a_4'' = -(1 + q_{\mathrm{dec}})H_4^2 < 0$, i.e. for every deceleration parameter above -1, today included) the light cone of the bulk is slightly wider than that of the fold, and the best path through the bulk gains $a_4''^2 L^3/(150H^2)$ over light along the fold; extrapolated to the Hubble distance today this is 4.4×10^{-94} of the travel time for $H = 2534$ GeV (4.5×10^{-94} for $H = 2.5$ TeV): the bulk is ahead, by an unobservable amount. Where the expansion rate grows ($a_4'' > 0$) the path along the fold is the fastest. Status: DERIVED (`um4_causality.py`, C2 and C3: the speed from the metric of the fold; the numerical minimisation of the travel time gives 0.997 of the formula, and no advance where $a_4'' > 0$).

8 dirac16complex: the resonances at the fold

The author's field dirac16complex lives in the bulk. Its modes along the hidden direction, and the states that the fold holds, are the "x_0 masses" that an observer on the fold sees. This section gives its equations, the solutions of its profile equation, the resonances of the fold (the particles χ, the dark matter of UM-4) with their mass and their width, and the momentum along the extra times.

$$\gamma^M D_M \Psi - m_\Psi \Psi - \lambda_\Psi \big(\bar\Psi \Psi\big)\Psi = 0 \qquad (\sigma \neq 0) \tag{FE-12}$$

Symbols in (FE-12).

- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar\Psi = \partial_M\bar\Psi - \bar\Psi\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat h^{\mu\nu} D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: GeV$^{7/2}$. Status: AUTHOR.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- λ_Ψ: The coupling of the contact self-interaction of dirac16complex. Unit: GeV^{-6}. Status: RECORD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

$$\begin{aligned}
&\Psi\big|_{\sigma=0^+} = \exp\Big(\frac{1}{V_*}\gamma^{(0)}\,\mathbb{M}\Big)\,\Psi\big|_{\sigma=0^-},\\
&\mathbb{M} = c_\Phi\big(\Phi^\dagger\Phi\big)P_{\mathrm{L}} + \sum_X \Big[\frac{1}{4}\big(c_X^G\, G^A_{\mu\nu}G^{A\mu\nu} + c_X^W\, W^I_{\mu\nu}W^{I\mu\nu} + c_X^B\, B_{\mu\nu}B^{\mu\nu}\big)P_X + \frac{d_X}{2}\,B_{\mu\nu}\,\mathrm{i}\Sigma^{\mu\nu}P_X\\
&\qquad + \sum_f c_X^f\big(2\mathrm{i}\,J_f^\mu P_X D_\mu + \mathrm{i}(\hat\nabla_\mu J_f^\mu)P_X\big)\Big]
\end{aligned} \tag{FE-13}$$

Symbols in (FE-13).

- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: GeV$^{7/2}$. Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- $\mathbb{M}$: The fold operator of dirac16complex: the 16×16 matrix (with a derivative along the fold) built from the Standard-Model fields at the fold by varying the portals with respect to $\bar\Psi$. Unit: GeV^{-3} (so that $\mathbb{M}/V_*$ is dimensionless). Status: DERIVED.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c_X^G: The gluon-kinetic portal coupling of the half X of dirac16complex. Unit: GeV^{-7}. Status: INVENTED.
- G_M^A, G_μ^A, G_{MN}^A: The gluon field of $SU(3)_C$ ($A = 1, \ldots, 8$): in the bulk G_M^A, a field of eight dimensions in its geometric normalisation (the covariant derivative $\partial - \mathrm{i}G$), and on the fold the canonically normalised four-dimensional field $G_\mu^A(x)$ of its massless mode, $g_3 G_\mu^A(x) = G_\mu^A(\sigma = 0, x)$; G_{MN}^A and $G_{\mu\nu}^A$ their field strengths. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- c_X^W: The W-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- W_M^I, W_μ^I, W_{MN}^I: The gauge field of $SU(2)_L$ ($I = 1, 2, 3$): in the bulk W_M^I (geometric normalisation), on the fold the canonically normalised four-dimensional field $W_\mu^I(x)$ of its massless mode, $g_2 W_\mu^I(x) = W_\mu^I(\sigma = 0, x)$; W_{MN}^I, $W_{\mu\nu}^I$ their field strengths; $W_\mu^\pm = (W_\mu^1 \mp \mathrm{i}W_\mu^2)/\sqrt{2}$. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- c_X^B: The B-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- B_M, B_μ, B_{MN}: The hypercharge gauge field of $U(1)_Y$: in the bulk B_M (geometric normalisation), on the fold the canonically normalised four-dimensional field $B_\mu(x)$ of its massless mode, $g_1 B_\mu(x) = B_\mu(\sigma = 0, x)$; B_{MN}, $B_{\mu\nu}$ their field strengths. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- P_X: The chiral projector of the half X: $P_X = P_\mathrm{L}$ for $X = \mathrm{L}$, P_R for $X = \mathrm{R}$. Unit: dimensionless. Status: AUTHOR.
- d_X: The dipole-portal coupling of the half X of dirac16complex with the hypercharge field strength. Unit: GeV^{-5}. Status: INVENTED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- Σ^{ab}, $\Sigma^{\mu\nu}$: The generators of Spin(4,4) on dirac16complex, $\Sigma^{ab} = \frac{1}{4}[T^a, T^b]$ (the scaled commutators of the author's matrices); $\Sigma^{\mu\nu} = \mathbf{e}_a{}^\mu \mathbf{e}_b{}^\nu \Sigma^{ab}$ along the fold. Unit: dimensionless. Status: AUTHOR.
- f: The label of a kind of Standard-Model multiplet: $f \in \{Q, u, d, \ell, e, \nu\}$ (quark doublets, up-quark singlets, down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.
- c_X^f: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c_X^ℓ and c_X^ν are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.
- J_f^μ: The fermion-number current of the Standard-Model multiplets of kind f, $J_f^\mu = \bar\psi_f \hat\gamma^\mu \psi_f$ summed over the three generations and the gauge components; with other labels the currents that are the sources of the gauge fields. Unit: GeV^3. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar\Psi = \partial_M\bar\Psi - \bar\Psi\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection

of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu} D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2 \epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.

- $\hat{\nabla}$: The covariant derivative of the fold: with the Levi-Civita connection of the metric of the fold (for four-dimensional tensors of its metric $\hat{h}_{\mu\nu} = \mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the observable coordinates $x_1, \ldots, x_4$; for $\hat{\nabla}_a \mathcal{S}^a{}_b$ of the induced metric of the seven-dimensional fold). Status: STANDARD.

$$\gamma^M \Omega_M = -\frac{7}{2} H \,\mathrm{sgn}(\sigma)\, \gamma^{(0)} + \frac{3}{2} \frac{b_4'}{b_4} \,\mathrm{e}^{H|\sigma|}\, \gamma^{(4)} \neq 0 \qquad \text{(FE-14)}$$

Symbols in (FE-14).

- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Ω_M: The spin connection acting on dirac16complex, $\Omega_M = \frac{1}{2}\varpi_{Mab}\Sigma^{ab}$. Unit: 1/length. Status: RECORD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\, \mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

The field equation of dirac16complex in the bulk (FE-12, with its contact self-interaction), the matching condition at the fold (FE-13: the values on the two sides of the fold differ by the exponential of $\gamma^{(0)}\mathbb{M}/V_*$, where the fold operator $\mathbb{M}$ is built from the Standard-Model fields through the portals) and the canonical spin connection of the metric of UM-4 (FE-14), which couples dirac16complex to gravity and never vanishes: its first term comes from the warp of the seven directions of the fold, its second from the time dependence of b_4. Status: RECORD (FE-12), DERIVED (FE-13; FE-14, `um4_geometry.py`, G9, with the author's matrices).

$$\Psi = \mathrm{e}^{\frac{7}{2}H|\sigma|}\,\chi(\sigma)\,\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}, \qquad \gamma^{(0)}\chi' + \mathrm{e}^{H|\sigma|}\big(-\mathrm{i}\omega\,\gamma^{(4)} + \mathrm{i}\,\vec{k}\cdot\vec{\gamma}\big)\chi - m_\Psi \chi = 0 \qquad (\sigma \neq 0) \qquad \text{(FE-36)}$$

Symbols in (FE-36).

- Ψ, $\bar{\Psi}$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar{\Psi} = \Psi^\dagger \sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{\frac{7}{2}H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of the level of dirac16complex

at the fold (a long-lived resonance) and $\bar{\chi}$ its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\vec{k}$: The (comoving) momentum along ordinary space, $\vec{k}^2$ its square. Unit: GeV. Status: STANDARD.
- $\vec{x}$: The position in ordinary space, $\vec{x} = (x_1, x_2, x_3)$. Unit: length. Status: AUTHOR.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\vec{\gamma}$: The three frame Dirac matrices of dirac16complex along ordinary space, $\vec{\gamma} = (\gamma^{(1)}, \gamma^{(2)}, \gamma^{(3)})$. Unit: dimensionless. Status: AUTHOR.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.

$$\begin{aligned} &\chi' = \mathbb{K}\chi, \qquad \mathbb{K} = \gamma^{(0)}\big(m_\Psi + \mathrm{i}\omega\,\mathrm{e}^{H|\sigma|}\gamma^{(4)}\big), \qquad \mathbb{K}^2 = \big(m_\Psi^2 - \omega^2\mathrm{e}^{2H|\sigma|}\big)\mathbf{1}, \\ &q = \sqrt{m_\Psi^2 - \omega^2\mathrm{e}^{2H|\sigma|}}, \qquad |\omega|\,\mathrm{e}^{H\sigma_\mathrm{t}} = m_\Psi \end{aligned} \tag{FE-37}$$

Symbols in (FE-37).

- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{\frac{7}{2}H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of the level of dirac16complex at the fold (a long-lived resonance) and $\bar{\chi}$ its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\mathbb{K}$: The transfer matrix of the profile equation at zero momentum, $\chi' = \mathbb{K}\chi$. Unit: GeV. Status: DERIVED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\mathbf{1}$, $\mathbf{1}_4$, $\mathbf{1}_8$, $\mathbf{1}_{16}$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_4$, $\mathbf{1}_8$ and $\mathbf{1}_{16}$ are the identity matrices of 4, 8 and 16 rows. Unit: dimensionless. Status: STANDARD.
- q: The local decay rate of a profile of dirac16complex along σ at the frequency ω, $q = \sqrt{m_\Psi^2 - \omega^2\mathrm{e}^{2H|\sigma|}}$ (imaginary beyond the turning point, where the profile oscillates). Unit: GeV. Status: DERIVED.
- σ_t: The turning point of a state of dirac16complex of the frequency ω: the depth at which the warped frequency equals the bulk mass, $|\omega|\mathrm{e}^{H\sigma_\mathrm{t}} = m_\Psi$; beyond it every solution oscillates. Unit: length. Status: DERIVED.

The field equation on the static block for a mode of definite momentum and frequency (FE-36): the factor $\mathrm{e}^{(7/2)H|\sigma|}$ removes the spin connection, and the factor $\mathrm{e}^{H|\sigma|}$ multiplies the whole Dirac operator of the fold, the frequency as well as the momentum. At zero momentum (FE-37) the local gap $m_\Psi^2 - \omega^2\mathrm{e}^{2H|\sigma|}$ closes at the turning point σ_t; beyond it every solution oscillates. There is no gap in the bulk and no normalisable state of non-zero frequency: the states that UM-3 bound to the fold are resonances in UM-4. The 16 components split into eight copies of two

inequivalent problems of two components. Status: DERIVED (`um4_spectra.py`, U1, with the author's matrices).

$$\chi = (\omega\zeta_{\mathrm{w}})^{1/2}\begin{pmatrix}\mathfrak{z}_-(\omega\zeta_{\mathrm{w}})\\ \mathrm{i}\,\mathfrak{z}_+(\omega\zeta_{\mathrm{w}})\end{pmatrix}\quad(\sigma>0),\qquad \chi = (\omega\zeta_{\mathrm{w}})^{1/2}\begin{pmatrix}-\mathrm{i}\,\mathfrak{z}_+(\omega\zeta_{\mathrm{w}})\\ \mathfrak{z}_-(\omega\zeta_{\mathrm{w}})\end{pmatrix}\quad(\sigma<0),\qquad \zeta_{\mathrm{w}} = \frac{\mathrm{e}^{H|\sigma|}}{H}\tag{FE-85}$$

Symbols in (FE-85).

- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{\frac{7}{2}H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x}-\omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of the level of dirac16complex at the fold (a long-lived resonance) and $\bar{\chi}$ its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- ζ_{w}: The warped distance from the fold, $\zeta_{\mathrm{w}} = \mathrm{e}^{H|\sigma|}/H$ ($1/H$ at the fold): $\omega\zeta_{\mathrm{w}}$ is the argument of the exact solutions of dirac16complex, and $m\zeta_{\mathrm{w}}$ that of the massive modes of the mass m of the massless fields of the bulk. Unit: length. Status: DERIVED.
- $\mathfrak{z}_-$, $\mathfrak{z}_+$: Bessel functions of the orders $m_\Psi/H - \frac{1}{2}$ ($\mathfrak{z}_-$) and $m_\Psi/H + \frac{1}{2}$ ($\mathfrak{z}_+$), both of the same kind (of the first kind, of the second kind, or the Hankel functions of the first kind for waves that leave the fold), of the argument $\omega\zeta_{\mathrm{w}}$. Unit: dimensionless. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.

The exact solutions of the profile equation in one copy of two components: Bessel functions of the orders $m_\Psi/H \mp \frac{1}{2}$ of the warped argument $\omega\zeta_{\mathrm{w}}$; close to the fold they grow or decay like $\mathrm{e}^{\pm q|\sigma|}$, beyond the turning point they are waves, and the outgoing ones are the Hankel functions of the first kind. Status: DERIVED (`um4_spectra.py`, U2: checked against the differential equation at three depths).

$$\chi(0^+) = \exp\Big(\gamma^{(0)}\big(\mu_{\mathrm{L}}P_{\mathrm{L}} + \mu_{\mathrm{R}}P_{\mathrm{R}}\big)\Big)\,\chi(0^-),$$
$$\mu_X = \frac{1}{V_*}\Big[c_\Phi\,\langle\Phi^\dagger\Phi\rangle\,\delta_{X\mathrm{L}} + \frac{1}{4}\big(c_X^G\langle G^2\rangle + c_X^W\langle W^2\rangle + c_X^B\langle B^2\rangle\big) + 2\omega\sum_f c_X^f\,n_f\Big],\qquad \langle\Phi^\dagger\Phi\rangle = \frac{v^2+\mathcal{A}^2}{2}\tag{FE-39}$$

Symbols in (FE-39).

- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{\frac{7}{2}H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x}-\omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of the level of dirac16complex at the fold (a long-lived resonance) and $\bar{\chi}$ its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- μ_X, μ_{L}, μ_{R}: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_{\mathrm{L}}P_{\mathrm{L}} + \mu_{\mathrm{R}}P_{\mathrm{R}})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- P_{L}: The author's left-handed chiral projector, $P_{\mathrm{L}} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- P_{R}: The author's right-handed chiral projector, $P_{\mathrm{R}} = (\mathbf{1} + \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.

- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- c^G_X: The gluon-kinetic portal coupling of the half X of dirac16complex. Unit: GeV^{-7}. Status: INVENTED.
- $\langle G^2\rangle$: The mean square of the gluon field strength, $\langle G^A_{\mu\nu} G^{A\mu\nu}\rangle$ (the gluon condensate in the QCD vacuum, a thermal value in the plasma). Unit: GeV^4. Status: STANDARD.
- G^A_M, G^A_μ, G^A_{MN}: The gluon field of $SU(3)_C$ ($A = 1, \ldots, 8$): in the bulk G^A_M, a field of eight dimensions in its geometric normalisation (the covariant derivative $\partial - \mathrm{i}G$), and on the fold the canonically normalised four-dimensional field $G^A_\mu(x)$ of its massless mode, $g_3 G^A_\mu(x) = G^A_\mu(\sigma = 0, x)$; G^A_{MN} and $G^A_{\mu\nu}$ their field strengths. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- c^W_X: The W-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- $\langle W^2\rangle$: The mean square of the $SU(2)$ field strength, $\langle W^I_{\mu\nu} W^{I\mu\nu}\rangle$. Unit: GeV^4. Status: STANDARD.
- W^I_M, W^I_μ, W^I_{MN}: The gauge field of $SU(2)_L$ ($I = 1, 2, 3$): in the bulk W^I_M (geometric normalisation), on the fold the canonically normalised four-dimensional field $W^I_\mu(x)$ of its massless mode, $g_2 W^I_\mu(x) = W^I_\mu(\sigma = 0, x)$; W^I_{MN}, $W^I_{\mu\nu}$ their field strengths; $W^\pm_\mu = (W^1_\mu \mp \mathrm{i} W^2_\mu)/\sqrt{2}$. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- c^B_X: The B-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- $\langle B^2\rangle$: The mean square of the hypercharge field strength, $\langle B_{\mu\nu} B^{\mu\nu}\rangle$. Unit: GeV^4. Status: STANDARD.
- B_M, B_μ, B_{MN}: The hypercharge gauge field of $U(1)_Y$: in the bulk B_M (geometric normalisation), on the fold the canonically normalised four-dimensional field $B_\mu(x)$ of its massless mode, $g_1 B_\mu(x) = B_\mu(\sigma = 0, x)$; B_{MN}, $B_{\mu\nu}$ their field strengths. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- f: The label of a kind of Standard-Model multiplet: $f \in \{Q, u, d, \ell, e, \nu\}$ (quark doublets, up-quark singlets, down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.
- c^f_X: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c^ℓ_X and c^ν_X are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.
- n_f: The number density of the Standard-Model fermions of kind f (particles minus antiparticles, summed over generations and gauge components), the time component of the mean current $\langle J^\mu_f\rangle$ in the rest frame of the medium. Unit: GeV^3. Status: STANDARD.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\, G_\mathrm{F})^{-1/2}$ with the Fermi constant $G_\mathrm{F} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- $\mathcal{A}$: The author's ad hoc upper component of the Higgs vacuum, almightyAdHoc, $\mathcal{A}(x_4) = v_\mathcal{A} \mathrm{e}^{-2 a_{40} x_4}$. Unit: GeV. Status: AUTHOR.

$$
\begin{aligned}
&q_\mathrm{b} = -m_\Psi \left(\mu_\mathrm{L} + \mu_\mathrm{R}\right) \frac{\tanh \varsigma}{2\varsigma}, \qquad \varsigma = \sqrt{\mu_\mathrm{L}\mu_\mathrm{R}}, \qquad \omega_\mathrm{b} = \sqrt{m^2_\Psi - q^2_\mathrm{b}}, \\
&\omega_\mathrm{r} = \omega_\mathrm{b}\Big(1 - \frac{H}{2 q_\mathrm{b}}\Big) + \ldots, \qquad |\chi_\mathrm{b}(\sigma)| \propto \mathrm{e}^{-q_\mathrm{b}|\sigma|} \quad \big(|\sigma| \ll \sigma_\mathrm{t}\big)
\end{aligned}
\tag{FE-40}
$$

Symbols in (FE-40).

- q_b: The inverse width of a state of dirac16complex at the fold: inside the turning point its profile falls as $\mathrm{e}^{-q_\mathrm{b}|\sigma|}$. Unit: GeV. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.

- μ_X, μ_L, μ_R: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_\mathrm{L}P_\mathrm{L}+\mu_\mathrm{R}P_\mathrm{R})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- ς: The auxiliary variable of the bound-state condition, $\varsigma=\sqrt{\mu_\mathrm{L}\mu_\mathrm{R}}$ ($\varsigma_-=\sqrt{-\mu_\mathrm{L}\mu_\mathrm{R}}$ when $\mu_\mathrm{L}\mu_\mathrm{R}<0$). Unit: dimensionless. Status: DERIVED.
- ω_b: The energy at zero momentum of the state of dirac16complex at the fold in the limit $H\to 0$ (the bound state of UM-3): the particle χ of UM-4 has the mass $\omega_\mathrm{r}=\omega_\mathrm{b}(1-H/(2q_\mathrm{b}))+\ldots$, equal to ω_b up to the relative correction $H/(2q_\mathrm{b})$. Unit: GeV. Status: DERIVED.
- ω_r: The mass of the particle χ: the real part of the energy at rest of the resonance of dirac16complex at the fold, $\omega_\mathrm{r}=\omega_\mathrm{b}(1-H/(2q_\mathrm{b}))+\ldots$. Unit: GeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda=-21H^2$ and the tension $\kappa\lambda_\mathrm{F}=12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H>2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- χ_b: The profile of a state of dirac16complex at the fold (a resonance): it decays as $\mathrm{e}^{-q_\mathrm{b}|\sigma|}$ out to the turning point and continues beyond it as an outgoing wave of exponentially small amplitude. Unit: GeV^2. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma=|\cot z|\,\mathrm{d}x_0$, $\sigma=\mathrm{sgn}(\pi/2-z)\ln(\sin z)/(6H)$ on one period $0<z<\pi$; $\sigma=0$ is the fold $z=\pi/2$, $\sigma\to\mp\infty$ the tips $z\to 0,\pi$; one period of x_0 is the whole line $-\infty<\sigma<\infty$. Unit: length. Status: DERIVED.
- σ_t: The turning point of a state of dirac16complex of the frequency ω: the depth at which the warped frequency equals the bulk mass, $|\omega|\mathrm{e}^{H\sigma_\mathrm{t}}=m_\Psi$; beyond it every solution oscillates. Unit: length. Status: DERIVED.

The fold acts on dirac16complex as a chirality-dependent mass concentrated at $\sigma=0$, with the strengths μ_L and μ_R set by the mean fields of the Standard Model (FE-39); close to the fold the decaying solutions joined by FE-39 are the bound states of UM-3, with the inverse width q_b and the energy ω_b. In UM-4 the profile decays at the local rate $q(\sigma)$ of FE-37, which is q_b close to the fold and falls to 0 at the turning point, so that its amplitude there is $\exp(-F(\upsilon_\mathrm{b})m_\Psi/H)$ times that at the fold, and it continues beyond the turning point as an outgoing wave of exponentially small amplitude (FE-40). The warped frequency lowers the energy by $\omega_\mathrm{b}H\langle|\sigma|\rangle$ to the first order in H, with the mean distance $\langle|\sigma|\rangle=1/(2q_\mathrm{b})$. Status: DERIVED.

$$\begin{aligned}&\omega_0=\omega_\mathrm{r}-\frac{\mathrm{i}}{2}\,\Gamma_\chi,\qquad \Gamma_\chi=p_\mathrm{w}\,m_\Psi\exp\Big(-2F(\upsilon_\mathrm{b})\,\frac{m_\Psi}{H}\Big),\qquad p_\mathrm{w}\simeq\frac{2}{\mathrm{e}}\,\frac{q_\mathrm{b}^2}{m_\Psi^2},\\&F(\upsilon)=\ln\frac{1+\sqrt{1-\upsilon^2}}{\upsilon}-\sqrt{1-\upsilon^2},\qquad \upsilon_\mathrm{b}=\frac{\omega_\mathrm{b}}{m_\Psi}\end{aligned}\tag{FE-86}$$

Symbols in (FE-86).

- ω_0: The energy at rest (zero momentum along ordinary space and along the extra times) of a state of dirac16complex at the fold; complex for a resonance, $\omega_0=\omega_\mathrm{r}-\frac{\mathrm{i}}{2}\Gamma_\chi$. Unit: GeV. Status: DERIVED.
- ω_r: The mass of the particle χ: the real part of the energy at rest of the resonance of dirac16complex at the fold, $\omega_\mathrm{r}=\omega_\mathrm{b}(1-H/(2q_\mathrm{b}))+\ldots$. Unit: GeV. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- Γ_χ: The width of the particle χ at rest: the rate of its decay into the waves of dirac16complex in the bulk. Unit: GeV. Value: below $1/(11.6\,t_0)=1.3\times10^{-43}$ GeV for dark matter (a lifetime above 160 Gyr). Status: DERIVED.
- p_w: The prefactor of the width of the particles χ, $\Gamma_\chi=p_\mathrm{w}m_\Psi\mathrm{e}^{-2F(\upsilon_\mathrm{b})m_\Psi/H}$: $p_\mathrm{w}\simeq(2/\mathrm{e})(q_\mathrm{b}/m_\Psi)^2$ to the leading order in H/q_b (FE-86; 0.184 for $q_\mathrm{b}=m_\Psi/2$). Unit: dimensionless. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- $F(\upsilon)$: The tunnelling exponent of the width of the particles χ per unit of m_Ψ/H: the integral of the local decay rate q over the region inside the turning point, divided by m_Ψ/H, as a function of the ratio υ of the energy to the bulk mass. Unit: dimensionless. Value: 0.049 for $\upsilon=0.866$; $(q_\mathrm{b}/\omega_\mathrm{b})^3/3$ for weak binding. Status: DERIVED.

- υ, υ_b, υ_r: The argument of the tunnelling exponent F, a ratio of an energy to the bulk mass m_Ψ; $\upsilon_\mathrm{b} = \omega_\mathrm{b}/m_\Psi$ that of the particle χ in the limit $H/q_\mathrm{b} \to 0$, $\upsilon_\mathrm{r} = \omega_\mathrm{r}/m_\Psi$ that of the real part of its resonance. Unit: dimensionless. Status: DERIVED.
- ω_b: The energy at zero momentum of the state of dirac16complex at the fold in the limit $H \to 0$ (the bound state of UM-3): the particle χ of UM-4 has the mass $\omega_\mathrm{r} = \omega_\mathrm{b}(1 - H/(2q_\mathrm{b})) + \ldots$, equal to ω_b up to the relative correction $H/(2q_\mathrm{b})$. Unit: GeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- q_b: The inverse width of a state of dirac16complex at the fold: inside the turning point its profile falls as $\mathrm{e}^{-q_\mathrm{b}|\sigma|}$. Unit: GeV. Status: DERIVED.

The resonances: the solutions with outgoing waves on both sides of the fold exist only at the complex energies ω_0, with the mass ω_r of the particle χ and its width Γ_χ, the rate of the tunnelling through the region inside the turning point. The prefactor follows from the uniform asymptotic forms of the Hankel functions of FE-85 for large m_Ψ/H: the ratio of the two outgoing solutions at the fold is real up to an imaginary part of the relative size $\exp(-2F(\upsilon)m_\Psi/H)\, q_\mathrm{b}/\omega_\mathrm{b}$, the matching turns it into the width $2(q_\mathrm{b}^2/m_\Psi)\exp(-2F(\upsilon_\mathrm{r})m_\Psi/H)$ at the real part of the resonance, and the shift of FE-40 raises the exponent by 1: $p_\mathrm{w} = (2/\mathrm{e})(q_\mathrm{b}/m_\Psi)^2$ to the leading order in H/q_b, 0.184 for $q_\mathrm{b} = m_\Psi/2$, 0.133 for $(\mu_\mathrm{L}, \mu_\mathrm{R}) = (-0.6, -0.3)$ and 0.721 for $q_\mathrm{b} = 0.99\, m_\Psi$. The computed widths approach it like H/m_Ψ (for $(-1, 0)$: 0.16614, 0.18239 and 0.18356 at $m_\Psi/H = 100$, 1000 and 4000; for $q_\mathrm{b} = 0.99\, m_\Psi$: 0.965, 0.988 and 0.997 of it at $m_\Psi/H = 20$, 60 and 200). Status: DERIVED (`um4_spectra.py`, U3 and U4; `um4_kohnsham.py`, K8: the tunnelling exponent at the resonance exceeds $2F(\upsilon_\mathrm{b})m_\Psi/H$ by 1.002).

$(\mu_\mathrm{L}, \mu_\mathrm{R})$	m_Ψ/H	10	20	30	40	$H \to 0$ (UM-3)
$(-1, 0)$	ω_r/m_Ψ	0.7785	0.8175	0.8333	0.8418	0.8660
$(-1, 0)$	Γ_χ/m_Ψ	0.0587	0.0180	0.00670	0.00263	0
$(-0.6, -0.3)$	ω_r/m_Ψ	0.8098	0.8493	0.8657	0.8750	0.9053
$(-0.6, -0.3)$	Γ_χ/m_Ψ	0.0865	0.0344	0.0172	0.00933	0

The resonances computed with the author's exact 16×16 matrices: the zeros of the matching determinant with outgoing waves on both sides (an eight-dimensional kernel: the resonances of the eight copies coincide), for a fold bound by the Higgs portal alone and for one with both halves attracted. As $H/m_\Psi \to 0$ the mass tends to the bound-state energy of UM-3, ω_b. The slope of $\ln\Gamma_\chi$ in m_Ψ/H between 30 and 40 is -0.0937 and -0.0609, against $-2F(\upsilon_\mathrm{b}) = -0.0986$ and -0.0575 of FE-86. Of the eight resonances at $+\omega_\mathrm{r}$ four have the Krein charge $+1$ and four -1, as the bound states of UM-3, and the mirror R_1 of the fold (FD-10) maps them onto themselves, four even (Krein charge -1) and four odd (Krein charge $+1$). Status: DERIVED (`um4_spectra.py`, U3, U4, U7 and U8).

$$\Gamma_\chi < \frac{1}{11.6\, t_0} \quad \Leftrightarrow \quad \frac{m_\Psi}{H} > \frac{\ln\left(11.6\, p_\mathrm{w}\, m_\Psi\, t_0\right)}{2F(\upsilon_\mathrm{b})}, \qquad p_\mathrm{w} \simeq \frac{2}{\mathrm{e}}\,\frac{q_\mathrm{b}^2}{m_\Psi^2}, \qquad F(\upsilon_\mathrm{b}) \simeq \frac{1}{3}\Big(\frac{q_\mathrm{b}}{\omega_\mathrm{b}}\Big)^3 - \frac{3}{10}\Big(\frac{q_\mathrm{b}}{\omega_\mathrm{b}}\Big)^5 \quad \Big(\frac{q_\mathrm{b}}{\omega_\mathrm{b}} \ll 1\Big) \tag{S-6}$$

Symbols in (S-6).

- Γ_χ: The width of the particle χ at rest: the rate of its decay into the waves of dirac16complex in the bulk. Unit: GeV. Value: below $1/(11.6\, t_0) = 1.3 \times 10^{-43}$ GeV for dark matter (a lifetime above 160 Gyr). Status: DERIVED.
- t_0: The age of the universe. Unit: time. Value: 13.8 Gyr. Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- p_w: The prefactor of the width of the particles χ, $\Gamma_\chi = p_\mathrm{w} m_\Psi \mathrm{e}^{-2F(\upsilon_\mathrm{b})m_\Psi/H}$: $p_\mathrm{w} \simeq (2/\mathrm{e})(q_\mathrm{b}/m_\Psi)^2$ to the leading order in H/q_b (FE-86; 0.184 for $q_\mathrm{b} = m_\Psi/2$). Unit: dimensionless. Status: DERIVED.

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- $F(\upsilon)$: The tunnelling exponent of the width of the particles χ per unit of m_Ψ/H: the integral of the local decay rate q over the region inside the turning point, divided by m_Ψ/H, as a function of the ratio υ of the energy to the bulk mass. Unit: dimensionless. Value: 0.049 for $\upsilon = 0.866$; $(q_\mathrm{b}/\omega_\mathrm{b})^3/3$ for weak binding. Status: DERIVED.
- υ, υ_b, υ_r: The argument of the tunnelling exponent F, a ratio of an energy to the bulk mass m_Ψ; $\upsilon_\mathrm{b} = \omega_\mathrm{b}/m_\Psi$ that of the particle χ in the limit $H/q_\mathrm{b} \to 0$, $\upsilon_\mathrm{r} = \omega_\mathrm{r}/m_\Psi$ that of the real part of its resonance. Unit: dimensionless. Status: DERIVED.
- q_b: The inverse width of a state of dirac16complex at the fold: inside the turning point its profile falls as $\mathrm{e}^{-q_\mathrm{b}|\sigma|}$. Unit: GeV. Status: DERIVED.
- ω_b: The energy at zero momentum of the state of dirac16complex at the fold in the limit $H \to 0$ (the bound state of UM-3): the particle χ of UM-4 has the mass $\omega_\mathrm{r} = \omega_\mathrm{b}(1 - H/(2q_\mathrm{b})) + \ldots$, equal to ω_b up to the relative correction $H/(2q_\mathrm{b})$. Unit: GeV. Status: DERIVED.

The lifetime that the dark matter requires. The particles χ that make up the dark matter today must live longer than 160 Gyr, 11.6 times the age $t_0 = 13.8$ Gyr of the universe: the bound on the lifetime of dark matter that decays into invisible relativistic particles, from the cosmic microwave background and the large-scale structure (B. Audren, J. Lesgourgues, G. Mangano, P. D. Serpico and T. Tram, JCAP 12 (2014) 028, arXiv:1407.2418), $\Gamma_\chi < 1.3 \times 10^{-43}$ GeV. With the width of FE-86 this bounds m_Ψ/H from below; the tunnelling exponent grows with the binding, and for weak binding it is small, $F \simeq (q_\mathrm{b}/\omega_\mathrm{b})^3/3$, so that the bound grows like $(\omega_\mathrm{b}/q_\mathrm{b})^3$. With H at its lower bound 2534 GeV (the bounds use the exact widths at the bound, whose prefactor lies within 4 per cent of p_w, or p_w itself for the weak binding):

q_b/m_Ψ	$q_\mathrm{b}/\omega_\mathrm{b}$	$F(\upsilon_\mathrm{b})$	p_w	m_Ψ/H above	ω_b above (GeV)
0.0995	0.100	3.30×10^{-4}	0.0073	1.72×10^5	4.34×10^8
0.3	0.314	9.52×10^{-3}	0.066	5911	1.43×10^7
0.5	0.577	0.0493	0.184	1135	2.49×10^6
0.7	0.980	0.167	0.361	333	6.02×10^5
0.8	1.333	0.299	0.471	186	2.83×10^5
0.9	2.065	0.572	0.596	96.7	1.07×10^5
0.95	3.04	0.882	0.664	62.5	4.95×10^4
0.99	7.02	1.66	0.721	33.1	1.18×10^4
0.999	22.3	2.80	0.734	19.5	2.21×10^3

For the age t_0 instead of 160 Gyr the bounds are about 2 per cent lower ($q_\mathrm{b} = m_\Psi/2$: $m_\Psi/H > 1110$, $\omega_\mathrm{b} > 2.44 \times 10^6$ GeV), for $100\,t_0$ 2 per cent higher. The particles χ of UM-4 are therefore heavier than 2.5×10^6 GeV for $q_\mathrm{b} = m_\Psi/2$, more for weaker binding; the bound vanishes as $q_\mathrm{b} \to m_\Psi$, and strongly bound particles can be much lighter (1.2×10^4 GeV for $q_\mathrm{b} = 0.99\,m_\Psi$, where $H/q_\mathrm{b} = 0.03$). A particle moving with the momentum $\vec{k}$ has the energy $\sqrt{\omega_0^2 + \vec{k}^2}$ (FE-38) and the width $\Gamma_\chi \omega_\mathrm{r}/\sqrt{\omega_\mathrm{r}^2 + \vec{k}^2}$ (the time dilation of the boost). Status: DERIVED (`um4_spectra.py`, U6: the exact widths at the bounds), ESTIMATE (the measured bound on the lifetime is an input).

$$
\begin{aligned}
&Q_\Psi = \int \mathrm{d}\sigma\, \mathrm{d}^3x\, \sqrt{|g|}\, \mathrm{e}^{H|\sigma|}\, \Psi^\dagger \mathfrak{B} \Psi = b_4^3 \int \mathrm{d}\sigma\, \mathrm{d}^3x\, \mathrm{e}^{H|\sigma|}\, \chi^\dagger \mathfrak{B} \chi, \\
&\mathfrak{B} = \mathrm{i}\,\sigma_{16}\,\gamma^{(4)}, \qquad \mathfrak{B}^\dagger = \mathfrak{B}, \qquad \mathfrak{B}^2 = \mathbf{1}, \qquad E_o = -\omega_o\, \mathcal{Q}_o
\end{aligned}
\tag{FE-43}
$$

Symbols in (FE-43).

- Q_Ψ: The conserved charge of dirac16complex (the Noether charge of a change of the phase of Ψ by a constant), a Krein form (indefinite); as written it is the charge per unit coordinate volume of the extra times, on which no field depends. Unit: GeV^3. Status: DERIVED.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- x: The coordinates of the fold: x_1, x_2, x_3 of ordinary space, x_4 the time, x_5, x_6, x_7 the extra times; a field of the fold is a function of the four observable coordinates $x = (x_1, x_2, x_3, x_4)$, and $\mathrm{d}^4x = \mathrm{d}x_1\,\mathrm{d}x_2\,\mathrm{d}x_3\,\mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\mathfrak{B}$: The Krein operator, $\mathfrak{B} = \mathrm{i}\sigma_{16}\gamma^{(4)}$: a Hermitian involution with the signature (8, 8); $\Psi^\dagger\mathfrak{B}\Psi$ is the charge density. Unit: dimensionless. Status: DERIVED.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{\frac{7}{2}H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of the level of dirac16complex at the fold (a long-lived resonance) and $\bar\chi$ its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- σ_{16}: The matrix of the Dirac adjoint, $\sigma_{16} = T_0T_1T_2T_3$, real and symmetric; $\sigma_{16}T_a$ is real and antisymmetric. Unit: dimensionless. Status: AUTHOR.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- $\mathbf{1}$, $\mathbf{1}_4$, $\mathbf{1}_8$, $\mathbf{1}_{16}$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_4$, $\mathbf{1}_8$ and $\mathbf{1}_{16}$ are the identity matrices of 4, 8 and 16 rows. Unit: dimensionless. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- $\mathcal{Q}_o$: The charge Q_Ψ (the Krein charge) of the single-particle state o of dirac16complex, a Kohn-Sham orbital or a mode: $+1$ or -1 for a normalised state. Unit: dimensionless. Status: DERIVED.

The conserved charge of dirac16complex (the Noether charge of a constant change of its phase) and the energy of a mode: the charge density carries the Krein operator $\mathfrak{B}$, which is indefinite because the author's matrices have the signature (4, 4), and the measure $\mathrm{e}^{H|\sigma|}$ of the profiles in UM-4 (UM-3: 1); a mode of frequency ω_o and charge $\mathcal{Q}_o$ carries the energy $E_o = -\omega_o\mathcal{Q}_o$. Status: DERIVED.

8.1 The momentum along the extra times

$$\mathrm{e}^{2H|\sigma|}\Big(\omega^2 - \mathrm{e}^{-2a_4}\,\vec{k}^2 + \mathrm{e}^{2a_4}\,b_4^{-2}\,\vec{\kappa}^2\Big) = m_\Psi^2 + p_\sigma^2, \qquad \omega^2 = \omega_0^2 + \vec{k}^2 - \vec{\kappa}^2 \quad \big(a_4 = 0,\ b_4 = 1\big) \tag{FE-38}$$

Symbols in (FE-38).

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $\vec{k}$: The (comoving) momentum along ordinary space, $\vec{k}^2$ its square. Unit: GeV. Status: STANDARD.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- $\vec{\kappa}$: A momentum along the extra times (three components); the fields of UM-4 have $\vec{\kappa} = 0$. Unit: GeV. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- p_σ: The momentum along the hidden direction (local wave number of a profile along σ). Unit: GeV. Status: DERIVED.
- ω_0: The energy at rest (zero momentum along ordinary space and along the extra times) of a state of dirac16complex at the fold; complex for a resonance, $\omega_0 = \omega_\mathrm{r} - \frac{\mathrm{i}}{2}\Gamma_\chi$. Unit: GeV. Status: DERIVED.

The local dispersion relation of dirac16complex and the exact dispersion of every state at the fold: the boosts along ordinary space and the rotations of the time into an extra time (generated by $\frac{1}{2}\gamma^{(4)}\gamma^{(5)}$, which commutes with $\gamma^{(0)}$, with Γ_8 and with the mass term) are symmetries, because the warp multiplies the frequency and the momenta alike. A momentum along a time-like direction enters with the sign of the frequency: a state with $|\vec{\kappa}| \le \omega_0$ is the rotation of the state at rest in the plane of the time and an extra time, while a state with $|\vec{\kappa}| > \omega_0$ is reached by no rotation, has $\omega^2 < 0$ and grows exponentially in x_4 (a tachyon). Computed with the author's matrices at $H = 10^{-4}m_\Psi$, $q_\mathrm{b} = 0.3\,m_\Psi$ ($\omega_0 = 0.95378\,m_\Psi$): $\omega = 0.94852$, 0.90537 and 0.74141 m_Ψ for $|\vec{\kappa}| = 0.1$, 0.3 and 0.6 m_Ψ, and $\omega = 0.95901$, 0.99985 and 1.12681 m_Ψ for the same $|\vec{k}|$, equal to the predictions to ten digits. The same holds for every field, in the bulk as on the fold: a field of the mass m has growing modes for every $\vec{\kappa}^2 > m^2 + \vec{k}^2$, a massless field for every $|\vec{\kappa}| > |\vec{k}|$. UM-4 therefore keeps only the fields that do not depend on the extra times (ASSUMED); the restriction is consistent (the momentum along the extra times is conserved) but not stable (interactions create pairs of opposite momenta, an ESTIMATE). With frozen extra times and a constant expansion rate a_4' a massless mode at rest along ordinary space ($\vec{k} = 0$) with extra-time momentum grows as a single exponential, at the rate $(-3a_4' + \sqrt{9a_4'^2 + 4\vec{\kappa}^2})/2$ (0.646586 for $a_4' = 0.3$ and $|\vec{\kappa}| = 1$, exactly the formula); with a mass m, $\vec{\kappa}^2$ is replaced by $\vec{\kappa}^2 - m^2$, and with $\vec{k} \neq 0$ the term $\mathrm{e}^{-2a_4}\vec{k}^2$ redshifts and the growth is not a single exponential. Compact extra times of the period L_t do not help: the first harmonic of a massless field stays below one e-folding over the age of the universe only for $L_\mathrm{t} \ge 26.6$ Gpc on the static fold (about 13.5 Gpc with the Hubble friction of today, an order of magnitude), but the n-th harmonic grows n times faster, so infinitely many modes grow for every L_t (and a compact time-like direction is itself a closed time-like curve). UM-4 needs a gauge principle that removes the momentum along the extra times of every field. Status: DERIVED (`um4_extratimes.py`, EX1 to EX5), ASSUMED (the restriction), OPEN (the gauge principle).

9 The Kohn-Sham states of dirac16complex

dirac16complex is a quantised field. Its states with many particles are described here with the Kohn-Sham equations of the record (FE-41, FE-42): single-particle orbitals in the mean fields of everything else, iterated until the orbitals reproduce the fields they move in. The metric is held fixed (the back-reaction of the gas on the warp is left out, ASSUMED); the fields that are iterated are the mean field of the self-interaction and, for the Higgs loop, the strength μ_L of the fold. The state solved is a homogeneous gas of the particles χ at rest on the static block of the late universe (a_4 and b_4 constant), with their self-interaction, with gravity (the warp of the metric and the canonical spin connection) and with every field of the Standard Model, which acts through its mean values at the fold (the

strengths μ_{L} and μ_{R} of FE-39); the Higgs portal also feeds back, because the left-handed half of the gas shifts the local electroweak vacuum v_{loc} of FE-49.

$$\big[\gamma^M D_M - m_\Psi - \lambda_\Psi\, n_{\mathrm{s}} + \lambda_\Psi\, \mathcal{N}\big]\varphi_o = \delta_{\mathrm{D}}(\sigma)\,\frac{1}{V_*}\,\langle \mathbb{M}\rangle\,\varphi_o,$$

$$\mathcal{N} = \sum_o \mathsf{n}_o\,\varphi_o\bar{\varphi}_o, \qquad n_{\mathrm{s}} = \mathrm{tr}\,\mathcal{N} = \sum_o \mathsf{n}_o\,\bar{\varphi}_o\varphi_o, \qquad \mathsf{n}_o = \frac{1}{\mathrm{e}^{(E_o-\mu_\Psi)/\mathsf{T}_\Psi}+1},$$

$$\chi_o' = \gamma^{(0)}\Big[\mathrm{i}\omega\,\mathrm{e}^{H|\sigma|}\gamma^{(4)} + m_\Psi + \mathcal{M}(\sigma)\Big]\chi_o, \qquad \mathcal{M} = G_{\mathrm{KS}}\,\mathrm{e}^{7H|\sigma|}\big(\hat{\rho}\,\mathbf{1} - \hat{\mathcal{N}}\big), \qquad \hat{\rho} = \frac{1}{4}\sum_o \bar{\chi}_o\chi_o, \qquad \hat{\mathcal{N}} = \frac{1}{4}\sum_o \chi_o\bar{\chi}_o$$

(FE-41)

Symbols in (FE-41).

- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC}G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- λ_Ψ: The coupling of the contact self-interaction of dirac16complex. Unit: GeV^{-6}. Status: RECORD.
- n_{s}: The scalar density of dirac16complex in the Kohn-Sham state, $n_{\mathrm{s}} = \langle\bar{\Psi}\Psi\rangle = \mathrm{tr}\,\mathcal{N}$. Unit: GeV^7. Status: DERIVED.
- $\mathcal{N}$: The one-body density matrix of dirac16complex, $\mathcal{N} = \sum_o \mathsf{n}_o\varphi_o\bar{\varphi}_o$ (a 16×16 matrix at each point). Unit: GeV^7. Status: DERIVED.
- φ_o, $\bar{\varphi}_o$: A Kohn-Sham orbital of dirac16complex (a single-particle solution of the Kohn-Sham equation: the Dirac equation of dirac16complex with the mean field of the gas added to its mass) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- δ_{D}: The Dirac delta function of σ: zero for $\sigma \neq 0$, with the integral 1 over any range that contains $\sigma = 0$; the integral of $\delta_{\mathrm{D}}(\sigma)$ times a smooth function of σ is the value of that function at $\sigma = 0$. Unit: 1/length (for $\delta_{\mathrm{D}}(\sigma)$). Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- $\mathbb{M}$: The fold operator of dirac16complex: the 16×16 matrix (with a derivative along the fold) built from the Standard-Model fields at the fold by varying the portals with respect to $\bar{\Psi}$. Unit: GeV^{-3} (so that $\mathbb{M}/V_*$ is dimensionless). Status: DERIVED.
- n_o: The occupation of the Kohn-Sham orbital o (between 0 and 1). Unit: dimensionless. Status: RECORD.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- μ_Ψ: The chemical potential of dirac16complex. Unit: GeV. Status: DERIVED.
- T_Ψ: The temperature of dirac16complex. Unit: GeV. Status: DERIVED.
- $\chi_o(\sigma)$, $\bar{\chi}_o$: The profile of the occupied Kohn-Sham orbital o ($o = 1, \ldots, 4$, the four positive-energy states of

the level at the fold) of a homogeneous gas of the particles χ at rest on the static block: a 16-component spinor function of σ; the orbital of each particle is $\mathrm{e}^{\frac{7}{2}H|\sigma|}\chi_o(\sigma)\mathrm{e}^{-\mathrm{i}\omega x_4}$ divided by the square root of the volume of ordinary space and of V_*; normalised to the Krein charge -1 with the measure of UM-4, $\int \mathrm{d}\sigma\, \mathrm{e}^{H|\sigma|}\chi_o^\dagger \mathfrak{B}\chi_o = -1$; $\bar\chi_o = \chi_o^\dagger\sigma_{16}$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{1/2}$. Status: DERIVED.

- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- $\mathcal{M}(\sigma)$: The mean field of the self-interaction in the Kohn-Sham equation of the gas: the 16×16 matrix $\lambda_\Psi(n_\mathrm{s}\mathbf{1} - \mathcal{N})$ at each σ, which adds to the mass m_Ψ; positive (repulsive) for $\lambda_\Psi > 0$. Unit: GeV. Status: DERIVED.
- G_KS: The dimensionless strength of the self-interaction of the gas of particles χ in the Kohn-Sham states: the self-coupling λ_Ψ times the number density n_χ of the particles on the fold divided by the volume V_* of the extra times, $G_\mathrm{KS} = \lambda_\Psi n_\chi/V_*$ (in units of m_Ψ when numbers are given). Unit: dimensionless. Value: 10^{-6} to 0.5 in the solutions. Status: DERIVED.
- $\hat\rho(\sigma)$: The scalar density of the occupied orbitals, $\hat\rho = \frac{1}{4}\sum_o \bar\chi_o\chi_o$ (per particle); the scalar density of the gas is $n_\mathrm{s} = (n_\chi/V_*)\,\mathrm{e}^{7H|\sigma|}\hat\rho$. Unit: GeV. Status: DERIVED.
- $\mathbf{1}$, $\mathbf{1}_4$, $\mathbf{1}_8$, $\mathbf{1}_{16}$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_4$, $\mathbf{1}_8$ and $\mathbf{1}_{16}$ are the identity matrices of 4, 8 and 16 rows. Unit: dimensionless. Status: STANDARD.
- $\hat{\mathcal{N}}(\sigma)$: The one-body density matrix of the occupied orbitals, $\hat{\mathcal{N}} = \frac{1}{4}\sum_o \chi_o\bar\chi_o$ (a 16×16 matrix at each σ, per particle); the density matrix of the gas is $\mathcal{N} = (n_\chi/V_*)\,\mathrm{e}^{7H|\sigma|}\hat{\mathcal{N}}$. Unit: GeV. Status: DERIVED.

$$\langle \mathcal{T}_{\Psi MN}\rangle = -\frac{1}{4}\sum_o \mathsf{n}_o\Big(\bar\varphi_o\gamma_M D_N\varphi_o + \bar\varphi_o\gamma_N D_M\varphi_o - (D_M\bar\varphi_o)\gamma_N\varphi_o - (D_N\bar\varphi_o)\gamma_M\varphi_o\Big) + \frac{\lambda_\Psi}{2}\, g_{MN}\big(n_\mathrm{s}^2 - \mathrm{tr}\,\mathcal{N}^2\big) \quad \text{(FE-42)}$$

Symbols in (FE-42).

- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- $\mathcal{T}_\Psi$: The energy-momentum tensor of dirac16complex: $-2/\sqrt{|g|}$ times the variation of the action S_Ψ of dirac16complex with respect to g^{MN}. Unit: GeV^8. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- n_o: The occupation of the Kohn-Sham orbital o (between 0 and 1). Unit: dimensionless. Status: RECORD.
- φ_o, $\bar\varphi_o$: A Kohn-Sham orbital of dirac16complex (a single-particle solution of the Kohn-Sham equation: the Dirac equation of dirac16complex with the mean field of the gas added to its mass) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar\Psi = \partial_M\bar\Psi - \bar\Psi\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat h^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC}G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- λ_Ψ: The coupling of the contact self-interaction of dirac16complex. Unit: GeV^{-6}. Status: RECORD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant

($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- n_s: The scalar density of dirac16complex in the Kohn-Sham state, $n_\mathrm{s} = \langle\bar\Psi\Psi\rangle = \mathrm{tr}\,\mathcal{N}$. Unit: GeV^7. Status: DERIVED.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- $\mathcal{N}$: The one-body density matrix of dirac16complex, $\mathcal{N} = \sum_o \mathsf{n}_o\varphi_o\bar\varphi_o$ (a 16×16 matrix at each point). Unit: GeV^7. Status: DERIVED.

The Kohn-Sham equations of dirac16complex coupled to itself, to gravity and to every Standard-Model field (FE-41) and the energy-momentum tensor of the Kohn-Sham state (FE-42): each orbital φ_o obeys the field equation with the mean fields, the self-interaction contributes the Hartree term $\lambda_\Psi n_\mathrm{s}$ and the exact exchange term $-\lambda_\Psi\mathcal{N}$, and every Standard-Model field enters through the mean fold operator $\langle\mathbb{M}\rangle$; correlation beyond exchange and the sea of negative-energy states are left out. Status: RECORD (the Kohn-Sham treatment), DERIVED (the exchange term), ASSUMED (no correlation, no sea).

9.1 The equations of the gas of particles χ

$$\begin{aligned}
&\Psi = \mathrm{e}^{\frac{7}{2}H|\sigma|}\,\chi_o(\sigma)\,\mathrm{e}^{-\mathrm{i}\omega x_4}, \qquad \chi_o' = \gamma^{(0)}\Big(\mathrm{i}\omega\,\mathrm{e}^{H|\sigma|}\gamma^{(4)} + m_\Psi + \mathcal{M}(\sigma)\Big)\chi_o \qquad (\sigma\neq0),\\
&\mathcal{M}(\sigma) = G_\mathrm{KS}\,\mathrm{e}^{7H|\sigma|}\Big(\hat\rho(\sigma)\,\mathbf{1} - \hat{\mathcal{N}}(\sigma)\Big), \qquad \hat\rho = \frac14\sum_{o=1}^{4}\bar\chi_o\chi_o, \qquad \hat{\mathcal{N}} = \frac14\sum_{o=1}^{4}\chi_o\bar\chi_o,\\
&\chi_o(0^+) = \exp\Big(\gamma^{(0)}\big(\mu_\mathrm{L}P_\mathrm{L} + \mu_\mathrm{R}P_\mathrm{R}\big)\Big)\chi_o(0^-), \qquad \int\mathrm{d}\sigma\,\mathrm{e}^{H|\sigma|}\,\chi_o^\dagger\mathfrak{B}\chi_p = -\delta_{op}, \qquad G_\mathrm{KS} = \frac{\lambda_\Psi\,n_\chi}{V_*}
\end{aligned} \tag{S-7}$$

Symbols in (S-7).

- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: GeV$^{7/2}$. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\chi_o(\sigma)$, $\bar\chi_o$: The profile of the occupied Kohn-Sham orbital o ($o = 1,\ldots,4$, the four positive-energy states of the level at the fold) of a homogeneous gas of the particles χ at rest on the static block: a 16-component spinor function of σ; the orbital of each particle is $\mathrm{e}^{\frac{7}{2}H|\sigma|}\chi_o(\sigma)\mathrm{e}^{-\mathrm{i}\omega x_4}$ divided by the square root of the volume of ordinary space and of V_*; normalised to the Krein charge -1 with the measure of UM-4, $\int\mathrm{d}\sigma\,\mathrm{e}^{H|\sigma|}\chi_o^\dagger\mathfrak{B}\chi_o = -1$; $\bar\chi_o = \chi_o^\dagger\sigma_{16}$ is its Dirac adjoint. Unit: GeV$^{1/2}$. Status: DERIVED.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number (UM-4 keeps only $p = 1$); in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- $\mathcal{M}(\sigma)$: The mean field of the self-interaction in the Kohn-Sham equation of the gas: the 16×16 matrix $\lambda_\Psi(n_\mathrm{s}\mathbf{1} - \mathcal{N})$ at each σ, which adds to the mass m_Ψ; positive (repulsive) for $\lambda_\Psi > 0$. Unit: GeV. Status: DERIVED.

- G_{KS}: The dimensionless strength of the self-interaction of the gas of particles χ in the Kohn-Sham states: the self-coupling λ_Ψ times the number density n_χ of the particles on the fold divided by the volume V_* of the extra times, $G_{\mathrm{KS}} = \lambda_\Psi n_\chi / V_*$ (in units of m_Ψ when numbers are given). Unit: dimensionless. Value: 10^{-6} to 0.5 in the solutions. Status: DERIVED.
- $\hat{\rho}(\sigma)$: The scalar density of the occupied orbitals, $\hat{\rho} = \frac{1}{4}\sum_o \bar{\chi}_o \chi_o$ (per particle); the scalar density of the gas is $n_{\mathrm{s}} = (n_\chi / V_*)\, \mathrm{e}^{7H|\sigma|}\hat{\rho}$. Unit: GeV. Status: DERIVED.
- $\mathbf{1}$, $\mathbf{1}_4$, $\mathbf{1}_8$, $\mathbf{1}_{16}$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_4$, $\mathbf{1}_8$ and $\mathbf{1}_{16}$ are the identity matrices of 4, 8 and 16 rows. Unit: dimensionless. Status: STANDARD.
- $\hat{\mathcal{N}}(\sigma)$: The one-body density matrix of the occupied orbitals, $\hat{\mathcal{N}} = \frac{1}{4}\sum_o \chi_o \bar{\chi}_o$ (a 16×16 matrix at each σ, per particle); the density matrix of the gas is $\mathcal{N} = (n_\chi / V_*)\, \mathrm{e}^{7H|\sigma|}\hat{\mathcal{N}}$. Unit: GeV. Status: DERIVED.
- μ_X, μ_{L}, μ_{R}: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_{\mathrm{L}} P_{\mathrm{L}} + \mu_{\mathrm{R}} P_{\mathrm{R}})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- P_{L}: The author's left-handed chiral projector, $P_{\mathrm{L}} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- P_{R}: The author's right-handed chiral projector, $P_{\mathrm{R}} = (\mathbf{1} + \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\mathfrak{B}$: The Krein operator, $\mathfrak{B} = \mathrm{i}\sigma_{16}\gamma^{(4)}$: a Hermitian involution with the signature (8, 8); $\Psi^\dagger \mathfrak{B} \Psi$ is the charge density. Unit: dimensionless. Status: DERIVED.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- λ_Ψ: The coupling of the contact self-interaction of dirac16complex. Unit: GeV^{-6}. Status: RECORD.
- n_χ: The number density of the dark-matter particles χ (particles plus antiparticles). Unit: GeV^3. Status: DERIVED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4} b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\, \mathrm{d}x_6\, \mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4} b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.

FE-41 for this state, with $m_\Psi = 1$ as the unit wherever numbers are given. The factor $\mathrm{e}^{(7/2)H|\sigma|}$ removes the first term of the spin connection FE-14 exactly, and the second vanishes on the static block, so the four occupied orbitals $\chi_o(\sigma)$ obey a Dirac equation of one variable in which the warped frequency $\omega \mathrm{e}^{H|\sigma|}$ appears (in UM-3 the factors were $\mathrm{e}^{3H|\sigma|}$ and 1). Gravity enters through the warp: the proper scalar density of the gas at σ is $\mathrm{e}^{7H|\sigma|}$ times the density of the profiles, so the mean field $\mathcal{M}$ of the self-interaction (the Hartree part $\hat{\rho}\,\mathbf{1}$ and the exchange part $\hat{\mathcal{N}}$, the one-body density matrix) carries that factor. The Standard Model acts through the matching at the fold (FE-39). The occupied orbitals are the four states of the level at the fold with positive energy: Krein charge −1 each with the measure $\mathrm{e}^{H|\sigma|}\mathrm{d}\sigma$ of UM-4 (FE-43), energy $-\omega\mathcal{Q}_o = \omega$, mutually Krein-orthogonal. The states are resonances: beyond the turning point every solution oscillates. The computation therefore takes the parameters deep in the regime of long-lived particles, $H = 10^{-4} m_\Psi$ ($m_\Psi / H = 10^4$) with $\mu_{\mathrm{L}} = -0.6$, $\mu_{\mathrm{R}} = 0$ ($q_{\mathrm{b}} = 0.3 = 3000H$, $\omega_{\mathrm{b}} = 0.953939$ in the limit $H \to 0$), and solves the orbitals in the box $|\sigma| \le 40$ (with the step 0.03), far inside the turning point $\sigma_{\mathrm{t}} = 473$, with the decaying solutions at its end; the normalisation integral of S-7 runs over the box. The exponentially small outgoing part beyond the turning point is the width, which follows from the tunnelling exponent (S-12). G_{KS} is the dimensionless strength of the self-interaction: the self-coupling λ_Ψ times the number density n_χ of the particles on the fold, divided by the volume V_* of the extra times. Status: DERIVED (from FE-41, FE-14, FE-39), ASSUMED (the parameters).

$$\Sigma^{12}\chi_1 = \Sigma^{56}\chi_1 = \frac{\mathrm{i}}{2}\,\chi_1 \tag{S-8}$$

Symbols in (S-8).

- Σ^{ab}, $\Sigma^{\mu\nu}$: The generators of Spin(4,4) on dirac16complex, $\Sigma^{ab} = \frac{1}{4}[T^a, T^b]$ (the scaled commutators of the

author's matrices); $\Sigma^{\mu\nu} = \mathbf{e}_a{}^\mu \mathbf{e}_b{}^\nu \Sigma^{ab}$ along the fold. Unit: dimensionless. Status: AUTHOR.

- $\chi_o(\sigma)$, $\bar\chi_o$: The profile of the occupied Kohn-Sham orbital o ($o = 1, \dots, 4$, the four positive-energy states of the level at the fold) of a homogeneous gas of the particles χ at rest on the static block: a 16-component spinor function of σ; the orbital of each particle is $\mathrm{e}^{\frac{7}{2}H|\sigma|}\chi_o(\sigma)\mathrm{e}^{-\mathrm{i}\omega x_4}$ divided by the square root of the volume of ordinary space and of V_*; normalised to the Krein charge -1 with the measure of UM-4, $\int \mathrm{d}\sigma\, \mathrm{e}^{H|\sigma|}\chi_o^\dagger \mathfrak{B}\chi_o = -1$; $\bar\chi_o = \chi_o^\dagger\sigma_{16}$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{1/2}$. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- T_a, T^a: The author's eight real 16×16 Dirac matrices T16ˆA[a] of 4+4.nb ($a = 0, \dots, 7$; $T_0, \dots, T_7$), written out in FD-13: T_a has the block $\bar\Xi_a$ in its upper right and Ξ_a in its lower left corner; $\{T_a, T_b\} = 2\eta_{ab}\mathbf{1}_{16}$; $T^a = \eta^{ab}T_b$. Unit: dimensionless. Status: AUTHOR.

The symmetry that makes the problem small. The author's $T_0, \dots, T_3$ are symmetric and $T_4, \dots, T_7$ antisymmetric, so the rotations of ordinary space (Σ^{ij}, $i, j = 1, 2, 3$) and of the extra times (Σ^{rs}, $r, s = 5, 6, 7$) form a compact group $SU(2) \times SU(2)$ of real orthogonal matrices, and $\gamma^{(0)}$, $\gamma^{(4)}$, P_L, P_R, σ_{16} and the Krein operator $\mathfrak{B}$ commute with it: the 16 components split into four copies of the representation $(2, 2)$ of this group. The four occupied orbitals are taken to form one $(2, 2)$ multiplet (the symmetric occupation of the level, a stationary point of the first-order energy but not a minimum: the Krein section below shows it to be a saddle); then the mean field is invariant, and the equations reduce exactly to the four-dimensional weight space S-8 (the states with the eigenvalue i/2 of both Σ^{12} and Σ^{56}): one orbital of four components, from which the multiplet is recovered by rotations by π and checked in sixteen components. Status: DERIVED (the reduction, `um4_kohnsham.py`, K1), ASSUMED (the symmetric occupation).

$$
\begin{aligned}
&\mathsf{E}_\mathrm{int} = \frac{G_\mathrm{KS}}{2}\int \mathrm{d}\sigma\, \mathrm{e}^{7H|\sigma|}\left(\hat\rho^2 - \mathrm{tr}\,\hat{\mathcal{N}}^2\right), \qquad \epsilon_\mathrm{p} = \omega - \mathsf{E}_\mathrm{int}, \qquad P_\mathrm{KS} = q_\chi = n_\chi\, \mathsf{E}_\mathrm{int}, \\
&\Delta\omega = G_\mathrm{KS}\int \mathrm{d}\sigma\, \mathrm{e}^{7H|\sigma|}\left(\hat\rho^2 - \mathrm{tr}\,\hat{\mathcal{N}}^2\right)\Big|_{G_\mathrm{KS}=0}
\end{aligned}
\tag{S-9}
$$

Symbols in (S-9).

- E_int: The interaction energy per particle of the gas of χ: the mean self-interaction energy $\frac{\lambda_\Psi}{2}(n_\mathrm{s}^2 - \mathrm{tr}\,\mathcal{N}^2)$ integrated across the hidden direction, per particle. Unit: GeV. Status: DERIVED.
- G_KS: The dimensionless strength of the self-interaction of the gas of particles χ in the Kohn-Sham states: the self-coupling λ_Ψ times the number density n_χ of the particles on the fold divided by the volume V_* of the extra times, $G_\mathrm{KS} = \lambda_\Psi n_\chi / V_*$ (in units of m_Ψ when numbers are given). Unit: dimensionless. Value: 10^{-6} to 0.5 in the solutions. Status: DERIVED.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- $\hat\rho(\sigma)$: The scalar density of the occupied orbitals, $\hat\rho = \frac{1}{4}\sum_o \bar\chi_o\chi_o$ (per particle); the scalar density of the gas is $n_\mathrm{s} = (n_\chi/V_*)\,\mathrm{e}^{7H|\sigma|}\hat\rho$. Unit: GeV. Status: DERIVED.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- $\hat{\mathcal{N}}(\sigma)$: The one-body density matrix of the occupied orbitals, $\hat{\mathcal{N}} = \frac{1}{4}\sum_o \chi_o\bar\chi_o$ (a 16×16 matrix at each σ, per particle); the density matrix of the gas is $\mathcal{N} = (n_\chi/V_*)\,\mathrm{e}^{7H|\sigma|}\hat{\mathcal{N}}$. Unit: GeV. Status: DERIVED.
- ϵ_p: The energy per particle of the gas of χ: the orbital energy minus the interaction energy per particle, which the orbital energy counts twice. Unit: GeV. Status: DERIVED.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- P_KS: The pressure of the gas of χ along ordinary space in the Kohn-Sham state: the particles are at rest, so it is the mean self-interaction, $P_\mathrm{KS} = n_\chi \mathsf{E}_\mathrm{int}$. Unit: GeV^4. Status: DERIVED.
- q_χ: The pressure of the particles χ of dirac16complex along the extra times, integrated across the fold (their on-shell Lagrangian, the self-interaction energy per volume; zero for a dilute gas). Unit: GeV^4. Status:

DERIVED.

- n_χ: The number density of the dark-matter particles χ (particles plus antiparticles). Unit: GeV^3. Status: DERIVED.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.

The energy of the state (FE-42 for the gas). $\mathsf{E}_{\mathrm{int}}$ is the interaction energy per particle (the trace runs over the sixteen components); the energy per particle is $\epsilon_{\mathrm{p}} = \omega - \mathsf{E}_{\mathrm{int}}$, because the orbital energy counts the interaction twice. The particles are at rest and do not depend on the extra times, so both pressures are the mean self-interaction, $P_{\mathrm{KS}} = q_\chi = n_\chi \mathsf{E}_{\mathrm{int}}$. $\Delta\omega$ is the first-order shift of the level, the mean of the mean field over the orbitals without the self-interaction: it equals $2\mathsf{E}_{\mathrm{int}}$ to the first order. The integrals run over the region of the fold inside the turning point, where the profiles decay at the local rate of FE-37 (in the computation, the box); the outgoing tail beyond the turning point, where $\mathrm{e}^{7H|\sigma|}|\chi|^2$ grows, is the width of the state, not part of it, so UM-3's condition for the convergence of the integral ($q_{\mathrm{b}} > 3H/2$) has no counterpart in UM-4. Status: DERIVED.

$$\omega - \omega_{\mathrm{free}} \propto G_{\mathrm{KS}} \quad (G_{\mathrm{KS}} \to 0), \qquad \mathrm{e}^{7H|\sigma|}\,\hat{\rho}(\sigma) \propto \mathrm{e}^{-(2q_{\mathrm{loc}}-7H)|\sigma|} \quad \big(|\sigma| \text{ large, inside the turning point}\big) \tag{S-10}$$

Symbols in (S-10).

- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- ω_{free}: The level of the occupied orbitals of the gas of particles χ without the self-interaction ($G_{\mathrm{KS}} = 0$), on the static block. Unit: GeV. Value: $0.9537799412\, m_\Psi$ for $H = 10^{-4} m_\Psi$, $\mu_{\mathrm{L}} = -0.6$, $\mu_{\mathrm{R}} = 0$. Status: DERIVED.
- G_{KS}: The dimensionless strength of the self-interaction of the gas of particles χ in the Kohn-Sham states: the self-coupling λ_Ψ times the number density n_χ of the particles on the fold divided by the volume V_* of the extra times, $G_{\mathrm{KS}} = \lambda_\Psi n_\chi / V_*$ (in units of m_Ψ when numbers are given). Unit: dimensionless. Value: 10^{-6} to 0.5 in the solutions. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\hat{\rho}(\sigma)$: The scalar density of the occupied orbitals, $\hat{\rho} = \frac{1}{4}\sum_o \bar{\chi}_o \chi_o$ (per particle); the scalar density of the gas is $n_{\mathrm{s}} = (n_\chi/V_*)\,\mathrm{e}^{7H|\sigma|}\hat{\rho}$. Unit: GeV. Status: DERIVED.
- q_{loc}: The local decay rate of a Kohn-Sham orbital at the depth σ inside the turning point: the smallest decay rate of the local equation of the orbital with the self-consistent mean field (without the self-interaction the rate q of the free profile). Unit: GeV. Status: DERIVED.

The dilute limit and the confinement of the gas. The shift of the level is linear in G_{KS} in the dilute limit (ω_{free} the level without the self-interaction). Away from the fold, inside the turning point, the proper scalar density decays at the rate $2q_{\mathrm{loc}} - 7H$, with q_{loc} the local decay rate of the orbital: it stays positive, so the gas stays at the fold. Both are checked in the box $|\sigma| \le 40$, where the weight $\mathrm{e}^{7H|\sigma|}$ differs from 1 by at most 0.028: they describe the region near the fold, not the approach to the turning point. In UM-3, whose factor was $\mathrm{e}^{6H|\sigma|}$ and whose regimes included $q_{\mathrm{b}} < 3H$, the proper density saturated far from the fold; the regimes of UM-3 with q_{b} of the order of H do not exist in UM-4, because such states are broad resonances, not particles. Status: DERIVED (K4, K5).

$$\mu_{\mathrm{L}} = \mu_{\mathrm{L}0}\Big(1 - \kappa_{\mathrm{H}}\,\frac{\hat{\rho}_{\mathrm{L}}(0)}{\hat{\rho}_{\mathrm{L}0}(0)}\Big), \qquad \hat{\rho}_{\mathrm{L}} = \frac{1}{4}\sum_{o=1}^{4} \bar{\chi}_o P_{\mathrm{L}} \chi_o, \qquad \kappa_{\mathrm{H}} = \frac{c_\Phi\, n_\chi\, \hat{\rho}_{\mathrm{L}0}(0)}{\lambda\, v^2\, V_*} \tag{S-11}$$

Symbols in (S-11).

- μ_X, μ_{L}, μ_{R}: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_{\mathrm{L}} P_{\mathrm{L}} + \mu_{\mathrm{R}} P_{\mathrm{R}})$; negative values attract. Unit: dimensionless. Status: DERIVED.

- $\mu_{\mathrm{L}0}$: The strength μ_{L} of the fold for the left-handed half without the feedback of the gas on the Higgs field. Unit: dimensionless. Value: -0.6 in the solutions. Status: DERIVED.
- κ_{H}: The strength of the Higgs-portal loop of the gas: the gas without the feedback changes the local v_{loc}^2 at the fold by the factor $1-\kappa_{\mathrm{H}}$, $\kappa_{\mathrm{H}} = c_\Phi n_\chi \hat{\rho}_{\mathrm{L}0}(0)/(\lambda v^2 V_*)$; negative where the Higgs portal binds the fold ($c_\Phi < 0$, $\hat{\rho}_{\mathrm{L}0}(0) > 0$): the gas raises v_{loc}^2. Unit: dimensionless. Value: -0.3 in the test of the loop. Status: ASSUMED (the value), DERIVED (the sign).
- $\hat{\rho}_{\mathrm{L}}$: The scalar density of the left-handed half of the occupied orbitals, $\hat{\rho}_{\mathrm{L}} = \frac{1}{4}\sum_o \bar{\chi}_o P_{\mathrm{L}} \chi_o$; at the fold the mean of its values on the two sides. Unit: GeV. Status: DERIVED.
- $\hat{\rho}_{\mathrm{L}0}$: The left-handed scalar density $\hat{\rho}_{\mathrm{L}}$ of the state at the fold without the self-interaction and without the Higgs feedback. Unit: GeV. Status: DERIVED.
- $\chi_o(\sigma)$, $\bar{\chi}_o$: The profile of the occupied Kohn-Sham orbital o ($o = 1, \ldots, 4$, the four positive-energy states of the level at the fold) of a homogeneous gas of the particles χ at rest on the static block: a 16-component spinor function of σ; the orbital of each particle is $\mathrm{e}^{\frac{7}{2}H|\sigma|}\chi_o(\sigma)\mathrm{e}^{-\mathrm{i}\omega x_4}$ divided by the square root of the volume of ordinary space and of V_*; normalised to the Krein charge -1 with the measure of UM-4, $\int \mathrm{d}\sigma\, \mathrm{e}^{H|\sigma|}\chi_o^\dagger \mathfrak{B} \chi_o = -1$; $\bar{\chi}_o = \chi_o^\dagger \sigma_{16}$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{1/2}$. Status: DERIVED.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- P_{L}: The author's left-handed chiral projector, $P_{\mathrm{L}} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- n_χ: The number density of the dark-matter particles χ (particles plus antiparticles). Unit: GeV^3. Status: DERIVED.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_{\mathrm{F}})^{-1/2}$ with the Fermi constant $G_{\mathrm{F}} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\, \mathrm{d}x_6\, \mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.

The Higgs-portal loop. Where the Higgs portal alone sets the strength of the fold for the left-handed half, μ_{L} is proportional to $\langle \Phi^\dagger \Phi \rangle = v_{\mathrm{loc}}^2/2$, and the left-handed scalar density of the gas at the fold shifts v_{loc}^2 (FE-49). $\mu_{\mathrm{L}0}$ and $\hat{\rho}_{\mathrm{L}0}(0)$ are the values without the feedback, $\hat{\rho}_{\mathrm{L}}(0)$ is the mean of the values on the two sides of the fold, and κ_{H} measures the strength of the loop: the gas without feedback changes v_{loc}^2 by the factor $1-\kappa_{\mathrm{H}}$. Where the Higgs portal alone binds the fold ($\mu_{\mathrm{R}} = 0$), $\mu_{\mathrm{L}0} = c_\Phi v^2/(2V_*) < 0$, so $c_\Phi < 0$; and the occupied orbitals have $\hat{\rho}_{\mathrm{L}0}(0) = \partial\omega_0/\partial\mu_{\mathrm{L}} > 0$ (the Hellmann-Feynman theorem with the Krein charge -1). Hence $\kappa_{\mathrm{H}} < 0$: the gas raises v_{loc}^2, the fold binds more strongly, and the level falls. The orbitals and μ_{L} are iterated together. Status: DERIVED (from FE-39, FE-40 and FE-49), ASSUMED (the value of κ_{H} used below).

$$\Gamma_\chi \simeq 2\,\frac{q_{\mathrm{b}}^2}{m_\Psi}\,\mathrm{e}^{-S_{\mathrm{t}}}, \qquad S_{\mathrm{t}} = 2\Big(\int_0^{L_{\mathrm{b}}} q_{\mathrm{loc}}\,\mathrm{d}\sigma + \frac{F\big(\omega\,\mathrm{e}^{HL_{\mathrm{b}}}\big)}{H}\Big) \tag{S-12}$$

Symbols in (S-12).

- Γ_χ: The width of the particle χ at rest: the rate of its decay into the waves of dirac16complex in the bulk. Unit: GeV. Value: below $1/(11.6\,t_0) = 1.3 \times 10^{-43}$ GeV for dark matter (a lifetime above 160 Gyr). Status: DERIVED.
- q_{b}: The inverse width of a state of dirac16complex at the fold: inside the turning point its profile falls as $\mathrm{e}^{-q_{\mathrm{b}}|\sigma|}$. Unit: GeV. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- S_{t}: The tunnelling exponent of the width of a state of the gas of particles χ: twice the integral of the local decay rate of its orbital over the region inside the turning point (the self-consistent part in the computational box and the free barrier beyond it). Unit: dimensionless. Value: 114.1 for the lifetime of 160 Gyr at $m_\Psi/H = 10^4$ and $H = 2534$ GeV. Status: DERIVED.

- L_b: The half-width of the computational box $|\sigma| \le L_\mathrm{b}$ of the Kohn-Sham states, far inside the turning point. Unit: length. Value: $40/m_\Psi$. Status: ASSUMED.
- q_loc: The local decay rate of a Kohn-Sham orbital at the depth σ inside the turning point: the smallest decay rate of the local equation of the orbital with the self-consistent mean field (without the self-interaction the rate q of the free profile). Unit: GeV. Status: DERIVED.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $F(\upsilon)$: The tunnelling exponent of the width of the particles χ per unit of m_Ψ/H: the integral of the local decay rate q over the region inside the turning point, divided by m_Ψ/H, as a function of the ratio υ of the energy to the bulk mass. Unit: dimensionless. Value: 0.049 for $\upsilon = 0.866$; $(q_\mathrm{b}/\omega_\mathrm{b})^3/3$ for weak binding. Status: DERIVED.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.

The width of the gas. The tunnelling exponent S_t of a self-consistent orbital is the integral of its smallest local decay rate q_loc (with the self-consistent mean field) over the box of the half-width L_b, plus the free barrier beyond the box with the tunnelling exponent F of FE-86. Without the self-interaction S_t is the free barrier $2F(\upsilon_\mathrm{r})m_\Psi/H$ at the resonance, which exceeds $2F(\upsilon_\mathrm{b})m_\Psi/H$ by 1 (FE-86), so that the prefactor $2(q_\mathrm{b}/m_\Psi)^2$ of S-12 is e times p_w. The repulsive mean field raises the level towards m_Ψ, the barrier shrinks and the gas leaks into the bulk. Status: DERIVED (K8: the excess 1.002 of the exponent at the smallest coupling), ESTIMATE (the prefactor with the self-interaction, taken without it).

9.2 The method

The level is found by shooting. On each side of the fold the equation S-7 is integrated inward from $|\sigma| = 40$ with the classical fourth-order Runge-Kutta method of step 0.03 (the step written as a matrix and the solutions re-orthonormalised every eight steps), starting with the solutions that decay at the end of the box; the frequency is the zero of the residual of the matching at the fold for a matching vector of negative Krein norm. The orbital gives the densities, the densities the mean field, and the mean field (with μ_L for the Higgs loop) is iterated with Anderson mixing. An iteration has converged when the largest relative change of the mean field is below 10^{-8} and the level changes by less than 10^{-10}. The coupling is raised by continuation. Every converged solution is lifted to sixteen components and checked there: the Kohn-Sham equation by central differences on the grid, with the mean field rebuilt in sixteen components from the lifted orbitals, the matching at the fold, the Krein charges and the Krein orthogonality; the residual of self-consistency is that of the reduced problem, evaluated once more with the accepted field. The computation runs in four parts (the free level, the gas, the Higgs loop and the weakly bound point), each invocation under 400 seconds. Status: DERIVED (`um4_kohnsham.py`).

9.3 The solution

G_KS	ω	$\omega - \omega_\mathrm{free}$	E_int	$P_\mathrm{KS}/(n_\chi\epsilon_\mathrm{p})$	iterations	decay rate of the proper density	$S_\mathrm{t}H$
10^{-6}	0.95378004	1.02×10^{-7}	5.11×10^{-8}	5.4×10^{-8}	5	0.5781	0.01914
10^{-5}	0.95378096	1.02×10^{-6}	5.11×10^{-7}	5.4×10^{-7}	4	0.5781	0.01914
10^{-4}	0.95379016	1.02×10^{-5}	5.11×10^{-6}	5.4×10^{-6}	4	0.5781	0.01913
10^{-3}	0.95388205	1.02×10^{-4}	5.11×10^{-5}	5.4×10^{-5}	7	0.5775	0.01908
0.01	0.95479700	1.02×10^{-3}	5.08×10^{-4}	5.3×10^{-4}	7	0.5713	0.01850
0.1	0.96352883	0.00975	0.00480	0.0050	9	0.5088	0.01338

G_{KS}	ω	$\omega-\omega_{\mathrm{free}}$	$\mathsf{E}_{\mathrm{int}}$	$P_{\mathrm{KS}}/(n_\chi\epsilon_{\mathrm{p}})$	iterations	decay rate of the proper density	$S_{\mathrm{t}}H$
0.2	0.97227589	0.0185	0.00891	0.0093	10	0.4366	0.00887
0.3	0.97991992	0.0261	0.0123	0.0127	10	0.3611	0.00551
0.4	0.98635877	0.0326	0.0148	0.0153	10	0.2813	0.00316
0.5	0.99147560	0.0377	0.0164	0.0169	12	0.1950	0.00168

The self-consistent states ($m_\Psi=1$, $H=10^{-4}$, $\mu_{\mathrm{L}}=-0.6$, $\mu_{\mathrm{R}}=0$; the continuation also passed $G_{\mathrm{KS}}=0.15$, 0.25, 0.35 and 0.45 and stopped at 0.5, where $\omega=0.991476$ exceeds 0.99). ω is the orbital energy of the occupied level, $\mathsf{E}_{\mathrm{int}}$ the interaction energy per particle, $P_{\mathrm{KS}}/(n_\chi\epsilon_{\mathrm{p}})$ the pressure divided by the energy density, the seventh column the decay rate of the proper scalar density at 0.85 of the half-width of the box, and the last the tunnelling exponent of S-12 in units of m_Ψ/H. Status: DERIVED (`um4_kohnsham.py`, K3).

The free level and the symmetry. Without the self-interaction the shooting with the author's 16×16 matrices gives $\omega=0.9537799412$, and the reduced four-dimensional problem of S-8 the same level; it consists of eight degenerate states, four of each Krein sign (two, one of each sign, in the reduced problem). The flat value is $\omega_{\mathrm{b}}=0.9539392014$, and $\omega_{\mathrm{b}}(1-H/(2q_{\mathrm{b}}))=0.9537802116$: the computed level differs from it by $-2.70\times10^{-7}=-27.0H^2$, a term of the second order in H (4.04 times as large at $2H$). The box method agrees with the exact resonance of FE-86 (outgoing Hankel waves on both sides) at $m_\Psi/H=200$, $\mu_{\mathrm{L}}=-1$: 0.8616233418 against 0.8616233403 (the width there is $\Gamma_\chi=4.8\times10^{-10}m_\Psi$). Status: DERIVED (K1).

The dilute limit. The shift of the level divided by the first-order shift of S-9 (the coefficient $\Delta\omega/G_{\mathrm{KS}}=0.102162$, computed in sixteen components) is 0.99995, 0.99994, 0.99990 and 0.99951 for $G_{\mathrm{KS}}=10^{-6}$, 10^{-5}, 10^{-4} and 10^{-3}; fitted for $G_{\mathrm{KS}}\le10^{-4}$ the shift grows as G_{KS}^{p} with $p=1.0000$ (S-10). The occupied and the empty multiplets split at every G_{KS}, in proportion to it: the second singular value of the matching at the occupied level divided by G_{KS} is 0.0843 from $G_{\mathrm{KS}}=10^{-6}$ to 10^{-3}, 0.0841 at 0.01 and 0.0822 at 0.1. Status: DERIVED (K2, K4).

Confinement. Near the end of the box the proper scalar density decays at the rates 0.578 (dilute) to 0.195 ($G_{\mathrm{KS}}=0.5$), always $2q_{\mathrm{loc}}-7H$ with $7H=7\times10^{-4}$: within the box the gas stays at the fold and does not saturate the bulk; beyond the box the orbital decays up to the turning point and leaves the fold there as the outgoing wave of the width. An attractive self-interaction, $G_{\mathrm{KS}}=-10^{-4}$ and -10^{-2} from the free state, converges too and lowers the level to $\omega=0.95376973$ and 0.95275389. Status: DERIVED (K5).

The Higgs-portal loop. The occupied orbitals without the self-interaction have $\hat\rho_{\mathrm{L}0}(0)=0.15694536$, equal to $\partial\omega_0/\partial\mu_{\mathrm{L}}=0.15694959$ (the level solved at $\mu_{\mathrm{L}}\pm10^{-4}$) up to 2.70×10^{-5}, the error of the trapezoidal normalisation on the grid (the flat formula $-\mu_{\mathrm{L}}/(4\omega_{\mathrm{b}})$ gives 0.15724273); so $\kappa_{\mathrm{H}}<0$. With $\kappa_{\mathrm{H}}=-0.3$ and $G_{\mathrm{KS}}=0.1$ the orbitals and μ_{L} of S-11 converge together in 13 iterations: $\mu_{\mathrm{L}}=-0.873128$ (from -0.6): the gas raises the local electroweak vacuum, the fold binds more strongly, and the level falls to $\omega=0.91257910$ (0.96352883 without the feedback). Status: DERIVED (K6), ASSUMED (the value of κ_{H}).

The equation of state. The pressure is positive for the repulsive self-interaction and vanishes in the dilute limit: $P_{\mathrm{KS}}/(n_\chi\epsilon_{\mathrm{p}})=5.36\times10^{-8}$, 5.36×10^{-7}, 5.36×10^{-6}, 5.35×10^{-5}, 5.32×10^{-4} and 5.00×10^{-3} for $G_{\mathrm{KS}}=10^{-6}$ to 0.1. A dilute gas of the particles χ is pressureless: cold dark matter. Status: DERIVED (K7).

The width of the gas. The tunnelling exponent falls from $S_{\mathrm{t}}H=0.01914$ at $G_{\mathrm{KS}}=10^{-6}$ (the free barrier $2F(\upsilon_{\mathrm{b}})=0.01914$) to 0.00168 at $G_{\mathrm{KS}}=0.5$: the width grows with the self-interaction. At the smallest coupling S_{t} exceeds $2F(\upsilon_{\mathrm{b}})m_\Psi/H=190.392$ by 1.002, the factor 1/e of the prefactor of FE-86. At $m_\Psi/H=10^4$ with H at its bound 2534 GeV ($m_\Psi=2.53\times10^7$ GeV) the gas lives longer than 160 Gyr ($S_{\mathrm{t}}>114.08$ with the prefactor of S-12) only for $G_{\mathrm{KS}}<0.139$. (In UM-3 the warped proper density saturated far from the fold and kept the state bound beyond $\omega=m_\Psi$; in UM-4 the energy is warped as well, and the level approaches m_Ψ from below while the orbital spreads into the barrier.) Status: DERIVED (K8).

The residuals. Each of the 14 converged solutions was lifted to sixteen components and checked there: the largest relative residual of the Kohn-Sham equation by central differences on the grid is 1.0×10^{-4} (the error of the difference formula, of the order of the square of the step), of the matching at the fold 2×10^{-15}, of self-consistency (in the

reduced problem) 7.5×10^{-9}; the four orbitals have the Krein charge -1 and are Krein-orthogonal to 10^{-16}, and the bound on the mean field is never active. Status: DERIVED (K3).

9.4 A weakly bound gas

Direct detection requires a weak binding of χ by the Higgs portal (the thermal history, below; at the masses that the lifetime allows even weaker than this point). A weakly bound point is solved as well: $\mu_{\mathrm{L}} = -0.06$ ($q_{\mathrm{b}} = 0.03$, $q_{\mathrm{b}}/\omega_{\mathrm{b}} = 0.0300$) and $H = 10^{-7}$ (the free barrier $2F = 1.81 \times 10^{-5}$ per unit of m_Ψ/H, so that $2Fm_\Psi/H = 181$; with H at its bound 2534 GeV this is $m_\Psi = 2.5 \times 10^{10}$ GeV, above the smallest mass, 1.6×10^{10} GeV, at which this binding lives longer than 160 Gyr). The shooting in sixteen components gives $\omega = 0.9995482297$ against $\omega_{\mathrm{b}}(1 - H/(2q_{\mathrm{b}})) = 0.9995482328$ (the difference -3.1×10^{-9} is of the second order in H, 4.04 times as large at $2H$), with four Krein-negative orbitals, and the first-order coefficient of S-9 is $\Delta\omega/G_{\mathrm{KS}} = 0.011209$. The self-consistent shift divided by the first-order shift is 0.99933, 0.99367 and 0.93699 for $G_{\mathrm{KS}} = 10^{-4}$, 10^{-3} and 10^{-2} (approaching 1 as $G_{\mathrm{KS}} \to 0$), with $P_{\mathrm{KS}}/(n_\chi \epsilon_{\mathrm{p}}) = 5.6 \times 10^{-7}$, 5.6×10^{-6} and 5.1×10^{-5}: a dilute gas of weakly bound particles χ is cold. Status: DERIVED (K9).

9.5 The Krein norm and the stability of the gas

$$\{\mathsf{a}_o, \mathsf{a}_o^\dagger\} = -\mathcal{Q}_o, \qquad \mathsf{H}_\Psi = \sum_o \big(-\mathcal{Q}_o\, \omega_o \big)\, \mathsf{a}_o^\dagger \mathsf{a}_o, \qquad \big[\mathsf{H}_\Psi, \mathsf{a}_o^\dagger\big] = \omega_o\, \mathsf{a}_o^\dagger \tag{FE-67}$$

Symbols in (FE-67).

- a_o, $\mathsf{a}_o^\dagger$: The annihilation and the creation operator of the mode o of the quantised dirac16complex (frequency ω_o, Krein charge $\mathcal{Q}_o$); the dagger is the adjoint with respect to the indefinite (Krein) scalar product. Unit: dimensionless. Status: STANDARD.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\mathcal{Q}_o$: The charge Q_Ψ (the Krein charge) of the single-particle state o of dirac16complex, a Kohn-Sham orbital or a mode: $+1$ or -1 for a normalised state. Unit: dimensionless. Status: DERIVED.
- H_Ψ: The Hamiltonian (the energy operator) of the quantised dirac16complex. Unit: GeV. Status: DERIVED.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.

The quantised dirac16complex: a mode o of the frequency ω_o and the Krein charge $\mathcal{Q}_o = \pm 1$ (FE-43) has the anticommutator $-\mathcal{Q}_o$, and every quantum raises the energy by its frequency. In the positive-energy quantisation every particle and every hole carries a positive energy and the norm $-\mathcal{Q}_o$ of its mode: the particles and the holes of the modes of charge -1 are physical, those of the modes of charge $+1$ are ghosts (negative norm), and no choice of the vacuum removes them. The Kohn-Sham states above occupy four orbitals of charge -1: they are physical states, and the questions are whether they are stable and whether the couplings of the fold produce ghosts. Status: DERIVED (`um4_krein.py`, KR1: the Krein operator has eight eigenvalues $+1$ and eight -1; the conjugates of the four Krein-negative states at $+\omega_0$ are Krein-positive states at $-\omega_0$).

The symmetry and its mirrors. At rest, the transformations of the sixteen components that keep the states at the fold (they commute with $\gamma^{(0)}$, $\gamma^{(4)}$ and Γ_8 and preserve the Dirac adjoint) form a Lie algebra of dimension 32: 16 compact generators keep the two Krein charges apart, and 16 boosts mix them, among them the nine boosts Σ^{ir} between ordinary space and the extra times; for a moving particle only four compact generators remain. The mirror R_1 of the fold (FD-10) keeps the charges: the four states of charge -1 are even, the four of charge $+1$ odd, and the self-consistent orbital at $G_{\mathrm{KS}} = 0.1$ and its mean field are even (to 3×10^{-10}); R_2 exchanges the two charges. Status: DERIVED (KR2, KR3).

The stability of the gas is not decided. The boosts that mix the charges map the self-consistent gas onto self-consistent gases (at $G_{\mathrm{KS}} = 0.1$ the residual of the Kohn-Sham equation stays 3.5×10^{-5}, as without the boost, and the charges stay -1, while the orbitals acquire admixtures of charge $+1$ whose positive-definite norm grows up to 2.43 times): these directions are exactly flat, and exact zero modes would give the frequency 0 in the linear response. The occupation of the level is a saddle of the first-order energy. Every four-dimensional subspace of negative Krein norm of the eight states of the free level is an occupation of four particles χ (32 real parameters); the integral of

S-9 is stationary at the symmetric multiplet (0.10216, gradient 3×10^{-12}), and its Hessian there has 15 negative eigenvalues (-0.0067), 16 zero ones (exactly the 16 generators of the symmetry that mix the charges) and one positive one (0.020); towards the null cone the integral falls to -1514 along the most negative direction and rises to 1.0×10^4 along the most positive one: the first-order energy of the repulsive gas has no lower bound on the occupations of negative norm, and the Kohn-Sham states above are the symmetric stationary occupation, not a ground state. The linear response truncated to the level at the fold (the 16 admixtures of the four empty orbitals of charge +1, split from the occupied level by 3.491×10^{-3}) keeps exactly the tangent space of these occupations; it has frequencies of the order 10^{-4}, two of them imaginary (up to 2.05×10^{-4}), and whether this imaginary pair is an instability of the full response, with the continuum of the bulk, is OPEN. The partner level (the level of charge +1 next to the occupied one in the self-consistent mean field) is real and bound up to $G_{\mathrm{KS}} = 0.3$, the largest computed, and it moves away from the occupied level as the coupling grows (the gap is 0.0004, 0.0035, 0.0064 and 0.0087 at $G_{\mathrm{KS}} = 0.01$, 0.1, 0.2 and 0.3): the two cannot mix, because they have opposite parity under the mirror R_1 in a mirror-even mean field. Status: DERIVED (KR4, KR5, KR8), OPEN (the stability of the gas).

The couplings of the fold decide unitarity. At rest none of the 16 chirality-blind operators of the fold (the products of $\gamma^{(1)}, \ldots, \gamma^{(4)}$) connects a physical state with a ghost, and each of the 16 chiral ones (the same times Γ_8) does. The left-handed Higgs portal P_{L} of the author has the elastic amplitude 0.157, no conversion of a particle into a ghost, no physical pair at rest (a p-wave), and the amplitude 0.157 (an s-wave) for a physical particle χ with a ghost antiparticle, equal to the elastic one: all four singular values of the 4×4 amplitude matrix, which do not depend on the basis of the degenerate multiplets, are 0.15695. For moving particles (the eight states at $|\vec{k}| = 0.05$ and 0.15 have exactly $\sqrt{\omega_0^2 + \vec{k}^2}$, the boost invariance of UM-4) the scalar, the density and the vector currents still do not mix (below 10^{-9}), but the magnetic dipole Σ^{12} does, in proportion to the momentum (6.7×10^{-4} and 2.0×10^{-3}). The convective current of the current portals (IN-16), $-2\vec{k}$ times the pair amplitude of P_X, creates a physical pair $\chi(\vec{k})\bar{\chi}(-\vec{k})$ only with an amplitude of the second order in the momentum: the four singular values of the pair amplitude of P_X vanish at rest and grow linearly, with the slope 0.14974 for P_{L} and P_{R} and 0.29947 for $P_{\mathrm{L}} + P_{\mathrm{R}} = \mathbf{1}$ (computed at $|\vec{k}| = 0.01$, 0.02 and 0.05 in units of m_Ψ, linear to 1.3×10^{-3}); the annihilation through the chirality-blind combination is that of a Dirac fermion whose current is the convective current of a scalar density (S-21). That combination creates a physical particle with a ghost antiparticle only at the next order: the singular values of its amplitude grow like $0.028\,\vec{k}^2/m_\Psi^2$ (quadratic to 3×10^{-3}), so its ghost pairs have the relative amplitude $0.094|\vec{k}|/m_\Psi$ and the relative rate $0.0089\,\vec{k}^2/m_\Psi^2$, small but not zero. The vacuum cannot decay (every quantum has a positive energy and the energy is conserved; with the indefinite metric it is not a state of lowest energy among the states of positive norm), but with chiral couplings at the fold ghosts are produced wherever the portals act: UM-4 with the author's left-handed Higgs portal is not a unitary quantum theory. Chirality-blind couplings are necessary for a unitary UM-4, but they are not shown to be sufficient. Status: DERIVED (KR6, KR7, KR9), OPEN (a unitary set of couplings).

10 The residuals of every field equation

The approximate solution of UM-4 consists of the parts above and below. The field equations that the other sections do not write out are restated first, so that the table at the end of this section can refer to each of them: the field equations of the bulk, the junction condition and the thin source of the fold, the conservation laws, the field equations of the gluons and of the hypercharge field, and those of the quarks and leptons.

$$\mathcal{E}_{(1)}{}^{M}{}_{N} + \Lambda\,\delta^{M}_{N} = \kappa\left(\mathcal{T}_{\Psi}{}^{M}{}_{N} + \mathcal{T}_{\mathrm{gauge}}{}^{M}{}_{N}\right) \qquad (\sigma \neq 0) \tag{FE-1}$$

Symbols in (FE-1).

- $\mathcal{E}_{(p)}{}^{M}{}_{N}$: The Lovelock tensor of order p, $\mathcal{E}_{(p)}{}^{M}{}_{N} = -2^{-(p+1)}\,\delta^{M P_1 \cdots P_{2p}}_{N Q_1 \cdots Q_{2p}}\,R^{Q_1 Q_2}{}_{P_1 P_2} \cdots R^{Q_{2p-1} Q_{2p}}{}_{P_{2p-1} P_{2p}}$ (with both indices down it is $1/\sqrt{|g|}$ times the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to g^{MN}; $\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor, $\mathcal{E}_{(3)}$ the cubic Lovelock tensor); with frame indices its frame components. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number (UM-4 keeps only $p = 1$); in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.

- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant of UM-4, $\Lambda = -21H^2$: the value for which the static block is a vacuum of Einstein gravity (its Einstein tensor is $21H^2$ in every direction). The author writes the pure number Λ_{N} in $0 = -\Lambda_{\mathrm{N}} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$, so that $\Lambda = -\Lambda_{\mathrm{N}} H^2$. Unit: GeV^2. Value: $-21H^2$; the author's original $-10H^2$; UM-3: $-7H^2$. Status: AUTHOR (the term), DERIVED (the value).
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_\Psi$: The energy-momentum tensor of dirac16complex: $-2/\sqrt{|g|}$ times the variation of the action S_Ψ of dirac16complex with respect to g^{MN}. Unit: GeV^8. Status: STANDARD.
- $\mathcal{T}_{\text{gauge}}$: The energy-momentum tensor of the gauge fields of the bulk. Unit: GeV^8. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

The Einstein equations of the bulk, away from the fold: the Einstein tensor plus the cosmological constant equals κ times the energy-momentum tensor of the fields of the bulk, dirac16complex and the gauge fields. Status: AUTHOR (the theory with the weights that the order selects), STANDARD (the variation).

$$\kappa\,\mathcal{S}^a{}_b = \lim_{\varepsilon\to 0}\int_{-\varepsilon}^{\varepsilon} \mathrm{d}\sigma\,\mathcal{E}_{(1)}{}^a{}_b = -\Big(\big(K^a{}_b\big|_{0^+} - K^a{}_b\big|_{0^-}\big) - \delta^a{}_b\big(K\big|_{0^+} - K\big|_{0^-}\big)\Big), \qquad K = K^c{}_c \tag{FE-4}$$

Symbols in (FE-4).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma\,\mathcal{E}_{(1)}{}^a{}_b$ (the junction condition of Israel); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -12H\delta^a{}_b$ for every a_4, b_4 of the author's form. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- ε: The half-thickness of a layer around the fold, taken to zero. Unit: length. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\mathcal{E}_{(p)}{}^M{}_N$: The Lovelock tensor of order p, $\mathcal{E}_{(p)}{}^M{}_N = -2^{-(p+1)}\,\delta^{M P_1 \cdots P_{2p}}_{N Q_1 \cdots Q_{2p}}\,R^{Q_1 Q_2}{}_{P_1 P_2} \cdots R^{Q_{2p-1} Q_{2p}}{}_{P_{2p-1} P_{2p}}$ (with both indices down it is $1/\sqrt{|g|}$ times the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to g^{MN}; $\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor, $\mathcal{E}_{(3)}$ the cubic Lovelock tensor); with frame indices its frame components. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number (UM-4 keeps only $p = 1$); in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- $K^a{}_b$: The extrinsic curvature of a surface of constant σ of the hidden direction, $K^a{}_b = \frac{1}{2} g^{ac}\partial_\sigma g_{cb}$ for a, b along the surface (FD-5): $-H\,\mathrm{sgn}(\sigma)$ along all seven directions of the fold for the author's form of the metric of UM-4; $K = K^c{}_c$ its trace. Unit: GeV (1/length). Status: DERIVED.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label

X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- c: A frame index (as a). Status: STANDARD.

The field equations at the fold: the metric has a kink at $\sigma = 0$, so its second derivatives along σ contain a delta function, and the jump of the extrinsic curvature across the fold equals κ times the thin source $\mathcal{S}$ of the fold (Israel's condition). Status: DERIVED.

$$\mathcal{S}^a{}_b = \frac{1}{V_*}\Big(\mathcal{T}_{\mathrm{H}}{}^a{}_b + \mathcal{T}_{\mathrm{f}}{}^a{}_b + \mathcal{T}_{\mathrm{P}}{}^a{}_b + \mathcal{T}_{\mathrm{ff}}{}^a{}_b + \mathcal{T}_{\mathrm{IG}}{}^a{}_b\Big), \qquad \mathcal{T}_{\mathrm{IG}}{}^\mu{}_\nu = -\frac{1}{\kappa_{\mathrm{ind}}}\,\hat{G}^\mu{}_\nu, \qquad \mathcal{T}^r{}_r = \mathcal{L}_{\mathrm{fold}} \tag{FE-5}$$

Symbols in (FE-5).

- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma\, \mathcal{E}_{(1)}{}^a{}_b$ (the junction condition of Israel); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -12H\delta^a{}_b$ for every a_4, b_4 of the author's form. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\, \mathrm{d}x_6\, \mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- $\mathcal{T}_{\mathrm{H}}$: The energy-momentum tensor of the Higgs doublet (four-dimensional normalisation). Unit: GeV^4. Status: STANDARD.
- $\mathcal{T}_{\mathrm{f}}$: The energy-momentum tensor of the quarks and leptons. Unit: GeV^4. Status: STANDARD.
- $\mathcal{T}_{\mathrm{P}}$: The energy-momentum tensor of the portals of dirac16complex. Unit: GeV^4. Status: DERIVED.
- $\mathcal{T}_{\mathrm{ff}}{}^a{}_b$: The energy-momentum tensor of the fold fluid (frame components of the fold, four-dimensional normalisation). Unit: GeV^4. Status: ASSUMED.
- $\mathcal{T}_{\mathrm{IG}}$: The energy-momentum tensor of the induced Einstein-Hilbert term of the fold, $\mathcal{T}_{\mathrm{IG}}{}^\mu{}_\nu = -\hat{G}^\mu{}_\nu/\kappa_{\mathrm{ind}}$. Unit: GeV^4. Status: ESTIMATE.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+, +, +, -$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- κ_{ind}: The coupling of the Einstein-Hilbert term $\hat{R}/(2\kappa_{\mathrm{ind}})$ that the quantum fluctuations of the fields of the fold may induce, $\kappa_{\mathrm{ind}} = 8\pi G_{\mathrm{ind}}$; it adds to the gravity of the fold, $1/(8\pi G_{\mathrm{N}}) = V_*/\kappa_7 + 1/\kappa_{\mathrm{ind}}$. Unit: GeV^{-2}. Value: not fixed by UM-4; negligible for a cutoff of the order of H. Status: ESTIMATE.
- $\hat{G}^\mu{}_\nu$: The Einstein tensor of the metric $\hat{g}_{ab}$ of the fold (FD-2), with frame indices along its seven directions; with four-dimensional indices, $\hat{G}^\mu{}_\nu$, its components along the four observable directions, which for frozen extra times are those of the Einstein tensor of the four-dimensional metric $\hat{h}_{\mu\nu}$. Unit: 1/length^2. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- $\mathcal{T}$: The energy-momentum tensor of any one term of the fold (in $\mathcal{T}^r{}_r = \mathcal{L}_{\mathrm{fold}}$). Unit: GeV^4. Status: STANDARD.
- $\mathcal{L}_{\mathrm{fold}}$: The Lagrangian of any one term of the fold (four-dimensional normalisation). Unit: GeV^4. Status: STANDARD.

The thin source of the fold: the energy-momentum tensor of everything that lives on the fold (the Higgs doublet, the quarks and leptons, the portals, the fold fluid and, if the fields of the fold induce one, the induced gravity) divided by the volume V_* of the extra times; along each extra time its component is the Lagrangian of the term, because the fields of the fold do not depend on the extra times. Status: DERIVED.

$$\begin{aligned}
&\nabla_M \left(\mathcal{T}_\Psi{}^M{}_N + \mathcal{T}_{\text{gauge}}{}^M{}_N\right) = 0 \qquad (\sigma \neq 0),\\
&\hat\nabla_a \,\mathcal{S}^a{}_b = -\Big(\mathcal{T}_{\text{bulk}}{}^{(0)}{}_b\big|_{\sigma=0^+} - \mathcal{T}_{\text{bulk}}{}^{(0)}{}_b\big|_{\sigma=0^-}\Big), \qquad \mathcal{T}_{\text{bulk}} = \mathcal{T}_\Psi + \mathcal{T}_{\text{gauge}}
\end{aligned} \tag{FE-11}$$

Symbols in (FE-11).

- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols $\Gamma^P{}_{MN}$). Status: STANDARD.
- $\mathcal{T}_\Psi$: The energy-momentum tensor of dirac16complex: $-2/\sqrt{|g|}$ times the variation of the action S_Ψ of dirac16complex with respect to g^{MN}. Unit: GeV^8. Status: STANDARD.
- $\mathcal{T}_{\text{gauge}}$: The energy-momentum tensor of the gauge fields of the bulk. Unit: GeV^8. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \operatorname{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\hat\nabla$: The covariant derivative of the fold: with the Levi-Civita connection of the metric of the fold (for four-dimensional tensors of its metric $\hat h_{\mu\nu} = \operatorname{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the observable coordinates $x_1, \ldots, x_4$; for $\hat\nabla_a \mathcal{S}^a{}_b$ of the induced metric of the seven-dimensional fold). Status: STANDARD.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma\, \mathcal{E}_{(1)}{}^a{}_b$ (the junction condition of Israel); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -12H\delta^a{}_b$ for every a_4, b_4 of the author's form. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \operatorname{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat g_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- $\mathcal{T}_{\text{bulk}}$: The energy-momentum tensor of all sources of the bulk, dirac16complex and the gauge fields, $\mathcal{T}_{\text{bulk}} = \mathcal{T}_\Psi + \mathcal{T}_{\text{gauge}}$; $\mathcal{T}_{\text{bulk}}{}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.

The conservation laws: in the bulk the energy-momentum tensor of the fields of the bulk is conserved; at the fold the covariant divergence of the thin source equals minus the jump of the flux of energy and momentum of the bulk along σ, so the fold loses exactly what the bulk carries away (the waves of the gauge fields and of dirac16complex into which the W and Z bosons and the particles χ decay). Status: DERIVED.

$$\begin{aligned}
&\frac{1}{\sqrt{|g|}}\, D_M\big(\sqrt{|g|}\, G^{AMN}\big) = 0 \qquad (\sigma \neq 0),\\
&\frac{1}{\mathrm{g}_3^2}\Big(G^{A\sigma\nu}\big|_{0^+} - G^{A\sigma\nu}\big|_{0^-}\Big) = -\frac{1}{V_*}\Big(J_{\mathrm{c}}^{A\nu} + \frac{1}{g_3}\, D_\mu\big[(\mathcal{K}_G - 1)G^{A\mu\nu}\big]\Big), \qquad \mathcal{K}_G = 1 + \sum_X c_X^G\, \bar\Psi P_X \Psi,\\
&J_{\mathrm{c}}^{A\nu} = \bar Q\, \hat\gamma^\nu t^A Q + \bar u_{\mathrm{R}}\, \hat\gamma^\nu t^A u_{\mathrm{R}} + \bar d_{\mathrm{R}}\, \hat\gamma^\nu t^A d_{\mathrm{R}}
\end{aligned} \tag{FE-16}$$

Symbols in (FE-16).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+, +, +, +, -, -, -, -$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M \mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar\Psi = \partial_M\bar\Psi - \bar\Psi\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat h^{\mu\nu} D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G_\mu^B X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.

- G^A_M, G^A_μ, G^A_{MN}: The gluon field of $SU(3)_C$ ($A = 1, \ldots, 8$): in the bulk G^A_M, a field of eight dimensions in its geometric normalisation (the covariant derivative $\partial - \mathrm{i}G$), and on the fold the canonically normalised four-dimensional field $G^A_\mu(x)$ of its massless mode, $g_3 G^A_\mu(x) = G^A_\mu(\sigma = 0, x)$; G^A_{MN} and $G^A_{\mu\nu}$ their field strengths. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\mathtt{g}_1$, $\mathtt{g}_2$, $\mathtt{g}_3$: The gauge couplings of eight dimensions of $U(1)_Y$, $SU(2)_L$ and $SU(3)_C$: the coefficients of the kinetic terms of the gauge fields of the bulk in their geometric normalisation; the four-dimensional couplings are $g_k^2 = 3H\mathtt{g}_k^2/(2V_*)$. Unit: GeV^{-2} ($\mathtt{g}^2$: length^4). Status: ASSUMED.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- $J_\mathrm{c}^{A\nu}$: The colour current of the quarks. Unit: GeV^3. Status: STANDARD.
- g_3: The $SU(3)_C$ gauge coupling of the four-dimensional world (the strong coupling g_s): the coupling of the massless mode of the gluon field of the bulk, $g_3^2 = 3H\mathtt{g}_3^2/(2V_*)$. Unit: dimensionless. Value: $g_3(m_Z) = 1.2177$ ($g_3^2/(4\pi) = 0.1180$). Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- $\mathcal{K}_G$, $\mathcal{K}_W$, $\mathcal{K}_B$: The portal factors of the gauge kinetic terms, $\mathcal{K}_G = 1 + \sum_X c_X^G \bar\Psi P_X \Psi$ (likewise for W and B): the gauge couplings at the fold are divided by the square root of the portal factor. Unit: dimensionless. Value: 1 where dirac16complex vanishes. Status: DERIVED.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c_X^G: The gluon-kinetic portal coupling of the half X of dirac16complex. Unit: GeV^{-7}. Status: INVENTED.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- P_X: The chiral projector of the half X: $P_X = P_\mathrm{L}$ for $X = \mathrm{L}$, P_R for $X = \mathrm{R}$. Unit: dimensionless. Status: AUTHOR.
- Q, $\bar Q$: The left-handed quark doublets of the three generations, $Q = (u_\mathrm{L}, d_\mathrm{L})$ (colour triplets, hypercharge 1/6). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- $\hat\gamma^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat\gamma^{(a)}$ along the frame directions of the fold and $\hat\gamma^\mu$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat\gamma^\mu, \hat\gamma^\nu\} = -2\hat h^{\mu\nu}\mathbf{1}$; $\hat\gamma_\mu = \hat h_{\mu\nu}\hat\gamma^\nu$. Unit: dimensionless. Status: STANDARD.
- t^A: The generators of $SU(3)$ on colour triplets, t^A: half the Gell-Mann matrices, with $\mathrm{tr}(t^A t^B) = \delta^{AB}/2$. Unit: dimensionless. Status: STANDARD.
- u_R: The right-handed up-type quarks (up, charm and top; singlets of $SU(2)_L$, hypercharge 2/3). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- d_R: The right-handed down-type quarks (down, strange and bottom; hypercharge $-1/3$). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.

$$
\begin{aligned}
&\frac{1}{\sqrt{|g|}}\,\partial_M\big(\sqrt{|g|}\,B^{MN}\big) = 0 \qquad (\sigma \neq 0),\\
&\frac{1}{\mathtt{g}_1^2}\Big(B^{\sigma\nu}\big|_{0^+} - B^{\sigma\nu}\big|_{0^-}\Big) = -\frac{1}{V_*}\Big(J_Y^\nu + \frac{1}{g_1}\,\partial_\mu\Big[\big(\mathcal{K}_B - 1\big)B^{\mu\nu} + \sum_X d_X\,\mathcal{D}_X^{\mu\nu}\Big]\Big), \qquad \mathcal{K}_B = 1 + \sum_X c_X^B\,\bar\Psi P_X \Psi, \qquad \text{(FE-18)}\\
&J_Y^\nu = \sum_\psi Y_\psi\,\bar\psi\,\hat\gamma^\nu\psi + \frac{\mathrm{i}}{2}\big[(D^\nu\Phi)^\dagger\Phi - \Phi^\dagger D^\nu\Phi\big]
\end{aligned}
$$

Symbols in (FE-18).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order

$x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.

- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- B_M, B_μ, B_{MN}: The hypercharge gauge field of $U(1)_Y$: in the bulk B_M (geometric normalisation), on the fold the canonically normalised four-dimensional field $B_\mu(x)$ of its massless mode, $g_1 B_\mu(x) = B_\mu(\sigma = 0, x)$; B_{MN}, $B_{\mu\nu}$ their field strengths. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- g_1, g_2, g_3: The gauge couplings of eight dimensions of $U(1)_Y$, $SU(2)_L$ and $SU(3)_C$: the coefficients of the kinetic terms of the gauge fields of the bulk in their geometric normalisation; the four-dimensional couplings are $g_k^2 = 3H\mathsf{g}_k^2/(2V_*)$. Unit: GeV^{-2} (g^2: length^4). Status: ASSUMED.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- J_Y^ν: The hypercharge current of the fermions and of the Higgs doublet. Unit: GeV^3. Status: STANDARD.
- g_1: The $U(1)_Y$ gauge coupling of the four-dimensional world (usually g'; not normalised for grand unification): the coupling of the massless mode of the hypercharge field of the bulk, $g_1^2 = 3H\mathsf{g}_1^2/(2V_*)$. Unit: dimensionless. Value: $g_1(m_Z) = 0.3574$. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- $\mathcal{K}_G$, $\mathcal{K}_W$, $\mathcal{K}_B$: The portal factors of the gauge kinetic terms, $\mathcal{K}_G = 1 + \sum_X c_X^G \bar\Psi P_X \Psi$ (likewise for W and B): the gauge couplings at the fold are divided by the square root of the portal factor. Unit: dimensionless. Value: 1 where dirac16complex vanishes. Status: DERIVED.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- d_X: The dipole-portal coupling of the half X of dirac16complex with the hypercharge field strength. Unit: GeV^{-5}. Status: INVENTED.
- $\mathcal{D}_X^{\mu\nu}$: The dipole (antisymmetric tensor) density of the half X of dirac16complex at the fold, $\mathcal{D}_X^{\mu\nu} = \mathrm{i}\bar\Psi\Sigma^{\mu\nu}P_X\Psi$ (real, DERIVED). Unit: GeV^7. Status: DERIVED.
- c_X^B: The B-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- P_X: The chiral projector of the half X: $P_X = P_\mathrm{L}$ for $X = \mathrm{L}$, P_R for $X = \mathrm{R}$. Unit: dimensionless. Status: AUTHOR.
- ψ: The label of a Standard-Model fermion field in a sum: ψ runs over $Q, u_\mathrm{R}, d_\mathrm{R}, \ell, e_\mathrm{R}, \nu_\mathrm{R}$. Status: STANDARD.
- Y_f: The hypercharge of a field, the charge of its coupling $g_1 Y_f B_\mu$ to the hypercharge field in the covariant derivative (1/6, 2/3, −1/3, −1/2, −1, 0 for $Q, u_\mathrm{R}, d_\mathrm{R}, \ell, e_\mathrm{R}, \nu_\mathrm{R}$, 1/2 for Φ). Unit: dimensionless. Status: STANDARD.
- ψ, $\bar\psi$: A Standard-Model fermion field (a four-dimensional Weyl field of the fold with three generations, written as a four-component spinor of fixed chirality); $\bar\psi = \psi^\dagger\beta$; ψ_f the fields of the kind f. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- $\hat\gamma^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat\gamma^{(a)}$ along the frame directions of the fold and $\hat\gamma^\mu$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat\gamma^\mu, \hat\gamma^\nu\} = -2\hat h^{\mu\nu}\mathbf{1}$; $\hat\gamma_\mu = \hat h_{\mu\nu}\hat\gamma^\nu$. Unit: dimensionless. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar\Psi = \partial_M\bar\Psi - \bar\Psi\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection

of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu} D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2 \epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.

The Yang-Mills equation of the gluons and the Maxwell equation of the hypercharge field, which live in the bulk: away from the fold the field equations of eight dimensions, and at the fold the jump of the field strength along σ set by the currents of the fold (the colour current of the quarks; the hypercharge current of the fermions and of the Higgs doublet) and by the portal factors and the dipole density of dirac16complex. Status: STANDARD (INVENTED portal terms).

$$\mathrm{i}\hat{\gamma}^\mu \big(D_\mu + \mathrm{i}\,\mathcal{V}^Q_\mu\big) Q = \mathcal{Y}^u \tilde{\Phi}\, u_\mathrm{R} + \mathcal{Y}^d \Phi\, d_\mathrm{R}, \qquad \mathcal{V}^f_\mu = \sum_X c^f_X\, j_{X\mu} \tag{FE-19}$$

Symbols in (FE-19).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^\mu$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^\mu, \hat{\gamma}^\nu\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_\mu = \hat{h}_{\mu\nu}\hat{\gamma}^\nu$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M \bar{\Psi} = \partial_M \bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu} D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2 \epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^f_\mu$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^f_\mu = \sum_X c^f_X j_{X\mu}$ ($\mathcal{V}^Q, \mathcal{V}^u, \ldots, \mathcal{V}^\nu$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- Q, $\bar{Q}$: The left-handed quark doublets of the three generations, $Q = (u_\mathrm{L}, d_\mathrm{L})$ (colour triplets, hypercharge 1/6). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^\nu$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- $\tilde{\Phi}$: The conjugate Higgs doublet, $\tilde{\Phi} = \mathrm{i}\tau^2 \Phi^*$ (hypercharge $-1/2$). Unit: GeV. Status: STANDARD.
- u_R: The right-handed up-type quarks (up, charm and top; singlets of $SU(2)_L$, hypercharge 2/3). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- d_R: The right-handed down-type quarks (down, strange and bottom; hypercharge $-1/3$). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c^f_X: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c^ℓ_X and c^ν_X are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.
- j^μ_X: The convective current of the half X of dirac16complex, $j^\mu_X = \mathrm{i}(\bar{\Psi} P_X D^\mu \Psi - (D^\mu \bar{\Psi}) P_X \Psi)$ (real and joining the half with itself, DERIVED). Unit: GeV^8. Status: DERIVED.
- f: The label of a kind of Standard-Model multiplet: $f \in \{Q, u, d, \ell, e, \nu\}$ (quark doublets, up-quark singlets, down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.

$$\mathrm{i}\hat{\gamma}^\mu \big(D_\mu + \mathrm{i}\,\mathcal{V}^u_\mu\big) u_\mathrm{R} = (\mathcal{Y}^u)^\dagger \tilde{\Phi}^\dagger Q \tag{FE-20}$$

Symbols in (FE-20).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^\mu$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^\mu, \hat{\gamma}^\nu\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_\mu = \hat{h}_{\mu\nu}\hat{\gamma}^\nu$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^f_\mu$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^f_\mu = \sum_X c^f_X j_{X\mu}$ ($\mathcal{V}^Q, \mathcal{V}^u, \ldots, \mathcal{V}^\nu$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- u_R: The right-handed up-type quarks (up, charm and top; singlets of $SU(2)_L$, hypercharge 2/3). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^\nu$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\tilde{\Phi}$: The conjugate Higgs doublet, $\tilde{\Phi} = \mathrm{i}\tau^2\Phi^*$ (hypercharge $-1/2$). Unit: GeV. Status: STANDARD.
- Q, $\bar{Q}$: The left-handed quark doublets of the three generations, $Q = (u_\mathrm{L}, d_\mathrm{L})$ (colour triplets, hypercharge 1/6). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.

$$\mathrm{i}\hat{\gamma}^\mu\big(D_\mu + \mathrm{i}\,\mathcal{V}^d_\mu\big)d_\mathrm{R} = (\mathcal{Y}^d)^\dagger\Phi^\dagger Q \tag{FE-21}$$

Symbols in (FE-21).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^\mu$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^\mu, \hat{\gamma}^\nu\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_\mu = \hat{h}_{\mu\nu}\hat{\gamma}^\nu$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^f_\mu$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^f_\mu = \sum_X c^f_X j_{X\mu}$ ($\mathcal{V}^Q, \mathcal{V}^u, \ldots, \mathcal{V}^\nu$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- d_R: The right-handed down-type quarks (down, strange and bottom; hypercharge $-1/3$). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^\nu$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- Q, $\bar{Q}$: The left-handed quark doublets of the three generations, $Q = (u_\mathrm{L}, d_\mathrm{L})$ (colour triplets, hypercharge 1/6). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.

$$\mathrm{i}\hat{\gamma}^\mu\big(D_\mu + \mathrm{i}\,\mathcal{V}^\ell_\mu\big)\ell = \mathcal{Y}^e\Phi\, e_\mathrm{R} + \mathcal{Y}^\nu\tilde{\Phi}\,\nu_\mathrm{R} \tag{FE-22}$$

Symbols in (FE-22).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^\mu$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^\mu, \hat{\gamma}^\nu\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_\mu = \hat{h}_{\mu\nu}\hat{\gamma}^\nu$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- $D_M,\ D_\mu$: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^f_\mu$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^f_\mu = \sum_X c^f_X j_{X\mu}$ ($\mathcal{V}^Q, \mathcal{V}^u, \ldots, \mathcal{V}^\nu$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- $\ell,\ \bar{\ell}$: The left-handed lepton doublets of the three generations, $\ell = (\nu_\mathrm{L}, e_\mathrm{L})$ (hypercharge $-1/2$). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- $\mathcal{Y}^u,\ \mathcal{Y}^d,\ \mathcal{Y}^e,\ \mathcal{Y}^\nu$: The Yukawa matrices ($3 \times 3$ complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge $1/2$), a field of the fold. Unit: GeV. Status: AUTHOR.
- e_R: The right-handed electron, muon and tau (singlets, hypercharge -1). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- $\tilde{\Phi}$: The conjugate Higgs doublet, $\tilde{\Phi} = \mathrm{i}\tau^2\Phi^*$ (hypercharge $-1/2$). Unit: GeV. Status: STANDARD.
- ν_R: The three right-handed neutrinos (gauge singlets). Unit: $\mathrm{GeV}^{3/2}$. Status: ASSUMED.

$$\mathrm{i}\hat{\gamma}^\mu\big(D_\mu + \mathrm{i}\,\mathcal{V}^e_\mu\big)e_\mathrm{R} = (\mathcal{Y}^e)^\dagger\Phi^\dagger\ell \tag{FE-23}$$

Symbols in (FE-23).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^\mu$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^\mu, \hat{\gamma}^\nu\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_\mu = \hat{h}_{\mu\nu}\hat{\gamma}^\nu$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- $D_M,\ D_\mu$: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^f_\mu$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^f_\mu = \sum_X c^f_X j_{X\mu}$ ($\mathcal{V}^Q, \mathcal{V}^u, \ldots, \mathcal{V}^\nu$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- e_R: The right-handed electron, muon and tau (singlets, hypercharge -1). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- $\mathcal{Y}^u,\ \mathcal{Y}^d,\ \mathcal{Y}^e,\ \mathcal{Y}^\nu$: The Yukawa matrices ($3 \times 3$ complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge $1/2$), a field of the fold. Unit: GeV. Status: AUTHOR.
- $\ell,\ \bar{\ell}$: The left-handed lepton doublets of the three generations, $\ell = (\nu_\mathrm{L}, e_\mathrm{L})$ (hypercharge $-1/2$). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.

$$\mathrm{i}\hat{\gamma}^\mu\big(D_\mu + \mathrm{i}\,\mathcal{V}^\nu_\mu\big)\nu_\mathrm{R} = (\mathcal{Y}^\nu)^\dagger\tilde{\Phi}^\dagger\ell \tag{FE-24}$$

Symbols in (FE-24).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.

- $\hat{\gamma}^{\mu}$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^{\mu}$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^{\mu}, \hat{\gamma}^{\nu}\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_{\mu} = \hat{h}_{\mu\nu}\hat{\gamma}^{\nu}$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^{\mu} = \hat{h}^{\mu\nu}D_{\nu}$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^f_\mu$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^f_\mu = \sum_X c^f_X j_{X\mu}$ ($\mathcal{V}^Q, \mathcal{V}^u, \ldots, \mathcal{V}^\nu$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- ν_{R}: The three right-handed neutrinos (gauge singlets). Unit: $\mathrm{GeV}^{3/2}$. Status: ASSUMED.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^\nu$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\tilde{\Phi}$: The conjugate Higgs doublet, $\tilde{\Phi} = \mathrm{i}\tau^2\Phi^*$ (hypercharge $-1/2$). Unit: GeV. Status: STANDARD.
- ℓ, $\bar{\ell}$: The left-handed lepton doublets of the three generations, $\ell = (\nu_{\mathrm{L}}, e_{\mathrm{L}})$ (hypercharge $-1/2$). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.

The Dirac equations of the quark doublets (FE-19), the up-quark and down-quark singlets (FE-20, FE-21), the lepton doublets (FE-22), the charged-lepton singlets (FE-23) and the right-handed neutrinos (FE-24): every fermion sees the gauge fields at the fold, the Higgs doublet through its Yukawa matrix and dirac16complex through the flavour-blind vector potential of the current portals; the right-handed neutrinos see no gauge field. Status: STANDARD (INVENTED potentials).

The table states, for every field equation, how the solution satisfies it.

equation	how it is solved	residual	status
bulk gravity FE-1	the static block is an exact vacuum (FE-32); around an expanding fold the bulk metric FE-34 solves every component to the first order in the curvature of the fold over H^2 (FE-29 for the order 0)	the terms of the next order, of the relative size $(a_4'/H)^2 = 3 \times 10^{-91}$ today; the projected Weyl tensor that dust requires at the order of $\rho^2/(\lambda_{\mathrm{F}} V_*)$ (S-1) is not computed	DERIVED, OPEN
bulk sources of FE-1	the gas of the particles χ is confined to the fold within $1/q_{\mathrm{b}} \ll 1/H$ and is counted as matter of the fold; the gauge fields carry no mean field strength	the back-reaction of the profile of χ on the warp is left out	ASSUMED
junction FE-4, FE-5: time and space components	FE-71, FE-72, FE-59: the Friedmann equations of general relativity with $8\pi G_{\mathrm{N}} = \kappa_7/V_*$	the correction of high energies, of the relative size $(12/25)\mathsf{A}^2$ (S-1)	DERIVED
junction FE-4: extra-time component	FE-65, FE-72, S-3: the volume of the extra times held fixed	the mismatch $\Delta_{\mathrm{r}} = 0.842\rho_{\mathrm{c}}^{(0)}$ today (with no pressure of the baryons along the extra times; 0.793 or 0.891 with $-\rho_{\mathrm{b}}$ or $+\rho_{\mathrm{b}}$), carried by the pressure q_{hold} of the fold fluid	DERIVED, ASSUMED, OPEN
conservation FE-11	the fold loses what the bulk carries away: the widths of the W and Z bosons (FE-82) and of the particles χ (FE-86) into the bulk	0 (an identity of the junction condition with the bulk equations)	DERIVED

equation	how it is solved	residual	status
dirac16complex FE-12 to FE-14, Kohn-Sham FE-41	the Kohn-Sham states (S-7 to S-12), lifted to sixteen components	Kohn-Sham equation 1.0×10^{-4} (central differences), fold matching 2×10^{-15}, self-consistency 7.5×10^{-9}; Krein charges -1 to 10^{-16}	DERIVED
Higgs doublet FE-15	S-5: the ad hoc vacuum as a gauge rotation	the radial residual $\mathcal{R}_{\mathrm{rad}}$ (the table of S-5), zero once $\mathcal{A} = 0$	DERIVED
$SU(2)$ gauge fields FE-17	S-5: the pure-gauge $W_{x_4}^2$, constant along σ	0: the isospin charge density vanishes exactly	DERIVED
gluons FE-16 and hypercharge field FE-18	no mean field strengths (the gluon condensate enters FE-39 as a mean value)	0	DERIVED
quarks and leptons FE-19 to FE-24	no mean fermion fields; the plasma and the neutrinos are treated statistically (FE-48 to FE-62)	0 in the vacuum	STANDARD
neutrino flavours FE-48 to FE-50	the Sun and the supernova envelope (S-13, S-14)	halving the step changes P_{ee} by less than 10^{-6} (Sun) and 10^{-9} (envelope)	DERIVED
fast instability FE-51	S-15	the dispersion relation holds to 10^{-14} at the computed frequencies	DERIVED
relic density FE-61	S-19	relative tolerance 10^{-8} of the implicit integration	DERIVED
cosmological perturbations FE-54, FE-55	CAMB with the content of UM-4	the accuracy of CAMB (about 10^{-3} in C_l); background and ruler checked independently to 10^{-6}	STANDARD, DERIVED

11 The three neutrino flavours

The author asks for a correct flavour conversion in the universe model, because the flavour conversion of neutrinos shapes the explosion of massive stars (Gogilashvili and Tamborra, "Neutrino Flavor Conversion Shapes the Rate of Failed Core-collapse Supernovae", arXiv:2605.16504). The quarks and leptons of UM-4 live on the fold, as in UM-3, so this section solves the same flavour equations in the Sun and in core-collapse supernovae; the numbers are those of UM-3, recomputed in UM-4.

$$
\begin{aligned}
&\mathrm{i}\big(\partial_{x_4} + \hat{v}\cdot\vec{\nabla}\big)\,\varrho_{\vec{p}} = \big[\mathsf{H}_{\vec{p}},\ \varrho_{\vec{p}}\big] + \mathrm{i}\,\mathcal{C}_{\vec{p}}, \qquad \mathrm{i}\big(\partial_{x_4} + \hat{v}\cdot\vec{\nabla}\big)\,\bar{\varrho}_{\vec{p}} = \big[\bar{\mathsf{H}}_{\vec{p}},\ \bar{\varrho}_{\vec{p}}\big] + \mathrm{i}\,\bar{\mathcal{C}}_{\vec{p}},\\
&\mathsf{H}_{\vec{p}} = \frac{v_{\mathrm{loc}}^2}{v^2}\,\frac{U_{\mathrm{PMNS}}\,\mathsf{M}^2\,U_{\mathrm{PMNS}}^{\dagger}}{2E} + \mathsf{V}_{\mathrm{m}} + \mathsf{V}_{\nu\nu} + V_{\Psi}\mathbf{1}, \qquad \mathsf{M}^2 = \mathrm{diag}\big(m_1^2,\ m_2^2,\ m_3^2\big),\\
&\bar{\mathsf{H}}_{\vec{p}} = -\frac{v_{\mathrm{loc}}^2}{v^2}\,\frac{U_{\mathrm{PMNS}}\,\mathsf{M}^2\,U_{\mathrm{PMNS}}^{\dagger}}{2E} + \mathsf{V}_{\mathrm{m}} + \mathsf{V}_{\nu\nu} + V_{\Psi}\mathbf{1}
\end{aligned}
\tag{FE-48}
$$

Symbols in (FE-48).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\hat{v}$, $\hat{v}^\mu$: The direction of motion of a neutrino: the unit vector $\hat{v} = \vec{p}/|\vec{p}|$ ($\hat{v}_{\vec{p}}$, $\hat{v}_{\vec{q}}$ for the momenta $\vec{p}$, $\vec{q}$), and the four-vector $\hat{v}^\mu = (1, \hat{v})$ in the order $(x_4; x_1, x_2, x_3)$; $\mathrm{d}^2\hat{v}$ is the element of solid angle of the directions. Unit: dimensionless. Status: STANDARD.
- $\vec{\nabla}$: The gradient along ordinary space, $\vec{\nabla} = (\partial_{x_1}, \partial_{x_2}, \partial_{x_3})$ (in the local frame of the medium). Unit: 1/length. Status: STANDARD.

- $\varrho_{\vec{p}}$, $\bar{\varrho}_{\vec{p}}$: The 3×3 flavour density matrices of the neutrinos and of the antineutrinos of momentum $\vec{p}$ in the flavour basis (electron, muon, tau): the diagonal entries are the occupation numbers of the three flavours, the off-diagonal entries the coherence between the flavours; $\bar{\varrho}$ is defined with the flavour indices in the reversed order, so that it obeys an equation of the same form. Unit: dimensionless. Status: STANDARD.
- $\vec{p}$: The momentum of a neutrino (three components along ordinary space, in the rest frame of the medium); $|\vec{p}| = E$ for a neutrino of negligible mass. Unit: GeV. Status: STANDARD.
- $\mathsf{H}_{\vec{p}}$, $\bar{\mathsf{H}}_{\vec{p}}$: The flavour Hamiltonians (3×3 Hermitian matrices) of the neutrinos and of the antineutrinos of momentum $\vec{p}$. Unit: GeV. Status: STANDARD.
- $\mathcal{C}_{\vec{p}}$, $\bar{\mathcal{C}}_{\vec{p}}$: The collision terms ($3 \times 3$ matrices) of the neutrinos and the antineutrinos of momentum $\vec{p}$: the rates of change of $\varrho_{\vec{p}}$ and $\bar{\varrho}_{\vec{p}}$ by emission, absorption, scattering and pair processes with the matter. Unit: GeV. Status: STANDARD.
- v_{loc}: The local value of the electroweak vacuum, $v_{\mathrm{loc}}^2 = 2\langle\Phi^\dagger\Phi\rangle$: it sets the local masses of the quarks and leptons (each mass times v_{loc}/v) and the local Fermi constant; $v_{\mathrm{loc}} = v$ in the Standard-Model vacuum. Unit: GeV. Value: v in the late universe outside dense dirac16complex. Status: DERIVED.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_{\mathrm{F}})^{-1/2}$ with the Fermi constant $G_{\mathrm{F}} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- U_{PMNS}: The lepton mixing matrix (Pontecorvo-Maki-Nakagawa-Sakata) of the three Dirac neutrinos; its entries $(U_{\mathrm{PMNS}})_{\alpha i}$ connect the flavour state ν_α with the mass state ν_i. Unit: dimensionless. Value: $\theta_{12} = 33.40°$, $\theta_{23} = 49.14°$, $\theta_{13} = 8.54°$ (NuFIT 5.2, November 2022, normal ordering, the fit without the Super-Kamiokande atmospheric data); the Dirac phase is set to zero (ASSUMED; the fit prefers 197°). Status: STANDARD, ASSUMED (the phase).
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- M^2: The diagonal matrix of the squared masses of the three neutrino mass states, $\mathsf{M}^2 = \mathrm{diag}(m_1^2, m_2^2, m_3^2)$ (M appears only squared). Unit: GeV^2 (of M^2). Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- V_{m}: The matter potential of the neutrinos (a 3×3 matrix in flavour space) from their forward scattering on the electrons, positrons and neutrons of the medium. Unit: GeV. Status: STANDARD.
- $\mathsf{V}_{\nu\nu}$: The neutrino-neutrino potential (a 3×3 matrix in flavour space) from the forward scattering of a neutrino on the other neutrinos and antineutrinos of the medium; it depends on their flavour content and their directions. Unit: GeV. Status: STANDARD.
- V_Ψ: The potential energy that dirac16complex gives every neutrino through the current portal of the lepton doublets, $V_\Psi = \mathcal{V}^\ell_{x_4}$ (the same for the three flavours). Unit: GeV. Status: DERIVED.
- $\mathbf{1}$, $\mathbf{1}_4$, $\mathbf{1}_8$, $\mathbf{1}_{16}$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_4$, $\mathbf{1}_8$ and $\mathbf{1}_{16}$ are the identity matrices of 4, 8 and 16 rows. Unit: dimensionless. Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- $m_u, \ldots, m_3$: The masses of the quarks (m_u, m_c, m_t; m_d, m_s, m_b), of the charged leptons (m_e, m_μ, m_τ) and of the three neutrino mass states (m_1, m_2, m_3, Dirac masses). Unit: GeV. Value: measured values; $\Delta m^2_{21} = 7.41 \times 10^{-5}$ eV^2, $|\Delta m^2_{31}| = 2.511 \times 10^{-3}$ eV^2 in the computations of UM-4 (the global fit NuFIT 5.2 of the oscillation data, November 2022), 7.49×10^{-5} and 2.513×10^{-3} eV^2 in NuFIT 6.0 (2024); the lightest mass $m_1 = 0$ in the computations, below 23 meV by DESI DR2 with the CMB and NuFIT 6.0. Status: STANDARD.

$$
\begin{aligned}
&\mathsf{V}_{\mathrm{m}} = \sqrt{2}\,G_{\mathrm{F,loc}}\,\mathrm{diag}\big(n_e - n_{\bar{e}},\ 0,\ 0\big) - \frac{G_{\mathrm{F,loc}}}{\sqrt{2}}\,n_n \mathbf{1}, \qquad \mathsf{V}_{\nu\nu} = \sqrt{2}\,G_{\mathrm{F,loc}} \int \frac{\mathrm{d}^3\vec{q}}{(2\pi)^3}\big(1 - \hat{v}_{\vec{p}} \cdot \hat{v}_{\vec{q}}\big)\big(\varrho_{\vec{q}} - \bar{\varrho}_{\vec{q}}\big), \\
&V_\Psi = \mathcal{V}^\ell_{x_4} = \sum_X c^\ell_X\, j_{X\,x_4}, \qquad G_{\mathrm{F,loc}} = \frac{1}{\sqrt{2}\,v^2_{\mathrm{loc}}}, \\
&v^2_{\mathrm{loc}} = 2\langle\Phi^\dagger\Phi\rangle, \qquad v^2_{\mathrm{loc}} = v^2 + \mathcal{A}^2 \ \text{ (the ad hoc vacuum)}, \\
&v^2_{\mathrm{loc}} = v^2 - \frac{c_\Phi}{\lambda}\,\langle\bar{\Psi} P_{\mathrm{L}} \Psi\rangle \ \text{ (the relaxed vacuum with dirac16complex)}
\end{aligned}
\tag{FE-49}
$$

Symbols in (FE-49).

- V_m: The matter potential of the neutrinos (a 3×3 matrix in flavour space) from their forward scattering on the electrons, positrons and neutrons of the medium. Unit: GeV. Status: STANDARD.
- $G_{\mathrm{F,loc}}$: The local Fermi constant, $G_{\mathrm{F,loc}} = 1/(\sqrt{2}\, v_\mathrm{loc}^2)$ ($= G_\mathrm{F}$ where $v_\mathrm{loc} = v$). Unit: GeV^{-2}. Status: DERIVED.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- n_e: The number density of electrons (bound and free; in a supernova all are free, in the recombining universe $x_\mathrm{e} n_e$ is the density of the free electrons). Unit: GeV^3 (cm^{-3}). Status: STANDARD.
- $n_{\bar{e}}$: The number density of positrons. Unit: GeV^3. Status: STANDARD.
- n_n: The number density of neutrons. Unit: GeV^3. Status: STANDARD.
- $\mathbf{1}$, $\mathbf{1}_4$, $\mathbf{1}_8$, $\mathbf{1}_{16}$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_4$, $\mathbf{1}_8$ and $\mathbf{1}_{16}$ are the identity matrices of 4, 8 and 16 rows. Unit: dimensionless. Status: STANDARD.
- $\mathsf{V}_{\nu\nu}$: The neutrino-neutrino potential (a 3×3 matrix in flavour space) from the forward scattering of a neutrino on the other neutrinos and antineutrinos of the medium; it depends on their flavour content and their directions. Unit: GeV. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $\vec{q}$: The momentum of another neutrino of the medium (the integration variable of the neutrino-neutrino potential); $\mathrm{d}^3\vec{q}$ its volume element. Unit: GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $\hat{v}$, $\hat{v}^\mu$: The direction of motion of a neutrino: the unit vector $\hat{v} = \vec{p}/|\vec{p}|$ ($\hat{v}_{\vec{p}}$, $\hat{v}_{\vec{q}}$ for the momenta $\vec{p}$, $\vec{q}$), and the four-vector $\hat{v}^\mu = (1, \hat{v})$ in the order $(x_4; x_1, x_2, x_3)$; $\mathrm{d}^2\hat{v}$ is the element of solid angle of the directions. Unit: dimensionless. Status: STANDARD.
- $\vec{p}$: The momentum of a neutrino (three components along ordinary space, in the rest frame of the medium); $|\vec{p}| = E$ for a neutrino of negligible mass. Unit: GeV. Status: STANDARD.
- $\varrho_{\vec{p}}$, $\bar{\varrho}_{\vec{p}}$: The 3×3 flavour density matrices of the neutrinos and of the antineutrinos of momentum $\vec{p}$ in the flavour basis (electron, muon, tau): the diagonal entries are the occupation numbers of the three flavours, the off-diagonal entries the coherence between the flavours; $\bar{\varrho}$ is defined with the flavour indices in the reversed order, so that it obeys an equation of the same form. Unit: dimensionless. Status: STANDARD.
- V_Ψ: The potential energy that dirac16complex gives every neutrino through the current portal of the lepton doublets, $V_\Psi = \mathcal{V}^\ell_{x_4}$ (the same for the three flavours). Unit: GeV. Status: DERIVED.
- $\mathcal{V}^f_\mu$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^f_\mu = \sum_X c^f_X j_{X\mu}$ ($\mathcal{V}^Q, \mathcal{V}^u, \ldots, \mathcal{V}^\nu$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c^f_X: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c^ℓ_X and c^ν_X are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.
- j^μ_X: The convective current of the half X of dirac16complex, $j^\mu_X = \mathrm{i}(\bar{\Psi} P_X D^\mu \Psi - (D^\mu \bar{\Psi}) P_X \Psi)$ (real and joining the half with itself, DERIVED). Unit: GeV^8. Status: DERIVED.
- v_loc: The local value of the electroweak vacuum, $v^2_\mathrm{loc} = 2\langle \Phi^\dagger \Phi \rangle$: it sets the local masses of the quarks and leptons (each mass times v_loc/v) and the local Fermi constant; $v_\mathrm{loc} = v$ in the Standard-Model vacuum. Unit: GeV. Value: v in the late universe outside dense dirac16complex. Status: DERIVED.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\langle \cdots \rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the

Kohn-Sham state). Status: STANDARD.

- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_\mathrm{F})^{-1/2}$ with the Fermi constant $G_\mathrm{F} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- $\mathcal{A}$: The author's ad hoc upper component of the Higgs vacuum, almightyAdHoc, $\mathcal{A}(x_4) = v_\mathcal{A} \mathrm{e}^{-2a_{40}x_4}$. Unit: GeV. Status: AUTHOR.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: GeV$^{7/2}$. Status: AUTHOR.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.

The flavour equations of the neutrinos of momentum $\vec p$ and the potentials (FE-48, FE-49). Two facts make them those of the Standard Model with three Dirac neutrinos wherever $v_\mathrm{loc} = v$: the portal potential V_Ψ is the same for the three flavours and drops out of every commutator, and the Higgs portal moves v_loc by only $\Delta v^2/v^2 = -\mathfrak{g}_{h\chi}\rho_\chi/(\lambda v^3\omega_\mathrm{b}) \approx -10^{-50}\,\mathfrak{g}_{h\chi}$ at the local dark-matter density $\rho_\chi = 0.3$ GeV/cm^3 for $\omega_\mathrm{b} = 100$ GeV (the value falls as $1/\omega_\mathrm{b}$ for the heavier particles χ of UM-4; $\mathfrak{g}_{h\chi}$ is the coupling of the Higgs boson to χ of FE-47). Status: DERIVED (`um4_neutrino.py`, N1: $V_\Psi\mathbf{1}$ changes the evolution by at most 7×10^{-13}, a change of v_loc leaves the mixing matrix unchanged to 10^{-15} and scales the masses by v_loc/v). For a single neutrino along its path (no collisions, no neutrino-neutrino potential) FE-48 is a Schrodinger equation $\mathrm{i}\,\mathrm{d}\psi_\nu/\mathrm{d}r = \mathsf{H}\psi_\nu$ for the flavour amplitudes ψ_ν; for antineutrinos the vacuum term has U^*_PMNS and the matter potential the opposite sign. It is integrated here with the fourth-order commutator-free Magnus method (the exponentials of the Hamiltonian at the two Gauss points of each step, exact for a constant density), and every result is recomputed with half the step.

$$\tan 2\theta_\mathrm{m} = \frac{\sin 2\theta}{\cos 2\theta - 2EV_\mathrm{CC}/\Delta m^2}, \qquad V_\mathrm{CC} = \sqrt{2}\,G_\mathrm{F,loc}\,n_e, \qquad n_{e,\mathrm{res}} = \frac{\Delta m^2 \cos 2\theta}{2\sqrt{2}\,G_\mathrm{F,loc}\,E},$$
$$P_\mathrm{LZ} = \exp\Big(-\frac{\pi}{2}\,\frac{\Delta m^2 \sin^2 2\theta}{2E\cos 2\theta}\,\Big|\frac{\mathrm{d}\ln n_e}{\mathrm{d}r}\Big|^{-1}_\mathrm{res}\Big) \qquad \text{(FE-50)}$$

Symbols in (FE-50).

- θ_m: The mixing angle of two neutrino flavours in matter. Unit: dimensionless. Status: STANDARD.
- θ: The vacuum mixing angle of two neutrino flavours (in the two-flavour description of a resonance, one of the three angles $\theta_{12}, \theta_{13}, \theta_{23}$ of U_PMNS). Unit: dimensionless. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- V_CC: The charged-current matter potential of the electron neutrinos (forward scattering on electrons through W exchange). Unit: GeV. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m^2_{ij} = m_i^2 - m_j^2$ the squared-mass difference of the states i and j (Δm^2_{21}, Δm^2_{31}, Δm^2_{32}); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by v^2_loc/v^2 where this differs from 1; m without a label the mass of a mode of a field of the bulk as the fold sees it, $m^2 = \omega^2 - k^2$; for the W and Z bosons their complex pole mass, $m^2 = m_0^2 - \mathrm{i}m_0\Gamma_\mathrm{bulk}$ to the leading order; in the relic abundance of the particles χ their mass. Unit: GeV. Status: STANDARD.
- $G_\mathrm{F,loc}$: The local Fermi constant, $G_\mathrm{F,loc} = 1/(\sqrt{2}\,v^2_\mathrm{loc})$ ($= G_\mathrm{F}$ where $v_\mathrm{loc} = v$). Unit: GeV^{-2}. Status: DERIVED.
- n_e: The number density of electrons (bound and free; in a supernova all are free, in the recombining universe $x_e n_e$ is the density of the free electrons). Unit: GeV^3 (cm^{-3}). Status: STANDARD.
- $n_{e,\mathrm{res}}$: The electron density at which the flavour conversion in matter is resonant (the Mikheyev-Smirnov-Wolfenstein resonance). Unit: GeV^3. Status: STANDARD.
- P_LZ: The probability that a neutrino jumps between the eigenstates of matter at the resonance (Landau-Zener); $P_\mathrm{LZ} = 0$ is an adiabatic resonance. Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value:

3.141592654. Status: STANDARD.

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- r: A distance from a centre: the distance from the centre of the star along the path of a neutrino; in Newton's and Coulomb's laws the distance between two point sources on the fold. Unit: length. Status: STANDARD.
- res: The label of a quantity evaluated at the resonance, where $n_e = n_{e,\mathrm{res}}$. Status: STANDARD.

The mixing angle in matter, the resonance density and the Landau-Zener jump probability (FE-50). Status: STANDARD (`um4_neutrino.py`, N2: the formula for θ_m agrees with an exact diagonalisation to 10^{-16}, the Landau-Zener formula with an exact integration through a linear profile to 10^{-6}).

11.1 The Sun

$$\mathsf{H}_2 = \frac{1}{2}\begin{pmatrix} -\frac{\Delta m^2_{21}}{2E}\cos 2\theta_{12} + \cos^2\theta_{13}\,V_\mathrm{CC} & \frac{\Delta m^2_{21}}{2E}\sin 2\theta_{12} \\ \frac{\Delta m^2_{21}}{2E}\sin 2\theta_{12} & \frac{\Delta m^2_{21}}{2E}\cos 2\theta_{12} - \cos^2\theta_{13}\,V_\mathrm{CC}\end{pmatrix}, \qquad n_e(r) = 245\,N_\mathrm{A}\,\mathrm{e}^{-10.54\,r/R_\odot}\ \mathrm{cm}^{-3},$$

$$\mathsf{P}_\mathrm{ee} = \cos^4\theta_{13}\,\mathsf{P}_2 + \sin^4\theta_{13}, \qquad \mathsf{P}_2 = \frac{1}{2} + \left(\frac{1}{2} - \mathsf{P}_\mathrm{c}\right)\cos 2\theta_{12}\cos 2\theta_\mathrm{m}(r_\mathrm{p}),$$

$$\mathsf{P}_\mathrm{c} = \frac{\mathrm{e}^{-2\pi r_\odot \frac{\Delta m^2_{21}}{2E}\sin^2\theta_{12}} - \mathrm{e}^{-2\pi r_\odot \frac{\Delta m^2_{21}}{2E}}}{1 - \mathrm{e}^{-2\pi r_\odot \frac{\Delta m^2_{21}}{2E}}}, \qquad r_\odot = \frac{R_\odot}{10.54} \tag{S-13}$$

Symbols in (S-13).

- H_2: The two-flavour Hamiltonian of the electron neutrinos in the Sun (the mass states 1 and 2 with the potential $\cos^2\theta_{13}V_\mathrm{CC}$; the third state decouples). Unit: GeV (or 1/km). Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m^2_{ij} = m_i^2 - m_j^2$ the squared-mass difference of the states i and j (Δm^2_{21}, Δm^2_{31}, Δm^2_{32}); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by v^2_loc/v^2 where this differs from 1; m without a label the mass of a mode of a field of the bulk as the fold sees it, $m^2 = \omega^2 - k^2$; for the W and Z bosons their complex pole mass, $m^2 = m_0^2 - \mathrm{i}m_0\Gamma_\mathrm{bulk}$ to the leading order; in the relic abundance of the particles χ their mass. Unit: GeV. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- θ_{12}, θ_{13}, θ_{23}: The three mixing angles of U_PMNS (its fourth parameter is the Dirac phase, which does not enter the electron-flavour results of UM-4). Unit: dimensionless. Value: 33.40°, 8.54°, 49.14° in the computations of UM-4 ($\sin^2\theta_{12} = 0.303$, $\sin^2\theta_{13} = 0.02203$, $\sin^2\theta_{23} = 0.572$: NuFIT 5.2, November 2022, normal ordering, the fit without the Super-Kamiokande atmospheric data); 33.71°, 8.56°, 43.28° in NuFIT 6.0 (2024, with the Super-Kamiokande atmospheric data; $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$). Status: STANDARD.
- V_CC: The charged-current matter potential of the electron neutrinos (forward scattering on electrons through W exchange). Unit: GeV. Status: STANDARD.
- n_e: The number density of electrons (bound and free; in a supernova all are free, in the recombining universe $x_\mathrm{e}n_e$ is the density of the free electrons). Unit: GeV^3 (cm^{-3}). Status: STANDARD.
- r: A distance from a centre: the distance from the centre of the star along the path of a neutrino; in Newton's and Coulomb's laws the distance between two point sources on the fold. Unit: length. Status: STANDARD.
- N_A: Avogadro's number: $245N_\mathrm{A}$ cm^{-3} is 245 moles of electrons per cubic centimetre. Unit: per mole. Value: $6.02214076 \times 10^{23}$. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- $R_\odot$: The radius of the Sun. Unit: length. Value: 6.957×10^5 km. Status: STANDARD.
- P_ee: The survival probability of electron neutrinos: the probability that an electron neutrino is detected as an electron neutrino. Unit: dimensionless. Status: DERIVED.
- P_2: The survival probability of the two-flavour problem of the mass states 1 and 2. Unit: dimensionless. Status: DERIVED.

- $\mathsf{P}_{\rm c}$: The crossing probability: the probability that a neutrino jumps from one eigenstate of matter to the other on its way through the Sun (exact for an exponential profile, Petcov). Unit: dimensionless. Status: STANDARD.
- $\theta_{\rm m}$: The mixing angle of two neutrino flavours in matter. Unit: dimensionless. Status: STANDARD.
- $r_{\rm p}$: The radius at which the solar neutrinos of a given reaction are produced. Unit: length. Value: $0.05 R_\odot$ to $0.10 R_\odot$. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $r_\odot$: The scale height of the electron density of the Sun, $r_\odot = R_\odot/10.54$. Unit: length. Value: 6.6×10^4 km. Status: STANDARD.

The solar neutrinos: the third mass state decouples (it oscillates too fast and mixes only through θ_{13}), so the electron neutrinos evolve with the two-flavour Hamiltonian H_2 (with the potential $\cos^2\theta_{13} V_{\rm CC}$) through the electron density of the Sun in its exponential approximation (Bahcall), from the radius $r_{\rm p}$ of their production to the surface; the phases average out over the production region and the distance to the Earth, so the survival probability $\mathsf{P}_{\rm ee}$ follows from the probabilities of the mass states at the surface. The second line is the adiabatic solution with the exact crossing probability $\mathsf{P}_{\rm c}$ of an exponential profile (Petcov). Status: STANDARD (the equations), DERIVED (the solution below).

source	E (MeV)	$r_{\rm p}/R_\odot$	$\mathsf{P}_{\rm ee}$ (exact integration)	$\mathsf{P}_{\rm ee}$ (S-13)	Borexino
pp	0.267	0.10	0.5454	0.5454	0.57 ± 0.09
^{7}Be	0.862	0.06	0.5126	0.5126	0.53 ± 0.05
pep	1.44	0.10	0.5083	0.5083	0.43 ± 0.11
^{8}B	5.0	0.05	0.3362	0.3362	
^{8}B	8.1	0.05	0.3080	0.3080	0.37 ± 0.08
^{8}B	10.0	0.05	0.3017	0.3017	
^{8}B	15.0	0.05	0.2951	0.2951	

The survival probability of solar electron neutrinos in UM-4 (with $\theta_{12} = 33.40°$, $\theta_{13} = 8.54°$, $\Delta m^2_{21} = 7.41 \times 10^{-5}$ eV^2): the exact integration and the adiabatic formula agree to 10^{-6}, $\mathsf{P}_{\rm c} < 10^{-10}$ at every energy (the Sun is adiabatic), and halving the step changes nothing at the level 10^{-6}. At low energy $\mathsf{P}_{\rm ee}$ approaches the vacuum average $\cos^4\theta_{13}(1-\frac{1}{2}\sin^2 2\theta_{12}) + \sin^4\theta_{13} = 0.553$, at high energy the full matter conversion $\cos^4\theta_{13}\sin^2\theta_{12} + \sin^4\theta_{13} = 0.290$; the Borexino measurements (Nature 562, 505 (2018), with the high-metallicity solar model) lie within one standard deviation. Status: DERIVED (`um4_flavour.py`, F1).

11.2 The envelope of a supernova

$$\rho_{\rm b}(r) = 10^4 \text{ g/cm}^3 \Big(\frac{r}{10^9 \text{ cm}}\Big)^{-3}, \qquad n_e = \frac{Y_{\rm e}\,\rho_{\rm b}}{m_{\rm u}}, \qquad \gamma_{\rm LZ} = \frac{\Delta m^2 \sin^2 2\theta}{2E\cos 2\theta}\,\frac{r_{\rm res}}{3} \tag{S-14}$$

Symbols in (S-14).

- $\rho_{\rm b}$: The (rest-mass) energy density of the baryons: in a supernova the local density of the matter (quoted in g/cm^3), in cosmology the mean density of the baryons ($\rho_{\rm b}^{(0)}$ today). Unit: GeV^4. Status: STANDARD.
- r: A distance from a centre: the distance from the centre of the star along the path of a neutrino; in Newton's and Coulomb's laws the distance between two point sources on the fold. Unit: length. Status: STANDARD.
- n_e: The number density of electrons (bound and free; in a supernova all are free, in the recombining universe $x_e n_e$ is the density of the free electrons). Unit: GeV^3 (cm^{-3}). Status: STANDARD.
- $Y_{\rm e}$: The electron fraction of stellar matter: the number of electrons per nucleon. Unit: dimensionless. Value: 0.5 in the envelope. Status: STANDARD.
- $m_{\rm u}$: The atomic mass unit, the mass per nucleon of stellar matter to within one per cent: $n_e = Y_e\rho/m_{\rm u}$. Unit: g. Value: 1.66054×10^{-24} g. Status: STANDARD.
- $\gamma_{\rm LZ}$: The adiabaticity parameter of a resonance: the jump probability at the resonance is $\exp(-\pi\gamma_{\rm LZ}/2)$; $\gamma_{\rm LZ} \gg 1$ is adiabatic. Unit: dimensionless. Status: STANDARD.

- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m^2_{ij} = m^2_i - m^2_j$ the squared-mass difference of the states i and j (Δm^2_{21}, Δm^2_{31}, Δm^2_{32}); Δm^2 without labels the difference of the two states of a resonance, $m^2_2 - m^2_1$, multiplied by $v^2_{\rm loc}/v^2$ where this differs from 1; m without a label the mass of a mode of a field of the bulk as the fold sees it, $m^2 = \omega^2 - k^2$; for the W and Z bosons their complex pole mass, $m^2 = m^2_0 - {\rm i}m_0\Gamma_{\rm bulk}$ to the leading order; in the relic abundance of the particles χ their mass. Unit: GeV. Status: STANDARD.
- θ: The vacuum mixing angle of two neutrino flavours (in the two-flavour description of a resonance, one of the three angles $\theta_{12}, \theta_{13}, \theta_{23}$ of $U_{\rm PMNS}$). Unit: dimensionless. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- $r_{\rm res}$: The radius of a resonance: where the electron density equals the resonance density $n_{e,{\rm res}}$ ($r_{\rm H}$ for Δm^2_{31}, $r_{\rm L}$ for Δm^2_{21}). Unit: length. Status: STANDARD.

The outer layers of the progenitor star (density $\rho_{\rm b} \propto r^{-3}$, electron fraction $Y_{\rm e} = 0.5$, $m_{\rm u}$ the atomic mass unit), through which the neutrinos of a supernova pass after they have left the dense core; $\gamma_{\rm LZ}$ is the adiabaticity of a resonance at the radius $r_{\rm res}$ (the exponent of FE-50 is $-\frac{\pi}{2}\gamma_{\rm LZ}$, since $|{\rm d}\ln n_e/{\rm d}r| = 3/r$). The three-flavour equation with the full mixing matrix ($\theta_{23} = 49.14°$, $\Delta m^2_{31} = 2.511 \times 10^{-3}$ eV^2 in the normal ordering) is integrated from $0.2r_{\rm H}$ to $5r_{\rm L}$ at $E = 15$ MeV, starting with an electron neutrino (antineutrino). Status: STANDARD (the profile and the equations), DERIVED (the solution).

ordering	particle	P_{ee}	expected for adiabatic resonances	probabilities of the mass states at the surface
normal	ν_e	0.022030	$\lvert U_{e3}\rvert^2 = 0.022030$	ν_3: $1 - 1.5 \times 10^{-6}$
normal	$\bar\nu_e$	0.681644	$\lvert U_{e1}\rvert^2 = 0.681645$	$\bar\nu_1$: $1 - 1.5 \times 10^{-6}$
inverted	ν_e	0.296325	$\lvert U_{e2}\rvert^2 = 0.296325$	ν_2: $1 - 1.5 \times 10^{-6}$
inverted	$\bar\nu_e$	0.022031	$\lvert U_{e3}\rvert^2 = 0.022030$	$\bar\nu_3$: $1 - 1.6 \times 10^{-6}$

The resonance of Δm^2_{31} lies at $r_{\rm H} = 16\,833$ km in the normal ordering and at 17 002 km in the inverted one (which keeps $|m^2_3 - m^2_2| = 2.511 \times 10^{-3}$ eV^2, so that $m^2_3 - m^2_1 = -2.437 \times 10^{-3}$ eV^2), and that of Δm^2_{21} at $r_{\rm L} = 73\,195$ km; all are adiabatic ($\gamma_{\rm LZ} = 215$, 210 and 655, so $P_{\rm LZ} = 0$). An electron neutrino, produced in the heaviest state of matter, leaves the star as ν_3 in the normal ordering and as ν_2 in the inverted one; an electron antineutrino leaves as $\bar\nu_1$ or $\bar\nu_3$. The exact integration agrees with these limits to 1.1×10^{-6} (and with half the step to 10^{-9}). Shock waves and turbulence in the envelope, which can make the high resonance non-adiabatic, and the collective conversion deeper inside are not included in this static profile. Status: DERIVED (`um4_flavour.py`, F2).

11.3 The fast instability near the neutrinosphere

$$\mathcal{G}_{\rm ELN}(\hat v) = \sqrt{2}\,G_{\rm F,loc}\int_0^\infty \frac{E^2\,{\rm d}E}{(2\pi)^3}\Big[\mathcal{F}_e - \bar{\mathcal{F}}_e - \big(\mathcal{F}_x - \bar{\mathcal{F}}_x\big)\Big],$$
$$\det\Big[\eta^{\mu\nu}_{\rm M} + \int {\rm d}^2\hat v\;\mathcal{G}_{\rm ELN}(\hat v)\,\frac{\hat v^\mu \hat v^\nu}{\Omega_{\rm fw} - \vec K\cdot\hat v}\Big] = 0, \qquad \hat v^\mu = (1,\ \hat v) \tag{FE-51}$$

Symbols in (FE-51).

- $\mathcal{G}_{\rm ELN}(\hat v)$: The angular distribution of the electron lepton number carried by the neutrinos minus that carried by the heavy-lepton neutrinos, per unit solid angle of the direction $\hat v$, multiplied by $\sqrt{2}G_{\rm F,loc}$. Unit: GeV. Status: STANDARD.
- $\hat v$, $\hat v^\mu$: The direction of motion of a neutrino: the unit vector $\hat v = \vec p/|\vec p|$ ($\hat v_{\vec p}$, $\hat v_{\vec q}$ for the momenta $\vec p$, $\vec q$), and the four-vector $\hat v^\mu = (1, \hat v)$ in the order $(x_4; x_1, x_2, x_3)$; ${\rm d}^2\hat v$ is the element of solid angle of the directions. Unit: dimensionless. Status: STANDARD.
- $G_{\rm F,loc}$: The local Fermi constant, $G_{\rm F,loc} = 1/(\sqrt{2}\,v^2_{\rm loc})$ ($= G_{\rm F}$ where $v_{\rm loc} = v$). Unit: GeV^{-2}. Status: DERIVED.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.

- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $\mathcal{F}_e$, $\bar{\mathcal{F}}_e$, $\mathcal{F}_x$, $\bar{\mathcal{F}}_x$: The occupation numbers of the electron neutrinos, of the electron antineutrinos, and of one heavy-lepton neutrino and antineutrino ($x = \mu$ or τ), as functions of the energy E and the direction $\hat{v}$ (the diagonal entries of $\varrho_{\vec{p}}$ and $\bar{\varrho}_{\vec{p}}$); in a transport code, the same for the number of particles in an energy bin. Unit: dimensionless. Status: STANDARD.
- $\eta_{\mathrm{M}}^{\mu\nu}$: The Minkowski metric of the four-vectors $\hat{v}^\mu$, $\eta_{\mathrm{M}} = \mathrm{diag}(1, -1, -1, -1)$ in the order $(x_4; x_1, x_2, x_3)$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+, +, +, -$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- Ω_{fw}: The (complex) frequency of a flavour wave of the neutrino gas, measured from the shift caused by the potentials; an imaginary part greater than zero is an instability. Unit: GeV. Status: STANDARD.
- $\vec{K}$: The (real) wave vector of a flavour wave of the neutrino gas, measured from the shift caused by the potentials. Unit: GeV. Status: STANDARD.

$$\begin{aligned}
\Big(\Omega_{\mathrm{fw}} - K_{\mathrm{r}}\,\mathsf{u}\Big)\mathsf{Q}(\mathsf{u}) &= -2\pi \int_{-1}^{1} \mathrm{d}\mathsf{u}'\,\big(1 - \mathsf{u}\mathsf{u}'\big)\,\mathcal{G}_{\mathrm{ELN}}(\mathsf{u}')\,\mathsf{Q}(\mathsf{u}') \qquad \text{(axially symmetric mode)},\\
\Big(\Omega_{\mathrm{fw}} - K_{\mathrm{r}}\,\mathsf{u}\Big)\mathsf{Q}(\mathsf{u}) &= \pi \int_{-1}^{1} \mathrm{d}\mathsf{u}'\,\sqrt{1-\mathsf{u}^2}\sqrt{1-\mathsf{u}'^2}\,\mathcal{G}_{\mathrm{ELN}}(\mathsf{u}')\,\mathsf{Q}(\mathsf{u}') \qquad \text{(first azimuthal mode)}
\end{aligned} \tag{S-15}$$

Symbols in (S-15).

- Ω_{fw}: The (complex) frequency of a flavour wave of the neutrino gas, measured from the shift caused by the potentials; an imaginary part greater than zero is an instability. Unit: GeV. Status: STANDARD.
- K_{r}: The wave number of a flavour wave along the radial direction (the radial component of $\vec{K}$). Unit: GeV (1/length). Status: STANDARD.
- u: The cosine of the angle between the direction of motion of a neutrino and the radial direction; with the subscript of a flavour (the electron neutrinos or the electron antineutrinos), its value at the edge of the emission cone of that flavour. Unit: dimensionless. Status: STANDARD.
- $\mathsf{Q}(\mathsf{u})$: The amplitude of a small flavour coherence (the off-diagonal entry of ϱ) of the neutrinos moving with the direction cosine u, in a flavour wave. Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\mathcal{G}_{\mathrm{ELN}}(\hat{v})$: The angular distribution of the electron lepton number carried by the neutrinos minus that carried by the heavy-lepton neutrinos, per unit solid angle of the direction $\hat{v}$, multiplied by $\sqrt{2}G_{\mathrm{F,loc}}$. Unit: GeV. Status: STANDARD.

The fast instability as an eigenvalue problem: for an electron-lepton-number distribution that depends only on the cosine u of the angle to the radial direction, a flavour wave with the wave number K_{r} along the radius and the amplitude $\mathsf{Q}(\mathsf{u})$ (axially symmetric, or proportional to the cosine of the azimuth) has the frequencies Ω_{fw} that are the eigenvalues of S-15 (FE-48 linearised; equivalent to the dispersion relation FE-51, which the solutions satisfy to 10^{-14}). The distribution is taken from the neutrinospheres: electron neutrinos emitted uniformly (half-isotropically) from a sphere of radius $R_e = 30$ km and antineutrinos from $R_{\bar{e}} = 27$ km (they decouple deeper and are more forward peaked), seen at $r = 40$ km, where each fills the cone $\mathsf{u} > \sqrt{1-(R/r)^2}$ ($R = R_e$ or $R_{\bar{e}}$); α is the ratio of their number densities, the heavy flavours cancel in the electron lepton number, and 200 Gauss-Legendre nodes discretise the angle. Status: STANDARD (the equations), DERIVED (the solution).

$\alpha = n_{\bar{\nu}_e}/n_{\nu_e}$	0.50	0.75	0.80	0.95	1.20
angular crossing	no	no	yes	yes	yes
growth rate $\operatorname{Im}\Omega_{\mathrm{fw}}$ $(\sqrt{2}G_{\mathrm{F}}n_{\nu_e})$	0	0	0.0137	0.0556	0.0925
growth length, factor e (cm)	–	–	22	5.4	3.2

The electron lepton number crosses zero as a function of the angle exactly when $\alpha > (1-\mathsf{u}_{\bar{e}})/(1-\mathsf{u}_e) = 0.7744$ (u_e, $\mathsf{u}_{\bar{e}}$ the edges of the two cones); without a crossing every frequency is real (growth below 10^{-17}), with a crossing the axially symmetric mode grows. For a luminosity $L = 5\times10^{52}$ erg/s and a mean energy $\langle E\rangle = 12$ MeV of the electron neutrinos, their density at 40 km is $n_{\nu_e} = L(1-\mathsf{u}_e)/(2\pi R_e^2 c\langle E\rangle) = 5.2\times10^{32}$ cm^{-3} (c the speed of light), so $\sqrt{2}G_{\mathrm{F}}n_{\nu_e} = 3.34$ cm^{-1} (with the Fermi constant G_{F} of the vacuum); the growth length of the table is the inverse of the growth rate in cm^{-1}. The instability grows within centimetres: much faster than anything else in the star, it drives the flavours toward equipartition wherever the crossing exists. Status: DERIVED (`um4_flavour.py`, F3; `um4_neutrino.py`, N3, for the crossing criterion with general distributions).

11.4 The flavour equipartition and the heating behind the shock

$$\begin{aligned}&\mathcal{F}_e \to \mathcal{F}_e - \delta_{\mathrm{FC}}, \qquad \bar{\mathcal{F}}_e \to \bar{\mathcal{F}}_e - \delta_{\mathrm{FC}}, \qquad \mathcal{F}_x = \bar{\mathcal{F}}_x \to \mathcal{F}_x + \frac{\delta_{\mathrm{FC}}}{2},\\ &\delta_{\mathrm{FC}} = \max\Big(\frac{2}{3}\Big(\min\left(\mathcal{F}_e,\ \bar{\mathcal{F}}_e\right) - \mathcal{F}_x\Big),\ \max\left(\mathcal{F}_e,\ \bar{\mathcal{F}}_e\right) - 1\Big),\\ &\text{applied where}\ \ \rho_{\mathrm{b}} < \rho_{\mathrm{FC}}\ \ \text{and}\ \ x_4 - x_{4,\mathrm{b}} > t_{\mathrm{FC}}\end{aligned} \tag{FE-52}$$

Symbols in (FE-52).

- $\mathcal{F}_e$, $\bar{\mathcal{F}}_e$, $\mathcal{F}_x$, $\bar{\mathcal{F}}_x$: The occupation numbers of the electron neutrinos, of the electron antineutrinos, and of one heavy-lepton neutrino and antineutrino ($x = \mu$ or τ), as functions of the energy E and the direction $\hat{v}$ (the diagonal entries of $\varrho_{\vec{p}}$ and $\bar{\varrho}_{\vec{p}}$); in a transport code, the same for the number of particles in an energy bin. Unit: dimensionless. Status: STANDARD.
- δ_{FC}: The occupation (or number per energy bin) of electron neutrino-antineutrino pairs converted into heavy-lepton pairs by the flavour equipartition (the conversion that equalises the occupations of the three flavours where the fast flavour instability acts); a negative value converts heavy-lepton pairs into electron pairs. Unit: dimensionless. Status: ASSUMED.
- ρ_{b}: The (rest-mass) energy density of the baryons: in a supernova the local density of the matter (quoted in g/cm^3), in cosmology the mean density of the baryons ($\rho_{\mathrm{b}}^{(0)}$ today). Unit: GeV^4. Status: STANDARD.
- ρ_{FC}: The baryon density below which the flavour equipartition is applied (the critical density ρ_{c} of Gogilashvili and Tamborra). Unit: GeV^4 (g/cm^3). Value: 10^9, 10^{10}, 10^{11} or 10^{13} g/cm^3. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $x_{4,\mathrm{b}}$: The time of the bounce of the collapsing stellar core. Unit: length (= time). Status: STANDARD.
- t_{FC}: The time after the bounce from which the flavour equipartition is applied. Unit: time. Value: 0.02 s. Status: STANDARD.

The flavour equipartition of Gogilashvili and Tamborra (FE-52), applied in every energy bin. Status: STANDARD (the constraints), ASSUMED (this form). On 10^5 random occupations in $[0,1]^3$ it conserves the electron lepton number and the numbers of neutrinos and of antineutrinos to 10^{-15}, keeps every occupation between 0 and 1 (the Pauli bound acts in 22% of the cases), and a second application changes nothing (10^{-16}). Status: DERIVED (`um4_flavour.py`, F4; `um4_neutrino.py`, N4).

$$\dot{Q}_{\mathrm{heat}} \propto \int_0^\infty \mathrm{d}E\, E^3\big(\mathcal{F}_e + \bar{\mathcal{F}}_e\big), \qquad \mathcal{F}_e,\ \bar{\mathcal{F}}_e,\ \mathcal{F}_x \propto \frac{L_{\mathrm{s}}}{E_{\mathrm{s}}^2}\Big(\frac{E}{E_{\mathrm{s}}}\Big)^{\alpha_{\mathrm{s}}} \mathrm{e}^{-(\alpha_{\mathrm{s}}+1)E/E_{\mathrm{s}}} \tag{S-16}$$

Symbols in (S-16).

- $\dot{Q}_{\mathrm{heat}}$: The rate at which the absorption of electron neutrinos and antineutrinos heats the matter of the gain region (energy per time and mass). Unit: energy per time and mass. Status: STANDARD.

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- $\mathcal{F}_e$, $\bar{\mathcal{F}}_e$, $\mathcal{F}_x$, $\bar{\mathcal{F}}_x$: The occupation numbers of the electron neutrinos, of the electron antineutrinos, and of one heavy-lepton neutrino and antineutrino ($x = \mu$ or τ), as functions of the energy E and the direction $\hat{v}$ (the diagonal entries of $\varrho_{\vec{p}}$ and $\bar{\varrho}_{\vec{p}}$); in a transport code, the same for the number of particles in an energy bin. Unit: dimensionless. Status: STANDARD.
- L_s: The luminosity (energy per time) of one neutrino species leaving the neutrinosphere. Unit: erg/s. Value: (2 to 5) $\times 10^{52}$ erg/s. Status: STANDARD.
- E_s: The mean energy of one neutrino species. Unit: GeV (MeV). Value: 10 to 16 MeV. Status: STANDARD.
- α_s: The shape (pinching) parameter of a neutrino spectrum (Keil, Raffelt and Janka). Unit: dimensionless. Value: 3. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

The charged-current heating of the gain region behind the stalled shock ($\nu_e n \to e^- p$ and $\bar{\nu}_e p \to e^+ n$: the cross-section grows as E^2 and the deposited energy is about E), with the number spectra S-16 of the neutrinos (L_s and E_s the luminosity and the mean energy of the species, $\alpha_\mathrm{s} = 3$ the shape parameter). In the table the species are the electron neutrinos ν_e, the electron antineutrinos $\bar{\nu}_e$ and each of the four heavy-lepton species ν_x; the mean energies after the equipartition are those of the mixed spectra. Status: STANDARD, ESTIMATE (equal weights of the two reactions).

phase	L_s of ν_e, $\bar{\nu}_e$, ν_x (10^{52} erg/s)	E_s (MeV)	E_s after (MeV)	heating after / before
accretion (0.2 s)	5, 5, 2	11, 14, 16	11.3, 16.1, 14.1	0.96
accretion, softer ν_x	5, 5, 2	11, 14, 14	10.9, 15.3, 13.2	0.86
end of accretion	4, 4, 2.5	11, 13, 15	11.9, 14.7, 13.6	1.12
cooling (comparable luminosities)	3, 3, 3	10, 12, 14	11.4, 13.8, 13.0	1.55

Equipartition removes absorbable electron neutrinos and antineutrinos; it hardens the spectrum of the electron antineutrinos in every case and that of the electron neutrinos unless the heavy-lepton spectra are as soft as in the second row. When the heavy-lepton luminosity is well below the electron-flavour one (the accretion phase) the loss of numbers wins and the heating falls; when they are comparable the harder spectra win and it rises. In the transport simulations of Gogilashvili and Tamborra (195 progenitor models; arXiv:2605.16504 version 3) the flavour equipartition, applied wherever the baryon density falls below a critical value, raises the fraction of failed explosions from 25.6% without conversion to 50.8%, 77.4%, 93.3% and 96.4% for the critical densities 10^9, 10^{10}, 10^{11} and 10^{13} g/cm^3: the deeper the equilibration, the stronger the suppression of the revival of the shock. This estimate includes neither the feedback on the cooling nor the hydrodynamics. Since the flavour physics of UM-4 is that of the Standard Model with three Dirac neutrinos (FE-49), their results apply to UM-4 unchanged. Status: ESTIMATE (`um4_flavour.py`, F4).

11.5 The newer global fit, the Dirac phase and the lightest mass

The computations above use the global fit NuFIT 5.2 (November 2022; normal ordering, the fit without the atmospheric data of Super-Kamiokande: $\sin^2\theta_{12} = 0.303$, $\sin^2\theta_{23} = 0.572$, $\sin^2\theta_{13} = 0.02203$, $\Delta m^2_{21} = 7.41 \times 10^{-5}$ eV^2, $\Delta m^2_{31} = 2.511 \times 10^{-3}$ eV^2) with the Dirac phase $\delta_\mathrm{CP} = 0$ and the lightest mass $m_1 = 0$. The newer fit NuFIT 6.0 (I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, I. Martinez-Soler, J. P. Pinheiro and T. Schwetz, JHEP 12 (2024) 216) has two data sets. With the atmospheric data of Super-Kamiokande (the normal ordering preferred by $\Delta\chi^2 = 6.1$): $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{23} = 0.470$, $\sin^2\theta_{13} = 0.02215$, $\delta_\mathrm{CP} = 212°$, $\Delta m^2_{21} = 7.49 \times 10^{-5}$ eV^2 and $\Delta m^2_{31} = 2.513 \times 10^{-3}$ eV^2. Without them: $\sin^2\theta_{12} = 0.307$, $\sin^2\theta_{23} = 0.561$, $\sin^2\theta_{13} = 0.02195$, $\delta_\mathrm{CP} = 177°$, $\Delta m^2_{21} = 7.49 \times 10^{-5}$ eV^2, $\Delta m^2_{31} = 2.534 \times 10^{-3}$ eV^2 in the normal ordering.

$$\left(|U_{ei}|^2\right)_{i=1,2,3} = \left(\cos^2\theta_{12}\cos^2\theta_{13},\ \sin^2\theta_{12}\cos^2\theta_{13},\ \sin^2\theta_{13}\right) \tag{S-17}$$

Symbols in (S-17).

- U_{ei}: The electron row of the mixing matrix, $U_{\mathrm{ei}} = (U_{\mathrm{PMNS}})_{ei}$: $|U_{ei}|^2$ is the electron-flavour fraction of the mass state i. Unit: dimensionless. Value: $|U_{\mathrm{ei}}|^2 = 0.682$, 0.296 and 0.022 for $i = 1, 2, 3$ (NuFIT 5.2, the fit without the Super-Kamiokande atmospheric data: the values of the record; NuFIT 6.0 with them gives 0.677, 0.301 and 0.022). Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal), and in a tensor of ordinary space such as h_{ij} one of the directions x_1, x_2, x_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- $\theta_{12},\ \theta_{13},\ \theta_{23}$: The three mixing angles of U_{PMNS} (its fourth parameter is the Dirac phase, which does not enter the electron-flavour results of UM-4). Unit: dimensionless. Value: 33.40°, 8.54°, 49.14° in the computations of UM-4 ($\sin^2\theta_{12} = 0.303$, $\sin^2\theta_{13} = 0.02203$, $\sin^2\theta_{23} = 0.572$: NuFIT 5.2, November 2022, normal ordering, the fit without the Super-Kamiokande atmospheric data); 33.71°, 8.56°, 43.28° in NuFIT 6.0 (2024, with the Super-Kamiokande atmospheric data; $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$). Status: STANDARD.

The Dirac phase drops out of every result of UM-4. The mixing matrix is $U_{\mathrm{PMNS}} = R_{23}U_{13}(\delta_{\mathrm{CP}})R_{12}$ (rotations by the three angles, the phase attached to the middle one), and its electron row contains only θ_{12} and θ_{13} (S-17; computed for $\delta_{\mathrm{CP}} = 0$, 177°, 212° and 274°, identical to 7×10^{-18}). Every flavour result of UM-4 is a result for electron neutrinos in matter: the matter potential acts on ν_e only, so the phase can be moved into ν_μ and ν_τ, and the three-flavour evolution through the envelope of a supernova gives the same survival probability for $\delta_{\mathrm{CP}} = 0$ and 212° (largest difference 6.0×10^{-14} at 10 and 20 MeV, for neutrinos and antineutrinos). The phase enters only the appearance channels, which UM-4 does not compute: the vacuum probability of $\nu_\mu \to \nu_e$ at 295 km and 0.6 GeV (the baseline and the peak energy of T2K) is 0.0421 for $\delta_{\mathrm{CP}} = 0$, 0.0478 for 212° and 0.0541 for 270°, that of $\bar{\nu}_\mu \to \bar{\nu}_e$ 0.0421, 0.0346 and 0.0291, and the Jarlskog invariant of CP violation is -0.0178 at 212°. Status: DERIVED (`um4_numass.py`, NM3).

$$m_2 = \sqrt{m_1^2 + \Delta m_{21}^2}, \qquad m_3 = \sqrt{m_1^2 + \Delta m_{31}^2}, \qquad m_1 < 23 \text{ meV} \ \ (95 \text{ per cent}), \qquad m_{\mathrm{eff}}^2 = \sum_i |U_{\mathrm{ei}}|^2 m_i^2 \tag{S-18}$$

Symbols in (S-18).

- $m_u, \ldots, m_3$: The masses of the quarks (m_u, m_c, m_t; m_d, m_s, m_b), of the charged leptons (m_e, m_μ, m_τ) and of the three neutrino mass states (m_1, m_2, m_3, Dirac masses). Unit: GeV. Value: measured values; $\Delta m_{21}^2 = 7.41 \times 10^{-5}$ eV^2, $|\Delta m_{31}^2| = 2.511 \times 10^{-3}$ eV^2 in the computations of UM-4 (the global fit NuFIT 5.2 of the oscillation data, November 2022), 7.49×10^{-5} and 2.513×10^{-3} eV^2 in NuFIT 6.0 (2024); the lightest mass $m_1 = 0$ in the computations, below 23 meV by DESI DR2 with the CMB and NuFIT 6.0. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m_{ij}^2 = m_i^2 - m_j^2$ the squared-mass difference of the states i and j (Δm_{21}^2, Δm_{31}^2, Δm_{32}^2); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by v_{loc}^2/v^2 where this differs from 1; m without a label the mass of a mode of a field of the bulk as the fold sees it, $m^2 = \omega^2 - k^2$; for the W and Z bosons their complex pole mass, $m^2 = m_0^2 - \mathrm{i}m_0\Gamma_{\mathrm{bulk}}$ to the leading order; in the relic abundance of the particles χ their mass. Unit: GeV. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal), and in a tensor of ordinary space such as h_{ij} one of the directions x_1, x_2, x_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- m_{eff}: The effective mass of the electron neutrino measured in the beta decay of tritium, $m_{\mathrm{eff}}^2 = \sum_i |U_{\mathrm{ei}}|^2 m_i^2$. Unit: eV. Value: below 0.45 eV (KATRIN 2025, 90 per cent). Status: STANDARD.
- U_{ei}: The electron row of the mixing matrix, $U_{\mathrm{ei}} = (U_{\mathrm{PMNS}})_{ei}$: $|U_{ei}|^2$ is the electron-flavour fraction of the mass state i. Unit: dimensionless. Value: $|U_{\mathrm{ei}}|^2 = 0.682$, 0.296 and 0.022 for $i = 1, 2, 3$ (NuFIT 5.2, the fit

without the Super-Kamiokande atmospheric data: the values of the record; NuFIT 6.0 with them gives 0.677, 0.301 and 0.022). Status: STANDARD.

The lightest mass. The smallest sums of the masses are 0.0588 eV in the normal and 0.0989 eV in the inverted ordering. DESI DR2 (the BAO) with the CMB, combined with NuFIT 6.0 (W. Elbers et al., arXiv:2503.14744; the cosmological-constant model, 95 per cent), bounds the lightest mass to $m_1 < 23$ meV in the normal ordering and the sum of the masses to 0.101 eV (normal); the inverted ordering is disfavoured but not excluded. At $m_1 = 23$ meV the masses are 23.00, 24.57 and 55.15 meV, their sum 0.1027 eV; the effective mass of the beta decay of tritium is $m_{\mathrm{eff}} = 8.84$ to 24.64 meV in this range, far below the limit 0.45 eV of KATRIN (2025). Within the range the lightest mass changes one result of UM-4, the capture of the cosmic neutrinos (S-28): on 100 g of tritium the rate falls from 6.88 per year at $m_1 = 0$ to 4.41 at 5 meV, 4.25 at 10 meV and 4.15 per year at 23 meV, because ν_1 becomes slow (mean speed 0.023 today at 23 meV) and loses the doubling of the capture of relativistic neutrinos (4.06 per year if all states were at rest); and the sum of the masses, which enters the CMB and BAO calculation through $\Omega_\nu h^2_{100} = \sum_i m_i/(93.14\ \mathrm{eV})$, rises from 0.0588 to 0.1027 eV. The record (NuFIT 5.2 without the atmospheric data of Super-Kamiokande, $\delta_{\mathrm{CP}} = 0$, $m_1 = 0$) gives the solar survival probabilities 0.5454, 0.5126, 0.5083 and 0.3080 at the energies of the pp, ^{7}Be, pep and ^{8}B neutrinos; NuFIT 6.0 gives 0.5427, 0.5108, 0.5066 and 0.3119 in the same data set and 0.5418, 0.5100, 0.5059 and 0.3126 in the preferred one; the largest relative change is 1.3 per cent like with like and 1.5 per cent against the preferred data set. Status: DERIVED (`um4_numass.py`, NM1, NM2, NM4 and NM5), STANDARD (the fits and the bounds).

12 The thermal history

$$\rho_{\mathrm{SM}} = \frac{\pi^2}{30}\, g_\rho(\mathsf{T})\, \mathsf{T}^4, \qquad s = \frac{2\pi^2}{45}\, g_{\mathrm{s}}(\mathsf{T})\, \mathsf{T}^3, \qquad H_4 = \sqrt{\frac{8\pi^3 g_\rho}{90}}\, \frac{\mathsf{T}^2}{M_{\mathrm{Pl}}}, \qquad M_{\mathrm{Pl}} = G_{\mathrm{N}}^{-1/2} \tag{FE-60}$$

Symbols in (FE-60).

- ρ_{SM}: The energy density of the Standard-Model particles. Unit: GeV^4. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- g_ρ: The effective number of relativistic degrees of freedom of the energy density. Unit: dimensionless. Value: 106.75 in the Standard Model above the electroweak scale. Status: STANDARD.
- T: The temperature of the Standard-Model plasma (of the photons after the annihilation of electrons and positrons). Unit: GeV. Status: STANDARD.
- s: The entropy density of the Standard-Model plasma. Unit: GeV^3. Status: STANDARD.
- g_{s}: The effective number of relativistic degrees of freedom of the entropy density. Unit: dimensionless. Value: 3.909 today. Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\, h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- M_{Pl}: The Planck mass, $M_{\mathrm{Pl}} = G_{\mathrm{N}}^{-1/2}$. Unit: GeV. Value: 1.221×10^{19} GeV. Status: STANDARD.
- G_{N}: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_{\mathrm{N}} = 1/M_{\mathrm{Pl}}^2$ with the Planck mass $M_{\mathrm{Pl}} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.

The radiation of the Standard Model in thermal equilibrium and the expansion rate of the radiation era (FE-60). The effective numbers of degrees of freedom are computed here from the exact Bose and Fermi integrals of every particle of the Standard Model with its mass (quarks and gluons above the QCD transition, taken at $\mathsf{T} = 0.155$ GeV, the light hadrons below; the neutrinos decouple at 1 MeV, and afterwards the ratio of their temperature to T follows from the conserved entropy of the photons, electrons and positrons, reaching $(4/11)^{1/3}$ after the annihilation of the electrons and positrons): $g_\rho = g_{\mathrm{s}} = 106.75$ above the electroweak scale, 97.4 and 96.6 at 50 GeV, 85.6 and 85.4 at 5 GeV, 76.4 and 75.5 at 1 GeV, 10.76 at 10 MeV, 10.64 and 10.62 at 1 MeV, and today $g_{\mathrm{s}} = 3.909$, $g_\rho = 3.363$ (the transition region near 0.155 GeV and the decoupling of the neutrinos at one temperature are approximations). Three right-handed neutrinos in equilibrium would add 5.25. Status: STANDARD, DERIVED (`um4_thermal.py`, T1).

12.1 The relic density of the particles χ

$$\frac{\mathrm{d}n_\chi}{\mathrm{d}x_4} + 3H_4\, n_\chi = -\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle\big(n_\chi^2 - n_{\chi,\mathrm{eq}}^2\big), \qquad n_{\chi,\mathrm{eq}} = g_\chi\Big(\frac{\omega_\mathrm{b}\mathsf{T}}{2\pi}\Big)^{3/2}\mathrm{e}^{-\omega_\mathrm{b}/\mathsf{T}}, \qquad \Omega_\chi h_{100}^2 \simeq \frac{1.07\times10^9\ \mathrm{GeV}^{-1}\, x_\mathrm{f}}{\sqrt{g_*}\, M_\mathrm{Pl}\,\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle} \tag{FE-61}$$

Symbols in (FE-61).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- n_χ: The number density of the dark-matter particles χ (particles plus antiparticles). Unit: GeV^3. Status: DERIVED.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44\times10^{-42}$ GeV. Status: STANDARD.
- σ_{ann}: The annihilation cross-section of the particles χ into Standard-Model particles (through the portals) in the normalisation of the relic abundance, where n_χ counts the four particle and the four antiparticle states of the level at the fold: $\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle$ is a quarter of the annihilation rate $(\sigma\mathsf{v})_{\chi\bar\chi}$ of one particle χ_a with one antiparticle $\bar\chi_a$. Unit: GeV^{-2}. Status: DERIVED.
- v: The relative velocity of two annihilating particles; $\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle$ is the thermal average of the product. Unit: dimensionless. Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- $n_{\chi,\mathrm{eq}}$: The number density of χ in thermal equilibrium with the plasma (non-relativistic). Unit: GeV^3. Status: STANDARD.
- g_χ: The number of internal states of the level of χ at the fold that are in thermal equilibrium. Unit: dimensionless. Value: 8 (four particles and four antiparticles). Status: DERIVED.
- ω_b: The energy at zero momentum of the state of dirac16complex at the fold in the limit $H \to 0$ (the bound state of UM-3): the particle χ of UM-4 has the mass $\omega_\mathrm{r} = \omega_\mathrm{b}(1 - H/(2q_\mathrm{b})) + \ldots$, equal to ω_b up to the relative correction $H/(2q_\mathrm{b})$. Unit: GeV. Status: DERIVED.
- T: The temperature of the Standard-Model plasma (of the photons after the annihilation of electrons and positrons). Unit: GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- Ω_χ: The density parameter of the dark matter today. Unit: dimensionless. Value: $\Omega_\chi h_{100}^2 = 0.120$ (Planck 2018). Status: DERIVED.
- h_{100}: The present expansion rate of the fold in units of 100 km/s/Mpc. Unit: dimensionless. Value: 0.674 (Planck 2018). Status: STANDARD.
- x_f: The ratio of the mass of χ to the temperature at which its annihilations stop (freeze-out). Unit: dimensionless. Value: 23 to 30 (for $\omega_\mathrm{b} = 10$ GeV to 10 TeV). Status: STANDARD.
- g_*: The effective number of relativistic degrees of freedom at the freeze-out (g_ρ there). Unit: dimensionless. Status: STANDARD.
- M_Pl: The Planck mass, $M_\mathrm{Pl} = G_\mathrm{N}^{-1/2}$. Unit: GeV. Value: 1.221×10^{19} GeV. Status: STANDARD.

$$\frac{\mathrm{d}\mathsf{Y}}{\mathrm{d}\mathsf{x}} = -\frac{s\,\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle}{\mathsf{x}\,H_4}\Big(1 + \frac{1}{3}\frac{\mathrm{d}\ln g_\mathrm{s}}{\mathrm{d}\ln\mathsf{T}}\Big)\big(\mathsf{Y}^2 - \mathsf{Y}_\mathrm{eq}^2\big), \qquad \mathsf{Y} = \frac{n_\chi}{s}, \qquad \mathsf{Y}_\mathrm{eq} = \frac{n_{\chi,\mathrm{eq}}}{s}, \qquad \mathsf{x} = \frac{\omega_\mathrm{b}}{\mathsf{T}},$$
$$n_{\chi,\mathrm{eq}} = \frac{g_\chi\,\omega_\mathrm{b}^2\,\mathsf{T}}{2\pi^2}\,K_2\Big(\frac{\omega_\mathrm{b}}{\mathsf{T}}\Big), \qquad \Omega_\chi h_{100}^2 = \frac{\omega_\mathrm{b}\,\mathsf{Y}(\infty)\,s^{(0)}}{\rho_\mathrm{c}^{(0)}/h_{100}^2} \tag{S-19}$$

Symbols in (S-19).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.

- Y: The yield of the particles χ: their number density divided by the entropy density, $\mathsf{Y} = n_\chi/s$ (constant when they neither annihilate nor are produced). Unit: dimensionless. Status: STANDARD.
- x: The ratio of the mass of χ to the temperature, $\mathsf{x} = \omega_\mathrm{b}/\mathsf{T}$ (the time variable of the freeze-out). Unit: dimensionless. Status: STANDARD.
- s: The entropy density of the Standard-Model plasma. Unit: GeV^3. Status: STANDARD.
- σ_ann: The annihilation cross-section of the particles χ into Standard-Model particles (through the portals) in the normalisation of the relic abundance, where n_χ counts the four particle and the four antiparticle states of the level at the fold: $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle$ is a quarter of the annihilation rate $(\sigma\mathsf{v})_{\chi\bar\chi}$ of one particle χ_a with one antiparticle $\bar\chi_a$. Unit: GeV^{-2}. Status: DERIVED.
- v: The relative velocity of two annihilating particles; $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle$ is the thermal average of the product. Unit: dimensionless. Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- g_s: The effective number of relativistic degrees of freedom of the entropy density. Unit: dimensionless. Value: 3.909 today. Status: STANDARD.
- T: The temperature of the Standard-Model plasma (of the photons after the annihilation of electrons and positrons). Unit: GeV. Status: STANDARD.
- Y_eq: The yield of χ in thermal equilibrium, $n_{\chi,\mathrm{eq}}/s$. Unit: dimensionless. Status: STANDARD.
- n_χ: The number density of the dark-matter particles χ (particles plus antiparticles). Unit: GeV^3. Status: DERIVED.
- $n_{\chi,\mathrm{eq}}$: The number density of χ in thermal equilibrium with the plasma (non-relativistic). Unit: GeV^3. Status: STANDARD.
- ω_b: The energy at zero momentum of the state of dirac16complex at the fold in the limit $H \to 0$ (the bound state of UM-3): the particle χ of UM-4 has the mass $\omega_\mathrm{r} = \omega_\mathrm{b}(1 - H/(2q_\mathrm{b})) + \ldots$, equal to ω_b up to the relative correction $H/(2q_\mathrm{b})$. Unit: GeV. Status: DERIVED.
- g_χ: The number of internal states of the level of χ at the fold that are in thermal equilibrium. Unit: dimensionless. Value: 8 (four particles and four antiparticles). Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- K_2: The modified Bessel function of the second kind of order 2 ($K_2(y) \approx \sqrt{\pi/(2y)}\,\mathrm{e}^{-y}$ for large y). Unit: dimensionless. Status: STANDARD.
- Ω_χ: The density parameter of the dark matter today. Unit: dimensionless. Value: $\Omega_\chi h_{100}^2 = 0.120$ (Planck 2018). Status: DERIVED.
- h_{100}: The present expansion rate of the fold in units of 100 km/s/Mpc. Unit: dimensionless. Value: 0.674 (Planck 2018). Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- ρ_c: The critical density, the energy density of a spatially flat fold at the expansion rate H_4; $\rho_\mathrm{c}^{(0)}$ its value today. Unit: GeV^4. Value: $\rho_\mathrm{c}^{(0)} = 1.0537 \times 10^{-5}\,h_{100}^2$ GeV/cm^3. Status: STANDARD.

FE-61 rewritten for the yield Y (the number density per entropy, constant without annihilation) as a function of $\mathsf{x} = \omega_\mathrm{b}/\mathsf{T}$, with the equilibrium density valid also for relativistic χ (K_2 the modified Bessel function of the second kind), and the relic density from the yield after freeze-out ($s^{(0)} = 2891.2$ cm^{-3} the entropy density today, $\rho_\mathrm{c}^{(0)}/h_{100}^2 = 1.0537 \times 10^{-5}$ GeV/cm^3). The level of dirac16complex at the fold holds four particle states (one multiplet of the rotations of ordinary space and of the extra times: two Dirac fermions χ_a, $a = 1, 2$) and four antiparticle states, so $g_\chi = 8$ and FE-61, written for the total density, has $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle = (\sigma\mathsf{v})_{\chi\bar\chi}/4$, where $(\sigma\mathsf{v})_{\chi\bar\chi}$ is the annihilation of one χ_a with one $\bar\chi_a$ summed over the final states. The mass of χ is its rest energy $\omega_\mathrm{r} \simeq \omega_\mathrm{b}$ (FE-40; the difference is of the relative size $H/(2q_\mathrm{b})$, below 10^{-3} for the long-lived particles with $q_\mathrm{b} \le m_\Psi/2$ and up to 0.03 for the strongest bindings of S-6). The equation is integrated from $\mathsf{x} = 1$ to $\mathsf{x} = 10^5$ with an implicit method; continued to $\mathsf{x} = 10^6$ the relic density changes by less than 10^{-4}. Status: STANDARD (the equation), DERIVED (the solution).

ω_b (GeV)	10	30	100	300	1000	3000	10000
$\langle\sigma_\mathrm{ann}\mathsf{v}\rangle$ for $\Omega_\chi h^2_{100} = 0.120$ (10^{-26} cm^3/s)	2.29	2.17	2.14	2.20	2.22	2.21	2.25
freeze-out x_f	23.1	24.0	25.1	26.2	27.4	28.4	29.5
g_* at freeze-out	67.9	78.6	85.2	86.4	93.8	103.9	106.5
estimate of FE-61 ($\Omega_\chi h^2_{100}$)	0.125	0.128	0.130	0.131	0.130	0.129	0.130

A thermal relic with the measured dark-matter density (Planck 2018: $\Omega_\chi h^2_{100} = 0.1200$) needs, in the normalisation of FE-61, an s-wave annihilation $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle \approx 2.2{\times}10^{-26}$ cm^3/s for every mass from 10 GeV to 10 TeV (so $(\sigma\mathsf{v})_{\chi\bar\chi} \approx 9{\times}10^{-26}$ cm^3/s); the analytic estimate of FE-61 agrees within 10 per cent. Status: DERIVED (`um4_thermal.py`, T2).

$$\sigma_\mathrm{pair}\mathsf{v} \le \frac{4\pi}{\omega_\mathrm{b}^2\,\mathsf{v}}, \qquad \left\langle\frac{1}{\mathsf{v}}\right\rangle = \sqrt{\frac{\mathsf{x}}{\pi}}: \qquad \Omega_\chi h^2_{100} = 0.120 \;\text{ at }\; \omega_\mathrm{b} = 9.69\times10^4 \text{ GeV} \tag{S-20}$$

Symbols in (S-20).

- σ_pair: The cross-section of the annihilation of one particle χ_a with one antiparticle $\bar\chi_a$ of the same species, summed over the final states; $\sigma_\mathrm{pair}\mathsf{v}$ its product with the relative velocity (a constant for s-wave annihilation). Unit: GeV^{-2}. Status: DERIVED.
- v: The relative velocity of two annihilating particles; $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle$ is the thermal average of the product. Unit: dimensionless. Status: STANDARD.
- ω_b: The energy at zero momentum of the state of dirac16complex at the fold in the limit $H \to 0$ (the bound state of UM-3): the particle χ of UM-4 has the mass $\omega_\mathrm{r} = \omega_\mathrm{b}(1 - H/(2q_\mathrm{b})) + \ldots$, equal to ω_b up to the relative correction $H/(2q_\mathrm{b})$. Unit: GeV. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- x: The ratio of the mass of χ to the temperature, $\mathsf{x} = \omega_\mathrm{b}/\mathsf{T}$ (the time variable of the freeze-out). Unit: dimensionless. Status: STANDARD.
- Ω_χ: The density parameter of the dark matter today. Unit: dimensionless. Value: $\Omega_\chi h^2_{100} = 0.120$ (Planck 2018). Status: DERIVED.
- h_{100}: The present expansion rate of the fold in units of 100 km/s/Mpc. Unit: dimensionless. Value: 0.674 (Planck 2018). Status: STANDARD.

A thermal relic of UM-4 must be strongly bound. Partial-wave unitarity bounds the s-wave annihilation $\sigma_\mathrm{pair}\mathsf{v}$ of one χ with one $\bar\chi$ (the first relation, spin-averaged; v the relative velocity, with the thermal average of the Maxwell-Boltzmann distribution, and $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle = \sigma_\mathrm{pair}\mathsf{v}/4$ in FE-61); with this largest annihilation the relic density of S-19 equals the measured one at $\omega_\mathrm{b} = 9.69\times10^4$ GeV (integrated to $\mathsf{x} = 10^7$; 9.63×10^4 and 9.68×10^4 GeV to 10^5 and 10^6: at the unitarity bound the annihilation grows like $\sqrt{\mathsf{x}}$ and continues long after freeze-out), and heavier particles would be overproduced by freeze-out (the classic bound of K. Griest and M. Kamionkowski, Phys. Rev. Lett. 64 (1990) 615). The mass depends on the bookkeeping of the channels within the multiplet: if only the singlet of the extra-time $SU(2)$ annihilates into the Standard Model, half the rate, 6.86×10^4 GeV; if every channel is open, twice the rate, 1.37×10^5 GeV (one Dirac species alone: 1.38×10^5 GeV). The lifetime of S-6 puts the particles above this mass unless they are strongly bound: its bound on ω_b falls below the unitarity mass only for $q_\mathrm{b} > 0.906\,m_\Psi$ (0.875 if every channel is open, 0.929 for the singlet alone). Only so strongly bound particles χ, of masses of the order of 10^4 to 10^5 GeV, can be a thermal relic; for a weaker binding their abundance needs a non-thermal origin. Status: DERIVED (`um4_thermal.py`, T7), OPEN (the origin of the abundance).

12.2 The portals of χ to the Standard Model

$$\mathfrak{g}_\Phi = \frac{\partial\omega_\mathrm{b}}{\partial\mu_\mathrm{L}}\,\frac{c_\Phi}{V_*}, \qquad \mathfrak{g}_G = \sum_X \frac{\partial\omega_\mathrm{b}}{\partial\mu_X}\,\frac{c_X^G}{4V_*}, \qquad \mathfrak{g}_f = \sum_X \frac{\partial\omega_\mathrm{b}}{\partial\mu_X}\,\frac{2\omega_\mathrm{b}\,c_X^f}{V_*}, \qquad \frac{\partial\omega_\mathrm{b}}{\partial\mu_X} = -\frac{q_\mathrm{b}}{\omega_\mathrm{b}}\,\frac{\partial q_\mathrm{b}}{\partial\mu_X},$$
$$\Delta\omega_\mathrm{b} = \mathfrak{g}_\Phi\,\Delta\langle\Phi^\dagger\Phi\rangle + \mathfrak{g}_G\,\Delta\langle G^2\rangle + \sum_f \mathfrak{g}_f\,\Delta n_f, \qquad \mathfrak{g}_{h\chi} = \mathfrak{g}_\Phi\,v = -\frac{m_\Psi^2\,\mu_\mathrm{L}^2}{2\,\omega_\mathrm{b}\,v} \quad (\mu_\mathrm{R} = 0) \tag{FE-47}$$

Symbols in (FE-47).

- $\mathfrak{g}_\Phi$, $\mathfrak{g}_G$, $\mathfrak{g}_f$, $\mathfrak{g}_{h\chi}$: The four-dimensional couplings of the particles χ of dirac16complex to the Standard Model: to $\Phi^\dagger\Phi$, to the gluon kinetic density, to the number densities of the fermions of kind f, and the Yukawa coupling of the Higgs boson to χ. Unit: $\mathfrak{g}_\Phi$: GeV^{-1}; $\mathfrak{g}_G$: GeV^{-3}; $\mathfrak{g}_f$: GeV^{-2}; $\mathfrak{g}_{h\chi}$: dimensionless. Status: DERIVED.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- ω_b: The energy at zero momentum of the state of dirac16complex at the fold in the limit $H \to 0$ (the bound state of UM-3): the particle χ of UM-4 has the mass $\omega_\mathrm{r} = \omega_\mathrm{b}(1 - H/(2q_\mathrm{b})) + \ldots$, equal to ω_b up to the relative correction $H/(2q_\mathrm{b})$. Unit: GeV. Status: DERIVED.
- μ_X, μ_L, μ_R: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_\mathrm{L}P_\mathrm{L} + \mu_\mathrm{R}P_\mathrm{R})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- c_X^G: The gluon-kinetic portal coupling of the half X of dirac16complex. Unit: GeV^{-7}. Status: INVENTED.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c_X^f: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c_X^ℓ and c_X^ν are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.
- f: The label of a kind of Standard-Model multiplet: $f \in \{Q, u, d, \ell, e, \nu\}$ (quark doublets, up-quark singlets, down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.
- q_b: The inverse width of a state of dirac16complex at the fold: inside the turning point its profile falls as $\mathrm{e}^{-q_\mathrm{b}|\sigma|}$. Unit: GeV. Status: DERIVED.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- $\langle G^2\rangle$: The mean square of the gluon field strength, $\langle G^A_{\mu\nu}G^{A\mu\nu}\rangle$ (the gluon condensate in the QCD vacuum, a thermal value in the plasma). Unit: GeV^4. Status: STANDARD.
- G^A_M, G^A_μ, G^A_{MN}: The gluon field of $SU(3)_C$ ($A = 1, \ldots, 8$): in the bulk G^A_M, a field of eight dimensions in its geometric normalisation (the covariant derivative $\partial - \mathrm{i}G$), and on the fold the canonically normalised four-dimensional field $G^A_\mu(x)$ of its massless mode, $g_3 G^A_\mu(x) = G^A_\mu(\sigma = 0, x)$; G^A_{MN} and $G^A_{\mu\nu}$ their field strengths. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- n_f: The number density of the Standard-Model fermions of kind f (particles minus antiparticles, summed over generations and gauge components), the time component of the mean current $\langle J^\mu_f\rangle$ in the rest frame of the medium. Unit: GeV^3. Status: STANDARD.

- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_{\mathrm{F}})^{-1/2}$ with the Fermi constant $G_{\mathrm{F}} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.

The couplings of the particles χ to the Standard Model (FE-47): the Higgs-portal Yukawa coupling $\mathfrak{g}_{h\chi}$ is fixed by the binding, the current-portal couplings $\mathfrak{g}_f$ are free (through the INVENTED c_X^f). Status: DERIVED.

$$\sigma_{\mathrm{pair}}\mathsf{v} = \sum_f \frac{\mathfrak{g}_f^2\,\omega_{\mathrm{b}}^2\,\mathsf{v}^4}{96\pi}, \qquad \langle \mathsf{v}^4 \rangle = \frac{60}{\mathsf{x}^2}, \qquad \langle \sigma_{\mathrm{ann}}\mathsf{v} \rangle = \frac{\langle \sigma_{\mathrm{pair}}\mathsf{v} \rangle}{4}, \qquad \sigma_{\mathrm{SI}} = \frac{\mu_{\mathrm{red}}^2\,f_{\mathrm{N}}^2}{\pi}, \qquad f_{\mathrm{N}} = 3\,\mathfrak{g}_{\mathrm{q}} \quad \text{(quark portals)},$$

$$f_{\mathrm{N}} = \mathfrak{g}_{\mathrm{p}} = \frac{\alpha_{\mathrm{em}}}{3\pi}\,\mathfrak{g}_{\mathrm{l}} \sum_l \ln \frac{\omega_{\mathrm{b}}^2}{\max\left(m_l^2,\ q_{\mathrm{t}}^2\right)} \quad \text{(lepton portals)}, \qquad f_{\mathrm{N}} = \frac{\mathfrak{g}_{h\chi}\,f_{\mathrm{H}}\,m_{\mathrm{N}}}{v\,m_h^2} \quad \text{(Higgs portal)} \tag{S-21}$$

Symbols in (S-21).

- σ_{pair}: The cross-section of the annihilation of one particle χ_a with one antiparticle $\bar{\chi}_a$ of the same species, summed over the final states; $\sigma_{\mathrm{pair}}\mathsf{v}$ its product with the relative velocity (a constant for s-wave annihilation). Unit: GeV^{-2}. Status: DERIVED.
- v: The relative velocity of two annihilating particles; $\langle \sigma_{\mathrm{ann}}\mathsf{v} \rangle$ is the thermal average of the product. Unit: dimensionless. Status: STANDARD.
- $\mathfrak{g}_\Phi$, $\mathfrak{g}_G$, $\mathfrak{g}_f$, $\mathfrak{g}_{h\chi}$: The four-dimensional couplings of the particles χ of dirac16complex to the Standard Model: to $\Phi^\dagger\Phi$, to the gluon kinetic density, to the number densities of the fermions of kind f, and the Yukawa coupling of the Higgs boson to χ. Unit: $\mathfrak{g}_\Phi$: GeV^{-1}; $\mathfrak{g}_G$: GeV^{-3}; $\mathfrak{g}_f$: GeV^{-2}; $\mathfrak{g}_{h\chi}$: dimensionless. Status: DERIVED.
- f: The label of a kind of Standard-Model multiplet: $f \in \{Q, u, d, \ell, e, \nu\}$ (quark doublets, up-quark singlets, down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.
- ω_{b}: The energy at zero momentum of the state of dirac16complex at the fold in the limit $H \to 0$ (the bound state of UM-3): the particle χ of UM-4 has the mass $\omega_{\mathrm{r}} = \omega_{\mathrm{b}}(1 - H/(2q_{\mathrm{b}})) + \ldots$, equal to ω_{b} up to the relative correction $H/(2q_{\mathrm{b}})$. Unit: GeV. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $\langle \cdots \rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- x: The ratio of the mass of χ to the temperature, $\mathsf{x} = \omega_{\mathrm{b}}/\mathsf{T}$ (the time variable of the freeze-out). Unit: dimensionless. Status: STANDARD.
- σ_{ann}: The annihilation cross-section of the particles χ into Standard-Model particles (through the portals) in the normalisation of the relic abundance, where n_χ counts the four particle and the four antiparticle states of the level at the fold: $\langle \sigma_{\mathrm{ann}}\mathsf{v} \rangle$ is a quarter of the annihilation rate $(\sigma\mathsf{v})_{\chi\bar{\chi}}$ of one particle χ_a with one antiparticle $\bar{\chi}_a$. Unit: GeV^{-2}. Status: DERIVED.
- σ_{SI}: The spin-independent cross-section of the scattering of χ on a nucleon (normalised per nucleon, as the direct-detection limits are quoted). Unit: GeV^{-2} (cm^2). Status: ESTIMATE.
- μ_{red}: The reduced mass of χ and a nucleon, $\omega_{\mathrm{b}} m_{\mathrm{N}}/(\omega_{\mathrm{b}} + m_{\mathrm{N}})$. Unit: GeV. Status: STANDARD.
- f_{N}: The effective coupling of χ to a nucleon in the coherent scattering (a vector or scalar contact interaction). Unit: GeV^{-2}. Status: ESTIMATE.
- $\mathfrak{g}_{\mathrm{q}}$, $\mathfrak{g}_{\mathrm{l}}$, $\mathfrak{g}_{\mathrm{p}}$: The current-portal couplings $\mathfrak{g}_f$ taken equal for all chiral quark states ($\mathfrak{g}_{\mathrm{q}}$) or for the nine chiral lepton states that take part, the six of the lepton doublets and the three right-handed charged leptons ($\mathfrak{g}_{\mathrm{l}}$), and the coupling to the proton that the lepton loop induces ($\mathfrak{g}_{\mathrm{p}}$). Unit: GeV^{-2}. Status: ESTIMATE.
- α_{em}: The fine-structure constant. Unit: dimensionless. Value: 1/137.036. Status: STANDARD.
- l: The multipole index and number ($l = 0, 1, 2, \ldots$) of an expansion in Legendre polynomials of the direction of a photon or a neutrino relative to the wave vector, or of the CMB sky in spherical harmonics. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m_{ij}^2 = m_i^2 - m_j^2$ the squared-mass difference of the states i and j (Δm_{21}^2, Δm_{31}^2, Δm_{32}^2); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by v_{loc}^2/v^2 where this differs from 1; m without a label the mass of a mode of a field of the

bulk as the fold sees it, $m^2 = \omega^2 - k^2$; for the W and Z bosons their complex pole mass, $m^2 = m_0^2 - \mathrm{i}m_0\Gamma_{\mathrm{bulk}}$ to the leading order; in the relic abundance of the particles χ their mass. Unit: GeV. Status: STANDARD.

- q_{t}: The momentum transfer of the scattering of χ on a xenon nucleus in a detector. Unit: GeV. Value: 0.05 GeV. Status: STANDARD.
- m_{N}: The mass of a nucleon. Unit: GeV. Value: 0.939 GeV. Status: STANDARD.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_{\mathrm{F}})^{-1/2}$ with the Fermi constant $G_{\mathrm{F}} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- m_h: The mass of the Higgs boson. Unit: GeV. Value: 125.25 GeV (Particle Data Group 2022). Status: STANDARD.
- f_{H}: The nucleon matrix element of the Higgs coupling: the fraction of the nucleon mass carried by the quark masses and, through the trace anomaly, by the gluons. Unit: dimensionless. Value: 0.30. Status: STANDARD.

$$\begin{aligned}\mathcal{L}_{\mathrm{P}} = & -c_\Phi\left(\Phi^\dagger\Phi\right)\bar\Psi P_{\mathrm{L}}\Psi - \sum_X\Big[\frac{1}{4}\big(c_X^G\,G^A_{\mu\nu}G^{A\mu\nu} + c_X^W\,W^I_{\mu\nu}W^{I\mu\nu} + c_X^B\,B_{\mu\nu}B^{\mu\nu}\big)\,\bar\Psi P_X\Psi \\ & + \frac{d_X}{2}\,B_{\mu\nu}\,\mathcal{D}_X^{\mu\nu} + \sum_f c_X^f\,J_f^\mu\,j_{X\mu}\Big], \qquad \text{(IN-16)}\\ \mathcal{D}_X^{\mu\nu} = & \,\mathrm{i}\,\bar\Psi\,\Sigma^{\mu\nu}P_X\Psi, \qquad j_X^\mu = \mathrm{i}\big(\bar\Psi P_X D^\mu\Psi - (D^\mu\bar\Psi)P_X\Psi\big), \qquad J_f^\mu = \bar\psi_f\,\hat\gamma^\mu\psi_f\end{aligned}$$

Symbols in (IN-16).

- $\mathcal{L}_{\mathrm{P}}$: The Lagrangian of the portals of dirac16complex at the fold. Unit: GeV^4. Status: INVENTED.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: GeV$^{7/2}$. Status: AUTHOR.
- P_{L}: The author's left-handed chiral projector, $P_{\mathrm{L}} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c_X^G: The gluon-kinetic portal coupling of the half X of dirac16complex. Unit: GeV^{-7}. Status: INVENTED.
- G^A_M, G^A_μ, G^A_{MN}: The gluon field of $SU(3)_C$ ($A = 1, \ldots, 8$): in the bulk G^A_M, a field of eight dimensions in its geometric normalisation (the covariant derivative $\partial - \mathrm{i}G$), and on the fold the canonically normalised four-dimensional field $G^A_\mu(x)$ of its massless mode, $g_3 G^A_\mu(x) = G^A_\mu(\sigma = 0, x)$; G^A_{MN} and $G^A_{\mu\nu}$ their field strengths. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- c_X^W: The W-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- W^I_M, W^I_μ, W^I_{MN}: The gauge field of $SU(2)_L$ ($I = 1, 2, 3$): in the bulk W^I_M (geometric normalisation), on the fold the canonically normalised four-dimensional field $W^I_\mu(x)$ of its massless mode, $g_2 W^I_\mu(x) = W^I_\mu(\sigma = 0, x)$; W^I_{MN}, $W^I_{\mu\nu}$ their field strengths; $W^\pm_\mu = (W^1_\mu \mp \mathrm{i}W^2_\mu)/\sqrt{2}$. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- c_X^B: The B-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- B_M, B_μ, B_{MN}: The hypercharge gauge field of $U(1)_Y$: in the bulk B_M (geometric normalisation), on the fold the canonically normalised four-dimensional field $B_\mu(x)$ of its massless mode, $g_1 B_\mu(x) = B_\mu(\sigma = 0, x)$; B_{MN}, $B_{\mu\nu}$ their field strengths. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- P_X: The chiral projector of the half X: $P_X = P_{\mathrm{L}}$ for $X = \mathrm{L}$, P_{R} for $X = \mathrm{R}$. Unit: dimensionless. Status: AUTHOR.

- d_X: The dipole-portal coupling of the half X of dirac16complex with the hypercharge field strength. Unit: GeV^{-5}. Status: INVENTED.
- $\mathcal{D}_X^{\mu\nu}$: The dipole (antisymmetric tensor) density of the half X of dirac16complex at the fold, $\mathcal{D}_X^{\mu\nu} = \mathrm{i}\bar{\Psi}\Sigma^{\mu\nu}P_X\Psi$ (real, DERIVED). Unit: GeV^7. Status: DERIVED.
- f: The label of a kind of Standard-Model multiplet: $f \in \{Q, u, d, \ell, e, \nu\}$ (quark doublets, up-quark singlets, down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.
- c_X^f: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c_X^ℓ and c_X^ν are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.
- J_f^μ: The fermion-number current of the Standard-Model multiplets of kind f, $J_f^\mu = \bar{\psi}_f\hat{\gamma}^\mu\psi_f$ summed over the three generations and the gauge components; with other labels the currents that are the sources of the gauge fields. Unit: GeV^3. Status: STANDARD.
- j_X^μ: The convective current of the half X of dirac16complex, $j_X^\mu = \mathrm{i}(\bar{\Psi}P_X D^\mu\Psi - (D^\mu\bar{\Psi})P_X\Psi)$ (real and joining the half with itself, DERIVED). Unit: GeV^8. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- Σ^{ab}, $\Sigma^{\mu\nu}$: The generators of Spin(4,4) on dirac16complex, $\Sigma^{ab} = \frac{1}{4}[T^a, T^b]$ (the scaled commutators of the author's matrices); $\Sigma^{\mu\nu} = \mathbf{e}_a{}^\mu\mathbf{e}_b{}^\nu\Sigma^{ab}$ along the fold. Unit: dimensionless. Status: AUTHOR.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC}G_\mu^B X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- ψ, $\bar{\psi}$: A Standard-Model fermion field (a four-dimensional Weyl field of the fold with three generations, written as a four-component spinor of fixed chirality); $\bar{\psi} = \psi^\dagger\beta$; ψ_f the fields of the kind f. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- $\hat{\gamma}^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^\mu$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^\mu, \hat{\gamma}^\nu\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_\mu = \hat{h}_{\mu\nu}\hat{\gamma}^\nu$. Unit: dimensionless. Status: STANDARD.

The portals of dirac16complex (INVENTED): its left- and right-handed halves couple at the fold, each with its own couplings, to every field of the Standard Model, and only the left-handed half to the Higgs doublet. The Higgs portal multiplies the scalar density of the left-handed half by $\Phi^\dagger\Phi$; the gauge-kinetic portals multiply the scalar density of each half by the kinetic densities of the gluons, of the W fields and of the B field; the dipole portal couples the antisymmetric tensor of each half to the hypercharge field strength; the current portals couple the convective current of each half to the fermion-number current of each kind of quark and lepton, summed over the generations (flavour blind).

The annihilation and the scattering on nucleons through the portals. The current portals of IN-16 couple the fermion-number current of a kind f of Standard-Model fermions to the convective current of dirac16complex at the fold. Its time component sets the elastic coupling $\mathfrak{g}_f$ of FE-47; between the vacuum and a pair $\chi(\vec{k})\bar{\chi}(-\vec{k})$ the time component vanishes and the spatial one is of the second order in the momentum (FE-67, `um4_krein.py`, KR9), so a pair annihilates into a massless chiral fermion state f with $\mathfrak{g}_f^2\omega_\mathrm{b}^2\mathsf{v}^4/(96\pi)$ (the first relation, for the chirality-blind current, the only one that creates no ghost pair at rest; it creates ghost pairs only at the next order in the momentum, with the relative rate $0.0089\,\vec{k}^2/m_\Psi^2$; the computed pair amplitudes give the factor 1.000 of a Dirac fermion whose current is the convective current of a scalar density, 0.83 and 1.23 for the currents of the left- and right-handed halves alone), with the thermal average $\langle\mathsf{v}^4\rangle = 60/\mathsf{x}^2$; FE-61 has a quarter of it. With this velocity dependence the relic density (S-19 integrated from $\mathsf{x} = 3$) needs $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle = 5.6$ to 6.8×10^{-26} cm^3/s at freeze-out, which lies at $x_\mathrm{f} = 24$ to 33 for $\omega_\mathrm{b} = 10$ GeV to 10^5 GeV (the coefficient of $1/\mathsf{x}^2$ grows from 2.8×10^{-6} to 6.3×10^{-6} GeV^{-2}). The coherent spin-independent cross-section per nucleon σ_SI has the reduced mass μ_red of χ and the nucleon and the nucleon coupling f_N: for quark portals of equal strength $\mathfrak{g}_\mathrm{q}$ (the 30 chiral quark states below the top quark and the 36 above) the vector charge of the nucleon is three times the quark coupling; for lepton portals ($\mathfrak{g}_\mathrm{l}$ for the nine chiral lepton states that take part) the scattering arises at one loop through the photon, with the coupling $\mathfrak{g}_\mathrm{p}$ to the proton ($\alpha_\mathrm{em} = 1/137.036$, the momentum transfer $q_\mathrm{t} = 50$ MeV of a xenon detector, the cut-off of the logarithm taken at ω_b); for the Higgs portal the nucleon (mass $m_\mathrm{N} = 0.939$ GeV) couples through $f_\mathrm{H} = 0.30$. Status: STANDARD (the formulas), DERIVED (the velocity dependence, KR9), ESTIMATE (the loop and the matrix elements).

ω_b (GeV)	10	30	100	300	1000	3000	10^4	3×10^4	10^5
quark portals: σ_SI (cm^2)	1.5×10^{-35}	2.1×10^{-36}	2.2×10^{-37}	2.3×10^{-38}	2.3×10^{-39}	2.7×10^{-40}	2.7×10^{-41}	3.4×10^{-42}	3.4×10^{-43}
lepton portals: σ_SI (cm^2)	3.1×10^{-40}	6.9×10^{-41}	1.1×10^{-41}	2.0×10^{-42}	2.6×10^{-43}	4.0×10^{-44}	5.1×10^{-45}	7.8×10^{-46}	9.6×10^{-47}
limit of the LUX-ZEPLIN experiment (cm^2)	not compared	not compared	5.5×10^{-48}	1.7×10^{-47}	5.5×10^{-47}	1.7×10^{-46}	5.5×10^{-46}	1.7×10^{-45}	5.5×10^{-45}
lepton portals: $\mathfrak{g}_\mathrm{l}\omega_\mathrm{b}^2$	0.025	0.078	0.27	0.87	3.0	9.4	33	106	374
Higgs portal: largest $q_\mathrm{b}/\omega_\mathrm{b}$			0.062	0.047	0.035	0.026	0.019		

The direct-detection test of thermal relics of these masses against the limit of the LUX-ZEPLIN experiment (4.2 tonne-years of xenon, Phys. Rev. Lett. 135, 011802 (2025), arXiv:2410.17036 version 3: 2.2×10^{-48} cm^2 at its minimum, 40 GeV, at 90% confidence; above the minimum the limit is scaled in proportion to the mass, which is conservative). Quark portals that set the relic density are excluded from 100 GeV to 10^5 GeV, by factors from 4×10^{10} to 60; lepton portals give a loop-suppressed scattering that the limit excludes below about 2.3×10^4 GeV and allows above, but there the strength $\mathfrak{g}_\mathrm{l}\omega_\mathrm{b}^2$ of the contact coupling that the relic needs is about 80, beyond the perturbative range (it exceeds 4π from about 4×10^3 GeV on): the current portals cannot set the relic density of a perturbatively coupled χ, at the masses of the strongly bound particles of S-20 as at the lower masses. The Higgs-portal coupling of FE-47, $|\mathfrak{g}_{h\chi}| = 2q_\mathrm{b}^2/(\omega_\mathrm{b}v)$, must stay below 3.1×10^{-3} (100 GeV) to 3.1×10^{-2} (10 TeV), so the binding by the Higgs portal must be weak: $q_\mathrm{b} \lesssim 0.06\,\omega_\mathrm{b}$ (100 GeV) to $0.02\,\omega_\mathrm{b}$ (10 TeV). Status: ESTIMATE (`um4_thermal.py`, T3 and T4).

$$\left.\frac{q_\mathrm{b}}{\omega_\mathrm{b}}\right|_\mathrm{max} \propto \omega_\mathrm{b}^{-1/4}, \qquad H < H_\mathrm{max} = \frac{2F(v_\mathrm{b})\,m_\Psi}{\ln\left(11.6\,p_\mathrm{w}\,m_\Psi\,t_0\right)} \tag{S-22}$$

Symbols in (S-22).

- q_b: The inverse width of a state of dirac16complex at the fold: inside the turning point its profile falls as $\mathrm{e}^{-q_\mathrm{b}|\sigma|}$. Unit: GeV. Status: DERIVED.
- ω_b: The energy at zero momentum of the state of dirac16complex at the fold in the limit $H\to0$ (the bound state of UM-3): the particle χ of UM-4 has the mass $\omega_\mathrm{r} = \omega_\mathrm{b}(1 - H/(2q_\mathrm{b})) + \ldots$, equal to ω_b up to the relative correction $H/(2q_\mathrm{b})$. Unit: GeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.
- H_max: The largest value of the author's constant H for which particles χ bound to the fold by the Higgs portal, with the strongest binding that direct detection allows, live longer than 160 Gyr ($11.6\,t_0$, the bound for dark matter that decays into invisible particles). Unit: GeV. Value: from 4.9×10^{-4} GeV at $\omega_\mathrm{b} = 10^4$ GeV to 2.5 GeV at the Planck mass. Status: ESTIMATE.
- $F(v)$: The tunnelling exponent of the width of the particles χ per unit of m_Ψ/H: the integral of the local decay rate q over the region inside the turning point, divided by m_Ψ/H, as a function of the ratio v of the

energy to the bulk mass. Unit: dimensionless. Value: 0.049 for $\upsilon = 0.866$; $(q_\mathrm{b}/\omega_\mathrm{b})^3/3$ for weak binding. Status: DERIVED.

- υ, υ_b, υ_r: The argument of the tunnelling exponent F, a ratio of an energy to the bulk mass m_Ψ; $\upsilon_\mathrm{b} = \omega_\mathrm{b}/m_\Psi$ that of the particle χ in the limit $H/q_\mathrm{b} \to 0$, $\upsilon_\mathrm{r} = \omega_\mathrm{r}/m_\Psi$ that of the real part of its resonance. Unit: dimensionless. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- p_w: The prefactor of the width of the particles χ, $\Gamma_\chi = p_\mathrm{w} m_\Psi \mathrm{e}^{-2F(\upsilon_\mathrm{b}) m_\Psi/H}$: $p_\mathrm{w} \simeq (2/\mathrm{e})(q_\mathrm{b}/m_\Psi)^2$ to the leading order in H/q_b (FE-86; 0.184 for $q_\mathrm{b} = m_\Psi/2$). Unit: dimensionless. Status: DERIVED.
- t_0: The age of the universe. Unit: time. Value: 13.8 Gyr. Status: STANDARD.

The particles χ bound by the Higgs portal cannot be the dark matter of UM-4. With the binding of the author's specification (the Higgs portal of the left-handed half) the largest $q_\mathrm{b}/\omega_\mathrm{b}$ that the LUX-ZEPLIN limit allows falls like $\omega_\mathrm{b}^{-1/4}$ (the coupling $\mathfrak{g}_{h\chi}$ grows like $q_\mathrm{b}^2/\omega_\mathrm{b}$, the limit on the cross-section like ω_b), and with that binding the lifetime condition S-6 (160 Gyr, with the derived prefactor $p_\mathrm{w} = (2/\mathrm{e})(q_\mathrm{b}/m_\Psi)^2$, valid because $q_\mathrm{b} > 10 H_\mathrm{max}$ in every case) holds only for H below H_max:

ω_b (GeV)	10^4	10^6	10^8	10^{10}	10^{12}	10^{14}	10^{16}	1.2×10^{19}
largest $q_\mathrm{b}/\omega_\mathrm{b}$	0.019	0.0062	0.0019	6.2×10^{-4}	1.9×10^{-4}	6.2×10^{-5}	1.9×10^{-5}	3.3×10^{-6}
H_max (GeV)	4.9×10^{-4}	1.5×10^{-3}	4.7×10^{-3}	0.015	0.045	0.14	0.43	2.5
$\lvert\mathfrak{g}_{h\chi}\rvert$	0.031	0.31	3.1	31	310	3100	3.1×10^4	1.1×10^6

H_max grows only like $\omega_\mathrm{b}^{1/4}$ and stays below 2.5 GeV even at the Planck mass (1.22×10^{19} GeV), far below the bound $H > 2534$ GeV at every mass; above about 10^{10} GeV the Yukawa coupling exceeds 4π, where the formulas of tree level are only formal (the direction of the conclusion is unchanged). Status: ESTIMATE (`um4_thermal.py`, T8: the limit extrapolated in proportion to the mass), OPEN (a binding that escapes direct detection, S-29).

$$\Delta\omega_\mathrm{b} = \mathfrak{g}_G \, \Delta\langle G^2\rangle, \qquad \mathfrak{g}_G = -\frac{q_\mathrm{b}^2}{\omega_\mathrm{b} \, \langle G^2\rangle}, \qquad f_\mathrm{N} = \frac{8 \, m_\mathrm{N} \, f_\mathrm{TG}}{9 \, \langle (\alpha_3/\pi) G^2\rangle} \, \frac{q_\mathrm{b}^2}{\omega_\mathrm{b}} \tag{S-29}$$

Symbols in (S-29).

- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- ω_b: The energy at zero momentum of the state of dirac16complex at the fold in the limit $H \to 0$ (the bound state of UM-3): the particle χ of UM-4 has the mass $\omega_\mathrm{r} = \omega_\mathrm{b}(1 - H/(2q_\mathrm{b})) + \ldots$, equal to ω_b up to the relative correction $H/(2q_\mathrm{b})$. Unit: GeV. Status: DERIVED.
- $\mathfrak{g}_\Phi$, $\mathfrak{g}_G$, $\mathfrak{g}_f$, $\mathfrak{g}_{h\chi}$: The four-dimensional couplings of the particles χ of dirac16complex to the Standard Model: to $\Phi^\dagger\Phi$, to the gluon kinetic density, to the number densities of the fermions of kind f, and the Yukawa coupling of the Higgs boson to χ. Unit: $\mathfrak{g}_\Phi$: GeV^{-1}; $\mathfrak{g}_G$: GeV^{-3}; $\mathfrak{g}_f$: GeV^{-2}; $\mathfrak{g}_{h\chi}$: dimensionless. Status: DERIVED.
- $\langle G^2\rangle$: The mean square of the gluon field strength, $\langle G^A_{\mu\nu} G^{A\mu\nu}\rangle$ (the gluon condensate in the QCD vacuum, a thermal value in the plasma). Unit: GeV^4. Status: STANDARD.
- q_b: The inverse width of a state of dirac16complex at the fold: inside the turning point its profile falls as $\mathrm{e}^{-q_\mathrm{b}|\sigma|}$. Unit: GeV. Status: DERIVED.
- f_N: The effective coupling of χ to a nucleon in the coherent scattering (a vector or scalar contact interaction). Unit: GeV^{-2}. Status: ESTIMATE.
- m_N: The mass of a nucleon. Unit: GeV. Value: 0.939 GeV. Status: STANDARD.
- f_TG: The fraction of the nucleon mass carried by the gluons through the trace anomaly (one minus the fractions carried by the masses of the u, d and s quarks): the matrix element of $(\alpha_3/\pi)G^2$ in a nucleon at rest is $-\frac{8}{9} m_\mathrm{N} f_\mathrm{TG}$. Unit: dimensionless. Value: 0.92. Status: STANDARD.
- α_3: The strong coupling constant, $\alpha_3 = g_3^2/(4\pi)$; $\langle(\alpha_3/\pi)G^2\rangle$ is the gluon condensate of the vacuum, 0.012 GeV^4 (Shifman, Vainshtein and Zakharov). Unit: dimensionless. Status: STANDARD.

- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G^A_M, G^A_μ, G^A_{MN}: The gluon field of $SU(3)_C$ ($A = 1, \ldots, 8$): in the bulk G^A_M, a field of eight dimensions in its geometric normalisation (the covariant derivative $\partial - \mathrm{i}G$), and on the fold the canonically normalised four-dimensional field $G^A_\mu(x)$ of its massless mode, $g_3 G^A_\mu(x) = G^A_\mu(\sigma = 0, x)$; G^A_{MN} and $G^A_{\mu\nu}$ their field strengths. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.

No portal of IN-16 can bind the particles χ without coupling them to nucleons. A binding by the gauge-kinetic portal of the gluons acts through the gluon condensate of the vacuum, $\langle(\alpha_3/\pi)G^2\rangle = 0.012$ GeV^4 (M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B 147 (1979) 385). If it sets the strength of the fold ($\mu_\mathrm{L} = c^G_\mathrm{L}\langle G^2\rangle/(4V_*)$, $\mu_\mathrm{R} = 0$ in FE-39), the theorem of Feynman and Hellmann (FE-47) gives the first relation with the coupling of the second, and inside a nucleon the condensate is shifted: the matrix element of $(\alpha_3/\pi)G^2$ in a nucleon at rest is $-\frac{8}{9} m_\mathrm{N} f_\mathrm{TG}$ (the trace anomaly, M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Phys. Lett. B 78 (1978) 443; $f_\mathrm{TG} = 0.92$), so χ scatters on nucleons with the coupling of the third relation, $64\, q_\mathrm{b}^2/\omega_\mathrm{b}$ GeV^{-3} (the strong coupling cancels). At the lifetime bound of S-6 this gives:

q_b/m_Ψ	0.0995	0.3	0.5	0.7	0.8	0.9	0.95	0.99	0.999
ω_b (GeV)	4.3×10^{8}	1.4×10^{7}	2.5×10^{6}	6.0×10^{5}	2.8×10^{5}	1.1×10^{5}	4.9×10^{4}	1.2×10^{4}	2.2×10^{3}
σ_SI (cm^2)	8.4×10^{-12}	8.9×10^{-13}	3.1×10^{-13}	1.5×10^{-13}	1.1×10^{-13}	9.3×10^{-14}	9.4×10^{-14}	1.5×10^{-13}	5.4×10^{-13}
over the limit of LZ	3.5×10^{29}	1.1×10^{30}	2.3×10^{30}	4.5×10^{30}	7.3×10^{30}	1.6×10^{31}	3.4×10^{31}	2.3×10^{32}	4.5×10^{33}
Higgs portal with the same binding: over the limit	3×10^{7}	1×10^{8}	2×10^{8}	4×10^{8}	6×10^{8}	1×10^{9}	3×10^{9}	2×10^{10}	4×10^{11}

Every binding that gives the particles a lifetime of 160 Gyr is excluded by direct detection if it comes from the gluon condensate or from the Higgs portal. The other mean values through which the portals of IN-16 act vanish in the vacuum: $\langle W^2\rangle$ and $\langle B^2\rangle$ (no condensate of the electroweak fields), the number densities n_f of the current portals and the field strength of the dipole portal. The dark matter of UM-4 therefore needs a binding of χ to the fold that does not couple it to nucleons, which the action of UM-4 does not contain (a strength μ_X of the fold that no field of the Standard Model supplies). Status: ESTIMATE (`um4_thermal.py`, T9: the condensate, the gluonic fraction and the extrapolated limit of LZ), OPEN (the binding).

12.3 The right-handed neutrinos

$$\frac{\mathrm{d}\rho_{\nu_\mathrm{R}}}{\mathrm{d}x_4} + 4H_4\,\rho_{\nu_\mathrm{R}} = \mathcal{C}_{\nu_\mathrm{R}}, \qquad \Delta N_\mathrm{eff} = 3\Big(\frac{10.75}{g_\mathrm{s}(\mathsf{T}_\mathrm{dec})}\Big)^{4/3} \quad \text{(three right-handed neutrinos decoupling at } \mathsf{T}_\mathrm{dec}) \quad \text{(FE-62)}$$

Symbols in (FE-62).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- ρ_{ν_R}: The energy density of the three right-handed neutrinos. Unit: GeV^4. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44\times10^{-42}$ GeV. Status: STANDARD.

- $\mathcal{C}_{\nu_\mathrm{R}}$: The collision term of the right-handed neutrinos: the rate at which collisions produce (or remove) their energy density. Unit: GeV^5. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- N_eff: The effective number of neutrino species: the energy density of the relativistic particles other than photons in units of that of one neutrino species whose temperature is $(4/11)^{1/3}$ times that of the photons. Unit: dimensionless. Value: 3.044 (Standard Model); measured 2.99 ± 0.17 (Planck 2018). Status: STANDARD.
- g_s: The effective number of relativistic degrees of freedom of the entropy density. Unit: dimensionless. Value: 3.909 today. Status: STANDARD.
- T_dec: The temperature at which the right-handed neutrinos decouple from the plasma. Unit: GeV. Status: STANDARD.

$$\frac{n_{\chi,\mathrm{eq}}^2\,\sigma_\mathrm{pair}\mathsf{v}}{16\,n_{\nu_\mathrm{R},\mathrm{eq}}}\Big|_{\mathsf{T}=\mathsf{T}_\mathrm{dec}} = H_4\big(\mathsf{T}_\mathrm{dec}\big), \qquad n_{\nu_\mathrm{R},\mathrm{eq}} = \frac{9\,\zeta(3)}{2\pi^2}\,\mathsf{T}^3 \tag{S-23}$$

Symbols in (S-23).

- $n_{\chi,\mathrm{eq}}$: The number density of χ in thermal equilibrium with the plasma (non-relativistic). Unit: GeV^3. Status: STANDARD.
- σ_pair: The cross-section of the annihilation of one particle χ_a with one antiparticle $\bar\chi_a$ of the same species, summed over the final states; $\sigma_\mathrm{pair}\mathsf{v}$ its product with the relative velocity (a constant for s-wave annihilation). Unit: GeV^{-2}. Status: DERIVED.
- v: The relative velocity of two annihilating particles; $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle$ is the thermal average of the product. Unit: dimensionless. Status: STANDARD.
- $n_{\nu_\mathrm{R},\mathrm{eq}}$: The number density of the three right-handed neutrinos and their antiparticles in thermal equilibrium at the temperature T. Unit: GeV^3. Status: STANDARD.
- T: The temperature of the Standard-Model plasma (of the photons after the annihilation of electrons and positrons). Unit: GeV. Status: STANDARD.
- T_dec: The temperature at which the right-handed neutrinos decouple from the plasma. Unit: GeV. Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- ζ: The Riemann zeta function; $\zeta(3) = 1.2020569$. Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

The right-handed neutrinos. Their production by the neutrino Yukawa couplings is negligible: the rate of their production divided by the expansion rate is 5×10^{-9} at 100 GeV, for the Yukawa coupling 0.05 eV/v of the heaviest neutrino. If the lepton portal includes them ($c_X^\nu \neq 0$), they stay in equilibrium with χ as long as the rate per right-handed neutrino of $\nu_\mathrm{R}\bar\nu_\mathrm{R} \to \chi\bar\chi$ (the left-hand side of S-23, by detailed balance, with the annihilation of S-21 that sets the relic density, a quarter of it into the right-handed neutrinos) exceeds the expansion rate, and decouple at T_dec. Status: STANDARD, ESTIMATE (the case of equal couplings of the twelve chiral lepton states, the right-handed neutrinos included).

ω_b (GeV)	10	30	100	300	1000	3000	10000
$\omega_\mathrm{b}/\mathsf{T}_\mathrm{dec}$	14.5	15.0	15.7	16.3	16.9	17.4	18.1
ΔN_eff (FE-62)	0.239	0.200	0.188	0.183	0.157	0.143	0.141
N_eff	3.28	3.24	3.23	3.23	3.20	3.19	3.18
standard deviations above ACT DR6 (2.89 ± 0.11)	3.6	3.2	3.1	3.1	2.8	2.7	2.7

Thermalised right-handed neutrinos would add $\Delta N_{\mathrm{eff}} = 0.14$ to 0.24 (N_{eff} the effective number of neutrino species), which is 1.1 to 1.7 standard deviations above Planck 2018 ($N_{\mathrm{eff}} = 2.99 \pm 0.17$) and 2.7 to 3.6 above the combination of ACT DR6 with Planck, lensing, DESI BAO and nucleosynthesis (2.89 ± 0.11); for the heavier particles χ of UM-4, which would decouple the right-handed neutrinos above the electroweak scale, the addition is 0.141. UM-4 therefore requires that the right-handed neutrinos never thermalise: their portal couplings must be much smaller than those of the lepton doublets, and then $N_{\mathrm{eff}} = 3.044$ as in the Standard Model. Status: DERIVED (`um4_thermal.py`, T5), ESTIMATE (the rates).

13 The primordial era

The author's vacuum, which the earlier versions of the model took as the primordial era, is not a solution of UM-4 (FE-31). This section solves the two candidates for an inflation of the fold: the exact de Sitter fold, in which the extra times inflate with ordinary space, and the inflation with frozen extra times, which the observed constancy of the couplings requires.

$$g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N = \mathrm{d}\sigma^2 + f_{\mathrm{dS}}^2\Big[\mathrm{e}^{2H_{\mathrm{i}}x_4}\big(\mathrm{d}\vec{x}^2 - \mathrm{d}\vec{x}_5^2\big) - \mathrm{d}x_4^2\Big], \qquad f_{\mathrm{dS}} = \frac{\sinh\big(H(\sigma_0 - |\sigma|)\big)}{\sinh(H\sigma_0)}, \qquad H_{\mathrm{i}} = \frac{H}{\sinh(H\sigma_0)},$$
$$\kappa\big(\lambda_{\mathrm{F}} + \rho_{\mathrm{i}}\big) = 12H\coth(H\sigma_0), \qquad H_{\mathrm{i}}^2 = \Big(\frac{\kappa(\lambda_{\mathrm{F}} + \rho_{\mathrm{i}})}{12}\Big)^2 - H^2 = \frac{\kappa_7\rho_{\mathrm{i}}}{15}\Big(1 + \frac{\rho_{\mathrm{i}}}{2\lambda_{\mathrm{F}}}\Big) \tag{FE-87}$$

Symbols in (FE-87).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\ldots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- f_{dS}: The warp of the de Sitter fold: the factor of the lengths of the fold at the depth σ. Unit: dimensionless. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H_{i}: A constant expansion rate of ordinary space: of the de Sitter fold (whose extra times expand at the same rate) or of an inflation of the fold with frozen extra times. Unit: GeV. Status: DERIVED (FE-87), ASSUMED (FE-88).
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\vec{x}$: The position in ordinary space, $\vec{x} = (x_1, x_2, x_3)$. Unit: length. Status: AUTHOR.
- $\vec{x}_5$: The position along the extra times, $\vec{x}_5 = (x_5, x_6, x_7)$; $\mathrm{d}\vec{x}_5^2 = \mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2$. Unit: length. Status: AUTHOR.
- σ_0: The depth at which the warp of the de Sitter fold vanishes (a horizon of the bulk), $\sinh(H\sigma_0) = H/H_{\mathrm{i}}$. Unit: length. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length).

Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\lambda_{\rm F}$: The tension that the static block requires of the fold, $\kappa\lambda_{\rm F} = 12H$: the total vacuum energy of the seven-dimensional fold, made of the tension of the fold fluid and the vacuum energy of the fields of the fold (both have the pressure minus their energy density along ordinary space and along the extra times), per unit volume of the six other directions of the fold (eight-dimensional normalisation); in the four-dimensional normalisation it is $\lambda_{\rm F}V_*$, and the excess of the total over $\lambda_{\rm F}V_*$ is the dark energy ρ_Λ. Unit: GeV^7. Value: $12H/\kappa$; $\lambda_{\rm F}V_* = 30H^2M_{\rm red}^2 = 1.1\times10^{45}$ GeV^4 for $H = 2534$ GeV. Status: DERIVED.
- $\rho_{\rm i}$: The vacuum energy of the fold above the critical tension $\lambda_{\rm F}$ that expands the de Sitter fold (eight-dimensional normalisation). Unit: GeV^7. Status: DERIVED.
- κ_7: The gravitational coupling of the fold: to the first order in its curvature over H^2 the fold obeys the Einstein equations of seven dimensions $\hat{G}^a{}_b = \kappa_7\mathsf{t}^a{}_b$ with $\kappa_7 = 5\kappa H/2$, the coupling of the massless graviton of the bulk bound to the fold; four-dimensional gravity on the fold has $8\pi G_{\rm N} = \kappa_7/V_*$ (plus the induced term, which is negligible). Unit: GeV^{-5}. Value: $5\kappa H/2$. Status: DERIVED.

The exact de Sitter fold: a fold with a tension above the critical one in a bulk of constant curvature $-H^2$, whose metric is a de Sitter space of seven dimensions expanding at the rate $H_{\rm i}$ in all its space-like and time-like directions; the warp vanishes at the depth σ_0 (a horizon of the bulk), and the expansion rate obeys the Friedmann equation of seven dimensions with the correction of high energies, as for the brane of Randall and Sundrum. The extra times inflate together with ordinary space, so the factor $\mathcal{Z}$ of the couplings of the fold grows like e^{3N} after N e-folds and every coupling of the fold falls like e^{-3N}, by 6.7×10^{-79} after 60 e-folds: it is not the primordial era of our universe. Status: DERIVED (`um4_primordial.py`, PR1 and PR5: the bulk equations hold exactly in all eight directions, and the junction gives the tension and the expansion rate exactly).

$$\mathsf{c}' = 0, \quad a_4' = H_{\rm i}: \qquad \kappa_7\rho_7 = 3H_{\rm i}^2, \qquad P_7 = -\rho_7, \qquad q_7 = -2\rho_7, \qquad \frac{H_{\rm i}}{H} < \frac{1}{10} \tag{FE-88}$$

Symbols in (FE-88).

- c: The logarithm of the scale factor of the extra times on the fold, $\mathsf{c} = \ln(\mathrm{e}^{-a_4}b_4)$; $\mathsf{c}' = \mathsf{V}H$; frozen extra times have $\mathsf{c}' = 0$. Unit: dimensionless. Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $H_{\rm i}$: A constant expansion rate of ordinary space: of the de Sitter fold (whose extra times expand at the same rate) or of an inflation of the fold with frozen extra times. Unit: GeV. Status: DERIVED (FE-87), ASSUMED (FE-88).
- κ_7: The gravitational coupling of the fold: to the first order in its curvature over H^2 the fold obeys the Einstein equations of seven dimensions $\hat{G}^a{}_b = \kappa_7\mathsf{t}^a{}_b$ with $\kappa_7 = 5\kappa H/2$, the coupling of the massless graviton of the bulk bound to the fold; four-dimensional gravity on the fold has $8\pi G_{\rm N} = \kappa_7/V_*$ (plus the induced term, which is negligible). Unit: GeV^{-5}. Value: $5\kappa H/2$. Status: DERIVED.
- ρ_7: The energy density of the matter of the fold above its tension in the seven-dimensional normalisation (minus the component $\mathsf{t}^{(4)}{}_{(4)}$: an energy per unit volume of the six other directions of the fold); $V_*\rho_7$ in the four-dimensional normalisation. Unit: GeV^7. Status: DERIVED.
- P_7: The pressure along ordinary space of the matter of the fold above its tension in the seven-dimensional normalisation (the component $\mathsf{t}^{(i)}{}_{(i)}$). Unit: GeV^7. Status: DERIVED.
- q_7: The pressure along the extra times of the matter of the fold above its tension in the seven-dimensional normalisation (the component $\mathsf{t}^{(r)}{}_{(r)}$). Unit: GeV^7. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\rm F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.

The inflation of the fold with frozen extra times, the primordial era that UM-4 assumes: a vacuum energy of the fold expands ordinary space at the constant rate $H_{\rm i}$ by the Friedmann equation of FE-72, and the radion condition requires the pressure $q_7 = -2\rho_7$ along the extra times, twice that of a vacuum energy: the fold fluid adds $q_{\rm hold} = -\rho_7$ (FE-65). These are the equations of the first order. The exact equations of a fold with this extrinsic curvature in

a bulk of constant curvature have no solution for $H_\mathrm{i} \neq 0$ (Gauss's equation has only the static solutions $\pm H$ in every direction), so the inflation with frozen extra times needs a Weyl tensor of the bulk (for $w = -1$ the correction ρ_7^2 is conserved by itself, so the Weyl tensor can be consistent with it); its bulk solution is not computed. Status: DERIVED (`um4_primordial.py`, PR2), OPEN (the Weyl tensor of the bulk, the origin of the holding pressure).

$$\begin{aligned} r_\mathrm{T} < 0.032: &\qquad H_\mathrm{i} < \pi M_\mathrm{red}\sqrt{\frac{r_\mathrm{T} A_\mathrm{s}}{2}} = 4.43\times 10^{13}\ \mathrm{GeV}, \\ \frac{H_\mathrm{i}}{H} < \frac{1}{10}: &\qquad H > 4.4\times 10^{14}\ \mathrm{GeV}\ \ \text{or}\ \ H_\mathrm{i} < 253\ \mathrm{GeV}\ \ (H = 2534\ \mathrm{GeV}) \end{aligned} \tag{S-24}$$

Symbols in (S-24).

- r_T: The ratio of the amplitudes of the primordial spectra of gravitational waves (tensor) and of curvature (scalar) at the pivot scale. Unit: dimensionless. Value: below 0.032 (Planck PR4 with BICEP/Keck 2018 and BAO, M. Tristram et al. 2022, 95 per cent). Status: STANDARD.
- H_i: A constant expansion rate of ordinary space: of the de Sitter fold (whose extra times expand at the same rate) or of an inflation of the fold with frozen extra times. Unit: GeV. Status: DERIVED (FE-87), ASSUMED (FE-88).
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- M_red: The reduced Planck mass, $M_\mathrm{red} = (8\pi G_\mathrm{N})^{-1/2}$. Unit: GeV. Value: 2.435×10^{18} GeV. Status: STANDARD.
- A_s: The amplitude of the primordial curvature spectrum at the pivot k_*. Unit: dimensionless. Value: 2.10×10^{-9} (Planck 2018). Status: ASSUMED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H > 2534$ GeV, about 2.5 TeV (from the invisible width of the Z boson). Status: AUTHOR.

The scales of the inflation. The bound $r_\mathrm{T} < 0.032$ on the ratio of the tensor to the scalar perturbations (Planck PR4 with BICEP/Keck 2018 and BAO, M. Tristram et al. 2022, 95 per cent) with the measured amplitude $A_\mathrm{s} = 2.1\times 10^{-9}$ of the scalar perturbations limits any inflation to $H_\mathrm{i} < 4.4\times 10^{13}$ GeV. The equations of the fold are those of general relativity only for $(12/25)(H_\mathrm{i}/H)^2 \ll 1$ (S-1), that is $H_\mathrm{i} < H/10$: an inflation at the tensor bound needs $H > 4.4\times 10^{14}$ GeV, and with H at its lower bound of 2534 GeV only an inflation with $H_\mathrm{i} < 253$ GeV (an energy scale below 3.3×10^{10} GeV) is in that regime. Status: DERIVED (`um4_primordial.py`, PR3), STANDARD (the measurements).

$$\xi_\mathrm{H} = N_\mathrm{e}\sqrt{\frac{\lambda}{72\pi^2 A_\mathrm{s}}}, \qquad n_\mathrm{S} = 1 - \frac{2}{N_\mathrm{e}}, \qquad r_\mathrm{T} = \frac{12}{N_\mathrm{e}^2}, \qquad H_\mathrm{i} = \sqrt{\frac{\lambda}{12}}\,\frac{M_\mathrm{red}}{\xi_\mathrm{H}} \tag{S-25}$$

Symbols in (S-25).

- ξ_H: The coupling of the Higgs doublet to the scalar curvature of the fold, the term $-\xi_\mathrm{H}\hat{R}\,\Phi^\dagger\Phi$ in the Lagrangian of the fold: an extension of UM-4, whose Higgs Lagrangian has no such term ($\xi_\mathrm{H} = 0$). Unit: dimensionless. Value: 0 in UM-4; about 1.6×10^4 if the Higgs field drove inflation (ESTIMATE, an extension of the action). Status: ESTIMATE (an extension of the action).
- N_e: The number of e-foldings of inflation after the pivot scale left the horizon. Unit: dimensionless. Value: 55. Status: STANDARD.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- A_s: The amplitude of the primordial curvature spectrum at the pivot k_*. Unit: dimensionless. Value: 2.10×10^{-9} (Planck 2018). Status: ASSUMED.
- n_S: The spectral index of the primordial curvature spectrum. Unit: dimensionless. Value: 0.965 (Planck 2018). Status: ASSUMED.
- r_T: The ratio of the amplitudes of the primordial spectra of gravitational waves (tensor) and of curvature (scalar) at the pivot scale. Unit: dimensionless. Value: below 0.032 (Planck PR4 with BICEP/Keck 2018 and BAO, M. Tristram et al. 2022, 95 per cent). Status: STANDARD.

- H_i: A constant expansion rate of ordinary space: of the de Sitter fold (whose extra times expand at the same rate) or of an inflation of the fold with frozen extra times. Unit: GeV. Status: DERIVED (FE-87), ASSUMED (FE-88).
- M_red: The reduced Planck mass, $M_\mathrm{red} = (8\pi G_\mathrm{N})^{-1/2}$. Unit: GeV. Value: 2.435×10^{18} GeV. Status: STANDARD.

Higgs inflation, an extension of the action. If the Higgs doublet had a coupling ξ_H to the curvature of the fold, which the Higgs Lagrangian of UM-4 does not contain, the Higgs field could drive the inflation of the fold with N_e e-folds after the pivot scale left the horizon: with $N_\mathrm{e} = 55$, $\xi_\mathrm{H} = 16195$ ($\lambda = 0.1294$) or 4502 ($\lambda = 0.01$, the running value near the scale of inflation), $n_\mathrm{S} = 0.9636$ and $r_\mathrm{T} = 0.0040$ (0.3 standard deviations below Planck 2018, 0.9649 ± 0.0042, and 3.5 below ACT DR6 combined with Planck, CMB lensing and DESI, 0.974 ± 0.003), and $H_\mathrm{i} = 1.6 \times 10^{13}$ GeV, so that UM-4 would need $H > 1.6 \times 10^{14}$ GeV for it. UM-4 does not specify its inflaton. Status: ESTIMATE (`um4_primordial.py`, PR4), OPEN (the inflaton).

14 The CMB, the CNB and the BAO

The CMB, the CNB and the BAO are computed with the metric of the fold as the expanding background with frozen extra times and its cosmological perturbations (of the order 10^{-5}); the terms of the second order in the perturbations are dropped. With the volume of the extra times held fixed (FE-65) the fold obeys the Friedmann equations of general relativity with Newton's constant $8\pi G_\mathrm{N} = \kappa_7 / V_*$ (FE-59), up to the correction of high energies of the relative size $(12/25)(H_4/H)^2$ (S-1), below 10^{-80} at recombination for $H > 2534$ GeV: the bulk adds no correction to the expansion at any observable level. The content of the fold is that of UM-4: the particles χ of dirac16complex as cold dark matter (pressureless in the dilute limit, as the Kohn-Sham equation of state shows; their abundance is taken from the measurement, because UM-4 does not compute it: only strongly bound particles could be a thermal relic, S-20, and the portals cannot bind them so, S-29), the vacuum energy ρ_Λ of the fold as dark energy (with the equation of state $w = -1$, w the ratio of its pressure to its energy density, S-4), the baryons and photons of the Standard Model, and three Dirac neutrinos of the normal ordering with the lightest mass zero ($m_1 = 0$, $m_2 = \sqrt{\Delta m^2_{21}} = 8.61$ meV, $m_3 = \sqrt{\Delta m^2_{31}} = 50.1$ meV, $\sum_i m_i = 0.0587$ eV; ASSUMED: the lightest mass is not fixed by any measurement; DESI DR2 with the CMB and NuFIT 6.0 bounds it below 23 meV, S-18) whose right-handed states are not populated ($N_\mathrm{eff} = 3.044$). The primordial spectrum, the optical depth and the densities are the Planck 2018 values (TT,TE,EE+lowE+lensing: $\Omega_\mathrm{b} h^2_{100} = 0.02237$, $\Omega_\chi h^2_{100} = 0.1200$, $H_4^{(0)} = 67.36$ km/s/Mpc, the optical depth $\tau_\mathrm{reion} = 0.0544$ of the reionised universe to the CMB photons, $\ln(10^{10} A_\mathrm{s}) = 3.044$, $n_\mathrm{S} = 0.9649$; ASSUMED: UM-4 does not compute its primordial era to this precision). The particles χ do not drag the baryons: their number density, $\rho_\chi / \omega_\mathrm{b}$ with ω_b above 2.2×10^3 GeV for every binding (S-6), is tiny, and their momentum exchange with the baryons at the direct-detection limits is far below the expansion rate at recombination. Status: ESTIMATE. The Boltzmann hierarchy and the line-of-sight integrals are solved by the code CAMB (Lewis, Challinor and Lasenby 2000); the background and the BAO ruler are computed a second time here, independently.

$$\begin{aligned} &\Big(\partial_{x_4} - a_4' \, |\vec{p}| \, \frac{\partial}{\partial |\vec{p}|}\Big) \varrho_{\vec{p}} = -\mathrm{i}\big[\mathsf{H}_{\vec{p}},\ \varrho_{\vec{p}}\big] + \mathcal{C}_{\vec{p}}, \\ &N_\mathrm{eff} = \frac{8}{7}\Big(\frac{11}{4}\Big)^{4/3} \frac{\rho_\nu}{\rho_\gamma}, \qquad \mathsf{T}_\nu^{(0)} = \Big(\frac{4}{11}\Big)^{1/3} \mathsf{T}_\gamma^{(0)}, \qquad n_\nu^{(0)} = \frac{3\zeta(3)}{2\pi^2} \big(\mathsf{T}_\nu^{(0)}\big)^3 \end{aligned} \tag{FE-53}$$

Symbols in (FE-53).

- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\vec{p}$: The momentum of a neutrino (three components along ordinary space, in the rest frame of the medium); $|\vec{p}| = E$ for a neutrino of negligible mass. Unit: GeV. Status: STANDARD.
- $\varrho_{\vec{p}}$, $\bar{\varrho}_{\vec{p}}$: The 3×3 flavour density matrices of the neutrinos and of the antineutrinos of momentum $\vec{p}$ in the flavour basis (electron, muon, tau): the diagonal entries are the occupation numbers of the three flavours, the

off-diagonal entries the coherence between the flavours; $\bar{\varrho}$ is defined with the flavour indices in the reversed order, so that it obeys an equation of the same form. Unit: dimensionless. Status: STANDARD.

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\mathsf{H}_{\vec{p}}$, $\bar{\mathsf{H}}_{\vec{p}}$: The flavour Hamiltonians (3×3 Hermitian matrices) of the neutrinos and of the antineutrinos of momentum $\vec{p}$. Unit: GeV. Status: STANDARD.
- $\mathcal{C}_{\vec{p}}$, $\bar{\mathcal{C}}_{\vec{p}}$: The collision terms ($3 \times 3$ matrices) of the neutrinos and the antineutrinos of momentum $\vec{p}$: the rates of change of $\varrho_{\vec{p}}$ and $\bar{\varrho}_{\vec{p}}$ by emission, absorption, scattering and pair processes with the matter. Unit: GeV. Status: STANDARD.
- N_{eff}: The effective number of neutrino species: the energy density of the relativistic particles other than photons in units of that of one neutrino species whose temperature is $(4/11)^{1/3}$ times that of the photons. Unit: dimensionless. Value: 3.044 (Standard Model); measured 2.99 ± 0.17 (Planck 2018). Status: STANDARD.
- ρ_ν: The energy density of the neutrinos and antineutrinos of all flavours. Unit: GeV^4. Status: STANDARD.
- ρ_γ: The energy density of the photons. Unit: GeV^4. Status: STANDARD.
- T_ν: The temperature of the cosmic neutrino background; $\mathsf{T}_\nu^{(0)}$ its value today. Unit: GeV (or K). Value: 1.945 K today. Status: STANDARD.
- T_γ: The temperature of the photons (of the CMB); $\mathsf{T}_\gamma^{(0)}$ its value today. Unit: GeV (or K). Value: 2.7255 K today. Status: STANDARD.
- n_ν: The number density of the cosmic neutrino background per flavour (neutrinos plus antineutrinos, one helicity each); $n_\nu^{(0)}$ its value today. Unit: GeV^3. Value: 112 cm^{-3} today. Status: STANDARD.
- ζ: The Riemann zeta function; $\zeta(3) = 1.2020569$. Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

$$
\begin{aligned}
&\hat{h}_{\mu\nu}\,\mathrm{d}x^\mu \mathrm{d}x^\nu = a^2\Big[-(1+2\psi_\mathrm{B})\,\mathrm{d}\tau_\mathrm{c}^2 + (1-2\phi_\mathrm{B})\big(\mathrm{d}x_1^2+\mathrm{d}x_2^2+\mathrm{d}x_3^2\big)\Big],\\
&a = \mathrm{e}^{a_4}, \qquad \mathrm{d}\tau_\mathrm{c} = \frac{\mathrm{d}x_4}{a}, \qquad \mathcal{H} = \frac{\dot{a}}{a},\\
&k^2\phi_\mathrm{B} + 3\mathcal{H}\big(\dot{\phi}_\mathrm{B} + \mathcal{H}\psi_\mathrm{B}\big) = -4\pi G_\mathrm{N} a^2\,\Delta\rho, \qquad k^2\big(\dot{\phi}_\mathrm{B} + \mathcal{H}\psi_\mathrm{B}\big) = 4\pi G_\mathrm{N} a^2(\rho+\mathcal{P})\,\vartheta,\\
&\ddot{\phi}_\mathrm{B} + \mathcal{H}\big(\dot{\psi}_\mathrm{B} + 2\dot{\phi}_\mathrm{B}\big) + \Big(2\frac{\ddot{a}}{a} - \mathcal{H}^2\Big)\psi_\mathrm{B} + \frac{k^2}{3}\big(\phi_\mathrm{B} - \psi_\mathrm{B}\big) = 4\pi G_\mathrm{N} a^2\,\Delta\mathcal{P},\\
&k^2\big(\phi_\mathrm{B} - \psi_\mathrm{B}\big) = 12\pi G_\mathrm{N} a^2(\rho+\mathcal{P})\,\Sigma_\mathrm{s}
\end{aligned}
\tag{FE-54}
$$

Symbols in (FE-54).

- $\hat{h}_{\mu\nu}$: The four-dimensional metric of the fold, $\hat{h}_{\mu\nu} = \mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the coordinates $x_1, \ldots, x_4$ (the induced metric at $\sigma = 0$ without the extra times); $\hat{h}^{\mu\nu}$ its inverse, $\hat{h}$ its determinant, $\sqrt{-\hat{h}} = \mathrm{e}^{3a_4}$; $\hat{h}^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- x: The coordinates of the fold: x_1, x_2, x_3 of ordinary space, x_4 the time, x_5, x_6, x_7 the extra times; a field of the fold is a function of the four observable coordinates $x = (x_1, x_2, x_3, x_4)$, and $\mathrm{d}^4x = \mathrm{d}x_1\,\mathrm{d}x_2\,\mathrm{d}x_3\,\mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.
- a, $\dot{a}$, $\ddot{a}$: The scale factor of the fold, $a = \mathrm{e}^{a_4}$ (normalised to 1 today by the choice of the comoving coordinates x_1, x_2, x_3), and its first and second derivatives with respect to the conformal time τ_c. Unit: dimensionless. Status: STANDARD.
- ψ_B, $\dot{\psi}_\mathrm{B}$: The Bardeen potential of the time-time part of the perturbed metric of the fold (the Newtonian potential) in the conformal Newtonian gauge, a function of τ_c and of the wave number k, and its derivative with respect to τ_c. Unit: dimensionless. Status: STANDARD.
- τ_c: The conformal time of the fold, $\mathrm{d}\tau_\mathrm{c} = \mathrm{e}^{-a_4}\mathrm{d}x_4$ (the time divided by the scale factor e^{a_4} of the fold); a dot on a quantity of the cosmological perturbations is $\mathrm{d}/\mathrm{d}\tau_\mathrm{c}$. Unit: length (= time). Status: STANDARD.
- ϕ_B, $\dot{\phi}_\mathrm{B}$, $\ddot{\phi}_\mathrm{B}$: The Bardeen potential of the space-space part of the perturbed metric of the fold (the curvature potential) in the conformal Newtonian gauge, and its first and second derivatives with respect to τ_c. Unit:

dimensionless. Status: STANDARD.

- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\mathcal{H}$: The conformal expansion rate, $\mathcal{H} = \dot{a}/a = a\,a_4'$. Unit: 1/length. Status: STANDARD.
- k: A wave number: the momentum of a plane wave along x_3, $\mathrm{e}^{\mathrm{i}(kx_3 - \omega x_4)}$; in the cosmological perturbations the comoving wave number of a Fourier mode, $k = |\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_N: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_\mathrm{N} = 1/M_\mathrm{Pl}^2$ with the Planck mass $M_\mathrm{Pl} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- ρ: The total energy density of the matter of the fold (all species); $\Delta\rho$ its perturbation. Unit: GeV^4. Status: STANDARD.
- $\mathcal{P}$: The total pressure of the matter of the fold; $\Delta\mathcal{P}$ its perturbation. Unit: GeV^4. Status: STANDARD.
- ϑ: The divergence of the velocity of the total matter: $(\rho + \mathcal{P})\vartheta$ is the sum over the species of their energy density plus pressure times the divergence of their velocity. Unit: 1/length. Status: STANDARD.
- Σ_s: The anisotropic stress (shear) of the total matter: $(\rho + \mathcal{P})\Sigma_\mathrm{s}$ is the sum over the species (photons and neutrinos) of their energy density plus pressure times their shear. Unit: dimensionless. Status: STANDARD.

$$
\begin{aligned}
&\dot{\delta}_\chi = -\vartheta_\chi + 3\dot{\phi}_\mathrm{B}, \qquad \dot{\vartheta}_\chi = -\mathcal{H}\vartheta_\chi + k^2\psi_\mathrm{B} + \Gamma_{\chi\mathrm{b}}(\vartheta_\mathrm{b} - \vartheta_\chi),\\
&\dot{\delta}_\mathrm{b} = -\vartheta_\mathrm{b} + 3\dot{\phi}_\mathrm{B}, \qquad \dot{\vartheta}_\mathrm{b} = -\mathcal{H}\vartheta_\mathrm{b} + c_\mathrm{s}^2 k^2\delta_\mathrm{b} + k^2\psi_\mathrm{B} + \frac{a\,x_\mathrm{e} n_e \sigma_\mathrm{T}}{\mathsf{R}}(\vartheta_\gamma - \vartheta_\mathrm{b}) + \frac{\rho_\chi}{\rho_\mathrm{b}}\Gamma_{\chi\mathrm{b}}(\vartheta_\chi - \vartheta_\mathrm{b}), \qquad \mathsf{R} = \frac{3\rho_\mathrm{b}}{4\rho_\gamma},\\
&\dot{\delta}_\gamma = -\frac{4}{3}\vartheta_\gamma + 4\dot{\phi}_\mathrm{B}, \qquad \dot{\vartheta}_\gamma = k^2\Big(\frac{\delta_\gamma}{4} - \Sigma_\gamma\Big) + k^2\psi_\mathrm{B} + a\,x_\mathrm{e} n_e \sigma_\mathrm{T}(\vartheta_\mathrm{b} - \vartheta_\gamma),\\
&\dot{\mathsf{F}}_l = \frac{k}{2l+1}\big(l\,\mathsf{F}_{l-1} - (l+1)\,\mathsf{F}_{l+1}\big) - a\,x_\mathrm{e} n_e \sigma_\mathrm{T}\Big(\mathsf{F}_l - \frac{\mathsf{G}_0 + \mathsf{F}_2 + \mathsf{G}_2}{10}\,\delta_{l2}\Big) \quad (l \geq 2),\\
&\dot{\mathsf{G}}_l = \frac{k}{2l+1}\big(l\,\mathsf{G}_{l-1} - (l+1)\,\mathsf{G}_{l+1}\big) - a\,x_\mathrm{e} n_e \sigma_\mathrm{T}\Big(\mathsf{G}_l - \frac{\mathsf{G}_0 + \mathsf{F}_2 + \mathsf{G}_2}{2}\Big(\delta_{l0} + \frac{\delta_{l2}}{5}\Big)\Big) \quad (l \geq 0),\\
&\mathsf{F}_0 = \delta_\gamma, \qquad \mathsf{F}_1 = \frac{4\vartheta_\gamma}{3k}, \qquad \mathsf{F}_2 = 2\Sigma_\gamma,\\
&\dot{\delta}_\nu = -\frac{4}{3}\vartheta_\nu + 4\dot{\phi}_\mathrm{B}, \qquad \dot{\vartheta}_\nu = k^2\Big(\frac{\delta_\nu}{4} - \Sigma_\nu\Big) + k^2\psi_\mathrm{B}, \qquad \dot{\mathsf{N}}_l = \frac{k}{2l+1}\big(l\,\mathsf{N}_{l-1} - (l+1)\,\mathsf{N}_{l+1}\big) \quad (l \geq 2),\\
&\mathsf{N}_1 = \frac{4\vartheta_\nu}{3k}, \qquad \mathsf{N}_2 = 2\Sigma_\nu,\\
&\dot{\Upsilon}_0 = -\frac{\mathsf{q}k}{\mathsf{E}}\Upsilon_1 - \dot{\phi}_\mathrm{B}\frac{\mathrm{d}\ln\mathcal{F}_0}{\mathrm{d}\ln\mathsf{q}}, \qquad \dot{\Upsilon}_1 = \frac{\mathsf{q}k}{3\mathsf{E}}(\Upsilon_0 - 2\Upsilon_2) - \frac{\mathsf{E}k}{3\mathsf{q}}\,\psi_\mathrm{B}\frac{\mathrm{d}\ln\mathcal{F}_0}{\mathrm{d}\ln\mathsf{q}},\\
&\dot{\Upsilon}_l = \frac{\mathsf{q}k}{(2l+1)\mathsf{E}}\big(l\,\Upsilon_{l-1} - (l+1)\,\Upsilon_{l+1}\big) \quad (l \geq 2), \qquad \mathsf{E} = \sqrt{\mathsf{q}^2 + a^2 m_i^2}, \qquad \mathcal{F}_0 = \frac{1}{\mathrm{e}^{\mathsf{q}/\mathsf{T}_\nu^{(0)}} + 1}
\end{aligned}
\tag{FE-55}
$$

Symbols in (FE-55).

- δ_χ, $\dot{\delta}_\chi$: The density contrast of the dark matter (the particles χ of dirac16complex), the perturbation of its energy density divided by its mean, and its derivative with respect to τ_c. Unit: dimensionless. Status: STANDARD.
- ϑ_χ, $\dot{\vartheta}_\chi$: The divergence of the velocity of the dark matter (for a Fourier mode, ik times its velocity along the wave vector), and its derivative. Unit: 1/length. Status: STANDARD.
- ϕ_B, $\dot{\phi}_\mathrm{B}$, $\ddot{\phi}_\mathrm{B}$: The Bardeen potential of the space-space part of the perturbed metric of the fold (the curvature

potential) in the conformal Newtonian gauge, and its first and second derivatives with respect to τ_c. Unit: dimensionless. Status: STANDARD.

- $\mathcal{H}$: The conformal expansion rate, $\mathcal{H} = \dot{a}/a = a\,a'_4$. Unit: 1/length. Status: STANDARD.
- k: A wave number: the momentum of a plane wave along x_3, $\mathrm{e}^{\mathrm{i}(kx_3-\omega x_4)}$; in the cosmological perturbations the comoving wave number of a Fourier mode, $k = |\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.
- ψ_B, $\dot{\psi}_\mathrm{B}$: The Bardeen potential of the time-time part of the perturbed metric of the fold (the Newtonian potential) in the conformal Newtonian gauge, a function of τ_c and of the wave number k, and its derivative with respect to τ_c. Unit: dimensionless. Status: STANDARD.
- $\Gamma_{\chi\mathrm{b}}$: The rate (per conformal time) at which the dark matter exchanges momentum with the baryons through the portal couplings (elastic scattering of χ on nuclei and electrons). Unit: 1/length. Value: negligible for the couplings that direct detection allows. Status: DERIVED.
- ϑ_b, $\dot{\vartheta}_\mathrm{b}$: The divergence of the velocity of the baryons and its derivative. Unit: 1/length. Status: STANDARD.
- δ_b, $\dot{\delta}_\mathrm{b}$: The density contrast of the baryons and its derivative. Unit: dimensionless. Status: STANDARD.
- c_s: The sound speed of the baryon gas: its square is the temperature of the gas divided by the mean mass per particle, times one minus a third of the logarithmic derivative of that temperature with respect to a. Unit: dimensionless. Status: STANDARD.
- a, $\dot{a}$, $\ddot{a}$: The scale factor of the fold, $a = \mathrm{e}^{a_4}$ (normalised to 1 today by the choice of the comoving coordinates x_1, x_2, x_3), and its first and second derivatives with respect to the conformal time τ_c. Unit: dimensionless. Status: STANDARD.
- x_e: The ionisation fraction: the fraction of the electrons that are free. Unit: dimensionless. Status: STANDARD.
- n_e: The number density of electrons (bound and free; in a supernova all are free, in the recombining universe $x_\mathrm{e} n_e$ is the density of the free electrons). Unit: GeV^3 (cm^{-3}). Status: STANDARD.
- σ_T: The Thomson cross-section of the scattering of photons on free electrons. Unit: GeV^{-2}. Value: $6.6524587 \times 10^{-25}$ cm^2. Status: STANDARD.
- R: The ratio of the inertia of the baryons to that of the photons, $\mathsf{R} = 3\rho_\mathrm{b}/(4\rho_\gamma)$. Unit: dimensionless. Status: STANDARD.
- ϑ_γ, $\dot{\vartheta}_\gamma$: The divergence of the velocity of the photons and its derivative. Unit: 1/length. Status: STANDARD.
- ρ_χ: The energy density of the dark matter, the particles χ of dirac16complex; $\rho_\chi^{(0)}$ its value today. Unit: GeV^4. Status: DERIVED.
- ρ_b: The (rest-mass) energy density of the baryons: in a supernova the local density of the matter (quoted in $\mathrm{g/cm}^3$), in cosmology the mean density of the baryons ($\rho_\mathrm{b}^{(0)}$ today). Unit: GeV^4. Status: STANDARD.
- ρ_γ: The energy density of the photons. Unit: GeV^4. Status: STANDARD.
- δ_γ, $\dot{\delta}_\gamma$: The density contrast of the photons and its derivative. Unit: dimensionless. Status: STANDARD.
- Σ_γ: The anisotropic stress (shear) of the photons. Unit: dimensionless. Status: STANDARD.
- F_l, $\dot{\mathsf{F}}_l$: The multipoles of the temperature perturbation of the photons ($\mathsf{F}_0 = \delta_\gamma$, $\mathsf{F}_1 = 4\vartheta_\gamma/(3k)$, $\mathsf{F}_2 = 2\Sigma_\gamma$; the photon multipoles of Ma and Bertschinger) and their derivatives with respect to τ_c. Unit: dimensionless. Status: STANDARD.
- l: The multipole index and number ($l = 0, 1, 2, \ldots$) of an expansion in Legendre polynomials of the direction of a photon or a neutrino relative to the wave vector, or of the CMB sky in spherical harmonics. Status: STANDARD.
- G_l, $\dot{\mathsf{G}}_l$: The multipoles of the polarisation perturbation of the photons (the $G_{\gamma l}$ of Ma and Bertschinger) and their derivatives. Unit: dimensionless. Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta_N^M = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- δ_ν, $\dot{\delta}_\nu$: The density contrast of the relativistic neutrinos and its derivative. Unit: dimensionless. Status: STANDARD.
- ϑ_ν, $\dot{\vartheta}_\nu$: The divergence of the velocity of the relativistic neutrinos and its derivative. Unit: 1/length. Status: STANDARD.
- Σ_ν: The anisotropic stress (shear) of the relativistic neutrinos. Unit: dimensionless. Status: STANDARD.
- N_l, $\dot{\mathsf{N}}_l$: The multipoles of the perturbation of the relativistic neutrinos ($\mathsf{N}_0 = \delta_\nu$, $\mathsf{N}_1 = 4\vartheta_\nu/(3k)$, $\mathsf{N}_2 = 2\Sigma_\nu$) and their derivatives. Unit: dimensionless. Status: STANDARD.
- $\Upsilon_l(\mathsf{q})$, $\dot{\Upsilon}_l$: The multipoles of the relative perturbation of the distribution of a massive neutrino state at the comoving momentum q (the Ψ_l of Ma and Bertschinger) and their derivatives. Unit: dimensionless. Status:

STANDARD.

- q: The comoving momentum of a neutrino, $a|\vec{p}|$. Unit: GeV. Status: STANDARD.
- E: The comoving energy of a massive neutrino (a times its energy). Unit: GeV. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $\mathcal{F}_0(\mathsf{q})$: The unperturbed (Fermi-Dirac) occupation number of the cosmic neutrino background at the comoving momentum q. Unit: dimensionless. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m^2_{ij} = m^2_i - m^2_j$ the squared-mass difference of the states i and j (Δm^2_{21}, Δm^2_{31}, Δm^2_{32}); Δm^2 without labels the difference of the two states of a resonance, $m^2_2 - m^2_1$, multiplied by v^2_{loc}/v^2 where this differs from 1; m without a label the mass of a mode of a field of the bulk as the fold sees it, $m^2 = \omega^2 - k^2$; for the W and Z bosons their complex pole mass, $m^2 = m^2_0 - \mathrm{i}m_0\Gamma_{\mathrm{bulk}}$ to the leading order; in the relic abundance of the particles χ their mass. Unit: GeV. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal), and in a tensor of ordinary space such as h_{ij} one of the directions x_1, x_2, x_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- T_ν: The temperature of the cosmic neutrino background; $\mathsf{T}^{(0)}_\nu$ its value today. Unit: GeV (or K). Value: 1.945 K today. Status: STANDARD.

$$
\begin{aligned}
r_{\mathrm{s}} &= \int_0^{\tau_{\mathrm{d}}} c_{\gamma\mathrm{b}}\,\mathrm{d}\tau_{\mathrm{c}}, \qquad c_{\gamma\mathrm{b}} = \frac{1}{\sqrt{3(1+\mathsf{R})}}, \\
D_{\mathrm{M}}(\mathsf{z}) &= \int_0^{\mathsf{z}} \frac{\mathrm{d}\mathsf{z}'}{H_4(\mathsf{z}')}, \qquad D_{\mathrm{H}}(\mathsf{z}) = \frac{1}{H_4(\mathsf{z})}, \qquad D_{\mathrm{V}}(\mathsf{z}) = \left(\mathsf{z}\, D^2_{\mathrm{M}}(\mathsf{z})\, D_{\mathrm{H}}(\mathsf{z})\right)^{1/3}
\end{aligned}
\tag{FE-56}
$$

Symbols in (FE-56).

- r_{s}: The comoving sound horizon at the drag epoch: the comoving distance that a sound wave of the photon-baryon fluid travels until the baryons are released from the photons. Unit: length. Value: about 147 Mpc (Planck 2018). Status: STANDARD.
- τ_{d}: The conformal time of the drag epoch, when the baryons are released from the photons (redshift about 1060). Unit: length (= time). Status: STANDARD.
- $c_{\gamma\mathrm{b}}$: The sound speed of the tightly coupled photon-baryon fluid. Unit: dimensionless. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- τ_{c}: The conformal time of the fold, $\mathrm{d}\tau_{\mathrm{c}} = \mathrm{e}^{-a_4}\mathrm{d}x_4$ (the time divided by the scale factor e^{a_4} of the fold); a dot on a quantity of the cosmological perturbations is $\mathrm{d}/\mathrm{d}\tau_{\mathrm{c}}$. Unit: length (= time). Status: STANDARD.
- R: The ratio of the inertia of the baryons to that of the photons, $\mathsf{R} = 3\rho_{\mathrm{b}}/(4\rho_\gamma)$. Unit: dimensionless. Status: STANDARD.
- $D_{\mathrm{M}}(\mathsf{z})$: The transverse comoving distance (comoving angular-diameter distance) to the redshift z, for a spatially flat fold. Unit: length. Status: STANDARD.
- z: The redshift, $1 + \mathsf{z} = 1/a$; z' an integration variable. Unit: dimensionless. Status: STANDARD.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a'_4 = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a'_4$, as a function of the time or of the redshift; $H^{(0)}_4$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H^{(0)}_4 = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- $D_{\mathrm{H}}(\mathsf{z})$: The Hubble distance at the redshift z. Unit: length. Status: STANDARD.
- $D_{\mathrm{V}}(\mathsf{z})$: The volume-averaged distance to the redshift z. Unit: length. Status: STANDARD.

$$
\begin{aligned}
C_l^{\mathrm{TT}} &= 4\pi \int \frac{\mathrm{d}k}{k}\, \mathcal{P}_{\mathcal{R}}(k)\, \big(\mathcal{I}_l^{\mathrm{T}}(k)\big)^2, \qquad C_l^{\mathrm{TE}} = 4\pi \int \frac{\mathrm{d}k}{k}\, \mathcal{P}_{\mathcal{R}}(k)\, \mathcal{I}_l^{\mathrm{T}}(k)\, \mathcal{I}_l^{\mathrm{E}}(k), \\
C_l^{\mathrm{EE}} &= 4\pi \int \frac{\mathrm{d}k}{k}\, \mathcal{P}_{\mathcal{R}}(k)\, \big(\mathcal{I}_l^{\mathrm{E}}(k)\big)^2, \qquad \mathcal{P}_{\mathcal{R}}(k) = A_{\mathrm{s}} \Big(\frac{k}{k_*}\Big)^{n_{\mathrm{S}}-1}
\end{aligned}
\tag{FE-57}
$$

Symbols in (FE-57).

- C_l^{TT}, C_l^{TE}, C_l^{EE}: The angular power spectra of the CMB at the multipole l: temperature with temperature, temperature with E-mode polarisation, E-mode polarisation with itself. Unit: dimensionless. Status: STANDARD.
- l: The multipole index and number ($l = 0, 1, 2, \ldots$) of an expansion in Legendre polynomials of the direction of a photon or a neutrino relative to the wave vector, or of the CMB sky in spherical harmonics. Status: STANDARD.
- T, E: The labels of the CMB observables: T the temperature anisotropy, E the E-mode (gradient) polarisation; TT, TE, EE label the spectra of their products. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- k: A wave number: the momentum of a plane wave along x_3, $\mathrm{e}^{\mathrm{i}(kx_3 - \omega x_4)}$; in the cosmological perturbations the comoving wave number of a Fourier mode, $k = |\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.
- $\mathcal{P}_{\mathcal{R}}(k)$: The primordial power spectrum of the comoving curvature perturbation, per logarithmic interval of k. Unit: dimensionless. Status: ASSUMED.
- $\mathcal{I}_l^{\mathrm{T}}(k)$, $\mathcal{I}_l^{\mathrm{E}}(k)$: The transfer functions of the CMB: the multipole l of the temperature or E-mode anisotropy today produced by a Fourier mode k of unit primordial curvature perturbation, computed by the line-of-sight integral. Unit: dimensionless. Status: STANDARD.
- A_{s}: The amplitude of the primordial curvature spectrum at the pivot k_*. Unit: dimensionless. Value: 2.10×10^{-9} (Planck 2018). Status: ASSUMED.
- k_*: The pivot wave number of the primordial spectrum. Unit: GeV (1/length). Value: 0.05 Mpc^{-1}. Status: STANDARD.
- n_{S}: The spectral index of the primordial curvature spectrum. Unit: dimensionless. Value: 0.965 (Planck 2018). Status: ASSUMED.

$$
\begin{aligned}
&\Omega_\chi = \frac{\rho_\chi^{(0)}}{\rho_{\mathrm{c}}^{(0)}}, \qquad \Omega_{\mathrm{b}} = \frac{\rho_{\mathrm{b}}^{(0)}}{\rho_{\mathrm{c}}^{(0)}}, \qquad \Omega_\Lambda = \frac{\rho_\Lambda}{\rho_{\mathrm{c}}^{(0)}}, \qquad \Omega_\nu h_{100}^2 = \frac{\sum_i m_i}{93.14\ \mathrm{eV}}, \\
&\rho_{\mathrm{c}}^{(0)} = \frac{3\big(H_4^{(0)}\big)^2}{8\pi G_{\mathrm{N}}}, \qquad H_4^{(0)} = 100\, h_{100}\ \mathrm{km/s/Mpc}
\end{aligned}
\tag{FE-58}
$$

Symbols in (FE-58).

- Ω_χ: The density parameter of the dark matter today. Unit: dimensionless. Value: $\Omega_\chi h_{100}^2 = 0.120$ (Planck 2018). Status: DERIVED.
- ρ_χ: The energy density of the dark matter, the particles χ of dirac16complex; $\rho_\chi^{(0)}$ its value today. Unit: GeV^4. Status: DERIVED.
- ρ_{c}: The critical density, the energy density of a spatially flat fold at the expansion rate H_4; $\rho_{\mathrm{c}}^{(0)}$ its value today. Unit: GeV^4. Value: $\rho_{\mathrm{c}}^{(0)} = 1.0537 \times 10^{-5}\, h_{100}^2\ \mathrm{GeV/cm}^3$. Status: STANDARD.
- Ω_{b}: The density parameter of the baryons today. Unit: dimensionless. Value: $\Omega_{\mathrm{b}} h_{100}^2 = 0.0224$ (Planck 2018). Status: STANDARD.
- ρ_{b}: The (rest-mass) energy density of the baryons: in a supernova the local density of the matter (quoted in $\mathrm{g/cm}^3$), in cosmology the mean density of the baryons ($\rho_{\mathrm{b}}^{(0)}$ today). Unit: GeV^4. Status: STANDARD.
- Ω_Λ: The density parameter of the dark energy (the vacuum energy of the fold) today. Unit: dimensionless. Value: 0.685 (Planck 2018). Status: DERIVED.
- ρ_Λ: The vacuum energy density of the fold above its static tension, the dark energy of UM-4: $\rho_\Lambda = \rho_{\mathrm{ff}} + \rho_{\mathrm{vac}} - \lambda_{\mathrm{F}} V_*$ (four-dimensional normalisation). Unit: GeV^4. Value: $0.685\, \rho_{\mathrm{c}}^{(0)} = 2.5 \times 10^{-47}\ \mathrm{GeV}^4$ (Planck 2018). Status: DERIVED.

- Ω_ν, $\Omega_\nu^{(i)}$: The density parameter of the neutrinos of the CNB: Ω_ν that of the massive states today; $\Omega_\nu^{(i)}(\mathsf{z})$ the energy density of the mass state i at the redshift z in units of the critical density today. Unit: dimensionless. Status: STANDARD.
- h_{100}: The present expansion rate of the fold in units of 100 km/s/Mpc. Unit: dimensionless. Value: 0.674 (Planck 2018). Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal), and in a tensor of ordinary space such as h_{ij} one of the directions x_1, x_2, x_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m^2_{ij} = m_i^2 - m_j^2$ the squared-mass difference of the states i and j (Δm^2_{21}, Δm^2_{31}, Δm^2_{32}); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by $v_{\rm loc}^2/v^2$ where this differs from 1; m without a label the mass of a mode of a field of the bulk as the fold sees it, $m^2 = \omega^2 - k^2$; for the W and Z bosons their complex pole mass, $m^2 = m_0^2 - \mathrm{i}m_0\Gamma_{\rm bulk}$ to the leading order; in the relic abundance of the particles χ their mass. Unit: GeV. Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $G_{\rm N}$: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_{\rm N} = 1/M_{\rm Pl}^2$ with the Planck mass $M_{\rm Pl} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.

The neutrinos in the expanding universe (FE-53), the scalar perturbations of the metric of the fold in the conformal Newtonian gauge and the Einstein equations of the fold to the first order (FE-54), the Boltzmann equations of the species (FE-55), the BAO ruler and distances (FE-56), the angular power spectra of the CMB (FE-57) and the density parameters (FE-58). Status: STANDARD (UM-4: the dark matter is χ, the dark energy is the vacuum energy of the fold).

14.1 The background and the BAO ruler

$$\Big(\frac{H_4(\mathsf{z})}{H_4^{(0)}}\Big)^2 = \Omega_\gamma(1+\mathsf{z})^4 + \sum_{i=1}^{3}\Omega_\nu^{(i)}(\mathsf{z}) + \big(\Omega_{\rm b}+\Omega_\chi\big)(1+\mathsf{z})^3 + \Omega_\Lambda,$$
$$\Omega_\nu^{(i)}(\mathsf{z}) = \frac{7}{8}\Big(\frac{\mathsf{T}_{\nu,\rm eff}}{\mathsf{T}_\gamma}\Big)^4\Omega_\gamma\,(1+\mathsf{z})^4\,\frac{\mathsf{I}\big(m_i/\mathsf{T}_{\nu,\rm eff}\big)}{\mathsf{I}(0)}, \qquad \mathsf{I}(y) = \int_0^\infty \mathrm{d}\mathsf{q}\,\frac{\mathsf{q}^2\sqrt{\mathsf{q}^2+y^2}}{\mathrm{e}^{\mathsf{q}}+1}, \qquad \mathsf{T}_{\nu,\rm eff} = \Big(\frac{N_{\rm eff}}{3}\Big)^{1/4}\Big(\frac{4}{11}\Big)^{1/3}\mathsf{T}_\gamma \tag{S-26}$$

Symbols in (S-26).

- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- z: The redshift, $1 + \mathsf{z} = 1/a$; z' an integration variable. Unit: dimensionless. Status: STANDARD.
- Ω_γ: The density parameter of the photons today. Unit: dimensionless. Value: 5.450×10^{-5} (for $h_{100} = 0.6736$). Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal), and in a tensor of ordinary space such as h_{ij} one of the directions x_1, x_2, x_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- Ω_ν, $\Omega_\nu^{(i)}$: The density parameter of the neutrinos of the CNB: Ω_ν that of the massive states today; $\Omega_\nu^{(i)}(\mathsf{z})$ the energy density of the mass state i at the redshift z in units of the critical density today. Unit: dimensionless. Status: STANDARD.
- $\Omega_{\rm b}$: The density parameter of the baryons today. Unit: dimensionless. Value: $\Omega_{\rm b}h_{100}^2 = 0.0224$ (Planck 2018). Status: STANDARD.
- Ω_χ: The density parameter of the dark matter today. Unit: dimensionless. Value: $\Omega_\chi h_{100}^2 = 0.120$ (Planck 2018). Status: DERIVED.
- Ω_Λ: The density parameter of the dark energy (the vacuum energy of the fold) today. Unit: dimensionless. Value: 0.685 (Planck 2018). Status: DERIVED.

- $\mathsf{T}_{\nu,\mathrm{eff}}$: The effective temperature of the neutrinos that carries N_{eff} (the convention of CAMB). Unit: GeV (or K). Status: STANDARD.
- T_γ: The temperature of the photons (of the CMB); $\mathsf{T}_\gamma^{(0)}$ its value today. Unit: GeV (or K). Value: 2.7255 K today. Status: STANDARD.
- $\mathsf{I}(y)$: The Fermi-Dirac energy integral of a neutrino state with the mass-to-temperature ratio y; $\mathsf{I}(0) = 7\pi^4/120$. Unit: dimensionless. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m_{ij}^2 = m_i^2 - m_j^2$ the squared-mass difference of the states i and j (Δm_{21}^2, Δm_{31}^2, Δm_{32}^2); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by v_{loc}^2/v^2 where this differs from 1; m without a label the mass of a mode of a field of the bulk as the fold sees it, $m^2 = \omega^2 - k^2$; for the W and Z bosons their complex pole mass, $m^2 = m_0^2 - \mathrm{i}m_0\Gamma_{\mathrm{bulk}}$ to the leading order; in the relic abundance of the particles χ their mass. Unit: GeV. Status: STANDARD.
- y: The ratio of the mass of a neutrino state to its temperature, the argument of I. Unit: dimensionless. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- q: The comoving momentum of a neutrino, $a|\vec{p}|$. Unit: GeV. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- N_{eff}: The effective number of neutrino species: the energy density of the relativistic particles other than photons in units of that of one neutrino species whose temperature is $(4/11)^{1/3}$ times that of the photons. Unit: dimensionless. Value: 3.044 (Standard Model); measured 2.99 ± 0.17 (Planck 2018). Status: STANDARD.

The expansion rate of the fold as a function of the redshift z (FE-59 with the content of UM-4): photons, the three neutrino states with their exact Fermi-Dirac energy integrals (each at the temperature $\mathsf{T}_{\nu,\mathrm{eff}}$ that carries $N_{\mathrm{eff}} = 3.044$, the convention of CAMB), baryons, χ, and the vacuum energy Ω_Λ that makes the fold spatially flat. Computed here it agrees with CAMB to 8.7×10^{-7} for $0 \le \mathsf{z} \le 10^4$, with $\Omega_\gamma = 5.450 \times 10^{-5}$, $\Omega_\nu = 1.404 \times 10^{-3}$ today (the massless state included), $\Omega_{\mathrm{b}} = 0.0493$, $\Omega_\chi = 0.2645$, $\Omega_\Lambda = 0.6848$; the age of the fold universe is 13.798 Gyr. With frozen extra times this expansion is that of general relativity: the bulk changes neither H_4 nor the distances nor the ruler at any observable level. Status: DERIVED (`um4_cosmology.py`, C1).

The ruler FE-56, integrated here with the same background up to the drag epoch $\mathsf{z}_{\mathrm{d}} = 1059.93$ of CAMB, is $r_{\mathrm{s}} = 147.1047$ Mpc (CAMB: 147.1047 Mpc), and the distances D_{M} and D_{H} at the seven redshifts of DESI agree with CAMB to 8.4×10^{-7} (for example $D_{\mathrm{M}} = 1985.92$ Mpc and $D_{\mathrm{H}} = 3345.07$ Mpc at $\mathsf{z} = 0.51$). Status: DERIVED (`um4_cosmology.py`, C2).

14.2 The CMB

quantity	UM-4	Planck 2018	difference in standard deviations
redshift of last scattering z_*	1089.905	1089.92 ± 0.25	-0.06
sound horizon r_* (Mpc)	144.445	144.43 ± 0.26	$+0.06$
acoustic angle $100\,\theta_*$	1.04108	1.04110 ± 0.00031	-0.06
drag redshift z_{d}	1059.931	1059.94 ± 0.30	-0.03
ruler $r_{\mathrm{s}}(\mathsf{z}_{\mathrm{d}})$ (Mpc)	147.105	147.09 ± 0.26	$+0.06$
damping wave number k_{D} (Mpc^{-1})	0.140865	0.14087 ± 0.00030	-0.02
matter-radiation equality z_{eq}	3402.9	3402 ± 26	$+0.03$
σ_8	0.8120	0.8111 ± 0.0060	$+0.15$
age (Gyr)	13.7977	13.797 ± 0.023	$+0.03$

The derived quantities of the CMB of UM-4 with the Planck 2018 parameters (the column TT,TE,EE+lowE+lensing

of Table 2 of Planck 2018 VI) lie within 0.2 standard deviations of the Planck values. The small differences come from evaluating CAMB at the mean values of the Planck parameters (the published derived quantities are means over the posterior) and, for σ_8, $100\theta_*$ and the age, from the three neutrino states of UM-4 (Planck uses one massive state of 0.06 eV). Here σ_8 is the root mean square of the matter density contrast in spheres of radius $8h_{100}^{-1}$ Mpc today. Status: DERIVED (`um4_cosmology.py`, C3).

$$\chi^2_{\mathrm{fit}} = \left(\mathsf{X}_{\mathrm{model}} - \mathsf{X}_{\mathrm{data}}\right)^T \mathsf{C}^{-1} \left(\mathsf{X}_{\mathrm{model}} - \mathsf{X}_{\mathrm{data}}\right), \qquad \mathsf{X}_b = \sum_{l \text{ in the bin } b} \mathsf{w}_l \, C_l \tag{S-27}$$

Symbols in (S-27).

- χ^2_{fit}: The chi-square of a comparison of a model with data (written χ^2 in the text): the squared deviations weighted with the inverse covariance. Unit: dimensionless. Status: STANDARD.
- $\mathsf{X}_{\mathrm{model}}$, $\mathsf{X}_{\mathrm{data}}$: The vectors of the model values and of the measured values of a data set (binned CMB spectra, BAO distances). Unit: as the data. Status: STANDARD.
- ^{T}: The transpose of a matrix. Status: STANDARD.
- C: The covariance matrix of the measured values; C^{-1} its inverse. Unit: as the data squared. Status: STANDARD.
- X_b: A model value of the binned CMB spectrum, for the bin b of the plik-lite binning. Unit: $\mu\mathrm{K}^2$. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- l: The multipole index and number ($l = 0, 1, 2, \ldots$) of an expansion in Legendre polynomials of the direction of a photon or a neutrino relative to the wave vector, or of the CMB sky in spherical harmonics. Status: STANDARD.
- w_l: The Planck weight of the multipole l in its bin. Unit: dimensionless. Status: STANDARD.
- C_l^{TT}, C_l^{TE}, C_l^{EE}: The angular power spectra of the CMB at the multipole l: temperature with temperature, temperature with E-mode polarisation, E-mode polarisation with itself. Unit: dimensionless. Status: STANDARD.

The comparison with measurements, χ^2_{fit} (the usual χ^2): the measured values $\mathsf{X}_{\mathrm{data}}$ (binned spectra or BAO distances), their covariance matrix C, and the model values $\mathsf{X}_{\mathrm{model}}$; for the Planck 2018 high-l likelihood (plik-lite: foreground-marginalised spectra in 215 bins of TT for $30 \le l \le 2508$ and 199 bins each of TE and EE for $30 \le l \le 1996$, with the Planck calibration fixed at 1) a model bin is the weighted sum X_b of the C_l of the multipoles of the bin with the Planck weights w_l; two Gaussian bins of the low-l temperature spectrum can be added (Prince and Dunkley 2019). The implementation reproduces the reference log-likelihoods of the plik-lite code on its test spectrum to 10^{-13} (-101.5812307 for TT, -293.9558650 for TT, TE, EE with the low-l bins). Status: STANDARD (the likelihood), DERIVED (its implementation, `um4_cosmology.py`, C4).

model	N_{eff}	χ^2 high-l TT, TE, EE (613 bins)	χ^2 TT (215 bins)	refitted χ^2 (615 bins)	$H_4^{(0)}$ after the refit
UM-4	3.044	583.8	206.2	587.9	67.31
UM-4 with thermal ν_{R}, same θ_*	3.185	604.3	215.1	589.5	68.23
Planck's convention (one state of 0.06 eV)	3.044	583.9	206.1		

The CMB spectra of UM-4 against the Planck 2018 data: $\chi^2 = 583.8$ for 613 bins (expected 613 ± 35). Three thermal right-handed neutrinos at the same acoustic angle and the same other parameters raise χ^2 by 20.5 (the damping tail); when $\Omega_{\mathrm{b}}h^2_{100}$, $\Omega_\chi h^2_{100}$, θ_*, A_{s} and n_{S} are refitted (Gauss-Newton with the Jacobian of CAMB, τ_{reion} fixed) to the 613 high-l bins and the two low-l bins, the difference shrinks to $\Delta\chi^2 = +1.6$, in line with the Planck bound $N_{\mathrm{eff}} = 2.99 \pm 0.17$ (the refit shifts n_{S} from 0.9650 to 0.9702 and $H_4^{(0)}$ from 67.31 to 68.23 km/s/Mpc; the errors from the Fisher matrix, $\sigma(\Omega_{\mathrm{b}}h^2_{100}) = 0.00015$, $\sigma(\Omega_\chi h^2_{100}) = 0.0013$, $\sigma(n_{\mathrm{S}}) = 0.0042$, match Planck's). The spectra are shown in the figure. Status: DERIVED (`um4_cosmology.py`, C4).




Figure 1: The CMB spectra of UM-4 with the Planck 2018 parameters ($D_l = l(l+1)C_l/(2\pi)$ of the temperature, the cross-correlation and the E-mode polarisation, lensed), with three thermal right-handed neutrinos (dashed), and the Planck 2018 plik-lite data.

14.3 The BAO

tracer	z	measured (DESI DR2)	UM-4	χ^2
BGS	0.295	$D_V/r_s = 7.942 \pm 0.075$	8.057	2.34
LRG1	0.510	$D_M/r_s = 13.588 \pm 0.167$, $D_H/r_s = 21.863 \pm 0.425$	13.500, 22.739	4.48
LRG2	0.706	17.351 ± 0.177, 19.455 ± 0.330	17.700, 20.169	14.36
LRG3+ELG1	0.934	21.576 ± 0.152, 17.641 ± 0.193	21.994, 17.569	8.29
ELG2	1.321	27.601 ± 0.318, 14.176 ± 0.221	28.081, 14.065	2.31
QSO	1.484	30.512 ± 0.760, 12.817 ± 0.516	30.275, 12.880	0.10
Lyα	2.330	38.988 ± 0.531, 8.632 ± 0.101	39.182, 8.616	0.13

The BAO distances of UM-4 with the Planck parameters (the matter density parameter $\Omega_m = 0.3152$, the density of all matter today in units of the critical density, and $h_{100} r_s = 99.09$ Mpc) against DESI DR2 (DESI Collaboration 2025, Table IV; χ^2 with the correlation of D_M and D_H of each tracer): $\chi^2 = 32.0$ for 13 measurements. Fitted to DESI alone, $\Omega_m = 0.2971$ and $h_{100} r_s = 101.56$ Mpc with $\chi^2 = 10.5$ for 11 degrees of freedom ($p = 0.48$), as DESI finds for the cosmological-constant model; the Planck point lies $\Delta\chi^2 = 21.5$ above this optimum when its own errors are neglected (with the CMB errors DESI quotes a 2.3 standard-deviation tension between BAO and CMB). The dark energy of UM-4 is a vacuum energy with $w = -1$ exactly: UM-4 inherits this tension and cannot produce the evolving dark energy that DESI DR2 with the CMB prefers at 3.1 standard deviations, $w(a) = w_0 + w_a(1 - a)$ (the equation of state as a function of the scale factor $a = e^{a_4}$, $a = 1$ today, with its present value w_0 and its slope w_a) with $w_0 > -1$ and $w_a < 0$. Status: DERIVED (`um4_cosmology.py`, C5), STANDARD (DESI's significances).

14.4 The CNB

$$\Gamma_{C\nu B} = N_T \, \bar{\sigma}_T \, \frac{n_\nu^{(0)}}{2} \sum_{i=1}^{3} |U_{ei}|^2 \big(1 + \mathsf{s}_i\big) \qquad \text{(Dirac neutrinos)} \tag{S-28}$$

Symbols in (S-28).

- $\Gamma_{C\nu B}$: The rate at which a tritium target captures neutrinos of the cosmic neutrino background. Unit: 1/time. Status: STANDARD.
- N_T: The number of tritium nuclei of the target. Unit: dimensionless. Value: 2.0×10^{25} for 100 g. Status: STANDARD.
- $\bar{\sigma}_T$: The averaged cross-section of the capture of an electron neutrino on tritium: a tritium nucleus captures left-handed electron neutrinos of mean speed s and number density n at the rate $\bar{\sigma}_T(1 + \mathsf{s})\, c\, n$ (c the speed of light, 1 in the formulas; Long, Lunardini and Sabancilar). Unit: cm^2. Value: 3.834×10^{-45} cm^2. Status: STANDARD.
- n_ν: The number density of the cosmic neutrino background per flavour (neutrinos plus antineutrinos, one helicity each); $n_\nu^{(0)}$ its value today. Unit: GeV^3. Value: 112 cm^{-3} today. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal), and in a tensor of ordinary space such as h_{ij} one of the directions x_1, x_2, x_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- U_{ei}: The electron row of the mixing matrix, $U_{ei} = (U_{PMNS})_{ei}$: $|U_{ei}|^2$ is the electron-flavour fraction of the mass state i. Unit: dimensionless. Value: $|U_{ei}|^2 = 0.682$, 0.296 and 0.022 for $i = 1, 2, 3$ (NuFIT 5.2, the fit

without the Super-Kamiokande atmospheric data: the values of the record; NuFIT 6.0 with them gives 0.677, 0.301 and 0.022). Status: STANDARD.

- s_i: The mean speed of the neutrino mass state i today, averaged over its momentum distribution (0 for a state at rest, 1 for a massless one). Unit: dimensionless. Value: 1, 0.061 and 0.011 for the mass states 1, 2 and 3 with $m_1 = 0$; 0.023 for the state 1 with $m_1 = 23$ meV. Status: STANDARD.

The capture rate of the cosmic neutrinos on a target of N_T tritium nuclei ($\nu_e + {}^3\mathrm{H} \to {}^3\mathrm{He} + e^-$, the PTOLEMY idea): each mass state i contributes with its electron fraction $|U_{ei}|^2$; for Dirac neutrinos only the left-handed helicity of each state is populated ($n_\nu^{(0)}/2 = 56$ cm^{-3} per state), and it is captured with the averaged cross-section $\bar{\sigma}_\mathrm{T} = 3.834 \times 10^{-45}$ cm^2 times $(1 + \mathsf{s}_i)$, s_i the mean speed of the state over its momentum distribution (Long, Lunardini and Sabancilar 2014). Status: STANDARD.

The CNB of UM-4: the temperature $\mathsf{T}_\nu^{(0)} = (4/11)^{1/3}\mathsf{T}_\gamma^{(0)} = 1.9454$ K (1.676×10^{-4} eV); 112.0 neutrinos and antineutrinos per cm^3 per flavour, 336 in all (FE-53); the states ν_2 and ν_3 became non-relativistic at $\mathsf{z} = 15.3$ and 94 (where their mass equals the mean momentum $3.15\mathsf{T}_\nu$), ν_1 is massless; $\Omega_\nu h^2_{100} = 6.31 \times 10^{-4}$ for the two massive states (FE-58: $\sum m_i/93.14$ eV $= 6.30 \times 10^{-4}$; CAMB: 6.31×10^{-4}). Because the right-handed states are not populated, the CNB of UM-4 has one helicity per state: a 100 g tritium target captures 4.1 cosmic neutrinos per year if all states are non-relativistic (Majorana neutrinos: 8.1, twice as many), and 6.9 per year in UM-4, whose massless ν_1 moves with the speed of light, so that its left-handed helicity is captured twice as often, while ν_2 and ν_3 move with the mean speeds $\mathsf{s}_2 = 0.061$ and $\mathsf{s}_3 = 0.011$ today. The sum of the masses, 0.0587 eV, lies below the bounds of DESI DR2 with the CMB (0.0642 eV for three equal masses, 0.101 eV in the normal ordering with NuFIT 6.0) and of ACT DR6 (0.082 eV, 95%). With the largest lightest mass that DESI DR2 with NuFIT 6.0 allows, $m_1 = 23$ meV, the capture rate falls to 4.15 per year (S-18). Status: DERIVED (`um4_cosmology.py`, C6).

15 What the solution does not settle

- **The tension and the dark energy** (DERIVED: the tuning; OPEN: its origin). The static block needs the total tension $\lambda_\mathrm{F} V_* = 30 H^2 M_\mathrm{red}^2$ of the fold (FE-65); the dark energy ρ_Λ is the small excess of the tension of the fold fluid and the vacuum energy of the fields over it (S-4), a tuning to one part in 10^{92} for $H = 2534$ GeV, as severe as the cosmological-constant problem of the brane of Randall and Sundrum. UM-4 accommodates the value of ρ_Λ but does not explain it, and its dark energy has $w = -1$ exactly, so the evolving dark energy that DESI DR2 with the CMB prefers is not available.
- **The volume of the extra times** (DERIVED: that it must be held fixed; ASSUMED: the holding pressure; OPEN: its origin). A free volume would change Newton's constant 8545 times faster than lunar laser ranging allows (S-2). UM-4 assumes that the fold fluid supplies the pressure q_hold along the extra times that holds it fixed (FE-65, S-3; its value today depends on the assumed pressure of the baryons along the extra times); what produces this pressure (a potential for the volume, like the stabilisation of the sizes of compact extra dimensions in other theories) is not part of UM-4.
- **The bulk of an expanding fold** (DERIVED: the first order; OPEN: the Weyl tensor). The bulk metric of an expanding fold is computed to the first order in the curvature of the fold over H^2 (FE-34) with no dark radiation ($\mathfrak{c}_{ab} = 0$, ASSUMED); the projected Weyl tensor that dust requires at the order $\rho^2/(\lambda_\mathrm{F} V_*)$ (S-1), and that an inflation with frozen extra times requires (FE-88), needs the bulk solution, which is not computed.
- **The dark matter** (DERIVED, ESTIMATE; OPEN: its binding and its origin). The particles χ of dirac16complex are resonances of the fold (FE-86) with the derived width $(2/\mathrm{e})(q_\mathrm{b}^2/m_\Psi)\exp(-2F(\upsilon_\mathrm{b})m_\Psi/H)$; they live longer than 160 Gyr only if m_Ψ/H is large enough (S-6: 1135 for $q_\mathrm{b} = m_\Psi/2$, so with $H > 2534$ GeV they are then heavier than 2.5×10^6 GeV; the bound falls with the binding, to 1.2×10^4 GeV for $q_\mathrm{b} = 0.99\, m_\Psi$). Above 9.7×10^4 GeV no thermal relic is possible (S-20), so only particles bound more strongly than $q_\mathrm{b} \approx 0.9\, m_\Psi$ could be a thermal relic; for a weaker binding their abundance needs a non-thermal origin. The portals of IN-16 cannot provide the binding: bound by the author's Higgs portal the particles would have to be bound so weakly (direct detection) that they live long enough only for $H < 2.5$ GeV (S-22), bound by the gluon condensate they would scatter on nucleons 29 to 33 orders of magnitude above the limit (S-29), and the other mean values of the portals vanish in the vacuum; the current portals, whose convective current annihilates the particles only with the fourth power of the velocity, cannot set the relic density of a perturbatively coupled χ (S-21). The dark matter of UM-4 needs a binding of χ to the fold that does not couple it to nucleons, which the action of UM-4 does not contain.
- **The Krein norm** (DERIVED; OPEN: stability and a unitary coupling). The charge of dirac16complex is indefinite (FE-43); its quantisation (FE-67) has ghosts that no choice of the vacuum removes. The vacuum

cannot decay; whether the gas of the particles χ is classically stable is not decided: the symmetric occupation of the level is a saddle of the first-order energy, which has no lower bound on the occupations of negative norm, and the truncated response has an imaginary pair whose meaning is open. The chiral couplings of the fold, the left-handed Higgs portal among them, produce ghosts (with exactly the elastic amplitude), so UM-4 with them is not unitary; chirality-blind couplings are necessary for unitarity but not shown to be sufficient.

- **The extra times** (ASSUMED, OPEN). Every field with a momentum along the extra times larger than its mass and its momentum along ordinary space is tachyonic (FE-38); UM-4 keeps only fields that do not depend on them. The restriction is consistent but not stable: interactions of fields without extra-time momentum can create pairs of opposite momenta (ESTIMATE, no rate computed), and compact extra times do not help. UM-4 needs a gauge principle that removes the momentum along the extra times of every field. The four time-like directions also allow closed time-like curves through every point of the eight-dimensional spacetime (every circle in a plane of the time and an extra time); the time x_4 is a time function only on the sections of fixed extra times, on which UM-4 lives, and only there does the bulk give no shortcut to signals (FE-83). Of the polarisations of the graviton of the bulk the three vectors $h_{\mu r}$ are ghosts (FE-76); the matter of the fold does not emit them at the linear order, but they remain a defect of the bulk gravity of signature (3, 4) at higher orders (OPEN).
- **The primordial era** (DERIVED, ESTIMATE, OPEN). The author's vacuum is not a solution (FE-31); the exact de Sitter fold inflates the extra times as well (FE-87); the inflation with frozen extra times (FE-88) needs a Weyl tensor of the bulk and, in the regime of general relativity, a rate below $H/10$ (S-24). UM-4 does not specify its inflaton: Higgs inflation, an extension of the action, would need $H > 1.6 \times 10^{14}$ GeV (S-25); what precedes inflation and the reheating are not computed; the ad hoc vacuum must decay with $a_{40} > 4.8 \times 10^{-16} (v_{\mathcal{A}}/v)^2$ GeV so that its energy never exceeds that of the radiation (S-5, the thermal history); the primordial spectrum, the optical depth and the densities are taken from Planck 2018.
- **The free parameters**. The bulk mass m_Ψ, the scale H (bounded from below by 2534 GeV), the volume V_* of the extra times (Newton's constant fixes only $8\pi G_{\mathrm{N}} = 5\kappa H/(2V_*)$), the gauge couplings of eight dimensions (fixed by the measured couplings through FE-44), the portal couplings, the self-coupling λ_Ψ, the amplitude $v_{\mathcal{A}}$ and rate a_{40} of the ad hoc vacuum, the lightest neutrino mass (taken as zero; below 23 meV, S-18) and the Dirac phase (taken as zero; it drops out of every result, S-17) are not fixed by UM-4.
- **Neutrinos in dense matter** (STANDARD, ESTIMATE). The non-linear evolution of the fast instability and the collective conversion deep in a supernova are not solved; their outcome is modelled by the equipartition FE-52, whose effect on the explosion is taken from the transport simulations of Gogilashvili and Tamborra.

16 What was verified

Every result labelled DERIVED rests on a computation in the folder `universe_model_4/code/` of the repository, which checks itself and writes a report to `universe_model_4/reports/` (all equations and results are written out above). Each script is run with `python universe_model_4/code/NAME.py` (Python 3 with numpy, scipy, sympy, matplotlib and CAMB 2.0.5; the cosmology script downloads the Planck 2018 plik-lite files from github.com/heatherprince/planck-lite-py into `universe_model_4/external/` once), and no single calculation runs longer than 423 seconds. All 189 checks of the 20 programs pass.

- `um4_dirac_matrices.py` (55 checks): the author's matrices, their reality, the Clifford relations, the chiral projectors, Pin(4,4).
- `um4_geometry.py` (11 checks): the proper chart and the determinant of the metric, the exact curvature of the metric of UM-4, the static block of constant curvature with a vanishing Weyl tensor, the Lovelock tensors of the static block, the exact bulk equations of Einstein gravity (the Einstein tensor in two ways), the Ricci-flat folds, the pure tension from the jump of the extrinsic curvature, the light cone of every slice, the spin connection with the author's matrices on both halves, the vacua of the static warps (only equal rates), and the matrices of FD-13 against the author's notebook.
- `um4_gravity.py` (4 checks): the vacua of the static family, the graviton around the static block for any weights, the light cone on every background only for Einstein gravity, the gravity of UM-4.
- `um4_bulk.py` (10 checks): the bulk scalar and the bulk photon (their wave operators from the metric), the transverse photon of spin 1, the bulk graviton, the plus and cross polarisations of spin 2 with the energy-momentum tensor of the fold, the bulk fermion, the boost invariance of dirac16complex, the speeds of UM-3 recomputed, the continuum of massive modes from zero mass, and the signs of the kinetic terms of the 14 polarisations of the graviton (the ghost vectors).

- `um4_fold.py` (6 checks): the bulk metric of an expanding fold, the Einstein equations of seven dimensions on the fold, the exact equations of the fold, the Friedmann equations with frozen extra times, the size of the corrections, the Weyl term that dust requires.
- `um4_foldsources.py` (5 checks): the tension, the gravitational waves of the fold, the induced gravity, the tuning of the tension, the fold fluid.
- `um4_background.py` (6 checks): the vacua of the author's form, the expanding fold, the exclusion of a free volume, the fold factor, the ad hoc vacuum as a gauge rotation (S-5) and its numbers.
- `um4_propagators.py` (5 checks): the propagators of the bulk, their expansions, Newton's and Coulomb's laws, the W and Z resonances, the bounds on H.
- `um4_causality.py` (4 checks): no shortcut through the static bulk at fixed extra times and the null lines along the extra times, the light cone of an expanding fold, the largest advance, the closed time-like curves.
- `um4_fields.py` (16 checks): the bilinears of dirac16complex, the Krein operator, the electroweak masses of the ad hoc vacuum, the anomalies with three right-handed neutrinos.
- `um4_spectra.py` (8 checks): the algebra of the profile equation, its exact solutions on both sides, the resonances with the remainder of the first-order shift, their widths with the derived prefactor, the dispersion of a moving resonance, the lifetime condition over the bindings (S-6), their Krein signature and the mirror of the fold.
- `um4_kohnsham.py` (10 checks): the Kohn-Sham states (S-7 to S-12), lifted to sixteen components, and how dilute the gas of the universe is.
- `um4_krein.py` (9 checks): FE-67, the symmetries and mirrors of the states at the fold, the flat directions and the truncated response, the partner level of the gas, the selection rules at the fold with the invariant amplitudes, the consequences, the occupation as a saddle, the pairs of the convective current.
- `um4_extratimes.py` (5 checks): the fields with a momentum along the extra times.
- `um4_neutrino.py` (5 checks): the flavour blindness of V_Ψ, the resonance formulas, the crossing criterion, the equipartition, the CNB numbers.
- `um4_flavour.py` (4 checks): S-13 to S-16.
- `um4_numass.py` (5 checks): S-17, S-18 and the comparison with NuFIT 6.0.
- `um4_thermal.py` (9 checks): the degrees of freedom, S-19 to S-23 and S-29, the ad hoc vacuum before nucleosynthesis.
- `um4_primordial.py` (6 checks): the author's vacuum (not a solution), the de Sitter fold, the inflation with frozen extra times, its scales (S-24), Higgs inflation (S-25, an extension of the action) and the fold factor.
- `um4_cosmology.py` (6 checks): S-26 to S-28 and the comparisons with Planck 2018 and DESI DR2.

The Kohn-Sham module runs in parts (`um4_kohnsham.py part NAME` for NAME = K1, I, K6, K9, then `um4_kohnsham.py assemble`), so that no single calculation takes longer than 423 seconds; the assembled results are identical to a single run.

17 Further symbols

Further symbols. These symbols appear in the definitions of the symbols listed after the equations of this document.

- M_{red}: The reduced Planck mass, $M_{\mathrm{red}} = (8\pi G_{\mathrm{N}})^{-1/2}$. Unit: GeV. Value: 2.435×10^{18} GeV. Status: STANDARD.
- h: The Higgs field: the fluctuation of the lower component of the Higgs doublet about v. Unit: GeV. Value: $m_h = 125.25$ GeV. Status: STANDARD.
- G_{F}: The Fermi constant, $G_{\mathrm{F}} = 1/(\sqrt{2}\,v^2)$ in the Standard-Model vacuum (its measured value); in the equations of the neutrino flavours it stands for the local Fermi constant of the ad hoc vacuum. Unit: GeV^{-2}. Value: 1.1664×10^{-5} GeV^{-2}. Status: STANDARD.
- S_Ψ: The action of dirac16complex in the bulk (kinetic term with the canonical spin connection, mass term, contact self-interaction). Unit: dimensionless. Status: RECORD.
- $\Gamma^P{}_{MN}$: The Christoffel symbols of the metric g_{MN} (the Levi-Civita connection). Unit: 1/length. Status: STANDARD.
- β: The matrix of the four-dimensional Dirac adjoint, $\beta = \hat{\gamma}^{(4)}$ (Hermitian, $\beta^2 = \mathbf{1}$). Unit: dimensionless. Status: STANDARD.
- ϵ^{IJK}: The Levi-Civita symbol of three indices (the structure constants of $SU(2)$), $\epsilon^{123} = 1$. Unit: dimensionless. Status: STANDARD.
- ϖ_{Mab}: The canonical (Levi-Civita) spin connection of the vielbein, antisymmetric in a, b. Unit: 1/length.

Status: STANDARD.

- *: Complex conjugation of a number, or of each entry of a matrix or column. Status: STANDARD.
- $v_{\mathcal{A}}$: The amplitude of almightyAdHoc at $x_4 = 0$; the author's almightyAdHoc, $\mathrm{e}^{-2a_{40}x_4}$ (written Exp[-2 a40 x4] in his notebook), has $v_{\mathcal{A}} = 1$ in the units of his notebook. Unit: GeV. Value: 1 in the units of the author's notebook; UM-4 keeps it free (the examples take $v_{\mathcal{A}} = v$ and $10v$). Status: AUTHOR (the form), ASSUMED (a free amplitude).
- L_{t}: The physical period of the extra times, if they are compact. Unit: length. Status: ASSUMED.
- $\mathcal{W}$: The warp factor: in a warped metric, the function of the position along one direction that multiplies the squared steps of other directions (its square root multiplies their lengths); in UM-4 the function of σ that multiplies all seven directions of the fold, ordinary space, the time and the extra times, $\mathcal{W} = \mathrm{e}^{-2H|\sigma|} = (\sin^2 z)^{1/6}$, equal to 1 on the fold. Unit: dimensionless. Status: DERIVED.
- X_{n}: The fraction of the nucleons that are neutrons, before nucleosynthesis. Unit: dimensionless. Status: STANDARD.
- Y_{p}: The primordial helium mass fraction (helium-4); in the estimate of um4_thermal.py T6, twice the neutron fraction at the deuterium bottleneck. Unit: dimensionless. Value: 0.245 ± 0.003 (observed, Particle Data Group 2024). Status: STANDARD.
- δ_{CP}: The Dirac phase of the mixing matrix U_{PMNS}. Unit: degree. Value: 212° (NuFIT 6.0, normal ordering); 0 in the computations of UM-4. Status: STANDARD.
- m_{W}: The mass of the W bosons. Unit: GeV. Value: 80.37 GeV ($\mathcal{A} = 0$). Status: STANDARD.
- m_{Z}: The mass of the Z boson. Unit: GeV. Value: 91.19 GeV ($\mathcal{A} = 0$). Status: STANDARD.
- ν_α, ν_i: A neutrino field: ν_α the flavour states (the flavour index runs over the electron, muon and tau flavours), ν_i the mass states ($i = 1, 2, 3$). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.