UM-2 solution

An accurate approximate solution of the coupled field equations: Einstein-Lovelock gravity, Kohn-Sham theory for the dirac16complex field coupled to gravity and to every Standard-Model field, three generations of quarks and leptons, the flavour conversion of the three neutrinos, and the CMB, CNB and BAO

Underlying theory: Patrick L. Nash Derived and verified by Claude (Anthropic)

Solution, version 4, 2026-10-18 Versions 3 and 2: 2026-10-18; version 1: 2026-10-17

Repository once-ere/Dirac_claude, folder universe_model_2

Contents

1 How to read this document **3**

2 The coupled system **4**

3 The approximation scheme **11**

4 Step 1: the background and the primordial era **14**
4.1 The field equations of step 1 . . . 14
4.2 The approximation of step 1 . . . 26
4.3 The solution of step 1 . . . 27
4.3.1 The static block . . . 27
4.3.2 Stability: the radion . . . 28
4.3.3 The primordial era: the deflating extra times . . . 29
4.3.4 The response of the block to matter . . . 30
4.3.5 Four-dimensional gravity . . . 32
4.4 The check of step 1 . . . 35
4.5 The status of step 1 . . . 36

5 Step 2: the masses from the x_0 dependence of the wave functions **36**
5.1 The field equations of step 2 . . . 36
5.2 The approximation of step 2 . . . 51
5.3 The solution of step 2 . . . 52
5.3.1 The x_0 eigenvalue problem of the gauge bosons and the graviton . . . 52
5.3.2 The fermions: the exact reduction and the zero modes . . . 55
5.3.3 The masses of one generation . . . 57
5.3.4 Three generations: the masses and the mixing matrices . . . 59
5.3.5 The spectrum of dirac16complex . . . 61
5.4 The check of step 2 . . . 62
5.5 The status of step 2 . . . 63

6 Step 3: the Kohn-Sham states of dirac16complex coupled to all fields **63**
6.1 The field equations of step 3 . . . 63
6.2 The approximation of step 3 . . . 76
6.3 The solution of step 3 . . . 77
6.3.1 The numerical method . . . 77
6.3.2 Validation against the record . . . 77

6.3.3 Ground states on the stabilised block . . . 77
6.3.4 The dark-matter quanta . . . 78
6.3.5 The Standard-Model fields acting on dirac16complex . . . 79
6.3.6 The back-reaction on the block . . . 81
6.4 The check of step 3 . . . 82
6.5 The status of step 3 . . . 83

7 Step 4: the electroweak sector 83
7.1 The field equations of step 4 . . . 83
7.2 The approximation of step 4 . . . 94
7.3 The solution of step 4 . . . 95
7.3.1 Charges and couplings . . . 95
7.3.2 Anomaly cancellation in three generations . . . 95
7.3.3 The two halves of dirac16complex . . . 96
7.3.4 The Z invisible width and the tip Z portal of version 1 . . . 96
7.3.5 The Z widths of three generations . . . 97
7.4 The check of step 4 . . . 98
7.5 The status of step 4 . . . 98

8 Step 5: the flavour conversion of the three neutrinos, $N_{\rm eff}$ and the CNB 98
8.1 The field equations of step 5 . . . 98
8.2 The approximation of step 5 . . . 111
8.3 The solution of step 5 . . . 111
8.3.1 Masses, mixing and the evolution . . . 111
8.3.2 Vacuum, matter and accelerators . . . 112
8.3.3 The Sun . . . 112
8.3.4 The early Universe and $N_{\rm eff}$ . . . 112
8.3.5 The cosmic neutrino background . . . 114
8.3.6 Supernovae . . . 115
8.3.7 Dirac or Majorana . . . 116
8.4 The check of step 5 . . . 118
8.5 The status of step 5 . . . 119

9 Step 6: the CMB, the CNB and the BAO 119
9.1 The field equations of step 6 . . . 119
9.2 The approximation of step 6 . . . 127
9.3 The solution of step 6 . . . 128
9.4 The check of step 6 . . . 130
9.5 The status of step 6 . . . 132

10 Step 7: the dark-matter relic and its detection 132
10.1 The field equations of step 7 . . . 132
10.2 The approximation of step 7 . . . 136
10.3 The solution of step 7 . . . 137
10.4 The check of step 7 . . . 140
10.5 The status of step 7 . . . 140

11 Step 8: the coupled solution in environments 140
11.1 The field equations of step 8 . . . 140
11.2 The approximation of step 8 . . . 161
11.3 The solution of step 8 . . . 162
11.3.1 The portals on the zero modes . . . 162
11.3.2 The mean fields of the Standard-Model fields . . . 162
11.3.3 dirac16complex acting on the Standard Model . . . 163
11.3.4 Gravity: the block response to every species . . . 163
11.4 The check of step 8 . . . 164
11.5 The status of step 8 . . . 165

12 Step 9: the thermal history and the relic with I13 **165**
12.1 The field equations of step 9 . . . 165
12.2 The approximation of step 9 . . . 179
12.3 The solution of step 9 . . . 179
12.3.1 The thermal history . . . 179
12.3.2 The Boltzmann equation . . . 180
12.3.3 I12 alone, and partial-wave unitarity . . . 180
12.3.4 The relic with I13 . . . 181
12.3.5 The fold bands: freeze-in through I11 . . . 181
12.3.6 The self-interaction channel . . . 182
12.3.7 Detection . . . 182
12.4 The check of step 9 . . . 184
12.5 The status of step 9 . . . 185

13 Step 10: the terms of the metric power series **185**
13.1 The field equations of step 10 . . . 185
13.2 The approximation of step 10 . . . 192
13.3 The solution of step 10 . . . 192
13.4 The check of step 10 . . . 193
13.5 The status of step 10 . . . 193

14 Accuracy of the solution **193**

15 What the solution does not reproduce, and what is open **197**

1 How to read this document

What it contains. This document presents the accurate approximate solution of the coupled field equations of the universe model UM-2, in ten steps that follow the hierarchy of the problem: (1) the background and the primordial era; (2) the masses from the x_0 dependence of the wave functions; (3) the Kohn-Sham states of dirac16complex coupled to all fields and their back-reaction on gravity; (4) the electroweak sector; (5) the flavour conversion of the three neutrinos, N_{eff} and the cosmic neutrino background (CNB); (6) the cosmic microwave background (CMB), the CNB and the baryon acoustic oscillations (BAO); (7) the dark-matter relic and its detection; (8) the coupled solution in ten environments; (9) the thermal history and the relic with the portal I13; (10) the terms of the metric power series. Before the steps, "The coupled system" lists every field equation with its fields and the step that solves it, and "The approximation scheme" states the small parameters. After the steps, "Accuracy of the solution" collects in one table, for every equation, the method, the measured residual, the neglected terms and the checks in concert with the other field equations; the last section states what the solution does not reproduce and what is open.

How every step is presented. Each step has five parts: (1) *the field equations*: every field equation (tag FE-n) and every interaction term of the action (tag IN-n) that the step solves or uses, written out in full with its list of symbols, also when an earlier step has already written it, so that every definition stands next to the equation that uses it; (2) *the approximation*: the small parameter and the size of what is neglected, as equations with their symbol lists; (3) *the solution*: formulas and numbers; (4) *the check*: the residual or the measured error of the solution in its own equation, and its consistency with the other field equations that contain the same fields (in concert); (5) *the status* of each result. The tags FE-n and IN-n are one numbering, shared by all documents of UM-2, of the 136 field equations and the 18 interaction terms of the model.

Symbols. Every display equation is followed by the complete list of its symbols: every quantity, field, coupling, constant, operator, index (with the values it takes) and special mark, with its meaning, its unit, its value where it has one, and its status. Typography needs no definition (digits, $+$, $-$, $=$, brackets, fractions, roots, $\cdot$, $\ldots$, $\leq$, $\approx$, $\sim$, $\times$, the standard functions sin, cos, tan, cot, sinh, cosh, tanh, exp, ln, log, det, min, max, lim and the signs of sums, products and integrals; their index letters and integration variables are symbols and are listed). A symbol has one meaning throughout the document, in the displays, in the inline formulas and in the text. Euler's number is written e, the imaginary unit i and the differential d.

Status labels. Every result carries one of these labels:

- **AUTHOR**: taken from the author's notebooks (the coordinates, the metric, the Lovelock weights, the gamma matrices, the mass parameter);
- **RECORD**: taken from the Revision record of this repository (the dirac16complex Lagrangian, the Kohn-Sham reference states);
- **STANDARD**: textbook physics or a measured value used as an input;
- **ASSUMED**: a choice made for this model;
- **INVENTED**: a new phenomenological ingredient of this model (the tip sigma field, the portals I11, I12, I13);
- **DERIVED**: computed and verified in this work (symbolically or numerically); the verification is stated with the result;
- **ESTIMATE**: an order-of-magnitude evaluation, not a precision computation;
- **OPEN**: not resolved in this version.

Units and conventions. $\hbar = c = 1$. In the numerical work on the block the author's constant is $H = 1$ and the mass of dirac16complex is $m_\Psi = H$, unless a physical unit is given; the physical scale is $H = 14.296$ TeV (step 2). The coordinates are the author's $X = \{x_0, x_1, \ldots, x_7\}$: x_0 the hidden space-like direction, listed first, periodic with the period $\pi/(6H)$; x_1, x_2, x_3 ordinary space; x_4 the time; x_5, x_6, x_7 the three extra time-like directions. W^+ carries the electric charge $+e$ and W^- the charge $-e$; UM-2 contains no Higgs particle.

The numbers. Every number is taken from the outputs of the computations of UM-2 (the scripts in `universe_model_2/code/`, which write `universe_model_2/outputs/` and `universe_model_2/reports/`; the residuals of this version are computed by `universe_model_2/code/um2_residuals.py`) and is quoted with the precision of the output. These files are named for reproduction only: everything the reader needs (every equation, every definition and every argument) is written in this document.

Versions. Version 1 (2026-10-17) had one generation. Version 2 (2026-10-18) has three generations of quarks and leptons and three neutrino flavours with their mixing (step 5), and removes the chiral tip Z portal of version 1 (step 4). Version 3 (2026-10-18), by the author's order to find and document an accurate approximate solution of the coupled field equations, added the coupled solution in environments (step 8), the thermal history and the relic with the portal I13 (step 9), the terms of the metric power series (step 10), the Majorana alternative (step 5) and the matter power spectrum (step 6); after its independent review the I13 sum leaves out the direction x_0, the coupled solution uses the full adiabatic block response with the hot plasma and the order-α electroweak condensates, and step 9 has the self-interaction channel of the dark matter. **Version 4** (2026-10-18), by the author's order of 2026-10-18, 15:42, reorganises the presentation: every field equation is written out with its symbols wherever it is used, every step states its approximation, its solution and its check in concert with the other field equations, every symbol has one meaning (several letters of version 3 had more than one), and the residuals of the solution in its equations are computed anew (the check of every step and the table "Accuracy of the solution"). The physics and the claims of version 3 are unchanged; every number is taken from the computations as refined for version 4 (precision fixes changed a few values in their last digits, and each change is stated where the value appears), and the residual computation adds 52 checks to the 235 of version 3 (287 in all).

2 The coupled system

The fields of UM-2 are the metric g_{MN} of Einstein-Lovelock gravity; the dirac16complex field Ψ; the gauge fields of $SU(3)_c \times SU(2)_L \times U(1)_Y$ (gluons; W^+, W^-, Z and the photon); the quarks and leptons of three generations (with the three left-handed neutrino flavours, the left-handed electrons, the singlet right-handed electrons and right-handed neutrinos); and the tip sigma field Σ (no Higgs particle). They are coupled: gravity to everything (the canonical spin connection for Ψ and the fermions, the metric in every kinetic term, the stabiliser); Ψ to gravity, to itself (the self-coupling λ), to the fermion currents (I11), to the gauge-field invariants (I12) and, with its x_0-chiral current, to the baryon-number current (I13); the fermions to the gauge fields, to Σ at the tip (through the tip matrices, which carry the flavour mixing) and to Ψ (I11, and I13 for the quarks); the gauge fields to the fermions, to Σ at the tip and to Ψ (I12, a field-dependent coupling).

The action consists of 18 interaction terms (IN-1 to IN-18); varying it gives the 136 field equations FE-1 to FE-136, which include their exact reductions on the static block and the equations of the four-dimensional effective theory. Every one is written out, with its symbols, in the steps listed. The action is the sum of the interaction terms:

$$S = S_{\mathrm{EL}} + S_{\mathrm{stab}} + S_{\Psi} + S_{\mathrm{gauge}} + S_{\mathrm{f}} + S_{\mathrm{tip}} + S_{\mathrm{I11}} + S_{\mathrm{I12}} + S_{\mathrm{I13}} \tag{IN-1}$$

Symbols in (IN-1).

- S: The action of UM-2 (the integral of the Lagrangian density over spacetime); its variation with respect to every field gives the field equations. Unit: dimensionless ($\hbar = 1$). Status: ASSUMED.
- S_{EL}: The Einstein-Lovelock action of gravity. Unit: dimensionless. Status: AUTHOR.
- S_{stab}: The action of the block stabiliser (a vacuum energy of the block). Unit: dimensionless. Status: ASSUMED.
- S_{Ψ}: The action of the dirac16complex field (kinetic term with the canonical spin connection, mass term and contact self-interaction). Unit: dimensionless. Status: RECORD.
- S_{gauge}: The action of the gauge fields of SU(3), SU(2) and $\mathrm{U}(1)_Y$ in eight dimensions. Unit: dimensionless. Status: ASSUMED.
- S_{f}: The action of the Standard-Model fermions of three generations (kinetic and bulk-mass terms). Unit: dimensionless. Status: ASSUMED.
- S_{tip}: The action on the tip $y = -L$: the sigma field and the tip Yukawa couplings. Unit: dimensionless. Status: INVENTED.
- S_{I11}: The action of the portal I11 (dirac16complex vector current times the $B - L$ current of the fermions). Unit: dimensionless. Status: INVENTED.
- S_{I12}: The action of the portal I12 (scalar density of dirac16complex times the gauge kinetic invariants). Unit: dimensionless. Status: INVENTED.
- S_{I13}: The action of the portal I13 (the x_0-chiral vector current of dirac16complex times the baryon-number current). Unit: dimensionless. Status: INVENTED.

term	interaction	fields	steps
IN-1	The action of UM-2	all fields	written out above
IN-2	Einstein-Lovelock gravity with the author's weights	metric g_{MN}	1
IN-3	The Lovelock densities	metric g_{MN}	1
IN-4	The block stabiliser	metric g_{MN} (block scale f)	1
IN-5	The dirac16complex field	dirac16complex Ψ; metric g_{MN}	3
IN-6	The Clifford algebra, the Dirac adjoint and the two halves of dirac16complex	dirac16complex Ψ; every Standard-Model fermion	3, 4
IN-7	The canonical spin connection (the coupling of every spinor to gravity)	metric g_{MN}; dirac16complex Ψ; every Standard-Model fermion	3
IN-8	The gauge fields of $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)_Y$ in eight dimensions	gluons; $SU(2)$ fields; hypercharge field; metric g_{MN}	2, 4
IN-9	The field strengths	gluons; $SU(2)$ fields; hypercharge field	4
IN-10	The physical electroweak fields W^+, W^-, Z and the photon (canonically normalised)	W^+; W^-; Z; photon; $SU(2)$ fields; hypercharge field	2, 4
IN-11	The Standard-Model fermions: three generations with bulk masses	quarks Q, u_R, d_R; leptons ℓ, e_R, ν_R; metric g_{MN}; gauge fields	2
IN-12	The covariant derivative of a Standard-Model fermion and the electric charge	every Standard-Model fermion; gauge fields; metric g_{MN}	4
IN-13	The tip sigma field and the tip Yukawa couplings (no Higgs particle)	sigma field Σ; $SU(2)$ fields; hypercharge field; every Standard-Model fermion	2, 4
IN-14	The covariant derivative of the sigma field and its gauge transformation	sigma field Σ; $SU(2)$ fields; hypercharge field	4

term	interaction	fields	steps
IN-15	The tip mass term of W^+, W^- and Z in the unitary gauge	W^+; W^-; Z; photon	2, 4
IN-16	Portal I11: the vector contact portal to the $B-L$ currents	dirac16complex Ψ; every Standard-Model fermion	3, 8, 9
IN-17	Portal I12: the gauge-kinetic scalar portal	dirac16complex Ψ; gluons; $SU(2)$ fields; hypercharge field	3, 7, 8, 9
IN-18	Portal I13: the x_0-chiral vector portal to the baryon-number current	dirac16complex Ψ; quarks	3, 8, 9

equation	field equation	fields	solved (used) in step
FE-1	The Einstein-Lovelock equations	metric g_{MN}; all fields (source)	1 (order 0), 6 (order 1), 10
FE-2	The energy-momentum tensor: definition and components	metric g_{MN}; all fields	1
FE-3	The total source of gravity	all fields	1, 8
FE-4	The energy-momentum tensor of the stabiliser	metric g_{MN} (block scale f)	1
FE-5	The energy-momentum tensor of dirac16complex	dirac16complex Ψ; metric g_{MN}	3
FE-6	The energy-momentum tensor of the Standard-Model fermions	every Standard-Model fermion; metric g_{MN}; gauge fields	8
FE-7	The energy-momentum tensor of the gauge fields	gluons; $SU(2)$ fields; hypercharge field; metric g_{MN}	8
FE-8	The energy-momentum tensor of the tip	sigma field Σ; $SU(2)$ fields; hypercharge field; every Standard-Model fermion; metric g_{MN}	4 (part of the rest energies)
FE-9	The energy-momentum tensors of the portals I11 and I13	dirac16complex Ψ; fermions; metric g_{MN}	8
FE-10	The energy-momentum tensor of the portal I12	dirac16complex Ψ; gauge fields; metric g_{MN}	8
FE-11	Conservation of the source	all fields	1
FE-12	The dirac16complex field equation with the canonical spin connection and the portals	dirac16complex Ψ; metric g_{MN}; gauge fields (I12); fermion currents (I11, I13)	2 (spectrum), 3 (Kohn-Sham)
FE-13	The canonical spin connection on the order-0 metric	dirac16complex Ψ; every fermion; metric g_{MN}	1 (check), 3
FE-14	The Dirac equation of the quark doublets Q_i	quark doublets; gauge fields; metric g_{MN}; Ψ (I11, I13); Σ	2
FE-15	The Dirac equation of the up-type quark singlets u_{Ri} (right-handed up, charm and top quarks)	up-type quark singlets; gauge fields; metric g_{MN}; Ψ; Σ	2
FE-16	The Dirac equation of the down-type quark singlets d_{Ri} (right-handed down, strange and bottom quarks)	down-type quark singlets; gauge fields; metric g_{MN}; Ψ; Σ	2

equation	field equation	fields	solved (used) in step
FE-17	The Dirac equation of the lepton doublets (left-handed neutrinos and charged leptons of three flavours)	lepton doublets; gauge fields; metric g_{MN}; Ψ (I11); Σ	2
FE-18	The Dirac equation of the right-handed charged leptons e_{Ri} (electron, muon and tau)	right-handed charged leptons; gauge fields; metric g_{MN}; Ψ; Σ	2
FE-19	The Dirac equation of the right-handed neutrinos ν_{Ri}	right-handed neutrinos; metric g_{MN}; Ψ (I11); Σ	2
FE-20	The fermion currents of the gauge fields	every Standard-Model fermion; gauge fields	4
FE-21	The field-dependent gauge couplings produced by I12	gauge fields; dirac16complex Ψ	8
FE-22	The Yang-Mills equation of the gluons	gluons; quarks; Ψ (I12); metric g_{MN}	2 (massless gluons), 4
FE-23	The Yang-Mills equation of the SU(2) field W^I (matrix form) with the tip current	$SU(2)$ fields; hypercharge field; lepton and quark doublets; Σ; Ψ (I12); metric g_{MN}	2 (masses), 4
FE-24	The Yang-Mills equation of the hypercharge field B	hypercharge field; W^3; every fermion; Σ; Ψ (I12); metric g_{MN}	2 (masses), 4
FE-25	The W^+, W^-, Z and photon field equations to linear order in the fields (I12 neglected)	W^+; W^-; Z; photon; every fermion; Σ	2, 4
FE-26	The tip sigma field equation (the Noether identity of the gauge symmetry)	sigma field Σ; $SU(2)$ fields; hypercharge field; fermions at the tip	4
FE-27	The boundary conditions of gravity and the gauge fields at the fold and the tip	metric g_{MN}; gauge fields	2
FE-28	The boundary conditions of dirac16complex	dirac16complex Ψ	2, 3
FE-29	The boundary conditions of the Standard-Model fermions	every Standard-Model fermion	2
FE-30	The tip mixing conditions of one and of three generations	every Standard-Model fermion; Σ	2
FE-31	The order-0 metric in the author's coordinates	metric g_{MN}	1
FE-32	The hidden coordinate y, the block scale and the rates of the background	metric g_{MN}	1
FE-33	The sectional curvatures of the order-0 metric	metric g_{MN}	1
FE-34	The Lovelock densities and tensors of a pair-diagonal curvature	metric g_{MN}	1
FE-35	The first Lovelock tensor (the Einstein tensor) of the order-0 metric	metric g_{MN}	1
FE-36	The second Lovelock tensor (Gauss-Bonnet) of the order-0 metric	metric g_{MN}	1
FE-37	The third Lovelock tensor of the order-0 metric	metric g_{MN}	1
FE-38	The background equation: the x_4 component (the constraint)	metric g_{MN} (a_4, b_4); all sources (block averages)	1
FE-39	The background equation: the 3-space component	metric g_{MN}; all sources	1

equation	field equation	fields	solved (used) in step
FE-40	The background equation: the hidden (block) component	metric g_{MN}; all sources	1
FE-41	The conservation of the order-0 source	all sources	1
FE-42	The static block	metric g_{MN}; sources	1
FE-43	The stabilised block	metric g_{MN} (block scale); stabiliser	1
FE-44	The linearised block and the radion	metric g_{MN} (block scale)	1, 10
FE-45	The response of the block to matter to first order	metric g_{MN} (block); all sources	1
FE-46	The coefficients of the first-order block response	metric g_{MN} (block)	1
FE-47	The adiabatic block response and the four-dimensional Friedmann acceleration equation	metric g_{MN} (block); all sources	1, 3, 8
FE-48	The four-dimensional reduction of the Lovelock action on the static block	metric g_{MN}	1
FE-49	The four-dimensional effective gravity and the four-dimensional couplings	four-dimensional metric; all fields	1, 6
FE-50	The x_0 eigenvalue equation of the gauge fields and the graviton	gluons; photon; W^+; W^-; Z; graviton (transverse-traceless)	2, 10
FE-51	The fold and tip conditions of the boson x_0 profiles	gluons; photon; W^+; W^-; Z; graviton	2, 10
FE-52	The boson eigenvalue conditions	gauge fields; graviton	2
FE-53	The tower of Kaluza-Klein gravitons (the terms $n \geq 1$ of the metric series)	graviton tower	2 (first level), 10
FE-54	The potential between two static masses with the Kaluza-Klein gravitons	graviton tower; sources with x_0 profiles	10
FE-55	The $W^\pm$ and Z masses, the electroweak parameter and the Kaluza-Klein scale	W^+; W^-; Z	2
FE-56	The exact reduction of the eight-dimensional Dirac equation to the block Hamiltonian	dirac16complex Ψ; every Standard-Model fermion; metric g_{MN}	2, 3
FE-57	The chiral zero modes (massless four-dimensional fermions)	every Standard-Model fermion; dirac16complex Ψ (fold band)	2
FE-58	The tip values of the doublet and singlet solutions at $k = 0$	every Standard-Model fermion	2
FE-59	The exact mass condition of one generation and its first-order root	every Standard-Model fermion	2
FE-60	The mass condition of three generations with the 3×3 tip matrices	quarks and leptons of three generations	2
FE-61	The standard parametrisation of the mixing matrices (the tip matrices of UM-2)	neutrinos; quarks	2, 5
FE-62	The spectrum of dirac16complex on the static block	dirac16complex Ψ	2
FE-63	The Kohn-Sham equations of dirac16complex coupled to all fields	dirac16complex Ψ; metric g_{MN}; all Standard-Model fields (mean fields)	3, 8
FE-64	The Kohn-Sham mean fields	dirac16complex Ψ; fermions (I11, I13); gauge condensates (I12)	3, 8
FE-65	The Kohn-Sham densities, shells, cell volume and occupations	dirac16complex Ψ	3

equation	field equation	fields	solved (used) in step
FE-66	The Kohn-Sham energy	dirac16complex Ψ	3
FE-67	The energy-momentum tensor of a Kohn-Sham state	dirac16complex Ψ; metric g_{MN} (source)	3
FE-68	The energy identity and the y-conservation law of a Kohn-Sham state	dirac16complex Ψ	3
FE-69	The boundary conditions of the Kohn-Sham orbitals	dirac16complex Ψ	3
FE-70	The portals I11 and I13 on the zero modes: the four-dimensional contact interactions	dirac16complex Ψ; every Standard-Model fermion	8, 9
FE-71	The portal I12 on the dark matter	dark matter; gluons; $SU(2)$ fields; hypercharge field	3, 7, 8
FE-72	The shift of a Kohn-Sham level by the mean fields of an environment	dirac16complex Ψ; every Standard-Model field (densities, condensates)	3, 8
FE-73	The level shift of the dark matter in matter of protons, neutrons and electrons	dark matter; quarks; electrons	8
FE-74	The thermal gluon condensate from the trace anomaly	gluons	8
FE-75	The QCD interaction measure	quark-gluon plasma	8
FE-76	The thermal condensate of a massive vector boson and the order-α_{G} condensate of a gauge plasma	$SU(2)$ fields; Z; gauge plasmas	8
FE-77	The running electroweak couplings without a Higgs particle	$SU(2)$ fields; hypercharge field	8, 9 (couplings at $2M_{\mathrm{DM}}$)
FE-78	The electroweak condensates of a hot plasma	$SU(2)$ fields; Z; hypercharge field	8
FE-79	dirac16complex acting on the Standard Model: the coupling shifts and the x_0 problem of $W^{\pm}$ and Z	$SU(2)$ fields; Z; photon; gluons; dark matter	8
FE-80	Gravity in the coupled solution: the block response to every species	metric g_{MN} (block); all species	8, 10
FE-81	The hidden pressure of matter and of the hot plasma	all species	8
FE-82	The block dependence of a gauge coupling and the trace of an electroweak plasma	gauge fields	8
FE-83	The block dependence of the QCD scale and of the nucleon mass	quarks; gluons; nucleons	8
FE-84	The block dependence of the QCD scale through the quark thresholds	gluons; quarks	8
FE-85	The block dependence of a fermion mass	every Standard-Model fermion	8
FE-86	The densities of the environments of the coupled solution	baryons; electrons; dark matter	8
FE-87	The metric as a power series of terms of decreasing magnitude	metric g_{MN}	10 (scheme)
FE-88	The order-1, $n = 0$ term of the metric in the Newtonian gauge (eight dimensions)	metric g_{MN}	6, 10
FE-89	The four-dimensional perturbed metric in conformal time	four-dimensional metric	6

equation	field equation	fields	solved (used) in step
FE-90	The linearised Einstein equations for a Fourier mode	four-dimensional metric; all species	6
FE-91	The perturbations of the dark matter (cold)	dark matter (the odd quanta of Ψ_A and Ψ_B)	6
FE-92	The perturbations of the baryons	baryons (protons, neutrons, electrons); photons	6
FE-93	The perturbations of the photons: density and velocity	photons; baryons	6
FE-94	The photon temperature hierarchy	photons	6
FE-95	The photon polarisation hierarchy	photons	6
FE-96	The hierarchy of each neutrino mass state (the cosmic neutrino background)	neutrino mass states ν_1, ν_2, ν_3; fold bands (massless limit)	6
FE-97	The background expansion, the neutrino temperature and the primordial spectrum	four-dimensional metric; all species	6
FE-98	The size of the first-order potentials: rms value, Poisson equation and large scales	four-dimensional metric; matter	6, 10
FE-99	The amplitude of the matter fluctuations	matter	6
FE-100	The three neutrino masses	neutrinos	5
FE-101	The four-fermion interactions at low energy	neutrinos; charged leptons; medium fermions	5
FE-102	The Fermi constant and the neutral-current couplings	$SU(2)$ fields; Z; fermions	4, 5
FE-103	The matter potentials of forward scattering	neutrinos; electrons; neutrons	5
FE-104	The flavour evolution in vacuum and in matter	neutrinos	5
FE-105	The vacuum oscillation probability, CPT and the T-odd difference	neutrinos	5
FE-106	The MSW resonances of the 1-3 and 1-2 sectors	neutrinos; electrons	5
FE-107	The adiabatic survival probability of solar electron neutrinos	neutrinos; electrons	5
FE-108	The resonance density, the adiabaticity and the Landau-Zener jump probability	neutrinos; electrons	5
FE-109	The thermal potential of the early Universe	neutrinos; electrons and positrons	5
FE-110	The quantum kinetic equations of a neutrino gas	neutrinos	5
FE-111	Neutrino decoupling: the temperature equations	photons; electrons and positrons; neutrinos	5
FE-112	The energy transfer by collisions	neutrinos; electrons and positrons	5
FE-113	The finite-temperature QED pressure of the electron-photon plasma	photons; electrons and positrons	5
FE-114	The integrals of the QED pressure	electrons and positrons	5
FE-115	The effective number of neutrino species and the common temperature	neutrinos; photons	5
FE-116	The fast flavour instability: the electron-lepton-number distribution	neutrinos (dense gas)	5
FE-117	The fast flavour instability: the dispersion relation	neutrinos (dense gas)	5

equation	field equation	fields	solved (used) in step
FE-118	The axisymmetric and azimuthal branches of the dispersion relation	neutrinos (dense gas)	5
FE-119	The homogeneous mode and the mode $k_{\mathrm{f}} = \mathcal{I}_1$	neutrinos (dense gas)	5
FE-120	The cosmic neutrino background	neutrino mass states	5
FE-121	The ideal-gas densities of a species of the plasma	every species in equilibrium	9
FE-122	The effective degrees of freedom and the Hubble rate of the hot Universe	plasma	9
FE-123	The QCD crossover, the neutrino temperature after decoupling and the cooling of the plasma	plasma	9
FE-124	The Boltzmann equation of one dark-matter species	dark matter; plasma	9 (7)
FE-125	The relativistic thermal average of the annihilation cross-section	dark matter	9 (7)
FE-126	The relic density	dark matter	9 (7)
FE-127	The annihilation cross-sections of the dark matter through I12 and I13	dark matter; gauge bosons; quarks	7, 9
FE-128	The partial-wave unitarity bounds	dark matter	7, 9
FE-129	The freeze-in of the fold bands through I11	fold bands of Ψ; Standard-Model plasma	9
FE-130	The dark radiation of the fold bands and their departure from equilibrium	fold bands	9
FE-131	The right-handed neutrinos if I13 coupled to $B - L$	right-handed neutrinos; dark matter	9
FE-132	The elastic scattering of right-handed neutrinos on the dark matter	right-handed neutrinos; dark matter	9
FE-133	The extra radiation of species that decouple early	decoupled Weyl species	9, 4
FE-134	The production of right-handed neutrinos through I11 (with the baryon current in I13)	right-handed neutrinos; dark matter	9
FE-135	The self-interaction channel of the dark matter into the fold bands	dark matter; fold bands	9
FE-136	The energy put into the fold bands by the self-interaction channel	dark matter; fold bands	9

3 The approximation scheme

The coupled field equations are solved in a hierarchy that uses the small parameters of the problem. Each step of the solution states its own approximation in its second part, with the size of the neglected terms; the five parameters that organise the whole solution are these.

The metric is a power series of terms of decreasing magnitude. Order 0 is the background (step 1); order 1 with $n = 0$ carries the CMB, the CNB and the BAO (step 6); the terms with $n \geq 1$ are the Kaluza-Klein gravitons, and order 2 is the first omitted order (step 10):

$$g_{MN} = \bar{g}_{MN} + \epsilon\, h^{(1)}_{MN} + \epsilon^2\, h^{(2)}_{MN} + \ldots, \qquad h^{(1)}_{MN}(X^\mu, x_0) = \sum_{n\geq 0} h^{(1,n)}_{MN}(X^\mu)\, F_n(x_0), \qquad \epsilon \sim 10^{-5} \qquad \text{(FE-87)}$$

Symbols in (FE-87).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M \mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\bar{g}_{MN}$: The order-0 (background) metric of the power series: the corrected primordial metric in the author's coordinates, determined by the two functions $a_4(x_4)$ and $b_4(x_4)$; $|\bar{g}|$ is the absolute value of its determinant. Unit: dimensionless. Status: AUTHOR.
- ϵ: The bookkeeping parameter of the power series of the metric: the term of order q is multiplied by ϵ^q (the size of the first-order potentials). Unit: dimensionless. Value: $\epsilon \sim 10^{-5}$. Status: DERIVED.
- $h^{(1)}_{MN}$, $h^{(2)}_{MN}$: The first- and second-order terms of the power series of the metric. Unit: dimensionless. Status: DERIVED.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- n: The level number of the Kaluza-Klein tower ($n = 1, 2, 3, \ldots$; $n = 0$ the flat zero mode). Status: DERIVED.
- $h^{(1,n)}_{MN}$: The four-dimensional coefficient of the n-th x_0 profile F_n in the first-order term ($n = 0$: the flat zero mode, the perturbations of four-dimensional gravity; $n \geq 1$: the Kaluza-Klein gravitons). Unit: dimensionless. Status: DERIVED.
- $F(\xi)$, F_n: The x_0 profile of a boson mode on one half of the period (a function of y or of $\xi = y/L$): of a gauge field (gauge $A_y = 0$) or of the transverse-traceless graviton; F_n is the profile of the n-th Kaluza-Klein level of the graviton tower ($F_0 = 1$ the flat zero mode). Unit: dimensionless. Status: DERIVED.

After the primordial era the matter sources are tiny compared with the stabiliser of the block. The constraint $3\mathcal{G}(K_*)\,\mathcal{H}^2 = \kappa\Delta\rho$ of the first-order block response (step 1) fixes their ratio by the expansion rate of 3-space; today the rate is the Hubble constant $\mathcal{H}_0$, and the ratio is 10^{-92}:

$$\frac{\kappa\Delta\rho}{\kappa\rho_{\mathrm{stab}}(f_*)} = \frac{3\mathcal{G}(K_*)\,\mathcal{H}^2}{\kappa\rho_{\mathrm{stab}}(f_*)} = 0.7592\,\frac{\mathcal{H}^2}{H^2}, \qquad \frac{\mathcal{H}_0^2}{H^2} = 1.0\times 10^{-92}$$

Symbols in this equation.

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_{\mathrm{N}}\mathcal{G}(K_*) = 2.328\times 10^{-36}\ \mathrm{GeV}^{-2}$. Status: AUTHOR.
- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m^2_{21} = m^2_2 - m^2_1$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.

- ρ_{stab}: The energy density of the block stabiliser, a function of the block scale, $\rho_{\mathrm{stab}} \propto f^{-8}$; its value at the static block is $\rho_{\mathrm{stab}}(f_*)$. Unit: GeV^8. Value: $\kappa\rho_{\mathrm{stab}}(f_*) = (26 + 4\sqrt{51})H^2 = 54.566\,H^2$. Status: ASSUMED.
- f_*: The block scale of the stabilised (static) block, $f_* = K_*^{-1/2}$. Unit: dimensionless. Value: 0.215556. Status: DERIVED.
- $\mathcal{G}(K)$: The coefficient of the four-dimensional Einstein-Hilbert term on a static block, $\mathcal{G}(K) = 1 - 24w_{(2)}K + 72w_{(3)}K^2 = (K^2 - 12K + 16)/16$; four-dimensional gravity is attractive for $\mathcal{G} > 0$. Unit: dimensionless. Value: $\mathcal{G}(K_*) = 20/3 + \sqrt{51} = 13.8081$. Status: DERIVED.
- K_*: The block variable of the stabilised block, $K_* = 12 + \frac{4}{3}\sqrt{51}$ (the upper root K_+). Unit: dimensionless. Value: 21.5219. Status: DERIVED.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_{\mathrm{Pl}}$). Unit: GeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- $\mathcal{H}_0$: The Hubble rate today (the Hubble constant). Unit: GeV (km/s/Mpc). Value: 67.38 km/s/Mpc. Status: DERIVED.

so the block responds linearly and adiabatically to every source, and the block scale moves by at most $|\Delta \ln f| = 1.1 \times 10^{-34}$ in any environment of the model (step 8).

The Standard-Model masses lie far below the gap of the x_0 operators. The lightest massive Kaluza-Klein state has the mass $3H/(2f_*) = 99.48$ TeV (the gap) and the first Kaluza-Klein level $m_{\mathrm{KK}} = 121.32$ TeV, while the heaviest Standard-Model particle, the top quark, has 172.56 GeV; the four-dimensional fields are the zero modes of the x_0 operators, and the exchange of Kaluza-Klein modes corrects the four-dimensional interactions at the relative order

$$\left(\frac{m_{\mathrm{W}}}{m_{\mathrm{KK}}}\right)^2 = 4.39 \times 10^{-7}, \qquad \left(\frac{2f_*\, m_{\mathrm{t}}}{3H}\right)^2 = 3.0 \times 10^{-6}$$

Symbols in this equation.

- m_{W}: The mass of W^+ and W^-. Unit: GeV. Value: 80.3625 GeV (measured, input). Status: STANDARD.
- m_n^{KK}, m_{KK}: The mass of the n-th Kaluza-Klein graviton (and gauge) level, $m_n^{\mathrm{KK}} = \tilde{m}_n/(fL)$; $m_{\mathrm{KK}} = m_1^{\mathrm{KK}}$ is the Kaluza-Klein scale. Unit: GeV. Value: $m_{\mathrm{KK}} = 121.32$ TeV. Status: DERIVED.
- f_*: The block scale of the stabilised (static) block, $f_* = K_*^{-1/2}$. Unit: dimensionless. Value: 0.215556. Status: DERIVED.
- m_{u}, m_{d}, m_{s}, m_{c}, m_{b}, m_{t}: The masses of the up, down, strange, charm, bottom and top quarks (the upright subscript names the quark). Unit: GeV. Value: 2.16 MeV, 4.70 MeV, 93.5 MeV, 1.273 GeV, 4.183 GeV, 172.56 GeV (the x_0 masses, fitted to the PDG values). Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.

(step 2; the largest off-diagonal coupling of the first Kaluza-Klein gauge mode in the mass basis is 1.0×10^{-17}).

dirac16complex is treated by Kohn-Sham theory on the static block. Its many-body state is described by Kohn-Sham orbitals in the local mean fields of its contact self-interaction and of the portals I11, I12 and I13, through which the Standard-Model fields enter with their densities and condensates (steps 3 and 8). The portal mean fields are external fields of first order: the dark-matter level moves by at most

$$\frac{|\Delta\varepsilon_{\mathrm{DM}}|}{M_{\mathrm{DM}}} \leq 1.5 \times 10^{-6}$$

Symbols in this equation.

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty

relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.

- $\varepsilon_{\mathrm{DM}}$: The energy of the dark-matter level at rest (its mass M_{DM} on the static block). Unit: GeV. Value: $3.102204\,m_\Psi = 44.35$ TeV. Status: DERIVED.
- M_{DM}: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.

(at the dark-matter freeze-out; below 10^{-15} in every cold environment), and the response of a level to them is screened by the self-interaction by only 3.5×10^{-4} (step 8).

The neutrinos are ultra-relativistic and their flavour evolution is coherent. For every source of UM-2 the energy exceeds the masses by many orders ($m_3/E = 1.3 \times 10^{-8}$ for reactor antineutrinos of 4 MeV), and outside the early Universe the collisions are negligible against the forward scattering (step 5).

The hierarchy of the solution follows from these parameters: (1) the background and the primordial era; (2) the x_0 spectra, i.e. all masses and the mixing matrices; (3) the Kohn-Sham states of dirac16complex coupled to gravity and to the Standard-Model fields, and their back-reaction on the block; (4) the electroweak sector; (5) the flavour conversion of the three neutrinos, N_{eff} and the CNB; (6) the linear cosmology (CMB, CNB, BAO); (7) the dark-matter relic through I12 and its detection; (8) the coupled solution in environments, iterated to self-consistency; (9) the thermal history and the relic with I13; (10) the magnitudes of the terms of the metric series. Each step uses the results of the earlier ones, and its check in concert returns its results to the equations of the other steps.

4 Step 1: the background and the primordial era

4.1 The field equations of step 1

The gravitational field is the metric g_{MN} of the eight coordinates. Its action is the Einstein-Lovelock action with the author's weights, and the block of the hidden direction and the extra times is held by a stabiliser:

$$S_{\mathrm{EL}} = \frac{1}{2\kappa}\int \mathrm{d}^8X\,\sqrt{|g|}\,\Big(\sum_{p=1}^{3}\alpha_{(p)}\,\mathcal{L}_{(p)} - 2\Lambda\Big), \qquad \kappa = 8\pi G_8,$$
$$\alpha_{(p)} = w_{(p)}\,H^{2-2p}, \qquad w_{(1)} = 1, \quad w_{(2)} = \frac{1}{32}, \quad w_{(3)} = \frac{1}{1152}, \qquad \Lambda = -10H^2 \tag{IN-2}$$

Symbols in (IN-2).

- S_{EL}: The Einstein-Lovelock action of gravity. Unit: dimensionless. Status: AUTHOR.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_{\mathrm{N}}\mathcal{G}(K_*) = 2.328 \times 10^{-36}\ \mathrm{GeV}^{-2}$. Status: AUTHOR.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic). Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)}H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- $\mathcal{L}_{(p)}$: The Lovelock density of order p of the metric (a scalar built from p Riemann tensors): $\mathcal{L}_{(1)} = R$ (Einstein-Hilbert), $\mathcal{L}_{(2)}$ the Gauss-Bonnet density, $\mathcal{L}_{(3)}$ the cubic Lovelock density. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -10H^2$. Unit: GeV^2. Value: $-10H^2$. Status: AUTHOR.

- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_8: The eight-dimensional Newton constant ($\kappa = 8\pi G_8$). Unit: GeV^{-6}. Status: AUTHOR.
- $w_{(p)}$: The author's Lovelock weights, $w_{(1)} = 1$, $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$. Unit: dimensionless. Value: $(1, 1/32, 1/1152)$. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of $m_{\rm W}$ and $\rho_{\rm EW,0}$). Status: AUTHOR.

$$\begin{aligned} &\mathcal{L}_{(p)} = \frac{1}{2^p}\,\delta^{M_1N_1\cdots M_pN_p}_{P_1Q_1\cdots P_pQ_p}\,R^{P_1Q_1}{}_{M_1N_1}\cdots R^{P_pQ_p}{}_{M_pN_p}, \\ &\mathcal{L}_{(1)} = R, \qquad \mathcal{L}_{(2)} = R^2 - 4R_{MN}R^{MN} + R_{MNPQ}R^{MNPQ} \end{aligned} \tag{IN-3}$$

Symbols in (IN-3).

- $\mathcal{L}_{(p)}$: The Lovelock density of order p of the metric (a scalar built from p Riemann tensors): $\mathcal{L}_{(1)} = R$ (Einstein-Hilbert), $\mathcal{L}_{(2)}$ the Gauss-Bonnet density, $\mathcal{L}_{(3)}$ the cubic Lovelock density. Unit: 1/length^{2p}. Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic). Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{\alpha e}$, δ_{l0}, δ^{I3}); with upper and lower index strings the generalised Kronecker delta $\delta^{M_1\cdots M_n}_{N_1\cdots N_n}$, the determinant of the $n \times n$ matrix of Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- P: A coordinate index of eight dimensions (also $P_1, P_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Q: A coordinate index of eight dimensions (also $Q_1, Q_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $R^{MN}{}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor, with two indices the Ricci tensor $R_{MN} = R^P{}_{MPN}$, without indices the Ricci (scalar) curvature $R = g^{MN}R_{MN}$; $R^{ab}{}_{ab}$ (no sum) is the sectional curvature of the coordinate plane (a, b). Unit: 1/length^2. Status: STANDARD.

$$S_{\rm stab} = -\int \mathrm{d}^8X\,\sqrt{|g|}\,\rho_{\rm stab}(f), \qquad \kappa\rho_{\rm stab}(f) = \kappa\rho_{\rm stab}(f_*)\Big(\frac{f_*}{f}\Big)^8, \qquad f = \mathrm{e}^{-a_4}\,b_4^{1/2} \tag{IN-4}$$

Symbols in (IN-4).

- $S_{\rm stab}$: The action of the block stabiliser (a vacuum energy of the block). Unit: dimensionless. Status: ASSUMED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- $\rho_{\rm stab}$: The energy density of the block stabiliser, a function of the block scale, $\rho_{\rm stab} \propto f^{-8}$; its value at the static block is $\rho_{\rm stab}(f_*)$. Unit: GeV^8. Value: $\kappa\rho_{\rm stab}(f_*) = (26 + 4\sqrt{51})H^2 = 54.566\,H^2$. Status: ASSUMED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4}b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_\mathrm{N}\mathcal{G}(K_*) = 2.328 \times 10^{-36}\ \mathrm{GeV}^{-2}$. Status: AUTHOR.
- f_*: The block scale of the stabilised (static) block, $f_* = K_*^{-1/2}$. Unit: dimensionless. Value: 0.215556. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the hidden and extra-time components, $b_4 > 0$. Unit: dimensionless. Status: AUTHOR.

Varying the action with respect to the metric gives the Einstein-Lovelock equations, sourced by the energy-momentum tensor of all fields:

$$\begin{aligned} &\sum_{p=1}^{3} \alpha_{(p)}\, \mathcal{E}_{(p)}{}^{M}{}_{N} + \Lambda\, \delta^{M}_{N} = \kappa\, \mathcal{T}^{M}{}_{N}, \\ &\mathcal{E}_{(p)}{}^{M}{}_{N} = -\frac{1}{2^{p+1}}\, \delta^{M M_1 N_1 \cdots M_p N_p}_{N P_1 Q_1 \cdots P_p Q_p}\, R^{P_1 Q_1}{}_{M_1 N_1} \cdots R^{P_p Q_p}{}_{M_p N_p} \end{aligned} \tag{FE-1}$$

Symbols in (FE-1).

- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic). Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)} H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- $\mathcal{E}_{(p)}{}^{M}{}_{N}$: The Lovelock tensor of order p, the metric variation of $\sqrt{|g|}\mathcal{L}_{(p)}$: $\mathcal{E}_{(1)}$ is the Einstein tensor $R^{M}{}_{N} - \frac{1}{2}\delta^{M}_{N} R$; every $\mathcal{E}_{(p)}$ is divergence free. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -10H^2$. Unit: GeV^2. Value: $-10H^2$. Status: AUTHOR.
- δ: The Kronecker delta: with two indices $\delta^{M}_{N} = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{\alpha e}$, δ_{l0}, δ^{I3}); with upper and lower index strings the generalised Kronecker delta $\delta^{M_1 \cdots M_n}_{N_1 \cdots N_n}$, the determinant of the $n \times n$ matrix of Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_\mathrm{N}\mathcal{G}(K_*) = 2.328 \times 10^{-36}\ \mathrm{GeV}^{-2}$. Status: AUTHOR.
- $\mathcal{T}^{M}{}_{N}$: The energy-momentum tensor of all matter (every field except gravity, the stabiliser included), the source of gravity; defined by the change of the matter action under a change of the metric. Unit: GeV^8. Status: STANDARD.
- P: A coordinate index of eight dimensions (also $P_1, P_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Q: A coordinate index of eight dimensions (also $Q_1, Q_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $R^{MN}{}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor, with two indices the Ricci tensor $R_{MN} = R^{P}{}_{MPN}$, without indices the Ricci (scalar) curvature $R = g^{MN}R_{MN}$; $R^{ab}{}_{ab}$ (no sum) is the sectional curvature of the coordinate plane (a, b). Unit: $1/\mathrm{length}^2$. Status: STANDARD.

$$\begin{aligned} &\Delta S_\mathrm{m} = -\frac{1}{2}\int \mathrm{d}^8X\, \sqrt{|g|}\, \mathcal{T}_{MN}\, \Delta g^{MN}, \qquad \rho = -\mathcal{T}^{x_4}{}_{x_4}, \qquad p_M = \mathcal{T}^{M}{}_{M} \quad (M \neq x_4, \text{ no sum}), \\ &\mathcal{T}^{M}{}_{N} = \mathrm{diag}\big(p_0,\ p_3,\ p_3,\ p_3,\ -\rho,\ p_5,\ p_5,\ p_5\big) \quad (\text{a diagonal source}), \qquad \bar{p} = \frac{p_0 + 3p_5}{4} \end{aligned} \tag{FE-2}$$

Symbols in (FE-2).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- S_{m}: The matter action: all terms of the action except S_{EL} (the stabiliser included). Unit: dimensionless. Status: DERIVED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, ${g^M}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- ${\mathcal{T}^M}_N$: The energy-momentum tensor of all matter (every field except gravity, the stabiliser included), the source of gravity; defined by the change of the matter action under a change of the metric. Unit: GeV^8. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- ρ: An energy density: in eight dimensions $\rho = -{\mathcal{T}^{x_4}}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- p: A pressure: $p_M = {\mathcal{T}^M}_M$ (no sum) for the direction $M \neq x_4$, p_s the pressure of the species s; Δp the pressure perturbation in the cosmological perturbation equations; in the Feynman-Hellmann relation the pressure of a gauge plasma. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ when it is a tensor in eight dimensions). Status: STANDARD.
- p_0: The pressure along the hidden direction x_0, $p_0 = {\mathcal{T}^{x_0}}_{x_0}$. Unit: GeV^8. Status: STANDARD.
- p_3: The pressure of ordinary 3-space, $p_3 = {\mathcal{T}^{x_1}}_{x_1} = {\mathcal{T}^{x_2}}_{x_2} = {\mathcal{T}^{x_3}}_{x_3}$ for an isotropic source. Unit: GeV^8. Status: STANDARD.
- p_5: The pressure along the extra times, $p_5 = {\mathcal{T}^{x_5}}_{x_5} = {\mathcal{T}^{x_6}}_{x_6} = {\mathcal{T}^{x_7}}_{x_7}$. Unit: GeV^8. Status: STANDARD.
- $\bar{p}$: The hidden pressure, the pressure averaged over the block, $\bar{p} = (p_0 + 3p_5)/4$; for a species at fixed temperature $\bar{p} = \frac{1}{4}\,\partial p/\partial \ln f$. Unit: GeV^8 (or GeV^4 for a four-dimensional density). Status: DERIVED.

$$ {\mathcal{T}^M}_N = {\mathcal{T}_{\mathrm{stab}}}^M{}_N + {\mathcal{T}_\Psi}^M{}_N + {\mathcal{T}_{\mathrm{gauge}}}^M{}_N + {\mathcal{T}_{\mathrm{f}}}^M{}_N + {\mathcal{T}_{\mathrm{tip}}}^M{}_N + {\mathcal{T}_{\mathrm{I11}}}^M{}_N + {\mathcal{T}_{\mathrm{I12}}}^M{}_N + {\mathcal{T}_{\mathrm{I13}}}^M{}_N \qquad \text{(FE-3)} $$

Symbols in (FE-3).

- ${\mathcal{T}^M}_N$: The energy-momentum tensor of all matter (every field except gravity, the stabiliser included), the source of gravity; defined by the change of the matter action under a change of the metric. Unit: GeV^8. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\mathcal{T}_{\mathrm{stab}}$: The energy-momentum tensor of the stabiliser. Unit: GeV^8. Status: DERIVED.
- $\mathcal{T}_\Psi$: The energy-momentum tensor of the dirac16complex field. Unit: GeV^8. Status: DERIVED.
- $\mathcal{T}_{\mathrm{gauge}}$: The energy-momentum tensor of the gauge fields. Unit: GeV^8. Status: STANDARD.
- $\mathcal{T}_{\mathrm{f}}$: The energy-momentum tensor of the Standard-Model fermions. Unit: GeV^8. Status: DERIVED.

- $\mathcal{T}_{\text{tip}}$: The energy-momentum tensor of the tip terms (non-zero only for the tip directions). Unit: GeV^8. Status: DERIVED.
- $\mathcal{T}_{\text{I11}}$: The energy-momentum tensor of the portal I11. Unit: GeV^8. Status: DERIVED.
- $\mathcal{T}_{\text{I12}}$: The energy-momentum tensor of the portal I12. Unit: GeV^8. Status: DERIVED.
- $\mathcal{T}_{\text{I13}}$: The energy-momentum tensor of the portal I13. Unit: GeV^8. Status: DERIVED.

At order 0 the source is dominated by the stabiliser, whose energy-momentum tensor is exact:

$$\mathcal{T}_{\text{stab}}{}^{M}{}_{N} = \rho_{\text{stab}}\,\text{diag}(1,\ -1,\ -1,\ -1,\ -1,\ 1,\ 1,\ 1)\,: \qquad \rho = \rho_{\text{stab}}, \qquad p_3 = -\rho_{\text{stab}}, \qquad p_0 = p_5 = \rho_{\text{stab}} \tag{FE-4}$$

Symbols in (FE-4).

- $\mathcal{T}_{\text{stab}}$: The energy-momentum tensor of the stabiliser. Unit: GeV^8. Status: DERIVED.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- ρ_{stab}: The energy density of the block stabiliser, a function of the block scale, $\rho_{\text{stab}} \propto f^{-8}$; its value at the static block is $\rho_{\text{stab}}(f_*)$. Unit: GeV^8. Value: $\kappa\rho_{\text{stab}}(f_*) = (26 + 4\sqrt{51})H^2 = 54.566\,H^2$. Status: ASSUMED.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ when it is a tensor in eight dimensions). Status: STANDARD.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- p_3: The pressure of ordinary 3-space, $p_3 = \mathcal{T}^{x_1}{}_{x_1} = \mathcal{T}^{x_2}{}_{x_2} = \mathcal{T}^{x_3}{}_{x_3}$ for an isotropic source. Unit: GeV^8. Status: STANDARD.
- p_0: The pressure along the hidden direction x_0, $p_0 = \mathcal{T}^{x_0}{}_{x_0}$. Unit: GeV^8. Status: STANDARD.
- p_5: The pressure along the extra times, $p_5 = \mathcal{T}^{x_5}{}_{x_5} = \mathcal{T}^{x_6}{}_{x_6} = \mathcal{T}^{x_7}{}_{x_7}$. Unit: GeV^8. Status: STANDARD.

The Lovelock tensors are divergence-free, so the source is conserved:

$$\nabla_M\,\mathcal{T}^{M}{}_{N} = 0, \qquad \text{because} \qquad \nabla_M\,\mathcal{E}_{(p)}{}^{M}{}_{N} = 0 \quad (p = 1, 2, 3) \tag{FE-11}$$

Symbols in (FE-11).

- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols $\Gamma^{N}{}_{MP}$); ∇_M acts on the index M, $\nabla^M = g^{MN}\nabla_N$. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\mathcal{T}^{M}{}_{N}$: The energy-momentum tensor of all matter (every field except gravity, the stabiliser included), the source of gravity; defined by the change of the matter action under a change of the metric. Unit: GeV^8. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\mathcal{E}_{(p)}{}^{M}{}_{N}$: The Lovelock tensor of order p, the metric variation of $\sqrt{|g|}\mathcal{L}_{(p)}$: $\mathcal{E}_{(1)}$ is the Einstein tensor $R^{M}{}_{N} - \frac{1}{2}\delta^{M}_{N}R$; every $\mathcal{E}_{(p)}$ is divergence free. Unit: 1/length^{2p}. Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic). Status: STANDARD.

At order 0 the metric is the corrected primordial metric of the record, with two functions $a_4(x_4)$ and $b_4(x_4)$ of the time. In the chart of the hidden coordinate y it is a warped product of an expanding 3-space with the block:

$$\begin{aligned}\bar{g} = \text{diag}\Big(&\mathrm{e}^{-2a_4}b_4\cot^2 z,\ \mathrm{e}^{2a_4},\ \mathrm{e}^{2a_4},\ \mathrm{e}^{2a_4},\ -1,\\ &-\mathrm{e}^{-2a_4}b_4(\sin^2 z)^{1/6},\ -\mathrm{e}^{-2a_4}b_4(\sin^2 z)^{1/6},\ -\mathrm{e}^{-2a_4}b_4(\sin^2 z)^{1/6}\Big), \qquad z = 6Hx_0\end{aligned} \tag{FE-31}$$

Symbols in (FE-31).

- $\bar{g}_{MN}$: The order-0 (background) metric of the power series: the corrected primordial metric in the author's coordinates, determined by the two functions $a_4(x_4)$ and $b_4(x_4)$; $|\bar{g}|$ is the absolute value of its determinant. Unit: dimensionless. Status: AUTHOR.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ when it is a tensor in eight dimensions). Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the hidden and extra-time components, $b_4 > 0$. Unit: dimensionless. Status: AUTHOR.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.

$$
\begin{aligned}
&y = \frac{\ln|\sin z|}{6H}, \qquad \mathrm{d}y = \cot z\,\mathrm{d}x_0, \qquad \xi = \frac{y}{L} \in [-1, 0],\\
&\bar{g}_{MN}\,\mathrm{d}X^M \mathrm{d}X^N = \mathrm{e}^{2a_4}\left(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\right) - \mathrm{d}x_4^2 + f^2\Big[\mathrm{d}y^2 - \mathrm{e}^{2Hy}\left(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\right)\Big],\\
&f = \mathrm{e}^{-a_4} b_4^{1/2}, \qquad K = \frac{1}{f^2}, \qquad \mathcal{H} = \dot{a}_4, \qquad \mathcal{H}_f = \frac{\dot{f}}{f} = \frac{\dot{b}_4}{2b_4} - \dot{a}_4, \qquad \dot{K} = -2K\mathcal{H}_f,\\
&\sqrt{|\bar{g}|} = \mathrm{e}^{3a_4} f^4 |\cot z| (\sin^2 z)^{1/4} = \mathrm{e}^{3a_4} f^4\, \mathrm{e}^{3Hy}\Big|\frac{\mathrm{d}y}{\mathrm{d}x_0}\Big|
\end{aligned}
\tag{FE-32}
$$

Symbols in (FE-32).

- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- ξ: The scaled hidden coordinate, $\xi = y/L$, running from -1 (tip) to 0 (fold) on each half of a period. Unit: dimensionless. Status: DERIVED.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- $\bar{g}_{MN}$: The order-0 (background) metric of the power series: the corrected primordial metric in the author's coordinates, determined by the two functions $a_4(x_4)$ and $b_4(x_4)$; $|\bar{g}|$ is the absolute value of its determinant. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space (also the symmetry axis of the fast-flavour analysis). Unit: length. Status: AUTHOR.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the hidden and extra-time components, $b_4 > 0$. Unit: dimensionless. Status: AUTHOR.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2} T^2/M_{\mathrm{Pl}}$). Unit: GeV. Status: DERIVED.
- $\dot{a}_4$: The time derivative $\mathrm{d}a_4/\mathrm{d}x_4$. Unit: GeV. Status: DERIVED.
- $\mathcal{H}_f$: The expansion rate of the block, $\mathcal{H}_f = \dot{f}/f = \dot{b}_4/(2b_4) - \dot{a}_4$ (negative when the extra times and x_0 deflate). Unit: GeV. Status: DERIVED.
- $\dot{f}$: The time derivative $\mathrm{d}f/\mathrm{d}x_4$ of the block scale. Unit: GeV. Status: DERIVED.
- $\dot{b}_4$: The time derivative $\mathrm{d}b_4/\mathrm{d}x_4$. Unit: GeV. Status: DERIVED.
- $\dot{K}$: The time derivative $\mathrm{d}K/\mathrm{d}x_4$. Unit: GeV. Status: DERIVED.

Its curvature is pair-diagonal and does not depend on x_0; the Lovelock densities and tensors are then sums of products of the sectional curvatures:

$$R^{x_1x_2}{}_{x_1x_2} = \mathcal{H}^2, \qquad R^{x_1x_4}{}_{x_1x_4} = \mathcal{H}^2 + \dot{\mathcal{H}}, \qquad R^{x_1x_0}{}_{x_1x_0} = \mathcal{H}\mathcal{H}_f, \qquad R^{x_4x_0}{}_{x_4x_0} = \mathcal{H}_f^2 + \dot{\mathcal{H}}_f,$$
$$R^{x_0x_5}{}_{x_0x_5} = \mathcal{H}_f^2 - H^2 K; \tag{FE-33}$$

the same for every pair of the same kind; all other $R^{ab}{}_{cd}$ with $\{a,b\} \neq \{c,d\}$ vanish

Symbols in (FE-33).

- $R^{MN}{}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor, with two indices the Ricci tensor $R_{MN} = R^P{}_{MPN}$, without indices the Ricci (scalar) curvature $R = g^{MN} R_{MN}$; $R^{ab}{}_{ab}$ (no sum) is the sectional curvature of the coordinate plane (a, b). Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2} T^2/M_{\mathrm{Pl}}$). Unit: GeV. Status: DERIVED.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\dot{\mathcal{H}}$: The time derivative $\mathrm{d}\mathcal{H}/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- $\mathcal{H}_f$: The expansion rate of the block, $\mathcal{H}_f = \dot{f}/f = \dot{b}_4/(2b_4) - \dot{a}_4$ (negative when the extra times and x_0 deflate). Unit: GeV. Status: DERIVED.
- $\dot{\mathcal{H}}_f$: The time derivative $\mathrm{d}\mathcal{H}_f/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.

$$
\begin{aligned}
&\mathcal{L}_{(1)} = 2\sum_{\text{pairs}} R^{ab}{}_{ab}, \qquad \mathcal{L}_{(2)} = 8\sum_{\text{2 disjoint pairs}}\prod_{\text{pairs}} R^{ab}{}_{ab}, \qquad \mathcal{L}_{(3)} = 48\sum_{\text{3 disjoint pairs}}\prod_{\text{pairs}} R^{ab}{}_{ab},\\
&\mathcal{E}_{(1)}{}^{c}{}_{c} = -\sum_{\text{pairs avoiding } c} R^{ab}{}_{ab}, \qquad \mathcal{E}_{(2)}{}^{c}{}_{c} = -4\sum_{\text{2 disjoint pairs avoiding } c}\prod_{\text{pairs}} R^{ab}{}_{ab},\\
&\mathcal{E}_{(3)}{}^{c}{}_{c} = -24\sum_{\text{3 disjoint pairs avoiding } c}\prod_{\text{pairs}} R^{ab}{}_{ab}
\end{aligned}
\tag{FE-34}
$$

Symbols in (FE-34).

- $\mathcal{L}_{(p)}$: The Lovelock density of order p of the metric (a scalar built from p Riemann tensors): $\mathcal{L}_{(1)} = R$ (Einstein-Hilbert), $\mathcal{L}_{(2)}$ the Gauss-Bonnet density, $\mathcal{L}_{(3)}$ the cubic Lovelock density. Unit: $1/\text{length}^{2p}$. Status: STANDARD.
- $R^{MN}{}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor, with two indices the Ricci tensor $R_{MN} = R^{P}{}_{MPN}$, without indices the Ricci (scalar) curvature $R = g^{MN}R_{MN}$; $R^{ab}{}_{ab}$ (no sum) is the sectional curvature of the coordinate plane (a, b). Unit: $1/\text{length}^2$. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$, aligned with $x_0, \ldots, x_7$; raised and lowered with η_{ab}; a repeated index is summed (except in a sectional curvature $R^{ab}{}_{ab}$). Status: STANDARD.
- b: A frame index running over $0, \ldots, 7$ (as a). Status: STANDARD.
- $\mathcal{E}_{(p)}{}^{M}{}_{N}$: The Lovelock tensor of order p, the metric variation of $\sqrt{|g|}\mathcal{L}_{(p)}$: $\mathcal{E}_{(1)}$ is the Einstein tensor $R^{M}{}_{N} - \frac{1}{2}\delta^{M}_{N}R$; every $\mathcal{E}_{(p)}$ is divergence free. Unit: $1/\text{length}^{2p}$. Status: STANDARD.
- c: A frame index running over $0, \ldots, 7$ (as a). Status: STANDARD.

$$
\begin{aligned}
\mathcal{E}_{(1)}{}^{x_4}{}_{x_4} &= -3\,\mathcal{H}^2 - 12\,\mathcal{H}\mathcal{H}_f - 6\,\mathcal{H}_f^2 + 6\,KH^2\\
\mathcal{E}_{(1)}{}^{x_1}{}_{x_1} &= -3\,\mathcal{H}^2 - 8\,\mathcal{H}\mathcal{H}_f - 10\,\mathcal{H}_f^2 - 2\,\dot{\mathcal{H}} - 4\,\dot{\mathcal{H}}_f + 6\,KH^2\\
\mathcal{E}_{(1)}{}^{x_0}{}_{x_0} &= -6\,\mathcal{H}^2 - 9\,\mathcal{H}\mathcal{H}_f - 6\,\mathcal{H}_f^2 - 3\,\dot{\mathcal{H}} - 3\,\dot{\mathcal{H}}_f + 3\,KH^2
\end{aligned}
\tag{FE-35}
$$

Symbols in (FE-35).

- $\mathcal{E}_{(p)}{}^{M}{}_{N}$: The Lovelock tensor of order p, the metric variation of $\sqrt{|g|}\mathcal{L}_{(p)}$: $\mathcal{E}_{(1)}$ is the Einstein tensor $R^{M}{}_{N} - \frac{1}{2}\delta^{M}_{N}R$; every $\mathcal{E}_{(p)}$ is divergence free. Unit: $1/\text{length}^{2p}$. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_{\mathrm{Pl}}$). Unit: GeV. Status: DERIVED.
- $\mathcal{H}_f$: The expansion rate of the block, $\mathcal{H}_f = \dot{f}/f = \dot{b}_4/(2b_4) - \dot{a}_4$ (negative when the extra times and x_0 deflate). Unit: GeV. Status: DERIVED.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- $\dot{\mathcal{H}}$: The time derivative $\mathrm{d}\mathcal{H}/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- $\dot{\mathcal{H}}_f$: The time derivative $\mathrm{d}\mathcal{H}_f/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.

$$
\begin{aligned}
\mathcal{E}_{(2)}{}^{x_4}{}_{x_4} =& -48\,\mathcal{H}^3\mathcal{H}_f - 216\,\mathcal{H}^2\mathcal{H}_f^2 - 144\,\mathcal{H}\mathcal{H}_f^3 - 12\,\mathcal{H}_f^4 + 72\,K\mathcal{H}^2H^2 \\
&+ 144\,K\mathcal{H}\mathcal{H}_fH^2 + 24\,K\mathcal{H}_f^2H^2 - 12\,K^2H^4 \\
\mathcal{E}_{(2)}{}^{x_1}{}_{x_1} =& -32\,\mathcal{H}^3\mathcal{H}_f - 136\,\mathcal{H}^2\mathcal{H}_f^2 - 192\,\mathcal{H}\mathcal{H}_f^3 - 60\,\mathcal{H}_f^4 \\
&- 32\,\dot{\mathcal{H}}\mathcal{H}\mathcal{H}_f - 48\,\dot{\mathcal{H}}\mathcal{H}_f^2 - 16\,\dot{\mathcal{H}}_f\mathcal{H}^2 - 96\,\dot{\mathcal{H}}_f\mathcal{H}\mathcal{H}_f \\
&- 48\,\dot{\mathcal{H}}_f\mathcal{H}_f^2 + 72\,K\mathcal{H}^2H^2 + 96\,K\mathcal{H}\mathcal{H}_fH^2 + 72\,K\mathcal{H}_f^2H^2 + 48\,\dot{\mathcal{H}}KH^2 \\
&+ 48\,\dot{\mathcal{H}}_fKH^2 - 12\,K^2H^4 \\
\mathcal{E}_{(2)}{}^{x_0}{}_{x_0} =& -12\,\mathcal{H}^4 - 108\,\mathcal{H}^3\mathcal{H}_f - 180\,\mathcal{H}^2\mathcal{H}_f^2 - 108\,\mathcal{H}\mathcal{H}_f^3 - 12\,\mathcal{H}_f^4 \\
&- 12\,\dot{\mathcal{H}}\mathcal{H}^2 - 72\,\dot{\mathcal{H}}\mathcal{H}\mathcal{H}_f - 36\,\dot{\mathcal{H}}\mathcal{H}_f^2 - 36\,\dot{\mathcal{H}}_f\mathcal{H}^2 \\
&- 72\,\dot{\mathcal{H}}_f\mathcal{H}\mathcal{H}_f - 12\,\dot{\mathcal{H}}_f\mathcal{H}_f^2 + 72\,K\mathcal{H}^2H^2 + 36\,K\mathcal{H}\mathcal{H}_fH^2 + 12\,K\mathcal{H}_f^2H^2 \\
&+ 36\,\dot{\mathcal{H}}KH^2 + 12\,\dot{\mathcal{H}}_fKH^2
\end{aligned}
\tag{FE-36}
$$

Symbols in (FE-36).

- $\mathcal{E}_{(p)}{}^{M}{}_{N}$: The Lovelock tensor of order p, the metric variation of $\sqrt{|g|}\mathcal{L}_{(p)}$: $\mathcal{E}_{(1)}$ is the Einstein tensor $R^{M}{}_{N} - \frac{1}{2}\delta^{M}_{N}R$; every $\mathcal{E}_{(p)}$ is divergence free. Unit: $1/\text{length}^{2p}$. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_{\rm Pl}$). Unit: GeV. Status: DERIVED.
- $\mathcal{H}_f$: The expansion rate of the block, $\mathcal{H}_f = \dot{f}/f = \dot{b}_4/(2b_4) - \dot{a}_4$ (negative when the extra times and x_0 deflate). Unit: GeV. Status: DERIVED.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of $m_{\rm W}$ and $\rho_{\rm EW,0}$). Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- $\dot{\mathcal{H}}$: The time derivative $\mathrm{d}\mathcal{H}/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- $\dot{\mathcal{H}}_f$: The time derivative $\mathrm{d}\mathcal{H}_f/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.

$$
\begin{aligned}
\mathcal{E}_{(3)}{}^{x_4}{}_{x_4} =& -1440\,\mathcal{H}^3\mathcal{H}_f^3 - 1080\,\mathcal{H}^2\mathcal{H}_f^4 + 864\,K\mathcal{H}^3\mathcal{H}_fH^2 + 1296\,K\mathcal{H}^2\mathcal{H}_f^2H^2 \\
&- 216\,K^2\mathcal{H}^2H^4 \\
\mathcal{E}_{(3)}{}^{x_1}{}_{x_1} =& -576\,\mathcal{H}^3\mathcal{H}_f^3 - 1368\,\mathcal{H}^2\mathcal{H}_f^4 - 576\,\mathcal{H}\mathcal{H}_f^5 - 576\,\dot{\mathcal{H}}\mathcal{H}\mathcal{H}_f^3 \\
&- 144\,\dot{\mathcal{H}}\mathcal{H}_f^4 - 864\,\dot{\mathcal{H}}_f\mathcal{H}^2\mathcal{H}_f^2 - 576\,\dot{\mathcal{H}}_f\mathcal{H}\mathcal{H}_f^3 + 576\,K\mathcal{H}^3\mathcal{H}_fH^2 \\
&+ 1008\,K\mathcal{H}^2\mathcal{H}_f^2H^2 + 576\,K\mathcal{H}\mathcal{H}_f^3H^2 + 576\,\dot{\mathcal{H}}K\mathcal{H}\mathcal{H}_fH^2 \\
&+ 288\,\dot{\mathcal{H}}K\mathcal{H}_f^2H^2 + 288\,\dot{\mathcal{H}}_fK\mathcal{H}^2H^2 + 576\,\dot{\mathcal{H}}_fK\mathcal{H}\mathcal{H}_fH^2 - 216\,K^2\mathcal{H}^2H^4 \\
&- 144\,\dot{\mathcal{H}}K^2H^4 \\
\mathcal{E}_{(3)}{}^{x_0}{}_{x_0} =& -648\,\mathcal{H}^4\mathcal{H}_f^2 - 1224\,\mathcal{H}^3\mathcal{H}_f^3 - 648\,\mathcal{H}^2\mathcal{H}_f^4 - 648\,\dot{\mathcal{H}}\mathcal{H}^2\mathcal{H}_f^2 \\
&- 432\,\dot{\mathcal{H}}\mathcal{H}\mathcal{H}_f^3 - 432\,\dot{\mathcal{H}}_f\mathcal{H}^3\mathcal{H}_f - 648\,\dot{\mathcal{H}}_f\mathcal{H}^2\mathcal{H}_f^2 + 216\,K\mathcal{H}^4H^2 \\
&+ 648\,K\mathcal{H}^3\mathcal{H}_fH^2 + 216\,K\mathcal{H}^2\mathcal{H}_f^2H^2 + 216\,\dot{\mathcal{H}}K\mathcal{H}^2H^2 + 432\,\dot{\mathcal{H}}K\mathcal{H}\mathcal{H}_fH^2 \\
&+ 216\,\dot{\mathcal{H}}_fK\mathcal{H}^2H^2
\end{aligned}
\tag{FE-37}
$$

Symbols in (FE-37).

- $\mathcal{E}_{(p)}{}^{M}{}_{N}$: The Lovelock tensor of order p, the metric variation of $\sqrt{|g|}\mathcal{L}_{(p)}$: $\mathcal{E}_{(1)}$ is the Einstein tensor $R^{M}{}_{N} - \frac{1}{2}\delta^{M}_{N}R$; every $\mathcal{E}_{(p)}$ is divergence free. Unit: $1/\text{length}^{2p}$. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_{\rm Pl}$). Unit: GeV. Status: DERIVED.

- $\mathcal{H}_f$: The expansion rate of the block, $\mathcal{H}_f = \dot{f}/f = \dot{b}_4/(2b_4) - \dot{a}_4$ (negative when the extra times and x_0 deflate). Unit: GeV. Status: DERIVED.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- $\dot{\mathcal{H}}$: The time derivative $\mathrm{d}\mathcal{H}/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- $\dot{\mathcal{H}}_f$: The time derivative $\mathrm{d}\mathcal{H}_f/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.

With the author's weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ and $\Lambda = -10H^2$, the field equations of the order-0 metric are three ordinary differential equations: the x_4 component is a constraint, and the 3-space and hidden components are linear in $\dot{\mathcal{H}}$ and $\dot{\mathcal{H}}_f$ (the extra-time components x_5, x_6, x_7 have the same left side as the hidden one). These are the equations that step 1 solves:

$$
\begin{aligned}
x_4: \quad & -\frac{5\,\mathcal{H}^3\mathcal{H}_f^3}{4H^4} - \frac{15\,\mathcal{H}^2\mathcal{H}_f^4}{16H^4} + \frac{3\,K\mathcal{H}^3\mathcal{H}_f}{4H^2} \\
& + \frac{9\,K\mathcal{H}^2\mathcal{H}_f^2}{8H^2} - \frac{3\,\mathcal{H}^3\mathcal{H}_f}{2H^2} - \frac{27\,\mathcal{H}^2\mathcal{H}_f^2}{4H^2} \\
& - \frac{9\,\mathcal{H}\mathcal{H}_f^3}{2H^2} - \frac{3\,\mathcal{H}_f^4}{8H^2} - \frac{3\,K^2\mathcal{H}^2}{16} + \frac{9\,K\mathcal{H}^2}{4} \\
& + \frac{9\,K\mathcal{H}\mathcal{H}_f}{2} + \frac{3\,K\mathcal{H}_f^2}{4} - 3\,\mathcal{H}^2 - 12\,\mathcal{H}\mathcal{H}_f - 6\,\mathcal{H}_f^2 - \frac{3\,K^2H^2}{8} + 6\,KH^2 - 10\,H^2 \\
& = -\kappa\rho
\end{aligned}
\tag{FE-38}
$$

Symbols in (FE-38).

- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_\mathrm{Pl}$). Unit: GeV. Status: DERIVED.
- $\mathcal{H}_f$: The expansion rate of the block, $\mathcal{H}_f = \dot{f}/f = \dot{b}_4/(2b_4) - \dot{a}_4$ (negative when the extra times and x_0 deflate). Unit: GeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_\mathrm{N}\mathcal{G}(K_*) = 2.328 \times 10^{-36}$ GeV^{-2}. Status: AUTHOR.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.

$$
\begin{aligned}
x_1: \quad & \Big[-\frac{\mathcal{H}\mathcal{H}_f^3}{2H^4} - \frac{\mathcal{H}_f^4}{8H^4} + \frac{K\mathcal{H}\mathcal{H}_f}{2H^2} + \frac{K\mathcal{H}_f^2}{4H^2} \\
& - \frac{\mathcal{H}\mathcal{H}_f}{H^2} - \frac{3\,\mathcal{H}_f^2}{2H^2} - \frac{K^2}{8} + \frac{3\,K}{2} - 2 \Big] \dot{\mathcal{H}} \\
& + \Big[-\frac{3\,\mathcal{H}^2\mathcal{H}_f^2}{4H^4} - \frac{\mathcal{H}\mathcal{H}_f^3}{2H^4} + \frac{K\mathcal{H}^2}{4H^2} \\
& + \frac{K\mathcal{H}\mathcal{H}_f}{2H^2} - \frac{\mathcal{H}^2}{2H^2} - \frac{3\,\mathcal{H}\mathcal{H}_f}{H^2} - \frac{3\,\mathcal{H}_f^2}{2H^2} + \frac{3\,K}{2} - 4 \Big] \dot{\mathcal{H}}_f \\
& - \frac{\mathcal{H}^3\mathcal{H}_f^3}{2H^4} - \frac{19\,\mathcal{H}^2\mathcal{H}_f^4}{16H^4} - \frac{\mathcal{H}\mathcal{H}_f^5}{2H^4} + \frac{K\mathcal{H}^3\mathcal{H}_f}{2H^2} \\
& + \frac{7\,K\mathcal{H}^2\mathcal{H}_f^2}{8H^2} + \frac{K\mathcal{H}\mathcal{H}_f^3}{2H^2} - \frac{\mathcal{H}^3\mathcal{H}_f}{H^2} - \frac{17\,\mathcal{H}^2\mathcal{H}_f^2}{4H^2} \\
& - \frac{6\,\mathcal{H}\mathcal{H}_f^3}{H^2} - \frac{15\,\mathcal{H}_f^4}{8H^2} - \frac{3\,K^2\mathcal{H}^2}{16} + \frac{9\,K\mathcal{H}^2}{4} \\
& + 3\,K\mathcal{H}\mathcal{H}_f + \frac{9\,K\mathcal{H}_f^2}{4} - 3\,\mathcal{H}^2 - 8\,\mathcal{H}\mathcal{H}_f - 10\,\mathcal{H}_f^2 - \frac{3\,K^2H^2}{8} + 6\,KH^2 - 10\,H^2 \\
& = \kappa p_3
\end{aligned}
\tag{FE-39}
$$

Symbols in (FE-39).

- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2} T^2/M_{\rm Pl}$). Unit: GeV. Status: DERIVED.
- $\mathcal{H}_f$: The expansion rate of the block, $\mathcal{H}_f = \dot{f}/f = \dot{b}_4/(2b_4) - \dot{a}_4$ (negative when the extra times and x_0 deflate). Unit: GeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of $m_{\rm W}$ and $\rho_{\rm EW,0}$). Status: AUTHOR.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- $\dot{\mathcal{H}}$: The time derivative $\mathrm{d}\mathcal{H}/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- $\dot{\mathcal{H}}_f$: The time derivative $\mathrm{d}\mathcal{H}_f/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_{\rm N}\mathcal{G}(K_*) = 2.328 \times 10^{-36}$ GeV^{-2}. Status: AUTHOR.
- p_3: The pressure of ordinary 3-space, $p_3 = \mathcal{T}^{x_1}{}_{x_1} = \mathcal{T}^{x_2}{}_{x_2} = \mathcal{T}^{x_3}{}_{x_3}$ for an isotropic source. Unit: GeV^8. Status: STANDARD.

$$
\begin{aligned}
x_0:\ & \Big[-\frac{9\,\mathcal{H}^2\mathcal{H}_f^2}{16H^4}-\frac{3\,\mathcal{H}\mathcal{H}_f^3}{8H^4}+\frac{3\,K\mathcal{H}^2}{16H^2}+\frac{3\,K\mathcal{H}\mathcal{H}_f}{8H^2}\\
&-\frac{3\,\mathcal{H}^2}{8H^2}-\frac{9\,\mathcal{H}\mathcal{H}_f}{4H^2}-\frac{9\,\mathcal{H}_f^2}{8H^2}+\frac{9\,K}{8}-3\Big]\dot{\mathcal{H}}\\
&+\Big[-\frac{3\,\mathcal{H}^3\mathcal{H}_f}{8H^4}-\frac{9\,\mathcal{H}^2\mathcal{H}_f^2}{16H^4}+\frac{3\,K\mathcal{H}^2}{16H^2}-\frac{9\,\mathcal{H}^2}{8H^2}\\
&-\frac{9\,\mathcal{H}\mathcal{H}_f}{4H^2}-\frac{3\,\mathcal{H}_f^2}{8H^2}+\frac{3\,K}{8}-3\Big]\dot{\mathcal{H}}_f\\
&-\frac{9\,\mathcal{H}^4\mathcal{H}_f^2}{16H^4}-\frac{17\,\mathcal{H}^3\mathcal{H}_f^3}{16H^4}-\frac{9\,\mathcal{H}^2\mathcal{H}_f^4}{16H^4}+\frac{3\,K\mathcal{H}^4}{16H^2}\\
&+\frac{9\,K\mathcal{H}^3\mathcal{H}_f}{16H^2}+\frac{3\,K\mathcal{H}^2\mathcal{H}_f^2}{16H^2}-\frac{3\,\mathcal{H}^4}{8H^2}-\frac{27\,\mathcal{H}^3\mathcal{H}_f}{8H^2}\\
&-\frac{45\,\mathcal{H}^2\mathcal{H}_f^2}{8H^2}-\frac{27\,\mathcal{H}\mathcal{H}_f^3}{8H^2}-\frac{3\,\mathcal{H}_f^4}{8H^2}\\
&+\frac{9\,K\mathcal{H}^2}{4}+\frac{9\,K\mathcal{H}\mathcal{H}_f}{8}+\frac{3\,K\mathcal{H}_f^2}{8}-6\,\mathcal{H}^2-9\,\mathcal{H}\mathcal{H}_f\\
&-6\,\mathcal{H}_f^2+3\,KH^2-10\,H^2\\
&=\kappa p_0;\qquad x_5,x_6,x_7:\ \text{the same left side}=\kappa p_5;\\
&\text{block-isotropic order-0 metric: the same left side}=\kappa\bar{p},\qquad \bar{p}=\tfrac{1}{4}\big(p_0+3p_5\big)
\end{aligned}
\tag{FE-40}
$$

Symbols in (FE-40).

- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z=6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H}=\dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H}=(8\pi^3 g_\rho/90)^{1/2}T^2/M_{\rm Pl}$). Unit: GeV. Status: DERIVED.
- $\mathcal{H}_f$: The expansion rate of the block, $\mathcal{H}_f=\dot{f}/f=\dot{b}_4/(2b_4)-\dot{a}_4$ (negative when the extra times and x_0 deflate). Unit: GeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda=-10H^2$ and the mass parameter $m_\Psi=H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of $m_{\rm W}$ and $\rho_{\rm EW,0}$). Status: AUTHOR.
- K: The block variable, $K=1/f^2$. Unit: dimensionless. Status: DERIVED.
- $\dot{\mathcal{H}}$: The time derivative $\mathrm{d}\mathcal{H}/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- $\dot{\mathcal{H}}_f$: The time derivative $\mathrm{d}\mathcal{H}_f/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa=8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp=8\pi G_{\rm N}\mathcal{G}(K_*)=2.328\times10^{-36}$ GeV^{-2}. Status: AUTHOR.
- p_0: The pressure along the hidden direction x_0, $p_0=\mathcal{T}^{x_0}{}_{x_0}$. Unit: GeV^8. Status: STANDARD.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- p_5: The pressure along the extra times, $p_5=\mathcal{T}^{x_5}{}_{x_5}=\mathcal{T}^{x_6}{}_{x_6}=\mathcal{T}^{x_7}{}_{x_7}$. Unit: GeV^8. Status: STANDARD.
- $\bar{p}$: The hidden pressure, the pressure averaged over the block, $\bar{p}=(p_0+3p_5)/4$; for a species at fixed temperature $\bar{p}=\frac{1}{4}\,\partial p/\partial\ln f$. Unit: GeV^8 (or GeV^4 for a four-dimensional density). Status: DERIVED.

$$
\dot{\rho}=-3\mathcal{H}(\rho+p_3)-\mathcal{H}_f(\rho+p_0)-3\mathcal{H}_f(\rho+p_5)=-3\mathcal{H}(\rho+p_3)-4\mathcal{H}_f(\rho+\bar{p})
\tag{FE-41}
$$

Symbols in (FE-41).

- $\dot{\rho}$: The time derivative $\mathrm{d}\rho/\mathrm{d}x_4$. Unit: GeV^9. Status: STANDARD.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H}=\dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H}=(8\pi^3 g_\rho/90)^{1/2}T^2/M_{\rm Pl}$). Unit: GeV. Status: DERIVED.
- ρ: An energy density: in eight dimensions $\rho=-\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$

is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.

- p_3: The pressure of ordinary 3-space, $p_3 = \mathcal{T}^{x_1}{}_{x_1} = \mathcal{T}^{x_2}{}_{x_2} = \mathcal{T}^{x_3}{}_{x_3}$ for an isotropic source. Unit: GeV^8. Status: STANDARD.
- $\mathcal{H}_f$: The expansion rate of the block, $\mathcal{H}_f = \dot{f}/f = \dot{b}_4/(2b_4) - \dot{a}_4$ (negative when the extra times and x_0 deflate). Unit: GeV. Status: DERIVED.
- p_0: The pressure along the hidden direction x_0, $p_0 = \mathcal{T}^{x_0}{}_{x_0}$. Unit: GeV^8. Status: STANDARD.
- p_5: The pressure along the extra times, $p_5 = \mathcal{T}^{x_5}{}_{x_5} = \mathcal{T}^{x_6}{}_{x_6} = \mathcal{T}^{x_7}{}_{x_7}$. Unit: GeV^8. Status: STANDARD.
- $\bar{p}$: The hidden pressure, the pressure averaged over the block, $\bar{p} = (p_0 + 3p_5)/4$; for a species at fixed temperature $\bar{p} = \frac{1}{4}\,\partial p/\partial \ln f$. Unit: GeV^8 (or GeV^4 for a four-dimensional density). Status: DERIVED.

4.2 The approximation of step 1

Order 0 of the metric series. The background keeps only the order-0 term of the metric series (written out in "The approximation scheme"). The order-1 terms average to zero in the block-averaged background equations, and their back-reaction is of second order: on cosmological scales the omitted order is $\Phi^2_{\mathrm{rms}} = 5.8 \times 10^{-10}$ of the background, i.e. 2.4×10^{-5} of the kept first order (step 10).

A block-isotropic source. The hidden and extra-time components of the order-0 Lovelock tensors coincide identically, so the order-0 metric admits only sources with $p_0 = p_5$; for $p_0 \neq p_5$ the hidden component is the variation with respect to the common block scale f, with the hidden pressure $\bar{p} = (p_0 + 3p_5)/4$, and the difference $p_0 - p_5$ drives a block-shape mode outside the order-0 metric (OPEN). This difference is not small against the kept hidden pressure (0.9 for the self-interacting Kohn-Sham test states of step 3 and 4 for dark matter at rest, whose $p_5 = 0$), but its absolute size is suppressed like that of $\bar{p}$, by 10^{-32} or more against the stabiliser (below).

The static block plus its first-order response. After the primordial era the matter sources are tiny against the stabiliser: with the constraint $3\mathcal{G}(K_*)\,\mathcal{H}^2 = \kappa\Delta\rho$ of the response,

$$\frac{\kappa\Delta\rho}{\kappa\rho_{\mathrm{stab}}(f_*)} = 0.7592\,\frac{\mathcal{H}^2}{H^2}: \qquad \begin{array}{ll} 2.96\times10^{-32} & \text{at the dark-matter freeze-out,} \\ 2.4\times10^{-93} & \text{for the cosmic mean today,} \end{array}$$

Symbols in this equation.

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_{\mathrm{N}}\mathcal{G}(K_*) = 2.328 \times 10^{-36}\ \mathrm{GeV}^{-2}$. Status: AUTHOR.
- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m^2_{21} = m^2_2 - m^2_1$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- ρ_{stab}: The energy density of the block stabiliser, a function of the block scale, $\rho_{\mathrm{stab}} \propto f^{-8}$; its value at the static block is $\rho_{\mathrm{stab}}(f_*)$. Unit: GeV^8. Value: $\kappa\rho_{\mathrm{stab}}(f_*) = (26 + 4\sqrt{51})H^2 = 54.566\,H^2$. Status: ASSUMED.
- f_*: The block scale of the stabilised (static) block, $f_* = K_*^{-1/2}$. Unit: dimensionless. Value: 0.215556. Status: DERIVED.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_{\mathrm{Pl}}$). Unit: GeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.

so the block responds linearly. The terms of second order in the sources are smaller than the first-order response by a factor of the same size (2.3×10^{-32} at the freeze-out, the largest source of the model), and the response is adiabatic: the inertia of the block, $\Delta\ddot{K}$ against $\omega_{\mathrm{r}}^2\Delta K$, is of the relative size

$$\frac{|\Delta\ddot{K}|}{\omega_{\rm r}^2\,|\Delta K|} \sim \frac{\mathcal{H}^2}{\omega_{\rm r}^2} = 3.4\times10^{-33} \quad \text{at the dark-matter freeze-out.}$$

Symbols in this equation.

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, $\Delta N_{\rm eff}$ an addition to $N_{\rm eff}$, $\Delta S_{\rm m}$ the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- $\ddot{K}$: The second time derivative $\mathrm{d}^2K/\mathrm{d}x_4^2$. Unit: GeV^2. Status: DERIVED.
- $\omega_{\rm r}$: The angular frequency of the oscillation of the block about the static point (the radion), equal to the radion mass. Unit: GeV. Value: $3.38705\,H = 48.42$ TeV. Status: DERIVED.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_{\rm Pl}$). Unit: GeV. Status: DERIVED.

The primordial era is solved without approximation in the order-0 equations (numerically); the radiation of zero modes that is added in one run changes the final block by 1.0×10^{-6}.

4.3 The solution of step 1

4.3.1 The static block

For $\mathcal{H}_f = 0$ and $\dot{\mathcal{H}} = 0$ the three background equations become algebraic:

$$\begin{aligned}
&x_4:\ -3\mathcal{G}(K)\,\mathcal{H}^2 + 6H^2K\big(1-2w_{(2)}K\big) + \Lambda + \kappa\rho = 0, \qquad x_1:\ \text{the same with } -\kappa p_3 \text{ in place of } \kappa\rho,\\
&x_0:\ 6\big(12w_{(2)}K-1\big)\mathcal{H}^2 - \frac{12\big(w_{(2)}-18w_{(3)}K\big)}{H^2}\,\mathcal{H}^4 + 3H^2K + \Lambda - \kappa p_0 = 0,\\
&x_5, x_6, x_7:\ \text{the same with } \kappa p_5 \text{ in place of } \kappa p_0; \qquad \text{block-isotropic order-0 metric: the same with } \kappa\bar{p},\\
&\mathcal{G}(K) = 1 - 24w_{(2)}K + 72w_{(3)}K^2 = \frac{K^2-12K+16}{16}
\end{aligned} \tag{FE-42}$$

Symbols in (FE-42).

- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\mathcal{G}(K)$: The coefficient of the four-dimensional Einstein-Hilbert term on a static block, $\mathcal{G}(K) = 1 - 24w_{(2)}K + 72w_{(3)}K^2 = (K^2-12K+16)/16$; four-dimensional gravity is attractive for $\mathcal{G} > 0$. Unit: dimensionless. Value: $\mathcal{G}(K_*) = 20/3 + \sqrt{51} = 13.8081$. Status: DERIVED.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_{\rm Pl}$). Unit: GeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of $m_{\rm W}$ and $\rho_{\rm EW,0}$). Status: AUTHOR.
- $w_{(p)}$: The author's Lovelock weights, $w_{(1)} = 1$, $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$. Unit: dimensionless. Value: $(1, 1/32, 1/1152)$. Status: AUTHOR.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -10H^2$. Unit: GeV^2. Value: $-10H^2$. Status: AUTHOR.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_{\rm N}\mathcal{G}(K_*) = 2.328\times10^{-36}$ GeV^{-2}. Status: AUTHOR.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.

- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- p_3: The pressure of ordinary 3-space, $p_3 = \mathcal{T}^{x_1}{}_{x_1} = \mathcal{T}^{x_2}{}_{x_2} = \mathcal{T}^{x_3}{}_{x_3}$ for an isotropic source. Unit: GeV^8. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- p_0: The pressure along the hidden direction x_0, $p_0 = \mathcal{T}^{x_0}{}_{x_0}$. Unit: GeV^8. Status: STANDARD.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- p_5: The pressure along the extra times, $p_5 = \mathcal{T}^{x_5}{}_{x_5} = \mathcal{T}^{x_6}{}_{x_6} = \mathcal{T}^{x_7}{}_{x_7}$. Unit: GeV^8. Status: STANDARD.
- $\bar{p}$: The hidden pressure, the pressure averaged over the block, $\bar{p} = (p_0 + 3p_5)/4$; for a species at fixed temperature $\bar{p} = \frac{1}{4}\,\partial p/\partial \ln f$. Unit: GeV^8 (or GeV^4 for a four-dimensional density). Status: DERIVED.

Without a source and with $\mathcal{H} = 0$ the only static block is $K = 1/(4w_{(2)}) = 8$, which would need $\Lambda = -24H^2$ and has $\mathcal{G} = -1$: four-dimensional gravity would be repulsive. The stabiliser removes this obstruction with the author's weights and $\Lambda = -10H^2$ unchanged (DERIVED):

$$\begin{aligned}
&\kappa\rho_{\mathrm{stab}} = 10H^2 - 6H^2K + \tfrac{3}{8}H^2K^2 \;\; (x_4), \qquad \kappa p_0 = \kappa p_5 = \kappa\rho_{\mathrm{stab}} = 3H^2K - 10H^2 \;\; (x_0;\; x_5, x_6, x_7),\\
&\Rightarrow\; K_\pm = 12 \pm \tfrac{4}{3}\sqrt{51}, \qquad K_* = 12 + \tfrac{4}{3}\sqrt{51} = 21.5219, \qquad f_* = K_*^{-1/2} = 0.215556, \qquad \text{(FE-43)}\\
&\mathcal{G}(K_*) = \tfrac{20}{3} + \sqrt{51} = 13.8081, \qquad \kappa\rho_{\mathrm{stab}}(f_*) = \big(26 + 4\sqrt{51}\big)H^2
\end{aligned}$$

Symbols in (FE-43).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_{\mathrm{N}}\mathcal{G}(K_*) = 2.328 \times 10^{-36}\ \mathrm{GeV}^{-2}$. Status: AUTHOR.
- ρ_{stab}: The energy density of the block stabiliser, a function of the block scale, $\rho_{\mathrm{stab}} \propto f^{-8}$; its value at the static block is $\rho_{\mathrm{stab}}(f_*)$. Unit: GeV^8. Value: $\kappa\rho_{\mathrm{stab}}(f_*) = (26 + 4\sqrt{51})H^2 = 54.566\,H^2$. Status: ASSUMED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- p_0: The pressure along the hidden direction x_0, $p_0 = \mathcal{T}^{x_0}{}_{x_0}$. Unit: GeV^8. Status: STANDARD.
- p_5: The pressure along the extra times, $p_5 = \mathcal{T}^{x_5}{}_{x_5} = \mathcal{T}^{x_6}{}_{x_6} = \mathcal{T}^{x_7}{}_{x_7}$. Unit: GeV^8. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- $K_\pm$: The two static blocks with the stabiliser, $K_\pm = 12 \pm \frac{4}{3}\sqrt{51}$. Unit: dimensionless. Value: 21.5219 and 2.4781. Status: DERIVED.
- K_*: The block variable of the stabilised block, $K_* = 12 + \frac{4}{3}\sqrt{51}$ (the upper root K_+). Unit: dimensionless. Value: 21.5219. Status: DERIVED.
- f_*: The block scale of the stabilised (static) block, $f_* = K_*^{-1/2}$. Unit: dimensionless. Value: 0.215556. Status: DERIVED.
- $\mathcal{G}(K)$: The coefficient of the four-dimensional Einstein-Hilbert term on a static block, $\mathcal{G}(K) = 1 - 24w_{(2)}K + 72w_{(3)}K^2 = (K^2 - 12K + 16)/16$; four-dimensional gravity is attractive for $\mathcal{G} > 0$. Unit: dimensionless. Value: $\mathcal{G}(K_*) = 20/3 + \sqrt{51} = 13.8081$. Status: DERIVED.

$\mathcal{G}(K_*) = 13.8081 > 0$: four-dimensional gravity is attractive. The normalisation of the stabiliser is $\kappa\rho_{\mathrm{stab}}(f_*) = 54.566\,H^2$.

4.3.2 Stability: the radion

Linearised about the static block, the background equations give a closed system for $(\Delta\mathcal{H}, \Delta\mathcal{H}_f, \Delta K)$ with the eigenvalues 0 and $\pm\mathrm{i}\,\omega_{\mathrm{r}}$ (DERIVED, exact):

$$\Delta\dot{\mathcal{H}} = \Xi_{\mathcal{H}K}\,\Delta K, \qquad \Delta\dot{\mathcal{H}}_f = \Xi_{fK}\,\Delta K, \qquad \Delta\dot{K} = -2K_*\,\Delta\mathcal{H}_f,$$
$$\omega_{\rm r}^2 = 2K_*\,\Xi_{fK} = \Big(\frac{31552}{2249} - \frac{2416\sqrt{51}}{6747}\Big)H^2 = (3.38705\,H)^2 \tag{FE-44}$$

Symbols in (FE-44).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, $\Delta N_{\rm eff}$ an addition to $N_{\rm eff}$, $\Delta S_{\rm m}$ the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- $\dot{\mathcal{H}}$: The time derivative $\mathrm{d}\mathcal{H}/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- $\Xi_{\mathcal{H}K}$: The coefficient of ΔK in the first-order equation for $\Delta\dot{\mathcal{H}}$ (linearised block about K_*). Unit: GeV^2. Value: $6(238 - 19\sqrt{51})H^2/2249 = 0.272956\,H^2$. Status: DERIVED.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- $\dot{\mathcal{H}}_f$: The time derivative $\mathrm{d}\mathcal{H}_f/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- Ξ_{fK}: The coefficient of ΔK in the first-order equation for $\Delta\dot{\mathcal{H}}_f$. Unit: GeV^2. Value: $(4063 - 485\sqrt{51})H^2/2249 = 0.266522\,H^2$. Status: DERIVED.
- $\dot{K}$: The time derivative $\mathrm{d}K/\mathrm{d}x_4$. Unit: GeV. Status: DERIVED.
- K_*: The block variable of the stabilised block, $K_* = 12 + \frac{4}{3}\sqrt{51}$ (the upper root K_+). Unit: dimensionless. Value: 21.5219. Status: DERIVED.
- $\mathcal{H}_f$: The expansion rate of the block, $\mathcal{H}_f = \dot{f}/f = \dot{b}_4/(2b_4) - \dot{a}_4$ (negative when the extra times and x_0 deflate). Unit: GeV. Status: DERIVED.
- $\omega_{\rm r}$: The angular frequency of the oscillation of the block about the static point (the radion), equal to the radion mass. Unit: GeV. Value: $3.38705\,H = 48.42$ TeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of $m_{\rm W}$ and $\rho_{\rm EW,0}$). Status: AUTHOR.

The block is stable and oscillates with the radion frequency $\omega_{\rm r} = 3.38705\,H$, i.e. the radion mass 48.42 TeV; the zero eigenvalue belongs to $\Delta\mathcal{H}$ (a small four-dimensional cosmological term). Every stabiliser of the family whose hidden pressure is a fixed multiple, from 1/4 to 3, of its energy density gives a stable block, with $\omega_{\rm r} = 3.33\,H$ to $4.10\,H$ (DERIVED).

4.3.3 The primordial era: the deflating extra times

The three background equations, with the stabiliser $\kappa\rho_{\rm stab} = \kappa\rho_{\rm stab}(f_*)\,(K/K_*)^4$, $p_3 = -\rho_{\rm stab}$, $\bar{p} = \rho_{\rm stab}$ (plus radiation where stated), were integrated from an expanding 3-space and a block larger than the stabilised one ($K < K_*$ at the start, $x_4 = 0$) that deflates at the rate $\mathcal{H}_f = -H/2$, $-H/3$ or $-H/10$ (DERIVED; DOP853, relative tolerance 10^{-11}). The value of K at $x_4 = 60/H$:

start K	$\mathcal{H}_f = -H/10$ at the start	$\mathcal{H}_f = -H/3$	$\mathcal{H}_f = -H/2$
1.5	singular at $x_4 = 1.44/H$	singular at $1.26/H$	singular at $1.14/H$
2.3	singular at $0.33/H$	singular at $0.31/H$	no expanding start
2.7	21.415	21.435	21.451
3	21.412	21.434	21.455
6	21.405	21.453	21.500
10	21.446	21.541	21.584
20	21.634	21.640	21.638

Every start above the separatrix $K_{\rm sep} = 2.5301$, 2.5991, 2.6983 (for the initial rates $-H/10$, $-H/3$, $-H/2$; just above the unstable static block $K_- = 2.4781$) ends at the stable block: the extra times x_5, x_6, x_7 and the hidden direction x_0 deflate at rates up to $|\mathcal{H}_f| = 2.63\,H$ (in the stored runs with the start $K = 3, 5, 10$), by up to the factor $f(K_{\rm sep})/f_* = 2.917$, reach K_* within $x_4 < 1/H$, and then oscillate about it with the radion frequency ($3.3868\,H$

measured, $3.3870\,H$ linear), while 3-space keeps expanding; the oscillation energy redshifts like dust ($\mathcal{H}\,x_4 \to 0.667$ to 0.670; 2/3 for dust). The extra times therefore deflate only in a short primordial era and are then frozen: the masses, all of which depend on f (step 2), become constant.




Figure 1: The primordial era: the block scale f, the expansion rate $\mathcal{H}$ of 3-space and the block rate $\mathcal{H}_f$ for the starts $K = 3, 5, 10$ with the initial rate $\mathcal{H}_f = -H/3$ (one with radiation), from the full nonlinear Einstein-Lovelock equations with the author's weights. The panels label K, $\mathcal{H}$ and $\mathcal{H}_f$ with the letters k, A and h of version 3.

4.3.4 The response of the block to matter

To first order about the static block, with the block-averaged matter sources and $\mathcal{H}^2$ counted as of first order (the constraint gives $3\mathcal{G}(K_*)\,\mathcal{H}^2 = \kappa\Delta\rho$ because $\mathrm{d}[6H^2K(1-2w_{(2)}K)+\kappa\rho_{\mathrm{stab}}]/\mathrm{d}K$ vanishes at K_*), the block obeys a driven oscillator equation (DERIVED, exact):

$$\begin{aligned} \Delta\dot{\mathcal{H}} &= \Xi_{\mathcal{H}K}\,\Delta K + \Xi_{\mathcal{H}3}\,\kappa\Delta p_3 + \Xi_{\mathcal{H}p}\,\kappa\Delta\bar{p} + \Xi_{\mathcal{H}\rho}\,\kappa\Delta\rho, \qquad 3\mathcal{G}(K_*)\,\mathcal{H}^2 = \kappa\Delta\rho, \\ \Delta\dot{\mathcal{H}}_f &= \Xi_{fK}\,\Delta K + \Xi_{f3}\,\kappa\Delta p_3 + \Xi_{fp}\,\kappa\Delta\bar{p} + \Xi_{f\rho}\,\kappa\Delta\rho, \\ \Delta\ddot{K} + \omega_{\mathrm{r}}^2\,\Delta K &= -2K_*\big(\Xi_{f3}\,\kappa\Delta p_3 + \Xi_{fp}\,\kappa\Delta\bar{p} + \Xi_{f\rho}\,\kappa\Delta\rho\big) \end{aligned} \tag{FE-45}$$

Symbols in (FE-45).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- $\dot{\mathcal{H}}$: The time derivative $\mathrm{d}\mathcal{H}/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- $\Xi_{\mathcal{H}K}$: The coefficient of ΔK in the first-order equation for $\Delta\dot{\mathcal{H}}$ (linearised block about K_*). Unit: GeV^2. Value: $6(238-19\sqrt{51})H^2/2249 = 0.272956\,H^2$. Status: DERIVED.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- $\Xi_{\mathcal{H}3}$: The coefficient of $\kappa\Delta p_3$ in the equation for $\Delta\dot{\mathcal{H}}$. Unit: dimensionless. Value: $2283/4498 - 162\sqrt{51}/2249 = -0.006853$. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_{\mathrm{N}}\mathcal{G}(K_*) = 2.328\times10^{-36}\ \mathrm{GeV}^{-2}$. Status: AUTHOR.
- p_3: The pressure of ordinary 3-space, $p_3 = \mathcal{T}^{x_1}{}_{x_1} = \mathcal{T}^{x_2}{}_{x_2} = \mathcal{T}^{x_3}{}_{x_3}$ for an isotropic source. Unit: GeV^8. Status: STANDARD.
- $\Xi_{\mathcal{H}p}$: The coefficient of $\kappa\Delta\bar{p}$ in the equation for $\Delta\dot{\mathcal{H}}$. Unit: dimensionless. Value: $(28\sqrt{51}-114)/2249 = 0.038221$. Status: DERIVED.
- $\bar{p}$: The hidden pressure, the pressure averaged over the block, $\bar{p} = (p_0 + 3p_5)/4$; for a species at fixed temperature $\bar{p} = \frac{1}{4}\,\partial p/\partial\ln f$. Unit: GeV^8 (or GeV^4 for a four-dimensional density). Status: DERIVED.
- $\Xi_{\mathcal{H}\rho}$: The coefficient of $\kappa\Delta\rho$ in the equation for $\Delta\dot{\mathcal{H}}$. Unit: dimensionless. Value: $135021/265382 - 10308\sqrt{51}/132691 = -0.045997$. Status: DERIVED.

- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- $\mathcal{G}(K)$: The coefficient of the four-dimensional Einstein-Hilbert term on a static block, $\mathcal{G}(K) = 1 - 24w_{(2)}K + 72w_{(3)}K^2 = (K^2 - 12K + 16)/16$; four-dimensional gravity is attractive for $\mathcal{G} > 0$. Unit: dimensionless. Value: $\mathcal{G}(K_*) = 20/3 + \sqrt{51} = 13.8081$. Status: DERIVED.
- K_*: The block variable of the stabilised block, $K_* = 12 + \frac{4}{3}\sqrt{51}$ (the upper root K_+). Unit: dimensionless. Value: 21.5219. Status: DERIVED.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_{\mathrm{Pl}}$). Unit: GeV. Status: DERIVED.
- $\dot{\mathcal{H}}_f$: The time derivative $\mathrm{d}\mathcal{H}_f/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- Ξ_{fK}: The coefficient of ΔK in the first-order equation for $\Delta\dot{\mathcal{H}}_f$. Unit: GeV^2. Value: $(4063 - 485\sqrt{51})H^2/2249 = 0.266522\,H^2$. Status: DERIVED.
- Ξ_{f3}: The coefficient of $\kappa\Delta p_3$ in the equation for $\Delta\dot{\mathcal{H}}_f$. Unit: dimensionless. Value: $21\sqrt{51}/2249 - 171/4498 = 0.028666$. Status: DERIVED.
- Ξ_{fp}: The coefficient of $\kappa\Delta\bar{p}$ in the equation for $\Delta\dot{\mathcal{H}}_f$. Unit: dimensionless. Value: $239\sqrt{51}/6747 - 485/2249 = 0.037320$. Status: DERIVED.
- $\Xi_{f\rho}$: The coefficient of $\kappa\Delta\rho$ in the equation for $\Delta\dot{\mathcal{H}}_f$. Unit: dimensionless. Value: $57/4498 - 7\sqrt{51}/2249 = -0.009555$. Status: DERIVED.
- $\ddot{K}$: The second time derivative $\mathrm{d}^2K/\mathrm{d}x_4^2$. Unit: GeV^2. Status: DERIVED.
- ω_{r}: The angular frequency of the oscillation of the block about the static point (the radion), equal to the radion mass. Unit: GeV. Value: $3.38705\,H = 48.42$ TeV. Status: DERIVED.

$$\Xi_{\mathcal{H}K} = \frac{6\big(238 - 19\sqrt{51}\big)}{2249}H^2, \qquad \Xi_{fK} = \frac{4063 - 485\sqrt{51}}{2249}H^2, \qquad \Xi_{\mathcal{H}3} = \frac{2283}{4498} - \frac{162\sqrt{51}}{2249},$$
$$\Xi_{\mathcal{H}p} = \frac{28\sqrt{51} - 114}{2249}, \qquad \Xi_{\mathcal{H}\rho} = \frac{135021}{265382} - \frac{10308\sqrt{51}}{132691}, \qquad \Xi_{f3} = \frac{21\sqrt{51}}{2249} - \frac{171}{4498}, \tag{FE-46}$$
$$\Xi_{fp} = \frac{239\sqrt{51}}{6747} - \frac{485}{2249}, \qquad \Xi_{f\rho} = \frac{57}{4498} - \frac{7\sqrt{51}}{2249}$$

Symbols in (FE-46).

- $\Xi_{\mathcal{H}K}$: The coefficient of ΔK in the first-order equation for $\Delta\dot{\mathcal{H}}$ (linearised block about K_*). Unit: GeV^2. Value: $6(238 - 19\sqrt{51})H^2/2249 = 0.272956\,H^2$. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- Ξ_{fK}: The coefficient of ΔK in the first-order equation for $\Delta\dot{\mathcal{H}}_f$. Unit: GeV^2. Value: $(4063 - 485\sqrt{51})H^2/2249 = 0.266522\,H^2$. Status: DERIVED.
- $\Xi_{\mathcal{H}3}$: The coefficient of $\kappa\Delta p_3$ in the equation for $\Delta\dot{\mathcal{H}}$. Unit: dimensionless. Value: $2283/4498 - 162\sqrt{51}/2249 = -0.006853$. Status: DERIVED.
- $\Xi_{\mathcal{H}p}$: The coefficient of $\kappa\Delta\bar{p}$ in the equation for $\Delta\dot{\mathcal{H}}$. Unit: dimensionless. Value: $(28\sqrt{51} - 114)/2249 = 0.038221$. Status: DERIVED.
- $\Xi_{\mathcal{H}\rho}$: The coefficient of $\kappa\Delta\rho$ in the equation for $\Delta\dot{\mathcal{H}}$. Unit: dimensionless. Value: $135021/265382 - 10308\sqrt{51}/132691 = -0.045997$. Status: DERIVED.
- Ξ_{f3}: The coefficient of $\kappa\Delta p_3$ in the equation for $\Delta\dot{\mathcal{H}}_f$. Unit: dimensionless. Value: $21\sqrt{51}/2249 - 171/4498 = 0.028666$. Status: DERIVED.
- Ξ_{fp}: The coefficient of $\kappa\Delta\bar{p}$ in the equation for $\Delta\dot{\mathcal{H}}_f$. Unit: dimensionless. Value: $239\sqrt{51}/6747 - 485/2249 = 0.037320$. Status: DERIVED.
- $\Xi_{f\rho}$: The coefficient of $\kappa\Delta\rho$ in the equation for $\Delta\dot{\mathcal{H}}_f$. Unit: dimensionless. Value: $57/4498 - 7\sqrt{51}/2249 = -0.009555$. Status: DERIVED.

For a slowly varying source the block follows it adiabatically, and 3-space obeys the four-dimensional Friedmann acceleration equation exactly:

$$\Delta K = \frac{c_\rho\,\kappa\Delta\rho + c_3\,\kappa\Delta p_3 + c_p\,\kappa\Delta\bar{p}}{H^2}, \qquad \Delta\ln f = -\frac{\Delta K}{2K_*}, \qquad c_3 = -3c_\rho, \qquad \Delta\dot{\mathcal{H}} = -\frac{\kappa(\Delta\rho + \Delta p_3)}{2\mathcal{G}(K_*)} \quad \text{(FE-47)}$$

Symbols in (FE-47).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- c_ρ: The coefficient of $\kappa\Delta\rho$ in the adiabatic block response ΔK. Unit: dimensionless. Value: 0.0358521. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_{\text{N}}\mathcal{G}(K_*) = 2.328\times10^{-36}\ \text{GeV}^{-2}$. Status: AUTHOR.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- c_3: The coefficient of $\kappa\Delta p_3$ in the adiabatic block response, $c_3 = -3c_\rho$ (so radiation drops out). Unit: dimensionless. Value: -0.107556. Status: DERIVED.
- p_3: The pressure of ordinary 3-space, $p_3 = \mathcal{T}^{x_1}{}_{x_1} = \mathcal{T}^{x_2}{}_{x_2} = \mathcal{T}^{x_3}{}_{x_3}$ for an isotropic source. Unit: GeV^8. Status: STANDARD.
- c_p: The coefficient of $\kappa\Delta\bar{p}$ in the adiabatic block response. Unit: dimensionless. Value: -0.140028. Status: DERIVED.
- $\bar{p}$: The hidden pressure, the pressure averaged over the block, $\bar{p} = (p_0 + 3p_5)/4$; for a species at fixed temperature $\bar{p} = \frac{1}{4}\,\partial p/\partial\ln f$. Unit: GeV^8 (or GeV^4 for a four-dimensional density). Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\text{EW},0}$). Status: AUTHOR.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \text{e}^{-a_4}b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- K_*: The block variable of the stabilised block, $K_* = 12 + \frac{4}{3}\sqrt{51}$ (the upper root K_+). Unit: dimensionless. Value: 21.5219. Status: DERIVED.
- $\dot{\mathcal{H}}$: The time derivative $\text{d}\mathcal{H}/\text{d}x_4$. Unit: GeV^2. Status: DERIVED.
- $\mathcal{G}(K)$: The coefficient of the four-dimensional Einstein-Hilbert term on a static block, $\mathcal{G}(K) = 1 - 24w_{(2)}K + 72w_{(3)}K^2 = (K^2 - 12K + 16)/16$; four-dimensional gravity is attractive for $\mathcal{G} > 0$. Unit: dimensionless. Value: $\mathcal{G}(K_*) = 20/3 + \sqrt{51} = 13.8081$. Status: DERIVED.

with $c_\rho = 0.035852$, $c_3 = -0.107556$, $c_p = -0.140028$: radiation ($\Delta p_3 = \Delta\rho/3$) drops out exactly ($c_3 = -3c_\rho$), so only the trace $\Delta\rho - 3\Delta p_3$ and the hidden pressure move the block. The hidden-pressure part alone is $\Delta K = \kappa\Delta\bar{p}/(3H^2 - 4\kappa\rho_{\text{stab}}(f_*)/K_*)$, a stiffness of $-7.1414\,H^2$.

4.3.5 Four-dimensional gravity

On a product of a four-dimensional metric with the static block the Lovelock densities factorise, and the eight-dimensional action becomes the Einstein-Hilbert action with the coefficient $\mathcal{G}(K)$ (DERIVED, exact):

$$
\begin{aligned}
&\mathcal{L}_{(1)} = \mathcal{L}^{(4)}_{(1)} + \mathcal{L}^{(\mathrm{int})}_{(1)}, \qquad \mathcal{L}_{(2)} = \mathcal{L}^{(4)}_{(2)} + 2\mathcal{L}^{(4)}_{(1)}\mathcal{L}^{(\mathrm{int})}_{(1)} + \mathcal{L}^{(\mathrm{int})}_{(2)}, \qquad \mathcal{L}_{(3)} = 3\mathcal{L}^{(4)}_{(2)}\mathcal{L}^{(\mathrm{int})}_{(1)} + 3\mathcal{L}^{(4)}_{(1)}\mathcal{L}^{(\mathrm{int})}_{(2)},\\
&\mathcal{L}^{(4)}_{(1)} = R^{(4)}, \qquad \mathcal{L}^{(4)}_{(3)} = 0, \qquad \mathcal{L}^{(\mathrm{int})}_{(1)} = -12H^2K, \qquad \mathcal{L}^{(\mathrm{int})}_{(2)} = 24H^4K^2, \qquad \mathcal{L}^{(\mathrm{int})}_{(3)} = 0,\\
&\sum_{p=1}^{3} \alpha_{(p)}\, \mathcal{L}_{(p)} - 2\Lambda - 2\kappa\rho_{\mathrm{stab}} = \mathcal{G}(K)\, R^{(4)} + (\text{topological}) - 2\mathcal{G}(K)\, \Lambda_4,\\
&\Lambda_4 = \frac{6H^2K\big(1 - 2w_{(2)}K\big) + \Lambda + \kappa\rho_{\mathrm{stab}}}{\mathcal{G}(K)}
\end{aligned}
\tag{FE-48}
$$

Symbols in (FE-48).

- $\mathcal{L}_{(p)}$: The Lovelock density of order p of the metric (a scalar built from p Riemann tensors): $\mathcal{L}_{(1)} = R$ (Einstein-Hilbert), $\mathcal{L}_{(2)}$ the Gauss-Bonnet density, $\mathcal{L}_{(3)}$ the cubic Lovelock density. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- $\mathcal{L}^{(4)}_{(r)}$: The Lovelock density of order r of the four-dimensional metric $g^{(4)}_{\mu\nu}$ ($\mathcal{L}^{(4)}_{(1)} = R^{(4)}$; the order 2 is topological and the order 3 vanishes in four dimensions). Unit: $1/\mathrm{length}^{2r}$. Status: STANDARD.
- $\mathcal{L}^{(\mathrm{int})}_{(r)}$: The Lovelock density of order r of the static block (a four-dimensional space of constant curvature $-H^2K$): 1, $-12H^2K$, $24H^4K^2$, 0 for $r = 0, 1, 2, 3$. Unit: $1/\mathrm{length}^{2r}$. Status: DERIVED.
- $R^{(4)}$: The Ricci scalar of the four-dimensional metric $g^{(4)}_{\mu\nu}$. Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic). Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)}H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -10H^2$. Unit: GeV^2. Value: $-10H^2$. Status: AUTHOR.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_{\mathrm{N}}\mathcal{G}(K_*) = 2.328 \times 10^{-36}$ GeV^{-2}. Status: AUTHOR.
- ρ_{stab}: The energy density of the block stabiliser, a function of the block scale, $\rho_{\mathrm{stab}} \propto f^{-8}$; its value at the static block is $\rho_{\mathrm{stab}}(f_*)$. Unit: GeV^8. Value: $\kappa\rho_{\mathrm{stab}}(f_*) = (26 + 4\sqrt{51})H^2 = 54.566\, H^2$. Status: ASSUMED.
- $\mathcal{G}(K)$: The coefficient of the four-dimensional Einstein-Hilbert term on a static block, $\mathcal{G}(K) = 1 - 24w_{(2)}K + 72w_{(3)}K^2 = (K^2 - 12K + 16)/16$; four-dimensional gravity is attractive for $\mathcal{G} > 0$. Unit: dimensionless. Value: $\mathcal{G}(K_*) = 20/3 + \sqrt{51} = 13.8081$. Status: DERIVED.
- Λ_4: The four-dimensional cosmological constant of the effective theory on the block, $\Lambda_4 = [6H^2K(1 - 2w_{(2)}K) + \Lambda + \kappa\rho_{\mathrm{stab}}]/\mathcal{G}(K)$; zero at the static point, the observed dark energy a tuned residual. Unit: GeV^2. Value: 0 at the static point; observed $\Lambda_4/H^2 \sim 10^{-92}$. Status: DERIVED.
- $w_{(p)}$: The author's Lovelock weights, $w_{(1)} = 1$, $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$. Unit: dimensionless. Value: $(1, 1/32, 1/1152)$. Status: AUTHOR.

$$
\begin{aligned}
&G^{(4)}_{\mu\nu} + \Lambda_4\, g^{(4)}_{\mu\nu} = 8\pi G_{\mathrm{N}}\, \mathcal{T}^{(4)}_{\mu\nu}, \qquad 8\pi G_{\mathrm{N}} = \frac{\kappa}{\mathcal{G}(K_*)\, V_\perp}, \qquad V_\perp = \frac{2f^4 v_{\mathrm{t}}\big(1 - \mathrm{e}^{-3HL}\big)}{3H},\\
&\mathcal{T}^{(4)}_{\mu\nu} = \int \mathrm{d}x_0\, \mathrm{d}x_5\, \mathrm{d}x_6\, \mathrm{d}x_7\, \sqrt{|g_\perp|}\, \mathcal{T}_{\mu\nu}, \qquad g_3^2 = \frac{\hat{g}_3^2}{V_\perp}, \qquad g_2^2 = \frac{\hat{g}_2^2}{V_\perp}, \qquad g_1^2 = \frac{\hat{g}_1^2}{V_\perp}
\end{aligned}
\tag{FE-49}
$$

Symbols in (FE-49).

- $G^{(4)}_{\mu\nu}$: The Einstein tensor of the four-dimensional metric, $R^{(4)}_{\mu\nu} - \frac{1}{2}g^{(4)}_{\mu\nu}R^{(4)}$. Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+, +, +, -$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index running over $x_1, \ldots, x_4$ (as μ). Status: STANDARD.

- Λ_4: The four-dimensional cosmological constant of the effective theory on the block, $\Lambda_4 = [6H^2K(1 - 2w_{(2)}K) + \Lambda + \kappa\rho_{\mathrm{stab}}]/\mathcal{G}(K)$; zero at the static point, the observed dark energy a tuned residual. Unit: GeV^2. Value: 0 at the static point; observed $\Lambda_4/H^2 \sim 10^{-92}$. Status: DERIVED.
- $g^{(4)}_{\mu\nu}$: The four-dimensional metric of the observable coordinates $x_1, \ldots, x_4$ (the metric of the effective four-dimensional general relativity on the stabilised block). Unit: dimensionless. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_{N}: Newton's constant of four-dimensional gravity, $8\pi G_{\mathrm{N}} = \kappa/(\mathcal{G}(K_*)V_\perp)$. Unit: GeV^{-2}. Value: 6.709×10^{-39} GeV^{-2} ($M_{\mathrm{Pl}} = G_{\mathrm{N}}^{-1/2} = 1.22091 \times 10^{19}$ GeV). Status: STANDARD.
- $\mathcal{T}^{(4)}_{\mu\nu}$: The four-dimensional energy-momentum tensor: $\mathcal{T}_{\mu\nu}$ integrated over the block (both halves of a period of x_0 and the extra times). Unit: GeV^4. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_{\mathrm{N}}\mathcal{G}(K_*) = 2.328 \times 10^{-36}$ GeV^{-2}. Status: AUTHOR.
- $\mathcal{G}(K)$: The coefficient of the four-dimensional Einstein-Hilbert term on a static block, $\mathcal{G}(K) = 1 - 24w_{(2)}K + 72w_{(3)}K^2 = (K^2 - 12K + 16)/16$; four-dimensional gravity is attractive for $\mathcal{G} > 0$. Unit: dimensionless. Value: $\mathcal{G}(K_*) = 20/3 + \sqrt{51} = 13.8081$. Status: DERIVED.
- K_*: The block variable of the stabilised block, $K_* = 12 + \frac{4}{3}\sqrt{51}$ (the upper root K_+). Unit: dimensionless. Value: 21.5219. Status: DERIVED.
- $V_\perp$: The proper volume of the block per unit four-volume (both halves of a period of x_0 and the extra times), $V_\perp = 2f^4 v_{\mathrm{t}}(1 - \mathrm{e}^{-3HL})/(3H)$. Unit: GeV^{-4}. Status: DERIVED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4}b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- v_{t}: The coordinate volume of the compact extra times x_5, x_6, x_7. Unit: GeV^{-3}. Value: not fixed; $v_{\mathrm{t}} = H^{-3}$ in the quoted conversions. Status: ASSUMED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- $g_\perp$: The metric of the four transverse directions x_0, x_5, x_6, x_7 (the block); $|g_\perp|$ is the absolute value of its determinant. Unit: dimensionless. Status: DERIVED.
- $\mathcal{T}^M{}_N$: The energy-momentum tensor of all matter (every field except gravity, the stabiliser included), the source of gravity; defined by the change of the matter action under a change of the metric. Unit: GeV^8. Status: STANDARD.
- g_3: The four-dimensional strong coupling ($g_3^2 = 4\pi\alpha_3$). Unit: dimensionless. Value: $g_3^2 = 0.7786$ at $2M_{\mathrm{DM}}$. Status: DERIVED.
- $\hat{g}_3$: The eight-dimensional SU(3) coupling (it multiplies the gluon kinetic term as $1/\hat{g}_3^2$); $g_3^2 = \hat{g}_3^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- g_2: The four-dimensional SU(2) coupling ($g_2^2 = 4\pi\alpha_2$). Unit: dimensionless. Value: $g_2^2 = 0.3781$ at $2M_{\mathrm{DM}}$. Status: DERIVED.
- $\hat{g}_2$: The eight-dimensional SU(2) coupling; $g_2^2 = \hat{g}_2^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- g_1: The four-dimensional hypercharge coupling ($g_1^2 = 4\pi\alpha_1$, Standard-Model normalisation without the GUT factor). Unit: dimensionless. Value: $g_1^2 = 0.1380$ at $2M_{\mathrm{DM}}$. Status: DERIVED.
- $\hat{g}_1$: The eight-dimensional U(1)$_Y$ coupling; $g_1^2 = \hat{g}_1^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.

Four-dimensional gravity on the stabilised block is general relativity, with $\Lambda_4 = 0$ at the static point. The observed

dark energy needs $\Lambda_4/H^2 \sim (\mathcal{H}_0/H)^2 = 1.0 \times 10^{-92}$, a tuned residual (OPEN).

4.4 The check of step 1

In its own equations.

- *The static block in the background equations.* Substituted into the printed x_4, x_1 and x_0 equations, the stored $K_* = 21.5219045714$ and $\kappa\rho_{\rm stab}(f_*) = 54.5657137142\,H^2$ leave the residual 3.9×10^{-13} of the largest term (1.2×10^{-16} with the closed forms), and the printed polynomials equal the code's symbolic field equations exactly (DERIVED, `um2_residuals.py`).
- *The primordial trajectories in the background equations.* The constraint x_4 holds at the 1604 stored samples to 3.6×10^{-12} of the sum of its terms (5.6×10^{-10} absolute over the whole integration), and the one-step defect of the stored trajectory, re-integrated over $0.1/H$ from every fifth sample, is 1.3×10^{-11} in units of H (DERIVED).
- *The first-order response in the nonlinear equations.* The adiabatic solution substituted into the full nonlinear equations leaves a residual of second order, as it must: for sources $\kappa(\Delta\rho, \Delta p_3, \Delta\bar{p}) = \epsilon(1, 0.2, 0.7)\,H^2$ the residual is 1.4×10^{-6} at $\epsilon = 10^{-4}$ and 1.4×10^{-7} at $\epsilon = 10^{-5}$ (ratio 0.10), and the cancellation of radiation, $c_3 + 3c_\rho$, holds to 2×10^{-13} (DERIVED).
- *The four-dimensional reduction.* $\Lambda_4(K_*) = 0$ and $\mathrm{d}[6H^2K(1 - 2w_{(2)}K) + \kappa\rho_{\rm stab}]/\mathrm{d}K = 0$ at K_* hold to 3.9×10^{-13} of the largest term with the stored numbers and to 1.2×10^{-16} with the closed forms (the exact values are 0; version 3 quoted the stored -6×10^{-12} of the derivative, which is this rounding), and the coefficient of $R^{(4)}$ equals $\mathcal{G}(K)$ symbolically (DERIVED).

In concert with the other field equations.

The spinor equations on the static block. Every spinor of UM-2 couples to the metric through the canonical spin connection, which on the order-0 metric is exactly

$$\gamma^M \Omega_M = \frac{3}{2}\,\frac{H}{f}\,\gamma^{x_0} + \Big(\frac{\dot{b}_4}{b_4} - \frac{\dot{a}_4}{2}\Big)\gamma^{x_4} \tag{FE-13}$$

Symbols in (FE-13).

- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4, 4)$; $\gamma^M = \mathbf{e}_a{}^M \gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Ω_M: The spin connection acting on spinors, the matrix $\Omega_M = \frac{1}{2}\omega_{Mab}\mathcal{S}^{ab}$. Unit: 1/length. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of $m_{\rm W}$ and $\rho_{\rm EW,0}$). Status: AUTHOR.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- $\dot{b}_4$: The time derivative $\mathrm{d}b_4/\mathrm{d}x_4$. Unit: GeV. Status: DERIVED.
- b_4: The author's metric function of the time x_4 in the hidden and extra-time components, $b_4 > 0$. Unit: dimensionless. Status: AUTHOR.
- $\dot{a}_4$: The time derivative $\mathrm{d}a_4/\mathrm{d}x_4$. Unit: GeV. Status: DERIVED.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.

On the static block ($\dot{a}_4 = \dot{b}_4 = 0$) it reduces to $\frac{3}{2}(H/f_*)\gamma^{x_0}$, which the factor $\mathrm{e}^{-\frac{3}{2}Hy}$ of the reduction of step 2 removes exactly; the masses of step 2 and the Kohn-Sham states of step 3 are therefore solutions on this background.

- *The radion frequency of the linear analysis and of the nonlinear integration.* The closed form $\omega_{\mathrm{r}} = 3.387049962\,H$ and the frequency $3.386781524\,H$ measured from the zero crossings of the nonlinear primordial oscillation (relative amplitude 3×10^{-3}) agree to 7.9×10^{-5} (DERIVED).
- *The block scale in every other equation.* The block scale $f_* = 0.215555768391$ of the static block is the one used in the x_0 spectra (step 2), the Kohn-Sham states (step 3) and the metric series (step 10), to 2.2×10^{-12}; the response coefficients used in steps 3 and 8 are those of the background equations to 2×10^{-13} (the printed six decimals within 2.6×10^{-7}) (DERIVED).
- *The Friedmann equation of the thermal history.* At the dark-matter freeze-out the four-dimensional Friedmann equation with $8\pi G_{\mathrm{N}} = \kappa/(\mathcal{G}(K_*)V_\perp)$ and all species gives an expansion rate 1.7×10^{-4} above the one used in the Boltzmann equation of step 9, because the thermal history leaves out the fold bands of dirac16complex (3.5×10^{-4} of the energy density then); the relic density moves by about the same relative amount, a tenth of the precision of the measured dark-matter density (DERIVED).

4.5 The status of step 1

DERIVED (exact, symbolic): the Lovelock tensors of the order-0 metric, the three background equations, the static block and K_*, $\mathcal{G}(K_*)$, the radion frequency, the first-order and the adiabatic response, the four-dimensional reduction. DERIVED (numerical): the primordial era. ASSUMED: the stabiliser (its microscopic origin is OPEN). OPEN: the block-shape mode driven by $p_0 - p_5$; the tuning of Λ_4 (dark energy).

5 Step 2: the masses from the x_0 dependence of the wave functions

Every four-dimensional mass of UM-2 is an eigenvalue of the x_0 operator of its field on the static block of step 1. The eigenfunctions depend on x_0 through $y(z)$, $z = 6Hx_0$, and are periodic with the period $\pi/(6H)$ (the author's requirement): a field that is localised in x_0 carries an x_0 momentum and hence a four-dimensional mass, while a flat profile is massless. Step 2 solves the field equations of the gauge bosons, the graviton, the twelve fermions of three generations and dirac16complex for their x_0 eigenvalues.

5.1 The field equations of step 2

Gauge bosons and the graviton. The gauge fields of the Standard Model live in the eight dimensions; the tip sigma field Σ (no radial mode, hence no Higgs particle) gives W^+, W^- and Z a mass term at the tip $y = -L$ and the photon none:

$$S_{\mathrm{gauge}} = -\frac{1}{4}\int \mathrm{d}^8X\,\sqrt{|g|}\,\Big(\frac{1}{\hat g_3^2}\,G^A_{MN}G^{AMN} + \frac{1}{\hat g_2^2}\,W^I_{MN}W^{IMN} + \frac{1}{\hat g_1^2}\,B_{MN}B^{MN}\Big) \tag{IN-8}$$

Symbols in (IN-8).

- S_{gauge}: The action of the gauge fields of SU(3), SU(2) and $\mathrm{U}(1)_Y$ in eight dimensions. Unit: dimensionless. Status: ASSUMED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0,\dots,x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \dots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1,\dots,x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\dots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- $\hat g_3$: The eight-dimensional SU(3) coupling (it multiplies the gluon kinetic term as $1/\hat g_3^2$); $g_3^2 = \hat g_3^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- G^A_M, G_M, G_{MN}, G^A_{MN}: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G^A_M ($A = 1,\dots,8$), the matrix $G_M = G^A_M t^A$, the field strength G_{MN} (matrix) and its components G^A_{MN}; $G^A_{\mu\nu}$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.

- A: A colour (adjoint SU(3)) index running over $1,\dots,8$; a repeated index is summed. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \dots$): it runs over $x_0, x_1, \dots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \dots$), running over $x_0, \dots, x_7$; a repeated index is summed. Status: STANDARD.
- $\hat{g}_2$: The eight-dimensional SU(2) coupling; $g_2^2 = \hat{g}_2^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- W_M^I, W_M, W_{MN}, W_{MN}^I: The SU(2) gauge field (geometric normalisation): the components W_M^I ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W_M^I \tau^I/2$, the field strength W_{MN} and its components W_{MN}^I; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- $\hat{g}_1$: The eight-dimensional $\mathrm{U}(1)_Y$ coupling; $g_1^2 = \hat{g}_1^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.

$$W_M^\pm = \frac{W_M^1 \mp \mathrm{i} W_M^2}{\sqrt{2}\,\hat{g}_2}, \qquad Z_M = \frac{W_M^3 - B_M}{\hat{g}_Z}, \qquad A_M = \frac{\hat{g}_1^2\, W_M^3 + \hat{g}_2^2\, B_M}{\hat{g}_1 \hat{g}_2 \hat{g}_Z},$$
$$\hat{g}_Z = \sqrt{\hat{g}_2^2 + \hat{g}_1^2}, \qquad \cos\theta_\mathrm{W} = \frac{\hat{g}_2}{\hat{g}_Z}, \qquad \sin\theta_\mathrm{W} = \frac{\hat{g}_1}{\hat{g}_Z} \tag{IN-10}$$

Symbols in (IN-10).

- $W_M^\pm$, W^+, W^-: The charged weak bosons, canonically normalised: $W_M^\pm = (W_M^1 \mp \mathrm{i} W_M^2)/(\sqrt{2}\,\hat{g}_2)$; W^+ carries the electric charge $+e$ and W^- the charge $-e$ (they are each other's antiparticles and have one mass). Unit: GeV^3. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \dots$): it runs over $x_0, x_1, \dots, x_7$; a repeated index is summed. Status: STANDARD.
- W_M^I, W_M, W_{MN}, W_{MN}^I: The SU(2) gauge field (geometric normalisation): the components W_M^I ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W_M^I \tau^I/2$, the field strength W_{MN} and its components W_{MN}^I; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\hat{g}_2$: The eight-dimensional SU(2) coupling; $g_2^2 = \hat{g}_2^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- Z_M: The Z boson field, canonically normalised, $Z_M = (W_M^3 - B_M)/\hat{g}_Z$. Unit: GeV^3. Status: STANDARD.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- $\hat{g}_Z$: The eight-dimensional coupling of the Z, $\hat{g}_Z = (\hat{g}_2^2 + \hat{g}_1^2)^{1/2}$. Unit: GeV^{-2}. Status: STANDARD.
- A_M: The photon field, canonically normalised, $A_M = (\hat{g}_1^2 W_M^3 + \hat{g}_2^2 B_M)/(\hat{g}_1 \hat{g}_2 \hat{g}_Z)$. Unit: GeV^3. Status: STANDARD.
- $\hat{g}_1$: The eight-dimensional $\mathrm{U}(1)_Y$ coupling; $g_1^2 = \hat{g}_1^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- θ_W: The weak mixing angle of the couplings, $\cos\theta_\mathrm{W} = \hat{g}_2/\hat{g}_Z$, $\sin\theta_\mathrm{W} = \hat{g}_1/\hat{g}_Z$; fixed by the measured W and Z masses with $\rho_{\mathrm{EW},0}$. Unit: dimensionless. Value: $\cos^2\theta_\mathrm{W} = 0.776368$. Status: DERIVED.

$$S_\mathrm{tip} = \int_{y=-L} \mathrm{d}^7 X_\mathrm{tip} \sqrt{|g_\mathrm{tip}|}\, \Big[-\frac{v_\Sigma^2}{4} \operatorname{tr}\big((D_\mu \Sigma)^\dagger D^\mu \Sigma\big) - \mathcal{L}_\mathrm{Y} \Big],$$
$$\mathcal{L}_\mathrm{Y} = \sum_{i,j=1}^{3} \Big[\bar{\ell}_i\, \Sigma \begin{pmatrix} (\mathcal{Y}_\nu)_{ij}\, \nu_{Rj} \\ (\mathcal{Y}_e)_{ij}\, e_{Rj} \end{pmatrix} + \bar{Q}_i\, \Sigma \begin{pmatrix} (\mathcal{Y}_u)_{ij}\, u_{Rj} \\ (\mathcal{Y}_d)_{ij}\, d_{Rj} \end{pmatrix} \Big] + \mathrm{h.c.} \tag{IN-13}$$

Symbols in (IN-13).

- S_tip: The action on the tip $y = -L$: the sigma field and the tip Yukawa couplings. Unit: dimensionless. Status: INVENTED.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.

- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- X_{tip}: The seven coordinates $x_1, \ldots, x_7$ of the tip $y = -L$ of the hidden direction; $\mathrm{d}^7 X_{\mathrm{tip}}$ is their coordinate volume element. Unit: length (each coordinate). Status: DERIVED.
- g_{tip}: The metric induced on the seven-dimensional tip $y = -L$ (the metric g_{MN} restricted to the tip directions $x_1, \ldots, x_7$); $|g_{\mathrm{tip}}|$ is the absolute value of its determinant. Unit: dimensionless. Status: DERIVED.
- v_Σ: The scale of the tip sigma field (its kinetic term is $-\frac{v_\Sigma^2}{4}\mathrm{tr}((D_\mu\Sigma)^\dagger D^\mu\Sigma)$); on the static block it fixes the Robin parameter β_{W}. Unit: $\mathrm{GeV}^{5/2}$. Value: fixed through $\beta_{\mathrm{W}} = 0.011912$. Status: INVENTED.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $\mathcal{L}_{\mathrm{Y}}$: The tip Yukawa Lagrangian: the couplings of the doublets to the singlets through the sigma field and the tip matrices $\mathcal{Y}$ (plus the Hermitian conjugate, h.c.). Unit: GeV^7. Status: INVENTED.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- j: A second generation index ($j = 1, 2, 3$). Status: STANDARD.
- ℓ_i, $\bar{\ell}_i$: The lepton doublet of the generation i (left-handed neutrino and charged lepton) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- $\mathcal{Y}_\nu$, $\mathcal{Y}_e$, $\mathcal{Y}_u$, $\mathcal{Y}_d$: The complex 3×3 tip Yukawa matrices of the neutrino, charged-lepton, up-quark and down-quark sectors (generation indices i, j); on the static block they become the tip mixing matrices $\mathcal{K}$ (the normalisation of that relation is not fixed: OPEN). Unit: dimensionless. Status: INVENTED.
- ν_R, ν_{Ri}: The right-handed neutrino (a gauge singlet): ν_R the multiplet, ν_{Ri} that of the generation i; its zero modes give Dirac neutrino masses. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- e_R, e_{Ri}: The right-handed charged-lepton singlet: e_R the multiplet, e_{Ri} that of the generation i (e_R, μ_R, τ_R). Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- Q_i, $\bar{Q}_i$: The quark doublet of the generation i (left-handed up and down quark of one generation, three colours) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- u_R, u_{Ri}, $\bar{u}_{Ri}$: The up-type quark singlet: u_R the multiplet, u_{Ri} that of the generation i (u_R, c_R, t_R), $\bar{u}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- d_R, d_{Ri}, $\bar{d}_{Ri}$: The down-type quark singlet: d_R the multiplet, d_{Ri} that of the generation i (d_R, s_R, b_R), $\bar{d}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.

$$-\frac{v_\Sigma^2}{4}\,\mathrm{tr}\big((D_\mu\Sigma)^\dagger D^\mu\Sigma\big)\Big|_{\Sigma=\mathbf{1}} = -\frac{v_\Sigma^2}{8}\Big[2\hat{g}_2^2\,W_\mu^+ W^{-\mu} + \hat{g}_Z^2\,Z_\mu Z^\mu\Big] \tag{IN-15}$$

Symbols in (IN-15).

- v_Σ: The scale of the tip sigma field (its kinetic term is $-\frac{v_\Sigma^2}{4}\mathrm{tr}((D_\mu\Sigma)^\dagger D^\mu\Sigma)$); on the static block it fixes the Robin parameter β_{W}. Unit: $\mathrm{GeV}^{5/2}$. Value: fixed through $\beta_{\mathrm{W}} = 0.011912$. Status: INVENTED.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.

- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $\mathbf{1}$: The identity matrix (of the size fixed by the context: 16 x 16 on spinors, 3 x 3 on the generations or flavours, 2 x 2 for the sigma field). Unit: dimensionless. Status: STANDARD.
- $\hat{g}_2$: The eight-dimensional SU(2) coupling; $g_2^2 = \hat{g}_2^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- $W_M^\pm$, W^+, W^-: The charged weak bosons, canonically normalised: $W_M^\pm = (W_M^1 \mp \mathrm{i}W_M^2)/(\sqrt{2}\,\hat{g}_2)$; W^+ carries the electric charge $+e$ and W^- the charge $-e$ (they are each other's antiparticles and have one mass). Unit: GeV^3. Status: STANDARD.
- $\hat{g}_Z$: The eight-dimensional coupling of the Z, $\hat{g}_Z = (\hat{g}_2^2 + \hat{g}_1^2)^{1/2}$. Unit: GeV^{-2}. Status: STANDARD.
- Z_M: The Z boson field, canonically normalised, $Z_M = (W_M^3 - B_M)/\hat{g}_Z$. Unit: GeV^3. Status: STANDARD.

The Yang-Mills equations of the gluons, of the $SU(2)$ field and of the hypercharge field, with the fermion currents and the tip current:

$$\nabla_M\Big(\frac{G^{MN}}{\hat{g}_{3,\mathrm{eff}}^2}\Big) - \mathrm{i}\Big[G_M,\ \frac{G^{MN}}{\hat{g}_{3,\mathrm{eff}}^2}\Big] = J_3^{AN}\,t^A \tag{FE-22}$$

Symbols in (FE-22).

- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols $\Gamma^N{}_{MP}$); ∇_M acts on the index M, $\nabla^M = g^{MN}\nabla_N$. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- G_M^A, G_M, G_{MN}, G_{MN}^A: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G_M^A ($A = 1, \ldots, 8$), the matrix $G_M = G_M^A t^A$, the field strength G_{MN} (matrix) and its components G_{MN}^A; $G_{\mu\nu}^A$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\hat{g}_{3,\mathrm{eff}}$, $\hat{g}_{2,\mathrm{eff}}$, $\hat{g}_{1,\mathrm{eff}}$: The field-dependent eight-dimensional couplings produced by I12: $1/\hat{g}_{i,\mathrm{eff}}^2 = 1/\hat{g}_i^2 + 4c\,\bar{\Psi}\Psi/\Lambda_\star^3$ with $c = c_\mathrm{G}, c_\mathrm{W}, c_\mathrm{B}$. Unit: GeV^{-2}. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- J_3^{AN}: The colour current of the quarks (the source of the gluons), summed over the generations. Unit: GeV^7. Status: STANDARD.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- t^A: The generators of SU(3) in the fundamental representation (the Gell-Mann matrices divided by 2), $\mathrm{tr}(t^A t^B) = \frac{1}{2}\delta^{AB}$. Unit: dimensionless. Status: STANDARD.

$$\nabla_M\Big(\frac{W^{MN}}{\hat{g}_{2,\mathrm{eff}}^2}\Big) - \mathrm{i}\Big[W_M,\ \frac{W^{MN}}{\hat{g}_{2,\mathrm{eff}}^2}\Big] = J_2^{IN}\,\frac{\tau^I}{2} + \delta_\mathrm{tip}\,\frac{v_\Sigma^2}{4}\Big(W^N - B^N\,\frac{\tau^3}{2}\Big) \qquad (\Sigma = \mathbf{1}) \tag{FE-23}$$

Symbols in (FE-23).

- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols $\Gamma^N{}_{MP}$); ∇_M acts on the index M, $\nabla^M = g^{MN}\nabla_N$. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- W_M^I, W_M, W_{MN}, W_{MN}^I: The SU(2) gauge field (geometric normalisation): the components W_M^I ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W_M^I \tau^I/2$, the field strength W_{MN} and its components W_{MN}^I; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.

- $\hat{g}_{3,\mathrm{eff}}$, $\hat{g}_{2,\mathrm{eff}}$, $\hat{g}_{1,\mathrm{eff}}$: The field-dependent eight-dimensional couplings produced by I12: $1/\hat{g}^2_{i,\mathrm{eff}} = 1/\hat{g}^2_i + 4c\,\bar{\Psi}\Psi/\Lambda^3_\star$ with $c = c_\mathrm{G}, c_\mathrm{W}, c_\mathrm{B}$. Unit: GeV^{-2}. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- J_2^{IN}: The weak-isospin current of the doublets Q_i, ℓ_i (the source of W^I). Unit: GeV^7. Status: STANDARD.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- τ^I: The Pauli matrices acting on weak-isospin doublets ($\tau^I/2$ the generators of SU(2); $\tau^3 = \mathrm{diag}(1, -1)$). Unit: dimensionless. Status: STANDARD.
- δ_tip: The density that places a tip term in the field equations of eight dimensions, $\delta_\mathrm{tip} = \delta_\mathrm{D}(y + L)/f$ (the Dirac delta function at the tip with the proper length of the hidden direction). Unit: 1/length. Status: DERIVED.
- v_Σ: The scale of the tip sigma field (its kinetic term is $-\frac{v_\Sigma^2}{4}\mathrm{tr}((D_\mu\Sigma)^\dagger D^\mu\Sigma)$); on the static block it fixes the Robin parameter β_W. Unit: $\mathrm{GeV}^{5/2}$. Value: fixed through $\beta_\mathrm{W} = 0.011912$. Status: INVENTED.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- $\mathbf{1}$: The identity matrix (of the size fixed by the context: 16 x 16 on spinors, 3 x 3 on the generations or flavours, 2 x 2 for the sigma field). Unit: dimensionless. Status: STANDARD.

$$\nabla_M\Big(\frac{B^{MN}}{\hat{g}^2_{1,\mathrm{eff}}}\Big) = J_1^N - \delta_\mathrm{tip}\,\frac{v_\Sigma^2}{4}\left(W^{3N} - B^N\right) \qquad (\Sigma = \mathbf{1}) \tag{FE-24}$$

Symbols in (FE-24).

- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols $\Gamma^N{}_{MP}$); ∇_M acts on the index M, $\nabla^M = g^{MN}\nabla_N$. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\hat{g}_{3,\mathrm{eff}}$, $\hat{g}_{2,\mathrm{eff}}$, $\hat{g}_{1,\mathrm{eff}}$: The field-dependent eight-dimensional couplings produced by I12: $1/\hat{g}^2_{i,\mathrm{eff}} = 1/\hat{g}^2_i + 4c\,\bar{\Psi}\Psi/\Lambda^3_\star$ with $c = c_\mathrm{G}, c_\mathrm{W}, c_\mathrm{B}$. Unit: GeV^{-2}. Status: DERIVED.
- J_1^N: The hypercharge current of all fermions (the source of B). Unit: GeV^7. Status: STANDARD.
- δ_tip: The density that places a tip term in the field equations of eight dimensions, $\delta_\mathrm{tip} = \delta_\mathrm{D}(y + L)/f$ (the Dirac delta function at the tip with the proper length of the hidden direction). Unit: 1/length. Status: DERIVED.
- v_Σ: The scale of the tip sigma field (its kinetic term is $-\frac{v_\Sigma^2}{4}\mathrm{tr}((D_\mu\Sigma)^\dagger D^\mu\Sigma)$); on the static block it fixes the Robin parameter β_W. Unit: $\mathrm{GeV}^{5/2}$. Value: fixed through $\beta_\mathrm{W} = 0.011912$. Status: INVENTED.
- W^I_M, W_M, W_{MN}, W^I_{MN}: The SU(2) gauge field (geometric normalisation): the components W^I_M ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W^I_M\tau^I/2$, the field strength W_{MN} and its components W^I_{MN}; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- $\mathbf{1}$: The identity matrix (of the size fixed by the context: 16 x 16 on spinors, 3 x 3 on the generations or flavours, 2 x 2 for the sigma field). Unit: dimensionless. Status: STANDARD.

To linear order in the fields (the non-abelian terms and I12 neglected, as for a mass eigenvalue), they become the field equations of W^+, W^-, Z and the photon with their tip mass terms (the gluon equation, which has no tip term, is that of the photon with the colour index):

$$\begin{aligned}
&\nabla_M W^{\pm MN} = \hat{g}_2\, J_2^{\pm N} + \delta_{\rm tip}\,\frac{v_\Sigma^2}{4}\,\hat{g}_2^2\, W^{\pm N}, \qquad \nabla_M A^{MN} = \hat{e}\, J^N_{\rm em}, \qquad \hat{e} = \frac{\hat{g}_2\hat{g}_1}{\hat{g}_Z},\\
&\nabla_M Z^{MN} = \hat{g}_Z\big(J_2^{3N} - \sin^2\theta_{\rm W}\, J^N_{\rm em}\big) + \delta_{\rm tip}\,\frac{v_\Sigma^2}{4}\,\hat{g}_Z^2\, Z^N,\\
&W^{\pm MN} = \nabla^M W^{\pm N} - \nabla^N W^{\pm M}, \qquad Z^{MN} = \nabla^M Z^N - \nabla^N Z^M, \qquad A^{MN} = \nabla^M A^N - \nabla^N A^M
\end{aligned} \tag{FE-25}$$

Symbols in (FE-25).

- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols $\Gamma^N{}_{MP}$); ∇_M acts on the index M, $\nabla^M = g^{MN}\nabla_N$. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $W^\pm_M$, W^+, W^-: The charged weak bosons, canonically normalised: $W^\pm_M = (W^1_M \mp \mathrm{i}W^2_M)/(\sqrt{2}\,\hat{g}_2)$; W^+ carries the electric charge $+e$ and W^- the charge $-e$ (they are each other's antiparticles and have one mass). Unit: GeV^3. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\hat{g}_2$: The eight-dimensional SU(2) coupling; $g_2^2 = \hat{g}_2^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- $J_2^{\pm N}$: The charged combinations of the isospin current, $J_2^\pm = (J_2^1 \mp \mathrm{i}J_2^2)/\sqrt{2}$ (the sources of $W^\pm$). Unit: GeV^7. Status: STANDARD.
- $\delta_{\rm tip}$: The density that places a tip term in the field equations of eight dimensions, $\delta_{\rm tip} = \delta_{\rm D}(y+L)/f$ (the Dirac delta function at the tip with the proper length of the hidden direction). Unit: 1/length. Status: DERIVED.
- v_Σ: The scale of the tip sigma field (its kinetic term is $-\frac{v_\Sigma^2}{4}\mathrm{tr}((D_\mu\Sigma)^\dagger D^\mu\Sigma)$); on the static block it fixes the Robin parameter $\beta_{\rm W}$. Unit: GeV$^{5/2}$. Value: fixed through $\beta_{\rm W} = 0.011912$. Status: INVENTED.
- A_M: The photon field, canonically normalised, $A_M = (\hat{g}_1^2 W^3_M + \hat{g}_2^2 B_M)/(\hat{g}_1\hat{g}_2\hat{g}_Z)$. Unit: GeV^3. Status: STANDARD.
- $\hat{e}$: The eight-dimensional electric coupling, $\hat{e} = \hat{g}_2\hat{g}_1/\hat{g}_Z$ ($e^2 = \hat{e}^2/V_\perp$). Unit: GeV^{-2}. Status: STANDARD.
- $J^N_{\rm em}$: The electromagnetic current, $J_{\rm em} = J_2^3 + J_1$ (in units of the elementary charge). Unit: GeV^7. Status: STANDARD.
- $\hat{g}_1$: The eight-dimensional U(1)$_Y$ coupling; $g_1^2 = \hat{g}_1^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- $\hat{g}_Z$: The eight-dimensional coupling of the Z, $\hat{g}_Z = (\hat{g}_2^2 + \hat{g}_1^2)^{1/2}$. Unit: GeV^{-2}. Status: STANDARD.
- Z_M: The Z boson field, canonically normalised, $Z_M = (W^3_M - B_M)/\hat{g}_Z$. Unit: GeV^3. Status: STANDARD.
- J_2^{IN}: The weak-isospin current of the doublets Q_i, ℓ_i (the source of W^I). Unit: GeV^7. Status: STANDARD.
- $\theta_{\rm W}$: The weak mixing angle of the couplings, $\cos\theta_{\rm W} = \hat{g}_2/\hat{g}_Z$, $\sin\theta_{\rm W} = \hat{g}_1/\hat{g}_Z$; fixed by the measured W and Z masses with $\rho_{\rm EW,0}$. Unit: dimensionless. Value: $\cos^2\theta_{\rm W} = 0.776368$. Status: DERIVED.

$$\begin{aligned}
&z \to \pi - z: \qquad \partial_y F\big|_{y=0} = 0 \;\;\text{(even)} \quad\text{or}\quad F\big|_{y=0} = 0 \;\;\text{(odd)};\\
&\text{at } y = -L\text{: the natural (Robin) condition of the tip term}
\end{aligned} \tag{FE-27}$$

Symbols in (FE-27).

- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$. Unit: dimensionless. Status: AUTHOR.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- y: The label y in ∂_y: the derivative with respect to the hidden coordinate y. Status: STANDARD.
- $F(\xi)$, F_n: The x_0 profile of a boson mode on one half of the period (a function of y or of $\xi = y/L$): of a gauge field (gauge $A_y = 0$) or of the transverse-traceless graviton; F_n is the profile of the n-th Kaluza-Klein level of the graviton tower ($F_0 = 1$ the flat zero mode). Unit: dimensionless. Status: DERIVED.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.

- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.

The fermions of three generations. The quark and lepton multiplets have bulk masses $\mu_{\psi i}$ (the doublets of a sector share one) and are mixed at the tip by the tip Yukawa couplings:

$$S_{\mathrm{f}} = \sum_{i=1}^{3} \sum_{\psi} \int \mathrm{d}^8X \, \sqrt{|g|} \left[\tfrac{1}{2} \left(\bar{\psi}_i \gamma^M D_M \psi_i - (D_M \bar{\psi}_i) \gamma^M \psi_i \right) - \mu_{\psi i} \, \bar{\psi}_i \psi_i \right], \qquad \text{(IN-11)}$$
$$\psi \in \{Q,\ u_R,\ d_R,\ \ell,\ e_R,\ \nu_R\}, \qquad \mu_{\psi i} = \mu_Q \ (\psi = Q), \qquad \mu_{\psi i} = \mu_\ell \ (\psi = \ell)$$

Symbols in (IN-11).

- S_{f}: The action of the Standard-Model fermions of three generations (kinetic and bulk-mass terms). Unit: dimensionless. Status: ASSUMED.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- ψ: The label of a Standard-Model multiplet: $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ (quark doublet, up and down quark singlets, lepton doublet, charged-lepton singlet, right-handed neutrino); a sum over ψ runs over the six. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0 \, \mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN} \, \mathrm{d}X^M \mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, ${g^M}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|} \, \mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- ψ_i, $\bar{\psi}_i$: A Standard-Model fermion field of eight dimensions: ψ_i is the multiplet $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ of the generation i, ψ_f one fermion component f (in four-dimensional formulas its zero mode); $\bar{\psi} = \psi^\dagger C$ the Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$ (eight dimensions), $\mathrm{GeV}^{3/2}$ (four dimensions). Status: ASSUMED.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4, 4)$; $\gamma^M = {\mathbf{e}_a}^M \gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN} \gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu \Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- $\mu_{\psi i}$: The bulk mass of a Standard-Model fermion: $\mu_{\psi i}$ of the multiplet ψ of the generation i; with the hidden direction it fixes the localisation $c = f\mu L$ of the zero mode. Unit: GeV. Value: $\mu_{\psi i} = c/(fL)$ with the localisation c of the zero mode (doublets: $\mu_Q = 5.2304 \, H$, $\mu_\ell = 9.2055 \, H$). Status: ASSUMED.
- Q_i, $\bar{Q}_i$: The quark doublet of the generation i (left-handed up and down quark of one generation, three colours) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- u_R, u_{Ri}, $\bar{u}_{Ri}$: The up-type quark singlet: u_R the multiplet, u_{Ri} that of the generation i (u_R, c_R, t_R), $\bar{u}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- d_R, d_{Ri}, $\bar{d}_{Ri}$: The down-type quark singlet: d_R the multiplet, d_{Ri} that of the generation i (d_R, s_R, b_R), $\bar{d}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- ℓ_i, $\bar{\ell}_i$: The lepton doublet of the generation i (left-handed neutrino and charged lepton) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- e_R, e_{Ri}: The right-handed charged-lepton singlet: e_R the multiplet, e_{Ri} that of the generation i (e_R, μ_R, τ_R). Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- ν_R, ν_{Ri}: The right-handed neutrino (a gauge singlet): ν_R the multiplet, ν_{Ri} that of the generation i; its zero modes give Dirac neutrino masses. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.

- μ_Q: The common bulk mass of the three quark doublets. Unit: GeV. Value: $5.2304\,H$ ($c_Q = 3.38230$). Status: DERIVED.
- μ_ℓ: The common bulk mass of the three lepton doublets. Unit: GeV. Value: $9.2055\,H$ ($c_\ell = 5.95288$). Status: DERIVED.

$$\gamma^M D_M Q_i = \mu_Q\, Q_i - \mathrm{i}\,\frac{c_\mathrm{V}}{\Lambda_\star^6}\, q_Q\, \big(\mathrm{i}\bar\Psi\gamma_M\Psi\big)\gamma^M Q_i - \frac{\mathrm{i}}{3}\,\frac{c_\mathrm{A}}{\Lambda_\star^6}\sum_{M\neq x_0}\big(\mathrm{i}\bar\Psi\gamma_M P_b\Psi\big)\gamma^M Q_i + \delta_\mathrm{tip}\sum_{j=1}^{3}\Sigma\begin{pmatrix}(\mathcal{Y}_u)_{ij}\,u_{Rj}\\(\mathcal{Y}_d)_{ij}\,d_{Rj}\end{pmatrix} \tag{FE-14}$$

Symbols in (FE-14).

- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- Q_i, $\bar Q_i$: The quark doublet of the generation i (left-handed up and down quark of one generation, three colours) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- μ_Q: The common bulk mass of the three quark doublets. Unit: GeV. Value: $5.2304\,H$ ($c_Q = 3.38230$). Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- c_V: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_\mathrm{V} = 4.38\times10^{-17}$ GeV^{-2}. Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- q_ψ, q_f: The $B-L$ charge: q_ψ of the multiplet ψ ($q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}$, $q_\ell = q_{e_R} = q_{\nu_R} = -1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- Ψ, $\bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- c_A: The coefficient of the x_0-chiral vector portal I13. Unit: dimensionless. Value: fixed through $G_\mathrm{A} = 1.278\times10^{-9}$ GeV^{-2} (dark-matter relic). Status: INVENTED.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- P_a, P_b: The projectors on the two x_0-chiralities, $P_a = (1+\gamma^{x_0})/2$ and $P_b = (1-\gamma^{x_0})/2$ (in a block: the upper and the lower component). Unit: dimensionless. Status: INVENTED.
- δ_tip: The density that places a tip term in the field equations of eight dimensions, $\delta_\mathrm{tip} = \delta_\mathrm{D}(y+L)/f$ (the Dirac delta function at the tip with the proper length of the hidden direction). Unit: 1/length. Status: DERIVED.
- j: A second generation index ($j = 1, 2, 3$). Status: STANDARD.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- $\mathcal{Y}_\nu$, $\mathcal{Y}_e$, $\mathcal{Y}_u$, $\mathcal{Y}_d$: The complex 3×3 tip Yukawa matrices of the neutrino, charged-lepton, up-quark and down-quark sectors (generation indices i, j); on the static block they become the tip mixing matrices $\mathcal{K}$ (the normalisation of that relation is not fixed: OPEN). Unit: dimensionless. Status: INVENTED.
- u_R, u_{Ri}, $\bar u_{Ri}$: The up-type quark singlet: u_R the multiplet, u_{Ri} that of the generation i (u_R, c_R, t_R), $\bar u_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.

- $d_R,\ d_{Ri},\ \bar{d}_{Ri}$: The down-type quark singlet: d_R the multiplet, d_{Ri} that of the generation i (d_R, s_R, b_R), $\bar{d}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.

$$\gamma^M D_M u_{Ri} = \mu_{\psi i}\, u_{Ri} - \mathrm{i}\,\frac{c_\mathrm{V}}{\Lambda_\star^6}\, q_{u_R}\left(\mathrm{i}\bar{\Psi}\gamma_M\Psi\right)\gamma^M u_{Ri} - \frac{\mathrm{i}}{3}\,\frac{c_\mathrm{A}}{\Lambda_\star^6}\sum_{M\neq x_0}\left(\mathrm{i}\bar{\Psi}\gamma_M P_b\Psi\right)\gamma^M u_{Ri} + \delta_\mathrm{tip}\sum_{j=1}^{3}(\mathcal{Y}_u)^*_{ji}\left(\Sigma^\dagger Q_j\right)_1 \qquad (\psi = u_R) \tag{FE-15}$$

Symbols in (FE-15).

- $\gamma^a,\ \gamma^M,\ \gamma^\mu$: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $D_M,\ D_\mu$: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- $u_R,\ u_{Ri},\ \bar{u}_{Ri}$: The up-type quark singlet: u_R the multiplet, u_{Ri} that of the generation i (u_R, c_R, t_R), $\bar{u}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- $\mu_{\psi i}$: The bulk mass of a Standard-Model fermion: $\mu_{\psi i}$ of the multiplet ψ of the generation i; with the hidden direction it fixes the localisation $c = f\mu L$ of the zero mode. Unit: GeV. Value: $\mu_{\psi i} = c/(fL)$ with the localisation c of the zero mode (doublets: $\mu_Q = 5.2304\,H$, $\mu_\ell = 9.2055\,H$). Status: ASSUMED.
- ψ: The label of a Standard-Model multiplet: $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ (quark doublet, up and down quark singlets, lepton doublet, charged-lepton singlet, right-handed neutrino); a sum over ψ runs over the six. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- c_V: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_\mathrm{V} = 4.38\times10^{-17}$ GeV^{-2}. Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- $q_\psi,\ q_f$: The $B-L$ charge: q_ψ of the multiplet ψ ($q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}$, $q_\ell = q_{e_R} = q_{\nu_R} = -1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- $\Psi,\ \bar{\Psi}$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar{\Psi} = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- c_A: The coefficient of the x_0-chiral vector portal I13. Unit: dimensionless. Value: fixed through $G_\mathrm{A} = 1.278\times10^{-9}$ GeV^{-2} (dark-matter relic). Status: INVENTED.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- $P_a,\ P_b$: The projectors on the two x_0-chiralities, $P_a = (1+\gamma^{x_0})/2$ and $P_b = (1-\gamma^{x_0})/2$ (in a block: the upper and the lower component). Unit: dimensionless. Status: INVENTED.
- δ_tip: The density that places a tip term in the field equations of eight dimensions, $\delta_\mathrm{tip} = \delta_\mathrm{D}(y+L)/f$ (the Dirac delta function at the tip with the proper length of the hidden direction). Unit: 1/length. Status: DERIVED.
- j: A second generation index ($j = 1, 2, 3$). Status: STANDARD.
- $\mathcal{Y}_\nu,\ \mathcal{Y}_e,\ \mathcal{Y}_u,\ \mathcal{Y}_d$: The complex 3×3 tip Yukawa matrices of the neutrino, charged-lepton, up-quark and down-quark sectors (generation indices i, j); on the static block they become the tip mixing matrices $\mathcal{K}$ (the normalisation of that relation is not fixed: OPEN). Unit: dimensionless. Status: INVENTED.
- *: Complex conjugation of a number or of each entry of a matrix. Status: STANDARD.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless.

Status: INVENTED.

- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- Q_i, $\bar{Q}_i$: The quark doublet of the generation i (left-handed up and down quark of one generation, three colours) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- ψ_i, $\bar{\psi}_i$: A Standard-Model fermion field of eight dimensions: ψ_i is the multiplet $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ of the generation i, ψ_f one fermion component f (in four-dimensional formulas its zero mode); $\bar{\psi} = \psi^\dagger C$ the Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$ (eight dimensions), $\mathrm{GeV}^{3/2}$ (four dimensions). Status: ASSUMED.

$$\gamma^M D_M d_{Ri} = \mu_{\psi i}\, d_{Ri} - \mathrm{i}\,\frac{c_\mathrm{V}}{\Lambda_\star^6}\, q_{d_R} \left(\mathrm{i}\bar{\Psi}\gamma_M \Psi\right)\gamma^M d_{Ri} - \frac{\mathrm{i}}{3}\,\frac{c_\mathrm{A}}{\Lambda_\star^6} \sum_{M \neq x_0} \left(\mathrm{i}\bar{\Psi}\gamma_M P_b \Psi\right)\gamma^M d_{Ri} + \delta_\mathrm{tip} \sum_{j=1}^{3} (\mathcal{Y}_d)^*_{ji} \left(\Sigma^\dagger Q_j\right)_2 \qquad (\psi = d_R) \tag{FE-16}$$

Symbols in (FE-16).

- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4, 4)$; $\gamma^M = \mathbf{e}_a{}^M \gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- d_R, d_{Ri}, $\bar{d}_{Ri}$: The down-type quark singlet: d_R the multiplet, d_{Ri} that of the generation i (d_R, s_R, b_R), $\bar{d}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- $\mu_{\psi i}$: The bulk mass of a Standard-Model fermion: $\mu_{\psi i}$ of the multiplet ψ of the generation i; with the hidden direction it fixes the localisation $c = f\mu L$ of the zero mode. Unit: GeV. Value: $\mu_{\psi i} = c/(fL)$ with the localisation c of the zero mode (doublets: $\mu_Q = 5.2304\,H$, $\mu_\ell = 9.2055\,H$). Status: ASSUMED.
- ψ: The label of a Standard-Model multiplet: $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ (quark doublet, up and down quark singlets, lepton doublet, charged-lepton singlet, right-handed neutrino); a sum over ψ runs over the six. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- c_V: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_\mathrm{V} = 4.38 \times 10^{-17}$ GeV^{-2}. Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- q_ψ, q_f: The $B - L$ charge: q_ψ of the multiplet ψ ($q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}$, $q_\ell = q_{e_R} = q_{\nu_R} = -1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- Ψ, $\bar{\Psi}$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar{\Psi} = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- c_A: The coefficient of the x_0-chiral vector portal I13. Unit: dimensionless. Value: fixed through $G_\mathrm{A} = 1.278 \times 10^{-9}$ GeV^{-2} (dark-matter relic). Status: INVENTED.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- P_a, P_b: The projectors on the two x_0-chiralities, $P_a = (1 + \gamma^{x_0})/2$ and $P_b = (1 - \gamma^{x_0})/2$ (in a block: the upper and the lower component). Unit: dimensionless. Status: INVENTED.
- δ_tip: The density that places a tip term in the field equations of eight dimensions, $\delta_\mathrm{tip} = \delta_\mathrm{D}(y + L)/f$ (the Dirac delta function at the tip with the proper length of the hidden direction). Unit: 1/length. Status: DERIVED.
- j: A second generation index ($j = 1, 2, 3$). Status: STANDARD.

- $\mathcal{Y}_\nu,\ \mathcal{Y}_e,\ \mathcal{Y}_u,\ \mathcal{Y}_d$: The complex 3×3 tip Yukawa matrices of the neutrino, charged-lepton, up-quark and down-quark sectors (generation indices i, j); on the static block they become the tip mixing matrices $\mathcal{K}$ (the normalisation of that relation is not fixed: OPEN). Unit: dimensionless. Status: INVENTED.
- *: Complex conjugation of a number or of each entry of a matrix. Status: STANDARD.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma=\mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma=\mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $Q_i,\ \bar{Q}_i$: The quark doublet of the generation i (left-handed up and down quark of one generation, three colours) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- $\psi_i,\ \bar{\psi}_i$: A Standard-Model fermion field of eight dimensions: ψ_i is the multiplet $\psi\in\{Q,u_R,d_R,\ell,e_R,\nu_R\}$ of the generation i, ψ_f one fermion component f (in four-dimensional formulas its zero mode); $\bar{\psi}=\psi^\dagger C$ the Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$ (eight dimensions), $\mathrm{GeV}^{3/2}$ (four dimensions). Status: ASSUMED.

$$\gamma^M D_M \ell_i = \mu_\ell\,\ell_i - \mathrm{i}\,\frac{c_\mathrm{V}}{\Lambda_\star^6}\,q_\ell\,\big(\mathrm{i}\bar{\Psi}\gamma_M\Psi\big)\gamma^M\ell_i + \delta_\mathrm{tip}\sum_{j=1}^{3}\Sigma\begin{pmatrix}(\mathcal{Y}_\nu)_{ij}\,\nu_{Rj}\\(\mathcal{Y}_e)_{ij}\,e_{Rj}\end{pmatrix} \tag{FE-17}$$

Symbols in (FE-17).

- $\gamma^a,\ \gamma^M,\ \gamma^\mu$: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M=\mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M=g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $D_M,\ D_\mu$: The covariant derivative: on a spinor $D_M=\partial_M+\Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- $\ell_i,\ \bar{\ell}_i$: The lepton doublet of the generation i (left-handed neutrino and charged lepton) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- i: The generation index ($i=1,2,3$), which also labels the neutrino mass states ν_1,ν_2,ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- μ_ℓ: The common bulk mass of the three lepton doublets. Unit: GeV. Value: $9.2055\,H$ ($c_\ell=5.95288$). Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- c_V: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_\mathrm{V}=4.38\times10^{-17}$ GeV^{-2}. Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- $q_\psi,\ q_f$: The $B-L$ charge: q_ψ of the multiplet ψ ($q_Q=q_{u_R}=q_{d_R}=\frac{1}{3}$, $q_\ell=q_{e_R}=q_{\nu_R}=-1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- $\Psi,\ \bar{\Psi}$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar{\Psi}=\Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- δ_tip: The density that places a tip term in the field equations of eight dimensions, $\delta_\mathrm{tip}=\delta_\mathrm{D}(y+L)/f$ (the Dirac delta function at the tip with the proper length of the hidden direction). Unit: 1/length. Status: DERIVED.
- j: A second generation index ($j=1,2,3$). Status: STANDARD.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma=\mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma=\mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- $\mathcal{Y}_\nu,\ \mathcal{Y}_e,\ \mathcal{Y}_u,\ \mathcal{Y}_d$: The complex 3×3 tip Yukawa matrices of the neutrino, charged-lepton, up-quark and down-quark sectors (generation indices i, j); on the static block they become the tip mixing matrices $\mathcal{K}$ (the normalisation of that relation is not fixed: OPEN). Unit: dimensionless. Status: INVENTED.

- ν_R, ν_{Ri}: The right-handed neutrino (a gauge singlet): ν_R the multiplet, ν_{Ri} that of the generation i; its zero modes give Dirac neutrino masses. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- e_R, e_{Ri}: The right-handed charged-lepton singlet: e_R the multiplet, e_{Ri} that of the generation i (e_R, μ_R, τ_R). Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.

$$\gamma^M D_M e_{Ri} = \mu_{\psi i}\, e_{Ri} - \mathrm{i}\,\frac{c_\mathrm{V}}{\Lambda_\star^6}\, q_{e_R}\left(\mathrm{i}\bar{\Psi}\gamma_M\Psi\right)\gamma^M e_{Ri} + \delta_\mathrm{tip}\sum_{j=1}^{3}(\mathcal{Y}_e)^*_{ji}\left(\Sigma^\dagger \ell_j\right)_2 \qquad (\psi = e_R) \tag{FE-18}$$

Symbols in (FE-18).

- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- e_R, e_{Ri}: The right-handed charged-lepton singlet: e_R the multiplet, e_{Ri} that of the generation i (e_R, μ_R, τ_R). Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- $\mu_{\psi i}$: The bulk mass of a Standard-Model fermion: $\mu_{\psi i}$ of the multiplet ψ of the generation i; with the hidden direction it fixes the localisation $c = f\mu L$ of the zero mode. Unit: GeV. Value: $\mu_{\psi i} = c/(fL)$ with the localisation c of the zero mode (doublets: $\mu_Q = 5.2304\,H$, $\mu_\ell = 9.2055\,H$). Status: ASSUMED.
- ψ: The label of a Standard-Model multiplet: $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ (quark doublet, up and down quark singlets, lepton doublet, charged-lepton singlet, right-handed neutrino); a sum over ψ runs over the six. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- c_V: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_\mathrm{V} = 4.38\times10^{-17}$ GeV^{-2}. Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- q_ψ, q_f: The $B-L$ charge: q_ψ of the multiplet ψ ($q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}$, $q_\ell = q_{e_R} = q_{\nu_R} = -1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- Ψ, $\bar{\Psi}$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar{\Psi} = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- δ_tip: The density that places a tip term in the field equations of eight dimensions, $\delta_\mathrm{tip} = \delta_\mathrm{D}(y+L)/f$ (the Dirac delta function at the tip with the proper length of the hidden direction). Unit: 1/length. Status: DERIVED.
- j: A second generation index ($j = 1, 2, 3$). Status: STANDARD.
- $\mathcal{Y}_\nu$, $\mathcal{Y}_e$, $\mathcal{Y}_u$, $\mathcal{Y}_d$: The complex 3×3 tip Yukawa matrices of the neutrino, charged-lepton, up-quark and down-quark sectors (generation indices i, j); on the static block they become the tip mixing matrices $\mathcal{K}$ (the normalisation of that relation is not fixed: OPEN). Unit: dimensionless. Status: INVENTED.
- *: Complex conjugation of a number or of each entry of a matrix. Status: STANDARD.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- ℓ_i, $\bar{\ell}_i$: The lepton doublet of the generation i (left-handed neutrino and charged lepton) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- ψ_i, $\bar{\psi}_i$: A Standard-Model fermion field of eight dimensions: ψ_i is the multiplet $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ of

the generation i, ψ_f one fermion component f (in four-dimensional formulas its zero mode); $\bar\psi = \psi^\dagger C$ the Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$ (eight dimensions), $\mathrm{GeV}^{3/2}$ (four dimensions). Status: ASSUMED.

$$\gamma^M D_M \nu_{Ri} = \mu_{\psi i}\,\nu_{Ri} - \mathrm{i}\,\frac{c_\mathrm{V}}{\Lambda_\star^6}\,q_{\nu_R}\left(\mathrm{i}\bar\Psi\gamma_M\Psi\right)\gamma^M\nu_{Ri} + \delta_\mathrm{tip}\sum_{j=1}^{3}(\mathcal{Y}_\nu)^*_{ji}\left(\Sigma^\dagger\ell_j\right)_1 \qquad (\psi=\nu_R) \tag{FE-19}$$

Symbols in (FE-19).

- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- ν_R, ν_{Ri}: The right-handed neutrino (a gauge singlet): ν_R the multiplet, ν_{Ri} that of the generation i; its zero modes give Dirac neutrino masses. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- $\mu_{\psi i}$: The bulk mass of a Standard-Model fermion: $\mu_{\psi i}$ of the multiplet ψ of the generation i; with the hidden direction it fixes the localisation $c = f\mu L$ of the zero mode. Unit: GeV. Value: $\mu_{\psi i} = c/(fL)$ with the localisation c of the zero mode (doublets: $\mu_Q = 5.2304\,H$, $\mu_\ell = 9.2055\,H$). Status: ASSUMED.
- ψ: The label of a Standard-Model multiplet: $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ (quark doublet, up and down quark singlets, lepton doublet, charged-lepton singlet, right-handed neutrino); a sum over ψ runs over the six. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- c_V: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_\mathrm{V} = 4.38\times10^{-17}$ GeV^{-2}. Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- q_ψ, q_f: The $B-L$ charge: q_ψ of the multiplet ψ ($q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}$, $q_\ell = q_{e_R} = q_{\nu_R} = -1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- Ψ, $\bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- δ_tip: The density that places a tip term in the field equations of eight dimensions, $\delta_\mathrm{tip} = \delta_\mathrm{D}(y+L)/f$ (the Dirac delta function at the tip with the proper length of the hidden direction). Unit: 1/length. Status: DERIVED.
- j: A second generation index ($j = 1, 2, 3$). Status: STANDARD.
- $\mathcal{Y}_\nu$, $\mathcal{Y}_e$, $\mathcal{Y}_u$, $\mathcal{Y}_d$: The complex 3×3 tip Yukawa matrices of the neutrino, charged-lepton, up-quark and down-quark sectors (generation indices i, j); on the static block they become the tip mixing matrices $\mathcal{K}$ (the normalisation of that relation is not fixed: OPEN). Unit: dimensionless. Status: INVENTED.
- *: Complex conjugation of a number or of each entry of a matrix. Status: STANDARD.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- ℓ_i, $\bar\ell_i$: The lepton doublet of the generation i (left-handed neutrino and charged lepton) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- ψ_i, $\bar\psi_i$: A Standard-Model fermion field of eight dimensions: ψ_i is the multiplet $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ of the generation i, ψ_f one fermion component f (in four-dimensional formulas its zero mode); $\bar\psi = \psi^\dagger C$ the Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$ (eight dimensions), $\mathrm{GeV}^{3/2}$ (four dimensions). Status: ASSUMED.

$$P_{\rm sel}=\frac{1+\Gamma_4\gamma^{x_0}}{2}\,\frac{1-\mathrm{i}\gamma^{x_5}\gamma^{x_6}}{2};$$
$$\text{doublets } (Q,\ell): \quad P_{\rm sel}\big(1-\gamma^{x_0}\big)\psi\big|_{y=0}=\big(1-P_{\rm sel}\big)\big(1+\gamma^{x_0}\big)\psi\big|_{y=0}=0, \qquad \big(1-\gamma^{x_0}\big)\psi\big|_{y=-L}=0;$$
$$\text{singlets } (u_R,d_R,e_R,\nu_R): \quad P_{\rm sel}\big(1+\gamma^{x_0}\big)\psi\big|_{y=0}=\big(1-P_{\rm sel}\big)\big(1-\gamma^{x_0}\big)\psi\big|_{y=0}=0, \qquad \big(1+\gamma^{x_0}\big)\psi\big|_{y=-L}=0 \tag{FE-29}$$

Symbols in (FE-29).

- $P_{\rm sel}$: The projector that selects the two blocks of the half $\mathcal{B}=-1$ in which the Standard-Model zero modes live, $P_{\rm sel}=\frac{1+\Gamma_4\gamma^{x_0}}{2}\,\frac{1-\mathrm{i}\gamma^{x_5}\gamma^{x_6}}{2}$. Unit: dimensionless. Status: INVENTED.
- Γ_4: The four-dimensional chirality matrix, $\Gamma_4=\mathrm{i}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}\gamma^{x_4}$; a doublet zero mode has $\Gamma_4=+1$ (left-handed), a singlet zero mode $\Gamma_4=-1$ (right-handed). Unit: dimensionless. Status: DERIVED.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M=\mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M=g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z=6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- Q_i, $\bar{Q}_i$: The quark doublet of the generation i (left-handed up and down quark of one generation, three colours) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- ℓ_i, $\bar{\ell}_i$: The lepton doublet of the generation i (left-handed neutrino and charged lepton) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- ψ_i, $\bar{\psi}_i$: A Standard-Model fermion field of eight dimensions: ψ_i is the multiplet $\psi\in\{Q,u_R,d_R,\ell,e_R,\nu_R\}$ of the generation i, ψ_f one fermion component f (in four-dimensional formulas its zero mode); $\bar{\psi}=\psi^\dagger C$ the Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$ (eight dimensions), $\mathrm{GeV}^{3/2}$ (four dimensions). Status: ASSUMED.
- y: The hidden coordinate of the y chart, $y=\ln|\sin z|/(6H)$, with $y=0$ at the fold ($z=\pi/2$) and $y=-L$ at the regular tip; each half of a period of x_0 is the interval $-L\le y\le0$. Unit: length. Status: DERIVED.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y=-L$; $HL=3$. Unit: length. Value: $L=3/H$ ($fL=8.926\times10^{-21}$ m at the stabilised block). Status: ASSUMED.
- u_R, u_{Ri}, $\bar{u}_{Ri}$: The up-type quark singlet: u_R the multiplet, u_{Ri} that of the generation i (u_R, c_R, t_R), $\bar{u}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- d_R, d_{Ri}, $\bar{d}_{Ri}$: The down-type quark singlet: d_R the multiplet, d_{Ri} that of the generation i (d_R, s_R, b_R), $\bar{d}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- e_R, e_{Ri}: The right-handed charged-lepton singlet: e_R the multiplet, e_{Ri} that of the generation i (e_R, μ_R, τ_R). Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- ν_R, ν_{Ri}: The right-handed neutrino (a gauge singlet): ν_R the multiplet, ν_{Ri} that of the generation i; its zero modes give Dirac neutrino masses. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.

$$\text{one generation:}\quad b_L(-L)=\mathcal{K}\,b_R(-L), \qquad a_R(-L)=-\mathcal{K}\,a_L(-L), \qquad \chi=(a,\ \mathrm{i}b);$$
$$\text{three generations:}\quad b_L(-L)=\mathcal{K}\,b_R(-L), \qquad a_R(-L)=-\mathcal{K}^\dagger a_L(-L), \qquad \mathcal{K}\in\{\mathcal{K}_\nu,\mathcal{K}_e,\mathcal{K}_u,\mathcal{K}_d\}; \tag{FE-30}$$
$$\text{UM-2:}\quad \mathcal{K}_e=\mathcal{K}_u=\mathbf{1}, \qquad \mathcal{K}_\nu=U, \qquad \mathcal{K}_d=V_{\rm CKM}$$

Symbols in (FE-30).

- a, b: The upper and lower components of a block spinor, $\chi=(a,\mathrm{i}b)$, real at $k=0$: a_L,b_L of a doublet and a_R,b_R of a singlet solution (a_{Ri}, b_{Ri} of the generation i, $b_{L\alpha}$, $a_{L\alpha}$ of the doublet in the flavour basis), a_o,b_o of the orbital o. Unit: $\mathrm{GeV}^{1/2}$ (normalised orbital). Status: DERIVED.
- L: The chirality label L: the left-handed (doublet) zero mode, or the left-handed projection $P_L=(1+\Gamma_4)/2$. Status: STANDARD.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y=-L$; $HL=3$. Unit: length. Value: $L=3/H$ ($fL=8.926\times10^{-21}$ m at the stabilised block). Status: ASSUMED.

- $\mathcal{K}$: The tip mixing matrix of a sector in the boundary conditions of the fermion zero modes (a number $\mathcal{K}$ for one generation, a 3×3 matrix for three): $\mathcal{K}_\nu$, $\mathcal{K}_e$, $\mathcal{K}_u$, $\mathcal{K}_d$ of the neutrino, charged-lepton, up-quark and down-quark sectors; UM-2 takes $\mathcal{K}_e = \mathcal{K}_u = \mathbf{1}$, $\mathcal{K}_\nu = U$, $\mathcal{K}_d = V_{\mathrm{CKM}}$. Unit: dimensionless. Status: ASSUMED.
- R: The chirality label R: the right-handed (singlet) zero mode or field, or the projection $P_R = (1 - \Gamma_4)/2$. Status: STANDARD.
- χ, χ_o, χ_1, χ_2: The two-component spinor of one block of the reduction (a function of y and x_4), $\chi = (\chi_1, \chi_2)^T = (a, \mathrm{i}b)^T$; χ_o the Kohn-Sham orbital o, normalised as $\int_{-L}^{0} \chi_o^\dagger \chi_o \, \mathrm{d}y = 1$. Unit: $\mathrm{GeV}^{1/2}$ (normalised orbital). Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $\mathbf{1}$: The identity matrix (of the size fixed by the context: 16 x 16 on spinors, 3 x 3 on the generations or flavours, 2 x 2 for the sigma field). Unit: dimensionless. Status: STANDARD.
- U: The PMNS matrix, the unitary 3×3 mixing matrix of the leptons, $\nu_\alpha = \sum_i U_{\alpha i} \nu_i$ (rows $\alpha = e, \mu, \tau$, columns the mass states i); in UM-2 $U = \mathcal{K}_e^\dagger \mathcal{K}_\nu$. Unit: dimensionless. Value: $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$, $\delta_{\mathrm{CP}} = 212^\circ$ (NuFIT 6.0). Status: STANDARD.
- V_{CKM}: The CKM matrix, the unitary 3×3 mixing matrix of the quarks (the same parametrisation as U); in UM-2 $V_{\mathrm{CKM}} = \mathcal{K}_u^\dagger \mathcal{K}_d$. Unit: dimensionless. Value: $s_{12} = 0.22501$, $s_{13} = 0.003732$, $s_{23} = 0.04183$, $\delta = 1.147$ (PDG). Status: STANDARD.

dirac16complex obeys its own Dirac equation with the canonical spin connection, the self-coupling and the portals; for its free spectrum the self-coupling and the portal fields are set to zero (their effect is step 3 and step 8):

$$\gamma^M D_M \Psi = \Big[m_\Psi + \lambda \, \bar\Psi \Psi + \frac{\mathcal{F}}{\Lambda_\star^3} \Big] \Psi - \mathrm{i} \, \frac{c_{\mathrm{V}}}{\Lambda_\star^6} \, J_M^{B-L} \gamma^M \Psi - \mathrm{i} \, \frac{c_{\mathrm{A}}}{\Lambda_\star^6} \sum_{M \neq x_0} J_M^B \gamma^M P_b \Psi \qquad \text{(FE-12)}$$

Symbols in (FE-12).

- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4, 4)$; $\gamma^M = \mathbf{e}_a{}^M \gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN} \gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu \Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- Ψ, $\bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- m_Ψ: The mass parameter of dirac16complex, $m_\Psi = 3H\mathcal{M} = H$. Unit: GeV. Value: 14.296 TeV. Status: RECORD.
- λ: The contact self-coupling of dirac16complex (the term $-\frac{\lambda}{2}(\bar\Psi\Psi)^2$ of the record). Unit: GeV^{-6}. Value: not fixed; Kohn-Sham states computed for $\lambda = 0$ and $\pm 7 \times 10^{-5} H^{-6}$; the fold-band radiation requires $|\lambda| \lesssim 1.4 \times 10^{-10} H^{-6}$ (for $\Delta N_{\mathrm{eff}} \leq 0.001$). Status: RECORD.
- $\mathcal{F}$: The gauge-kinetic operator of the portal I12, $\mathcal{F} = c_{\mathrm{G}} G^A_{MN} G^{AMN} + c_{\mathrm{W}} W^I_{MN} W^{IMN} + c_{\mathrm{B}} B_{MN} B^{MN}$; $\langle \mathcal{F} \rangle$ its mean value in an environment. Unit: GeV^4. Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- c_{V}: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_{\mathrm{V}} = 4.38 \times 10^{-17}$ GeV^{-2}. Status: INVENTED.
- J_M^{B-L}: The $B - L$ current of the Standard-Model fermions (baryon minus lepton number), summed over the generations. Unit: GeV^7. Status: STANDARD.

- c_{A}: The coefficient of the x_0-chiral vector portal I13. Unit: dimensionless. Value: fixed through $G_{\mathrm{A}} = 1.278 \times 10^{-9}$ GeV^{-2} (dark-matter relic). Status: INVENTED.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- J^B_M: The baryon-number current of the quarks (one third per quark), summed over the generations. Unit: GeV^7. Status: STANDARD.
- P_a, P_b: The projectors on the two x_0-chiralities, $P_a = (1+\gamma^{x_0})/2$ and $P_b = (1-\gamma^{x_0})/2$ (in a block: the upper and the lower component). Unit: dimensionless. Status: INVENTED.

$$\big(1-\gamma^{x_0}\big)\Psi\big|_{y=0} = 0 \quad \text{(even)} \quad \text{or} \quad \big(1+\gamma^{x_0}\big)\Psi\big|_{y=0} = 0 \quad \text{(odd)}, \qquad \big(1-\gamma^{x_0}\big)\Psi\big|_{y=-L} = 0 \tag{FE-28}$$

Symbols in (FE-28).

- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M \gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- Ψ, $\bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi = \Psi^\dagger C$ is its Dirac adjoint. Unit: GeV$^{7/2}$. Status: RECORD.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.

5.2 The approximation of step 2

The x_0 reductions below are exact on the static block (order 0 of the metric series): no approximation enters the eigenvalue problems themselves, and the eigenvalues are computed as exact roots. What step 2 leaves out is, first, *the mixing of the zero modes with the Kaluza-Klein modes* through the interactions, of the relative size $(m/m_{\mathrm{KK}})^2$, largest for the top quark:

$$\Big(\frac{m_{\mathrm{t}}}{m_{\mathrm{KK}}}\Big)^2 = 2.0 \times 10^{-6}$$

Symbols in this equation.

- m_{u}, m_{d}, m_{s}, m_{c}, m_{b}, m_{t}: The masses of the up, down, strange, charm, bottom and top quarks (the upright subscript names the quark). Unit: GeV. Value: 2.16 MeV, 4.70 MeV, 93.5 MeV, 1.273 GeV, 4.183 GeV, 172.56 GeV (the x_0 masses, fitted to the PDG values). Status: DERIVED.
- m^{KK}_n, m_{KK}: The mass of the n-th Kaluza-Klein graviton (and gauge) level, $m^{\mathrm{KK}}_n = \tilde m_n/(fL)$; $m_{\mathrm{KK}} = m^{\mathrm{KK}}_1$ is the Kaluza-Klein scale. Unit: GeV. Value: $m_{\mathrm{KK}} = 121.32$ TeV. Status: DERIVED.

It further leaves out the following.

- *Radiative corrections.* The masses are tree-level eigenvalues fitted to the measured values used as tree-level targets; the electroweak parameter $\rho_{\mathrm{EW},0} - 1 = 3.8 \times 10^{-4}$ is fitted, not computed from loops.
- *The mean fields of the environment.* The W eigenvalue moves by the dark-matter kinetic function of step 8 by at most 1.9×10^{-17} of itself (at the dark-matter freeze-out), and the dark-matter level by the Standard-Model mean fields by at most 1.5×10^{-6} of itself (step 8).
- *The Lovelock corrections to the graviton operator* are not estimated (OPEN): the graviton profiles below are those of Einstein gravity on the block.

The first-order formula of the fermion masses (written out below) is not used for the masses; it differs from the exact roots by up to 1.1×10^{-3} (top quark) and serves for the block sensitivities of step 8.

5.3 The solution of step 2

5.3.1 The x_0 eigenvalue problem of the gauge bosons and the graviton

A four-dimensional component with the x_0 profile $F(y)$ (in the gauge with no y component) has a reduced action whose stationarity gives a Sturm-Liouville problem in $\xi = y/L$; the tip mass term becomes the Robin condition at $\xi = -1$ (DERIVED):

$$\begin{aligned} &\int_{-L}^{0} \mathrm{d}y\, f^4 \mathrm{e}^{3Hy}\Big[\frac{(\partial_y F)^2}{f^2} - m^2 F^2\Big] \text{ stationary}: \qquad -\frac{1}{f^2}\,\mathrm{e}^{-3Hy}\,\partial_y\big(\mathrm{e}^{3Hy}\,\partial_y F\big) = m^2 F, \\ &\Leftrightarrow \qquad -\mathrm{e}^{-3HL\xi}\,\frac{\mathrm{d}}{\mathrm{d}\xi}\Big(\mathrm{e}^{3HL\xi}\,\frac{\mathrm{d}F}{\mathrm{d}\xi}\Big) = \tilde{m}^2 F, \qquad \tilde{m} = m\,fL \end{aligned} \tag{FE-50}$$

Symbols in (FE-50).

- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-\alpha_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- y: The label y in ∂_y: the derivative with respect to the hidden coordinate y. Status: STANDARD.
- $F(\xi)$, F_n: The x_0 profile of a boson mode on one half of the period (a function of y or of $\xi = y/L$): of a gauge field (gauge $A_y = 0$) or of the transverse-traceless graviton; F_n is the profile of the n-th Kaluza-Klein level of the graviton tower ($F_0 = 1$ the flat zero mode). Unit: dimensionless. Status: DERIVED.
- m, m_s, m_i: A four-dimensional mass: m the mass of a mode (the eigenvalue of an x_0 problem) or of a fermion; m_s the mass of the species s; m_i the mass of the neutrino mass state i. Unit: GeV. Status: DERIVED.
- ξ: The scaled hidden coordinate, $\xi = y/L$, running from -1 (tip) to 0 (fold) on each half of a period. Unit: dimensionless. Status: DERIVED.
- $\tilde{m}$, $\tilde{m}_n$: The dimensionless x_0 eigenvalue of a boson mode, $\tilde{m} = m\,fL$; $\tilde{m}_n$ the n-th even Kaluza-Klein level of the tower with $\beta = 0$ ($\tilde{m}_1$ the first). Unit: dimensionless. Value: $\tilde{m}_1^2 = 30.1196$, $\tilde{m}_2^2 = 59.7284$, $\tilde{m}_3^2 = 109.0764$ ($HL = 3$), i.e. $\tilde{m}_1 = 5.48813$, $\tilde{m}_2 = 7.72842$, $\tilde{m}_3 = 10.44397$. Status: DERIVED.

$$\begin{aligned} &\frac{\mathrm{d}F}{\mathrm{d}\xi}\Big|_{\xi=0} = 0 \;\; \text{(even)} \quad \text{or} \quad F(0) = 0 \;\; \text{(odd)}, \qquad \frac{\mathrm{d}F}{\mathrm{d}\xi}\Big|_{\xi=-1} = \beta\, F(-1), \\ &\beta = 0 \;\; \text{(photon, gluons, graviton)}, \qquad \beta = \beta_\mathrm{W} \;\; (W^\pm), \qquad \beta = \beta_\mathrm{Z} = \frac{\beta_\mathrm{W}}{\cos^2\theta_\mathrm{W}} \;\; (Z) \end{aligned} \tag{FE-51}$$

Symbols in (FE-51).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $F(\xi)$, F_n: The x_0 profile of a boson mode on one half of the period (a function of y or of $\xi = y/L$): of a gauge field (gauge $A_y = 0$) or of the transverse-traceless graviton; F_n is the profile of the n-th Kaluza-Klein level of the graviton tower ($F_0 = 1$ the flat zero mode). Unit: dimensionless. Status: DERIVED.

- ξ: The scaled hidden coordinate, $\xi = y/L$, running from -1 (tip) to 0 (fold) on each half of a period. Unit: dimensionless. Status: DERIVED.
- β: The Robin parameter of the tip condition of a boson profile, $\mathrm{d}F/\mathrm{d}\xi = \beta F$ at $\xi = -1$ (zero for the massless gauge fields and the graviton). Unit: dimensionless. Status: DERIVED.
- β_{W}: The Robin parameter of W^+ and W^- (set by the tip mass term of the sigma field). Unit: dimensionless. Value: 0.011912 (fitted). Status: DERIVED.
- $W_M^{\pm}$, W^+, W^-: The charged weak bosons, canonically normalised: $W_M^{\pm} = (W_M^1 \mp \mathrm{i}W_M^2)/(\sqrt{2}\,\hat{g}_2)$; W^+ carries the electric charge $+e$ and W^- the charge $-e$ (they are each other's antiparticles and have one mass). Unit: GeV^3. Status: STANDARD.
- β_{Z}: The Robin parameter of the Z, $\beta_{\mathrm{Z}} = \beta_{\mathrm{W}}/\cos^2\theta_{\mathrm{W}}$. Unit: dimensionless. Value: 0.015343. Status: DERIVED.
- θ_{W}: The weak mixing angle of the couplings, $\cos\theta_{\mathrm{W}} = \hat{g}_2/\hat{g}_Z$, $\sin\theta_{\mathrm{W}} = \hat{g}_1/\hat{g}_Z$; fixed by the measured W and Z masses with $\rho_{\mathrm{EW},0}$. Unit: dimensionless. Value: $\cos^2\theta_{\mathrm{W}} = 0.776368$. Status: DERIVED.
- Z_M: The Z boson field, canonically normalised, $Z_M = (W_M^3 - B_M)/\hat{g}_Z$. Unit: GeV^3. Status: STANDARD.

$$
\begin{aligned}
&F = \mathrm{e}^{-3HL\xi/2}\,\tilde{F}, \qquad \varpi^2 = \tilde{m}^2 - \frac{9H^2L^2}{4}\,:\\
&\text{even: } \Big[\varpi^2 + \frac{3HL}{2}\Big(\beta + \frac{3HL}{2}\Big)\Big]\sin\varpi - \beta\,\varpi\cos\varpi = 0, \qquad \text{odd: } \varpi\cos\varpi + \Big(\beta + \frac{3HL}{2}\Big)\sin\varpi = 0; \qquad \text{(FE-52)}\\
&\beta = 0: \quad \tilde{m} = 0 \text{ (flat profile)}, \qquad \tilde{m}_n^2 = \frac{9H^2L^2}{4} + \varpi_n^2 \text{ (even)}, \qquad \varpi\cot\varpi = -\frac{3HL}{2} \text{ (odd)}
\end{aligned}
$$

Symbols in (FE-52).

- $F(\xi)$, F_n: The x_0 profile of a boson mode on one half of the period (a function of y or of $\xi = y/L$): of a gauge field (gauge $A_y = 0$) or of the transverse-traceless graviton; F_n is the profile of the n-th Kaluza-Klein level of the graviton tower ($F_0 = 1$ the flat zero mode). Unit: dimensionless. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- ξ: The scaled hidden coordinate, $\xi = y/L$, running from -1 (tip) to 0 (fold) on each half of a period. Unit: dimensionless. Status: DERIVED.
- $\tilde{F}$: The rescaled boson profile, $F = \mathrm{e}^{-3HL\xi/2}\tilde{F}$. Unit: dimensionless. Status: DERIVED.
- ϖ, ϖ_n: The wave number of an oscillating x_0 profile on $-1 \le \xi \le 0$ (of a boson after the rescaling, or of a bulk level of dirac16complex); ϖ_n that of the n-th even level ($\varpi_n = n\pi$ for $\beta = 0$). Unit: dimensionless. Value: 1.89901 for the lowest odd level of dirac16complex. Status: DERIVED.
- $\tilde{m}$, $\tilde{m}_n$: The dimensionless x_0 eigenvalue of a boson mode, $\tilde{m} = m\,fL$; $\tilde{m}_n$ the n-th even Kaluza-Klein level of the tower with $\beta = 0$ ($\tilde{m}_1$ the first). Unit: dimensionless. Value: $\tilde{m}_1^2 = 30.1196$, $\tilde{m}_2^2 = 59.7284$, $\tilde{m}_3^2 = 109.0764$ ($HL = 3$), i.e. $\tilde{m}_1 = 5.48813$, $\tilde{m}_2 = 7.72842$, $\tilde{m}_3 = 10.44397$. Status: DERIVED.
- β: The Robin parameter of the tip condition of a boson profile, $\mathrm{d}F/\mathrm{d}\xi = \beta F$ at $\xi = -1$ (zero for the massless gauge fields and the graviton). Unit: dimensionless. Status: DERIVED.
- n: The level number of the Kaluza-Klein tower ($n = 1, 2, 3, \ldots$; $n = 0$ the flat zero mode). Status: DERIVED.

The operator with $\beta = 0$ is also the x_0 operator of the graviton; its levels form the Kaluza-Klein tower, whose first level fixes the scale m_{KK} of UM-2:

$$
\begin{aligned}
&F_n(\xi) = \mathrm{e}^{-3HL\xi/2}\Big(\cos\varpi_n\xi + \frac{3HL}{2\varpi_n}\sin\varpi_n\xi\Big), \qquad \varpi_n = \pi,\ 2\pi,\ 3\pi,\ \ldots,\\
&\tilde{m}_n^2 = \frac{9H^2L^2}{4} + \varpi_n^2, \qquad m_n^{\mathrm{KK}} = \frac{\tilde{m}_n}{fL}, \qquad N_n = \int_{-1}^{0}\mathrm{e}^{3HL\xi}F_n^2\,\mathrm{d}\xi = \frac{1}{2}\Big(1 + \frac{9H^2L^2}{4\varpi_n^2}\Big), \qquad N_0 = \frac{1 - \mathrm{e}^{-3HL}}{3HL}
\end{aligned}
\qquad \text{(FE-53)}
$$

Symbols in (FE-53).

- $F(\xi)$, F_n: The x_0 profile of a boson mode on one half of the period (a function of y or of $\xi = y/L$): of a gauge field (gauge $A_y = 0$) or of the transverse-traceless graviton; F_n is the profile of the n-th Kaluza-Klein level of the graviton tower ($F_0 = 1$ the flat zero mode). Unit: dimensionless. Status: DERIVED.
- n: The level number of the Kaluza-Klein tower ($n = 1, 2, 3, \ldots$; $n = 0$ the flat zero mode). Status: DERIVED.
- ξ: The scaled hidden coordinate, $\xi = y/L$, running from -1 (tip) to 0 (fold) on each half of a period. Unit: dimensionless. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- ϖ, ϖ_n: The wave number of an oscillating x_0 profile on $-1 \le \xi \le 0$ (of a boson after the rescaling, or of a bulk level of dirac16complex); ϖ_n that of the n-th even level ($\varpi_n = n\pi$ for $\beta = 0$). Unit: dimensionless. Value: 1.89901 for the lowest odd level of dirac16complex. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $\tilde{m}$, $\tilde{m}_n$: The dimensionless x_0 eigenvalue of a boson mode, $\tilde{m} = m\,fL$; $\tilde{m}_n$ the n-th even Kaluza-Klein level of the tower with $\beta = 0$ ($\tilde{m}_1$ the first). Unit: dimensionless. Value: $\tilde{m}_1^2 = 30.1196$, $\tilde{m}_2^2 = 59.7284$, $\tilde{m}_3^2 = 109.0764$ ($HL = 3$), i.e. $\tilde{m}_1 = 5.48813$, $\tilde{m}_2 = 7.72842$, $\tilde{m}_3 = 10.44397$. Status: DERIVED.
- m_n^KK, m_KK: The mass of the n-th Kaluza-Klein graviton (and gauge) level, $m_n^\mathrm{KK} = \tilde{m}_n/(fL)$; $m_\mathrm{KK} = m_1^\mathrm{KK}$ is the Kaluza-Klein scale. Unit: GeV. Value: $m_\mathrm{KK} = 121.32$ TeV. Status: DERIVED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- N_n, N_0: The normalisation integral of the profile F_n of the n-th Kaluza-Klein level, $N_n = \int_{-1}^{0} \mathrm{e}^{3HL\xi} F_n^2 \,\mathrm{d}\xi$; N_0 that of the flat zero mode. Unit: dimensionless. Value: $N_0 = (1 - \mathrm{e}^{-3HL})/(3HL) = 0.1111$. Status: DERIVED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.

With the depth of the cell $HL = 3$ (ASSUMED) the eigenvalues are (DERIVED):

mode ($HL = 3$)	$\tilde{m}^2 = (m\,fL)^2$	status
photon, gluons, graviton ($\beta = 0$, flat profile)	0 exactly	DERIVED
first even Kaluza-Klein level	$9H^2L^2/4 + \pi^2 = 30.1196$	DERIVED (finite elements 1.9×10^{-10})
first odd level	27.0890 ($\varpi \cot \varpi = -9/2$)	DERIVED (finite elements 4.5×10^{-10})
gap	$9H^2L^2/4 = 20.25$	DERIVED
W^+, W^- ($\beta_\mathrm{W} = 0.011912$)	1.32149×10^{-5}	DERIVED (fit; finite elements 3×10^{-14})
Z ($\beta_\mathrm{Z} = \beta_\mathrm{W}/\cos^2\theta_\mathrm{W} = 0.015343$)	1.70150×10^{-5}	DERIVED
Dirichlet bound ($\beta \to \infty$)	0.010016: $m_\mathrm{W} \le 0.01824\, m_\mathrm{KK}$	DERIVED

Here θ_W is the angle of the couplings, $\cos^2\theta_\mathrm{W} = \hat{g}_2^2/(\hat{g}_2^2 + \hat{g}_1^2)$. The bending of the tip-localised profiles moves the mass ratio $m_\mathrm{W}^2/m_\mathrm{Z}^2 = \tilde{m}_\mathrm{W}^2/\tilde{m}_\mathrm{Z}^2$ away from $\cos^2\theta_\mathrm{W}$, so the electroweak parameter exceeds 1: $\rho_\mathrm{EW} - 1 = 3.2 \times 10^{-8}$ at $\beta_\mathrm{W} = 10^{-6}$ (custodial symmetry at leading order), 6.4×10^{-5} at $\beta_\mathrm{W} = 0.002$, 3.2×10^{-3} at 0.1. This tree-level custodial violation is fitted to the measured $\rho_{\mathrm{EW},0} - 1 = 0.00038$ (the violation beyond the Standard-Model loops), and the angle of the couplings is fixed by the measured masses:

$$\rho_{\mathrm{EW}} = \frac{\tilde{m}_{\mathrm{W}}^2}{\tilde{m}_{\mathrm{Z}}^2 \cos^2\theta_{\mathrm{W}}} = \rho_{\mathrm{EW},0}, \qquad \cos^2\theta_{\mathrm{W}} = \frac{m_{\mathrm{W}}^2/m_{\mathrm{Z}}^2}{\rho_{\mathrm{EW},0}}, \qquad m_{\mathrm{KK}} = m_{\mathrm{W}}\,\frac{\tilde{m}_1}{\tilde{m}_{\mathrm{W}}}, \qquad m_{\mathrm{Z}} = m_{\mathrm{W}}\,\frac{\tilde{m}_{\mathrm{Z}}}{\tilde{m}_{\mathrm{W}}} \tag{FE-55}$$

Symbols in (FE-55).

- ρ_{EW}: The electroweak parameter, $\rho_{\mathrm{EW}} = \tilde{m}_{\mathrm{W}}^2/(\tilde{m}_{\mathrm{Z}}^2\cos^2\theta_{\mathrm{W}})$. Unit: dimensionless. Status: DERIVED.
- $\tilde{m}_{\mathrm{W}}$: The dimensionless x_0 eigenvalue of the lowest $W^{\pm}$ mode ($\beta = \beta_{\mathrm{W}}$). Unit: dimensionless. Value: $\tilde{m}_{\mathrm{W}}^2 = 1.32149 \times 10^{-5}$. Status: DERIVED.
- $\tilde{m}_{\mathrm{Z}}$: The dimensionless x_0 eigenvalue of the lowest Z mode ($\beta = \beta_{\mathrm{Z}}$). Unit: dimensionless. Value: $\tilde{m}_{\mathrm{Z}}^2 = 1.70150 \times 10^{-5}$. Status: DERIVED.
- θ_{W}: The weak mixing angle of the couplings, $\cos\theta_{\mathrm{W}} = \hat{g}_2/\hat{g}_Z$, $\sin\theta_{\mathrm{W}} = \hat{g}_1/\hat{g}_Z$; fixed by the measured W and Z masses with $\rho_{\mathrm{EW},0}$. Unit: dimensionless. Value: $\cos^2\theta_{\mathrm{W}} = 0.776368$. Status: DERIVED.
- $\rho_{\mathrm{EW},0}$: The measured value of the electroweak parameter (PDG fit). Unit: dimensionless. Value: 1.00038. Status: STANDARD.
- m_{W}: The mass of W^+ and W^-. Unit: GeV. Value: 80.3625 GeV (measured, input). Status: STANDARD.
- m_{Z}: The mass of the Z. Unit: GeV. Value: 91.1879 GeV (reproduced). Status: STANDARD.
- m_n^{KK}, m_{KK}: The mass of the n-th Kaluza-Klein graviton (and gauge) level, $m_n^{\mathrm{KK}} = \tilde{m}_n/(fL)$; $m_{\mathrm{KK}} = m_1^{\mathrm{KK}}$ is the Kaluza-Klein scale. Unit: GeV. Value: $m_{\mathrm{KK}} = 121.32$ TeV. Status: DERIVED.
- $\tilde{m}$, $\tilde{m}_n$: The dimensionless x_0 eigenvalue of a boson mode, $\tilde{m} = m\,fL$; $\tilde{m}_n$ the n-th even Kaluza-Klein level of the tower with $\beta = 0$ ($\tilde{m}_1$ the first). Unit: dimensionless. Value: $\tilde{m}_1^2 = 30.1196$, $\tilde{m}_2^2 = 59.7284$, $\tilde{m}_3^2 = 109.0764$ ($HL = 3$), i.e. $\tilde{m}_1 = 5.48813$, $\tilde{m}_2 = 7.72842$, $\tilde{m}_3 = 10.44397$. Status: DERIVED.

i.e. $\cos^2\theta_{\mathrm{W}} = 0.776368$ (the on-shell ratio $m_{\mathrm{W}}^2/m_{\mathrm{Z}}^2 = 0.776663$); both masses are then reproduced. With $m_{\mathrm{W}} = 80.3625$ GeV (STANDARD, PDG 2026) the scales of UM-2 are (DERIVED):

quantity	value
first Kaluza-Klein level $m_{\mathrm{KK}} = m_{\mathrm{W}}\,\tilde{m}_1/\tilde{m}_{\mathrm{W}}$	121.32 TeV
the author's constant $H = 3f_*\,m_{\mathrm{KK}}/\tilde{m}_1$	14.296 TeV
proper half cell fL; period of x_0, $\pi/(6H)$	8.926×10^{-21} m; 7.227×10^{-21} m
gap $3H/(2f_*)$	99.48 TeV
radion mass $\omega_{\mathrm{r}} = 3.387\,H$	48.42 TeV
$m_{\mathrm{Z}} = m_{\mathrm{W}}\,\tilde{m}_{\mathrm{Z}}/\tilde{m}_{\mathrm{W}}$	91.1879 GeV (the measured value, reproduced)

5.3.2 The fermions: the exact reduction and the zero modes

With the factors of the ansatz below, the eight-dimensional Dirac equation, the canonical spin connection included, is EXACTLY a two-component equation in each of the eight blocks of the spinor (DERIVED, all blocks, general $a_4(x_4)$, $b_4(x_4)$, $M(y)$, $v(y)$):

$$\begin{aligned}
&\Psi = \mathrm{e}^{\mathrm{i}kx_1}\,\mathrm{e}^{-\frac{3}{2}Hy}\,\mathrm{e}^{a_4/2}\,b_4^{-1}\,\mathcal{V}\chi(y,x_4)\,:\\
&\gamma^M D_M\Psi - \big(M(y) - \mathrm{i}v(y)\gamma^{x_4}\big)\Psi = \mathrm{e}^{\mathrm{i}kx_1}\,\mathrm{e}^{-\frac{3}{2}Hy}\,\mathrm{e}^{a_4/2}\,b_4^{-1}\,\gamma^{x_4}\mathcal{V}\big[\partial_{x_4}\chi + \mathrm{i}h\chi\big],\\
&h = \varsigma\Big[-\frac{\mathrm{i}}{f}\sigma_1\partial_y + M(y)\,\sigma_2 + \mathrm{e}^{-a_4}k\,\sigma_3\Big] + v(y), \qquad \mathrm{i}\,\partial_{x_4}\chi = h\chi;\\
&\text{in each block: } \gamma^{x_0} = \sigma_3, \qquad \mathcal{B} = \varsigma\varsigma_2, \qquad C = \varsigma_2\sigma_2, \qquad \Gamma_4 = -\varsigma\varsigma_2\sigma_3
\end{aligned} \tag{FE-56}$$

Symbols in (FE-56).

- Ψ, $\bar{\Psi}$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar{\Psi} = \Psi^{\dagger}C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.

- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the hidden and extra-time components, $b_4 > 0$. Unit: dimensionless. Status: AUTHOR.
- $\mathcal{V}$: The constant 16×2 matrix of the reduction ansatz that places the two-component spinor χ of one block into the 16 components of the eight-dimensional spinor. Unit: dimensionless. Status: DERIVED.
- χ, χ_o, χ_1, χ_2: The two-component spinor of one block of the reduction (a function of y and x_4), $\chi = (\chi_1, \chi_2)^T = (a, \mathrm{i}b)^T$; χ_o the Kohn-Sham orbital o, normalised as $\int_{-L}^{0} \chi_o^\dagger \chi_o \,\mathrm{d}y = 1$. Unit: $\mathrm{GeV}^{1/2}$ (normalised orbital). Status: DERIVED.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4, 4)$; $\gamma^M = \mathbf{e}_a{}^M \gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- $M(y)$: The mass function of the block Hamiltonian (the scalar potential): $M = \mu_{\psi i}$ for a Standard-Model fermion, $M = M_\mathrm{eff}$ for dirac16complex. Unit: GeV. Status: DERIVED.
- $v(y)$: The vector potential of the block Hamiltonian (the mean field along γ^{x_4}): for dirac16complex the contact term and the mean field of I11; zero for a Standard-Model fermion. Unit: GeV. Status: DERIVED.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- h: The block Hamiltonian: the 2×2 differential operator of one block that generates the time evolution of χ, $\mathrm{i}\,\partial_{x_4}\chi = h\chi$; its eigenvalues are the energies ε. Unit: GeV. Status: DERIVED.
- ς: The first of the three signs $(\varsigma, \varsigma_2, \varsigma_3)$, each ± 1, that label the eight blocks of the reduction. Unit: dimensionless. Status: DERIVED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- σ_1, σ_2, σ_3: The Pauli matrices acting on the two components of a block spinor χ ($\sigma_3 = \mathrm{diag}(1, -1)$). Unit: dimensionless. Status: STANDARD.
- y: The label y in ∂_y: the derivative with respect to the hidden coordinate y. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- $\mathcal{B}$: The author's operator $\mathcal{B} = -\mathrm{i}C\gamma^{x_4}$; its eigenvalues ∓ 1 define the halves Ψ_A and Ψ_B, and it commutes with every interaction of the good sector. Unit: dimensionless. Status: AUTHOR.
- ς_2: The second sign that labels the blocks (± 1). Unit: dimensionless. Status: DERIVED.
- C: The matrix of the Dirac adjoint, $C = \gamma^{x_0}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}$ ($\bar\Psi = \Psi^\dagger C$); in each block of the reduction $C = \varsigma_2\sigma_2$. Unit: dimensionless. Status: AUTHOR.
- Γ_4: The four-dimensional chirality matrix, $\Gamma_4 = \mathrm{i}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}\gamma^{x_4}$; a doublet zero mode has $\Gamma_4 = +1$ (left-handed), a singlet zero mode $\Gamma_4 = -1$ (right-handed). Unit: dimensionless. Status: DERIVED.

For a Standard-Model multiplet $M = \mu_{\psi i}$ and $v = 0$; the zero modes are massless four-dimensional particles:

$$\chi = \big(\mathrm{e}^{c\xi},\ 0\big)^T \ \text{(doublet, } \Gamma_4 = +1\text{)}, \qquad \chi = \big(0,\ \mathrm{e}^{-c\xi}\big)^T \ \text{(singlet, } \Gamma_4 = -1\text{)}, \qquad c = f\mu L,$$
$$h\big(\mathrm{e}^{c\xi},\ 0\big)^T = \varsigma\,\mathrm{e}^{-a_4} k\big(\mathrm{e}^{c\xi},\ 0\big)^T \tag{FE-57}$$

Symbols in (FE-57).

- $\chi,\ \chi_o,\ \chi_1,\ \chi_2$: The two-component spinor of one block of the reduction (a function of y and x_4), $\chi = (\chi_1, \chi_2)^T = (a, \mathrm{i}b)^T$; χ_o the Kohn-Sham orbital o, normalised as $\int_{-L}^{0} \chi_o^\dagger \chi_o \,\mathrm{d}y = 1$. Unit: $\mathrm{GeV}^{1/2}$ (normalised orbital). Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- $c,\ c_L,\ c_R$: The localisation of a zero mode, $c = f\mu L$ (bulk mass times the proper depth); c_L of a doublet (localised at the fold for $c_L > 0$), c_R of a singlet (localised at the fold for $c_R < 0$). Unit: dimensionless. Value: doublets $c_L = 3.38230$ (quarks) and 5.95288 (leptons); singlets c_R: up -15.4300, down -14.6258, strange -11.5158, charm -8.7683, bottom -7.5005, top -3.3823, electron -14.5539, muon -8.9809, tau -5.9529, neutrinos -35.0452, -32.8481, -31.0702 (fixed by the measured masses). Status: DERIVED.
- ξ: The scaled hidden coordinate, $\xi = y/L$, running from -1 (tip) to 0 (fold) on each half of a period. Unit: dimensionless. Status: DERIVED.
- ^{T}: The transpose: a row written with ^{T} is a column. Status: STANDARD.
- Γ_4: The four-dimensional chirality matrix, $\Gamma_4 = \mathrm{i}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}\gamma^{x_4}$; a doublet zero mode has $\Gamma_4 = +1$ (left-handed), a singlet zero mode $\Gamma_4 = -1$ (right-handed). Unit: dimensionless. Status: DERIVED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- $\mu_{\psi i}$: The bulk mass of a Standard-Model fermion: $\mu_{\psi i}$ of the multiplet ψ of the generation i; with the hidden direction it fixes the localisation $c = f\mu L$ of the zero mode. Unit: GeV. Value: $\mu_{\psi i} = c/(fL)$ with the localisation c of the zero mode (doublets: $\mu_Q = 5.2304\,H$, $\mu_\ell = 9.2055\,H$). Status: ASSUMED.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- h: The block Hamiltonian: the 2×2 differential operator of one block that generates the time evolution of χ, $\mathrm{i}\,\partial_{x_4}\chi = h\chi$; its eigenvalues are the energies ε. Unit: GeV. Status: DERIVED.
- ς: The first of the three signs $(\varsigma, \varsigma_2, \varsigma_3)$, each ± 1, that label the eight blocks of the reduction. Unit: dimensionless. Status: DERIVED.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.

5.3.3 The masses of one generation

At $k = 0$ the doublet and singlet solutions that satisfy the fold conditions reach the tip with closed-form values, and the tip mixing conditions give the exact mass condition (DERIVED):

$$\frac{\mathrm{d}a}{\mathrm{d}\xi} = c\,a - \tilde{\varepsilon}\,b, \qquad \frac{\mathrm{d}b}{\mathrm{d}\xi} = \tilde{\varepsilon}\,a - c\,b, \qquad \chi = (a,\ \mathrm{i}b), \qquad \tilde{\varepsilon} = \varepsilon f L, \qquad \upsilon_L = \sqrt{c_L^2 - \tilde{\varepsilon}^2}, \qquad \upsilon_R = \sqrt{c_R^2 - \tilde{\varepsilon}^2},$$
$$(a_L,\ b_L)\big|_{\xi=-1} = \Big(\cosh\upsilon_L - c_L\frac{\sinh\upsilon_L}{\upsilon_L}, -\tilde{\varepsilon}\,\frac{\sinh\upsilon_L}{\upsilon_L}\Big), \qquad (a_R,\ b_R)\big|_{\xi=-1} = \Big(\tilde{\varepsilon}\,\frac{\sinh\upsilon_R}{\upsilon_R}, \cosh\upsilon_R + c_R\frac{\sinh\upsilon_R}{\upsilon_R}\Big) \tag{FE-58}$$

Symbols in (FE-58).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.

- $a,\ b$: The upper and lower components of a block spinor, $\chi = (a, \mathrm{i}b)$, real at $k = 0$: a_L, b_L of a doublet and a_R, b_R of a singlet solution (a_{Ri}, b_{Ri} of the generation i, $b_{L\alpha}$, $a_{L\alpha}$ of the doublet in the flavour basis), a_o, b_o of the orbital o. Unit: $\mathrm{GeV}^{1/2}$ (normalised orbital). Status: DERIVED.
- ξ: The scaled hidden coordinate, $\xi = y/L$, running from -1 (tip) to 0 (fold) on each half of a period. Unit: dimensionless. Status: DERIVED.
- $c,\ c_L,\ c_R$: The localisation of a zero mode, $c = f\mu L$ (bulk mass times the proper depth); c_L of a doublet (localised at the fold for $c_L > 0$), c_R of a singlet (localised at the fold for $c_R < 0$). Unit: dimensionless. Value: doublets $c_L = 3.38230$ (quarks) and 5.95288 (leptons); singlets c_R: up -15.4300, down -14.6258, strange -11.5158, charm -8.7683, bottom -7.5005, top -3.3823, electron -14.5539, muon -8.9809, tau -5.9529, neutrinos -35.0452, -32.8481, -31.0702 (fixed by the measured masses). Status: DERIVED.
- $\tilde{\varepsilon}$: The dimensionless energy (mass) of a fermion mode, $\tilde{\varepsilon} = \varepsilon f L$. Unit: dimensionless. Status: DERIVED.
- $\chi,\ \chi_o,\ \chi_1,\ \chi_2$: The two-component spinor of one block of the reduction (a function of y and x_4), $\chi = (\chi_1, \chi_2)^T = (a, \mathrm{i}b)^T$; χ_o the Kohn-Sham orbital o, normalised as $\int_{-L}^{0} \chi_o^\dagger \chi_o \,\mathrm{d}y = 1$. Unit: $\mathrm{GeV}^{1/2}$ (normalised orbital). Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\varepsilon,\ \varepsilon_o$: The energy of a fermion mode (an eigenvalue of the block Hamiltonian; at $k = 0$ the four-dimensional mass); ε_o the Kohn-Sham level of the orbital o. Unit: GeV. Status: DERIVED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- $\upsilon_L,\ \upsilon_R$: The auxiliary roots $\upsilon = (c^2 - \tilde{\varepsilon}^2)^{1/2}$ of the doublet (υ_L) and singlet (υ_R) solutions at $k = 0$. Unit: dimensionless. Status: DERIVED.
- L: The chirality label L: the left-handed (doublet) zero mode, or the left-handed projection $P_L = (1 + \Gamma_4)/2$. Status: STANDARD.
- R: The chirality label R: the right-handed (singlet) zero mode or field, or the projection $P_R = (1 - \Gamma_4)/2$. Status: STANDARD.

$$\mathcal{D}(\tilde{\varepsilon}) = \tilde{\varepsilon}^2 \frac{\sinh \upsilon_L}{\upsilon_L} \frac{\sinh \upsilon_R}{\upsilon_R} - \mathcal{K}^2 \Big(\cosh \upsilon_L - c_L \frac{\sinh \upsilon_L}{\upsilon_L} \Big) \Big(\cosh \upsilon_R + c_R \frac{\sinh \upsilon_R}{\upsilon_R} \Big) = 0,$$
$$\tilde{\varepsilon} \simeq \mathcal{K}\, \mathcal{Z}(c_L)\, \mathcal{Z}(-c_R), \qquad \mathcal{Z}(c)^2 = \frac{c\, \mathrm{e}^{-c}}{\sinh c} \tag{FE-59}$$

Symbols in (FE-59).

- $\mathcal{D}(\tilde{\varepsilon})$: The mass function of one generation: its zeros are the dimensionless masses $\tilde{\varepsilon}$. Unit: dimensionless. Status: DERIVED.
- $\tilde{\varepsilon}$: The dimensionless energy (mass) of a fermion mode, $\tilde{\varepsilon} = \varepsilon f L$. Unit: dimensionless. Status: DERIVED.
- $\upsilon_L,\ \upsilon_R$: The auxiliary roots $\upsilon = (c^2 - \tilde{\varepsilon}^2)^{1/2}$ of the doublet (υ_L) and singlet (υ_R) solutions at $k = 0$. Unit: dimensionless. Status: DERIVED.
- L: The chirality label L: the left-handed (doublet) zero mode, or the left-handed projection $P_L = (1 + \Gamma_4)/2$. Status: STANDARD.
- R: The chirality label R: the right-handed (singlet) zero mode or field, or the projection $P_R = (1 - \Gamma_4)/2$. Status: STANDARD.
- $\mathcal{K}$: The tip mixing matrix of a sector in the boundary conditions of the fermion zero modes (a number $\mathcal{K}$ for one generation, a 3×3 matrix for three): $\mathcal{K}_\nu$, $\mathcal{K}_e$, $\mathcal{K}_u$, $\mathcal{K}_d$ of the neutrino, charged-lepton, up-quark and down-quark sectors; UM-2 takes $\mathcal{K}_e = \mathcal{K}_u = \mathbf{1}$, $\mathcal{K}_\nu = U$, $\mathcal{K}_d = V_{\mathrm{CKM}}$. Unit: dimensionless. Status: ASSUMED.
- $c,\ c_L,\ c_R$: The localisation of a zero mode, $c = f\mu L$ (bulk mass times the proper depth); c_L of a doublet (localised at the fold for $c_L > 0$), c_R of a singlet (localised at the fold for $c_R < 0$). Unit: dimensionless. Value: doublets $c_L = 3.38230$ (quarks) and 5.95288 (leptons); singlets c_R: up -15.4300, down -14.6258, strange -11.5158, charm -8.7683, bottom -7.5005, top -3.3823, electron -14.5539, muon -8.9809, tau -5.9529, neutrinos -35.0452, -32.8481, -31.0702 (fixed by the measured masses). Status: DERIVED.
- $\mathcal{Z}(c)$: The tip amplitude of a normalised zero mode with the localisation c, $\mathcal{Z}(c)^2 = c\,\mathrm{e}^{-c}/\sinh c$. Unit: dimensionless. Status: DERIVED.

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

5.3.4 Three generations: the masses and the mixing matrices

For three generations the tip conditions hold for the vectors of boundary values over the generations; for one doublet bulk mass per sector and a unitary tip matrix the determinant factorises into the one-generation conditions with $\mathcal{K} = 1$ (DERIVED, mpmath with 50 digits):

$$\Big[\mathrm{diag}\big(b_{L\alpha}\big) + \mathcal{K}\,\mathrm{diag}\Big(\frac{b_{Ri}}{a_{Ri}}\Big)\mathcal{K}^\dagger\,\mathrm{diag}\big(a_{L\alpha}\big)\Big]\boldsymbol{\alpha} = 0; \qquad \det\big[\,\cdots\big] = \prod_{i=1}^{3}\frac{b_L a_{Ri} + a_L b_{Ri}}{a_{Ri}},$$
$$b_L a_R + a_L b_R = -\mathcal{D}(\tilde{\varepsilon})\big|_{\mathcal{K}=1}, \qquad \boldsymbol{\alpha}^{(i)} = \mathcal{K}\,\mathbf{u}_i, \qquad \boldsymbol{\beta}^{(i)} \propto \mathbf{u}_i, \qquad U = \mathcal{K}_e^\dagger\mathcal{K}_\nu, \qquad V_{\mathrm{CKM}} = \mathcal{K}_u^\dagger\mathcal{K}_d \tag{FE-60}$$

Symbols in (FE-60).

- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ when it is a tensor in eight dimensions). Status: STANDARD.
- a, b: The upper and lower components of a block spinor, $\chi = (a, \mathrm{i}b)$, real at $k = 0$: a_L, b_L of a doublet and a_R, b_R of a singlet solution (a_{Ri}, b_{Ri} of the generation i, $b_{L\alpha}$, $a_{L\alpha}$ of the doublet in the flavour basis), a_o, b_o of the orbital o. Unit: $\mathrm{GeV}^{1/2}$ (normalised orbital). Status: DERIVED.
- L: The chirality label L: the left-handed (doublet) zero mode, or the left-handed projection $P_L = (1 + \Gamma_4)/2$. Status: STANDARD.
- α: A flavour index running over e, μ, τ (the flavour states of the leptons and neutrinos). Status: STANDARD.
- $\mathcal{K}$: The tip mixing matrix of a sector in the boundary conditions of the fermion zero modes (a number $\mathcal{K}$ for one generation, a 3×3 matrix for three): $\mathcal{K}_\nu$, $\mathcal{K}_e$, $\mathcal{K}_u$, $\mathcal{K}_d$ of the neutrino, charged-lepton, up-quark and down-quark sectors; UM-2 takes $\mathcal{K}_e = \mathcal{K}_u = \mathbf{1}$, $\mathcal{K}_\nu = U$, $\mathcal{K}_d = V_{\mathrm{CKM}}$. Unit: dimensionless. Status: ASSUMED.
- R: The chirality label R: the right-handed (singlet) zero mode or field, or the projection $P_R = (1 - \Gamma_4)/2$. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $\boldsymbol{\alpha}$, $\boldsymbol{\alpha}^{(i)}$: The vector of the three doublet amplitudes at the tip in the mass condition of three generations; $\boldsymbol{\alpha}^{(i)}$ its null vector for the mass state i (the left-handed flavour content). Unit: dimensionless. Status: DERIVED.
- $\mathcal{D}(\tilde{\varepsilon})$: The mass function of one generation: its zeros are the dimensionless masses $\tilde{\varepsilon}$. Unit: dimensionless. Status: DERIVED.
- $\tilde{\varepsilon}$: The dimensionless energy (mass) of a fermion mode, $\tilde{\varepsilon} = \varepsilon f L$. Unit: dimensionless. Status: DERIVED.
- $\mathbf{u}_i$: The unit vector of the generation i (the i-th column of the 3×3 identity). Unit: dimensionless. Status: STANDARD.
- $\boldsymbol{\beta}^{(i)}$: The vector of the three singlet amplitudes of the mass state i. Unit: dimensionless. Status: DERIVED.
- U: The PMNS matrix, the unitary 3×3 mixing matrix of the leptons, $\nu_\alpha = \sum_i U_{\alpha i}\nu_i$ (rows $\alpha = e, \mu, \tau$, columns the mass states i); in UM-2 $U = \mathcal{K}_e^\dagger\mathcal{K}_\nu$. Unit: dimensionless. Value: $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$, $\delta_{\mathrm{CP}} = 212^\circ$ (NuFIT 6.0). Status: STANDARD.
- V_{CKM}: The CKM matrix, the unitary 3×3 mixing matrix of the quarks (the same parametrisation as U); in UM-2 $V_{\mathrm{CKM}} = \mathcal{K}_u^\dagger\mathcal{K}_d$. Unit: dimensionless. Value: $s_{12} = 0.22501$, $s_{13} = 0.003732$, $s_{23} = 0.04183$, $\delta = 1.147$ (PDG). Status: STANDARD.

The masses are therefore the one-generation roots with $\mathcal{K} = 1$, the left-handed rotation of each sector is its tip matrix, and $U = \mathcal{K}_e^\dagger\mathcal{K}_\nu$, $V_{\mathrm{CKM}} = \mathcal{K}_u^\dagger\mathcal{K}_d$; UM-2 takes $\mathcal{K}_e = \mathcal{K}_u = 1$, $\mathcal{K}_\nu = U$, $\mathcal{K}_d = V_{\mathrm{CKM}}$ (ASSUMED). With one doublet bulk mass per sector, fixed by the heaviest member with $c_R = -c_L$ (τ, t), the masses require (DERIVED, exact roots):

fermion	mass	doublet c_L	singlet c_R	$\mu_{\psi i}/H$ (singlet)	singlet width $fL/(2\lvert c_R\rvert)$	$\tilde{\varepsilon} = m\,fL$	first order
ν_1	1.000 meV	5.95288	-35.04518	-54.194	1.27×10^{-22} m	4.5235×10^{-17}	5.7×10^{-7}
ν_2	8.712 meV	5.95288	-32.84810	-50.796	1.36×10^{-22} m	3.9409×10^{-16}	6.1×10^{-7}
ν_3	50.140 meV	5.95288	-31.07017	-48.047	1.44×10^{-22} m	2.2681×10^{-15}	6.5×10^{-7}
e	0.51100 MeV	5.95288	-14.55391	-22.506	3.07×10^{-22} m	2.3115×10^{-8}	1.4×10^{-6}
μ	105.658 MeV	5.95288	-8.98093	-13.888	4.97×10^{-22} m	4.7795×10^{-6}	2.2×10^{-6}
τ	1776.93 MeV	5.95288	-5.95288	-9.205	7.50×10^{-22} m	8.0380×10^{-5}	6.8×10^{-6}
u	2.16 MeV	3.38230	-15.43001	-23.861	2.89×10^{-22} m	9.7708×10^{-8}	1.3×10^{-4}
c	1.2730 GeV	3.38230	-8.76830	-13.559	5.09×10^{-22} m	5.7585×10^{-5}	2.2×10^{-4}
t	172.56 GeV	3.38230	-3.38230	-5.230	1.32×10^{-21} m	7.8058×10^{-3}	1.1×10^{-3}
d	4.70 MeV	3.38230	-14.62578	-22.617	3.05×10^{-22} m	2.1261×10^{-7}	1.3×10^{-4}
s	93.5 MeV	3.38230	-11.51582	-17.808	3.88×10^{-22} m	4.2295×10^{-6}	1.7×10^{-4}
b	4.183 GeV	3.38230	-7.50053	-11.599	5.95×10^{-22} m	1.8922×10^{-4}	2.6×10^{-4}

(the last column is the relative difference of the first-order root $\mathcal{K}\,\mathcal{Z}(c_L)\mathcal{Z}(-c_R)$ from the exact root; the doublets have $\mu_\ell = 9.2055\,H$ and $\mu_Q = 5.2304\,H$; the masses are the PDG values, the light-quark masses in $\overline{\text{MS}}$ at 2 GeV, $m_{\text{c}}(m_{\text{c}})$, $m_{\text{b}}(m_{\text{b}})$ and the directly measured m_{t}, used as tree-level targets; m_1 is ASSUMED and m_2, m_3 follow from the NuFIT 6.0 splittings of step 5; the masses are STANDARD inputs, PDG 2025.) The mixing matrices have the standard form

$$U = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}\mathrm{e}^{-\mathrm{i}\delta_{\text{CP}}} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}\mathrm{e}^{\mathrm{i}\delta_{\text{CP}}} & c_{12}c_{23} - s_{12}s_{23}s_{13}\mathrm{e}^{\mathrm{i}\delta_{\text{CP}}} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}\mathrm{e}^{\mathrm{i}\delta_{\text{CP}}} & -c_{12}s_{23} - s_{12}c_{23}s_{13}\mathrm{e}^{\mathrm{i}\delta_{\text{CP}}} & c_{23}c_{13} \end{pmatrix}, \qquad J_{\text{CP}} = c_{12}s_{12}c_{23}s_{23}c_{13}^2 s_{13}\sin\delta_{\text{CP}} \tag{FE-61}$$

Symbols in (FE-61).

- U: The PMNS matrix, the unitary 3×3 mixing matrix of the leptons, $\nu_\alpha = \sum_i U_{\alpha i}\nu_i$ (rows $\alpha = e, \mu, \tau$, columns the mass states i); in UM-2 $U = \mathcal{K}_e^\dagger \mathcal{K}_\nu$. Unit: dimensionless. Value: $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$, $\delta_{\text{CP}} = 212^\circ$ (NuFIT 6.0). Status: STANDARD.
- $c_{12},\ c_{13},\ c_{23}$: The cosines of the mixing angles, $c_{ij} = \cos\theta_{ij}$. Unit: dimensionless. Status: STANDARD.
- $s_{12},\ s_{13},\ s_{23}$: The sines of the mixing angles, $s_{ij} = \sin\theta_{ij}$. Unit: dimensionless. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- δ_{CP}: The CP-violating phase of a mixing matrix. Unit: dimensionless. Value: 212° (PMNS), 1.147 rad (CKM). Status: STANDARD.
- J_{CP}: The Jarlskog invariant of a mixing matrix (the measure of CP violation). Unit: dimensionless. Value: -0.01777 (PMNS), 3.117×10^{-5} (CKM). Status: STANDARD.

with the NuFIT 6.0 values $\sin^2\theta_{12}=0.308$, $\sin^2\theta_{23}=0.470$, $\sin^2\theta_{13}=0.02215$, $\delta_{\mathrm{CP}}=212^\circ$ ($J_{\mathrm{CP}}=-0.01777$; its largest value over the phase, 0.0335 for these best-fit angles, against NuFIT's marginalised 0.0333 ± 0.0007) for the PMNS matrix U, and the PDG values $s_{12}=0.22501$, $s_{13}=0.003732$, $s_{23}=0.04183$, $\delta_{\mathrm{CP}}=1.147$ ($J_{\mathrm{CP}}=3.117\times10^{-5}$) for the CKM matrix V_{CKM}, both unitary to 1.1×10^{-16} (DERIVED). The first even Kaluza-Klein gauge mode couples to the zero modes with the overlaps 0.14434 (every left-handed lepton) and -0.30066 (every left-handed quark) relative to the zero mode, and in the mass basis its largest off-diagonal coupling is 1.0×10^{-17}: no tree-level flavour-changing neutral current (with the three charged-lepton $|c_R|$ used as different doublet localisations it would be 0.033). The rare decay $\mu\to e\gamma$, through the neutrino masses alone, has the branching ratio (flavour indices $\alpha=\mu$, $\beta=e$)

$$\mathrm{BR}=\frac{3\alpha_{\mathrm{em},0}}{32\pi}\Big|\sum_{i=1}^{3}U^{*}_{\alpha i}U_{\beta i}\,\frac{m_i^2}{m_{\mathrm{W}}^2}\Big|^2=2.8\times10^{-55}$$

Symbols in this equation.

- BR: The branching ratio of the lepton-flavour-violating decay of a charged lepton of the flavour α into one of the flavour β and a photon, the fraction of all its decays that end in this final state; in step 2 it is the decay $\mu\to e\gamma$ ($\alpha=\mu$, $\beta=e$) through the neutrino masses alone. Unit: dimensionless. Value: 2.8×10^{-55} for $\mu\to e\gamma$. Status: DERIVED.
- $\alpha_{\mathrm{em},0}$: The electromagnetic fine-structure constant at zero momentum transfer, $e^2/(4\pi)$ measured at low energy (distinct from its value at the Z mass). Unit: dimensionless. Value: 1/137.036. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- i: The generation index ($i=1,2,3$), which also labels the neutrino mass states ν_1,ν_2,ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- U: The PMNS matrix, the unitary 3×3 mixing matrix of the leptons, $\nu_\alpha=\sum_i U_{\alpha i}\nu_i$ (rows $\alpha=e,\mu,\tau$, columns the mass states i); in UM-2 $U=\mathcal{K}_e^{\dagger}\mathcal{K}_\nu$. Unit: dimensionless. Value: $\sin^2\theta_{12}=0.308$, $\sin^2\theta_{13}=0.02215$, $\sin^2\theta_{23}=0.470$, $\delta_{\mathrm{CP}}=212^\circ$ (NuFIT 6.0). Status: STANDARD.
- α: A flavour index running over e,μ,τ (the flavour states of the leptons and neutrinos). Status: STANDARD.
- *: Complex conjugation of a number or of each entry of a matrix. Status: STANDARD.
- β: A second flavour index running over e,μ,τ. Status: STANDARD.
- m, m_s, m_i: A four-dimensional mass: m the mass of a mode (the eigenvalue of an x_0 problem) or of a fermion; m_s the mass of the species s; m_i the mass of the neutrino mass state i. Unit: GeV. Status: DERIVED.
- m_{W}: The mass of W^+ and W^-. Unit: GeV. Value: 80.3625 GeV (measured, input). Status: STANDARD.

(DERIVED).

5.3.5 The spectrum of dirac16complex

With $M=m_\Psi$ and $v=0$ the exact spectrum at $k=0$ is (DERIVED):

$$\begin{aligned}&\text{even: } \chi=\big(\mathrm{e}^{c_\Psi\xi},\,0\big)^T,\ \varepsilon=0\ \text{(all eight blocks)},\qquad \varepsilon=\pm\sqrt{m_\Psi^2+\Big(\frac{\varpi_n}{fL}\Big)^2},\ \varpi_n=\pi,\ 2\pi,\ \ldots;\\ &\text{odd: } \varepsilon=\pm\sqrt{m_\Psi^2+\Big(\frac{\varpi}{fL}\Big)^2},\quad \tan\varpi=-\frac{\varpi}{c_\Psi};\qquad c_\Psi=f\,m_\Psi L;\qquad \text{fold band: } \varepsilon=\mathrm{e}^{-a_4}|k|\end{aligned}\tag{FE-62}$$

Symbols in (FE-62).

- χ, χ_o, χ_1, χ_2: The two-component spinor of one block of the reduction (a function of y and x_4), $\chi=(\chi_1,\chi_2)^T=(a,\mathrm{i}b)^T$; χ_o the Kohn-Sham orbital o, normalised as $\int_{-L}^{0}\chi_o^{\dagger}\chi_o\,\mathrm{d}y=1$. Unit: $\mathrm{GeV}^{1/2}$ (normalised orbital). Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- c_Ψ: The localisation parameter of dirac16complex, $c_\Psi=f\,m_\Psi L$. Unit: dimensionless. Value: 0.646667 at the stabilised block. Status: DERIVED.

- ξ: The scaled hidden coordinate, $\xi = y/L$, running from -1 (tip) to 0 (fold) on each half of a period. Unit: dimensionless. Status: DERIVED.
- ^{T}: The transpose: a row written with ^{T} is a column. Status: STANDARD.
- ε, ε_o: The energy of a fermion mode (an eigenvalue of the block Hamiltonian; at $k = 0$ the four-dimensional mass); ε_o the Kohn-Sham level of the orbital o. Unit: GeV. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex, $m_\Psi = 3H\mathcal{M} = H$. Unit: GeV. Value: 14.296 TeV. Status: RECORD.
- ϖ, ϖ_n: The wave number of an oscillating x_0 profile on $-1 \le \xi \le 0$ (of a boson after the rescaling, or of a bulk level of dirac16complex); ϖ_n that of the n-th even level ($\varpi_n = n\pi$ for $\beta = 0$). Unit: dimensionless. Value: 1.89901 for the lowest odd level of dirac16complex. Status: DERIVED.
- n: The level number of the Kaluza-Klein tower ($n = 1, 2, 3, \ldots$; $n = 0$ the flat zero mode). Status: DERIVED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.

At the stabilised block $c_\Psi = f_* m_\Psi L = 0.646667$: the fold zero mode ($\varepsilon = 0$, all eight blocks); the even bulk levels, the lowest $4.959981\, m_\Psi$; the odd levels, the lowest $3.102204\, m_\Psi$ ($\varpi/(fL) = 2.93661\, m_\Psi$, $\varpi = 1.89901$), which is the dark matter of UM-2 (steps 3, 7, 9); and the fold band $\varepsilon = \mathrm{e}^{-a_4}|k|$, massless, $\varepsilon/|k| = 1$ to 6×10^{-14}.

5.4 The check of step 2

In its own equations.

- *The boson eigenvalues.* Every stored eigenvalue, substituted into its exact characteristic equation, has the implied relative error $|F/(\tilde{m}^2\, \partial F/\partial \tilde{m}^2)|$ (for the condition $F(\tilde{m}^2) = 0$) of at most 5.9×10^{-12} (W; Z 1.4×10^{-12}, the Kaluza-Klein levels 2.4×10^{-13} to 1.8×10^{-12}, the Dirichlet bound 2.0×10^{-12}); the flat profile solves the problem with $\beta = 0$ and $\tilde{m} = 0$ exactly; finite elements agree to 5×10^{-10} (DERIVED).
- *The fermion roots.* The implied relative root error $|\mathcal{D}/(\tilde{\varepsilon}\, \mathrm{d}\mathcal{D}/\mathrm{d}\tilde{\varepsilon})|$ of the twelve stored roots is at most 5.4×10^{-11} (ν_1; 10^{-13} for τ), and the three-generation determinant at the stored roots of the four sectors is zero to 1.2×10^{-11} in the same measure; the roots of the 3×3 determinant agree with the one-generation roots, both refined in the same computation, to 3.0×10^{-42}, and the null vectors with the columns of the tip matrix to 1.1×10^{-16}; direct shooting of the coupled Dirac system agrees with the determinant to 6.1×10^{-10} (e), 1.5×10^{-11} (τ), 5.6×10^{-12} (t); for a random complex non-unitary tip matrix the exact roots agree with the singular values of $\mathcal{Z}(c_L)\, \mathcal{K}\, \mathrm{diag}(\mathcal{Z}(-c_{Ri}))$ to 3.5×10^{-5}, 2.8×10^{-6}, 2.0×10^{-6} (DERIVED).
- *The spectrum of dirac16complex.* The stored odd level, its x_0 momentum and the lowest even level agree with the closed forms to 8.5×10^{-13}, and the Kohn-Sham solver of step 3 reproduces them to 10^{-12} (DERIVED).

In concert with the other field equations.

- *The Z mass in the Z widths.* The Z mass of the x_0 eigenvalues, $m_\mathrm{W}\tilde{m}_\mathrm{Z}/\tilde{m}_\mathrm{W} = 91.187900$ GeV, is the measured value used in the Z widths of step 4, to 2.0×10^{-12}; the angle of the couplings of the x_0 problem ($\sin^2\theta_\mathrm{W} = 0.223632$) and the $\overline{\mathrm{MS}}$ value used in the widths (0.23122) are two values of one angle, and the difference changes the total width by 0.82 % at tree level (0.83 % with the QCD factor on the quarks), the size of the electroweak loop corrections that UM-2 does not compute (step 4) (DERIVED).
- *The fermion masses in the thermal history.* The twelve masses of the exact roots are the masses of the ideal gases of step 9, of the quark components of I13 (step 9) and of the top threshold of step 4, to 3.1×10^{-12} (DERIVED).
- *The neutrino masses and the PMNS matrix.* The three neutrino masses of the tip conditions, their sum and

their splittings are those of the flavour evolution (step 5), the CNB (step 5) and the CMB computation (step 6), to 5.7×10^{-12}, and the neutrino tip matrix is the PMNS matrix of the flavour evolution and of the flavour content of the CNB, to 8.2×10^{-13} (DERIVED).

- *The dark-matter mass.* The odd level $M_{\rm DM}=3.102204\,m_\Psi=44347.904991$ GeV ($m_\Psi=H$) is the mass of the Kohn-Sham quanta (step 3) and of the Boltzmann equation (steps 7 and 9), to 8.1×10^{-13}, and its block sensitivity $-\mathrm{d}\ln\varepsilon_{\rm DM}/\mathrm{d}\ln f=0.772034343$ equals the hidden pressure of the Kohn-Sham quanta (step 3) to 6.1×10^{-12} (DERIVED).

5.5 The status of step 2

DERIVED: the exact x_0 reductions, the eigenvalue conditions and their roots, the factorisation of the three-generation condition, the absence of tree-level flavour-changing neutral currents, the spectrum of dirac16complex. ASSUMED: the depth $HL=3$, one doublet bulk mass per sector, the tip matrices $\mathcal{K}_e=\mathcal{K}_u=1$, $\mathcal{K}_\nu=U$, $\mathcal{K}_d=V_{\rm CKM}$, the lightest neutrino mass. STANDARD: the measured masses, mixing angles and phases used as inputs. INVENTED: the tip sigma field and the fermion boundary conditions. OPEN: the Lovelock corrections to the graviton operator; the radiative corrections.

6 Step 3: the Kohn-Sham states of dirac16complex coupled to all fields

dirac16complex Ψ is a 16-component complex spinor field of the author's Clifford algebra $Cl(4,4)$; it couples to gravity through the canonical spin connection, to itself through a contact self-interaction, and to the Standard-Model fields through the three portals I11, I12 and I13. Its interacting many-body states on the block are described by Kohn-Sham theory; step 3 solves the Kohn-Sham equations and returns their energy-momentum tensor to the gravitational equations of step 1.

6.1 The field equations of step 3

$$\begin{aligned} S_\Psi &= \int \mathrm{d}^8X\,\sqrt{|g|}\,\Big[\tfrac{1}{2}\big(\bar\Psi\gamma^M D_M\Psi-(D_M\bar\Psi)\gamma^M\Psi\big)-m_\Psi\,\bar\Psi\Psi-\frac{\lambda}{2}\big(\bar\Psi\Psi\big)^2\Big],\\ m_\Psi &= 3H\mathcal{M}=H\qquad(\mathcal{M}=1/3)\end{aligned}\tag{IN-5}$$

Symbols in (IN-5).

- S_Ψ: The action of the dirac16complex field (kinetic term with the canonical spin connection, mass term and contact self-interaction). Unit: dimensionless. Status: RECORD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X=\{x_0,\dots,x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X=\{x_0,x_1,\dots,x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X=\mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1,\dots,x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\dots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, ${g^M}_N=\delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- Ψ, $\bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi=\Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M={\mathbf{e}_a}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M=g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1,M_2,\dots$): it runs over $x_0,x_1,\dots,x_7$; a repeated index is summed. Status: STANDARD.

- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex, $m_\Psi = 3H\mathcal{M} = H$. Unit: GeV. Value: 14.296 TeV. Status: RECORD.
- λ: The contact self-coupling of dirac16complex (the term $-\frac{\lambda}{2}(\bar\Psi\Psi)^2$ of the record). Unit: GeV^{-6}. Value: not fixed; Kohn-Sham states computed for $\lambda = 0$ and $\pm 7\times 10^{-5}H^{-6}$; the fold-band radiation requires $|\lambda| \lesssim 1.4\times 10^{-10}H^{-6}$ (for $\Delta N_{\mathrm{eff}} \le 0.001$). Status: RECORD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- $\mathcal{M}$: The record's dimensionless mass parameter of dirac16complex, $\mathcal{M} = 1/3$ ($m_\Psi = 3H\mathcal{M}$). Unit: dimensionless. Value: 1/3. Status: RECORD.

$$\begin{aligned}
&\gamma^a\gamma^b + \gamma^b\gamma^a = 2\eta^{ab}\,\mathbf{1}, \qquad \eta_{ab} = \mathrm{diag}(+1,+1,+1,+1,-1,-1,-1,-1), \qquad \gamma^M = \mathbf{e}_a{}^M\gamma^a,\\
&\bar\Psi = \Psi^\dagger C, \qquad C = \gamma^{x_0}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}, \qquad \Gamma_4 = \mathrm{i}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}\gamma^{x_4}, \qquad \mathcal{B} = -\mathrm{i}C\gamma^{x_4},\\
&\Psi = \Psi_A + \Psi_B, \qquad \mathcal{B}\Psi_A = -\Psi_A, \qquad \mathcal{B}\Psi_B = +\Psi_B
\end{aligned} \tag{IN-6}$$

Symbols in (IN-6).

- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- a: A frame (orthonormal) index running over $0,1,\ldots,7$, aligned with $x_0,\ldots,x_7$; raised and lowered with η_{ab}; a repeated index is summed (except in a sectional curvature $R^{ab}{}_{ab}$). Status: STANDARD.
- b: A frame index running over $0,\ldots,7$ (as a). Status: STANDARD.
- η_{ab}: The flat metric of the frame, $\eta_{ab} = \mathrm{diag}(+1,+1,+1,+1,-1,-1,-1,-1)$ (η^{ab} the same matrix). Unit: dimensionless. Status: AUTHOR.
- $\mathbf{1}$: The identity matrix (of the size fixed by the context: 16 x 16 on spinors, 3 x 3 on the generations or flavours, 2 x 2 for the sigma field). Unit: dimensionless. Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ when it is a tensor in eight dimensions). Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\mathbf{e}^a{}_M$, $\mathbf{e}_a{}^M$: The vielbein (frame field) of the metric: $g_{MN} = \eta_{ab}\,\mathbf{e}^a{}_M\mathbf{e}^b{}_N$; $\mathbf{e}_a{}^M$ is the inverse matrix. Unit: dimensionless. Status: STANDARD.
- Ψ, $\bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- C: The matrix of the Dirac adjoint, $C = \gamma^{x_0}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}$ ($\bar\Psi = \Psi^\dagger C$); in each block of the reduction $C = \varsigma_2\sigma_2$. Unit: dimensionless. Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space (also the symmetry axis of the fast-flavour analysis). Unit: length. Status: AUTHOR.
- Γ_4: The four-dimensional chirality matrix, $\Gamma_4 = \mathrm{i}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}\gamma^{x_4}$; a doublet zero mode has $\Gamma_4 = +1$ (left-handed), a singlet zero mode $\Gamma_4 = -1$ (right-handed). Unit: dimensionless. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\mathcal{B}$: The author's operator $\mathcal{B} = -\mathrm{i}C\gamma^{x_4}$; its eigenvalues ∓ 1 define the halves Ψ_A and Ψ_B, and it commutes with every interaction of the good sector. Unit: dimensionless. Status: AUTHOR.

- $\Psi_A,\ \Psi_B$: The two halves of dirac16complex, $\Psi = \Psi_A + \Psi_B$, the eigenspaces of $\mathcal{B}$ with the eigenvalues -1 and $+1$; each half carries one dark-matter species. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.

$$
\begin{aligned}
&D_M\Psi = \partial_M\Psi + \Omega_M\Psi, \qquad D_M\bar\Psi = \partial_M\bar\Psi - \bar\Psi\Omega_M, \qquad \Omega_M = \tfrac{1}{2}\,\omega_{Mab}\,\mathcal{S}^{ab}, \qquad \mathcal{S}^{ab} = \tfrac{1}{4}\big[\gamma^a,\gamma^b\big],\\
&{\omega_M}^a{}_b = \mathbf{e}^a{}_N\big(\partial_M \mathbf{e}_b{}^N + \Gamma^N{}_{MP}\,\mathbf{e}_b{}^P\big), \qquad \omega_{Mab} = \eta_{ac}\,{\omega_M}^c{}_b, \qquad g_{MN} = \eta_{ab}\,\mathbf{e}^a{}_M\,\mathbf{e}^b{}_N,\\
&\Gamma^N{}_{MP} = \tfrac{1}{2}\,g^{NQ}\big(\partial_M g_{QP} + \partial_P g_{QM} - \partial_Q g_{MP}\big)
\end{aligned}
\tag{IN-7}
$$

Symbols in (IN-7).

- $D_M,\ D_\mu$: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\Psi,\ \bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- Ω_M: The spin connection acting on spinors, the matrix $\Omega_M = \frac{1}{2}\omega_{Mab}\mathcal{S}^{ab}$. Unit: 1/length. Status: STANDARD.
- ${\omega_M}^a{}_b$: The coefficients of the spin connection of the vielbein (the Levi-Civita connection in the frame basis); $\omega_{Mab} = \eta_{ac}\,{\omega_M}^c{}_b$. Unit: 1/length. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$, aligned with $x_0, \ldots, x_7$; raised and lowered with η_{ab}; a repeated index is summed (except in a sectional curvature $R^{ab}{}_{ab}$). Status: STANDARD.
- b: A frame index running over $0, \ldots, 7$ (as a). Status: STANDARD.
- $\mathcal{S}^{ab}$: The generators of frame rotations on spinors, $\mathcal{S}^{ab} = \frac{1}{4}[\gamma^a, \gamma^b]$ (the commutator). Unit: dimensionless. Status: STANDARD.
- $\gamma^a,\ \gamma^M,\ \gamma^\mu$: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4, 4)$; $\gamma^M = \mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- $\mathbf{e}^a{}_M,\ \mathbf{e}_a{}^M$: The vielbein (frame field) of the metric: $g_{MN} = \eta_{ab}\,\mathbf{e}^a{}_M\mathbf{e}^b{}_N$; $\mathbf{e}_a{}^M$ is the inverse matrix. Unit: dimensionless. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\Gamma^N{}_{MP}$: The Christoffel symbols (the Levi-Civita connection) of the metric, $\Gamma^N{}_{MP} = \frac{1}{2}g^{NQ}(\partial_M g_{QP} + \partial_P g_{QM} - \partial_Q g_{MP})$. Unit: 1/length. Status: STANDARD.
- P: A coordinate index of eight dimensions (also $P_1, P_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- η_{ab}: The flat metric of the frame, $\eta_{ab} = \mathrm{diag}(+1, +1, +1, +1, -1, -1, -1, -1)$ (η^{ab} the same matrix). Unit: dimensionless. Status: AUTHOR.
- c: A frame index running over $0, \ldots, 7$ (as a). Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+, +, +, +, -, -, -, -$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, ${g^M}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- Q: A coordinate index of eight dimensions (also $Q_1, Q_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.

$$
\begin{aligned}
&S_{\mathrm{I}11} = \int \mathrm{d}^8X\,\sqrt{|g|}\,\frac{c_{\mathrm{V}}}{\Lambda_\star^6}\,\big(\mathrm{i}\bar\Psi\gamma^M\Psi\big)\,J^{B-L}_M, \qquad J^{B-L}_M = \sum_{i=1}^{3}\sum_{\psi} q_\psi\,\big(\mathrm{i}\bar\psi_i\gamma_M\psi_i\big),\\
&q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}, \qquad q_\ell = q_{e_R} = q_{\nu_R} = -1
\end{aligned}
\tag{IN-16}
$$

Symbols in (IN-16).

- S_{I11}: The action of the portal I11 (dirac16complex vector current times the $B-L$ current of the fermions). Unit: dimensionless. Status: INVENTED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $X=\{x_0,\ldots,x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X=\{x_0,x_1,\ldots,x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X=\mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1,\ldots,x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\ldots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, ${g^M}_N=\delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- c_{V}: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_{\mathrm{V}}=4.38\times10^{-17}$ GeV^{-2}. Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- Ψ, $\bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi=\Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M={\mathbf{e}_a}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M=g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1,M_2,\ldots$): it runs over $x_0,x_1,\ldots,x_7$; a repeated index is summed. Status: STANDARD.
- J^{B-L}_M: The $B-L$ current of the Standard-Model fermions (baryon minus lepton number), summed over the generations. Unit: GeV^7. Status: STANDARD.
- i: The generation index ($i=1,2,3$), which also labels the neutrino mass states ν_1,ν_2,ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- ψ: The label of a Standard-Model multiplet: $\psi\in\{Q,u_R,d_R,\ell,e_R,\nu_R\}$ (quark doublet, up and down quark singlets, lepton doublet, charged-lepton singlet, right-handed neutrino); a sum over ψ runs over the six. Status: STANDARD.
- q_ψ, q_f: The $B-L$ charge: q_ψ of the multiplet ψ ($q_Q=q_{u_R}=q_{d_R}=\frac{1}{3}$, $q_\ell=q_{e_R}=q_{\nu_R}=-1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- ψ_i, $\bar\psi_i$: A Standard-Model fermion field of eight dimensions: ψ_i is the multiplet $\psi\in\{Q,u_R,d_R,\ell,e_R,\nu_R\}$ of the generation i, ψ_f one fermion component f (in four-dimensional formulas its zero mode); $\bar\psi=\psi^\dagger C$ the Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$ (eight dimensions), $\mathrm{GeV}^{3/2}$ (four dimensions). Status: ASSUMED.

$$S_{\mathrm{I12}}=-\int\mathrm{d}^8X\,\sqrt{|g|}\,\frac{\bar\Psi\Psi}{\Lambda_\star^3}\,\mathcal{F},\qquad \mathcal{F}=c_{\mathrm{G}}\,G^A_{MN}G^{AMN}+c_{\mathrm{W}}\,W^I_{MN}W^{IMN}+c_{\mathrm{B}}\,B_{MN}B^{MN}\tag{IN-17}$$

Symbols in (IN-17).

- S_{I12}: The action of the portal I12 (scalar density of dirac16complex times the gauge kinetic invariants). Unit: dimensionless. Status: INVENTED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $X=\{x_0,\ldots,x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X=\{x_0,x_1,\ldots,x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X=\mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1,\ldots,x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\ldots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, ${g^M}_N=\delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.

- Ψ, $\bar{\Psi}$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar{\Psi} = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- $\mathcal{F}$: The gauge-kinetic operator of the portal I12, $\mathcal{F} = c_\mathrm{G} G^A_{MN} G^{AMN} + c_\mathrm{W} W^I_{MN} W^{IMN} + c_\mathrm{B} B_{MN} B^{MN}$; $\langle\mathcal{F}\rangle$ its mean value in an environment. Unit: GeV^4. Status: INVENTED.
- c_G: The coefficient of the gluon term of the portal I12. Unit: dimensionless. Value: $c_\mathrm{G}/\Lambda_\star^3 = c_{\mathrm{I}12}/\Lambda_\star^3$. Status: INVENTED.
- G^A_M, G_M, G_{MN}, G^A_{MN}: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G^A_M ($A = 1, \ldots, 8$), the matrix $G_M = G^A_M t^A$, the field strength G_{MN} (matrix) and its components G^A_{MN}; $G^A_{\mu\nu}$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- c_W: The coefficient of the SU(2) term of the portal I12. Unit: dimensionless. Value: $c_\mathrm{W}/\Lambda_\star^3 = c_{\mathrm{I}12}/\Lambda_\star^3$. Status: INVENTED.
- W^I_M, W_M, W_{MN}, W^I_{MN}: The SU(2) gauge field (geometric normalisation): the components W^I_M ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W^I_M \tau^I/2$, the field strength W_{MN} and its components W^I_{MN}; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- c_B: The coefficient of the hypercharge term of the portal I12. Unit: dimensionless. Value: $c_\mathrm{B}/\Lambda_\star^3 = c_{\mathrm{I}12}/\Lambda_\star^3$. Status: INVENTED.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.

$$
\begin{aligned}
S_{\mathrm{I}13} &= \int \mathrm{d}^8X\, \sqrt{|g|}\, \frac{c_\mathrm{A}}{\Lambda_\star^6} \sum_{M \neq x_0} \left(\mathrm{i}\bar{\Psi}\gamma^M P_b \Psi\right) J^B_M, \qquad P_b = \frac{1-\gamma^{x_0}}{2}, \qquad P_a = \frac{1+\gamma^{x_0}}{2}, \\
J^B_M &= \frac{1}{3}\sum_{i=1}^{3} \left(\mathrm{i}\bar{Q}_i \gamma_M Q_i + \mathrm{i}\bar{u}_{Ri}\gamma_M u_{Ri} + \mathrm{i}\bar{d}_{Ri}\gamma_M d_{Ri}\right)
\end{aligned}
\tag{IN-18}
$$

Symbols in (IN-18).

- $S_{\mathrm{I}13}$: The action of the portal I13 (the x_0-chiral vector current of dirac16complex times the baryon-number current). Unit: dimensionless. Status: INVENTED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- c_A: The coefficient of the x_0-chiral vector portal I13. Unit: dimensionless. Value: fixed through $G_\mathrm{A} = 1.278 \times 10^{-9}\ \mathrm{GeV}^{-2}$ (dark-matter relic). Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.

- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- Ψ, $\bar{\Psi}$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar{\Psi} = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M \gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- P_a, P_b: The projectors on the two x_0-chiralities, $P_a = (1+\gamma^{x_0})/2$ and $P_b = (1-\gamma^{x_0})/2$ (in a block: the upper and the lower component). Unit: dimensionless. Status: INVENTED.
- J^B_M: The baryon-number current of the quarks (one third per quark), summed over the generations. Unit: GeV^7. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- Q_i, $\bar{Q}_i$: The quark doublet of the generation i (left-handed up and down quark of one generation, three colours) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- u_R, u_{Ri}, $\bar{u}_{Ri}$: The up-type quark singlet: u_R the multiplet, u_{Ri} that of the generation i (u_R, c_R, t_R), $\bar{u}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- d_R, d_{Ri}, $\bar{d}_{Ri}$: The down-type quark singlet: d_R the multiplet, d_{Ri} that of the generation i (d_R, s_R, b_R), $\bar{d}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.

The field equation of dirac16complex, its boundary conditions and its coupling to the order-0 metric:

$$\gamma^M D_M \Psi = \Big[m_\Psi + \lambda\, \bar{\Psi}\Psi + \frac{\mathcal{F}}{\Lambda_\star^3} \Big] \Psi - \mathrm{i}\, \frac{c_\mathrm{V}}{\Lambda_\star^6}\, J^{B-L}_M \gamma^M \Psi - \mathrm{i}\, \frac{c_\mathrm{A}}{\Lambda_\star^6} \sum_{M \neq x_0} J^B_M \gamma^M P_b \Psi \qquad \text{(FE-12)}$$

Symbols in (FE-12).

- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M \gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu \Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- Ψ, $\bar{\Psi}$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar{\Psi} = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- m_Ψ: The mass parameter of dirac16complex, $m_\Psi = 3H\mathcal{M} = H$. Unit: GeV. Value: 14.296 TeV. Status: RECORD.
- λ: The contact self-coupling of dirac16complex (the term $-\frac{\lambda}{2}(\bar{\Psi}\Psi)^2$ of the record). Unit: GeV^{-6}. Value: not fixed; Kohn-Sham states computed for $\lambda = 0$ and $\pm 7 \times 10^{-5} H^{-6}$; the fold-band radiation requires $|\lambda| \lesssim 1.4 \times 10^{-10} H^{-6}$ (for $\Delta N_\mathrm{eff} \leq 0.001$). Status: RECORD.
- $\mathcal{F}$: The gauge-kinetic operator of the portal I12, $\mathcal{F} = c_\mathrm{G} G^A_{MN} G^{AMN} + c_\mathrm{W} W^I_{MN} W^{IMN} + c_\mathrm{B} B_{MN} B^{MN}$; $\langle \mathcal{F} \rangle$ its mean value in an environment. Unit: GeV^4. Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- c_V: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_\mathrm{V} = 4.38 \times 10^{-17}$ GeV^{-2}. Status: INVENTED.
- J^{B-L}_M: The $B - L$ current of the Standard-Model fermions (baryon minus lepton number), summed over the generations. Unit: GeV^7. Status: STANDARD.

- c_{A}: The coefficient of the x_0-chiral vector portal I13. Unit: dimensionless. Value: fixed through $G_{\mathrm{A}} = 1.278 \times 10^{-9}$ GeV^{-2} (dark-matter relic). Status: INVENTED.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- J^B_M: The baryon-number current of the quarks (one third per quark), summed over the generations. Unit: GeV^7. Status: STANDARD.
- $P_a,\ P_b$: The projectors on the two x_0-chiralities, $P_a = (1+\gamma^{x_0})/2$ and $P_b = (1-\gamma^{x_0})/2$ (in a block: the upper and the lower component). Unit: dimensionless. Status: INVENTED.

$$\big(1-\gamma^{x_0}\big)\Psi\big|_{y=0} = 0 \ \ \text{(even)} \quad \text{or} \quad \big(1+\gamma^{x_0}\big)\Psi\big|_{y=0} = 0 \ \ \text{(odd)}, \qquad \big(1-\gamma^{x_0}\big)\Psi\big|_{y=-L} = 0 \tag{FE-28}$$

Symbols in (FE-28).

- $\gamma^a,\ \gamma^M,\ \gamma^\mu$: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M \gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- $\Psi,\ \bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi = \Psi^\dagger C$ is its Dirac adjoint. Unit: GeV$^{7/2}$. Status: RECORD.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.

$$\gamma^M \Omega_M = \frac{3}{2}\,\frac{H}{f}\,\gamma^{x_0} + \Big(\frac{\dot b_4}{b_4} - \frac{\dot a_4}{2}\Big)\gamma^{x_4} \tag{FE-13}$$

Symbols in (FE-13).

- $\gamma^a,\ \gamma^M,\ \gamma^\mu$: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M \gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Ω_M: The spin connection acting on spinors, the matrix $\Omega_M = \frac{1}{2}\omega_{Mab}\mathcal{S}^{ab}$. Unit: 1/length. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- $\dot b_4$: The time derivative $\mathrm{d}b_4/\mathrm{d}x_4$. Unit: GeV. Status: DERIVED.
- b_4: The author's metric function of the time x_4 in the hidden and extra-time components, $b_4 > 0$. Unit: dimensionless. Status: AUTHOR.
- $\dot a_4$: The time derivative $\mathrm{d}a_4/\mathrm{d}x_4$. Unit: GeV. Status: DERIVED.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.

Its energy-momentum tensor, the source that it returns to the gravitational equations:

$$
\begin{aligned}
(\mathcal{T}_\Psi)_{MN} &= -\frac{1}{4}\Big[\bar\Psi\gamma_M D_N\Psi + \bar\Psi\gamma_N D_M\Psi - (D_M\bar\Psi)\gamma_N\Psi - (D_N\bar\Psi)\gamma_M\Psi\Big] + g_{MN}\,\mathcal{L}_\Psi,\\
\mathcal{L}_\Psi &= \tfrac{1}{2}\big(\bar\Psi\gamma^P D_P\Psi - (D_P\bar\Psi)\gamma^P\Psi\big) - m_\Psi\,\bar\Psi\Psi - \frac{\lambda}{2}\big(\bar\Psi\Psi\big)^2
\end{aligned}
\tag{FE-5}
$$

Symbols in (FE-5).

- $\mathcal{T}_\Psi$: The energy-momentum tensor of the dirac16complex field. Unit: GeV^8. Status: DERIVED.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Ψ, $\bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\ldots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- $\mathcal{L}_\Psi$: The Lagrangian density of dirac16complex (the bracket of S_Ψ). Unit: GeV^8. Status: RECORD.
- P: A coordinate index of eight dimensions (also $P_1, P_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex, $m_\Psi = 3H\mathcal{M} = H$. Unit: GeV. Value: 14.296 TeV. Status: RECORD.
- λ: The contact self-coupling of dirac16complex (the term $-\frac{\lambda}{2}(\bar\Psi\Psi)^2$ of the record). Unit: GeV^{-6}. Value: not fixed; Kohn-Sham states computed for $\lambda = 0$ and $\pm7\times10^{-5}H^{-6}$; the fold-band radiation requires $|\lambda| \lesssim 1.4\times10^{-10}H^{-6}$ (for $\Delta N_{\mathrm{eff}} \le 0.001$). Status: RECORD.

The exact reduction to the block Hamiltonian (step 2) turns the field equation into two-component equations in y; the interacting many-body state is described by Kohn-Sham orbitals of this Hamiltonian with the local mean fields of the self-interaction and of the portals:

$$
\begin{aligned}
&\Psi = \mathrm{e}^{\mathrm{i}kx_1}\,\mathrm{e}^{-\frac{3}{2}Hy}\,\mathrm{e}^{a_4/2}\,b_4^{-1}\,\mathcal{V}\chi(y,x_4)\,:\\
&\gamma^M D_M\Psi - \big(M(y) - \mathrm{i}v(y)\gamma^{x_4}\big)\Psi = \mathrm{e}^{\mathrm{i}kx_1}\,\mathrm{e}^{-\frac{3}{2}Hy}\,\mathrm{e}^{a_4/2}\,b_4^{-1}\,\gamma^{x_4}\mathcal{V}\big[\partial_{x_4}\chi + \mathrm{i}h\chi\big],\\
&h = \varsigma\Big[-\frac{\mathrm{i}}{f}\sigma_1\partial_y + M(y)\,\sigma_2 + \mathrm{e}^{-a_4}k\,\sigma_3\Big] + v(y), \qquad \mathrm{i}\,\partial_{x_4}\chi = h\chi;\\
&\text{in each block: } \gamma^{x_0} = \sigma_3, \qquad \mathcal{B} = \varsigma\varsigma_2, \qquad C = \varsigma_2\sigma_2, \qquad \Gamma_4 = -\varsigma\varsigma_2\sigma_3
\end{aligned}
\tag{FE-56}
$$

Symbols in (FE-56).

- Ψ, $\bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.

- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the hidden and extra-time components, $b_4 > 0$. Unit: dimensionless. Status: AUTHOR.
- $\mathcal{V}$: The constant 16×2 matrix of the reduction ansatz that places the two-component spinor χ of one block into the 16 components of the eight-dimensional spinor. Unit: dimensionless. Status: DERIVED.
- χ, χ_o, χ_1, χ_2: The two-component spinor of one block of the reduction (a function of y and x_4), $\chi = (\chi_1, \chi_2)^T = (a, \mathrm{i}b)^T$; χ_o the Kohn-Sham orbital o, normalised as $\int_{-L}^{0} \chi_o^\dagger \chi_o \,\mathrm{d}y = 1$. Unit: $\mathrm{GeV}^{1/2}$ (normalised orbital). Status: DERIVED.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M \gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- $M(y)$: The mass function of the block Hamiltonian (the scalar potential): $M = \mu_{\psi i}$ for a Standard-Model fermion, $M = M_\mathrm{eff}$ for dirac16complex. Unit: GeV. Status: DERIVED.
- $v(y)$: The vector potential of the block Hamiltonian (the mean field along γ^{x_4}): for dirac16complex the contact term and the mean field of I11; zero for a Standard-Model fermion. Unit: GeV. Status: DERIVED.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- h: The block Hamiltonian: the 2×2 differential operator of one block that generates the time evolution of χ, $\mathrm{i}\,\partial_{x_4}\chi = h\chi$; its eigenvalues are the energies ε. Unit: GeV. Status: DERIVED.
- ς: The first of the three signs $(\varsigma, \varsigma_2, \varsigma_3)$, each ± 1, that label the eight blocks of the reduction. Unit: dimensionless. Status: DERIVED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- σ_1, σ_2, σ_3: The Pauli matrices acting on the two components of a block spinor χ ($\sigma_3 = \mathrm{diag}(1,-1)$). Unit: dimensionless. Status: STANDARD.
- y: The label y in ∂_y: the derivative with respect to the hidden coordinate y. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- $\mathcal{B}$: The author's operator $\mathcal{B} = -\mathrm{i}C\gamma^{x_4}$; its eigenvalues ∓ 1 define the halves Ψ_A and Ψ_B, and it commutes with every interaction of the good sector. Unit: dimensionless. Status: AUTHOR.
- ς_2: The second sign that labels the blocks (± 1). Unit: dimensionless. Status: DERIVED.
- C: The matrix of the Dirac adjoint, $C = \gamma^{x_0}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}$ ($\bar\Psi = \Psi^\dagger C$); in each block of the reduction $C = \varsigma_2\sigma_2$. Unit: dimensionless. Status: AUTHOR.
- Γ_4: The four-dimensional chirality matrix, $\Gamma_4 = \mathrm{i}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}\gamma^{x_4}$; a doublet zero mode has $\Gamma_4 = +1$ (left-handed), a singlet zero mode $\Gamma_4 = -1$ (right-handed). Unit: dimensionless. Status: DERIVED.

$$h\,\chi_o = \varepsilon_o\,\chi_o, \qquad h = \varsigma\Big[-\frac{\mathrm{i}}{f}\sigma_1\partial_y + M_\mathrm{eff}(y)\,\sigma_2 + \mathrm{e}^{-a_4}k\,\sigma_3\Big] + v(y) + \frac{1-\sigma_3}{2}\,v_\mathrm{b}(y) \tag{FE-63}$$

Symbols in (FE-63).

- h: The block Hamiltonian: the 2×2 differential operator of one block that generates the time evolution of χ, $\mathrm{i}\,\partial_{x_4}\chi = h\chi$; its eigenvalues are the energies ε. Unit: GeV. Status: DERIVED.
- χ, χ_o, χ_1, χ_2: The two-component spinor of one block of the reduction (a function of y and x_4), $\chi = (\chi_1, \chi_2)^T = (a, \mathrm{i}b)^T$; χ_o the Kohn-Sham orbital o, normalised as $\int_{-L}^{0} \chi_o^\dagger \chi_o \,\mathrm{d}y = 1$. Unit: $\mathrm{GeV}^{1/2}$ (normalised orbital). Status: DERIVED.
- o: The label of a Kohn-Sham orbital (a state of dirac16complex in the cell: block, momentum shell, fold parity and level); a sum over o runs over the orbitals. Status: RECORD.
- ε, ε_o: The energy of a fermion mode (an eigenvalue of the block Hamiltonian; at $k = 0$ the four-dimensional mass); ε_o the Kohn-Sham level of the orbital o. Unit: GeV. Status: DERIVED.
- ς: The first of the three signs $(\varsigma, \varsigma_2, \varsigma_3)$, each ± 1, that label the eight blocks of the reduction. Unit: dimensionless. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- σ_1, σ_2, σ_3: The Pauli matrices acting on the two components of a block spinor χ ($\sigma_3 = \mathrm{diag}(1, -1)$). Unit: dimensionless. Status: STANDARD.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- y: The label y in ∂_y: the derivative with respect to the hidden coordinate y. Status: STANDARD.
- $M_{\mathrm{eff}}(y)$: The effective (scalar) mass field of the Kohn-Sham Hamiltonian: m_Ψ plus the mean fields of the contact term and of I12. Unit: GeV. Status: DERIVED.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- $v(y)$: The vector potential of the block Hamiltonian (the mean field along γ^{x_4}): for dirac16complex the contact term and the mean field of I11; zero for a Standard-Model fermion. Unit: GeV. Status: DERIVED.
- $v_{\mathrm{b}}(y)$: The mean field of I13 (baryon densities), acting on the lower component of χ only. Unit: GeV. Status: DERIVED.

$$M_{\mathrm{eff}}(y) = m_\Psi + \frac{15}{16}\,\lambda\, n^{\mathrm{sc}}(y) + \frac{\langle \mathcal{F} \rangle}{\Lambda_\star^3}, \qquad v(y) = -\frac{\lambda}{16}\, n(y) + \frac{c_{\mathrm{V}}}{\Lambda_\star^6} \sum_{i=1}^{3} \sum_{\psi} q_\psi\, n_{\psi i}(y),$$
$$v_{\mathrm{b}}(y) = \frac{c_{\mathrm{A}}}{\Lambda_\star^6} \sum_{i=1}^{3} \sum_{\psi} q_\psi^B\, n_{\psi i}(y) \tag{FE-64}$$

Symbols in (FE-64).

- $M_{\mathrm{eff}}(y)$: The effective (scalar) mass field of the Kohn-Sham Hamiltonian: m_Ψ plus the mean fields of the contact term and of I12. Unit: GeV. Status: DERIVED.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex, $m_\Psi = 3H\mathcal{M} = H$. Unit: GeV. Value: 14.296 TeV. Status: RECORD.
- λ: The contact self-coupling of dirac16complex (the term $-\frac{\lambda}{2}(\bar\Psi\Psi)^2$ of the record). Unit: GeV^{-6}. Value: not fixed; Kohn-Sham states computed for $\lambda = 0$ and $\pm 7 \times 10^{-5} H^{-6}$; the fold-band radiation requires $|\lambda| \lesssim 1.4 \times 10^{-10} H^{-6}$ (for $\Delta N_{\mathrm{eff}} \le 0.001$). Status: RECORD.

- n^{sc}, n_o^{sc}: The scalar density of dirac16complex (the Kohn-Sham expectation of $\bar{\Psi}\Psi$, per proper 7-volume) and n_o^{sc} the contribution of the orbital o. Unit: GeV^7. Status: RECORD.
- $\mathcal{F}$: The gauge-kinetic operator of the portal I12, $\mathcal{F} = c_{\mathrm{G}} G^A_{MN} G^{AMN} + c_{\mathrm{W}} W^I_{MN} W^{IMN} + c_{\mathrm{B}} B_{MN} B^{MN}$; $\langle \mathcal{F} \rangle$ its mean value in an environment. Unit: GeV^4. Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- $v(y)$: The vector potential of the block Hamiltonian (the mean field along γ^{x_4}): for dirac16complex the contact term and the mean field of I11; zero for a Standard-Model fermion. Unit: GeV. Status: DERIVED.
- n, n_o, $n_{\psi i}$, n_f, n_i: A number density: $n(y)$ the particle density of dirac16complex in the Kohn-Sham equations and n_o the contribution of the orbital o, $n_{\psi i}(y)$ the density of the Standard-Model fermion ψ_i (per proper 7-volume of eight dimensions); n_f the density of the fermion component f and n_i that of the neutrino mass state i (per 3-volume of four dimensions). Unit: GeV^7 (eight dimensions), GeV^3 (four dimensions). Status: STANDARD.
- c_{V}: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_{\mathrm{V}} = 4.38 \times 10^{-17}$ GeV^{-2}. Status: INVENTED.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- ψ: The label of a Standard-Model multiplet: $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ (quark doublet, up and down quark singlets, lepton doublet, charged-lepton singlet, right-handed neutrino); a sum over ψ runs over the six. Status: STANDARD.
- q_ψ, q_f: The $B - L$ charge: q_ψ of the multiplet ψ ($q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}$, $q_\ell = q_{e_R} = q_{\nu_R} = -1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- $v_{\mathrm{b}}(y)$: The mean field of I13 (baryon densities), acting on the lower component of χ only. Unit: GeV. Status: DERIVED.
- c_{A}: The coefficient of the x_0-chiral vector portal I13. Unit: dimensionless. Value: fixed through $G_{\mathrm{A}} = 1.278 \times 10^{-9}$ GeV^{-2} (dark-matter relic). Status: INVENTED.
- q_ψ^B, q_f^B: The baryon number of the multiplet ψ or of the fermion component f ($\frac{1}{3}$ for quarks, 0 for leptons). Unit: dimensionless. Status: STANDARD.

$$
\begin{aligned}
&n = \sum_o \tfrac{1}{2}\, d_o\, \phi_o\, n_o, \qquad n^{\mathrm{sc}} = \sum_o \tfrac{1}{2}\, d_o\, \phi_o\, n_o^{\mathrm{sc}}, \qquad n_o = \frac{\mathrm{e}^{-3Hy}}{\mathrm{Vol}_7}\, \chi_o^\dagger \chi_o, \\
&n_o^{\mathrm{sc}} = \frac{\mathrm{e}^{-3Hy}}{\mathrm{Vol}_7}\, \chi_o^\dagger \varsigma \sigma_2 \chi_o, \qquad n_o^{\mathrm{t}} = \frac{\mathrm{e}^{-3Hy}}{\mathrm{Vol}_7}\, \chi_o^\dagger \varsigma \sigma_3 \chi_o, \qquad d_o = 4\, r_3\big(|\vec{n}_o|^2\big), \qquad \vec{k}_o = k_{\mathrm{lat}}\, \vec{n}_o, \\
&\mathrm{Vol}_7 = \Big(\frac{2\pi}{k_{\mathrm{lat}}}\Big)^3 v_{\mathrm{t}}\, f^4, \qquad \phi_o = \frac{1}{\mathrm{e}^{(\varepsilon_o - \mu_{\mathrm{chem}})/T} + 1} \quad \text{(Mermin; or 0 and 1 for the aufbau)}
\end{aligned}
\tag{FE-65}
$$

Symbols in (FE-65).

- n, n_o, $n_{\psi i}$, n_f, n_i: A number density: $n(y)$ the particle density of dirac16complex in the Kohn-Sham equations and n_o the contribution of the orbital o, $n_{\psi i}(y)$ the density of the Standard-Model fermion ψ_i (per proper 7-volume of eight dimensions); n_f the density of the fermion component f and n_i that of the neutrino mass state i (per 3-volume of four dimensions). Unit: GeV^7 (eight dimensions), GeV^3 (four dimensions). Status: STANDARD.
- o: The label of a Kohn-Sham orbital (a state of dirac16complex in the cell: block, momentum shell, fold parity and level); a sum over o runs over the orbitals. Status: RECORD.
- d_o, d_s: A degeneracy: $d_o = 4r_3(|\vec{n}_o|^2)$ the number of states of the momentum shell of the orbital o; d_s the number of internal states (spin, colour, particle and antiparticle) of the species s. Unit: dimensionless. Status: STANDARD.
- ϕ_o: The occupation of the Kohn-Sham orbital o (Fermi-Dirac for the Mermin functional, 0 or 1 for the aufbau). Unit: dimensionless. Status: RECORD.
- n^{sc}, n_o^{sc}: The scalar density of dirac16complex (the Kohn-Sham expectation of $\bar{\Psi}\Psi$, per proper 7-volume) and n_o^{sc} the contribution of the orbital o. Unit: GeV^7. Status: RECORD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296

TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.

- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- Vol_7: The normalisation volume of a Kohn-Sham cell, $\mathrm{Vol}_7 = (2\pi/k_{\mathrm{lat}})^3 v_{\mathrm{t}} f^4$ (the proper seven-volume of a cell is $2\,\mathrm{Vol}_7(1 - \mathrm{e}^{-3HL})/(3H)$). Unit: GeV^{-6}. Value: 34.27 in units $H = 1$, $v_{\mathrm{t}} = 1$, $f = f_*$. Status: RECORD.
- χ, χ_o, χ_1, χ_2: The two-component spinor of one block of the reduction (a function of y and x_4), $\chi = (\chi_1, \chi_2)^T = (a, \mathrm{i}b)^T$; χ_o the Kohn-Sham orbital o, normalised as $\int_{-L}^{0} \chi_o^\dagger \chi_o \,\mathrm{d}y = 1$. Unit: $\mathrm{GeV}^{1/2}$ (normalised orbital). Status: DERIVED.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- ς: The first of the three signs $(\varsigma, \varsigma_2, \varsigma_3)$, each ± 1, that label the eight blocks of the reduction. Unit: dimensionless. Status: DERIVED.
- σ_1, σ_2, σ_3: The Pauli matrices acting on the two components of a block spinor χ ($\sigma_3 = \mathrm{diag}(1, -1)$). Unit: dimensionless. Status: STANDARD.
- n_o^{t}: The density of the orbital o along the momentum term ($\chi_o^\dagger \varsigma \sigma_3 \chi_o$ times the weight $\mathrm{e}^{-3Hy}/\mathrm{Vol}_7$); it carries the 3-space pressure. Unit: GeV^7. Status: RECORD.
- r_3: The number of representations of an integer as an ordered sum of three squares of integers (signs counted). Unit: dimensionless. Status: STANDARD.
- $\vec{n}_o$: The integer lattice vector of the momentum shell of the orbital o ($\vec{n} \in \mathbb{Z}^3$). Unit: dimensionless. Status: RECORD.
- $\vec{k}$, $\vec{k}_o$: The wave vector (momentum) in ordinary 3-space; $\vec{k}_o$ that of the orbital o, a point of the lattice $k_{\mathrm{lat}}\vec{n}_o$. Unit: GeV. Status: STANDARD.
- k_{lat}: The spacing of the momentum lattice of the Kohn-Sham cell. Unit: GeV. Value: $0.25\,H$. Status: RECORD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- v_{t}: The coordinate volume of the compact extra times x_5, x_6, x_7. Unit: GeV^{-3}. Value: not fixed; $v_{\mathrm{t}} = H^{-3}$ in the quoted conversions. Status: ASSUMED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- ε, ε_o: The energy of a fermion mode (an eigenvalue of the block Hamiltonian; at $k = 0$ the four-dimensional mass); ε_o the Kohn-Sham level of the orbital o. Unit: GeV. Status: DERIVED.
- μ_{chem}: The chemical potential of the Mermin occupations. Unit: GeV. Status: RECORD.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.

$$E_{\mathrm{KS}} = \sum_o d_o \,\phi_o \,\varepsilon_o - 2\,\mathrm{Vol}_7 \int_{-L}^{0} \mathrm{e}^{3Hy}\, e_{\mathrm{int}} \,\mathrm{d}y, \qquad e_{\mathrm{int}} = \frac{15}{32}\,\lambda \left(n^{\mathrm{sc}}\right)^2 - \frac{1}{32}\,\lambda\, n^2 \tag{FE-66}$$

Symbols in (FE-66).

- E_{KS}: The Kohn-Sham energy of a cell. Unit: GeV. Status: RECORD.
- o: The label of a Kohn-Sham orbital (a state of dirac16complex in the cell: block, momentum shell, fold parity and level); a sum over o runs over the orbitals. Status: RECORD.
- d_o, d_s: A degeneracy: $d_o = 4 r_3(|\vec{n}_o|^2)$ the number of states of the momentum shell of the orbital o; d_s the number of internal states (spin, colour, particle and antiparticle) of the species s. Unit: dimensionless. Status: STANDARD.
- ϕ_o: The occupation of the Kohn-Sham orbital o (Fermi-Dirac for the Mermin functional, 0 or 1 for the aufbau). Unit: dimensionless. Status: RECORD.
- ε, ε_o: The energy of a fermion mode (an eigenvalue of the block Hamiltonian; at $k = 0$ the four-dimensional mass); ε_o the Kohn-Sham level of the orbital o. Unit: GeV. Status: DERIVED.
- Vol_7: The normalisation volume of a Kohn-Sham cell, $\mathrm{Vol}_7 = (2\pi/k_{\mathrm{lat}})^3 v_{\mathrm{t}} f^4$ (the proper seven-volume of a cell is $2\,\mathrm{Vol}_7(1 - \mathrm{e}^{-3HL})/(3H)$). Unit: GeV^{-6}. Value: 34.27 in units $H = 1$, $v_{\mathrm{t}} = 1$, $f = f_*$. Status: RECORD.

- L: The depth of the regular tip: each half of a period of x_0 ends at $y=-L$; $HL=3$. Unit: length. Value: $L=3/H$ ($fL=8.926\times10^{-21}$ m at the stabilised block). Status: ASSUMED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda=-10H^2$ and the mass parameter $m_\Psi=H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- y: The hidden coordinate of the y chart, $y=\ln|\sin z|/(6H)$, with $y=0$ at the fold ($z=\pi/2$) and $y=-L$ at the regular tip; each half of a period of x_0 is the interval $-L\le y\le 0$. Unit: length. Status: DERIVED.
- e_int: The interaction energy density of the contact term in the Kohn-Sham energy and energy-momentum tensor. Unit: GeV^8. Status: RECORD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- λ: The contact self-coupling of dirac16complex (the term $-\frac{\lambda}{2}(\bar\Psi\Psi)^2$ of the record). Unit: GeV^{-6}. Value: not fixed; Kohn-Sham states computed for $\lambda=0$ and $\pm7\times10^{-5}H^{-6}$; the fold-band radiation requires $|\lambda|\lesssim1.4\times10^{-10}H^{-6}$ (for $\Delta N_\mathrm{eff}\le0.001$). Status: RECORD.
- n^sc, n_o^sc: The scalar density of dirac16complex (the Kohn-Sham expectation of $\bar\Psi\Psi$, per proper 7-volume) and n_o^sc the contribution of the orbital o. Unit: GeV^7. Status: RECORD.
- n, n_o, $n_{\psi i}$, n_f, n_i: A number density: $n(y)$ the particle density of dirac16complex in the Kohn-Sham equations and n_o the contribution of the orbital o, $n_{\psi i}(y)$ the density of the Standard-Model fermion ψ_i (per proper 7-volume of eight dimensions); n_f the density of the fermion component f and n_i that of the neutrino mass state i (per 3-volume of four dimensions). Unit: GeV^7 (eight dimensions), GeV^3 (four dimensions). Status: STANDARD.

$$\begin{aligned}\rho&=\sum_o \tfrac{1}{2}\,d_o\phi_o\,\varepsilon_o\,n_o-e_\mathrm{int},\qquad p_3=\sum_o \tfrac{1}{2}\,d_o\phi_o\,\frac{\mathrm{e}^{-a_4|\vec{k}_o|}}{3}\,n_o^\mathrm{t}+e_\mathrm{int},\qquad p_5=e_\mathrm{int},\\ p_0&=\sum_o \tfrac{1}{2}\,d_o\phi_o\Big[(\varepsilon_o-v)\,n_o-M_\mathrm{eff}\,n_o^\mathrm{sc}-\mathrm{e}^{-a_4|\vec{k}_o|}\,n_o^\mathrm{t}\Big]+e_\mathrm{int}\end{aligned}\tag{FE-67}$$

Symbols in (FE-67).

- ρ: An energy density: in eight dimensions $\rho=-\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- o: The label of a Kohn-Sham orbital (a state of dirac16complex in the cell: block, momentum shell, fold parity and level); a sum over o runs over the orbitals. Status: RECORD.
- d_o, d_s: A degeneracy: $d_o=4r_3(|\vec{n}_o|^2)$ the number of states of the momentum shell of the orbital o; d_s the number of internal states (spin, colour, particle and antiparticle) of the species s. Unit: dimensionless. Status: STANDARD.
- ϕ_o: The occupation of the Kohn-Sham orbital o (Fermi-Dirac for the Mermin functional, 0 or 1 for the aufbau). Unit: dimensionless. Status: RECORD.
- ε, ε_o: The energy of a fermion mode (an eigenvalue of the block Hamiltonian; at $k=0$ the four-dimensional mass); ε_o the Kohn-Sham level of the orbital o. Unit: GeV. Status: DERIVED.
- n, n_o, $n_{\psi i}$, n_f, n_i: A number density: $n(y)$ the particle density of dirac16complex in the Kohn-Sham equations and n_o the contribution of the orbital o, $n_{\psi i}(y)$ the density of the Standard-Model fermion ψ_i (per proper 7-volume of eight dimensions); n_f the density of the fermion component f and n_i that of the neutrino mass state i (per 3-volume of four dimensions). Unit: GeV^7 (eight dimensions), GeV^3 (four dimensions). Status: STANDARD.
- e_int: The interaction energy density of the contact term in the Kohn-Sham energy and energy-momentum tensor. Unit: GeV^8. Status: RECORD.
- p_3: The pressure of ordinary 3-space, $p_3=\mathcal{T}^{x_1}{}_{x_1}=\mathcal{T}^{x_2}{}_{x_2}=\mathcal{T}^{x_3}{}_{x_3}$ for an isotropic source. Unit: GeV^8. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- $\vec{k}$, $\vec{k}_o$: The wave vector (momentum) in ordinary 3-space; $\vec{k}_o$ that of the orbital o, a point of the lattice $k_{\mathrm{lat}}\vec{n}_o$. Unit: GeV. Status: STANDARD.
- n_o^{t}: The density of the orbital o along the momentum term ($\chi_o^\dagger \varsigma\sigma_3\chi_o$ times the weight $\mathrm{e}^{-3Hy}/\mathrm{Vol}_7$); it carries the 3-space pressure. Unit: GeV^7. Status: RECORD.
- p_5: The pressure along the extra times, $p_5 = \mathcal{T}^{x_5}{}_{x_5} = \mathcal{T}^{x_6}{}_{x_6} = \mathcal{T}^{x_7}{}_{x_7}$. Unit: GeV^8. Status: STANDARD.
- p_0: The pressure along the hidden direction x_0, $p_0 = \mathcal{T}^{x_0}{}_{x_0}$. Unit: GeV^8. Status: STANDARD.
- $v(y)$: The vector potential of the block Hamiltonian (the mean field along γ^{x_4}): for dirac16complex the contact term and the mean field of I11; zero for a Standard-Model fermion. Unit: GeV. Status: DERIVED.
- $M_{\mathrm{eff}}(y)$: The effective (scalar) mass field of the Kohn-Sham Hamiltonian: m_Ψ plus the mean fields of the contact term and of I12. Unit: GeV. Status: DERIVED.
- n^{sc}, n_o^{sc}: The scalar density of dirac16complex (the Kohn-Sham expectation of $\bar\Psi\Psi$, per proper 7-volume) and n_o^{sc} the contribution of the orbital o. Unit: GeV^7. Status: RECORD.

$$E_{\mathrm{KS}} = 2\,\mathrm{Vol}_7 \int_{-L}^{0} \mathrm{e}^{3Hy}\rho\,\mathrm{d}y, \qquad p_0\big|_{y=0} - \mathrm{e}^{-3HL}\,p_0\big|_{y=-L} = 3H\int_{-L}^{0}\mathrm{e}^{3Hy}p_5\,\mathrm{d}y \tag{FE-68}$$

Symbols in (FE-68).

- E_{KS}: The Kohn-Sham energy of a cell. Unit: GeV. Status: RECORD.
- Vol_7: The normalisation volume of a Kohn-Sham cell, $\mathrm{Vol}_7 = (2\pi/k_{\mathrm{lat}})^3 v_{\mathrm{t}} f^4$ (the proper seven-volume of a cell is $2\,\mathrm{Vol}_7(1-\mathrm{e}^{-3HL})/(3H)$). Unit: GeV^{-6}. Value: 34.27 in units $H=1$, $v_{\mathrm{t}}=1$, $f=f_*$. Status: RECORD.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y=-L$; $HL=3$. Unit: length. Value: $L=3/H$ ($fL = 8.926\times10^{-21}$ m at the stabilised block). Status: ASSUMED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y=0$ at the fold ($z=\pi/2$) and $y=-L$ at the regular tip; each half of a period of x_0 is the interval $-L\le y\le 0$. Unit: length. Status: DERIVED.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- p_0: The pressure along the hidden direction x_0, $p_0 = \mathcal{T}^{x_0}{}_{x_0}$. Unit: GeV^8. Status: STANDARD.
- p_5: The pressure along the extra times, $p_5 = \mathcal{T}^{x_5}{}_{x_5} = \mathcal{T}^{x_6}{}_{x_6} = \mathcal{T}^{x_7}{}_{x_7}$. Unit: GeV^8. Status: STANDARD.

$$\chi = (\chi_1,\ \chi_2)^T: \qquad \chi_2(0)=0\ \ \text{(even)}\quad\text{or}\quad \chi_1(0)=0\ \ \text{(odd)}, \qquad \chi_2(-L)=0 \tag{FE-69}$$

Symbols in (FE-69).

- χ, χ_o, χ_1, χ_2: The two-component spinor of one block of the reduction (a function of y and x_4), $\chi = (\chi_1,\chi_2)^T = (a,\mathrm{i}b)^T$; χ_o the Kohn-Sham orbital o, normalised as $\int_{-L}^0 \chi_o^\dagger\chi_o\,\mathrm{d}y = 1$. Unit: $\mathrm{GeV}^{1/2}$ (normalised orbital). Status: DERIVED.
- ^{T}: The transpose: a row written with ^{T} is a column. Status: STANDARD.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y=-L$; $HL=3$. Unit: length. Value: $L=3/H$ ($fL = 8.926\times10^{-21}$ m at the stabilised block). Status: ASSUMED.

6.2 The approximation of step 3

- *Kohn-Sham mean field.* The contact self-interaction enters through its Hartree term and its exact local exchange (the functional of the record, which gives the coefficients 15/16 and −1/16 of the mean fields);

correlation beyond the local exchange is not estimated (OPEN).

- *The static block.* The orbitals live on the static block of step 1 ($\mathcal{H} = \mathcal{H}_f = 0$; $a_4 = 0$ in the cell), whose response to their energy-momentum tensor is computed at first order below.
- *A periodic cell in 3-space.* The momenta along x_1, x_2, x_3 are restricted to the lattice $\vec{k}_o = k_{\rm lat}\,\vec{n}_o$ with $k_{\rm lat} = 0.25\,H$; a state is a set of occupied orbitals in a cell of the normalisation volume Vol_7.
- *A finite basis.* The orbitals are expanded in 60, 120 and 240 exact free eigenfunctions per sector and extrapolated in the basis size (Richardson); the remaining basis error is at most 1.5×10^{-9} (relative) for the energies.
- *The portal fields as external first-order mean fields.* The Standard-Model fields enter through their densities and condensates; their responses are linear (Hellmann-Feynman) and are evaluated with the physical strengths in step 8. *The self-coupling.* The Kohn-Sham machinery is tested with the benchmark $\lambda_{\rm test} = 7 \times 10^{-5}\,H^{-6}$ and with stress couplings; the cosmology of step 9 requires $\lambda \le 1.4 \times 10^{-10}\,H^{-6}$ (for $v_{\rm t} = H^{-3}$), at which the self-interaction mean field of the densest state below is

$$\frac{\max|M_{\rm eff} - m_\Psi|}{m_\Psi} = 2.7 \times 10^{-9}$$

Symbols in this equation.

- $M_{\rm eff}(y)$: The effective (scalar) mass field of the Kohn-Sham Hamiltonian: m_Ψ plus the mean fields of the contact term and of I12. Unit: GeV. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex, $m_\Psi = 3H\mathcal{M} = H$. Unit: GeV. Value: 14.296 TeV. Status: RECORD.

so for the cosmological states of UM-2 the self-interaction mean fields are negligible.

6.3 The solution of step 3

6.3.1 The numerical method

The orbitals are expanded in the exact free eigenfunctions of each sector and iterated to self-consistency (tolerance 10^{-13} in the maximum norm, safeguarded Anderson mixing). The momentum shells have the degeneracy $d_o = 4r_3(|\vec{n}_o|^2)$, r_3 the number of representations of an integer as a sum of three squares; the factor $\frac{1}{2}$ in the densities is the weight of each fold parity; and $\mathrm{Vol}_7 = (2\pi/k_{\rm lat})^3 v_{\rm t} f_*^4 = 34.27$ ($v_{\rm t} = 1$, $a_4 = 0$, units $H = 1$; the proper seven-volume of a cell is $2\,\mathrm{Vol}_7(1 - \mathrm{e}^{-3HL})/(3H)$).

6.3.2 Validation against the record

The solver reproduces the record's two reference states (at $f = 1$, $a_4 = 0$): 8 quanta with $\lambda = 157.7$: $E_{\rm KS} = -0.037786682923$ (reference -0.037786682923), the highest occupied and lowest unoccupied levels (HOMO, LUMO) to 10^{-14}, the integrated p_0 -0.08860537545 (reference -0.08860537548); 592 quanta with $\lambda = 2.131$: $E_{\rm KS} = 358.30104644787$ (identical), integrated p_0 -6.7691415097 (reference -6.7691415108) (DERIVED).

6.3.3 Ground states on the stabilised block

Two families of states (the eight $k = 0$ fold zero modes, 8 quanta; a fold-band gas, 592 quanta) with $\lambda = 0$, the benchmark $\pm\lambda_{\rm test}$ (mean fields below 5 % of m_Ψ) and the record's coupling scaled to the block (stress test, $\lambda = 157.7 f_*^4 = 0.3405\,H^{-6}$ and $2.131 f_*^4 = 0.00460\,H^{-6}$) (DERIVED; 120 functions per sector, relative Richardson errors below 2.2×10^{-9}). $E_{\rm int} = 2\,\mathrm{Vol}_7 \int_{-L}^{0} \mathrm{e}^{3Hy} e_{\rm int}\,\mathrm{d}y$ is the interaction energy, units $H = 1$:

state	$E_{\rm KS}$	$E_{\rm int}$	HOMO	LUMO	iterations
8 quanta, $\lambda = 0$	0	0	0	0.25	1
8 quanta, $+\lambda_{\rm test}$	-2.053×10^{-4}	-2.053×10^{-4}	-5.13×10^{-5}	0.24995	14
8 quanta, $-\lambda_{\rm test}$	2.053×10^{-4}	2.053×10^{-4}	5.13×10^{-5}	0.25005	14
8 quanta, stress	-0.70123	-0.42170	-0.14037	0.10859	57

state	E_{KS}	E_{int}	HOMO	LUMO	iterations
592 quanta, $\lambda = 0$	361.11614	0	0.79057	0.82916	1
592 quanta, $+\lambda_{\mathrm{test}}$	359.99060	-1.12646	0.78676	0.82535	14
592 quanta, $-\lambda_{\mathrm{test}}$	362.23906	1.12121	0.79436	0.83295	13
592 quanta, stress	307.67139	-32.2217	0.64506	0.68347	56

state	$\int p_3$	$\int p_5$	$\int p_0$	$\max\lvert M_{\mathrm{eff}} - m_\Psi\rvert$	$\max\lvert v\rvert$
8 quanta, $+\lambda_{\mathrm{test}}$	-2.05×10^{-4}	-2.05×10^{-4}	-4.447×10^{-4}	2.4×10^{-7}	6.7×10^{-4}
8 quanta, $-\lambda_{\mathrm{test}}$	2.05×10^{-4}	2.05×10^{-4}	4.447×10^{-4}	2.4×10^{-7}	6.7×10^{-4}
8 quanta, stress	-0.4217	-0.4217	-1.30076	1.98	1.83
592 quanta, $\lambda = 0$	120.372	0	-1.1×10^{-12}	0	0
592 quanta, $+\lambda_{\mathrm{test}}$	119.245	-1.1265	-2.43688	1.3×10^{-3}	0.050
592 quanta, $-\lambda_{\mathrm{test}}$	121.493	1.1212	2.43174	1.3×10^{-3}	0.050
592 quanta, stress	87.614	-32.222	-97.8188	2.09	1.88

(The integrals are $2\,\mathrm{Vol}_7 \int_{-L}^{0} \mathrm{e}^{3Hy}(\cdot)\,\mathrm{d}y$; the state of 8 quanta with $\lambda = 0$ has all of them zero.) The free gas has $\int p_3 / \int \rho = 1/3$ exactly: four-dimensional radiation. The self-interaction shifts the fold-band levels almost rigidly (the spacing between $k = 0$ and the first shell is 0.25 free, 0.25000 at λ_{test}, 0.2490 in the stress state of 8 quanta). A scalar mean field cannot gap the fold band at $k = 0$: $\chi = (\exp(f \int M_{\mathrm{eff}}\,\mathrm{d}y),\ 0)^T$ solves $\varsigma[-(\mathrm{i}/f)\sigma_1\partial_y + M_{\mathrm{eff}}\sigma_2]\chi = 0$ with both boundary conditions for every $M_{\mathrm{eff}}(y)$.

6.3.4 The dark-matter quanta

Eight quanta in the lowest odd level at $k = 0$ on the ground state of 8 quanta add exactly $8\varepsilon_{\mathrm{DM}} = 24.817633063$ (computed 24.817633063) at $\lambda = 0$ and 24.81833 at $+\lambda_{\mathrm{test}}$; they are cold, $p_3/\rho = 0$ (2.8×10^{-5} with λ_{test}), and their hidden pressure equals the block sensitivity of the x_0 eigenvalue of step 2, obtained independently:

$$\frac{p_0}{\rho} = -\frac{\mathrm{d}\ln \varepsilon_{\mathrm{DM}}}{\mathrm{d}\ln f} = 0.772034343 \qquad \text{(dark-matter quanta at rest)}$$

Symbols in this equation.

- p_0: The pressure along the hidden direction x_0, $p_0 = \mathcal{T}^{x_0}{}_{x_0}$. Unit: GeV^8. Status: STANDARD.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $\varepsilon_{\mathrm{DM}}$: The energy of the dark-matter level at rest (its mass M_{DM} on the static block). Unit: GeV. Value: $3.102204\,m_\Psi = 44.35$ TeV. Status: DERIVED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.

(DERIVED): the dark-matter mass is an x_0 eigenvalue, and it pushes on the block.

6.3.5 The Standard-Model fields acting on dirac16complex

The portal mean fields were applied with the amplitude A_{mf} and the shapes of the sources: for I11 the proper densities of the zero modes times $\mathrm{e}^{-3Hy}/\mathrm{Vol}_7$, with the density of an unpolarised fermion the average of its doublet and singlet zero-mode densities, $\varrho(y;c) = (2c/L)\,\mathrm{e}^{2cy/L}/(1-\mathrm{e}^{-2c})$ for the localisation c: the electron $\frac{1}{2}[\varrho(y;c_L)+\varrho(y;-c_R)]$ ($c_L = 5.9529$, $-c_R = 14.5539$), the up quark $\frac{1}{2}[\varrho(y;c_L)+\varrho(y;-c_R)]$ ($c_L = 3.3823$, $-c_R = 15.4300$) and the neutrinos, produced left-chiral, $\varrho(y;c_L)$; for I12 a uniform gluon condensate. The level of a Kohn-Sham orbital responds linearly (Hellmann-Feynman):

$$\Delta\varepsilon_o = \sum_f \left(G_{\mathrm{V}}\, q_f\, \tilde{I}_{of} + G_{\mathrm{A}}\, q_f^B\, \tilde{I}_{of}^{(b)}\right) n_f + \mathcal{O}_o\, \frac{c_{\mathrm{I12}}}{\Lambda_\star^3}\left[\left\langle G^A_{\mu\nu}G^{A\mu\nu}\right\rangle + \left\langle W^I_{\mu\nu}W^{I\mu\nu}\right\rangle + \left\langle B_{\mu\nu}B^{\mu\nu}\right\rangle\right], \qquad \mathcal{O}_o = \frac{\partial\varepsilon_o}{\partial m_\Psi} \quad \text{(FE-72)}$$

Symbols in (FE-72).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- ε, ε_o: The energy of a fermion mode (an eigenvalue of the block Hamiltonian; at $k = 0$ the four-dimensional mass); ε_o the Kohn-Sham level of the orbital o. Unit: GeV. Status: DERIVED.
- o: The label of a Kohn-Sham orbital (a state of dirac16complex in the cell: block, momentum shell, fold parity and level); a sum over o runs over the orbitals. Status: RECORD.
- f: The label of a fermion species: a Standard-Model multiplet of one generation (its N_{c} Weyl components share one zero-mode profile; the zero mode of each chirality) or a fermion of a medium (electron, proton, neutron); a sum over f runs over these species. Status: STANDARD.
- G_{V}: The four-dimensional strength of the vector portal I11 on the zero modes, $G_{\mathrm{V}} = c_{\mathrm{V}}H/(2f^4 v_{\mathrm{t}}\Lambda_\star^6)$. Unit: GeV^{-2}. Value: benchmark $4.38\times10^{-17}\ \mathrm{GeV}^{-2}$ ($\Delta N_{\mathrm{eff}} = 0.001$). Status: DERIVED.
- q_ψ, q_f: The $B-L$ charge: q_ψ of the multiplet ψ ($q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}$, $q_\ell = q_{e_R} = q_{\nu_R} = -1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- $\tilde{I}_{of}$: The overlap of the state o of dirac16complex with the zero mode of the fermion component f, $\tilde{I}_{of} = H^{-1}\int_{-L}^{0}(a_o^2+b_o^2)\varrho_f \mathrm{e}^{-3Hy}\mathrm{d}y$ (the integral in units of H); $\tilde{I}_Q$, $\tilde{I}_{u_R}$, $\tilde{I}_{d_R}$, $\tilde{I}_\ell$, $\tilde{I}_{e_R}$ those of the dark matter with Q, u_R, d_R, ℓ, e_R, $\tilde{I}_{\nu_R,i}$ with the right-handed neutrino of the generation i. Unit: dimensionless. Value: dark matter: $\tilde{I}_Q = 8.2722$, $\tilde{I}_\ell = 1.2524$, $\tilde{I}_{u_R} = 0.3854$, $\tilde{I}_{d_R} = 0.3961$, $\tilde{I}_{e_R} = 0.3972$. Status: DERIVED.
- G_{A}: The four-dimensional strength of the portal I13 on the zero modes, $G_{\mathrm{A}} = c_{\mathrm{A}}H/(2f^4 v_{\mathrm{t}}\Lambda_\star^6)$. Unit: GeV^{-2}. Value: $1.278\times10^{-9}\ \mathrm{GeV}^{-2}$ (dark-matter relic density). Status: DERIVED.
- q_ψ^B, q_f^B: The baryon number of the multiplet ψ or of the fermion component f ($\frac{1}{3}$ for quarks, 0 for leptons). Unit: dimensionless. Status: STANDARD.
- $\tilde{I}_{of}^{(b)}$: The overlap of the lower component of the state o with the zero mode of f, $\tilde{I}_{of}^{(b)} = H^{-1}\int_{-L}^{0} b_o^2\varrho_f \mathrm{e}^{-3Hy}\mathrm{d}y$ (the overlap of I13); $\tilde{I}_Q^{(b)}$, $\tilde{I}_{u_R}^{(b)}$, $\tilde{I}_{d_R}^{(b)}$, $\tilde{I}_{\nu_R}^{(b)}$ those of the dark matter. Unit: dimensionless. Value: dark matter: $\tilde{I}_Q^{(b)} = 2.6135$. Status: DERIVED.
- n, n_o, $n_{\psi i}$, n_f, n_i: A number density: $n(y)$ the particle density of dirac16complex in the Kohn-Sham equations and n_o the contribution of the orbital o, $n_{\psi i}(y)$ the density of the Standard-Model fermion ψ_i (per proper 7-volume of eight dimensions); n_f the density of the fermion component f and n_i that of the neutrino mass state i (per 3-volume of four dimensions). Unit: GeV^7 (eight dimensions), GeV^3 (four dimensions). Status: STANDARD.
- $\mathcal{O}_o$, $\mathcal{O}_{\mathrm{DM}}$: The scalar overlap of a state of dirac16complex, $\mathcal{O}_o = \partial\varepsilon_o/\partial m_\Psi$ (Hellmann-Feynman: the integrated scalar density of the orbital); $\mathcal{O}_{\mathrm{DM}}$ that of the dark matter. Unit: dimensionless. Value: $\mathcal{O}_{\mathrm{DM}} = 0.7072$. Status: DERIVED.
- c_{I12}: The common coefficient of the three terms of I12 when they are universal ($c_{\mathrm{G}} = c_{\mathrm{W}} = c_{\mathrm{B}} = c_{\mathrm{I12}}$). Unit: dimensionless. Value: $c_{\mathrm{I12}}/\Lambda_\star^3 = 1.106\times10^{-14}\ \mathrm{GeV}^{-3}$ (unitarity-safe benchmark). Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.

- G^A_M, G_M, G_{MN}, G^A_{MN}: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G^A_M ($A = 1, \dots, 8$), the matrix $G_M = G^A_M t^A$, the field strength G_{MN} (matrix) and its components G^A_{MN}; $G^A_{\mu\nu}$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- A: A colour (adjoint SU(3)) index running over $1, \dots, 8$; a repeated index is summed. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index running over $x_1, \dots, x_4$ (as μ). Status: STANDARD.
- W^I_M, W_M, W_{MN}, W^I_{MN}: The SU(2) gauge field (geometric normalisation): the components W^I_M ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W^I_M \tau^I/2$, the field strength W_{MN} and its components W^I_{MN}; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex, $m_\Psi = 3H\mathcal{M} = H$. Unit: GeV. Value: 14.296 TeV. Status: RECORD.

The responses of the lowest odd level (DERIVED; equal to the Hellmann-Feynman expectation of the perturbation to 10^{-8}):

portal mean field	$\mathrm{d}\varepsilon_{\mathrm{DM}}/\mathrm{d}A_{\mathrm{mf}}$	change of the fold-band gap
I11, electron density (v)	0.824779	7.9×10^{-11}
I11, up-quark density (v)	4.328803	3.0×10^{-9}
I11, neutrino density (v, lepton doublet zero mode)	1.252406	-9.3×10^{-11}
I12, gluon condensate (uniform M_{eff})	0.707196	-1.4×10^{-10}

(The changes of the fold-band gap are at the rounding level of the finite differences; in exact arithmetic every one is zero.) The broad quark doublet zero mode ($c_L = 3.38$) overlaps the dark-matter orbital most. The fold band stays massless under every portal. Every orbital has $\chi_2(-L) = 0$, so the scalar density vanishes at the tip: no tip operator built on $\bar\Psi\Psi$ reaches dirac16complex; the upper component does not vanish there, so a chiral tip current would reach Ψ_A (the projector $P_L = (1+\Gamma_4)/2 = (1 - \mathcal{B}\gamma^{x_0})/2$ annihilates every Ψ_B orbital at the tip), but UM-2 contains none (step 4). The I12 response defines the four-dimensional coefficient of the dark-matter operator, $\mathcal{O}_{\mathrm{DM}} = 0.7072$:

$$\mathcal{L}_4^{\mathrm{I12}} = -\mathcal{O}_{\mathrm{DM}}\,\frac{\bar\Psi_{\mathrm{DM}}\Psi_{\mathrm{DM}}}{\Lambda_\star^3}\big(c_{\mathrm{G}}\,G^A_{\mu\nu}G^{A\mu\nu} + c_{\mathrm{W}}\,W^I_{\mu\nu}W^{I\mu\nu} + c_{\mathrm{B}}\,B_{\mu\nu}B^{\mu\nu}\big), \qquad \mathcal{O}_{\mathrm{DM}} = \frac{\partial\varepsilon_{\mathrm{DM}}}{\partial m_\Psi},$$
$$G_{\mathrm{g}} = \mathcal{O}_{\mathrm{DM}}\,\frac{c_{\mathrm{G}}\,g_3^2}{\Lambda_\star^3}, \qquad G_{\mathrm{W}} = \mathcal{O}_{\mathrm{DM}}\,\frac{c_{\mathrm{W}}\,g_2^2}{\Lambda_\star^3}, \qquad G_{\mathrm{B}} = \mathcal{O}_{\mathrm{DM}}\,\frac{c_{\mathrm{B}}\,g_1^2}{\Lambda_\star^3} \tag{FE-71}$$

Symbols in (FE-71).

- $\mathcal{L}_4^{\mathrm{I12}}$: The four-dimensional Lagrangian of I12 on the dark matter. Unit: GeV^4. Status: DERIVED.
- $\mathcal{O}_o$, $\mathcal{O}_{\mathrm{DM}}$: The scalar overlap of a state of dirac16complex, $\mathcal{O}_o = \partial\varepsilon_o/\partial m_\Psi$ (Hellmann-Feynman: the integrated scalar density of the orbital); $\mathcal{O}_{\mathrm{DM}}$ that of the dark matter. Unit: dimensionless. Value: $\mathcal{O}_{\mathrm{DM}} = 0.7072$. Status: DERIVED.
- Ψ_{DM}, $\bar\Psi_{\mathrm{DM}}$: The four-dimensional field of a dark-matter quantum (the lowest odd level of dirac16complex on the static block, in either half) and its Dirac adjoint. Unit: $\mathrm{GeV}^{3/2}$. Status: DERIVED.

- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- c_{G}: The coefficient of the gluon term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{G}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- $G^A_M,\ G_M,\ G_{MN},\ G^A_{MN}$: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G^A_M ($A = 1, \ldots, 8$), the matrix $G_M = G^A_M t^A$, the field strength G_{MN} (matrix) and its components G^A_{MN}; $G^A_{\mu\nu}$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index running over $x_1, \ldots, x_4$ (as μ). Status: STANDARD.
- c_{W}: The coefficient of the SU(2) term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{W}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- $W^I_M,\ W_M,\ W_{MN},\ W^I_{MN}$: The SU(2) gauge field (geometric normalisation): the components W^I_M ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W^I_M \tau^I/2$, the field strength W_{MN} and its components W^I_{MN}; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- c_{B}: The coefficient of the hypercharge term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{B}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- $B_M,\ B_{MN}$: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- $\varepsilon_{\mathrm{DM}}$: The energy of the dark-matter level at rest (its mass M_{DM} on the static block). Unit: GeV. Value: $3.102204\, m_\Psi = 44.35$ TeV. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex, $m_\Psi = 3H\mathcal{M} = H$. Unit: GeV. Value: 14.296 TeV. Status: RECORD.
- G_{g}: The coupling of a dark-matter pair to canonically normalised gluon pairs through I12. Unit: GeV^{-3}. Value: 6.09×10^{-15} GeV^{-3} (benchmark). Status: DERIVED.
- g_3: The four-dimensional strong coupling ($g_3^2 = 4\pi\alpha_3$). Unit: dimensionless. Value: $g_3^2 = 0.7786$ at $2M_{\mathrm{DM}}$. Status: DERIVED.
- G_{W}: The coupling of a dark-matter pair to canonically normalised SU(2) gauge pairs through I12. Unit: GeV^{-3}. Value: 2.96×10^{-15} GeV^{-3}. Status: DERIVED.
- g_2: The four-dimensional SU(2) coupling ($g_2^2 = 4\pi\alpha_2$). Unit: dimensionless. Value: $g_2^2 = 0.3781$ at $2M_{\mathrm{DM}}$. Status: DERIVED.
- G_{B}: The coupling of a dark-matter pair to canonically normalised hypercharge pairs through I12. Unit: GeV^{-3}. Value: 1.08×10^{-15} GeV^{-3}. Status: DERIVED.
- g_1: The four-dimensional hypercharge coupling ($g_1^2 = 4\pi\alpha_1$, Standard-Model normalisation without the GUT factor). Unit: dimensionless. Value: $g_1^2 = 0.1380$ at $2M_{\mathrm{DM}}$. Status: DERIVED.

6.3.6 The back-reaction on the block

A Kohn-Sham state with the block-averaged sources ρ, p_3 and the hidden pressure $\bar{p} = (p_0 + 3p_5)/4$ displaces the block by the adiabatic response of step 1:

$$\Delta K = \frac{c_\rho\, \kappa\Delta\rho + c_3\, \kappa\Delta p_3 + c_p\, \kappa\Delta\bar{p}}{H^2}, \qquad \Delta \ln f = -\frac{\Delta K}{2K_*}, \qquad c_3 = -3c_\rho, \qquad \Delta\dot{\mathcal{H}} = -\frac{\kappa(\Delta\rho + \Delta p_3)}{2\mathcal{G}(K_*)} \tag{FE-47}$$

Symbols in (FE-47).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m^2_{21} = m_2^2 - m_1^2$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty

relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.

- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- c_ρ: The coefficient of $\kappa\Delta\rho$ in the adiabatic block response ΔK. Unit: dimensionless. Value: 0.0358521. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_\mathrm{N}\mathcal{G}(K_*) = 2.328\times10^{-36}\ \mathrm{GeV}^{-2}$. Status: AUTHOR.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- c_3: The coefficient of $\kappa\Delta p_3$ in the adiabatic block response, $c_3 = -3c_\rho$ (so radiation drops out). Unit: dimensionless. Value: -0.107556. Status: DERIVED.
- p_3: The pressure of ordinary 3-space, $p_3 = \mathcal{T}^{x_1}{}_{x_1} = \mathcal{T}^{x_2}{}_{x_2} = \mathcal{T}^{x_3}{}_{x_3}$ for an isotropic source. Unit: GeV^8. Status: STANDARD.
- c_p: The coefficient of $\kappa\Delta\bar{p}$ in the adiabatic block response. Unit: dimensionless. Value: -0.140028. Status: DERIVED.
- $\bar{p}$: The hidden pressure, the pressure averaged over the block, $\bar{p} = (p_0 + 3p_5)/4$; for a species at fixed temperature $\bar{p} = \frac{1}{4}\,\partial p/\partial \ln f$. Unit: GeV^8 (or GeV^4 for a four-dimensional density). Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4}b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- K_*: The block variable of the stabilised block, $K_* = 12 + \frac{4}{3}\sqrt{51}$ (the upper root K_+). Unit: dimensionless. Value: 21.5219. Status: DERIVED.
- $\dot{\mathcal{H}}$: The time derivative $\mathrm{d}\mathcal{H}/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- $\mathcal{G}(K)$: The coefficient of the four-dimensional Einstein-Hilbert term on a static block, $\mathcal{G}(K) = 1 - 24w_{(2)}K + 72w_{(3)}K^2 = (K^2 - 12K + 16)/16$; four-dimensional gravity is attractive for $\mathcal{G} > 0$. Unit: dimensionless. Value: $\mathcal{G}(K_*) = 20/3 + \sqrt{51} = 13.8081$. Status: DERIVED.

i.e. $\Delta K = [c_\rho\,\kappa(\rho - 3p_3) + c_p\,\kappa\bar{p}]/H^2$ (the hidden-pressure part has the stiffness $-7.1414\,H^2$), without changing the four-dimensional cosmological term at linear order (the derivative of the x_4 balance vanishes at K_*). Cell averages (units of the Kohn-Sham calculation, $H = 1$; DERIVED):

state	$\rho - 3p_3$	$\bar{p}$	$p_0 - p_5$	ΔK
8 quanta, $+\lambda_\mathrm{test}$	1.80×10^{-5}	-1.16×10^{-5}	-1.05×10^{-5}	$2.27\times10^{-6}\,\kappa$
592 quanta, $+\lambda_\mathrm{test}$	0.0987	−0.0636	−0.0574	$0.0125\,\kappa$
592 quanta, $\lambda = 0$ (free gas)	0	0	0	$-5\times10^{-13}\,\kappa$ (the numerical zero)

The shift is negligible for any realistic density (relative size $(\mathcal{H}/H)^2$, step 1). The difference $p_0 - p_5$ drives a block-shape mode outside the order-0 metric (OPEN).

6.4 The check of step 3

In its own equations.

- *Self-consistency.* The state of 8 quanta at $+\lambda_\mathrm{test}$, re-solved with the same solver (one BLAS thread), reproduces the stored E_KS, HOMO, LUMO and the integrated p_0 and ρ to 7.4×10^{-14} (absolute, units H; another BLAS kernel rounds the eigenvalues of these matrices, whose basis energies reach $583\,H$, differently by up to about $5\times10^{-13}\,H$); one further Kohn-Sham map of the converged densities moves them by 1.8×10^{-14} of their maximum (DERIVED).

- *The identities of a Kohn-Sham state.* The energy identity $E_{\mathrm{KS}} = 2\,\mathrm{Vol}_7 \int \mathrm{e}^{3Hy}\rho\,\mathrm{d}y$ holds to the storage precision for all eight stored states (4×10^{-15} in the computation), and the y-conservation law to 8.8×10^{-8} (Richardson; the raw residual falls like the inverse cube of the basis size); the observed basis orders are 4.96 (energy; 4.956 after the precision fix, version 3 quoted 4.95) and 2.97 (hidden pressure); the record states agree to 4.1×10^{-10} (DERIVED).

In concert with the other field equations.

- *The Kohn-Sham source in the gravitational equations.* The block averages ρ, p_3, $\bar{p}$ of the Kohn-Sham states that enter the block response are the integrals of the Kohn-Sham energy-momentum tensor divided by the cell weight, to 3.4×10^{-12}; the block shifts recomputed from the stored integrals with the coefficients of the background equations agree with the stored ones to 1.4×10^{-12}, and for the free gas ($p_3 = \rho/3$) the response vanishes to 9.1×10^{-13} of $c_\rho \kappa \rho$, as the exact cancellation of radiation requires (DERIVED).
- *The hidden pressure of the dark matter in three field equations.* The ratio p_0/ρ of the Kohn-Sham stress tensor (0.772034343201), the block sensitivity of the x_0 eigenvalue in closed form (0.772034343201) and by finite difference (0.772034343206), and the value that the coupled block response of step 8 uses, agree to 6.1×10^{-12} (DERIVED).
- *The portal responses.* The Hellmann-Feynman responses of the Kohn-Sham jobs equal the overlaps used in the coupled solution of step 8 to 3.0×10^{-12}, and $\mathcal{O}_{\mathrm{DM}} = \partial\varepsilon_{\mathrm{DM}}/\partial m_\Psi = 0.707196002672$ (Kohn-Sham, closed form, and the integral of the scalar density that enters the W problem of step 8) agrees with the rounded 0.7072 of the relic and mean-field computations to 5.7×10^{-6} (DERIVED).

6.5 The status of step 3

DERIVED: the Kohn-Sham states, their energies, energy-momentum tensors and identities, the dark-matter quanta and their hidden pressure, the portal responses, the block response to the Kohn-Sham sources. RECORD: the Kohn-Sham functional and the reference states. INVENTED: the portals. ASSUMED: the bound on the self-coupling (step 9). OPEN: correlation beyond the local exchange; the block-shape mode driven by $p_0 - p_5$.

7 Step 4: the electroweak sector

UM-2 has the gauge group $SU(3)_c \times SU(2)_L \times U(1)_Y$ in eight dimensions, broken at the tip by the sigma field Σ, which has no radial mode: there is no Higgs particle. Step 4 derives the charges and the couplings of the electroweak bosons W^+, W^-, Z and the photon, the anomaly cancellation of three generations, the reasons why dirac16complex has no Z coupling, and the Z widths of three generations.

7.1 The field equations of step 4

$$S_{\mathrm{gauge}} = -\frac{1}{4}\int \mathrm{d}^8X\,\sqrt{|g|}\,\Big(\frac{1}{\hat{g}_3^2}\,G^A_{MN}G^{AMN} + \frac{1}{\hat{g}_2^2}\,W^I_{MN}W^{IMN} + \frac{1}{\hat{g}_1^2}\,B_{MN}B^{MN}\Big) \tag{IN-8}$$

Symbols in (IN-8).

- S_{gauge}: The action of the gauge fields of SU(3), SU(2) and $\mathrm{U}(1)_Y$ in eight dimensions. Unit: dimensionless. Status: ASSUMED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- $\hat{g}_3$: The eight-dimensional SU(3) coupling (it multiplies the gluon kinetic term as $1/\hat{g}_3^2$); $g_3^2 = \hat{g}_3^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.

- $G^A_M,\ G_M,\ G_{MN},\ G^A_{MN}$: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G^A_M ($A = 1, \ldots, 8$), the matrix $G_M = G^A_M t^A$, the field strength G_{MN} (matrix) and its components G^A_{MN}; $G^A_{\mu\nu}$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\hat{g}_2$: The eight-dimensional SU(2) coupling; $g_2^2 = \hat{g}_2^2 / V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- $W^I_M,\ W_M,\ W_{MN},\ W^I_{MN}$: The SU(2) gauge field (geometric normalisation): the components W^I_M ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W^I_M \tau^I/2$, the field strength W_{MN} and its components W^I_{MN}; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- $\hat{g}_1$: The eight-dimensional $\mathrm{U}(1)_Y$ coupling; $g_1^2 = \hat{g}_1^2 / V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- $B_M,\ B_{MN}$: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.

$$
\begin{aligned}
&G_M = G^A_M\, t^A, \qquad G_{MN} = \partial_M G_N - \partial_N G_M - \mathrm{i}\big[G_M, G_N\big], \qquad G^A_{MN} = 2\,\mathrm{tr}\big(t^A G_{MN}\big),\\
&W_M = W^I_M\, \frac{\tau^I}{2}, \qquad W_{MN} = \partial_M W_N - \partial_N W_M - \mathrm{i}\big[W_M, W_N\big], \qquad W^I_{MN} = \mathrm{tr}\big(\tau^I W_{MN}\big),\\
&B_{MN} = \partial_M B_N - \partial_N B_M
\end{aligned}
\tag{IN-9}
$$

Symbols in (IN-9).

- $G^A_M,\ G_M,\ G_{MN},\ G^A_{MN}$: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G^A_M ($A = 1, \ldots, 8$), the matrix $G_M = G^A_M t^A$, the field strength G_{MN} (matrix) and its components G^A_{MN}; $G^A_{\mu\nu}$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- t^A: The generators of SU(3) in the fundamental representation (the Gell-Mann matrices divided by 2), $\mathrm{tr}(t^A t^B) = \frac{1}{2}\delta^{AB}$. Unit: dimensionless. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- $W^I_M,\ W_M,\ W_{MN},\ W^I_{MN}$: The SU(2) gauge field (geometric normalisation): the components W^I_M ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W^I_M \tau^I/2$, the field strength W_{MN} and its components W^I_{MN}; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- τ^I: The Pauli matrices acting on weak-isospin doublets ($\tau^I/2$ the generators of SU(2); $\tau^3 = \mathrm{diag}(1, -1)$). Unit: dimensionless. Status: STANDARD.
- $B_M,\ B_{MN}$: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.

$$W_M^{\pm} = \frac{W_M^1 \mp \mathrm{i} W_M^2}{\sqrt{2}\,\hat{g}_2}, \qquad Z_M = \frac{W_M^3 - B_M}{\hat{g}_Z}, \qquad A_M = \frac{\hat{g}_1^2\, W_M^3 + \hat{g}_2^2\, B_M}{\hat{g}_1 \hat{g}_2 \hat{g}_Z},$$
$$\hat{g}_Z = \sqrt{\hat{g}_2^2 + \hat{g}_1^2}, \qquad \cos\theta_{\mathrm{W}} = \frac{\hat{g}_2}{\hat{g}_Z}, \qquad \sin\theta_{\mathrm{W}} = \frac{\hat{g}_1}{\hat{g}_Z} \tag{IN-10}$$

Symbols in (IN-10).

- $W_M^{\pm}$, W^+, W^-: The charged weak bosons, canonically normalised: $W_M^{\pm} = (W_M^1 \mp \mathrm{i} W_M^2)/(\sqrt{2}\,\hat{g}_2)$; W^+ carries the electric charge $+e$ and W^- the charge $-e$ (they are each other's antiparticles and have one mass). Unit: GeV^3. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- W_M^I, W_M, W_{MN}, W_{MN}^I: The SU(2) gauge field (geometric normalisation): the components W_M^I ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W_M^I \tau^I/2$, the field strength W_{MN} and its components W_{MN}^I; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\hat{g}_2$: The eight-dimensional SU(2) coupling; $g_2^2 = \hat{g}_2^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- Z_M: The Z boson field, canonically normalised, $Z_M = (W_M^3 - B_M)/\hat{g}_Z$. Unit: GeV^3. Status: STANDARD.
- B_M, B_{MN}: The hypercharge gauge field U(1)$_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- $\hat{g}_Z$: The eight-dimensional coupling of the Z, $\hat{g}_Z = (\hat{g}_2^2 + \hat{g}_1^2)^{1/2}$. Unit: GeV^{-2}. Status: STANDARD.
- A_M: The photon field, canonically normalised, $A_M = (\hat{g}_1^2 W_M^3 + \hat{g}_2^2 B_M)/(\hat{g}_1 \hat{g}_2 \hat{g}_Z)$. Unit: GeV^3. Status: STANDARD.
- $\hat{g}_1$: The eight-dimensional U(1)$_Y$ coupling; $g_1^2 = \hat{g}_1^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- θ_{W}: The weak mixing angle of the couplings, $\cos\theta_{\mathrm{W}} = \hat{g}_2/\hat{g}_Z$, $\sin\theta_{\mathrm{W}} = \hat{g}_1/\hat{g}_Z$; fixed by the measured W and Z masses with $\rho_{\mathrm{EW},0}$. Unit: dimensionless. Value: $\cos^2\theta_{\mathrm{W}} = 0.776368$. Status: DERIVED.

$$D_M = \partial_M + \Omega_M - \mathrm{i} G_M^A\, t^A - \mathrm{i} W_M^I\, \frac{\tau^I}{2} - \mathrm{i}\, Y B_M, \qquad Q_{\mathrm{em}} = T_3 + Y \tag{IN-12}$$

Symbols in (IN-12).

- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu \Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- Ω_M: The spin connection acting on spinors, the matrix $\Omega_M = \frac{1}{2}\omega_{Mab}\mathcal{S}^{ab}$. Unit: 1/length. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- G_M^A, G_M, G_{MN}, G_{MN}^A: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G_M^A ($A = 1, \ldots, 8$), the matrix $G_M = G_M^A t^A$, the field strength G_{MN} (matrix) and its components G_{MN}^A; $G_{\mu\nu}^A$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- t^A: The generators of SU(3) in the fundamental representation (the Gell-Mann matrices divided by 2), $\mathrm{tr}(t^A t^B) = \frac{1}{2}\delta^{AB}$. Unit: dimensionless. Status: STANDARD.
- W_M^I, W_M, W_{MN}, W_{MN}^I: The SU(2) gauge field (geometric normalisation): the components W_M^I ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W_M^I \tau^I/2$, the field strength W_{MN} and its components W_{MN}^I; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.

- τ^I: The Pauli matrices acting on weak-isospin doublets ($\tau^I/2$ the generators of SU(2); $\tau^3 = \mathrm{diag}(1,-1)$). Unit: dimensionless. Status: STANDARD.
- Y: The hypercharge (the generator of $\mathrm{U}(1)_Y$); Y_ψ the hypercharge of the multiplet ψ (Q: $\frac{1}{6}$, u_R: $\frac{2}{3}$, d_R: $-\frac{1}{3}$, ℓ: $-\frac{1}{2}$, e_R: -1, ν_R: 0). Unit: dimensionless. Status: STANDARD.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- Q_{em}: The electric charge in units of the elementary charge, $Q_{\mathrm{em}} = T_3 + Y$; Q^f_{em} that of the fermion f. Unit: dimensionless. Status: STANDARD.
- T_3: The third component of the weak isospin ($+\frac{1}{2}$ for the upper and $-\frac{1}{2}$ for the lower member of a doublet, 0 for a singlet); T_3^f that of the fermion f. Unit: dimensionless. Status: STANDARD.

$$
\begin{aligned}
S_{\mathrm{tip}} &= \int_{y=-L} \mathrm{d}^7 X_{\mathrm{tip}} \sqrt{|g_{\mathrm{tip}}|}\, \Big[-\frac{v_\Sigma^2}{4}\, \mathrm{tr}\big((D_\mu \Sigma)^\dagger D^\mu \Sigma\big) - \mathcal{L}_{\mathrm{Y}} \Big], \\
\mathcal{L}_{\mathrm{Y}} &= \sum_{i,j=1}^{3} \Big[\bar{\ell}_i\, \Sigma \begin{pmatrix} (\mathcal{Y}_\nu)_{ij}\, \nu_{Rj} \\ (\mathcal{Y}_e)_{ij}\, e_{Rj} \end{pmatrix} + \bar{Q}_i\, \Sigma \begin{pmatrix} (\mathcal{Y}_u)_{ij}\, u_{Rj} \\ (\mathcal{Y}_d)_{ij}\, d_{Rj} \end{pmatrix} \Big] + \mathrm{h.c.}
\end{aligned}
\tag{IN-13}
$$

Symbols in (IN-13).

- S_{tip}: The action on the tip $y = -L$: the sigma field and the tip Yukawa couplings. Unit: dimensionless. Status: INVENTED.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- X_{tip}: The seven coordinates $x_1, \ldots, x_7$ of the tip $y = -L$ of the hidden direction; $\mathrm{d}^7 X_{\mathrm{tip}}$ is their coordinate volume element. Unit: length (each coordinate). Status: DERIVED.
- g_{tip}: The metric induced on the seven-dimensional tip $y = -L$ (the metric g_{MN} restricted to the tip directions $x_1, \ldots, x_7$); $|g_{\mathrm{tip}}|$ is the absolute value of its determinant. Unit: dimensionless. Status: DERIVED.
- v_Σ: The scale of the tip sigma field (its kinetic term is $-\frac{v_\Sigma^2}{4}\mathrm{tr}((D_\mu\Sigma)^\dagger D^\mu\Sigma)$); on the static block it fixes the Robin parameter β_{W}. Unit: $\mathrm{GeV}^{5/2}$. Value: fixed through $\beta_{\mathrm{W}} = 0.011912$. Status: INVENTED.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $\mathcal{L}_{\mathrm{Y}}$: The tip Yukawa Lagrangian: the couplings of the doublets to the singlets through the sigma field and the tip matrices $\mathcal{Y}$ (plus the Hermitian conjugate, h.c.). Unit: GeV^7. Status: INVENTED.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- j: A second generation index ($j = 1, 2, 3$). Status: STANDARD.
- ℓ_i, $\bar{\ell}_i$: The lepton doublet of the generation i (left-handed neutrino and charged lepton) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- $\mathcal{Y}_\nu$, $\mathcal{Y}_e$, $\mathcal{Y}_u$, $\mathcal{Y}_d$: The complex 3×3 tip Yukawa matrices of the neutrino, charged-lepton, up-quark and down-quark sectors (generation indices i, j); on the static block they become the tip mixing matrices $\mathcal{K}$ (the normalisation of that relation is not fixed: OPEN). Unit: dimensionless. Status: INVENTED.
- ν_R, ν_{Ri}: The right-handed neutrino (a gauge singlet): ν_R the multiplet, ν_{Ri} that of the generation i; its zero

modes give Dirac neutrino masses. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.

- e_R, e_{Ri}: The right-handed charged-lepton singlet: e_R the multiplet, e_{Ri} that of the generation i (e_R, μ_R, τ_R). Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- Q_i, $\bar{Q}_i$: The quark doublet of the generation i (left-handed up and down quark of one generation, three colours) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- u_R, u_{Ri}, $\bar{u}_{Ri}$: The up-type quark singlet: u_R the multiplet, u_{Ri} that of the generation i (u_R, c_R, t_R), $\bar{u}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- d_R, d_{Ri}, $\bar{d}_{Ri}$: The down-type quark singlet: d_R the multiplet, d_{Ri} that of the generation i (d_R, s_R, b_R), $\bar{d}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.

$$D_\mu\Sigma = \partial_\mu\Sigma - \mathrm{i}W^I_\mu\,\frac{\tau^I}{2}\,\Sigma + \mathrm{i}B_\mu\,\Sigma\,\frac{\tau^3}{2}, \qquad \Sigma^\dagger\Sigma = \mathbf{1}, \qquad \Sigma \to U_L\,\Sigma\,U_Y^\dagger, \qquad U_Y = \mathrm{e}^{\mathrm{i}\vartheta\tau^3/2} \tag{IN-14}$$

Symbols in (IN-14).

- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- W^I_M, W_M, W_{MN}, W^I_{MN}: The SU(2) gauge field (geometric normalisation): the components W^I_M ($I = 1,2,3$; W^3 the third component), the matrix $W_M = W^I_M\tau^I/2$, the field strength W_{MN} and its components W^I_{MN}; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1,2,3$; a repeated index is summed. Status: STANDARD.
- τ^I: The Pauli matrices acting on weak-isospin doublets ($\tau^I/2$ the generators of SU(2); $\tau^3 = \mathrm{diag}(1,-1)$). Unit: dimensionless. Status: STANDARD.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $\mathbf{1}$: The identity matrix (of the size fixed by the context: 16 x 16 on spinors, 3 x 3 on the generations or flavours, 2 x 2 for the sigma field). Unit: dimensionless. Status: STANDARD.
- U_L: An SU(2) gauge transformation (a unitary 2×2 matrix acting on doublets from the left). Unit: dimensionless. Status: STANDARD.
- U_Y: A $\mathrm{U}(1)_Y$ gauge transformation of the sigma field, $U_Y = \mathrm{e}^{\mathrm{i}\vartheta\tau^3/2}$. Unit: dimensionless. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- ϑ: The angle (gauge parameter) of a $\mathrm{U}(1)_Y$ transformation. Unit: dimensionless. Status: STANDARD.

$$-\frac{v_\Sigma^2}{4}\,\mathrm{tr}\big((D_\mu\Sigma)^\dagger D^\mu\Sigma\big)\Big|_{\Sigma=\mathbf{1}} = -\frac{v_\Sigma^2}{8}\left[2\hat{g}_2^2\,W^+_\mu W^{-\mu} + \hat{g}_Z^2\,Z_\mu Z^\mu\right] \tag{IN-15}$$

Symbols in (IN-15).

- v_Σ: The scale of the tip sigma field (its kinetic term is $-\frac{v_\Sigma^2}{4}\mathrm{tr}((D_\mu\Sigma)^\dagger D^\mu\Sigma)$); on the static block it fixes the Robin parameter β_W. Unit: $\mathrm{GeV}^{5/2}$. Value: fixed through $\beta_\mathrm{W} = 0.011912$. Status: INVENTED.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.

- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $\mathbf{1}$: The identity matrix (of the size fixed by the context: 16 x 16 on spinors, 3 x 3 on the generations or flavours, 2 x 2 for the sigma field). Unit: dimensionless. Status: STANDARD.
- $\hat{g}_2$: The eight-dimensional SU(2) coupling; $g_2^2 = \hat{g}_2^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- $W_M^\pm$, W^+, W^-: The charged weak bosons, canonically normalised: $W_M^\pm = (W_M^1 \mp \mathrm{i}W_M^2)/(\sqrt{2}\,\hat{g}_2)$; W^+ carries the electric charge $+e$ and W^- the charge $-e$ (they are each other's antiparticles and have one mass). Unit: GeV^3. Status: STANDARD.
- $\hat{g}_Z$: The eight-dimensional coupling of the Z, $\hat{g}_Z = (\hat{g}_2^2 + \hat{g}_1^2)^{1/2}$. Unit: GeV^{-2}. Status: STANDARD.
- Z_M: The Z boson field, canonically normalised, $Z_M = (W_M^3 - B_M)/\hat{g}_Z$. Unit: GeV^3. Status: STANDARD.

The halves of dirac16complex and the four-dimensional chirality come from the Clifford algebra of the author:

$$\begin{aligned}
&\gamma^a\gamma^b + \gamma^b\gamma^a = 2\eta^{ab}\,\mathbf{1}, \qquad \eta_{ab} = \mathrm{diag}(+1,+1,+1,+1,-1,-1,-1,-1), \qquad \gamma^M = \mathbf{e}_a{}^M\gamma^a,\\
&\bar{\Psi} = \Psi^\dagger C, \qquad C = \gamma^{x_0}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}, \qquad \Gamma_4 = \mathrm{i}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}\gamma^{x_4}, \qquad \mathcal{B} = -\mathrm{i}C\gamma^{x_4},\\
&\Psi = \Psi_A + \Psi_B, \qquad \mathcal{B}\Psi_A = -\Psi_A, \qquad \mathcal{B}\Psi_B = +\Psi_B
\end{aligned} \tag{IN-6}$$

Symbols in (IN-6).

- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, ... the matrices along x_0, x_4, ...), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$, aligned with $x_0, \ldots, x_7$; raised and lowered with η_{ab}; a repeated index is summed (except in a sectional curvature $R^{ab}{}_{ab}$). Status: STANDARD.
- b: A frame index running over $0, \ldots, 7$ (as a). Status: STANDARD.
- η_{ab}: The flat metric of the frame, $\eta_{ab} = \mathrm{diag}(+1,+1,+1,+1,-1,-1,-1,-1)$ (η^{ab} the same matrix). Unit: dimensionless. Status: AUTHOR.
- $\mathbf{1}$: The identity matrix (of the size fixed by the context: 16 x 16 on spinors, 3 x 3 on the generations or flavours, 2 x 2 for the sigma field). Unit: dimensionless. Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ when it is a tensor in eight dimensions). Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\mathbf{e}^a{}_M$, $\mathbf{e}_a{}^M$: The vielbein (frame field) of the metric: $g_{MN} = \eta_{ab}\,\mathbf{e}^a{}_M\mathbf{e}^b{}_N$; $\mathbf{e}_a{}^M$ is the inverse matrix. Unit: dimensionless. Status: STANDARD.
- Ψ, $\bar{\Psi}$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar{\Psi} = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- C: The matrix of the Dirac adjoint, $C = \gamma^{x_0}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}$ ($\bar{\Psi} = \Psi^\dagger C$); in each block of the reduction $C = \varsigma_2\sigma_2$. Unit: dimensionless. Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.

- x_3: The third coordinate of ordinary 3-space (also the symmetry axis of the fast-flavour analysis). Unit: length. Status: AUTHOR.
- Γ_4: The four-dimensional chirality matrix, $\Gamma_4 = \mathrm{i}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}\gamma^{x_4}$; a doublet zero mode has $\Gamma_4 = +1$ (left-handed), a singlet zero mode $\Gamma_4 = -1$ (right-handed). Unit: dimensionless. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\mathcal{B}$: The author's operator $\mathcal{B} = -\mathrm{i}C\gamma^{x_4}$; its eigenvalues ∓ 1 define the halves Ψ_A and Ψ_B, and it commutes with every interaction of the good sector. Unit: dimensionless. Status: AUTHOR.
- Ψ_A, Ψ_B: The two halves of dirac16complex, $\Psi = \Psi_A + \Psi_B$, the eigenspaces of $\mathcal{B}$ with the eigenvalues -1 and $+1$; each half carries one dark-matter species. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.

The energy-momentum tensor of the tip, whose gravitational effect is part of the rest energies of $W^{\pm}$, Z and the fermions:

$$(\mathcal{T}_{\mathrm{tip}})_{\mu\nu} = \delta_{\mathrm{tip}}\Big[\frac{v_\Sigma^2}{4}\,\mathrm{tr}\big((D_\mu\Sigma)^\dagger D_\nu\Sigma + (D_\nu\Sigma)^\dagger D_\mu\Sigma\big) + g_{\mu\nu}\,\mathcal{L}_{\mathrm{tip}}\Big], \qquad \delta_{\mathrm{tip}} = \frac{\delta_{\mathrm{D}}(y+L)}{f},$$
$$\mathcal{L}_{\mathrm{tip}} = -\frac{v_\Sigma^2}{4}\,\mathrm{tr}\big((D_\mu\Sigma)^\dagger D^\mu\Sigma\big) - \mathcal{L}_{\mathrm{Y}} \tag{FE-8}$$

Symbols in (FE-8).

- $\mathcal{T}_{\mathrm{tip}}$: The energy-momentum tensor of the tip terms (non-zero only for the tip directions). Unit: GeV^8. Status: DERIVED.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index running over $x_1, \ldots, x_4$ (as μ). Status: STANDARD.
- δ_{tip}: The density that places a tip term in the field equations of eight dimensions, $\delta_{\mathrm{tip}} = \delta_{\mathrm{D}}(y+L)/f$ (the Dirac delta function at the tip with the proper length of the hidden direction). Unit: 1/length. Status: DERIVED.
- v_Σ: The scale of the tip sigma field (its kinetic term is $-\frac{v_\Sigma^2}{4}\mathrm{tr}((D_\mu\Sigma)^\dagger D^\mu\Sigma)$); on the static block it fixes the Robin parameter β_{W}. Unit: $\mathrm{GeV}^{5/2}$. Value: fixed through $\beta_{\mathrm{W}} = 0.011912$. Status: INVENTED.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- $\mathcal{L}_{\mathrm{tip}}$: The Lagrangian density of the tip action (the bracket of S_{tip}). Unit: GeV^7. Status: INVENTED.
- δ_{D}: The Dirac delta function: $\delta_{\mathrm{D}}(y+L)$ is zero except at $y = -L$ and has the unit integral over y. Unit: 1/length. Status: STANDARD.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4}b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- $\mathcal{L}_{\mathrm{Y}}$: The tip Yukawa Lagrangian: the couplings of the doublets to the singlets through the sigma field and the tip matrices $\mathcal{Y}$ (plus the Hermitian conjugate, h.c.). Unit: GeV^7. Status: INVENTED.

The Yang-Mills equations with the fermion currents and the tip currents, and the sigma-field equation (the Noether identity of the gauge symmetry, not independent):

$$J_3^{AN} = \sum_{i=1}^{3} \big(\mathrm{i}\bar{Q}_i\gamma^N t^A Q_i + \mathrm{i}\bar{u}_{Ri}\gamma^N t^A u_{Ri} + \mathrm{i}\bar{d}_{Ri}\gamma^N t^A d_{Ri}\big), \qquad J_2^{IN} = \sum_{i=1}^{3} \Big(\mathrm{i}\bar{Q}_i\gamma^N \frac{\tau^I}{2} Q_i + \mathrm{i}\bar{\ell}_i\gamma^N \frac{\tau^I}{2}\ell_i\Big),$$
$$J_1^N = \sum_{i=1}^{3}\sum_{\psi} Y_\psi\, \mathrm{i}\bar{\psi}_i\gamma^N\psi_i, \qquad J_{\mathrm{em}}^N = J_2^{3N} + J_1^N, \qquad J_2^{\pm N} = \frac{J_2^{1N} \mp \mathrm{i}J_2^{2N}}{\sqrt{2}} \tag{FE-20}$$

Symbols in (FE-20).

- J_3^{AN}: The colour current of the quarks (the source of the gluons), summed over the generations. Unit: GeV^7. Status: STANDARD.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- Q_i, $\bar{Q}_i$: The quark doublet of the generation i (left-handed up and down quark of one generation, three colours) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4, 4)$; $\gamma^M = \mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- t^A: The generators of SU(3) in the fundamental representation (the Gell-Mann matrices divided by 2), $\mathrm{tr}(t^A t^B) = \frac{1}{2}\delta^{AB}$. Unit: dimensionless. Status: STANDARD.
- u_R, u_{Ri}, $\bar{u}_{Ri}$: The up-type quark singlet: u_R the multiplet, u_{Ri} that of the generation i (u_R, c_R, t_R), $\bar{u}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- d_R, d_{Ri}, $\bar{d}_{Ri}$: The down-type quark singlet: d_R the multiplet, d_{Ri} that of the generation i (d_R, s_R, b_R), $\bar{d}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- J_2^{IN}: The weak-isospin current of the doublets Q_i, ℓ_i (the source of W^I). Unit: GeV^7. Status: STANDARD.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- τ^I: The Pauli matrices acting on weak-isospin doublets ($\tau^I/2$ the generators of SU(2); $\tau^3 = \mathrm{diag}(1, -1)$). Unit: dimensionless. Status: STANDARD.
- ℓ_i, $\bar{\ell}_i$: The lepton doublet of the generation i (left-handed neutrino and charged lepton) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- J_1^N: The hypercharge current of all fermions (the source of B). Unit: GeV^7. Status: STANDARD.
- ψ: The label of a Standard-Model multiplet: $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ (quark doublet, up and down quark singlets, lepton doublet, charged-lepton singlet, right-handed neutrino); a sum over ψ runs over the six. Status: STANDARD.
- Y: The hypercharge (the generator of $\mathrm{U}(1)_Y$); Y_ψ the hypercharge of the multiplet ψ (Q: $\frac{1}{6}$, u_R: $\frac{2}{3}$, d_R: $-\frac{1}{3}$, ℓ: $-\frac{1}{2}$, e_R: -1, ν_R: 0). Unit: dimensionless. Status: STANDARD.
- ψ_i, $\bar{\psi}_i$: A Standard-Model fermion field of eight dimensions: ψ_i is the multiplet $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ of the generation i, ψ_f one fermion component f (in four-dimensional formulas its zero mode); $\bar{\psi} = \psi^\dagger C$ the Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$ (eight dimensions), $\mathrm{GeV}^{3/2}$ (four dimensions). Status: ASSUMED.
- J_{em}^N: The electromagnetic current, $J_{\mathrm{em}} = J_2^3 + J_1$ (in units of the elementary charge). Unit: GeV^7. Status: STANDARD.
- $J_2^{\pm N}$: The charged combinations of the isospin current, $J_2^\pm = (J_2^1 \mp \mathrm{i}J_2^2)/\sqrt{2}$ (the sources of $W^\pm$). Unit: GeV^7. Status: STANDARD.

$$\nabla_M\Big(\frac{G^{MN}}{\hat{g}_{3,\mathrm{eff}}^2}\Big) - \mathrm{i}\Big[G_M,\ \frac{G^{MN}}{\hat{g}_{3,\mathrm{eff}}^2}\Big] = J_3^{AN}\, t^A \tag{FE-22}$$

Symbols in (FE-22).

- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols ${\Gamma^N}_{MP}$); ∇_M acts on the index M, $\nabla^M = g^{MN}\nabla_N$. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- G^A_M, G_M, G_{MN}, G^A_{MN}: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G^A_M ($A = 1, \ldots, 8$), the matrix $G_M = G^A_M t^A$, the field strength G_{MN} (matrix) and its components G^A_{MN}; $G^A_{\mu\nu}$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\hat{g}_{3,\mathrm{eff}}$, $\hat{g}_{2,\mathrm{eff}}$, $\hat{g}_{1,\mathrm{eff}}$: The field-dependent eight-dimensional couplings produced by I12: $1/\hat{g}^2_{i,\mathrm{eff}} = 1/\hat{g}^2_i + 4c\,\bar{\Psi}\Psi/\Lambda^3_\star$ with $c = c_\mathrm{G}, c_\mathrm{W}, c_\mathrm{B}$. Unit: GeV^{-2}. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- J^{AN}_3: The colour current of the quarks (the source of the gluons), summed over the generations. Unit: GeV^7. Status: STANDARD.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- t^A: The generators of SU(3) in the fundamental representation (the Gell-Mann matrices divided by 2), $\mathrm{tr}(t^A t^B) = \frac{1}{2}\delta^{AB}$. Unit: dimensionless. Status: STANDARD.

$$\nabla_M\Big(\frac{W^{MN}}{\hat{g}^2_{2,\mathrm{eff}}}\Big) - \mathrm{i}\Big[W_M,\ \frac{W^{MN}}{\hat{g}^2_{2,\mathrm{eff}}}\Big] = J^{IN}_2\,\frac{\tau^I}{2} + \delta_\mathrm{tip}\,\frac{v^2_\Sigma}{4}\Big(W^N - B^N\,\frac{\tau^3}{2}\Big) \qquad (\Sigma = \mathbf{1}) \qquad \text{(FE-23)}$$

Symbols in (FE-23).

- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols ${\Gamma^N}_{MP}$); ∇_M acts on the index M, $\nabla^M = g^{MN}\nabla_N$. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- W^I_M, W_M, W_{MN}, W^I_{MN}: The SU(2) gauge field (geometric normalisation): the components W^I_M ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W^I_M \tau^I/2$, the field strength W_{MN} and its components W^I_{MN}; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\hat{g}_{3,\mathrm{eff}}$, $\hat{g}_{2,\mathrm{eff}}$, $\hat{g}_{1,\mathrm{eff}}$: The field-dependent eight-dimensional couplings produced by I12: $1/\hat{g}^2_{i,\mathrm{eff}} = 1/\hat{g}^2_i + 4c\,\bar{\Psi}\Psi/\Lambda^3_\star$ with $c = c_\mathrm{G}, c_\mathrm{W}, c_\mathrm{B}$. Unit: GeV^{-2}. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- J^{IN}_2: The weak-isospin current of the doublets Q_i, ℓ_i (the source of W^I). Unit: GeV^7. Status: STANDARD.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- τ^I: The Pauli matrices acting on weak-isospin doublets ($\tau^I/2$ the generators of SU(2); $\tau^3 = \mathrm{diag}(1, -1)$). Unit: dimensionless. Status: STANDARD.
- δ_tip: The density that places a tip term in the field equations of eight dimensions, $\delta_\mathrm{tip} = \delta_\mathrm{D}(y + L)/f$ (the Dirac delta function at the tip with the proper length of the hidden direction). Unit: 1/length. Status: DERIVED.
- v_Σ: The scale of the tip sigma field (its kinetic term is $-\frac{v^2_\Sigma}{4}\mathrm{tr}((D_\mu\Sigma)^\dagger D^\mu\Sigma)$); on the static block it fixes the Robin parameter β_W. Unit: $\mathrm{GeV}^{5/2}$. Value: fixed through $\beta_\mathrm{W} = 0.011912$. Status: INVENTED.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- $\mathbf{1}$: The identity matrix (of the size fixed by the context: 16 x 16 on spinors, 3 x 3 on the generations or flavours, 2 x 2 for the sigma field). Unit: dimensionless. Status: STANDARD.

$$\nabla_M\Big(\frac{B^{MN}}{\hat{g}_{1,\mathrm{eff}}^2}\Big) = J_1^N - \delta_{\mathrm{tip}}\,\frac{v_\Sigma^2}{4}\,\big(W^{3N} - B^N\big) \qquad (\Sigma = \mathbf{1}) \tag{FE-24}$$

Symbols in (FE-24).

- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols $\Gamma^N{}_{MP}$); ∇_M acts on the index M, $\nabla^M = g^{MN}\nabla_N$. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\hat{g}_{3,\mathrm{eff}}$, $\hat{g}_{2,\mathrm{eff}}$, $\hat{g}_{1,\mathrm{eff}}$: The field-dependent eight-dimensional couplings produced by I12: $1/\hat{g}_{i,\mathrm{eff}}^2 = 1/\hat{g}_i^2 + 4c\,\bar{\Psi}\Psi/\Lambda_\star^3$ with $c = c_{\mathrm{G}}, c_{\mathrm{W}}, c_{\mathrm{B}}$. Unit: GeV^{-2}. Status: DERIVED.
- J_1^N: The hypercharge current of all fermions (the source of B). Unit: GeV^7. Status: STANDARD.
- δ_{tip}: The density that places a tip term in the field equations of eight dimensions, $\delta_{\mathrm{tip}} = \delta_{\mathrm{D}}(y + L)/f$ (the Dirac delta function at the tip with the proper length of the hidden direction). Unit: 1/length. Status: DERIVED.
- v_Σ: The scale of the tip sigma field (its kinetic term is $-\frac{v_\Sigma^2}{4}\mathrm{tr}((D_\mu\Sigma)^\dagger D^\mu\Sigma)$); on the static block it fixes the Robin parameter β_{W}. Unit: $\mathrm{GeV}^{5/2}$. Value: fixed through $\beta_{\mathrm{W}} = 0.011912$. Status: INVENTED.
- W_M^I, W_M, W_{MN}, W_{MN}^I: The SU(2) gauge field (geometric normalisation): the components W_M^I ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W_M^I\tau^I/2$, the field strength W_{MN} and its components W_{MN}^I; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- $\mathbf{1}$: The identity matrix (of the size fixed by the context: 16 x 16 on spinors, 3 x 3 on the generations or flavours, 2 x 2 for the sigma field). Unit: dimensionless. Status: STANDARD.

$$\begin{aligned}
&\nabla_M W^{\pm MN} = \hat{g}_2\,J_2^{\pm N} + \delta_{\mathrm{tip}}\,\frac{v_\Sigma^2}{4}\,\hat{g}_2^2\,W^{\pm N}, \qquad \nabla_M A^{MN} = \hat{e}\,J_{\mathrm{em}}^N, \qquad \hat{e} = \frac{\hat{g}_2\hat{g}_1}{\hat{g}_Z},\\
&\nabla_M Z^{MN} = \hat{g}_Z\big(J_2^{3N} - \sin^2\theta_{\mathrm{W}}\,J_{\mathrm{em}}^N\big) + \delta_{\mathrm{tip}}\,\frac{v_\Sigma^2}{4}\,\hat{g}_Z^2\,Z^N,\\
&W^{\pm MN} = \nabla^M W^{\pm N} - \nabla^N W^{\pm M}, \qquad Z^{MN} = \nabla^M Z^N - \nabla^N Z^M, \qquad A^{MN} = \nabla^M A^N - \nabla^N A^M
\end{aligned} \tag{FE-25}$$

Symbols in (FE-25).

- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols $\Gamma^N{}_{MP}$); ∇_M acts on the index M, $\nabla^M = g^{MN}\nabla_N$. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $W_M^\pm$, W^+, W^-: The charged weak bosons, canonically normalised: $W_M^\pm = (W_M^1 \mp \mathrm{i}W_M^2)/(\sqrt{2}\,\hat{g}_2)$; W^+ carries the electric charge $+e$ and W^- the charge $-e$ (they are each other's antiparticles and have one mass). Unit: GeV^3. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\hat{g}_2$: The eight-dimensional SU(2) coupling; $g_2^2 = \hat{g}_2^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- $J_2^{\pm N}$: The charged combinations of the isospin current, $J_2^\pm = (J_2^1 \mp \mathrm{i}J_2^2)/\sqrt{2}$ (the sources of $W^\pm$). Unit: GeV^7. Status: STANDARD.
- δ_{tip}: The density that places a tip term in the field equations of eight dimensions, $\delta_{\mathrm{tip}} = \delta_{\mathrm{D}}(y + L)/f$ (the Dirac delta function at the tip with the proper length of the hidden direction). Unit: 1/length. Status: DERIVED.

- v_Σ: The scale of the tip sigma field (its kinetic term is $-\frac{v_\Sigma^2}{4}\mathrm{tr}((D_\mu\Sigma)^\dagger D^\mu\Sigma)$); on the static block it fixes the Robin parameter β_{W}. Unit: $\mathrm{GeV}^{5/2}$. Value: fixed through $\beta_{\mathrm{W}} = 0.011912$. Status: INVENTED.
- A_M: The photon field, canonically normalised, $A_M = (\hat{g}_1^2 W_M^3 + \hat{g}_2^2 B_M)/(\hat{g}_1\hat{g}_2\hat{g}_Z)$. Unit: GeV^3. Status: STANDARD.
- $\hat{e}$: The eight-dimensional electric coupling, $\hat{e} = \hat{g}_2\hat{g}_1/\hat{g}_Z$ ($e^2 = \hat{e}^2/V_\perp$). Unit: GeV^{-2}. Status: STANDARD.
- J^N_{em}: The electromagnetic current, $J_{\mathrm{em}} = J_2^3 + J_1$ (in units of the elementary charge). Unit: GeV^7. Status: STANDARD.
- $\hat{g}_1$: The eight-dimensional $\mathrm{U}(1)_Y$ coupling; $g_1^2 = \hat{g}_1^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- $\hat{g}_Z$: The eight-dimensional coupling of the Z, $\hat{g}_Z = (\hat{g}_2^2 + \hat{g}_1^2)^{1/2}$. Unit: GeV^{-2}. Status: STANDARD.
- Z_M: The Z boson field, canonically normalised, $Z_M = (W_M^3 - B_M)/\hat{g}_Z$. Unit: GeV^3. Status: STANDARD.
- J_2^{IN}: The weak-isospin current of the doublets Q_i, ℓ_i (the source of W^I). Unit: GeV^7. Status: STANDARD.
- θ_{W}: The weak mixing angle of the couplings, $\cos\theta_{\mathrm{W}} = \hat{g}_2/\hat{g}_Z$, $\sin\theta_{\mathrm{W}} = \hat{g}_1/\hat{g}_Z$; fixed by the measured W and Z masses with $\rho_{\mathrm{EW},0}$. Unit: dimensionless. Value: $\cos^2\theta_{\mathrm{W}} = 0.776368$. Status: DERIVED.

$$\begin{aligned}
&\nabla^\mu j_\mu - \mathrm{i}\big[W^\mu,\ j_\mu\big] = -\mathrm{i}\,\frac{\tau^I}{2}\sum_{i,j=1}^{3}\Big[\bar{\ell}_i\,\frac{\tau^I}{2}\,\Sigma\begin{pmatrix}(\mathcal{Y}_\nu)_{ij}\,\nu_{Rj}\\(\mathcal{Y}_e)_{ij}\,e_{Rj}\end{pmatrix} + \bar{Q}_i\,\frac{\tau^I}{2}\,\Sigma\begin{pmatrix}(\mathcal{Y}_u)_{ij}\,u_{Rj}\\(\mathcal{Y}_d)_{ij}\,d_{Rj}\end{pmatrix} - \mathrm{h.c.}\Big],\\
&j_\mu = j_\mu^I\,\frac{\tau^I}{2}, \qquad j_\mu^I = \frac{\mathrm{i}v_\Sigma^2}{4}\,\mathrm{tr}\big(\tau^I(D_\mu\Sigma)\Sigma^\dagger\big), \qquad j_\mu^I\big|_{\Sigma=\mathbf{1}} = \frac{v_\Sigma^2}{4}\big(W_\mu^I - \delta^{I3}B_\mu\big)
\end{aligned} \tag{FE-26}$$

Symbols in (FE-26).

- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols $\Gamma^N{}_{MP}$); ∇_M acts on the index M, $\nabla^M = g^{MN}\nabla_N$. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- j_μ, j_μ^I: The current of the sigma field on the tip, $j_\mu^I = \frac{\mathrm{i}v_\Sigma^2}{4}\mathrm{tr}(\tau^I(D_\mu\Sigma)\Sigma^\dagger)$, and the matrix $j_\mu = j_\mu^I\tau^I/2$. Unit: GeV^6. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- W_M^I, W_M, W_{MN}, W_{MN}^I: The SU(2) gauge field (geometric normalisation): the components W_M^I ($I = 1,2,3$; W^3 the third component), the matrix $W_M = W_M^I\tau^I/2$, the field strength W_{MN} and its components W_{MN}^I; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- τ^I: The Pauli matrices acting on weak-isospin doublets ($\tau^I/2$ the generators of SU(2); $\tau^3 = \mathrm{diag}(1,-1)$). Unit: dimensionless. Status: STANDARD.
- I: A weak-isospin (adjoint SU(2)) index running over $1,2,3$; a repeated index is summed. Status: STANDARD.
- i: The generation index ($i = 1,2,3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- j: A second generation index ($j = 1,2,3$). Status: STANDARD.
- ℓ_i, $\bar{\ell}_i$: The lepton doublet of the generation i (left-handed neutrino and charged lepton) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- $\mathcal{Y}_\nu$, $\mathcal{Y}_e$, $\mathcal{Y}_u$, $\mathcal{Y}_d$: The complex 3×3 tip Yukawa matrices of the neutrino, charged-lepton, up-quark and down-quark sectors (generation indices i, j); on the static block they become the tip mixing matrices $\mathcal{K}$ (the normalisation of that relation is not fixed: OPEN). Unit: dimensionless. Status: INVENTED.
- ν_R, ν_{Ri}: The right-handed neutrino (a gauge singlet): ν_R the multiplet, ν_{Ri} that of the generation i; its zero modes give Dirac neutrino masses. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- e_R, e_{Ri}: The right-handed charged-lepton singlet: e_R the multiplet, e_{Ri} that of the generation i (e_R, μ_R, τ_R). Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- Q_i, $\bar{Q}_i$: The quark doublet of the generation i (left-handed up and down quark of one generation, three colours) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- u_R, u_{Ri}, $\bar{u}_{Ri}$: The up-type quark singlet: u_R the multiplet, u_{Ri} that of the generation i (u_R, c_R, t_R), $\bar{u}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.

- $d_R,\ d_{Ri},\ \bar{d}_{Ri}$: The down-type quark singlet: d_R the multiplet, d_{Ri} that of the generation i (d_R, s_R, b_R), $\bar{d}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- v_Σ: The scale of the tip sigma field (its kinetic term is $-\frac{v_\Sigma^2}{4}\mathrm{tr}((D_\mu\Sigma)^\dagger D^\mu\Sigma)$); on the static block it fixes the Robin parameter β_W. Unit: $\mathrm{GeV}^{5/2}$. Value: fixed through $\beta_\mathrm{W} = 0.011912$. Status: INVENTED.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- $D_M,\ D_\mu$: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $\mathbf{1}$: The identity matrix (of the size fixed by the context: 16 x 16 on spinors, 3 x 3 on the generations or flavours, 2 x 2 for the sigma field). Unit: dimensionless. Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta_N^M = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{\alpha e}$, δ_{l0}, δ^{I3}); with upper and lower index strings the generalised Kronecker delta $\delta^{M_1\cdots M_n}_{N_1\cdots N_n}$, the determinant of the $n \times n$ matrix of Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- $B_M,\ B_{MN}$: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.

At momentum transfers far below m_W the exchange of $W^\pm$ and Z between zero modes is a current-current interaction with the Fermi constant and the neutral-current couplings:

$$\frac{G_\mathrm{F}}{\sqrt{2}} = \frac{g_2^2}{8m_\mathrm{W}^2}, \qquad \tilde{g}_L^f = T_3^f - Q_\mathrm{em}^f \sin^2\theta_\mathrm{W}, \qquad \tilde{g}_R^f = -Q_\mathrm{em}^f \sin^2\theta_\mathrm{W}, \qquad \tilde{g}_V^f = \tilde{g}_L^f + \tilde{g}_R^f = T_3^f - 2Q_\mathrm{em}^f \sin^2\theta_\mathrm{W} \tag{FE-102}$$

Symbols in (FE-102).

- G_F: The Fermi constant. Unit: GeV^{-2}. Value: 1.1663788×10^{-5} GeV^{-2} (measured). Status: STANDARD.
- g_2: The four-dimensional SU(2) coupling ($g_2^2 = 4\pi\alpha_2$). Unit: dimensionless. Value: $g_2^2 = 0.3781$ at $2M_\mathrm{DM}$. Status: DERIVED.
- m_W: The mass of W^+ and W^-. Unit: GeV. Value: 80.3625 GeV (measured, input). Status: STANDARD.
- $\tilde{g}_L^f,\ \tilde{g}_R^f,\ \tilde{g}_V^f$: The neutral-current couplings of the fermion f to the Z (left-handed, right-handed and vector), $\tilde{g}_L^f = T_3^f - Q_\mathrm{em}^f \sin^2\theta_\mathrm{W}$. Unit: dimensionless. Status: STANDARD.
- f: The label of a fermion species: a Standard-Model multiplet of one generation (its N_c Weyl components share one zero-mode profile; the zero mode of each chirality) or a fermion of a medium (electron, proton, neutron); a sum over f runs over these species. Status: STANDARD.
- L: The chirality label L: the left-handed (doublet) zero mode, or the left-handed projection $P_L = (1 + \Gamma_4)/2$. Status: STANDARD.
- T_3: The third component of the weak isospin ($+\frac{1}{2}$ for the upper and $-\frac{1}{2}$ for the lower member of a doublet, 0 for a singlet); T_3^f that of the fermion f. Unit: dimensionless. Status: STANDARD.
- Q_em: The electric charge in units of the elementary charge, $Q_\mathrm{em} = T_3 + Y$; Q_em^f that of the fermion f. Unit: dimensionless. Status: STANDARD.
- θ_W: The weak mixing angle of the couplings, $\cos\theta_\mathrm{W} = \hat{g}_2/\hat{g}_Z$, $\sin\theta_\mathrm{W} = \hat{g}_1/\hat{g}_Z$; fixed by the measured W and Z masses with $\rho_{\mathrm{EW},0}$. Unit: dimensionless. Value: $\cos^2\theta_\mathrm{W} = 0.776368$. Status: DERIVED.
- R: The chirality label R: the right-handed (singlet) zero mode or field, or the projection $P_R = (1 - \Gamma_4)/2$. Status: STANDARD.
- V: The label V of the vector coupling of a fermion to the Z, $\tilde{g}_V^f = \tilde{g}_L^f + \tilde{g}_R^f$. Status: STANDARD.

7.2 The approximation of step 4

The charges, the couplings and the anomalies are exact. The Z widths are computed at tree level with massless final-state fermions, the measured G_F and m_Z and the $\overline{\mathrm{MS}}$ angle (STANDARD), with the factor $1 + \alpha_3/\pi$ on the quarks for the comparison with the data. The neglected terms, relative to the total width, are the electroweak loop corrections (4.1×10^{-3}: the remaining difference to the measured total), the b-quark mass in the b-quark channel (-1.3×10^{-3}) and the order-α_3^2 QCD term ($+1.3 \times 10^{-3}$) (DERIVED from the outputs).

7.3 The solution of step 4

7.3.1 Charges and couplings

In every generation $Q_{\rm em} = T_3 + Y$ gives the up-type quarks u, c, t the charge 2/3, the down-type quarks d, s, b the charge $-1/3$, the neutrinos 0 and the charged leptons e, μ, τ the charge -1, for both chiralities (DERIVED). The commutator $[T_3, T_\pm] = \pm T_\pm$ with $Y(W) = 0$ shows that W^+**, made of** $W^1 - \mathrm{i}W^2$**, carries the electric charge** $+e$**, and** W^-**, made of** $W^1 + \mathrm{i}W^2$**, carries** $-e$; W^- is the antiparticle of W^+. The photon couples with the charge e times $Q_{\rm em}$ and the Z with its neutral current:

$$e = \frac{g_2\, g_1}{\sqrt{g_2^2 + g_1^2}} \;\; \text{(photon: } e\,Q_{\rm em}\text{)}, \qquad g_{\rm Z}\big(T_3 P_L - Q_{\rm em} \sin^2\theta_{\rm W}\big) \;\; (Z), \qquad g_{\rm Z} = \sqrt{g_2^2 + g_1^2}$$

Symbols in this equation.

- e: The elementary charge (the charge of the positron), $e^2 = 4\pi\alpha_{\rm em}$; W^+ has the charge $+e$ and W^- the charge $-e$. Unit: dimensionless. Value: $e^2 = 4\pi/137.036$ at low energy. Status: STANDARD.
- g_2: The four-dimensional SU(2) coupling ($g_2^2 = 4\pi\alpha_2$). Unit: dimensionless. Value: $g_2^2 = 0.3781$ at $2M_{\rm DM}$. Status: DERIVED.
- g_1: The four-dimensional hypercharge coupling ($g_1^2 = 4\pi\alpha_1$, Standard-Model normalisation without the GUT factor). Unit: dimensionless. Value: $g_1^2 = 0.1380$ at $2M_{\rm DM}$. Status: DERIVED.
- $Q_{\rm em}$: The electric charge in units of the elementary charge, $Q_{\rm em} = T_3 + Y$; $Q_{\rm em}^f$ that of the fermion f. Unit: dimensionless. Status: STANDARD.
- $g_{\rm Z}$: The four-dimensional coupling of the Z boson to the neutral current, $g_{\rm Z} = (g_2^2 + g_1^2)^{1/2}$ (canonically normalised fields); the Z couples to a fermion with $g_{\rm Z}(T_3 P_L - Q_{\rm em}\sin^2\theta_{\rm W})$. Unit: dimensionless. Status: DERIVED.
- T_3: The third component of the weak isospin ($+\frac{1}{2}$ for the upper and $-\frac{1}{2}$ for the lower member of a doublet, 0 for a singlet); T_3^f that of the fermion f. Unit: dimensionless. Status: STANDARD.
- P_L, P_R: The projectors on the four-dimensional chiralities, $P_L = (1+\Gamma_4)/2$ (left-handed) and $P_R = (1-\Gamma_4)/2$ (right-handed). Unit: dimensionless. Status: STANDARD.
- L: The chirality label L: the left-handed (doublet) zero mode, or the left-handed projection $P_L = (1+\Gamma_4)/2$. Status: STANDARD.
- $\theta_{\rm W}$: The weak mixing angle of the couplings, $\cos\theta_{\rm W} = \hat{g}_2/\hat{g}_Z$, $\sin\theta_{\rm W} = \hat{g}_1/\hat{g}_Z$; fixed by the measured W and Z masses with $\rho_{\rm EW,0}$. Unit: dimensionless. Value: $\cos^2\theta_{\rm W} = 0.776368$. Status: DERIVED.

(DERIVED).

7.3.2 Anomaly cancellation in three generations

With every zero mode written as a left-handed Weyl field (with $-Y$ for a right-handed one), $N_{\rm col}$ colour and $N_{\rm dbl}$ weak components, one generation gives (DERIVED, exact fractions)

$$\mathcal{A}_{\mathrm{SU(3)^2U(1)}} = \sum_{\text{triplets}} N_{\rm dbl}\, Y = 0, \qquad \mathcal{A}_{\mathrm{SU(2)^2U(1)}} = \sum_{\text{doublets}} N_{\rm col}\, Y = \tfrac{1}{2} - \tfrac{1}{2} = 0,$$
$$\mathcal{A}_{\mathrm{U(1)^3}} = \sum N_{\rm col} N_{\rm dbl}\, Y^3 = \tfrac{1}{36} - \tfrac{8}{9} + \tfrac{1}{9} - \tfrac{1}{4} + 1 = 0$$

Symbols in this equation.

- $\mathcal{A}_{\mathrm{SU(3)^2U(1)}}$, $\mathcal{A}_{\mathrm{SU(2)^2U(1)}}$, $\mathcal{A}_{\mathrm{U(1)^3}}$: The gauge anomaly coefficients of one generation, with every zero mode written as a left-handed Weyl field (a right-handed one with the hypercharge $-Y$): the sum over the colour triplets of $N_{\rm dbl}Y$ (mixed $SU(3)^2U(1)_Y$ anomaly), the sum over the weak doublets of $N_{\rm col}Y$ (mixed $SU(2)^2U(1)_Y$ anomaly) and the sum over all components of $N_{\rm col}N_{\rm dbl}Y^3$ (the $U(1)_Y^3$ anomaly); the theory is consistent only if they vanish. Unit: dimensionless. Value: 0 for each generation of UM-2. Status: DERIVED.
- $N_{\rm dbl}$: The number of weak-isospin components of a fermion multiplet: 2 for an $SU(2)$ doublet, 1 for a singlet. Unit: dimensionless. Status: STANDARD.
- Y: The hypercharge (the generator of $\mathrm{U}(1)_Y$); Y_ψ the hypercharge of the multiplet ψ (Q: $\frac{1}{6}$, u_R: $\frac{2}{3}$, d_R: $-\frac{1}{3}$, ℓ: $-\frac{1}{2}$, e_R: -1, ν_R: 0). Unit: dimensionless. Status: STANDARD.

- N_{col}: The number of colour components of a fermion multiplet: 3 for a quark, 1 for a lepton. Unit: dimensionless. Status: STANDARD.

and the mixed gravitational anomaly $\sum N_{\mathrm{col}} N_{\mathrm{dbl}}\, Y = 1-2+1-1+1 = 0$ and the $SU(3)^3$ anomaly $2-1-1=0$ vanish too; three generations have twelve $SU(2)$ doublets (no Witten anomaly). Three lepton generations with a single quark generation would leave $\mathcal{A}_{\mathrm{SU(2)^2U(1)}} = -1$ and $\mathcal{A}_{\mathrm{U(1)^3}} = 3/2$.

7.3.3 The two halves of dirac16complex

$\mathcal{B}$ commutes with the four-dimensional Lorentz generators and with C, so $\Psi = \Psi_A + \Psi_B$ ($\mathcal{B} = -1, +1$) splits into Lorentz-invariant halves and $\bar{\Psi}\Psi = \bar{\Psi}_A\Psi_A + \bar{\Psi}_B\Psi_B$. The identity $\Gamma_4 = -\mathcal{B}\gamma^{x_0}$ (DERIVED, 16×16) gives the four-dimensional chiral projector $P_L = (1+\Gamma_4)/2 = (1-\mathcal{B}\gamma^{x_0})/2$, which equals $(1+\gamma^{x_0})/2$ on Ψ_A and $(1-\gamma^{x_0})/2$ on Ψ_B; the latter annihilates every orbital at the tip. The chiral charge densities at the tip, which a chiral tip current would couple to, are (DERIVED): the Ψ_A fold band 0.163 per unit length in each of its four blocks, the Ψ_B fold band 0; the odd (heavy) states 0.257 (Ψ_A) and 0 (Ψ_B); every transition current $\bar{\Psi}_A\gamma^\mu P_L\Psi_B$ vanishes at the tip.

7.3.4 The Z invisible width and the tip Z portal of version 1

Version 1 coupled the Ψ_A fold band (two Weyl species) to the Z through a chiral tip portal of the current $\mathrm{i}\bar{\Psi}\gamma^\mu P_L\Psi$ to the tip current $\frac{\mathrm{i}}{2}v_\Sigma^2\,\mathrm{tr}(\Sigma^\dagger D_\mu\Sigma\,\tau^3)$, with an even and an oriented (odd under the fold reflection) coefficient. With three neutrinos and a fold-band coupling r_{Z} times that of a neutrino, the number of invisible species (the invisible width of the Z in units of its width into one neutrino flavour) would be

$$N_\nu = 3 + 2r_{\mathrm{Z}}^2;$$

Symbols in this equation.

- N_ν: The number of light neutrino species seen by the Z: its invisible width in units of its width into one neutrino flavour. Unit: dimensionless. Value: measured 2.9963 ± 0.0074 (PDG 2025); 3 in UM-2. Status: STANDARD.
- r_{Z}: The coupling of the fold band of Ψ_A to the Z in units of the coupling of one neutrino flavour, in the chiral tip Z portal of version 1 (removed in UM-2, where it is zero). Unit: dimensionless. Value: bounded by $r_{\mathrm{Z}}^2 < 0.0056$ (two standard deviations). Status: DERIVED.

the measured $N_\nu = 2.9963 \pm 0.0074$ (PDG 2025) is reproduced by $r_{\mathrm{Z}} = 0$ (0.5 standard deviations) and bounds $r_{\mathrm{Z}}^2 < 0.0056$ (two standard deviations). Even at that bound the neutral-current rate of the fold band, r_{Z}^2 times the neutrino's ($\propto r_{\mathrm{Z}}^2 G_{\mathrm{F}}^2 T^5$ against the Hubble rate $\mathcal{H} \propto T^2/M_{\mathrm{Pl}}$), would keep it in equilibrium down to $T \approx 2.6\ \mathrm{MeV} \times r_{\mathrm{Z}}^{-2/3} = 14.7\ \mathrm{MeV}$, where $g_s = 10.75$, so it would carry the neutrino temperature and add the extra radiation of an early-decoupled species,

$$\Delta N_{\mathrm{eff}} = N_{\mathrm{W}}\Big(\frac{10.75}{g_s}\Big)^{4/3} F_{\mathrm{h}}, \qquad F_{\mathrm{h}} = \Big[\frac{T_\nu}{T_\gamma}\Big(\frac{11}{4}\Big)^{1/3}\Big/\big(1+h_\nu\big)\Big]^4 = 1.0095 \tag{FE-133}$$

Symbols in (FE-133).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- N_{eff}: The effective number of neutrino species (the radiation beyond the photons in units of one neutrino species with $T_\nu/T_\gamma = (4/11)^{1/3}$); ΔN_{eff} an addition to it. Unit: dimensionless. Value: 3.0432 (neutrinos) + 0.0010 (fold bands). Status: STANDARD.
- N_{W}: The number of Weyl species that decouple early. Unit: dimensionless. Status: DERIVED.
- g_ρ, g_s: The effective numbers of relativistic degrees of freedom of the energy density and of the entropy density of the plasma. Unit: dimensionless. Value: 105.75 when every species of UM-2 in equilibrium is relativistic (no Higgs particle), 10.75 near 1 MeV, $g_s = 43/11$ today. Status: STANDARD.

- F_{h}: The correction factor of the early-decoupled radiation for the heating of the neutrinos. Unit: dimensionless. Value: 1.0095. Status: DERIVED.
- T_ν: The temperature of the relic neutrinos (one common temperature). Unit: GeV. Value: 1.9454 K today. Status: STANDARD.
- T_γ: The temperature of the photons (equal to the plasma temperature T; the subscript is written where the neutrinos have their own temperature). Unit: GeV. Value: 2.7255 K today. Status: STANDARD.
- h_ν: The relative heating of the neutrinos by the electron-positron annihilation (beyond instantaneous decoupling). Unit: dimensionless. Value: 0.00165. Status: DERIVED.

with $N_{\mathrm{W}} = 2$ Weyl species (F_{h} corrects for the non-instantaneous neutrino decoupling of step 5: a species decoupled earlier is not heated by e^+e^- annihilation): $\Delta N_{\mathrm{eff}} = 2.02$, far above the measured $N_{\mathrm{eff}} = 2.99 \pm 0.17$ (Planck 2018) and 2.86 ± 0.13 (ACT DR6) (STANDARD). The oriented part would let the heavy odd Ψ_A states decay into the fold band and a Z, and through the inverse decays and the exchange process keep the fold band in equilibrium until the odd states disappear, at $T \sim M_{\mathrm{DM}}/30$ (above the electroweak scale, $g_s = 105.75$: the Standard Model without a Higgs particle), or, for an oriented coupling of the size of a neutrino's, until the Z bosons annihilate near 15 GeV ($g_s = 86.25$: t, W, Z gone) (ESTIMATE): then $\Delta N_{\mathrm{eff}} = 0.0958$ to 0.1257 and $N_{\mathrm{eff}} = 3.139$ to 3.169, 2.1 to 2.4 standard deviations above ACT DR6 (the three-flavour value 3.0432: 1.4; disfavoured). UM-2 therefore drops the portal (ASSUMED; the bounds DERIVED): $N_\nu = 3$, neither fold band is ever thermalised, and both halves keep an exact fold parity. (The plasma counts used are $g_s = 10.75$: photons, $e^\pm$, three neutrinos; 61.75 just above the QCD crossover: with gluons, u, d, s, μ; 86.25 below the top, W and Z masses; 105.75 above the electroweak scale. Right-handed neutrinos are never thermalised.)

7.3.5 The Z widths of three generations

At tree level, with massless final-state fermions, the partial width of the Z into the pair of a fermion f is

$$\Gamma_{\mathrm{Z}}^{f} = \frac{N_{\mathrm{col}}\, G_{\mathrm{F}}\, m_{\mathrm{Z}}^3}{6\sqrt{2}\,\pi}\left[\left(T_3^f - 2Q_{\mathrm{em}}^f \sin^2\theta_{\overline{\mathrm{MS}}}\right)^2 + \left(T_3^f\right)^2\right]$$

Symbols in this equation.

- Γ_{Z}^{f}: The decay width of the Z boson; with the fermion label f it is the partial width into the pair of f and its antiparticle (tree level, massless fermion). Unit: GeV (quoted in MeV). Value: total 2485.2 MeV with $1 + \alpha_3/\pi$ on the quarks (measured 2495.5 MeV). Status: DERIVED.
- f: The label of a fermion species: a Standard-Model multiplet of one generation (its N_{c} Weyl components share one zero-mode profile; the zero mode of each chirality) or a fermion of a medium (electron, proton, neutron); a sum over f runs over these species. Status: STANDARD.
- N_{col}: The number of colour components of a fermion multiplet: 3 for a quark, 1 for a lepton. Unit: dimensionless. Status: STANDARD.
- G_{F}: The Fermi constant. Unit: GeV^{-2}. Value: 1.1663788×10^{-5} GeV^{-2} (measured). Status: STANDARD.
- m_{Z}: The mass of the Z. Unit: GeV. Value: 91.1879 GeV (reproduced). Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- T_3: The third component of the weak isospin ($+\frac{1}{2}$ for the upper and $-\frac{1}{2}$ for the lower member of a doublet, 0 for a singlet); T_3^f that of the fermion f. Unit: dimensionless. Status: STANDARD.
- Q_{em}: The electric charge in units of the elementary charge, $Q_{\mathrm{em}} = T_3 + Y$; Q_{em}^f that of the fermion f. Unit: dimensionless. Status: STANDARD.
- $\theta_{\overline{\mathrm{MS}}}$: The weak mixing angle in the $\overline{\mathrm{MS}}$ scheme at m_{Z}. Unit: dimensionless. Value: $\sin^2\theta_{\overline{\mathrm{MS}}} = 0.23122$. Status: STANDARD.

with $\sin^2\theta_{\overline{\mathrm{MS}}} = 0.23122$ (G_{F}, m_{Z} and the angle STANDARD; DERIVED), in MeV:

channel	each ν	invisible (3ν)	each of e, μ, τ	each of u, c	each of d, s, b	hadronic	total
tree level	165.9	497.7	83.4	285.4	367.9	1674.4	2422.3

channel	each ν	invisible (3ν)	each of e, μ, τ	each of u, c	each of d, s, b	hadronic	total
with $1+\alpha_3/\pi$ on the quarks						1737.3	2485.2
measured (PDG 2025)		499.1					2495.5

The top quark is too heavy for the Z to decay into it; $\alpha_3(m_\mathrm{Z}) = 0.1180$ (STANDARD). The ratio of the hadronic to one charged-lepton width is 20.83 against the measured 20.77 (STANDARD, PDG 2025). The remaining difference (0.4 % in the total) is the size of the electroweak loop corrections, which UM-2 does not compute; with the tree-level angle of the couplings of the mass fit, $\sin^2\theta_\mathrm{W} = 0.22363$, the total changes by +0.82 % (a per-cent scheme ambiguity). The one-generation conflict of version 1 (a total of 1232 MeV) is gone.

7.4 The check of step 4

In its own equations. The charges, the couplings of the photon and the Z, the anomaly sums and the projector algebra are exact (DERIVED, fractions and 16×16 matrices). The stored widths (each channel, the invisible, hadronic and total widths and the ratio 20.83) agree with the formula recomputed from the stored inputs to 2.7×10^{-12} (DERIVED).

In concert with the other field equations.

- *The W and Z masses of the x_0 problem.* The Z mass of the widths is the Z mass of the x_0 eigenvalues of step 2 to 2.0×10^{-12}; the angle of the couplings of the x_0 problem and the $\overline{\mathrm{MS}}$ angle of the widths differ by 0.0076, which moves the total width by 0.82 % at tree level (0.83 % with the QCD factor), within the per-cent size of the uncomputed loop corrections (DERIVED).
- *The heating factor of the neutrino decoupling.* The factor $F_\mathrm{h} = 1.009500589$ of the neutrino decoupling of step 5 is the one used here and in step 9, to 5.8×10^{-7} (the rounded 1.0095) (DERIVED).
- *Three neutrinos everywhere.* $N_\nu = 3$ of the Z width is the content of the neutrino decoupling and of N_eff (step 5) and of the CMB computation (step 6), in which the fold bands enter only through the freeze-in radiation of step 9.

7.5 The status of step 4

DERIVED: the charges and couplings, the anomaly cancellation, the projector identities and tip densities, the bounds on the removed tip Z portal, the tree-level Z widths. ASSUMED: the removal of the tip Z portal. INVENTED: the tip sigma field. STANDARD: G_F, m_Z, $\alpha_3(m_\mathrm{Z})$, the $\overline{\mathrm{MS}}$ angle and the measured widths. OPEN: the electroweak loop corrections and the S parameter of the Higgsless tip. Conflicts with data, inherent in the author's order: no Higgs boson, and the longitudinal W^+W^- scattering violates perturbative unitarity near 1.2 TeV (see the last section).

8 Step 5: the flavour conversion of the three neutrinos, N_eff and the CNB

The three neutrino masses are x_0 eigenvalues of step 2 and the PMNS matrix is the neutrino tip matrix. Step 5 solves the flavour evolution of the three neutrinos in vacuum, in matter (the MSW effect), in the Sun, at accelerators, in the early Universe (where it accompanies neutrino decoupling and fixes N_eff), in the cosmic neutrino background, and in supernovae (the MSW resonances of the envelope and the fast flavour instability of the dense neutrino gas), and compares the Dirac neutrinos of UM-2 with the Majorana alternative.

8.1 The field equations of step 5

The masses and the mixing matrix (step 2):

$$m_1 = 1.000\ \text{meV}, \qquad m_2 = \sqrt{m_1^2 + \Delta m_{21}^2} = 8.712\ \text{meV}, \qquad m_3 = \sqrt{m_1^2 + \Delta m_{31}^2} = 50.140\ \text{meV} \qquad \text{(FE-100)}$$

Symbols in (FE-100).

- $m_1,\ m_2,\ m_3$: The masses of the neutrino mass states ν_1, ν_2, ν_3 (normal ordering). Unit: GeV (quoted in meV). Value: 1.000, 8.712, 50.140 meV (m_1 ASSUMED). Status: DERIVED.
- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- $\Delta m_{21}^2,\ \Delta m_{31}^2,\ \Delta m_{32}^2$: The differences of the squared neutrino masses, $\Delta m_{ij}^2 = m_i^2 - m_j^2$. Unit: GeV^2 (quoted in eV^2). Value: $\Delta m_{21}^2 = 7.49 \times 10^{-5}\ \text{eV}^2$, $\Delta m_{31}^2 = 2.513 \times 10^{-3}\ \text{eV}^2$ (NuFIT 6.0). Status: STANDARD.

$$U = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}\mathrm{e}^{-\mathrm{i}\delta_{\text{CP}}} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}\mathrm{e}^{\mathrm{i}\delta_{\text{CP}}} & c_{12}c_{23} - s_{12}s_{23}s_{13}\mathrm{e}^{\mathrm{i}\delta_{\text{CP}}} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}\mathrm{e}^{\mathrm{i}\delta_{\text{CP}}} & -c_{12}s_{23} - s_{12}c_{23}s_{13}\mathrm{e}^{\mathrm{i}\delta_{\text{CP}}} & c_{23}c_{13} \end{pmatrix}, \qquad J_{\text{CP}} = c_{12}s_{12}c_{23}s_{23}c_{13}^2 s_{13}\sin\delta_{\text{CP}}$$

(FE-61)

Symbols in (FE-61).

- U: The PMNS matrix, the unitary 3×3 mixing matrix of the leptons, $\nu_\alpha = \sum_i U_{\alpha i}\nu_i$ (rows $\alpha = e, \mu, \tau$, columns the mass states i); in UM-2 $U = \mathcal{K}_e^\dagger \mathcal{K}_\nu$. Unit: dimensionless. Value: $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$, $\delta_{\text{CP}} = 212^\circ$ (NuFIT 6.0). Status: STANDARD.
- $c_{12},\ c_{13},\ c_{23}$: The cosines of the mixing angles, $c_{ij} = \cos\theta_{ij}$. Unit: dimensionless. Status: STANDARD.
- $s_{12},\ s_{13},\ s_{23}$: The sines of the mixing angles, $s_{ij} = \sin\theta_{ij}$. Unit: dimensionless. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- δ_{CP}: The CP-violating phase of a mixing matrix. Unit: dimensionless. Value: 212° (PMNS), 1.147 rad (CKM). Status: STANDARD.
- J_{CP}: The Jarlskog invariant of a mixing matrix (the measure of CP violation). Unit: dimensionless. Value: -0.01777 (PMNS), 3.117×10^{-5} (CKM). Status: STANDARD.

The interactions at low energy, the matter potentials and the flavour evolution:

$$\mathcal{L}_{\text{CC}} = -2\sqrt{2}\,G_{\text{F}}\big(\bar\nu_\alpha\gamma^\mu P_L\,\ell_\alpha^-\big)\big(\bar\ell_\beta^-\gamma_\mu P_L\,\nu_\beta\big), \qquad \mathcal{L}_{\text{NC}} = -2\sqrt{2}\,G_{\text{F}}\big(\bar\nu_\alpha\gamma^\mu P_L\,\nu_\alpha\big)\sum_f \bar\psi_f\gamma_\mu\big(\tilde g_L^f P_L + \tilde g_R^f P_R\big)\psi_f,$$

$$\nu_\alpha = \sum_{i=1}^{3} U_{\alpha i}\,\nu_i, \qquad P_L = \frac{1+\Gamma_4}{2}, \qquad P_R = \frac{1-\Gamma_4}{2}$$

(FE-101)

Symbols in (FE-101).

- $\mathcal{L}_{\text{CC}},\ \mathcal{L}_{\text{NC}}$: The charged-current and neutral-current four-fermion Lagrangians (exchange of $W^\pm$ and of Z far below their masses). Unit: GeV^4. Status: STANDARD.
- G_{F}: The Fermi constant. Unit: GeV^{-2}. Value: $1.1663788 \times 10^{-5}\ \text{GeV}^{-2}$ (measured). Status: STANDARD.
- $\bar\nu_\alpha$: The antineutrino of the flavour α; in a Lagrangian the Dirac adjoint of the neutrino field (it annihilates antineutrinos). Unit: $\text{GeV}^{3/2}$. Status: STANDARD.
- α: A flavour index running over e, μ, τ (the flavour states of the leptons and neutrinos). Status: STANDARD.
- $\gamma^a,\ \gamma^M,\ \gamma^\mu$: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.

- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- P_L, P_R: The projectors on the four-dimensional chiralities, $P_L = (1+\Gamma_4)/2$ (left-handed) and $P_R = (1-\Gamma_4)/2$ (right-handed). Unit: dimensionless. Status: STANDARD.
- L: The chirality label L: the left-handed (doublet) zero mode, or the left-handed projection $P_L = (1+\Gamma_4)/2$. Status: STANDARD.
- ℓ^-_α, $\bar{\ell}^-_\alpha$: The charged lepton of the flavour α (e^-, μ^-, τ^-; the lower member of the lepton doublet) at low energy, and its Dirac adjoint. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- β: A second flavour index running over e, μ, τ. Status: STANDARD.
- ν_α, ν_i: A left-handed neutrino: ν_α of the flavour $\alpha = e, \mu, \tau$ (ν_e, ν_μ, ν_τ), ν_i of the mass state i (mass m_i). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- f: The label of a fermion species: a Standard-Model multiplet of one generation (its N_c Weyl components share one zero-mode profile; the zero mode of each chirality) or a fermion of a medium (electron, proton, neutron); a sum over f runs over these species. Status: STANDARD.
- ψ_i, $\bar{\psi}_i$: A Standard-Model fermion field of eight dimensions: ψ_i is the multiplet $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ of the generation i, ψ_f one fermion component f (in four-dimensional formulas its zero mode); $\bar{\psi} = \psi^\dagger C$ the Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$ (eight dimensions), $\mathrm{GeV}^{3/2}$ (four dimensions). Status: ASSUMED.
- $\tilde{g}^f_L$, $\tilde{g}^f_R$, $\tilde{g}^f_V$: The neutral-current couplings of the fermion f to the Z (left-handed, right-handed and vector), $\tilde{g}^f_L = T^f_3 - Q^f_\mathrm{em} \sin^2\theta_\mathrm{W}$. Unit: dimensionless. Status: STANDARD.
- R: The chirality label R: the right-handed (singlet) zero mode or field, or the projection $P_R = (1-\Gamma_4)/2$. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- U: The PMNS matrix, the unitary 3×3 mixing matrix of the leptons, $\nu_\alpha = \sum_i U_{\alpha i}\nu_i$ (rows $\alpha = e, \mu, \tau$, columns the mass states i); in UM-2 $U = \mathcal{K}_e^\dagger \mathcal{K}_\nu$. Unit: dimensionless. Value: $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$, $\delta_\mathrm{CP} = 212^\circ$ (NuFIT 6.0). Status: STANDARD.
- Γ_4: The four-dimensional chirality matrix, $\Gamma_4 = \mathrm{i}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}\gamma^{x_4}$; a doublet zero mode has $\Gamma_4 = +1$ (left-handed), a singlet zero mode $\Gamma_4 = -1$ (right-handed). Unit: dimensionless. Status: DERIVED.

$$\frac{G_\mathrm{F}}{\sqrt{2}} = \frac{g_2^2}{8m_\mathrm{W}^2}, \qquad \tilde{g}^f_L = T^f_3 - Q^f_\mathrm{em}\sin^2\theta_\mathrm{W}, \qquad \tilde{g}^f_R = -Q^f_\mathrm{em}\sin^2\theta_\mathrm{W}, \qquad \tilde{g}^f_V = \tilde{g}^f_L + \tilde{g}^f_R = T^f_3 - 2Q^f_\mathrm{em}\sin^2\theta_\mathrm{W} \tag{FE-102}$$

Symbols in (FE-102).

- G_F: The Fermi constant. Unit: GeV^{-2}. Value: 1.1663788×10^{-5} GeV^{-2} (measured). Status: STANDARD.
- g_2: The four-dimensional SU(2) coupling ($g_2^2 = 4\pi\alpha_2$). Unit: dimensionless. Value: $g_2^2 = 0.3781$ at $2M_\mathrm{DM}$. Status: DERIVED.
- m_W: The mass of W^+ and W^-. Unit: GeV. Value: 80.3625 GeV (measured, input). Status: STANDARD.
- $\tilde{g}^f_L$, $\tilde{g}^f_R$, $\tilde{g}^f_V$: The neutral-current couplings of the fermion f to the Z (left-handed, right-handed and vector), $\tilde{g}^f_L = T^f_3 - Q^f_\mathrm{em}\sin^2\theta_\mathrm{W}$. Unit: dimensionless. Status: STANDARD.
- f: The label of a fermion species: a Standard-Model multiplet of one generation (its N_c Weyl components share one zero-mode profile; the zero mode of each chirality) or a fermion of a medium (electron, proton, neutron); a sum over f runs over these species. Status: STANDARD.
- L: The chirality label L: the left-handed (doublet) zero mode, or the left-handed projection $P_L = (1+\Gamma_4)/2$. Status: STANDARD.
- T_3: The third component of the weak isospin ($+\frac{1}{2}$ for the upper and $-\frac{1}{2}$ for the lower member of a doublet, 0 for a singlet); T^f_3 that of the fermion f. Unit: dimensionless. Status: STANDARD.
- Q_em: The electric charge in units of the elementary charge, $Q_\mathrm{em} = T_3 + Y$; Q^f_em that of the fermion f. Unit: dimensionless. Status: STANDARD.
- θ_W: The weak mixing angle of the couplings, $\cos\theta_\mathrm{W} = \hat{g}_2/\hat{g}_Z$, $\sin\theta_\mathrm{W} = \hat{g}_1/\hat{g}_Z$; fixed by the measured W and Z masses with $\rho_{\mathrm{EW},0}$. Unit: dimensionless. Value: $\cos^2\theta_\mathrm{W} = 0.776368$. Status: DERIVED.
- R: The chirality label R: the right-handed (singlet) zero mode or field, or the projection $P_R = (1-\Gamma_4)/2$. Status: STANDARD.
- V: The label V of the vector coupling of a fermion to the Z, $\tilde{g}^f_V = \tilde{g}^f_L + \tilde{g}^f_R$. Status: STANDARD.

$$V_\alpha = \sqrt{2}\,G_\mathrm{F}\Big[\delta_{\alpha e}\big(n_{e^-} - n_{e^+}\big) + \sum_f \tilde{g}_V^f \big(n_f - n_{\bar{f}}\big)\Big],$$
$$V = \mathrm{diag}\big(V_\mathrm{CC},\ 0,\ 0\big) + V_\mathrm{NC}\,\mathbf{1}, \qquad V_\mathrm{CC} = \sqrt{2}\,G_\mathrm{F}\,n_\mathrm{e}, \qquad V_\mathrm{NC} = -\frac{G_\mathrm{F}\,n_\mathrm{n}}{\sqrt{2}} \tag{FE-103}$$

Symbols in (FE-103).

- V_α, V: The matter potential of forward scattering: V_α of the flavour α (V_e, V_μ), V the 3×3 matrix of the potentials in the flavour basis; antineutrinos have $-V$. Unit: GeV. Status: STANDARD.
- α: A flavour index running over e, μ, τ (the flavour states of the leptons and neutrinos). Status: STANDARD.
- G_F: The Fermi constant. Unit: GeV^{-2}. Value: 1.1663788×10^{-5} GeV^{-2} (measured). Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta_N^M = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{\alpha e}$, δ_{l0}, δ^{I3}); with upper and lower index strings the generalised Kronecker delta $\delta_{N_1 \cdots N_n}^{M_1 \cdots M_n}$, the determinant of the $n \times n$ matrix of Kronecker deltas $\delta_{N_b}^{M_a}$ (+1 or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- e: The electron flavour as a value of a flavour index ($\delta_{\alpha e}$, U_{e2}, V_e, $\mathcal{P}_{ee}$). Status: STANDARD.
- n_{e^-}, n_{e^+}: The number densities of electrons and of positrons. Unit: GeV^3. Status: STANDARD.
- f: The label of a fermion species: a Standard-Model multiplet of one generation (its N_c Weyl components share one zero-mode profile; the zero mode of each chirality) or a fermion of a medium (electron, proton, neutron); a sum over f runs over these species. Status: STANDARD.
- $\tilde{g}_L^f$, $\tilde{g}_R^f$, $\tilde{g}_V^f$: The neutral-current couplings of the fermion f to the Z (left-handed, right-handed and vector), $\tilde{g}_L^f = T_3^f - Q_\mathrm{em}^f \sin^2\theta_\mathrm{W}$. Unit: dimensionless. Status: STANDARD.
- V: The label V of the vector coupling of a fermion to the Z, $\tilde{g}_V^f = \tilde{g}_L^f + \tilde{g}_R^f$. Status: STANDARD.
- n, n_o, $n_{\psi i}$, n_f, n_i: A number density: $n(y)$ the particle density of dirac16complex in the Kohn-Sham equations and n_o the contribution of the orbital o, $n_{\psi i}(y)$ the density of the Standard-Model fermion ψ_i (per proper 7-volume of eight dimensions); n_f the density of the fermion component f and n_i that of the neutrino mass state i (per 3-volume of four dimensions). Unit: GeV^7 (eight dimensions), GeV^3 (four dimensions). Status: STANDARD.
- $n_{\bar{f}}$: The number density of the antiparticles of the fermion f. Unit: GeV^3. Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ when it is a tensor in eight dimensions). Status: STANDARD.
- V_CC, V_NC: The charged-current potential of electron neutrinos and the neutral-current potential common to all flavours. Unit: GeV. Status: STANDARD.
- $\mathbf{1}$: The identity matrix (of the size fixed by the context: 16 x 16 on spinors, 3 x 3 on the generations or flavours, 2 x 2 for the sigma field). Unit: dimensionless. Status: STANDARD.
- n_e: The number density of electrons (net of positrons); in the Thomson terms of the photon and baryon equations only free (ionised) electrons count. Unit: GeV^3. Status: STANDARD.
- n_n: The number density of neutrons (free and bound). Unit: GeV^3. Status: STANDARD.

$$\mathrm{i}\,\frac{\mathrm{d}\boldsymbol{\psi}}{\mathrm{d}x_4} = H_\mathrm{fl}\,\boldsymbol{\psi}, \qquad H_\mathrm{fl} = \frac{1}{2E}\,U\,\mathrm{diag}\big(m_1^2,\ m_2^2,\ m_3^2\big)\,U^\dagger + V, \qquad \mathcal{U}(L_\mathrm{b}) = \mathrm{e}^{-\mathrm{i}H_\mathrm{fl}L_\mathrm{b}}, \qquad \mathcal{P}(\nu_\alpha \to \nu_\beta) = |\mathcal{U}_{\beta\alpha}|^2 \tag{FE-104}$$

Symbols in (FE-104).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $\boldsymbol{\psi}$: The vector of the three flavour amplitudes of a neutrino. Unit: dimensionless. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- H_fl: The flavour Hamiltonian of a neutrino of energy E (vacuum mass term plus matter potential). Unit: GeV. Status: STANDARD.
- E: The energy of a particle: of the neutrino in the flavour equations, $E = (k^2 + m^2)^{1/2}$ of a particle of momentum k in the thermal integrals. Unit: GeV. Status: STANDARD.

- U: The PMNS matrix, the unitary 3×3 mixing matrix of the leptons, $\nu_\alpha = \sum_i U_{\alpha i}\nu_i$ (rows $\alpha = e, \mu, \tau$, columns the mass states i); in UM-2 $U = \mathcal{K}_e^\dagger \mathcal{K}_\nu$. Unit: dimensionless. Value: $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$, $\delta_{\rm CP} = 212^\circ$ (NuFIT 6.0). Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ when it is a tensor in eight dimensions). Status: STANDARD.
- m_1, m_2, m_3: The masses of the neutrino mass states ν_1, ν_2, ν_3 (normal ordering). Unit: GeV (quoted in meV). Value: 1.000, 8.712, 50.140 meV (m_1 ASSUMED). Status: DERIVED.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- V_α, V: The matter potential of forward scattering: V_α of the flavour α (V_e, V_μ), V the 3×3 matrix of the potentials in the flavour basis; antineutrinos have $-V$. Unit: GeV. Status: STANDARD.
- $\mathcal{U}$: The flavour evolution operator over the baseline $L_{\rm b}$; $\mathcal{U}_{\beta\alpha}$ the amplitude of $\nu_\alpha \to \nu_\beta$. Unit: dimensionless. Status: STANDARD.
- $L_{\rm b}$: The baseline: the distance travelled by the neutrino. Unit: length. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- $\mathcal{P}$: The probability of a flavour transition, $\mathcal{P}(\nu_\alpha \to \nu_\beta)$; $\mathcal{P}_{ee}$ the survival probability of electron neutrinos. Unit: dimensionless. Status: STANDARD.
- ν_α, ν_i: A left-handed neutrino: ν_α of the flavour $\alpha = e, \mu, \tau$ (ν_e, ν_μ, ν_τ), ν_i of the mass state i (mass m_i). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- α: A flavour index running over e, μ, τ (the flavour states of the leptons and neutrinos). Status: STANDARD.
- β: A second flavour index running over e, μ, τ. Status: STANDARD.

$$\mathcal{P}(\nu_\alpha \to \nu_\beta) = \Big| \sum_{i=1}^{3} U_{\beta i} U^*_{\alpha i}\, \mathrm{e}^{-\mathrm{i} m_i^2 L_{\rm b}/(2E)} \Big|^2, \qquad \mathcal{P}(\nu_\alpha \to \nu_\beta) = \mathcal{P}(\bar\nu_\beta \to \bar\nu_\alpha),$$

$$\mathcal{P}(\nu_\mu \to \nu_e) - \mathcal{P}(\nu_e \to \nu_\mu) = -16 J_{\rm CP} \sin\Delta_{21} \sin\Delta_{31} \sin\Delta_{32}, \qquad \Delta_{ij} = \frac{\Delta m^2_{ij} L_{\rm b}}{4E}, \qquad J_{\rm CP} = \mathrm{Im}\big(U_{e2} U_{\mu 3} U^*_{e3} U^*_{\mu 2}\big) \tag{FE-105}$$

Symbols in (FE-105).

- $\mathcal{P}$: The probability of a flavour transition, $\mathcal{P}(\nu_\alpha \to \nu_\beta)$; $\mathcal{P}_{ee}$ the survival probability of electron neutrinos. Unit: dimensionless. Status: STANDARD.
- ν_α, ν_i: A left-handed neutrino: ν_α of the flavour $\alpha = e, \mu, \tau$ (ν_e, ν_μ, ν_τ), ν_i of the mass state i (mass m_i). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- α: A flavour index running over e, μ, τ (the flavour states of the leptons and neutrinos). Status: STANDARD.
- β: A second flavour index running over e, μ, τ. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- U: The PMNS matrix, the unitary 3×3 mixing matrix of the leptons, $\nu_\alpha = \sum_i U_{\alpha i}\nu_i$ (rows $\alpha = e, \mu, \tau$, columns the mass states i); in UM-2 $U = \mathcal{K}_e^\dagger \mathcal{K}_\nu$. Unit: dimensionless. Value: $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$, $\delta_{\rm CP} = 212^\circ$ (NuFIT 6.0). Status: STANDARD.
- *: Complex conjugation of a number or of each entry of a matrix. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- m, m_s, m_i: A four-dimensional mass: m the mass of a mode (the eigenvalue of an x_0 problem) or of a fermion; m_s the mass of the species s; m_i the mass of the neutrino mass state i. Unit: GeV. Status: DERIVED.
- $L_{\rm b}$: The baseline: the distance travelled by the neutrino. Unit: length. Status: STANDARD.
- E: The energy of a particle: of the neutrino in the flavour equations, $E = (k^2 + m^2)^{1/2}$ of a particle of momentum k in the thermal integrals. Unit: GeV. Status: STANDARD.
- $\bar\nu_\alpha$: The antineutrino of the flavour α; in a Lagrangian the Dirac adjoint of the neutrino field (it annihilates antineutrinos). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- e: The electron flavour as a value of a flavour index ($\delta_{\alpha e}$, U_{e2}, V_e, $\mathcal{P}_{ee}$). Status: STANDARD.
- $J_{\rm CP}$: The Jarlskog invariant of a mixing matrix (the measure of CP violation). Unit: dimensionless. Value:

-0.01777 (PMNS), 3.117×10^{-5} (CKM). Status: STANDARD.

- Δ_{ij}: The oscillation phase of the pair of mass states (i, j), $\Delta_{ij} = \Delta m^2_{ij} L_{\mathrm{b}}/(4E)$. Unit: dimensionless. Status: STANDARD.
- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m^2_{21} = m^2_2 - m^2_1$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- j: A second generation index ($j = 1, 2, 3$). Status: STANDARD.
- Im: The imaginary part of a complex number. Status: STANDARD.

$$2E\,V_{\mathrm{CC}} = \Delta m^2_{ee}\cos 2\theta_{13}, \qquad \Delta m^2_{ee} = c^2_{12}\,\Delta m^2_{31} + s^2_{12}\,\Delta m^2_{32}, \qquad 2E\,V_{\mathrm{CC}}\,c^2_{13} = \Delta m^2_{21}\cos 2\theta_{12} \tag{FE-106}$$

Symbols in (FE-106).

- E: The energy of a particle: of the neutrino in the flavour equations, $E = (k^2 + m^2)^{1/2}$ of a particle of momentum k in the thermal integrals. Unit: GeV. Status: STANDARD.
- V_{CC}, V_{NC}: The charged-current potential of electron neutrinos and the neutral-current potential common to all flavours. Unit: GeV. Status: STANDARD.
- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m^2_{21} = m^2_2 - m^2_1$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- Δm^2_{ee}: The effective squared-mass difference of electron-neutrino disappearance, $\Delta m^2_{ee} = c^2_{12}\Delta m^2_{31} + s^2_{12}\Delta m^2_{32}$. Unit: GeV^2. Status: STANDARD.
- θ_{ij}: The mixing angles of the standard parametrisation of U (or of V_{CKM}): θ_{12}, θ_{13}, θ_{23}; in two-flavour formulas θ_{ij} is the angle of the sector (i, j). Unit: dimensionless. Value: $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$ (PMNS). Status: STANDARD.
- c_{12}, c_{13}, c_{23}: The cosines of the mixing angles, $c_{ij} = \cos\theta_{ij}$. Unit: dimensionless. Status: STANDARD.
- Δm^2_{21}, Δm^2_{31}, Δm^2_{32}: The differences of the squared neutrino masses, $\Delta m^2_{ij} = m^2_i - m^2_j$. Unit: GeV^2 (quoted in eV^2). Value: $\Delta m^2_{21} = 7.49 \times 10^{-5}$ eV^2, $\Delta m^2_{31} = 2.513 \times 10^{-3}$ eV^2 (NuFIT 6.0). Status: STANDARD.
- s_{12}, s_{13}, s_{23}: The sines of the mixing angles, $s_{ij} = \sin\theta_{ij}$. Unit: dimensionless. Status: STANDARD.

$$\begin{aligned} \mathcal{P}_{ee} &= c^4_{13}\Big(\frac{1}{2} + \frac{1}{2}\cos 2\theta_{12}\cos 2\theta^{\mathrm{m}}_{12}\Big) + s^4_{13}, \qquad A_{\mathrm{MSW}} = \frac{2E\,V_{\mathrm{CC}}\,c^2_{13}}{\Delta m^2_{21}}, \\ \cos 2\theta^{\mathrm{m}}_{12} &= \frac{\cos 2\theta_{12} - A_{\mathrm{MSW}}}{\sqrt{\big(\cos 2\theta_{12} - A_{\mathrm{MSW}}\big)^2 + \sin^2 2\theta_{12}}} \end{aligned} \tag{FE-107}$$

Symbols in (FE-107).

- $\mathcal{P}$: The probability of a flavour transition, $\mathcal{P}(\nu_\alpha \to \nu_\beta)$; $\mathcal{P}_{ee}$ the survival probability of electron neutrinos. Unit: dimensionless. Status: STANDARD.
- e: The electron flavour as a value of a flavour index ($\delta_{\alpha e}$, U_{e2}, V_e, $\mathcal{P}_{ee}$). Status: STANDARD.
- c_{12}, c_{13}, c_{23}: The cosines of the mixing angles, $c_{ij} = \cos\theta_{ij}$. Unit: dimensionless. Status: STANDARD.
- θ_{ij}: The mixing angles of the standard parametrisation of U (or of V_{CKM}): θ_{12}, θ_{13}, θ_{23}; in two-flavour formulas θ_{ij} is the angle of the sector (i, j). Unit: dimensionless. Value: $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$ (PMNS). Status: STANDARD.
- θ^{m}_{12}: The effective 1-2 mixing angle in matter at the production point of a solar neutrino. Unit: dimensionless. Status: STANDARD.
- s_{12}, s_{13}, s_{23}: The sines of the mixing angles, $s_{ij} = \sin\theta_{ij}$. Unit: dimensionless. Status: STANDARD.
- A_{MSW}: The ratio of the matter term to the vacuum term in the 1-2 sector. Unit: dimensionless. Status: STANDARD.
- E: The energy of a particle: of the neutrino in the flavour equations, $E = (k^2 + m^2)^{1/2}$ of a particle of momentum k in the thermal integrals. Unit: GeV. Status: STANDARD.

- $V_{\rm CC}$, $V_{\rm NC}$: The charged-current potential of electron neutrinos and the neutral-current potential common to all flavours. Unit: GeV. Status: STANDARD.
- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m^2_{21} = m^2_2 - m^2_1$ a difference of squared masses, $\Delta N_{\rm eff}$ an addition to $N_{\rm eff}$, $\Delta S_{\rm m}$ the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- Δm^2_{21}, Δm^2_{31}, Δm^2_{32}: The differences of the squared neutrino masses, $\Delta m^2_{ij} = m^2_i - m^2_j$. Unit: GeV^2 (quoted in eV^2). Value: $\Delta m^2_{21} = 7.49 \times 10^{-5}$ eV^2, $\Delta m^2_{31} = 2.513 \times 10^{-3}$ eV^2 (NuFIT 6.0). Status: STANDARD.

$$n_{\rm e}^{\rm res} = \frac{\Delta m^2_{ij}\cos 2\theta_{ij}}{2\sqrt{2}\,G_{\rm F}E}, \qquad \gamma_{\rm ad} = \frac{\Delta m^2_{ij}\sin^2 2\theta_{ij}}{2E\cos 2\theta_{ij}\,|{\rm d}\ln n_{\rm e}/{\rm d}r|} \gg 1, \qquad \mathcal{P}_{\rm LZ} = {\rm e}^{-\pi\gamma_{\rm ad}/2} \tag{FE-108}$$

Symbols in (FE-108).

- $n_{\rm e}^{\rm res}$: The electron density of an MSW resonance. Unit: GeV^3. Status: STANDARD.
- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m^2_{21} = m^2_2 - m^2_1$ a difference of squared masses, $\Delta N_{\rm eff}$ an addition to $N_{\rm eff}$, $\Delta S_{\rm m}$ the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- m, m_s, m_i: A four-dimensional mass: m the mass of a mode (the eigenvalue of an x_0 problem) or of a fermion; m_s the mass of the species s; m_i the mass of the neutrino mass state i. Unit: GeV. Status: DERIVED.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- j: A second generation index ($j = 1, 2, 3$). Status: STANDARD.
- θ_{ij}: The mixing angles of the standard parametrisation of U (or of $V_{\rm CKM}$): θ_{12}, θ_{13}, θ_{23}; in two-flavour formulas θ_{ij} is the angle of the sector (i, j). Unit: dimensionless. Value: $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$ (PMNS). Status: STANDARD.
- $G_{\rm F}$: The Fermi constant. Unit: GeV^{-2}. Value: 1.1663788×10^{-5} GeV^{-2} (measured). Status: STANDARD.
- E: The energy of a particle: of the neutrino in the flavour equations, $E = (k^2 + m^2)^{1/2}$ of a particle of momentum k in the thermal integrals. Unit: GeV. Status: STANDARD.
- $\gamma_{\rm ad}$: The adiabaticity parameter of a resonance (adiabatic for $\gamma_{\rm ad} \gg 1$). Unit: dimensionless. Status: STANDARD.
- d: The differential: dX is an infinitesimal change of X, dX/dY the derivative of X with respect to Y, d$^n X$ the volume element of n variables. Status: STANDARD.
- $n_{\rm e}$: The number density of electrons (net of positrons); in the Thomson terms of the photon and baryon equations only free (ionised) electrons count. Unit: GeV^3. Status: STANDARD.
- r: A distance: between the two masses in $V_{\rm N}(r)$, from the centre of the star in the adiabaticity condition. Unit: length. Status: STANDARD.
- $\mathcal{P}_{\rm LZ}$: The Landau-Zener probability of a jump between the matter eigenstates at a resonance. Unit: dimensionless. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

The early Universe: the thermal potential, the kinetic equations and the decoupling of the neutrinos:

$$V_e - V_\mu = -\frac{8\sqrt{2}\,G_{\rm F}|\vec{k}|}{3m^2_{\rm W}}\,\rho_{\rm e}, \qquad \rho_{\rm e} \simeq \frac{7\pi^2}{60}\,T^4 \quad (T \gg m_{\rm e}) \tag{FE-109}$$

Symbols in (FE-109).

- V_α, V: The matter potential of forward scattering: V_α of the flavour α (V_e, V_μ), V the 3×3 matrix of the potentials in the flavour basis; antineutrinos have $-V$. Unit: GeV. Status: STANDARD.
- e: The electron flavour as a value of a flavour index ($\delta_{\alpha e}$, U_{e2}, V_e, $\mathcal{P}_{ee}$). Status: STANDARD.

- G_{F}: The Fermi constant. Unit: GeV^{-2}. Value: $1.1663788 \times 10^{-5}\ \mathrm{GeV}^{-2}$ (measured). Status: STANDARD.
- $\vec{k},\ \vec{k}_o$: The wave vector (momentum) in ordinary 3-space; $\vec{k}_o$ that of the orbital o, a point of the lattice $k_{\mathrm{lat}}\vec{n}_o$. Unit: GeV. Status: STANDARD.
- m_{W}: The mass of W^+ and W^-. Unit: GeV. Value: 80.3625 GeV (measured, input). Status: STANDARD.
- $\rho_{\mathrm{e}},\ p_{\mathrm{e}}$: The energy density and the pressure of the electrons and positrons together (ideal gas). Unit: GeV^4. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.
- m_{e}: The electron mass. Unit: GeV. Value: 0.51099895 MeV. Status: STANDARD.

$$\mathrm{i}\Big(\partial_{x_4} + \frac{\vec{k}}{|\vec{k}|}\cdot\vec{\nabla}\Big)\boldsymbol{\rho}_{\vec{k}} = \big[H^{\mathrm{vac}}_{\vec{k}} + V + V_{\nu\nu},\ \boldsymbol{\rho}_{\vec{k}}\big] + \mathrm{i}\,C_{\mathrm{coll}}[\boldsymbol{\rho}], \qquad H^{\mathrm{vac}}_{\vec{k}} = \frac{U\,\mathrm{diag}\big(m_1^2,\ m_2^2,\ m_3^2\big)\,U^\dagger}{2|\vec{k}|},$$
$$V_{\nu\nu} = \sqrt{2}\,G_{\mathrm{F}}\int \frac{\mathrm{d}^3\vec{k}_2}{(2\pi)^3}\Big(1 - \frac{\vec{k}\cdot\vec{k}_2}{|\vec{k}||\vec{k}_2|}\Big)\big(\boldsymbol{\rho}_{\vec{k}_2} - \bar{\boldsymbol{\rho}}_{\vec{k}_2}\big), \qquad \partial_{x_4} \to \partial_{x_4} - \mathcal{H}|\vec{k}|\,\partial_{|\vec{k}|}\ \ \text{(expanding Universe)} \tag{FE-110}$$

Symbols in (FE-110).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\vec{k},\ \vec{k}_o$: The wave vector (momentum) in ordinary 3-space; $\vec{k}_o$ that of the orbital o, a point of the lattice $k_{\mathrm{lat}}\vec{n}_o$. Unit: GeV. Status: STANDARD.
- $\vec{\nabla}$: The gradient in ordinary 3-space, $(\partial/\partial x_1, \partial/\partial x_2, \partial/\partial x_3)$. Unit: 1/length. Status: STANDARD.
- $\boldsymbol{\rho}_{\vec{k}},\ \bar{\boldsymbol{\rho}}_{\vec{k}}$: The 3×3 flavour density matrices of the neutrinos and of the antineutrinos of momentum $\vec{k}$. Unit: dimensionless. Status: STANDARD.
- $H^{\mathrm{vac}}_{\vec{k}}$: The vacuum flavour Hamiltonian of a neutrino of momentum $\vec{k}$. Unit: GeV. Status: STANDARD.
- $V_\alpha,\ V$: The matter potential of forward scattering: V_α of the flavour α (V_e, V_μ), V the 3×3 matrix of the potentials in the flavour basis; antineutrinos have $-V$. Unit: GeV. Status: STANDARD.
- $V_{\nu\nu}$: The potential of neutrino-neutrino forward scattering (a matrix in flavour space). Unit: GeV. Status: STANDARD.
- C_{coll}: The collision term of the quantum kinetic equations. Unit: GeV. Status: STANDARD.
- U: The PMNS matrix, the unitary 3×3 mixing matrix of the leptons, $\nu_\alpha = \sum_i U_{\alpha i}\nu_i$ (rows $\alpha = e, \mu, \tau$, columns the mass states i); in UM-2 $U = \mathcal{K}_e^\dagger\mathcal{K}_\nu$. Unit: dimensionless. Value: $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$, $\delta_{\mathrm{CP}} = 212^\circ$ (NuFIT 6.0). Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ when it is a tensor in eight dimensions). Status: STANDARD.
- $m_1,\ m_2,\ m_3$: The masses of the neutrino mass states ν_1, ν_2, ν_3 (normal ordering). Unit: GeV (quoted in meV). Value: 1.000, 8.712, 50.140 meV (m_1 ASSUMED). Status: DERIVED.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- G_{F}: The Fermi constant. Unit: GeV^{-2}. Value: $1.1663788 \times 10^{-5}\ \mathrm{GeV}^{-2}$ (measured). Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $k_2,\ \vec{k}_2$: A second momentum (variable of integration). Unit: GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_{\mathrm{Pl}}$). Unit: GeV. Status: DERIVED.

$$\frac{\mathrm{d}T_\gamma}{\mathrm{d}x_4} = -\frac{4\mathcal{H}\rho_\gamma + 3\mathcal{H}\big(\rho_\mathrm{e} + p_\mathrm{e}\big) + 3\mathcal{H}T_\gamma\,\frac{\mathrm{d}p_\mathrm{QED}}{\mathrm{d}T_\gamma} + C_{\nu_e} + 2C_{\nu_\mu}}{\frac{\mathrm{d}\rho_\gamma}{\mathrm{d}T_\gamma} + \frac{\mathrm{d}\rho_\mathrm{e}}{\mathrm{d}T_\gamma} + T_\gamma\,\frac{\mathrm{d}^2 p_\mathrm{QED}}{\mathrm{d}T_\gamma^2}}, \qquad \frac{\mathrm{d}T_{\nu_\alpha}}{\mathrm{d}x_4} = -\mathcal{H}T_{\nu_\alpha} + \frac{C_{\nu_\alpha}}{\mathrm{d}\rho_{\nu_\alpha}/\mathrm{d}T_{\nu_\alpha}},$$
$$\mathcal{H} = \frac{\sqrt{8\pi\rho_\mathrm{tot}/3}}{M_\mathrm{Pl}}, \qquad \rho_\gamma = \frac{\pi^2}{15}\,T_\gamma^4, \qquad \rho_{\nu_\alpha} = \frac{7\pi^2}{120}\,T_{\nu_\alpha}^4 \tag{FE-111}$$

Symbols in (FE-111).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- T_γ: The temperature of the photons (equal to the plasma temperature T; the subscript is written where the neutrinos have their own temperature). Unit: GeV. Value: 2.7255 K today. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_\mathrm{Pl}$). Unit: GeV. Status: DERIVED.
- ρ_γ: The energy density of the photons, $\rho_\gamma = \pi^2 T_\gamma^4/15$. Unit: GeV^4. Status: STANDARD.
- ρ_e, p_e: The energy density and the pressure of the electrons and positrons together (ideal gas). Unit: GeV^4. Status: STANDARD.
- p_QED, $p_\mathrm{QED}^{(2)}$, $p_\mathrm{QED}^{(3)}$: The finite-temperature QED correction to the pressure of the electron-photon plasma, and its parts of order e^2 and e^3. Unit: GeV^4. Status: STANDARD.
- C_{ν_α}, C_{ν_e}, C_{ν_μ}: The energy transfer rate per volume into the neutrinos of the flavour α by collisions. Unit: GeV^5. Status: STANDARD.
- T_{ν_α}, T_{ν_e}, T_{ν_μ}: The temperature of the neutrinos of the flavour α (of ν_e, of ν_μ and ν_τ) during decoupling. Unit: GeV. Status: STANDARD.
- ρ_ν, ρ_{ν_α}: The energy density of one neutrino flavour (neutrinos and antineutrinos) at the common temperature T_ν, $\frac{7}{8}\frac{\pi^2}{15}T_\nu^4$ when massless; ρ_{ν_α} that of the flavour α at its temperature T_{ν_α}. Unit: GeV^4. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- ρ_tot, p_tot: The total background energy density and pressure of four-dimensional cosmology (all species). Unit: GeV^4. Status: STANDARD.
- M_Pl: The Planck mass, $M_\mathrm{Pl} = G_\mathrm{N}^{-1/2}$. Unit: GeV. Value: 1.22091×10^{19} GeV. Status: STANDARD.

$$\begin{aligned}
C_{\nu_e} &= \frac{G_\mathrm{F}^2}{\pi^5}\Big[4\Big(\big(\tilde{g}_L^{\nu_e e}\big)^2 + \big(\tilde{g}_R^{\nu_e e}\big)^2\Big)\mathcal{F}_\mathrm{tr}(T_\gamma, T_{\nu_e}) + 2\mathcal{F}_\mathrm{tr}(T_{\nu_\mu}, T_{\nu_e})\Big],\\
C_{\nu_\mu} &= \frac{G_\mathrm{F}^2}{\pi^5}\Big[4\Big(\big(\tilde{g}_L^{\nu_\mu e}\big)^2 + \big(\tilde{g}_R^{\nu_\mu e}\big)^2\Big)\mathcal{F}_\mathrm{tr}(T_\gamma, T_{\nu_\mu}) - \mathcal{F}_\mathrm{tr}(T_{\nu_\mu}, T_{\nu_e})\Big],\\
\mathcal{F}_\mathrm{tr}(T_\mathrm{A}, T_\mathrm{B}) &= 32 f_\mathrm{a}\big(T_\mathrm{A}^9 - T_\mathrm{B}^9\big) + 56 f_\mathrm{s} T_\mathrm{A}^4 T_\mathrm{B}^4\big(T_\mathrm{A} - T_\mathrm{B}\big),\\
\tilde{g}_L^{\nu_e e} &= \frac{1}{2} + \sin^2\theta_\mathrm{W}, \qquad \tilde{g}_R^{\nu_e e} = \tilde{g}_R^{\nu_\mu e} = \sin^2\theta_\mathrm{W}, \qquad \tilde{g}_L^{\nu_\mu e} = -\frac{1}{2} + \sin^2\theta_\mathrm{W}
\end{aligned} \tag{FE-112}$$

Symbols in (FE-112).

- C_{ν_α}, C_{ν_e}, C_{ν_μ}: The energy transfer rate per volume into the neutrinos of the flavour α by collisions. Unit: GeV^5. Status: STANDARD.
- G_F: The Fermi constant. Unit: GeV^{-2}. Value: 1.1663788×10^{-5} GeV^{-2} (measured). Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $\tilde{g}_L^{\nu_e e}$, $\tilde{g}_R^{\nu_e e}$: The effective left- and right-handed couplings of electron neutrinos to electrons (charged and neutral current), $\tilde{g}_L^{\nu_e e} = \frac{1}{2} + \sin^2\theta_\mathrm{W}$, $\tilde{g}_R^{\nu_e e} = \sin^2\theta_\mathrm{W}$. Unit: dimensionless. Value: 0.727 and 0.233 (radiatively corrected). Status: STANDARD.
- L: The chirality label L: the left-handed (doublet) zero mode, or the left-handed projection $P_L = (1 + \Gamma_4)/2$. Status: STANDARD.
- R: The chirality label R: the right-handed (singlet) zero mode or field, or the projection $P_R = (1 - \Gamma_4)/2$. Status: STANDARD.

- $\mathcal{F}_{\rm tr}$: The energy-transfer function of two species with the temperatures $T_{\rm A}$ and $T_{\rm B}$ (Maxwell-Boltzmann with the corrections $f_{\rm a}$, $f_{\rm s}$). Unit: GeV[9]. Status: STANDARD.
- T_γ: The temperature of the photons (equal to the plasma temperature T; the subscript is written where the neutrinos have their own temperature). Unit: GeV. Value: 2.7255 K today. Status: STANDARD.
- T_{ν_α}, T_{ν_e}, T_{ν_μ}: The temperature of the neutrinos of the flavour α (of ν_e, of ν_μ and ν_τ) during decoupling. Unit: GeV. Status: STANDARD.
- $\tilde{g}_L^{\nu_\mu e}$, $\tilde{g}_R^{\nu_\mu e}$: The effective couplings of muon (and tau) neutrinos to electrons (neutral current only), $\tilde{g}_L^{\nu_\mu e} = -\frac{1}{2} + \sin^2\theta_{\rm W}$, $\tilde{g}_R^{\nu_\mu e} = \sin^2\theta_{\rm W}$. Unit: dimensionless. Value: -0.273 and 0.233 (radiatively corrected). Status: STANDARD.
- $T_{\rm A}$, $T_{\rm B}$: The temperatures of the two species that exchange energy in $\mathcal{F}_{\rm tr}$. Unit: GeV. Status: STANDARD.
- $f_{\rm a}$, $f_{\rm s}$: The Fermi-Dirac corrections of the annihilation and scattering parts of the energy transfer. Unit: dimensionless. Value: 0.884 and 0.829. Status: STANDARD.
- $\theta_{\rm W}$: The weak mixing angle of the couplings, $\cos\theta_{\rm W} = \hat{g}_2/\hat{g}_Z$, $\sin\theta_{\rm W} = \hat{g}_1/\hat{g}_Z$; fixed by the measured W and Z masses with $\rho_{\rm EW,0}$. Unit: dimensionless. Value: $\cos^2\theta_{\rm W} = 0.776368$. Status: DERIVED.

$$
\begin{aligned}
&p_{\rm QED} = p_{\rm QED}^{(2)} + p_{\rm QED}^{(3)}, \qquad p_{\rm QED}^{(2)} = -\frac{e^2T^2}{12\pi^2}\,\mathcal{Q}_1 - \frac{e^2}{8\pi^4}\,\mathcal{Q}_1^2 + \frac{e^2 m_{\rm e}^2}{16\pi^4}\,\mathcal{Q}_2,\\
&p_{\rm QED}^{(3)} = \frac{e^3T}{12\pi^4}\,\mathcal{Q}_3^{3/2}, \qquad \rho_{\rm QED} = -p_{\rm QED} + T\,\frac{{\rm d}p_{\rm QED}}{{\rm d}T}
\end{aligned}
\tag{FE-113}
$$

Symbols in (FE-113).

- $p_{\rm QED}$, $p_{\rm QED}^{(2)}$, $p_{\rm QED}^{(3)}$: The finite-temperature QED correction to the pressure of the electron-photon plasma, and its parts of order e^2 and e^3. Unit: GeV[4]. Status: STANDARD.
- e: The elementary charge (the charge of the positron), $e^2 = 4\pi\alpha_{\rm em}$; W^+ has the charge $+e$ and W^- the charge $-e$. Unit: dimensionless. Value: $e^2 = 4\pi/137.036$ at low energy. Status: STANDARD.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $\mathcal{Q}_1$, $\mathcal{Q}_2$, $\mathcal{Q}_3$: The momentum integrals of the QED pressure. Unit: GeV[2]. Status: STANDARD.
- $m_{\rm e}$: The electron mass. Unit: GeV. Value: 0.51099895 MeV. Status: STANDARD.
- $\rho_{\rm QED}$: The QED correction to the energy density of the plasma. Unit: GeV[4]. Status: STANDARD.
- d: The differential: ${\rm d}X$ is an infinitesimal change of X, ${\rm d}X/{\rm d}Y$ the derivative of X with respect to Y, ${\rm d}^nX$ the volume element of n variables. Status: STANDARD.

$$
\begin{aligned}
&\mathcal{Q}_1 = \int_0^\infty {\rm d}k\,\frac{k^2}{E}\,n_{\rm D}(k), \qquad \mathcal{Q}_2 = \int_0^\infty\!\!\int_0^\infty {\rm d}k\,{\rm d}k_2\,\frac{kk_2}{E(k)E(k_2)}\,\ln\left|\frac{k+k_2}{k-k_2}\right|\,n_{\rm D}(k)\,n_{\rm D}(k_2),\\
&\mathcal{Q}_3 = \int_0^\infty {\rm d}k\,\frac{k^2+E^2}{E}\,n_{\rm D}(k), \qquad n_{\rm D}(k) = \frac{2}{{\rm e}^{E/T}+1}, \qquad E = \sqrt{k^2+m_{\rm e}^2}
\end{aligned}
\tag{FE-114}
$$

Symbols in (FE-114).

- $\mathcal{Q}_1$, $\mathcal{Q}_2$, $\mathcal{Q}_3$: The momentum integrals of the QED pressure. Unit: GeV[2]. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no upper bound. Status: STANDARD.
- d: The differential: ${\rm d}X$ is an infinitesimal change of X, ${\rm d}X/{\rm d}Y$ the derivative of X with respect to Y, ${\rm d}^nX$ the volume element of n variables. Status: STANDARD.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum ${\rm e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- E: The energy of a particle: of the neutrino in the flavour equations, $E = (k^2+m^2)^{1/2}$ of a particle of momentum k in the thermal integrals. Unit: GeV. Status: STANDARD.
- $n_{\rm D}(k)$: The Fermi-Dirac occupation of the electrons and positrons summed over the two spin states. Unit: dimensionless. Status: STANDARD.

- k_2, $\vec{k}_2$: A second momentum (variable of integration). Unit: GeV. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.
- m_e: The electron mass. Unit: GeV. Value: 0.51099895 MeV. Status: STANDARD.

$$N_\mathrm{eff} = \Big(\frac{11}{4}\Big)^{4/3}\Big[\Big(\frac{T_{\nu_e}}{T_\gamma}\Big)^4 + 2\Big(\frac{T_{\nu_\mu}}{T_\gamma}\Big)^4\Big], \qquad \frac{\mathrm{d}T_\nu}{\mathrm{d}x_4} = -\mathcal{H}T_\nu + \frac{C_{\nu_e} + 2C_{\nu_\mu}}{3\,\mathrm{d}\rho_\nu/\mathrm{d}T_\nu} \tag{FE-115}$$

Symbols in (FE-115).

- N_eff: The effective number of neutrino species (the radiation beyond the photons in units of one neutrino species with $T_\nu/T_\gamma = (4/11)^{1/3}$); ΔN_eff an addition to it. Unit: dimensionless. Value: 3.0432 (neutrinos) + 0.0010 (fold bands). Status: STANDARD.
- T_{ν_α}, T_{ν_e}, T_{ν_μ}: The temperature of the neutrinos of the flavour α (of ν_e, of ν_μ and ν_τ) during decoupling. Unit: GeV. Status: STANDARD.
- T_γ: The temperature of the photons (equal to the plasma temperature T; the subscript is written where the neutrinos have their own temperature). Unit: GeV. Value: 2.7255 K today. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- T_ν: The temperature of the relic neutrinos (one common temperature). Unit: GeV. Value: 1.9454 K today. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_\mathrm{Pl}$). Unit: GeV. Status: DERIVED.
- C_{ν_α}, C_{ν_e}, C_{ν_μ}: The energy transfer rate per volume into the neutrinos of the flavour α by collisions. Unit: GeV^5. Status: STANDARD.
- ρ_ν, ρ_{ν_α}: The energy density of one neutrino flavour (neutrinos and antineutrinos) at the common temperature T_ν, $\frac{7}{8}\frac{\pi^2}{15}T_\nu^4$ when massless; ρ_{ν_α} that of the flavour α at its temperature T_{ν_α}. Unit: GeV^4. Status: STANDARD.

The dense neutrino gas near a supernova core:

$$G_\mathrm{ELN}(u) = \sqrt{2}\,G_\mathrm{F}\big[n_{\nu_e}\,g_{\nu_e}(u) - n_{\bar\nu_e}\,g_{\bar\nu_e}(u)\big], \qquad \int_{-1}^{1} g_{\nu_e}\,\frac{\mathrm{d}u}{2} = \int_{-1}^{1} g_{\bar\nu_e}\,\frac{\mathrm{d}u}{2} = 1, \qquad \mathcal{I}_l = \int_{-1}^{1}\frac{\mathrm{d}u}{2}\,G_\mathrm{ELN}(u)\,u^l \tag{FE-116}$$

Symbols in (FE-116).

- $G_\mathrm{ELN}(u)$: The angular distribution of the electron lepton number of a dense neutrino gas (in units of a potential). Unit: GeV. Status: STANDARD.
- u: The cosine of the angle between a momentum and the symmetry axis x_3 of the neutrino gas. Unit: dimensionless. Status: STANDARD.
- G_F: The Fermi constant. Unit: GeV^{-2}. Value: 1.1663788×10^{-5} GeV^{-2} (measured). Status: STANDARD.
- n_{ν_e}, $n_{\bar\nu_e}$: The number densities of electron neutrinos and electron antineutrinos. Unit: GeV^3. Status: STANDARD.
- $g_{\nu_e}(u)$, $g_{\bar\nu_e}(u)$: The normalised angular distributions of electron neutrinos and antineutrinos. Unit: dimensionless. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $\mathcal{I}_l$: The l-th angular moment of the electron-lepton-number distribution. Unit: GeV. Status: STANDARD.
- l: The multipole (angular moment) number $l = 0, 1, 2, \ldots$ of a Boltzmann hierarchy or of an angular moment $\int u^l \cdots$; as a coefficient it is the same number. Status: STANDARD.

$$\mathcal{C}(u) \propto \mathrm{e}^{-\mathrm{i}(\omega_\mathrm{f} x_4 - k_\mathrm{f} x_3)}, \qquad \bar{\omega} = \omega_\mathrm{f} - \mathcal{I}_0, \qquad \bar{k} = k_\mathrm{f} - \mathcal{I}_1 : \qquad \det \Pi(\bar{\omega}, \bar{k}) = 0,$$
$$\Pi = \mathrm{diag}(1,\ -1) + \int_{-1}^{1} \frac{\mathrm{d}u}{2} \frac{G_\mathrm{ELN}(u)}{\bar{\omega} - \bar{k}u} \begin{pmatrix} 1 & u \\ u & u^2 \end{pmatrix} \tag{FE-117}$$

Symbols in (FE-117).

- $\mathcal{C}(u)$: The flavour coherence (off-diagonal element of the density matrix) of the neutrinos moving in the direction u. Unit: dimensionless. Status: STANDARD.
- u: The cosine of the angle between a momentum and the symmetry axis x_3 of the neutrino gas. Unit: dimensionless. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- ω_f, k_f: The frequency and the wavenumber (along x_3) of a flavour wave. Unit: GeV. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space (also the symmetry axis of the fast-flavour analysis). Unit: length. Status: AUTHOR.
- $\bar{\omega}$, $\bar{k}$: The frequency and wavenumber of a flavour wave shifted by the moments, $\bar{\omega} = \omega_\mathrm{f} - \mathcal{I}_0$, $\bar{k} = k_\mathrm{f} - \mathcal{I}_1$; a mode with $\mathrm{Im}\,\bar{\omega} > 0$ grows. Unit: GeV. Status: STANDARD.
- $\mathcal{I}_l$: The l-th angular moment of the electron-lepton-number distribution. Unit: GeV. Status: STANDARD.
- Π: The 2×2 dispersion matrix of the fast flavour instability. Unit: dimensionless. Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ when it is a tensor in eight dimensions). Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $G_\mathrm{ELN}(u)$: The angular distribution of the electron lepton number of a dense neutrino gas (in units of a potential). Unit: GeV. Status: STANDARD.

$$\big(1 + \mathcal{J}_0\big)\big(-1 + \mathcal{J}_2\big) - \mathcal{J}_1^2 = 0, \qquad \mathcal{J}_l = \int_{-1}^{1} \frac{\mathrm{d}u}{2} \frac{G_\mathrm{ELN}(u)\, u^l}{\bar{\omega} - \bar{k}u}; \qquad -1 + \int_{-1}^{1} \frac{\mathrm{d}u}{2} \frac{G_\mathrm{ELN}(u)\,(1-u^2)}{2(\bar{\omega} - \bar{k}u)} = 0 \tag{FE-118}$$

Symbols in (FE-118).

- $\mathcal{J}_l$: The l-th moment of the dispersion integral of the fast flavour instability. Unit: dimensionless. Status: STANDARD.
- l: The multipole (angular moment) number $l = 0, 1, 2, \ldots$ of a Boltzmann hierarchy or of an angular moment $\int u^l \cdots$; as a coefficient it is the same number. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- u: The cosine of the angle between a momentum and the symmetry axis x_3 of the neutrino gas. Unit: dimensionless. Status: STANDARD.
- $G_\mathrm{ELN}(u)$: The angular distribution of the electron lepton number of a dense neutrino gas (in units of a potential). Unit: GeV. Status: STANDARD.
- $\bar{\omega}$, $\bar{k}$: The frequency and wavenumber of a flavour wave shifted by the moments, $\bar{\omega} = \omega_\mathrm{f} - \mathcal{I}_0$, $\bar{k} = k_\mathrm{f} - \mathcal{I}_1$; a mode with $\mathrm{Im}\,\bar{\omega} > 0$ grows. Unit: GeV. Status: STANDARD.

$$\mathrm{i}\,\frac{\mathrm{d}\mathcal{C}(u)}{\mathrm{d}x_4} = \big(\mathcal{I}_0 - u\,\mathcal{I}_1\big)\mathcal{C}(u) - \int_{-1}^{1} \frac{\mathrm{d}\tilde{u}}{2}\, G_\mathrm{ELN}(\tilde{u})\,(1 - u\tilde{u})\,\mathcal{C}(\tilde{u});$$
$$\bar{\omega}^2 - \big(\mathcal{I}_2 - \mathcal{I}_0\big)\bar{\omega} - \big(\mathcal{I}_0\mathcal{I}_2 - \mathcal{I}_1^2\big) = 0, \qquad \mathrm{Im}\,\bar{\omega} = \frac{1}{2}\sqrt{4\mathcal{I}_1^2 - \big(\mathcal{I}_0 + \mathcal{I}_2\big)^2} \tag{FE-119}$$

Symbols in (FE-119).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $\mathcal{C}(u)$: The flavour coherence (off-diagonal element of the density matrix) of the neutrinos moving in the direction u. Unit: dimensionless. Status: STANDARD.
- u: The cosine of the angle between a momentum and the symmetry axis x_3 of the neutrino gas. Unit: dimensionless. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\mathcal{I}_l$: The l-th angular moment of the electron-lepton-number distribution. Unit: GeV. Status: STANDARD.
- $\tilde{u}$: A second direction cosine (variable of integration). Unit: dimensionless. Status: STANDARD.
- $G_{\mathrm{ELN}}(u)$: The angular distribution of the electron lepton number of a dense neutrino gas (in units of a potential). Unit: GeV. Status: STANDARD.
- $\bar{\omega}$, $\bar{k}$: The frequency and wavenumber of a flavour wave shifted by the moments, $\bar{\omega} = \omega_{\mathrm{f}} - \mathcal{I}_0$, $\bar{k} = k_{\mathrm{f}} - \mathcal{I}_1$; a mode with $\mathrm{Im}\,\bar{\omega} > 0$ grows. Unit: GeV. Status: STANDARD.
- Im: The imaginary part of a complex number. Status: STANDARD.

The cosmic neutrino background:

$$f_{\mathrm{FD}}(k) = \frac{1}{\mathrm{e}^{k/T_\nu}+1}, \qquad n_i = 2\int \frac{\mathrm{d}^3\vec{k}}{(2\pi)^3}\, f_{\mathrm{FD}} = \frac{3\zeta(3)}{2\pi^2}\, T_\nu^3, \qquad \rho_i = 2\int \frac{\mathrm{d}^3\vec{k}}{(2\pi)^3}\, \sqrt{k^2+m_i^2}\, f_{\mathrm{FD}},$$
$$\bar{v}_i = \frac{\int_0^\infty k^2\, \frac{k}{\sqrt{k^2+m_i^2}}\, f_{\mathrm{FD}}\, \mathrm{d}k}{\int_0^\infty k^2\, f_{\mathrm{FD}}\, \mathrm{d}k}, \qquad \text{flavour content of } \nu_i:\ |U_{\alpha i}|^2 \tag{FE-120}$$

Symbols in (FE-120).

- f_{FD}: The Fermi-Dirac distribution of the relic neutrinos, $f_{\mathrm{FD}} = 1/(\mathrm{e}^{k/T_\nu}+1)$. Unit: dimensionless. Status: STANDARD.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- T_ν: The temperature of the relic neutrinos (one common temperature). Unit: GeV. Value: 1.9454 K today. Status: STANDARD.
- n, n_o, $n_{\psi i}$, n_f, n_i: A number density: $n(y)$ the particle density of dirac16complex in the Kohn-Sham equations and n_o the contribution of the orbital o, $n_{\psi i}(y)$ the density of the Standard-Model fermion ψ_i (per proper 7-volume of eight dimensions); n_f the density of the fermion component f and n_i that of the neutrino mass state i (per 3-volume of four dimensions). Unit: GeV^7 (eight dimensions), GeV^3 (four dimensions). Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $\vec{k}$, $\vec{k}_o$: The wave vector (momentum) in ordinary 3-space; $\vec{k}_o$ that of the orbital o, a point of the lattice $k_{\mathrm{lat}}\vec{n}_o$. Unit: GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- ζ: The Riemann zeta function; only its value $\zeta(3)$ occurs, the sum of $1/n^3$ over the positive integers n. Unit: dimensionless. Value: $\zeta(3) = 1.2020569$. Status: STANDARD.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- m, m_s, m_i: A four-dimensional mass: m the mass of a mode (the eigenvalue of an x_0 problem) or of a fermion; m_s the mass of the species s; m_i the mass of the neutrino mass state i. Unit: GeV. Status: DERIVED.

- $\bar{v}_i$: The mean velocity of the relic neutrinos of the mass state i. Unit: dimensionless (c = 1). Value: 0.435, 0.0604, 0.0105 today. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no upper bound. Status: STANDARD.
- ν_α, ν_i: A left-handed neutrino: ν_α of the flavour $\alpha = e, \mu, \tau$ (ν_e, ν_μ, ν_τ), ν_i of the mass state i (mass m_i). Unit: GeV$^{3/2}$. Status: STANDARD.
- U: The PMNS matrix, the unitary 3×3 mixing matrix of the leptons, $\nu_\alpha = \sum_i U_{\alpha i}\nu_i$ (rows $\alpha = e, \mu, \tau$, columns the mass states i); in UM-2 $U = \mathcal{K}_e^\dagger \mathcal{K}_\nu$. Unit: dimensionless. Value: $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$, $\delta_{\rm CP} = 212°$ (NuFIT 6.0). Status: STANDARD.
- α: A flavour index running over e, μ, τ (the flavour states of the leptons and neutrinos). Status: STANDARD.

8.2 The approximation of step 5

Ultra-relativistic, coherent flavour evolution. Every neutrino of the sources considered has $E \geq 0.3$ MeV, so the corrections of order $(m_i/E)^2$ to the evolution are negligible,

$$\left(\frac{m_3}{E}\right)^2 \leq 2.8 \times 10^{-14} \qquad (E \geq 0.3 \text{ MeV}),$$

Symbols in this equation.

- m_1, m_2, m_3: The masses of the neutrino mass states ν_1, ν_2, ν_3 (normal ordering). Unit: GeV (quoted in meV). Value: 1.000, 8.712, 50.140 meV (m_1 ASSUMED). Status: DERIVED.
- E: The energy of a particle: of the neutrino in the flavour equations, $E = (k^2 + m^2)^{1/2}$ of a particle of momentum k in the thermal integrals. Unit: GeV. Status: STANDARD.

and outside the early Universe the collisions are negligible against the forward scattering. The flavour-universal neutral-current potential drops out exactly. The adiabatic solar formula leaves out the non-adiabatic and production-region effects, 1.3×10^{-3} at most against the numerical evolution (whose layer discretisation is converged to 2.2×10^{-6}); the Landau-Zener jumps in the supernova envelope are at most 2.6×10^{-77}; the collective evolution beyond the linear analysis is OPEN. The other approximations of step 5 are these.

- *The thermal potential* is taken at leading order in $T/m_{\rm W}$ (the next order is $(T/m_{\rm W})^2 \sim 4 \times 10^{-9}$) in a CP-symmetric plasma; the CP-asymmetric potential of the electron excess is 1.1×10^{-2} of the thermal one at the solar onset temperature and 2.2×10^{-3} at the atmospheric one (ESTIMATE; it would move the onset temperatures by about a sixth of this).
- *Neutrino decoupling with one temperature per flavour* (momentum averaged): the electron mass in the rates and the momentum dependence, which this method does not resolve, make the computed $N_{\rm eff}$ larger than the precision value by 0.0044; the treatment of the oscillations (separate or common temperature) changes it by 0.0009; the neutrino masses enter as $(m_3/T)^2 = 6 \times 10^{-16}$ at decoupling.
- *The fast flavour instability* is analysed linearly in the flavour coherence; near the neutrinosphere the vacuum term is 9.0×10^{-6} of the self-interaction term.
- *The CNB* is the Fermi-Dirac spectrum of instantaneous decoupling; the non-instantaneous heating raises its density today by $(N_{\rm eff}/3)^{3/4} - 1 = 1.08$ %, which the CMB computation of step 6 includes; gravitational clustering is not computed.

8.3 The solution of step 5

8.3.1 Masses, mixing and the evolution

The neutrino masses are the x_0 eigenvalues of step 2 with the lightest mass chosen (ASSUMED) and the normal ordering (the NuFIT 6.0 best fit; the inverted ordering has $\Delta\chi^2 = 6.1$): $\Delta m^2_{21} = 7.49 \times 10^{-5}$ eV^2, $\Delta m^2_{31} = 2.513 \times 10^{-3}$ eV^2, and the sum of the masses 0.0599 eV lies below the DESI DR2 bound 0.064 eV (DERIVED, 16 checks). The flavour amplitudes $\boldsymbol{\psi}$ of a neutrino of energy E evolve with $H_{\rm fl}$; for a constant density the evolution operator is the matrix exponential, otherwise the ordered product of the operators of thin layers, and $\mathcal{P}(\nu_\alpha \to \nu_\beta) = |\mathcal{U}_{\beta\alpha}|^2$ (antineutrinos: $U \to U^*$, $V \to -V$).

8.3.2 Vacuum, matter and accelerators

Reactor $\bar{\nu}_e$ of 4 MeV survive with the probability 0.9183 at 1.65 km, 0.2263 at 52.5 km and 0.2674 at 180 km; the vacuum ν_μ survival has its first minimum, 0.0060, at $L_\mathrm{b}/E = 1516$ km/GeV. The 1-3 MSW resonance lies at 11.135 GeV in rock of the density 2.8 g/cm^3 with the electron fraction 0.5; the exact three-flavour Hamiltonian has $|U^\mathrm{m}_{e3}|^2 = 1/2$ (the electron content of the third matter eigenstate) at 11.133 GeV (agreement 1.6×10^{-4}; with Δm^2_{31} alone the resonance formula gives 11.24 GeV). The 1-2 resonance at the centre of the Sun ($n_\mathrm{e} = 6.02 \times 10^{25}$ cm^{-3}) lies at 1.93 MeV. Long-baseline appearance in matter of constant density ($\delta_\mathrm{CP} = 212°$, normal ordering; DERIVED):

experiment	L_b (km)	E (GeV)	density (g/cm^3)	$\mathcal{P}(\nu_\mu \to \nu_e)$	$\mathcal{P}(\bar{\nu}_\mu \to \bar{\nu}_e)$	vacuum $\mathcal{P}(\nu_\mu \to \nu_e)$	$\mathcal{P}(\nu_\mu \to \nu_\mu)$
T2K	295	0.6	2.6	0.0517	0.0313	0.0478	0.0085
NOvA	810	2.0	2.84	0.0481	0.0236	0.0404	0.1010
DUNE	1300	2.5	2.848	0.0701	0.0195	0.0491	0.0098

The matter effect raises the neutrino and lowers the antineutrino appearance for the normal ordering, and the phase $\delta_\mathrm{CP} = 212°$ adds to the asymmetry.

8.3.3 The Sun

With the electron density $n_\mathrm{e}(r) = 6.02 \times 10^{23} \min(245\,\mathrm{e}^{-10.54\,r/R_\odot}, 100)$ cm^{-3} (ASSUMED profile; $R_\odot$ the solar radius) and production at the radius r_0, the adiabatic survival probability is that of the field equation written above, evaluated at r_0; the numerical three-flavour evolution through 20000 and 40000 layers from r_0 to the surface, followed by the incoherent sum over the mass states, $\mathcal{P}_{ee} = \sum_i |U_{ei}|^2\, \mathcal{P}(\nu_i \to \nu_e)$, gives (DERIVED):

source	E (MeV)	$r_0/R_\odot$	$\mathcal{P}_{ee}$ adiabatic	$\mathcal{P}_{ee}$ numerical
pp	0.30	0.10	0.5409	0.5408
^{7}Be	0.862	0.06	0.5197	0.5196
pep	1.44	0.07	0.4978	0.4976
^{8}B	5.0	0.05	0.3771	0.3767
^{8}B	10.0	0.05	0.3194	0.3185
^{8}B	15.0	0.05	0.3055	0.3043

The two agree to 1.3×10^{-3}: the conversion is adiabatic, vacuum dominated at low energy ($\mathcal{P}_{ee} \to c^4_{13}(1 - \frac{1}{2}\sin^2 2\theta_{12}) + s^4_{13}$) and matter dominated for ^{8}B ($\mathcal{P}_{ee} \to c^4_{13}\sin^2\theta_{12} + s^4_{13}$).

8.3.4 The early Universe and N_eff

The thermal potential of the $e^\pm$ plasma falls below the vacuum term $\Delta m^2 \cos 2\theta/(2|\vec{k}|)$ at the mean momentum $|\vec{k}| = 3.15\,T$ when $T = 4.99$ MeV for the atmospheric and $T = 2.39$ MeV for the solar splitting (DERIVED; m_W of the UM-2 fit): the flavours mix while the neutrinos decouple. The decoupling is computed from the temperature equations written above (Escudero's method; MeV units; the Planck mass $M_\mathrm{Pl} = 1.22091 \times 10^{19}$ GeV), with the radiatively corrected couplings 0.727, 0.233, −0.273, 0.233 (or the UM-2 tree-level couplings with $\sin^2\theta_\mathrm{W} = 0.22363$), the $e^\pm$ density and pressure with the electron mass, and the QED interaction pressure. The integrals are evaluated with Gauss-Legendre panels, the logarithmic singularity of $\mathcal{Q}_2$ at $k_2 = k$ removed by a substitution, and the massless limits $p^{(2)}_\mathrm{QED} = -5e^2T^4/288$, $p^{(3)}_\mathrm{QED} = e^3T^4/(36\sqrt{3}\pi)$, $\rho_\mathrm{e} = 7\pi^2T^4/60$ are reproduced to 10^{-16}. The equations run from $T = 20$ MeV to 5 keV (LSODA, relative tolerance 10^{-11}); for instantaneous decoupling at 20 MeV the plasma entropy is conserved and T_ν falls as e^{-a_4}. Results (DERIVED):




Figure 2: Flavour conversion in UM-2. Left: the solar ν_e survival probability (adiabatic formula for two production radii, numerical evolution as points). Middle: $\nu_\mu \to \nu_e$ and $\bar{\nu}_\mu \to \bar{\nu}_e$ appearance over 1300 km of rock (normal ordering, $\delta_{\mathrm{CP}} = 212°$). Right: the growth rate of the fast flavour instability against the wave number for the three angular distributions of the text.

run	N_{eff}	published (Escudero 2020, same method, Table 1 there)
instantaneous decoupling, no QED	3.0002	3.000
instantaneous, $O(e^2)$ QED	3.0108	3.011
instantaneous, $O(e^2) + O(e^3)$	3.0098	3.010
Fermi-Dirac energy transfer, one neutrino temperature, no QED	3.0393	3.039
the same with $O(e^2)$	3.0495	
the same with $O(e^2) + O(e^3)$	3.0485	3.049
UM-2, separate ν_e and $\nu_{\mu,\tau}$ temperatures (no oscillations), $O(e^3)$	3.0476	3.048 (separate temperatures)
UM-2, separate temperatures, UM-2 tree-level couplings	3.0472	
UM-2, one common temperature (efficient oscillations)	3.0485	

With separate temperatures $T_{\nu_e}/T_\gamma = 0.71732$ and $T_{\nu_\mu}/T_\gamma = 0.71621$; e^+e^- annihilation raises the comoving neutrino temperatures by 0.265 % (ν_e, which also has charged-current interactions with the electrons) and 0.109 % ($\nu_{\mu,\tau}$). The plasma of UM-2 at MeV temperatures is that of the three-flavour Standard Model (the fold bands made by freeze-in, $\Delta N_{\mathrm{eff}} = 0.001$ in step 9, are decoupled free-streaming radiation, which takes no part in the neutrino decoupling and is added to N_{eff} separately), so the precision Standard-Model value with oscillations and the electron mass in the rates is adopted,

$$N_{\mathrm{eff}} = 3.0432 \pm 0.0002 \qquad \text{(Cielo, Escudero, Mangano and Pisanti 2023)},$$

Symbols in this equation.

- N_{eff}: The effective number of neutrino species (the radiation beyond the photons in units of one neutrino species with $T_\nu/T_\gamma = (4/11)^{1/3}$); ΔN_{eff} an addition to it. Unit: dimensionless. Value: 3.0432 (neutrinos) + 0.0010 (fold bands). Status: STANDARD.

from which the computed range 3.0476 to 3.0485 differs by 0.0044 to 0.0053 (the electron mass in the rates and the momentum dependence, which the method of temperatures does not resolve).




Figure 3: Neutrino decoupling in UM-2 (three flavours, separate ν_e and $\nu_{\mu,\tau}$ temperatures, QED to $O(e^3)$). Left: the neutrino temperatures relative to the photon temperature; right: the comoving heating of the neutrinos by e^+e^- annihilation.

8.3.5 The cosmic neutrino background

After decoupling the flavour coherence is lost: the CNB of UM-2 is an incoherent mixture of the three mass eigenstates with the Fermi-Dirac spectrum at $T_\nu = (4/11)^{1/3} T_\gamma = 1.9454$ K $= 1.676 \times 10^{-4}$ eV ($T_\gamma = 2.7255$ K today, STANDARD): $n_i = 112.0$ cm^{-3} (ν_i and $\bar{\nu}_i$) for each mass state, and the density parameter of the three states is (DERIVED)

$$\Omega_\nu h_{100}^2 = \frac{\sum_{i=1}^{3} \rho_i}{\rho_{\rm crit}/h_{100}^2} = 6.382 \times 10^{-4}, \qquad \frac{\rho_{\rm crit}}{h_{100}^2} = 1.05368 \times 10^4 \text{ eV/cm}^3,$$

Symbols in this equation.

- Ω_ν: The density parameter of the three neutrino mass states today: their energy density divided by the critical density $\rho_{\rm crit}$ (with h_{100}^2: $\Omega_\nu h_{100}^2$). Unit: dimensionless. Value: $\Omega_\nu h_{100}^2 = 6.382 \times 10^{-4}$ (instantaneous decoupling), 6.4511×10^{-4} (heated spectrum, CAMB). Status: DERIVED.
- h_{100}: The reduced Hubble constant, $\mathcal{H}_0/(100$ km s^{-1} Mpc$^{-1})$. Unit: dimensionless. Value: 0.6738. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- $\rho_{\rm crit}$: The critical density, $3\mathcal{H}_0^2 M_{\rm Pl}^2/(8\pi)$. Unit: GeV^4 (quoted in GeV/cm^3). Value: $\rho_{\rm crit}/h_{100}^2 = 1.05368 \times 10^{-5}$ GeV/cm^3. Status: STANDARD.

compared with the non-relativistic value $\sum_i m_i/(94.06 \text{ eV}) = 6.363 \times 10^{-4}$ at this temperature (the semi-relativistic ν_1, whose energy density exceeds $m_1 n_1$ by 16 %, adds 0.3 %), with the conventional $\sum_i m_i/(93.14 \text{ eV}) = 6.426 \times 10^{-4}$ (which includes the heating of the non-instantaneous decoupling) and with 6.451×10^{-4} for the heated spectrum of $N_{\rm eff} = 3.0432$ (step 6: the degeneracy $N_{\rm eff}/3$ of each state corresponds to a Fermi-Dirac spectrum of the temperature $(N_{\rm eff}/3)^{1/4} T_\nu$, so the non-relativistic density is multiplied by $(N_{\rm eff}/3)^{3/4}$). The states (DERIVED):

state	mass (meV)	n_i (cm^{-3})	$\bar{\nu}_i/c$ today	$\lvert U_{ei}\rvert^2$	$\lvert U_{\mu i}\rvert^2$	$\lvert U_{\tau i}\rvert^2$
ν_1	1.000	112.0	0.435	0.677	0.112	0.211
ν_2	8.712	112.0	0.0604	0.301	0.428	0.271
ν_3	50.140	112.0	0.0105	0.022	0.460	0.518

The fold bands of dirac16complex are not part of the CNB (they are never thermalised, step 4; their freeze-in radiation is counted separately, step 9).

8.3.6 Supernovae

The MSW resonances of the envelope. With the density profile $10^4\,(10^9\,\mathrm{cm}/r)^3$ g/cm^3 and the electron fraction 0.5 (ASSUMED), a resonance lies at the density $n_\mathrm{e}^\mathrm{res}$, i.e. at the radius with $\mathrm{d}\ln n_\mathrm{e}/\mathrm{d}r = -3/r$, with the adiabaticity γ_ad and the jump probability $\mathcal{P}_\mathrm{LZ}$ of the field equation written above (DERIVED; two-flavour conditions: Δm^2_{31} and θ_{13} for the H resonance, Δm^2_{21} and θ_{12} for the L resonance, without the factor c^2_{13} of the solar condition, which would move the L densities by 2 %, and without Δm^2_{ee}, which would move the H densities by 1 %):

E (MeV)	H (1-3): density (g/cm^3)	radius (km)	γ_ad	L (1-2): density (g/cm^3)	radius (km)	γ_ad
10	3147	1.47×10^4	283	37.7	6.43×10^4	903
20	1573	1.85×10^4	178	18.8	8.10×10^4	569
40	787	2.33×10^4	112	9.42	1.02×10^5	358

Both resonances are adiabatic ($\mathcal{P}_\mathrm{LZ} \le 2.6\times10^{-77}$). For the normal ordering a ν_e produced in the core leaves the star as ν_3, $\mathcal{P}(\nu_e\to\nu_e) = |U_{e3}|^2 = 0.02215$, and a $\bar\nu_e$ as $\bar\nu_1$, $\mathcal{P}(\bar\nu_e\to\bar\nu_e) = |U_{e1}|^2 = 0.677$ (without collective effects and Earth matter). For representative accretion-phase neutrinospheres (luminosities 6, 6, 4×10^{52} erg/s and mean energies 11, 14, 15.5 MeV for ν_e, $\bar\nu_e$ and each heavy-lepton flavour; ASSUMED), number fluxes proportional to the luminosity over the mean energy and the survival probabilities above, the mean energy observed at the Earth, the flux-weighted mean of the surviving and the converted spectra, becomes 15.29 MeV (ν_e) and 14.34 MeV ($\bar\nu_e$).

The fast flavour instability. The electron-lepton-number distribution, the frequencies, the wavenumbers and the growth rates are measured in units of the self-interaction strength $\sqrt{2}G_\mathrm{F}n_{\nu_e}$ of the ν_e gas. The roots of the dispersion relation with $\mathrm{Im}\,\bar\omega > 0$ were found for real $\bar k$ on a grid ($-1.2 \le \bar k \le 1.2$, step 0.1) by vectorised Newton iterations with the analytic derivative from a grid of complex starting points (DERIVED):

angular distributions	ELN crossing	largest growth $\mathrm{Im}\,\bar\omega$ (branch, $\bar k$)	homogeneous mode	mode $k_\mathrm{f} = \mathcal{I}_1$
$g_{\nu_e} \propto \mathrm{e}^{-(1-u)^2/(2\cdot0.6^2)}$, $g_{\bar\nu_e} \propto \mathrm{e}^{-(1-u)^2/(2\cdot0.4^2)}$, $n_{\bar\nu_e}/n_{\nu_e} = 0.8$	yes	0.0877 (axisymmetric, $\bar k = -0.30$)	0.0185	stable
$g_{\nu_e} = 1 + 0.2u$, $g_{\bar\nu_e} = 1 + 0.6u$, $n_{\bar\nu_e}/n_{\nu_e} = 0.9$	yes	0.0979 (axisymmetric, $\bar k = -0.10$)	stable	0.0917
as the first, $n_{\bar\nu_e}/n_{\nu_e} = 0.3$	no	none	stable	stable

(the largest azimuthal growth rates are 0.0200 and 0.0050 for the first two cases). Three independent evaluations agree to 1.2×10^{-14}: the analytic growth of the mode $k_\mathrm{f} = \mathcal{I}_1$ against the root of the dispersion relation at $\bar k = 0$, and the eigenvalues of the discretised homogeneous equation (300 nodes) against the root at $\bar k = -\mathcal{I}_1$. With an ELN crossing the gas is unstable, without one it is not, as Morinaga's theorem (2022) states; the second case shows that a stable homogeneous mode does not exclude growing inhomogeneous modes. Near the neutrinosphere, $n_{\nu_e} = 10^{32}$ to 10^{33} cm^{-3} ($\sqrt{2}G_\mathrm{F}n_{\nu_e} = 1.27\times10^{-5}$ to 1.27×10^{-4} eV), the first case grows by the factor e in 5.9×10^{-10} to 5.9×10^{-11} s over 17.8 to 1.78 cm: a million times faster than the hydrodynamics, which is why explosion models treat the conversion as instantaneous.

Flavour conversion and the explosions. The charged-current heating behind the shock is, schematically, proportional to the sum over ν_e and $\bar\nu_e$ of the number flux times the mean squared energy, with the mean squared

energy $\frac{5}{4}$ times the squared mean energy (spectra with the pinching parameter 3). Full three-flavour equipartition reduces it to 0.884 of its value, and pair conversion $\nu_e\bar{\nu}_e \to \nu_x\bar{\nu}_x$, which conserves the electron lepton number until $\bar{\nu}_e$ is equipartitioned, to 0.760 (DERIVED, schematic): the heavy-lepton neutrinos are more energetic but fewer and do not heat through charged currents. Gogilashvili and Tamborra (2026, arXiv:2605.16504) simulated 195 progenitors of 9 to 120 solar masses (GR1D, M1 transport of ν_e, $\bar{\nu}_e$, ν_x in 18 energy bins, SFHo equation of state) with instantaneous flavour equipartition, conserving the electron lepton number and the number and momentum of the neutrinos in each energy bin, imposed at 0.02 s after bounce where the baryon density is below a threshold (STANDARD, cited):

threshold density (g/cm^3)	no conversion	10^9	10^{10}	10^{11}	10^{13}
failed explosions	25.6 %	50.8 %	77.4 %	93.3 %	96.4 %
weighted with the Salpeter mass function	27.0 %	47.6 %	61.9 %	73.9 %	88.3 %

The effect is strongest between 16 and 30 solar masses (easing the red-supergiant and supernova-rate problems), and the successful explosions leave lighter neutron stars (gravitational masses around 1.2 to 1.4 solar masses); the authors caution that the large fractions for deep conversion are in tension with observations and call for dynamical, multi-dimensional treatments.

8.3.7 Dirac or Majorana

UM-2 gives the neutrinos Dirac masses (the lepton-doublet zero mode mixed at the tip with three right-handed singlets localised at $|c_R| = 31$ to 35, tip couplings of order one). The alternative without right-handed neutrinos is a Majorana mass from the gauge-invariant tip operator

$$S_{\mathrm{M}} = -\frac{1}{2}\int_{y=-L} \mathrm{d}^7X_{\mathrm{tip}}\sqrt{|g_{\mathrm{tip}}|}\sum_{i,j=1}^{3}(\mathcal{K}_{\mathrm{M}})_{ij}\,\tilde{\nu}_i\tilde{\nu}_j + \mathrm{h.c.}, \qquad \tilde{\nu}_i = \left(\Sigma^{\dagger}\ell_i\right)_1$$

Symbols in this equation.

- S_{M}: The action of the gauge-invariant Majorana mass operator on the tip, the alternative to the Dirac neutrinos of UM-2 without right-handed neutrinos (not part of UM-2). Unit: dimensionless (an action, $\hbar = 1$). Status: INVENTED.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- X_{tip}: The seven coordinates $x_1, \ldots, x_7$ of the tip $y = -L$ of the hidden direction; $\mathrm{d}^7X_{\mathrm{tip}}$ is their coordinate volume element. Unit: length (each coordinate). Status: DERIVED.
- g_{tip}: The metric induced on the seven-dimensional tip $y = -L$ (the metric g_{MN} restricted to the tip directions $x_1, \ldots, x_7$); $|g_{\mathrm{tip}}|$ is the absolute value of its determinant. Unit: dimensionless. Status: DERIVED.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- j: A second generation index ($j = 1, 2, 3$). Status: STANDARD.
- $\mathcal{K}_{\mathrm{M}}^{(i)}$: The dimensionless Majorana tip coefficients of the alternative tip operator S_{M}: the complex symmetric 3×3 matrix $(\mathcal{K}_{\mathrm{M}})_{ij}$ (generation indices i, j; dimensionless because the tip operator is integrated over seven dimensions and $\tilde{\nu}_i\tilde{\nu}_j$ has mass dimension 7); its eigenvalues $\mathcal{K}_{\mathrm{M}}^{(i)}$ give the four-dimensional masses $m_i = \mathcal{K}_{\mathrm{M}}^{(i)}\,\mathcal{Z}(c_L)^2/(fL)$ to first order in the tip value of the doublet zero mode. Unit: dimensionless. Value: 5.63×10^{-13}, 4.90×10^{-12}, 2.82×10^{-11} for $i = 1, 2, 3$. Status: DERIVED.

- $\tilde{\nu}_i$: The neutral upper component $(\Sigma^\dagger \ell_i)_1$ of the lepton doublet of generation i dressed with the tip sigma field: a left-handed field that is neutral under $SU(2)_L \times U(1)_Y$; $\tilde{\nu}_i\tilde{\nu}_j$ is the Lorentz-invariant (Majorana) product of two such fields. Unit: $\mathrm{GeV}^{7/2}$ (an eight-dimensional spinor field at the tip). Status: INVENTED.
- Σ: The tip sigma field: a unitary 2×2 matrix field on the tip ($\Sigma^\dagger\Sigma = \mathbf{1}$) that replaces the Higgs doublet; its three modes are the longitudinal W^+, W^- and Z, and $\Sigma = \mathbf{1}$ is the unitary gauge. Unit: dimensionless. Status: INVENTED.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- ℓ_i, $\bar{\ell}_i$: The lepton doublet of the generation i (left-handed neutrino and charged lepton) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.

(the upper component $\tilde{\nu}_i$ of $\Sigma^\dagger\ell_i$ is a left-handed field neutral under $SU(2)_L \times U(1)_Y$, because $\Sigma \to U_L \Sigma U_Y^\dagger$, and $\tilde{\nu}_i\tilde{\nu}_j$ is the Lorentz-invariant Majorana product of two such fields), whose four-dimensional mass is, to first order in the tip value of the doublet zero mode,

$$m_i = \mathcal{K}_{\mathrm{M}}^{(i)} \frac{\mathcal{Z}(c_L)^2}{fL}, \qquad \mathcal{Z}(c_L)^2 = \frac{c_L\, \mathrm{e}^{-c_L}}{\sinh c_L} = 8.038 \times 10^{-5}, \qquad \frac{1}{fL} = 22.1 \text{ TeV}$$

Symbols in this equation.

- m, m_s, m_i: A four-dimensional mass: m the mass of a mode (the eigenvalue of an x_0 problem) or of a fermion; m_s the mass of the species s; m_i the mass of the neutrino mass state i. Unit: GeV. Status: DERIVED.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- $\mathcal{K}_{\mathrm{M}}^{(i)}$: The dimensionless Majorana tip coefficients of the alternative tip operator S_{M}: the complex symmetric 3×3 matrix $(\mathcal{K}_{\mathrm{M}})_{ij}$ (generation indices i, j; dimensionless because the tip operator is integrated over seven dimensions and $\tilde{\nu}_i\tilde{\nu}_j$ has mass dimension 7); its eigenvalues $\mathcal{K}_{\mathrm{M}}^{(i)}$ give the four-dimensional masses $m_i = \mathcal{K}_{\mathrm{M}}^{(i)}\, \mathcal{Z}(c_L)^2/(fL)$ to first order in the tip value of the doublet zero mode. Unit: dimensionless. Value: 5.63×10^{-13}, 4.90×10^{-12}, 2.82×10^{-11} for $i = 1, 2, 3$. Status: DERIVED.
- $\mathcal{Z}(c)$: The tip amplitude of a normalised zero mode with the localisation c, $\mathcal{Z}(c)^2 = c\,\mathrm{e}^{-c}/\sinh c$. Unit: dimensionless. Status: DERIVED.
- c, c_L, c_R: The localisation of a zero mode, $c = f\mu L$ (bulk mass times the proper depth); c_L of a doublet (localised at the fold for $c_L > 0$), c_R of a singlet (localised at the fold for $c_R < 0$). Unit: dimensionless. Value: doublets $c_L = 3.38230$ (quarks) and 5.95288 (leptons); singlets c_R: up -15.4300, down -14.6258, strange -11.5158, charm -8.7683, bottom -7.5005, top -3.3823, electron -14.5539, muon -8.9809, tau -5.9529, neutrinos -35.0452, -32.8481, -31.0702 (fixed by the measured masses). Status: DERIVED.
- L: The chirality label L: the left-handed (doublet) zero mode, or the left-handed projection $P_L = (1 + \Gamma_4)/2$. Status: STANDARD.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

(DERIVED): the measured masses need $\mathcal{K}_{\mathrm{M}}^{(i)} = 5.63 \times 10^{-13}$, 4.90×10^{-12}, 2.82×10^{-11}, a hierarchy that the x_0 mechanism does not explain, whereas the Dirac masses come from the localisation with couplings of order one: the mechanism favours Dirac neutrinos. Neutrinoless double-beta decay would have

$$m_{\beta\beta} = \Big| \sum_{i=1}^{3} U_{ei}^2\, m_i\, \mathrm{e}^{\mathrm{i}\varphi_i^{\mathrm{M}}} \Big|, \qquad 0.84 \text{ meV} \le m_{\beta\beta} \le 4.41 \text{ meV}$$

Symbols in this equation.

- $m_{\beta\beta}$: The effective Majorana mass of neutrinoless double-beta decay, $|\sum_i U_{ei}^2 m_i \mathrm{e}^{\mathrm{i}\varphi_i^{\mathrm{M}}}|$. Unit: eV. Value: 0.84 to 4.41 meV over all Majorana phases. Status: DERIVED.

- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- U: The PMNS matrix, the unitary 3×3 mixing matrix of the leptons, $\nu_\alpha = \sum_i U_{\alpha i}\nu_i$ (rows $\alpha = e, \mu, \tau$, columns the mass states i); in UM-2 $U = \mathcal{K}_e^\dagger \mathcal{K}_\nu$. Unit: dimensionless. Value: $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$, $\delta_{\mathrm{CP}} = 212°$ (NuFIT 6.0). Status: STANDARD.
- e: The electron flavour as a value of a flavour index ($\delta_{\alpha e}$, U_{e2}, V_e, $\mathcal{P}_{ee}$). Status: STANDARD.
- m, m_s, m_i: A four-dimensional mass: m the mass of a mode (the eigenvalue of an x_0 problem) or of a fermion; m_s the mass of the species s; m_i the mass of the neutrino mass state i. Unit: GeV. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- φ_i^{M}: The Majorana phase of the neutrino mass state i in the Majorana alternative (a free parameter; all values are scanned). Unit: dimensionless (radians). Status: ASSUMED.

for all Majorana phases φ_i^{M} ($|U_{ei}|^2 m_i = 0.677$, 2.624, 1.111 meV; the extremes $\sum_i |U_{ei}|^2 m_i$ and twice the largest term minus the sum agree with a scan of the phases), below the reach of present experiments; the non-relativistic CNB would be captured twice as often as for Dirac neutrinos (both helicities active); the flavour conversion is the same.

8.4 The check of step 5

In its own equations.

- *The flavour evolution.* With the x_0 masses and the stored PMNS matrix the evolution operators are unitary to 1.3×10^{-15}; the exact vacuum evolution reproduces the analytic sum formula to 7.1×10^{-12} (with the masses of the roots, which differ from the stored ones by the fit residual; 3.9×10^{-15} with the same masses), CPT holds to 2.2×10^{-16} and the T-odd difference to 5.9×10^{-14}; the stored T2K, NOvA and DUNE probabilities and the adiabatic solar probabilities are reproduced to 4.9×10^{-13} and 4.4×10^{-13} (DERIVED).
- *The onset of the mixing.* The thermal potential equals the vacuum term at the stored temperatures to 8.7×10^{-12} (DERIVED).
- *The decoupling.* N_{eff} of instantaneous decoupling without QED, recomputed independently from the separate entropy conservation of the $e^\pm\gamma$ plasma with exact Fermi-Dirac integrals, equals the stored 3.00018038904 to 2.7×10^{-12}, and the method reproduces the six published values of the same method to 4.6×10^{-4} (DERIVED).
- *The fast flavour instability.* The moments of the distributions, the analytic growth and the eigenvalues of the discretised homogeneous equation, recomputed, agree with the stored values to 3.3×10^{-13} (in units of $\sqrt{2} G_{\mathrm{F}} n_{\nu_e}$) (DERIVED).
- *The CNB.* The number densities, energy densities, mean velocities, flavour contents and $\Omega_\nu h^2_{100}$, recomputed with the x_0 masses by an independent adaptive quadrature, agree with the stored values to 2.9×10^{-12} (DERIVED).

In concert with the other field equations.

- *The masses and the mixing of step 2.* The neutrino masses of the tip conditions and their splittings are those of the flavour evolution, the CNB and the CMB computation to 5.7×10^{-12}; the PMNS matrix of the flavour evolution and of the flavour content of the CNB is the neutrino tip matrix to 8.2×10^{-13} (DERIVED).
- *N_{eff} and $\Omega_\nu h^2_{100}$ in the CMB.* The CMB computation of step 6 uses $N_{\mathrm{eff}} = 3.0432$ plus the freeze-in radiation 0.0010 of step 9 exactly (5×10^{-16}), and its neutrino density today, 6.4511×10^{-4}, equals the CNB density times $(N_{\mathrm{eff}}/3)^{3/4}$, 6.4510×10^{-4}, to 1.9×10^{-5} (the scaling is exact only for non-relativistic states; the semi-relativistic ν_1 leaves this) (DERIVED).
- *The heating factor and the radiation today.* The heating factor F_{h} of the decoupling is the one used in steps 4 and 9 to 5.8×10^{-7}; the radiation density after decoupling in the thermal history of step 9, which decouples the neutrinos instantaneously at 2 MeV ($g_\rho = 3.3626$ today), differs from that of the CMB computation with $N_{\mathrm{eff}} = 3.0442$ (3.3827) by 0.59 %, a documented approximation of step 9 that enters only the entropy today (taken as the standard $g_s = 43/11$) and the nucleosynthesis environment of step 8 (effects below 10^{-58}) (DERIVED).
- *The Z width of step 4.* $N_\nu = 3$, the number of neutrinos of the decoupling, is the measured invisible width of the Z.

8.5 The status of step 5

DERIVED: the flavour evolution in vacuum, matter, the Sun, at accelerators and in the supernova envelope, the onset of the mixing, the decoupling and the computed $N_{\rm eff}$, the CNB, the fast-flavour growth rates, the schematic heating ratios, the Majorana alternative. ASSUMED: the lightest mass and the ordering, the solar and supernova profiles, the neutrinospheres. STANDARD: the splittings, the mixing angles and the phase (NuFIT 6.0), the adopted precision $N_{\rm eff} = 3.0432$, the cited supernova simulations. OPEN: the absolute mass scale and the ordering (chosen here); the collective flavour evolution beyond the linear analysis.

9 Step 6: the CMB, the CNB and the BAO

On the stabilised block the four-dimensional effective theory is general relativity (step 1) with photons, baryons, the three neutrino mass eigenstates (step 5), the dark matter (the odd quanta of Ψ_A and Ψ_B, steps 2 and 3), the fold bands of dirac16complex made by freeze-in (massless free-streaming radiation with $\Delta N_{\rm eff} = 0.001$, step 9) and the four-dimensional cosmological term Λ_4. Step 6 solves the order-1, $n = 0$ term of the metric series: the linear Einstein-Boltzmann equations that give the CMB, the CNB and the BAO.

9.1 The field equations of step 6

The Einstein-Lovelock equations, whose order-1, $n = 0$ part this step solves through the four-dimensional gravity of the stabilised block (step 1):

$$\begin{aligned} &\sum_{p=1}^{3} \alpha_{(p)}\, {\mathcal{E}_{(p)}}^{M}{}_{N} + \Lambda\, \delta^{M}_{N} = \kappa\, {\mathcal{T}}^{M}{}_{N}, \\ &{\mathcal{E}_{(p)}}^{M}{}_{N} = -\frac{1}{2^{p+1}}\, \delta^{M M_1 N_1 \cdots M_p N_p}_{N P_1 Q_1 \cdots P_p Q_p}\, R^{P_1 Q_1}{}_{M_1 N_1} \cdots R^{P_p Q_p}{}_{M_p N_p} \end{aligned} \tag{FE-1}$$

Symbols in (FE-1).

- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic). Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)} H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- ${\mathcal{E}_{(p)}}^{M}{}_{N}$: The Lovelock tensor of order p, the metric variation of $\sqrt{|g|}\mathcal{L}_{(p)}$: $\mathcal{E}_{(1)}$ is the Einstein tensor $R^{M}{}_{N} - \frac{1}{2}\delta^{M}_{N} R$; every $\mathcal{E}_{(p)}$ is divergence free. Unit: 1/length^{2p}. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -10H^2$. Unit: GeV^2. Value: $-10H^2$. Status: AUTHOR.
- δ: The Kronecker delta: with two indices $\delta^{M}_{N} = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{\alpha e}$, δ_{l0}, δ^{I3}); with upper and lower index strings the generalised Kronecker delta $\delta^{M_1 \cdots M_n}_{N_1 \cdots N_n}$, the determinant of the $n \times n$ matrix of Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_{\rm N}\mathcal{G}(K_*) = 2.328 \times 10^{-36}$ GeV^{-2}. Status: AUTHOR.
- ${\mathcal{T}}^{M}{}_{N}$: The energy-momentum tensor of all matter (every field except gravity, the stabiliser included), the source of gravity; defined by the change of the matter action under a change of the metric. Unit: GeV^8. Status: STANDARD.
- P: A coordinate index of eight dimensions (also $P_1, P_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Q: A coordinate index of eight dimensions (also $Q_1, Q_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $R^{MN}{}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor, with two indices the Ricci tensor $R_{MN} = R^{P}{}_{MPN}$, without indices the Ricci (scalar) curvature $R = g^{MN}R_{MN}$; $R^{ab}{}_{ab}$ (no sum) is the sectional curvature of the coordinate plane (a, b). Unit: 1/length^2. Status: STANDARD.

$$
\begin{aligned}
&G^{(4)}_{\mu\nu} + \Lambda_4\, g^{(4)}_{\mu\nu} = 8\pi G_{\mathrm{N}}\, \mathcal{T}^{(4)}_{\mu\nu}, \qquad 8\pi G_{\mathrm{N}} = \frac{\kappa}{\mathcal{G}(K_*)\, V_\perp}, \qquad V_\perp = \frac{2 f^4 v_{\mathrm{t}} \bigl(1 - \mathrm{e}^{-3HL}\bigr)}{3H}, \\
&\mathcal{T}^{(4)}_{\mu\nu} = \int \mathrm{d}x_0\, \mathrm{d}x_5\, \mathrm{d}x_6\, \mathrm{d}x_7\, \sqrt{|g_\perp|}\, \mathcal{T}_{\mu\nu}, \qquad g_3^2 = \frac{\hat{g}_3^2}{V_\perp}, \qquad g_2^2 = \frac{\hat{g}_2^2}{V_\perp}, \qquad g_1^2 = \frac{\hat{g}_1^2}{V_\perp}
\end{aligned}
\tag{FE-49}
$$

Symbols in (FE-49).

- $G^{(4)}_{\mu\nu}$: The Einstein tensor of the four-dimensional metric, $R^{(4)}_{\mu\nu} - \frac{1}{2} g^{(4)}_{\mu\nu} R^{(4)}$. Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index running over $x_1, \ldots, x_4$ (as μ). Status: STANDARD.
- Λ_4: The four-dimensional cosmological constant of the effective theory on the block, $\Lambda_4 = [6H^2 K(1 - 2w_{(2)}K) + \Lambda + \kappa\rho_{\mathrm{stab}}]/\mathcal{G}(K)$; zero at the static point, the observed dark energy a tuned residual. Unit: GeV^2. Value: 0 at the static point; observed $\Lambda_4/H^2 \sim 10^{-92}$. Status: DERIVED.
- $g^{(4)}_{\mu\nu}$: The four-dimensional metric of the observable coordinates $x_1, \ldots, x_4$ (the metric of the effective four-dimensional general relativity on the stabilised block). Unit: dimensionless. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_{N}: Newton's constant of four-dimensional gravity, $8\pi G_{\mathrm{N}} = \kappa/(\mathcal{G}(K_*)V_\perp)$. Unit: GeV^{-2}. Value: 6.709×10^{-39} GeV^{-2} ($M_{\mathrm{Pl}} = G_{\mathrm{N}}^{-1/2} = 1.22091 \times 10^{19}$ GeV). Status: STANDARD.
- $\mathcal{T}^{(4)}_{\mu\nu}$: The four-dimensional energy-momentum tensor: $\mathcal{T}_{\mu\nu}$ integrated over the block (both halves of a period of x_0 and the extra times). Unit: GeV^4. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_{\mathrm{N}}\mathcal{G}(K_*) = 2.328 \times 10^{-36}$ GeV^{-2}. Status: AUTHOR.
- $\mathcal{G}(K)$: The coefficient of the four-dimensional Einstein-Hilbert term on a static block, $\mathcal{G}(K) = 1 - 24w_{(2)}K + 72w_{(3)}K^2 = (K^2 - 12K + 16)/16$; four-dimensional gravity is attractive for $\mathcal{G} > 0$. Unit: dimensionless. Value: $\mathcal{G}(K_*) = 20/3 + \sqrt{51} = 13.8081$. Status: DERIVED.
- K_*: The block variable of the stabilised block, $K_* = 12 + \frac{4}{3}\sqrt{51}$ (the upper root K_+). Unit: dimensionless. Value: 21.5219. Status: DERIVED.
- $V_\perp$: The proper volume of the block per unit four-volume (both halves of a period of x_0 and the extra times), $V_\perp = 2f^4 v_{\mathrm{t}}(1 - \mathrm{e}^{-3HL})/(3H)$. Unit: GeV^{-4}. Status: DERIVED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- v_{t}: The coordinate volume of the compact extra times x_5, x_6, x_7. Unit: GeV^{-3}. Value: not fixed; $v_{\mathrm{t}} = H^{-3}$ in the quoted conversions. Status: ASSUMED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- $g_\perp$: The metric of the four transverse directions x_0, x_5, x_6, x_7 (the block); $|g_\perp|$ is the absolute value of its determinant. Unit: dimensionless. Status: DERIVED.
- $\mathcal{T}^M{}_N$: The energy-momentum tensor of all matter (every field except gravity, the stabiliser included), the source of gravity; defined by the change of the matter action under a change of the metric. Unit: GeV^8.

Status: STANDARD.

- g_3: The four-dimensional strong coupling ($g_3^2 = 4\pi\alpha_3$). Unit: dimensionless. Value: $g_3^2 = 0.7786$ at $2M_{\mathrm{DM}}$. Status: DERIVED.
- $\hat{g}_3$: The eight-dimensional SU(3) coupling (it multiplies the gluon kinetic term as $1/\hat{g}_3^2$); $g_3^2 = \hat{g}_3^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- g_2: The four-dimensional SU(2) coupling ($g_2^2 = 4\pi\alpha_2$). Unit: dimensionless. Value: $g_2^2 = 0.3781$ at $2M_{\mathrm{DM}}$. Status: DERIVED.
- $\hat{g}_2$: The eight-dimensional SU(2) coupling; $g_2^2 = \hat{g}_2^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- g_1: The four-dimensional hypercharge coupling ($g_1^2 = 4\pi\alpha_1$, Standard-Model normalisation without the GUT factor). Unit: dimensionless. Value: $g_1^2 = 0.1380$ at $2M_{\mathrm{DM}}$. Status: DERIVED.
- $\hat{g}_1$: The eight-dimensional $\mathrm{U}(1)_Y$ coupling; $g_1^2 = \hat{g}_1^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.

The order-1, $n = 0$ term of the metric in the Newtonian gauge, in eight dimensions and in the four-dimensional conformal time:

$$g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N = \mathrm{e}^{2a_4}\Big[\big(1-2\Psi_{\mathrm{N}}\big)\big(\mathrm{d}x_1^2+\mathrm{d}x_2^2+\mathrm{d}x_3^2\big)+\mathrm{d}\vec{x}\cdot h^{\mathrm{TT}}\cdot\mathrm{d}\vec{x}\Big]-\big(1+2\Phi\big)\,\mathrm{d}x_4^2 + f^2\big(1+2\sigma_{\mathrm{r}}\big)\Big[\mathrm{d}y^2-\mathrm{e}^{2Hy}\big(\mathrm{d}x_5^2+\mathrm{d}x_6^2+\mathrm{d}x_7^2\big)\Big] \tag{FE-88}$$

Symbols in (FE-88).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\ldots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- Ψ_{N}: The curvature potential of the Newtonian gauge (the perturbation of the 3-space metric, $1-2\Psi_{\mathrm{N}}$); not related to the field Ψ. Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space (also the symmetry axis of the fast-flavour analysis). Unit: length. Status: AUTHOR.
- $\vec{x}$: The position in ordinary 3-space, $\vec{x} = (x_1, x_2, x_3)$; $\mathrm{d}\vec{x}^{\,2} = \mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2$. Unit: length. Status: STANDARD.
- h^{TT}: The transverse-traceless (tensor) part of the first-order metric of 3-space (gravitational waves). Unit: dimensionless. Value: an input (tensor-to-scalar ratio below 0.036). Status: STANDARD.
- Φ: The lapse potential of the Newtonian gauge (the perturbation of the time component, $1+2\Phi$). Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4}b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- σ_{r}: The radion perturbation: the first-order change of the block scale, $f^2 \to f^2(1+2\sigma_{\mathrm{r}})$. Unit: dimensionless. Value: at most 1.1×10^{-34} (static response). Status: DERIVED.

- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.

$$g^{(4)}_{\mu\nu}\,\mathrm{d}X^\mu \mathrm{d}X^\nu = \mathrm{e}^{2a_4}\Big[-\big(1+2\Phi\big)\,\mathrm{d}\tau_\mathrm{c}^2 + \big(1-2\Psi_\mathrm{N}\big)\,\mathrm{d}\vec{x}^{\,2}\Big], \qquad \mathrm{d}x_4 = \mathrm{e}^{a_4}\,\mathrm{d}\tau_\mathrm{c} \tag{FE-89}$$

Symbols in (FE-89).

- $g^{(4)}_{\mu\nu}$: The four-dimensional metric of the observable coordinates $x_1, \ldots, x_4$ (the metric of the effective four-dimensional general relativity on the stabilised block). Unit: dimensionless. Status: DERIVED.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index running over $x_1, \ldots, x_4$ (as μ). Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- Φ: The lapse potential of the Newtonian gauge (the perturbation of the time component, $1 + 2\Phi$). Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- τ_c: The conformal time, $\mathrm{d}x_4 = \mathrm{e}^{a_4}\mathrm{d}\tau_\mathrm{c}$; primes denote $\mathrm{d}/\mathrm{d}\tau_\mathrm{c}$. Unit: length. Status: STANDARD.
- Ψ_N: The curvature potential of the Newtonian gauge (the perturbation of the 3-space metric, $1 - 2\Psi_\mathrm{N}$); not related to the field Ψ. Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- $\vec{x}$: The position in ordinary 3-space, $\vec{x} = (x_1, x_2, x_3)$; $\mathrm{d}\vec{x}^{\,2} = \mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2$. Unit: length. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.

The linearised Einstein equations and the equations of every species of UM-2:

$$\begin{aligned}
&k^2\Psi_\mathrm{N} + 3a_4'\big(\Psi_\mathrm{N}' + a_4'\Phi\big) = -4\pi G_\mathrm{N}\,\mathrm{e}^{2a_4}\,\Delta\rho, \qquad k^2\big(\Psi_\mathrm{N}' + a_4'\Phi\big) = 4\pi G_\mathrm{N}\,\mathrm{e}^{2a_4}\big(\rho_\mathrm{tot} + p_\mathrm{tot}\big)\theta_\mathrm{tot},\\
&\Psi_\mathrm{N}'' + a_4'\big(\Phi' + 2\Psi_\mathrm{N}'\big) + \big(2a_4'' + a_4'^2\big)\Phi + \frac{k^2}{3}\big(\Psi_\mathrm{N} - \Phi\big) = 4\pi G_\mathrm{N}\,\mathrm{e}^{2a_4}\,\Delta p,\\
&k^2\big(\Psi_\mathrm{N} - \Phi\big) = 12\pi G_\mathrm{N}\,\mathrm{e}^{2a_4}\big(\rho_\mathrm{tot} + p_\mathrm{tot}\big)\sigma_\mathrm{an}
\end{aligned} \tag{FE-90}$$

Symbols in (FE-90).

- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- Ψ_N: The curvature potential of the Newtonian gauge (the perturbation of the 3-space metric, $1 - 2\Psi_\mathrm{N}$); not related to the field Ψ. Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- $'$: The prime on a function of the conformal time τ_c is its derivative $\mathrm{d}/\mathrm{d}\tau_\mathrm{c}$ (it is used only in the cosmological perturbation equations; two primes are the second derivative). Status: STANDARD.

- Φ: The lapse potential of the Newtonian gauge (the perturbation of the time component, $1 + 2\Phi$). Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_N: Newton's constant of four-dimensional gravity, $8\pi G_\mathrm{N} = \kappa/(\mathcal{G}(K_*)V_\perp)$. Unit: GeV^{-2}. Value: 6.709×10^{-39} GeV^{-2} ($M_\mathrm{Pl} = G_\mathrm{N}^{-1/2} = 1.22091 \times 10^{19}$ GeV). Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_eff an addition to N_eff, ΔS_m the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- ρ_tot, p_tot: The total background energy density and pressure of four-dimensional cosmology (all species). Unit: GeV^4. Status: STANDARD.
- θ_tot: The velocity divergence of the total matter, weighted with $\rho + p$ of each species. Unit: 1/length. Status: STANDARD.
- p: A pressure: $p_M = \mathcal{T}^M{}_M$ (no sum) for the direction $M \neq x_4$, p_s the pressure of the species s; Δp the pressure perturbation in the cosmological perturbation equations; in the Feynman-Hellmann relation the pressure of a gauge plasma. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- σ_an: The anisotropic stress (shear) of the total matter (the quantity σ of Ma and Bertschinger), weighted with $\rho + p$. Unit: dimensionless. Status: STANDARD.

$$\delta'_\mathrm{DM} = -\theta_\mathrm{DM} + 3\Psi'_\mathrm{N}, \qquad \theta'_\mathrm{DM} = -a'_4\,\theta_\mathrm{DM} + k^2\Phi \tag{FE-91}$$

Symbols in (FE-91).

- δ_DM, θ_DM: The density contrast and the velocity divergence of the dark matter. Unit: dimensionless; 1/length. Status: STANDARD.
- $'$: The prime on a function of the conformal time τ_c is its derivative $\mathrm{d}/\mathrm{d}\tau_\mathrm{c}$ (it is used only in the cosmological perturbation equations; two primes are the second derivative). Status: STANDARD.
- Ψ_N: The curvature potential of the Newtonian gauge (the perturbation of the 3-space metric, $1 - 2\Psi_\mathrm{N}$); not related to the field Ψ. Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- Φ: The lapse potential of the Newtonian gauge (the perturbation of the time component, $1 + 2\Phi$). Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.

$$\delta'_\mathrm{b} = -\theta_\mathrm{b} + 3\Psi'_\mathrm{N}, \qquad \theta'_\mathrm{b} = -a'_4\,\theta_\mathrm{b} + c_\mathrm{s}^2 k^2\delta_\mathrm{b} + R_\mathrm{b}\,\mathrm{e}^{a_4} n_\mathrm{e}\sigma_\mathrm{T}\big(\theta_\gamma - \theta_\mathrm{b}\big) + k^2\Phi, \qquad R_\mathrm{b} = \frac{4\rho_\gamma}{3\rho_\mathrm{b}} \tag{FE-92}$$

Symbols in (FE-92).

- δ_b, θ_b: The density contrast and the velocity divergence of the baryons (nucleons and electrons). Unit: dimensionless; 1/length. Status: STANDARD.
- $'$: The prime on a function of the conformal time τ_c is its derivative $\mathrm{d}/\mathrm{d}\tau_\mathrm{c}$ (it is used only in the cosmological perturbation equations; two primes are the second derivative). Status: STANDARD.
- Ψ_N: The curvature potential of the Newtonian gauge (the perturbation of the 3-space metric, $1 - 2\Psi_\mathrm{N}$); not related to the field Ψ. Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.

- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- c_s: The sound speed of the baryon fluid. Unit: dimensionless ($\mathrm{c} = 1$). Status: STANDARD.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- R_b: The photon-to-baryon momentum ratio of the Thomson drag, $R_\mathrm{b} = 4\rho_\gamma/(3\rho_\mathrm{b})$. Unit: dimensionless. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- n_e: The number density of electrons (net of positrons); in the Thomson terms of the photon and baryon equations only free (ionised) electrons count. Unit: GeV^3. Status: STANDARD.
- σ_T: The Thomson cross-section. Unit: GeV^{-2}. Value: $6.6524587 \times 10^{-29}$ m^2. Status: STANDARD.
- δ_γ, θ_γ: The density contrast and the velocity divergence of the photons. Unit: dimensionless; 1/length. Status: STANDARD.
- Φ: The lapse potential of the Newtonian gauge (the perturbation of the time component, $1 + 2\Phi$). Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- ρ_γ: The energy density of the photons, $\rho_\gamma = \pi^2 T_\gamma^4/15$. Unit: GeV^4. Status: STANDARD.
- ρ_b: The energy density of the baryons. Unit: GeV^4. Status: STANDARD.

$$\delta_\gamma' = -\frac{4}{3}\,\theta_\gamma + 4\Psi_\mathrm{N}', \qquad \theta_\gamma' = k^2\Big(\frac{1}{4}\,\delta_\gamma - \sigma_{\mathrm{an},\gamma}\Big) + k^2\Phi + \mathrm{e}^{a_4} n_\mathrm{e}\sigma_\mathrm{T}\big(\theta_\mathrm{b} - \theta_\gamma\big) \tag{FE-93}$$

Symbols in (FE-93).

- δ_γ, θ_γ: The density contrast and the velocity divergence of the photons. Unit: dimensionless; 1/length. Status: STANDARD.
- $'$: The prime on a function of the conformal time τ_c is its derivative $\mathrm{d}/\mathrm{d}\tau_\mathrm{c}$ (it is used only in the cosmological perturbation equations; two primes are the second derivative). Status: STANDARD.
- Ψ_N: The curvature potential of the Newtonian gauge (the perturbation of the 3-space metric, $1 - 2\Psi_\mathrm{N}$); not related to the field Ψ. Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- $\sigma_{\mathrm{an},\gamma}$: The anisotropic stress (shear) of the photons, $F_2^\gamma/2$. Unit: dimensionless. Status: STANDARD.
- Φ: The lapse potential of the Newtonian gauge (the perturbation of the time component, $1 + 2\Phi$). Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- n_e: The number density of electrons (net of positrons); in the Thomson terms of the photon and baryon equations only free (ionised) electrons count. Unit: GeV^3. Status: STANDARD.
- σ_T: The Thomson cross-section. Unit: GeV^{-2}. Value: $6.6524587 \times 10^{-29}$ m^2. Status: STANDARD.
- δ_b, θ_b: The density contrast and the velocity divergence of the baryons (nucleons and electrons). Unit: dimensionless; 1/length. Status: STANDARD.

$$\begin{aligned} &F_l^{\gamma\prime} = \frac{k}{2l+1}\big[l\,F_{l-1}^\gamma - (l+1)F_{l+1}^\gamma\big] - \mathrm{e}^{a_4} n_\mathrm{e}\sigma_\mathrm{T}\Big[F_l^\gamma - \frac{\delta_{l2}}{10}\big(F_2^\gamma + G_0^\gamma + G_2^\gamma\big)\Big] \quad (l \geq 2), \\ &F_0^\gamma = \delta_\gamma, \qquad F_1^\gamma = \frac{4\theta_\gamma}{3k}, \qquad F_2^\gamma = 2\sigma_{\mathrm{an},\gamma} \end{aligned} \tag{FE-94}$$

Symbols in (FE-94).

- F_l^γ: The multipole l of the photon temperature perturbation (the brightness function of Ma and Bertschinger). Unit: dimensionless. Status: STANDARD.
- l: The multipole (angular moment) number $l = 0, 1, 2, \ldots$ of a Boltzmann hierarchy or of an angular moment $\int u^l \cdots$; as a coefficient it is the same number. Status: STANDARD.
- $'$: The prime on a function of the conformal time τ_c is its derivative $\mathrm{d}/\mathrm{d}\tau_\mathrm{c}$ (it is used only in the cosmological perturbation equations; two primes are the second derivative). Status: STANDARD.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- n_e: The number density of electrons (net of positrons); in the Thomson terms of the photon and baryon equations only free (ionised) electrons count. Unit: GeV^3. Status: STANDARD.
- σ_T: The Thomson cross-section. Unit: GeV^{-2}. Value: $6.6524587 \times 10^{-29}$ m^2. Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta_N^M = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{\alpha e}$, δ_{l0}, δ^{I3}); with upper and lower index strings the generalised Kronecker delta $\delta_{N_1 \cdots N_n}^{M_1 \cdots M_n}$, the determinant of the $n \times n$ matrix of Kronecker deltas $\delta_{N_b}^{M_a}$ (+1 or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- G_l^γ: The multipole l of the photon polarisation perturbation. Unit: dimensionless. Status: STANDARD.
- δ_γ, θ_γ: The density contrast and the velocity divergence of the photons. Unit: dimensionless; 1/length. Status: STANDARD.
- $\sigma_{\mathrm{an},\gamma}$: The anisotropic stress (shear) of the photons, $F_2^\gamma/2$. Unit: dimensionless. Status: STANDARD.

$$G_l^{\gamma\prime} = \frac{k}{2l+1}\left[l\, G_{l-1}^\gamma - (l+1)G_{l+1}^\gamma\right] + \mathrm{e}^{a_4} n_\mathrm{e} \sigma_\mathrm{T}\Big[- G_l^\gamma + \frac{1}{2}\big(F_2^\gamma + G_0^\gamma + G_2^\gamma\big)\Big(\delta_{l0} + \frac{\delta_{l2}}{5}\Big)\Big] \tag{FE-95}$$

Symbols in (FE-95).

- G_l^γ: The multipole l of the photon polarisation perturbation. Unit: dimensionless. Status: STANDARD.
- l: The multipole (angular moment) number $l = 0, 1, 2, \ldots$ of a Boltzmann hierarchy or of an angular moment $\int u^l \cdots$; as a coefficient it is the same number. Status: STANDARD.
- $'$: The prime on a function of the conformal time τ_c is its derivative $\mathrm{d}/\mathrm{d}\tau_\mathrm{c}$ (it is used only in the cosmological perturbation equations; two primes are the second derivative). Status: STANDARD.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- n_e: The number density of electrons (net of positrons); in the Thomson terms of the photon and baryon equations only free (ionised) electrons count. Unit: GeV^3. Status: STANDARD.
- σ_T: The Thomson cross-section. Unit: GeV^{-2}. Value: $6.6524587 \times 10^{-29}$ m^2. Status: STANDARD.
- F_l^γ: The multipole l of the photon temperature perturbation (the brightness function of Ma and Bertschinger). Unit: dimensionless. Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta_N^M = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{\alpha e}$, δ_{l0}, δ^{I3}); with upper and lower index strings the generalised Kronecker delta $\delta_{N_1 \cdots N_n}^{M_1 \cdots M_n}$, the determinant of the $n \times n$ matrix of Kronecker deltas $\delta_{N_b}^{M_a}$ (+1 or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.

$$\mathcal{N}_0^{(i)\prime} = -\frac{k_\nu k}{E_i^{\mathrm{c}}}\,\mathcal{N}_1^{(i)} - \Psi_{\mathrm{N}}'\,\frac{\mathrm{d}\ln f_{\mathrm{FD}}}{\mathrm{d}\ln k_\nu}, \qquad \mathcal{N}_1^{(i)\prime} = \frac{k_\nu k}{3E_i^{\mathrm{c}}}\big(\mathcal{N}_0^{(i)} - 2\mathcal{N}_2^{(i)}\big) - \frac{E_i^{\mathrm{c}} k}{3k_\nu}\,\Phi\,\frac{\mathrm{d}\ln f_{\mathrm{FD}}}{\mathrm{d}\ln k_\nu},$$
$$\mathcal{N}_l^{(i)\prime} = \frac{k_\nu k}{(2l+1)E_i^{\mathrm{c}}}\big[l\,\mathcal{N}_{l-1}^{(i)} - (l+1)\mathcal{N}_{l+1}^{(i)}\big] \quad (l \geq 2), \qquad E_i^{\mathrm{c}} = \sqrt{k_\nu^2 + \mathrm{e}^{2a_4} m_i^2} \tag{FE-96}$$

Symbols in (FE-96).

- $\mathcal{N}_l^{(i)}$: The multipole l of the perturbation of the distribution of the neutrino mass state i (a function of k, k_ν and τ_{c}; the quantity Ψ_l of Ma and Bertschinger). Unit: dimensionless. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- $'$: The prime on a function of the conformal time τ_{c} is its derivative $\mathrm{d}/\mathrm{d}\tau_{\mathrm{c}}$ (it is used only in the cosmological perturbation equations; two primes are the second derivative). Status: STANDARD.
- k_ν: The comoving momentum of a neutrino (the physical momentum times e^{a_4}). Unit: GeV. Status: STANDARD.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- E_i^{c}: The comoving energy of a neutrino of the mass state i, $E_i^{\mathrm{c}} = (k_\nu^2 + \mathrm{e}^{2a_4} m_i^2)^{1/2}$. Unit: GeV. Status: STANDARD.
- Ψ_{N}: The curvature potential of the Newtonian gauge (the perturbation of the 3-space metric, $1 - 2\Psi_{\mathrm{N}}$); not related to the field Ψ. Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- f_{FD}: The Fermi-Dirac distribution of the relic neutrinos, $f_{\mathrm{FD}} = 1/(\mathrm{e}^{k/T_\nu} + 1)$. Unit: dimensionless. Status: STANDARD.
- Φ: The lapse potential of the Newtonian gauge (the perturbation of the time component, $1 + 2\Phi$). Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- l: The multipole (angular moment) number $l = 0, 1, 2, \ldots$ of a Boltzmann hierarchy or of an angular moment $\int u^l \cdots$; as a coefficient it is the same number. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- m, m_s, m_i: A four-dimensional mass: m the mass of a mode (the eigenvalue of an x_0 problem) or of a fermion; m_s the mass of the species s; m_i the mass of the neutrino mass state i. Unit: GeV. Status: DERIVED.

$$a_4'^2 = \frac{8\pi G_{\mathrm{N}}}{3}\,\mathrm{e}^{2a_4}\rho_{\mathrm{tot}} + \frac{\Lambda_4}{3}\,\mathrm{e}^{2a_4}, \qquad T_\nu = \Big(\frac{4}{11}\Big)^{1/3} T_\gamma, \qquad P_{\mathcal{R}}(k) = A_{\mathrm{s}}\Big(\frac{k}{k_0}\Big)^{n_{\mathrm{s}}-1} \tag{FE-97}$$

Symbols in (FE-97).

- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- $'$: The prime on a function of the conformal time τ_{c} is its derivative $\mathrm{d}/\mathrm{d}\tau_{\mathrm{c}}$ (it is used only in the cosmological perturbation equations; two primes are the second derivative). Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_{N}: Newton's constant of four-dimensional gravity, $8\pi G_{\mathrm{N}} = \kappa/(\mathcal{G}(K_*)V_\perp)$. Unit: GeV^{-2}. Value: 6.709×10^{-39} GeV^{-2} ($M_{\mathrm{Pl}} = G_{\mathrm{N}}^{-1/2} = 1.22091 \times 10^{19}$ GeV). Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- ρ_{tot}, p_{tot}: The total background energy density and pressure of four-dimensional cosmology (all species). Unit: GeV^4. Status: STANDARD.

- Λ_4: The four-dimensional cosmological constant of the effective theory on the block, $\Lambda_4 = [6H^2K(1 - 2w_{(2)}K) + \Lambda + \kappa\rho_{\mathrm{stab}}]/\mathcal{G}(K)$; zero at the static point, the observed dark energy a tuned residual. Unit: GeV^2. Value: 0 at the static point; observed $\Lambda_4/H^2 \sim 10^{-92}$. Status: DERIVED.
- T_ν: The temperature of the relic neutrinos (one common temperature). Unit: GeV. Value: 1.9454 K today. Status: STANDARD.
- T_γ: The temperature of the photons (equal to the plasma temperature T; the subscript is written where the neutrinos have their own temperature). Unit: GeV. Value: 2.7255 K today. Status: STANDARD.
- $P_{\mathcal{R}}(k)$: The dimensionless power spectrum of the primordial curvature perturbation $\mathcal{R}$. Unit: dimensionless. Status: STANDARD.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- A_{s}: The amplitude of the primordial spectrum at k_0. Unit: dimensionless. Value: 2.099×10^{-9} ($\ln(10^{10}A_{\mathrm{s}}) = 3.044$, Planck 2018). Status: STANDARD.
- k_0: The pivot wavenumber of the primordial spectrum. Unit: 1/length. Value: 0.05 Mpc^{-1}. Status: STANDARD.
- n_{s}: The spectral index of the primordial spectrum. Unit: dimensionless. Value: 0.9649 (Planck 2018). Status: STANDARD.

The amplitude of the matter fluctuations:

$$\sigma_8^2 = \int_0^\infty \frac{\mathrm{d}k}{k}\,\frac{k^3 P_{\mathrm{m}}(k)}{2\pi^2}\,W_{\mathrm{th}}(kR_8)^2, \qquad W_{\mathrm{th}}(kR_8) = \frac{3\big[\sin(kR_8) - kR_8\cos(kR_8)\big]}{(kR_8)^3}, \qquad R_8 = 8\,h_{100}^{-1}\,\mathrm{Mpc} \quad \text{(FE-99)}$$

Symbols in (FE-99).

- σ_8: The root-mean-square linear matter fluctuation in spheres of radius R_8 today. Unit: dimensionless. Value: 0.8118. Status: DERIVED.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no upper bound. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- $P_{\mathrm{m}}(k)$: The linear matter power spectrum today. Unit: length^3. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- W_{th}: The Fourier transform of a spherical top-hat window. Unit: dimensionless. Status: STANDARD.
- R_8: The radius of the spheres of σ_8, $R_8 = 8\,h_{100}^{-1}$ Mpc. Unit: length. Status: STANDARD.
- h_{100}: The reduced Hubble constant, $\mathcal{H}_0/(100\ \mathrm{km\ s^{-1}\ Mpc^{-1}})$. Unit: dimensionless. Value: 0.6738. Status: STANDARD.

9.2 The approximation of step 6

- *First order in ϵ.* The omitted order 2 is $\Phi_{\mathrm{rms}} = 2.4 \times 10^{-5}$ of the kept order on the largest scales (step 10).
- *Only the $n = 0$ term.* The Kaluza-Klein gravitons and gauge bosons (≥ 99 TeV) and the radion (48 TeV) are absent at cosmological scales; the radion responds statically, by at most $\Delta \ln f = 1.1 \times 10^{-34}$ (step 8).
- *The standard numerical solution.* CAMB 2.0.5 at its default accuracy (about 10^{-3}), lensing included; the three neutrino masses are iterated to 7.8×10^{-12}.
- *The primordial spectrum is an input* (Planck 2018), not predicted by UM-2.

9.3 The solution of step 6

The equations were solved with CAMB 2.0.5 at the Planck 2018 values (STANDARD) $\Omega_{\rm b}h^2_{100} = 0.02237$, $\Omega_{\rm DM}h^2_{100} = 0.1200$, $100\,\theta_{\rm MC} = 1.04092$ (the angular size of the sound horizon at last scattering, held fixed, so that $\mathcal{H}_0$ is an output, in km/s/Mpc), the optical depth to reionisation 0.0544, $\ln(10^{10}A_{\rm s}) = 3.044$ and $n_{\rm s} = 0.9649$. In UM-2 the three mass states are evolved separately, each with the degeneracy $N_{\rm eff}/3 = 1.0144$ (the heated spectrum of step 5); CAMB gives each the density of a Fermi-Dirac gas of the temperature $T_\nu = (4/11)^{1/3}T_\gamma$ times the degeneracy, so its masses are the UM-2 masses times the degeneracy to the power $-1/4$, which reproduce the late-time density of the heated spectrum; CAMB finds them by inverting the full (relativistic) density from the total neutrino density and the mass fractions, which were therefore iterated (13 iterations) until the masses agree to 7.8×10^{-12} (DERIVED). The models are ΛCDM with the Planck 2018 parameters (one neutrino of 0.06 eV, $N_{\rm eff} = 3.044$: the CAMB defaults), UM-2 (the three mass states and the freeze-in fold bands, $N_{\rm eff} = 3.0432 + 0.0010 = 3.0442$), UM-2 without the freeze-in fold bands (the neutrino content of version 2, $N_{\rm eff} = 3.0432$), and two variants with a thermalised fold band of dirac16complex instead (two Weyl species, which UM-2 does not contain; steps 4 and 9), decoupled above the electroweak scale ($\Delta N_{\rm eff} = 0.0958$) or just above the QCD crossover ($\Delta N_{\rm eff} = 2(10.75/61.75)^{4/3} \times 1.0095 = 0.1963$) (DERIVED, 8 checks).

The BAO. The distances (flat space, $c = 1$) in units of the sound horizon at the drag epoch $r_{\rm d}$,

$$d_{\rm M}(z_{\rm red}) = \int_0^{z_{\rm red}} \frac{{\rm d}\tilde{z}_{\rm red}}{\mathcal{H}(\tilde{z}_{\rm red})}, \qquad d_{\rm H}(z_{\rm red}) = \frac{1}{\mathcal{H}(z_{\rm red})}, \qquad d_{\rm V}(z_{\rm red}) = \left(z_{\rm red}\, d_{\rm M}^2 d_{\rm H}\right)^{1/3},$$

Symbols in this equation.

- $d_{\rm M}$, $d_{\rm H}$, $d_{\rm V}$: The distances of the BAO analysis at the redshift $z_{\rm red}$ (flat space, $c = 1$): the comoving transverse distance $d_{\rm M}$, the Hubble distance $d_{\rm H} = 1/\mathcal{H}$ at that redshift, and the volume-averaged distance $d_{\rm V} = (z_{\rm red} d_{\rm M}^2 d_{\rm H})^{1/3}$ (written with a lower-case d because D_M is the covariant derivative). Unit: length (Mpc). Status: STANDARD.
- $z_{\rm red}$, $\tilde{z}_{\rm red}$: The cosmological redshift, $1 + z_{\rm red} = {\rm e}^{-a_4}$ with $a_4 = 0$ today (not $z = 6Hx_0$, the angle variable of the hidden direction); $\tilde{z}_{\rm red}$ is the redshift as the variable of an integral. Unit: dimensionless. Status: STANDARD.
- d: The differential: ${\rm d}X$ is an infinitesimal change of X, ${\rm d}X/{\rm d}Y$ the derivative of X with respect to Y, ${\rm d}^nX$ the volume element of n variables. Status: STANDARD.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_{\rm Pl}$). Unit: GeV. Status: DERIVED.

are compared with the DESI DR2 measurements (STANDARD, arXiv:2503.14738; the last column is the published correlation coefficient of the pair $d_{\rm M}/r_{\rm d}$, $d_{\rm H}/r_{\rm d}$):

tracer	$z_{\rm red}$	$d_{\rm V}/r_{\rm d}$	$d_{\rm M}/r_{\rm d}$	$d_{\rm H}/r_{\rm d}$	correlation
BGS	0.295	7.942 ± 0.075			
LRG1	0.510		13.588 ± 0.167	21.863 ± 0.425	-0.459
LRG2	0.706		17.351 ± 0.177	19.455 ± 0.330	-0.404
LRG3+ELG1	0.934		21.576 ± 0.152	17.641 ± 0.193	-0.416
ELG2	1.321		27.601 ± 0.318	14.176 ± 0.221	-0.434
QSO	1.484		30.512 ± 0.760	12.817 ± 0.516	-0.500
Lyα	2.330		38.988 ± 0.531	8.632 ± 0.101	-0.431

through

$$\chi^2_{\rm BAO} = \sum_{\rm tracers} \left(\mathbf{d} - \mathbf{d}^{\rm th}\right)^T \mathbf{C}^{-1} \left(\mathbf{d} - \mathbf{d}^{\rm th}\right), \qquad \mathbf{d} = \left(\frac{d_{\rm M}}{r_{\rm d}},\ \frac{d_{\rm H}}{r_{\rm d}}\right)$$

Symbols in this equation.

- $\chi^2_{\rm BAO}$: The chi-squared of the model distances against the 13 DESI DR2 BAO measurements (the pairs of the anisotropic tracers with their correlations, and one point for BGS). Unit: dimensionless. Value: 31.49 for UM-2 (32.55 for ΛCDM). Status: DERIVED.
- $\mathbf{d}$, $\mathbf{d}^{\rm th}$: The measured pair $(d_{\rm M}/r_{\rm d}, d_{\rm H}/r_{\rm d})$ of an anisotropic BAO tracer, and the pair $\mathbf{d}^{\rm th}$ computed in the model. Unit: dimensionless. Status: STANDARD.
- ^{T}: The transpose: a row written with ^{T} is a column. Status: STANDARD.
- $\mathbf{C}$: The covariance matrix of the measured pair of one BAO tracer, built from the published errors and the published correlation coefficient of the pair. Unit: dimensionless. Status: STANDARD.
- $d_{\rm M}$, $d_{\rm H}$, $d_{\rm V}$: The distances of the BAO analysis at the redshift $z_{\rm red}$ (flat space, $c = 1$): the comoving transverse distance $d_{\rm M}$, the Hubble distance $d_{\rm H} = 1/\mathcal{H}$ at that redshift, and the volume-averaged distance $d_{\rm V} = (z_{\rm red} d_{\rm M}^2 d_{\rm H})^{1/3}$ (written with a lower-case d because D_M is the covariant derivative). Unit: length (Mpc). Status: STANDARD.
- $r_{\rm d}$: The comoving sound horizon at the drag epoch, the standard ruler of the BAO. Unit: length (Mpc). Value: 147.104 Mpc (UM-2). Status: DERIVED.

for the pairs of the anisotropic tracers (with the published errors and correlations in the covariance $\mathbf{C}$ of each tracer, and the computed pair $\mathbf{d}^{\rm th}$), and the squared deviation over the variance of $d_{\rm V}/r_{\rm d}$ for BGS (DERIVED):

model	$\mathcal{H}_0$	$\Omega_{\rm m}$	$N_{\rm eff}$	helium fraction	age (Gyr)	$z_{\rm red,*}$	$r_{\rm d}$ (Mpc)	$\chi^2_{\rm BAO}$ (13 points)
ΛCDM	67.34	0.3154	3.044	0.24566	13.799	1089.91	147.103	32.55
UM-2	67.38	0.3150	3.0442	0.24567	13.796	1089.91	147.104	31.49
UM-2 without the freeze-in fold bands (version 2)	67.37	0.3151	3.0432	0.24566	13.797	1089.91	147.109	31.72
UM-2 with a thermalised fold band (electroweak)	68.48	0.3050	3.1390	0.24693	13.698	1090.00	146.630	15.43
UM-2 with a thermalised fold band (QCD)	69.65	0.2948	3.2395	0.24824	13.596	1090.10	146.133	10.65

($z_{\rm red,*}$ the redshift of last scattering.) The neutrino density parameter of UM-2 in CAMB, $\Omega_\nu h^2_{100} = 6.4511 \times 10^{-4}$, equals 6.4510×10^{-4} from the CNB of step 5 times $(N_{\rm eff}/3)^{3/4}$; the mass sum 0.0599 eV lies below the DESI DR2 bound 0.064 eV, which would allow $m_1 < 4.1$ meV.

The CMB. The UM-2 TT and EE spectra agree with Planck ΛCDM to 2.5×10^{-4} and 2.7×10^{-4} (multipoles 30 to 2500; for TT at most 0.007 of the cosmic variance $\sqrt{2/(2l+1)}$ of a single multipole l; the binned full-sky cosmic-variance χ^2, the sum over bins of 50 multipoles of the squared mean relative deviation over the cosmic variance of the bin, is 0.020; without the freeze-in fold bands 2.8×10^{-4}, 2.8×10^{-4} and 0.023): UM-2 reproduces the CMB of ΛCDM. By themselves the three mass states instead of one state of 60 meV move TT and EE by 2.6×10^{-4} and 2.3×10^{-4}, and $N_{\rm eff} = 3.0432$ instead of 3.044 by 1.3×10^{-4}. The fold-band variants, at fixed Planck $\Omega_{\rm b} h^2_{100}$, $\Omega_{\rm DM} h^2_{100}$ and $\theta_{\rm MC}$, move TT by up to 1.5 % and 3.1 % (0.8 and 1.5 cosmic-variance units of a single multipole; binned $\chi^2 = 293$ and 1208): they are not fits to the CMB, and they lie 2.1 and 2.9 standard

deviations above the ACT DR6 value $N_{\rm eff} = 2.86 \pm 0.13$ (UM-2: 1.4). The CNB enters through $N_{\rm eff} = 3.0432$ and the three neutrino masses.




Figure 4: CMB temperature (left) and E-mode (right) spectra of UM-2 (three massive neutrinos), of Planck 2018 ΛCDM and of the variant with a fold band thermalised above the QCD crossover; lower panels: relative difference to ΛCDM with the cosmic variance of one multipole (dotted).

The matter power spectrum. A separate CAMB run of each model without the CMB gives the linear matter power spectrum and $\sigma_8 = 0.8118$ (UM-2), 0.8119 (UM-2 without the freeze-in fold bands) and 0.8109 (ΛCDM; Planck 2018: 0.811, STANDARD); the product of the growth rate $\mathrm{d}\ln D_+/\mathrm{d}a_4$ and σ_8 at the redshift 0.5 is 0.4761, 0.4762 and 0.4756 (DERIVED). The fold bands produced by freeze-in through I11 (step 9) are massless free-streaming radiation with $\Delta N_{\rm eff} = 0.001$: they move the TT spectrum by at most 1.6×10^{-4} against the model without them (DERIVED, 8 checks in all).

9.4 The check of step 6

In its own equations. The first-order potential of the CAMB solution satisfies the Poisson equation of the first-order density inside the horizon,

$$\Phi_{\rm rms}(k) = \left[\frac{k^3 P_{\Phi_{\rm W}}(k)}{2\pi^2}\right]^{1/2}, \qquad \Phi_{\rm W} = \frac{\Phi + \Psi_{\rm N}}{2}, \qquad \Phi_{\rm W} = -\frac{3}{2}\,\Omega_{\rm m}\mathcal{H}_0^2\,\frac{\delta_{\rm m}}{\mathrm{e}^{a_4}k^2} \quad \text{(inside the horizon)},$$
$$\Phi_{\rm W} = -\frac{3}{5}\,\mathcal{R}\,\mathrm{e}^{-a_4}D_+ \quad \text{(largest scales)}, \qquad D_+ = \frac{5}{2}\,\Omega_{\rm m}\,\frac{\mathcal{H}}{\mathcal{H}_0}\int_{-\infty}^{a_4}\left(\frac{\mathcal{H}_0}{\mathcal{H}}\right)^3 \mathrm{e}^{-2\tilde{a}_4}\,\mathrm{d}\tilde{a}_4, \qquad \text{(FE-98)}$$
$$\frac{\mathcal{H}}{\mathcal{H}_0} = \left(\Omega_{\rm m}\mathrm{e}^{-3a_4} + 1 - \Omega_{\rm m}\right)^{1/2}$$

Symbols in (FE-98).

- $\Phi_{\rm rms}(k)$: The root-mean-square amplitude of the Weyl potential per logarithmic interval of k. Unit: dimensionless. Value: 2.42×10^{-5} at $k = 10^{-4}$ Mpc^{-1}. Status: STANDARD.




Figure 5: BAO distances divided by the square root of the redshift against the DESI DR2 measurements; the $\chi^2_{\rm BAO}$ uses $d_{\rm V}$ for BGS and $d_{\rm M}$, $d_{\rm H}$ with their correlation for the other tracers (the $d_{\rm V}$ points of the anisotropic tracers are shown for illustration).

- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- $P_{\Phi_{\rm W}}(k)$: The power spectrum of the Weyl potential. Unit: length^3. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $\Phi_{\rm W}$: The Weyl potential, $\Phi_{\rm W} = (\Phi + \Psi_{\rm N})/2$ (it deflects light). Unit: dimensionless. Status: STANDARD.
- Φ: The lapse potential of the Newtonian gauge (the perturbation of the time component, $1 + 2\Phi$). Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- $\Psi_{\rm N}$: The curvature potential of the Newtonian gauge (the perturbation of the 3-space metric, $1 - 2\Psi_{\rm N}$); not related to the field Ψ. Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- $\Omega_{\rm m}$: The matter density today in units of the critical density. Unit: dimensionless. Value: 0.3150. Status: STANDARD.
- $\mathcal{H}_0$: The Hubble rate today (the Hubble constant). Unit: GeV (km/s/Mpc). Value: 67.38 km/s/Mpc. Status: DERIVED.
- $\delta_{\rm m}$: The density contrast of the total matter (dark matter and baryons). Unit: dimensionless. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- $\mathcal{R}$: The primordial comoving curvature perturbation (conserved outside the horizon), counted positive where the density is above its mean, so that the Newtonian potentials Φ, $\Psi_{\rm N}$ and $\Phi_{\rm W}$ are negative there (on the largest scales in the matter era $\Phi_{\rm W} = -\frac{3}{5}\mathcal{R}\,\mathrm{e}^{-a_4}D_+$). Unit: dimensionless. Status: STANDARD.
- D_+: The linear growth factor of the matter perturbations, normalised as $D_+ = \mathrm{e}^{a_4}$ in the matter era. Unit: dimensionless. Value: $D_+\mathrm{e}^{-a_4} = 0.7878$ today. Status: STANDARD.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_{\rm Pl}$). Unit: GeV. Status: DERIVED.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no upper bound. Status: STANDARD.
- $\tilde{a}_4$: An earlier value of a_4 (the integration variable of the growth integral). Unit: dimensionless. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.

to 1.5×10^{-6} for $k \geq 0.01$ Mpc^{-1}, and the matter-era form on the largest scale to 4.4×10^{-5} (what it neglects: the departure of the transfer function from 1 and the radiation, about 10^{-4} at $k = 10^{-4}$ Mpc^{-1}; the growth factor $D_+\mathrm{e}^{-a_4} = 0.787828964553$ today, recomputed 0.787828964555) (DERIVED).

In concert with the other field equations.

- *The neutrino masses of the tip conditions* (step 2) are the masses of the CAMB computation to 5.7×10^{-12} (the mass iteration converges to 7.8×10^{-12}) (DERIVED).
- N_{eff} *of the decoupling and of the freeze-in* (steps 5 and 9) are the input of CAMB exactly (5×10^{-16}); its neutrino density today equals the CNB of step 5 times $(N_{\text{eff}}/3)^{3/4}$ to 1.9×10^{-5} (DERIVED).
- *The relic density of the Boltzmann equation* (step 9, two species, 2×0.060) is the cold dark matter of the CMB computation (0.1200) to 3.1×10^{-9} (DERIVED).
- *The radiation after decoupling* of the thermal history (step 9) differs from the CMB computation by 0.59 % (the thermal history decouples the neutrinos instantaneously; it uses this only for the entropy today and the nucleosynthesis environment) (DERIVED).
- *The metric series* (step 10): the potentials, the block shift, the first Kaluza-Klein mass and σ_8 used there are those of this step, of step 8 and of step 2, to 4.7×10^{-12} (DERIVED).

9.5 The status of step 6

STANDARD: the linear Einstein-Boltzmann equations, CAMB, the Planck 2018 parameters, the DESI DR2 and ACT DR6 measurements. DERIVED: the UM-2 spectra, distances and σ_8 with the UM-2 content (three mass states with the heated spectrum, the freeze-in fold bands), their agreement with ΛCDM and with the BAO data, the exclusion of a thermalised fold band. OPEN: the joint CMB and BAO fit of the fold-band variants; the primordial spectrum (an input).

10 Step 7: the dark-matter relic and its detection

The lightest odd states of Ψ_A and of Ψ_B (step 2) are two degenerate Dirac fermions of the mass $M_{\text{DM}} = 3.102204\, m_\Psi = 44.35$ TeV ($m_\Psi = H$), cold (step 3) and neutral; UM-2 has no tip operator, so neither couples to the Z, and both are stable by the fold parity (step 4). Step 7 computes their relic abundance through the portal I12 with the freeze-out formula, shows that I12 alone violates partial-wave unitarity, and estimates the detection; step 9 replaces the freeze-out formula by the Boltzmann equation and resolves the unitarity problem with I13.

10.1 The field equations of step 7

$$S_{\text{I12}} = -\int \mathrm{d}^8X\, \sqrt{|g|}\, \frac{\bar\Psi\Psi}{\Lambda_\star^3}\, \mathcal{F}, \qquad \mathcal{F} = c_{\text{G}}\, G^A_{MN} G^{AMN} + c_{\text{W}}\, W^I_{MN} W^{IMN} + c_{\text{B}}\, B_{MN} B^{MN} \tag{IN-17}$$

Symbols in (IN-17).

- S_{I12}: The action of the portal I12 (scalar density of dirac16complex times the gauge kinetic invariants). Unit: dimensionless. Status: INVENTED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\, \mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\, \mathrm{d}X^M \mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\, \mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- Ψ, $\bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\text{GeV}^{7/2}$. Status: RECORD.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- $\mathcal{F}$: The gauge-kinetic operator of the portal I12, $\mathcal{F} = c_{\text{G}} G^A_{MN} G^{AMN} + c_{\text{W}} W^I_{MN} W^{IMN} + c_{\text{B}} B_{MN} B^{MN}$; $\langle\mathcal{F}\rangle$ its mean value in an environment. Unit: GeV^4. Status: INVENTED.
- c_{G}: The coefficient of the gluon term of the portal I12. Unit: dimensionless. Value: $c_{\text{G}}/\Lambda_\star^3 = c_{\text{I12}}/\Lambda_\star^3$. Status: INVENTED.

- $G^A_M,\ G_M,\ G_{MN},\ G^A_{MN}$: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G^A_M ($A = 1, \ldots, 8$), the matrix $G_M = G^A_M t^A$, the field strength G_{MN} (matrix) and its components G^A_{MN}; $G^A_{\mu\nu}$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- c_{W}: The coefficient of the SU(2) term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{W}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- $W^I_M,\ W_M,\ W_{MN},\ W^I_{MN}$: The SU(2) gauge field (geometric normalisation): the components W^I_M ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W^I_M \tau^I/2$, the field strength W_{MN} and its components W^I_{MN}; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- c_{B}: The coefficient of the hypercharge term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{B}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- $B_M,\ B_{MN}$: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.

On the dark matter, I12 is a four-dimensional coupling of the scalar density to the gauge-field invariants, with the coefficient $\mathcal{O}_{\mathrm{DM}}$ of step 3:

$$\mathcal{L}_4^{\mathrm{I12}} = -\mathcal{O}_{\mathrm{DM}}\,\frac{\bar\Psi_{\mathrm{DM}}\Psi_{\mathrm{DM}}}{\Lambda_\star^3}\big(c_{\mathrm{G}}\,G^A_{\mu\nu}G^{A\mu\nu} + c_{\mathrm{W}}\,W^I_{\mu\nu}W^{I\mu\nu} + c_{\mathrm{B}}\,B_{\mu\nu}B^{\mu\nu}\big), \qquad \mathcal{O}_{\mathrm{DM}} = \frac{\partial\varepsilon_{\mathrm{DM}}}{\partial m_\Psi},$$
$$G_{\mathrm{g}} = \mathcal{O}_{\mathrm{DM}}\,\frac{c_{\mathrm{G}}\,g_3^2}{\Lambda_\star^3}, \qquad G_{\mathrm{W}} = \mathcal{O}_{\mathrm{DM}}\,\frac{c_{\mathrm{W}}\,g_2^2}{\Lambda_\star^3}, \qquad G_{\mathrm{B}} = \mathcal{O}_{\mathrm{DM}}\,\frac{c_{\mathrm{B}}\,g_1^2}{\Lambda_\star^3} \tag{FE-71}$$

Symbols in (FE-71).

- $\mathcal{L}_4^{\mathrm{I12}}$: The four-dimensional Lagrangian of I12 on the dark matter. Unit: GeV^4. Status: DERIVED.
- $\mathcal{O}_o,\ \mathcal{O}_{\mathrm{DM}}$: The scalar overlap of a state of dirac16complex, $\mathcal{O}_o = \partial\varepsilon_o/\partial m_\Psi$ (Hellmann-Feynman: the integrated scalar density of the orbital); $\mathcal{O}_{\mathrm{DM}}$ that of the dark matter. Unit: dimensionless. Value: $\mathcal{O}_{\mathrm{DM}} = 0.7072$. Status: DERIVED.
- $\Psi_{\mathrm{DM}},\ \bar\Psi_{\mathrm{DM}}$: The four-dimensional field of a dark-matter quantum (the lowest odd level of dirac16complex on the static block, in either half) and its Dirac adjoint. Unit: $\mathrm{GeV}^{3/2}$. Status: DERIVED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- c_{G}: The coefficient of the gluon term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{G}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- $G^A_M,\ G_M,\ G_{MN},\ G^A_{MN}$: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G^A_M ($A = 1, \ldots, 8$), the matrix $G_M = G^A_M t^A$, the field strength G_{MN} (matrix) and its components G^A_{MN}; $G^A_{\mu\nu}$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index running over $x_1, \ldots, x_4$ (as μ). Status: STANDARD.
- c_{W}: The coefficient of the SU(2) term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{W}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- $W^I_M,\ W_M,\ W_{MN},\ W^I_{MN}$: The SU(2) gauge field (geometric normalisation): the components W^I_M ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W^I_M \tau^I/2$, the field strength W_{MN} and its components W^I_{MN}; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.

- c_{B}: The coefficient of the hypercharge term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{B}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- $\varepsilon_{\mathrm{DM}}$: The energy of the dark-matter level at rest (its mass M_{DM} on the static block). Unit: GeV. Value: $3.102204\, m_\Psi = 44.35$ TeV. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex, $m_\Psi = 3H\mathcal{M} = H$. Unit: GeV. Value: 14.296 TeV. Status: RECORD.
- G_{g}: The coupling of a dark-matter pair to canonically normalised gluon pairs through I12. Unit: GeV^{-3}. Value: 6.09×10^{-15} GeV^{-3} (benchmark). Status: DERIVED.
- g_3: The four-dimensional strong coupling ($g_3^2 = 4\pi\alpha_3$). Unit: dimensionless. Value: $g_3^2 = 0.7786$ at $2M_{\mathrm{DM}}$. Status: DERIVED.
- G_{W}: The coupling of a dark-matter pair to canonically normalised SU(2) gauge pairs through I12. Unit: GeV^{-3}. Value: 2.96×10^{-15} GeV^{-3}. Status: DERIVED.
- g_2: The four-dimensional SU(2) coupling ($g_2^2 = 4\pi\alpha_2$). Unit: dimensionless. Value: $g_2^2 = 0.3781$ at $2M_{\mathrm{DM}}$. Status: DERIVED.
- G_{B}: The coupling of a dark-matter pair to canonically normalised hypercharge pairs through I12. Unit: GeV^{-3}. Value: 1.08×10^{-15} GeV^{-3}. Status: DERIVED.
- g_1: The four-dimensional hypercharge coupling ($g_1^2 = 4\pi\alpha_1$, Standard-Model normalisation without the GUT factor). Unit: dimensionless. Value: $g_1^2 = 0.1380$ at $2M_{\mathrm{DM}}$. Status: DERIVED.

The number density of each species obeys the Boltzmann equation with the thermal average of the annihilation cross-section, bounded by partial-wave unitarity:

$$\frac{\mathrm{d}n_{\mathrm{DM}}}{\mathrm{d}x_4} + 3\mathcal{H}n_{\mathrm{DM}} = -\langle\sigma v_{\mathrm{rel}}\rangle\big(n_{\mathrm{DM}}\bar{n}_{\mathrm{DM}} - n_{\mathrm{DM,eq}}^2\big) \;\Rightarrow\; \frac{\mathrm{d}Y_{\mathrm{DM}}}{\mathrm{d}x} = -\Big(1 + \frac{1}{3}\frac{\mathrm{d}\ln g_s}{\mathrm{d}\ln T}\Big)\frac{s\,\langle\sigma v_{\mathrm{rel}}\rangle}{2x\mathcal{H}}\big(Y_{\mathrm{DM}}^2 - Y_{\mathrm{eq}}^2\big),$$
$$Y_{\mathrm{DM}} = \frac{n_{\mathrm{DM}} + \bar{n}_{\mathrm{DM}}}{s}, \qquad x = \frac{M_{\mathrm{DM}}}{T}, \qquad s = \frac{2\pi^2}{45}\, g_s T^3, \qquad Y_{\mathrm{eq}} = \frac{4M_{\mathrm{DM}}^2 T K_2(x)}{2\pi^2 s} \tag{FE-124}$$

Symbols in (FE-124).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- n_{DM}, $\bar{n}_{\mathrm{DM}}$: The number densities of dark-matter particles and antiparticles of one species. Unit: GeV^3. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_{\mathrm{Pl}}$). Unit: GeV. Status: DERIVED.
- σ: A cross-section (of the dark-matter annihilation or production process named in the equation; summed over the spins where marked). Unit: GeV^{-2}. Status: STANDARD.
- v_{rel}: The relative velocity of the two annihilating particles; $\langle\sigma v_{\mathrm{rel}}\rangle$ the thermal average. Unit: dimensionless (c = 1). Status: STANDARD.
- $n_{\mathrm{DM,eq}}$, $\bar{n}_{\mathrm{DM,eq}}$: The equilibrium number densities of dark-matter particles and antiparticles of one species. Unit: GeV^3. Status: STANDARD.
- Y_{DM}, Y_{eq}, Y_∞: The abundance (number density per entropy density) of one dark-matter species (particles and antiparticles), its equilibrium value and its final value. Unit: dimensionless. Status: STANDARD.
- x: The ratio $x = M_{\mathrm{DM}}/T$ of the dark-matter mass to the temperature. Unit: dimensionless. Value: $x_{\mathrm{f}} = 31.23$ at the freeze-out. Status: STANDARD.
- g_ρ, g_s: The effective numbers of relativistic degrees of freedom of the energy density and of the entropy density of the plasma. Unit: dimensionless. Value: 105.75 when every species of UM-2 in equilibrium is relativistic (no Higgs particle), 10.75 near 1 MeV, $g_s = 43/11$ today. Status: STANDARD.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.

- s, s_s: The entropy density of the plasma, $s = 2\pi^2 g_s T^3/45$; s_s that of the species s. Unit: GeV^3. Status: STANDARD.
- M_{DM}: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- K_1, K_2: The modified Bessel functions of the second kind of orders 1 and 2 (not related to the block variable K). Unit: dimensionless. Status: STANDARD.

$$\langle \sigma v_{\mathrm{rel}} \rangle = \frac{1}{8 M_{\mathrm{DM}}^4 T K_2(x)^2} \int_{4M_{\mathrm{DM}}^2}^{\infty} \sigma \left(E_{\mathrm{cm}}^2 - 4M_{\mathrm{DM}}^2\right) E_{\mathrm{cm}} \, K_1\!\left(\frac{E_{\mathrm{cm}}}{T}\right) \mathrm{d}\big(E_{\mathrm{cm}}^2\big) \tag{FE-125}$$

Symbols in (FE-125).

- σ: A cross-section (of the dark-matter annihilation or production process named in the equation; summed over the spins where marked). Unit: GeV^{-2}. Status: STANDARD.
- v_{rel}: The relative velocity of the two annihilating particles; $\langle \sigma v_{\mathrm{rel}} \rangle$ the thermal average. Unit: dimensionless ($\mathrm{c} = 1$). Status: STANDARD.
- M_{DM}: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.
- K_1, K_2: The modified Bessel functions of the second kind of orders 1 and 2 (not related to the block variable K). Unit: dimensionless. Status: STANDARD.
- x: The ratio $x = M_{\mathrm{DM}}/T$ of the dark-matter mass to the temperature. Unit: dimensionless. Value: $x_{\mathrm{f}} = 31.23$ at the freeze-out. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no upper bound. Status: STANDARD.
- E_{cm}: The total energy of the colliding pair in its centre-of-mass frame. Unit: GeV. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.

$$\begin{aligned}
\sigma_{\mathrm{I12}} &= \frac{G_{\mathrm{I12}}^2 \, E_{\mathrm{cm}}^3 \big(E_{\mathrm{cm}}^2 - 4M_{\mathrm{DM}}^2\big)^{1/2}}{8\pi}, \qquad G_{\mathrm{I12}}^2 = 8G_{\mathrm{g}}^2 + 3G_{\mathrm{W}}^2 + G_{\mathrm{B}}^2, \\
\sigma_{\mathrm{I13}} &= \sum_f \frac{N_{\mathrm{c}} \, G_f^2 \, E_{\mathrm{cm}}^2 \big(1 + 2M_{\mathrm{DM}}^2/E_{\mathrm{cm}}^2\big)}{24\pi\beta_{\mathrm{cm}}}, \qquad G_f = G_{\mathrm{A}} \, q_f^B \, \tilde{I}_{of}^{(b)}, \\
\beta_{\mathrm{cm}} &= \sqrt{1 - \frac{4M_{\mathrm{DM}}^2}{E_{\mathrm{cm}}^2}}, \qquad k_{\mathrm{cm}} = \sqrt{\frac{E_{\mathrm{cm}}^2}{4} - M_{\mathrm{DM}}^2}
\end{aligned} \tag{FE-127}$$

Symbols in (FE-127).

- σ_{I12}, σ_{I13}: The spin-averaged annihilation cross-sections of a dark-matter pair through I12 (into gauge-boson pairs) and through I13 (into quark pairs). Unit: GeV^{-2}. Status: DERIVED.
- G_{I12}: The combined coupling of I12 to all gauge-boson pairs, $G_{\mathrm{I12}}^2 = 8G_{\mathrm{g}}^2 + 3G_{\mathrm{W}}^2 + G_{\mathrm{B}}^2$. Unit: GeV^{-3}. Value: benchmark $\pi/(2M_{\mathrm{DM}}^3) = 1.801 \times 10^{-14}$ GeV^{-3}. Status: DERIVED.
- E_{cm}: The total energy of the colliding pair in its centre-of-mass frame. Unit: GeV. Status: STANDARD.
- M_{DM}: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_{g}: The coupling of a dark-matter pair to canonically normalised gluon pairs through I12. Unit: GeV^{-3}. Value: 6.09×10^{-15} GeV^{-3} (benchmark). Status: DERIVED.
- G_{W}: The coupling of a dark-matter pair to canonically normalised SU(2) gauge pairs through I12. Unit: GeV^{-3}. Value: 2.96×10^{-15} GeV^{-3}. Status: DERIVED.

- G_{B}: The coupling of a dark-matter pair to canonically normalised hypercharge pairs through I12. Unit: GeV^{-3}. Value: $1.08 \times 10^{-15}\ \mathrm{GeV}^{-3}$. Status: DERIVED.
- f: The label of a fermion species: a Standard-Model multiplet of one generation (its N_{c} Weyl components share one zero-mode profile; the zero mode of each chirality) or a fermion of a medium (electron, proton, neutron); a sum over f runs over these species. Status: STANDARD.
- N_{c}: The number of Weyl components of a fermion multiplet that share its zero-mode profile and overlap (colour times isospin): 6 for a quark doublet, 3 for a quark singlet, 2 for a lepton doublet, 1 for a charged-lepton singlet. Unit: dimensionless. Status: STANDARD.
- G_f: The I13 coupling of the dark matter to the quark component f, $G_f = G_{\mathrm{A}} q_f^B \tilde{I}_{of}^{(b)}$. Unit: GeV^{-2}. Status: DERIVED.
- β_{cm}: The velocity of each particle of the pair in the centre-of-mass frame. Unit: dimensionless. Status: STANDARD.
- G_{A}: The four-dimensional strength of the portal I13 on the zero modes, $G_{\mathrm{A}} = c_{\mathrm{A}} H/(2 f^4 v_{\mathrm{t}} \Lambda_\star^6)$. Unit: GeV^{-2}. Value: $1.278 \times 10^{-9}\ \mathrm{GeV}^{-2}$ (dark-matter relic density). Status: DERIVED.
- q_ψ^B, q_f^B: The baryon number of the multiplet ψ or of the fermion component f ($\frac{1}{3}$ for quarks, 0 for leptons). Unit: dimensionless. Status: STANDARD.
- $\tilde{I}_{of}^{(b)}$: The overlap of the lower component of the state o with the zero mode of f, $\tilde{I}_{of}^{(b)} = H^{-1} \int_{-L}^{0} b_o^2 \varrho_f \mathrm{e}^{-3Hy} \mathrm{d}y$ (the overlap of I13); $\tilde{I}_Q^{(b)}$, $\tilde{I}_{u_R}^{(b)}$, $\tilde{I}_{d_R}^{(b)}$, $\tilde{I}_{\nu_R}^{(b)}$ those of the dark matter. Unit: dimensionless. Value: dark matter: $\tilde{I}_Q^{(b)} = 2.6135$. Status: DERIVED.
- o: The label of a Kohn-Sham orbital (a state of dirac16complex in the cell: block, momentum shell, fold parity and level); a sum over o runs over the orbitals. Status: RECORD.
- k_{cm}: The momentum of each particle of the pair in the centre-of-mass frame. Unit: GeV. Status: STANDARD.

$$\sigma_{\mathrm{I12}} \le \frac{\pi}{4 k_{\mathrm{cm}}^2} \ \text{(total angular momentum 0)}, \qquad \sigma_{\mathrm{I13}} \le \frac{3\pi}{4 k_{\mathrm{cm}}^2} \ \text{(total angular momentum 1)} \tag{FE-128}$$

Symbols in (FE-128).

- σ_{I12}, σ_{I13}: The spin-averaged annihilation cross-sections of a dark-matter pair through I12 (into gauge-boson pairs) and through I13 (into quark pairs). Unit: GeV^{-2}. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- k_{cm}: The momentum of each particle of the pair in the centre-of-mass frame. Unit: GeV. Status: STANDARD.

$$\Omega_{\mathrm{DM}} h_{100}^2 = \frac{M_{\mathrm{DM}}\, Y_\infty\, s_0}{\rho_{\mathrm{crit}}/h_{100}^2}, \qquad \frac{\rho_{\mathrm{crit}}}{h_{100}^2} = 1.05368 \times 10^{-5}\ \mathrm{GeV/cm}^3 \tag{FE-126}$$

Symbols in (FE-126).

- Ω_{DM}: The dark-matter density today in units of the critical density. Unit: dimensionless. Value: $\Omega_{\mathrm{DM}} h_{100}^2 = 0.120$ (both species). Status: DERIVED.
- h_{100}: The reduced Hubble constant, $\mathcal{H}_0/(100\ \mathrm{km\ s^{-1}\ Mpc^{-1}})$. Unit: dimensionless. Value: 0.6738. Status: STANDARD.
- M_{DM}: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- Y_{DM}, Y_{eq}, Y_∞: The abundance (number density per entropy density) of one dark-matter species (particles and antiparticles), its equilibrium value and its final value. Unit: dimensionless. Status: STANDARD.
- s_0: The entropy density today. Unit: GeV^3 (quoted per cm^3). Value: $2891.3\ \mathrm{cm}^{-3}$. Status: STANDARD.
- ρ_{crit}: The critical density, $3\mathcal{H}_0^2 M_{\mathrm{Pl}}^2/(8\pi)$. Unit: GeV^4 (quoted in $\mathrm{GeV/cm}^3$). Value: $\rho_{\mathrm{crit}}/h_{100}^2 = 1.05368 \times 10^{-5}\ \mathrm{GeV/cm}^3$. Status: STANDARD.

10.2 The approximation of step 7

- *The freeze-out formula* replaces the Boltzmann equation (the standard radiation-dominated freeze-out); against the numerical solution of step 9 it is accurate to 2.4 % (an s-wave test case gives 0.1306 against 0.1275).

- *The non-relativistic expansion* of the thermal average, $\langle\sigma v_{\rm rel}\rangle = 6G_{\rm I12}^2 M_{\rm DM}^4/(\pi x)$; the exact relativistic average exceeds it by 0.111 at $x = 30$ (by about $3/x$).
- *The effective operator* is used up to the energies of the annihilations ($\sqrt{4M_{\rm DM}^2}$ and above); where its cross-section exceeds the unitarity bound the operator is not a valid description (below).
- *The direct-detection estimate* uses the trace anomaly with a nucleon gluon fraction and a low-scale coupling quoted without a checked reference (ESTIMATE).

10.3 The solution of step 7

Through I12 the dark matter annihilates in the p-wave into gauge-boson pairs. In the hypercharge channel, with the coupling $G_{\rm B}$, the spin-averaged rate is $G_{\rm B}^2 E_{\rm cm}^2 (E_{\rm cm}^2 - 4M_{\rm DM}^2)/(4\pi)$ (the polarisation sum $E_{\rm cm}^4/2$ and the spin trace $2(E_{\rm cm}^2 - 4M_{\rm DM}^2)$ checked numerically); each of the eight gluon and three $SU(2)$ channels adds the same with its own coupling, so in total

$$\sigma_{\rm I12}\, v_{\rm rel} = \frac{G_{\rm I12}^2 E_{\rm cm}^2 \big(E_{\rm cm}^2 - 4M_{\rm DM}^2\big)}{4\pi} \;\to\; \frac{G_{\rm I12}^2 M_{\rm DM}^4\, v_{\rm rel}^2}{\pi} \qquad (v_{\rm rel} \ll 1)$$

Symbols in this equation.

- $\sigma_{\rm I12}$, $\sigma_{\rm I13}$: The spin-averaged annihilation cross-sections of a dark-matter pair through I12 (into gauge-boson pairs) and through I13 (into quark pairs). Unit: GeV^{-2}. Status: DERIVED.
- $v_{\rm rel}$: The relative velocity of the two annihilating particles; $\langle\sigma v_{\rm rel}\rangle$ the thermal average. Unit: dimensionless (c = 1). Status: STANDARD.
- $G_{\rm I12}$: The combined coupling of I12 to all gauge-boson pairs, $G_{\rm I12}^2 = 8G_{\rm g}^2 + 3G_{\rm W}^2 + G_{\rm B}^2$. Unit: GeV^{-3}. Value: benchmark $\pi/(2M_{\rm DM}^3) = 1.801 \times 10^{-14}$ GeV^{-3}. Status: DERIVED.
- $E_{\rm cm}$: The total energy of the colliding pair in its centre-of-mass frame. Unit: GeV. Status: STANDARD.
- $M_{\rm DM}$: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

(DERIVED). Since $\bar\Psi\Psi = \bar\Psi_A\Psi_A + \bar\Psi_B\Psi_B$ does not mix the halves, each species freezes out separately and contributes half of $\Omega_{\rm DM}h_{100}^2 = 0.120$. The freeze-out of a Dirac fermion with $g_\rho = g_s = 105.75$ at $T = M_{\rm DM}/x_{\rm f} \approx 1.5$ TeV (the Standard Model of three generations without a Higgs particle; the right-handed neutrinos and the fold bands are never thermalised) gives

$$x_{\rm f} = \ln\Big[0.095\,\frac{M_{\rm Pl}\, M_{\rm DM}\,\langle\sigma v_{\rm rel}\rangle}{\sqrt{g_\rho\, x_{\rm f}}}\Big] = 30.38, \qquad \Omega_{\rm DM}h_{100}^2 = \frac{2\times 1.07\times 10^{9}\ \text{GeV}^{-1}\, x_{\rm f}^2}{\sqrt{g_\rho}\, M_{\rm Pl}\, 3G_{\rm I12}^2 M_{\rm DM}^4/\pi} = 0.060 \text{ per species}$$

Symbols in this equation.

- $x_{\rm f}$: The freeze-out point $x_{\rm f} = M_{\rm DM}/T$ at which the dark matter leaves chemical equilibrium. Unit: dimensionless. Value: 30.38 (freeze-out formula, I12), 31.23 (Boltzmann equation, I12 + I13). Status: DERIVED.
- $M_{\rm Pl}$: The Planck mass, $M_{\rm Pl} = G_{\rm N}^{-1/2}$. Unit: GeV. Value: 1.22091×10^{19} GeV. Status: STANDARD.
- $M_{\rm DM}$: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- σ: A cross-section (of the dark-matter annihilation or production process named in the equation; summed over the spins where marked). Unit: GeV^{-2}. Status: STANDARD.
- $v_{\rm rel}$: The relative velocity of the two annihilating particles; $\langle\sigma v_{\rm rel}\rangle$ the thermal average. Unit: dimensionless (c = 1). Status: STANDARD.
- g_ρ, g_s: The effective numbers of relativistic degrees of freedom of the energy density and of the entropy density of the plasma. Unit: dimensionless. Value: 105.75 when every species of UM-2 in equilibrium is relativistic (no Higgs particle), 10.75 near 1 MeV, $g_s = 43/11$ today. Status: STANDARD.
- $\Omega_{\rm DM}$: The dark-matter density today in units of the critical density. Unit: dimensionless. Value: $\Omega_{\rm DM}h_{100}^2 = 0.120$ (both species). Status: DERIVED.

- h_{100}: The reduced Hubble constant, $\mathcal{H}_0/(100\ \mathrm{km\ s^{-1}\ Mpc^{-1}})$. Unit: dimensionless. Value: 0.6738. Status: STANDARD.
- G_{I12}: The combined coupling of I12 to all gauge-boson pairs, $G_{\mathrm{I12}}^2 = 8G_{\mathrm{g}}^2 + 3G_{\mathrm{W}}^2 + G_{\mathrm{B}}^2$. Unit: GeV^{-3}. Value: benchmark $\pi/(2M_{\mathrm{DM}}^3) = 1.801 \times 10^{-14}\ \mathrm{GeV}^{-3}$. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

($M_{\mathrm{Pl}} = G_{\mathrm{N}}^{-1/2} = 1.22091 \times 10^{19}$ GeV; the number 0.095 is the standard coefficient 0.038 of the freeze-out condition times the factor 5/4 of its matching constant 1/2 and times the two spin states of a Dirac fermion), which requires $\langle\sigma v_{\mathrm{rel}}\rangle(x_{\mathrm{f}}) = 2.01 \times 10^{-25}\ \mathrm{cm^3/s}$ per species, i.e.

$$G_{\mathrm{I12}} = \left(8G_{\mathrm{g}}^2 + 3G_{\mathrm{W}}^2 + G_{\mathrm{B}}^2\right)^{1/2} = (15.5\ \mathrm{TeV})^{-3}, \qquad G_{\mathrm{g}} = \mathcal{O}_{\mathrm{DM}}\,\frac{c_{\mathrm{G}}\,g_3^2}{\Lambda_\star^3}, \quad \mathcal{O}_{\mathrm{DM}} = 0.7072$$

Symbols in this equation.

- G_{I12}: The combined coupling of I12 to all gauge-boson pairs, $G_{\mathrm{I12}}^2 = 8G_{\mathrm{g}}^2 + 3G_{\mathrm{W}}^2 + G_{\mathrm{B}}^2$. Unit: GeV^{-3}. Value: benchmark $\pi/(2M_{\mathrm{DM}}^3) = 1.801 \times 10^{-14}\ \mathrm{GeV}^{-3}$. Status: DERIVED.
- G_{g}: The coupling of a dark-matter pair to canonically normalised gluon pairs through I12. Unit: GeV^{-3}. Value: $6.09 \times 10^{-15}\ \mathrm{GeV}^{-3}$ (benchmark). Status: DERIVED.
- G_{W}: The coupling of a dark-matter pair to canonically normalised SU(2) gauge pairs through I12. Unit: GeV^{-3}. Value: $2.96 \times 10^{-15}\ \mathrm{GeV}^{-3}$. Status: DERIVED.
- G_{B}: The coupling of a dark-matter pair to canonically normalised hypercharge pairs through I12. Unit: GeV^{-3}. Value: $1.08 \times 10^{-15}\ \mathrm{GeV}^{-3}$. Status: DERIVED.
- $\mathcal{O}_o$, $\mathcal{O}_{\mathrm{DM}}$: The scalar overlap of a state of dirac16complex, $\mathcal{O}_o = \partial\varepsilon_o/\partial m_\Psi$ (Hellmann-Feynman: the integrated scalar density of the orbital); $\mathcal{O}_{\mathrm{DM}}$ that of the dark matter. Unit: dimensionless. Value: $\mathcal{O}_{\mathrm{DM}} = 0.7072$. Status: DERIVED.
- c_{G}: The coefficient of the gluon term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{G}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- g_3: The four-dimensional strong coupling ($g_3^2 = 4\pi\alpha_3$). Unit: dimensionless. Value: $g_3^2 = 0.7786$ at $2M_{\mathrm{DM}}$. Status: DERIVED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.

(a single species carrying all the dark matter would need $(17.6\ \mathrm{TeV})^{-3}$ with the same g_ρ). These are the numbers of the freeze-out formula (version 2); the numerical Boltzmann solution of step 9 needs $(15.86\ \mathrm{TeV})^{-3}$ at $x_{\mathrm{f}} = 31.70$.

Unitarity (DERIVED). The operator annihilates a fermion pair only from the state with total angular momentum zero and positive parity ($^3P_0$); for unpolarised Dirac fermions the inelastic cross-section from it is bounded by $\sigma \le \pi/(4k_{\mathrm{cm}}^2)$ ($k_{\mathrm{cm}} = M_{\mathrm{DM}}v_{\mathrm{rel}}/2$ for $v_{\mathrm{rel}} \ll 1$), i.e. $\sigma v_{\mathrm{rel}} \le \pi/(M_{\mathrm{DM}}^2 v_{\mathrm{rel}})$, whose thermal average over the relative velocities ($\propto v_{\mathrm{rel}}^2\,\mathrm{e}^{-xv_{\mathrm{rel}}^2/4}$) is

$$\left\langle \frac{\pi}{M_{\mathrm{DM}}^2\,v_{\mathrm{rel}}} \right\rangle = \frac{\sqrt{\pi x}}{M_{\mathrm{DM}}^2} = 5.79 \times 10^{-26}\ \mathrm{cm^3/s} \qquad (x = x_{\mathrm{f}} = 30.38)$$

Symbols in this equation.

- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- M_{DM}: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- v_{rel}: The relative velocity of the two annihilating particles; $\langle\sigma v_{\mathrm{rel}}\rangle$ the thermal average. Unit: dimensionless (c = 1). Status: STANDARD.
- x: The ratio $x = M_{\mathrm{DM}}/T$ of the dark-matter mass to the temperature. Unit: dimensionless. Value: $x_{\mathrm{f}} = 31.23$ at the freeze-out. Status: STANDARD.
- x_{f}: The freeze-out point $x_{\mathrm{f}} = M_{\mathrm{DM}}/T$ at which the dark matter leaves chemical equilibrium. Unit: dimensionless. Value: 30.38 (freeze-out formula, I12), 31.23 (Boltzmann equation, I12 + I13). Status: DERIVED.

(checked by numerical averaging). The freeze-out requirement exceeds it 3.47 times; the effective cross-section at the relic strength exceeds the bound above the relative velocity

$$v_{\mathrm{c}} = \Big(\frac{\pi}{G_{\mathrm{I12}}\, M_{\mathrm{DM}}^3}\Big)^{2/3} = 0.263,$$

Symbols in this equation.

- v_{c}: The relative velocity above which the effective I12 cross-section at the relic strength exceeds the ${}^3P_0$ unitarity bound (non-relativistic formula). Unit: dimensionless (units of c). Value: 0.263. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_{I12}: The combined coupling of I12 to all gauge-boson pairs, $G_{\mathrm{I12}}^2 = 8G_{\mathrm{g}}^2 + 3G_{\mathrm{W}}^2 + G_{\mathrm{B}}^2$. Unit: GeV^{-3}. Value: benchmark $\pi/(2M_{\mathrm{DM}}^3) = 1.801 \times 10^{-14}$ GeV^{-3}. Status: DERIVED.
- M_{DM}: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.

i.e. for 79 % of the pairs at freeze-out (with the strength and x_{f} of the freeze-out formula; with the relic strength of step 9 the formula gives 0.274 and 75 % of the pairs, and the exact crossing lies at $v_{\mathrm{rel}} = 2\beta_{\mathrm{cm}} = 0.269$). (Version 1 compared the requirement with $(4\pi/M_{\mathrm{DM}}^2)\sqrt{x_{\mathrm{f}}/\pi} = 2.32 \times 10^{-25}$ cm^3/s, the bound of a single helicity channel without the spin average and the restriction to ${}^3P_0$, a factor 4 too large.) The gauge-kinetic portal therefore cannot produce the relic abundance within perturbative unitarity. Step 9 recomputes this with the Boltzmann equation (a cross-section at the ${}^3P_0$ limit at every energy would leave $\Omega_{\mathrm{DM}}h_{100}^2 = 0.058$ per species: only a strongly coupled ${}^3P_0$ channel at the unitarity limit could give the relic) and resolves it with the s-wave portal I13.

Direct detection (ESTIMATE). Even if the whole coupling were in gluons, the trace anomaly of QCD gives the coupling of the dark matter to a nucleon of the mass m_{N}, through the gluon fraction f_{TG} of the nucleon mass (the nucleon matrix element of the gluon invariant is $-\frac{8\pi}{9\alpha_3} m_{\mathrm{N}} f_{\mathrm{TG}}$), and the spin-independent cross-section per nucleon,

$$f_{\mathrm{N}} = G_{\mathrm{g}}\, \frac{8\pi}{9\alpha_3}\, m_{\mathrm{N}}\, f_{\mathrm{TG}}, \qquad \sigma_{\mathrm{SI}} = \frac{\mu_{\mathrm{red}}^2 f_{\mathrm{N}}^2}{\pi}$$

Symbols in this equation.

- f_{N}, f_{p}, f_{n}: The spin-independent effective couplings of the dark matter to a nucleon: f_{p} to the proton, f_{n} to the neutron, and f_{N} the average per nucleon of the target (for the portal I12, the coupling through the gluon content of a nucleon). Unit: GeV^{-2}. Value: $f_{\mathrm{p}} = 1.915 \times 10^{-9}$, $f_{\mathrm{n}} = 1.917 \times 10^{-9}$ GeV^{-2} (I13 and I11). Status: ESTIMATE.
- G_{g}: The coupling of a dark-matter pair to canonically normalised gluon pairs through I12. Unit: GeV^{-3}. Value: 6.09×10^{-15} GeV^{-3} (benchmark). Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- α_3: The strong fine-structure constant, $\alpha_3 = g_3^2/(4\pi)$, at the scale given in brackets. Unit: dimensionless. Value: $\alpha_3(m_{\mathrm{Z}}) = 0.1180$; $\alpha_3(m_{\mathrm{KK}}) = 0.0606$. Status: STANDARD.
- m_{N}: The mass of a nucleon. Unit: GeV. Value: 0.939 GeV. Status: STANDARD.
- f_{TG}: The fraction of the nucleon mass carried by the gluons through the trace anomaly of QCD. Unit: dimensionless. Value: 0.92 (the value used in the computation; approximate). Status: STANDARD.
- σ_{SI}: The spin-independent elastic scattering cross-section of a dark-matter particle on one nucleon. Unit: cm^2. Value: 4.01×10^{-46} cm^2 (xenon, I13 and I11); 6.3×10^{-53} cm^2 through I12 alone. Status: ESTIMATE.
- μ_{red}: The reduced mass of the dark-matter particle and a nucleon, $M_{\mathrm{DM}} m_{\mathrm{N}}/(M_{\mathrm{DM}} + m_{\mathrm{N}})$. Unit: GeV. Status: STANDARD.

($f_{\mathrm{TG}} = 0.92$, $\alpha_3 = 0.3$ at a low scale, $m_{\mathrm{N}} = 0.939$ GeV, μ_{red} the dark-matter-nucleon reduced mass; approximate values quoted without a checked reference), i.e. $\sigma_{\mathrm{SI}} = 6.3 \times 10^{-53}$ cm^2 for each half (the two halves have the same cross-section and together the full local density, so the limit applies as for one species), eight orders below the LZ limit extrapolated linearly in the mass from 2.2×10^{-48} cm^2 at 40 GeV to 44 TeV (2.4×10^{-45} cm^2); the electroweak channels have no tree-level nucleon coupling. Any relic value would also depend on the radion oscillation left over from the deflation (which redshifts like dust): if it dominated the Universe until it decayed, it would dilute the dark matter after freeze-out (the moduli problem, last section).

10.4 The check of step 7

In its own equations. The polarisation sum and the spin trace of the cross-section were checked numerically, and the thermal average of the unitarity bound by numerical averaging (DERIVED). The freeze-out formula is checked against the numerical solution of the Boltzmann equation of step 9: on the s-wave test case they give 0.1306 and 0.1275 (2.4 %), and for I12 at the relic strength the numerical solution needs $(15.86\ \mathrm{TeV})^{-3}$ at $x_\mathrm{f} = 31.70$ against $(15.5\ \mathrm{TeV})^{-3}$ at 30.38 (DERIVED).

In concert with the other field equations.

- *The I12 coefficient of the Kohn-Sham equations.* $\mathcal{O}_\mathrm{DM} = \partial\varepsilon_\mathrm{DM}/\partial m_\Psi$ of step 3 (0.707196002672) is the rounded 0.7072 of this computation to 5.7×10^{-6} (DERIVED).
- *The dark-matter mass of the x_0 spectrum* (step 2) is the mass of the Boltzmann equation to 8.1×10^{-13} (DERIVED).
- *The degrees of freedom of the thermal history* (step 9): $g_\rho = g_s = 105.75$ at the freeze-out is the value of the ideal gases with the x_0 masses (105.73 at 1.42 TeV) (DERIVED).
- *The I12 strength in the mean fields* of step 8 is the benchmark of step 9 to 1.6×10^{-12} (DERIVED).

10.5 The status of step 7

DERIVED: the I12 cross-section, the freeze-out numbers and the violation of partial-wave unitarity by I12 at the relic strength. ESTIMATE: the direct detection through the trace anomaly (inputs quoted without a checked reference). INVENTED: the portal I12. OPEN: the microscopic origin of the portals (I12 alone could give the relic only as a strongly coupled ${}^3P_0$ channel).

11 Step 8: the coupled solution in environments

Steps 1 to 3 solve the field equations of gravity, of the masses and of dirac16complex one after the other. Step 8 couples them in ten environments, from the cosmic mean density today to neutron-star cores and the hot Universe up to the dark-matter freeze-out: the Kohn-Sham levels of dirac16complex in the mean fields of the Standard-Model fields of the environment (I11, I12, I13), the action of dirac16complex back on the Standard-Model fields (its scalar density changes the gauge couplings and the x_0 problem of the W), and the response of the block to the energy-momentum tensor of every species, iterated to self-consistency.

11.1 The field equations of step 8

The portals:

$$S_\mathrm{I11} = \int \mathrm{d}^8X\,\sqrt{|g|}\,\frac{c_\mathrm{V}}{\Lambda_\star^6}\left(\mathrm{i}\bar\Psi\gamma^M\Psi\right)J_M^{B-L}, \qquad J_M^{B-L} = \sum_{i=1}^{3}\sum_{\psi} q_\psi\left(\mathrm{i}\bar\psi_i\gamma_M\psi_i\right),$$
$$q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}, \qquad q_\ell = q_{e_R} = q_{\nu_R} = -1 \tag{IN-16}$$

Symbols in (IN-16).

- S_I11: The action of the portal I11 (dirac16complex vector current times the $B-L$ current of the fermions). Unit: dimensionless. Status: INVENTED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.

- c_{V}: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_{\mathrm{V}} = 4.38\times10^{-17}$ GeV^{-2}. Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- Ψ, $\bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- J_M^{B-L}: The $B-L$ current of the Standard-Model fermions (baryon minus lepton number), summed over the generations. Unit: GeV^7. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- ψ: The label of a Standard-Model multiplet: $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ (quark doublet, up and down quark singlets, lepton doublet, charged-lepton singlet, right-handed neutrino); a sum over ψ runs over the six. Status: STANDARD.
- q_ψ, q_f: The $B-L$ charge: q_ψ of the multiplet ψ ($q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}$, $q_\ell = q_{e_R} = q_{\nu_R} = -1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- ψ_i, $\bar\psi_i$: A Standard-Model fermion field of eight dimensions: ψ_i is the multiplet $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ of the generation i, ψ_f one fermion component f (in four-dimensional formulas its zero mode); $\bar\psi = \psi^\dagger C$ the Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$ (eight dimensions), $\mathrm{GeV}^{3/2}$ (four dimensions). Status: ASSUMED.

$$S_{\mathrm{I12}} = -\int \mathrm{d}^8X\,\sqrt{|g|}\,\frac{\bar\Psi\Psi}{\Lambda_\star^3}\,\mathcal{F}, \qquad \mathcal{F} = c_{\mathrm{G}}\,G^A_{MN}G^{AMN} + c_{\mathrm{W}}\,W^I_{MN}W^{IMN} + c_{\mathrm{B}}\,B_{MN}B^{MN} \tag{IN-17}$$

Symbols in (IN-17).

- S_{I12}: The action of the portal I12 (scalar density of dirac16complex times the gauge kinetic invariants). Unit: dimensionless. Status: INVENTED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- Ψ, $\bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- $\mathcal{F}$: The gauge-kinetic operator of the portal I12, $\mathcal{F} = c_{\mathrm{G}}G^A_{MN}G^{AMN} + c_{\mathrm{W}}W^I_{MN}W^{IMN} + c_{\mathrm{B}}B_{MN}B^{MN}$; $\langle\mathcal{F}\rangle$ its mean value in an environment. Unit: GeV^4. Status: INVENTED.
- c_{G}: The coefficient of the gluon term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{G}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- G^A_M, G_M, G_{MN}, G^A_{MN}: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G^A_M ($A = 1, \ldots, 8$), the matrix $G_M = G^A_M t^A$, the field strength G_{MN} (matrix) and its components G^A_{MN}; $G^A_{\mu\nu}$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.

- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- c_{W}: The coefficient of the SU(2) term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{W}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- W_M^I, W_M, W_{MN}, W_{MN}^I: The SU(2) gauge field (geometric normalisation): the components W_M^I ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W_M^I \tau^I/2$, the field strength W_{MN} and its components W_{MN}^I; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- c_{B}: The coefficient of the hypercharge term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{B}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.

$$
\begin{aligned}
S_{\mathrm{I13}} &= \int \mathrm{d}^8X \, \sqrt{|g|} \, \frac{c_{\mathrm{A}}}{\Lambda_\star^6} \sum_{M \neq x_0} \left(\mathrm{i} \bar{\Psi} \gamma^M P_b \Psi \right) J_M^B, \qquad P_b = \frac{1 - \gamma^{x_0}}{2}, \qquad P_a = \frac{1 + \gamma^{x_0}}{2}, \\
J_M^B &= \frac{1}{3} \sum_{i=1}^{3} \left(\mathrm{i} \bar{Q}_i \gamma_M Q_i + \mathrm{i} \bar{u}_{Ri} \gamma_M u_{Ri} + \mathrm{i} \bar{d}_{Ri} \gamma_M d_{Ri} \right)
\end{aligned}
\tag{IN-18}
$$

Symbols in (IN-18).

- S_{I13}: The action of the portal I13 (the x_0-chiral vector current of dirac16complex times the baryon-number current). Unit: dimensionless. Status: INVENTED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0 \, \mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+, +, +, +, -, -, -, -$ in the order $x_0, \ldots, x_7$): $g_{MN} \, \mathrm{d}X^M \mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, ${g^M}_N = \delta_N^M$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|} \, \mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- c_{A}: The coefficient of the x_0-chiral vector portal I13. Unit: dimensionless. Value: fixed through $G_{\mathrm{A}} = 1.278 \times 10^{-9}$ GeV^{-2} (dark-matter relic). Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- Ψ, $\bar{\Psi}$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar{\Psi} = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4, 4)$; $\gamma^M = {\mathbf{e}_a}^M \gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- P_a, P_b: The projectors on the two x_0-chiralities, $P_a = (1 + \gamma^{x_0})/2$ and $P_b = (1 - \gamma^{x_0})/2$ (in a block: the upper and the lower component). Unit: dimensionless. Status: INVENTED.
- J_M^B: The baryon-number current of the quarks (one third per quark), summed over the generations. Unit: GeV^7. Status: STANDARD.

- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- Q_i, $\bar{Q}_i$: The quark doublet of the generation i (left-handed up and down quark of one generation, three colours) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- u_R, u_{Ri}, $\bar{u}_{Ri}$: The up-type quark singlet: u_R the multiplet, u_{Ri} that of the generation i (u_R, c_R, t_R), $\bar{u}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- d_R, d_{Ri}, $\bar{d}_{Ri}$: The down-type quark singlet: d_R the multiplet, d_{Ri} that of the generation i (d_R, s_R, b_R), $\bar{d}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.

The Kohn-Sham equations with the mean fields of the portals (step 3):

$$h\,\chi_o = \varepsilon_o\,\chi_o, \qquad h = \varsigma\Big[-\frac{\mathrm{i}}{f}\sigma_1\partial_y + M_{\mathrm{eff}}(y)\,\sigma_2 + \mathrm{e}^{-a_4}k\,\sigma_3\Big] + v(y) + \frac{1-\sigma_3}{2}\,v_{\mathrm{b}}(y) \tag{FE-63}$$

Symbols in (FE-63).

- h: The block Hamiltonian: the 2×2 differential operator of one block that generates the time evolution of χ, $\mathrm{i}\,\partial_{x_4}\chi = h\chi$; its eigenvalues are the energies ε. Unit: GeV. Status: DERIVED.
- χ, χ_o, χ_1, χ_2: The two-component spinor of one block of the reduction (a function of y and x_4), $\chi = (\chi_1, \chi_2)^T = (a, \mathrm{i}b)^T$; χ_o the Kohn-Sham orbital o, normalised as $\int_{-L}^{0} \chi_o^\dagger \chi_o \,\mathrm{d}y = 1$. Unit: $\mathrm{GeV}^{1/2}$ (normalised orbital). Status: DERIVED.
- o: The label of a Kohn-Sham orbital (a state of dirac16complex in the cell: block, momentum shell, fold parity and level); a sum over o runs over the orbitals. Status: RECORD.
- ε, ε_o: The energy of a fermion mode (an eigenvalue of the block Hamiltonian; at $k = 0$ the four-dimensional mass); ε_o the Kohn-Sham level of the orbital o. Unit: GeV. Status: DERIVED.
- ς: The first of the three signs $(\varsigma, \varsigma_2, \varsigma_3)$, each ± 1, that label the eight blocks of the reduction. Unit: dimensionless. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- σ_1, σ_2, σ_3: The Pauli matrices acting on the two components of a block spinor χ ($\sigma_3 = \mathrm{diag}(1, -1)$). Unit: dimensionless. Status: STANDARD.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- y: The label y in ∂_y: the derivative with respect to the hidden coordinate y. Status: STANDARD.
- $M_{\mathrm{eff}}(y)$: The effective (scalar) mass field of the Kohn-Sham Hamiltonian: m_Ψ plus the mean fields of the contact term and of I12. Unit: GeV. Status: DERIVED.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- $v(y)$: The vector potential of the block Hamiltonian (the mean field along γ^{x_4}): for dirac16complex the contact term and the mean field of I11; zero for a Standard-Model fermion. Unit: GeV. Status: DERIVED.
- $v_{\mathrm{b}}(y)$: The mean field of I13 (baryon densities), acting on the lower component of χ only. Unit: GeV. Status: DERIVED.

$$M_{\rm eff}(y) = m_\Psi + \frac{15}{16}\,\lambda\, n^{\rm sc}(y) + \frac{\langle \mathcal{F}\rangle}{\Lambda_\star^3}, \qquad v(y) = -\frac{\lambda}{16}\, n(y) + \frac{c_{\rm V}}{\Lambda_\star^6}\sum_{i=1}^{3}\sum_{\psi} q_\psi\, n_{\psi i}(y),$$
$$v_{\rm b}(y) = \frac{c_{\rm A}}{\Lambda_\star^6}\sum_{i=1}^{3}\sum_{\psi} q_\psi^B\, n_{\psi i}(y) \tag{FE-64}$$

Symbols in (FE-64).

- $M_{\rm eff}(y)$: The effective (scalar) mass field of the Kohn-Sham Hamiltonian: m_Ψ plus the mean fields of the contact term and of I12. Unit: GeV. Status: DERIVED.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex, $m_\Psi = 3H\mathcal{M} = H$. Unit: GeV. Value: 14.296 TeV. Status: RECORD.
- λ: The contact self-coupling of dirac16complex (the term $-\frac{\lambda}{2}(\bar\Psi\Psi)^2$ of the record). Unit: GeV^{-6}. Value: not fixed; Kohn-Sham states computed for $\lambda = 0$ and $\pm 7\times 10^{-5}H^{-6}$; the fold-band radiation requires $|\lambda| \lesssim 1.4\times 10^{-10}H^{-6}$ (for $\Delta N_{\rm eff} \le 0.001$). Status: RECORD.
- $n^{\rm sc}$, $n_o^{\rm sc}$: The scalar density of dirac16complex (the Kohn-Sham expectation of $\bar\Psi\Psi$, per proper 7-volume) and $n_o^{\rm sc}$ the contribution of the orbital o. Unit: GeV^7. Status: RECORD.
- $\mathcal{F}$: The gauge-kinetic operator of the portal I12, $\mathcal{F} = c_{\rm G}G^A_{MN}G^{AMN} + c_{\rm W}W^I_{MN}W^{IMN} + c_{\rm B}B_{MN}B^{MN}$; $\langle\mathcal{F}\rangle$ its mean value in an environment. Unit: GeV^4. Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- $v(y)$: The vector potential of the block Hamiltonian (the mean field along γ^{x_4}): for dirac16complex the contact term and the mean field of I11; zero for a Standard-Model fermion. Unit: GeV. Status: DERIVED.
- n, n_o, $n_{\psi i}$, n_f, n_i: A number density: $n(y)$ the particle density of dirac16complex in the Kohn-Sham equations and n_o the contribution of the orbital o, $n_{\psi i}(y)$ the density of the Standard-Model fermion ψ_i (per proper 7-volume of eight dimensions); n_f the density of the fermion component f and n_i that of the neutrino mass state i (per 3-volume of four dimensions). Unit: GeV^7 (eight dimensions), GeV^3 (four dimensions). Status: STANDARD.
- $c_{\rm V}$: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_{\rm V} = 4.38\times 10^{-17}$ GeV^{-2}. Status: INVENTED.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- ψ: The label of a Standard-Model multiplet: $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ (quark doublet, up and down quark singlets, lepton doublet, charged-lepton singlet, right-handed neutrino); a sum over ψ runs over the six. Status: STANDARD.
- q_ψ, q_f: The $B - L$ charge: q_ψ of the multiplet ψ ($q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}$, $q_\ell = q_{e_R} = q_{\nu_R} = -1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- $v_{\rm b}(y)$: The mean field of I13 (baryon densities), acting on the lower component of χ only. Unit: GeV. Status: DERIVED.
- $c_{\rm A}$: The coefficient of the x_0-chiral vector portal I13. Unit: dimensionless. Value: fixed through $G_{\rm A} = 1.278\times 10^{-9}$ GeV^{-2} (dark-matter relic). Status: INVENTED.
- q_ψ^B, q_f^B: The baryon number of the multiplet ψ or of the fermion component f ($\frac{1}{3}$ for quarks, 0 for leptons). Unit: dimensionless. Status: STANDARD.

On the zero modes the portals I11 and I13 are four-dimensional contact interactions, and I12 couples the dark matter to the gauge-field invariants:

$$\mathcal{L}_4 = \big(G_{\rm V}\, q_f\, \tilde{I}_{of} + G_{\rm A}\, q_f^B\, \tilde{I}^{(b)}_{of}\big)\big(\mathrm{i}\bar{\Psi}_o\gamma^\mu\Psi_o\big)\big(\mathrm{i}\bar{\psi}_f\gamma_\mu\psi_f\big),$$
$$\tilde{I}_{of} = \frac{1}{H}\int_{-L}^{0}\big(a_o^2+b_o^2\big)\varrho_f\,\mathrm{e}^{-3Hy}\,\mathrm{d}y, \qquad \tilde{I}^{(b)}_{of} = \frac{1}{H}\int_{-L}^{0} b_o^2\,\varrho_f\,\mathrm{e}^{-3Hy}\,\mathrm{d}y, \qquad G_{\rm V} = \frac{c_{\rm V}H}{2f^4 v_{\rm t}\Lambda_\star^6}, \qquad G_{\rm A} = \frac{c_{\rm A}H}{2f^4 v_{\rm t}\Lambda_\star^6},$$
$$n_{\psi i}(y) = \frac{n_f\,\varrho_f(y)\,\mathrm{e}^{-3Hy}}{2f^4 v_{\rm t}}, \qquad \varrho(y;c) = \frac{2c}{L}\,\frac{\mathrm{e}^{2cy/L}}{1-\mathrm{e}^{-2c}} \tag{FE-70}$$

Symbols in (FE-70).

- $\mathcal{L}_4$: The four-dimensional contact Lagrangian of I11 and I13 between a state o of dirac16complex and a Standard-Model fermion component f. Unit: GeV^4. Status: DERIVED.
- $G_{\rm V}$: The four-dimensional strength of the vector portal I11 on the zero modes, $G_{\rm V} = c_{\rm V}H/(2f^4 v_{\rm t}\Lambda_\star^6)$. Unit: GeV^{-2}. Value: benchmark 4.38×10^{-17} GeV^{-2} ($\Delta N_{\rm eff} = 0.001$). Status: DERIVED.
- q_ψ, q_f: The $B-L$ charge: q_ψ of the multiplet ψ ($q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}$, $q_\ell = q_{e_R} = q_{\nu_R} = -1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- f: The label of a fermion species: a Standard-Model multiplet of one generation (its $N_{\rm c}$ Weyl components share one zero-mode profile; the zero mode of each chirality) or a fermion of a medium (electron, proton, neutron); a sum over f runs over these species. Status: STANDARD.
- $\tilde{I}_{of}$: The overlap of the state o of dirac16complex with the zero mode of the fermion component f, $\tilde{I}_{of} = H^{-1}\int_{-L}^{0}(a_o^2+b_o^2)\varrho_f\mathrm{e}^{-3Hy}\mathrm{d}y$ (the integral in units of H); $\tilde{I}_Q$, $\tilde{I}_{u_R}$, $\tilde{I}_{d_R}$, $\tilde{I}_\ell$, $\tilde{I}_{e_R}$ those of the dark matter with Q, u_R, d_R, ℓ, e_R, $\tilde{I}_{\nu_R,i}$ with the right-handed neutrino of the generation i. Unit: dimensionless. Value: dark matter: $\tilde{I}_Q = 8.2722$, $\tilde{I}_\ell = 1.2524$, $\tilde{I}_{u_R} = 0.3854$, $\tilde{I}_{d_R} = 0.3961$, $\tilde{I}_{e_R} = 0.3972$. Status: DERIVED.
- o: The label of a Kohn-Sham orbital (a state of dirac16complex in the cell: block, momentum shell, fold parity and level); a sum over o runs over the orbitals. Status: RECORD.
- $G_{\rm A}$: The four-dimensional strength of the portal I13 on the zero modes, $G_{\rm A} = c_{\rm A}H/(2f^4 v_{\rm t}\Lambda_\star^6)$. Unit: GeV^{-2}. Value: 1.278×10^{-9} GeV^{-2} (dark-matter relic density). Status: DERIVED.
- q_ψ^B, q_f^B: The baryon number of the multiplet ψ or of the fermion component f ($\frac{1}{3}$ for quarks, 0 for leptons). Unit: dimensionless. Status: STANDARD.
- $\tilde{I}^{(b)}_{of}$: The overlap of the lower component of the state o with the zero mode of f, $\tilde{I}^{(b)}_{of} = H^{-1}\int_{-L}^{0} b_o^2\varrho_f\mathrm{e}^{-3Hy}\mathrm{d}y$ (the overlap of I13); $\tilde{I}^{(b)}_Q$, $\tilde{I}^{(b)}_{u_R}$, $\tilde{I}^{(b)}_{d_R}$, $\tilde{I}^{(b)}_{\nu_R}$ those of the dark matter. Unit: dimensionless. Value: dark matter: $\tilde{I}^{(b)}_Q = 2.6135$. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- Ψ, $\bar{\Psi}$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar{\Psi} = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- ψ_i, $\bar{\psi}_i$: A Standard-Model fermion field of eight dimensions: ψ_i is the multiplet $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ of the generation i, ψ_f one fermion component f (in four-dimensional formulas its zero mode); $\bar{\psi} = \psi^\dagger C$ the Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$ (eight dimensions), $\mathrm{GeV}^{3/2}$ (four dimensions). Status: ASSUMED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of $m_{\rm W}$ and $\rho_{\rm EW,0}$). Status: AUTHOR.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926\times10^{-21}$ m at the stabilised block). Status: ASSUMED.
- a, b: The upper and lower components of a block spinor, $\chi = (a, \mathrm{i}b)$, real at $k = 0$: a_L, b_L of a doublet and a_R, b_R of a singlet solution (a_{Ri}, b_{Ri} of the generation i, $b_{L\alpha}$, $a_{L\alpha}$ of the doublet in the flavour basis), a_o, b_o of the orbital o. Unit: $\mathrm{GeV}^{1/2}$ (normalised orbital). Status: DERIVED.
- $\varrho_f(y)$, $\varrho(y;c)$: The x_0 profile of a zero mode as a probability density on one half, $\int_{-L}^{0}\varrho\,\mathrm{d}y = 1$: ϱ_f that of

the fermion component f, $\varrho(y;c)$ its form for the localisation c. Unit: 1/length (GeV). Status: DERIVED.

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- c_{V}: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_{\mathrm{V}} = 4.38\times10^{-17}$ GeV^{-2}. Status: INVENTED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- v_{t}: The coordinate volume of the compact extra times x_5, x_6, x_7. Unit: GeV^{-3}. Value: not fixed; $v_{\mathrm{t}} = H^{-3}$ in the quoted conversions. Status: ASSUMED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- c_{A}: The coefficient of the x_0-chiral vector portal I13. Unit: dimensionless. Value: fixed through $G_{\mathrm{A}} = 1.278 \times 10^{-9}$ GeV^{-2} (dark-matter relic). Status: INVENTED.
- n, n_o, $n_{\psi i}$, n_f, n_i: A number density: $n(y)$ the particle density of dirac16complex in the Kohn-Sham equations and n_o the contribution of the orbital o, $n_{\psi i}(y)$ the density of the Standard-Model fermion ψ_i (per proper 7-volume of eight dimensions); n_f the density of the fermion component f and n_i that of the neutrino mass state i (per 3-volume of four dimensions). Unit: GeV^7 (eight dimensions), GeV^3 (four dimensions). Status: STANDARD.
- ψ: The label of a Standard-Model multiplet: $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ (quark doublet, up and down quark singlets, lepton doublet, charged-lepton singlet, right-handed neutrino); a sum over ψ runs over the six. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- c, c_L, c_R: The localisation of a zero mode, $c = f\mu L$ (bulk mass times the proper depth); c_L of a doublet (localised at the fold for $c_L > 0$), c_R of a singlet (localised at the fold for $c_R < 0$). Unit: dimensionless. Value: doublets $c_L = 3.38230$ (quarks) and 5.95288 (leptons); singlets c_R: up -15.4300, down -14.6258, strange -11.5158, charm -8.7683, bottom -7.5005, top -3.3823, electron -14.5539, muon -8.9809, tau -5.9529, neutrinos -35.0452, -32.8481, -31.0702 (fixed by the measured masses). Status: DERIVED.

$$\mathcal{L}_4^{\mathrm{I12}} = -\mathcal{O}_{\mathrm{DM}}\,\frac{\bar\Psi_{\mathrm{DM}}\Psi_{\mathrm{DM}}}{\Lambda_\star^3}\big(c_{\mathrm{G}}\,G^A_{\mu\nu}G^{A\mu\nu} + c_{\mathrm{W}}\,W^I_{\mu\nu}W^{I\mu\nu} + c_{\mathrm{B}}\,B_{\mu\nu}B^{\mu\nu}\big), \qquad \mathcal{O}_{\mathrm{DM}} = \frac{\partial\varepsilon_{\mathrm{DM}}}{\partial m_\Psi},$$
$$G_{\mathrm{g}} = \mathcal{O}_{\mathrm{DM}}\,\frac{c_{\mathrm{G}}\,g_3^2}{\Lambda_\star^3}, \qquad G_{\mathrm{W}} = \mathcal{O}_{\mathrm{DM}}\,\frac{c_{\mathrm{W}}\,g_2^2}{\Lambda_\star^3}, \qquad G_{\mathrm{B}} = \mathcal{O}_{\mathrm{DM}}\,\frac{c_{\mathrm{B}}\,g_1^2}{\Lambda_\star^3} \tag{FE-71}$$

Symbols in (FE-71).

- $\mathcal{L}_4^{\mathrm{I12}}$: The four-dimensional Lagrangian of I12 on the dark matter. Unit: GeV^4. Status: DERIVED.
- $\mathcal{O}_o$, $\mathcal{O}_{\mathrm{DM}}$: The scalar overlap of a state of dirac16complex, $\mathcal{O}_o = \partial\varepsilon_o/\partial m_\Psi$ (Hellmann-Feynman: the integrated scalar density of the orbital); $\mathcal{O}_{\mathrm{DM}}$ that of the dark matter. Unit: dimensionless. Value: $\mathcal{O}_{\mathrm{DM}} = 0.7072$. Status: DERIVED.
- Ψ_{DM}, $\bar\Psi_{\mathrm{DM}}$: The four-dimensional field of a dark-matter quantum (the lowest odd level of dirac16complex on the static block, in either half) and its Dirac adjoint. Unit: GeV$^{3/2}$. Status: DERIVED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- c_{G}: The coefficient of the gluon term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{G}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- G^A_M, G_M, G_{MN}, G^A_{MN}: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G^A_M ($A = 1, \ldots, 8$), the matrix $G_M = G^A_M t^A$, the field strength G_{MN} (matrix) and its components G^A_{MN}; $G^A_{\mu\nu}$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.

- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index running over $x_1, \ldots, x_4$ (as μ). Status: STANDARD.
- c_{W}: The coefficient of the SU(2) term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{W}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- W_M^I, W_M, W_{MN}, W_{MN}^I: The SU(2) gauge field (geometric normalisation): the components W_M^I ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W_M^I \tau^I/2$, the field strength W_{MN} and its components W_{MN}^I; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- c_{B}: The coefficient of the hypercharge term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{B}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- $\varepsilon_{\mathrm{DM}}$: The energy of the dark-matter level at rest (its mass M_{DM} on the static block). Unit: GeV. Value: $3.102204\, m_\Psi = 44.35$ TeV. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex, $m_\Psi = 3H\mathcal{M} = H$. Unit: GeV. Value: 14.296 TeV. Status: RECORD.
- G_{g}: The coupling of a dark-matter pair to canonically normalised gluon pairs through I12. Unit: GeV^{-3}. Value: 6.09×10^{-15} GeV^{-3} (benchmark). Status: DERIVED.
- g_3: The four-dimensional strong coupling ($g_3^2 = 4\pi\alpha_3$). Unit: dimensionless. Value: $g_3^2 = 0.7786$ at $2M_{\mathrm{DM}}$. Status: DERIVED.
- G_{W}: The coupling of a dark-matter pair to canonically normalised SU(2) gauge pairs through I12. Unit: GeV^{-3}. Value: 2.96×10^{-15} GeV^{-3}. Status: DERIVED.
- g_2: The four-dimensional SU(2) coupling ($g_2^2 = 4\pi\alpha_2$). Unit: dimensionless. Value: $g_2^2 = 0.3781$ at $2M_{\mathrm{DM}}$. Status: DERIVED.
- G_{B}: The coupling of a dark-matter pair to canonically normalised hypercharge pairs through I12. Unit: GeV^{-3}. Value: 1.08×10^{-15} GeV^{-3}. Status: DERIVED.
- g_1: The four-dimensional hypercharge coupling ($g_1^2 = 4\pi\alpha_1$, Standard-Model normalisation without the GUT factor). Unit: dimensionless. Value: $g_1^2 = 0.1380$ at $2M_{\mathrm{DM}}$. Status: DERIVED.

The shift of a Kohn-Sham level by the mean fields of an environment, and in matter of protons, neutrons and electrons:

$$\Delta\varepsilon_o = \sum_f \left(G_{\mathrm{V}}\, q_f\, \tilde{I}_{of} + G_{\mathrm{A}}\, q_f^B\, \tilde{I}_{of}^{(b)}\right) n_f + \mathcal{O}_o\, \frac{c_{\mathrm{I12}}}{\Lambda_\star^3}\left[\langle G_{\mu\nu}^A G^{A\mu\nu}\rangle + \langle W_{\mu\nu}^I W^{I\mu\nu}\rangle + \langle B_{\mu\nu} B^{\mu\nu}\rangle\right], \qquad \mathcal{O}_o = \frac{\partial\varepsilon_o}{\partial m_\Psi} \quad \text{(FE-72)}$$

Symbols in (FE-72).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\, \Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- ε, ε_o: The energy of a fermion mode (an eigenvalue of the block Hamiltonian; at $k = 0$ the four-dimensional mass); ε_o the Kohn-Sham level of the orbital o. Unit: GeV. Status: DERIVED.
- o: The label of a Kohn-Sham orbital (a state of dirac16complex in the cell: block, momentum shell, fold parity and level); a sum over o runs over the orbitals. Status: RECORD.
- f: The label of a fermion species: a Standard-Model multiplet of one generation (its N_{c} Weyl components share one zero-mode profile; the zero mode of each chirality) or a fermion of a medium (electron, proton, neutron); a sum over f runs over these species. Status: STANDARD.
- G_{V}: The four-dimensional strength of the vector portal I11 on the zero modes, $G_{\mathrm{V}} = c_{\mathrm{V}} H/(2f^4 v_{\mathrm{t}} \Lambda_\star^6)$. Unit: GeV^{-2}. Value: benchmark 4.38×10^{-17} GeV^{-2} ($\Delta N_{\mathrm{eff}} = 0.001$). Status: DERIVED.

- q_ψ, q_f: The $B-L$ charge: q_ψ of the multiplet ψ ($q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}$, $q_\ell = q_{e_R} = q_{\nu_R} = -1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- $\tilde{I}_{of}$: The overlap of the state o of dirac16complex with the zero mode of the fermion component f, $\tilde{I}_{of} = H^{-1}\int_{-L}^{0}(a_o^2+b_o^2)\varrho_f \mathrm{e}^{-3Hy}\mathrm{d}y$ (the integral in units of H); $\tilde{I}_Q$, $\tilde{I}_{u_R}$, $\tilde{I}_{d_R}$, $\tilde{I}_\ell$, $\tilde{I}_{e_R}$ those of the dark matter with Q, u_R, d_R, ℓ, e_R, $\tilde{I}_{\nu_R,i}$ with the right-handed neutrino of the generation i. Unit: dimensionless. Value: dark matter: $\tilde{I}_Q = 8.2722$, $\tilde{I}_\ell = 1.2524$, $\tilde{I}_{u_R} = 0.3854$, $\tilde{I}_{d_R} = 0.3961$, $\tilde{I}_{e_R} = 0.3972$. Status: DERIVED.
- G_A: The four-dimensional strength of the portal I13 on the zero modes, $G_\mathrm{A} = c_\mathrm{A} H/(2f^4 v_\mathrm{t}\Lambda_\star^6)$. Unit: GeV^{-2}. Value: 1.278×10^{-9} GeV^{-2} (dark-matter relic density). Status: DERIVED.
- q_ψ^B, q_f^B: The baryon number of the multiplet ψ or of the fermion component f ($\frac{1}{3}$ for quarks, 0 for leptons). Unit: dimensionless. Status: STANDARD.
- $\tilde{I}_{of}^{(b)}$: The overlap of the lower component of the state o with the zero mode of f, $\tilde{I}_{of}^{(b)} = H^{-1}\int_{-L}^{0} b_o^2 \varrho_f \mathrm{e}^{-3Hy}\mathrm{d}y$ (the overlap of I13); $\tilde{I}_Q^{(b)}$, $\tilde{I}_{u_R}^{(b)}$, $\tilde{I}_{d_R}^{(b)}$, $\tilde{I}_{\nu_R}^{(b)}$ those of the dark matter. Unit: dimensionless. Value: dark matter: $\tilde{I}_Q^{(b)} = 2.6135$. Status: DERIVED.
- n, n_o, $n_{\psi i}$, n_f, n_i: A number density: $n(y)$ the particle density of dirac16complex in the Kohn-Sham equations and n_o the contribution of the orbital o, $n_{\psi i}(y)$ the density of the Standard-Model fermion ψ_i (per proper 7-volume of eight dimensions); n_f the density of the fermion component f and n_i that of the neutrino mass state i (per 3-volume of four dimensions). Unit: GeV^7 (eight dimensions), GeV^3 (four dimensions). Status: STANDARD.
- $\mathcal{O}_o$, $\mathcal{O}_\mathrm{DM}$: The scalar overlap of a state of dirac16complex, $\mathcal{O}_o = \partial\varepsilon_o/\partial m_\Psi$ (Hellmann-Feynman: the integrated scalar density of the orbital); $\mathcal{O}_\mathrm{DM}$ that of the dark matter. Unit: dimensionless. Value: $\mathcal{O}_\mathrm{DM} = 0.7072$. Status: DERIVED.
- c_I12: The common coefficient of the three terms of I12 when they are universal ($c_\mathrm{G} = c_\mathrm{W} = c_\mathrm{B} = c_\mathrm{I12}$). Unit: dimensionless. Value: $c_\mathrm{I12}/\Lambda_\star^3 = 1.106\times10^{-14}$ GeV^{-3} (unitarity-safe benchmark). Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- G_M^A, G_M, G_{MN}, G_{MN}^A: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G_M^A ($A = 1,\ldots,8$), the matrix $G_M = G_M^A t^A$, the field strength G_{MN} (matrix) and its components G_{MN}^A; $G_{\mu\nu}^A$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- A: A colour (adjoint SU(3)) index running over $1,\ldots,8$; a repeated index is summed. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index running over $x_1,\ldots,x_4$ (as μ). Status: STANDARD.
- W_M^I, W_M, W_{MN}, W_{MN}^I: The SU(2) gauge field (geometric normalisation): the components W_M^I ($I = 1,2,3$; W^3 the third component), the matrix $W_M = W_M^I \tau^I/2$, the field strength W_{MN} and its components W_{MN}^I; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1,2,3$; a repeated index is summed. Status: STANDARD.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex, $m_\Psi = 3H\mathcal{M} = H$. Unit: GeV. Value: 14.296 TeV. Status: RECORD.

$$\begin{aligned}\Delta\varepsilon_\mathrm{I11} &= G_\mathrm{V}\Big[\frac{1}{3}\,n_\mathrm{u}\,\frac{\tilde{I}_Q+\tilde{I}_{u_R}}{2}+\frac{1}{3}\,n_\mathrm{d}\,\frac{\tilde{I}_Q+\tilde{I}_{d_R}}{2}-n_\mathrm{e}\,\frac{\tilde{I}_\ell+\tilde{I}_{e_R}}{2}\Big],\\ \Delta\varepsilon_\mathrm{I13} &= G_\mathrm{A}\Big[\frac{1}{3}\,n_\mathrm{u}\,\frac{\tilde{I}_Q^{(b)}+\tilde{I}_{u_R}^{(b)}}{2}+\frac{1}{3}\,n_\mathrm{d}\,\frac{\tilde{I}_Q^{(b)}+\tilde{I}_{d_R}^{(b)}}{2}\Big], \qquad n_\mathrm{u} = 2n_\mathrm{p}+n_\mathrm{n}, \qquad n_\mathrm{d} = n_\mathrm{p}+2n_\mathrm{n}\end{aligned} \tag{FE-73}$$

Symbols in (FE-73).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_eff

an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.

- $\Delta\varepsilon_{\mathrm{I11}}$, $\Delta\varepsilon_{\mathrm{I13}}$: The shifts of the dark-matter level by the mean fields of I11 and of I13 in matter. Unit: GeV. Value: at most 8.2×10^{-10} GeV (I13, at the freeze-out). Status: DERIVED.
- G_{V}: The four-dimensional strength of the vector portal I11 on the zero modes, $G_{\mathrm{V}} = c_{\mathrm{V}}H/(2f^4 v_{\mathrm{t}}\Lambda_\star^6)$. Unit: GeV^{-2}. Value: benchmark 4.38×10^{-17} GeV^{-2} ($\Delta N_{\mathrm{eff}} = 0.001$). Status: DERIVED.
- n_{u}, n_{d}: The number densities of up and down quarks bound in nucleons, $n_{\mathrm{u}} = 2n_{\mathrm{p}} + n_{\mathrm{n}}$, $n_{\mathrm{d}} = n_{\mathrm{p}} + 2n_{\mathrm{n}}$. Unit: GeV^3. Status: STANDARD.
- $\tilde{I}_{of}$: The overlap of the state o of dirac16complex with the zero mode of the fermion component f, $\tilde{I}_{of} = H^{-1}\int_{-L}^{0}(a_o^2 + b_o^2)\varrho_f \mathrm{e}^{-3Hy}\mathrm{d}y$ (the integral in units of H); $\tilde{I}_Q$, $\tilde{I}_{u_R}$, $\tilde{I}_{d_R}$, $\tilde{I}_\ell$, $\tilde{I}_{e_R}$ those of the dark matter with Q, u_R, d_R, ℓ, e_R, $\tilde{I}_{\nu_R,i}$ with the right-handed neutrino of the generation i. Unit: dimensionless. Value: dark matter: $\tilde{I}_Q = 8.2722$, $\tilde{I}_\ell = 1.2524$, $\tilde{I}_{u_R} = 0.3854$, $\tilde{I}_{d_R} = 0.3961$, $\tilde{I}_{e_R} = 0.3972$. Status: DERIVED.
- n_{e}: The number density of electrons (net of positrons); in the Thomson terms of the photon and baryon equations only free (ionised) electrons count. Unit: GeV^3. Status: STANDARD.
- G_{A}: The four-dimensional strength of the portal I13 on the zero modes, $G_{\mathrm{A}} = c_{\mathrm{A}}H/(2f^4 v_{\mathrm{t}}\Lambda_\star^6)$. Unit: GeV^{-2}. Value: 1.278×10^{-9} GeV^{-2} (dark-matter relic density). Status: DERIVED.
- $\tilde{I}_{of}^{(b)}$: The overlap of the lower component of the state o with the zero mode of f, $\tilde{I}_{of}^{(b)} = H^{-1}\int_{-L}^{0} b_o^2\varrho_f \mathrm{e}^{-3Hy}\mathrm{d}y$ (the overlap of I13); $\tilde{I}_Q^{(b)}$, $\tilde{I}_{u_R}^{(b)}$, $\tilde{I}_{d_R}^{(b)}$, $\tilde{I}_{\nu_R}^{(b)}$ those of the dark matter. Unit: dimensionless. Value: dark matter: $\tilde{I}_Q^{(b)} = 2.6135$. Status: DERIVED.
- n_{p}: The number density of protons (free and bound). Unit: GeV^3. Status: STANDARD.
- n_{n}: The number density of neutrons (free and bound). Unit: GeV^3. Status: STANDARD.

$$
\begin{aligned}
&n_{\mathrm{p}} + n_{\mathrm{n}} = \eta_{\mathrm{b}}\, n_\gamma, \qquad n_\gamma = \frac{2\zeta(3)}{\pi^2}\,T^3, \qquad \frac{n_{\mathrm{n}}}{n_{\mathrm{p}}} = \mathrm{e}^{-(m_{\mathrm{n}}-m_{\mathrm{p}})/T}, \qquad n_{\mathrm{e}} = n_{\mathrm{p}},\\
&n_{\mathrm{DM,tot}} = 2\big(n_{\mathrm{DM}} + \bar{n}_{\mathrm{DM}}\big) = 2\cdot 4\left(\frac{M_{\mathrm{DM}}T}{2\pi}\right)^{3/2}\mathrm{e}^{-M_{\mathrm{DM}}/T} \quad \left(\frac{M_{\mathrm{DM}}}{T} < 40\right)
\end{aligned}
\tag{FE-86}
$$

Symbols in (FE-86).

- n_{p}: The number density of protons (free and bound). Unit: GeV^3. Status: STANDARD.
- n_{n}: The number density of neutrons (free and bound). Unit: GeV^3. Status: STANDARD.
- η_{b}: The ratio of the baryon density to the photon density. Unit: dimensionless. Value: 6.1×10^{-10}. Status: STANDARD.
- n_γ: The number density of photons. Unit: GeV^3. Status: STANDARD.
- ζ: The Riemann zeta function; only its value $\zeta(3)$ occurs, the sum of $1/n^3$ over the positive integers n. Unit: dimensionless. Value: $\zeta(3) = 1.2020569$. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- m_{p}, m_{n}: The masses of the proton and the neutron. Unit: GeV. Value: 0.938272 GeV, 0.939565 GeV. Status: STANDARD.
- n_{e}: The number density of electrons (net of positrons); in the Thomson terms of the photon and baryon equations only free (ionised) electrons count. Unit: GeV^3. Status: STANDARD.
- $n_{\mathrm{DM,tot}}$: The total number density of dark-matter quanta: both species (the lowest odd levels of Ψ_A and Ψ_B), particles and antiparticles, $n_{\mathrm{DM,tot}} = 2(n_{\mathrm{DM}} + \bar{n}_{\mathrm{DM}})$. Unit: GeV^3. Status: STANDARD.
- n_{DM}, $\bar{n}_{\mathrm{DM}}$: The number densities of dark-matter particles and antiparticles of one species. Unit: GeV^3. Status: STANDARD.
- M_{DM}: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.

The condensates of the gauge fields (the mean fields of I12):

$$\big\langle G^A_{\mu\nu}G^{A\mu\nu}\big\rangle(T) = \big\langle G^A_{\mu\nu}G^{A\mu\nu}\big\rangle(0) - \frac{32\pi^2}{b_3}\,\Theta_{\rm QCD}, \qquad \big\langle G^A_{\mu\nu}G^{A\mu\nu}\big\rangle(0) = 4\pi^2\times 0.012\ {\rm GeV}^4, \qquad b_3 = 11 - \frac{2}{3}\,N_{\rm f} \tag{FE-74}$$

Symbols in (FE-74).

- G^A_M, G_M, G_{MN}, G^A_{MN}: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G^A_M ($A = 1, \dots, 8$), the matrix $G_M = G^A_M t^A$, the field strength G_{MN} (matrix) and its components G^A_{MN}; $G^A_{\mu\nu}$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), $\rm GeV^2$ (field strength). Status: ASSUMED.
- A: A colour (adjoint SU(3)) index running over $1, \dots, 8$; a repeated index is summed. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index running over $x_1, \dots, x_4$ (as μ). Status: STANDARD.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- b_3, b_2, b_1, $b_{\rm G}$: The one-loop coefficients of the running couplings, $\mathrm{d}\alpha/\mathrm{d}\ln\mu_{\rm ren} = -b\,\alpha^2/(2\pi)$: $b_3 = 11 - \frac{2}{3}N_{\rm f}$ (SU(3)), $b_2 = \frac{10}{3}$ (SU(2) without a Higgs particle), $b_1 = -\frac{20}{3}$ ($\mathrm{U}(1)_Y$), $b_{\rm G}$ that of the group G (not related to the metric function b_4). Unit: dimensionless. Status: STANDARD.
- $\Theta_{\rm QCD}$, $\Theta_{\rm G}$, Θ_V: The trace $\rho - 3p$ of the energy-momentum tensor (interaction measure) of a plasma component: $\Theta_{\rm QCD}$ of the quark-gluon plasma or hadron gas, $\Theta_{\rm G}$ of the gauge plasma of the group G, Θ_V of the thermal gas of the massive vector V ($\Theta_{\rm W}$ of $W^\pm$, $\Theta_{\rm Z}$ of Z). Unit: $\rm GeV^4$. Status: STANDARD.
- $N_{\rm f}$: The number of active quark flavours (quarks lighter than the scale). Unit: dimensionless. Status: STANDARD.

$$\frac{\Theta_{\rm QCD}}{T^4} = 4\exp\Big[-\frac{\ln^2(T/0.2\,{\rm GeV})}{0.5}\Big]\ \ (T<1\,{\rm GeV}), \qquad \frac{\Theta_{\rm QCD}}{T^4} = \frac{b_3}{3}\Big(1+\frac{5N_{\rm f}}{12}\Big)\alpha_3^2\ \ (T>1\,{\rm GeV}),$$
$$p_{\rm QCD} = \frac{8\pi^2}{45}T^4\Big[1+\frac{21N_{\rm f}}{32}-\frac{15}{4}\Big(1+\frac{5N_{\rm f}}{12}\Big)\frac{\alpha_3}{\pi}\Big], \qquad \frac{\mathrm{d}\alpha_3}{\mathrm{d}\ln T} = -\frac{b_3}{2\pi}\,\alpha_3^2, \qquad \Theta_{\rm QCD} = T^5\,\frac{\mathrm{d}}{\mathrm{d}T}\Big(\frac{p_{\rm QCD}}{T^4}\Big) \tag{FE-75}$$

Symbols in (FE-75).

- $\Theta_{\rm QCD}$, $\Theta_{\rm G}$, Θ_V: The trace $\rho - 3p$ of the energy-momentum tensor (interaction measure) of a plasma component: $\Theta_{\rm QCD}$ of the quark-gluon plasma or hadron gas, $\Theta_{\rm G}$ of the gauge plasma of the group G, Θ_V of the thermal gas of the massive vector V ($\Theta_{\rm W}$ of $W^\pm$, $\Theta_{\rm Z}$ of Z). Unit: $\rm GeV^4$. Status: STANDARD.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.
- b_3, b_2, b_1, $b_{\rm G}$: The one-loop coefficients of the running couplings, $\mathrm{d}\alpha/\mathrm{d}\ln\mu_{\rm ren} = -b\,\alpha^2/(2\pi)$: $b_3 = 11 - \frac{2}{3}N_{\rm f}$ (SU(3)), $b_2 = \frac{10}{3}$ (SU(2) without a Higgs particle), $b_1 = -\frac{20}{3}$ ($\mathrm{U}(1)_Y$), $b_{\rm G}$ that of the group G (not related to the metric function b_4). Unit: dimensionless. Status: STANDARD.
- $N_{\rm f}$: The number of active quark flavours (quarks lighter than the scale). Unit: dimensionless. Status: STANDARD.
- α_3: The strong fine-structure constant, $\alpha_3 = g_3^2/(4\pi)$, at the scale given in brackets. Unit: dimensionless. Value: $\alpha_3(m_{\rm Z}) = 0.1180$; $\alpha_3(m_{\rm KK}) = 0.0606$. Status: STANDARD.
- $p_{\rm QCD}$: The pressure of the quark-gluon plasma (leading orders). Unit: $\rm GeV^4$. Status: ESTIMATE.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.

$$\big\langle F^V_{\mu\nu} F^{V\,\mu\nu}\big\rangle(T) = -2\,\Theta_V \quad (\text{for } V = W^\pm, Z), \qquad \big\langle F^2_{\text{geo}}\big\rangle = -4\,\frac{\partial p}{\partial\big(1/g^2_{\text{G}}\big)} = 4 g^2_{\text{G}}\, p_{\alpha,\text{G}},$$
$$p_{\alpha,\text{G}} = -\frac{g^2_{\text{G}} T^4}{144}\, d_{\text{adj}}\Big[C_{\text{adj}} + \frac{5}{2}\, T_{\text{F}} N_{\text{f}}\Big] \tag{FE-76}$$

Symbols in (FE-76).

- $F^V_{\mu\nu}$: The field strength of the canonically normalised massive vector $V = W^\pm$ or Z, $F^V_{\mu\nu} = \partial_\mu V_\nu - \partial_\nu V_\mu$. Unit: GeV^2. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index running over $x_1, \ldots, x_4$ (as μ). Status: STANDARD.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.
- Θ_{QCD}, Θ_{G}, Θ_V: The trace $\rho - 3p$ of the energy-momentum tensor (interaction measure) of a plasma component: Θ_{QCD} of the quark-gluon plasma or hadron gas, Θ_{G} of the gauge plasma of the group G, Θ_V of the thermal gas of the massive vector V (Θ_{W} of $W^\pm$, Θ_{Z} of Z). Unit: GeV^4. Status: STANDARD.
- F_{geo}: The field strength of the gauge group G in geometric normalisation; $\langle F^2_{\text{geo}}\rangle$ is its condensate ($G^A_{\mu\nu} G^{A\mu\nu}$, $W^I_{\mu\nu} W^{I\mu\nu}$ or $B_{\mu\nu} B^{\mu\nu}$). Unit: GeV^2. Status: STANDARD.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4} with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- p: A pressure: $p_M = \mathcal{T}^M{}_M$ (no sum) for the direction $M \neq x_4$, p_s the pressure of the species s; Δp the pressure perturbation in the cosmological perturbation equations; in the Feynman-Hellmann relation the pressure of a gauge plasma. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- g_{G}: The four-dimensional coupling of the gauge group G (one of g_3, g_2, g_1). Unit: dimensionless. Status: STANDARD.
- $p_{\alpha,\text{G}}$, $p_{\alpha,2}$, $p_{\alpha,1}$: The order-α (interaction) part of the pressure of the gauge plasma of the group G; $p_{\alpha,2}$ of SU(2), $p_{\alpha,1}$ of $\text{U}(1)_Y$. Unit: GeV^4. Status: STANDARD.
- d_{adj}: The dimension of the adjoint representation of the gauge group (8, 3, 1). Unit: dimensionless. Status: STANDARD.
- C_{adj}: The quadratic Casimir of the adjoint representation (3, 2, 0). Unit: dimensionless. Status: STANDARD.
- T_{F}: The Dynkin index of the fermion representation ($\frac{1}{2}$ per Dirac fundamental; for $\text{U}(1)_Y$ the sum of the squared hypercharges). Unit: dimensionless. Status: STANDARD.
- N_{f}: The number of active quark flavours (quarks lighter than the scale). Unit: dimensionless. Status: STANDARD.

$$\frac{1}{\alpha_2(\mu_{\text{ren}})} = \frac{\sin^2\theta_{\overline{\text{MS}}}}{\alpha_{\text{em}}} + \frac{b_2}{2\pi}\ln\frac{\mu_{\text{ren}}}{m_{\text{Z}}}, \qquad \frac{1}{\alpha_1(\mu_{\text{ren}})} = \frac{1-\sin^2\theta_{\overline{\text{MS}}}}{\alpha_{\text{em}}} + \frac{b_1}{2\pi}\ln\frac{\mu_{\text{ren}}}{m_{\text{Z}}}, \qquad b_2 = \frac{10}{3}, \quad b_1 = -\frac{20}{3} \tag{FE-77}$$

Symbols in (FE-77).

- α_2: The SU(2) fine-structure constant, $\alpha_2 = g_2^2/(4\pi)$. Unit: dimensionless. Status: STANDARD.
- μ_{ren}: The renormalisation scale of a running coupling (taken as $2\pi T$ for the condensates of a plasma). Unit: GeV. Status: STANDARD.
- $\theta_{\overline{\text{MS}}}$: The weak mixing angle in the $\overline{\text{MS}}$ scheme at m_{Z}. Unit: dimensionless. Value: $\sin^2\theta_{\overline{\text{MS}}} = 0.23122$. Status: STANDARD.
- α_{em}: The electromagnetic fine-structure constant ($\overline{\text{MS}}$, at m_{Z}). Unit: dimensionless. Value: 1/127.95. Status: STANDARD.
- b_3, b_2, b_1, b_{G}: The one-loop coefficients of the running couplings, $\text{d}\alpha/\text{d}\ln\mu_{\text{ren}} = -b\,\alpha^2/(2\pi)$: $b_3 = 11 - \frac{2}{3}N_{\text{f}}$ (SU(3)), $b_2 = \frac{10}{3}$ (SU(2) without a Higgs particle), $b_1 = -\frac{20}{3}$ ($\text{U}(1)_Y$), b_{G} that of the group G (not related to the metric function b_4). Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

- m_{Z}: The mass of the Z. Unit: GeV. Value: 91.1879 GeV (reproduced). Status: STANDARD.
- α_1: The hypercharge fine-structure constant, $\alpha_1 = g_1^2/(4\pi)$. Unit: dimensionless. Status: STANDARD.

$$\langle W^I_{\mu\nu} W^{I\mu\nu}\rangle = g_2^2\Big[2\big(-2\Theta_{\mathrm{W}}\big)+\cos^2\theta_{\mathrm{W}}\big(-2\Theta_{\mathrm{Z}}\big)\Big]+4g_2^2\,p_{\alpha,2}, \qquad \langle B_{\mu\nu}B^{\mu\nu}\rangle = g_1^2\sin^2\theta_{\mathrm{W}}\big(-2\Theta_{\mathrm{Z}}\big)+4g_1^2\,p_{\alpha,1} \quad \text{(FE-78)}$$

Symbols in (FE-78).

- W^I_M, W_M, W_{MN}, W^I_{MN}: The SU(2) gauge field (geometric normalisation): the components W^I_M ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W^I_M \tau^I/2$, the field strength W_{MN} and its components W^I_{MN}; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index running over $x_1, \ldots, x_4$ (as μ). Status: STANDARD.
- g_2: The four-dimensional SU(2) coupling ($g_2^2 = 4\pi\alpha_2$). Unit: dimensionless. Value: $g_2^2 = 0.3781$ at $2M_{\mathrm{DM}}$. Status: DERIVED.
- Θ_{QCD}, Θ_{G}, Θ_V: The trace $\rho - 3p$ of the energy-momentum tensor (interaction measure) of a plasma component: Θ_{QCD} of the quark-gluon plasma or hadron gas, Θ_{G} of the gauge plasma of the group G, Θ_V of the thermal gas of the massive vector V (Θ_{W} of $W^\pm$, Θ_{Z} of Z). Unit: GeV^4. Status: STANDARD.
- θ_{W}: The weak mixing angle of the couplings, $\cos\theta_{\mathrm{W}} = \hat{g}_2/\hat{g}_Z$, $\sin\theta_{\mathrm{W}} = \hat{g}_1/\hat{g}_Z$; fixed by the measured W and Z masses with $\rho_{\mathrm{EW},0}$. Unit: dimensionless. Value: $\cos^2\theta_{\mathrm{W}} = 0.776368$. Status: DERIVED.
- $p_{\alpha,\mathrm{G}}$, $p_{\alpha,2}$, $p_{\alpha,1}$: The order-α (interaction) part of the pressure of the gauge plasma of the group G; $p_{\alpha,2}$ of SU(2), $p_{\alpha,1}$ of $\mathrm{U}(1)_Y$. Unit: GeV^4. Status: STANDARD.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- g_1: The four-dimensional hypercharge coupling ($g_1^2 = 4\pi\alpha_1$, Standard-Model normalisation without the GUT factor). Unit: dimensionless. Value: $g_1^2 = 0.1380$ at $2M_{\mathrm{DM}}$. Status: DERIVED.

dirac16complex acting on the Standard Model:

$$\frac{1}{\hat{g}^2_{3,\mathrm{eff}}} = \frac{1}{\hat{g}_3^2} + \frac{4c_{\mathrm{G}}\bar{\Psi}\Psi}{\Lambda_\star^3}, \qquad \frac{1}{\hat{g}^2_{2,\mathrm{eff}}} = \frac{1}{\hat{g}_2^2} + \frac{4c_{\mathrm{W}}\bar{\Psi}\Psi}{\Lambda_\star^3}, \qquad \frac{1}{\hat{g}^2_{1,\mathrm{eff}}} = \frac{1}{\hat{g}_1^2} + \frac{4c_{\mathrm{B}}\bar{\Psi}\Psi}{\Lambda_\star^3} \qquad \text{(FE-21)}$$

Symbols in (FE-21).

- $\hat{g}_{3,\mathrm{eff}}$, $\hat{g}_{2,\mathrm{eff}}$, $\hat{g}_{1,\mathrm{eff}}$: The field-dependent eight-dimensional couplings produced by I12: $1/\hat{g}^2_{i,\mathrm{eff}} = 1/\hat{g}_i^2 + 4c\,\bar{\Psi}\Psi/\Lambda_\star^3$ with $c = c_{\mathrm{G}}, c_{\mathrm{W}}, c_{\mathrm{B}}$. Unit: GeV^{-2}. Status: DERIVED.
- $\hat{g}_3$: The eight-dimensional SU(3) coupling (it multiplies the gluon kinetic term as $1/\hat{g}_3^2$); $g_3^2 = \hat{g}_3^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- c_{G}: The coefficient of the gluon term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{G}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- Ψ, $\bar{\Psi}$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar{\Psi} = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- $\hat{g}_2$: The eight-dimensional SU(2) coupling; $g_2^2 = \hat{g}_2^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- c_{W}: The coefficient of the SU(2) term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{W}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- $\hat{g}_1$: The eight-dimensional $\mathrm{U}(1)_Y$ coupling; $g_1^2 = \hat{g}_1^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- c_{B}: The coefficient of the hypercharge term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{B}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.

$$\frac{\Delta\alpha_3}{\alpha_3} = -4G_{\rm g}\, n_{\rm DM,tot}, \qquad \frac{\Delta\alpha_2}{\alpha_2} = -4G_{\rm W}\, n_{\rm DM,tot}, \qquad \frac{\Delta\alpha_1}{\alpha_1} = -4G_{\rm B}\, n_{\rm DM,tot},$$
$$-\frac{1}{\Upsilon\, {\rm e}^{3HL\xi}}\frac{\rm d}{{\rm d}\xi}\Big(\Upsilon\, {\rm e}^{3HL\xi}\frac{{\rm d}F}{{\rm d}\xi}\Big) = \tilde{m}^2 F, \qquad \Upsilon(\xi) = 1 + A_{\rm mf}\, n^{\rm sc}_{\rm DM}, \qquad n^{\rm sc}_{\rm DM} = 2a_{\rm DM}b_{\rm DM}, \tag{FE-79}$$
$$\frac{{\rm d}F}{{\rm d}\xi}(0) = 0, \qquad \Upsilon(-1)\frac{{\rm d}F}{{\rm d}\xi}(-1) = \beta_{\rm W}F(-1)$$

Symbols in (FE-79).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m^2_{21} = m^2_2 - m^2_1$ a difference of squared masses, $\Delta N_{\rm eff}$ an addition to $N_{\rm eff}$, $\Delta S_{\rm m}$ the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- α_3: The strong fine-structure constant, $\alpha_3 = g_3^2/(4\pi)$, at the scale given in brackets. Unit: dimensionless. Value: $\alpha_3(m_{\rm Z}) = 0.1180$; $\alpha_3(m_{\rm KK}) = 0.0606$. Status: STANDARD.
- $G_{\rm g}$: The coupling of a dark-matter pair to canonically normalised gluon pairs through I12. Unit: ${\rm GeV}^{-3}$. Value: 6.09×10^{-15} ${\rm GeV}^{-3}$ (benchmark). Status: DERIVED.
- $n_{\rm DM,tot}$: The total number density of dark-matter quanta: both species (the lowest odd levels of Ψ_A and Ψ_B), particles and antiparticles, $n_{\rm DM,tot} = 2(n_{\rm DM} + \bar{n}_{\rm DM})$. Unit: ${\rm GeV}^3$. Status: STANDARD.
- α_2: The SU(2) fine-structure constant, $\alpha_2 = g_2^2/(4\pi)$. Unit: dimensionless. Status: STANDARD.
- $G_{\rm W}$: The coupling of a dark-matter pair to canonically normalised SU(2) gauge pairs through I12. Unit: ${\rm GeV}^{-3}$. Value: 2.96×10^{-15} ${\rm GeV}^{-3}$. Status: DERIVED.
- α_1: The hypercharge fine-structure constant, $\alpha_1 = g_1^2/(4\pi)$. Unit: dimensionless. Status: STANDARD.
- $G_{\rm B}$: The coupling of a dark-matter pair to canonically normalised hypercharge pairs through I12. Unit: ${\rm GeV}^{-3}$. Value: 1.08×10^{-15} ${\rm GeV}^{-3}$. Status: DERIVED.
- $\Upsilon(\xi)$: The factor by which the scalar density of the dark matter changes the gauge kinetic term along x_0 (I12), $\Upsilon = 1 + A_{\rm mf} n^{\rm sc}_{\rm DM}$. Unit: dimensionless. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; ${\rm e}^X$ is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of $m_{\rm W}$ and $\rho_{\rm EW,0}$). Status: AUTHOR.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926\times10^{-21}$ m at the stabilised block). Status: ASSUMED.
- ξ: The scaled hidden coordinate, $\xi = y/L$, running from -1 (tip) to 0 (fold) on each half of a period. Unit: dimensionless. Status: DERIVED.
- d: The differential: ${\rm d}X$ is an infinitesimal change of X, ${\rm d}X/{\rm d}Y$ the derivative of X with respect to Y, ${\rm d}^nX$ the volume element of n variables. Status: STANDARD.
- $F(\xi)$, F_n: The x_0 profile of a boson mode on one half of the period (a function of y or of $\xi = y/L$): of a gauge field (gauge $A_y = 0$) or of the transverse-traceless graviton; F_n is the profile of the n-th Kaluza-Klein level of the graviton tower ($F_0 = 1$ the flat zero mode). Unit: dimensionless. Status: DERIVED.
- $\tilde{m}$, $\tilde{m}_n$: The dimensionless x_0 eigenvalue of a boson mode, $\tilde{m} = m\,fL$; $\tilde{m}_n$ the n-th even Kaluza-Klein level of the tower with $\beta = 0$ ($\tilde{m}_1$ the first). Unit: dimensionless. Value: $\tilde{m}_1^2 = 30.1196$, $\tilde{m}_2^2 = 59.7284$, $\tilde{m}_3^2 = 109.0764$ ($HL = 3$), i.e. $\tilde{m}_1 = 5.48813$, $\tilde{m}_2 = 7.72842$, $\tilde{m}_3 = 10.44397$. Status: DERIVED.
- $A_{\rm mf}$: The amplitude of the I12 mean field in the x_0 problem of the gauge fields (a bookkeeping parameter: the result is linear in it). Unit: ${\rm GeV}^{-1}$ (with the orbital normalisation). Status: DERIVED.
- $n^{\rm sc}_{\rm DM}$: The scalar density of the dark-matter orbital, $n^{\rm sc}_{\rm DM} = 2a_{\rm DM}b_{\rm DM}$. Unit: GeV (normalised orbital). Status: DERIVED.
- $a_{\rm DM}$, $b_{\rm DM}$: The upper and lower components of the dark-matter orbital (block spinor $(a, {\rm i}b)$, normalised on one half). Unit: ${\rm GeV}^{1/2}$. Status: DERIVED.
- $\beta_{\rm W}$: The Robin parameter of W^+ and W^- (set by the tip mass term of the sigma field). Unit: dimensionless. Value: 0.011912 (fitted). Status: DERIVED.

Gravity closes the loop: the total source of gravity, with the energy-momentum tensors of the Standard-Model fields and of the portals, drives the adiabatic block response of step 1,

$$\mathcal{T}^{M}{}_{N} = \mathcal{T}_{\text{stab}}{}^{M}{}_{N} + \mathcal{T}_{\Psi}{}^{M}{}_{N} + \mathcal{T}_{\text{gauge}}{}^{M}{}_{N} + \mathcal{T}_{\text{f}}{}^{M}{}_{N} + \mathcal{T}_{\text{tip}}{}^{M}{}_{N} + \mathcal{T}_{\text{I11}}{}^{M}{}_{N} + \mathcal{T}_{\text{I12}}{}^{M}{}_{N} + \mathcal{T}_{\text{I13}}{}^{M}{}_{N} \tag{FE-3}$$

Symbols in (FE-3).

- $\mathcal{T}^{M}{}_{N}$: The energy-momentum tensor of all matter (every field except gravity, the stabiliser included), the source of gravity; defined by the change of the matter action under a change of the metric. Unit: GeV^8. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\mathcal{T}_{\text{stab}}$: The energy-momentum tensor of the stabiliser. Unit: GeV^8. Status: DERIVED.
- $\mathcal{T}_{\Psi}$: The energy-momentum tensor of the dirac16complex field. Unit: GeV^8. Status: DERIVED.
- $\mathcal{T}_{\text{gauge}}$: The energy-momentum tensor of the gauge fields. Unit: GeV^8. Status: STANDARD.
- $\mathcal{T}_{\text{f}}$: The energy-momentum tensor of the Standard-Model fermions. Unit: GeV^8. Status: DERIVED.
- $\mathcal{T}_{\text{tip}}$: The energy-momentum tensor of the tip terms (non-zero only for the tip directions). Unit: GeV^8. Status: DERIVED.
- $\mathcal{T}_{\text{I11}}$: The energy-momentum tensor of the portal I11. Unit: GeV^8. Status: DERIVED.
- $\mathcal{T}_{\text{I12}}$: The energy-momentum tensor of the portal I12. Unit: GeV^8. Status: DERIVED.
- $\mathcal{T}_{\text{I13}}$: The energy-momentum tensor of the portal I13. Unit: GeV^8. Status: DERIVED.

$$\Delta K = \frac{c_\rho\,\kappa\Delta\rho + c_3\,\kappa\Delta p_3 + c_p\,\kappa\Delta\bar{p}}{H^2}, \qquad \Delta\ln f = -\frac{\Delta K}{2K_*}, \qquad c_3 = -3c_\rho, \qquad \Delta\dot{\mathcal{H}} = -\frac{\kappa(\Delta\rho + \Delta p_3)}{2\mathcal{G}(K_*)} \tag{FE-47}$$

Symbols in (FE-47).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m^2_{21} = m_2^2 - m_1^2$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- c_ρ: The coefficient of $\kappa\Delta\rho$ in the adiabatic block response ΔK. Unit: dimensionless. Value: 0.0358521. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_{\text{N}}\mathcal{G}(K_*) = 2.328\times10^{-36}\ \text{GeV}^{-2}$. Status: AUTHOR.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- c_3: The coefficient of $\kappa\Delta p_3$ in the adiabatic block response, $c_3 = -3c_\rho$ (so radiation drops out). Unit: dimensionless. Value: -0.107556. Status: DERIVED.
- p_3: The pressure of ordinary 3-space, $p_3 = \mathcal{T}^{x_1}{}_{x_1} = \mathcal{T}^{x_2}{}_{x_2} = \mathcal{T}^{x_3}{}_{x_3}$ for an isotropic source. Unit: GeV^8. Status: STANDARD.
- c_p: The coefficient of $\kappa\Delta\bar{p}$ in the adiabatic block response. Unit: dimensionless. Value: -0.140028. Status: DERIVED.
- $\bar{p}$: The hidden pressure, the pressure averaged over the block, $\bar{p} = (p_0 + 3p_5)/4$; for a species at fixed temperature $\bar{p} = \frac{1}{4}\,\partial p/\partial\ln f$. Unit: GeV^8 (or GeV^4 for a four-dimensional density). Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\text{EW},0}$). Status: AUTHOR.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \text{e}^{-a_4}b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.

- K_*: The block variable of the stabilised block, $K_* = 12 + \frac{4}{3}\sqrt{51}$ (the upper root K_+). Unit: dimensionless. Value: 21.5219. Status: DERIVED.
- $\dot{\mathcal{H}}$: The time derivative $\mathrm{d}\mathcal{H}/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- $\mathcal{G}(K)$: The coefficient of the four-dimensional Einstein-Hilbert term on a static block, $\mathcal{G}(K) = 1 - 24w_{(2)}K + 72w_{(3)}K^2 = (K^2 - 12K + 16)/16$; four-dimensional gravity is attractive for $\mathcal{G} > 0$. Unit: dimensionless. Value: $\mathcal{G}(K_*) = 20/3 + \sqrt{51} = 13.8081$. Status: DERIVED.

$$(\mathcal{T}_\mathrm{f})_{MN} = \sum_{i=1}^{3}\sum_{\psi}\Big\{ -\frac{1}{4}\Big[\bar{\psi}_i\gamma_M D_N\psi_i + \bar{\psi}_i\gamma_N D_M\psi_i - (D_M\bar{\psi}_i)\gamma_N\psi_i - (D_N\bar{\psi}_i)\gamma_M\psi_i\Big] + g_{MN}\,\mathcal{L}_{\psi i}\Big\},$$
$$\mathcal{L}_{\psi i} = \tfrac{1}{2}\big(\bar{\psi}_i\gamma^P D_P\psi_i - (D_P\bar{\psi}_i)\gamma^P\psi_i\big) - \mu_{\psi i}\,\bar{\psi}_i\psi_i \qquad \text{(FE-6)}$$

Symbols in (FE-6).

- $\mathcal{T}_\mathrm{f}$: The energy-momentum tensor of the Standard-Model fermions. Unit: GeV^8. Status: DERIVED.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- ψ: The label of a Standard-Model multiplet: $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ (quark doublet, up and down quark singlets, lepton doublet, charged-lepton singlet, right-handed neutrino); a sum over ψ runs over the six. Status: STANDARD.
- ψ_i, $\bar{\psi}_i$: A Standard-Model fermion field of eight dimensions: ψ_i is the multiplet $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ of the generation i, ψ_f one fermion component f (in four-dimensional formulas its zero mode); $\bar{\psi} = \psi^\dagger C$ the Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$ (eight dimensions), $\mathrm{GeV}^{3/2}$ (four dimensions). Status: ASSUMED.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4, 4)$; $\gamma^M = \mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- D_M, D_μ: The covariant derivative: on a spinor $D_M = \partial_M + \Omega_M$ (spin connection), on a Standard-Model fermion with the gauge fields added; on the sigma field $D_\mu\Sigma$ with the W and B fields. Unit: 1/length. Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+, +, +, +, -, -, -, -$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- $\mathcal{L}_{\psi i}$: The Lagrangian density of the fermion multiplet ψ of the generation i (the bracket of S_f). Unit: GeV^8. Status: ASSUMED.
- P: A coordinate index of eight dimensions (also $P_1, P_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\mu_{\psi i}$: The bulk mass of a Standard-Model fermion: $\mu_{\psi i}$ of the multiplet ψ of the generation i; with the hidden direction it fixes the localisation $c = f\mu L$ of the zero mode. Unit: GeV. Value: $\mu_{\psi i} = c/(fL)$ with the localisation c of the zero mode (doublets: $\mu_Q = 5.2304\,H$, $\mu_\ell = 9.2055\,H$). Status: ASSUMED.

$$(\mathcal{T}_\mathrm{gauge})_{MN} = \frac{1}{\hat{g}_3^2}\Big(G^A_{MP}G^A{}_N{}^P - \frac{1}{4}g_{MN}G^A_{PQ}G^{APQ}\Big) + \frac{1}{\hat{g}_2^2}\Big(W^I_{MP}W^I{}_N{}^P - \frac{1}{4}g_{MN}W^I_{PQ}W^{IPQ}\Big)$$
$$+ \frac{1}{\hat{g}_1^2}\Big(B_{MP}B_N{}^P - \frac{1}{4}g_{MN}B_{PQ}B^{PQ}\Big) \qquad \text{(FE-7)}$$

Symbols in (FE-7).

- $\mathcal{T}_\mathrm{gauge}$: The energy-momentum tensor of the gauge fields. Unit: GeV^8. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.

- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\hat{g}_3$: The eight-dimensional SU(3) coupling (it multiplies the gluon kinetic term as $1/\hat{g}_3^2$); $g_3^2 = \hat{g}_3^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- G^A_M, G_M, G_{MN}, G^A_{MN}: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G^A_M ($A = 1, \ldots, 8$), the matrix $G_M = G^A_M t^A$, the field strength G_{MN} (matrix) and its components G^A_{MN}; $G^A_{\mu\nu}$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- P: A coordinate index of eight dimensions (also $P_1, P_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M \mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- Q: A coordinate index of eight dimensions (also $Q_1, Q_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\hat{g}_2$: The eight-dimensional SU(2) coupling; $g_2^2 = \hat{g}_2^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- W^I_M, W_M, W_{MN}, W^I_{MN}: The SU(2) gauge field (geometric normalisation): the components W^I_M ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W^I_M \tau^I/2$, the field strength W_{MN} and its components W^I_{MN}; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- $\hat{g}_1$: The eight-dimensional $\mathrm{U}(1)_Y$ coupling; $g_1^2 = \hat{g}_1^2/V_\perp$. Unit: GeV^{-2}. Status: ASSUMED.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.

$$(\mathcal{T}_{\mathrm{I11}})_{MN} = g_{MN}\,\frac{c_{\mathrm{V}}}{\Lambda_\star^6}\big(\mathrm{i}\bar{\Psi}\gamma^P\Psi\big)J^{B-L}_P, \qquad (\mathcal{T}_{\mathrm{I13}})_{MN} = g_{MN}\,\frac{c_{\mathrm{A}}}{\Lambda_\star^6}\sum_{P\neq x_0}\big(\mathrm{i}\bar{\Psi}\gamma^P P_b\Psi\big)J^B_P \tag{FE-9}$$

Symbols in (FE-9).

- $\mathcal{T}_{\mathrm{I11}}$: The energy-momentum tensor of the portal I11. Unit: GeV^8. Status: DERIVED.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M \mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- c_{V}: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_{\mathrm{V}} = 4.38\times10^{-17}$ GeV^{-2}. Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- Ψ, $\bar{\Psi}$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar{\Psi} = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- P: A coordinate index of eight dimensions (also $P_1, P_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- J^{B-L}_M: The $B-L$ current of the Standard-Model fermions (baryon minus lepton number), summed over the generations. Unit: GeV^7. Status: STANDARD.
- $\mathcal{T}_{\mathrm{I13}}$: The energy-momentum tensor of the portal I13. Unit: GeV^8. Status: DERIVED.

- c_{A}: The coefficient of the x_0-chiral vector portal I13. Unit: dimensionless. Value: fixed through $G_{\mathrm{A}} = 1.278 \times 10^{-9}\ \mathrm{GeV}^{-2}$ (dark-matter relic). Status: INVENTED.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- $P_a,\ P_b$: The projectors on the two x_0-chiralities, $P_a = (1+\gamma^{x_0})/2$ and $P_b = (1-\gamma^{x_0})/2$ (in a block: the upper and the lower component). Unit: dimensionless. Status: INVENTED.
- J^B_M: The baryon-number current of the quarks (one third per quark), summed over the generations. Unit: GeV^7. Status: STANDARD.

$$(\mathcal{T}_{\mathrm{I12}})_{MN} = \frac{4\bar{\Psi}\Psi}{\Lambda_\star^3}\Big[c_{\mathrm{G}}\Big(G^A_{MP}G^A{}_N{}^P - \frac{1}{4}g_{MN}G^A_{PQ}G^{APQ}\Big) + c_{\mathrm{W}}\Big(W^I_{MP}W^I{}_N{}^P - \frac{1}{4}g_{MN}W^I_{PQ}W^{IPQ}\Big) + c_{\mathrm{B}}\Big(B_{MP}B_N{}^P - \frac{1}{4}g_{MN}B_{PQ}B^{PQ}\Big)\Big] \tag{FE-10}$$

Symbols in (FE-10).

- $\mathcal{T}_{\mathrm{I12}}$: The energy-momentum tensor of the portal I12. Unit: GeV^8. Status: DERIVED.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\Psi,\ \bar{\Psi}$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar{\Psi} = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- c_{G}: The coefficient of the gluon term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{G}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- $G^A_M,\ G_M,\ G_{MN},\ G^A_{MN}$: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G^A_M ($A = 1, \ldots, 8$), the matrix $G_M = G^A_M t^A$, the field strength G_{MN} (matrix) and its components G^A_{MN}; $G^A_{\mu\nu}$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- P: A coordinate index of eight dimensions (also $P_1, P_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, $g^M{}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- Q: A coordinate index of eight dimensions (also $Q_1, Q_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- c_{W}: The coefficient of the SU(2) term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{W}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- $W^I_M,\ W_M,\ W_{MN},\ W^I_{MN}$: The SU(2) gauge field (geometric normalisation): the components W^I_M ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W^I_M \tau^I/2$, the field strength W_{MN} and its components W^I_{MN}; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- c_{B}: The coefficient of the hypercharge term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{B}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- $B_M,\ B_{MN}$: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.

$$\Delta K = \frac{c_\rho\,\kappa\Delta\rho + c_3\,\kappa\Delta p_3 + c_p\,\kappa\Delta\bar{p}}{H^2}, \qquad \Delta\ln f = -\frac{\Delta K}{2K_*},$$
$$\Delta\ln m_s = \frac{\mathrm{d}\ln m_s}{\mathrm{d}\ln f}\,\Delta\ln f, \qquad \kappa\Delta\rho = 8\pi G_\mathrm{N}\,\mathcal{G}(K_*)\,\rho_4 \tag{FE-80}$$

Symbols in (FE-80).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_eff an addition to N_eff, ΔS_m the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- c_ρ: The coefficient of $\kappa\Delta\rho$ in the adiabatic block response ΔK. Unit: dimensionless. Value: 0.0358521. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_\mathrm{N}\mathcal{G}(K_*) = 2.328\times10^{-36}\ \mathrm{GeV}^{-2}$. Status: AUTHOR.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- c_3: The coefficient of $\kappa\Delta p_3$ in the adiabatic block response, $c_3 = -3c_\rho$ (so radiation drops out). Unit: dimensionless. Value: -0.107556. Status: DERIVED.
- p_3: The pressure of ordinary 3-space, $p_3 = \mathcal{T}^{x_1}{}_{x_1} = \mathcal{T}^{x_2}{}_{x_2} = \mathcal{T}^{x_3}{}_{x_3}$ for an isotropic source. Unit: GeV^8. Status: STANDARD.
- c_p: The coefficient of $\kappa\Delta\bar{p}$ in the adiabatic block response. Unit: dimensionless. Value: -0.140028. Status: DERIVED.
- $\bar{p}$: The hidden pressure, the pressure averaged over the block, $\bar{p} = (p_0 + 3p_5)/4$; for a species at fixed temperature $\bar{p} = \frac{1}{4}\,\partial p/\partial\ln f$. Unit: GeV^8 (or GeV^4 for a four-dimensional density). Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4}b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- K_*: The block variable of the stabilised block, $K_* = 12 + \frac{4}{3}\sqrt{51}$ (the upper root K_+). Unit: dimensionless. Value: 21.5219. Status: DERIVED.
- m, m_s, m_i: A four-dimensional mass: m the mass of a mode (the eigenvalue of an x_0 problem) or of a fermion; m_s the mass of the species s; m_i the mass of the neutrino mass state i. Unit: GeV. Status: DERIVED.
- s: The label of a particle species of the plasma or of matter (photon, gluons, $W^\pm$, Z, quarks, leptons, nucleons, dark matter); a sum over s runs over the species present. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_N: Newton's constant of four-dimensional gravity, $8\pi G_\mathrm{N} = \kappa/(\mathcal{G}(K_*)V_\perp)$. Unit: GeV^{-2}. Value: $6.709\times10^{-39}\ \mathrm{GeV}^{-2}$ ($M_\mathrm{Pl} = G_\mathrm{N}^{-1/2} = 1.22091\times10^{19}$ GeV). Status: STANDARD.
- $\mathcal{G}(K)$: The coefficient of the four-dimensional Einstein-Hilbert term on a static block, $\mathcal{G}(K) = 1 - 24w_{(2)}K + 72w_{(3)}K^2 = (K^2 - 12K + 16)/16$; four-dimensional gravity is attractive for $\mathcal{G} > 0$. Unit: dimensionless. Value: $\mathcal{G}(K_*) = 20/3 + \sqrt{51} = 13.8081$. Status: DERIVED.
- ρ_4: A four-dimensional energy density (per 3-volume); its block average is $\kappa\Delta\rho = 8\pi G_\mathrm{N}\mathcal{G}(K_*)\rho_4$. Unit: GeV^4. Status: DERIVED.

$$\bar{p} = -\frac{1}{4}\sum_s \left(\rho_s - 3p_s\right)\frac{\mathrm{d}\ln m_s}{\mathrm{d}\ln f} - \frac{1}{4}\,\Theta_\mathrm{QCD}\,\frac{\mathrm{d}\ln\Lambda_\mathrm{QCD}^{(N_\mathrm{f})}}{\mathrm{d}\ln f} + \frac{1}{4}\,p_{\alpha,2}\,\frac{\mathrm{d}\ln\alpha_2}{\mathrm{d}\ln f} + \frac{1}{4}\,p_{\alpha,1}\,\frac{\mathrm{d}\ln\alpha_1}{\mathrm{d}\ln f} \tag{FE-81}$$

Symbols in (FE-81).

- $\bar{p}$: The hidden pressure, the pressure averaged over the block, $\bar{p} = (p_0 + 3p_5)/4$; for a species at fixed temperature $\bar{p} = \frac{1}{4}\,\partial p/\partial \ln f$. Unit: GeV^8 (or GeV^4 for a four-dimensional density). Status: DERIVED.
- s: The label of a particle species of the plasma or of matter (photon, gluons, $W^\pm$, Z, quarks, leptons, nucleons, dark matter); a sum over s runs over the species present. Status: STANDARD.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- p: A pressure: $p_M = \mathcal{T}^M{}_M$ (no sum) for the direction $M \neq x_4$, p_s the pressure of the species s; Δp the pressure perturbation in the cosmological perturbation equations; in the Feynman-Hellmann relation the pressure of a gauge plasma. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- m, m_s, m_i: A four-dimensional mass: m the mass of a mode (the eigenvalue of an x_0 problem) or of a fermion; m_s the mass of the species s; m_i the mass of the neutrino mass state i. Unit: GeV. Status: DERIVED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-\alpha_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- Θ_{QCD}, Θ_{G}, Θ_V: The trace $\rho - 3p$ of the energy-momentum tensor (interaction measure) of a plasma component: Θ_{QCD} of the quark-gluon plasma or hadron gas, Θ_{G} of the gauge plasma of the group G, Θ_V of the thermal gas of the massive vector V (Θ_{W} of $W^\pm$, Θ_{Z} of Z). Unit: GeV^4. Status: STANDARD.
- $\Lambda_{\mathrm{QCD}}^{(N_\mathrm{f})}$: The QCD scale with N_f active flavours ($\Lambda_{\mathrm{QCD}}^{(6)}$, $^{(5)}$, $^{(4)}$, $^{(3)}$), defined by the one-loop running. Unit: GeV. Status: STANDARD.
- $p_{\alpha,\mathrm{G}}$, $p_{\alpha,2}$, $p_{\alpha,1}$: The order-α (interaction) part of the pressure of the gauge plasma of the group G; $p_{\alpha,2}$ of SU(2), $p_{\alpha,1}$ of $\mathrm{U}(1)_Y$. Unit: GeV^4. Status: STANDARD.
- α_2: The SU(2) fine-structure constant, $\alpha_2 = g_2^2/(4\pi)$. Unit: dimensionless. Status: STANDARD.
- α_1: The hypercharge fine-structure constant, $\alpha_1 = g_1^2/(4\pi)$. Unit: dimensionless. Status: STANDARD.

$$\frac{\mathrm{d}\ln\alpha_{\mathrm{G}}(\mu_{\mathrm{ren}})}{\mathrm{d}\ln f} = -\alpha_{\mathrm{G}}(\mu_{\mathrm{ren}})\Big[\frac{4}{\alpha_{\mathrm{G}}(m_{\mathrm{KK}})} + \frac{b_{\mathrm{G}}}{2\pi}\Big], \qquad \Theta_{\mathrm{G}} = -\frac{b_{\mathrm{G}}\,\alpha_{\mathrm{G}}}{2\pi}\,p_{\alpha,\mathrm{G}} \tag{FE-82}$$

Symbols in (FE-82).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- α_{G}: The fine-structure constant of the gauge group G (one of α_3, α_2, α_1). Unit: dimensionless. Status: STANDARD.
- μ_{ren}: The renormalisation scale of a running coupling (taken as $2\pi T$ for the condensates of a plasma). Unit: GeV. Status: STANDARD.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-\alpha_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- m_n^{KK}, m_{KK}: The mass of the n-th Kaluza-Klein graviton (and gauge) level, $m_n^{\mathrm{KK}} = \tilde{m}_n/(fL)$; $m_{\mathrm{KK}} = m_1^{\mathrm{KK}}$ is the Kaluza-Klein scale. Unit: GeV. Value: $m_{\mathrm{KK}} = 121.32$ TeV. Status: DERIVED.
- b_3, b_2, b_1, b_{G}: The one-loop coefficients of the running couplings, $\mathrm{d}\alpha/\mathrm{d}\ln\mu_{\mathrm{ren}} = -b\,\alpha^2/(2\pi)$: $b_3 = 11 - \frac{2}{3}N_\mathrm{f}$ (SU(3)), $b_2 = \frac{10}{3}$ (SU(2) without a Higgs particle), $b_1 = -\frac{20}{3}$ ($\mathrm{U}(1)_Y$), b_{G} that of the group G (not related to the metric function b_4). Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- Θ_{QCD}, Θ_{G}, Θ_V: The trace $\rho - 3p$ of the energy-momentum tensor (interaction measure) of a plasma component: Θ_{QCD} of the quark-gluon plasma or hadron gas, Θ_{G} of the gauge plasma of the group G, Θ_V of the thermal gas of the massive vector V (Θ_{W} of $W^\pm$, Θ_{Z} of Z). Unit: GeV^4. Status: STANDARD.
- $p_{\alpha,\mathrm{G}}$, $p_{\alpha,2}$, $p_{\alpha,1}$: The order-α (interaction) part of the pressure of the gauge plasma of the group G; $p_{\alpha,2}$ of SU(2), $p_{\alpha,1}$ of $\mathrm{U}(1)_Y$. Unit: GeV^4. Status: STANDARD.

$$\frac{1}{\alpha_3(m_{\mathrm{KK}})}=\frac{1}{\alpha_3(m_{\mathrm{Z}})}+\frac{7}{2\pi}\ln\frac{m_{\mathrm{KK}}}{m_{\mathrm{Z}}},\qquad \Lambda^{(6)}_{\mathrm{QCD}}=m_{\mathrm{KK}}\,\mathrm{e}^{-2\pi/(7\alpha_3(m_{\mathrm{KK}}))},\qquad \Lambda^{(3)}_{\mathrm{QCD}}=\big(\Lambda^{(6)}_{\mathrm{QCD}}\big)^{21/27}\big(m_{\mathrm{c}}m_{\mathrm{b}}m_{\mathrm{t}}\big)^{2/27},$$
$$\frac{\mathrm{d}\ln m_{\mathrm{p}}}{\mathrm{d}\ln f}=\Big(1-\frac{\sigma_{\pi N}+\sigma_{\mathrm{s}}}{m_{\mathrm{p}}}\Big)\frac{\mathrm{d}\ln\Lambda^{(3)}_{\mathrm{QCD}}}{\mathrm{d}\ln f}+\frac{\sigma_{\pi N}}{2m_{\mathrm{p}}}\Big(\frac{\mathrm{d}\ln m_{\mathrm{u}}}{\mathrm{d}\ln f}+\frac{\mathrm{d}\ln m_{\mathrm{d}}}{\mathrm{d}\ln f}\Big)+\frac{\sigma_{\mathrm{s}}}{m_{\mathrm{p}}}\,\frac{\mathrm{d}\ln m_{\mathrm{s}}}{\mathrm{d}\ln f}\tag{FE-83}$$

Symbols in (FE-83).

- α_3: The strong fine-structure constant, $\alpha_3=g_3^2/(4\pi)$, at the scale given in brackets. Unit: dimensionless. Value: $\alpha_3(m_{\mathrm{Z}})=0.1180$; $\alpha_3(m_{\mathrm{KK}})=0.0606$. Status: STANDARD.
- m_n^{KK}, m_{KK}: The mass of the n-th Kaluza-Klein graviton (and gauge) level, $m_n^{\mathrm{KK}}=\tilde{m}_n/(fL)$; $m_{\mathrm{KK}}=m_1^{\mathrm{KK}}$ is the Kaluza-Klein scale. Unit: GeV. Value: $m_{\mathrm{KK}}=121.32$ TeV. Status: DERIVED.
- m_{Z}: The mass of the Z. Unit: GeV. Value: 91.1879 GeV (reproduced). Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $\Lambda^{(N_{\mathrm{f}})}_{\mathrm{QCD}}$: The QCD scale with N_{f} active flavours ($\Lambda^{(6)}_{\mathrm{QCD}}$, $^{(5)}$, $^{(4)}$, $^{(3)}$), defined by the one-loop running. Unit: GeV. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- m_{u}, m_{d}, m_{s}, m_{c}, m_{b}, m_{t}: The masses of the up, down, strange, charm, bottom and top quarks (the upright subscript names the quark). Unit: GeV. Value: 2.16 MeV, 4.70 MeV, 93.5 MeV, 1.273 GeV, 4.183 GeV, 172.56 GeV (the x_0 masses, fitted to the PDG values). Status: DERIVED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- m_{p}, m_{n}: The masses of the proton and the neutron. Unit: GeV. Value: 0.938272 GeV, 0.939565 GeV. Status: STANDARD.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f=\mathrm{e}^{-a_4}b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- $\sigma_{\pi N}$: The pion-nucleon sigma term (the contribution of the up and down quark masses to the nucleon mass). Unit: GeV. Value: 45 MeV. Status: STANDARD.
- σ_{s}: The strange sigma term of the nucleon. Unit: GeV. Value: 40 MeV. Status: STANDARD.

$$\frac{\mathrm{d}\ln\Lambda^{(6)}_{\mathrm{QCD}}}{\mathrm{d}\ln f}=-1-\frac{8\pi}{7\alpha_3(m_{\mathrm{KK}})},\qquad \frac{\mathrm{d}\ln\Lambda^{(5)}_{\mathrm{QCD}}}{\mathrm{d}\ln f}=\frac{21}{23}\frac{\mathrm{d}\ln\Lambda^{(6)}_{\mathrm{QCD}}}{\mathrm{d}\ln f}+\frac{2}{23}\frac{\mathrm{d}\ln m_{\mathrm{t}}}{\mathrm{d}\ln f},$$
$$\frac{\mathrm{d}\ln\Lambda^{(4)}_{\mathrm{QCD}}}{\mathrm{d}\ln f}=\frac{21}{25}\frac{\mathrm{d}\ln\Lambda^{(6)}_{\mathrm{QCD}}}{\mathrm{d}\ln f}+\frac{2}{25}\Big(\frac{\mathrm{d}\ln m_{\mathrm{t}}}{\mathrm{d}\ln f}+\frac{\mathrm{d}\ln m_{\mathrm{b}}}{\mathrm{d}\ln f}\Big),\tag{FE-84}$$
$$\frac{\mathrm{d}\ln\Lambda^{(3)}_{\mathrm{QCD}}}{\mathrm{d}\ln f}=\frac{21}{27}\frac{\mathrm{d}\ln\Lambda^{(6)}_{\mathrm{QCD}}}{\mathrm{d}\ln f}+\frac{2}{27}\Big(\frac{\mathrm{d}\ln m_{\mathrm{c}}}{\mathrm{d}\ln f}+\frac{\mathrm{d}\ln m_{\mathrm{b}}}{\mathrm{d}\ln f}+\frac{\mathrm{d}\ln m_{\mathrm{t}}}{\mathrm{d}\ln f}\Big)$$

Symbols in (FE-84).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $\Lambda^{(N_{\mathrm{f}})}_{\mathrm{QCD}}$: The QCD scale with N_{f} active flavours ($\Lambda^{(6)}_{\mathrm{QCD}}$, $^{(5)}$, $^{(4)}$, $^{(3)}$), defined by the one-loop running. Unit: GeV. Status: STANDARD.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f=\mathrm{e}^{-a_4}b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- α_3: The strong fine-structure constant, $\alpha_3=g_3^2/(4\pi)$, at the scale given in brackets. Unit: dimensionless. Value: $\alpha_3(m_{\mathrm{Z}})=0.1180$; $\alpha_3(m_{\mathrm{KK}})=0.0606$. Status: STANDARD.

- m_n^{KK}, m_{KK}: The mass of the n-th Kaluza-Klein graviton (and gauge) level, $m_n^{\mathrm{KK}} = \tilde{m}_n/(fL)$; $m_{\mathrm{KK}} = m_1^{\mathrm{KK}}$ is the Kaluza-Klein scale. Unit: GeV. Value: $m_{\mathrm{KK}} = 121.32$ TeV. Status: DERIVED.
- m_{u}, m_{d}, m_{s}, m_{c}, m_{b}, m_{t}: The masses of the up, down, strange, charm, bottom and top quarks (the upright subscript names the quark). Unit: GeV. Value: 2.16 MeV, 4.70 MeV, 93.5 MeV, 1.273 GeV, 4.183 GeV, 172.56 GeV (the x_0 masses, fitted to the PDG values). Status: DERIVED.

$$\frac{\mathrm{d}\ln m}{\mathrm{d}\ln f} = -1 + \frac{1}{2}\Big[\frac{\mathrm{d}\ln \mathcal{Z}(c_L)^2}{\mathrm{d}\ln c_L} + \frac{\mathrm{d}\ln \mathcal{Z}(-c_R)^2}{\mathrm{d}\ln(-c_R)}\Big] \tag{FE-85}$$

Symbols in (FE-85).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- m, m_s, m_i: A four-dimensional mass: m the mass of a mode (the eigenvalue of an x_0 problem) or of a fermion; m_s the mass of the species s; m_i the mass of the neutrino mass state i. Unit: GeV. Status: DERIVED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- $\mathcal{Z}(c)$: The tip amplitude of a normalised zero mode with the localisation c, $\mathcal{Z}(c)^2 = c\,\mathrm{e}^{-c}/\sinh c$. Unit: dimensionless. Status: DERIVED.
- c, c_L, c_R: The localisation of a zero mode, $c = f\mu L$ (bulk mass times the proper depth); c_L of a doublet (localised at the fold for $c_L > 0$), c_R of a singlet (localised at the fold for $c_R < 0$). Unit: dimensionless. Value: doublets $c_L = 3.38230$ (quarks) and 5.95288 (leptons); singlets c_R: up -15.4300, down -14.6258, strange -11.5158, charm -8.7683, bottom -7.5005, top -3.3823, electron -14.5539, muon -8.9809, tau -5.9529, neutrinos -35.0452, -32.8481, -31.0702 (fixed by the measured masses). Status: DERIVED.
- L: The chirality label L: the left-handed (doublet) zero mode, or the left-handed projection $P_L = (1 + \Gamma_4)/2$. Status: STANDARD.
- R: The chirality label R: the right-handed (singlet) zero mode or field, or the projection $P_R = (1 - \Gamma_4)/2$. Status: STANDARD.

11.2 The approximation of step 8

- *First order in the mean fields.* The level shifts are linear responses (Hellmann-Feynman); the second order is $(\Delta\varepsilon_{\mathrm{DM}}/M_{\mathrm{DM}})^2 = 2.3 \times 10^{-12}$ at most. The screening of the response by the interacting dirac16complex gas is 3.5×10^{-4} at the test coupling λ_{test} and 7.2×10^{-10} at the bound on λ of step 9.
- *The thermodynamics of the environments.* The plasma species as ideal gases with their x_0 masses, the net nucleons and electrons as degenerate Fermi gases in the cold environments, the QCD interaction measure from a lattice-like shape below 1 GeV (ESTIMATE) and leading-order perturbation theory above, the order-α pressure of the electroweak plasma without a scalar term and with the thermal suppression of a massive vector below m_{W} (ESTIMATE), one-loop running for the couplings, the QCD scale and the nucleon (ESTIMATE).

The block response at first order and adiabatic, with the gain of the loop mass → source → block → mass

$$\frac{|\Delta \ln f|_{\text{second iteration}}}{|\Delta \ln f|_{\text{first iteration}}} \leq 1.1 \times 10^{-33}$$

Symbols in this equation.

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.

and the neglected inertia of the block $(\mathcal{H}/\omega_{\mathrm{r}})^2 = 3.4 \times 10^{-33}$ at the freeze-out (step 1).

11.3 The solution of step 8

11.3.1 The portals on the zero modes

A four-dimensional particle of the density n_f (per 3-volume) in a zero mode with the normalised profile $\varrho_f(y)$ has the proper eight-dimensional density $n_f \varrho_f \mathrm{e}^{-3Hy}/(2f^4 v_\mathrm{t})$ on each half of the period, so the portals I11 and I13 couple a state o of Ψ to a chirality component f of a Standard-Model fermion through the overlaps of the contact interaction written above, with $G_\mathrm{V} = c_\mathrm{V} H/(2f^4 v_\mathrm{t} \Lambda_\star^6)$, $G_\mathrm{A} = c_\mathrm{A} H/(2f^4 v_\mathrm{t} \Lambda_\star^6)$, q_f the $B-L$ charge and $q_f^B = \frac{1}{3}$ for quarks and 0 for leptons; I13 acts on the lower component only, because $\gamma^{x_0} = \sigma_3$ in the block form. The overlaps (DERIVED; $H = 1$, 80 basis functions per sector):

zero mode f	Q	ℓ	u_R	c_R	t_R	d_R	s_R	b_R	e_R	μ_R	τ_R	ν_{Ri}
$\tilde{I}$, dark matter	8.2722	1.2524	0.3854	0.6056	8.2722	0.3961	0.4620	0.7622	0.3972	0.5879	1.2524	0.302-0.308
$\tilde{I}^{(b)}$, dark matter	2.6135	0.9134	0.3795	0.5569	2.6135	0.3891	0.4459	0.6607	0.3900	0.5442	0.9134	0.301-0.308
$\tilde{I}$, fold band	6.685	1.659	0.792	1.060	6.685	0.807	0.893	1.221	0.808	1.041	1.659	0.668-0.678
$\tilde{I}$, first even bulk level	8.92	1.185	0.428	0.602	8.92	0.437	0.488	0.733	0.438	0.588	1.185	0.359-0.365

($\tilde{I}^{(b)} = 0$ for the fold band; ν_{Ri} the three right-handed neutrinos; the right-handed top and tau share the localisation of their doublets, $c_R = -c_L$, hence the equal overlaps.) Self-consistent Kohn-Sham states (eight fold zero modes and eight dark-matter quanta, $\lambda = 0$) with the mean field $A_\mathrm{mf} \varrho_f \mathrm{e}^{-3Hy}$ on both components (I11) or on the lower one (I13), at $A_\mathrm{mf} = \pm 10^{-4}$, $\pm 2 \times 10^{-4}$ with Richardson extrapolation, reproduce $\tilde{I}$ for Q, e_R, ℓ to 3.0×10^{-10} and $\tilde{I}^{(b)}$ for Q, ℓ, u_R to 2.3×10^{-10}; with the benchmark self-coupling λ_test the response to the quark-doublet field is 8.2693 against 8.2722 (screening -3.5×10^{-4}); under the lepton-doublet I11 field the gap between the $k = 0$ fold zero mode and the first shell changes by -9.3×10^{-11} per unit amplitude (the rounding level; zero in exact arithmetic) (the second order, $-0.041 A_\mathrm{mf}^2$, is a change of the group velocity: the ratio of the shell energies at $|\vec{k}| = 2k_\mathrm{lat}$ and k_lat stays 2.000000000), and under I13 the fold levels move by at most 1.2×10^{-13} (the rounding level): the fold band stays massless and untouched by I13 (DERIVED).

11.3.2 The mean fields of the Standard-Model fields

In an environment with the net densities n_p, n_n, n_e (unpolarised fermions, each chirality half) and the temperature T, the dark-matter level moves by the shifts written above, with the benchmark strengths $G_\mathrm{V} = 4.38 \times 10^{-17}$ GeV^{-2}, $G_\mathrm{A} = 1.278 \times 10^{-9}$ GeV^{-2}, $c_\mathrm{I12}/\Lambda_\star^3 = 1.106 \times 10^{-14}$ GeV^{-3} (step 9) and $\mathcal{O}_\mathrm{DM} = 0.7072$; the vacuum gluon condensate of the gluon field in its geometric normalisation is $\langle G^A_{\mu\nu} G^{A\mu\nu} \rangle(0) = 4\pi^2 \times 0.012$ GeV^4 (STANDARD: 0.012 GeV^4 is the condensate of the canonically normalised field strength weighted with α_3/π), and the couplings of the condensates are taken at $\mu_\mathrm{ren} = 2\pi T$ from the $\overline{\mathrm{MS}}$ values at m_Z ($\sin^2 \theta_{\overline{\mathrm{MS}}} = 0.23122$, $\alpha_\mathrm{em} = 1/127.95$). The trace anomaly is written in the signature $(+,-,-,-)$, in which the trace of the energy-momentum tensor of a fluid is $\rho - 3p$; in the signature $(+,+,+,-)$ of UM-2 both sides change sign and the relation is the same; likewise on shell the Proca Lagrangian of a massive vector is a total derivative in either signature. The cold environments: the cosmic mean today (the Planck baryon density with the helium fraction 0.2457, $\Omega_\mathrm{DM} h_{100}^2 = 0.120$), the Galaxy at the Sun ($n_\mathrm{p} = n_\mathrm{e} = 1$ cm^{-3}), the Earth (mean density 5.51 g/cm^3, 48 % protons), the centre of the Sun (150

g/cm^3, 66 % protons, 32 % neutrons), a white dwarf (10^6 g/cm^3 of carbon and oxygen, equal numbers of protons and neutrons) and a neutron-star core (0.3 fm^{-3}, 10 % protons, $n_\mathrm{e} = n_\mathrm{p}$), with the local dark-matter density 0.4 GeV/cm^3 in the bound objects. In the early-Universe environments the densities are those of the field equation written above ($\eta_\mathrm{b} = 6.1 \times 10^{-10}$, $m_\mathrm{n} - m_\mathrm{p} = 1.293$ MeV), with the dark matter in equilibrium for $M_\mathrm{DM}/T < 40$ and its relic density scaled with the entropy afterwards; the freeze-out environment is at $T = M_\mathrm{DM}/x_\mathrm{f}$ with the $x_\mathrm{f} = 31.23$ of step 9. Results (DERIVED):

environment	n_p (cm^{-3})	$\Delta\varepsilon_\mathrm{I11}$ (GeV)	$\Delta\varepsilon_\mathrm{I13}$ (GeV)	$\Delta\varepsilon_\mathrm{I12}$ (GeV)	$\Delta\varepsilon/M_\mathrm{DM}$	$\Delta \ln f$
cosmic mean today	2.2×10^{-7}	3×10^{-64}	4×10^{-57}	3.7×10^{-15}	8.4×10^{-20}	7×10^{-94}
the Galaxy at the Sun	1	1×10^{-57}	1.5×10^{-50}	3.7×10^{-15}	8.4×10^{-20}	3×10^{-87}
the Earth (5.51 g/cm^3)	1.6×10^{24}	4.4×10^{-33}	4.9×10^{-26}	3.7×10^{-15}	8.4×10^{-20}	1×10^{-62}
centre of the Sun (150 g/cm^3)	6.0×10^{25}	1.1×10^{-31}	1.3×10^{-24}	3.7×10^{-15}	8.4×10^{-20}	3×10^{-61}
white dwarf (10^6 g/cm^3)	3.0×10^{29}	7.9×10^{-28}	8.9×10^{-21}	3.7×10^{-15}	8.4×10^{-20}	2×10^{-57}
neutron-star core (0.3 fm^{-3})	3.0×10^{37}	4.3×10^{-19}	4.4×10^{-12}	3.7×10^{-15}	1.0×10^{-16}	9×10^{-49}
nucleosynthesis ($T = 1$ MeV)	1.5×10^{22}	2.4×10^{-35}	2.8×10^{-28}	3.7×10^{-15}	8.4×10^{-20}	7×10^{-60}
QCD crossover ($T = 200$ MeV)	7.8×10^{28}	2.0×10^{-28}	2.3×10^{-21}	1.9×10^{-15}	4.4×10^{-20}	3×10^{-48}
electroweak era ($T = 100$ GeV)	9.7×10^{36}	2.6×10^{-20}	2.8×10^{-13}	-2.4×10^{-6}	-5.5×10^{-11}	7×10^{-39}
dark-matter freeze-out ($T = 1.42$ TeV)	2.8×10^{40}	7.3×10^{-17}	8.2×10^{-10}	-6.8×10^{-2}	-1.5×10^{-6}	1.1×10^{-34}

The QCD vacuum condensate dominates at low temperature ($\langle G^A_{\mu\nu} G^{A\mu\nu} \rangle = 0.474$ GeV^4, half of it left at 200 MeV); the thermal condensates dominate in the electroweak era and at the freeze-out, where the shift reaches 1.5×10^{-6} of the mass ($\langle G^A_{\mu\nu} G^{A\mu\nu} \rangle = -8.1 \times 10^{12}$, $\langle W^I_{\mu\nu} W^{I\mu\nu} \rangle = -5.0 \times 10^{11}$, almost all from the order-α interaction, $\langle B_{\mu\nu} B^{\mu\nu} \rangle = -2.6 \times 10^{10}$ GeV^4; all condensates in the geometric normalisation of the gauge fields).

11.3.3 dirac16complex acting on the Standard Model

Through I12 the scalar density of the dark matter changes the gauge couplings: $\Delta\alpha_3/\alpha_3 = -1.7 \times 10^{-16}$ at the freeze-out, -1.1×10^{-20} in the electroweak era, at most -1.1×10^{-36} from nucleosynthesis on, -1.7×10^{-60} in the Galaxy today. In the x_0 problem of the W with the kinetic function $\Upsilon = 1 + A_\mathrm{mf} n^\mathrm{sc}_\mathrm{DM}$ written above, first-order perturbation theory of the Sturm-Liouville problem with the exact W eigenfunction and the exact dark-matter orbital gives $\mathrm{d} \ln \tilde{m}^2_\mathrm{W} / \mathrm{d} A_\mathrm{mf} = -0.111253$, and direct shooting of the perturbed problem agrees to 3×10^{-11} (DERIVED; version 3 quoted -0.1126 from quadratic finite elements, whose finite differences in the amplitude are limited by rounding). The photon and gluon zero mode keeps $\tilde{m} = 0$ exactly: its profile is flat, so the kinetic function does not act on it (version 3 quoted the finite-element rounding -2.5×10^{-11}). The physical amplitude is of the size of the relative shift of the $SU(2)$ coupling, $A_\mathrm{mf} \sim |\Delta\alpha_2/\alpha_2| = 4 G_\mathrm{W} n_\mathrm{DM,tot}$, which is at most the shift of the strong coupling quoted above (with universal coefficients smaller by g_2^2/g_3^2): at most 1.7×10^{-16} at the freeze-out, 1.1×10^{-20} in the electroweak era, 1.1×10^{-36} from nucleosynthesis on and 5×10^{-66} for the cosmic mean today, so the squared W mass (the eigenvalue $\tilde{m}^2_\mathrm{W}$) moves by at most 1.9×10^{-17} of itself.

11.3.4 Gravity: the block response to every species

The block responds to the block-averaged sources of every species with the adiabatic response written above, in which radiation drops out; the hidden pressure is the response of the pressure to the block scale at fixed temperature,

through the species of mass $m_s(f)$ as ideal gases with their x_0 masses (for an ideal gas $m\,\partial p/\partial m = -(\rho - 3p)$; non-relativistic matter gives $-\frac{1}{4}\rho\,\mathrm{d}\ln m/\mathrm{d}\ln f$), the QCD plasma through $\partial p/\partial \ln \Lambda_{\mathrm{QCD}} = -\Theta_{\mathrm{QCD}}$ and the electroweak plasma through its order-α pressure, with the bulk couplings scaling as $\alpha_{\mathrm{G}}(m_{\mathrm{KK}}) \propto f^{-4}$ and $m_{\mathrm{KK}} \propto 1/f$; in the trace the QCD and electroweak plasmas add Θ_{QCD} and Θ_{G}. The sensitivities (DERIVED, and one loop where stated): $\mathrm{d}\ln m_{\mathrm{e}}/\mathrm{d}\ln f = -20.5$, $\mathrm{d}\ln M_{\mathrm{DM}}/\mathrm{d}\ln f = -0.772$, and for the quarks u, d, s, c, b, t -18.82, -18.01, -14.90, -12.15, -10.89, -6.77; through the QCD scale ($\alpha_3(m_{\mathrm{KK}}) = 0.0606$, $\sigma_{\pi N} = 45$ MeV, $\sigma_{\mathrm{s}} = 40$ MeV; ESTIMATE, one loop) -60.2 ($\Lambda^{(6)}_{\mathrm{QCD}}$), -55.6, -52.0, -49.0 ($\Lambda^{(3)}_{\mathrm{QCD}}$), and for the nucleon $\mathrm{d}\ln m_{\mathrm{p}}/\mathrm{d}\ln f = -46.1$, the most block-sensitive mass of UM-2. The block moves by at most $\Delta \ln f = 1.1 \times 10^{-34}$, at the freeze-out, where the hot plasma has $\bar{p} = 3.1 \times 10^{12}$ GeV^4 (mostly the QCD interaction measure with $\mathrm{d}\ln\Lambda_{\mathrm{QCD}}/\mathrm{d}\ln f = -60$) and $\rho - 3p_3 = 2.2 \times 10^{11}$ GeV^4 (the nucleon mass moves by 5.3×10^{-33}); by 7×10^{-39} in the electroweak era; by at most 9×10^{-49} from nucleosynthesis to today. The gain of the loop is at most 1.1×10^{-33}, so the iteration converges after one step in every environment. (Version 3 as first written kept only the hidden pressure of non-relativistic matter and only the $\Delta\bar{p}$ term of the response and found 5.6×10^{-45}; the conclusion is unchanged.)

The coupled solution is therefore the stabilised background with the x_0 spectra and the Kohn-Sham states of steps 1 to 3, corrected by the mean fields of this step; the corrections are computed and below 1.5×10^{-6}. The I12 benchmark changes the strong coupling by 10 % through the thermal dark-matter density only at $T = 27.4$ TeV ($M_{\mathrm{DM}}/T = 1.62$), so the coupled effective description holds through the freeze-out and everything later (DERIVED, 16 checks with step 9).




Figure 6: The coupled solution: the x_0 profiles that enter the overlaps (left) and the shift of the dark-matter level by the Standard-Model mean fields in the ten environments (right).

11.4 The check of step 8

In its own equations.

- *The overlaps.* Recomputed from the closed-form orbitals (the odd level, the fold zero mode, the first even level) and the zero-mode profiles with 2400 Gauss nodes refined at the fold, $\tilde{I}$, $\tilde{I}^{(b)}$ and $\tilde{I}_\lambda$ agree with the stored values to 3.8×10^{-12}, 3.5×10^{-12} and 1.3×10^{-12}; they equal the responses of self-consistent Kohn-Sham levels to finite mean fields to 3.0×10^{-10} (I11) and 2.3×10^{-10} (I13) (DERIVED).
- *The mean fields and the block response.* Every stored shift, condensate and block response of the ten environments, recomputed from the stored overlaps, strengths, densities, condensates and sources, agrees to 2.0×10^{-11} (relative); the vacuum condensate $4\pi^2 \times 0.012$ GeV^4 to 6×10^{-13}; the iterated and the one-step block shifts are equal (the loop gain is 1.1×10^{-33}) (DERIVED).
- *The condensate of a gauge plasma.* The Feynman-Hellmann condensate of the order-α pressure reproduces the trace-anomaly gluon condensate to 1.1×10^{-16} (DERIVED).
- *The W problem with the kinetic function.* The stored coefficient is reproduced by the same first-order integral on an independent quadrature to 3.3×10^{-12}, and the shooting of the perturbed problem agrees to 3.0×10^{-11} (DERIVED).

In concert with the other field equations.

- *The Kohn-Sham equations of step 3.* The portal responses of the Kohn-Sham jobs are the overlaps of the coupled solution to 3.0×10^{-12}; the hidden pressure of the dark matter in the block response is the Kohn-Sham stress and the x_0 sensitivity to 6.1×10^{-12}; the dark-matter mass agrees to 8.1×10^{-13} (DERIVED).
- *The relic and freeze-in couplings of step 9.* The strengths G_{A} and G_{V} implied by the stored shifts at the freeze-out are those of the relic and of the freeze-in to 6.7×10^{-13}; the I12 benchmark is that of the relic computation to 1.6×10^{-12}; the freeze-out temperature of the environment is $M_{\mathrm{DM}}/x_{\mathrm{f}}$ of the relic to 2.1×10^{-12} (DERIVED).
- *The thermal history of step 9.* The plasma energy density at the freeze-out in the block sources (1.416139×10^{14} GeV^4) differs from the ideal gases of the thermal history (1.414292×10^{14} GeV^4) by 1.3×10^{-3}, by design: the block sources add the trace anomaly of the interacting plasma (of order $\alpha_3^2 T^4$), which enters only $\rho - 3p_3$; the order-α interaction energy (5 %) is absent from both (DERIVED).
- *The x_0 problems of step 2.* The W eigenvalue of the finite elements at zero amplitude, stored with the kinetic-function computation, is the exact eigenvalue to 2.3×10^{-6} (the rounding of the matrix Rayleigh quotient), and the shooting of the perturbed problem at zero amplitude to 1.4×10^{-11}; the block sensitivities of the masses used in the hidden pressure equal the first-order x_0 mass formula to 4.5×10^{-11}; the response coefficients are those of step 1 to 2×10^{-13} (DERIVED).

11.5 The status of step 8

DERIVED: the overlaps and their Hellmann-Feynman checks, the mean-field shifts in the ten environments, the coupling shifts and the W problem, the block response to every species and the convergence of the coupled iteration. ESTIMATE: the QCD interaction measure below 1 GeV, the electroweak condensate below m_{W}, the one-loop QCD scale and nucleon sensitivity. STANDARD: the vacuum gluon condensate, the trace anomaly and the leading-order plasma pressures. ASSUMED: the compositions of the environments and the benchmark strengths (fixed in step 9). INVENTED: the portals.

12 Step 9: the thermal history and the relic with I13

Step 9 computes the thermal history of UM-2 with the x_0 masses of step 2, solves the Boltzmann equation of the dark matter with the cross-sections of I12 and I13 and their partial-wave unitarity bounds, fixes the benchmark strengths of the three portals (I13 by the relic, I11 by the freeze-in of the fold bands, I12 by unitarity), shows why I13 couples to the baryon current and not to $B - L$, bounds the self-coupling of dirac16complex by the dark radiation it would make, and estimates the detection of the dark matter.

12.1 The field equations of step 9

The portals whose strengths step 9 fixes:

$$S_{\mathrm{I11}} = \int \mathrm{d}^8X \, \sqrt{|g|} \, \frac{c_{\mathrm{V}}}{\Lambda_\star^6} \left(\mathrm{i}\bar{\Psi}\gamma^M\Psi \right) J_M^{B-L}, \qquad J_M^{B-L} = \sum_{i=1}^{3} \sum_{\psi} q_\psi \left(\mathrm{i}\bar{\psi}_i \gamma_M \psi_i \right),$$
$$q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}, \qquad q_\ell = q_{e_R} = q_{\nu_R} = -1 \tag{IN-16}$$

Symbols in (IN-16).

- S_{I11}: The action of the portal I11 (dirac16complex vector current times the $B - L$ current of the fermions). Unit: dimensionless. Status: INVENTED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0 \, \mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\dots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, ${g^M}_N=\delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- c_{V}: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_{\mathrm{V}}=4.38\times10^{-17}$ GeV^{-2}. Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- Ψ, $\bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi=\Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M={\mathbf{e}_a}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M=g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1,M_2,\dots$): it runs over $x_0,x_1,\dots,x_7$; a repeated index is summed. Status: STANDARD.
- J^{B-L}_M: The $B-L$ current of the Standard-Model fermions (baryon minus lepton number), summed over the generations. Unit: GeV^7. Status: STANDARD.
- i: The generation index ($i=1,2,3$), which also labels the neutrino mass states ν_1,ν_2,ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- ψ: The label of a Standard-Model multiplet: $\psi\in\{Q,u_R,d_R,\ell,e_R,\nu_R\}$ (quark doublet, up and down quark singlets, lepton doublet, charged-lepton singlet, right-handed neutrino); a sum over ψ runs over the six. Status: STANDARD.
- q_ψ, q_f: The $B-L$ charge: q_ψ of the multiplet ψ ($q_Q=q_{u_R}=q_{d_R}=\frac13$, $q_\ell=q_{e_R}=q_{\nu_R}=-1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- ψ_i, $\bar\psi_i$: A Standard-Model fermion field of eight dimensions: ψ_i is the multiplet $\psi\in\{Q,u_R,d_R,\ell,e_R,\nu_R\}$ of the generation i, ψ_f one fermion component f (in four-dimensional formulas its zero mode); $\bar\psi=\psi^\dagger C$ the Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$ (eight dimensions), $\mathrm{GeV}^{3/2}$ (four dimensions). Status: ASSUMED.

$$S_{\mathrm{I12}}=-\int\mathrm{d}^8X\,\sqrt{|g|}\,\frac{\bar\Psi\Psi}{\Lambda_\star^3}\,\mathcal{F},\qquad \mathcal{F}=c_{\mathrm{G}}\,G^A_{MN}G^{AMN}+c_{\mathrm{W}}\,W^I_{MN}W^{IMN}+c_{\mathrm{B}}\,B_{MN}B^{MN}\tag{IN-17}$$

Symbols in (IN-17).

- S_{I12}: The action of the portal I12 (scalar density of dirac16complex times the gauge kinetic invariants). Unit: dimensionless. Status: INVENTED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X=\{x_0,\dots,x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X=\{x_0,x_1,\dots,x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X=\mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1,\dots,x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\dots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, ${g^M}_N=\delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- Ψ, $\bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi=\Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- $\mathcal{F}$: The gauge-kinetic operator of the portal I12, $\mathcal{F}=c_{\mathrm{G}}G^A_{MN}G^{AMN}+c_{\mathrm{W}}W^I_{MN}W^{IMN}+c_{\mathrm{B}}B_{MN}B^{MN}$; $\langle\mathcal{F}\rangle$ its mean value in an environment. Unit: GeV^4. Status: INVENTED.
- c_{G}: The coefficient of the gluon term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{G}}/\Lambda_\star^3=c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.

- G^A_M, G_M, G_{MN}, G^A_{MN}: The gluon field (SU(3)), in the geometric normalisation of the action (couplings in front of the kinetic terms): the components G^A_M ($A = 1, \ldots, 8$), the matrix $G_M = G^A_M t^A$, the field strength G_{MN} (matrix) and its components G^A_{MN}; $G^A_{\mu\nu}$ are its components along the observable directions (the zero mode, flat in x_0). Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- c_{W}: The coefficient of the SU(2) term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{W}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- W^I_M, W_M, W_{MN}, W^I_{MN}: The SU(2) gauge field (geometric normalisation): the components W^I_M ($I = 1, 2, 3$; W^3 the third component), the matrix $W_M = W^I_M \tau^I/2$, the field strength W_{MN} and its components W^I_{MN}; with Greek indices the components along the observable directions. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- c_{B}: The coefficient of the hypercharge term of the portal I12. Unit: dimensionless. Value: $c_{\mathrm{B}}/\Lambda_\star^3 = c_{\mathrm{I12}}/\Lambda_\star^3$. Status: INVENTED.
- B_M, B_{MN}: The hypercharge gauge field $\mathrm{U}(1)_Y$ (geometric normalisation) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: ASSUMED.

$$S_{\mathrm{I13}} = \int \mathrm{d}^8X\, \sqrt{|g|}\, \frac{c_{\mathrm{A}}}{\Lambda_\star^6} \sum_{M \neq x_0} \left(\mathrm{i}\bar\Psi \gamma^M P_b \Psi\right) J^B_M, \qquad P_b = \frac{1-\gamma^{x_0}}{2}, \qquad P_a = \frac{1+\gamma^{x_0}}{2},$$
$$J^B_M = \frac{1}{3}\sum_{i=1}^{3} \left(\mathrm{i}\bar Q_i \gamma_M Q_i + \mathrm{i}\bar u_{Ri}\gamma_M u_{Ri} + \mathrm{i}\bar d_{Ri}\gamma_M d_{Ri}\right) \tag{IN-18}$$

Symbols in (IN-18).

- S_{I13}: The action of the portal I13 (the x_0-chiral vector current of dirac16complex times the baryon-number current). Unit: dimensionless. Status: INVENTED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\, \mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\, \mathrm{d}X^M \mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, ${g^M}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\, \mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- c_{A}: The coefficient of the x_0-chiral vector portal I13. Unit: dimensionless. Value: fixed through $G_{\mathrm{A}} = 1.278 \times 10^{-9}\ \mathrm{GeV}^{-2}$ (dark-matter relic). Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- Ψ, $\bar\Psi$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar\Psi = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = {\mathbf{e}_a}^M \gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.

- P_a, P_b: The projectors on the two x_0-chiralities, $P_a = (1+\gamma^{x_0})/2$ and $P_b = (1-\gamma^{x_0})/2$ (in a block: the upper and the lower component). Unit: dimensionless. Status: INVENTED.
- J^B_M: The baryon-number current of the quarks (one third per quark), summed over the generations. Unit: GeV^7. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- Q_i, $\bar{Q}_i$: The quark doublet of the generation i (left-handed up and down quark of one generation, three colours) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- u_R, u_{Ri}, $\bar{u}_{Ri}$: The up-type quark singlet: u_R the multiplet, u_{Ri} that of the generation i (u_R, c_R, t_R), $\bar{u}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- d_R, d_{Ri}, $\bar{d}_{Ri}$: The down-type quark singlet: d_R the multiplet, d_{Ri} that of the generation i (d_R, s_R, b_R), $\bar{d}_{Ri}$ its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.

The degrees of freedom of the hot Universe:

$$\rho_s = \frac{d_s}{2\pi^2}\int_0^\infty \frac{k^2 E\,\mathrm{d}k}{\mathrm{e}^{E/T}\pm 1}, \qquad p_s = \frac{d_s}{6\pi^2}\int_0^\infty \frac{k^4\,\mathrm{d}k}{E\,(\mathrm{e}^{E/T}\pm 1)}, \qquad s_s = \frac{\rho_s + p_s}{T}, \qquad E = \sqrt{k^2+m_s^2} \qquad \text{(FE-121)}$$

Symbols in (FE-121).

- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- s: The label of a particle species of the plasma or of matter (photon, gluons, $W^\pm$, Z, quarks, leptons, nucleons, dark matter); a sum over s runs over the species present. Status: STANDARD.
- d_o, d_s: A degeneracy: $d_o = 4r_3(|\vec{n}_o|^2)$ the number of states of the momentum shell of the orbital o; d_s the number of internal states (spin, colour, particle and antiparticle) of the species s. Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no upper bound. Status: STANDARD.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- E: The energy of a particle: of the neutrino in the flavour equations, $E = (k^2+m^2)^{1/2}$ of a particle of momentum k in the thermal integrals. Unit: GeV. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.
- p: A pressure: $p_M = \mathcal{T}^M{}_M$ (no sum) for the direction $M \neq x_4$, p_s the pressure of the species s; Δp the pressure perturbation in the cosmological perturbation equations; in the Feynman-Hellmann relation the pressure of a gauge plasma. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- s, s_s: The entropy density of the plasma, $s = 2\pi^2 g_s T^3/45$; s_s that of the species s. Unit: GeV^3. Status: STANDARD.
- m, m_s, m_i: A four-dimensional mass: m the mass of a mode (the eigenvalue of an x_0 problem) or of a fermion; m_s the mass of the species s; m_i the mass of the neutrino mass state i. Unit: GeV. Status: DERIVED.

$$g_\rho = \frac{30}{\pi^2 T^4}\sum_s \rho_s, \qquad g_s = \frac{45}{2\pi^2 T^3}\sum_s s_s, \qquad \mathcal{H} = \Big(\frac{8\pi^3 g_\rho}{90}\Big)^{1/2}\frac{T^2}{M_{\mathrm{Pl}}} \qquad \text{(FE-122)}$$

Symbols in (FE-122).

- $g_\rho,\ g_s$: The effective numbers of relativistic degrees of freedom of the energy density and of the entropy density of the plasma. Unit: dimensionless. Value: 105.75 when every species of UM-2 in equilibrium is relativistic (no Higgs particle), 10.75 near 1 MeV, $g_s = 43/11$ today. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.
- s: The label of a particle species of the plasma or of matter (photon, gluons, $W^\pm$, Z, quarks, leptons, nucleons, dark matter); a sum over s runs over the species present. Status: STANDARD.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.
- $s,\ s_s$: The entropy density of the plasma, $s = 2\pi^2 g_s T^3/45$; s_s that of the species s. Unit: GeV^3. Status: STANDARD.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2} T^2/M_{\mathrm{Pl}}$). Unit: GeV. Status: DERIVED.
- M_{Pl}: The Planck mass, $M_{\mathrm{Pl}} = G_{\mathrm{N}}^{-1/2}$. Unit: GeV. Value: 1.22091×10^{19} GeV. Status: STANDARD.

$$w_{\mathrm{QGP}}(T) = \frac{1}{2}\Big[1 + \tanh\frac{T - T_{\mathrm{c}}}{T_{\mathrm{w}}}\Big], \qquad \frac{T_\nu}{T} = \Big(\frac{g_{s,e\gamma}(T)}{11/2}\Big)^{1/3}, \qquad \frac{\mathrm{d}\ln T}{\mathrm{d}x_4} = -\frac{\mathcal{H}}{1 + \frac{1}{3}\,\mathrm{d}\ln g_s/\mathrm{d}\ln T} \tag{FE-123}$$

Symbols in (FE-123).

- $w_{\mathrm{QGP}}(T)$: The weight of the quark-gluon plasma in the model of the QCD crossover (1 above, 0 below). Unit: dimensionless. Status: ASSUMED.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.
- $T_{\mathrm{c}},\ T_{\mathrm{w}}$: The temperature and the width of the QCD crossover. Unit: GeV. Value: 155 MeV and 15 MeV. Status: ASSUMED.
- T_ν: The temperature of the relic neutrinos (one common temperature). Unit: GeV. Value: 1.9454 K today. Status: STANDARD.
- $g_{s,e\gamma}$: The entropy degrees of freedom of the electrons, positrons and photons only (after neutrino decoupling). Unit: dimensionless. Value: 11/2 before and 2 after the annihilation of $e^\pm$. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2} T^2/M_{\mathrm{Pl}}$). Unit: GeV. Status: DERIVED.
- $g_\rho,\ g_s$: The effective numbers of relativistic degrees of freedom of the energy density and of the entropy density of the plasma. Unit: dimensionless. Value: 105.75 when every species of UM-2 in equilibrium is relativistic (no Higgs particle), 10.75 near 1 MeV, $g_s = 43/11$ today. Status: STANDARD.

The electroweak couplings of the unitarity-safe I12 benchmark are run to $2M_{\mathrm{DM}}$ with the one-loop equations of step 8:

$$\frac{1}{\alpha_2(\mu_{\mathrm{ren}})} = \frac{\sin^2\theta_{\overline{\mathrm{MS}}}}{\alpha_{\mathrm{em}}} + \frac{b_2}{2\pi}\ln\frac{\mu_{\mathrm{ren}}}{m_{\mathrm{Z}}}, \qquad \frac{1}{\alpha_1(\mu_{\mathrm{ren}})} = \frac{1 - \sin^2\theta_{\overline{\mathrm{MS}}}}{\alpha_{\mathrm{em}}} + \frac{b_1}{2\pi}\ln\frac{\mu_{\mathrm{ren}}}{m_{\mathrm{Z}}}, \qquad b_2 = \frac{10}{3}, \quad b_1 = -\frac{20}{3} \tag{FE-77}$$

Symbols in (FE-77).

- α_2: The SU(2) fine-structure constant, $\alpha_2 = g_2^2/(4\pi)$. Unit: dimensionless. Status: STANDARD.
- μ_{ren}: The renormalisation scale of a running coupling (taken as $2\pi T$ for the condensates of a plasma). Unit: GeV. Status: STANDARD.

- $\theta_{\overline{\rm MS}}$: The weak mixing angle in the $\overline{\rm MS}$ scheme at $m_{\rm Z}$. Unit: dimensionless. Value: $\sin^2\theta_{\overline{\rm MS}} = 0.23122$. Status: STANDARD.
- $\alpha_{\rm em}$: The electromagnetic fine-structure constant ($\overline{\rm MS}$, at $m_{\rm Z}$). Unit: dimensionless. Value: 1/127.95. Status: STANDARD.
- $b_3,\ b_2,\ b_1,\ b_{\rm G}$: The one-loop coefficients of the running couplings, $\mathrm{d}\alpha/\mathrm{d}\ln\mu_{\rm ren} = -b\,\alpha^2/(2\pi)$: $b_3 = 11 - \frac{2}{3}N_{\rm f}$ (SU(3)), $b_2 = \frac{10}{3}$ (SU(2) without a Higgs particle), $b_1 = -\frac{20}{3}$ ($\mathrm{U}(1)_Y$), $b_{\rm G}$ that of the group G (not related to the metric function b_4). Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $m_{\rm Z}$: The mass of the Z. Unit: GeV. Value: 91.1879 GeV (reproduced). Status: STANDARD.
- α_1: The hypercharge fine-structure constant, $\alpha_1 = g_1^2/(4\pi)$. Unit: dimensionless. Status: STANDARD.

The dark matter: the Boltzmann equation, the thermal average, the cross-sections and their unitarity bounds, with the four-dimensional couplings of the portals on the zero modes (step 8):

$$\frac{\mathrm{d}n_{\rm DM}}{\mathrm{d}x_4} + 3\mathcal{H}n_{\rm DM} = -\langle\sigma v_{\rm rel}\rangle\big(n_{\rm DM}\bar{n}_{\rm DM} - n_{\rm DM,eq}^2\big) \ \Rightarrow \frac{\mathrm{d}Y_{\rm DM}}{\mathrm{d}x} = -\Big(1 + \frac{1}{3}\frac{\mathrm{d}\ln g_s}{\mathrm{d}\ln T}\Big)\frac{s\,\langle\sigma v_{\rm rel}\rangle}{2x\mathcal{H}}\big(Y_{\rm DM}^2 - Y_{\rm eq}^2\big),$$
$$Y_{\rm DM} = \frac{n_{\rm DM} + \bar{n}_{\rm DM}}{s}, \qquad x = \frac{M_{\rm DM}}{T}, \qquad s = \frac{2\pi^2}{45}\,g_s T^3, \qquad Y_{\rm eq} = \frac{4M_{\rm DM}^2 T K_2(x)}{2\pi^2 s} \tag{FE-124}$$

Symbols in (FE-124).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $n_{\rm DM},\ \bar{n}_{\rm DM}$: The number densities of dark-matter particles and antiparticles of one species. Unit: GeV^3. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_{\rm Pl}$). Unit: GeV. Status: DERIVED.
- σ: A cross-section (of the dark-matter annihilation or production process named in the equation; summed over the spins where marked). Unit: GeV^{-2}. Status: STANDARD.
- $v_{\rm rel}$: The relative velocity of the two annihilating particles; $\langle\sigma v_{\rm rel}\rangle$ the thermal average. Unit: dimensionless (c = 1). Status: STANDARD.
- $n_{\rm DM,eq},\ \bar{n}_{\rm DM,eq}$: The equilibrium number densities of dark-matter particles and antiparticles of one species. Unit: GeV^3. Status: STANDARD.
- $Y_{\rm DM},\ Y_{\rm eq},\ Y_\infty$: The abundance (number density per entropy density) of one dark-matter species (particles and antiparticles), its equilibrium value and its final value. Unit: dimensionless. Status: STANDARD.
- x: The ratio $x = M_{\rm DM}/T$ of the dark-matter mass to the temperature. Unit: dimensionless. Value: $x_{\rm f} = 31.23$ at the freeze-out. Status: STANDARD.
- $g_\rho,\ g_s$: The effective numbers of relativistic degrees of freedom of the energy density and of the entropy density of the plasma. Unit: dimensionless. Value: 105.75 when every species of UM-2 in equilibrium is relativistic (no Higgs particle), 10.75 near 1 MeV, $g_s = 43/11$ today. Status: STANDARD.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.
- $s,\ s_s$: The entropy density of the plasma, $s = 2\pi^2 g_s T^3/45$; s_s that of the species s. Unit: GeV^3. Status: STANDARD.
- $M_{\rm DM}$: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $K_1,\ K_2$: The modified Bessel functions of the second kind of orders 1 and 2 (not related to the block variable K). Unit: dimensionless. Status: STANDARD.

$$\langle\sigma v_{\rm rel}\rangle = \frac{1}{8M_{\rm DM}^4 T K_2(x)^2}\int_{4M_{\rm DM}^2}^{\infty}\sigma\left(E_{\rm cm}^2 - 4M_{\rm DM}^2\right)E_{\rm cm}\,K_1\Big(\frac{E_{\rm cm}}{T}\Big)\,\mathrm{d}\big(E_{\rm cm}^2\big) \tag{FE-125}$$

Symbols in (FE-125).

- σ: A cross-section (of the dark-matter annihilation or production process named in the equation; summed over the spins where marked). Unit: GeV^{-2}. Status: STANDARD.
- v_{rel}: The relative velocity of the two annihilating particles; $\langle \sigma v_{\mathrm{rel}} \rangle$ the thermal average. Unit: dimensionless ($c = 1$). Status: STANDARD.
- M_{DM}: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.
- $K_1,\ K_2$: The modified Bessel functions of the second kind of orders 1 and 2 (not related to the block variable K). Unit: dimensionless. Status: STANDARD.
- x: The ratio $x = M_{\mathrm{DM}}/T$ of the dark-matter mass to the temperature. Unit: dimensionless. Value: $x_{\mathrm{f}} = 31.23$ at the freeze-out. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no upper bound. Status: STANDARD.
- E_{cm}: The total energy of the colliding pair in its centre-of-mass frame. Unit: GeV. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.

$$\Omega_{\mathrm{DM}} h_{100}^2 = \frac{M_{\mathrm{DM}}\, Y_\infty\, s_0}{\rho_{\mathrm{crit}}/h_{100}^2}, \qquad \frac{\rho_{\mathrm{crit}}}{h_{100}^2} = 1.05368 \times 10^{-5}\ \mathrm{GeV/cm^3} \tag{FE-126}$$

Symbols in (FE-126).

- Ω_{DM}: The dark-matter density today in units of the critical density. Unit: dimensionless. Value: $\Omega_{\mathrm{DM}} h_{100}^2 = 0.120$ (both species). Status: DERIVED.
- h_{100}: The reduced Hubble constant, $\mathcal{H}_0/(100\ \mathrm{km\ s^{-1}\ Mpc^{-1}})$. Unit: dimensionless. Value: 0.6738. Status: STANDARD.
- M_{DM}: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- $Y_{\mathrm{DM}},\ Y_{\mathrm{eq}},\ Y_\infty$: The abundance (number density per entropy density) of one dark-matter species (particles and antiparticles), its equilibrium value and its final value. Unit: dimensionless. Status: STANDARD.
- s_0: The entropy density today. Unit: GeV^3 (quoted per cm^3). Value: $2891.3\ \mathrm{cm}^{-3}$. Status: STANDARD.
- ρ_{crit}: The critical density, $3\mathcal{H}_0^2 M_{\mathrm{Pl}}^2/(8\pi)$. Unit: GeV^4 (quoted in $\mathrm{GeV/cm^3}$). Value: $\rho_{\mathrm{crit}}/h_{100}^2 = 1.05368 \times 10^{-5}\ \mathrm{GeV/cm^3}$. Status: STANDARD.

$$\begin{aligned}
&\mathcal{L}_4 = \big(G_{\mathrm{V}}\, q_f\, \tilde{I}_{of} + G_{\mathrm{A}}\, q_f^B\, \tilde{I}_{of}^{(b)}\big)\big(\mathrm{i}\bar{\Psi}_o \gamma^\mu \Psi_o\big)\big(\mathrm{i}\bar{\psi}_f \gamma_\mu \psi_f\big),\\
&\tilde{I}_{of} = \frac{1}{H}\int_{-L}^{0} \big(a_o^2 + b_o^2\big)\varrho_f\, \mathrm{e}^{-3Hy}\,\mathrm{d}y, \qquad \tilde{I}_{of}^{(b)} = \frac{1}{H}\int_{-L}^{0} b_o^2\, \varrho_f\, \mathrm{e}^{-3Hy}\,\mathrm{d}y, \qquad G_{\mathrm{V}} = \frac{c_{\mathrm{V}} H}{2 f^4 v_{\mathrm{t}} \Lambda_\star^6}, \qquad G_{\mathrm{A}} = \frac{c_{\mathrm{A}} H}{2 f^4 v_{\mathrm{t}} \Lambda_\star^6},\\
&n_{\psi i}(y) = \frac{n_f\, \varrho_f(y)\, \mathrm{e}^{-3Hy}}{2 f^4 v_{\mathrm{t}}}, \qquad \varrho(y;c) = \frac{2c}{L}\,\frac{\mathrm{e}^{2cy/L}}{1-\mathrm{e}^{-2c}}
\end{aligned} \tag{FE-70}$$

Symbols in (FE-70).

- $\mathcal{L}_4$: The four-dimensional contact Lagrangian of I11 and I13 between a state o of dirac16complex and a Standard-Model fermion component f. Unit: GeV^4. Status: DERIVED.
- G_{V}: The four-dimensional strength of the vector portal I11 on the zero modes, $G_{\mathrm{V}} = c_{\mathrm{V}} H/(2 f^4 v_{\mathrm{t}} \Lambda_\star^6)$. Unit: GeV^{-2}. Value: benchmark $4.38 \times 10^{-17}\ \mathrm{GeV}^{-2}$ ($\Delta N_{\mathrm{eff}} = 0.001$). Status: DERIVED.
- $q_\psi,\ q_f$: The $B - L$ charge: q_ψ of the multiplet ψ ($q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}$, $q_\ell = q_{e_R} = q_{\nu_R} = -1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- f: The label of a fermion species: a Standard-Model multiplet of one generation (its N_{c} Weyl components share one zero-mode profile; the zero mode of each chirality) or a fermion of a medium (electron, proton, neutron); a sum over f runs over these species. Status: STANDARD.

- $\tilde{I}_{of}$: The overlap of the state o of dirac16complex with the zero mode of the fermion component f, $\tilde{I}_{of} = H^{-1}\int_{-L}^{0}(a_o^2+b_o^2)\varrho_f \mathrm{e}^{-3Hy}\mathrm{d}y$ (the integral in units of H); $\tilde{I}_Q$, $\tilde{I}_{u_R}$, $\tilde{I}_{d_R}$, $\tilde{I}_\ell$, $\tilde{I}_{e_R}$ those of the dark matter with Q, u_R, d_R, ℓ, e_R, $\tilde{I}_{\nu_R,i}$ with the right-handed neutrino of the generation i. Unit: dimensionless. Value: dark matter: $\tilde{I}_Q = 8.2722$, $\tilde{I}_\ell = 1.2524$, $\tilde{I}_{u_R} = 0.3854$, $\tilde{I}_{d_R} = 0.3961$, $\tilde{I}_{e_R} = 0.3972$. Status: DERIVED.
- o: The label of a Kohn-Sham orbital (a state of dirac16complex in the cell: block, momentum shell, fold parity and level); a sum over o runs over the orbitals. Status: RECORD.
- G_A: The four-dimensional strength of the portal I13 on the zero modes, $G_\mathrm{A} = c_\mathrm{A} H/(2f^4 v_\mathrm{t}\Lambda_\star^6)$. Unit: GeV^{-2}. Value: 1.278×10^{-9} GeV^{-2} (dark-matter relic density). Status: DERIVED.
- q_ψ^B, q_f^B: The baryon number of the multiplet ψ or of the fermion component f ($\frac{1}{3}$ for quarks, 0 for leptons). Unit: dimensionless. Status: STANDARD.
- $\tilde{I}_{of}^{(b)}$: The overlap of the lower component of the state o with the zero mode of f, $\tilde{I}_{of}^{(b)} = H^{-1}\int_{-L}^{0} b_o^2 \varrho_f \mathrm{e}^{-3Hy}\mathrm{d}y$ (the overlap of I13); $\tilde{I}_Q^{(b)}$, $\tilde{I}_{u_R}^{(b)}$, $\tilde{I}_{d_R}^{(b)}$, $\tilde{I}_{\nu_R}^{(b)}$ those of the dark matter. Unit: dimensionless. Value: dark matter: $\tilde{I}_Q^{(b)} = 2.6135$. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- Ψ, $\bar{\Psi}$: dirac16complex: the 16-component complex spinor field of eight dimensions of the record (the author's field), the source of the dark matter (its lowest odd level) and of the massless fold bands; $\bar{\Psi} = \Psi^\dagger C$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- γ^a, γ^M, γ^μ: The Dirac matrices: γ^a with a frame index are the author's real 16×16 matrices of the Clifford algebra of signature $(4,4)$; $\gamma^M = \mathbf{e}_a{}^M\gamma^a$ along a coordinate (γ^{x_0}, γ^{x_4}, … the matrices along x_0, x_4, …), $\gamma_M = g_{MN}\gamma^N$; γ^μ the matrices of the four observable directions in four-dimensional formulas. Unit: dimensionless. Status: AUTHOR.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- ψ_i, $\bar{\psi}_i$: A Standard-Model fermion field of eight dimensions: ψ_i is the multiplet $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ of the generation i, ψ_f one fermion component f (in four-dimensional formulas its zero mode); $\bar{\psi} = \psi^\dagger C$ the Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$ (eight dimensions), $\mathrm{GeV}^{3/2}$ (four dimensions). Status: ASSUMED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926\times10^{-21}$ m at the stabilised block). Status: ASSUMED.
- a, b: The upper and lower components of a block spinor, $\chi = (a, \mathrm{i}b)$, real at $k = 0$: a_L, b_L of a doublet and a_R, b_R of a singlet solution (a_{Ri}, b_{Ri} of the generation i, $b_{L\alpha}$, $a_{L\alpha}$ of the doublet in the flavour basis), a_o, b_o of the orbital o. Unit: $\mathrm{GeV}^{1/2}$ (normalised orbital). Status: DERIVED.
- $\varrho_f(y)$, $\varrho(y;c)$: The x_0 profile of a zero mode as a probability density on one half, $\int_{-L}^{0}\varrho\,\mathrm{d}y = 1$: ϱ_f that of the fermion component f, $\varrho(y;c)$ its form for the localisation c. Unit: 1/length (GeV). Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- c_V: The coefficient of the vector portal I11. Unit: dimensionless. Value: benchmark through $G_\mathrm{V} = 4.38\times10^{-17}$ GeV^{-2}. Status: INVENTED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- v_t: The coordinate volume of the compact extra times x_5, x_6, x_7. Unit: GeV^{-3}. Value: not fixed; $v_\mathrm{t} = H^{-3}$ in the quoted conversions. Status: ASSUMED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda_\star^n$ enter. Status: INVENTED.
- c_A: The coefficient of the x_0-chiral vector portal I13. Unit: dimensionless. Value: fixed through $G_\mathrm{A} = 1.278\times10^{-9}$ GeV^{-2} (dark-matter relic). Status: INVENTED.
- n, n_o, $n_{\psi i}$, n_f, n_i: A number density: $n(y)$ the particle density of dirac16complex in the Kohn-Sham

equations and n_o the contribution of the orbital o, $n_{\psi i}(y)$ the density of the Standard-Model fermion ψ_i (per proper 7-volume of eight dimensions); n_f the density of the fermion component f and n_i that of the neutrino mass state i (per 3-volume of four dimensions). Unit: GeV^7 (eight dimensions), GeV^3 (four dimensions). Status: STANDARD.

- ψ: The label of a Standard-Model multiplet: $\psi \in \{Q, u_R, d_R, \ell, e_R, \nu_R\}$ (quark doublet, up and down quark singlets, lepton doublet, charged-lepton singlet, right-handed neutrino); a sum over ψ runs over the six. Status: STANDARD.
- i: The generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- c, c_L, c_R: The localisation of a zero mode, $c = f\mu L$ (bulk mass times the proper depth); c_L of a doublet (localised at the fold for $c_L > 0$), c_R of a singlet (localised at the fold for $c_R < 0$). Unit: dimensionless. Value: doublets $c_L = 3.38230$ (quarks) and 5.95288 (leptons); singlets c_R: up -15.4300, down -14.6258, strange -11.5158, charm -8.7683, bottom -7.5005, top -3.3823, electron -14.5539, muon -8.9809, tau -5.9529, neutrinos -35.0452, -32.8481, -31.0702 (fixed by the measured masses). Status: DERIVED.

$$\sigma_{\mathrm{I12}} = \frac{G_{\mathrm{I12}}^2\, E_{\mathrm{cm}}^3 \big(E_{\mathrm{cm}}^2 - 4M_{\mathrm{DM}}^2\big)^{1/2}}{8\pi}, \qquad G_{\mathrm{I12}}^2 = 8G_{\mathrm{g}}^2 + 3G_{\mathrm{W}}^2 + G_{\mathrm{B}}^2,$$
$$\sigma_{\mathrm{I13}} = \sum_f \frac{N_{\mathrm{c}}\, G_f^2\, E_{\mathrm{cm}}^2 \big(1 + 2M_{\mathrm{DM}}^2/E_{\mathrm{cm}}^2\big)}{24\pi\beta_{\mathrm{cm}}}, \qquad G_f = G_{\mathrm{A}}\, q_f^B\, \tilde{I}_{of}^{(b)}, \tag{FE-127}$$
$$\beta_{\mathrm{cm}} = \sqrt{1 - \frac{4M_{\mathrm{DM}}^2}{E_{\mathrm{cm}}^2}}, \qquad k_{\mathrm{cm}} = \sqrt{\frac{E_{\mathrm{cm}}^2}{4} - M_{\mathrm{DM}}^2}$$

Symbols in (FE-127).

- σ_{I12}, σ_{I13}: The spin-averaged annihilation cross-sections of a dark-matter pair through I12 (into gauge-boson pairs) and through I13 (into quark pairs). Unit: GeV^{-2}. Status: DERIVED.
- G_{I12}: The combined coupling of I12 to all gauge-boson pairs, $G_{\mathrm{I12}}^2 = 8G_{\mathrm{g}}^2 + 3G_{\mathrm{W}}^2 + G_{\mathrm{B}}^2$. Unit: GeV^{-3}. Value: benchmark $\pi/(2M_{\mathrm{DM}}^3) = 1.801 \times 10^{-14}\ \mathrm{GeV}^{-3}$. Status: DERIVED.
- E_{cm}: The total energy of the colliding pair in its centre-of-mass frame. Unit: GeV. Status: STANDARD.
- M_{DM}: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_{g}: The coupling of a dark-matter pair to canonically normalised gluon pairs through I12. Unit: GeV^{-3}. Value: $6.09 \times 10^{-15}\ \mathrm{GeV}^{-3}$ (benchmark). Status: DERIVED.
- G_{W}: The coupling of a dark-matter pair to canonically normalised SU(2) gauge pairs through I12. Unit: GeV^{-3}. Value: $2.96 \times 10^{-15}\ \mathrm{GeV}^{-3}$. Status: DERIVED.
- G_{B}: The coupling of a dark-matter pair to canonically normalised hypercharge pairs through I12. Unit: GeV^{-3}. Value: $1.08 \times 10^{-15}\ \mathrm{GeV}^{-3}$. Status: DERIVED.
- f: The label of a fermion species: a Standard-Model multiplet of one generation (its N_{c} Weyl components share one zero-mode profile; the zero mode of each chirality) or a fermion of a medium (electron, proton, neutron); a sum over f runs over these species. Status: STANDARD.
- N_{c}: The number of Weyl components of a fermion multiplet that share its zero-mode profile and overlap (colour times isospin): 6 for a quark doublet, 3 for a quark singlet, 2 for a lepton doublet, 1 for a charged-lepton singlet. Unit: dimensionless. Status: STANDARD.
- G_f: The I13 coupling of the dark matter to the quark component f, $G_f = G_{\mathrm{A}} q_f^B \tilde{I}_{of}^{(b)}$. Unit: GeV^{-2}. Status: DERIVED.
- β_{cm}: The velocity of each particle of the pair in the centre-of-mass frame. Unit: dimensionless. Status: STANDARD.
- G_{A}: The four-dimensional strength of the portal I13 on the zero modes, $G_{\mathrm{A}} = c_{\mathrm{A}} H/(2f^4 v_{\mathrm{t}} \Lambda_\star^6)$. Unit: GeV^{-2}. Value: $1.278 \times 10^{-9}\ \mathrm{GeV}^{-2}$ (dark-matter relic density). Status: DERIVED.
- q_ψ^B, q_f^B: The baryon number of the multiplet ψ or of the fermion component f ($\frac{1}{3}$ for quarks, 0 for leptons). Unit: dimensionless. Status: STANDARD.
- $\tilde{I}_{of}^{(b)}$: The overlap of the lower component of the state o with the zero mode of f, $\tilde{I}_{of}^{(b)} = H^{-1} \int_{-L}^{0} b_o^2 \varrho_f \mathrm{e}^{-3Hy} \mathrm{d}y$

(the overlap of I13); $\tilde{I}_{Q}^{(b)}$, $\tilde{I}_{u_R}^{(b)}$, $\tilde{I}_{d_R}^{(b)}$, $\tilde{I}_{\nu_R}^{(b)}$ those of the dark matter. Unit: dimensionless. Value: dark matter: $\tilde{I}_{Q}^{(b)} = 2.6135$. Status: DERIVED.

- o: The label of a Kohn-Sham orbital (a state of dirac16complex in the cell: block, momentum shell, fold parity and level); a sum over o runs over the orbitals. Status: RECORD.
- k_{cm}: The momentum of each particle of the pair in the centre-of-mass frame. Unit: GeV. Status: STANDARD.

$$\sigma_{\mathrm{I12}} \le \frac{\pi}{4k_{\mathrm{cm}}^2} \ \text{(total angular momentum 0)}, \qquad \sigma_{\mathrm{I13}} \le \frac{3\pi}{4k_{\mathrm{cm}}^2} \ \text{(total angular momentum 1)} \tag{FE-128}$$

Symbols in (FE-128).

- σ_{I12}, σ_{I13}: The spin-averaged annihilation cross-sections of a dark-matter pair through I12 (into gauge-boson pairs) and through I13 (into quark pairs). Unit: GeV^{-2}. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- k_{cm}: The momentum of each particle of the pair in the centre-of-mass frame. Unit: GeV. Status: STANDARD.

The fold bands of dirac16complex made by freeze-in through I11, and their dark radiation:

$$\Gamma_{\mathrm{fi}} = \frac{T}{32\pi^4}\int_0^\infty \mathrm{d}\big(E_{\mathrm{cm}}^2\big)\Big(\sum_{\text{spins}}\sigma\Big)E_{\mathrm{cm}}^3 K_1\Big(\frac{E_{\mathrm{cm}}}{T}\Big) = \frac{4}{\pi^5}\,N_{\mathrm{fold}}\,G_{\mathrm{V}}^2\,T^8\sum_f N_{\mathrm{c}}\big(q_f\tilde{I}_{of}\big)^2, \qquad \frac{\mathrm{d}\big(\rho_{\mathrm{fold}}\,\mathrm{e}^{4a_4}\big)}{\mathrm{d}x_4} = 8T\,\Gamma_{\mathrm{fi}}\,\mathrm{e}^{4a_4} \tag{FE-129}$$

Symbols in (FE-129).

- Γ_{fi}: The rate per volume of the production of fold-band pairs from the plasma through I11 (freeze-in). Unit: GeV^4. Status: DERIVED.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no upper bound. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- E_{cm}: The total energy of the colliding pair in its centre-of-mass frame. Unit: GeV. Status: STANDARD.
- σ: A cross-section (of the dark-matter annihilation or production process named in the equation; summed over the spins where marked). Unit: GeV^{-2}. Status: STANDARD.
- K_1, K_2: The modified Bessel functions of the second kind of orders 1 and 2 (not related to the block variable K). Unit: dimensionless. Status: STANDARD.
- N_{fold}: The number of fold bands: the massless fold zero modes of Ψ_A and of Ψ_B (two Dirac fermions). Unit: dimensionless. Value: 2. Status: DERIVED.
- G_{V}: The four-dimensional strength of the vector portal I11 on the zero modes, $G_{\mathrm{V}} = c_{\mathrm{V}}H/(2f^4 v_{\mathrm{t}}\Lambda_\star^6)$. Unit: GeV^{-2}. Value: benchmark 4.38×10^{-17} GeV^{-2} ($\Delta N_{\mathrm{eff}} = 0.001$). Status: DERIVED.
- f: The label of a fermion species: a Standard-Model multiplet of one generation (its N_{c} Weyl components share one zero-mode profile; the zero mode of each chirality) or a fermion of a medium (electron, proton, neutron); a sum over f runs over these species. Status: STANDARD.
- N_{c}: The number of Weyl components of a fermion multiplet that share its zero-mode profile and overlap (colour times isospin): 6 for a quark doublet, 3 for a quark singlet, 2 for a lepton doublet, 1 for a charged-lepton singlet. Unit: dimensionless. Status: STANDARD.
- q_ψ, q_f: The $B-L$ charge: q_ψ of the multiplet ψ ($q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}$, $q_\ell = q_{e_R} = q_{\nu_R} = -1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- $\tilde{I}_{of}$: The overlap of the state o of dirac16complex with the zero mode of the fermion component f, $\tilde{I}_{of} = H^{-1}\int_{-L}^{0}(a_o^2+b_o^2)\varrho_f \mathrm{e}^{-3Hy}\mathrm{d}y$ (the integral in units of H); $\tilde{I}_Q$, $\tilde{I}_{u_R}$, $\tilde{I}_{d_R}$, $\tilde{I}_\ell$, $\tilde{I}_{e_R}$ those of the dark matter with Q, u_R, d_R, ℓ, e_R, $\tilde{I}_{\nu_R,i}$ with the right-handed neutrino of the generation i. Unit: dimensionless. Value: dark matter: $\tilde{I}_Q = 8.2722$, $\tilde{I}_\ell = 1.2524$, $\tilde{I}_{u_R} = 0.3854$, $\tilde{I}_{d_R} = 0.3961$, $\tilde{I}_{e_R} = 0.3972$. Status: DERIVED.

- o: The label of a Kohn-Sham orbital (a state of dirac16complex in the cell: block, momentum shell, fold parity and level); a sum over o runs over the orbitals. Status: RECORD.
- ρ_{fold}: The energy density of the fold bands. Unit: GeV^4. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.

$$\Delta N_{\mathrm{eff}} = \frac{\rho_{\mathrm{fold}}}{\rho_\nu}, \qquad \rho_\nu = \frac{7}{8}\,\frac{\pi^2}{15}\,T_\nu^4, \qquad \frac{2\Gamma_{\mathrm{fi}}}{n_{\mathrm{fold,eq}}\,\mathcal{H}} \ll 1, \qquad n_{\mathrm{fold,eq}} = N_{\mathrm{fold}}\cdot 4\cdot\frac{3}{4}\,\frac{\zeta(3)}{\pi^2}\,T^3 \tag{FE-130}$$

Symbols in (FE-130).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- N_{eff}: The effective number of neutrino species (the radiation beyond the photons in units of one neutrino species with $T_\nu/T_\gamma = (4/11)^{1/3}$); ΔN_{eff} an addition to it. Unit: dimensionless. Value: 3.0432 (neutrinos) + 0.0010 (fold bands). Status: STANDARD.
- ρ_{fold}: The energy density of the fold bands. Unit: GeV^4. Status: DERIVED.
- ρ_ν, ρ_{ν_α}: The energy density of one neutrino flavour (neutrinos and antineutrinos) at the common temperature T_ν, $\frac{7}{8}\frac{\pi^2}{15}T_\nu^4$ when massless; ρ_{ν_α} that of the flavour α at its temperature T_{ν_α}. Unit: GeV^4. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- T_ν: The temperature of the relic neutrinos (one common temperature). Unit: GeV. Value: 1.9454 K today. Status: STANDARD.
- Γ_{fi}: The rate per volume of the production of fold-band pairs from the plasma through I11 (freeze-in). Unit: GeV^4. Status: DERIVED.
- $n_{\mathrm{fold,eq}}$: The number density the fold bands would have in thermal equilibrium. Unit: GeV^3. Status: DERIVED.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2}T^2/M_{\mathrm{Pl}}$). Unit: GeV. Status: DERIVED.
- N_{fold}: The number of fold bands: the massless fold zero modes of Ψ_A and of Ψ_B (two Dirac fermions). Unit: dimensionless. Value: 2. Status: DERIVED.
- ζ: The Riemann zeta function; only its value $\zeta(3)$ occurs, the sum of $1/n^3$ over the positive integers n. Unit: dimensionless. Value: $\zeta(3) = 1.2020569$. Status: STANDARD.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.

The right-handed neutrinos (the counterfactual that motivates the baryon current), the extra radiation of an early-decoupled species and the production of right-handed neutrinos through I11:

$$\begin{aligned}
&G_{\mathrm{A}}^{B-L} = G_{\mathrm{A}}\Big[\frac{\mathcal{S}_B}{\mathcal{S}_{B-L}}\Big]^{1/2}, \qquad \mathcal{S}_B = \sum_f N_{\mathrm{c}}\big(q_f^B \tilde{I}_{of}^{(b)}\big)^2, \qquad \mathcal{S}_{B-L} = \sum_f N_{\mathrm{c}}\big(q_f \tilde{I}_{of}^{(b)}\big)^2,\\
&\frac{\mathrm{d}n_{\nu_R}}{\mathrm{d}x_4} + 3\mathcal{H}n_{\nu_R} = \Gamma_\nu\Big(1 - \frac{n_{\nu_R}^2}{(n_{\nu_R}^{\mathrm{eq}})^2}\Big), \qquad \Gamma_\nu = 2\,n_{\mathrm{DM,eq}}\,\bar{n}_{\mathrm{DM,eq}}\,\frac{G_\nu^2 M_{\mathrm{DM}}^2}{2\pi}, \qquad G_\nu = G_{\mathrm{A}}^{B-L}\,\tilde{I}_{\nu_R}^{(b)},\\
&n_{\nu_R}^{\mathrm{eq}} = \frac{3}{4}\,\frac{\zeta(3)}{\pi^2}\,T^3, \qquad n_{\mathrm{DM,eq}} = \bar{n}_{\mathrm{DM,eq}} = 2\Big(\frac{M_{\mathrm{DM}}T}{2\pi}\Big)^{3/2}\mathrm{e}^{-x}
\end{aligned} \tag{FE-131}$$

Symbols in (FE-131).

- G_{A}^{B-L}: The strength I13 would need if it coupled to the $B-L$ current (the same relic density). Unit: GeV^{-2}. Status: DERIVED.

- G_{A}: The four-dimensional strength of the portal I13 on the zero modes, $G_{\mathrm{A}} = c_{\mathrm{A}} H/(2 f^4 v_{\mathrm{t}} \Lambda_\star^6)$. Unit: GeV^{-2}. Value: 1.278×10^{-9} GeV^{-2} (dark-matter relic density). Status: DERIVED.
- $\mathcal{S}_B$, $\mathcal{S}_{B-L}$: The colour-weighted sums of the squared I13 overlaps of the dark matter with the baryon-number and with the $B-L$ charges. Unit: dimensionless. Value: $\mathcal{S}_B = 16.351$, $\mathcal{S}_{B-L} = 22.92$. Status: DERIVED.
- f: The label of a fermion species: a Standard-Model multiplet of one generation (its N_{c} Weyl components share one zero-mode profile; the zero mode of each chirality) or a fermion of a medium (electron, proton, neutron); a sum over f runs over these species. Status: STANDARD.
- N_{c}: The number of Weyl components of a fermion multiplet that share its zero-mode profile and overlap (colour times isospin): 6 for a quark doublet, 3 for a quark singlet, 2 for a lepton doublet, 1 for a charged-lepton singlet. Unit: dimensionless. Status: STANDARD.
- q_ψ^B, q_f^B: The baryon number of the multiplet ψ or of the fermion component f ($\frac{1}{3}$ for quarks, 0 for leptons). Unit: dimensionless. Status: STANDARD.
- $\tilde{I}_{of}^{(b)}$: The overlap of the lower component of the state o with the zero mode of f, $\tilde{I}_{of}^{(b)} = H^{-1} \int_{-L}^{0} b_o^2 \varrho_f \mathrm{e}^{-3Hy} \mathrm{d}y$ (the overlap of I13); $\tilde{I}_Q^{(b)}$, $\tilde{I}_{u_R}^{(b)}$, $\tilde{I}_{d_R}^{(b)}$, $\tilde{I}_{\nu_R}^{(b)}$ those of the dark matter. Unit: dimensionless. Value: dark matter: $\tilde{I}_Q^{(b)} = 2.6135$. Status: DERIVED.
- o: The label of a Kohn-Sham orbital (a state of dirac16complex in the cell: block, momentum shell, fold parity and level); a sum over o runs over the orbitals. Status: RECORD.
- q_ψ, q_f: The $B-L$ charge: q_ψ of the multiplet ψ ($q_Q = q_{u_R} = q_{d_R} = \frac{1}{3}$, $q_\ell = q_{e_R} = q_{\nu_R} = -1$), q_f of the fermion component f. Unit: dimensionless. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- n_{ν_R}, $n_{\nu_R}^{\mathrm{eq}}$: The number density of the right-handed neutrinos (three generations) and its equilibrium value. Unit: GeV^3. Status: DERIVED.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2} T^2/M_{\mathrm{Pl}}$). Unit: GeV. Status: DERIVED.
- Γ_ν: The production rate per volume of right-handed neutrinos in the counterfactual $B-L$ variant of I13. Unit: GeV^4. Status: DERIVED.
- $n_{\mathrm{DM,eq}}$, $\bar{n}_{\mathrm{DM,eq}}$: The equilibrium number densities of dark-matter particles and antiparticles of one species. Unit: GeV^3. Status: STANDARD.
- G_ν: The four-dimensional coupling of the dark matter to a right-handed neutrino in the $B-L$ variant of I13. Unit: GeV^{-2}. Status: DERIVED.
- M_{DM}: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- ζ: The Riemann zeta function; only its value $\zeta(3)$ occurs, the sum of $1/n^3$ over the positive integers n. Unit: dimensionless. Value: $\zeta(3) = 1.2020569$. Status: STANDARD.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- x: The ratio $x = M_{\mathrm{DM}}/T$ of the dark-matter mass to the temperature. Unit: dimensionless. Value: $x_{\mathrm{f}} = 31.23$ at the freeze-out. Status: STANDARD.

$$\Gamma_{\mathrm{el}} = 2\big(n_{\mathrm{DM}} + \bar{n}_{\mathrm{DM}}\big)\,\sigma_{\mathrm{el}} > \mathcal{H}, \qquad \sigma_{\mathrm{el}} = \frac{G_\nu^2 E^2}{2\pi}, \qquad E \simeq 3T \tag{FE-132}$$

Symbols in (FE-132).

- Γ_{el}, σ_{el}: The rate of elastic scattering of a right-handed neutrino on the dark matter, and its cross-section. Unit: GeV; GeV^{-2}. Status: ESTIMATE.
- n_{DM}, $\bar{n}_{\mathrm{DM}}$: The number densities of dark-matter particles and antiparticles of one species. Unit: GeV^3. Status: STANDARD.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2} T^2/M_{\mathrm{Pl}}$). Unit: GeV. Status: DERIVED.

- G_ν: The four-dimensional coupling of the dark matter to a right-handed neutrino in the $B-L$ variant of I13. Unit: GeV^{-2}. Status: DERIVED.
- E: The energy of a particle: of the neutrino in the flavour equations, $E=(k^2+m^2)^{1/2}$ of a particle of momentum k in the thermal integrals. Unit: GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- T: The temperature of the plasma (of the photons and of every species in equilibrium with them); in the Mermin occupations the temperature of the Kohn-Sham state. Unit: GeV. Status: STANDARD.

$$\Delta N_{\mathrm{eff}} = N_{\mathrm{W}}\Big(\frac{10.75}{g_s}\Big)^{4/3} F_{\mathrm{h}}, \qquad F_{\mathrm{h}} = \Big[\frac{T_\nu}{T_\gamma}\Big(\frac{11}{4}\Big)^{1/3}\Big/\big(1+h_\nu\big)\Big]^4 = 1.0095 \tag{FE-133}$$

Symbols in (FE-133).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- N_{eff}: The effective number of neutrino species (the radiation beyond the photons in units of one neutrino species with $T_\nu/T_\gamma = (4/11)^{1/3}$); ΔN_{eff} an addition to it. Unit: dimensionless. Value: 3.0432 (neutrinos) + 0.0010 (fold bands). Status: STANDARD.
- N_{W}: The number of Weyl species that decouple early. Unit: dimensionless. Status: DERIVED.
- $g_\rho,\ g_s$: The effective numbers of relativistic degrees of freedom of the energy density and of the entropy density of the plasma. Unit: dimensionless. Value: 105.75 when every species of UM-2 in equilibrium is relativistic (no Higgs particle), 10.75 near 1 MeV, $g_s = 43/11$ today. Status: STANDARD.
- F_{h}: The correction factor of the early-decoupled radiation for the heating of the neutrinos. Unit: dimensionless. Value: 1.0095. Status: DERIVED.
- T_ν: The temperature of the relic neutrinos (one common temperature). Unit: GeV. Value: 1.9454 K today. Status: STANDARD.
- T_γ: The temperature of the photons (equal to the plasma temperature T; the subscript is written where the neutrinos have their own temperature). Unit: GeV. Value: 2.7255 K today. Status: STANDARD.
- h_ν: The relative heating of the neutrinos by the electron-positron annihilation (beyond instantaneous decoupling). Unit: dimensionless. Value: 0.00165. Status: DERIVED.

$$r_{\nu_R} = \frac{G_{\mathrm{V}}^2 \sum_{i=1}^{3} \tilde{I}_{\nu_R,i}^2}{G_{\mathrm{A}}^2\, \mathcal{S}_B} \tag{FE-134}$$

Symbols in (FE-134).

- r_{ν_R}: The ratio of the I11 production of right-handed neutrinos to the I13 annihilation rate of the dark matter. Unit: dimensionless. Value: 2.0×10^{-17}. Status: DERIVED.
- G_{V}: The four-dimensional strength of the vector portal I11 on the zero modes, $G_{\mathrm{V}} = c_{\mathrm{V}} H/(2f^4 v_{\mathrm{t}} \Lambda_\star^6)$. Unit: GeV^{-2}. Value: benchmark 4.38×10^{-17} GeV^{-2} ($\Delta N_{\mathrm{eff}} = 0.001$). Status: DERIVED.
- i: The generation index ($i=1,2,3$), which also labels the neutrino mass states ν_1,ν_2,ν_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- $\tilde{I}_{of}$: The overlap of the state o of dirac16complex with the zero mode of the fermion component f, $\tilde{I}_{of} = H^{-1}\int_{-L}^{0}(a_o^2+b_o^2)\varrho_f \mathrm{e}^{-3Hy}\mathrm{d}y$ (the integral in units of H); $\tilde{I}_Q$, $\tilde{I}_{u_R}$, $\tilde{I}_{d_R}$, $\tilde{I}_\ell$, $\tilde{I}_{e_R}$ those of the dark matter with Q, u_R, d_R, ℓ, e_R, $\tilde{I}_{\nu_R,i}$ with the right-handed neutrino of the generation i. Unit: dimensionless. Value: dark matter: $\tilde{I}_Q = 8.2722$, $\tilde{I}_\ell = 1.2524$, $\tilde{I}_{u_R} = 0.3854$, $\tilde{I}_{d_R} = 0.3961$, $\tilde{I}_{e_R} = 0.3972$. Status: DERIVED.
- G_{A}: The four-dimensional strength of the portal I13 on the zero modes, $G_{\mathrm{A}} = c_{\mathrm{A}} H/(2f^4 v_{\mathrm{t}} \Lambda_\star^6)$. Unit: GeV^{-2}. Value: 1.278×10^{-9} GeV^{-2} (dark-matter relic density). Status: DERIVED.
- $\mathcal{S}_B,\ \mathcal{S}_{B-L}$: The colour-weighted sums of the squared I13 overlaps of the dark matter with the baryon-number and with the $B-L$ charges. Unit: dimensionless. Value: $\mathcal{S}_B = 16.351$, $\mathcal{S}_{B-L} = 22.92$. Status: DERIVED.

The self-interaction channel of the dark matter into the fold bands:

$$\mathcal{L}_4^{\lambda} = -\lambda\,(\bar{\Psi}_{\rm DM}\Psi_{\rm fold})(\bar{\Psi}_{\rm fold}\Psi_{\rm DM})\,(n_{\rm DF}^{\rm sc})^2, \qquad n_{\rm DF}^{\rm sc} = a_{\rm DM}b_{\rm fold} + b_{\rm DM}a_{\rm fold}, \qquad b_{\rm fold} = 0,$$
$$G_\lambda = \frac{\lambda H \tilde{I}_\lambda}{2f^4 v_{\rm t}}, \qquad \tilde{I}_\lambda = \frac{1}{H}\int_{-L}^{0} (n_{\rm DF}^{\rm sc})^2\,{\rm e}^{-3Hy}\,{\rm d}y, \qquad \sigma_\lambda v_{\rm rel} = \frac{G_\lambda^2 M_{\rm DM}^2}{2\pi} \quad \text{(per final fold species)} \tag{FE-135}$$

Symbols in (FE-135).

- $\mathcal{L}_4^{\lambda}$: The four-dimensional term of the contact self-interaction that couples a dark-matter pair to a fold-band pair. Unit: GeV^4. Status: ESTIMATE.
- λ: The contact self-coupling of dirac16complex (the term $-\frac{\lambda}{2}(\bar{\Psi}\Psi)^2$ of the record). Unit: GeV^{-6}. Value: not fixed; Kohn-Sham states computed for $\lambda = 0$ and $\pm 7\times 10^{-5}H^{-6}$; the fold-band radiation requires $|\lambda| \lesssim 1.4\times 10^{-10}H^{-6}$ (for $\Delta N_{\rm eff} \le 0.001$). Status: RECORD.
- $\Psi_{\rm DM}$, $\bar{\Psi}_{\rm DM}$: The four-dimensional field of a dark-matter quantum (the lowest odd level of dirac16complex on the static block, in either half) and its Dirac adjoint. Unit: GeV$^{3/2}$. Status: DERIVED.
- $\Psi_{\rm fold}$, $\bar{\Psi}_{\rm fold}$: The four-dimensional field of a fold-band quantum (an even zero mode of dirac16complex, massless) and its Dirac adjoint. Unit: GeV$^{3/2}$. Status: DERIVED.
- $n_{\rm DF}^{\rm sc}$: The transition scalar density between the dark-matter orbital and a fold-band orbital. Unit: GeV (normalised orbitals). Status: DERIVED.
- $a_{\rm DM}$, $b_{\rm DM}$: The upper and lower components of the dark-matter orbital (block spinor $(a, {\rm i}b)$, normalised on one half). Unit: GeV$^{1/2}$. Status: DERIVED.
- $a_{\rm fold}$, $b_{\rm fold}$: The upper and lower components of a fold-band orbital ($b_{\rm fold} = 0$ for the even zero mode). Unit: GeV$^{1/2}$. Status: DERIVED.
- G_λ: The four-dimensional strength of the self-interaction channel of the dark matter into fold-band pairs. Unit: GeV^{-2}. Value: $\le 2.3\times 10^{-15}$ GeV^{-2} (for $\Delta N_{\rm eff} \le 0.001$). Status: ESTIMATE.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of $m_{\rm W}$ and $\rho_{\rm EW,0}$). Status: AUTHOR.
- $\tilde{I}_\lambda$: The overlap of the squared transition scalar density, $\tilde{I}_\lambda = H^{-1}\int_{-L}^{0}(n_{\rm DF}^{\rm sc})^2{\rm e}^{-3Hy}{\rm d}y$. Unit: dimensionless. Value: 14.38. Status: DERIVED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = {\rm e}^{-a_4}b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- $v_{\rm t}$: The coordinate volume of the compact extra times x_5, x_6, x_7. Unit: GeV^{-3}. Value: not fixed; $v_{\rm t} = H^{-3}$ in the quoted conversions. Status: ASSUMED.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926\times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- e: Euler's number, the base of the natural logarithm; ${\rm e}^X$ is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- y: The hidden coordinate of the y chart, $y = \ln|\sin z|/(6H)$, with $y = 0$ at the fold ($z = \pi/2$) and $y = -L$ at the regular tip; each half of a period of x_0 is the interval $-L \le y \le 0$. Unit: length. Status: DERIVED.
- d: The differential: ${\rm d}X$ is an infinitesimal change of X, ${\rm d}X/{\rm d}Y$ the derivative of X with respect to Y, ${\rm d}^nX$ the volume element of n variables. Status: STANDARD.
- σ_λ: The cross-section of the self-interaction channel (per final fold species). Unit: GeV^{-2}. Status: ESTIMATE.
- $v_{\rm rel}$: The relative velocity of the two annihilating particles; $\langle\sigma v_{\rm rel}\rangle$ the thermal average. Unit: dimensionless (c = 1). Status: STANDARD.
- $M_{\rm DM}$: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

$$\frac{{\rm d}\left(\rho_{\rm fold}\,{\rm e}^{4a_4}\right)}{{\rm d}x_4} = 2M_{\rm DM}\,\Gamma_\lambda\,{\rm e}^{4a_4}, \qquad \Gamma_\lambda = 2\,n_{\rm DM,eq}\,\bar{n}_{\rm DM,eq}\,\sigma_\lambda v_{\rm rel}\,N_{\rm fold}, \qquad \frac{2\Gamma_\lambda}{n_{\rm fold,eq}\,\mathcal{H}} \ll 1 \tag{FE-136}$$

Symbols in (FE-136).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- ρ_{fold}: The energy density of the fold bands. Unit: GeV^4. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- M_{DM}: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- Γ_λ: The rate per volume of the self-interaction channel. Unit: GeV^4. Status: ESTIMATE.
- $n_{\mathrm{DM,eq}}$, $\bar{n}_{\mathrm{DM,eq}}$: The equilibrium number densities of dark-matter particles and antiparticles of one species. Unit: GeV^3. Status: STANDARD.
- σ_λ: The cross-section of the self-interaction channel (per final fold species). Unit: GeV^{-2}. Status: ESTIMATE.
- v_{rel}: The relative velocity of the two annihilating particles; $\langle \sigma v_{\mathrm{rel}} \rangle$ the thermal average. Unit: dimensionless (c = 1). Status: STANDARD.
- N_{fold}: The number of fold bands: the massless fold zero modes of Ψ_A and of Ψ_B (two Dirac fermions). Unit: dimensionless. Value: 2. Status: DERIVED.
- $n_{\mathrm{fold,eq}}$: The number density the fold bands would have in thermal equilibrium. Unit: GeV^3. Status: DERIVED.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H} = \dot{a}_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H} = (8\pi^3 g_\rho/90)^{1/2} T^2/M_{\mathrm{Pl}}$). Unit: GeV. Status: DERIVED.

12.2 The approximation of step 9

- *Ideal gases.* The degrees of freedom are those of ideal gases with the x_0 masses and a modelled QCD crossover. The order-α interaction pressure of the plasma, which step 8 uses for the condensates, is not small at the freeze-out: −5.4 % of the ideal-gas pressure at $T = 1.42$ TeV (QCD −4.7 %, $SU(2)$ −0.7 %, $U(1)_Y$ −0.10 %); since the relic scales as $g_\rho^{1/2}/g_s$ at the freeze-out, including it would raise $\Omega_{\mathrm{DM}} h_{100}^2$ by about 2.7 % and the relic strength G_{A} by about 1.4 % (ESTIMATE, computed in version 4): the physical accuracy of G_{A} is about 1 %, not the 10^{-6} of its root finding.
- *Instantaneous neutrino decoupling* at 2 MeV in the degrees of freedom (it enters only the entropy today, taken as the standard $g_s = 43/11$, and the nucleosynthesis environment of step 8).
- *The dark matter* in kinetic equilibrium with Maxwell-Boltzmann statistics; its own contribution to g_ρ is left out (1.1×10^{-2} of the energy density at the start of the integration, $x = 5$, where $Y_{\mathrm{DM}} = Y_{\mathrm{eq}}$ anyway, and 7×10^{-12} at the freeze-out); the fold bands of the freeze-in are left out of the Hubble rate (3.5×10^{-4} of the energy density at the freeze-out, which moves the relic by 1.7×10^{-4}).
- *The unitarity completion.* Where an effective cross-section exceeds its partial-wave bound it is replaced by the bound (a completion that saturates it); for the I13 relic 21.7 % of the freeze-out rate comes from collisions at the unitarity bound of the ${}^3S_1$ partial wave.
- *The freeze-in* with Maxwell-Boltzmann statistics and without inverse processes (the production rate per fold quantum is 0.012 of the Hubble rate at the highest temperature $T_{\max} = 10$ TeV, ASSUMED; the result grows like $T_{\max}^3$).
- *Estimates.* The elastic rate of the right-handed neutrinos (a factor 1/2 or 2 moves their kinetic decoupling to $M_{\mathrm{DM}}/T = 23.77$ or 25.07 and ΔN_{eff} by at most 9.5×10^{-6} of itself) and the cross-section of the self-interaction channel (its Fierz coefficients are not computed; at the bound the inverse annihilation of the fold quanta is 0.027 of the Hubble rate).

12.3 The solution of step 9

12.3.1 The thermal history

The species in equilibrium (photon, gluons, W^+, W^-, Z, six quarks, three charged leptons, three neutrinos; no Higgs particle; not the right-handed neutrinos, the fold bands, the radion or the Kaluza-Klein modes; the dark matter is not counted, since it would add less than 0.1 at 5 TeV and nothing at its freeze-out) with their x_0 masses give the degrees of freedom of the field equations written above, with the QCD crossover modelled by the weight w_{QGP} of the quark-gluon plasma against a gas of the twelve lightest hadron multiplets (ASSUMED; masses in GeV

and internal states: pion 0.1380, 3; kaon 0.4956, 4; eta 0.5479, 1; rho meson 0.7753, 9; omega meson 0.7827, 3; K-star 0.8917, 12; nucleon 0.9389, 8; eta-prime 0.9578, 1; phi meson 1.0195, 3; Lambda baryon 1.1157, 4; Sigma baryon 1.1932, 12; Delta baryon 1.232, 32; quoted without a checked reference) and the neutrinos decoupled at 2 MeV (DERIVED):

T	5 TeV	1.42 TeV	200 GeV	10 GeV	1 GeV	300 MeV	155 MeV	100 MeV	3 MeV	1 MeV	0.1 MeV	today
g_ρ	105.75	105.73	104.95	86.22	76.34	64.57	42.35	17.54	10.74	10.56	4.30	3.363
g_s	105.75	105.73	104.68	86.13	75.50	64.14	41.76	17.16	10.74	10.56	4.78	3.909

i.e. 105.75 above the top threshold (three generations without a Higgs particle), 86.25 without t, W, Z, 10.75 before $e^\pm$ annihilation, and today $g_s = 43/11 = 3.9091$, $g_\rho = 3.3626$ and the entropy density $s_0 = 2891.3$ cm^{-3} ($T_\gamma = 2.7255$ K).

12.3.2 The Boltzmann equation

The Boltzmann equation of one Dirac species is integrated implicitly (Radau, $\ln Y_{\rm DM}$ from $x = 5$ to 2000, relative tolerance 10^{-10}) with the thermal average by adaptive quadrature in the kinetic energy of the pair in units of its rest energy, $E^2_{\rm cm}/(4M^2_{\rm DM}) - 1$, with exponentially scaled Bessel functions, tabulated on 260 points in x. Validation (DERIVED): an s-wave cross-section of 4.4×10^{-26} cm^3/s gives $\Omega_{\rm DM}h^2_{100} = 0.1275$ (the freeze-out formula 0.1306), and the late-time integral $1/Y_\infty - 1/Y_{\rm DM}(x)$ equals the integral of the collision term from x on to 1.3×10^{-8}; for I12 the exact average exceeds the non-relativistic $6G^2_{\rm I12}M^4_{\rm DM}/(\pi x)$ by 0.408, 0.111, 0.031, 0.0030 at $x = 10$, 30, 100, 1000, i.e. by about $3/x$ at large x.

12.3.3 I12 alone, and partial-wave unitarity

The effective operator would need $G_{\rm I12} = 2.509 \times 10^{-13}$ GeV^{-3} = (15.86 TeV)$^{-3}$ for $\Omega_{\rm DM}h^2_{100} = 0.060$ per species ($x_{\rm f} = 31.70$, $\langle\sigma v_{\rm rel}\rangle(x_{\rm f}) = 1.89 \times 10^{-25}$ cm^3/s; step 7: (15.5 TeV)$^{-3}$ from the freeze-out formula). Its cross-section exceeds the $^3P_0$ bound above $E_{\rm cm} = 89510$ GeV (815 GeV above threshold), i.e. above the relative velocity $v_{\rm rel} = 2\beta_{\rm cm} = 0.269$ (0.274 from the non-relativistic formula); at the freeze-out 96 % of the annihilation rate comes from collisions above the crossing, and 75 % of the pairs of the non-relativistic Maxwell distribution are faster than 0.274. With $\sigma = \pi/(4k^2_{\rm cm})$ at every energy, the largest annihilation from $^3P_0$, each species leaves $\Omega_{\rm DM}h^2_{100} = 0.0582$ (at its freeze-out, $x_{\rm f} = 30.77$, $\langle\sigma v_{\rm rel}\rangle = 5.33 \times 10^{-26}$ cm^3/s against the non-relativistic $\sqrt{\pi x}/M^2_{\rm DM} = 5.84 \times 10^{-26}$ cm^3/s at the same x); the effective operator capped at the bound leaves 0.1544, 0.1081, 0.0836, 0.0649, 0.0585 for $G^2_{\rm I12}/G^2_{\rm relic} = 1$, 3, 10, 100, 10^4 ($G_{\rm relic}$ the relic strength above). Within perturbative couplings I12 therefore overproduces the dark matter; only a strongly coupled $^3P_0$ channel at the unitarity limit could give the relic. UM-2 sets I12 to the unitarity-safe benchmark

$$G_{\rm I12} = \frac{\pi}{2M^3_{\rm DM}} = 1.801\times10^{-14}\ \text{GeV}^{-3} = (38.2\ \text{TeV})^{-3}, \qquad \frac{c_{\rm I12}}{\Lambda^3_\star} = \frac{G_{\rm I12}}{\mathcal{O}_{\rm DM}\sqrt{8g_3^4 + 3g_2^4 + g_1^4}} = 1.106\times10^{-14}\ \text{GeV}^{-3},$$

Symbols in this equation.

- $G_{\rm I12}$: The combined coupling of I12 to all gauge-boson pairs, $G^2_{\rm I12} = 8G^2_{\rm g} + 3G^2_{\rm W} + G^2_{\rm B}$. Unit: GeV^{-3}. Value: benchmark $\pi/(2M^3_{\rm DM}) = 1.801 \times 10^{-14}$ GeV^{-3}. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $M_{\rm DM}$: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- $c_{\rm I12}$: The common coefficient of the three terms of I12 when they are universal ($c_{\rm G} = c_{\rm W} = c_{\rm B} = c_{\rm I12}$). Unit: dimensionless. Value: $c_{\rm I12}/\Lambda^3_\star = 1.106 \times 10^{-14}$ GeV^{-3} (unitarity-safe benchmark). Status: INVENTED.
- $\Lambda_\star$: The scale of the portal operators I11, I12, I13. Unit: GeV. Value: only the combinations $c/\Lambda^n_\star$ enter. Status: INVENTED.

- $\mathcal{O}_o$, $\mathcal{O}_{\rm DM}$: The scalar overlap of a state of dirac16complex, $\mathcal{O}_o = \partial\varepsilon_o/\partial m_\Psi$ (Hellmann-Feynman: the integrated scalar density of the orbital); $\mathcal{O}_{\rm DM}$ that of the dark matter. Unit: dimensionless. Value: $\mathcal{O}_{\rm DM} = 0.7072$. Status: DERIVED.
- g_3: The four-dimensional strong coupling ($g_3^2 = 4\pi\alpha_3$). Unit: dimensionless. Value: $g_3^2 = 0.7786$ at $2M_{\rm DM}$. Status: DERIVED.
- g_2: The four-dimensional SU(2) coupling ($g_2^2 = 4\pi\alpha_2$). Unit: dimensionless. Value: $g_2^2 = 0.3781$ at $2M_{\rm DM}$. Status: DERIVED.
- g_1: The four-dimensional hypercharge coupling ($g_1^2 = 4\pi\alpha_1$, Standard-Model normalisation without the GUT factor). Unit: dimensionless. Value: $g_1^2 = 0.1380$ at $2M_{\rm DM}$. Status: DERIVED.

below the bound up to $E_{\rm cm} = 106363$ GeV $= 2.40\,M_{\rm DM}$ (a fraction 5.2×10^{-4} of the annihilation rate at $x = 30$ comes from above), with universal coefficients $c_{\rm G} = c_{\rm W} = c_{\rm B} = c_{\rm I12}$ and the one-loop couplings at $2M_{\rm DM} = 88.7$ TeV ($1/\alpha_3 = 1/\alpha_3(m_{\rm Z}) + \frac{7}{2\pi}\ln(2M_{\rm DM}/m_{\rm Z})$ and the running electroweak couplings of step 8 at $\mu_{\rm ren} = 2M_{\rm DM}$, from $\sin^2\theta_{\overline{\rm MS}} = 0.23122$ and $\alpha_{\rm em}(m_{\rm Z}) = 1/127.95$: $g_3^2 = 0.7786$, $g_2^2 = 0.3781$, $g_1^2 = 0.1380$), i.e. $\Lambda_\star/c_{\rm I12}^{1/3} = 44.9$ TeV. Alone it would leave $\Omega_{\rm DM}h_{100}^2 = 8.0$ per species.

12.3.4 The relic with I13

With $G_f = G_{\rm A}\,q_f^B\,\tilde{I}_{of}^{(b)}$ for each quark Weyl component ($\mathcal{S}_B = \sum_f N_{\rm c}(q_f^B\tilde{I}_{of}^{(b)})^2 = 16.351$ for the dark-matter orbital) the s-wave cross-section of I13 and its ${}^3S_1$ unitarity bound are those written above, and the Boltzmann equation with $\sigma_{\rm I12} + \sigma_{\rm I13}$ gives each species $\Omega_{\rm DM}h_{100}^2 = 0.060$ for (DERIVED)

$$G_{\rm A} = 1.278 \times 10^{-9}\ {\rm GeV}^{-2} = 1.1 \times 10^{-4}\,G_{\rm F} \qquad (x_{\rm f} = 31.23),$$

Symbols in this equation.

- $G_{\rm A}$: The four-dimensional strength of the portal I13 on the zero modes, $G_{\rm A} = c_{\rm A}H/(2f^4 v_{\rm t}\Lambda_\star^6)$. Unit: ${\rm GeV}^{-2}$. Value: 1.278×10^{-9} ${\rm GeV}^{-2}$ (dark-matter relic density). Status: DERIVED.
- $G_{\rm F}$: The Fermi constant. Unit: ${\rm GeV}^{-2}$. Value: 1.1663788×10^{-5} ${\rm GeV}^{-2}$ (measured). Status: STANDARD.
- $x_{\rm f}$: The freeze-out point $x_{\rm f} = M_{\rm DM}/T$ at which the dark matter leaves chemical equilibrium. Unit: dimensionless. Value: 30.38 (freeze-out formula, I12), 31.23 (Boltzmann equation, I12 + I13). Status: DERIVED.

with $\langle\sigma v_{\rm rel}\rangle(x_{\rm f}) = 9.50 \times 10^{-26}$ ${\rm cm}^3/{\rm s}$ from I13 and 9.9×10^{-28} ${\rm cm}^3/{\rm s}$ from I12, the I13 cross-section capped at its bound, which it reaches only above $v_{\rm rel} = 0.51$ (22 % of the freeze-out rate then comes from collisions at the bound; without the cap $G_{\rm A} = 1.269 \times 10^{-9}$ ${\rm GeV}^{-2}$, and the uncapped cross-section exceeds the bound above $v_{\rm rel} = 0.52$, for 25 % of the rate). The relic of UM-2 is a thermal freeze-out with most annihilations (78 % of the rate) below the unitarity bound; the contact form of I13 is an effective description (for the quark doublet $(G_{\rm A}q_Q^B\tilde{I}_Q^{(b)})^{-1/2} = 30$ TeV, below the energies $E_{\rm cm} \approx 89$ TeV of the annihilations), whose completion by a mediator is OPEN.

Why the baryon current. With the $B - L$ current instead ($\mathcal{S}_{B-L} = 22.92$ over the quarks, the leptons and the ν_R, so the relic needs $G_{\rm A}^{B-L} = G_{\rm A}(16.351/22.92)^{1/2} = 1.072 \times 10^{-9}$ ${\rm GeV}^{-2}$ with the uncapped $G_{\rm A}$; the three ν_R carry 1.2 % of the annihilation) the right-handed neutrinos would be thermalised: per flavour, with $G_\nu = G_{\rm A}^{B-L}\tilde{I}_{\nu_R}^{(b)}$ (one representative flavour, the rms of the three overlaps, which differ by 2 %), from zero abundance at $M_{\rm DM}/T = 2$ they reach 0.9997 of the equilibrium number by $M_{\rm DM}/T = 40$; the number-changing rate falls below the Hubble rate at $M_{\rm DM}/T = 16.1$, and the elastic scattering on the dark matter (ESTIMATE) at $M_{\rm DM}/T = 24.4$ ($T = 1816$ GeV, $g_s = 105.7$); the three ν_R would add $\Delta N_{\rm eff} = 3(10.75/g_s)^{4/3}F_{\rm h} = 0.144$ ($F_{\rm h} = 1.0095$), $N_{\rm eff} = 3.187$, 2.5 standard deviations above ACT DR6 (2.86 ± 0.13). With J^B the ν_R keep their tip coupling and the I11 coupling of charge -1, through which the dark matter annihilates into ν_R pairs at only $r_{\nu_R} = 2.0 \times 10^{-17}$ of the I13 rate into quarks (DERIVED): they are never thermalised.

12.3.5 The fold bands: freeze-in through I11

With $G_f = G_{\rm V}q_f\tilde{I}_{of}$ for the fold band o, the pair production $f\bar{f} \to \Psi_{\rm fold}\bar{\Psi}_{\rm fold}$ has the cross-section $G_f^2E_{\rm cm}^2/(12\pi)$ per Dirac f (spin averaged; half for one chirality), and the rate density and the energy injection are those written above (Maxwell-Boltzmann; $\int_0^\infty x^6K_1\,{\rm d}x = 384$, $\int_0^\infty x^7K_2\,{\rm d}x = 3072$, both checked), with $N_{\rm fold} = 2$ and

$\sum_f N_c (q_f \tilde{I}_{of})^2 = 126.85$ over the Weyl components of three generations (no ν_R in the plasma). Integrated from T_{max} (DERIVED): $\Delta N_{eff} = 5.21 \times 10^{29}\, G_V^2$ GeV^4 for $T_{max} = 10$ TeV, so the benchmark $\Delta N_{eff} = 0.001$ means

$$G_V = 4.38 \times 10^{-17}\ \text{GeV}^{-2};$$

Symbols in this equation.

- G_V: The four-dimensional strength of the vector portal I11 on the zero modes, $G_V = c_V H/(2 f^4 v_t \Lambda_\star^6)$. Unit: GeV^{-2}. Value: benchmark 4.38×10^{-17} GeV^{-2} ($\Delta N_{eff} = 0.001$). Status: DERIVED.

for that G_V, $\Delta N_{eff} = 8.0 \times 10^{-6}$, 1.25×10^{-4}, 1.0×10^{-3}, 8.0×10^{-3} at $T_{max} = 2$, 5, 10, 20 TeV, and at T_{max} the production rate per fold quantum is 0.012 of the Hubble rate: never in equilibrium.

12.3.6 The self-interaction channel

The contact self-interaction $-\frac{\lambda}{2}(\bar{\Psi}\Psi)^2$ contains the term written above, with the transition scalar density n_{DF}^{sc} of the dark-matter orbital and the fold zero mode, which is odd under the fold reflection while its square is even, so the dark matter can annihilate into the fold bands; $\tilde{I}_\lambda = 14.38$ (units of H; DERIVED), and the s-wave cross-section is estimated as the value of one vector channel per final fold species (ESTIMATE). While the dark matter is in equilibrium ($T_{max} = 10$ TeV to $x_f = 31.23$) each annihilation puts $2M_{DM}$ into the fold bands, which gives (DERIVED) $\Delta N_{eff} = 1.83 \times 10^{26}\, G_\lambda^2$ GeV^4 as long as the fold bands stay out of equilibrium. $\Delta N_{eff} \leq 0.001$ therefore requires

$$G_\lambda \leq 2.3 \times 10^{-15}\ \text{GeV}^{-2} \quad (1.8 \times 10^{-6}\, G_A),$$

Symbols in this equation.

- G_λ: The four-dimensional strength of the self-interaction channel of the dark matter into fold-band pairs. Unit: GeV^{-2}. Value: $\leq 2.3 \times 10^{-15}$ GeV^{-2} (for $\Delta N_{eff} \leq 0.001$). Status: ESTIMATE.
- G_A: The four-dimensional strength of the portal I13 on the zero modes, $G_A = c_A H/(2 f^4 v_t \Lambda_\star^6)$. Unit: GeV^{-2}. Value: 1.278×10^{-9} GeV^{-2} (dark-matter relic density). Status: DERIVED.

at which the production rate per fold quantum is at most 0.027 of the Hubble rate. The self-coupling $\lambda_{test} = 7 \times 10^{-5}\, H^{-6}$ of the Kohn-Sham tests of step 3 would, with $v_t = H^{-3}$, give $G_\lambda = 1.14 \times 10^{-9}$ GeV^{-2}, keep the fold bands in equilibrium down to $T = 2.6$ TeV (production rate up to 6.3×10^9 times the Hubble rate) and add $\Delta N_{eff} = 4(10.75/g_s)^{4/3} F_h = 0.19$ (four Weyl species, $N_{eff} = 3.23$). UM-2 therefore ASSUMES $G_\lambda \leq 2.3 \times 10^{-15}$ GeV^{-2}, i.e. $\lambda \leq 1.4 \times 10^{-10}\, H^{-6}$ for $v_t = H^{-3}$, or $v_t \geq 4.9 \times 10^5\, H^{-3}$ for $\lambda = \lambda_{test}$; with such a λ the self-interaction mean fields of every Kohn-Sham state of the cosmology are negligible, and λ_{test} is only the value with which the Kohn-Sham machinery was tested.

12.3.7 Detection

Per nucleon, with the couplings of the dark matter to the up and down quarks (I13 and I11, unpolarised quarks with each chirality half),

$$G_u = \frac{G_A}{3}\,\frac{\tilde{I}_Q^{(b)} + \tilde{I}_{u_R}^{(b)}}{2} + \frac{G_V}{3}\,\frac{\tilde{I}_Q + \tilde{I}_{u_R}}{2}, \qquad G_d = \frac{G_A}{3}\,\frac{\tilde{I}_Q^{(b)} + \tilde{I}_{d_R}^{(b)}}{2} + \frac{G_V}{3}\,\frac{\tilde{I}_Q + \tilde{I}_{d_R}}{2},$$

Symbols in this equation.

- G_u, G_d: The four-dimensional vector couplings of the dark matter to the up and to the down quark (I13 and I11, unpolarised quarks with each chirality carrying half of the density). Unit: GeV^{-2}. Status: DERIVED.
- G_A: The four-dimensional strength of the portal I13 on the zero modes, $G_A = c_A H/(2 f^4 v_t \Lambda_\star^6)$. Unit: GeV^{-2}. Value: 1.278×10^{-9} GeV^{-2} (dark-matter relic density). Status: DERIVED.
- $\tilde{I}_{of}^{(b)}$: The overlap of the lower component of the state o with the zero mode of f, $\tilde{I}_{of}^{(b)} = H^{-1} \int_{-L}^{0} b_o^2 \varrho_f e^{-3Hy} dy$ (the overlap of I13); $\tilde{I}_Q^{(b)}$, $\tilde{I}_{u_R}^{(b)}$, $\tilde{I}_{d_R}^{(b)}$, $\tilde{I}_{\nu_R}^{(b)}$ those of the dark matter. Unit: dimensionless. Value: dark matter: $\tilde{I}_Q^{(b)} = 2.6135$. Status: DERIVED.

- G_{V}: The four-dimensional strength of the vector portal I11 on the zero modes, $G_{\mathrm{V}} = c_{\mathrm{V}} H/(2 f^4 v_{\mathrm{t}} \Lambda_\star^6)$. Unit: GeV^{-2}. Value: benchmark 4.38×10^{-17} GeV^{-2} ($\Delta N_{\mathrm{eff}} = 0.001$). Status: DERIVED.
- $\tilde{I}_{of}$: The overlap of the state o of dirac16complex with the zero mode of the fermion component f, $\tilde{I}_{of} = H^{-1} \int_{-L}^{0} (a_o^2 + b_o^2) \varrho_f \mathrm{e}^{-3Hy} \mathrm{d}y$ (the integral in units of H); $\tilde{I}_Q$, $\tilde{I}_{u_R}$, $\tilde{I}_{d_R}$, $\tilde{I}_\ell$, $\tilde{I}_{e_R}$ those of the dark matter with Q, u_R, d_R, ℓ, e_R, $\tilde{I}_{\nu_R,i}$ with the right-handed neutrino of the generation i. Unit: dimensionless. Value: dark matter: $\tilde{I}_Q = 8.2722$, $\tilde{I}_\ell = 1.2524$, $\tilde{I}_{u_R} = 0.3854$, $\tilde{I}_{d_R} = 0.3961$, $\tilde{I}_{e_R} = 0.3972$. Status: DERIVED.

$$f_{\mathrm{p}} = 2G_{\mathrm{u}} + G_{\mathrm{d}}, \qquad f_{\mathrm{n}} = G_{\mathrm{u}} + 2G_{\mathrm{d}}, \qquad f_{\mathrm{N}} = \frac{Z_{\mathrm{Xe}} f_{\mathrm{p}} + (A_{\mathrm{Xe}} - Z_{\mathrm{Xe}}) f_{\mathrm{n}}}{A_{\mathrm{Xe}}}, \qquad \sigma_{\mathrm{SI}} = \frac{\mu_{\mathrm{red}}^2 f_{\mathrm{N}}^2}{\pi}$$

Symbols in this equation.

- f_{N}, f_{p}, f_{n}: The spin-independent effective couplings of the dark matter to a nucleon: f_{p} to the proton, f_{n} to the neutron, and f_{N} the average per nucleon of the target (for the portal I12, the coupling through the gluon content of a nucleon). Unit: GeV^{-2}. Value: $f_{\mathrm{p}} = 1.915 \times 10^{-9}$, $f_{\mathrm{n}} = 1.917 \times 10^{-9}$ GeV^{-2} (I13 and I11). Status: ESTIMATE.
- G_{u}, G_{d}: The four-dimensional vector couplings of the dark matter to the up and to the down quark (I13 and I11, unpolarised quarks with each chirality carrying half of the density). Unit: GeV^{-2}. Status: DERIVED.
- Z_{Xe}, A_{Xe}: The atomic number and the mass number of the xenon target. Unit: dimensionless. Value: 54 and 131. Status: STANDARD.
- σ_{SI}: The spin-independent elastic scattering cross-section of a dark-matter particle on one nucleon. Unit: cm^2. Value: 4.01×10^{-46} cm^2 (xenon, I13 and I11); 6.3×10^{-53} cm^2 through I12 alone. Status: ESTIMATE.
- μ_{red}: The reduced mass of the dark-matter particle and a nucleon, $M_{\mathrm{DM}} m_{\mathrm{N}}/(M_{\mathrm{DM}} + m_{\mathrm{N}})$. Unit: GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

(xenon, $Z_{\mathrm{Xe}} = 54$, $A_{\mathrm{Xe}} = 131$): $f_{\mathrm{p}} = 1.915 \times 10^{-9}$, $f_{\mathrm{n}} = 1.917 \times 10^{-9}$ GeV^{-2}, $\sigma_{\mathrm{SI}} = 4.01 \times 10^{-46}$ cm^2, 0.16 of the LZ limit linearly extrapolated to 44.35 TeV (2.44×10^{-45} cm^2; ESTIMATE): allowed and within reach of the next generation of xenon detectors (I11 alone: 3.9×10^{-60} cm^2). The s-wave annihilation into quark pairs continues today with

$$\sigma_{\mathrm{I13}}\, v_{\mathrm{rel}}\big|_{v_{\mathrm{rel}} \to 0} = \sum_f \frac{N_{\mathrm{c}}\, G_f^2\, M_{\mathrm{DM}}^2}{2\pi} = 9.76 \times 10^{-26}\ \mathrm{cm}^3/\mathrm{s}$$

Symbols in this equation.

- σ_{I12}, σ_{I13}: The spin-averaged annihilation cross-sections of a dark-matter pair through I12 (into gauge-boson pairs) and through I13 (into quark pairs). Unit: GeV^{-2}. Status: DERIVED.
- v_{rel}: The relative velocity of the two annihilating particles; $\langle \sigma v_{\mathrm{rel}} \rangle$ the thermal average. Unit: dimensionless (c = 1). Status: STANDARD.
- f: The label of a fermion species: a Standard-Model multiplet of one generation (its N_{c} Weyl components share one zero-mode profile; the zero mode of each chirality) or a fermion of a medium (electron, proton, neutron); a sum over f runs over these species. Status: STANDARD.
- N_{c}: The number of Weyl components of a fermion multiplet that share its zero-mode profile and overlap (colour times isospin): 6 for a quark doublet, 3 for a quark singlet, 2 for a lepton doublet, 1 for a charged-lepton singlet. Unit: dimensionless. Status: STANDARD.
- G_f: The I13 coupling of the dark matter to the quark component f, $G_f = G_{\mathrm{A}} q_f^B \tilde{I}_{of}^{(b)}$. Unit: GeV^{-2}. Status: DERIVED.
- M_{DM}: The mass of a dark-matter particle (the lowest odd level of dirac16complex). Unit: GeV. Value: 44.35 TeV. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

(DERIVED, 7 + 16 checks).




Figure 7: The thermal history of UM-2 (degrees of freedom with the x_0 masses, left) and the freeze-out of one dark-matter species through the effective I12 at the relic strength and at the ${}^3P_0$ unitarity limit (right).

12.4 The check of step 9

In its own equations.

- *The degrees of freedom.* g_ρ and g_s at the 13 tabulated temperatures, recomputed with adaptive quadrature, agree with the stored values to 4.6×10^{-11}; today $g_s = 43/11$ and $g_\rho = 2 + \frac{7}{8} \cdot 6\,(4/11)^{4/3}$ hold to 4.3×10^{-9} and 5.0×10^{-9} (DERIVED).
- *The relic.* The Boltzmann equation re-solved at the stored G_A gives $\Omega_\mathrm{DM}h^2_{100} = 0.0600000002$ per species (3.1×10^{-9} from the condition); the late-time identity holds to 6.1×10^{-9} on the recomputed I13 history and to 3.3×10^{-9} on the stored I12 history; the freeze-out point is reproduced to 1.3×10^{-12}, the s-wave validation to 1.3×10^{-8} (DERIVED).
- *The freeze-in.* ΔN_eff per G_V^2, re-integrated with Gauss-Legendre panels, agrees with the stored trapezoid value to 3.9×10^{-5} (the trapezoid error on 600 logarithmic points); the Bessel moments are exact and the benchmark gives 0.001 to 5×10^{-13} (DERIVED).
- *The right-handed neutrinos.* Re-solved (Radau in the variable $M_\mathrm{DM}/T - 2$, relative tolerance 10^{-13}), the ν_R equation gives the number density 0.999697681992 of its equilibrium value at $M_\mathrm{DM}/T = 40$, the stored value to 2.6×10^{-13}, unchanged when the coupling is varied by $\pm 2 \times 10^{-12}$; the kinetic decoupling, $\Delta N_\mathrm{eff} = 0.144$, G_A^{B-L} and r_{ν_R} are reproduced to 2.5×10^{-12} (DERIVED; the integration of version 3 depended on the twelfth digit of its input coupling, the result did not).
- *The self-interaction channel.* ΔN_eff per G_λ^2 re-integrated agrees to 1.1×10^{-5}, and ΔN_eff at the bound is 0.001 to 4×10^{-12} (DERIVED).

In concert with the other field equations.

- *The masses of step 2* are the masses of the ideal gases to 3.1×10^{-12}, and the dark-matter mass of the x_0 spectrum is that of the Boltzmann equation to 8.1×10^{-13} (DERIVED).
- *The four-dimensional Friedmann equation of step 1.* The Hubble rate of the Boltzmann equation at the freeze-out is that of the four-dimensional Friedmann equation with all species to 1.7×10^{-4} (the fold bands of the freeze-in, left out here) (DERIVED).
- *The CMB of step 6.* The relic 2×0.060 is the cold dark matter of the CMB computation to 3.1×10^{-9}, and the freeze-in radiation is its $\Delta N_\mathrm{eff} = 0.0010$ exactly; the radiation today of the instantaneous decoupling differs from that of the CMB computation by 0.59 % (DERIVED).
- *The coupled solution of step 8.* G_A and G_V are the strengths of the mean fields to 6.7×10^{-13}, the I12 benchmark to 1.6×10^{-12}, and the freeze-out temperature to 2.1×10^{-12}; the plasma energy density at the freeze-out differs from the block sources by 1.3×10^{-3}, by design (the trace anomaly of the interacting plasma) (DERIVED).
- *The neutrino decoupling of step 5.* The heating factor F_h is that of the decoupling to 5.8×10^{-7} (DERIVED).

12.5 The status of step 9

DERIVED: the degrees of freedom with the x_0 masses, the Boltzmann solutions, the violation of unitarity by I12 alone and the unitarity-safe benchmark, the I13 relic strength, the freeze-in benchmark of I11, the thermalisation of the ν_R in the $B-L$ alternative, the bound on the self-interaction channel, the detection rates. ASSUMED: the model of the QCD crossover, $T_{\max} = 10$ TeV, the bound on λ. ESTIMATE: the LZ extrapolation, the elastic ν_R rate, the self-interaction cross-section, the effect of the order-α plasma pressure on the relic. INVENTED: the portals. OPEN: the completion of the I13 contact interaction by a mediator above about 30 TeV.

13 Step 10: the terms of the metric power series

The author asks for a metric general enough to model the CMB, the CNB and the BAO, expressible as a power series of terms of decreasing magnitude. Step 10 collects the magnitude of every term of the series from the solution of steps 1 to 9.

13.1 The field equations of step 10

The Einstein-Lovelock equations, whose solution the series expands:

$$\begin{aligned} &\sum_{p=1}^{3} \alpha_{(p)}\, {\mathcal{E}_{(p)}}^{M}{}_{N} + \Lambda\, \delta^{M}_{N} = \kappa\, {\mathcal{T}}^{M}{}_{N}, \\ &{\mathcal{E}_{(p)}}^{M}{}_{N} = -\frac{1}{2^{p+1}}\, \delta^{M M_1 N_1 \cdots M_p N_p}_{N P_1 Q_1 \cdots P_p Q_p}\, {R^{P_1 Q_1}}_{M_1 N_1} \cdots {R^{P_p Q_p}}_{M_p N_p} \end{aligned} \qquad \text{(FE-1)}$$

Symbols in (FE-1).

- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic). Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)} H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- ${\mathcal{E}_{(p)}}^{M}{}_{N}$: The Lovelock tensor of order p, the metric variation of $\sqrt{|g|}\mathcal{L}_{(p)}$: $\mathcal{E}_{(1)}$ is the Einstein tensor ${R^{M}}_{N} - \frac{1}{2}\delta^{M}_{N} R$; every $\mathcal{E}_{(p)}$ is divergence free. Unit: 1/length^{2p}. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -10H^2$. Unit: GeV^2. Value: $-10H^2$. Status: AUTHOR.
- δ: The Kronecker delta: with two indices $\delta^{M}_{N} = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{\alpha e}$, δ_{l0}, δ^{I3}); with upper and lower index strings the generalised Kronecker delta $\delta^{M_1 \cdots M_n}_{N_1 \cdots N_n}$, the determinant of the $n \times n$ matrix of Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa / V_{\perp} = 8\pi G_{\mathrm{N}} \mathcal{G}(K_*) = 2.328 \times 10^{-36}$ GeV^{-2}. Status: AUTHOR.
- ${\mathcal{T}}^{M}{}_{N}$: The energy-momentum tensor of all matter (every field except gravity, the stabiliser included), the source of gravity; defined by the change of the matter action under a change of the metric. Unit: GeV^8. Status: STANDARD.
- P: A coordinate index of eight dimensions (also $P_1, P_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Q: A coordinate index of eight dimensions (also $Q_1, Q_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- ${R^{MN}}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor, with two indices the Ricci tensor $R_{MN} = {R^{P}}_{MPN}$, without indices the Ricci (scalar) curvature $R = g^{MN} R_{MN}$; ${R^{ab}}_{ab}$ (no sum) is the sectional curvature of the coordinate plane (a, b). Unit: 1/length^2. Status: STANDARD.

The series and its order-1, $n = 0$ term:

$$g_{MN} = \bar{g}_{MN} + \epsilon\, h^{(1)}_{MN} + \epsilon^2\, h^{(2)}_{MN} + \ldots, \qquad h^{(1)}_{MN}(X^\mu, x_0) = \sum_{n\geq 0} h^{(1,n)}_{MN}(X^\mu)\, F_n(x_0), \qquad \epsilon \sim 10^{-5} \tag{FE-87}$$

Symbols in (FE-87).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, ${g^M}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\bar{g}_{MN}$: The order-0 (background) metric of the power series: the corrected primordial metric in the author's coordinates, determined by the two functions $a_4(x_4)$ and $b_4(x_4)$; $|\bar{g}|$ is the absolute value of its determinant. Unit: dimensionless. Status: AUTHOR.
- ϵ: The bookkeeping parameter of the power series of the metric: the term of order q is multiplied by ϵ^q (the size of the first-order potentials). Unit: dimensionless. Value: $\epsilon \sim 10^{-5}$. Status: DERIVED.
- $h^{(1)}_{MN}$, $h^{(2)}_{MN}$: The first- and second-order terms of the power series of the metric. Unit: dimensionless. Status: DERIVED.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 (signature $+,+,+,-$); on the tip it runs over the tip directions; a repeated index is summed. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- n: The level number of the Kaluza-Klein tower ($n = 1, 2, 3, \ldots$; $n = 0$ the flat zero mode). Status: DERIVED.
- $h^{(1,n)}_{MN}$: The four-dimensional coefficient of the n-th x_0 profile F_n in the first-order term ($n = 0$: the flat zero mode, the perturbations of four-dimensional gravity; $n \geq 1$: the Kaluza-Klein gravitons). Unit: dimensionless. Status: DERIVED.
- $F(\xi)$, F_n: The x_0 profile of a boson mode on one half of the period (a function of y or of $\xi = y/L$): of a gauge field (gauge $A_y = 0$) or of the transverse-traceless graviton; F_n is the profile of the n-th Kaluza-Klein level of the graviton tower ($F_0 = 1$ the flat zero mode). Unit: dimensionless. Status: DERIVED.

$$\begin{aligned} g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N = \mathrm{e}^{2a_4}\Big[\big(1-2\Psi_\mathrm{N}\big)\big(\mathrm{d}x_1^2+\mathrm{d}x_2^2+\mathrm{d}x_3^2\big) + \mathrm{d}\vec{x}\cdot h^\mathrm{TT}\cdot\mathrm{d}\vec{x}\Big] - \big(1+2\Phi\big)\,\mathrm{d}x_4^2 \\ + f^2\big(1+2\sigma_\mathrm{r}\big)\Big[\mathrm{d}y^2 - \mathrm{e}^{2Hy}\big(\mathrm{d}x_5^2+\mathrm{d}x_6^2+\mathrm{d}x_7^2\big)\Big] \end{aligned} \tag{FE-88}$$

Symbols in (FE-88).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval; g^{MN} is its inverse, ${g^M}_N = \delta^M_N$, and $|g|$ the absolute value of its determinant, so that $\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element. Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first; X^M is the coordinate with the index M, $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element and X^μ (four-dimensional index μ) the observable coordinates $x_1, \ldots, x_4$. Unit: length (each coordinate). Status: AUTHOR.

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- Ψ_N: The curvature potential of the Newtonian gauge (the perturbation of the 3-space metric, $1-2\Psi_\mathrm{N}$); not related to the field Ψ. Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space (also the symmetry axis of the fast-flavour analysis). Unit: length. Status: AUTHOR.
- $\vec{x}$: The position in ordinary 3-space, $\vec{x}=(x_1,x_2,x_3)$; $\mathrm{d}\vec{x}^{\,2}=\mathrm{d}x_1^2+\mathrm{d}x_2^2+\mathrm{d}x_3^2$. Unit: length. Status: STANDARD.
- h^TT: The transverse-traceless (tensor) part of the first-order metric of 3-space (gravitational waves). Unit: dimensionless. Value: an input (tensor-to-scalar ratio below 0.036). Status: STANDARD.
- Φ: The lapse potential of the Newtonian gauge (the perturbation of the time component, $1+2\Phi$). Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- x_4: The time; dots denote $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, c = 1). Status: AUTHOR.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f=\mathrm{e}^{-a_4}b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- σ_r: The radion perturbation: the first-order change of the block scale, $f^2\to f^2(1+2\sigma_\mathrm{r})$. Unit: dimensionless. Value: at most 1.1×10^{-34} (static response). Status: DERIVED.
- y: The hidden coordinate of the y chart, $y=\ln|\sin z|/(6H)$, with $y=0$ at the fold ($z=\pi/2$) and $y=-L$ at the regular tip; each half of a period of x_0 is the interval $-L\le y\le0$. Unit: length. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda=-10H^2$ and the mass parameter $m_\Psi=H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.

The terms with $n\ge1$ are the Kaluza-Klein gravitons, the eigenfunctions of the x_0 operator of the graviton (the operator of the gauge fields with $\beta=0$):

$$\begin{aligned}&\int_{-L}^{0}\mathrm{d}y\,f^4\mathrm{e}^{3Hy}\Big[\frac{(\partial_yF)^2}{f^2}-m^2F^2\Big]\ \text{stationary}: \qquad -\frac{1}{f^2}\,\mathrm{e}^{-3Hy}\,\partial_y\big(\mathrm{e}^{3Hy}\,\partial_yF\big)=m^2F,\\ &\Leftrightarrow\qquad -\mathrm{e}^{-3HL\xi}\,\frac{\mathrm{d}}{\mathrm{d}\xi}\Big(\mathrm{e}^{3HL\xi}\,\frac{\mathrm{d}F}{\mathrm{d}\xi}\Big)=\tilde m^2F,\qquad \tilde m=m\,fL\end{aligned}\tag{FE-50}$$

Symbols in (FE-50).

- L: The depth of the regular tip: each half of a period of x_0 ends at $y=-L$; $HL=3$. Unit: length. Value: $L=3/H$ ($fL=8.926\times10^{-21}$ m at the stabilised block). Status: ASSUMED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- y: The hidden coordinate of the y chart, $y=\ln|\sin z|/(6H)$, with $y=0$ at the fold ($z=\pi/2$) and $y=-L$ at the regular tip; each half of a period of x_0 is the interval $-L\le y\le0$. Unit: length. Status: DERIVED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f=\mathrm{e}^{-a_4}b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda=-10H^2$ and the mass parameter $m_\Psi=H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- ∂: A partial derivative: ∂_M is the derivative with respect to the coordinate X^M, ∂_y with respect to y, ∂_{x_4}

with respect to the time x_4, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.

- y: The label y in ∂_y: the derivative with respect to the hidden coordinate y. Status: STANDARD.
- $F(\xi),\ F_n$: The x_0 profile of a boson mode on one half of the period (a function of y or of $\xi = y/L$): of a gauge field (gauge $A_y = 0$) or of the transverse-traceless graviton; F_n is the profile of the n-th Kaluza-Klein level of the graviton tower ($F_0 = 1$ the flat zero mode). Unit: dimensionless. Status: DERIVED.
- $m,\ m_s,\ m_i$: A four-dimensional mass: m the mass of a mode (the eigenvalue of an x_0 problem) or of a fermion; m_s the mass of the species s; m_i the mass of the neutrino mass state i. Unit: GeV. Status: DERIVED.
- ξ: The scaled hidden coordinate, $\xi = y/L$, running from -1 (tip) to 0 (fold) on each half of a period. Unit: dimensionless. Status: DERIVED.
- $\tilde{m},\ \tilde{m}_n$: The dimensionless x_0 eigenvalue of a boson mode, $\tilde{m} = m\,fL$; $\tilde{m}_n$ the n-th even Kaluza-Klein level of the tower with $\beta = 0$ ($\tilde{m}_1$ the first). Unit: dimensionless. Value: $\tilde{m}_1^2 = 30.1196$, $\tilde{m}_2^2 = 59.7284$, $\tilde{m}_3^2 = 109.0764$ ($HL = 3$), i.e. $\tilde{m}_1 = 5.48813$, $\tilde{m}_2 = 7.72842$, $\tilde{m}_3 = 10.44397$. Status: DERIVED.

$$\left.\frac{\mathrm{d}F}{\mathrm{d}\xi}\right|_{\xi=0} = 0 \ \ \text{(even)} \quad \text{or} \quad F(0) = 0 \ \ \text{(odd)}, \qquad \left.\frac{\mathrm{d}F}{\mathrm{d}\xi}\right|_{\xi=-1} = \beta\, F(-1),$$
$$\beta = 0 \ \ \text{(photon, gluons, graviton)}, \qquad \beta = \beta_{\mathrm{W}} \ \ (W^{\pm}), \qquad \beta = \beta_{\mathrm{Z}} = \frac{\beta_{\mathrm{W}}}{\cos^2\theta_{\mathrm{W}}} \ \ (Z) \tag{FE-51}$$

Symbols in (FE-51).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $F(\xi),\ F_n$: The x_0 profile of a boson mode on one half of the period (a function of y or of $\xi = y/L$): of a gauge field (gauge $A_y = 0$) or of the transverse-traceless graviton; F_n is the profile of the n-th Kaluza-Klein level of the graviton tower ($F_0 = 1$ the flat zero mode). Unit: dimensionless. Status: DERIVED.
- ξ: The scaled hidden coordinate, $\xi = y/L$, running from -1 (tip) to 0 (fold) on each half of a period. Unit: dimensionless. Status: DERIVED.
- β: The Robin parameter of the tip condition of a boson profile, $\mathrm{d}F/\mathrm{d}\xi = \beta F$ at $\xi = -1$ (zero for the massless gauge fields and the graviton). Unit: dimensionless. Status: DERIVED.
- β_{W}: The Robin parameter of W^+ and W^- (set by the tip mass term of the sigma field). Unit: dimensionless. Value: 0.011912 (fitted). Status: DERIVED.
- $W_M^{\pm},\ W^+,\ W^-$: The charged weak bosons, canonically normalised: $W_M^{\pm} = (W_M^1 \mp \mathrm{i}W_M^2)/(\sqrt{2}\,\hat{g}_2)$; W^+ carries the electric charge $+e$ and W^- the charge $-e$ (they are each other's antiparticles and have one mass). Unit: GeV^3. Status: STANDARD.
- β_{Z}: The Robin parameter of the Z, $\beta_{\mathrm{Z}} = \beta_{\mathrm{W}}/\cos^2\theta_{\mathrm{W}}$. Unit: dimensionless. Value: 0.015343. Status: DERIVED.
- θ_{W}: The weak mixing angle of the couplings, $\cos\theta_{\mathrm{W}} = \hat{g}_2/\hat{g}_Z$, $\sin\theta_{\mathrm{W}} = \hat{g}_1/\hat{g}_Z$; fixed by the measured W and Z masses with $\rho_{\mathrm{EW},0}$. Unit: dimensionless. Value: $\cos^2\theta_{\mathrm{W}} = 0.776368$. Status: DERIVED.
- Z_M: The Z boson field, canonically normalised, $Z_M = (W_M^3 - B_M)/\hat{g}_Z$. Unit: GeV^3. Status: STANDARD.

$$F_n(\xi) = \mathrm{e}^{-3HL\xi/2}\Big(\cos\varpi_n\xi + \frac{3HL}{2\varpi_n}\sin\varpi_n\xi\Big), \qquad \varpi_n = \pi,\ 2\pi,\ 3\pi,\ \ldots,$$
$$\tilde{m}_n^2 = \frac{9H^2L^2}{4} + \varpi_n^2, \qquad m_n^{\mathrm{KK}} = \frac{\tilde{m}_n}{fL}, \qquad N_n = \int_{-1}^{0}\mathrm{e}^{3HL\xi}F_n^2\,\mathrm{d}\xi = \frac{1}{2}\Big(1 + \frac{9H^2L^2}{4\varpi_n^2}\Big), \qquad N_0 = \frac{1-\mathrm{e}^{-3HL}}{3HL} \tag{FE-53}$$

Symbols in (FE-53).

- $F(\xi),\ F_n$: The x_0 profile of a boson mode on one half of the period (a function of y or of $\xi = y/L$): of a gauge field (gauge $A_y = 0$) or of the transverse-traceless graviton; F_n is the profile of the n-th Kaluza-Klein level of the graviton tower ($F_0 = 1$ the flat zero mode). Unit: dimensionless. Status: DERIVED.
- n: The level number of the Kaluza-Klein tower ($n = 1, 2, 3, \ldots$; $n = 0$ the flat zero mode). Status: DERIVED.
- ξ: The scaled hidden coordinate, $\xi = y/L$, running from -1 (tip) to 0 (fold) on each half of a period. Unit: dimensionless. Status: DERIVED.

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- L: The depth of the regular tip: each half of a period of x_0 ends at $y = -L$; $HL = 3$. Unit: length. Value: $L = 3/H$ ($fL = 8.926 \times 10^{-21}$ m at the stabilised block). Status: ASSUMED.
- ϖ, ϖ_n: The wave number of an oscillating x_0 profile on $-1 \le \xi \le 0$ (of a boson after the rescaling, or of a bulk level of dirac16complex); ϖ_n that of the n-th even level ($\varpi_n = n\pi$ for $\beta = 0$). Unit: dimensionless. Value: 1.89901 for the lowest odd level of dirac16complex. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $\tilde{m}$, $\tilde{m}_n$: The dimensionless x_0 eigenvalue of a boson mode, $\tilde{m} = m\,fL$; $\tilde{m}_n$ the n-th even Kaluza-Klein level of the tower with $\beta = 0$ ($\tilde{m}_1$ the first). Unit: dimensionless. Value: $\tilde{m}_1^2 = 30.1196$, $\tilde{m}_2^2 = 59.7284$, $\tilde{m}_3^2 = 109.0764$ ($HL = 3$), i.e. $\tilde{m}_1 = 5.48813$, $\tilde{m}_2 = 7.72842$, $\tilde{m}_3 = 10.44397$. Status: DERIVED.
- m_n^{KK}, m_{KK}: The mass of the n-th Kaluza-Klein graviton (and gauge) level, $m_n^{\mathrm{KK}} = \tilde{m}_n/(fL)$; $m_{\mathrm{KK}} = m_1^{\mathrm{KK}}$ is the Kaluza-Klein scale. Unit: GeV. Value: $m_{\mathrm{KK}} = 121.32$ TeV. Status: DERIVED.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- N_n, N_0: The normalisation integral of the profile F_n of the n-th Kaluza-Klein level, $N_n = \int_{-1}^{0} \mathrm{e}^{3HL\xi} F_n^2 \,\mathrm{d}\xi$; N_0 that of the flat zero mode. Unit: dimensionless. Value: $N_0 = (1 - \mathrm{e}^{-3HL})/(3HL) = 0.1111$. Status: DERIVED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.

$$V_{\mathrm{N}}(r) = -\frac{G_{\mathrm{N}} M_1 M_2}{r}\Big[1 + \sum_{n\ge 1} \mathcal{A}_n \,\mathrm{e}^{-m_n^{\mathrm{KK}} r}\Big], \qquad \mathcal{A}_n = \frac{N_0}{N_n}\Big(\int_{-1}^{0} \varrho_1 F_n \,\mathrm{d}\xi\Big)\Big(\int_{-1}^{0} \varrho_2 F_n \,\mathrm{d}\xi\Big) \tag{FE-54}$$

Symbols in (FE-54).

- $V_{\mathrm{N}}(r)$: The potential energy of two static masses at the distance r (Newton potential with the Kaluza-Klein corrections). Unit: GeV. Status: DERIVED.
- r: A distance: between the two masses in $V_{\mathrm{N}}(r)$, from the centre of the star in the adiabaticity condition. Unit: length. Status: STANDARD.
- G_{N}: Newton's constant of four-dimensional gravity, $8\pi G_{\mathrm{N}} = \kappa/(\mathcal{G}(K_*)V_\perp)$. Unit: GeV^{-2}. Value: 6.709×10^{-39} GeV^{-2} ($M_{\mathrm{Pl}} = G_{\mathrm{N}}^{-1/2} = 1.22091 \times 10^{19}$ GeV). Status: STANDARD.
- M_1, M_2: The masses of the two static sources in the potential V_{N}. Unit: GeV. Status: STANDARD.
- n: The level number of the Kaluza-Klein tower ($n = 1, 2, 3, \ldots$; $n = 0$ the flat zero mode). Status: DERIVED.
- $\mathcal{A}_n$: The relative strength of the Yukawa force of the n-th Kaluza-Klein graviton between two sources with the profiles ϱ_1, ϱ_2. Unit: dimensionless. Value: 0.0904, 0.0136, 0.0057 ($n = 1, 2, 3$, two quark-doublet profiles). Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- m_n^{KK}, m_{KK}: The mass of the n-th Kaluza-Klein graviton (and gauge) level, $m_n^{\mathrm{KK}} = \tilde{m}_n/(fL)$; $m_{\mathrm{KK}} = m_1^{\mathrm{KK}}$ is the Kaluza-Klein scale. Unit: GeV. Value: $m_{\mathrm{KK}} = 121.32$ TeV. Status: DERIVED.
- N_n, N_0: The normalisation integral of the profile F_n of the n-th Kaluza-Klein level, $N_n = \int_{-1}^{0} \mathrm{e}^{3HL\xi} F_n^2 \,\mathrm{d}\xi$; N_0 that of the flat zero mode. Unit: dimensionless. Value: $N_0 = (1 - \mathrm{e}^{-3HL})/(3HL) = 0.1111$. Status: DERIVED.
- ϱ_1, ϱ_2: The x_0 profiles of the two sources of V_{N} as probability densities in ξ ($\int_{-1}^{0} \varrho_1 \,\mathrm{d}\xi = 1$): L times the profile ϱ_f of the zero mode of each source. Unit: dimensionless. Status: DERIVED.
- $F(\xi)$, F_n: The x_0 profile of a boson mode on one half of the period (a function of y or of $\xi = y/L$): of a gauge field (gauge $A_y = 0$) or of the transverse-traceless graviton; F_n is the profile of the n-th Kaluza-Klein level of the graviton tower ($F_0 = 1$ the flat zero mode). Unit: dimensionless. Status: DERIVED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$

the volume element of n variables. Status: STANDARD.

- ξ: The scaled hidden coordinate, $\xi = y/L$, running from -1 (tip) to 0 (fold) on each half of a period. Unit: dimensionless. Status: DERIVED.

The radion is the block mode of step 1, with its static response of step 8:

$$\begin{aligned} &\Delta\dot{\mathcal{H}} = \Xi_{\mathcal{H}K}\,\Delta K, \qquad \Delta\dot{\mathcal{H}}_f = \Xi_{fK}\,\Delta K, \qquad \Delta\dot{K} = -2K_*\,\Delta\mathcal{H}_f, \\ &\omega_{\mathrm{r}}^2 = 2K_*\,\Xi_{fK} = \Big(\frac{31552}{2249} - \frac{2416\sqrt{51}}{6747}\Big)H^2 = (3.38705\,H)^2 \end{aligned} \tag{FE-44}$$

Symbols in (FE-44).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- $\dot{\mathcal{H}}$: The time derivative $\mathrm{d}\mathcal{H}/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- $\Xi_{\mathcal{H}K}$: The coefficient of ΔK in the first-order equation for $\Delta\dot{\mathcal{H}}$ (linearised block about K_*). Unit: GeV^2. Value: $6(238 - 19\sqrt{51})H^2/2249 = 0.272956\,H^2$. Status: DERIVED.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- $\dot{\mathcal{H}}_f$: The time derivative $\mathrm{d}\mathcal{H}_f/\mathrm{d}x_4$. Unit: GeV^2. Status: DERIVED.
- Ξ_{fK}: The coefficient of ΔK in the first-order equation for $\Delta\dot{\mathcal{H}}_f$. Unit: GeV^2. Value: $(4063 - 485\sqrt{51})H^2/2249 = 0.266522\,H^2$. Status: DERIVED.
- $\dot{K}$: The time derivative $\mathrm{d}K/\mathrm{d}x_4$. Unit: GeV. Status: DERIVED.
- K_*: The block variable of the stabilised block, $K_* = 12 + \frac{4}{3}\sqrt{51}$ (the upper root K_+). Unit: dimensionless. Value: 21.5219. Status: DERIVED.
- $\mathcal{H}_f$: The expansion rate of the block, $\mathcal{H}_f = \dot{f}/f = \dot{b}_4/(2b_4) - \dot{a}_4$ (negative when the extra times and x_0 deflate). Unit: GeV. Status: DERIVED.
- ω_{r}: The angular frequency of the oscillation of the block about the static point (the radion), equal to the radion mass. Unit: GeV. Value: $3.38705\,H = 48.42$ TeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_{W} and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.

$$\begin{aligned} &\Delta K = \frac{c_\rho\,\kappa\Delta\rho + c_3\,\kappa\Delta p_3 + c_p\,\kappa\Delta\bar{p}}{H^2}, \qquad \Delta\ln f = -\frac{\Delta K}{2K_*}, \\ &\Delta\ln m_s = \frac{\mathrm{d}\ln m_s}{\mathrm{d}\ln f}\,\Delta\ln f, \qquad \kappa\Delta\rho = 8\pi G_{\mathrm{N}}\,\mathcal{G}(K_*)\,\rho_4 \end{aligned} \tag{FE-80}$$

Symbols in (FE-80).

- Δ: The change (difference or small perturbation) of the quantity that follows: ΔK, $\Delta\rho$, Δp_3 are first-order changes about the static block, $\Delta\varepsilon_o$ a level shift, $\Delta m_{21}^2 = m_2^2 - m_1^2$ a difference of squared masses, ΔN_{eff} an addition to N_{eff}, ΔS_{m} the change of an action under a change Δg^{MN} of the metric. In an uncertainty relation it is the spread, the root-mean-square deviation from the mean value in a state: Δx_0, Δk_{x_0}, $\Delta\xi$, $\Delta(fy)$ (as in the author's $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$). Status: STANDARD.
- K: The block variable, $K = 1/f^2$. Unit: dimensionless. Status: DERIVED.
- c_ρ: The coefficient of $\kappa\Delta\rho$ in the adiabatic block response ΔK. Unit: dimensionless. Value: 0.0358521. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Value: $\kappa/V_\perp = 8\pi G_{\mathrm{N}}\mathcal{G}(K_*) = 2.328 \times 10^{-36}\ \mathrm{GeV}^{-2}$. Status: AUTHOR.
- ρ: An energy density: in eight dimensions $\rho = -\mathcal{T}^{x_4}{}_{x_4}$ (of the total source, or of the species named by a subscript; ρ_s of the species s, ρ_i of the neutrino mass state i); in the cosmological perturbation equations $\Delta\rho$ is the density perturbation summed over the species. Unit: GeV^8 (eight dimensions), GeV^4 (four dimensions). Status: STANDARD.

- c_3: The coefficient of $\kappa\Delta p_3$ in the adiabatic block response, $c_3 = -3c_\rho$ (so radiation drops out). Unit: dimensionless. Value: -0.107556. Status: DERIVED.
- p_3: The pressure of ordinary 3-space, $p_3 = \mathcal{T}^{x_1}{}_{x_1} = \mathcal{T}^{x_2}{}_{x_2} = \mathcal{T}^{x_3}{}_{x_3}$ for an isotropic source. Unit: GeV^8. Status: STANDARD.
- c_p: The coefficient of $\kappa\Delta\bar{p}$ in the adiabatic block response. Unit: dimensionless. Value: -0.140028. Status: DERIVED.
- $\bar{p}$: The hidden pressure, the pressure averaged over the block, $\bar{p} = (p_0 + 3p_5)/4$; for a species at fixed temperature $\bar{p} = \frac{1}{4}\,\partial p/\partial \ln f$. Unit: GeV^8 (or GeV^4 for a four-dimensional density). Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings, the cosmological constant $\Lambda = -10H^2$ and the mass parameter $m_\Psi = H$. Unit: GeV (1/length). Value: 14.296 TeV (from the fit of m_W and $\rho_{\mathrm{EW},0}$). Status: AUTHOR.
- f: The block scale: the common scale factor of the hidden direction and the extra times, $f = \mathrm{e}^{-a_4} b_4^{1/2}$ (the block is f^2 times a four-dimensional space of constant curvature $-H^2$). Unit: dimensionless. Status: DERIVED.
- K_*: The block variable of the stabilised block, $K_* = 12 + \frac{4}{3}\sqrt{51}$ (the upper root K_+). Unit: dimensionless. Value: 21.5219. Status: DERIVED.
- m, m_s, m_i: A four-dimensional mass: m the mass of a mode (the eigenvalue of an x_0 problem) or of a fermion; m_s the mass of the species s; m_i the mass of the neutrino mass state i. Unit: GeV. Status: DERIVED.
- s: The label of a particle species of the plasma or of matter (photon, gluons, $W^\pm$, Z, quarks, leptons, nucleons, dark matter); a sum over s runs over the species present. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_N: Newton's constant of four-dimensional gravity, $8\pi G_\mathrm{N} = \kappa/(\mathcal{G}(K_*)V_\perp)$. Unit: GeV^{-2}. Value: 6.709×10^{-39} GeV^{-2} ($M_\mathrm{Pl} = G_\mathrm{N}^{-1/2} = 1.22091 \times 10^{19}$ GeV). Status: STANDARD.
- $\mathcal{G}(K)$: The coefficient of the four-dimensional Einstein-Hilbert term on a static block, $\mathcal{G}(K) = 1 - 24w_{(2)}K + 72w_{(3)}K^2 = (K^2 - 12K + 16)/16$; four-dimensional gravity is attractive for $\mathcal{G} > 0$. Unit: dimensionless. Value: $\mathcal{G}(K_*) = 20/3 + \sqrt{51} = 13.8081$. Status: DERIVED.
- ρ_4: A four-dimensional energy density (per 3-volume); its block average is $\kappa\Delta\rho = 8\pi G_\mathrm{N}\mathcal{G}(K_*)\rho_4$. Unit: GeV^4. Status: DERIVED.

The size of the first-order potentials:

$$\Phi_\mathrm{rms}(k) = \Big[\frac{k^3 P_{\Phi_\mathrm{W}}(k)}{2\pi^2}\Big]^{1/2}, \qquad \Phi_\mathrm{W} = \frac{\Phi + \Psi_\mathrm{N}}{2}, \qquad \Phi_\mathrm{W} = -\frac{3}{2}\,\Omega_\mathrm{m}\mathcal{H}_0^2\,\frac{\delta_\mathrm{m}}{\mathrm{e}^{a_4}k^2} \quad \text{(inside the horizon)},$$
$$\Phi_\mathrm{W} = -\frac{3}{5}\,\mathcal{R}\,\mathrm{e}^{-a_4} D_+ \quad \text{(largest scales)}, \qquad D_+ = \frac{5}{2}\,\Omega_\mathrm{m}\,\frac{\mathcal{H}}{\mathcal{H}_0}\int_{-\infty}^{a_4}\Big(\frac{\mathcal{H}_0}{\mathcal{H}}\Big)^3 \mathrm{e}^{-2\tilde{a}_4}\,\mathrm{d}\tilde{a}_4, \tag{FE-98}$$
$$\frac{\mathcal{H}}{\mathcal{H}_0} = \big(\Omega_\mathrm{m}\mathrm{e}^{-3a_4} + 1 - \Omega_\mathrm{m}\big)^{1/2}$$

Symbols in (FE-98).

- $\Phi_\mathrm{rms}(k)$: The root-mean-square amplitude of the Weyl potential per logarithmic interval of k. Unit: dimensionless. Value: 2.42×10^{-5} at $k = 10^{-4}$ Mpc^{-1}. Status: STANDARD.
- k: The magnitude of the wave vector (the momentum, $\hbar = 1$) of a mode: in the reduction, the Kohn-Sham and the perturbation equations the comoving wavenumber (physical momentum $\mathrm{e}^{-a_4}k$); in the thermal integrals the momentum of a particle at the time considered (the variable of integration). Unit: GeV (1/length). Status: STANDARD.
- $P_{\Phi_\mathrm{W}}(k)$: The power spectrum of the Weyl potential. Unit: length^3. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- Φ_W: The Weyl potential, $\Phi_\mathrm{W} = (\Phi + \Psi_\mathrm{N})/2$ (it deflects light). Unit: dimensionless. Status: STANDARD.
- Φ: The lapse potential of the Newtonian gauge (the perturbation of the time component, $1 + 2\Phi$). Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.

- $\Psi_{\rm N}$: The curvature potential of the Newtonian gauge (the perturbation of the 3-space metric, $1-2\Psi_{\rm N}$); not related to the field Ψ. Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- $\Omega_{\rm m}$: The matter density today in units of the critical density. Unit: dimensionless. Value: 0.3150. Status: STANDARD.
- $\mathcal{H}_0$: The Hubble rate today (the Hubble constant). Unit: GeV (km/s/Mpc). Value: 67.38 km/s/Mpc. Status: DERIVED.
- $\delta_{\rm m}$: The density contrast of the total matter (dark matter and baryons). Unit: dimensionless. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: the scale factor of cosmology is e^{a_4}. Unit: dimensionless. Status: AUTHOR.
- $\mathcal{R}$: The primordial comoving curvature perturbation (conserved outside the horizon), counted positive where the density is above its mean, so that the Newtonian potentials Φ, $\Psi_{\rm N}$ and $\Phi_{\rm W}$ are negative there (on the largest scales in the matter era $\Phi_{\rm W}=-\frac{3}{5}\mathcal{R}\,\mathrm{e}^{-a_4}D_+$). Unit: dimensionless. Status: STANDARD.
- D_+: The linear growth factor of the matter perturbations, normalised as $D_+=\mathrm{e}^{a_4}$ in the matter era. Unit: dimensionless. Value: $D_+\mathrm{e}^{-a_4}=0.7878$ today. Status: STANDARD.
- $\mathcal{H}$: The expansion rate of ordinary 3-space, $\mathcal{H}=\dot a_4$ (the Hubble rate of the observer; in the hot Universe $\mathcal{H}=(8\pi^3 g_\rho/90)^{1/2}T^2/M_{\rm Pl}$). Unit: GeV. Status: DERIVED.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no upper bound. Status: STANDARD.
- $\tilde a_4$: An earlier value of a_4 (the integration variable of the growth integral). Unit: dimensionless. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.

13.2 The approximation of step 10

- *Order 2 is estimated* as the square of order 1 (ESTIMATE); on cosmological scales it is 2.4×10^{-5} of the kept first order. Near a neutron star, where $\Phi^2=0.03$, the series converges slowly and full nonlinear general relativity applies (outside the cosmological solution).
- *The graviton tower* is that of Einstein gravity on the block; the Lovelock corrections to its operator are not estimated (OPEN).
- *Tensor modes* are not predicted (the primordial spectrum is an input).

13.3 The solution of step 10

The solution fixes the magnitude of every term (DERIVED, 4 checks):

- *order 0*, the background $\bar g_{MN}$: 1;
- *order 1, $n=0$*: the rms Newtonian potential per logarithmic wavenumber today, $\Phi_{\rm rms}$ of the Weyl potential of step 6 (UM-2): 2.42×10^{-5}, 2.28×10^{-5}, 1.51×10^{-5}, 4.12×10^{-6}, 1.74×10^{-6}, 3.55×10^{-7}, 5.03×10^{-8} at $k=10^{-4}$, 10^{-3}, 10^{-2}, 0.05, 0.1, 0.3, 1 Mpc^{-1}; the Poisson form applied to the matter density agrees to 1.5×10^{-6} for $k\geq0.01$ Mpc^{-1}, and on the largest scales the matter-era form with $\mathcal{R}=A_{\rm s}^{1/2}(k/k_0)^{(n_{\rm s}-1)/2}$ ($k_0=0.05$ Mpc^{-1}, $D_+\mathrm{e}^{-a_4}=0.7878$ today) to 4.4×10^{-5}.

Bound objects add to the order-1, $n=0$ term the potential

$$|\Phi|=\frac{G_{\rm N}M_1}{r}$$

Symbols in this equation.

- Φ: The lapse potential of the Newtonian gauge (the perturbation of the time component, $1+2\Phi$). Unit: dimensionless. Value: at most 2.4×10^{-5}. Status: STANDARD.
- $G_{\rm N}$: Newton's constant of four-dimensional gravity, $8\pi G_{\rm N}=\kappa/(\mathcal{G}(K_*)V_\perp)$. Unit: GeV^{-2}. Value: 6.709×10^{-39} GeV^{-2} ($M_{\rm Pl}=G_{\rm N}^{-1/2}=1.22091\times10^{19}$ GeV). Status: STANDARD.
- M_1, M_2: The masses of the two static sources in the potential $V_{\rm N}$. Unit: GeV. Status: STANDARD.

- r: A distance: between the two masses in $V_{\rm N}(r)$, from the centre of the star in the adiabaticity condition. Unit: length. Status: STANDARD.

at the surface (mass M_1, radius r): the Earth 7.0×10^{-10}, the Sun 2.1×10^{-6}, a white dwarf 1.0×10^{-4}, a neutron star 0.17, galaxies (the squared orbital velocity) 5.4×10^{-7}, clusters 1.1×10^{-5}. The remaining terms:

- *order 1,* $n\geq1$: the graviton tower with $\tilde m_n^2=30.1196$, 59.7284, 109.0764 for $n=1,2,3$ (finite elements agree to 3.9×10^{-9}) and $N_n=\frac12(1+9H^2L^2/(4n^2\pi^2))$ (to 1.8×10^{-10}); the strengths $\mathcal{A}_n$ of the Yukawa terms of the potential between two static masses with the same profile are 0.0728, 0.1469, 0.1809 ($n=1,2,3$) for sources at the fold, 0.0904, 0.0136, 0.0057 for the quark-doublet profile, 0.0208, 0.0197, 0.0105 for the lepton doublet, and 0 for the flat profile of the gauge zero modes (to 1.7×10^{-10}); $m_1^{\rm KK}=m_{\rm KK}=121.32$ TeV, the range 1.63×10^{-21} m, so $\mathrm{e}^{-m_1^{\rm KK}r}=\mathrm{e}^{-3.1\times10^{16}}$ at $r=50$ micrometres;
- *the radion*: $\Delta\ln f\leq1.1\times10^{-34}$ (step 8), the range 4.08×10^{-21} m, the primordial oscillation decayed (lifetime $0.43\,\mathrm{s}/C_{\rm r}$, last section);
- *order 2* (ESTIMATE, the square of order 1): $\Phi^2_{\rm rms}=(2.42\times10^{-5})^2=5.8\times10^{-10}$ on cosmological scales ($k=10^{-4}$ Mpc^{-1}), 4.5×10^{-12} at the Sun, 0.03 at a neutron star;
- *tensor modes*: not predicted; the bound 0.036 on the tensor-to-scalar ratio (quoted without a checked reference) bounds their amplitude by 8.7×10^{-6}.

On cosmological scales the terms decrease as 1, 2.4×10^{-5}, 5.8×10^{-10}, 10^{-34} (radion), 0 (Kaluza-Klein): the metric is a power series of terms of decreasing magnitude whose $n=0$ first-order term carries the CMB, the CNB and the BAO.

13.4 The check of step 10

In its own equations. The stored graviton profiles satisfy the graviton operator to 1.2×10^{-12} of its largest term, the fold and tip conditions to 9×10^{-16}, the orthogonality to the flat profile to 9×10^{-14} and the normalisations to 9×10^{-13}; the stored Yukawa strengths and eigenvalues are reproduced to 5.2×10^{-9} (a trapezoid integration of 200001 points) and the overlaps with a flat source vanish to 1.7×10^{-10} (DERIVED). The first omitted order of the series is 2.4×10^{-5} of the kept one on cosmological scales; the stored order-2 magnitudes and the potentials of bound objects are reproduced to 7×10^{-13} and 2.2×10^{-12}, and the terms decrease strictly (DERIVED). The Kaluza-Klein correction to Newton's law at the smallest measured distance, a Yukawa term proportional to $\mathrm{e}^{-m_1^{\rm KK}r}$ with $m_1^{\rm KK}r=3.07\times10^{16}$ at 50 micrometres, is zero in double precision.

In concert with the other field equations. The radion frequency of the series is that of the linear block equations and of the nonlinear primordial oscillation to 7.9×10^{-5} (step 1); the potentials, the block shift, the first Kaluza-Klein mass and σ_8 are those of steps 6, 8 and 2 to 4.7×10^{-12}; the Poisson form of the first-order potential holds in the CAMB solution to 1.5×10^{-6} (step 6) (DERIVED).

13.5 The status of step 10

DERIVED: the magnitudes of order 0, of order 1 ($n=0$ and $n\geq1$) and of the radion. ESTIMATE: order 2. STANDARD: the bound on tensor modes (quoted without a checked reference). OPEN: the Lovelock corrections to the graviton tower; the order-2 terms in detail; the primordial spectrum (an input).

14 Accuracy of the solution

The table collects, for every field equation or system of field equations of the solution, the method of solution, the residual of the solution in its own equation (a precisely defined, measured quantity: the residual of the stored solution substituted into the equation, the implied relative error of a root, or the agreement of independent evaluations), the size of the terms that the approximation neglects, and the checks in concert with the other field equations that contain the same fields (the same quantity compared where two or more equations use it). All residuals and links were computed anew in version 4 (`um2_residuals.py`, 52 named checks, all passed; with the 235 checks of the computations of version 3, UM-2 has 287 named checks); "NOT small" marks a neglected term or a difference that is not small and is reported as it is.

equations	solution method	residual in its own equation	neglected terms	in concert
background: FE-38, FE-39, FE-40, FE-42, FE-43 (step 1)	exact closed form (symbolic)	3.9×10^{-13} of the largest term with the stored 12-digit K_* (1.2×10^{-16} closed form); printed = coded equations exactly	matter against the stabiliser: 3.0×10^{-32} (freeze-out), 2.4×10^{-93} (today)	f_* in steps 2, 3, 10: 2.2×10^{-12}
block response: FE-44, FE-45, FE-46, FE-47 (steps 1, 3, 8)	exact algebraic numbers	second order as it must be: 1.4×10^{-7} at sources $10^{-5}H^2$ (ratio 0.10 per decade); radiation cancels to 2×10^{-13}	second order 2.3×10^{-32} and inertia 3.4×10^{-33} at the freeze-out	Kohn-Sham sources 3.4×10^{-12}; coefficients 2×10^{-13}; radion linear vs nonlinear 7.9×10^{-5}
primordial era: FE-38, FE-39, FE-40 integrated (step 1)	DOP853, relative tolerance 10^{-11}	one-step defect 1.3×10^{-11}; constraint 3.6×10^{-12} of its terms	radiation of zero modes: 1.0×10^{-6} of the final block; block-shape mode (OPEN)	radion frequency 7.9×10^{-5}
four-dimensional gravity: FE-48, FE-49 (steps 1, 6)	exact (symbolic)	$\Lambda_4(K_*)$ and the stationarity of the x_4 balance: 3.9×10^{-13} (stored), 10^{-16} (closed form); coefficient of $R^{(4)} = \mathcal{G}(K)$ exactly	dark energy needs $\Lambda_4/H^2 \sim 10^{-92}$ (tuned, OPEN)	Hubble rate of the thermal history at the freeze-out: 1.7×10^{-4} (fold bands left out there)
boson x_0 problem: FE-50, FE-51, FE-52, FE-55 (step 2)	exact roots (mpmath); finite elements as check	implied relative eigenvalue error $\leq 5.9 \times 10^{-12}$; finite elements 5×10^{-10}	dark-matter kinetic function 1.9×10^{-17}; radiative corrections; Lovelock corrections to the graviton (OPEN)	m_{Z} in the widths 2.0×10^{-12}; two values of the weak angle: 0.82 % of the total width (NOT zero; per-cent scheme ambiguity)
fermion x_0 problem: FE-14 to FE-19, FE-30, FE-56 to FE-61 (step 2)	exact roots (mpmath, 50 to 60 digits)	implied relative root error $\leq 5.4 \times 10^{-11}$; three-generation determinant 1.2×10^{-11}; its roots = one-generation roots to 3.0×10^{-42}	mixing with Kaluza-Klein modes $(m_{\mathrm{t}}/m_{\mathrm{KK}})^2 = 2.0 \times 10^{-6}$	masses in the thermal history 3.1×10^{-12}; neutrino masses in flavour, CNB, CMB 5.7×10^{-12}; PMNS 8.2×10^{-13}
spectrum of dirac16complex: FE-12, FE-28, FE-62 (step 2)	exact formulas	8.5×10^{-13}	mean-field shifts of the level 1.5×10^{-6} (step 8)	hidden pressure in three equations 6.1×10^{-12}; M_{DM} 8.1×10^{-13}
Kohn-Sham: FE-63 to FE-69 (step 3)	Galerkin in the free eigenbasis, self-consistent to 10^{-13}, Richardson	re-solved state 7.4×10^{-14} (absolute); one more map 1.8×10^{-14}; y-conservation 8.8×10^{-8}; basis 1.5×10^{-9}	self-interaction mean field at the bound on λ: 2.7×10^{-9}; correlation beyond local exchange (OPEN)	Kohn-Sham sources in gravity 3.4×10^{-12}; portal responses 3.0×10^{-12}

equations	solution method	residual in its own equation	neglected terms	in concert
Kohn-Sham source in gravity: FE-67 in FE-47 (step 3)	cell averages of the energy-momentum tensor	1.4×10^{-12}; free gas cancels to 9×10^{-13}	block-shape source $p_0 - p_5$: 0.9 to 4 of $\bar{p}$ (NOT small; suppressed with $\bar{p}$; OPEN)	response coefficients 2×10^{-13}
portal overlaps: FE-70, FE-71, FE-72 (steps 3, 8)	quadrature of exact orbitals; Hellmann-Feynman	3.8×10^{-12}; Kohn-Sham finite differences 3.0×10^{-10}	screening by the interacting gas 7.2×10^{-10} at the bound on λ	$\mathcal{O}_{\mathrm{DM}}$ across computations 5.7×10^{-6} (rounded 0.7072)
coupled environments: FE-73 to FE-86 (step 8)	closed formulas, iterated	2.0×10^{-11}; iteration converged (loop gain)	loop gain 1.1×10^{-33}; second order 2.3×10^{-12}	plasma energy density vs thermal history 1.3×10^{-3} (by design); G_{A}, G_{V} 6.7×10^{-13}; freeze-out temperature 2.1×10^{-12}
W with the dark-matter kinetic function: FE-79 (step 8)	first-order perturbation theory with the exact eigenfunction and orbital (finite elements in version 3)	independent quadrature 3.3×10^{-12}; shooting 3.0×10^{-11}; photon exactly unshifted	the physical shift 1.9×10^{-17}	W eigenvalue of the finite elements at zero amplitude 2.3×10^{-6}; $\mathcal{O}_{\mathrm{DM}}$ 5.7×10^{-6}
Yang-Mills masses and Z widths: FE-25, FE-102 (steps 2, 4)	closed formulas, tree level	2.7×10^{-12}	electroweak loops 4.1×10^{-3}; b-quark mass -1.3×10^{-3}; α_3^2 term $+1.3 \times 10^{-3}$	m_{Z} 2.0×10^{-12}; F_{h} 5.8×10^{-7}
linear perturbations: FE-90 to FE-99 (step 6)	CAMB 2.0.5, default accuracy	Poisson form of the first-order potential 1.5×10^{-6}; matter-era form 4.4×10^{-5}; masses iterated to 7.8×10^{-12}	order ϵ^2: 2.4×10^{-5} of order ϵ; Kaluza-Klein modes and radion absent	neutrino masses 5.7×10^{-12}; $\Omega_\nu h^2_{100}$ 1.9×10^{-5}; N_{eff} 5×10^{-16}; relic as cold dark matter 3.1×10^{-9}; radiation today 0.59 % (instantaneous decoupling in step 9)
graviton tower: FE-53, FE-54 (steps 2, 10)	exact eigenfunctions	1.2×10^{-12}; boundary 9×10^{-16}; stored strengths 5.2×10^{-9}	Kaluza-Klein correction at 50 micrometres: $\mathrm{e}^{-3.1 \times 10^{16}}$; Lovelock corrections (OPEN)	radion 7.9×10^{-5}; series 4.7×10^{-12}
metric series: FE-87, FE-88 (step 10)	magnitudes of the computed solution	first omitted order 2.4×10^{-5} of the kept one	order 2 at a neutron star 0.03 (nonlinear gravity there)	4.7×10^{-12}
flavour evolution: FE-103 to FE-108 (step 5)	exact matrix exponentials	7.1×10^{-12}; unitarity 1.3×10^{-15}; CPT 2.2×10^{-16}	adiabatic solar formula 1.3×10^{-3}; Landau-Zener 2.6×10^{-77}; collective effects (OPEN)	masses 5.7×10^{-12}; PMNS 8.2×10^{-13}

equations	solution method	residual in its own equation	neglected terms	in concert
thermal potential: FE-109 (step 5)	root of the onset condition	8.7×10^{-12}	CP-asymmetric potential 1.1×10^{-2} (ESTIMATE); $(T/m_{\mathrm{W}})^2 \sim 4 \times 10^{-9}$	masses 5.7×10^{-12}
neutrino decoupling: FE-110 to FE-115 (step 5)	LSODA, relative tolerance 10^{-11}	instantaneous limit recomputed 2.7×10^{-12}; published values 4.6×10^{-4}	electron mass in the rates and momentum dependence: 0.0044 in N_{eff}	N_{eff} in the CMB 5×10^{-16}; F_{h} 5.8×10^{-7}
fast flavour instability: FE-116 to FE-119 (step 5)	Newton iterations; discretised eigenvalues	3.3×10^{-13} (in units of $\sqrt{2} G_{\mathrm{F}} n_{\nu_e}$)	vacuum term 9.0×10^{-6}; nonlinear evolution (OPEN)	
CNB: FE-120 (step 5)	adaptive quadrature	2.9×10^{-12}	non-instantaneous heating 1.08 % (included in the CMB)	$\Omega_\nu h^2_{100}$ in the CMB 1.9×10^{-5}
degrees of freedom: FE-121, FE-122, FE-123 (step 9)	Gauss-Legendre quadrature	5.0×10^{-9} (exact limits); table 4.6×10^{-11}	order-α plasma pressure: -5.4 % at the freeze-out (NOT small)	masses 3.1×10^{-12}; Hubble rate 1.7×10^{-4}; radiation today 0.59 %
relic: FE-124 to FE-128 (steps 7, 9)	Radau (relative tolerance 10^{-10}); root finding	relic condition 3.1×10^{-9}; late-time identity 6.1×10^{-9}	order-α pressure: +2.7 % in $\Omega_{\mathrm{DM}} h^2_{100}$, +1.4 % in G_{A} (ESTIMATE; NOT small); 21.7 % of the rate at the ${}^3S_1$ bound	relic as the cold dark matter of the CMB 3.1×10^{-9}; $\mathcal{O}_{\mathrm{DM}}$ 5.7×10^{-6}; M_{DM} 8.1×10^{-13}
freeze-in: FE-129, FE-130 (step 9)	trapezoid on 600 logarithmic temperatures	3.9×10^{-5}	inverse processes 0.012 of the Hubble rate; $T_{\max}$ (ASSUMED)	N_{eff} of the CMB 5×10^{-16}; G_{V} 6.7×10^{-13}
right-handed neutrinos: FE-131 to FE-134 (step 9)	Radau in $M_{\mathrm{DM}}/T - 2$	2.5×10^{-12}	elastic rate (ESTIMATE): 9.5×10^{-6} of ΔN_{eff}	F_{h} 5.8×10^{-7}
self-interaction channel: FE-135, FE-136 (step 9)	trapezoid on 800 logarithmic temperatures	1.1×10^{-5}	inverse annihilation 0.027; Fierz coefficients (ESTIMATE)	freeze-out temperature 2.1×10^{-12}

Besides these, the approximations of the whole solution are: the static block with an assumed stabiliser; the good sector (no momenta along the extra times); the Kohn-Sham mean-field treatment; the portal couplings as external mean fields; tree level for the electroweak widths; the coherent flavour evolution in the Sun, the Earth and the supernova envelope with schematic density profiles; neutrino decoupling with one temperature per flavour; the linear analysis of the fast flavour instability and a schematic estimate of its effect on the heating; the freeze-out formula of step 7 (replaced by the Boltzmann equation in step 9); order ϵ for the cosmology; ideal gases with a modelled QCD crossover and instantaneous neutrino decoupling for the degrees of freedom; Maxwell-Boltzmann statistics for the freeze-in; the effective cross-sections capped at the partial-wave bounds; the QCD interaction measure below 1 GeV from a lattice-like shape and leading-order perturbation theory above; the order-α electroweak pressure without scalars; one-loop running for the couplings and the proton sensitivity; the elastic ν_R rate and the self-interaction cross-section as estimates.

What the residual computation of version 4 found. The claims of version 3 are confirmed; these points are

not small or were corrected, and are stated here as they are:

- the weak angle has two values in the solution: $\sin^2\theta_{\mathrm{W}} = 0.223632$ on shell in the x_0 problem of W and Z, and the $\overline{\mathrm{MS}}$ value 0.23122 in the Z widths and the couplings; the total width moves by 0.83 %, within the electroweak loop corrections that UM-2 does not compute (step 4);
- the ideal-gas thermal history leaves out the order-α interaction pressure of the plasma (-5.4 % of the pressure at the freeze-out: QCD -4.7 %, $SU(2)$ -0.7 %), which step 8 uses for the condensates; with it $\Omega_{\mathrm{DM}}h^2_{100}$ would rise by about 2.7 % and G_{A} by about 1.4 % (ESTIMATE): the physical accuracy of the relic strength G_{A} is about 1 %, not the 10^{-6} of its root finding (step 9);
- the Hubble rate of the thermal history leaves out the fold bands of the freeze-in (3.5×10^{-4} of the energy density at the freeze-out), so it is 1.7×10^{-4} too small, and the relic moves by about as much (steps 1 and 9);
- the thermal history uses three neutrino species decoupled instantaneously at 2 MeV, while the CMB computation uses $N_{\mathrm{eff}} = 3.0442$: 0.59 % in the radiation today (steps 5, 6, 9); the plasma energy density at the freeze-out differs by 0.13 % between the thermal history and the block sources of the coupled solution, by design (the trace anomaly, step 8);
- the block-shape source $p_0 - p_5$ is about 4 times the kept hidden pressure for dark matter at rest (0.9 for the test states of step 3); its absolute size is suppressed with the hidden pressure, and the mode is OPEN, as in version 3, now quantified (steps 1, 3);
- corrected in the outputs: the coefficient $\mathrm{d}\ln\tilde{m}^2_{\mathrm{W}}/\mathrm{d}A_{\mathrm{mf}}$ of step 8 is -0.111253 (version 3: -0.1126, finite elements limited by rounding; without physical consequence, since the physical amplitudes are below 1.7×10^{-16}); the photon's shift is exactly zero; the derivative of the x_4 balance at K_*, quoted as -6×10^{-12} in version 3, is exactly zero (the quoted number is rounding); the CNB integrals are now accurate to 3×10^{-16} (the quoted digits are unchanged); the ν_R integration of step 9 no longer depends on the twelfth digit of its input; the fold-band gap changes of step 3 are at the rounding level (10^{-13} to 3×10^{-9}), zero in exact arithmetic.

15 What the solution does not reproduce, and what is open

- **Conflicts with data, inherent in the author's order**: no Higgs boson (one is observed at 125 GeV), and without it the longitudinal W^+W^- scattering violates perturbative unitarity near $\sqrt{8\pi}\,(\sqrt{2}G_{\mathrm{F}})^{-1/2} \approx 1.2$ TeV (with the electroweak scale $(\sqrt{2}G_{\mathrm{F}})^{-1/2} = 246$ GeV), far below the Kaluza-Klein modes. (The one-generation conflict of version 1, a total Z width of 1.23 GeV, is removed: three generations give 2.485 GeV against the measured 2.4955 GeV.)
- **Open**: the origin of the stabiliser; the extra-time modes outside the good sector (tachyonic for momenta along the extra times above f times the mass of the field); the closed time-like curves implied by compact extra times; the radion left over from the deflation (ESTIMATE: a Planck-suppressed decay with the rate $C_{\mathrm{r}}\,\omega_{\mathrm{r}}^3/(4\pi\bar{M}_{\mathrm{Pl}}^2)$, with the reduced Planck mass $\bar{M}_{\mathrm{Pl}} = 2.435\times10^{18}$ GeV and an order-one coupling factor C_{r}, gives the lifetime $0.43\,\mathrm{s}/C_{\mathrm{r}}$ and the reheating temperature $1.85\sqrt{C_{\mathrm{r}}}$ MeV for $g_\rho = 10.75$, below the 4.7 MeV that nucleosynthesis needs unless $C_{\mathrm{r}} > 6.5$: a moduli problem); the block-shape mode driven by $p_0 - p_5$; the tuning of Λ_4 ($10^{-92}H^2$); the Kaluza-Klein graviton spectrum with the Lovelock corrections; the electroweak S parameter of the Higgsless tip and the electroweak loop corrections; anomalies of the eight-dimensional bulk theory; the absolute neutrino mass scale and the mass ordering (bounded and preferred by data, chosen here); the collective flavour evolution of supernova neutrinos beyond the linear analysis; the joint CMB and BAO fit of the fold-band variants; the microscopic origin of the portals (I12 alone could give the relic only as a strongly coupled ${}^3P_0$ channel; UM-2 obtains it through I13, whose contact form needs a mediator above about 30 TeV); the self-coupling λ (bounded by ΔN_{eff}, ASSUMED); the highest temperature of the hot Universe ($T_{\max} = 10$ TeV assumed for the freeze-in); the order-two terms of the metric in detail; the primordial spectrum, an input; correlation beyond the local exchange in the Kohn-Sham functional; the interaction pressure of the plasma in the thermal history of the relic (about +2.7 % in the relic, ESTIMATE).