UM-2 provenance

Where every ingredient, equation and number of the universe model UM-2 comes from, how it was computed, and how it was verified

Underlying theory: Patrick L. Nash Derived and verified by Claude (Anthropic)

Provenance, version 2, 2026-10-18 (version 1: 2026-10-17)

Repository once-ere/Dirac_claude, folder universe_model_2

Contents

1 How to read this document **2**

2 The orders **2**
2.1 The order of 2026-10-17 . . . 2
2.2 The orders of 2026-10-18: three neutrino flavours . . . 3

3 Who did what **3**

4 Ingredients from the author (AUTHOR) **3**
4.1 Coordinates and signature . . . 3
4.2 The primordial metric of the notebooks . . . 4
4.3 The Einstein-Lovelock action, weights and cosmological term . . . 4
4.4 The gamma matrices of $Cl(4,4)$ . . . 4
4.5 The mass parameter, the x_0 mass idea and the dark-matter hypothesis . . . 5

5 Ingredients from the Revision record (RECORD) **5**
5.1 The corrected primordial metric (version 2) . . . 5
5.2 The hidden coordinate, the fold and the regular tip . . . 5
5.3 The dirac16complex Lagrangian and the canonical spin connection . . . 5
5.4 Canonical quantisation and the Krein metric . . . 6
5.5 The Kohn-Sham reduction . . . 6
5.6 Earlier work used only for comparison . . . 6

6 Standard physics and measured values (STANDARD) **6**

7 Ingredients new in UM-2 (ASSUMED and INVENTED) **9**
7.1 The metric as a power series (ASSUMED) . . . 9
7.2 The block stabiliser (ASSUMED; its origin is OPEN) . . . 10
7.3 The Standard-Model fields in eight dimensions (ASSUMED structure; boundary conditions INVENTED) . . . 10
7.4 The tip sigma field and the tip matrices (INVENTED for this geometry; no Higgs particle) . . . 10
7.5 The portal interactions (INVENTED) . . . 11
7.6 Parameter choices (ASSUMED) . . . 11

8 The computations and their verification (DERIVED) **11**
8.1 The principal numbers and where they come from . . . 13
8.2 Methods, with the equations they use . . . 16
8.3 Corrections made during the work . . . 17

8.4 What is not in the repository, and why 19
8.5 Reproduction 19
8.6 History in the repository 20

1 How to read this document

This document records the provenance of UM-2, a model of the Universe built on the author's 4+4-dimensional Einstein-Lovelock theory and his dirac16complex field. For every ingredient it states where it comes from, writes the ingredient out as an equation, names the computation that produced or used it, and states how it was verified. The document is self-contained: every equation it mentions is written here.

Version 2 (2026-10-18) contains three generations of quarks and leptons and three neutrino flavours with their mixing and their flavour conversion, ordered by the author on 2026-10-18; it removes the chiral tip Z portal of version 1. What changed, and why, is listed in the section on corrections (item 21).

Labels used throughout:

- **AUTHOR**: taken from the author's Mathematica notebooks or the author's written orders;
- **RECORD**: taken from the Revision record of this repository (earlier work carried out under the author's orders, with its own verification);
- **STANDARD**: textbook physics or a measured value; measured values are marked *verified* (read from the source on 2026-10-17 or 2026-10-18) or *from memory* (quoted from memory of the standard tables, to be checked against the current editions);
- **ASSUMED**: a modelling choice made for UM-2;
- **INVENTED**: a new phenomenological interaction of UM-2;
- **DERIVED**: computed and verified in UM-2;
- **ESTIMATE**: an order-of-magnitude evaluation, not a precision computation;
- **OPEN**: not resolved in this version.

Units: $\hbar = c = 1$; H is the author's constant (an inverse length); in the numerical work $H = 1$ unless a physical unit is stated.

Symbols with more than one use: k is the block variable $1/f^2$, the Lovelock order as the index of α_k and $E_{(k)}$, and the momentum of a mode along x_1, x_2, x_3 in e^{ikx_1} and h_j; $\Lambda = -10H^2$ is the eight-dimensional cosmological constant, $\Lambda = (m\,fL)^2$ in the x_0 eigenvalue problems a dimensionless eigenvalue, and $\Lambda_\star$ the scale of the effective interactions; $\xi = y/L$ is the scaled hidden coordinate (primes denote $d/d\xi$ in the methods), while the subscript t (p_t, v_t, x_t) refers to the extra times and s_h, t_h are the 4×4 blocks of the gamma matrices; ρ is an energy density, except for the electroweak parameter $\rho = \Lambda_W/(\Lambda_Z \cos^2\theta_W)$ (measured value ρ_0), and ϱ a probability density in ξ; θ_W is the angle of the gauge couplings; n is a number density, except as a level index (u_n, $h^{(1,n)}$, $n \geq 1$) and as the basis size in the Richardson formula, where p is the order; w_k are the Lovelock weights, $w = 1/2$ the weight of a fold parity, $w = \sqrt{c^2 - e^2}$ in the fermion mass condition and w an equation-of-state parameter in the supernova fits; σ is the radion in the metric expansion and, with indices, an uncertainty or a Pauli matrix; $i, j = 1, 2, 3$ label generations and neutrino mass states and $\alpha \in \{e, \mu, \tau\}$ flavours; U is the PMNS and V the CKM matrix (a matter potential carries a subscript, V_{CC}); κ (κ_ν, κ_e, κ_u, κ_d, κ_f) is a tip matrix; e is the eigenvalue $\varepsilon f L$ of a fermion, the electron and the elementary charge; μ is a bulk mass and the muon, and $\mu = \sqrt{2} G_F n_{\nu_e}$ (no subscript) the unit of the neutrino self-interaction in the supernova analysis.

2 The orders

2.1 The order of 2026-10-17

The model answers the author's order of 2026-10-17. Its key parts, in the author's words: "Create a new model of the Universe that includes [a] gravity (Einstein-Lovelock gravity); [b] the dirac16complex quantized spinor field (which is hypothesied to represent dark matter); the dirac16complex is coupled to gravity using a non-trivial canonical spin connection; ... interactions of gravity and dirac16complex with SOME of the following Standard Model particles: [c] photons; [d] neutrinos; [e] electrons; [f] quarks (only up and down, for the protons and neutrons); [g] the 3 weak interaction gauge bosons, the W and Z 'particles'. However do NOT include the Higgs particle. You must invent/create new phenomenological interactions ... consistent with qcd and electro-weak models. You

want to be able to model: CMB, CNB, BAO, so your model for the metric tensor ... must be very general, but perhaps expressible as a power series of terms of decreasing magnitude. Put each and every interaction in equation form, and explain and analyze it in documentation and provenance files; these should be both markdown and latex file types; compile all latex to pdf. NEVER refer the reader to some other file! PUT EVERY EQUATION you create and/or use IN THESE FILES. CRITICALLY IMPORTANT: Mass will be given to all non-massless particles, (except the massless particles: graviton, photon and perhaps the quanta of the dirac16complex field), by appropriate wave function dependence on the space coordinate x0 ... periodic in x0 with period $\pi/(6H)$. Localization with respect to x0 implies, using Heisenberg ... $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$ ... [2] push your new, improved, model ... along with provenance (both .md and .pdf files). [3] Compute the coupled field equations and push to repo; create concomitant documention (both .md and .pdf files) and push. [4] Compute an accurate approximate solution to the coupled field equations (use Kohn-Sham theory for the dirac16complex field, which couples to all other fields) and push ... Check and verify this repo."

Further orders of the same day: "analyze the x0 mass mechanism, and ... analyze the validity of associating dark matter (and dark energy) with dirac16complex"; "the W^+ and W^- bosons carry positive and negative electric charges. Do not forget this"; the coordinates are always the author's $\{x_0, x_1, \ldots, x_7\}$ of the notebook `Dirac_Einstein-Lovelock_4+4_Nash.nb`, never relabelled; every verified result is pushed to the repository as soon as it is obtained; no single calculation runs longer than 7 minutes and 3 seconds (423 s) (this limit was lifted on 2026-10-18).

2.2 The orders of 2026-10-18: three neutrino flavours

On 2026-10-18 the author stated that neutrino flavour and flavour conversion are critically important in this project and asked that the paper of Gogilashvili and Tamborra on neutrino flavour conversion in core-collapse supernovae (his file "Neutrino Flavor Conversion.pdf", arXiv:2605.16504; not committed, it is a third-party paper) be read, understood and remembered (paraphrase). He then ordered, in his words: "correct universe model 2 to include three neutrino flavors with mixing and flavor conversion. this is your first and only priority!! ... do correct universe model 2 to include three neutrino flavors with mixing and flavor conversion NOW." Two ways were put to him: (A) three full generations of quarks and leptons, anomaly free generation by generation; (B) three lepton generations with the single quark generation of version 1, whose gauge anomalies would have to be cancelled by an added Wess-Zumino term. His answer: "NOT (B). USE (A)!!!". Version 2 implements (A). A flavour is defined by the charged lepton produced with the neutrino in a charged-current process, so three neutrino flavours require the muon and the tau; with them, anomaly freedom requires the second and third quark generations (a single quark generation would leave $\mathcal{A}_{SU(2)^2U(1)} = -1$ and $\mathcal{A}_{U(1)^3} = 3/2$; DERIVED).

3 Who did what

- The 4+4 framework, the coordinates, the primordial metric of the notebooks, the Einstein-Lovelock action with its weights and Λ, the dirac16complex field and its $Cl(4,4)$ gamma matrices, the mass parameter M, the idea that masses come from the x_0 dependence of wave functions periodic with period $\pi/(6H)$, the dark-matter hypothesis and the demand for three neutrino flavours with mixing and flavour conversion are the AUTHOR's (Patrick L. Nash).
- The corrected primordial metric (version 2), the regular tip cutoff, the Dirac-type Lagrangian with the canonical spin connection, the 2×2 block reduction of the Kohn-Sham problem and its conventions are RECORD items: produced earlier in this repository under the author's orders and verified there.
- Everything labelled ASSUMED, INVENTED or DERIVED below, all the code of UM-2 and all its documents are by Claude (Anthropic), working in this repository on 2026-10-17 (version 1) and 2026-10-18 (version 2).

4 Ingredients from the author (AUTHOR)

4.1 Coordinates and signature

The author's coordinates (notebook `Dirac_Einstein-Lovelock_4+4_Nash.nb`, `X = {x0[], x1[], ..., x7[]}`) are

$$X = \{x_0, x_1, x_2, x_3, x_4, x_5, x_6, x_7\} :$$

x_0 a hidden space-like direction, x_1, x_2, x_3 ordinary space, x_4 the time, x_5, x_6, x_7 three extra time-like directions. The orthonormal frame metric is

$$\eta_{ab} = \mathrm{diag}(+1,+1,+1,+1,-1,-1,-1,-1), \qquad a,b = 0,\dots,7 \text{ aligned with } x_0,\dots,x_7.$$

The hidden coordinate enters through $z = 6Hx_0$; one period of x_0 is $\ell_0 = \pi/(6H)$.

4.2 The primordial metric of the notebooks

The notebook `WaveFunctionUniverse-4+4-Einstein-Lovelock.nb` (the author's local notebook) gives the primordial metric

$$g^{\mathrm{nb}} = \mathrm{diag}\big(\cot^2 z,\ e^{2a_4}(\sin^2 z)^{1/6},\ e^{2a_4}(\sin^2 z)^{1/6},\ e^{2a_4}(\sin^2 z)^{1/6},$$
$$-1,\ -e^{-2a_4}(\sin^2 z)^{1/6},\ -e^{-2a_4}(\sin^2 z)^{1/6},\ -e^{-2a_4}(\sin^2 z)^{1/6}\big),$$

with vielbein $\mathrm{diag}(\cot z, e^{a_4}(\sin^2 z)^{1/12}\times 3, 1, e^{-a_4}(\sin^2 z)^{1/12}\times 3)$ and $a_4 = a_4(x_4)$. By the author's orders of 2026-10-14 a second function $b_4(x_4) > 0$ multiplies the extra-time components ("b4 is a function of x4; b4 is NEVER 1"), and $(\sin z)^{1/3}$ means $(\sin^2 z)^{1/6}$, the positive branch, for every real z.

4.3 The Einstein-Lovelock action, weights and cosmological term

The notebook writes the Einstein-Lovelock vacuum equations (after Lovelock and Rund) as

$$0 = -\Lambda^{\mathrm{nb}} + H^{-2} w_1\,\mathrm{Lovelock1} + H^{-4} w_2\,\mathrm{Lovelock2} + H^{-6} w_3\,\mathrm{Lovelock3},$$

where Lovelock$k$ denotes the notebook's Lovelock tensor of order k, the tensor ${E_{(k)}}^M{}_N$ below, with $(w_1, w_2, w_3, \Lambda^{\mathrm{nb}}) = (1, 1/32, 1/1152, 10)$: in the notebook `Dirac_Einstein-Lovelock_4+4_Nash.nb` (In[133] to In[136]) the author sets $w_1 = 1$, $w_2 = 1/2^5$ and solves the vacuum equations on the history $a_4 = Hx_4$, $b_4 = 1$ of the notebook metric for w_3 and Λ^{nb} (In[134]); In[136] factors $1152 = 2^7\,3^2$; the Revision specification records the solution $w_3 = 1/1152$, $\Lambda^{\mathrm{nb}} = 10$. Multiplying by H^2 and writing the equations as $\sum_k \alpha_k {E_{(k)}}^M{}_N + \Lambda\,\delta^M_N = \kappa {T}^M{}_N$ gives the UM-2 form

$$\alpha_k = w_k H^{2-2k}, \qquad w = \Big(1, \frac{1}{32}, \frac{1}{1152}\Big), \qquad \Lambda = -\Lambda^{\mathrm{nb}} H^2 = -10H^2,$$

where

$${E_{(k)}}^M{}_N = -\frac{1}{2^{k+1}}\,\delta^{M A_1 B_1 \cdots A_k B_k}_{N C_1 D_1 \cdots C_k D_k}\, R^{C_1 D_1}{}_{A_1 B_1} \cdots R^{C_k D_k}{}_{A_k B_k},$$
$$\mathcal{L}_{(k)} = \frac{1}{2^k}\,\delta^{A_1 B_1 \cdots A_k B_k}_{C_1 D_1 \cdots C_k D_k}\, R^{C_1 D_1}{}_{A_1 B_1} \cdots R^{C_k D_k}{}_{A_k B_k},$$

$\delta^{\cdots}_{\cdots}$ being the generalised Kronecker delta (the determinant of ordinary deltas); $E_{(1)}$ is the Einstein tensor, $\mathcal{L}_{(1)} = R$, $\mathcal{L}_{(2)} = R^2 - 4R_{AB}R^{AB} + R_{ABCD}R^{ABCD}$. That the UM-2 normalisation reproduces the record's equations (which were checked against the notebook) is verified exactly (script `um2_geometry.py`, checks `record_equation`). UM-2 keeps the author's weights and Λ unchanged; an earlier model of this repository (UM-1) had changed w_3 and Λ.

4.4 The gamma matrices of $Cl(4,4)$

The author's notebooks (*Dirac_Einstein-Lovelock_4+4_Nash.nb* and *WaveFunctionUniverse-4+4-Einstein-Lovelock.nb*) build real 16×16 matrices from ($h = 1,2,3$; $p,q = 1,\dots,4$; ϵ the Levi-Civita symbol)

$$Q_a(h,p,q) = \epsilon_{hpq4}, \qquad Q_b(h,p,q) = \delta_{p4}\delta_{qh} - \delta_{ph}\delta_{q4}, \qquad s_h = Q_a - Q_b, \quad t_h = Q_a + Q_b,$$
$$\zeta_0 = I_8, \quad \zeta_{7-h} = \begin{pmatrix} 0 & t_h \\ -t_h & 0 \end{pmatrix}, \quad \zeta_h = \begin{pmatrix} 0 & s_h \\ s_h & 0 \end{pmatrix}, \quad \zeta_7 = \zeta_1\zeta_2\zeta_3\zeta_4\zeta_5\zeta_6, \quad \sigma = \begin{pmatrix} 0 & I_4 \\ I_4 & 0 \end{pmatrix},$$
$$\tau_0 = I_8, \quad \tau_A = \zeta_A, \quad \bar\tau_A = \sigma\,\tau_A^T\,\sigma, \qquad \gamma^{x_A} \equiv T16[A] = \begin{pmatrix} 0 & \bar\tau_A \\ \tau_A & 0 \end{pmatrix} \quad (A = 0,\dots,7).$$

They satisfy $\{\gamma^a, \gamma^b\} = 2\eta^{ab} I_{16}$ and the notebook identities $\frac{1}{2}(\tau_A\bar\tau_B + \tau_B\bar\tau_A) = \eta_{AB} I_8$, $\sigma = \tau_1\tau_2\tau_3 = \tau_4\tau_5\tau_6\tau_7$ (notebook Out[322] to Out[324]). Derived objects used by UM-2:

$$C = \gamma^{x_0}\gamma^{x_1}\gamma^{x_2}\gamma^{x_3} = \begin{pmatrix} -\sigma & 0 \\ 0 & \sigma \end{pmatrix}, \qquad \Gamma = \gamma^{x_0}\cdots\gamma^{x_7} = \mathrm{diag}(-I_8, I_8), \qquad B = -iC\gamma^{x_4}, \qquad S^{ab} = \tfrac{1}{4}[\gamma^a, \gamma^b].$$

4.5 The mass parameter, the x_0 mass idea and the dark-matter hypothesis

The dirac16complex mass is written $m = 3HM$ in the author's notebooks. The order requires every non-massless particle to get its mass from the dependence of its wave function on x_0, periodic with period $\pi/(6H)$, with the uncertainty relation $\Delta x_0\,\Delta p_{x_0} \geq \hbar/2$ as the physical reason, and states the hypothesis that the quanta of dirac16complex represent dark matter (and asks for the validity of a dark-energy association to be analysed).

5 Ingredients from the Revision record (RECORD)

5.1 The corrected primordial metric (version 2)

The author's order of 2026-10-16: "a diagonal form for the metric of the primordial gravitational field is only self-consistent if all off-diagonal terms of all source terms in the Einstein field equations ARE ZERO. This is not the case here. I task you to: [1] find the simplest modification for the current metric of the primordial gravitational field that is probably self-consistent with the Einstein field equations". The metric of 2026-10-14 (the notebook metric with b_4) has the off-diagonal component ${E_{(k)}}^{x_4}{}_{x_0} \propto \cot z\,\dot b_4/b_4$ at every Lovelock order. In the family (with $g_{44} = -1$)

$$g_{00} = b_4^{2p} e^{2qa_4}\cot^2 z, \qquad g_{ii} = e^{2a_4}(\sin^2 z)^{3\alpha}, \qquad g_{tt} = -e^{-2a_4} b_4 (\sin^2 z)^{3\beta}$$

the mixed curvature is proportional to $\alpha\big((q-1)\dot a_4 + p\,\dot b_4/b_4\big)$ (3-space) and $\beta\big((q+1)\dot a_4 + (p-\tfrac12)\dot b_4/b_4\big)$ (extra times). Keeping the warp of the extra times ($\beta = 1/6$) forces $\alpha = 0$, $q = -1$, $p = 1/2$: the record's current metric, used by UM-2,

$$\bar g = \mathrm{diag}\big(e^{-2a_4} b_4 \cot^2 z,\ e^{2a_4},\ e^{2a_4},\ e^{2a_4},\ -1,\ -e^{-2a_4} b_4 (\sin^2 z)^{1/6},\ -e^{-2a_4} b_4 (\sin^2 z)^{1/6},\ -e^{-2a_4} b_4 (\sin^2 z)^{1/6}\big).$$

Every off-diagonal component of $E_{(1)}, E_{(2)}, E_{(3)}$ vanishes identically for all $a_4(x_4)$, $b_4(x_4) > 0$ and every real z (record: sympy 14/14, Wolfram 13/13; re-derived in UM-2, `um2_geometry.py`). The alternative K of the record ($g_{00} = e^{2a_4}\cot^2 z$, $g_{ii} = e^{2a_4}(\sin^2 z)^{1/6}$, $g_{tt} = -e^{-2a_4} b_4$) is also diagonal at every order; the record leaves the choice to the author, and UM-2 uses version 2.

5.2 The hidden coordinate, the fold and the regular tip

On every interval with $\sin z \neq 0$,

$$y = \frac{\ln|\sin z|}{6H} \in (-\infty, 0], \qquad dy = \cot z\, dx_0, \qquad f = e^{-a_4} b_4^{1/2},$$

$$ds^2 = e^{2a_4} d\vec{x}^{\,2} - dx_4^2 + f^2\big[dy^2 - e^{2Hy}(dx_5^2 + dx_6^2 + dx_7^2)\big].$$

The fold is $z = \pi/2$ ($y = 0$); the tips are $z \to 0^+$ and $z \to \pi^-$ ($y \to -\infty$). The record replaces the tip by a regular cutoff at $y = -L$ with

$$\eta \equiv HL = 3,$$

a chosen cutoff (the extended levels depend on it like those of a box of proper length fL). UM-2 keeps $\eta = 3$ (ASSUMED).

5.3 The dirac16complex Lagrangian and the canonical spin connection

The specification of the record (under the author's orders) fixes the Dirac-type Lagrangian

$$\mathcal{L}_\Psi = \sqrt{|g|}\Big[\tfrac12\big(\bar\Psi\gamma^M D_M\Psi - (D_M\bar\Psi)\gamma^M\Psi\big) - m\,\bar\Psi\Psi - U(S)\Big], \qquad S = \bar\Psi\Psi, \quad U(S) = \frac{\lambda}{2}S^2, \quad \bar\Psi = \Psi^\dagger C,$$

$$D_M\Psi = \partial_M\Psi + \Omega_M\Psi, \quad \Omega_M = \tfrac12\omega_{Mab}S^{ab}, \quad {\omega_M}^a{}_b = e^a{}_N\big(\partial_M e_b{}^N + \Gamma^N{}_{ML} e_b{}^L\big), \quad \gamma^M = e_a{}^M\gamma^a,$$

the canonical (vielbein-postulate, torsion-free) spin connection. The record chose the Dirac-type form because the notebook's real Majorana-type Lagrangian is a total derivative for anticommuting real fields (it has no field equations), and it uses the canonical connection rather than the notebook's contraction of the mixed ${\omega_M}^a{}_b$, which lacks a metric factor. The field equation is $\gamma^M D_M\Psi = (m + U'(S))\Psi$.

5.4 Canonical quantisation and the Krein metric

With x_4 as the evolution time, $\{\Psi_r(x), \Psi_s^\dagger(x')\} = B_{rs}\,\delta^7(x-x')/\sqrt{|g|}$ (r, s spinor indices) on a slice $x_4 = \text{const}$; B has eigenvalues $+1$ (8 times) and -1 (8 times), so the inner product is indefinite (a Krein space). The record works in the good sector (no extra-time momentum), counts the positive-energy branch of every block as particles, and evaluates expectation values as $\langle \Psi^\dagger M \Psi \rangle = u^\dagger B M u$.

5.5 The Kohn-Sham reduction

For the corrected metric the record reduces the 16-component equation, with $\Psi = e^{ikx_1}\, e^{-\frac{3}{2}Hy}\, e^{a_4/2} b_4^{-1}\, V_\beta\, \chi(y, x_4)$ (k the momentum along x_1), to the 2×2 block Hamiltonians

$$h_j = j\Big[-\frac{i}{f}\sigma_1 \partial_y + M(y)\,\sigma_2 + e^{-a_4} k\,\sigma_3 \Big] + v(y),$$

in the blocks (j, s_2, s_3) of the commuting set $J = \gamma^{x_0}\gamma^{x_1}\gamma^{x_4}$, $K_1 = \gamma^{x_2}\gamma^{x_3}$, $K_2 = \gamma^{x_5}\gamma^{x_6}$ (eigenvalues j, is_2, is_3), with the contact-interaction mean fields of the record's Hartree-plus-exchange treatment of the contact interaction

$$e_{int} = \frac{15}{32}\lambda S^2 - \frac{1}{32}\lambda n^2, \qquad M_{eff} = m + \frac{15}{16}\lambda S, \qquad v = -\frac{\lambda}{16} n,$$

the fold conditions $\theta \in \{0, \pi\}$ (both parities filled, weight $w = 1/2$ each), the regular tip, and the densities carrying $P = e^{-3Hy}/\mathrm{Vol}_7$, with $\mathrm{Vol}_7 = (2\pi/\Delta k)^3 v_t f^4$ the normalisation volume of a cell (Δk the spacing of the momentum lattice, v_t the coordinate volume of the extra times, $a_4 = 0$; the proper seven-volume of a cell is $2\,\mathrm{Vol}_7(1 - e^{-3\eta})/(3H)$). The record's mass is $m = 3HM$ with $M = 1/3$, i.e. $m = H$; UM-2 keeps it (ASSUMED). Two reference states of the record (at $f = 1$, $a_4 = 0$: $N = 8$ with $\lambda = 157.7$, and $N = 592$ with $\lambda = 2.131$) serve as validation targets of the independent UM-2 solver.

5.6 Earlier work used only for comparison

The first universe model of this repository (UM-1, folder `Revision/universe_model`) is not an input of UM-2. UM-2 differs from it in keeping the author's weights and Λ, in the electroweak sector, in the number of generations and in the dark-matter candidate. Version 1 of UM-2 (2026-10-17, one generation) is superseded by version 2; its differences are listed in item 21 of the corrections.

6 Standard physics and measured values (STANDARD)

Every measured value enters UM-2 only as the target of a fit or as a comparison, never as a derived result.

input	value used	source	status
$W^\pm$ mass	80.3625 ± 0.0077 GeV	PDG 2026, OUR EVALUATION (pdgLive S043M)	verified
Z mass	91.1879 ± 0.0020 GeV	PDG 2026, OUR AVERAGE (pdgLive S044M)	verified
$\sin^2\theta_W$ ($\overline{MS}$, M_Z)	0.23122 ± 0.00004	PDG electroweak review	from memory
electroweak ρ parameter	$\rho_0 - 1 = 0.00038 \pm 0.00020$	PDG electroweak review (global fit)	from memory
charged-lepton masses	$m_e = 0.51099895000$ MeV, $m_\mu = 105.6583755$ MeV, $m_\tau = 1776.93 \pm 0.09$ MeV	PDG 2025 summary tables	verified
light-quark masses ($\overline{MS}$, 2 GeV)	$m_u = 2.16 \pm 0.07$ MeV, $m_d = 4.70 \pm 0.07$ MeV, $m_s = 93.5 \pm 0.8$ MeV	PDG 2025 summary tables	verified

input	value used	source	status
heavy-quark masses	$m_c(m_c) = 1.2730 \pm 0.0046$ GeV, $m_b(m_b) = 4.183 \pm 0.007$ GeV, $m_t = 172.56 \pm 0.31$ GeV (direct)	PDG 2025 summary tables	verified
neutrino oscillation parameters (normal ordering)	$\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{23} = 0.470$, $\sin^2\theta_{13} = 0.02215$, $\delta = 212°$, $\Delta m^2_{21} = 7.49 \times 10^{-5}$ eV^2, $\Delta m^2_{31} = 2.513 \times 10^{-3}$ eV^2; inverted ordering $\Delta\chi^2 = 6.1$; $J_{max} = 0.0333 \pm 0.0007$	NuFIT 6.0, arXiv:2410.05380 (JHEP 12 (2024) 216), Table 1, IC24 with SK atmospheric data	verified
lightest neutrino mass	$m_1 = 1.000$ meV	chosen	ASSUMED
sum of neutrino masses	$\sum m_\nu < 0.064$ eV (95 %, ΛCDM)	DESI DR2 Results II, arXiv:2503.14738	verified
CKM parameters	$s_{12} = 0.22501$, $s_{13} = 0.003732$, $s_{23} = 0.04183$, $\delta = 1.147$ rad; $J = 3.12 \times 10^{-5}$	PDG 2024 review "CKM quark-mixing matrix", global fit	verified
strong coupling	$\alpha_s(m_Z) = 0.1180$	PDG world average	from memory
Fermi constant, electroweak scale	$G_F = 1.1663788 \times 10^{-5}$ GeV^{-2}, $v = 246.22$ GeV	PDG	from memory
fine-structure constant	$\alpha^{-1}(0) = 137.035999$	CODATA	from memory
number of light neutrinos	$N_\nu = 2.9963 \pm 0.0074$	PDG 2025 (Janot and Jadach, Phys. Lett. B803 (2020) 135319)	verified
(superseded, version 1)	$N_\nu = 2.9840 \pm 0.0082$	LEP/SLD, Phys. Rept. 427 (2006) 257	verified
Z width and branching fractions	$\Gamma_Z = 2495.5 \pm 2.3$ MeV; invisible 20.000 %, hadrons 69.911 %, $\ell^+\ell^-$ 3.3658 %	PDG 2025 Z listing	verified
nucleon masses	$m_p = 938.272$ MeV, $m_n = 939.565$ MeV	PDG	from memory
$\hbar c$, k_B, N_A	$1.973269804 \times 10^{-16}$ GeV m, $8.617333262 \times 10^{-5}$ eV/K, $6.02214076 \times 10^{23}$	CODATA, SI	from memory
CMB temperature	$T_0 = 2.7255$ K	Fixsen 2009	from memory
N_{eff} in the Standard Model	3.0432 ± 0.0002 (three flavours, oscillations, NLO)	Cielo, Escudero, Mangano, Pisanti, arXiv:2306.05460	verified
N_{eff} by the method of temperatures	3.000, 3.011, 3.010, 3.039, 3.049, 3.048 (Table 1); couplings $g_{eL} = 0.727$, $g_{eR} = 0.233$, $g_{\mu L} = -0.273$, $g_{\mu R} = 0.233$; $f_a = 0.884$, $f_s = 0.829$	Escudero, arXiv:2001.04466	verified (2026-10-18)
finite-temperature QED pressure	P_2, P_3 of the $e^\pm\gamma$ plasma (formulas below)	Bennett et al., arXiv:1911.04504	from memory; massless limits reproduced to 10^{-16}

input	value used	source	status
N_{eff} measured	2.99 ± 0.17 (Planck 2018 + BAO); 2.86 ± 0.13 (ACT DR6)	Planck 2018 VI, A&A 641 (2020) A6; Calabrese et al., arXiv:2503.14454	verified
Planck 2018 base parameters	$\omega_b = 0.02237$, $\omega_c = 0.1200$, $100\,\theta_{MC} = 1.04092$, $\tau_{reio} = 0.0544$, $\ln(10^{10} A_s) = 3.044$, $n_s = 0.9649$	Planck 2018 results VI, Table 2	from memory
BAO	DESI DR2: 13 measurements of D_V/r_d, D_M/r_d, D_H/r_d (table below)	DESI collaboration, arXiv:2503.14738	values verified (2026-10-17) against Table 2 of arXiv:2603.26861; correlations from memory
dark-matter density	$\Omega_c h^2 = 0.120$	Planck 2018	from memory
Planck mass (freeze-out)	$M_{Pl} = G_N^{-1/2} = 1.22091 \times 10^{19}$ GeV		from memory
reduced Planck mass (radion decay)	$\bar{M}_{Pl} = (8\pi G_N)^{-1/2} = 2.435 \times 10^{18}$ GeV		from memory
angle of the gauge couplings	$\cos^2\theta_W = (m_W^2/m_Z^2)/\rho_0 = 0.776368$ (used for $\beta_Z = \beta_W/\cos^2\theta_W$ and in $\rho = \Lambda_W/(\Lambda_Z \cos^2\theta_W)$); the on-shell value $m_W^2/m_Z^2 = 0.776663$ for comparison	computed from the $W^\pm$, Z masses and ρ_0 above	computed
solar electron density	$n_e = 245\,N_A\,e^{-10.54\,r/R_\odot}$ cm^{-3} outside the core, at most $100\,N_A$ cm^{-3} (exponential fit)	after Bahcall	ASSUMED profile
matter of the Earth	$\rho = 2.6$ (T2K), 2.84 (NOvA), 2.848 (DUNE), 2.8 g/cm^3 (resonance); $Y_e = 0.5$	representative constant densities	ASSUMED
supernova envelope and neutrinospheres	$\rho(r) = 10^4\,(10^9\,\text{cm}/r)^3$ g/cm^3, $Y_e = 0.5$; $L = 6, 6, 4 \times 10^{52}$ erg/s, $\langle E\rangle = 11, 14, 15.5$ MeV (ν_e, $\bar{\nu}_e$, ν_x); $n_{\nu_e} = 10^{32}$ to 10^{33} cm^{-3}	representative values	ASSUMED
supernova explosion statistics with flavour conversion	failed fractions 25.6 % (none), 50.8, 77.4, 93.3, 96.4 % ($\rho_c = 10^9$ to 10^{13} g/cm^3); Salpeter-weighted 27.0, 47.6, 61.9, 73.9, 88.3 %	Gogilashvili and Tamborra, arXiv:2605.16504 (the author's file)	verified (read 2026-10-18)
fast flavour instability criterion	instability if and only if the ELN distribution has a crossing	Morinaga, Phys. Rev. D 105 (2022) L101301	from memory
direct-detection limit	LZ: 2.2×10^{-48} cm^2 at 40 GeV, scaled linearly with the mass	LZ 2024	from memory; the extrapolation is an ESTIMATE

input	value used	source	status
nucleon gluon content, strong coupling	$f_{TG}=0.92$, $\alpha_s=0.3$ (low scale)	standard	from memory; ESTIMATE
dark-energy density	$\rho_\Lambda \approx 2.5\times10^{-47}$ GeV^4		from memory
supernova dark-energy fits	constant $w=-0.764$; $(w_0,w_a)=(-0.861,-0.60)$	the author's private file (not in the repository)	AUTHOR

The DESI DR2 BAO measurements used (z; D_V/r_d; D_M/r_d; D_H/r_d; correlation r_{MH} of D_M and D_H):

tracer	z	D_V/r_d	D_M/r_d	D_H/r_d	r_{MH}
BGS	0.295	7.942 ± 0.075			
LRG1	0.510		13.588 ± 0.167	21.863 ± 0.425	-0.459
LRG2	0.706		17.351 ± 0.177	19.455 ± 0.330	-0.404
LRG3+ELG1	0.934		21.576 ± 0.152	17.641 ± 0.193	-0.416
ELG2	1.321		27.601 ± 0.318	14.176 ± 0.221	-0.434
QSO	1.484		30.512 ± 0.760	12.817 ± 0.516	-0.500
Lyα	2.330		38.988 ± 0.531	8.632 ± 0.101	-0.431

with

$$\chi^2=\sum_i (d_i-d_i^{th})^T C_i^{-1}(d_i-d_i^{th}),\qquad C_i=\begin{pmatrix}\sigma_M^2 & r_{MH}\sigma_M\sigma_H\\ r_{MH}\sigma_M\sigma_H & \sigma_H^2\end{pmatrix}$$

(d_i the measured and d_i^{th} the computed pair of D_M/r_d, D_H/r_d) for the anisotropic tracers and $(d-d^{th})^2/\sigma_V^2$ for BGS, where $D_M(z)=\int_0^z c\,dz'/H(z')$ (flat), $D_H=c/H(z)$, $D_V=(zD_M^2D_H)^{1/3}$.

Standard theory used: the Standard-Model gauge theory $SU(3)_c\times SU(2)_L\times U(1)_Y$ with three generations; QCD confinement of the quarks into hadrons (protons and neutrons); the PMNS and CKM matrices in the standard parametrisation

$$U=\begin{pmatrix}c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta}\\ -s_{12}c_{23}-c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23}-s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13}\\ s_{12}s_{23}-c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23}-s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13}\end{pmatrix}$$

($s_{ij}=\sin\theta_{ij}$, $c_{ij}=\cos\theta_{ij}$; the CKM matrix V has the same form) with the Jarlskog invariant $J=c_{12}s_{12}c_{23}s_{23}c_{13}^2s_{13}\sin\delta$; the flavour evolution $i\,d\psi/dt=[U\,\mathrm{diag}(m_i^2)U^\dagger/(2E)+\mathrm{diag}(V_{CC},0,0)]\psi$ with $V_{CC}=\sqrt{2}G_Fn_e$ (Wolfenstein, Mikheyev and Smirnov); the tree-level Z partial width $\Gamma(Z\to f\bar f)=N_c\,G_Fm_Z^3(g_V^2+g_A^2)/(6\sqrt{2}\pi)$, $g_V=T_3-2Q\sin^2\theta_W$ (with the $\overline{MS}$ value $\sin^2\theta_W=0.23122$), $g_A=T_3$; the radiative decay $\mathrm{BR}(\mu\to e\gamma)=\frac{3\alpha}{32\pi}|\sum_i U_{\mu i}^*U_{ei}m_i^2/m_W^2|^2$ for Dirac neutrinos; the freeze-out of a relic (Kolb and Turner); the trace-anomaly nucleon matrix element; the Einstein-Boltzmann equations of linear cosmological perturbations (solved by CAMB); the linearised equations of a dense neutrino gas (fast flavour conversion); Higgsless electroweak symmetry breaking by a non-linear sigma field and fermion localisation in an extra dimension (concepts only, no numbers).

7 Ingredients new in UM-2 (ASSUMED and INVENTED)

7.1 The metric as a power series (ASSUMED)

$$g_{MN}=\bar g_{MN}+\epsilon\,h^{(1)}_{MN}+\epsilon^2h^{(2)}_{MN}+\ldots,\qquad \epsilon\sim10^{-5}\ \text{(the CMB anisotropy)},$$
$$h^{(1)}_{MN}(x^\mu,x_0)=\sum_{n\ge0}h^{(1,n)}_{MN}(x^\mu)\,u_n(x_0),$$

with u_n the eigenfunctions of the x_0 operator; the flat $n=0$ term in the Newtonian gauge,

$$ds^2 = e^{2a_4}\big[(1-2\epsilon\Psi_N)\delta_{ij}+\epsilon h_{ij}^{TT}\big]dx_i dx_j - (1+2\epsilon\Phi)dx_4^2 + f^2(1+2\epsilon\sigma)\big[dy^2 - e^{2Hy}d\vec{x}_t^{\,2}\big],$$

carries the CMB, CNB and BAO physics; the radion σ and the $n\geq 1$ Kaluza-Klein modes are massive.

7.2 The block stabiliser (ASSUMED; its origin is OPEN)

$$S_{stab} = -\int d^8x\sqrt{|g|}\;\rho_b(f), \qquad \kappa\rho_b(f) = \kappa\rho_b^*\Big(\frac{f_*}{f}\Big)^8,$$

$$T^M{}_N[\text{stab}] = \text{diag}(p_0,p_3,p_3,p_3,-\rho_b,p_t,p_t,p_t), \qquad p_3=-\rho_b, \qquad p_0=p_t=\rho_b.$$

It is conserved ($\dot\rho_b = -3A(\rho_b+p_3) - 4h(\rho_b+p_0) = -8h\rho_b$, so $\rho_b\propto f^{-8}$), four-dimensionally Lorentz invariant ($p_3=-\rho_b$) and isotropic in the block ($p_0=p_t$). It was introduced because the author's weights and Λ admit no static block with attractive four-dimensional gravity without a source: the only sourceless static block with $A=0$ is $k_0 = 1/(4w_2)=8$, where the coefficient of the four-dimensional Einstein-Hilbert term is $G(k_0)=1-24w_2k_0+72w_3k_0^2=-1$ (repulsive), with $\Lambda_0=-24H^2\neq -10H^2$ (`um2_background.py`). The family $p_0=p_t=\nu\rho_b$, $\rho_b\propto f^{-4(1+\nu)}$, was also computed at $\nu = 1/4, 1/2, 3/4, 1, 5/4, 3/2, 2, 3$.

7.3 The Standard-Model fields in eight dimensions (ASSUMED structure; boundary conditions INVENTED)

$$S_{gauge} = -\frac{1}{4}\int d^8x\sqrt{|g|}\,g^{MP}g^{NQ}\Big(\frac{1}{\hat g_s^2}G^\alpha_{MN}G^\alpha_{PQ} + \frac{1}{\hat g^2}W^a_{MN}W^a_{PQ} + \frac{1}{\hat g'^2}B_{MN}B_{PQ}\Big),$$

$$S_f = \sum_{i=1}^{3}\sum_{\psi}\int d^8x\sqrt{|g|}\Big[\tfrac{1}{2}\big(\bar\psi_i\gamma^M D_M\psi_i - (D_M\bar\psi_i)\gamma^M\psi_i\big) - \mu_{\psi i}\bar\psi_i\psi_i\Big],$$

$$D_M = \partial_M + \Omega_M - iG^\alpha_M T^\alpha - iW^a_M\frac{\tau^a}{2} - iYB_M,$$

for three generations $Q_i=(u_i,d_i)_L$, u_{Ri}, d_{Ri}, $L_i=(\nu_i,\ell_i)_L$, ℓ_{Ri}, ν_{Ri} ($u_i=u,c,t$; $d_i=d,s,b$; $\ell_i=e,\mu,\tau$; hypercharges $1/6, 2/3, -1/3, -1/2, -1, 0$), each a 16-component spinor of the same Clifford algebra times its gauge indices, with one doublet bulk mass per sector ($\mu_{Li}=\mu_L$, $\mu_{Qi}=\mu_Q$; ASSUMED). With the selection projector

$$P_{sel} = \frac{1+\Gamma_4\gamma^{x_0}}{2}\,\frac{1-i\gamma^{x_5}\gamma^{x_6}}{2}, \qquad \Gamma_4 = i\gamma^{x_1}\gamma^{x_2}\gamma^{x_3}\gamma^{x_4},$$

the conditions at the fold and the tip are, for doublets ($\mu_\psi>0$), $P_{sel}(1-\gamma^{x_0})\psi=0$, $(1-P_{sel})(1+\gamma^{x_0})\psi=0$ at the fold and $(1-\gamma^{x_0})\psi=0$ at the tip, and for singlets ($\mu_\psi<0$) the same with $\gamma^{x_0}\to-\gamma^{x_0}$. The weak bosons are $W^\pm = (W^1\mp iW^2)/\sqrt{2}$, with electric charges $+e$ (W^+) and $-e$ (W^-).

7.4 The tip sigma field and the tip matrices (INVENTED for this geometry; no Higgs particle)

$$S_{tip} = \int_{y=-L} d^7x\sqrt{|g_{tip}|}\Big[-\frac{v_\Sigma^2}{4}\text{tr}\big((D_\mu\Sigma)^\dagger D^\mu\Sigma\big) - \sum_{i,j=1}^{3}\big(\bar L_i\,\Sigma\,(\Lambda_\ell)_{ij}\,\ell_{Rj} + \bar Q_i\,\Sigma\,(\Lambda_q)_{ij}\,q_{Rj} + \text{h.c.}\big)\Big], \qquad \Sigma\in SU(2),$$

$$D_\mu\Sigma = \partial_\mu\Sigma - iW^a_\mu\frac{\tau^a}{2}\Sigma + iB_\mu\Sigma\frac{\tau^3}{2}, \qquad \Sigma\to U_L\Sigma U_Y^\dagger, \quad U_Y = e^{i\theta\tau^3/2},$$

with $\ell_R=(\nu_R,\ell_R)$, $q_R=(u_R,d_R)$ arranged as $SU(2)_R$-type doublets and $\Lambda_\ell=\text{diag}(\Lambda_\nu,\Lambda_e)$, $\Lambda_q=\text{diag}(\Lambda_u,\Lambda_d)$ complex 3×3 generation matrices. On the zero modes they become the tip conditions $b_L=\kappa b_R$, $a_R=-\kappa^\dagger a_L$ with the tip matrices (ASSUMED values)

$$\kappa_\nu = U_{PMNS}, \qquad \kappa_e = 1, \qquad \kappa_u = 1, \qquad \kappa_d = V_{CKM}.$$

7.5 The portal interactions (INVENTED)

$$S_{I11} = \int d^8x\sqrt{|g|}\,\frac{c_V}{\Lambda_\star^6}(i\bar\Psi\gamma^M\Psi)\sum_{i=1}^{3}\sum_{\psi} q_\psi(i\bar\psi_i\gamma_M\psi_i),$$

$$S_{I12} = -\int d^8x\sqrt{|g|}\,\frac{\bar\Psi\Psi}{\Lambda_\star^3}\big[c_G G^\alpha_{MN}G^{\alpha MN} + c_W W^a_{MN}W^{aMN} + c_B B_{MN}B^{MN}\big].$$

Removed in version 2 (DERIVED reasons in item 21): the chiral tip Z portal of version 1,

$$S_{I9} = \int_{y=-L} d^7x\sqrt{|g_{tip}|}\big[c_Z + c'_Z\varsigma\big](i\bar\Psi\gamma^\mu P_+\Psi)\,\frac{iv_\Sigma^2}{2}\mathrm{tr}\big(\Sigma^\dagger D_\mu\Sigma\,\tau^3\big), \qquad P_+ = \frac{1+\Gamma_4}{2}, \quad \varsigma = \pm 1,$$

now $c_Z = c'_Z = 0$. Operators considered and rejected: a dipole portal $\bar\Psi S^{MN}\Psi B_{MN}$ (it would thermalise the sterile fold bands through the photon: too much dark radiation) and any operator coupling the neutral Ψ to a single W^+ or W^- (none is gauge invariant).

7.6 Parameter choices (ASSUMED)

$\eta = 3$ and $m = H$ (from the record); the universal tip mixing $\kappa_f = 1$ (unitary tip matrices); the doublet localisations fixed by the heaviest member of each sector, $c_R(\tau) = -c_L$ and $c_R(t) = -c_Q$; the lightest neutrino mass 1 meV and the normal ordering; $c_V = 0$ (benchmark); the Kohn-Sham benchmark self-coupling $\lambda_B = 7\times10^{-5}H^{-6}$ (mean fields below 5 % of m in the densest benchmark state); the extra-time coordinate volume v_t (it enters only $V_\perp$).

8 The computations and their verification (DERIVED)

All computations are Python scripts in the folder `universe_model_2/code`; each writes its numbers to `universe_model_2/outputs` and a report of named checks (name, verdict, detail) to `universe_model_2/reports`. In total 205 checks, none failed.

script	what it computes	method	checks
`um2_algebra.py`	the author's gamma matrices rebuilt from the notebook formulas; C, Γ, B; Pin(4,4) irreducibility; the two inequivalent Spin(4,4) halves; the 2×2 block basis and the block forms of every matrix the reductions use	sympy, exact	31/31
`um2_geometry.py`	Riemann tensor of $\bar g$ in two charts; Lovelock tensors by four routes (pairing formula, Ricci, explicit Gauss-Bonnet tensor, minisuperspace variation with a lapse); Bianchi identities; the record's equations; the static block; the canonical spin connection	sympy, exact	25/25

script	what it computes	method	checks
`um2_background.py`	the class-form background equations; the static block; the stabiliser and its family; the radion; the linear response of the block to matter sources; the dimensional reduction	sympy, exact; eigenvalues numerically	22/22
`um2_fields.py`	the exact x_0 reduction of the eight-dimensional Dirac equation (all blocks, general a_4, b_4, $M(y)$, $v(y)$); boundary conditions; zero modes; tip mixing for one and for three generations (self-adjointness with an arbitrary complex 3×3 matrix); the gauge x_0 operator; gauge invariance of the three-generation tip terms	sympy and numpy	16/16
`um2_spectra.py`	the x_0 spectra of gauge bosons and graviton; the $W^{\pm}$, Z fit to ρ_0 with the angle of the couplings fixed by the measured masses (both masses reproduced); the twelve fermion masses as exact x_0 eigenvalues	mpmath root finding; quadratic finite elements with Richardson; ODE shooting	14/14
`um2_flavour.py`	the PMNS and CKM matrices; the exact three-generation tip determinant and its null vectors; a general complex tip matrix; flavour-changing couplings of the Kaluza-Klein gauge modes; $\mu \to e\gamma$; lepton universality	mpmath (50 digits), numpy	7/7
`um2_neutrino.py`	vacuum and matter oscillations, the MSW resonances, long-baseline appearance, the Sun, the onset of oscillations in the early Universe, neutrino decoupling and N_{eff}, the CNB, supernova resonances, the fast flavour instability, the heating behind the shock	exact matrix exponentials and layer products; LSODA; Gauss-Legendre panels; vectorised Newton iterations	15/15

script	what it computes	method	checks
`um2_ks_jobs.py`	Kohn-Sham: validation against the record (2/2), free spectrum (3/3), interacting ground states $N = 8$ (3/3) and $N = 592$ (4/4), dark-matter quanta (3/3), portal responses with the three-generation sources (5/5), block feedback (2/2)	Galerkin in the free basis, safeguarded Anderson self-consistency, Richardson in the basis size	22/22
`um2_electroweak.py`	charges (W^+: $+e$, W^-: $-e$), anomalies of three generations, couplings, P_+, $\Gamma_4 = -B\gamma^{x_0}$, chiral tip charges of Ψ_A, Ψ_B, N_ν, the removal of the tip Z portal, the Z widths of three generations and their scheme dependence	sympy and numerics	19/19
`um2_primordial.py`	the nonlinear primordial evolution, the basin, the separatrix, the radion oscillation	DOP853 integration, bisection	5/5
`um2_cosmology.py`	TT and EE spectra with three massive neutrinos, BAO χ^2, N_{eff}, helium, the attribution of the differences, the variants with a thermalised fold band	CAMB 2.0.5	6/6
`um2_analysis.py`	the x_0 decompositions and the uncertainty relation on the cell (with its tip term); the gap theorem; sum rules; the twelve fermions against the uncertainty estimate; the conservation of the halves Ψ_A, Ψ_B; relic abundance of two species; direct detection; the condensate; the vacuum energy; the sensitivities to the block size (with the neutrino splittings); the bound on the block rate today; the radion decay	mpmath, numpy, sympy	23/23

8.1 The principal numbers and where they come from

quantity	value	computed by
static block k_*, f_*, $G(k_*)$, $\kappa\rho_b^*$	$12+\frac{4}{3}\sqrt{51}=21.5219$, 0.215556, $\frac{20}{3}+\sqrt{51}=13.8081$, $(26+4\sqrt{51})H^2$	`um2_background.py`
radion	$\omega_r^2/H^2=31552/2249-2416\sqrt{51}/6747=11.4721$, $m_r=3.387H=48.42$ TeV	`um2_background.py`, `um2_spectra.py`
unbroken x_0 spectrum ($\eta=3$)	$\Lambda=0$; $\Lambda_1=30.1196$; odd 27.0890; gap 20.25	`um2_spectra.py`
$W^{\pm}$, Z	$\beta_W=0.011912$, $\beta_Z=0.015343$ ($\cos^2\theta_W=0.776368$), $\Lambda_W=1.32149\times10^{-5}$, $\Lambda_Z=1.70150\times10^{-5}$, $\rho-1=0.00038$, model $m_Z=91.1879$ GeV; Dirichlet bound $\Lambda_D=0.010016$	`um2_spectra.py`
scales	$m_{KK}=121.32$ TeV, $H=14.296$ TeV, proper half cell $fL=8.926\times10^{-21}$ m, x_0 period 7.227×10^{-21} m, gap 99.48 TeV	`um2_spectra.py`
fermion localisations ($\kappa_f=1$)	doublets $c_L=5.9529$, $c_Q=3.3823$; singlets c_R: ν_1 -35.0452, ν_2 -32.8481, ν_3 -31.0702, e -14.5539, μ -8.9809, τ -5.9529, u -15.4300, c -8.7683, t -3.3823, d -14.6258, s -11.5158, b -7.5005	`um2_spectra.py`
three generations	tip determinant roots equal the one-generation roots to 3.0×10^{-42}, null vectors the columns of κ to 1.1×10^{-16}; general complex tip matrix: singular values to 3.5×10^{-5}	`um2_flavour.py`
mixing	PMNS ($\theta_{12}=33.71^\circ$, $\theta_{13}=8.56^\circ$, $\theta_{23}=43.28^\circ$, $\delta=212^\circ$) unitary to 1.1×10^{-16}, $J=-0.01777$; CKM $J=3.117\times10^{-5}$; Kaluza-Klein gauge overlaps 0.14434 (leptons), -0.30066 (quarks), off-diagonal 1.0×10^{-17}; $\mathrm{BR}(\mu\to e\gamma)=2.8\times10^{-55}$	`um2_flavour.py`
neutrino masses	1.000, 8.712, 50.140 meV; $\sum m_\nu=0.0599$ eV	`um2_spectra.py`, `um2_neutrino.py`
flavour conversion	1-3 MSW resonance 11.135 GeV (exact 11.133 GeV) at 2.8 g/cm^3; solar 1-2 resonance 1.93 MeV; $P(\nu_\mu\to\nu_e)$: T2K 0.0517, NOvA 0.0481, DUNE 0.0701 (antineutrinos 0.0313, 0.0236, 0.0195); solar P_{ee} from 0.5408 (pp) to 0.3043 (^{8}B, 15 MeV)	`um2_neutrino.py`
early Universe	oscillation onset $T=4.99$ MeV (atmospheric), 2.39 MeV (solar); $N_{eff}=3.0476$ (separate temperatures), 3.0485 (common); adopted 3.0432	`um2_neutrino.py`

quantity	value	computed by
CNB	$T_\nu = 1.9454$ K, 112.0 cm^{-3} per mass state, $\langle v \rangle = 0.435$, 0.0604, 0.0105; $\Omega_\nu h^2 = 6.382 \times 10^{-4}$	`um2_neutrino.py`
supernovae	H resonance (20 MeV) at 1573 g/cm^3, $\gamma = 178$; $P_{ee} = 0.02215$, $\bar{P}_{ee} = 0.677$; fast instability growth 0.0877μ and 0.0979μ, e-folding 5.9×10^{-10} to 5.9×10^{-11} s; heating factors 0.884, 0.760	`um2_neutrino.py`
dirac16complex at f_*	$c_\psi = 0.646667$; $\varepsilon_{DM} = 3.102204\,m$ ($p = 2.93661\,m$); even bulk $4.959981\,m$; x_0-momentum fraction 0.896	`um2_ks_jobs.py spectrum`
dark matter	$M_{DM} = 44.35$ TeV (both halves); $w = 0$; $p_0/\rho = 0.772034343$ $= -d\ln\varepsilon_{DM}/d\ln f$	`um2_ks_jobs.py darkmatter`
portal responses $d\varepsilon_{odd}/de$	I11 electron 0.824779, up quark 4.328803, neutrino 1.252406; I12 gluon condensate 0.707196	`um2_ks_jobs.py portals`
tip Z portal (removed)	$N_\nu = 3 + 2r^2$, $r^2 < 0.0056$; at the bound decoupling at 14.7 MeV and $\Delta N_{eff} = 2.02$; oriented part $\Delta N_{eff} = 0.0958$ to 0.1257	`um2_electroweak.py`
Z widths	invisible 497.7 MeV (measured 499.1), total 2422.3 MeV tree level, 2485.2 MeV with $1 + \alpha_s/\pi$ (measured 2495.5); $R_\ell = 20.83$ (measured 20.77)	`um2_electroweak.py`
primordial era	separatrix $k_c = 2.6983, 2.5991, 2.5301$; deflation factor up to 2.9166; radion $3.3868H$ measured	`um2_primordial.py`
cosmology	UM-2: $H_0 = 67.37$ km/s/Mpc, $\Omega_m = 0.3151$, $r_{drag} = 147.109$ Mpc, BAO $\chi^2 = 31.72$ (13 points; ΛCDM 32.55), TT/EE within 2.8×10^{-4} of ΛCDM	`um2_cosmology.py`
relic	two species, $x_f = 30.38$, $g_* = 105.75$, $\langle \sigma v \rangle = 2.01 \times 10^{-25}$ cm^3/s per species, $(8G_g^2 + 3G_W^2 + G_B^2)^{1/2} =$ (15.5 TeV)$^{-3}$	`um2_analysis.py`
direct detection	$\sigma_{SI} = 6.3 \times 10^{-53}$ cm^2 (gluon-only) against 2.4×10^{-45} cm^2	`um2_analysis.py`
dark energy	condensate tuned to $w_0 = -1$: $w_a = -6$, energy zero at $z = 0.260$; vacuum-energy ratio 5.5×10^{64}; $(H_0/H)^2 = 1.0 \times 10^{-92}$	`um2_analysis.py`

quantity	value	computed by
uncertainty relation	m/m_H: first KK 7.49, first odd 6.47, dark matter 2.29, $W^\pm$ 1.14×10^{-3}, Z 1.29×10^{-3}, top quark 3.2×10^{-3}, electron 5.5×10^{-9}, ν_1 1.1×10^{-17}; $\langle p^2\rangle\Delta\xi^2$: photon zero mode 0.497, Ψ fold zero mode 0.097 ($\varrho(\text{tip})=0.489$); $\sqrt{\langle p^2\rangle}\Delta\xi \geq (1-\varrho(\text{tip}))/2$ for every tabulated state	`um2_analysis.py`
sensitivities $d\ln X/d\ln f$	KK -1, m_W -0.501, fermions from -6.8 (t) to -41.0 (ν_1), m_e -20.5, M_{DM} -0.772, α and the gauge couplings -4, G_N $+2.05$; splittings Δm^2_{21} -77.5, Δm^2_{31} -74.0	`um2_analysis.py`
block rate today (ESTIMATE)	$\lvert h\rvert < 3.6\times10^{-8}H_0$ (from $\lvert\dot\alpha/\alpha\rvert<10^{-17}$/yr), $<7.1\times10^{-4}H_0$ (from $\lvert\dot G/G\rvert<10^{-13}$/yr)	`um2_analysis.py`
radion decay (ESTIMATE)	$\Gamma = C_r\, m_r^3/(4\pi\bar M_{Pl}^2)$, $T_{RH}=(90/(\pi^2 g_*))^{1/4}\sqrt{\Gamma\bar M_{Pl}}$ ($g_*=10.75$): $\tau = 0.43\,\text{s}/C_r$, $T_{RH}=1.85\sqrt{C_r}$ MeV (<4.7 MeV unless $C_r>6.5$)	`um2_analysis.py`
first-order block response	$s_{h3}=0.028666$, $s_{hp}=0.037320$, $q_h=-0.009555$; adiabatic $\delta k=(-0.1076\,\kappa\delta p_3-0.1400\,\kappa\delta\bar p+0.0359\,\kappa\delta\rho)/H^2$	`um2_background.py`

8.2 Methods, with the equations they use

- **Exact checks.** Symbolic identities are verified with sympy by simplifying the difference to zero; matrix identities entry by entry.
- **Boson eigenvalues.** With $\xi = y/L$, $u = e^{-3\eta\xi/2}\tilde u$, $q^2 = \Lambda - 9\eta^2/4$, the operator $-e^{-3\eta\xi}(e^{3\eta\xi}u')' = \Lambda u$ with $u'(0)=0$ (even) or $u(0)=0$ (odd) and $u'(-1)=\beta u(-1)$ has the eigenvalue conditions $\left[q^2+\frac{3\eta}{2}(\beta+\frac{3\eta}{2})\right]\sin q - \beta q\cos q = 0$ (even) and $q\cos q + (\beta+\frac{3\eta}{2})\sin q = 0$ (odd), solved with mpmath at 30 digits. Cross-check: quadratic finite elements for $\int e^{3\eta\xi}u'^2 d\xi + \beta e^{-3\eta}u(-1)^2 = \Lambda\int e^{3\eta\xi}u^2 d\xi$ on 100, 200, 400 elements, the eigenvalue taken from the Rayleigh quotient of the gradient form and extrapolated by Richardson.
- **Richardson extrapolation.** For a quantity with error $C\,n^{-p}$: $x \approx x_{2n} + (x_{2n}-x_n)/(2^p-1)$; the observed order is $\log_2[(x_n-x_{2n})/(x_{2n}-x_{4n})]$.
- **Fermion eigenvalues.** The exact condition $D(e) = e^2 s_L s_R - \kappa_f^2(C_L - c_L s_L)(C_R + c_R s_R)$, $s=\sinh w/w$, $C=\cosh w$, $w=\sqrt{c^2-e^2}$, solved with mpmath; the first-order solution $e = \kappa_f\zeta(c_L)\zeta(-c_R)$, $\zeta(c)^2 = c\,e^{-c}/\sinh c$ ($c_L>0$, $c_R<0$); cross-check by direct integration of the coupled (L,R) Dirac system (DOP853, relative tolerance 10^{-13}). With the doublet value fixed by the heaviest member of a sector ($c_R=-c_L$ for τ and t), each lighter c_R is found by root finding on the mass.
- **Three generations.** With the boundary values $(a_L,b_L) = (C_L - c_L s_L, -s_L e)$, $(a_R,b_R) = (s_R e, C_R + c_R s_R)$ at the tip, the conditions $b_L = \kappa b_R$, $a_R = -\kappa^\dagger a_L$ give the 3×3 problem $\left[\text{diag}(b_{L\alpha}) + \kappa\,\text{diag}(b_{Ri}/a_{Ri})\,\kappa^\dagger\,\text{diag}(a_{L\alpha})\right]\mathcal{A} = 0$, whose determinant is evaluated with mpmath at 50 digits and whose roots are found by the Illinois (modified regula falsi) method in a ±2 % bracket around the one-generation roots; the null vectors (right singular vectors of the smallest singular value) are compared with the columns of κ. For a non-unitary random tip matrix the masses are compared with the singular values of $\zeta(c_L)\,\kappa\,\text{diag}(\zeta(-c_{Ri}))$.
- **Flavour evolution.** $S = e^{-i\mathcal{H}L}$ by diagonalisation of the Hermitian $\mathcal{H}$ for constant density; the ordered product of layer operators (20000 and 40000 layers) through the Sun, followed by the incoherent sum $P_{ee} = \sum_i |U_{ei}|^2 P(\nu_i)$; the exact 1-3 resonance from $|U^m_{e3}|^2 = 1/2$ by Brent's method.

- **Neutrino decoupling.** The temperature equations of Escudero with the energy-transfer kernel $F(T_1,T_2) = 32 f_a (T_1^9 - T_2^9) + 56 f_s T_1^4 T_2^4 (T_1 - T_2)$ and the QED pressure $P_2 = -\frac{e^2 T^2}{12\pi^2} J_1 - \frac{e^2}{8\pi^4} J_1^2 + \frac{e^2 m_e^2}{16\pi^4} J_2$, $P_3 = \frac{e^3 T}{12\pi^4} I^{3/2}$, with $J_1 = \int dp\,(p^2/E) n_D$, $J_2 = \iint dp\,dq\,\frac{pq}{E_p E_q} \ln\!\left|\frac{p+q}{p-q}\right| n_D(p) n_D(q)$, $I = \int dp\,\frac{p^2+E^2}{E} n_D$, $n_D = 2/(e^{E/T}+1)$; Gauss-Legendre panels, the logarithmic singularity of J_2 removed by $q = p(1-s^2)$; LSODA from 20 MeV to 5 keV (relative tolerance 10^{-11}); $N_{eff} = (11/4)^{4/3}[(T_{\nu_e}/T_\gamma)^4 + 2(T_{\nu_\mu}/T_\gamma)^4]$.
- **Fast flavour instability.** The dispersion relation $\det \Pi = 0$, $\Pi^{\mu\nu} = \eta^{\mu\nu} + \int du/2\, G\, v^\mu v^\nu/(\omega - ku)$, solved for real k by vectorised Newton iterations with the analytic derivative from a grid of complex starting points; cross-checks against the analytic growth $\frac{1}{2}\sqrt{4I_1^2 - (I_0+I_2)^2}$ of the mode $K = \Lambda^3$ and the eigenvalues of the discretised homogeneous equations (300 nodes).
- **Kohn-Sham.** Orbitals of h_j are expanded in the exact eigenfunctions of the free problem of each sector (60, 120 or 240 functions, quadrature with ten points per function), the mean fields $M_{eff} = m + \frac{15}{16}\lambda S$, $v = -\frac{1}{16}\lambda n$ are iterated to self-consistency (tolerance 10^{-13}) with a safeguarded Anderson mixing and a fixed labelling of the occupied set; checks: the energy identity $E_{KS} = 2\,\mathrm{Vol}_7 \int_{-L}^{0} e^{3Hy}\rho\,dy$, the y-conservation law $\left[e^{3Hy} p_0\right]_{-L}^{0} = 3H \int_{-L}^{0} e^{3Hy} p_t\,dy$, Richardson in the basis size, the Hellmann-Feynman theorem $d\varepsilon/de = \langle\chi|\partial h/\partial e|\chi\rangle$ for every portal mean field, and the reproduction of the record's two reference states to 10^{-13} (energies, HOMO, LUMO) and 10^{-9} (integrated hidden pressure). The portal sources are the zero-mode densities $\varrho_c(y) = (2c/L) e^{2cy/L}/(1-e^{-2c})$: the electron $\frac{1}{2}(\varrho_{c_L} + \varrho_{-c_R(e)})$, the up quark $\frac{1}{2}(\varrho_{c_Q} + \varrho_{-c_R(u)})$, the neutrino ϱ_{c_L}.
- **Primordial era.** The 3-space and block components of the field equations are solved for $\dot A$, $\dot h$, with $\dot k = -2kh$ and the stabiliser $\kappa\rho_b \propto k^4$; the x_4 constraint fixes the initial A and is monitored (residual below 6×10^{-10}); the separatrix is found by bisection in the initial k.
- **Cosmology.** CAMB 2.0.5 with the Planck 2018 parameters above, θ_{MC} fixed (so H_0 is an output), $\ell_{max} = 2600$, lensing included. UM-2 has three massive eigenstates, each with the degeneracy $N_{eff}/3 = 1.0144$; CAMB gives each the density deg $\times \rho_{FD}(m'; T_\nu)$, so the masses passed are $m_i' = m_i\,\mathrm{deg}^{-1/4}$; because CAMB recovers them by inverting the relativistic density from ω_ν and the mass fractions, ω_ν and the fractions are iterated (damped exponent 0.8, 13 iterations) until the masses agree to 7.8×10^{-12}. The comparison model is the CAMB default (one neutrino of 0.06 eV, $N_{eff} = 3.044$).
- **Relic abundance.** For $\mathcal{L} = G\,\bar\chi\chi F_{\mu\nu}F^{\mu\nu}$ per abelian channel, $\sigma v = G^2 s(s-4M^2)/(4\pi) \to G^2 M^4 v^2/\pi$ (p-wave); in total $\sigma v = (M^4 v^2/\pi)(8G_g^2 + 3G_W^2 + G_B^2)$, $\langle\sigma v\rangle(x) = 6b/x$ with $b = (M^4/\pi)(8G_g^2 + 3G_W^2 + G_B^2)$; freeze-out (Kolb and Turner, Dirac fermion, $g=2$, $c=1/2$): $x_f = \ln\left[0.038\,c(c+2)\,g\,M_{Pl} M \langle\sigma v\rangle(x_f)/\sqrt{g_* x_f}\right]$, $\Omega h^2 = 2\times1.07\times10^9\,\mathrm{GeV}^{-1}\,x_f/\left(\sqrt{g_*}\,M_{Pl}\,3b/x_f\right) = 0.060$ per species (two species), with $M_{Pl} = 1.22091\times10^{19}$ GeV and $g_* = 105.75$ (three generations without ν_R, the gauge bosons; no Higgs; the fold bands not thermalised); the comparison scale is the s-wave thermal unitarity estimate $\langle\sigma v\rangle_{max} = (4\pi/M^2)\sqrt{x_f/\pi}$ (applied to a p-wave process).
- **Direct detection** (ESTIMATE). Trace anomaly: $\langle N|G^\alpha G^\alpha|N\rangle = -\frac{8\pi}{9\alpha_s} m_N f_{TG}$, $f_N = G_g \frac{8\pi}{9\alpha_s} m_N f_{TG}$, $\sigma_{SI} = \mu_N^2 f_N^2/\pi$ (μ_N the reduced mass).
- **Uncertainty relation.** For bosons, with $u = e^{-3\eta\xi/2}\tilde u$: $e^{3\eta\xi}u'^2 = \tilde u'^2 + \frac{9}{4}\eta^2\tilde u^2 - \frac{3}{2}\eta(\tilde u^2)'$, hence $(m\,fL)^2 = \langle p^2\rangle + \frac{9\eta^2}{4} - \frac{3\eta}{2}\varrho(\mathrm{fold}) + (\frac{3\eta}{2} + \beta)\varrho(\mathrm{tip})$ with $\langle p^2\rangle = \int \tilde u'^2 / \int \tilde u^2$ and the normalised density $\varrho = \tilde u^2/\int\tilde u^2$ ($\varrho = |\chi|^2/\int|\chi|^2$ and $\langle p^2\rangle = \int|\chi'|^2/\int|\chi|^2$ for fermions); the Heisenberg estimate is $m_H = \hbar c/(2\Delta s)$ with $\Delta s = fL\,\Delta\xi$, $\Delta\xi^2 = \int \xi^2 \varrho\,d\xi / \int \varrho\,d\xi$ the mean square distance from the fold. On the cell the relation reads $\sqrt{\langle p^2\rangle}\,\Delta\xi \geq (1 - \varrho(\mathrm{tip}))/2$, from $1 = \varrho(\mathrm{tip}) - \int \xi\varrho'\,d\xi$ and the Cauchy-Schwarz inequality (a theorem); it is also checked numerically for each of the 21 tabulated states (five boson modes, two doublet and twelve singlet zero modes, two states of Ψ).
- **$W^\pm$, Z fit.** $\rho(\beta_W) - 1 = \Lambda_W(\beta_W)/[\Lambda_Z(\beta_W/\cos^2\theta_W)\cos^2\theta_W] - 1$ with $\cos^2\theta_W = (m_W^2/m_Z^2)/\rho_0$ is solved for $\rho - 1 = \rho_0 - 1$ by bisection in $\ln\beta_W$ (80 steps); the model's $m_Z = m_W\sqrt{\Lambda_Z/\Lambda_W}$ then equals the measured value to 10^{-8} (checked).
- **Cosmic variance.** The relative error of a single multipole is $\sqrt{2/(2\ell+1)}$; the binned full-sky $\chi^2 = \sum_b (\bar\delta_b/\sigma_b)^2$ uses bins of $\Delta\ell = 50$, the mean relative deviation $\bar\delta_b$ and $\sigma_b = \sqrt{2/\sum_{\ell\in b}(2\ell+1)}$.

8.3 Corrections made during the work

1. The gauge-invariance test of the tip terms first used the wrong sign of the hypercharge phase; with $\psi \to e^{iY\theta}\psi$ every term is invariant to 4×10^{-15}.
2. The finite-element cross-check of Λ_W was first limited by round-off; the gradient-form Rayleigh quotient with

Richardson extrapolation (observed order 4.01) agrees with the exact value to 3×10^{-14}.

3. The Kohn-Sham basis convergence and the conservation law were first checked without extrapolation; Richardson in the basis size (observed orders 4.95 for the energy, 2.97 for the hidden pressure) was added.
4. The self-consistency loop did not converge for $N = 592$ at the record's coupling; a fixed labelling of the occupied set and a safeguarded Anderson mixing were introduced, the benchmark coupling λ_B was adopted, and the record's coupling is kept as a stress test, which now converges.
5. A first statement that every primordial start above the unstable static block reaches the stable one was false in detail; the separatrix $k_c(h_0)$ was computed by bisection instead.
6. The primordial figure was first sampled too coarsely (aliasing of the radion oscillation) and is now densely sampled.
7. The sum rules of the uncertainty analysis were first imprecise for the sharp fermion zero modes; exact mpmath normalisation and the tip term fixed them (10^{-8}).
8. The radion quantity was stored under a wrong sign label; it is now stored as $\omega_r^2 =$ det of the block Jacobian.
9. Portal names of the first universe model (P1 to P6) in two check texts were replaced by I9, I11, I12.
10. The gauge-kinetic portal was first written with the scale to the power 5, which is dimensionally wrong in the geometric normalisation of the gauge fields; it is $\bar{\Psi}\Psi\, F_{MN}F^{MN}/\Lambda_\star^3$, and the symbol of the portal scale was changed so that it no longer clashes with the Dirichlet bound Λ_D.
11. Before publication every number of the documents was compared with the computed outputs; four statements were corrected (the dark-matter momentum $p\,fL = 1.89901$, the finite-element agreement 5×10^{-10}, the TT and EE agreements of version 1, and the restriction $A = 0$ of the sourceless static block).
12. The kinetic term of the tip sigma field was first written with the sign of the signature $(+,-,-,-)$; in the signature $(+,+,+,-)$ of $x_1, \ldots, x_4$ it is $-\frac{v_\Sigma^2}{4}\mathrm{tr}((D_\mu\Sigma)^\dagger D^\mu\Sigma)$, which gives the mass terms $-\frac{1}{2}m^2 V_\mu V^\mu$ with $m^2 > 0$ (the computations always used $m^2 > 0$).
13. With the real matrices $C\gamma^a$ antisymmetric, $\bar{\Psi}\gamma^a\Psi$ is anti-Hermitian; the portal currents are now written as the Hermitian currents $i\bar{\Psi}\gamma^M\Psi$ and the Standard-Model currents as $i\bar{\psi}\gamma^M\psi$.
14. The block response to the hidden pressure of a Kohn-Sham state first used p_0 alone; the common block scale is driven by the block-averaged pressure $\bar{p} = (p_0 + 3p_t)/4$, and $p_0 - p_t$ drives a block-shape mode outside the order-0 metric (OPEN). The linearised sourced equations were added (`um2_background.py`, 2 checks); they first omitted the density, which enters at first order through $3G(k_*)A^2 = \kappa\delta\rho$; with it the adiabatic response reproduces the four-dimensional Friedmann acceleration equation exactly.
15. The statement that N_A and N_B are separately conserved was first recorded as an asserted check; it is now computed with the author's matrices (commutators of B with C, γ^a, $C\gamma^a$, S^{ab}), which also shows that extra-time momenta, outside the good sector, would mix the halves.
16. The radion lifetime was first quoted as an unverified estimate; it is now computed (ESTIMATE, with the coupling factor C_r explicit), as is the bound on the block rate today.
17. Figures: the BAO legend showed the data label twice and two labels of the uncertainty figure overlapped; the condensate figure no longer draws the vertical lines where w diverges.
18. An independent review of the five documents of version 1 against the outputs found, and the documents were corrected: the first-order fermion mass formula (it needs $\zeta(-c_R)$ for the singlet, $c_R < 0$; the code always used $|c|$); the gap statement, which holds for the boson operators but not for Ψ (the dark matter, 44.35 TeV, lies below the boson gap of 99.5 TeV); the sensitivity of G_N to the block, which is $d\ln G_N/d\ln f = -4 + 2k_*G'(k_*)/G(k_*) = +2.05$ and not -4 (recomputed; the lunar-ranging bound becomes $7.1 \times 10^{-4} H_0$); ΔN_{eff} of a thermalised fold band (recomputed); a level spacing (0.25000); a rounding (1.948); the definitions of M_{Pl}, M_{Pl}, the radion formulas, Δs, the unitarity estimate and the cosmic-variance measure; the statement on proper and coordinate localisation; and several notational clashes ($\kappa = e^{-a_4}$ is now written out).
19. A second review of version 1 found, and the documents were corrected: the electroweak parameter was written $\rho = m_W^2/(m_Z^2\cos^2\theta_W)$ next to the on-shell definition $\cos^2\theta_W = m_W^2/m_Z^2$, which makes it identically 1; the computation always used $\rho = \Lambda_W/(\Lambda_Z\cos^2\theta_W)$. Checking the uncertainty relation for every state then showed that the bound $\langle p^2\rangle \geq 1/(4\Delta\xi^2)$ quoted for the hidden kinetic energy is not valid on the cell: the tip, where the cell ends, adds a boundary term, and the valid relation is $\sqrt{\langle p^2\rangle}\,\Delta\xi \geq (1 - \varrho(\mathrm{tip}))/2$ (the broad fold zero mode of Ψ has $\langle p^2\rangle\Delta\xi^2 = 0.097$); a computed check of it was added. Further: the $\overline{MS}$ angle in the Z widths, the shooting accuracy, the normalisation volume Vol_7 (not the proper seven-volume), the occupations f_o, the coupling matrix Λ_ℓ, and the notation (with a list of the symbols with more than one use in each document).
20. A third review of version 1 found, and the computations and documents were corrected: (a) the $W^\pm$, Z fit set the angle of the gauge couplings equal to the on-shell angle of the measured masses and fitted the

tree-level $\rho - 1$ to $\rho_0 - 1$, which put the model's $m_Z = 91.1706$ GeV 17 MeV below the measured value; since ρ_0 measures the custodial violation beyond the Standard-Model loops, the angle of the couplings is now fixed by the measured masses, $\cos^2\theta_W = (m_W^2/m_Z^2)/\rho_0 = 0.776368$, and the model reproduces both masses (check `mZ_reproduced`); $\beta_W = 0.011912$ and every scale in GeV moved by $+8.50 \times 10^{-4}$ ($m_{KK} = 121.32$ TeV, $H = 14.296$ TeV, gap 99.48 TeV, radion 48.42 TeV, $M_{DM} = 44.35$ TeV). The tree-level Z widths with the angle of the couplings were added (check `Z_width_scheme`). (b) The familiar bound $\langle p^2 \rangle \geq 1/(4\Delta\xi^2)$ is guaranteed only for profiles negligible at the tip. (c) Symbol lists and wording.

21. **Version 2 (2026-10-18): three generations and three neutrino flavours.** By the author's orders of 2026-10-18 (option A):
    (a) three generations of quarks and leptons with one doublet bulk mass per sector, the twelve masses as exact x_0 eigenvalues and the PMNS and CKM matrices as the tip matrices, proved exact for three generations (`um2_flavour.py`, new; `um2_spectra.py`, `um2_fields.py` extended); (b) the flavour conversion of the three neutrinos (`um2_neutrino.py`, new): vacuum, matter, accelerators, the Sun, the early Universe and N_{eff}, the CNB of three mass states, supernovae; (c) the chiral tip Z portal I9 is removed: with three neutrinos it has no role left, its even part is bounded by the invisible width ($N_\nu = 3 + 2r^2$, $r^2 < 0.0056$) and would still keep the fold band in equilibrium to 14.7 MeV ($\Delta N_{eff} = 2.02$), and its oriented part would thermalise the fold band through the decays of the odd Ψ_A states ($\Delta N_{eff} = 0.0958$ to 0.1257, 2.1 to 2.4 standard deviations above ACT DR6); consequently both halves of Ψ keep an exact fold parity and the dark matter consists of the lightest odd states of both halves (two species, relic recomputed: $x_f = 30.38$, $(15.5\ \text{TeV})^{-3}$); (d) the invisible width uses the PDG 2025 value $N_\nu = 2.9963 \pm 0.0074$ instead of the LEP 2006 value; (e) N_{eff} is computed for three flavours (3.0476 to 3.0485) and the precision value 3.0432 (Cielo et al.) replaces 3.044; (f) the cosmology has three massive neutrinos ($\sum m_\nu = 0.0599$ eV) instead of one of 0.05 eV and no dark radiation; (g) the Z widths of three generations (2485.2 MeV with the QCD factor) remove the one-generation conflict (1232 MeV); (h) the Kohn-Sham portal sources use the three-generation localisations. Errors found and corrected during the version-2 work: the 1-3 MSW resonance energy was first evaluated with a spurious factor $1/c_{13}^2$ (11.49 GeV); it is $E = \Delta m_{ee}^2 \cos 2\theta_{13}/(2V_{CC}) = 11.135$ GeV with $\Delta m_{ee}^2 = c_{12}^2\Delta m_{31}^2 + s_{12}^2\Delta m_{32}^2$, confirmed by the exact three-flavour root 11.133 GeV (now a check); CAMB's built-in normal-hierarchy option put the mass in one eigenstate, so the three eigenstates are now set up explicitly with the iteration described in the methods; a check of the analysis (fermions below the uncertainty estimate) first used a threshold that the muon and the strange quark exceed and now tests the documented statement ($m/m_H < 0.01$ for every fermion); in the documents the muon eigenvalue (4.7795×10^{-6}), the Landau-Zener bound ($P_{LZ} \leq 2.6 \times 10^{-77}$) and the explanation of $\Omega_\nu h^2$ (the non-relativistic coefficient 94.06 eV at the instantaneous-decoupling temperature, 93.14 eV with the heating) were corrected; the published values of the decoupling method were read from the source (Escudero 2020, Table 1) and the separate-temperature value 3.048 added to the validation.

8.4 What is not in the repository, and why

By the author's instruction the following private inputs are never committed (they are git-ignored): the prompt files, the author's private PDF with the supernova dark-energy fits, the folders `dirac-main/` and `vendor/`, the archive `dirac-main_2.zip`, the files `Generalized_Kronecker_Delta.*`, and the notebook `WaveFunctionUniverse-4+4-Einstein-Lovelock.nb` with its `.mx` files. The paper "Neutrino Flavor Conversion.pdf" (Gogilashvili and Tamborra, arXiv:2605.16504) is a third-party publication and is not committed either; UM-2 cites its results. UM-2 uses from these inputs only what is written out in this document (the notebook formulas of the metric and of the gamma matrices, the two supernova fits, the explosion statistics).

8.5 Reproduction

From the repository root, with Python 3 and the packages sympy, numpy, scipy, mpmath, matplotlib and camb installed (`pip install sympy numpy scipy mpmath matplotlib camb`), the commands

```
python universe_model_2/code/um2_algebra.py
python universe_model_2/code/um2_geometry.py
python universe_model_2/code/um2_background.py
python universe_model_2/code/um2_fields.py
python universe_model_2/code/um2_spectra.py
```

```
python universe_model_2/code/um2_flavour.py
python universe_model_2/code/um2_neutrino.py
python universe_model_2/code/um2_ks_jobs.py validate
python universe_model_2/code/um2_ks_jobs.py spectrum
python universe_model_2/code/um2_ks_jobs.py ground8
python universe_model_2/code/um2_ks_jobs.py ground592
python universe_model_2/code/um2_ks_jobs.py darkmatter
python universe_model_2/code/um2_ks_jobs.py portals
python universe_model_2/code/um2_ks_jobs.py feedback
python universe_model_2/code/um2_electroweak.py
python universe_model_2/code/um2_primordial.py
python universe_model_2/code/um2_cosmology.py
python universe_model_2/code/um2_analysis.py
python universe_model_2/tools/build_docs.py
```

rerun every computation in dependency order (each exits with code 0 if and only if all its checks pass) and rebuild the PDF documents from the Markdown sources (pandoc and a TeX Live installation with the `lmodern` package are needed for the last command). Version 1 was rerun from fresh clones of the repository (2026-10-17 at commit `b3c3c1a`; 2026-10-18 at commits `fa7305f`, `9285bb6` and `5b73d16`): every check passed and every stored number was reproduced to a relative 10^{-9} or better (quantities at the round-off level differ in their last digits because the threaded linear algebra does not sum in a fixed order), and the LaTeX sources, the PDF documents and the figures were rebuilt byte-identical. The document build is deterministic: no dates in the LaTeX sources, PDF dates and trailer identifiers suppressed, a fixed `SOURCE_DATE_EPOCH`.

Software used: Python 3.13.16, sympy 1.14.0, numpy 2.5.3, scipy 1.18.1, mpmath 1.3.0, matplotlib 3.11.2, camb 2.0.5, pandoc 3.1.3, pdfTeX 1.40.25 (TeX Live 2023).

8.6 History in the repository

UM-2 was started on top of commit `0cb4916`. Version 1: its computations were committed in the order algebra and geometry (`ca523d9`), background gravity (`0b3e10d`), field reductions (`6fc35a0`), mass spectra (`dcfd528`), Kohn-Sham (`2f3018e`), electroweak sector (`2e4c48d`), primordial era (`a502e62`), cosmology (`717b46e`), analysis (`a06b55e`), followed by the corrections listed above and the documents MODEL, PROVENANCE, FIELD_EQUATIONS, SOLUTION and ANALYSIS (Markdown, LaTeX and PDF each); the fixes of the three independent reviews are in `b386446`, `fa7305f`, `b0e686e`, `9285bb6`, `d92a61b`, `c98de95`, `1561566`, `df358db` and `5b73d16`; a checklist for manual reruns and the comparison tool are in `7b86239`. Version 2: three generations and the masses and mixing (`94621af`), the neutrino flavour conversion (`a0d750a`), the electroweak sector of three generations (`cf824d5`), the removal of the tip Z portal and the cosmology with three massive neutrinos (`53c5139`), the dark matter of both halves, the twelve-fermion tables and the three-generation tip and gauge checks (`b755990`), the corrected 1-3 resonance (`b999f57`), and the documents MODEL (`9a5b92b`), FIELD_EQUATIONS (`79d9d31`), SOLUTION (`a3dcc6a`), ANALYSIS (`df6d94a`) and PROVENANCE (this version).