The stabiliser of the bulk

What the stabiliser of UM-3 is, why the field equations of the author's metric require it, what it does and what it does not do, and why UM-3 needs it: a complete account for a newcomer, with its provenance

The stabiliser: found and named by Claude (Anthropic), 2026-10-24

This account and its checks: Claude (Anthropic), 2026-10-25

Underlying theory, metric and Lovelock weights: Patrick L. Nash

Repository once-ere/Dirac_claude, folder universe_model_3

Contents

1 The fold: definition and examples **2**

2 How to read this document **9**
2.1 Who this document is for . . . 9
2.2 The status labels . . . 10
2.3 Every equation is followed by its symbols . . . 10
2.4 The checks . . . 10

3 The stabiliser in brief **10**

4 The language **11**
4.1 Coordinates, distances and the metric . . . 11
4.2 Warp and warp factor . . . 12
4.3 The bulk . . . 13
4.4 Energy, pressure and the energy-momentum tensor . . . 13
4.5 The field equations of gravity . . . 13

5 The static bulk **17**
5.1 What static means . . . 17
5.2 Static is not the same as empty . . . 18
5.3 Examples of static spacetimes and static media . . . 18

6 The stabiliser of the author's original weights **19**
6.1 How the static block fixes the source . . . 19
6.2 The stabiliser . . . 21
6.3 Why it is called the stabiliser, and who found it . . . 22
6.4 What kind of medium it was . . . 23
6.5 How it would have evolved on its own . . . 24
6.6 What the original equations required when the universe expanded . . . 24
6.7 The constraints of the original equations . . . 27

7 The correction of the weights removes the stabiliser **28**
7.1 Gravitons must travel at the speed of light . . . 28
7.2 The static block with the corrected weights . . . 30
7.3 The expanding fold with the corrected weights . . . 31
7.4 What takes the place of the stabiliser . . . 34

8 The five points, verified **35**

9 The importance of the stabiliser for UM-3 **36**

10 Provenance **38**

1 The fold: definition and examples

Definition. The *fold* is the seven-dimensional surface of the author's eight-dimensional spacetime on which the cosine of the angle $z = 6Hx_0$ of the hidden direction vanishes:

$$\mathcal{F} = \big\{X = (x_0, x_1, \ldots, x_7) :\ \cos(6Hx_0) = 0\big\}, \qquad \text{in one period } 0 < z < \pi \text{ of the metric: } z = \frac{\pi}{2}, \quad x_0 = \frac{\pi}{12H} \tag{FD-1}$$

Symbols in (FD-1).

- $\mathcal{F}$: The fold: the seven-dimensional surface $\cos(6Hx_0) = 0$ of the eight-dimensional spacetime (FD-1), in one period $0 < z < \pi$ the surface $z = \pi/2$, $x_0 = \pi/(12H)$, where $\sigma = 0$; $|_{\mathcal{F}}$ means restricted to the fold. Status: DERIVED.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$, the fold is $z = \pi/2$. Unit: dimensionless. Status: AUTHOR.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

$X = \{x_0, \ldots, x_7\}$ are the author's coordinates: x_0 the hidden space-like direction, x_1, x_2, x_3 ordinary space, x_4 the time, x_5, x_6, x_7 the three extra times; H is the author's constant (an inverse length). The author's metric depends on x_0 only through $\cot^2 z$ and $(\sin^2 z)^{1/6}$, so it repeats with the period $\pi/(6H)$ of x_0, and every period $0 < z < \pi$ contains exactly one surface of $\mathcal{F}$, at $z = \pi/2$: $\mathcal{F}$ is the union of these copies, one per period (example 2), and "the fold" without a qualifier is the copy in the period that contains our world. A point of the fold is labelled by the seven coordinates $x_1, \ldots, x_7$, and the metric restricted to the fold ($\mathrm{d}x_0 = 0$, $z = \pi/2$) is

$$g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N\big|_{\mathcal{F}} = \mathrm{e}^{2a_4}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{d}x_4^2 - \mathrm{e}^{-2a_4}b_4^2\big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big) \tag{FD-2}$$

Symbols in (FD-2).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate

with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.

- $\mathcal{F}$: The fold: the seven-dimensional surface $\cos(6Hx_0) = 0$ of the eight-dimensional spacetime (FD-1), in one period $0 < z < \pi$ the surface $z = \pi/2$, $x_0 = \pi/(12H)$, where $\sigma = 0$; $|_{\mathcal{F}}$ means restricted to the fold. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.

with three space-like and four time-like directions; a_4 and b_4 are the author's two functions of the time x_4. Notation: ln is the natural logarithm, e Euler's number, $\exp(y) = \mathrm{e}^y$, sin, cos and $\cot = \cos/\sin$ the trigonometric functions, tanh the hyperbolic tangent, $|y|$ the absolute value; $f(0^+)$ and $f(0^-)$ are the limits of $f(\sigma)$ as σ approaches 0 from above and from below.

Why this surface is called a fold. Measured by the proper distance along the hidden direction,

$$\sigma = \mathrm{sgn}\Big(\frac{\pi}{2} - z\Big)\frac{\ln\sin z}{6H}, \qquad \mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0, \qquad \sin z = \mathrm{e}^{-6H|\sigma|} \tag{FD-3}$$

Symbols in (FD-3).

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- sgn: The sign function: $\mathrm{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$, the fold is $z = \pi/2$. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

(the fold is $\sigma = 0$; the half-period $0 < z < \pi/2$ is $\sigma < 0$, the half-period $\pi/2 < z < \pi$ is $\sigma > 0$; the two ends of the period, the tips $z \to 0$ and $z \to \pi$, lie at $\sigma \to -\infty$ and $\sigma \to +\infty$), the author's metric is

$$g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N = \mathrm{d}\sigma^2 + \mathrm{e}^{-2H|\sigma|}\Big[\mathrm{e}^{2a_4}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{e}^{-2a_4}b_4^2\big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big)\Big] - \mathrm{d}x_4^2 \tag{FD-4}$$

Symbols in (FD-4).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\ldots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.

It depends on σ only through $|\sigma|$. The function $\mathcal{W}(\sigma) = \mathrm{e}^{-2H|\sigma|}$, which multiplies the squared steps of ordinary space and of the extra times, is the warp factor of the metric (its square root $\mathrm{e}^{-H|\sigma|}$ multiplies the lengths); the time x_4 is not warped. Four properties make the surface $\sigma = 0$ a fold, and together they single it out within the period:

1. *It is a mirror.* The reflection $\sigma \to -\sigma$ (that is $z \to \pi - z$, or $x_0 \to \pi/(6H) - x_0$) maps the metric to itself, and the fold is the only surface of the period that the reflection leaves in place: the two halves of the period are mirror images of each other, joined along the fold, as the two halves of a folded sheet of paper are joined along the crease.
2. *The warp factor is largest on it.* The warp factor $(\sin^2 z)^{1/6} = \mathrm{e}^{-2H|\sigma|}$ equals 1 on the fold and decreases on both sides.
3. *The metric has a kink on it.* The metric is continuous across the fold, but its first derivative along σ jumps. The extrinsic curvature of the surfaces $\sigma = \text{const}$ changes sign across the fold,

$$K^a{}_b = \frac{1}{2}\,g^{ac}\,\partial_\sigma g_{cb}: \qquad K^1{}_1 = K^2{}_2 = K^3{}_3 = K^5{}_5 = K^6{}_6 = K^7{}_7 = -H\,\mathrm{sgn}(\sigma), \qquad K^4{}_4 = 0 \tag{FD-5}$$

Symbols in (FD-5).

- $K^a{}_b$: The extrinsic curvature of a surface $\sigma = \text{const}$ of the hidden direction, $K^a{}_b = \frac{1}{2}g^{ac}\partial_\sigma g_{cb}$ for a, b along the surface (FD-5): $-H\,\mathrm{sgn}(\sigma)$ along ordinary space and the extra times, 0 along the time. Unit: GeV (1/length). Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric

$\eta_{ab} = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1,\dots,7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.

- b: A frame index (as a). Status: STANDARD.
- c: A frame index (as a). Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\dots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- ∂: A partial derivative: ∂_M with respect to the coordinate X^M, ∂_μ with respect to x^μ, ∂_σ with respect to σ, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- sgn: The sign function: $\mathrm{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.

(a, b frame indices along the fold); it jumps by $-2H$ along ordinary space and along the extra times. The Einstein-Lovelock equations of UM-3, $\sum_p \alpha_{(p)}\mathcal{E}_{(p)}{}^a{}_b + \Lambda\delta^a{}_b = \kappa\mathcal{T}^a{}_b$ ($\mathcal{E}_{(p)}$ the Lovelock tensor of order $p = 1, 2, 3$, $\alpha_{(p)} = w_{(p)}H^{2-2p}$ its coupling with the weights $w = (1, 1/20, 1/360)$, $\Lambda = -7H^2$ the cosmological constant, $\kappa = 8\pi G_8$ the gravitational coupling of eight dimensions, $\mathcal{T}^a{}_b$ the energy-momentum tensor, $\delta^a{}_b$ the Kronecker delta), are linear in the second σ-derivatives of the metric, so the jump puts a Dirac delta function $\delta_{\mathrm{D}}(\sigma)$ into them: the metric solves them only if the fold carries a thin source $\mathcal{S}^a{}_b$, an energy-momentum tensor concentrated on $\sigma = 0$, given by the integral of the equations across the fold (the junction conditions),

$$\kappa\,\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_{p=1}^{3} \alpha_{(p)}\,\mathcal{E}_{(p)}{}^a{}_b \tag{FD-11}$$

Symbols in (FD-11).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+}\mathrm{d}\sigma\sum_p \alpha_{(p)}\mathcal{E}_{(p)}{}^a{}_b$ (FD-11, the junction conditions); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -\frac{32}{5}H\delta^a{}_b$ at the static block. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \dots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1,\dots,7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations a second label of a Kohn-Sham orbital (as o, with δ_{op} the Kronecker delta of the two labels). Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)}H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- $\mathcal{E}_{(p)}{}^M{}_N$: The Lovelock tensor of order p: the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to the metric ($\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor); with frame indices its frame components. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.

with $\mathcal{S}^4{}_4 = -$(its energy density), $\mathcal{S}^1{}_1 = \mathcal{S}^2{}_2 = \mathcal{S}^3{}_3$ its pressure along ordinary space and $\mathcal{S}^5{}_5 = \mathcal{S}^6{}_6 = \mathcal{S}^7{}_7$ its pressure along the extra times; $\mathcal{S}$ is $\mathcal{T}$ integrated across the fold, an energy per unit volume of the six other directions

of the fold (unit GeV^7). On the static block (a_4 and b_4 constant) and on an expanding fold whose extra times are frozen (b_4 proportional to e^{a_4}; $\mathsf{A} = a_4'/H$, $\mathsf{A}_1 = a_4''/H^2$)

$$\begin{aligned}
&\text{static block:}\quad \kappa\mathcal{S}^a{}_b = -\frac{32}{5}H\,\delta^a{}_b \quad \text{(a pure tension)},\\
&\text{frozen extra times:}\quad \kappa\mathcal{S}^1{}_1 = -\frac{16}{15}\big(6+3\mathsf{A}^2+2\mathsf{A}_1\big)H,\quad \kappa\mathcal{S}^4{}_4 = -\frac{16}{5}\big(2+\mathsf{A}^2\big)H,\\
&\qquad\qquad \kappa\mathcal{S}^5{}_5 = -\frac{4}{5}\big(8+8\mathsf{A}^2+\mathsf{A}^4+4\mathsf{A}_1+\mathsf{A}^2\mathsf{A}_1\big)H
\end{aligned} \tag{FD-12}$$

Symbols in (FD-12).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)}\mathcal{E}_{(p)}{}^a{}_b$ (FD-11, the junction conditions); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -\frac{32}{5}H\delta^a{}_b$ at the static block. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with upper and lower index strings the generalised Kronecker delta $\delta^{M_1\cdots M_n}_{N_1\cdots N_n}$, the determinant of the $n \times n$ matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); 1 at the author's vacuum, which is not a solution with the corrected weights.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless.

the tension plus matter that obeys the Friedmann equations of general relativity. The fold is therefore not only a place but a physical object: a thin layer that carries energy and momentum (a brane). 4. *The author's coordinate x_0 degenerates on it.* The metric component $g_{x_0x_0} = \cot^2(6Hx_0)$ vanishes on the fold and nowhere else in the period; near the fold the proper distance grows with the square of the coordinate distance,

$$|\sigma| = 3H\Big(x_0 - \frac{\pi}{12H}\Big)^2\Big[1 + 6H^2\Big(x_0 - \frac{\pi}{12H}\Big)^2 + \ldots\Big] \tag{FD-6}$$

Symbols in (FD-6).

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

so the two coordinate values on either side of $x_0 = \pi/(12H)$ at the same coordinate distance lie at the same proper distance from the fold: the coordinate line of x_0 is folded onto the line of proper distances (a fold map, example 5).

What lives on the fold. In UM-3 the Standard Model (the Higgs doublet with the author's ad hoc vacuum, a time-dependent choice of its vacuum value; the gauge fields of $SU(3)_C \times SU(2)_L \times U(1)_Y$; three generations of quarks and leptons and three right-handed neutrinos) and the fold fluid (its tension and the pressure along the extra times that keeps their volume fixed) live on the fold: they are fields at $\sigma = 0$, functions of $x_1, \ldots, x_4$ (no field depends on the extra times). Integrated over the extra times, the fold is the four-dimensional universe that we observe, with the metric $\mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in $x_1, \ldots, x_4$. dirac16complex (the author's spinor field of eight dimensions) and gravity live in the bulk $\sigma \neq 0$ as well as on the fold; the bound states χ of dirac16complex are the part of it that the fold holds, and the portals (couplings between dirac16complex and Standard-Model fields) act at $\sigma = 0$. In UM-3 files "the fold" means this surface; the examples below use the word for other surfaces with the same properties. The word is not used in its other senses: a set of equal energy levels is called degenerate, and a growth by the factor e is called an e-folding.

Examples of folds that are meaningful for UM-3.

1. *Our fold.* The fold $z = \pi/2$ of the period that contains our world (FD-1): the seven-dimensional world-volume of the Standard Model, and after the integration over the extra times the four-dimensional universe that we observe.
2. *The copies of the fold.* Because the metric repeats with the period $\pi/(6H)$, every surface $x_0 = \pi/(12H) +$ (an integer multiple of $\pi/(6H)$) is a copy of the fold, one per period. Neighbouring copies are separated by a tip (z a multiple of π), where the warp factor $(\sin^2 z)^{1/6}$ vanishes and $g_{x_0x_0} = \cot^2 z$ diverges, and which lies at an infinite proper distance from both: in proper distance every copy has its own complete line $-\infty < \sigma < \infty$. A compact direction of length L would give every field a tower of particles with the masses $2\pi n/L$, $n = 1, 2, \ldots$ (a Kaluza-Klein tower); because the copies are infinitely far apart, the period $\pi/(6H)$ of x_0 produces no such tower with the spacing $12H$, and the x_0 mass of the bound states is their binding to the fold.
3. *A folded sheet of paper.* The crease of a folded sheet is a fold in this sense: the two halves are mirror images (property 1) joined along the crease, the sheet is continuous there and its direction changes abruptly (property 3). In UM-3 the direction across the crease is the hidden direction σ, the seven directions along the crease are $x_1, \ldots, x_7$, and the crease is the fold.
4. *A folded function.* The warp factor $\mathrm{e}^{-2H|\sigma|}$ is the simplest folded function: continuous, mirror symmetric, largest at $\sigma = 0$, with a jump of its derivative,

$$\frac{\mathrm{d}}{\mathrm{d}\sigma}\mathrm{e}^{-2H|\sigma|} = -2H\,\mathrm{sgn}(\sigma)\,\mathrm{e}^{-2H|\sigma|}, \qquad \frac{\mathrm{d}^2}{\mathrm{d}\sigma^2}\mathrm{e}^{-2H|\sigma|} = 4H^2\,\mathrm{e}^{-2H|\sigma|} - 4H\,\delta_{\mathrm{D}}(\sigma) \tag{FD-7}$$

Symbols in (FD-7).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- sgn: The sign function: $\mathrm{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.
- δ_{D}: The Dirac delta function: $\int \mathrm{d}\sigma\,\delta_{\mathrm{D}}(\sigma)F(\sigma) = F(0)$. Unit: 1/length (for $\delta_{\mathrm{D}}(\sigma)$). Status: STANDARD.

whose delta function is the origin of the thin source of property 3. Every metric function of FD-4 depends on σ through $|\sigma|$ and is folded at $\sigma = 0$ in this sense. 5. *A fold map.* In the mathematics of smooth maps the fold (Whitney's fold singularity) is the map $u \mapsto u^2$ of a line onto a half-line: it folds the line at $u = 0$, two points $\pm u$ land on the same point, and the derivative of the map vanishes on the fold. Near the fold the map from the author's coordinate x_0 to the distance $|\sigma|$ from the fold is of this kind (FD-6), and $g_{x_0x_0} = \cot^2 z$, the square of the derivative $\mathrm{d}\sigma/\mathrm{d}x_0$, vanishes there. 6. *The brane of Randall and Sundrum.* The five-dimensional metric

$$\mathrm{d}y_{\mathrm{RS}}^2 + \mathrm{e}^{-2k_{\mathrm{RS}}|y_{\mathrm{RS}}|}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2 - \mathrm{d}x_4^2\big), \qquad \lambda_{\mathrm{RS}} = \frac{6k_{\mathrm{RS}}}{\kappa_5}, \qquad \Lambda_{\mathrm{RS}} = -\frac{6k_{\mathrm{RS}}^2}{\kappa_5} \tag{FD-8}$$

Symbols in (FD-8).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- y_{RS}: The proper coordinate across the brane of the five-dimensional model of Randall and Sundrum (FD-8, an example, not UM-3). Unit: length. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- k_{RS}: The inverse curvature length of the bulk of the model of Randall and Sundrum: the warp factor of the brane is $\mathrm{e}^{-2k_{\mathrm{RS}}|y_{\mathrm{RS}}|}$ (FD-8, an example, not UM-3). Unit: GeV (1/length). Status: STANDARD.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- λ_{RS}: The tension (energy per unit volume of ordinary space) of the brane of the model of Randall and Sundrum, $\lambda_{\mathrm{RS}} = 6k_{\mathrm{RS}}/\kappa_5$ (FD-8, an example, not UM-3). Unit: GeV^4. Status: STANDARD.
- κ_5: The gravitational coupling of five dimensions in the model of Randall and Sundrum, $\kappa_5 = 8\pi G_5$ (FD-8, an example, not UM-3). Unit: GeV^{-3}. Status: STANDARD.
- Λ_{RS}: The vacuum energy density of the bulk of the model of Randall and Sundrum, $\Lambda_{\mathrm{RS}} = -6k_{\mathrm{RS}}^2/\kappa_5$ (FD-8, an example, not UM-3). Unit: GeV^5. Status: STANDARD.

has a fold at $y_{\mathrm{RS}} = 0$ with the properties 1 to 3 (its coordinate y_{RS} is already the proper distance, so property 4 does not arise); Einstein's equations require there a brane of the tension λ_{RS} (Israel's junction condition) and in the bulk the vacuum energy Λ_{RS}, and the bulk is otherwise empty. The fold of UM-3 is of the same kind: its static block is a vacuum of the bulk with a fold of pure tension (FD-12). It differs in that the time x_4 is not warped, three extra times are warped together with ordinary space, the Lovelock terms add to the junction conditions, and the space is not cut at the brane: Randall and Sundrum identify the two sides of their brane (one physical side), while the fold of UM-3 has two distinct sides (example 7). 7. *A particle bound to a thin layer.* A particle held by an attractive thin layer has a wave function that falls off exponentially with the distance on both sides, and the layer makes its derivative jump (for the first-order Dirac equation, the wave function itself). In UM-3 the particles χ of dark matter are the states of dirac16complex bound to the fold; at rest and without their self-interaction (the field is $\Psi = \mathrm{e}^{3H|\sigma|}\chi(\sigma)\mathrm{e}^{-\mathrm{i}\omega_{\mathrm{b}}x_4}$)

$$\begin{aligned} &\chi(\sigma) = \mathrm{e}^{-q_{\mathrm{b}}|\sigma|}\chi(0^+) \quad (\sigma > 0), \qquad \chi(\sigma) = \mathrm{e}^{-q_{\mathrm{b}}|\sigma|}\chi(0^-) \quad (\sigma < 0), \\ &\chi(0^+) = \exp\Big(\gamma^{(0)}\big(\mu_{\mathrm{L}}P_{\mathrm{L}} + \mu_{\mathrm{R}}P_{\mathrm{R}}\big)\Big)\chi(0^-), \qquad \omega_{\mathrm{b}} = \sqrt{m_\Psi^2 - q_{\mathrm{b}}^2} \end{aligned} \tag{FD-9}$$

Symbols in (FD-9).

- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{3H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of a bound state. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- q_{b}: The inverse width of a state of dirac16complex bound to the fold: its profile falls as $\mathrm{e}^{-q_{\mathrm{b}}|\sigma|}$. Unit: GeV. Status: DERIVED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- μ_X, μ_{L}, μ_{R}: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_{\mathrm{L}}P_{\mathrm{L}} + \mu_{\mathrm{R}}P_{\mathrm{R}})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- P_{L}: The author's left-handed chiral projector, $P_{\mathrm{L}} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- P_{R}: The author's right-handed chiral projector, $P_{\mathrm{R}} = (\mathbf{1} + \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- Γ_8: The chirality matrix of eight dimensions, $\Gamma_8 = T_0T_1\cdots T_7$ (the author's T16^A[8]) $= \mathrm{diag}(-\mathbf{1}_8, \mathbf{1}_8)$; $\Gamma_8^2 = \mathbf{1}$; it anticommutes with every T_a. Unit: dimensionless. Status: AUTHOR.

- $\mathbf{1}$, $\mathbf{1}_8$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_8$ is the 8×8 identity matrix. Unit: dimensionless. Status: STANDARD.
- ω_{b}: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.

where μ_{L}, μ_{R} are the strengths of the fold for the left- and right-handed halves of dirac16complex (set by the Standard-Model fields on the fold) and $q_{\mathrm{b}}=-m_\Psi(\mu_{\mathrm{L}}+\mu_{\mathrm{R}})\tanh\varsigma/(2\varsigma)$, $\varsigma=\sqrt{\mu_{\mathrm{L}}\mu_{\mathrm{R}}}$ ($q_{\mathrm{b}}=-m_\Psi\mu_{\mathrm{L}}/2$ for $\mu_{\mathrm{R}}=0$). The profile is not a function of $|\sigma|$: it jumps at the fold, $\chi(0^+)\neq\chi(0^-)$. 8. *The fold as a mirror for dirac16complex.* The reflection across the fold acts on the profiles $\chi(\sigma)$ in two ways,

$$(\mathsf{R}_1\chi)(\sigma)=T_0T_1T_2T_3\,\chi(-\sigma),\qquad(\mathsf{R}_2\chi)(\sigma)=T_0T_5T_6T_7\,\chi(-\sigma)\tag{FD-10}$$

Symbols in (FD-10).

- R_1: The reflection across the fold acting on a profile of dirac16complex together with the reflection of ordinary space, $(\mathsf{R}_1\chi)(\sigma)=T_0T_1T_2T_3\,\chi(-\sigma)$ (FD-10): a symmetry of dirac16complex, of gravity and of the mean field of the gas of the particles χ; the bound states of Krein charge -1 are even, those of Krein charge $+1$ odd. Unit: dimensionless. Status: DERIVED.
- R_2: The reflection across the fold acting on a profile of dirac16complex together with the reflection of the extra times, $(\mathsf{R}_2\chi)(\sigma)=T_0T_5T_6T_7\,\chi(-\sigma)$ (FD-10): it maps the bound states of one Krein charge onto those of the other. Unit: dimensionless. Status: DERIVED.
- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi=\mathrm{e}^{3H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x}-\omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of a bound state. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma=|\cot z|\,\mathrm{d}x_0$, $\sigma=\mathrm{sgn}(\pi/2-z)\ln(\sin z)/(6H)$ on one period $0<z<\pi$; $\sigma=0$ is the fold $z=\pi/2$, $\sigma\to\mp\infty$ the tips $z\to0,\pi$; one period of x_0 is the whole line $-\infty<\sigma<\infty$. Unit: length. Status: DERIVED.
- T_a, T^a: The author's eight real 16×16 Dirac matrices T16ˆA[a] of 4+4.nb ($a=0,\ldots,7$; $T_0,\ldots,T_7$), with $\{T_a,T_b\}=2\eta_{ab}\mathbf{1}$; $T^a=\eta^{ab}T_b$. Unit: dimensionless. Status: AUTHOR.

(T_a the author's Dirac matrices; for states at rest R_1 is the reflection across the fold combined with the reflection of ordinary space, R_2 the reflection across the fold combined with the reflection of the extra times). The Krein charge of a state is its share of the conserved charge of dirac16complex, which is indefinite: a state can carry the charge -1 or $+1$. R_1 is a symmetry of dirac16complex, of gravity and of the mean field of a gas of the particles χ (the average field that the particles of the gas create and feel): the four bound states of Krein charge -1 (the particles χ) are even under it and the four of Krein charge $+1$ are odd, so the mean field of the gas cannot mix the two signs of the indefinite charge. R_2 maps the bound states of one Krein charge onto those of the other and makes the two sets degenerate without the self-interaction. 9. *The planes of Horava and Witten.* In M-theory (the eleven-dimensional theory whose limits include the superstring theories) on an interval (a circle whose points are identified with their mirror images), the two ten-dimensional planes left in place by the reflection are mirror surfaces (property 1); each carries one E_8 gauge group while gravity fills the bulk. As in UM-3, the gauge fields live on the mirror surface and gravity in the bulk; unlike UM-3, the two sides of each plane are identified (one physical side).

What is not a fold.

- *The tips* $z=0$ and $z=\pi$ (mod π): the mirror exchanges them within a period, but there the warp factor vanishes, $g_{x_0x_0}$ diverges, and they lie at an infinite proper distance; they carry no thin source.
- *A smooth maximum of a warp factor*, such as $1/\cosh^2(H\sigma)$: mirror symmetric and largest at $\sigma=0$, but without a kink and without a thin source.

2 How to read this document

2.1 Who this document is for

It is written for a reader who is clever but new to the subject: someone who knows what a coordinate, a derivative, an energy and a pressure are, but not general relativity, warped geometries or the universe model UM-3. Every term

is explained where it is first used, every equation that the document uses is written out in it, and every symbol of every display equation is listed and defined right after the display. Nothing has to be looked up anywhere else.

2.2 The status labels

Every statement carries one of the labels **AUTHOR** (from the author, Patrick L. Nash: his notebooks `Dirac_Einstein-Lovelock_4+4_Nash.nb` and `4+4.nb` and his orders of 2026-10-24 and 2026-10-25), **RECORD** (from the earlier record of this repository, such as the Lagrangian of dirac16complex), **STANDARD** (textbook physics or a published measurement, named in the provenance), **ASSUMED** (a choice made in UM-3 without derivation), **DERIVED** (computed and verified in UM-3; the check that verifies it is named, e.g. ST1), **ESTIMATE** (an order-of-magnitude evaluation) and **OPEN** (not settled). A result that holds only for the author's original Lovelock weights, which UM-3 no longer uses, is called a record of the original weights.

2.3 Every equation is followed by its symbols

A display equation carries a tag. SB-n is an equation of this document; FD-n belongs to the definition of the fold at the beginning; FE-n (a field equation) and IN-n (a term of the action) are equations of UM-3 that are written identically, with the same tag, in every UM-3 document that uses them. After every display comes the list "Symbols in ..." of every symbol in it, with its meaning, its unit, its value where one exists, and its status. The units are natural units: the reduced Planck constant and the speed of light are 1, so that a time is a length, and energies, masses and inverse lengths are all measured in GeV.

2.4 The checks

Most statements labelled DERIVED in this document are verified by the nine checks ST1 to ST9 of a program that computes the field equations exactly with computer algebra (ST1 to ST8 with the author's original weights, ST9 with the corrected ones). The others are verified by checks of the other programs of UM-3, named where they are used: the gravitational waves and the corrected weights (W1 to W8), the curvature and the bulk equations of the author's metric (G2 to G4), the junction conditions and the fold sources (J4 to J7), and the proper distance of the definition of the fold. The provenance at the end says what each check does and how to run it.

3 The stabiliser in brief

UM-3 is a model universe built on a metric (a rule for measuring distances and durations, explained below) that the author wrote down for an eight-dimensional spacetime. The metric has a fold, the surface defined at the beginning of this document, and around it the bulk: everything that is not on the fold. Gravity in UM-3 obeys the Einstein-Lovelock field equations, which say how the curvature of spacetime is fixed by the energy that it contains. These equations contain three weights and a cosmological constant.

The finding (with the author's original weights). When the universe is old and expands slowly, as ours does, the bulk of the author's metric is very nearly static: nothing in it changes with time. With the weights and the cosmological constant of the author's notebook, a static bulk of this metric is **not** a solution of the field equations with empty space. The equations demanded a definite energy content: a negative energy density, and a negative pressure five times as large, the same in every direction of the bulk except the time (SB-9). UM-3 called whatever supplies it **the stabiliser of the bulk**, because without it the bulk could not stay static. It had no free parameter: its energy density and its pressure were fixed by the author's constant H and the gravitational coupling κ.

Who found it. The requirement and the stabiliser are not part of the author's specification. They were found, introduced and named by Claude, an AI model made by Anthropic, working in Claude Code sessions configured for the model claude-opus-5-5, on 2026-10-24 while it derived the complete field equations of UM-3. The idea of something that holds a static block of the bulk is older in this repository: the two earlier universe models assumed a "block stabiliser" of their own (the provenance gives the lineage); the stabiliser of UM-3, for the author's new metric, was a different object with a definite, negative energy density.

The correction. On 2026-10-25 the author ordered: "Gravitons must travel at c. If these incorrect claims come from my choice of the Lovelock couplings and my choice of the cosmological constant Λ THEN my choices are wrong, and you MUST correct them." With the original weights, gravitational waves in the bulk moved $\sqrt{5} = 2.24$ times faster than light. There is exactly one choice of the two higher weights for which gravitons travel at the speed of

light, and with it exactly one cosmological constant makes the static bulk empty (FE-68 to FE-70). UM-3 now uses these values. **The stabiliser was a consequence of the author's original weights and cosmological constant; with the corrected ones the static bulk is a vacuum, and the stabiliser vanishes** (IN-7, ST9).

The five points of the author's order of 2026-10-25, in brief. 1. *The stabiliser couples only to gravity.* That is how UM-3 defined it, and on the static bulk it was consistent: the stabiliser was conserved by itself (ST4). In an expanding universe the bulk source that the original equations required released energy that flowed along the hidden direction into the fold (ST6); reading this as an exchange of energy between the stabiliser and the matter of the fold is an interpretation (ASSUMED). With the corrected weights the question disappears: there is no stabiliser, and the bulk of an expanding fold with frozen extra times is a vacuum without any flow (FE-33). 2. *It is absent in the primordial era, whose bulk is the author's vacuum (FE-31).* With the original weights this was true: the author's vacuum solved the bulk equations with no source (ST2), and a medium that couples only to gravity with a pressure five times its energy density could not appear later out of nothing (ST6). With the corrected weights the author's vacuum is not a solution any more, the primordial era is an inflation of the fold, and no era of UM-3 has a stabiliser. 3. *Its existence and value are DERIVED from the requirement of a static bulk.* True: the static bulk fixed every entry (ST1), and a bulk without any source could only be the author's vacuum or its time reverse at every instant (ST3, ST8). That the source was a separate medium was ASSUMED. With the corrected weights the same derivation gives the value zero (ST9). 4. *The static bulk* is explained, with five examples, in the part "The static bulk". 5. *Warp and warp factor* are defined, with four examples, in the part "Warp and warp factor".

Its importance. With the original weights the stabiliser was what made a slowly expanding late universe with constant couplings possible in the author's metric; it was also the most exotic ingredient of the model, because it violated the null energy condition. Its removal by the corrected weights changes UM-3 in four ways: gravitational waves travel at the speed of light at every wavelength; the fold obeys the Friedmann equations of general relativity exactly; the bulk needs no exotic medium; and a new requirement appears on the fold, a pressure along the extra times that holds their volume fixed, a role that the "block stabiliser" of the first universe model of this repository had played for its own metric.

4 The language

4.1 Coordinates, distances and the metric

A point of space and time is labelled by numbers, its coordinates. Ordinary space needs three (x_1, x_2, x_3) and time one (x_4). The spacetime of UM-3 has eight: x_0, a hidden direction that we do not see, x_1, x_2, x_3 ordinary space, x_4 the time, and x_5, x_6, x_7 three extra directions that behave like time (the extra times). The metric is the rule that turns a small step of the coordinates into the squared length (or the squared duration) of the step. A familiar example is the surface of a globe of radius R_g, on which a point is labelled by its angle θ_g from the north pole and its longitude ϕ_g:

$$\mathrm{d}\ell_\mathrm{g}^2 = R_\mathrm{g}^2\,\mathrm{d}\theta_\mathrm{g}^2 + R_\mathrm{g}^2 \sin^2\theta_\mathrm{g}\,\mathrm{d}\phi_\mathrm{g}^2 \tag{SB-1}$$

Symbols in (SB-1).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- ℓ_g: The length measured along the surface of a globe; $\mathrm{d}\ell_\mathrm{g}$ is the length of a small step on it (an example of a metric, not UM-3). Unit: length. Status: STANDARD.
- R_g: The radius of a globe (an example, not UM-3). Unit: length. Status: STANDARD.
- θ_g: The angle of a point of a globe from its north pole: 0 at the north pole, $\pi/2$ on the equator, π at the south pole (an example, not UM-3). Unit: radian. Status: STANDARD.
- ϕ_g: The longitude of a point of a globe (an example, not UM-3). Unit: radian. Status: STANDARD.

A step of the longitude by $\mathrm{d}\phi_\mathrm{g}$ is long on the equator ($\sin\theta_\mathrm{g} = 1$) and short near the poles: the factor $R_\mathrm{g}^2 \sin^2\theta_\mathrm{g}$ in front of $\mathrm{d}\phi_\mathrm{g}^2$ says how large one unit of longitude is at each latitude (sin is the sine function; d marks a small step). Status: STANDARD.

The metric of the eight-dimensional spacetime is written $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$: the indices M and N run over the eight coordinates $X = \{x_0, \ldots, x_7\}$, a repeated index is summed, and the numbers g_{MN} are the components of the metric. In spacetime a step can be space-like (a positive squared interval: a distance) or time-like (a negative one: minus

the square of a duration). UM-3 counts x_0, x_1, x_2, x_3 as space-like and x_4, x_5, x_6, x_7 as time-like (the signature $+,+,+,+,-,-,-,-$). The author's metric is written in FD-4 of the definition of the fold with the proper distance σ from the fold: $\mathrm{d}\sigma^2$ for the hidden direction, a factor that depends on σ in front of ordinary space and of the extra times, and $-\mathrm{d}x_4^2$ for the time. The two functions $a_4(x_4)$ and $b_4(x_4)$ of the time are the author's: at the depth $|\sigma|$, a coordinate step of ordinary space has the length $\mathrm{e}^{a_4}\mathrm{e}^{-H|\sigma|}$ times the step, and a coordinate step of an extra time the duration $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ times the step; on the fold, e^{a_4} is the size of ordinary space and $\mathrm{e}^{-a_4}b_4$ that of the extra times (e is Euler's number, $|\sigma|$ the absolute value). Status: AUTHOR (the metric), DERIVED (its form in σ, FD-3 and FD-4).

4.2 Warp and warp factor

Definition. A metric is *warped* along a direction when the size of other directions depends on the position along it. In a metric of the form $\mathrm{d}\sigma^2 + \mathcal{W}(\sigma)\,\mathrm{d}y^2 + \ldots$ the function $\mathcal{W}(\sigma)$ that multiplies the squared steps $\mathrm{d}y^2$ of the other directions is the **warp factor**. The length of a fixed coordinate step $\mathrm{d}y$ is $\sqrt{\mathcal{W}}\,\mathrm{d}y$, so lengths along those directions scale with $\sqrt{\mathcal{W}}$. The **warp** is the dependence itself, measured by the rate at which the natural logarithm $\ln\mathcal{W}$ changes along σ. In UM-3:

$$\mathcal{W}(\sigma) = \mathrm{e}^{-2H|\sigma|} = (\sin^2 z)^{1/6}, \qquad \frac{\mathrm{d}\ln\mathcal{W}}{\mathrm{d}\sigma} = -2H\,\mathrm{sgn}(\sigma) \tag{SB-2}$$

Symbols in (SB-2).

- $\mathcal{W}$: The warp factor: in a warped metric, the function of the position along one direction that multiplies the squared steps of other directions (its square root multiplies their lengths); in UM-3 the function of σ that multiplies ordinary space and the extra times, $\mathcal{W} = \mathrm{e}^{-2H|\sigma|} = (\sin^2 z)^{1/6}$, equal to 1 on the fold. Unit: dimensionless. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$, the fold is $z = \pi/2$. Unit: dimensionless. Status: AUTHOR.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- sgn: The sign function: $\mathrm{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.

The warp factor multiplies ordinary space (together with e^{2a_4}) and the extra times (together with $\mathrm{e}^{-2a_4}b_4^2$), but not the time x_4 and not σ itself. The warp factor is largest, 1, on the fold and falls on both sides. A coordinate step of ordinary space at the depth $|\sigma|$ in the bulk is $\mathrm{e}^{-H|\sigma|}$ times as long as the same coordinate step on the fold: the deeper in the bulk, the smaller ordinary space and the extra times become. The warp factor is the same on both sides of the fold; the warp, its logarithmic rate of change, has the constant size $2H$ and changes its sign on the fold (sgn is the sign function): this jump is the kink of the fold (property 3 of the definition). Status: AUTHOR (the metric), DERIVED (the form in σ).

Examples of warps.

- *A globe* (SB-1): the longitude is warped along the angle from the pole, with the warp factor $R_\mathrm{g}^2\sin^2\theta_\mathrm{g}$ in the proper coordinate $R_\mathrm{g}\theta_\mathrm{g}$ (the distance from the pole along a meridian).
- *The fold of UM-3* (SB-2): ordinary space and the extra times are warped along the hidden direction.
- *The brane of Randall and Sundrum* (FD-8, example 6 of the definition of the fold): the four directions of our world are warped by the warp factor $\mathrm{e}^{-2k_\mathrm{RS}|y_\mathrm{RS}|}$ along a fifth direction. Unlike in UM-3, the time is warped as well.
- *Time near the Earth.* Near a mass the time is warped. In the weak gravity of the Earth the factor in front of $-\mathrm{d}x_4^2$ is $1 + 2\Phi_\mathrm{N}$, where Φ_N is the Newtonian gravitational potential (negative, and larger in size closer to the Earth). A clock lower down therefore runs slower. Pound and Rebka measured this in 1960 over the 22.5 m

height of a tower, and the clocks of the GPS satellites are corrected for it. In UM-3 the time is not warped along the hidden direction: a clock deep in the bulk runs at the same rate as a clock on the fold.

4.3 The bulk

Definition. The bulk is all of the eight-dimensional spacetime except the fold: the points with $\sigma \neq 0$. It extends on both sides of the fold to the tips, which lie at an infinite proper distance. Gravity and dirac16complex live in the bulk as well as on the fold; dirac16complex is the author's field of matter of spin one half in eight dimensions, a spinor with sixteen complex components. The Standard Model of particle physics lives only on the fold. Status: ASSUMED (where the fields live), AUTHOR (the metric and dirac16complex).

4.4 Energy, pressure and the energy-momentum tensor

Matter and fields carry energy and momentum. Their densities and flows are collected in the energy-momentum tensor, an eight-by-eight array whose entries are measured along the eight directions of an orthonormal frame: the frame direction (0) along σ, $(1), (2), (3)$ along ordinary space, (4) along the time and $(5), (6), (7)$ along the extra times. For a medium at rest the entry along the time is minus its energy density, the entries along the other directions are its pressures, and an entry that mixes the time with another direction is a flow of energy along that direction. A medium at rest with the same pressure p_{med} in every direction but the time has

$$\mathcal{T}_{\mathrm{med}}{}^{a}{}_{b} = \mathrm{diag}\big(p_{\mathrm{med}}, p_{\mathrm{med}}, p_{\mathrm{med}}, p_{\mathrm{med}}, -\rho_{\mathrm{med}}, p_{\mathrm{med}}, p_{\mathrm{med}}, p_{\mathrm{med}}\big) \tag{SB-3}$$

Symbols in (SB-3).

- $\mathcal{T}_{\mathrm{med}}{}^{a}{}_{b}$: The energy-momentum tensor of one medium (frame components): for a medium at rest the entry along the time is minus its energy density, the entries along the other directions are its pressures, and the entries that mix the time with another direction are its flows of energy along that direction. Unit: GeV^8 (energy per volume in eight dimensions). Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, $(1), (2), (3)$ along ordinary space, (4) along the time, $(5), (6), (7)$ along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- p_{med}: The pressure of a medium at rest, the same along every direction but the time (for a medium of the bulk: along σ, ordinary space and the extra times). Unit: GeV^8. Status: STANDARD.
- ρ_{med}: The energy density (energy per volume) of a medium at rest. Unit: GeV^8. Status: STANDARD.

Three examples in the four dimensions of our world: dust (pressureless matter, $p_{\mathrm{med}} = 0$), light ($p_{\mathrm{med}} = \rho_{\mathrm{med}}/3$) and the energy of the vacuum ($p_{\mathrm{med}} = -\rho_{\mathrm{med}}$). A positive vacuum energy is a tension: it pulls instead of pushing, like a stretched skin. Yet its gravity is repulsive, because in general relativity pressure gravitates too: the source of the attraction is $\rho + 3p$ in four dimensions, and a tension with $p = -\rho$ makes it negative. This is why a positive cosmological constant makes a universe expand faster. Two energy conditions classify media: the weak energy condition (the energy density is never negative) and the null energy condition (the energy density plus the pressure along any space-like direction is never negative; for the frame of UM-3, which has four time-like directions, the condition is taken for light moving along the time x_4 and one space-like direction). Ordinary matter, light and a positive vacuum energy satisfy both. A negative vacuum energy, such as the negative cosmological constant of the bulk of UM-3, violates the weak condition but not the null one, since its energy density plus its pressure is zero. A medium that violates the null condition is called exotic. Status: STANDARD.

4.5 The field equations of gravity

Einstein's insight was that gravity is the curvature of spacetime, and that the curvature is fixed by the energy-momentum tensor. One side of his field equations is a combination of derivatives of the metric (the Einstein tensor); the other is the energy-momentum tensor times the gravitational coupling. UM-3 uses the generalisation found by

Lovelock, which in eight dimensions adds two more combinations of the curvature, each with a weight. The action (the quantity whose variation gives the equations) and its curvature terms are

$$S_{\mathrm{EL}} = \frac{1}{2\kappa}\int \mathrm{d}^8X\,\sqrt{|g|}\,\Big(\sum_{p=1}^{3}\alpha_{(p)}\,\mathcal{L}_{(p)} - 2\Lambda\Big), \qquad \kappa = 8\pi G_8,$$
$$\alpha_{(p)} = w_{(p)}\,H^{2-2p}, \qquad w_{(1)} = 1, \quad w_{(2)} = \frac{1}{20}, \quad w_{(3)} = \frac{1}{360}, \qquad \Lambda = -7H^2 \tag{IN-2}$$

Symbols in (IN-2).

- S_{EL}: The Einstein-Lovelock action of gravity in eight dimensions. Unit: dimensionless. Status: AUTHOR.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\ldots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations a second label of a Kohn-Sham orbital (as o, with δ_{op} the Kronecker delta of the two labels). Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)}H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- $\mathcal{L}_{(p)}$: The Lovelock density of order p (a scalar built from p Riemann tensors): $\mathcal{L}_{(1)} = R$ (Einstein-Hilbert), $\mathcal{L}_{(2)}$ the Gauss-Bonnet density, $\mathcal{L}_{(3)}$ the cubic Lovelock density. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -7H^2$: with the corrected weights the value for which the static block is a vacuum (its Lovelock tensors add to $7H^2$ in every direction). The author's original value was $\Lambda = -10H^2$ (he writes the pure number $\Lambda_{\mathrm{N}} = 10$ in $0 = -\Lambda_{\mathrm{N}} + \sum_p H^{-2p}w_{(p)}\mathcal{E}_{(p)}$). Unit: GeV^2. Value: $-7H^2$; the author's original $-10H^2$. Status: AUTHOR (the term), DERIVED (the value).
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_8: The eight-dimensional Newton constant ($\kappa = 8\pi G_8$). Unit: GeV^{-6}. Status: AUTHOR.
- $w_{(p)}$: The Lovelock weights, $w_{(1)} = 1$, $w_{(2)} = 1/20$, $w_{(3)} = 1/360$: the only weights for which gravitational waves travel at the speed of light in the bulk and on the fold (both coefficients of the bulk wave, $1 - 40w_{(2)} + 360w_{(3)}$ and $1 - 24w_{(2)} + 72w_{(3)}$, vanish). The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) made bulk gravitons faster than light; they were corrected at the author's order of 2026-10-25. Unit: dimensionless. Value: $(1, 1/20, 1/360)$; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), DERIVED (the values).
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

$$\mathcal{L}_{(p)} = \frac{1}{2^p}\,\delta^{M_1N_1\cdots M_pN_p}_{P_1Q_1\cdots P_pQ_p}\,R^{P_1Q_1}{}_{M_1N_1}\cdots R^{P_pQ_p}{}_{M_pN_p},$$
$$\mathcal{L}_{(1)} = R, \qquad \mathcal{L}_{(2)} = R^2 - 4R_{MN}R^{MN} + R_{MNPQ}R^{MNPQ} \tag{IN-3}$$

Symbols in (IN-3).

- $\mathcal{L}_{(p)}$: The Lovelock density of order p (a scalar built from p Riemann tensors): $\mathcal{L}_{(1)} = R$ (Einstein-Hilbert), $\mathcal{L}_{(2)}$ the Gauss-Bonnet density, $\mathcal{L}_{(3)}$ the cubic Lovelock density. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations a second label of a Kohn-Sham orbital (as o, with δ_{op} the Kronecker delta of the two labels). Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with upper and lower index strings the generalised Kronecker

delta $\delta^{M_1\cdots M_n}_{N_1\cdots N_n}$, the determinant of the $n\times n$ matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- P: A coordinate index of eight dimensions (also $P_1, P_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Q: A coordinate index of eight dimensions (also $Q_1, Q_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- ${R^{MN}}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor (convention $[\nabla_P, \nabla_Q]V^M = {R^M}_{NPQ}V^N$), with two indices the Ricci tensor $R_{MN} = {R^P}_{MPN}$, without indices the scalar curvature $R = g^{MN}R_{MN}$; with frame indices ${R^{(a)(b)}}_{(c)(d)}$ its frame components, ${R^{(a)(b)}}_{(a)(b)}$ (no sum) the sectional curvature of the frame plane (a, b). Unit: $1/\text{length}^2$. Status: STANDARD.

The curvature is measured by the Riemann tensor R, which says how much a direction turns when it is carried around a small loop; the generalised Kronecker delta δ is the totally antisymmetric combination of ordinary Kronecker deltas, $+1$ or -1 when its lower indices are an even or odd rearrangement of its upper ones and 0 otherwise. The weights in IN-2 are the corrected ones; the author's notebook had $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ and $\Lambda = -10H^2$ (he writes the pure number 10 in his vacuum equations). Varying the action with respect to the inverse metric g^{MN} and dividing by $\sqrt{|g|}$ gives the Lovelock tensors

$$
\begin{aligned}
{\mathcal{E}_{(p)}}^M{}_N &= -\frac{1}{2^{p+1}}\,\delta^{MM_1N_1\cdots M_pN_p}_{NP_1Q_1\cdots P_pQ_p}\,{R^{P_1Q_1}}_{M_1N_1}\cdots{R^{P_pQ_p}}_{M_pN_p},\\
{\mathcal{E}_{(1)}}^M{}_N &= {R^M}_N - \frac{1}{2}\,\delta^M_N\,R,\\
{\mathcal{E}_{(2)}}^M{}_N &= 2\big(R\,{R^M}_N - 2{R^M}_P{R^P}_N - 2{R^{MP}}_{NQ}{R^Q}_P + R^{MPQS}R_{NPQS}\big) - \frac{1}{2}\,\delta^M_N\,\mathcal{L}_{(2)}
\end{aligned}
\tag{FE-2}
$$

Symbols in (FE-2).

- ${\mathcal{E}_{(p)}}^M{}_N$: The Lovelock tensor of order p: the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to the metric ($\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor); with frame indices its frame components. Unit: $1/\text{length}^{2p}$. Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations a second label of a Kohn-Sham orbital (as o, with δ_{op} the Kronecker delta of the two labels). Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with upper and lower index strings the generalised Kronecker delta $\delta^{M_1\cdots M_n}_{N_1\cdots N_n}$, the determinant of the $n\times n$ matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- P: A coordinate index of eight dimensions (also $P_1, P_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Q: A coordinate index of eight dimensions (also $Q_1, Q_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- ${R^{MN}}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor (convention $[\nabla_P, \nabla_Q]V^M = {R^M}_{NPQ}V^N$), with two indices the Ricci tensor $R_{MN} = {R^P}_{MPN}$, without indices the scalar curvature $R = g^{MN}R_{MN}$; with frame indices ${R^{(a)(b)}}_{(c)(d)}$ its frame components, ${R^{(a)(b)}}_{(a)(b)}$ (no sum) the sectional curvature of the frame plane (a, b). Unit: $1/\text{length}^2$. Status: STANDARD.
- S: A coordinate index of eight dimensions, running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- $\mathcal{L}_{(p)}$: The Lovelock density of order p (a scalar built from p Riemann tensors): $\mathcal{L}_{(1)} = R$ (Einstein-Hilbert), $\mathcal{L}_{(2)}$ the Gauss-Bonnet density, $\mathcal{L}_{(3)}$ the cubic Lovelock density. Unit: $1/\text{length}^{2p}$. Status: STANDARD.

($R^M{}_N$ the Ricci tensor and R the scalar curvature, contractions of the Riemann tensor), and the field equations of the bulk are

$$\sum_{p=1}^{3} \alpha_{(p)}\, \mathcal{E}_{(p)}{}^M{}_N + \Lambda\, \delta^M_N = \kappa\, \mathcal{T}_\Psi{}^M{}_N \qquad (\sigma \neq 0) \tag{FE-1}$$

Symbols in (FE-1).

- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations a second label of a Kohn-Sham orbital (as o, with δ_{op} the Kronecker delta of the two labels). Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)} H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- $\mathcal{E}_{(p)}{}^M{}_N$: The Lovelock tensor of order p: the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to the metric ($\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor); with frame indices its frame components. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -7H^2$: with the corrected weights the value for which the static block is a vacuum (its Lovelock tensors add to $7H^2$ in every direction). The author's original value was $\Lambda = -10H^2$ (he writes the pure number $\Lambda_\mathrm{N} = 10$ in $0 = -\Lambda_\mathrm{N} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$). Unit: GeV^2. Value: $-7H^2$; the author's original $-10H^2$. Status: AUTHOR (the term), DERIVED (the value).
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with upper and lower index strings the generalised Kronecker delta $\delta^{M_1 \cdots M_n}_{N_1 \cdots N_n}$, the determinant of the $n \times n$ matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_\Psi$: The energy-momentum tensor of dirac16complex, $\mathcal{T}_{MN} = -(2/\sqrt{|g|})\, \delta S_\Psi / \delta g^{MN}$. Unit: GeV^8. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\, \mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z) \ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

The left-hand side is built from the metric alone. The right-hand side is the energy content of the bulk: dirac16complex, and, with the original weights, the stabiliser. A metric solves the equations only if the right-hand side has exactly the value that the left-hand side computes; a solution with nothing on the right is a *vacuum*. Status: AUTHOR (the theory), STANDARD (Lovelock gravity), DERIVED (the corrected weights).

The divergence of every Lovelock tensor vanishes identically (the Bianchi identity). The right-hand side must therefore be conserved: the energy and momentum of all sources together cannot be created or destroyed, only moved. A single medium that interacts with nothing but gravity is conserved by itself:

$$\nabla_M\, \mathcal{T}_\mathrm{med}{}^M{}_N = 0 \tag{SB-4}$$

Symbols in (SB-4).

- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols $\Gamma^P{}_{MN}$). Status: STANDARD.
- $\mathcal{T}_\mathrm{med}{}^a{}_b$: The energy-momentum tensor of one medium (frame components): for a medium at rest the entry along the time is minus its energy density, the entries along the other directions are its pressures, and the entries that mix the time with another direction are its flows of energy along that direction. Unit: GeV^8 (energy per volume in eight dimensions). Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.

(∇ is the covariant derivative, the derivative that takes into account how the directions of the coordinates change from point to point.) Status: STANDARD.

The junction conditions. The metric has a kink at the fold, so its second derivatives along σ contain a Dirac delta function there, and the field equations hold across the fold only if the fold carries a thin source $\mathcal{S}$, its energy-momentum tensor integrated across it (unit GeV^7: an energy density times a length): the integral FD-11 of the definition of the fold. For the Einstein term alone this is Israel's condition (1966): $\kappa\mathcal{S}^a{}_b = -([K^a{}_b] - \delta^a{}_b[K])$, with the jump $[K^a{}_b]$ of the extrinsic curvature of the fold (FD-5) and its trace $[K]$; for the Gauss-Bonnet and cubic terms the integral adds terms of higher order in the extrinsic curvature and the curvature of the fold (Davis 2003; Gravanis and Willison 2003). Status: STANDARD.

5 The static bulk

5.1 What static means

Definition. A spacetime is *static* when nothing in it changes with the time and nothing flows: its metric does not depend on the time, and no motion that the reversal of time would change takes place in it. In the author's metric the time dependence sits in the two functions $a_4(x_4)$ and $b_4(x_4)$. The bulk is static when both are constant, that is when the rates $\mathsf{A} = a_4'/H$ (the rate at which ordinary space grows) and $\mathsf{U} = b_4'/(Hb_4)$ (the rate at which b_4 changes) both vanish:

$$\begin{aligned} &\mathsf{A} = \frac{a_4'}{H} = 0, \qquad \mathsf{U} = \frac{b_4'}{Hb_4} = 0 : \\ &g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N = \mathrm{d}\sigma^2 + \mathrm{e}^{-2H|\sigma|}\Big[\mathrm{e}^{2a_4}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{e}^{-2a_4}b_4^2\big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big)\Big] - \mathrm{d}x_4^2 \end{aligned} \tag{SB-5}$$

Symbols in (SB-5).

- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); 1 at the author's vacuum, which is not a solution with the corrected weights.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\ldots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0,\ldots,x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0,\ldots,x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.

with constant a_4 and b_4 (a prime is the derivative with respect to the time x_4). UM-3 calls this the *static block*. Status: DERIVED (the form), ASSUMED (that the late bulk is the static block to first approximation).

5.2 Static is not the same as empty

A static spacetime may be empty or it may need a source, and an empty spacetime may be static or not:

spacetime	static?	empty (a vacuum)?
the static block of UM-3 (SB-5), corrected weights	yes	yes (IN-7)
the static block of UM-3 (SB-5), the author's original weights	yes	no: it needed the stabiliser (SB-9)
the author's vacuum (FE-31)	no: ordinary space grows at the rate H	with the original weights yes; with the corrected weights no
the space outside a star	yes	yes
Einstein's static universe (SB-6)	yes	no: it needs dust and a cosmological constant

5.3 Examples of static spacetimes and static media

Example 1: the static block of UM-3 (SB-5). It is the late universe to first approximation. Today ordinary space expands at the Hubble rate $H_4^{(0)} = 1.44 \times 10^{-42}$ GeV. If the author's constant H is much larger, the rate $\mathsf{A} = H_4^{(0)}/H$ is tiny and the bulk is static up to corrections of order A. The extra times are frozen ($\mathrm{e}^{-a_4}b_4$ constant, $\mathsf{U} = \mathsf{A}$), so that the couplings of the Standard Model do not change. With the corrected weights the static block is an exact vacuum, and so is the bulk of an expanding fold with frozen extra times for every expansion history (FE-33). Status: ASSUMED (the late universe), DERIVED (the vacuum).

Example 2: the space outside a star or a planet (Schwarzschild's solution of 1916). It is static and empty: a static vacuum. Not every static spacetime needs a source. Status: STANDARD.

Example 3: Einstein's static universe (1917). Einstein wanted a universe that does not change. With dust of the mass density ρ_E and nothing else his equations do not allow it, because gravity pulls the dust together. He added a positive cosmological constant Λ_E, whose gravity is repulsive (it acts like a positive vacuum energy, a tension), and found a static universe, a three-dimensional sphere of radius R_E, but only for one exact balance:

$$\Lambda_\mathrm{E} = 4\pi G_\mathrm{N}\,\rho_\mathrm{E} = \frac{1}{R_\mathrm{E}^2} \qquad \text{(SB-6)}$$

Symbols in (SB-6).

- Λ_E: The cosmological constant of Einstein's static universe of 1917, in four dimensions (an example, not UM-3). Unit: 1/length^2. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

- G_{N}: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_{\mathrm{N}} = 1/M_{\mathrm{Pl}}^2$ with the Planck mass $M_{\mathrm{Pl}} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- ρ_{E}: The mass density of the dust that fills Einstein's static universe (an example, not UM-3). Unit: mass per volume. Status: STANDARD.
- R_{E}: The radius of the three-dimensional sphere that is the space of Einstein's static universe (an example, not UM-3). Unit: length. Status: STANDARD.

This is the classic example of a static spacetime that requires a definite source. It is also unstable (Eddington, 1930): a small excess of matter makes it collapse, a small deficit makes it expand for ever. Einstein's Λ_{E} played for his universe the part that the stabiliser played for the bulk of UM-3 with the original weights. Status: STANDARD.

Example 4: the air at rest above the ground. The atmosphere stays at rest only because its pressure falls with the altitude exactly so fast that the difference of pressure carries the weight of the air in between:

$$\frac{\mathrm{d}P_{\mathrm{air}}}{\mathrm{d}h_{\mathrm{alt}}} = -\rho_{\mathrm{air}}\, g_{\oplus} \tag{SB-7}$$

Symbols in (SB-7).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- P_{air}: The pressure of the air at the altitude h_{alt} (an example, not UM-3). Unit: Pa. Value: about 1.0×10^5 Pa at the ground. Status: STANDARD.
- h_{alt}: The altitude above the ground (an example, not UM-3). Unit: m. Status: STANDARD.
- ρ_{air}: The mass density of the air at the altitude h_{alt} (an example, not UM-3). Unit: $\mathrm{kg/m}^3$. Value: about 1.2 $\mathrm{kg/m}^3$ at the ground. Status: STANDARD.
- $g_{\oplus}$: The acceleration of free fall at the surface of the Earth (an example, not UM-3). Unit: $\mathrm{m/s}^2$. Value: 9.81 $\mathrm{m/s}^2$. Status: STANDARD.

Again, the static state needs a definite distribution of pressure; any other distribution sets the air in motion. Status: STANDARD.

Example 5: a drum skin at rest. A stretched skin stays flat and at rest because its tension is the same everywhere, so that the pulls on every small piece of it balance. Pressing it or loosening it locally sets it in motion. The fold of UM-3 is a skin of this kind: at the static block its thin source is a pure tension (FE-32). Status: STANDARD.

In all these examples except the empty space outside a star, staying static requires a definite content: a definite balance of energy and pressure. Whether the static block of UM-3 is of this kind or a vacuum depends on the Lovelock weights.

6 The stabiliser of the author's original weights

This part is the record of the stabiliser as it was found, with the weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ and the cosmological constant $\Lambda = -10H^2$ of the author's notebook. The next part shows what the corrected weights change.

6.1 How the static block fixes the source

The static block SB-5 has a simple curvature. In each half of the bulk, the seven directions other than the time (σ, ordinary space and the extra times) form a space of constant negative curvature $-H^2$, like a slice of anti-de Sitter space: every pair of these directions has the frame curvature $R^{ab}{}_{ab} = -H^2$, and no pair with the time is curved (the time is not warped). For such a space times a flat line, the Lovelock tensor of order p (FE-2) has only diagonal entries, and the entry along a direction depends only on the number m_{c} of the other curved directions:

$$\mathcal{E}_{(p)}{}^{(a)}{}_{(a)}\Big|_{\text{static block}} = -\frac{1}{2}\left(-H^2\right)^p \frac{m_{\mathrm{c}}!}{(m_{\mathrm{c}}-2p)!} \tag{SB-8}$$

$\left(m_{\mathrm{c}} = 6 \text{ along the hidden direction, ordinary space and the extra times, } m_{\mathrm{c}} = 7 \text{ along the time}\right)$

Symbols in (SB-8).

- ${\mathcal{E}_{(p)}}^{M}{}_{N}$: The Lovelock tensor of order p: the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to the metric ($\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor); with frame indices its frame components. Unit: $1/\text{length}^{2p}$. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \text{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations a second label of a Kohn-Sham orbital (as o, with δ_{op} the Kronecker delta of the two labels). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- m_{c}: The number of the other curved directions of the static block seen from one direction: 6 for σ, ordinary space and the extra times (the seven curved directions without the given one), 7 for the time (which is not curved itself). Unit: dimensionless. Value: 6 or 7. Status: DERIVED.

(no sum over the frame index a; $m_{\mathrm{c}}!$ is the factorial $1 \cdot 2 \cdots m_{\mathrm{c}}$.) Each Riemann tensor in FE-2 contributes the curvature $-H^2$ of a pair of directions, and the generalised Kronecker delta counts the ordered ways of choosing the $2p$ other directions of the p pairs among the m_{c} curved ones, $m_{\mathrm{c}}!/(m_{\mathrm{c}} - 2p)!$. Multiplied by the couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$, the entries along a warped direction and along the time are, in units of H^2:

order p	per unit weight (warped, time)	original weights (warped, time)	corrected weights (warped, time)
1, Einstein	$(15, 21)$	$(15, 21)$	$(15, 21)$
2, Gauss-Bonnet	$(-180, -420)$	$(-45/8, -105/8)$	$(-9, -21)$
3, cubic	$(360, 2520)$	$(5/16, 35/16)$	$(1, 7)$
the cosmological constant		$(-10, -10)$	$(-7, -7)$
total		$(-5/16, 1/16)$	$(0, 0)$

For general weights this is FE-70:

$$
\begin{aligned}
&\sum_{p=1}^{3} \alpha_{(p)}\, {\mathcal{E}_{(p)}}^{1}{}_{1} + \Lambda = \big(15 - 180w_{(2)} + 360w_{(3)}\big)H^2 + \Lambda, \\
&\sum_{p=1}^{3} \alpha_{(p)}\, {\mathcal{E}_{(p)}}^{4}{}_{4} + \Lambda = \big(21 - 420w_{(2)} + 2520w_{(3)}\big)H^2 + \Lambda, \\
&w_{(2)} = \frac{1}{20},\ w_{(3)} = \frac{1}{360}: \quad \text{both } 7H^2 + \Lambda, \qquad \Lambda = -7H^2; \\
&\text{curvature } -\mathsf{y}H^2: \quad (\mathsf{y}-1)^2(\mathsf{y}-7)\,H^2, \quad 7(\mathsf{y}-1)^3H^2
\end{aligned}
\tag{FE-70}
$$

Symbols in (FE-70).

- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations a second label of a Kohn-Sham orbital (as o, with δ_{op} the Kronecker delta of the two labels). Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)}H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- ${\mathcal{E}_{(p)}}^{M}{}_{N}$: The Lovelock tensor of order p: the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to the metric ($\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor); with frame indices its frame components. Unit: $1/\text{length}^{2p}$. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -7H^2$: with the corrected weights the value for which the static block is a vacuum (its Lovelock tensors add to $7H^2$ in every direction). The author's original value was $\Lambda = -10H^2$ (he writes the pure number $\Lambda_{\mathrm{N}} = 10$ in $0 = -\Lambda_{\mathrm{N}} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$). Unit: GeV^2. Value: $-7H^2$; the author's original $-10H^2$. Status: AUTHOR (the term), DERIVED (the value).

- $w_{(p)}$: The Lovelock weights, $w_{(1)}=1$, $w_{(2)}=1/20$, $w_{(3)}=1/360$: the only weights for which gravitational waves travel at the speed of light in the bulk and on the fold (both coefficients of the bulk wave, $1-40w_{(2)}+360w_{(3)}$ and $1-24w_{(2)}+72w_{(3)}$, vanish). The author's original weights $w_{(2)}=1/32$, $w_{(3)}=1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) made bulk gravitons faster than light; they were corrected at the author's order of 2026-10-25. Unit: dimensionless. Value: $(1, 1/20, 1/360)$; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), DERIVED (the values).
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)}=w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda=-7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- y: The curvature of a seven-dimensional space of constant curvature $-\mathsf{y}H^2$ (in units of $-H^2$), times the time line: $\mathsf{y}=1$ is the static block. Unit: dimensionless.

A cosmological constant adds the same number to every direction, so it can cancel the left-hand side only if the warped and the time entries are equal. With the original weights they are not (155/16 along the warped directions and 161/16 along the time before Λ), and no value of Λ makes the static block a vacuum. Status: DERIVED (ST1 by computer algebra on the static block, ST9; `um3_weights.py`, W4, for general weights).

6.2 The stabiliser

With the original weights and $\Lambda=-10H^2$ the left-hand side of FE-1 at the static block is therefore, on both sides of the fold and with no entry that mixes two directions,

$$
\begin{aligned}
&w_{(2)}=\frac{1}{32},\ w_{(3)}=\frac{1}{1152},\ \Lambda=-10H^2:\\
&\sum_{p=1}^{3}\alpha_{(p)}\,\mathcal{E}_{(p)}{}^{a}{}_{b}+\Lambda\,\delta^{a}{}_{b}=H^2\,\mathrm{diag}\Big(-\frac{5}{16},-\frac{5}{16},-\frac{5}{16},-\frac{5}{16},\frac{1}{16},-\frac{5}{16},-\frac{5}{16},-\frac{5}{16}\Big)=\kappa\,\mathcal{T}_{\mathrm{stab}}{}^{a}{}_{b},\qquad\text{(SB-9)}\\
&\mathcal{T}_{\mathrm{stab}}{}^{a}{}_{b}=\mathrm{diag}\big(p_{\mathrm{s}},p_{\mathrm{s}},p_{\mathrm{s}},p_{\mathrm{s}},-\rho_{\mathrm{s}},p_{\mathrm{s}},p_{\mathrm{s}},p_{\mathrm{s}}\big),\qquad \kappa\rho_{\mathrm{s}}=-\frac{H^2}{16},\qquad \kappa p_{\mathrm{s}}=-\frac{5H^2}{16}
\end{aligned}
$$

Symbols in (SB-9).

- $w_{(p)}$: The Lovelock weights, $w_{(1)}=1$, $w_{(2)}=1/20$, $w_{(3)}=1/360$: the only weights for which gravitational waves travel at the speed of light in the bulk and on the fold (both coefficients of the bulk wave, $1-40w_{(2)}+360w_{(3)}$ and $1-24w_{(2)}+72w_{(3)}$, vanish). The author's original weights $w_{(2)}=1/32$, $w_{(3)}=1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) made bulk gravitons faster than light; they were corrected at the author's order of 2026-10-25. Unit: dimensionless. Value: $(1, 1/20, 1/360)$; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), DERIVED (the values).
- Λ: The eight-dimensional cosmological constant, $\Lambda=-7H^2$: with the corrected weights the value for which the static block is a vacuum (its Lovelock tensors add to $7H^2$ in every direction). The author's original value was $\Lambda=-10H^2$ (he writes the pure number $\Lambda_{\mathrm{N}}=10$ in $0=-\Lambda_{\mathrm{N}}+\sum_p H^{-2p}w_{(p)}\mathcal{E}_{(p)}$). Unit: GeV^2. Value: $-7H^2$; the author's original $-10H^2$. Status: AUTHOR (the term), DERIVED (the value).
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)}=w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda=-7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- p: The Lovelock order $p=1,2,3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations a second label of a Kohn-Sham orbital (as o, with δ_{op} the Kronecker delta of the two labels). Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)}=w_{(p)}H^{2-2p}$ ($\alpha_{(1)}=1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- $\mathcal{E}_{(p)}{}^{M}{}_{N}$: The Lovelock tensor of order p: the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to the metric ($\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor); with frame indices its frame components. Unit: 1/length^{2p}. Status: STANDARD.
- a: A frame (orthonormal) index running over $0,1,\ldots,7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab}=\mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a,b\in\{1,\ldots,7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.

- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with upper and lower index strings the generalised Kronecker delta $\delta^{M_1\cdots M_n}_{N_1\cdots N_n}$, the determinant of the $n \times n$ matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\mathrm{stab}}{}^a{}_b$: The energy-momentum tensor of the stabiliser of the bulk (frame components): the source that a static bulk of the author's metric required with the author's original Lovelock weights, found and named by Claude (Anthropic) on 2026-10-24; with the corrected weights and $\Lambda = -7H^2$ it is zero (IN-7). Unit: GeV^8. Value: 0 with the corrected weights. Status: DERIVED.
- p_s: The pressure of the stabiliser of the bulk with the author's original weights (equal along σ, ordinary space and the extra times). Unit: GeV^8. Value: $\kappa p_\mathrm{s} = -5H^2/16$ with the author's original weights; 0 with the corrected weights. Status: DERIVED.
- ρ_s: The energy density of the stabiliser of the bulk with the author's original weights; with frozen extra times it was the energy density of the bulk at which ordinary space stands still. Unit: GeV^8. Value: $\kappa\rho_\mathrm{s} = -H^2/16$ with the author's original weights; 0 with the corrected weights. Status: DERIVED.

in the order (σ, x_1, x_2, x_3, x_4, x_5, x_6, x_7). The static bulk was therefore not a vacuum: the field equations held only if the bulk contained a source whose energy-momentum tensor is this left-hand side divided by κ, a medium at rest in the language of SB-3, without any flow, with the energy density ρ_s and the same pressure $p_\mathrm{s} = 5\rho_\mathrm{s}$ in every direction but the time. Calling this source a medium of its own, the stabiliser, assumes that dirac16complex contributes nothing to the static bulk (its energy-momentum tensor $\mathcal{T}_\Psi$ zero or negligible there; ASSUMED). Status: DERIVED (ST1).

The kink of the static block needed, with the original weights, the source of the fold

$$\kappa\,\mathcal{S}^{(i)}{}_{(i)} = \kappa\,\mathcal{S}^{(r)}{}_{(r)} = -\frac{61}{8}H, \qquad \kappa\,\mathcal{S}^{(4)}{}_{(4)} = -\frac{31}{4}H \qquad \text{(original weights, static block)} \tag{SB-10}$$

Symbols in (SB-10).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)} \mathcal{E}_{(p)}{}^a{}_b$ (FD-11, the junction conditions); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -\frac{32}{5}H\delta^a{}_b$ at the static block. Status: DERIVED.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

along ordinary space, the time and the extra times. Order by order (in units of H): Einstein $(-10, -12, -10)$, which is Israel's $-([K^a{}_b] - \delta^a{}_b[K])$ with the jump $-2H$ of the extrinsic curvature along the six warped directions; Gauss-Bonnet $(5/2, 5, 5/2)$; cubic $(-1/8, -3/4, -1/8)$. It splits into a tension $61H/8$ (pressures equal to minus the energy density) and a pressureless part $H/8$. With the corrected weights the Gauss-Bonnet and cubic parts are $(4, 8, 4)$ and $(-2/5, -12/5, -2/5)$, and the total is the pure tension $-32H/5$ in all seven directions (FE-32 below). Status: DERIVED (ST1 and ST9, from the junction conditions FD-11; `um3_fold.py`, J5).

6.3 Why it is called the stabiliser, and who found it

It was the energy content without which the bulk of the author's metric could not stay static: it *stabilised* the bulk in the sense of keeping it static. (Whether the static bulk with the stabiliser was also stable, that is whether small

disturbances die away, is a separate question that was not analysed.) The stabiliser was not part of the author's specification. Claude (Anthropic's AI model, working in Claude Code) found the requirement SB-9 on 2026-10-24 while deriving the complete coupled field equations of UM-3, introduced the medium to meet it, and named it the stabiliser of the bulk. The name and the idea of a medium that holds a static block had appeared before in this repository, in the two earlier universe models, for their own metrics and with other properties (the provenance gives the lineage). Status: DERIVED (the requirement), ASSUMED (the medium).

6.4 What kind of medium it was

Its energy density and its pressure were negative. $\rho_\mathrm{s} = -H^2/(16\kappa)$, and the pressure five times as large, $p_\mathrm{s} = 5\rho_\mathrm{s}$, the same along the hidden direction, ordinary space and the extra times. It did not flow.

It violated both energy conditions.

$$\kappa\rho_\mathrm{s} = -\frac{H^2}{16} < 0, \qquad \kappa\big(\rho_\mathrm{s} + p_\mathrm{s}\big) = -\frac{3H^2}{8} < 0 \tag{SB-11}$$

Symbols in (SB-11).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- ρ_s: The energy density of the stabiliser of the bulk with the author's original weights; with frozen extra times it was the energy density of the bulk at which ordinary space stands still. Unit: GeV^8. Value: $\kappa\rho_\mathrm{s} = -H^2/16$ with the author's original weights; 0 with the corrected weights. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- p_s: The pressure of the stabiliser of the bulk with the author's original weights (equal along σ, ordinary space and the extra times). Unit: GeV^8. Value: $\kappa p_\mathrm{s} = -5H^2/16$ with the author's original weights; 0 with the corrected weights. Status: DERIVED.

The second inequality, for light moving along the time and one space-like direction, is the violation of the null energy condition; it is what made the stabiliser exotic. The negative cosmological constant of the bulk violates only the weak condition, since its energy density plus its pressure is zero. Status: DERIVED (ST5).

Among homogeneous scalar fields, only a ghost reproduced it. A scalar field that fills the bulk uniformly and changes only with the time has a kinetic energy K_f and a potential energy V_f per volume, and its energy density and pressure (the same in every direction but the time) are

$$\rho_\mathrm{med} = K_\mathrm{f} + V_\mathrm{f}, \qquad p_\mathrm{med} = K_\mathrm{f} - V_\mathrm{f}; \qquad p_\mathrm{med} = 5\rho_\mathrm{med}: \quad V_\mathrm{f} = -\frac{2}{3}K_\mathrm{f}, \quad \rho_\mathrm{med} = \frac{K_\mathrm{f}}{3} \tag{SB-12}$$

Symbols in (SB-12).

- ρ_med: The energy density (energy per volume) of a medium at rest. Unit: GeV^8. Status: STANDARD.
- K_f: The kinetic energy per volume of a field that fills the bulk uniformly and changes only with the time x_4: half the square of its rate of change for an ordinary field (never negative), minus that for a field whose kinetic term has the opposite sign (a ghost). Unit: GeV^8. Status: STANDARD.
- V_f: The potential energy per volume of a field that fills the bulk uniformly and changes only with the time x_4. Unit: GeV^8. Status: STANDARD.
- p_med: The pressure of a medium at rest, the same along every direction but the time (for a medium of the bulk: along σ, ordinary space and the extra times). Unit: GeV^8. Status: STANDARD.

(derived by varying the action of the field with respect to the metric, ST5). An ordinary field has $K_\mathrm{f} \geq 0$ and so, with a pressure five times its energy density, a positive energy density. Only a field whose kinetic energy is negative, a so-called ghost, with $K_\mathrm{f} = 3\rho_\mathrm{s}$ and $V_\mathrm{f} = -2\rho_\mathrm{s}$, reproduced the stabiliser; on the static bulk it would roll at a constant speed on a flat stretch of its potential. A scalar field that depends on σ instead of the time could not do it, because its pressure along σ differs from its pressure along the other directions unless it is constant, and a constant field is a vacuum energy with $p = -\rho$. Vector fields and other kinds of fields were not examined. Status: DERIVED (ST5, for homogeneous scalar fields).

If it had been a simple fluid, its sound would have been faster than light. For a fluid whose pressure is a fixed multiple of its energy density, small compressions travel at the speed of sound c_s with $c_\mathrm{s}^2 = \mathrm{d}p/\mathrm{d}\rho$; for the stabiliser this is 5, so its sound would have travelled $\sqrt{5} \approx 2.24$ times faster than light. The ghost field instead propagates at the speed of light, but its quanta carry negative energy. A general theorem (Dubovsky, Grégoire, Nicolis and Rattazzi, 2006) states that, under broad conditions, a medium that violates the null energy condition has either signals faster than light or unstable excitations. Status: STANDARD (the theorem and the fluid formula), DERIVED (their application).

It was conserved by itself on the static bulk. The covariant divergence SB-4 of the constant tensor of SB-9 vanishes on the static block, in every component (ST4). The reason: its pressure along the hidden direction equals its pressure along every warped direction, and the time, along which its entry differs, is not warped. Unlike the air of SB-7, the stabiliser needed no change of pressure along the hidden direction: because the time is not warped there is no gravitational pull along σ. Status: DERIVED (ST4).

6.5 How it would have evolved on its own

A medium that interacts only through gravity obeys SB-4. For a medium that fills the bulk uniformly, with the energy density $\rho_\mathrm{med}(x_4)$ and the pressure $p_\mathrm{med}(x_4)$ in every warped direction, in the author's metric with arbitrary $a_4(x_4)$ and $b_4(x_4)$, the time component of SB-4 reads (ST6)

$$\rho'_\mathrm{med} = -3\,\frac{b'_4}{b_4}\,\big(\rho_\mathrm{med} + p_\mathrm{med}\big); \qquad p_\mathrm{med} = 5\rho_\mathrm{med}: \quad \rho_\mathrm{med}\, b_4^{18} = \text{constant} \tag{SB-13}$$

Symbols in (SB-13).

- ρ_med: The energy density (energy per volume) of a medium at rest. Unit: GeV^8. Status: STANDARD.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a'_4 = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.
- p_med: The pressure of a medium at rest, the same along every direction but the time (for a medium of the bulk: along σ, ordinary space and the extra times). Unit: GeV^8. Status: STANDARD.

The growth of ordinary space drops out, because ordinary space grows with e^{a_4} and the extra times with $\mathrm{e}^{-a_4} b_4$, so that the volume of all six warped directions together grows with b_4^3. Two consequences follow. A stabiliser that is absent ($\rho = 0$) stays absent. And with frozen extra times (b_4 proportional to e^{a_4}) a stabiliser on its own would have been diluted as $\mathrm{e}^{-18 a_4}$ while the universe expands. Status: DERIVED (ST6).

6.6 What the original equations required when the universe expanded

With frozen extra times, the condition of the late universe ($\mathsf{U} = \mathsf{A}$: the size $\mathrm{e}^{-a_4} b_4$ of the extra times constant, so that the couplings of the Standard Model stay constant), the original equations required in the bulk and on the fold (with $\mathsf{A}_1 = a''_4/H^2$ the rate of change of A):

$$\begin{aligned}
\mathsf{U} = \mathsf{A}: \qquad & \frac{\kappa\rho_\mathrm{bulk}}{H^2} = \frac{3\mathsf{A}^2 - 1}{16}, & & \frac{\kappa\,{\mathcal{T}_\mathrm{bulk}}^{(0)}{}_{(0)}}{H^2} = -\frac{3\mathsf{A}^4 + 3\mathsf{A}^2\mathsf{A}_1 + 18\mathsf{A}^2 + 3\mathsf{A}_1 + 5}{16}, \\
& \frac{\kappa\,{\mathcal{T}_\mathrm{bulk}}^{(i)}{}_{(i)}}{H^2} = -\frac{9\mathsf{A}^2 + 2\mathsf{A}_1 + 5}{16}, & & \frac{\kappa\,{\mathcal{T}_\mathrm{bulk}}^{(r)}{}_{(r)}}{H^2} = -\frac{3\mathsf{A}^4 + 3\mathsf{A}^2\mathsf{A}_1 + 24\mathsf{A}^2 + 3\mathsf{A}_1 + 5}{16}, \\
& \frac{\kappa\,{\mathcal{T}_\mathrm{bulk}}^{(0)}{}_{(4)}}{H^2} = \mathrm{sgn}(\sigma)\,\frac{3\mathsf{A}\big(1 + \mathsf{A}^2\big)}{16}, & & \frac{\kappa\,{\mathcal{S}}^{(4)}{}_{(4)}}{H} = -\frac{10\mathsf{A}^2 + 31}{4}, \\
& \frac{\kappa\,{\mathcal{S}}^{(i)}{}_{(i)}}{H} = -\frac{57\mathsf{A}^2 + 40\mathsf{A}_1 + 183}{24}, & & \frac{\kappa\,{\mathcal{S}}^{(r)}{}_{(r)}}{H} = -\frac{2\mathsf{A}^4 + 2\mathsf{A}^2\mathsf{A}_1 + 37\mathsf{A}^2 + 20\mathsf{A}_1 + 61}{8}
\end{aligned} \tag{SB-14}$$

Symbols in (SB-14).

- U: The rate of change of b_4 in units of H, $\mathsf{U} = b'_4/(H b_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times.

- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); 1 at the author's vacuum, which is not a solution with the corrected weights.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- ρ_{bulk}: The energy density of the bulk sources, $\rho_{\mathrm{bulk}} = -{\mathcal{T}_{\mathrm{bulk}}}^{(4)}{}_{(4)}$. Unit: GeV^8. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- $\mathcal{T}_{\mathrm{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); ${\mathcal{T}_{\mathrm{bulk}}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- sgn: The sign function: $\mathrm{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)}{\mathcal{E}_{(p)}}^a{}_b$ (FD-11, the junction conditions); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -\frac{32}{5}H\delta^a{}_b$ at the static block. Status: DERIVED.

($\rho_{\mathrm{bulk}} = -{\mathcal{T}_{\mathrm{bulk}}}^{(4)}{}_{(4)}$ is the energy density of all bulk sources together.) For $\mathsf{A} \to 0$ they reduce to SB-9 and SB-10. The entry ${\mathcal{T}_{\mathrm{bulk}}}^{(0)}{}_{(4)}$ is a flow of energy along the hidden direction, equal in size on the two sides of the fold and opposite in sign, so it jumps across the fold by $\frac{3}{8}\mathsf{A}(1+\mathsf{A}^2)H^2/\kappa$; the energy current, minus this entry, points towards the fold on both sides. Status: DERIVED (ST6, with its A_1 terms; `um3_fold.py`, J5 and J7).

Where the energy came from. With frozen extra times only ordinary space grows, and the volume of the bulk grows at the rate $3a_4'$. The required bulk source had, for constant A, an energy density plus pressure along ordinary space that is negative:

$$\begin{aligned} &\kappa\big(\rho_{\mathrm{bulk}} + {\mathcal{T}_{\mathrm{bulk}}}^{(1)}{}_{(1)}\big) = -\frac{3}{8}\big(1+\mathsf{A}^2\big)H^2 \qquad (\mathsf{U} = \mathsf{A},\ \mathsf{A}_1 = 0), \\ &\int_{-\infty}^{\infty} \mathrm{d}\sigma\ \mathrm{e}^{-6H|\sigma|}\ \big(-3a_4'\big)\,\kappa\big(\rho_{\mathrm{bulk}} + {\mathcal{T}_{\mathrm{bulk}}}^{(1)}{}_{(1)}\big) = \frac{3}{8}\,\mathsf{A}\big(1+\mathsf{A}^2\big)H^2 \end{aligned} \tag{SB-15}$$

Symbols in (SB-15).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- ρ_{bulk}: The energy density of the bulk sources, $\rho_{\mathrm{bulk}} = -{\mathcal{T}_{\mathrm{bulk}}}^{(4)}{}_{(4)}$. Unit: GeV^8. Status: DERIVED.
- $\mathcal{T}_{\mathrm{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); ${\mathcal{T}_{\mathrm{bulk}}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); 1 at the author's vacuum, which is not a solution with the corrected weights.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless.

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.

Left to itself, a medium of the bulk changes its energy density at the rate $-3a_4'(\rho + p)$ while only ordinary space grows. The required source, with $\rho + p < 0$, would have seen its negative energy density rise towards zero; since the field equations kept its energy density fixed, it had to give off the energy $-3a_4'(\rho_{\mathrm{bulk}} + \mathcal{T}_{\mathrm{bulk}}{}^{(1)}{}_{(1)})$ per volume and time. Integrated over the hidden direction with the volume factor $\mathrm{e}^{-6H|\sigma|}$ of the six warped directions, this is exactly the jump of the flow: all the energy that the bulk gave off arrived at the fold. This is a consistency identity (it follows from the Bianchi identity), not an independent result. Status: DERIVED (ST6).

Where the energy went. The source of the fold took up exactly the energy that arrived from the bulk, for any $a_4(x_4)$:

$$-\big(\kappa\,\mathcal{S}^{(4)}{}_{(4)}\big)' + 3a_4'\big(\kappa\,\mathcal{S}^{(i)}{}_{(i)} - \kappa\,\mathcal{S}^{(4)}{}_{(4)}\big) = \kappa\Big(\mathcal{T}_{\mathrm{bulk}}{}^{(0)}{}_{(4)}\big|_{\sigma=0^+} - \mathcal{T}_{\mathrm{bulk}}{}^{(0)}{}_{(4)}\big|_{\sigma=0^-}\Big) \qquad (\mathsf{U} = \mathsf{A}) \qquad \text{(SB-16)}$$

Symbols in (SB-16).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)}\mathcal{E}_{(p)}{}^a{}_b$ (FD-11, the junction conditions); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -\frac{32}{5}H\delta^a{}_b$ at the static block. Status: DERIVED.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\mathcal{T}_{\mathrm{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); $\mathcal{T}_{\mathrm{bulk}}{}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); 1 at the author's vacuum, which is not a solution with the corrected weights.

($-\mathcal{S}^{(4)}{}_{(4)}$ is the energy density of the source of the fold and $\mathcal{S}^{(i)}{}_{(i)}$ its pressure along ordinary space; 0^+ and 0^- mark the limits from above and from below.) This is the conservation law at the fold, FE-11, again an identity of the junction conditions. At the static block the earlier version of UM-3 split the source of the fold into a tension $\lambda_{\mathrm{F}} = 61HV_*/(8\kappa)$ and a pressureless part $\rho_{\mathrm{d}} = HV_*/(8\kappa)$ (per unit volume of ordinary space, V_* the volume of the extra times), and the pressureless part needed exactly the energy $3a_4'\rho_{\mathrm{d}} = \frac{3}{8}\mathsf{A}H^2V_*/\kappa$, the part of the jump linear in A. Reading this as an exchange of energy between the stabiliser and a fold fluid is an interpretation: what the

equations fix is only the total bulk source and its flow; a medium of the bulk that couples only to gravity could also carry its flow into a thin layer of its own (ASSUMED). Moreover, ordinary matter on the fold needs metric functions that depend on σ, which the separable metric of the author does not have (`um3_fold.py`, J7), so these energy balances describe the fold sources that the separable metric requires, not ordinary matter. Status: DERIVED (ST6), ASSUMED (the interpretation). The conservation law at the fold, for every metric of UM-3:

$$\begin{aligned} &\nabla_M \,\mathcal{T}_\Psi{}^M{}_N = 0 \qquad (\sigma \neq 0), \\ &\hat{\nabla}_a \,\mathcal{S}^a{}_b = -\Big(\mathcal{T}_{\text{bulk}}{}^{(0)}{}_b\big|_{\sigma=0^+} - \mathcal{T}_{\text{bulk}}{}^{(0)}{}_b\big|_{\sigma=0^-}\Big) \end{aligned} \tag{FE-11}$$

Symbols in (FE-11).

- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols $\Gamma^P{}_{MN}$). Status: STANDARD.
- $\mathcal{T}_\Psi$: The energy-momentum tensor of dirac16complex, $\mathcal{T}_{MN} = -(2/\sqrt{|g|})\,\delta S_\Psi/\delta g^{MN}$. Unit: GeV^8. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \operatorname{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\hat{\nabla}$: The covariant derivative of the fold: with the Levi-Civita connection of the metric of the fold (for four-dimensional tensors of its metric $\hat{h}_{\mu\nu} = \operatorname{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the observable coordinates $x_1, \ldots, x_4$; for $\hat{\nabla}_a \mathcal{S}^a{}_b$ of the induced metric of the seven-dimensional fold). Status: STANDARD.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)} \mathcal{E}_{(p)}{}^a{}_b$ (FD-11, the junction conditions); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -\frac{32}{5} H \delta^a{}_b$ at the static block. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \operatorname{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- $\mathcal{T}_{\text{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); $\mathcal{T}_{\text{bulk}}{}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.

6.7 The constraints of the original equations

Two components of the bulk equations contain no second time derivative of the metric, for any $a_4(x_4)$ and $b_4(x_4)$: the energy constraint (the time component, like the Friedmann equation of cosmology) and the flow of energy along the hidden direction. With the original weights they were

$$\begin{aligned} &\kappa\,\mathcal{T}_{\text{bulk}}{}^{(4)}{}_{(4)} = \frac{H^2}{16}\,\mathsf{P}, \qquad \kappa\,\mathcal{T}_{\text{bulk}}{}^{(0)}{}_{(4)} = \frac{3}{16}\,\mathsf{U}\,\mathcal{G}\,H^2\operatorname{sgn}(\sigma), \qquad \mathsf{V} = \mathsf{U} - \mathsf{A}, \\ &\mathsf{P} = 1 - 3\mathsf{A}^2 - 3\mathsf{V}^2 - 9\mathsf{AV} - 27\mathsf{A}^2\mathsf{V}^2 - 9\mathsf{A}^3\mathsf{V} - 9\mathsf{AV}^3 - 5\mathsf{A}^3\mathsf{V}^3, \qquad \mathcal{G} = 1 + \mathsf{A}^2 + 8\mathsf{AV} + \mathsf{V}^2 + 5\mathsf{A}^2\mathsf{V}^2 \end{aligned} \tag{SB-17}$$

Symbols in (SB-17).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\text{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); $\mathcal{T}_{\text{bulk}}{}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.

- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- $\mathsf{P}(\mathsf{A},\mathsf{U})$: The polynomial of the energy constraint of the bulk with the author's original Lovelock weights: the (x_4, x_4) component of the bulk equations of the author's metric was $\kappa {\mathcal{T}_{\text{bulk}}}^{(4)}{}_{(4)} = (H^2/16)\,\mathsf{P}(\mathsf{A},\mathsf{U})$, whatever the time derivatives of the rates (with the corrected weights it is $\mathsf{A}^3(\mathsf{A}-\mathsf{U})^3H^2$). Unit: dimensionless. Value: $\mathsf{P}(0,0) = 1$, $\mathsf{P}(\pm1, 0) = 0$. Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times.
- $\mathcal{G}$: The factor of the (σ, x_4) bulk equation of the author's metric with the author's original weights, $\mathcal{G} = 5\mathsf{A}^2(\mathsf{A}-\mathsf{U})^2 - 6\mathsf{A}^2 + 6\mathsf{A}\mathsf{U} + \mathsf{U}^2 + 1$. Unit: dimensionless. Status: DERIVED.
- sgn: The sign function: $\operatorname{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \operatorname{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- V: The rate of change of the scale factor $\mathrm{e}^{-a_4}b_4$ of the extra times on the fold in units of H, $\mathsf{V} = \mathsf{U} - \mathsf{A}$; zero with frozen extra times. Unit: dimensionless.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); 1 at the author's vacuum, which is not a solution with the corrected weights.

written with the rate A of ordinary space and the rate V of the scale factor $\mathrm{e}^{-a_4}b_4$ of the extra times; both polynomials are symmetric under the exchange of A and V. A bulk without any source needs $\mathsf{P} = 0$ and $\mathsf{U}\mathcal{G} = 0$ at every instant. For $\mathsf{U} = 0$, $\mathsf{P} = (\mathsf{A}^2-1)^2(5\mathsf{A}^2+1)$ vanishes only at $\mathsf{A} = \pm1$; for $\mathsf{U} \neq 0$ both polynomials would need a common root, but their resultant has no real root except $\mathsf{U} = 0$. **With the original weights, a bulk without any source was therefore at every instant the author's vacuum** $(\mathsf{A},\mathsf{U}) = (1,0)$ **or its time reverse** $(-1,0)$, whatever the time dependence of the rates. A late universe that expands slowly with constant couplings needed a bulk source. With frozen extra times ($\mathsf{V} = 0$) $\mathsf{P} = 1 - 3\mathsf{A}^2$ for any $a_4(x_4)$, and the energy constraint became a Friedmann equation of the bulk, $3a_4'^2 = H^2 + 16\kappa\rho_{\text{bulk}} = 16\kappa(\rho_{\text{bulk}} - \rho_{\text{s}})$: the energy density of the stabiliser was the energy density of the bulk at which ordinary space stands still. Status: DERIVED (ST3, ST8).

7 The correction of the weights removes the stabiliser

7.1 Gravitons must travel at the speed of light

A gravitational wave is a small ripple of the metric that travels. The simplest one, of the "+" polarisation, multiplies the metric components along x_1 and x_2 by $\mathrm{e}^{h_{\text{gw}}}$ and $\mathrm{e}^{-h_{\text{gw}}}$ and travels along x_3. Put into the field equations of the static bulk and kept to first order in h_{gw}, the Lovelock terms give it the wave equation

$$\frac{c_{\text{t}}}{2}\frac{\partial^2 h_{\text{gw}}}{\partial x_4^2} - \frac{c_{\text{x}}}{2}\left(\mathrm{e}^{2H|\sigma|}\frac{\partial^2 h_{\text{gw}}}{\partial x_3^2} + \frac{\partial^2 h_{\text{gw}}}{\partial \sigma^2} - 6H\operatorname{sgn}(\sigma)\frac{\partial h_{\text{gw}}}{\partial \sigma}\right) = 0, \qquad \text{(FE-68)}$$
$$c_{\text{t}} = 1 - 40w_{(2)} + 360w_{(3)}, \qquad c_{\text{x}} = 1 - 24w_{(2)} + 72w_{(3)}$$

Symbols in (FE-68).

- c_{t}: The coefficient of the second time derivative of a transverse traceless gravitational wave in the bulk at the static block, in units of its value with Einstein gravity alone: $c_{\text{t}} = 1 - 40w_{(2)} + 360w_{(3)}$ (FE-68). Unit: dimensionless. Value: 0 with the corrected weights; 1/16 with the author's original weights. Status: DERIVED.
- ∂: A partial derivative: ∂_M with respect to the coordinate X^M, ∂_μ with respect to x^μ, ∂_σ with respect to σ, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- h_{gw}: The amplitude of a gravitational wave of the + polarisation travelling along x_3: the metric components along x_1 and x_2 multiplied by $\mathrm{e}^{h_{\text{gw}}}$ and $\mathrm{e}^{-h_{\text{gw}}}$; a function of σ, x_3 and x_4 in the bulk, and on the fold the gravitational wave of the four-dimensional world. Unit: dimensionless. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- c_{x}: The coefficient of the second derivatives along ordinary space and along σ of the same wave, in the same units: $c_{\text{x}} = 1 - 24w_{(2)} + 72w_{(3)}$ (FE-68); bulk gravitons move with the local speed $\sqrt{c_{\text{x}}/c_{\text{t}}}$. Unit: dimensionless. Value: 0 with the corrected weights; 5/16 with the author's original weights. Status: DERIVED.

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- sgn: The sign function: $\mathrm{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.
- $w_{(p)}$: The Lovelock weights, $w_{(1)} = 1$, $w_{(2)} = 1/20$, $w_{(3)} = 1/360$: the only weights for which gravitational waves travel at the speed of light in the bulk and on the fold (both coefficients of the bulk wave, $1 - 40w_{(2)} + 360w_{(3)}$ and $1 - 24w_{(2)} + 72w_{(3)}$, vanish). The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) made bulk gravitons faster than light; they were corrected at the author's order of 2026-10-25. Unit: dimensionless. Value: $(1, 1/20, 1/360)$; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), DERIVED (the values).

with a time-kinetic coefficient c_t and a gradient coefficient c_x (∂ marks a partial derivative). In the bulk the wave moves with the local speed $\sqrt{c_\mathrm{x}/c_\mathrm{t}}$ times the speed of light. Einstein gravity alone gives $c_\mathrm{t} = c_\mathrm{x} = 1$, the speed of light; the Gauss-Bonnet and cubic terms change the two coefficients differently, because the static block is curved along σ, ordinary space and the extra times but not along the time. With the author's original weights $c_\mathrm{t} = 1/16$ and $c_\mathrm{x} = 5/16$: gravitational waves in the bulk moved $\sqrt{5} = 2.24$ times faster than light. At the fold the wave has a thin part of its own, and a wave of long wavelength collects both:

$$\int_{0^-}^{0^+} \mathrm{d}\sigma\,(\dots) = \frac{16\big(w_{(2)} - 6w_{(3)}\big)}{H}\,\frac{\partial^2 h_\mathrm{gw}}{\partial x_4^2} - \frac{12\big(w_{(2)} - 2w_{(3)}\big)}{H}\,\frac{\partial^2 h_\mathrm{gw}}{\partial x_3^2} - \frac{c_\mathrm{x}}{2}\Big(\frac{\partial h_\mathrm{gw}}{\partial\sigma}\Big|_{0^+} - \frac{\partial h_\mathrm{gw}}{\partial\sigma}\Big|_{0^-}\Big) = \kappa\,\Pi_\mathrm{gw},$$

$$\kappa K_{44} = \frac{c_\mathrm{t}}{6H} + \frac{16\big(w_{(2)} - 6w_{(3)}\big)}{H}, \qquad \kappa K_{33} = \frac{c_\mathrm{x}}{4H} + \frac{12\big(w_{(2)} - 2w_{(3)}\big)}{H}, \qquad \kappa\big(K_{44} - K_{33}\big) = -\frac{c_\mathrm{t}}{12H}, \qquad c_\mathrm{g}^2 = \frac{K_{33}}{K_{44}} \tag{FE-69}$$

Symbols in (FE-69).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $w_{(p)}$: The Lovelock weights, $w_{(1)} = 1$, $w_{(2)} = 1/20$, $w_{(3)} = 1/360$: the only weights for which gravitational waves travel at the speed of light in the bulk and on the fold (both coefficients of the bulk wave, $1 - 40w_{(2)} + 360w_{(3)}$ and $1 - 24w_{(2)} + 72w_{(3)}$, vanish). The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) made bulk gravitons faster than light; they were corrected at the author's order of 2026-10-25. Unit: dimensionless. Value: $(1, 1/20, 1/360)$; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), DERIVED (the values).
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- ∂: A partial derivative: ∂_M with respect to the coordinate X^M, ∂_μ with respect to x^μ, ∂_σ with respect to σ, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- h_gw: The amplitude of a gravitational wave of the + polarisation travelling along x_3: the metric components along x_1 and x_2 multiplied by $\mathrm{e}^{h_\mathrm{gw}}$ and $\mathrm{e}^{-h_\mathrm{gw}}$; a function of σ, x_3 and x_4 in the bulk, and on the fold the gravitational wave of the four-dimensional world. Unit: dimensionless. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- c_x: The coefficient of the second derivatives along ordinary space and along σ of the same wave, in the same units: $c_\mathrm{x} = 1 - 24w_{(2)} + 72w_{(3)}$ (FE-68); bulk gravitons move with the local speed $\sqrt{c_\mathrm{x}/c_\mathrm{t}}$. Unit: dimensionless. Value: 0 with the corrected weights; 5/16 with the author's original weights. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.

- $\Pi_{\rm gw}$: The transverse traceless stress of the matter of the fold that drives the gravitational wave $h_{\rm gw}$ (half the difference of its components along x_1 and x_2), integrated across the fold (eight-dimensional normalisation). Unit: GeV^7. Status: STANDARD.
- K_{44}: The coefficient of $\partial^2 h_{\rm gw}/\partial x_4^2$ in the equation of the long-wavelength gravitational wave (the same at every σ) at the static block, integrated across the hidden direction with the volume factor $\mathrm{e}^{-6H|\sigma|}$: the part of the bulk plus the thin part at the fold. Unit: GeV^5 (κK_{44} is a length). Status: DERIVED.
- $c_{\rm t}$: The coefficient of the second time derivative of a transverse traceless gravitational wave in the bulk at the static block, in units of its value with Einstein gravity alone: $c_{\rm t} = 1 - 40w_{(2)} + 360w_{(3)}$ (FE-68). Unit: dimensionless. Value: 0 with the corrected weights; 1/16 with the author's original weights. Status: DERIVED.
- K_{33}: The coefficient of $-\partial^2 h_{\rm gw}/\partial x_3^2$ in the same equation, with the same two parts; the wave travels at $c_{\rm g}^2 = K_{33}/K_{44}$. Unit: GeV^5. Status: DERIVED.
- $c_{\rm g}$: The speed of gravitational waves on the fold, in units of the speed of light. Unit: dimensionless. Value: 1 at every wavelength with the corrected weights; with the author's original weights the long-wavelength mode had $c_{\rm g}^2 = 83/82$ and shorter waves were slower than light. Status: DERIVED.

For every pair of weights the difference of the two coefficients of the long wave is $-c_{\rm t}/(12\kappa H)$: it travels at the speed of light only if $c_{\rm t} = 0$. If $c_{\rm t}$ were zero but not $c_{\rm x}$, waves in the bulk would have no time-kinetic term at all, an infinite local speed. Gravitons travel at the speed of light everywhere only if both coefficients vanish: $1-40w_{(2)}+360w_{(3)} = 0$ and $1 - 24w_{(2)} + 72w_{(3)} = 0$, two linear equations with the single solution $w_{(2)} = 1/20$, $w_{(3)} = 1/360$. With the original weights the long wave had $c_{\rm g}^2 = 83/82$, faster than light (the bulk part alone 15/2: $c_{\rm g} = 2.74$, a figure that an earlier version of UM-3 quoted), and the wave bound to the fold was slower than light at short wavelengths. With the corrected weights the bulk part vanishes, $\kappa K_{44} = \kappa K_{33} = 8/(15H)$, and $c_{\rm g} = 1$ at every wavelength. Status: DERIVED (`um3_weights.py`, W1 to W3, W7; `um3_bulk.py`, B5).

7.2 The static block with the corrected weights

With $w_{(2)} = 1/20$ and $w_{(3)} = 1/360$ the warped and the time entries of the static block are both $7H^2$ before the cosmological constant (the last column of the table of SB-8; FE-70), so the cosmological constant $\Lambda = -7H^2$ makes the static block a vacuum:

$$\kappa\,\mathcal{T}_{\rm stab}{}^{a}{}_{b} = \Big(\sum_{p=1}^{3}\alpha_{(p)}\,\mathcal{E}_{(p)}{}^{a}{}_{b} + \Lambda\,\delta^{a}{}_{b}\Big)\Big|_{\text{static block}} = 0 : \qquad \text{(IN-7)}$$
$$(15-9+1-7)H^2 = 0 \text{ (warped directions)}, \qquad (21-21+7-7)H^2 = 0 \text{ (the time)}$$

Symbols in (IN-7).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\rm stab}{}^{a}{}_{b}$: The energy-momentum tensor of the stabiliser of the bulk (frame components): the source that a static bulk of the author's metric required with the author's original Lovelock weights, found and named by Claude (Anthropic) on 2026-10-24; with the corrected weights and $\Lambda = -7H^2$ it is zero (IN-7). Unit: GeV^8. Value: 0 with the corrected weights. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations a second label of a Kohn-Sham orbital (as o, with δ_{op} the Kronecker delta of the two labels). Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)}H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- $\mathcal{E}_{(p)}{}^{M}{}_{N}$: The Lovelock tensor of order p: the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to the metric ($\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor); with frame indices its frame components. Unit: 1/length^{2p}. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -7H^2$: with the corrected weights the value for which the static block is a vacuum (its Lovelock tensors add to $7H^2$ in every direction). The author's original value was $\Lambda = -10H^2$ (he writes the pure number $\Lambda_{\rm N} = 10$ in $0 = -\Lambda_{\rm N} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$). Unit: GeV^2. Value: $-7H^2$; the author's original $-10H^2$. Status: AUTHOR (the term), DERIVED (the value).

- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with upper and lower index strings the generalised Kronecker delta $\delta^{M_1\cdots M_n}_{N_1\cdots N_n}$, the determinant of the $n \times n$ matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

The stabiliser of SB-9 is zero: order by order, Einstein $(15, 21)$, Gauss-Bonnet $(-9, -21)$, cubic $(1, 7)$ and the cosmological constant $(-7, -7)$ add to zero along the warped directions and along the time. The source of the fold of the static block becomes a pure tension, the same in all seven directions of the fold:

$$\mathsf{A} = \mathsf{U} = \mathsf{A}_1 = \mathsf{U}_1 = 0: \qquad \kappa\,{\mathcal{T}_{\mathrm{bulk}}}^a{}_b = 0, \qquad \kappa\,{\mathcal{S}^{(i)}}_{(i)} = \kappa\,{\mathcal{S}^{(4)}}_{(4)} = \kappa\,{\mathcal{S}^{(r)}}_{(r)} = -\frac{32}{5}H \tag{FE-32}$$

Symbols in (FE-32).

- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); 1 at the author's vacuum, which is not a solution with the corrected weights.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\mathrm{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); ${\mathcal{T}_{\mathrm{bulk}}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)}{\mathcal{E}_{(p)}}^a{}_b$ (FD-11, the junction conditions); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -\frac{32}{5}H\delta^a{}_b$ at the static block. Status: DERIVED.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as ${\mathcal{S}^{(i)}}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

This is the arrangement of Randall and Sundrum (FD-8): an empty bulk with a negative cosmological constant and a fold of pure tension, like the drum skin of example 5. Status: DERIVED (ST9; `um3_weights.py`, W4; `um3_fold.py`, J5).

7.3 The expanding fold with the corrected weights

The corrected equations have no need of a stabiliser in any era. Their two constraints are

$$\begin{aligned} &\kappa\,{\mathcal{T}_{\mathrm{bulk}}}^{(4)}{}_{(4)} = \mathsf{A}^3(\mathsf{A} - \mathsf{U})^3H^2, \qquad \kappa\,{\mathcal{T}_{\mathrm{bulk}}}^{(0)}{}_{(4)} = 3\mathsf{A}^2\mathsf{U}(\mathsf{A} - \mathsf{U})^2H^2\,\mathrm{sgn}(\sigma), \\ &\text{no bulk source:} \quad \mathsf{A} = 0 \ \text{ or } \ \mathsf{U} = \mathsf{A} \quad \text{at every instant} \end{aligned} \tag{FE-66}$$

Symbols in (FE-66).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\text{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); ${\mathcal{T}_{\text{bulk}}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); 1 at the author's vacuum, which is not a solution with the corrected weights.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- sgn: The sign function: $\operatorname{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \operatorname{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

so a bulk without any source has, at every instant, either static ordinary space ($\mathsf{A} = 0$) or frozen extra times ($\mathsf{U} = \mathsf{A}$), and on both branches every component of the bulk equations vanishes. The expanding fold of our universe is on the second branch, where the bulk is an exact vacuum for every expansion history and the fold obeys exactly the Friedmann equations of four-dimensional general relativity (the equations of the expansion of a homogeneous universe, $3a_4'^2 = \kappa_7\rho$ for the energy density ρ and $-(2a_4'' + 3a_4'^2) = \kappa_7 p$ for the pressure p of its matter above the tension, both in the normalisation of the thin source):

$$
\begin{gathered}
\mathsf{U} = \mathsf{A}, \quad \mathsf{U}_1 = \mathsf{A}_1 \ \left(\mathrm{e}^{-a_4}b_4 = \text{constant},\ \mathcal{Z} = 1\right): \qquad \kappa\,{\mathcal{T}_{\text{bulk}}}^{a}{}_{b} = 0 \ \text{ for any } a_4(x_4), \\
\frac{\kappa\,{\mathcal{S}^{(i)}}_{(i)}}{H} = -\frac{16}{15}\left(6 + 3\mathsf{A}^2 + 2\mathsf{A}_1\right), \qquad \frac{\kappa\,{\mathcal{S}^{(4)}}_{(4)}}{H} = -\frac{16}{5}\left(2 + \mathsf{A}^2\right), \qquad \frac{\kappa\,{\mathcal{S}^{(r)}}_{(r)}}{H} = -\frac{4}{5}\left(8 + 8\mathsf{A}^2 + \mathsf{A}^4 + 4\mathsf{A}_1 + \mathsf{A}^2\mathsf{A}_1\right), \\
3a_4'^2 = -\kappa_7\left({\mathcal{S}^{(4)}}_{(4)} + \frac{32H}{5\kappa}\right), \qquad -\left(2a_4'' + 3a_4'^2\right) = \kappa_7\left({\mathcal{S}^{(i)}}_{(i)} + \frac{32H}{5\kappa}\right), \qquad \kappa_7 = \frac{15}{16}\,\kappa H
\end{gathered}
\tag{FE-33}
$$

Symbols in (FE-33).

- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); 1 at the author's vacuum, which is not a solution with the corrected weights.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4) = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z} = 1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\text{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); ${\mathcal{T}_{\text{bulk}}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric

$\eta_{ab} = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1,\dots,7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.

- b: A frame index (as a). Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)} \mathcal{E}_{(p)}{}^a{}_b$ (FD-11, the junction conditions); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -\frac{32}{5}H\delta^a{}_b$ at the static block. Status: DERIVED.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- κ_7: The gravitational coupling of the fold: with the corrected Lovelock weights the thin source of the fold equals, to second order in the rates of expansion, the Einstein tensor of the seven-dimensional metric of the fold divided by $\kappa_7 = 15\kappa H/16$, and exactly so in the Friedmann equations of a fold with frozen extra times; four-dimensional gravity on the fold has $8\pi G_\mathrm{N} = \kappa_7/V_*$ (plus the induced term, if the fields of the fold induce one). Unit: GeV^{-5}. Value: $15\kappa H/16$. Status: DERIVED.

There is no flow of energy along the hidden direction, so the matter of the fold is conserved by itself (FE-11 with a vanishing right-hand side). The author's vacuum, in which the original weights needed no source, now needs one:

$$a_4 = Hx_4, \quad b_4 = \text{constant } (\mathsf{A} = 1,\ \mathsf{U} = \mathsf{A}_1 = \mathsf{U}_1 = 0): \qquad \frac{\kappa\,\mathcal{T}_\mathrm{bulk}{}^a{}_b}{H^2} = \mathrm{diag}\Big(-\frac{7}{5}, -\frac{7}{5}, -\frac{7}{5}, -\frac{7}{5}, 1, -\frac{7}{5}, -\frac{7}{5}, -\frac{7}{5}\Big),$$
$$\kappa\rho_\mathrm{bulk} = -H^2, \qquad \kappa\,\mathcal{S}^{(i)}{}_{(i)} = \kappa\,\mathcal{S}^{(r)}{}_{(r)} = -\frac{42}{5}H, \qquad \kappa\,\mathcal{S}^{(4)}{}_{(4)} = -\frac{28}{5}H \tag{FE-31}$$

Symbols in (FE-31).

- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); 1 at the author's vacuum, which is not a solution with the corrected weights.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_\mathrm{bulk}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); $\mathcal{T}_\mathrm{bulk}{}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.

- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- ρ_{bulk}: The energy density of the bulk sources, $\rho_{\mathrm{bulk}} = -\mathcal{T}_{\mathrm{bulk}}{}^{(4)}{}_{(4)}$. Unit: GeV^8. Status: DERIVED.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)} \mathcal{E}_{(p)}{}^a{}_b$ (FD-11, the junction conditions); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -\frac{32}{5} H \delta^a{}_b$ at the static block. Status: DERIVED.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.

so the primordial era of UM-3 is no longer the author's vacuum but an inflation of the fold on the branch of frozen extra times, driven by the vacuum energy of a field of the fold, as in general relativity. Status: DERIVED (ST9; `um3_weights.py`, W5 and W6; `um3_geometry.py`, G4).

7.4 What takes the place of the stabiliser

Nothing in the bulk: the corrected bulk is a vacuum on both branches. But one requirement remains, on the fold. The component of the junction along the extra times (the third line of FE-33) asks the matter of the fold for a definite pressure along the extra times, which radiation has and dust and vacuum energy do not. Without it the volume of the extra times would change, and with it Newton's constant and every coupling of the fold: since the time of nucleosynthesis Newton's constant would have fallen by the factor 15.8, and today it would change 6800 times faster than lunar laser ranging allows. UM-3 therefore assumes that the fold fluid, the thin medium of the fold, supplies the missing pressure q_{hold}:

$$
\begin{aligned}
&p_{\mathrm{ff}} = -\rho_{\mathrm{ff}}, \qquad q_{\mathrm{ff}} = -\rho_{\mathrm{ff}} + q_{\mathrm{hold}}, \qquad \lambda_{\mathrm{F}} = \frac{32 H V_*}{5\kappa}, \\
&q_{\mathrm{hold}} = -\Delta_{\mathrm{r}} - \frac{4}{5}\big(\mathsf{A}^4 + \mathsf{A}^2\mathsf{A}_1\big)\frac{H V_*}{\kappa}, \qquad \Delta_{\mathrm{r}} = q_{\mathrm{SM}} + q_\chi + q_{\mathrm{P}} - \big(\rho_{\mathrm{ff}} - \lambda_{\mathrm{F}}\big) + \frac{\rho - 3\mathcal{P}}{2}
\end{aligned}
\tag{FE-65}
$$

Symbols in (FE-65).

- p_{ff}: The pressure of the fold fluid along ordinary space. Unit: GeV^4. Value: $-\lambda_{\mathrm{F}} = -32HV_*/(5\kappa)$. Status: ASSUMED (the fluid), DERIVED (the value).
- ρ_{ff}: The energy density of the fold fluid. Unit: GeV^4. Value: $\lambda_{\mathrm{F}} = 32HV_*/(5\kappa)$ (its tension). Status: ASSUMED (the fluid), DERIVED (the value).
- q_{ff}: The pressure of the fold fluid along the extra times: minus its tension plus the pressure q_{hold} that holds the volume of the extra times fixed. Unit: GeV^4. Value: $-\lambda_{\mathrm{F}} + q_{\mathrm{hold}}$; $-32HV_*/(5\kappa)$ at the static block. Status: ASSUMED (the fluid), DERIVED (the value).
- q_{hold}: The pressure along the extra times that the fold fluid adds to its tension to hold the volume of the extra times fixed (four-dimensional normalisation). Unit: GeV^4. Value: $-\Delta_{\mathrm{r}} = -0.793\,\rho_{\mathrm{c}}^{(0)}$ today. Status: DERIVED (the value), ASSUMED (that the fold fluid supplies it).
- λ_{F}: The tension of the fold: the part of the fold fluid whose pressures along ordinary space and along the extra times are minus its energy density (a vacuum energy of the seven-dimensional fold), per unit volume of ordinary space of the fold (four-dimensional normalisation). Unit: GeV^4. Value: $32HV_*/(5\kappa)$. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- Δ_r: The mismatch of the radion condition: the sum over the matter of the fold of its pressure along the extra times and $(\rho - 3\mathcal{P})/2$ (zero for radiation). Unit: GeV^4. Value: $0.793\,\rho_\mathrm{c}^{(0)}$ today. Status: DERIVED.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); 1 at the author's vacuum, which is not a solution with the corrected weights.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless.
- q_SM: The pressure of the Standard Model along the extra times at the fold: the mean on-shell Lagrangian density of the Standard-Model fields (the extra-time component of the energy-momentum tensor of every term of the fold is its Lagrangian, because the fields of the fold do not depend on the extra times). Unit: GeV^4. Status: DERIVED.
- q_χ: The pressure of the bound states of dirac16complex along the extra times, integrated across the fold (their on-shell Lagrangian, the self-interaction energy per volume; zero for a dilute gas). Unit: GeV^4. Status: DERIVED.
- q_P: The pressure of the portals along the extra times: their mean on-shell Lagrangian density. Unit: GeV^4. Status: DERIVED.
- ρ: The total energy density of the matter of the fold (all species); $\Delta\rho$ its perturbation. Unit: GeV^4. Status: STANDARD.
- $\mathcal{P}$: The total pressure of the matter of the fold; $\Delta\mathcal{P}$ its perturbation. Unit: GeV^4. Status: STANDARD.

The equation of the volume of the extra times (the radion) shows what q_hold does:

$$H^2\big(\mathsf{V}_1 + 3\mathsf{V}(\mathsf{A}+\mathsf{V})\big) = \frac{8\pi G_\mathrm{N}}{5}\big(\rho - 3\mathcal{P} + 2q_\mathrm{tot}\big) \tag{FE-72}$$

Symbols in (FE-72).

- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- V_1: The time derivative of V in units of H, $\mathsf{V}_1 = \mathsf{V}'/H$. Unit: dimensionless.
- V: The rate of change of the scale factor $\mathrm{e}^{-a_4}b_4$ of the extra times on the fold in units of H, $\mathsf{V} = \mathsf{U} - \mathsf{A}$; zero with frozen extra times. Unit: dimensionless.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); 1 at the author's vacuum, which is not a solution with the corrected weights.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_N: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_\mathrm{N} = 1/M_\mathrm{Pl}^2$ with the Planck mass $M_\mathrm{Pl} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- ρ: The total energy density of the matter of the fold (all species); $\Delta\rho$ its perturbation. Unit: GeV^4. Status: STANDARD.
- $\mathcal{P}$: The total pressure of the matter of the fold; $\Delta\mathcal{P}$ its perturbation. Unit: GeV^4. Status: STANDARD.
- q_tot: The total pressure along the extra times of the matter of the fold (the Standard Model, the bound states of dirac16complex, the portals and the fold fluid without its tension). Unit: GeV^4. Status: STANDARD.

The rate V of the scale factor of the extra times stays zero only if $\rho - 3\mathcal{P} + 2q_\mathrm{tot} = 0$, and q_hold makes this so. It acts as a potential that holds the size of the extra times, a role that the "block stabiliser" of the first universe model of this repository played for its own metric; what produces it is OPEN. Status: DERIVED (the requirement and its value, `um3_series.py`, SR4 to SR6), ASSUMED (that the fold fluid supplies it), OPEN (its origin).

8 The five points, verified

[1] The stabiliser couples only to gravity. By its definition it had no term in the action of UM-3 other than its own: it entered the field equations only through its energy-momentum tensor in FE-1, and it exerted no force on

dirac16complex or on the Standard Model except through the curvature that it caused. On the static bulk this was consistent: the stabiliser was conserved by itself (ST4). In an expanding universe the bulk source that the original equations required released energy (SB-15) that flowed into the fold (SB-14) and was taken up there (SB-16). What the equations fixed was the total bulk source and its flow; whether this meant an exchange of energy between the stabiliser and the matter of the fold, or a second medium of the bulk, or a thin layer of the stabiliser itself, they did not say (ASSUMED, an interpretation). **Verdict, original weights:** exact on the static bulk; in an expanding universe a medium that couples only to gravity and keeps $p = 5\rho$ was not enough to supply the required source (SB-13 against SB-14). **Verdict, corrected weights:** there is no stabiliser, and the bulk of an expanding fold with frozen extra times is a vacuum without any flow (FE-33). Status: DERIVED (ST4, ST6, ST9).

[2] It is absent in the primordial era, whose bulk is the author's vacuum (FE-31). With the original weights the author's vacuum, $a_4 = Hx_4$ with constant b_4 ($\mathsf{A} = 1$, $\mathsf{U} = 0$), solved the field equations of the bulk with nothing on the right (ST2), and with the stabiliser present it would not have been a solution: the stabiliser was absent there. But a stabiliser that couples only to gravity and keeps $p = 5\rho$ cannot appear later out of nothing (SB-13). Points [1] and [2] together required a mechanism that created it between the primordial era and the late universe: an exchange of energy with the fold (which breaks [1]), a conversion from or into dirac16complex in the bulk (the bulk source is the sum of both), or a change of its equation of state (which a single medium with $p = 5\rho$ does not have); UM-3 contained none. **Verdict, original weights:** [2] true, with the creation of the stabiliser OPEN. **Verdict, corrected weights:** the author's vacuum is not a solution (FE-31); the primordial era is an inflation of the fold with frozen extra times, whose bulk is a vacuum; no era of UM-3 has a stabiliser, so the question of its creation no longer arises. Status: DERIVED (ST2, ST6, ST9; `um3_primordial.py`, PR0).

[3] Its existence and value are DERIVED from the requirement of a static bulk. With the original weights the value was DERIVED: SB-9 fixed every entry (ST1). The existence of a bulk source was DERIVED more generally than for an exactly static bulk: without any bulk source the bulk was at every instant the author's vacuum or its time reverse (SB-17, ST3, ST8), so a slowly expanding bulk needed one. What was ASSUMED is that the source is a separate medium, with dirac16complex contributing nothing; its microscopic nature was OPEN, and among homogeneous scalar fields only a ghost reproduced it (SB-12, ST5). **Verdict:** value DERIVED; the existence of a bulk source DERIVED; its identification with a separate medium ASSUMED. With the corrected weights the same derivation gives the value zero (IN-7, ST9): the existence of the stabiliser was a consequence of the original weights.

[4] The static bulk is defined in SB-5 and illustrated by five examples and a comparison of static and empty spacetimes in the part "The static bulk".

[5] Warp and warp factor are defined in SB-2 and illustrated by four examples in the part "Warp and warp factor".

9 The importance of the stabiliser for UM-3

With the original weights, no late universe like ours without it. A bulk without any source was at every instant the author's vacuum or its time reverse (SB-17): ordinary space had to grow or shrink at the full rate H. The only alternative to the stabiliser would have been to make the author's vacuum itself the late universe, with H equal to today's expansion rate $H_4^{(0)}$. But in the author's vacuum the extra times shrink, as $\mathrm{e}^{-a_4} b_4 = \mathrm{e}^{-Hx_4} b_4$, and every coupling of the fold is divided by the volume of the extra times, so every squared coupling of the Standard Model grows as e^{3Hx_4}. The natural nuclear reactor of Oklo (Gabon), which ran about 1.8 billion years ago, bounds the change of the fine-structure constant α_{em} since then to about one part in ten million ($-0.9 \times 10^{-7} < \Delta\alpha_{\mathrm{em}}/\alpha_{\mathrm{em}} < 1.2 \times 10^{-7}$). The author's vacuum would change it by

$$\Delta \ln \alpha_{\mathrm{em}} = 3H_4^{(0)}\, \Delta x_4 = 0.372 \qquad \text{(SB-18)}$$

Symbols in (SB-18).

- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- α_{em}: The fine-structure constant. Unit: dimensionless. Value: 1/137.036. Status: STANDARD.

- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.

($H_4^{(0)} = 67.4$ km/s/Mpc, $\Delta x_4 = 1.8$ Gyr; Δ marks a change): 3.1 million times the bound. With the stabiliser and frozen extra times the couplings stayed constant. Status: DERIVED (ST7), STANDARD (the bound of Oklo).

With the original weights, the zero point of the expansion. With frozen extra times the energy constraint was a Friedmann equation of the bulk, $3a_4'^2 = 16\kappa(\rho_{\mathrm{bulk}} - \rho_{\mathrm{s}})$ (SB-17): the energy density of the stabiliser was the energy density of the bulk at which ordinary space stands still, and the expansion measured the excess of the bulk energy density over it. Integrated over the hidden direction and the extra times, per unit volume of ordinary space on the fold, the stabiliser carried the energy

$$\rho_{4,\mathrm{s}} = \frac{\rho_{\mathrm{s}} V_*}{3H} = -\frac{H V_*}{48\kappa} \tag{SB-19}$$

Symbols in (SB-19).

- $\rho_{4,\mathrm{s}}$: The energy of the stabiliser per unit volume of ordinary space of the fold with the author's original weights: its energy density integrated over the hidden direction and the extra times. Unit: GeV^4. Value: $-HV_*/(48\kappa)$ with the author's original weights; 0 with the corrected weights. Status: DERIVED.
- ρ_{s}: The energy density of the stabiliser of the bulk with the author's original weights; with frozen extra times it was the energy density of the bulk at which ordinary space stands still. Unit: GeV^8. Value: $\kappa\rho_{\mathrm{s}} = -H^2/16$ with the author's original weights; 0 with the corrected weights. Status: DERIVED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.

(the volume factor $\mathrm{e}^{-6H|\sigma|}$ of the warped directions, integrated over both sides of the fold, gives $1/(3H)$, and V_* is the physical volume of the extra times on the fold in the late universe). It was not a vacuum energy that makes the fold expand: the static block as a whole did not expand at all. Status: DERIVED (ST7, ST8).

With the original weights, the most exotic ingredient. Its violation of the null energy condition (SB-11) put it among the sources that come with instabilities or with signals faster than light; Einstein's static universe, the example most like it, is unstable. If the stabiliser had been a ghost field, its quanta would carry negative energy, and even through gravity alone a ghost lets the vacuum decay into pairs of ghosts and pairs of ordinary particles, at a rate that is finite only if the momenta of the ghosts are cut off (Cline, Jeon and Moore, 2004). Whether the static block with the stabiliser was stable was never analysed. Status: STANDARD (the general results).

With the corrected weights, its absence changes UM-3. The weights for which gravitons travel at the speed of light make the static bulk a vacuum, and the stabiliser disappears with them. Four consequences:

1. Gravitational waves travel at the speed of light at every wavelength, and the gravitational waves of the neutron-star merger GW170817, which arrived together with its light, are explained for every value of H (FE-69).
2. The fold obeys the Friedmann equations of four-dimensional general relativity exactly, with Newton's constant $8\pi G_{\mathrm{N}} = \kappa_7/V_*$, $\kappa_7 = 15\kappa H/16$ (FE-33): the Gauss-Bonnet and cubic terms give the fold an Einstein-Hilbert term of its own.
3. The bulk needs no exotic medium in any era: no violation of the null energy condition, no ghost, and no question of how the stabiliser would be created after the primordial era.
4. Two new tasks appear. The fold fluid must hold the volume of the extra times fixed with the pressure q_{hold} (FE-65, FE-72), whose origin is OPEN; and the static block is a degenerate vacuum (the double and triple roots of FE-70), around which the bulk equations have no linear part, so the response of the bulk to sources that depend on σ is strongly coupled and not computed (OPEN).

Summary. The stabiliser was the price of a slowly expanding late universe with constant couplings in the author's metric with the author's original weights and cosmological constant. Its value was fixed exactly and introduced no new parameter. It was not a property of the author's metric but of those weights: the weights that make gravitons travel at the speed of light make the static bulk empty, and the stabiliser vanishes with them.

10 Provenance

What comes from the author (AUTHOR). The coordinates $X = \{x_0, \ldots, x_7\}$ and their roles (`Dirac_Einstein-Lovelock_4+4_Nash.nb`, section DEFCHART); the metric with $b_4[x_4]^2$ and its two functions $a_4(x_4)$ and $b_4(x_4)$ (the author's order of 2026-10-24); the Einstein-Lovelock action with its original weights $w_{(1)} = 1$, $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ and the cosmological constant $\Lambda = -10H^2$ (`Dirac_Einstein-Lovelock_4+4_Nash.nb`, In[134], where the author writes the pure number 10); the author's vacuum $a_4 = Hx_4$ with constant b_4; the field dirac16complex; the order of 2026-10-25, 05:09 PDT, to document the stabiliser with proper attribution and to verify the five points; and the order of 2026-10-25, 07:55 PDT: "Gravitons must travel at c. If these incorrect claims come from my choice of the Lovelock couplings and my choice of the cosmological constant Λ THEN my choices are wrong, and you MUST correct them."

What UM-3 adds. The proper distance σ (DERIVED from the metric); the static block as the first approximation of the late universe (ASSUMED); the stabiliser of the original weights (found and named by Claude on 2026-10-24; its value DERIVED, the medium ASSUMED); the corrected weights $w_{(2)} = 1/20$, $w_{(3)} = 1/360$ and $\Lambda = -7H^2$ (DERIVED at the order of 07:55); the fold fluid with its tension and its holding pressure (ASSUMED, with DERIVED values).

The lineage. The first universe model of this repository (the folder `Revision/universe_model`, version 1.1, present at the first commit of the repository, `0fdf375` of 2026-10-19) assumed a "block stabiliser": a potential for the size $\mathrm{e}^{-a_4} b_4^{1/2}$ of the block of the hidden direction and the extra times of its metric (whose extra-time components carried b_4 instead of b_4^2), with an origin left open; that version had itself changed the third weight to 1/768. The second universe model (`universe_model_2`, version 3 of 2026-10-18) assumed a block stabiliser whose energy density falls as the inverse eighth power of the block scale, $\kappa\rho_{\mathrm{stab}} = (26 + 4\sqrt{51})H^2 = 54.566H^2$ at its static point. The stabiliser of UM-3 was a new object, for the author's new metric with b_4^2, with a negative energy density and $p = 5\rho$.

Who found what, and when. On 2026-10-24 Claude computed, in the gravity module of UM-3 (`um3_geometry.py`, check G4, commit `4448938` of 2026-10-24 17:54 UTC-7), that a static bulk of the author's metric needs the energy density $-H^2/(16\kappa)$ and that the bulk with frozen extra times needs the energy density $(3\mathsf{A}^2 - 1)H^2/(16\kappa)$ and a flow of energy along the hidden direction; the pressures $-5H^2/(16\kappa)$ were added when the medium was introduced and named "the stabiliser of the bulk", as the term IN-7 of the action of UM-3, in the commit `170d8d3` of 2026-10-24 19:05 UTC-7 (the field equations of UM-3, where its existence was labelled ASSUMED and its value DERIVED). On 2026-10-25, at the author's order of 05:09, Claude wrote the first version of this document with its LaTeX source and its PDF and the checks ST1 to ST8 (commit `cd5709d` of 2026-10-25 06:55 UTC-7), and an independent referee read it; its corrections (the energy conditions of a negative vacuum energy, the use of "warp", the constraints of a sourceless bulk, the rates of the extra times, the derivation of SB-9, the junction conditions, the lineage and the references) are in this version. After the author's order of 07:55 Claude derived the weights for which gravitons travel at the speed of light (`um3_weights.py`), found that with them the static bulk is a vacuum and the stabiliser vanishes (check ST9), and rewrote this document as the record of the stabiliser and of its removal. All of this work was done in Claude Code sessions configured for the model claude-opus-5-5 (session https://claude.ai/code/session_01AyG7BZ5rkLxBARRjLHHquB).

How it was computed. The left-hand side of the field equations was computed exactly with the computer algebra system sympy for the general diagonal warped metric $\mathrm{d}\sigma^2 + \mathrm{e}^{2\mathfrak{a}}(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2) - \mathrm{e}^{2\mathfrak{n}}\mathrm{d}x_4^2 - \mathrm{e}^{2\mathfrak{b}}(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2)$, with functions $\mathfrak{a}$, $\mathfrak{n}$, $\mathfrak{b}$ of σ and x_4: the curvature in an orthonormal frame, and the Lovelock tensors of orders 1, 2 and 3 contracted with the generalised Kronecker delta. It was then specialised to the author's metric, $\mathfrak{a} = a_4 - H|\sigma|$, $\mathfrak{b} = \ln b_4 - a_4 - H|\sigma|$, $\mathfrak{n} = 0$. The source of the fold was computed from the jump of the first σ-derivatives of the metric across the fold (the junction conditions FD-11), as the difference of a junction potential on the two sides, and independently by smoothing the kink. The program `universe_model_3/code/um3_stabiliser.py` of the repository computes and checks the following:

check	what it verifies	result
ST1	original weights: the static bulk requires the source SB-9 on both sides, without flow; the order-by-order values of SB-8 and its table; the source of the fold SB-10 order by order	PASS
ST2	original weights: the author's vacuum needs no source; its fold source $(-10, -6, -10)H$; the expansion around it	PASS
ST3	original weights: without a source the bulk is at every instant the author's vacuum or its time reverse (the two constraints and their resultant, SB-17)	PASS
ST4	the stabiliser is conserved by itself on the static bulk	PASS
ST5	from the values of ST1: the weak and the null energy condition fail (SB-11); among homogeneous scalar fields only a ghost reproduces it (SB-12)	PASS
ST6	SB-13; the frozen branch SB-14 with its A_1 terms and its fold source; the energy budget SB-15; the tension and pressureless part of the fold source; the balance of the fold SB-16	PASS
ST7	the drift SB-18 in the author's vacuum, and the share SB-19	PASS
ST8	the two constraints contain no second time derivative; their form in (A, V); the Friedmann equation of the bulk	PASS
ST9	corrected weights: the static bulk is a vacuum (IN-7), the fold source is a pure tension (FE-32), the frozen branch and static ordinary space are vacua without flow, the constraints FE-66, the author's vacuum is no longer a solution	PASS

The other checks used here are in the programs `um3_weights.py` (W1 to W8: the gravitational waves FE-68 and FE-69 for any weights, the unique weights, FE-70, the vacua and the fold of the corrected weights), `um3_geometry.py` (G2 to G4: the curvature and the bulk equations of the author's metric), `um3_fold.py` (J4 to J7: the junction conditions and the sources of the fold), `um3_series.py` (SR4 to SR6: the volume of the extra times) and `um3_primordial.py` (PR0). `um3_stabiliser.py` writes `outputs/stabiliser.json` and `reports/stabiliser_checks.json`, reads the general source of the fold from the output of `um3_fold.py`, and runs in about 45 seconds (`python universe_model_3/code/um3_stabiliser.py` from the root of the repository).

Sources outside UM-3 (STANDARD). A. Einstein, "Kosmologische Betrachtungen zur allgemeinen Relativitätstheorie", Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften (1917) 142 (the static universe, SB-6); A. S. Eddington, "On the instability of Einstein's spherical world", Monthly Notices of the Royal Astronomical Society 90 (1930) 668; K. Schwarzschild, "Über das Gravitationsfeld eines Massenpunktes nach der

Einsteinschen Theorie", Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften (1916) 189 (the static field outside a mass); D. Lovelock, "The Einstein tensor and its generalizations", Journal of Mathematical Physics 12 (1971) 498; W. Israel, "Singular hypersurfaces and thin shells in general relativity", Il Nuovo Cimento B 44 (1966) 1 (the junction conditions); S. C. Davis, "Generalised Israel junction conditions for a Gauss-Bonnet brane world", Physical Review D 67 (2003) 024030; E. Gravanis and S. Willison, "Israel conditions for the Gauss-Bonnet theory and the Friedmann equation on the brane universe", Physics Letters B 562 (2003) 118; R. V. Pound and G. A. Rebka, "Apparent weight of photons", Physical Review Letters 4 (1960) 337; L. Randall and R. Sundrum, "A large mass hierarchy from a small extra dimension", Physical Review Letters 83 (1999) 3370, and "An alternative to compactification", Physical Review Letters 83 (1999) 4690 (FD-8); P. Hořava and E. Witten, "Heterotic and type I string dynamics from eleven dimensions", Nuclear Physics B 460 (1996) 506, and "Eleven-dimensional supergravity on a manifold with boundary", Nuclear Physics B 475 (1996) 94 (example 9 of the definition of the fold); H. Whitney, "On singularities of mappings of Euclidean spaces. I. Mappings of the plane into the plane", Annals of Mathematics 62 (1955) 374 (the fold map of example 5); S. W. Hawking and G. F. R. Ellis, "The Large Scale Structure of Space-Time" (Cambridge University Press, 1973) (the energy conditions); S. Dubovsky, T. Grégoire, A. Nicolis and R. Rattazzi, "Null energy condition and superluminal propagation", Journal of High Energy Physics 03 (2006) 025; J. M. Cline, S. Jeon and G. D. Moore, "The phantom menaced: constraints on low-energy effective ghosts", Physical Review D 70 (2004) 043543; the bound of Oklo on the drift of the fine-structure constant over 1.8 Gyr (A. I. Shlyakhter, "Direct test of the constancy of fundamental nuclear constants", Nature 264 (1976) 340; T. Damour and F. Dyson, "The Oklo bound on the time variation of the fine-structure constant revisited", Nuclear Physics B 480 (1996) 37); the Hubble constant $H_4^{(0)} = 67.4$ km/s/Mpc (Planck Collaboration, "Planck 2018 results. VI. Cosmological parameters", Astronomy and Astrophysics 641 (2020) A6); the speed of gravitational waves (B. P. Abbott et al., "Gravitational waves and gamma-rays from a binary neutron star merger: GW170817 and GRB 170817A", Astrophysical Journal Letters 848 (2017) L13).