

dirac16complex in the author's primordial gravitational field

Lagrangian, covariant field equations, non-triviality [1], energy-momentum tensor operator,
equations of state and the field equations for a4[x4] (Revision record)

dirac16complex project (with Claude Opus 5.5)

September 2026

Contents

1	Scope, sources and status of the record	3
2	The primordial gravitational field	6
3	The field dirac16complex: Clifford algebra, Pin(4,4) and Spin(4,4)	7
4	Coupling through the canonical spin connection	8
5	The Lagrangian	10
6	The exact covariant field equations (Euler-Lagrange)	11
7	Non-triviality [1]	13
8	Self-consistency	14
9	The energy-momentum tensor	16
10	Kinetic energy, potential energy, energy density, pressures and equations of state	17
11	Canonical quantisation in 4 + 4 dimensions and the Krein structure	18
12	The energy-momentum tensor operator	20
13	The field equations for a4[x4] with dirac16complex as the source	22
14	Relation to the pairing theorems	26

15 What this document does not claim	27
16 Reproduction	28
17 Index of formulas and checks	29

Abstract

This document records, for the fermion field `dirac16complex` coupled to the author's primordial gravitational field, the Lagrangian, its coupling through the canonical spin connection, the exact covariant field equations (Euler-Lagrange) written out component by component, the non-triviality test [1] with its proof, the self-consistency of the field equations, the energy-momentum tensor (as a Grassmann-algebra expression and as a normal-ordered operator after canonical quantisation in $4 + 4$ dimensions, with the Krein structure that this quantisation forces), the kinetic and potential energy, the energy density, the pressures and the equations of state, and the Einstein-Lovelock field equations for `a4[x4]` with `dirac16complex` as the source. Every formula is an output of Revision code (a WolframScript verifier and, for most statements, an independent sympy verifier) and is listed with the names of the checks that verify it. The main exact results are: in the diagonal vielbein the spin-connection term of the field equation is $\gamma^\mu \Omega_\mu = 3H\gamma^{(8)}$ for every a_4 (the time-direction terms of the three inflating and the three deflating directions cancel, the hidden-direction terms add); this value belongs to the diagonal frame and to the field variables Ψ (it vanishes identically in a frame boosted in the (x_4, x_8) plane, and the rescaling $\Psi = \sin^{-1/2} z \chi$ removes it), while the frame-independent content of non-triviality [1] is that Ω_μ vanishes in no frame (the metric is never flat for $H > 0$) and that the vielbein factors enter every derivative term; when $\Psi^\dagger$ is realised as the Hilbert adjoint, the canonical anticommutator $\{\Psi, \Psi^\dagger\} = B \delta^7 / \sqrt{|g|}$ with B of signature (8,8) forces an indefinite (Krein) inner product (a positive Fock realisation with $\Psi^\dagger = \chi B$ is constructed and checked only for single good-sector momenta with frozen coefficients, `mode_hamiltonian_good_sector`, `Fock_space_good_sector_example`, `good_sector_positive_fock_realisation`; in the author's metric the curved good-sector mode operator is Hermitian only up to the boundary term at $z = \pi/2$, where this record imposes no boundary condition, and without one the good sector has complex frequencies, for $U = 0$ whenever $m^2 < 9H^2$, `good_sector_hermiticity_up_to_the_branes_flux`, `good_sector_x8_independent_modes_without_boundary_condition`; a positive-norm Hilbert space for the full field is not established); homogeneous on-shell configurations have $\rho = mS + U$ and $p = SU' - U$ in every direction; the field equations for a_4 reduce to a constraint, an evolution equation $a_4'' F(a_4') = \kappa(p_3 - p_t)$, a hidden-direction equation and the condition $p_3 + p_t = 2p_8$. Section 15 states what is not claimed, including the frame dependence of $3H\gamma^{(8)}$, the boundary terms at the patch end $z = \pi/2$ and the ill-posedness of the Cauchy problem for data that depend on the extra times. Added on 2026-10-08 from the second wave of the record: the field equations for a_4 with the Kohn-Sham source (section 13: no recorded Kohn-Sham state is an admissible source, and in a stated approximation the source does not start or select the deflation), the completion of the Kohn-Sham-level pairing T3 (section 14), the operator form of the on-shell identity for $\lambda \neq 0$ in a finite fermionic Fock model of one good-sector mode set with frozen coefficients, where, of the tested operator orderings, it holds exactly only when the potential and the energy-momentum tensor are both Wick ordered (section 12; nothing is proved for the field on a whole slice), the verdict of the dark-sector investigation for `dirac16complex` (section 15) and pointers to the documents on the Kohn-Sham model, the dark-sector hypotheses and the Lovelock tensors (section 1).

1 Scope, sources and status of the record

The binding plan is `Revision/SPEC.md`; the author's task is quoted in `Revision/README.md`. This document belongs to the new record under `Revision/` and uses nothing from the earlier stages of the repository: every statement below is an output of Revision code and is cited by the name of the check that verifies it. Numbers and formulas are taken only from the following files (verdicts as recorded in them when this document was written):

part	files	checks
algebra (SPEC section 2)	Revision/algebra /reports/wolfram -algebra.json, Revision /algebra/reports/python- algebra.json	Wolfram: 45 of 45 checks pass; sympy: 35 of 35 checks pass
field theory (SPEC sections 3, 4, 6)	Revision/theory/ field-theory.json, Revision/theory/reports/ wolfram-field-th eory.json, Revision /theory/reports/ python-field-the ory.json	Wolfram: 84 of 84 checks pass; sympy: 70 of 70 checks pass
field equations for a_4 (SPEC section 5)	Revision/field_e quations_a4/a4-e quations.json, Revision /field_equations _a4/reports/wolfram-a4-r eport.json, Revision /field_equations _a4/reports/pyth on-a4-report.json	Wolfram: 52 of 52 checks pass; sympy: 63 of 63 checks pass
pairing (SPEC section 9)	Revision/pairing /pairing-theory.json, Revision/pairing /reports/wolfram -pairing.json, Revision /pairing/reports/python- pairing.json	Wolfram: 101 of 101 checks pass; sympy: 66 of 66 checks pass
scope of the statements (frame dependence, boundary terms, growth, sign of the energy)	Revision/theory/reports/ wolfram-scope.json, Revision/theory/reports/ python-scope.json	Wolfram: 15 of 15 checks pass; sympy: 14 of 14 checks pass
Kohn-Sham pairing T3 (SPEC section 9)	Revision/pairing /kohn_sham/t3-th eory.json, Revision /pairing/kohn_sham/repor ts/wolfram-t3.json, Revision/pairing /kohn_sham/repor ts/python-t3.json	Wolfram: 10 of 10 checks pass; sympy: 13 of 13 checks pass
the Kohn-Sham states as a source of the a_4 equations	Revision/field_e quations_a4/repo rts/ks-source-co nditions.json	Python: 5 of 5 checks pass
the a_4 equations with the Kohn-Sham source (2026-10-08)	Revision/field_e quations_a4/ks_s ource/reports/ks-source- a4.json	Python: 23 of 23 checks pass

part	files	checks
completion of T3 (2026-10-08)	Revision/pairing/ kohn_sham/t3-completion.json, Revision/ pairing/kohn_sham/reports/wolfram-t3-completion.json, Revision/ pairing/kohn_sham/reports/python-t3-completion.json	Wolfram: 3 of 3 checks pass; sympy: 7 of 7 checks pass
the energy-momentum tensor operator for $\lambda \neq 0$ in a finite Fock model (SPEC sections 4, 6; 2026-10-08)	Revision/theory/ fock_quartic/reports/fock-quartic.json	Python (sympy and its own exact engine for the anticommutation relations; no Wolfram counterpart): 21 of 21 checks pass
dark sector, Hypothesis (SPEC section 8)	Revision/dark_sector/dirac16complex/reports/derivation-checks.json, Revision/dark_sector/dirac16complex/reports/ks-history-run.json, Revision/dark_sector/dirac16complex/reports/eos-checks.json, Revision/ dark_sector/dirac16complex/reports/independent-checks.json	derivation: 30 of 30 checks pass; Kohn-Sham history: 5 of 5 checks pass; equation of state: 13 of 13 checks pass; independent: 9 of 9 checks pass

The parts with a Wolfram and a sympy count have two verifiers that share no code: a WolframScript verifier (exact symbolic arithmetic) and a sympy verifier (an independent re-derivation; the gamma matrices are rebuilt from the author’s formulas on both sides and compared entry by entry). Where only one engine verifies a statement, only that engine’s check is cited. A check is cited in typewriter type, for example `nontriviality_1_dirac16complex` (Wolfram) or `gamma_mu_Omega_mu_equals_3H_gamma_x8` (sympy); in the field-theory reports the Wolfram names end in `_G` for `dirac16complex` (explicit Grassmann algebra) and the sympy names begin with `grassmann_`. The proofs are exact identities in the stated generality (symbolic H , $a_4(x_4)$, m , λ , and a general field configuration on its jet space); exact numerical examples are labelled as examples.

Related Revision documents. The Kohn-Sham model of `dirac16complex` in the deflating field (SPEC section 7: the reduction to 2×2 blocks in the hidden coordinate, the instantaneous ground and first excited states along the prescribed history $a_4 = AHx_4$, Mermin thermodynamics, and the full cross-check of the Rust solver against the independent reference, `Revision/kohn_sham/reports/ks-crosscheck.json`: 31 of 31 checks pass) is described in `Revision/docs/KOHN_SHAM_DEFLATING_FIELD`. The generalized Kronecker delta and the Lovelock tensors $P_{(1)}$, $P_{(2)}$, $P_{(3)}$ behind the field equations for a_4 of section 13 are described in `Revision/docs/LOVELOCK_GKD`. The dark-sector hypotheses of SPEC section 8 are investigated in `Revision/docs/DARK_SECTO`

R_HYPOTHESES (its verdict for dirac16complex is summarised in section 15). The pairing theorems, with the completion of T3, are in Revision/docs/PAIR_CREATION_PROOFS.

Status of the comparison record of the field-theory branch. The sympy report contains a comparison with the current Wolfram side (comparison_with_wolfram): 32 of 32 formula records and 63 of 63 check pairs agree, against a Wolfram run of 84 checks. Every formula record of Revision/theory/field-theory.json is compared (the record energy_exchange, three Wolfram expressions, component by component against the divergence recomputed from the sympy Christoffel symbols; the prose records nontriviality, majorana_negative_control, exact_solutions, equation_of_state_definitions and hidden_direction_hermiticity by re-deriving their formulas in sympy), and every one of the 84 Wolfram checks is paired with a sympy check (the list wolfram_checks_without_sympy_counterpart of the comparison is empty; the Grassmann and the commuting form of the nonlinear homogeneous solution, exact_solution_nonlinear_homogeneous_G and exact_solution_nonlinear_homogeneous_C, are both paired with exact_nonlinear_homogeneous_solution, which checks both statistics). The sympy verifier exits with an error unless the comparison agrees entirely, so a stale comparison cannot pass unnoticed. This document still cites each formula by the checks of each engine separately.

Conventions (SPEC sections 1 and 2): coordinates $x_1, \dots, x_8$ as the author names them; x_1, x_2, x_3 are ordinary 3-space, x_4 is the time, x_5, x_6, x_7 are the three extra times (time-like, deflating exponentially as a_4 increases), x_8 is the hidden space direction; $z = 6Hx_8 \in (0, \pi/2)$, $H > 0$; $a_4 = a_4(x_4)$, $a'_4 = da_4/dx_4$; $\partial_a = \partial/\partial x_a$. Frame indices are aligned with the coordinates and written in parentheses: $\gamma^{(a)}$ is the frame gamma matrix of direction x_a . Throughout, $i \in \{1, 2, 3\}$ labels the 3-space directions and $t \in \{5, 6, 7\}$ the extra times.

2 The primordial gravitational field

The author's metric (SPEC section 1), with signature (4,4) (space-like x_1, x_2, x_3, x_8 ; time-like $x_4, \dots, x_7$):

$$ds^2 = e^{2a_4} \sin^{1/3} z (dx_1^2 + dx_2^2 + dx_3^2) - dx_4^2 - e^{-2a_4} \sin^{1/3} z (dx_5^2 + dx_6^2 + dx_7^2) + \cot^2 z dx_8^2, \quad z = 6Hx_8 \in (0, \pi/2).$$

The scale factor of 3-space is $e^{a_4} \sin^{1/6} z$ (inflating as a_4 increases), that of the extra times $e^{-a_4} \sin^{1/6} z$ (exponentially deflating). The diagonal vielbein $e^a{}_\mu = f_a \delta^a_\mu$ with frame metric $\eta_{ab} = \text{diag}(+1, +1, +1, -1, -1, -1, -1, +1)$ has

$$f_1 = f_2 = f_3 = e^{a_4} \sin^{1/6} z, \quad f_4 = 1, \quad f_5 = f_6 = f_7 = e^{-a_4} \sin^{1/6} z, \quad f_8 = \cot z$$

and reproduces the metric entry by entry (metric_is_the_authors, vielbein_inverse; sympy metric_from_vielbein_equals_SPEC). The volume element is

$$\sqrt{|g|} = f_1 f_2 \cdots f_8 = \sin z \cot z = \cos z,$$

independent of x_4 : the inflation e^{3a_4} of 3-space and the deflation e^{-3a_4} of the extra times compensate (sqrt_det_g_is_cos_z, sqrt_det_g_equals_cos_z). $H = 0$ is not a member of the family but

a degenerate limit ($g_{11} \rightarrow 0$, $g_{88} \rightarrow \infty$ at fixed x_8 ; `degenerate_at_H_0`); every statement below assumes $H > 0$ and $0 < z < \pi/2$.

The Levi-Civita connection has 25 independent nonzero symbols (`christoffel_count`, `christoffel_symmetric_metric_compatible`; formula key `christoffel_nonzero`):

$$\begin{aligned}\Gamma^{x_i}_{x_i x_4} &= a'_4, & \Gamma^{x_i}_{x_i x_8} &= H \cot z, \\ \Gamma^{x_t}_{x_4 x_t} &= -a'_4, & \Gamma^{x_t}_{x_t x_8} &= H \cot z, \\ \Gamma^{x_4}_{x_i x_i} &= e^{2a_4} \sin^{1/3} z a'_4, & \Gamma^{x_4}_{x_t x_t} &= e^{-2a_4} \sin^{1/3} z a'_4, \\ \Gamma^{x_8}_{x_i x_i} &= -e^{2a_4} H \frac{\sin^{4/3} z}{\cos z}, & \Gamma^{x_8}_{x_t x_t} &= e^{-2a_4} H \frac{\sin^{4/3} z}{\cos z}, \\ \Gamma^{x_8}_{x_8 x_8} &= -\frac{6H}{\sin z \cos z}.\end{aligned}$$

The mixed Ricci tensor is diagonal (`ricci_mixed_components`, `ricci_scalar`, `curvature_nonzero_flat_only_formally`):

$$R^{x_i}_{x_i} = a''_4 - 6H^2, \quad R^{x_4}_{x_4} = 6(a'_4)^2, \quad R^{x_t}_{x_t} = -a''_4 - 6H^2, \quad R^{x_8}_{x_8} = -6H^2,$$

$$R = 6((a'_4)^2 - 7H^2).$$

Since $R^{x_8}_{x_8} = -6H^2 < 0$ for every $H > 0$ and every function a_4 (a constant a_4 included), the metric is never flat for $H > 0$ (`never_flat_for_H_positive`).

3 The field `dirac16complex`: Clifford algebra, $\text{Pin}(4,4)$ and $\text{Spin}(4,4)$

`dirac16complex` is a field Ψ with 16 complex anticommuting components $\Psi_A(x)$, $A = 1, \dots, 16$ (elements of a Grassmann algebra; after quantisation, operators with canonical anticommutators, section 11). It transforms under a 16-dimensional irreducible representation of $\text{Pin}(4,4)$, the double cover of $\text{O}(4,4)$; for the transformations of determinant 1 it transforms under the direct sum of two inequivalent irreducible 8-dimensional representations of $\text{Spin}(4,4)$, the double cover of $\text{SO}(4,4)$. These representation facts are proved exactly below for the author's gamma matrices.

The gamma matrices are the author's real 16×16 matrices `T16`, rebuilt by Revision code from the author's formulas (the tau matrices of the notebook) in WolframScript and, independently, in Python, with the notebook's frame order mapped to the author's coordinates (`SPEC` section 2): $\gamma^{(8)} = \text{T16}[0]$, $\gamma^{(1,2,3)} = \text{T16}[1..3]$, $\gamma^{(4)} = \text{T16}[4]$, $\gamma^{(5,6,7)} = \text{T16}[5..7]$ (`T16_block_form`, `coordinate_map`; the two constructions agree entry by entry, `fixture_comparison_gammas_json`). Exact properties:

- Clifford relation $\{\gamma^{(a)}, \gamma^{(b)}\} = 2\eta^{ab} I_{16}$ for all 64 pairs; the squares are +1 for x_1, x_2, x_3, x_8 and -1 for $x_4, \dots, x_7$ (`Clifford_relation`, `clifford_relation`, `clifford_relation_sympy`). Every $\gamma^{(a)}$ is a real signed permutation matrix, symmetric for the space-like and antisymmetric

for the time-like directions, $(\gamma^{(a)})^T = \eta_{aa}\gamma^{(a)}$ (reality, signed_permutation_matrices, symmetry_pattern).

- $C = \gamma^{(8)}\gamma^{(1)}\gamma^{(2)}\gamma^{(3)}$ (the notebook's sigma16) is real symmetric with $C^2 = 1$, and every $C\gamma^{(a)}$ is real antisymmetric, equivalently $C\gamma^{(a)}C^{-1} = -(\gamma^{(a)})^T$ (C_real_symmetric, C_squared_identity, C_gamma_real_antisymmetric; sympy C_conjugation). The Dirac adjoint is $\bar{\Psi} = \Psi^\dagger C$ and $S = \bar{\Psi}\Psi$.
- The chirality $\Gamma = \gamma^{(8)}\gamma^{(1)}\dots\gamma^{(7)} = \text{diag}(-I_8, I_8)$ anticommutes with every $\gamma^{(a)}$, $\Gamma^2 = 1$, $[\Gamma, C] = 0$ (Gamma_definition, Gamma_diag, Gamma_anticommutes_with_gammas, Gamma_commutes_with_C; sympy chirality_diag).
- $B = -iC\gamma^{(4)}$ is Hermitian with $B^2 = 1$ and eigenvalues $+1$ and -1 , eight each: signature (8,8) (B_definition, B_Hermitian, B_squared_identity, B_signature_8_8; sympy B_hermitian_involution_signature). B commutes with $\gamma^{(1)}, \gamma^{(2)}, \gamma^{(3)}, \gamma^{(4)}, \gamma^{(8)}$ and anticommutes with $\gamma^{(5)}, \gamma^{(6)}, \gamma^{(7)}$ (B_gamma_relations).
- The 28 generators $S^{ab} = \frac{1}{4}[\gamma^{(a)}, \gamma^{(b)}]$ satisfy the relations of $\text{so}(4,4)$ (S_Lorentz_algebra, S_lorentz_algebra), rotate the gammas as a vector (S_gamma_commutator, S_vector_action), satisfy $(S^{ab})^T C + CS^{ab} = 0$ so that $\bar{\Psi}\Psi$ is $\text{Spin}(4,4)$ -invariant (S_preserves_C), and commute with Γ (S_commutes_with_Gamma). The Hermitian form $\Psi^\dagger B\Psi$ is invariant only under the $\text{Spin}(4,3)$ that fixes x_4 (S_preserves_B_only_off_x4).

Representation theorem (exact). The 256 ordered products of gammas are linearly independent and span the full matrix algebra $\text{Mat}(16)$ (Clifford_basis_spans_full_matrix_algebra, clifford_products_span_M16), and the commutant of the gammas is one-dimensional (Pin44_irreducible_commutant_dim_1, pin_commutant_dimension_1): the 16-dimensional representation is irreducible under $\text{Pin}(4,4)$. The commutant of the 28 generators S^{ab} is two-dimensional, spanned by the chiral projectors $(1 \mp \Gamma)/2$ (Spin44_commutant_dim_2_chiral_projectors, spin_commutant_dimension_2); each chiral half (rows 1 to 8 with $\Gamma = -1$, rows 9 to 16 with $\Gamma = +1$) is irreducible (chiral_halves_irreducible, spin_halves_irreducible); the spaces of intertwiners between the halves are zero (chiral_halves_inequivalent_intertwiners_0, spin_halves_inequivalent); the even subalgebra is $\text{Mat}(8) \oplus \text{Mat}(8)$ (even_subalgebra_dimension, even_products_span_M8_plus_M8). Every $\gamma^{(a)}$, the lift of a reflection (an element of $\text{Pin}(4,4)$ of determinant -1 in $\text{O}(4,4)$), exchanges the two halves (Spin_Pin_reflection_swaps_halves, reflections_exchange_halves). Hence under $\text{Spin}(4,4)$ the 16 splits into $8_- \oplus 8_+$, two inequivalent irreducible 8-dimensional representations, which the reflections combine into the one irreducible representation of $\text{Pin}(4,4)$.

4 Coupling through the canonical spin connection

The coupling to gravity uses the vielbein and the canonical spin connection (SPEC section 3): $\omega_\mu{}^a{}_b$ is defined by the vielbein postulate $\partial_\mu e^a{}_\nu - \Gamma^\lambda_{\mu\nu} e^a{}_\lambda + \omega_\mu{}^a{}_b e^b{}_\nu = 0$, $\omega_{\mu ab} = \eta_{ac}\omega_\mu{}^c{}_b$ is antisymmetric in a, b , and

$$\begin{aligned}\Omega_\mu &= \frac{1}{2}\omega_{\mu ab}S^{ab}, & D_\mu\Psi &= \partial_\mu\Psi + \Omega_\mu\Psi, \\ \gamma^\mu &= e_a{}^\mu\gamma^{(a)} = \gamma^{(\mu)}/f_\mu, & D_\mu\bar{\Psi} &= \partial_\mu\bar{\Psi} - \bar{\Psi}\Omega_\mu.\end{aligned}$$

(This is the canonical connection with the frame metric; the notebook's contraction of the mixed $\omega_\mu{}^a{}_b$, which lacks a metric factor, is not used.) The postulate holds for all 512 components and $\omega_{\mu ab} = -\omega_{\mu ba}$ (`vielbein_postulate` in both field-theory reports, `omega_antisymmetric`, `spin_connection_antisymmetric`). Exactly 12 components with $a < b$ are nonzero (`omega_components`; formula key `omega_nonzero`):

$$\begin{aligned}\omega_{x_i(i)(4)} &= a'_4 e^{a_4} \sin^{1/6} z, & \omega_{x_i(i)(8)} &= H e^{a_4} \sin^{1/6} z, \\ \omega_{x_t(4)(t)} &= -a'_4 e^{-a_4} \sin^{1/6} z, & \omega_{x_t(t)(8)} &= -H e^{-a_4} \sin^{1/6} z,\end{aligned}$$

and $\omega_{x_4} = \omega_{x_8} = 0$. The spinor connection is (`Omega_components`; sympy `Omega_x4_and_Omega_x8_vanish`)

$$\begin{aligned}\Omega_{x_i} &= \frac{1}{2} e^{a_4} \sin^{1/6} z (a'_4 \gamma^{(i)} \gamma^{(4)} + H \gamma^{(i)} \gamma^{(8)}) & (i = 1, 2, 3), \\ \Omega_{x_t} &= -\frac{1}{2} e^{-a_4} \sin^{1/6} z (a'_4 \gamma^{(4)} \gamma^{(t)} + H \gamma^{(t)} \gamma^{(8)}) & (t = 5, 6, 7), \\ \Omega_{x_4} &= \Omega_{x_8} = 0.\end{aligned}$$

It makes the gammas covariantly constant, $D_\mu \gamma^\nu = \partial_\mu \gamma^\nu + \Gamma^\nu_{\mu\lambda} \gamma^\lambda + [\Omega_\mu, \gamma^\nu] = 0$ for all 64 pairs (`gamma_covariantly_constant`, `covariant_constancy_D_mu_gamma_nu`), and its curvature is the Riemann tensor, $[D_\mu, D_\nu] \Psi = \frac{1}{2} R_{ab\mu\nu} S^{ab} \Psi$ (`spin_curvature_equals_Riemann`, `spinor_curvature_equals_riemann`).

The contraction that enters the field equation, per direction (no sum; formula key `gammaOmega_per_direction`):

$$\begin{aligned}\gamma^{x_i} \Omega_{x_i} &= \frac{1}{2} a'_4 \gamma^{(4)} + \frac{1}{2} H \gamma^{(8)} & (i = 1, 2, 3), \\ \gamma^{x_t} \Omega_{x_t} &= -\frac{1}{2} a'_4 \gamma^{(4)} + \frac{1}{2} H \gamma^{(8)} & (t = 5, 6, 7), \\ \gamma^{x_4} \Omega_{x_4} &= \gamma^{x_8} \Omega_{x_8} = 0,\end{aligned}$$

and summed:

$$\gamma^\mu \Omega_\mu = 3H \gamma^{(8)} = \frac{1}{2\sqrt{|g|}} \partial_\mu (\sqrt{|g|} \gamma^\mu)$$

(`gammaOmega_equals_3H_gamma_x8`, `gammaOmega_x4_terms_cancel`, `gammaOmega_divergence_form`). The sympy verifier confirms each of these statements independently (`gamma_mu_Omega_mu_equals_3H_gamma_x8`, `time_terms_cancel_hidden_term_survives` and `divergence_of_sqrtg_gamma`). The field-equations branch verifies the same value again, in a Clifford representation of its own and with T16 (`ownrep_gravity_term`, `authorT16_gravity_term`). For each direction separately $\{\gamma^\mu, \Omega_\mu\} = 0$ (no sum), because Ω_μ contains only generators $S^{(\mu)b}$ (`gamma_mu_anticomutes_with_Omega_mu_no_sum`, `anticommutator_gamma_Omega_vanishes`).

The value $3H\gamma^{(8)}$ belongs to the diagonal vielbein. The canonical spin connection depends on the choice of the local frame, and so does the contraction $\gamma^\mu \Omega_\mu$. Boost the frame in the (x_4, x_8) plane, $e'^{(4)} = \cosh b e^{(4)} + \sinh b e^{(8)}$, $e'^{(8)} = \sinh b e^{(4)} + \cosh b e^{(8)}$, the other $e'^{(a)} = e^{(a)}$, with rapidity $b = \beta x_4 + b_0$: the frame reproduces the author's metric, its canonical connection satisfies the vielbein postulate, and exactly, for every a_4 and H ,

$$\gamma'^\mu \Omega'_\mu = \frac{6H - \beta}{2} (\cosh b \gamma^{(8)} - \sinh b \gamma^{(4)}),$$

which is $3H\gamma^{(8)}$ for the unboosted frame and vanishes identically for $\beta = 6H$. Both engines verify the frame and its connection (`boosted_frame_reproduces_metric` and `boosted_frame_canonical_connection`) and the contraction (`boosted_frame_gammaOmega_formula` and `boosted_frame_gammaOmega_vanishes`). In that frame Ω'_μ is nonzero for $\mu = x_1, \dots, x_7$ and its curvature is nonzero (`boosted_frame_curvature_nonzero`): what no frame removes is Ω_μ itself, whose curvature is the Riemann tensor, not the contraction $\gamma^\mu \Omega_\mu$.

5 The Lagrangian

The Lagrangian density (SPEC section 3), with an explicit mass term linear in m and quadratic in the spinor:

$$\begin{aligned} \mathcal{L} &= \sqrt{|g|} \left[\frac{1}{2} (\bar{\Psi} \gamma^\mu D_\mu \Psi - (D_\mu \bar{\Psi}) \gamma^\mu \Psi) - m \bar{\Psi} \Psi - U(\bar{\Psi} \Psi) \right], \\ \bar{\Psi} &= \Psi^\dagger C, \quad S = \bar{\Psi} \Psi, \quad U(S) = \frac{\lambda}{2} S^2 \quad (\text{general } U \text{ where stated}). \end{aligned}$$

Properties, proved exactly for an explicit Grassmann algebra (Wolfram: generators θ_1, θ_2 and their conjugates, $\Psi_A = \sum_k \phi_{A,k}(x) \theta_k$; sympy: an own jet super-algebra; `grassmann_algebra_structure`, `superalgebra_axioms`):

- $\mathcal{L}$ is real: $\mathcal{L}^* = \mathcal{L}$, where conjugation reverses the order of Grassmann factors and m, λ are real (`L_real_G`, `grassmann_lagrangian_real`; the control `grassmann_controls_not_vacuuous` shows that the unsymmetrised form is not real off shell).
- It differs from the unsymmetrised form by a total divergence (`L_total_divergence_to_unsymmetrised_G`, `grassmann_total_divergence_relation`):

$$\mathcal{L} = \sqrt{|g|} [\bar{\Psi} \gamma^\mu D_\mu \Psi - mS - U(S)] - \frac{1}{2} \partial_\mu (\sqrt{|g|} \bar{\Psi} \gamma^\mu \Psi).$$

- In this metric the spin connection drops out of $\mathcal{L}$ itself, because $\bar{\Psi} \{\gamma^\mu, \Omega_\mu\} \Psi = 0$ term by term (`L_spin_connection_drops_out_G`):

$$\mathcal{L} = \cos z \left[\frac{1}{2} \sum_{a=1}^8 \frac{1}{f_a} (\bar{\Psi} \gamma^{(a)} \partial_a \Psi - \partial_a \bar{\Psi} \gamma^{(a)} \Psi) - mS - U(S) \right].$$

It reaches the field equations only through this identity: varying the symmetrised kinetic term produces $\partial_\mu (\sqrt{|g|} \gamma^\mu) = 2\sqrt{|g|} \gamma^\mu \Omega_\mu$ (section 7). Hence the Euler-Lagrange equations of $\mathcal{L}$ are those of the connection-free symmetric Lagrangian (the display above), and the term $3H\gamma^{(8)}$ is its half-density (volume and vielbein divergence) term (`connection_free_lagrangian_same_equations`, both engines).

- S is an even element of the Grassmann algebra and U is a polynomial in S ; in the two-generator test algebra $S^3 = 0$, so $U = u_1 S + \frac{\lambda}{2} S^2$ is the most general potential there (`grassmann_algebra_structure`).

Negative control: the notebook's real Majorana-type Lagrangian $L_g = \sqrt{|g|} \Theta^T C \gamma^\mu D_\mu \Theta$ with 16 real anticommuting components is a total derivative, $L_g = \frac{1}{2} \partial_\mu (\sqrt{|g|} \Theta^T C \gamma^\mu \Theta)$: all its Euler-Lagrange expressions vanish identically and it has no field equations; moreover $\Theta^T C \Theta = 0$ (`Majorana_Lg_total_derivative_grassmann`, `negative_control_majorana_grassmann_total_derivative`). For real commuting components the same expression is not a total derivative (`Majorana_Lg_commuting_control`, `negative_control_majorana_commuting_contrast`). This is why the Dirac-type Lagrangian with $\bar{\Psi} = \Psi^\dagger C$ is used.

6 The exact covariant field equations (Euler-Lagrange)

Theorem (Euler-Lagrange equations). For the Lagrangian of section 5 with an arbitrary potential U , the variation with respect to $\Psi^\dagger$ (left derivatives) gives $\sqrt{|g|} C [\gamma^\mu D_\mu \Psi - (m + U'(S))\Psi]$ and the variation with respect to Ψ (right derivatives) gives the adjoint expression, for every component and every Grassmann generator. The field equations are

$$\gamma^\mu D_\mu \Psi = (m + U'(S))\Psi, \quad (D_\mu \bar{\Psi})\gamma^\mu = -(m + U'(S))\bar{\Psi},$$

with $U'(S) = \lambda S$ for the quadratic potential (`EL_Psibar_G`, `EL_Psi_G`, `grassmann_euler_lagrange_psibar_variation` with the control that $m \rightarrow -m$ fails, `grassmann_euler_lagrange_psi_variation`). The adjoint equation is the Dirac conjugate of the field equation, so the two form one system (`adjoint_equation_is_Dirac_conjugate_G`, `grassmann_adjoint_equation_is_conjugate`).

Written out in the primordial field (`Dirac_operator_explicit_G`; formula key `field_equation`):

$$\begin{aligned} & e^{-a_4} \sin^{-1/6} z (\gamma^{(1)} \partial_1 + \gamma^{(2)} \partial_2 + \gamma^{(3)} \partial_3) \Psi + \gamma^{(4)} \partial_4 \Psi \\ & + e^{a_4} \sin^{-1/6} z (\gamma^{(5)} \partial_5 + \gamma^{(6)} \partial_6 + \gamma^{(7)} \partial_7) \Psi \\ & + \tan z \gamma^{(8)} \partial_8 \Psi + 3H \gamma^{(8)} \Psi = (m + U'(S))\Psi, \end{aligned}$$

and the adjoint equation (formula key `adjoint_equation`):

$$\begin{aligned} & e^{-a_4} \sin^{-1/6} z (\partial_1 \bar{\Psi} \gamma^{(1)} + \partial_2 \bar{\Psi} \gamma^{(2)} + \partial_3 \bar{\Psi} \gamma^{(3)}) + \partial_4 \bar{\Psi} \gamma^{(4)} \\ & + e^{a_4} \sin^{-1/6} z (\partial_5 \bar{\Psi} \gamma^{(5)} + \partial_6 \bar{\Psi} \gamma^{(6)} + \partial_7 \bar{\Psi} \gamma^{(7)}) \\ & + \tan z \partial_8 \bar{\Psi} \gamma^{(8)} + 3H \bar{\Psi} \gamma^{(8)} = -(m + U'(S))\bar{\Psi}. \end{aligned}$$

Since $(\gamma^{(4)})^2 = -1$, the field equation is equivalent to the first-order evolution equation (`evolution_form_G`)

$$\partial_4 \Psi = -\gamma^{(4)} \left[(m + U'(S))\Psi - \sum_{a \neq 4} \frac{1}{f_a} \gamma^{(a)} \partial_a \Psi - 3H \gamma^{(8)} \Psi \right],$$

so the slices $x_4 = \text{const}$ are non-characteristic ($g^{44} = -1$). In the chiral split $\Psi = (\psi_-, \psi_+)$ (rows 1 to 8 with $\Gamma = -1$, rows 9 to 16 with $\Gamma = +1$) every gamma is block off-diagonal, $\gamma^{(a)} = \begin{pmatrix} 0 & \bar{\tau}_a \\ \tau_a & 0 \end{pmatrix}$ with $\bar{\tau}_8 = \tau_8 = I_8$, and the field equation is the pair (`block_form`; formula key `field_equation_blocks`)

$$\sum_a \frac{1}{f_a} \bar{\tau}_a \partial_a \psi_+ + 3H \psi_+ = V \psi_-, \quad \sum_a \frac{1}{f_a} \tau_a \partial_a \psi_- + 3H \psi_- = V \psi_+, \quad V = m + U'(S).$$

The 16 component equations, $(\gamma^\mu D_\mu \Psi)_A = V \Psi_A$ with $\psi_A = \Psi_A$ the components in the basis of the fixture gammas, $\epsilon_s = e^{-a_4} \sin^{-1/6} z = 1/f_1$ and $\epsilon_t = e^{a_4} \sin^{-1/6} z = 1/f_5$ (formula key `field_equation_components`; `Dirac_operator_explicit_G`, `EL_Psibar_G`; the publication test of this document re-derives them from the formula file and compares them with the independent sympy expressions `dirac_equation_components` of the sympy report): the eight equations of the $\Gamma = -1$ rows,

$$\begin{aligned} \epsilon_s(-\partial_1 \psi_{16} + \partial_2 \psi_{15} - \partial_3 \psi_{14}) - \partial_4 \psi_{14} + \epsilon_t(\partial_5 \psi_{15} + \partial_6 \psi_{16} + \partial_7 \psi_9) + (\tan z \partial_8 + 3H) \psi_9 &= V \psi_1 \\ \epsilon_s(-\partial_1 \psi_{15} - \partial_2 \psi_{16} + \partial_3 \psi_{13}) + \partial_4 \psi_{13} + \epsilon_t(\partial_5 \psi_{16} - \partial_6 \psi_{15} + \partial_7 \psi_{10}) + (\tan z \partial_8 + 3H) \psi_{10} &= V \psi_2 \\ \epsilon_s(\partial_1 \psi_{14} - \partial_2 \psi_{13} - \partial_3 \psi_{16}) + \partial_4 \psi_{16} + \epsilon_t(-\partial_5 \psi_{13} + \partial_6 \psi_{14} + \partial_7 \psi_{11}) + (\tan z \partial_8 + 3H) \psi_{11} &= V \psi_3 \\ \epsilon_s(\partial_1 \psi_{13} + \partial_2 \psi_{14} + \partial_3 \psi_{15}) - \partial_4 \psi_{15} + \epsilon_t(-\partial_5 \psi_{14} - \partial_6 \psi_{13} + \partial_7 \psi_{12}) + (\tan z \partial_8 + 3H) \psi_{12} &= V \psi_4 \\ \epsilon_s(-\partial_1 \psi_{12} + \partial_2 \psi_{11} - \partial_3 \psi_{10}) + \partial_4 \psi_{10} + \epsilon_t(-\partial_5 \psi_{11} - \partial_6 \psi_{12} - \partial_7 \psi_{13}) + (\tan z \partial_8 + 3H) \psi_{13} &= V \psi_5 \\ \epsilon_s(-\partial_1 \psi_{11} - \partial_2 \psi_{12} + \partial_3 \psi_9) - \partial_4 \psi_9 + \epsilon_t(-\partial_5 \psi_{12} + \partial_6 \psi_{11} - \partial_7 \psi_{14}) + (\tan z \partial_8 + 3H) \psi_{14} &= V \psi_6 \\ \epsilon_s(\partial_1 \psi_{10} - \partial_2 \psi_9 - \partial_3 \psi_{12}) - \partial_4 \psi_{12} + \epsilon_t(\partial_5 \psi_9 - \partial_6 \psi_{10} - \partial_7 \psi_{15}) + (\tan z \partial_8 + 3H) \psi_{15} &= V \psi_7 \\ \epsilon_s(\partial_1 \psi_9 + \partial_2 \psi_{10} + \partial_3 \psi_{11}) + \partial_4 \psi_{11} + \epsilon_t(\partial_5 \psi_{10} + \partial_6 \psi_9 - \partial_7 \psi_{16}) + (\tan z \partial_8 + 3H) \psi_{16} &= V \psi_8 \end{aligned}$$

and the eight equations of the $\Gamma = +1$ rows,

$$\begin{aligned} \epsilon_s(\partial_1 \psi_8 - \partial_2 \psi_7 + \partial_3 \psi_6) + \partial_4 \psi_6 + \epsilon_t(-\partial_5 \psi_7 - \partial_6 \psi_8 - \partial_7 \psi_1) + (\tan z \partial_8 + 3H) \psi_1 &= V \psi_9 \\ \epsilon_s(\partial_1 \psi_7 + \partial_2 \psi_8 - \partial_3 \psi_5) - \partial_4 \psi_5 + \epsilon_t(-\partial_5 \psi_8 + \partial_6 \psi_7 - \partial_7 \psi_2) + (\tan z \partial_8 + 3H) \psi_2 &= V \psi_{10} \\ \epsilon_s(-\partial_1 \psi_6 + \partial_2 \psi_5 + \partial_3 \psi_8) - \partial_4 \psi_8 + \epsilon_t(\partial_5 \psi_5 - \partial_6 \psi_6 - \partial_7 \psi_3) + (\tan z \partial_8 + 3H) \psi_3 &= V \psi_{11} \\ \epsilon_s(-\partial_1 \psi_5 - \partial_2 \psi_6 - \partial_3 \psi_7) + \partial_4 \psi_7 + \epsilon_t(\partial_5 \psi_6 + \partial_6 \psi_5 - \partial_7 \psi_4) + (\tan z \partial_8 + 3H) \psi_4 &= V \psi_{12} \\ \epsilon_s(\partial_1 \psi_4 - \partial_2 \psi_3 + \partial_3 \psi_2) - \partial_4 \psi_2 + \epsilon_t(\partial_5 \psi_3 + \partial_6 \psi_4 + \partial_7 \psi_5) + (\tan z \partial_8 + 3H) \psi_5 &= V \psi_{13} \\ \epsilon_s(\partial_1 \psi_3 + \partial_2 \psi_4 - \partial_3 \psi_1) + \partial_4 \psi_1 + \epsilon_t(\partial_5 \psi_4 - \partial_6 \psi_3 + \partial_7 \psi_6) + (\tan z \partial_8 + 3H) \psi_6 &= V \psi_{14} \\ \epsilon_s(-\partial_1 \psi_2 + \partial_2 \psi_1 + \partial_3 \psi_4) + \partial_4 \psi_4 + \epsilon_t(-\partial_5 \psi_1 + \partial_6 \psi_2 + \partial_7 \psi_7) + (\tan z \partial_8 + 3H) \psi_7 &= V \psi_{15} \\ \epsilon_s(-\partial_1 \psi_1 - \partial_2 \psi_2 - \partial_3 \psi_3) - \partial_4 \psi_3 + \epsilon_t(-\partial_5 \psi_2 - \partial_6 \psi_1 + \partial_7 \psi_8) + (\tan z \partial_8 + 3H) \psi_8 &= V \psi_{16} \end{aligned}$$

For the quadratic potential $V = m + \lambda S$ with $S = \Psi^\dagger C \Psi$, an even (bilinear) element; the equations hold in the Grassmann algebra, and after quantisation as operator equations (section 11).

7 Non-triviality [1]

Theorem 1 (non-triviality [1]). Hypotheses: the metric of section 2 with $H > 0$ and $0 < z < \pi/2$; a_4 any twice differentiable function of x_4 (a constant a_4 included); real m and λ (or any real potential U); the canonical spin connection of the diagonal vielbein; Ψ any dirac16complex configuration (Grassmann-valued components) that is not identically zero. Then:

1. in the diagonal vielbein the Euler-Lagrange operator contains the gravitational term $\gamma^\mu D_\mu \Psi - \gamma^\mu \partial_\mu \Psi = \gamma^\mu \Omega_\mu \Psi = 3H\gamma^{(8)}\Psi$, and $3H\gamma^{(8)}\Psi \neq 0$ (this value depends on the frame and on the field variables, see the Remarks);
2. per direction, the time-direction terms of the three inflating and the three deflating directions cancel ($\frac{3}{2}a'_4 - \frac{3}{2}a'_4 = 0$), while the hidden-direction terms of all six directions add ($6 \cdot \frac{H}{2} = 3H$);
3. in addition the vielbein enters every derivative term, $\gamma^\mu \partial_\mu = \epsilon_s \gamma^{(i)} \partial_i + \gamma^{(4)} \partial_4 + \epsilon_t \gamma^{(t)} \partial_t + \tan z \gamma^{(8)} \partial_8$, none of whose factors is identically 1; because of the cancellation in 2, a_4 enters the field equations through these factors $e^{\mp a_4}$;
4. all gravitational terms vanish together only in flat 4+4 space: $\Omega_\mu = 0$ for every μ if and only if $a'_4 = 0$ and $H = 0$, and $H = 0$ lies outside the family (a degenerate limit); no choice of local frame removes Ω_μ , because its curvature is the Riemann tensor and $R^{x_8}_{x_8} = -6H^2 \neq 0$: the metric is never flat for $H > 0$.

Proof. (a) The connection components of section 4 are each a'_4 or H times a factor $\pm e^{\pm a_4} \sin^{1/6} z$ that does not vanish on $0 < z < \pi/2$; since the S^{ab} are linearly independent, the a'_4 -part of Ω_μ vanishes if and only if a_4 is constant and its H -part if and only if $H = 0$ (`Omega_vanishes_iff_a4prime_and_H_vanish`, `nontriviality_Omega_zero_iff_flat`). (b) Contracting with $\gamma^\mu = \gamma^{(\mu)}/f_\mu$ and using $(\gamma^{(i)})^2 = 1$, $(\gamma^{(t)})^2 = -1$ gives the per-direction values of section 4: the $\gamma^{(4)}$ coefficients are $+\frac{1}{2}a'_4$ for each of the three inflating and $-\frac{1}{2}a'_4$ for each of the three deflating directions and cancel, the $\gamma^{(8)}$ coefficients are $\frac{1}{2}H$ for all six and add to $3H$; hence $\gamma^\mu \Omega_\mu = 3H\gamma^{(8)}$, a constant matrix independent of a_4 , x_4 and x_8 (`gammaOmega_x4_terms_cancel`, `gammaOmega_equals_3H_gamma_x8`, `time_terms_cancel_hidden_term_survives`, `gamma_mu_Omega_mu_equals_3H_gamma_x8`). The same value follows from the divergence form $\frac{1}{2\cos z} \partial_8 (\cos z \tan z) \gamma^{(8)} = \frac{6H \cos z}{2\cos z} \gamma^{(8)} = 3H\gamma^{(8)}$ (`gammaOmega_divergence_form`, `divergence_of_sqrtg_gamma`). (c) $(\gamma^{(8)})^2 = I_{16}$ and $\gamma^{(8)}$ is a signed permutation matrix, so each component of $\gamma^{(8)}\Psi$ is $\pm$ a component of Ψ ; therefore $3H\gamma^{(8)}\Psi = 0$ implies $\Psi = 0$ for $H > 0$ (`nontriviality_1_dirac16complex`). (d) The term is part of the Euler-Lagrange equation although the connection drops out of $\mathcal{L}$ in this metric: the variation of the symmetrised kinetic term produces $\partial_\mu(\sqrt{|g|}\gamma^\mu) = 2\sqrt{|g|}\gamma^\mu \Omega_\mu$, and the Euler-Lagrange expressions equal $\sqrt{|g|}C(\gamma^\mu D_\mu \Psi - V\Psi)$ with the full covariant derivative (`L_spin_connection_drops_out_G`, `EL_Psibar_G`). (e) The curvature of Ω_μ is $\frac{1}{2}R_{ab\mu\nu}S^{ab}$ (`spin_curvature_equals_Riemann`) and $R^{x_8}_{x_8} = -6H^2$ (`never_flat_for_H_positive`); if Ω_μ vanished in some frame, the Riemann tensor would vanish. $H = 0$ is a degenerate limit of the metric (`degenerate_at_H_0`). This completes the proof.

Remarks (the exact scope of Theorem 1). In the formal flat limit ($a'_4 = 0$ and $H = 0$ with the factors frozen) all gravitational terms vanish and the equation becomes the flat 4+4 Dirac-type equation, as the test [1] requires ("unless we are in flat 4+4 spacetime"). Items 1 and 2 are statements about the diagonal vielbein and the field variables Ψ ; read without these hypotheses they would overstate [1]:

- The canonical connection drops out of $\mathcal{L}$, so the Euler-Lagrange equations are those of the connection-free symmetric Lagrangian; $3H\gamma^{(8)} = \frac{1}{2\sqrt{|g|}}\partial_\mu(\sqrt{|g|}\gamma^\mu)$ is its half-density term (`connection_free_lagrangian_same_equations`).
- In the frame boosted in the (x_4, x_8) plane with rapidity $6Hx_4 + b_0$ the term $\gamma'^\mu\Omega'_\mu$ is identically zero (section 4; `boosted_frame_gammaOmega_vanishes`); gravity then enters the Euler-Lagrange operator only through the frame vectors $e'_a{}^\mu$.
- In the diagonal frame the rescaling $\Psi = \sin^{-1/2} z \chi$ removes it: $\gamma^\mu D_\mu \Psi = \sin^{-1/2} z \gamma^\mu \partial_\mu \chi$ exactly, so $\gamma^\mu \partial_\mu \chi = (m + U')\chi$; for $U = \frac{\lambda}{2}S^2$ the term turns into the x_8 -dependent coupling $\lambda S[\chi]/\sin z$. The checks are `rescaling_removes_the_connection_term` and `rescaled_equation_quadratic_potential`. The exact $U = 0$ family of section 8 at $\alpha = -\frac{1}{2}$ is this statement ($M = -m\gamma^{(4)}$, no H).
- The deflation of the extra times contributes nothing to $\gamma^\mu\Omega_\mu$, although $R^{x_4}{}_{x_4} = 6(a'_4)^2 \neq 0$: a_4 enters only through the factors $e^{\mp a_4}$ of the derivative terms (`gammaOmega_blind_to_the_deflation`).
- The frame-independent content of [1] is item 4 with item 3: Ω_μ vanishes in no frame (its curvature is the Riemann tensor and $R^{x_8}{}_{x_8} = -6H^2$), the vielbein factors $e^{\mp a_4} \sin^{-1/6} z$ and $\tan z$ enter every derivative term, and the metric is curved for every $H > 0$. In the diagonal frame the connection also enters the energy-momentum tensor, for example the off-diagonal $K^{x_4}{}_{x_1}$ of a homogeneous configuration (`spin_connection_in_the_energy_momentum_tensor`).
- Interpretation (labelled): the hidden-direction operator $\tan z \partial_8 + 3H$ is antisymmetric for the measure $\cos z dx_8$ in the variables Ψ up to a boundary term, $\cos z [p(\tan z \partial_8 + 3H)q + ((\tan z \partial_8 + 3H)p)q] = \partial_8(\sin z pq)$ (`good_sector_hermiticity_curved`; formula key `hidden_direction_hermiticity`); in the variables χ with the measure $dy = \cot z dx_8$ the operator ∂_y has the same property without any H term. This role of $3H$ therefore depends on the variables and the measure, and the boundary term does not vanish at $z = \pi/2$ (section 11).

Non-triviality [2], for the commuting field `dirac16complex00`, has the same algebraic content and the same scope (`nontriviality_2_dirac16complex00`) and is the subject of the companion document `Revision/docs/DIRAC16COMPLEX00_FIELD_THEORY.md`.

8 Self-consistency

The field equations of section 6 are self-consistent in the following exact senses.

- One system. The adjoint equation is the Dirac conjugate of the field equation: with $E = \gamma^\mu D_\mu \Psi - (m + U')\Psi$ and $\bar{E} = (D_\mu \bar{\Psi})\gamma^\mu + (m + U')\bar{\Psi}$, $E^\dagger C = -\bar{E}$ exactly (`adjoint_equation_is_Dirac_conjugate_G`, `grassmann_adjoint_equation_is_conjugate`).
- Evolution form, not a well-posed Cauchy problem. The evolution form of section 6 holds exactly (`evolution_form_G`), and the Heisenberg equation of the quantised field reproduces it: see the Wolfram check `Heisenberg_equation_reproduces_field_equation` and the

sympy check `hamiltonian_form_and_heisenberg_equation`. The slices $x_4 = \text{const}$ are non-characteristic, but the Cauchy problem is not well posed in Hadamard's sense for data that depend on x_5, x_6, x_7 : with frozen coefficients an extra-time momentum $k_5 = K$ gives the growth rate $\sqrt{K^2 - m^2 - k_1^2 - k_2^2 - k_3^2 - k_8^2}$, unbounded as K grows (`extra_time_growths_rates_unbounded`), and in the metric the frame momentum $e^{a_4} \sin^{-1/6} z k_{x_5}$ grows as the extra times deflate. A well-posed evolution is available at most in the good sector (no extra-time dependence) and only with a boundary condition at $z = \pi/2$ (section 11).

- Integrability. The gammas are covariantly constant and the spinor curvature is the Riemann tensor (section 4). The square of the Dirac operator obeys the Lichnerowicz identity (`Lichnerowicz_identity_G`, `lichnerowicz_identity_on_fields`, `lichnerowicz_contraction`):

$$(\gamma^\mu D_\mu)^2 \Psi = g^{\mu\nu} (D_\mu D_\nu \Psi - \Gamma^\lambda_{\mu\nu} D_\lambda \Psi) - \frac{R}{4} \Psi, \quad R = 6((a'_4)^2 - 7H^2),$$

so the second-order consequence of the field equations is consistent with the curvature.

- Conserved current. $J^\mu = -i\bar{\Psi}\gamma^\mu\Psi$ is real, and off shell $\partial_\mu(\sqrt{|g|}J^\mu) = -i\sqrt{|g|}(\bar{E}\Psi + \bar{\Psi}E)$, so $\nabla_\mu J^\mu = 0$ on shell (`current_real_G`, `current_conservation_identity_G`, `grassmann_n_current_conservation`). The charge density is $J^{x_4} = \Psi^\dagger B \Psi$, and the charge of a slice of constant x_4 , $Q = \int \cos z \Psi^\dagger B \Psi d^7x$, is an indefinite form (`charge_density_is_Krein_form_G`). Only the local law $\nabla_\mu J^\mu = 0$ is established: Q is constant in x_4 only if no charge flows through the boundary of the slice, and at the brane $z = \pi/2$ the flow is not zero in general (`good_sector_hermiticity_up_to_the_brane_flux`, both engines; the lead's independent check `u1_noether_matrix_identity` of `Revision/lead_checks/reports/charge-conjugation-and-u1.json` verifies the local law for general $a_4(x_4)$ and states that the constancy of Q is not established). This record imposes no boundary condition at $z = \pi/2$ (sections 11 and 15).
- Bianchi-type identity of the energy-momentum tensor. For the tensor $T^\mu{}_\nu$ of section 9 the invariance of the action under coordinate changes gives, off shell and for each λ , $\partial_\mu(\sqrt{|g|}T^\mu{}_\lambda) - \sqrt{|g|}T^\mu{}_\alpha \partial_\lambda e^a{}_\mu = \sum_\phi \text{EL}_\phi \partial_\lambda \phi$ (`Noether_identity_diffeomorphisms_G`), and the invariance under the 28 local frame rotations makes the antisymmetric part of T a combination of Euler-Lagrange expressions (`Noether_identity_local_Lorentz_G`). On shell, therefore, T is symmetric and $\nabla_\mu T^\mu{}_\nu = 0$ for general solutions (`conservation_on_shell_general`); the sympy side verifies $\nabla_\mu T^\mu{}_\nu = 0$ for all eight ν after the on-shell substitution, with a negative control (`grassmann_empt_conservation_on_shell`, `grassmann_empt_conservation_negative_control`). The gravitational side satisfies the matching identities $\nabla_\mu E_{(k)}{}^\mu{}_\nu = 0$ (section 13).
- Exact solutions as test points. For $U = 0$ and every a_4 , $\Psi = \sin^\alpha z \left(\cosh(kx_4) + \frac{\sinh(kx_4)}{k} M \right) \chi$ with constant χ , $M = -m\gamma^{(4)} + 3H(2\alpha + 1)\gamma^{(4)}\gamma^{(8)}$, $k^2 = 9H^2(2\alpha + 1)^2 - m^2$, solves the field equation, and conservation of T and J holds exactly there (`solution_matrix_square`, `exact_solution_x4_x8_G`, `exact_solution_family_x4_x8`); the scope of this test is stated in the report: at these solutions the pressures and the momentum density vanish and $\rho = mS$ is constant, so the general proof is the pair of Noether identities. An exact nonlinear homogeneous solution with $U = \frac{\lambda}{2}S^2$ exists in the explicit Grassmann algebra,

$\Psi = \sum_k [e^{M_m x_4} + \lambda S_0 \partial_m e^{M_m x_4}] \chi_k \theta_k$ with $M_m = -m\gamma^{(4)} + 3H\gamma^{(4)}\gamma^{(8)}$ (`exact_solution_nonlinearmhomogeneous_G`; formula key `exact_solutions`).

9 The energy-momentum tensor

The energy-momentum tensor is defined by the vielbein variation of the action and has the closed form (`T_vielbein_variation_closed_form_G`; formula key `T_variation`; sympy `grassmann_empt_equals_general_vielbein_variation_on_shell` with all 64 components of a general first-order vielbein variation)

$$T^\nu{}_\mu = e^b{}_\mu \frac{1}{\sqrt{|g|}} \frac{\delta S}{\delta e^b{}_\nu} = \delta^\nu{}_\mu \mathcal{L}_0 - \frac{1}{2} (\bar{\Psi} \gamma^\nu D_\mu \Psi - D_\mu \bar{\Psi} \gamma^\nu \Psi) \\ - \frac{1}{4} g_{\mu\rho} \nabla_\lambda (\bar{\Psi} \{\gamma^\lambda, \Sigma^{\nu\rho}\} \Psi), \quad \Sigma^{\nu\rho} = \frac{1}{4} [\gamma^\nu, \gamma^\rho], \quad \mathcal{L}_0 = \mathcal{L}/\sqrt{|g|},$$

where the last term (the totally antisymmetric spin density) comes from the variation of the spin connection. Its symmetric part is the Belinfante tensor (`T_symmetric_part_Belinfante_G`, `grassmann_empt_symmetric`; formula key `T_symmetric`),

$$T^{\nu\rho} = g^{\nu\rho} \mathcal{L}_0 - \frac{1}{4} (\bar{\Psi} \gamma^\nu D^\rho \Psi + \bar{\Psi} \gamma^\rho D^\nu \Psi - D^\rho \bar{\Psi} \gamma^\nu \Psi - D^\nu \bar{\Psi} \gamma^\rho \Psi),$$

which equals the variation tensor on shell (section 8). Sign convention (SPEC section 4): the energy density is $\rho = -T^{x_4}{}_{x_4}$ and the pressures are $p_\mu = T^\mu{}_\mu$ (no sum) for $\mu \neq x_4$; with this convention $\rho = mS + U$ for homogeneous on-shell configurations (section 10).

Kinetic and potential parts (formula key `EMT_kinetic_potential`): $T^\nu{}_\mu = T_{\text{kin}}{}^\nu{}_\mu + T_{\text{pot}}{}^\nu{}_\mu$ with

$$T_{\text{pot}}{}^\nu{}_\mu = -\delta^\nu{}_\mu (mS + U(S)), \\ T_{\text{kin}}{}^\nu{}_\mu = \delta^\nu{}_\mu \sum_\lambda K_\lambda - \frac{1}{4} (\bar{\Psi} \gamma^\nu D_\mu \Psi - D_\mu \bar{\Psi} \gamma^\nu \Psi + \bar{\Psi} \gamma_\mu D^\nu \Psi - D^\nu \bar{\Psi} \gamma_\mu \Psi).$$

Diagonal components in the primordial field (no sum; `T_diagonal_components_G`, `grassmann_empt_equals_vielbein_variation_diagonal`; formula key `EMT_diagonal`): with the directional kinetic terms

$$K_\mu = \frac{1}{2f_\mu} (\bar{\Psi} \gamma^{(\mu)} \partial_\mu \Psi - \partial_\mu \bar{\Psi} \gamma^{(\mu)} \Psi), \quad \mathcal{L}_0 = \sum_\mu K_\mu - mS - U(S),$$

$$T^\mu{}_\mu = \mathcal{L}_0 - K_\mu.$$

No spin-connection term survives in the diagonal components, because $\{\gamma^\mu, \Omega_\mu\} = 0$ for each μ . The off-diagonal component x_4 - x_8 , which SPEC section 5 lists among the components of the field equations for a_4 , is (`T_x4_x8_component_G`; formula key `EMT_offdiagonal_x4_x8`)

$$T^{x_4}{}_{x_8} = -\frac{1}{4} (B_{48} - \cot z B_{84}), \quad T^{x_8}{}_{x_4} = -\tan^2 z T^{x_4}{}_{x_8},$$

with $B_{48} = \bar{\Psi}\gamma^{(4)}\partial_8\Psi - \partial_8\bar{\Psi}\gamma^{(4)}\Psi$ and $B_{84} = \bar{\Psi}\gamma^{(8)}\partial_4\Psi - \partial_4\bar{\Psi}\gamma^{(8)}\Psi$. For homogeneous on-shell configurations both $T^{x_4}_{x_8}$ and $T^{x_8}_{x_4}$ vanish identically (`grassmann_T_x4x8_homogeneous`), so these configurations do not source the x_4 - x_8 component of the a_4 equations; this is not claimed for the other off-diagonal components (42 of the remaining 54 are nonzero bilinears in general), whose vanishing is a separate condition on the state (section 13). The trace (off shell, an exact identity in both engines) and the on-shell kinetic sum are (`EMT_trace_G`, `grassmann_trace_on_shell`, `kinetic_sum_on_shell_G`)

$$T^\mu{}_\mu = 7 \sum_\mu K_\mu - 8(mS + U), \quad \sum_\mu K_\mu = (m + U'(S))S + \frac{1}{2}(\bar{\Psi}E - \bar{E}\Psi),$$

so on shell $\mathcal{L}_0 = SU' - U$ and $T^\mu{}_\mu = -mS + 7SU' - 8U = -mS + 3\lambda S^2$. For a diagonal tensor $T = \text{diag}(p_3, p_3, p_3, -\rho, p_t, p_t, p_t, p_8)$ whose entries are functions of x_4 and x_8 , the covariant divergence is, in all eight components (`energy_exchange_equation` in both field-theory reports; formula key `energy_exchange`),

$$\nabla_\mu T^\mu{}_{x_4} = -\partial_4\rho - 3a'_4(p_3 - p_t), \quad \nabla_\mu T^\mu{}_{x_8} = \partial_8 p_8 + 3H \cot z (2p_8 - p_3 - p_t),$$

and $\nabla_\mu T^\mu{}_\nu = 0$ identically for $\nu = x_1, x_2, x_3, x_5, x_6, x_7$. Conservation is therefore exactly

$$\partial_4\rho = -3a'_4(p_3 - p_t), \quad \partial_8 p_8 = -3H \cot z (2p_8 - p_3 - p_t) :$$

energy flows between 3-space and the extra times unless $p_3 = p_t$ (the volume $\sqrt{|g|} = \cos z$ does not depend on x_4); the x_4 component contains no x_8 derivative and no H , the x_8 component no x_4 derivative and no a'_4 . For entries independent of x_8 the two conditions become $d\rho/dx_4 = -3a'_4(p_3 - p_t)$ and $p_8 = (p_3 + p_t)/2$.

For `dirac16complex` these expressions are even elements of the Grassmann algebra; they are the classical counterpart of the energy-momentum tensor operator of section 12, from which all numbers come.

10 Kinetic energy, potential energy, energy density, pressures and equations of state

From the diagonal components of section 9 (formula keys `EMT_diagonal`, `EMT_kinetic_potential`, `equation_of_state_definitions`):

$$\begin{aligned} \rho &= -T^{x_4}_{x_4} = \rho_{\text{kin}} + \rho_{\text{pot}}, & \rho_{\text{kin}} &= - \sum_{\mu \neq x_4} K_\mu, & \rho_{\text{pot}} &= mS + U(S), \\ p_3 &= T^{x_1}_{x_1} = \sum_{\mu \neq x_1} K_\mu - mS - U(S), \\ p_t &= T^{x_5}_{x_5} = \sum_{\mu \neq x_5} K_\mu - mS - U(S), \\ p_8 &= T^{x_8}_{x_8} = \sum_{\mu \neq x_8} K_\mu - mS - U(S), \end{aligned}$$

with $p_3 = T^{x_1}_{x_1} = T^{x_2}_{x_2} = T^{x_3}_{x_3}$ for configurations isotropic in 3-space and $p_t = T^{x_5}_{x_5} = T^{x_6}_{x_6} = T^{x_7}_{x_7}$ for configurations isotropic in the extra times. The kinetic energy density is ρ_{kin} , the potential energy density $\rho_{\text{pot}} = mS + U$; the kinetic pressure in direction μ is $\sum_{\nu \neq \mu} K_\nu$ and the potential pressure is $-(mS + U)$ in every direction. The equations of state are

$$w_3 = \frac{p_3}{\rho}, \quad w_t = \frac{p_t}{\rho}, \quad w_8 = \frac{p_8}{\rho}.$$

Homogeneous on-shell configurations (depending on x_4 only): $K_\mu = 0$ for $\mu \neq x_4$ and $K_{x_4} = (m + U')S$, hence (grassmann_homogeneous_on_shell_rho_p, exact_solution_nonlinear_homogeneous_G; formula key EMT_homogeneous_on_shell)

$$\rho = mS + U, \quad p_3 = p_t = p_8 = SU' - U, \quad \rho_{\text{kin}} = 0, \quad p_{\text{kin}} = (m + U')S,$$

$$w_3 = w_t = w_8 = \frac{SU' - U}{mS + U},$$

and for $U = \frac{\lambda}{2}S^2$

$$\rho = mS + \frac{\lambda}{2}S^2, \quad p = \frac{\lambda}{2}S^2, \quad w_3 = w_t = w_8 = \frac{\lambda S}{2m + \lambda S}.$$

For the exact homogeneous solutions $S = S_0$ is constant along x_4 (exact_solution_nonlinear_homogeneous_G), so ρ , p and w are constant in x_4 , in agreement with the energy-exchange equation ($p_3 = p_t$ gives $d\rho/dx_4 = 0$). For dirac16complex all these quantities are Grassmann-algebra elements until they are evaluated as expectation values of the normal-ordered operators of section 12. The equation of state seen by a 3-space observer and its time dependence (the dark-sector hypotheses of SPEC section 8) are not computed in this document; they are investigated in Revision/docs/DARK_SECTOR_HYPOTHESES, whose verdict for dirac16complex is summarised in section 15.

11 Canonical quantisation in 4 + 4 dimensions and the Krein structure

The evolution time is x_4 . The momentum conjugate to Ψ_A (right derivative) is $\pi_A = \frac{i}{2} \cos z (\Psi^\dagger B)_A$ for the symmetric Lagrangian and $i \cos z (\Psi^\dagger B)_A$ for the first-order form; it depends on $\Psi^\dagger$ only, a second-class constraint as for Dirac's field (canonical_momentum in both reports). Up to a total divergence $\mathcal{L} = \Psi^\dagger K \partial_4 \Psi -$ (terms without ∂_4) with $K = \cos z C \gamma^{(4)} = i \cos z B$, and the canonical (Dirac-bracket) anticommutator on a slice $x_4 = \text{const}$ is (first_order_form_and_anticommutator, canonical_anticommutator_B)

$$\{\Psi_A(x), \Psi_C^\dagger(y)\}_{x_4=y_4} = B_{AC} \frac{\delta^7(x-y)}{\sqrt{|g|}} = B_{AC} \frac{\delta^7(x-y)}{\cos z}.$$

With the Hamiltonian density $\mathcal{H} = \cos z [\Psi^\dagger (mC - \sum_{\mu \neq x_4} C \gamma^\mu D_\mu) \Psi + U(S)]$ (the unsymmetrised $\mathcal{L} = \Psi^\dagger K \partial_4 \Psi - \mathcal{H}$ exactly), the Heisenberg equation $\partial_4 \Psi = i[H, \Psi]$ equals the field equation

solved for $\partial_4 \Psi$, with the operator ordering of the classical expression (`Heisenberg_equation_reproduces_field_equation`, `hamiltonian_form_and_heisenberg_equation`).

Theorem 2 (the canonical anticommutator forces a Krein structure). Hypotheses: the canonical anticommutator above, and $\Psi^\dagger$ realised as the adjoint of Ψ in a space with a positive-definite inner product. These are incompatible: there is no positive inner product with $\Psi^\dagger$ as the adjoint. *Proof.* B has the eigenvalue -1 with multiplicity 8; for the exact unit vector u with components $u_7 = -i/\sqrt{2}$, $u_{16} = 1/\sqrt{2}$ and all others zero, $Bu = -u$, and the smeared operator $X = \sum_A u_A^* \Psi_A(f)$ has $\{X, X^\dagger\} = (u^\dagger Bu) \|f\|^2 = -\|f\|^2 < 0$, whereas in a space with positive inner product $\{X, X^\dagger\} = XX^\dagger + X^\dagger X$ is a positive operator (`no_positive_inner_product` in both reports). Hence the state space carries an indefinite (Krein) inner product with fundamental symmetry B ($B = B^\dagger$, $B^2 = 1$, signature (8,8)); equivalently, in a positive Hilbert space the canonical conjugate $\Psi^\dagger$ is not the Hilbert adjoint χ but $\Psi^\dagger = \chi B$, with $\{\Psi_A, \chi_C\} = \delta_{AC}$.

The one-particle Krein structure. The curved mode operator h ($i\partial_4 \Psi = h\Psi$ for $U = 0$) is formally self-adjoint for the Krein form $\int \cos z u^\dagger B v d^7 x$, that is up to boundary terms (`Krein_form_conserved_curved`). The boundary term of the hidden direction does not vanish at the patch end: $\cos z [u^\dagger (hv) - (hu)^\dagger v] = \partial_8 (\sin z u^\dagger M_8 v)$ with $M_8 = i\gamma^{(4)}\gamma^{(8)}$ in the good sector at $k = 0$, which vanishes at the tip but not at $z = \pi/2$, a surface at finite proper distance where $\sin z = 1$ (`good_sector_hermiticity_up_to_the_brane_flux`, both engines). The Krein form is therefore conserved, and h self-adjoint, only with a boundary condition at $z = \pi/2$ (for example the ASSUMED brane condition of the Kohn-Sham record); this record imposes none. For frozen coefficients (frame momenta k_a), plane waves $u e^{i(k \cdot x - E x_4)}$ obey $Eu = h_k u$ with $h_k = m\beta + \sum_a k_a \alpha^a$, $\beta = -i\gamma^{(4)}$, $\alpha^a = -\gamma^{(4)}\gamma^{(a)}$; β and $\alpha^{1,2,3,8}$ are Hermitian, $\alpha^{5,6,7}$ anti-Hermitian, $Bh_k = h_k^\dagger B$ (Krein self-adjoint), and (`mode_hamiltonian_good_sector`, `mode_hamiltonian_Krein_selfadjoint`, `mode_hamiltonian_B_selfadjoint_dispersion`)

$$h_k^2 = (m^2 + k_1^2 + k_2^2 + k_3^2 + k_8^2 - k_5^2 - k_6^2 - k_7^2) I_{16}.$$

Good sector (no dependence on the extra times, $k_5 = k_6 = k_7 = 0$): h_k is Hermitian with eigenvalues $\pm E$, eight each, and commutes with B ; B also commutes with the curved term $3iH\gamma^{(4)}\gamma^{(8)}$ (`good_sector_spectrum_and_B_sectors`). The curved mode operator is symmetric for the measure $\cos z d^7 x$ in this sector only up to the boundary term at $z = \pi/2$ (`good_sector_hermiticity_curved`, `good_sector_hermiticity_up_to_the_brane_flux`): on the good-sector functions that do not depend on x_8 (finite norm, $\int_0^{\pi/(12H)} \cos(6Hx_8) dx_8 = 1/(6H)$) it acts as the matrix $-im\gamma^{(4)} + 3iH\gamma^{(4)}\gamma^{(8)}$, which is not Hermitian and has the square $(m^2 - 9H^2)I_{16}$; at $m = H = 1$ its eigenvalues are $\pm 2\sqrt{2}i$ (`good_sector_x8_independent_modes_without_boundary_condition`). Without a boundary condition at $z = \pi/2$ the good sector of the author's metric therefore contains growing modes, for $U = 0$ whenever $m^2 < 9H^2$; the exact solutions of section 8 with $\alpha = 0$ ($k^2 = 9H^2 - m^2$) are such modes. Positive Fock realisation: $\Psi = \sum_s (b_s u_s + d_s^* v_s)$ with orthonormal eigenvectors u_s (energy $+E$) and v_s (energy $-E$), $\{b, b^*\} = \{d, d^*\} = 1$ on a positive Fock space, and $\Psi^\dagger = \chi B$, so that $\{\Psi_A, \Psi_C^\dagger\} = B_{AC}$. Exact examples (one momentum each): Wolfram, $m = 3$, $k = (4, 0, 0, 0)$ along x_1 , $E = 5$, on the explicit fermionic Fock space of 2^{16} occupation states; sympy, $m = 2$, $(k_1, k_2, k_3, k_8) = (1, 2, 0, 4)$, $E = 5$. The Hamiltonian $\chi h_k \Psi$ has the vacuum value $-8E$ (the filled sea; -40 in the sympy example), and after normal ordering the energy is $+E$ for the particle $b_0^*|0\rangle$ and $+E$ for the antiparticle $d_0^*|0\rangle$; the charge $\Psi^\dagger B \Psi = \chi \Psi$ is $+1$ and -1 (`Fock_space_good_sector_example`, `good_sector_positive_fock_realisation`).

Expectation-value rule. In the positive realisation, $\langle u | : \Psi^\dagger M \Psi : | u \rangle = u^\dagger B M u$ for a particle and $-v^\dagger B M v$ for an antiparticle (verified for $M = C$, $-iC\gamma^{(4)}$, $-iC\gamma^{(1)}$, $C\gamma^{(2)}\gamma^{(3)}$, a dense integer matrix and a generic symbolic matrix). In the Krein-Fock realisation ($\Psi^\dagger$ the Krein adjoint, modes with $u_n^\dagger B u_n = \epsilon_n$) the normalised expectation of $: \Psi^\dagger M \Psi :$ is $\epsilon_n u_n^\dagger M u_n$, which equals $u_n^\dagger B M u_n$ when $B u_n = \epsilon_n u_n$ and not for a Krein-boosted mode (Fock_space_good_sector_example, good_sector_positive_fock_realisation, expectation_value_rule).

Extra-time modes. With extra-time momentum, $E^2 = m^2 + k_s^2 - k_t^2$ is negative when $k_t^2 > m^2 + k_s^2$; exact example $m = 1$, $k_5 = 2$: $E = \pm i\sqrt{3}$, modes growing like $e^{\sqrt{3}x_4}$. The growth rate has no upper bound as the extra-time momentum grows (extra_time_growth_rates_unbounded). In the metric the frame momentum $k_{(5)} = e^{a_4} \sin^{-1/6} z k_{x_5}$ grows as the extra times deflate (a_4 increasing), so every extra-time mode eventually enters the growing regime (a local-frame statement); extra-time momenta mix the $B = \pm 1$ sectors (extra_time_modes_grow, mode_hamiltonian_good_sector, good_sector_spectrum_and_B_sectors). Krein structure of the one-particle eigenspaces in flat 4+4 space (pairing record): every eigenspace of REAL frequency, $E^2 > 0$, with or without extra-time momentum, has Krein inertia (4,4) (proved for every real frequency: Q_one_particle_Krein_inertia_real_frequencies, Q_one_particle_Krein_inertia_proof), while the eigenspaces of imaginary or zero frequency, those of the growing modes, are Krein-neutral, B vanishes on them identically (Q_one_particle_complex_and_zero_frequencies_Krein_neutral, Q_one_particle_complex_frequency_Krein_neutral; for $m = 1$, $k_5 = 2$ both 8-dimensional eigenspaces of $\pm i\sqrt{3}$).

Scope: the positive Fock realisation is constructed and checked for single good-sector momenta with frozen coefficients; together with the formal (up to the boundary term at $z = \pi/2$) Hermiticity of the curved good-sector mode operator and the formal conservation of the Krein form this is the quantisation recorded here. A complete construction for the interacting field ($\lambda \neq 0$), for the extra-time sector with its growing modes, and of the field algebra on a whole slice is not part of this record (section 15).

12 The energy-momentum tensor operator

The energy-momentum tensor operator of dirac16complex is the tensor of section 9 with the field replaced by the quantised field and normal ordered with respect to the good-sector modes (formula key quantisation):

$$\hat{T}^\nu{}_\mu(x) = : T^\nu{}_\mu[\hat{\Psi}, \hat{\Psi}] :, \quad \hat{\hat{\Psi}} = \hat{\Psi}^\dagger C, \quad \hat{\Psi}^\dagger = \hat{\chi} B \quad (\text{positive realisation}).$$

Explicitly, with $\hat{S} = \hat{\Psi}^\dagger C \hat{\Psi}$ and $\hat{K}_\mu = \frac{1}{2f_\mu} (\hat{\hat{\Psi}} \gamma^{(\mu)} \partial_\mu \hat{\Psi} - \partial_\mu \hat{\hat{\Psi}} \gamma^{(\mu)} \hat{\Psi})$:

$$\begin{aligned} \hat{\rho} &= -\hat{T}^{x_4}{}_{x_4} = : \left(- \sum_{\mu \neq x_4} \hat{K}_\mu + m \hat{S} + U(\hat{S}) \right) :, \\ \hat{p}_\mu &= \hat{T}^\mu{}_\mu = : \left(\sum_{\nu \neq \mu} \hat{K}_\nu - m \hat{S} - U(\hat{S}) \right) : \quad (\mu \neq x_4), \\ \hat{T}^{\nu\rho} &= : \left(g^{\nu\rho} \hat{\mathcal{L}}_0 - \frac{1}{4} (\hat{\hat{\Psi}} \gamma^\nu D^\rho \hat{\Psi} + \hat{\hat{\Psi}} \gamma^\rho D^\nu \hat{\Psi} - D^\rho \hat{\hat{\Psi}} \gamma^\nu \hat{\Psi} - D^\nu \hat{\hat{\Psi}} \gamma^\rho \hat{\Psi}) \right) :, \end{aligned}$$

with the kinetic part $:(-\sum_{\mu \neq x_4} \hat{K}_\mu):$ and the potential part $:(m\hat{S} + U(\hat{S})):$ of the energy density. Properties and status:

- Normal ordering removes the vacuum value of the filled sea ($-8E$ per good-sector momentum in the exact examples of section 11) and gives positive energy $+E$ to particles and antiparticles (`Fock_space_good_sector_example`, `good_sector_positive_fock_realisation`).
- For $\lambda = 0$ every component of $\hat{T}$ is a normal-ordered bilinear $\hat{\Psi}^\dagger M \hat{\Psi}$ (with M containing the derivatives, which act on the mode functions), and its one-particle expectation values follow the expectation-value rule of section 11; in the sympy formulation the one-particle expectation of $\hat{T}$ is ϵ_n times the classical bilinear evaluated on the mode u_n (formula key `quantisation` of the sympy report).
- For $\lambda \neq 0$ the potential term is quartic. The field-equations record states the rule used there: every bilinear is replaced by the expectation value of its normal-ordered operator in the state, with $\langle :U(S): \rangle \neq U(\langle :S: \rangle)$ in general, and it records the operator form of the on-shell identity $\sum_\mu \langle :K_\mu: \rangle = \langle :(m + U')S: \rangle$ as an assumption to be confirmed by the theory branch (key `dirac16complex` of `Revision/field_equations_a4/a4-equations.json`). The identity is verified exactly for the Grassmann-algebra field (`kinetic_sum_on_shell_G`).
- The operator form for $\lambda \neq 0$ in a finite model (record `Revision/theory/fock_quartic/`, 2026-10-08; one Python checker with sympy and its own exact engine for the canonical anticommutation relations, no Wolfram counterpart; 21 of 21 checks pass). The model is the positive fermionic Fock space ($\Psi^\dagger = \chi B$) of one good-sector plane-wave mode set with frozen coefficients: flat frame, volume 1, 16 modes, 2^{16} states, at rest ($k = 0$, symbolic m) and at $m = 3$, $k = (4, 0, 0, 0)$; at $\lambda = 0$ it reproduces the Fock example of section 11 (`control_record_Fock_example`). With the Wick-ordered potential $U = \frac{\lambda}{2} :SS:$ in the Hamiltonian and K_μ defined as Wick products with the Heisenberg derivative of the field, $:\sum_\mu K_\mu: = :(m + U'(S))S:$ holds as an exact operator identity on every state of the model, for the solution of the interacting operator field equation (`trace_identity_operator_identity_wick`), and at $k = 0$ the homogeneous values $\hat{\rho} = :mS + U:$ and $\hat{p} = :SU' - U:$ are operator identities in the same sense (`homogeneous_rho_p_operator_identities_wick`). Of the tested choices (four orderings of the potential, Wick product or vacuum subtraction on the left, two right-hand sides) only this one works: every other combination (30 of them on the two mode sets) fails by a nonzero one-body operator (sometimes plus a constant) linear in λ , and already in expectation values, at $k = 0$ on every state except the vacuum of the mode set (`trace_identity_fails_for_every_other_combination`, `expectation_classes_M0`, `expectation_values_M1`); so the identity holds in this model if and only if the potential and the energy-momentum tensor are both Wick ordered. It also fails for the free dynamics, so it is a statement about solutions of the interacting operator field equation (`negative_control_free_dynamics`). On states with n quanta at $k = 0$, $\rho = mn + \frac{\lambda}{2}n(n-1)$ and $p = \frac{\lambda}{2}n(n-1)$, so $\langle :U(S): \rangle \neq U(\langle :S: \rangle)$ (`homogeneous_expectation_values_are_not_U_of_expectation`). Nothing is proved for the field on a whole slice (many momenta), for the curved x_8 dependence with its boundary term at $z = \pi/2$, or for the extra-time sector; the model gives no numerical expectation value for a physical many-fermion state.
- Numerical expectation values of $\hat{T}$ for many-fermion states are computed only in the Kohn-Sham approximation (`Revision/kohn_sham/`, document `Revision/docs/KOHN_SHAM_DEFLATING_FIELD`); this document quotes only quantities derived from them, in section 13

(the source-condition ratios, the relative defects and the runs of the a_4 equations with the Kohn-Sham source) and in section 15 (the equation-of-state values w_{eff} of the Kohn-Sham gas from the dark-sector record).

- The Grassmann-algebra tensor of sections 9 and 10 obeys the identities of section 8 (symmetry and conservation on shell); the operator inherits them for the bilinear ($\lambda = 0$) part, where they are linear in the field equations; for the quartic part they are not verified as operator statements (the finite model above decides the trace identity and the homogeneous ρ and p , not symmetry or conservation).

13 The field equations for a4[x4] with dirac16complex as the source

Einstein-Lovelock gravity (SPEC section 5), with the Lovelock tensors computed with the generalised Kronecker delta in `Revision/gkd_lovelock` and recomputed in both verifiers of the field-equations branch (for example `P3_direct_equals_gkd_branch_monomials`, `P1_direct_equals_minus_4_Einstein` and `P4_vanishes_pigeonhole`; `sympy P3_equals_gkd_branch_monomials`):

$$\sum_{k=1}^3 \alpha_k E_{(k)}{}^\mu{}_\nu + \Lambda \delta^\mu_\nu = \kappa T^\mu{}_\nu, \quad E_{(k)} = -\frac{P_{(k)}}{2^{k+1}}, \quad E_{(1)} = G,$$

with $\alpha_1 = 1$ for Einstein gravity and the source $T^\mu{}_\nu = \text{diag}(p_3, p_3, p_3, -\rho, p_t, p_t, p_t, p_8) + q_{48} + q_{84}$ (q_{48} at $\mu = x_4, \nu = x_8$; q_{84} at $\mu = x_8, \nu = x_4$). The left-hand sides are divergence-free (`E1_divergence_free`, `E2_divergence_free`, `E3_divergence_free` in both reports). Their nonzero components are diagonal, with $x_1 = x_2 = x_3$ and $x_5 = x_6 = x_7$, free of x_8 and of a_4 itself; every off-diagonal left-hand side, including x_4 - x_8 , is zero (`independent_components`, `P1_structure`, `P2_structure`, `P3_structure`; `sympy other_components_vanish`). The diagonal components (formula key `lovelockTensors`; `json_lovelock_components`):

$$\begin{aligned} E_{(1)}{}^{x_1}{}_{x_1} &= -3(a'_4)^2 + a''_4 + 15H^2 \\ E_{(1)}{}^{x_4}{}_{x_4} &= 3(a'_4)^2 + 21H^2 \\ E_{(1)}{}^{x_5}{}_{x_5} &= -3(a'_4)^2 - a''_4 + 15H^2 \\ E_{(1)}{}^{x_8}{}_{x_8} &= -3(a'_4)^2 + 15H^2 \\ E_{(2)}{}^{x_1}{}_{x_1} &= 12(a'_4)^4 - 24(a'_4)^2 a''_4 + 168(a'_4)^2 H^2 - 40a''_4 H^2 - 180H^4 \\ E_{(2)}{}^{x_4}{}_{x_4} &= -36(a'_4)^4 - 120(a'_4)^2 H^2 - 420H^4 \\ E_{(2)}{}^{x_5}{}_{x_5} &= 12(a'_4)^4 + 24(a'_4)^2 a''_4 + 168(a'_4)^2 H^2 + 40a''_4 H^2 - 180H^4 \\ E_{(2)}{}^{x_8}{}_{x_8} &= 12(a'_4)^4 + 168(a'_4)^2 H^2 - 180H^4 \\ E_{(3)}{}^{x_1}{}_{x_1} &= -72(a'_4)^6 + 360(a'_4)^4 a''_4 - 648(a'_4)^4 H^2 + 432(a'_4)^2 a''_4 H^2 \\ &\quad - 1944(a'_4)^2 H^4 + 360a''_4 H^4 + 360H^6 \\ E_{(3)}{}^{x_4}{}_{x_4} &= 360(a'_4)^6 + 648(a'_4)^4 H^2 + 1080(a'_4)^2 H^4 + 2520H^6 \\ E_{(3)}{}^{x_5}{}_{x_5} &= -72(a'_4)^6 - 360(a'_4)^4 a''_4 - 648(a'_4)^4 H^2 - 432(a'_4)^2 a''_4 H^2 \\ &\quad - 1944(a'_4)^2 H^4 - 360a''_4 H^4 + 360H^6 \\ E_{(3)}{}^{x_8}{}_{x_8} &= -72(a'_4)^6 - 648(a'_4)^4 H^2 - 1944(a'_4)^2 H^4 + 360H^6 \end{aligned}$$

The independent field equations (formula key `generalSource`):

- constraint (x_4): $\sum_k \alpha_k E_{(k)}^{x_4}{}_{x_4} + \Lambda = -\kappa\rho$;
- 3-space ($x_1 = x_2 = x_3$): $\sum_k \alpha_k E_{(k)}^{x_1}{}_{x_1} + \Lambda = \kappa p_3$;
- extra times ($x_5 = x_6 = x_7$): $\sum_k \alpha_k E_{(k)}^{x_5}{}_{x_5} + \Lambda = \kappa p_t$;
- hidden direction (x_8): $\sum_k \alpha_k E_{(k)}^{x_8}{}_{x_8} + \Lambda = \kappa p_8$;
- off-diagonal: $0 = \kappa T^\mu{}_\nu$ for $\mu \neq \nu$, in particular $q_{48} = q_{84} = 0$.

The evolution equation is the 3-space minus extra-time combination; it contains a_4'' and factorises (`evolution_factorises_a4pp_times_F`, `evolution_factorises`; formula key `evolution_F`):

$$a_4'' F(a_4') = \kappa (p_3 - p_t),$$

$$F(a_4') = 2\alpha_1 + \alpha_2 (-48(a_4')^2 - 80H^2) + \alpha_3 (720(a_4')^4 + 864(a_4')^2 H^2 + 720H^4)$$

and F vanishes identically only for $\alpha_1 = \alpha_2 = \alpha_3 = 0$ (`evolution_F_not_identically_zero` in both reports). The constraint and the x_8 equation are first order (`x8_component_contains_no_a4pp`, `constraint_and_x8_first_order`). The remaining consistency conditions:

- algebraic condition: $\sum_k \alpha_k (E^{x_1}{}_{x_1} + E^{x_5}{}_{x_5} - 2E^{x_8}{}_{x_8}) = 0$ identically, so the source must satisfy $p_3 + p_t = 2p_8$ (`algebraic_identity_x1_plus_x5_minus_2x8`, `algebraic_identity`);
- x_8 -dependence: since the left-hand sides do not depend on x_8 , every component of $T^\mu{}_\nu$ must be independent of x_8 , and $q_{48} = q_{84} = 0$ (key `x8_dependence`);
- constraint propagation (Bianchi identity): $\frac{d}{dx_4} \sum_k \alpha_k E_{(k)}^{x_4}{}_{x_4} = 3a_4' \sum_k \alpha_k (E^{x_1}{}_{x_1} - E^{x_5}{}_{x_5})$ identically (`constraint_propagation_bianchi`, `bianchi_x4`); with the conservation law $\rho' = -3a_4'(p_3 - p_t)$ the evolution equation follows from the constraint wherever $a_4' \neq 0$. The conservation components of a general source are $\nabla_\mu T^\mu{}_{x_4} = -\partial_4 \rho - 3a_4'(p_3 - p_t) + \partial_8 q_{84} - 6H \tan z q_{84}$ and $\nabla_\mu T^\mu{}_{x_8} = \partial_8 p_8 + 3H \cot z (2p_8 - p_3 - p_t) + \partial_4 q_{48}$ (`conservation_components` in both reports).

Einstein gravity ($\alpha_1 = 1$, $\alpha_2 = \alpha_3 = 0$; `einstein_components`; formula key `einstein`):

$$\begin{aligned} 3(a_4')^2 + 21H^2 + \Lambda &= -\kappa\rho \\ -3(a_4')^2 + a_4'' + 15H^2 + \Lambda &= \kappa p_3 \\ -3(a_4')^2 - a_4'' + 15H^2 + \Lambda &= \kappa p_t \\ -3(a_4')^2 + 15H^2 + \Lambda &= \kappa p_8 \end{aligned}$$

so $2a_4'' = \kappa(p_3 - p_t)$ and $\kappa(\rho + p_8) = -6((a_4')^2 + H^2) < 0$ for every a_4 and every Λ : the source must violate the null energy condition along the null direction $x_4 + x_8$ (`einstein_null_energy_x8`), and there is no vacuum solution for $H > 0$ (`einstein_no_vacuum_solution`, `einstein_no_vacu`).

um). Every twice differentiable a_4 is allowed if the source is the one these equations require (key `sourceRequiredByGivenA4`); $\kappa\rho > 0$ needs $\Lambda < -(3(a'_4)^2 + 21H^2)$.

The linear member $a_4 = AHx_4 + a_0$ (3-space scale factor e^{AHx_4} , extra-time scale factor e^{-AHx_4} : exponential deflation for $A > 0$). The field equations then require $p_3 = p_t = p_8 = p$, with ρ and p constant, for any α_k (`linear_member_equal_pressures`; key `linearMember`):

$$\begin{aligned}\kappa\rho &= \alpha_1(-3A^2H^2 - 21H^2) + \alpha_2(36A^4H^4 + 120A^2H^4 + 420H^4) \\ &\quad + \alpha_3(-360A^6H^6 - 648A^4H^6 - 1080A^2H^6 - 2520H^6) - \Lambda \\ \kappa p &= \alpha_1(-3A^2H^2 + 15H^2) + \alpha_2(12A^4H^4 + 168A^2H^4 - 180H^4) \\ &\quad + \alpha_3(-72A^6H^6 - 648A^4H^6 - 1944A^2H^6 + 360H^6) + \Lambda \\ \mathcal{V}(A) &= \alpha_1 + \alpha_2(-8A^2H^2 - 40H^2) + \alpha_3(72A^4H^4 + 144A^2H^4 + 360H^4)\end{aligned}$$

A vacuum ($\rho = p = 0$) needs $\mathcal{V}(A) = 0$ (`linear_member_vacuum_factor`); in Einstein-Gauss-Bonnet gravity ($\alpha_3 = 0$) this gives $A^2 = (1 - 40\alpha_2H^2)/(8\alpha_2H^2)$, real for $0 < \alpha_2H^2 \leq 1/40$ (`einstein_gauss_bonnet_vacuum_linear`). For Einstein gravity $\kappa\rho = -(3A^2 + 21)H^2 - \Lambda$, $\kappa p = (15 - 3A^2)H^2 + \Lambda$ and $\kappa(\rho + p) = -6(A^2 + 1)H^2$.

Specialisation to `dirac16complex`. The source is the energy-momentum tensor of section 9 with the declared sign $\sigma_T = +1$ (key `emtConvention`: $T^\mu{}_\nu = \sigma_T(-K^{(\mu}{}_\nu) + \delta^\mu{}_\nu \mathcal{L}_0)$, the convention $\rho = -T^{x_4}{}_{x_4}$ of SPEC section 4; the relative sign of the gravitational and matter actions is carried as the parameter σ_T in the field-equations record and is not fixed by a Revision check). For the quantised field every bilinear is replaced by the expectation value of its normal-ordered operator in the state (section 12), with $k_\mu = K_\mu$:

$$\begin{aligned}\rho &= \langle :k_4: \rangle - \langle :\mathcal{L}_0: \rangle, & p_3 &= \langle :\mathcal{L}_0: \rangle - \langle :k_1: \rangle, \\ p_t &= \langle :\mathcal{L}_0: \rangle - \langle :k_5: \rangle, & p_8 &= \langle :\mathcal{L}_0: \rangle - \langle :k_8: \rangle,\end{aligned}$$

and the equations become (key `dirac16complex` of the field-equations record):

- constraint: $\sum_k \alpha_k E_{(k)}^{x_4 x_4} + \Lambda = -\kappa(\langle :k_4: \rangle - \langle :\mathcal{L}_0: \rangle)$;
- evolution: $a_4'' F(a_4') = \kappa(\langle :k_5: \rangle - \langle :k_1: \rangle)$: a state with $\langle :k_1: \rangle \neq \langle :k_5: \rangle$ would drive a_4'' if it also satisfied the conditions below ($p_3 + p_t = 2p_8$, every expectation value independent of x_8 , vanishing off-diagonal components); no such state is constructed in Revision;
- hidden direction: $\sum_k \alpha_k E_{(k)}^{x_8 x_8} + \Lambda = \kappa(\langle :\mathcal{L}_0: \rangle - \langle :k_8: \rangle)$, and the algebraic condition $2\langle :k_8: \rangle = \langle :k_1: \rangle + \langle :k_5: \rangle$;
- every off-diagonal expectation value $\langle :T^\mu{}_\nu: \rangle$, $\mu \neq \nu$, must vanish, and every expectation value must be independent of x_8 . For a homogeneous configuration the off-diagonal components are multiples of the 15 three-gamma bilinears $\bar{\Psi}\gamma^{(a)}\gamma^{(b)}\gamma^{(c)}\Psi$ with $\{a, b, c\} = \{i, 4, 8\}$ ($i = 1, 2, 3, 5, 6, 7$) and $\{i, 4, j\}$ ($i = 1, 2, 3, j = 5, 6, 7$), derived for the classical homogeneous condensate (`condensate_offdiagonal_are_three_gamma_bilinears`, `authorT16_condensate_offdiagonal_three_gamma`; key `offDiagonalConditions`); for `dirac16complex` their expectation values must vanish.

Statement (states with equal 3-space and extra-time kinetic terms; key homogeneousSingleMode). Hypotheses: a state whose normal-ordered expectation values depend on x_4 only, satisfy the off-diagonal conditions and have $\langle :k_1: \rangle = \langle :k_5: \rangle$ for every x_4 (for example all quanta at zero 3-momentum and zero extra-time momentum); a_4 twice continuously differentiable; $(\alpha_1, \alpha_2, \alpha_3) \neq (0, 0, 0)$. Then $a_4'' F(a_4') = 0$; F is a nonzero polynomial in a_4' with finitely many roots (**evolution_F_not_identically_zero**) and a_4' is continuous, so $a_4'' = 0$ on every interval and $a_4 = A H x_4 + a_0$ with real A : only the linear member is allowed (the same argument as the theorem for the classical condensate, key **theoremLinear**, with **linear_member_equal_pressures**). The equations contain A only through A^2 , so they are invariant under $A \rightarrow -A$: the extra times deflate for $A > 0$, which is a choice of sign (an initial condition) and is not selected by the equations; $A < 0$ is allowed on the same footing, and $A = 0$ is the static case. If in addition $\langle :k_\mu: \rangle = 0$ for every $\mu \neq x_4$, then $\rho = \langle :mS + U: \rangle$ exactly and $p = \langle :k_4: \rangle - \langle :mS + U: \rangle$, which equals $\langle :SU' - U: \rangle$ under the operator form of the on-shell identity of section 12 (an assumption for $\lambda \neq 0$). With it, Einstein gravity requires $\kappa \langle :mS + U: \rangle = -(3A^2 + 21)H^2 - \Lambda$ and $\kappa \langle :SU' - U: \rangle = (15 - 3A^2)H^2 + \Lambda$, hence $\kappa \langle :m + U' : S: \rangle = -6(A^2 + 1)H^2 < 0$ (key **einsteinConditions**; for the classical quadratic case **condensate_einstein_quadratic_U** in both reports).

What a_4 the equations allow, in summary: for a prescribed a_4 the equations fix the required source (ρ, p_3, p_t, p_8 above, with $p_3 + p_t = 2p_8$ and all components independent of x_8), and whether a state of **dirac16complex** supplies it is a separate question; states with $\langle :k_1: \rangle = \langle :k_5: \rangle$ allow only the linear member $a_4 = A H x_4 + a_0$ with constant ρ and p (the sign of A , and with it deflation or inflation of the extra times, is not selected); a non-linear a_4 needs $p_3 \neq p_t$, that is $\langle :k_1: \rangle \neq \langle :k_5: \rangle$, together with all the other conditions. The Kohn-Sham states computed in **Revision/kohn_sham/** (key **kohnSham** of the field-equations record) are not such a source: every recorded state with a nonzero energy-momentum tensor depends on x_8 and violates $p_3 + p_t = 2p_8$ pointwise and also after integration over x_8 (for $N = 136$, $\lambda = 0$ the ratio $(\int p_3 + \int p_t) / (2 \int p_8)$ is 0.339767, 0.25969 and 0.239714 at $a_{4,0} = 0, 1, 2$, where the equations require 1), so the author's metric is not a solution with them (**ks_profiles_depend_on_x8**, **ks_profiles_violate_algebraic_condition**, **ks_integrals_violate_algebraic_condition**). The Kohn-Sham history $a_4 = A H x_4$ is a prescribed test-field background without back-reaction: the linear member needs $p_3 = p_t = p_8$ and constant ρ , which the Kohn-Sham gas along it does not have (**ks_history_is_a_prescribed_background**).

The field equations for a_4 with the Kohn-Sham source (2026-10-08, Revision/field_equations_a4/ks_source). A separate analysis (Python: 23 of 23 checks pass) makes the last statement exact and asks what the equations would do with the Kohn-Sham source. Exact: for every Lovelock order $E_{(k)}^{x_1 x_1} + E_{(k)}^{x_5 x_5} = 2E_{(k)}^{x_8 x_8}$, so the $x_1 + x_5 - 2x_8$ combination of the field equations reads $0 = \kappa(p_3 + p_t - 2p_8)$ for every a_4 (**A1_lovelock_identity_x1_plus_x5_equals_2x8**); in the Kohn-Sham coordinate y the conservation law along x_8 gives $p_3 + p_t - 2p_8 = \partial_y p_8 / (3H)$ on shell (**A6_x8_conservation_in_y**), and the recorded states obey that conservation law (**B2_x8_conservation_on_every_profile**), so their violation of the algebraic condition is exactly the x_8 dependence of p_8 ; after averaging over the hidden direction the relative defect stays between 0.414086 and 0.806379 over the 70 nonzero states (**C1_averaged_algebraic_condition_fails**). The coupled problem therefore has no solution, and nothing about a_4 is derived from it. In a stated approximation (the x_4 moment and the $x_1 - x_5$ moment of the field equations kept, the x_8 moment and the $x_1 + x_5 - 2x_8$ moment dropped; their residuals are of order one, not small, **D7_dropped_x8_moment_residual_and_adiabaticity**) the Kohn-Sham source does not start or select exponential deflation of the extra times: the initial rate $a_4'(0)$ and its sign are initial data, and the gas changes $(a_4')^2$ only by a bounded amount set by the energy it loses (**D5_einstein_sign_of_the_effect**);

for the series $N = 136$, $\lambda = 0$ in Einstein gravity $a'_4(2)/a'_4(0) = 1.95362$, 1.1321 and 0.847549 for $\sigma_0 = \kappa\bar{\rho}(0)/H^2 = 10$, 1 and -1 . With $\Lambda = 0$ and the initial rate $a'_4(0) = H$ the constraint forces $\sigma_0 < 0$, that is $\kappa < 0$ for this gas of positive energy density ($\bar{\rho} > 0$, `C4_source_signs`): a negative gravitational coupling in the sign convention $\sigma_T = +1$ of the field-equations record. In the four series with 3-momentum that were integrated ($N = 136$ with $\lambda = 0$; $N = 688$ with $\lambda = 0, \pm\lambda_2$; the other six series with 3-momentum were not integrated) the deflation then halts ($a'_4 = 0$) at $a_4 = 0.149623$ to 0.151697 (Einstein), 0.0724197 to 0.0733808 (Einstein-Gauss-Bonnet) and 0.102772 to 0.104159 (cubic Lovelock), where the integration stops (`D3_lambda_zero_cases_halt`); what follows the turning point is not computed; a source that does not change with a_4 (the interacting $N = 8$ states) gives exactly the linear member $a_4 = AHx_4 + a_0$, allowed but not selected (`D4_constant_source_gives_linear_member`).

14 Relation to the pairing theorems

The pairing theorems of SPEC section 9 use the Lagrangian, the field equations and the energy-momentum tensor of this document; their full statements, hypotheses and limits are in `Revision/docs/PAIR_CREATION_PROOFS.md`. For `dirac16complex`, in brief:

- T1 (chirality): $\mathcal{L}_{m,\lambda}[\Gamma\Psi] = -\mathcal{L}_{-m,-\lambda}[\Psi]$, solutions of the (m, λ) equations are mapped to solutions of the $(-m, -\lambda)$ equations, $T \rightarrow -T$ and $J \rightarrow -J$, in every gravitational field; the total energy-momentum and the total current of the pair vanish identically (`Wolfram T1_Lagrangian_primordial_grassmann` and `T1_energy_momentum_primordial_grassmann`, `sympy T1.metric.grassmann.pair_total_empt_zero`).
- T2 (chirality combined with a $\text{Pin}(4,4)$ reflection of character -1 , the mirror $z \rightarrow \pi - z$ across the assumed Z2 brane): $(m, \lambda) \rightarrow (-m, \lambda)$, $\mathcal{L} \rightarrow +\mathcal{L}$, with equal, not opposite, energy-momentum (`T2_mirror_Lagrangian_grassmann`, `T2_mirror_energy_momentum_and_current_grassmann`; `sympy T2.metric.grassmann.empt`).
- Quantum reading: the chirality image $\Gamma\Psi$ carries the Krein metric $-B$; there is no cancellation between two independently quantised universes (`Q_Krein_metric_of_images`, `Q_no_identification_of_independent_universes`; `sympy Q.no_cancellation_independent_universes`).
- T3 (the Kohn-Sham level): the block map of Γ with the brane parities exchanged and the tip angle $\theta \rightarrow \pi - \theta$ maps every self-consistent instantaneous Kohn-Sham state with (m, λ, θ) onto one with $(-m, +\lambda, \pi - \theta)$, with equal levels, occupations, Kohn-Sham energies and energy-momentum profiles, and $S \rightarrow -S$; the Z2 brane is ASSUMED (`T3_energies_and_empt_profiles_equal`, `T3_exact_k0_spectra`; `sympy T3.energies_and_empt_profiles_equal`).
- Completion of T3 (2026-10-08): with the expectation rule of the canonical quantisation (section 11) the image state has $\rho' = -\Gamma\rho\Gamma$, so n is unchanged, S and Q change sign, and every component of the 16-component energy-momentum tensor and of the current of the $(-m, +\lambda)$ image state equals that of the original state (statement S6; `T3C_krein_rule_16_component`, `sympy T3C.krein_rule_16_component`); the filling convention of the $+m$ member is carried onto the $-m$ member (`T3C_filling_convention_mapped`). Two numerical demonstrations,

with the Rust solver (210 states) and with the independent reference solver (18 states), are demonstrations, not proofs.

These are correspondences between solutions; no creation process, rate or amplitude follows from them (`not_established` in the sympy pairing report, in the T3 record and in the completion record).

15 What this document does not claim

- It derives no creation process, rate or amplitude for universes, and no statement that universes of masses $+m$ and $-m$ are created in pairs; the pairing theorems (section 14) map solutions to solutions.
- Non-triviality [1] is a statement about the Euler-Lagrange equations in the author's metric with $H > 0$; it is not a statement about observable effects of the gravitational term. The value $\gamma^\mu \Omega_\mu = 3H\gamma^{(8)}$ belongs to the diagonal vielbein and to the field variables Ψ : it vanishes identically in the frame boosted in the (x_4, x_8) plane with rapidity $6Hx_4 + b_0$, and the rescaling $\Psi = \sin^{-1/2} z \chi$ removes it; the canonical connection drops out of $\mathcal{L}$, and the deflation a_4 makes no contribution to $\gamma^\mu \Omega_\mu$. What is frame-independent is that Ω_μ vanishes in no frame (curvature $R^{x_8}_{x_8} = -6H^2$), together with the vielbein factors of the derivative terms (section 7, Remarks). That the term is "not cosmetic" is not claimed; its role for the Hermiticity of the mode operator depends on the measure and the variables (an interpretation).
- The quantisation recorded here is: the canonical anticommutator, the theorem that it forces a Krein structure when $\Psi^\dagger$ is realised as the Hilbert adjoint, the formal Krein self-adjointness of the evolution (up to the boundary term at $z = \pi/2$), and positive Fock realisations checked for single good-sector momenta with frozen coefficients. A positive-norm Hilbert space for the full field, a stable vacuum in the presence of the growing extra-time modes, and the interacting ($\lambda \neq 0$) operator theory beyond the finite model of section 12 are not constructed. In flat 4+4 space every eigenspace of REAL frequency of the one-particle generator has Krein inertia (4,4) (`Q_one_particle_Krein_signatures`, `Q_one_particle_Krein_inertia`, proved for every real frequency in `Q_one_particle_Krein_inertia_real_frequencies` and `Q_one_particle_Krein_inertia_proof`), and every eigenspace of imaginary or zero frequency is Krein-neutral (`Q_one_particle_complex_and_zero_frequencies_Krein_neutral`). In flat space the good-sector restriction removes the complex frequencies, not the indefiniteness of the real-frequency eigenspaces; in the curved good sector without a boundary condition at $z = \pi/2$ complex frequencies remain (section 11).
- Boundary conditions: none is imposed at $z = \pi/2$ in this record. The Hermiticity of the curved good-sector mode operator and the conservation of the Krein form hold up to the boundary term there; with no boundary condition the good sector contains growing x_8 -independent modes (for $U = 0$ whenever $m^2 < 9H^2$).
- No well-posed Cauchy problem is claimed: the slices $x_4 = \text{const}$ are non-characteristic, but for data that depend on the extra times the growth rates are unbounded (Hadamard ill-posedness).
- The normal-ordered operator form of the on-shell trace identity for $\lambda \neq 0$ is an assumption of the field-equations record. It is verified only in the finite model of section 12 (one good-sector plane-wave mode set with frozen coefficients, record `Revision/theory/fock_quartic/`), where it

holds as an exact operator identity if and only if the potential and the energy-momentum tensor are both Wick ordered and fails for every other tested ordering already in expectation values; for the field on a whole slice, the curved x_8 dependence and the extra-time sector it is not verified, and symmetry and conservation of the quartic operator are not verified. No numerical expectation value of the energy-momentum tensor operator for a physical many-fermion state is computed in this document (the finite model gives only its exact values on the occupation states of one mode set); the Kohn-Sham quantities it quotes, in section 13 (the source-condition ratios, the relative defects and the runs of the a_4 equations with the Kohn-Sham source) and in this section (the equation-of-state values w_{eff} of the Kohn-Sham gas from the dark-sector record, below), are computed from Kohn-Sham expectation values.

- The relative sign σ_T between the gravitational and the matter side of the a_4 equations is a declared convention ($\sigma_T = +1$), not derived.
- No solution $a_4(x_4)$ with a `dirac16complex` source is computed here beyond the statements of section 13; it is not claimed that any state of `dirac16complex` supplies a source that satisfies all the conditions of section 13 (in Einstein gravity such a source must violate the null energy condition along $x_4 + x_8$). The recorded Kohn-Sham states fail the conditions (section 13), and the Kohn-Sham history is a prescribed background without back-reaction; the equations do not select deflation ($A > 0$) over inflation of the extra times.
- The equation of state seen by a 3-space observer and its time dependence are not derived in this document; Hypothesis (`dirac16complex` as a mechanism for a time-varying dark-energy or dark-matter equation of state) is investigated in `Revision/docs/DARK_SECTOR_HYPOTHESES`. Its verdict for `dirac16complex`, under its stated observer assumption (three definitions (A), (B), (C) of the 4-dimensional density; $w_{\text{eff}}(C) = w_{\text{eff}}(A) - 1$ exactly): the computed Kohn-Sham gas along the prescribed history is radiation-like, with w_{eff} rising toward $1/3$ under (A) and (B); the dark-matter-like fall from $1/3$ toward 0 occurs for quanta of the massive bulk band, which the computed ground states do not populate; a condensate is dust-like; dark-energy-like values occur only under (C), where the gas has w_{eff} between -0.707107 and -0.671895 with a thawing-sign slope and mixtures with a condensate are freezing; nothing computed comes near the Unite pair $(w_0, w_a) = (-0.861, -0.60)$, the Unite constant $w = -0.764$ is matched only with tuned parameters, and no computed state crosses $w = -1$. That record establishes neither Hypothesis nor Hypothesis00.
- $H = 0$ is a degenerate limit, not a member of the family; the surface $z = \pi/2$ is degenerate ($g_{88} = 0$, $\sqrt{|g|} = 0$); nothing is claimed there.
- The comparison record of the field-theory branch agrees entirely (section 1); every formula is nevertheless cited by the separate checks of the two engines.

16 Reproduction

From the repository root (each verifier writes its outputs to the paths listed in section 1; the Wolfram verifier of each part runs first because the sympy verifier compares with its output at the end):

```
wolframscript -file Revision/algebra/wolfram/verify_algebra.wls
python Revision/algebra/python/check_algebra.py
```

```

wolframscript -file Revision/theory/wolfram/verify_field_theory.wls
python Revision/theory/python/check_field_theory.py
wolframscript -file Revision/field_equations_a4/wolfram/verify_field_equations_a4.wls
python Revision/field_equations_a4/python/check_field_equations_a4.py
wolframscript -file Revision/pairing/wolfram/verify_pairing.wls
python Revision/pairing/python/check_pairing.py
wolframscript -file Revision/theory/wolfram/verify_scope.wls
python Revision/theory/python/check_scope.py
wolframscript -file Revision/pairing/kohn_sham/wolfram/verify_t3.wls
python Revision/pairing/kohn_sham/python/check_t3.py
python Revision/field_equations_a4/python/check_ks_source_conditions.py
python Revision/field_equations_a4/ks_source/ks_source_a4.py
wolframscript -file Revision/pairing/kohn_sham/wolfram/verify_t3_completion.wls
python Revision/theory/fock_quartic/check_fock_quartic.py

```

The analysis of the a_4 equations with the Kohn-Sham source re-derives the source-conditions report, so it runs after that checker (3 s on an idle development machine, up to about 20 s under load); the Wolfram completion verifier of T3 ran in about 15 s, and the checker of the finite Fock model for $\lambda \neq 0$ (it reads only `Revision/algebra/gammas.json`) in 6.7 s by its README. The sympy completion checker reads the reports of the two numerical demonstrations, so it runs after them, in the order given in section 10 of `Revision/docs/PAIR_CREATION_PROOFS`; the programs of the dark-sector record are listed in section 12 of `Revision/docs/DARK_SECTOR_HYPOTHESES`.

The sympy field-theory checker exits with an error if its comparison with the Wolfram record does not agree entirely. It can write its report to another path, which leaves the recorded report untouched; it ran in about two minutes in this session (the scope checkers in about 17 s and 1 s, the T3 verifiers in about 3 s and 1 s):

```
python Revision/theory/python/check_field_theory.py --out <path>
```

This document (md, tex, pdf) is built and verified with the first command below (one command line); with the optional last argument it records the edition `dirac16complex-field-theory` in `Revision/pdf-specifications.json`. The second command runs the publication test, the third prints the blocks that the test generates:

```

python scripts/build_provenance_pdf.py Revision/docs/DIRAC16COMPLEX_FIELD_THEORY.md
--developer-layout --specifications Revision/pdf-specifications.json [--register]
python -m unittest Revision/tests/test_dirac16complex_field_theory_publication.py -v
python Revision/tests/test_dirac16complex_field_theory_publication.py --emit

```

The test checks that the `.tex` is the builder output of the `.md`, that the registered PDF is the PDF file next to it, that every cited check exists and passes in the reports, that the quoted check counts are those of the reports, and that the component field equations and the a_4 equations above are the output of its renderers applied to `Revision/theory/field-theory.json` and `Revision/field_equations_a4/a4-equations.json`.

17 Index of formulas and checks

formula	section	Wolfram checks	sympy checks
metric, vielbein, volume element	2	metric_is_the_au thors, sqrt_det _g_is_cos_z	metric_from_viel bein_equals_SPEC, sqrt_det_g_equal s_cos_z
Christoffel symbols, Ricci tensor, curvature	2	christoffel_count, ricci_mixed_comp onents, ricci_sc alar, never_fl at_for_H_positive	christoffel_symm etric_metric_com patible, curvatur e_nonzero_flat_o nly_formally
Clifford algebra, C, Gamma, B	3	Clifford_relation, C_gamma_real_ant isymmetric, Gamma_diag, B_signature_8_8	clifford_relation, C_conjugation, chirality_diag, B_hermitian_invo lution_signature
Pin(4,4) irreducible; Spin(4,4): two inequivalent 8-dimensional irreducible halves	3	Pin44_irreducibl e_commutant_dim_1, chiral_halves_ir reducible, chiral_halves_in equivalent_inter twiners_0	pin_commutant_di mension_1, spin_halves_irre ducible, spin_hal ves_inequivalent
canonical spin connection, covariant constancy	4	vielbein_postula te, omega_co mponents, Omega_co mponents, gamma_co variantly_consta nt	vielbein_postula te, Omega_x4 _and_Omega_x8_va nish, covarian t_constancy_D_mu _gamma_nu
gravitational term 3H gamma8 (diagonal frame)	4, 7	gammaOmega_equal s_3H_gamma_x8, gammaOmega_x4_te rms_cancel	gamma_mu_Omega_m u_equals_3H_gamm a_x8, time_ter ms_cancel_hidden _term_survives
frame and variable dependence of the term; frame-independent content	4, 7	boosted_frame_ga mmaOmega_vanishes, boosted_frame_cu rvature_nonzero, rescaling_remove s_the_connection _term, connecti on_free_lagrangi an_same_equations	boosted_frame_ga mmaOmega_formula, rescaled_equatio n_quadratic_pote ntial, gammaOme ga_blind_to_the_ deflation, spin_connection_ in_the_energy_mo mentum_tensor
Lagrangian: reality, total divergence	5	L_real_G, L_total_ divergence_to_un symmetrised_G	grassmann_lagran gian_real, grassmann_total_ divergence_relat ion
Majorana-type negative control	5	Majorana_Lg_tota l_derivative_gra ssmann	negative_control _majorana_grassm ann_total_deriva tive

formula	section	Wolfram checks	sympy checks
Euler-Lagrange equations, explicit form	6	EL_Psibar_G, EL_Psi_G, Dirac_operator_explicit_G, evolution_form_G, block_form	grassmann_euler_lagrange_psibar_variation, grassmann_euler_lagrange_psi_variation
non-triviality [1]	7	nontriviality_1_dirac16complex, Omega_vanishes_iff_a4prime_and_H_vanish, spin_curvature_equals_Riemann	nontriviality_Omega_zero_iff_flat, spinor_curvature_equals_riemann
self-consistency	8	adjoint_equation_is_Dirac_conjugate_G, Lichnerowicz_identity_G, current_conservation_identity_G, Noether_identity_diffeomorphisms_G, Noether_identity_local_Lorentz_G	grassmann_adjoint_equation_is_conjugate, lichnerowicz_identity_on_fields, grassmann_current_conservation, grassmann_nemt_conservation_on_shell
energy-momentum tensor	9	T_vielbein_variation_closed_form_G, T_symmetric_part_Belinfante_G, T_diagonal_components_G, T_x4_x8_components_G, EMT_trace_G	grassmann_nemt_equals_general_vielbein_variation_on_shell, grassmann_nemt_symmetric, grassmann_trace_on_shell
energy density, pressures, equations of state	10	kinetic_sum_on_shell_G, exact_solution_nonlinear_homogeneous_G, energy_exchange_equation	grassmann_homogeneous_on_shell_rho_p, energy_exchange_equation
canonical quantisation, Krein structure	11	canonical_momentum, first_order_form_and_anticommutator, no_positive_inner_product, Krein_form_conservated_curved, good_sector_hermicity_curved, Q_one_particle_Krein_inertia_real_frequencies	canonical_anticommutator_B, no_positive_inner_product, good_sector_spectrum_and_B_sectors, Q.one_particle_Krein_inertia_proof, Q.one_particle_complex_frequency_Krein_neutral

formula	section	Wolfram checks	sympy checks
boundary term at $z = \pi/2$, growing modes, ill-posedness	8, 11	good_sector_herm iticity_up_to_th e_brane_flux, good_sector_x8_i ndependent_modes _without_boundar y_condition	extra_time_growt h_rates_unbounded
Fock space, normal ordering, expectation values	11, 12	Fock_space_good_ sector_example, mode_hamiltonian _Krein_selfadjoi nt	good_sector_posi tive_fock_realis ation, expectat ion_value_rule, extra_time_modes _grow
energy-momentum tensor operator for $\lambda \neq 0$ in a finite Fock model	12		trace_identity_o perator_identity _wick, trace_id entity_fails_for _every_other_com bination, homogene ous_rho_p_operat or_identities_wi ck, negative _control_free_dy namics (Python)
field equations for a4	13	evolution_factor ises_a4pp_times_F, algebraic_identi ty_x1_plus_x5_mi nus_2x8, constrai nt_propagation_b ianchi, einstein _null_energy_x8, linear_member_eq ual_pressures	evolution_factor ises, algebrai c_identity, bianchi_x4, einstein_null_en ergy_x8, linear_m ember_equal_pres sures
the Kohn-Sham states as a source	13		ks_profiles_depe nd_on_x8, ks_integ rals_violate_alg ebraic_condition, ks_history_is_a_ prescribed_backg round (Python)

formula	section	Wolfram checks	sympy checks
the a_4 equations with the Kohn-Sham source	13		A1_lovelock_iden tity_x1_plus_x5_ equals_2x8, A6_x8_conservati on_in_y, B2_x8_co nservation_on_ev ery_profile, C1_averaged_alge braic_condition_ fails, D3_lambd a_zero_cases_halt, D4_constant_sour ce_gives_linear_ member, D5_einst ein_sign_of_the_ effect (Python)
completion of T3 (S6, filling convention)	14	T3C_krein_rule_1 6_component, T3C_filling_conv ention_mapped	T3C.krein_rule_1 6_component, T3C.filling_conv ention_mapped
pairing (section 14)	14	T1_Lagrangian_pr imordial_grassma nn, T2_mirro r_Lagrangian_gra ssmann, Q_Krein_ metric_of_images, T3_energies_and_ emt_profiles_equ al	T1.metric.grassm ann.lagrangian, T2.metric.grassm ann.emt, Q.no_can cellation_indepe ndent_universes, T3.energies_and_ emt_profiles_equ al