UM-4 propagation

The answer to the author's order of 2026-10-25, 13:15 PDT: the metric in which the time is warped like every other direction of the fold, photons and gravitons at the speed of light in the bulk, the photon a transverse wave of spin 1, the gravitational wave a wave of spin 2 with its plus and cross polarisations

Underlying theory: Patrick L. Nash Formulated and verified by Claude (Anthropic)

Propagation in UM-4, 2026-10-25

Repository once-ere/Dirac_claude, folder universe_model_4

Contents

1 The fold: definition and examples **1**

2 How to read this document **12**
2.1 What this document contains . . . 12
2.2 The status labels . . . 12
2.3 Coordinates, signature and units . . . 12

3 The order and the answer **12**

4 Why UM-3 was physically incorrect **13**

5 The metric of UM-4 **15**

6 Gravitons at the speed of light: spin 2 with the plus and cross polarisations **19**
6.1 Which gravity . . . 19
6.2 The gravitational wave . . . 23

7 Photons at the speed of light: transverse waves of spin 1 **26**

8 The other fields of the bulk **30**

9 From one point of the fold to another **34**

10 Causality around an expanding fold **37**

11 What the change of the metric changes elsewhere **38**

12 What is open **38**

13 What was verified **39**

14 Further symbols **40**

1 The fold: definition and examples

Definition. The *fold* is the seven-dimensional surface of the author's eight-dimensional spacetime on which the cosine of the angle $z = 6Hx_0$ of the hidden direction vanishes:

$$\mathcal{F} = \big\{X = (x_0, x_1, \ldots, x_7) :\ \cos(6Hx_0) = 0\big\}, \qquad \text{in one period } 0 < z < \pi \text{ of the metric: } z = \frac{\pi}{2}, \quad x_0 = \frac{\pi}{12H} \tag{FD-1}$$

Symbols in (FD-1).

- $\mathcal{F}$: The fold: the seven-dimensional surface $\cos(6Hx_0) = 0$ of the eight-dimensional spacetime (FD-1), in one period $0 < z < \pi$ the surface $z = \pi/2$, $x_0 = \pi/(12H)$, where $\sigma = 0$; $|_{\mathcal{F}}$ means restricted to the fold. Status: DERIVED.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$, the fold is $z = \pi/2$. Unit: dimensionless. Status: AUTHOR.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant of UM-4, $\Lambda = -21H^2$: the value for which the static block is a vacuum of Einstein gravity (its Einstein tensor is $21H^2$ in every direction). The author writes the pure number Λ_{N} in $0 = -\Lambda_{\mathrm{N}} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$, so that $\Lambda = -\Lambda_{\mathrm{N}} H^2$. Unit: GeV^2. Value: $-21H^2$; the author's original $-10H^2$; UM-3: $-7H^2$. Status: AUTHOR (the term), DERIVED (the value).
- Λ_{N}: The author's pure number in the vacuum equations of his notebook, $\Lambda = -\Lambda_{\mathrm{N}} H^2$: his original value 10; 21 in UM-4 (Einstein gravity), 7 in UM-3. Unit: dimensionless. Value: 21 (the author's original 10). Status: AUTHOR.
- $w_{(p)}$: The Lovelock weights of the author's Einstein-Lovelock action, $w_{(1)} = 1$ and $w_{(2)}$, $w_{(3)}$ of the Gauss-Bonnet and the cubic densities. UM-4 sets $w_{(2)} = w_{(3)} = 0$ (Einstein gravity): the only weights for which gravitons propagate on the light cone on every background, not only around the static block. The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) keep the gravitons on the light cone only around the static block (with $\Lambda = -161H^2/16$); UM-3 used $(1, 1/20, 1/360)$. Unit: dimensionless. Value: $(1, 0, 0)$ in UM-4; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), DERIVED (the values that the order of 2026-10-25 selects).

$X = \{x_0, \ldots, x_7\}$ are the author's coordinates: x_0 the hidden space-like direction, x_1, x_2, x_3 ordinary space, x_4 the time, x_5, x_6, x_7 the three extra times; H is the author's constant (an inverse length). The metric of UM-4 depends on x_0 only through $\cot^2 z$ and $(\sin^2 z)^{1/6}$, so it repeats with the period $\pi/(6H)$ of x_0, and every period $0 < z < \pi$ contains exactly one surface of $\mathcal{F}$, at $z = \pi/2$: $\mathcal{F}$ is the union of these copies, one per period (example 2), and "the fold" without a qualifier is the copy in the period that contains our world. A point of the fold is labelled by the seven coordinates $x_1, \ldots, x_7$, and the metric restricted to the fold ($\mathrm{d}x_0 = 0$, $z = \pi/2$) is

$$g_{MN}\,\mathrm{d}X^M \mathrm{d}X^N\big|_{\mathcal{F}} = \mathrm{e}^{2a_4}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{d}x_4^2 - \mathrm{e}^{-2a_4} b_4^2 \big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big) \tag{FD-2}$$

Symbols in (FD-2).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M \mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- $\mathcal{F}$: The fold: the seven-dimensional surface $\cos(6Hx_0) = 0$ of the eight-dimensional spacetime (FD-1), in one period $0 < z < \pi$ the surface $z = \pi/2$, $x_0 = \pi/(12H)$, where $\sigma = 0$; $|_{\mathcal{F}}$ means restricted to the fold. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.

with three space-like and four time-like directions; a_4 and b_4 are the author's two functions of the time x_4. Notation: ln is the natural logarithm, e Euler's number, $\exp(y) = \mathrm{e}^y$, sin, cos and $\cot = \cos / \sin$ the trigonometric functions, sinh and tanh the hyperbolic functions, $|y|$ the absolute value; $f(0^+)$ and $f(0^-)$ are the limits of $f(\sigma)$ as σ approaches 0 from above and from below.

Why this surface is called a fold. Measured by the proper distance along the hidden direction,

$$\sigma = \mathrm{sgn}\Big(\frac{\pi}{2} - z\Big)\frac{\ln\sin z}{6H}, \qquad \mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0, \qquad \sin z = \mathrm{e}^{-6H|\sigma|} \tag{FD-3}$$

Symbols in (FD-3).

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$, the fold is $z = \pi/2$. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.

(the fold is $\sigma = 0$; the half-period $0 < z < \pi/2$ is $\sigma < 0$, the half-period $\pi/2 < z < \pi$ is $\sigma > 0$; the two ends of the period, the tips $z \to 0$ and $z \to \pi$, lie at $\sigma \to -\infty$ and $\sigma \to +\infty$), the metric of UM-4 is

$$g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N = \mathrm{d}\sigma^2 + \mathrm{e}^{-2H|\sigma|}\Big[\mathrm{e}^{2a_4}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{d}x_4^2 - \mathrm{e}^{-2a_4}b_4^2\big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big)\Big] \qquad \text{(FD-4)}$$

Symbols in (FD-4).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.

It depends on σ only through $|\sigma|$. The function $\mathcal{W}(\sigma) = \mathrm{e}^{-2H|\sigma|}$, which multiplies the squared steps of all seven directions of the fold (ordinary space, the time and the extra times), is the warp factor of the metric (its square root $\mathrm{e}^{-H|\sigma|}$ multiplies the lengths). The time x_4 is warped like the other directions (the author's order of 2026-10-25, 13:15 PDT; in UM-3 it was not), so that every surface $\sigma = \text{const}$ has the light cone of the fold. Four properties make the surface $\sigma = 0$ a fold, and together they single it out within the period:

1. *It is a mirror.* The reflection $\sigma \to -\sigma$ (that is $z \to \pi - z$, or $x_0 \to \pi/(6H) - x_0$) maps the metric to itself, and the fold is the only surface of the period that the reflection leaves in place: the two halves of the period are mirror images of each other, joined along the fold, as the two halves of a folded sheet of paper are joined along the crease.
2. *The warp factor is largest on it.* The warp factor $(\sin^2 z)^{1/6} = \mathrm{e}^{-2H|\sigma|}$ equals 1 on the fold and decreases on both sides.

3. *The metric has a kink on it.* The metric is continuous across the fold, but its first derivative along σ jumps. The extrinsic curvature of the surfaces $\sigma = \text{const}$ changes sign across the fold,

$$K^a{}_b = \frac{1}{2}\, g^{ac}\, \partial_\sigma g_{cb} = -H\, \mathrm{sgn}(\sigma)\, \delta^a{}_b \quad \text{(all seven directions of the fold)} \tag{FD-5}$$

Symbols in (FD-5).

- $K^a{}_b$: The extrinsic curvature of a surface of constant σ of the hidden direction, $K^a{}_b = \frac{1}{2} g^{ac} \partial_\sigma g_{cb}$ for a, b along the surface (FD-5): $-H\,\mathrm{sgn}(\sigma)$ along all seven directions of the fold for the author's form of the metric of UM-4; $K = K^c{}_c$ its trace. Unit: GeV (1/length). Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- c: A frame index (as a). Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+, +, +, +, -, -, -, -$ in the order $x_0, \ldots, x_7$): $g_{MN}\, \mathrm{d}X^M \mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\, \mathrm{d}^8 X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\, \mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z) \ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- η_{ab}: The flat metric of the frame, $\eta = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$ (eta4488 of 4+4.nb); η^{ab} is its inverse (the same matrix). Unit: dimensionless. Status: AUTHOR.
- $\hat{g}_{ab}$: The metric of the seven-dimensional fold (FD-2) in its coordinates $x_1, \ldots, x_7$ (or in its frame), $\hat{g}^{ab}$ its inverse; $\hat{h}_{\mu\nu}$ is its part along the four observable directions. Unit: dimensionless. Status: AUTHOR.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- x: The coordinates of the fold: x_1, x_2, x_3 of ordinary space, x_4 the time, x_5, x_6, x_7 the extra times; a field of the fold is a function of the four observable coordinates $x = (x_1, x_2, x_3, x_4)$, and $\mathrm{d}^4 x = \mathrm{d}x_1\, \mathrm{d}x_2\, \mathrm{d}x_3\, \mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.

(a, b frame indices along the fold); it jumps by $-2H$ in every direction of the fold. The field equations of gravity of UM-4, $\mathcal{E}_{(1)}{}^M{}_N + \Lambda \delta^M_N = \kappa \mathcal{T}^M{}_N$ (Einstein gravity: the author's Einstein-Lovelock equations with the weights of the Gauss-Bonnet and cubic terms set to zero; $\mathcal{E}_{(1)}$ the Einstein tensor, $\Lambda = -21H^2$ the cosmological constant, $\kappa = 8\pi G_8$ the gravitational coupling of eight dimensions, $\mathcal{T}^M{}_N$ the energy-momentum tensor, δ^M_N the Kronecker delta), are linear in the second σ-derivatives of the metric, so the jump puts a Dirac delta function $\delta_\mathrm{D}(\sigma)$ into them: the metric solves them only if the fold carries a thin source $\mathcal{S}^a{}_b$, an energy-momentum tensor concentrated on $\sigma = 0$, given by the integral of the equations across the fold (the junction condition of Israel),

$$\kappa\, \mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma\, \mathcal{E}_{(1)}{}^a{}_b = -\Big(\big(K^a{}_b\big|_{0^+} - K^a{}_b\big|_{0^-}\big) - \delta^a{}_b \big(K\big|_{0^+} - K\big|_{0^-}\big) \Big) \tag{FD-11}$$

Symbols in (FD-11).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- ${\mathcal{S}^a}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa {\mathcal{S}^a}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \, {\mathcal{E}_{(1)}}^a{}_b$ (the junction condition of Israel); ${\mathcal{S}^4}_4$ is minus its energy density, ${\mathcal{S}^1}_1$ and ${\mathcal{S}^5}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa {\mathcal{S}^a}_b = -12H{\delta^a}_b$ for every a_4, b_4 of the author's form. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z| \, \mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z) \ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- ${\mathcal{E}_{(p)}}^M{}_N$: The Lovelock tensor of order p, ${\mathcal{E}_{(p)}}^M{}_N = -2^{-(p+1)} \, \delta^{MP_1\cdots P_{2p}}_{NQ_1\cdots Q_{2p}} \, {R^{Q_1Q_2}}_{P_1P_2} \cdots {R^{Q_{2p-1}Q_{2p}}}_{P_{2p-1}P_{2p}}$ (with both indices down it is $1/\sqrt{|g|}$ times the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to g^{MN}; $\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor, $\mathcal{E}_{(3)}$ the cubic Lovelock tensor); with frame indices its frame components. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number (UM-4 keeps only $p = 1$); in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- ${K^a}_b$: The extrinsic curvature of a surface of constant σ of the hidden direction, ${K^a}_b = \frac{1}{2} g^{ac} \partial_\sigma g_{cb}$ for a, b along the surface (FD-5): $-H \, \mathrm{sgn}(\sigma)$ along all seven directions of the fold for the author's form of the metric of UM-4; $K = {K^c}_c$ its trace. Unit: GeV (1/length). Status: DERIVED.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1M_2\cdots}_{N_1N_2\cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- G_8: The eight-dimensional Newton constant ($\kappa = 8\pi G_8$). Unit: GeV^{-6}. Status: AUTHOR.
- ${R^{MN}}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor ${R^M}_{NPQ}$ (in the convention in which the commutator of two covariant derivatives, acting on a vector, is the Riemann tensor, with its last two indices those of the two derivatives, contracted with the vector), with two indices the Ricci tensor $R_{MN} = {R^P}_{MPN}$, without indices the scalar curvature $R = g^{MN} R_{MN}$; with frame indices ${R^{(a)(b)}}_{(c)(d)}$ its frame components, ${R^{(a)(b)}}_{(a)(b)}$ (no sum) the sectional curvature of the frame plane (a, b). Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- $\mathcal{L}_{(p)}$: The Lovelock density of order p, $\mathcal{L}_{(p)} = 2^{-p} \, \delta^{P_1\cdots P_{2p}}_{Q_1\cdots Q_{2p}} \, {R^{Q_1Q_2}}_{P_1P_2} \cdots {R^{Q_{2p-1}Q_{2p}}}_{P_{2p-1}P_{2p}}$, a scalar built from p curvature tensors: $\mathcal{L}_{(1)} = R$ (Einstein-Hilbert), $\mathcal{L}_{(2)}$ the Gauss-Bonnet density, $\mathcal{L}_{(3)}$ the cubic Lovelock density. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.

($K = {K^c}_c$ the trace of the extrinsic curvature), with ${\mathcal{S}^4}_4 = -$(its energy density), ${\mathcal{S}^1}_1 = {\mathcal{S}^2}_2 = {\mathcal{S}^3}_3$ its pressure along ordinary space and ${\mathcal{S}^5}_5 = {\mathcal{S}^6}_6 = {\mathcal{S}^7}_7$ its pressure along the extra times; $\mathcal{S}$ is $\mathcal{T}$ integrated across the fold, an energy per unit volume of the six other directions of the fold (unit GeV^7). For the static block (a_4 and b_4 constant) and, to the first order in the curvature of the fold divided by H^2, for every fold

$$\begin{aligned} &\text{static block:} \quad \kappa {\mathcal{S}^a}_b = -12H \, {\delta^a}_b \quad \text{(a pure tension)}, \qquad \kappa\lambda_{\mathrm{F}} = 12H, \\ &\text{any fold:} \quad \kappa {\mathcal{S}^a}_b = -12H \, {\delta^a}_b + \frac{2}{5H} \, {\hat{G}^a}_b + \ldots \end{aligned} \tag{FD-12}$$

Symbols in (FD-12).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.

- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma\, \mathcal{E}_{(1)}{}^a{}_b$ (the junction condition of Israel); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -12H\delta^a{}_b$ for every a_4, b_4 of the author's form. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- λ_F: The tension that the static block requires of the fold, $\kappa\lambda_\mathrm{F} = 12H$: the total vacuum energy of the seven-dimensional fold, made of the tension of the fold fluid and the vacuum energy of the fields of the fold (both have the pressure minus their energy density along ordinary space and along the extra times), per unit volume of the six other directions of the fold (eight-dimensional normalisation); in the four-dimensional normalisation it is $\lambda_\mathrm{F} V_*$, and the excess of the total over $\lambda_\mathrm{F} V_*$ is the dark energy ρ_Λ. Unit: GeV^7. Value: $12H/\kappa$; $\lambda_\mathrm{F} V_* = 30H^2 M_\mathrm{red}^2 = 1.1 \times 10^{45}$ GeV^4 for $H = 2.5$ TeV. Status: DERIVED.
- $\hat{G}^\mu{}_\nu$: The Einstein tensor of the metric $\hat{g}_{ab}$ of the fold (FD-2), with frame indices along its seven directions; with four-dimensional indices, $\hat{G}^\mu{}_\nu$, its components along the four observable directions, which for frozen extra times are those of the Einstein tensor of the four-dimensional metric $\hat{h}_{\mu\nu}$. Unit: 1/length^2. Status: STANDARD.
- M_red: The reduced Planck mass, $M_\mathrm{red} = (8\pi G_\mathrm{N})^{-1/2}$. Unit: GeV. Value: 2.435×10^{18} GeV. Status: STANDARD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4} b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\, \mathrm{d}x_6\, \mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4} b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- $\hat{h}_{\mu\nu}$: The four-dimensional metric of the fold, $\hat{h}_{\mu\nu} = \mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the coordinates $x_1, \ldots, x_4$ (the induced metric at $\sigma = 0$ without the extra times); $\hat{h}^{\mu\nu}$ its inverse, $\hat{h}$ its determinant, $\sqrt{-\hat{h}} = \mathrm{e}^{3a_4}$; $\hat{h}^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- ρ_Λ: The vacuum energy density of the fold above its static tension, the dark energy of UM-4: $\rho_\Lambda = \rho_\mathrm{ff} + \rho_\mathrm{vac} - \lambda_\mathrm{F} V_*$ (four-dimensional normalisation). Unit: GeV^4. Value: $0.685\, \rho_\mathrm{c}^{(0)} = 2.5 \times 10^{-47}$ GeV^4 (Planck 2018). Status: DERIVED.
- G_N: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_\mathrm{N} = 1/M_\mathrm{Pl}^2$ with the Planck mass $M_\mathrm{Pl} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- V_t: The coordinate volume of the (compact) extra times, $V_\mathrm{t} = \int \mathrm{d}x_5\, \mathrm{d}x_6\, \mathrm{d}x_7$. Unit: length^3. Status: ASSUMED.
- ρ_c: The critical density, the energy density of a spatially flat fold at the expansion rate H_4; $\rho_\mathrm{c}^{(0)}$ its value today. Unit: GeV^4. Value: $\rho_\mathrm{c}^{(0)} = 1.0537 \times 10^{-5}\, h_{100}^2$ GeV/cm^3. Status: STANDARD.
- ρ_ff: The energy density of the fold fluid. Unit: GeV^4. Value: $\lambda_\mathrm{F} V_* + \rho_\Lambda - \rho_\mathrm{vac}$: together with the vacuum energy ρ_vac of the fields of the fold it makes up the tension $\lambda_\mathrm{F} V_*$ that the static block requires plus the dark energy ρ_Λ. Status: ASSUMED (the fluid), DERIVED (the value).
- ρ_vac: The vacuum energy of the fields of the fold: the minimum of their potential (for the Standard Model the minimum of the Higgs potential with its loop corrections), a tension of the seven-dimensional fold. Unit: GeV^4. Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\, h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.

- M_{Pl}: The Planck mass, $M_{\mathrm{Pl}} = G_{\mathrm{N}}^{-1/2}$. Unit: GeV. Value: 1.221×10^{19} GeV. Status: STANDARD.
- h_{100}: The present expansion rate of the fold in units of 100 km/s/Mpc. Unit: dimensionless. Value: 0.674 (Planck 2018). Status: STANDARD.

($\hat{G}^a{}_b$ the Einstein tensor of the metric FD-2 of the fold): the tension plus matter that obeys the Einstein equations of seven dimensions, $\hat{G}^a{}_b = \kappa_7 \mathtt{t}^a{}_b$ with $\kappa_7 = 5\kappa H/2$ ($\mathtt{t}^a{}_b$ the matter of the fold), which for frozen extra times are the Friedmann equations of general relativity. The fold is therefore not only a place but a physical object: a thin layer that carries energy and momentum (a brane). 4. *The author's coordinate x_0 degenerates on it.* The metric component $g_{x_0x_0} = \cot^2(6Hx_0)$ vanishes on the fold and nowhere else in the period; near the fold the proper distance grows with the square of the coordinate distance,

$$|\sigma| = 3H\Big(x_0 - \frac{\pi}{12H}\Big)^2 \Big[1 + 6H^2\Big(x_0 - \frac{\pi}{12H}\Big)^2 + \dots\Big] \tag{FD-6}$$

Symbols in (FD-6).

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

so the two coordinate values on either side of $x_0 = \pi/(12H)$ at the same coordinate distance lie at the same proper distance from the fold: the coordinate line of x_0 is folded onto the line of proper distances (a fold map, example 5).

What lives on the fold. In UM-4 the Higgs doublet (with the author's ad hoc vacuum, a time-dependent choice of its vacuum value), three generations of quarks and leptons, three right-handed neutrinos and the fold fluid (its tension and the pressure along the extra times that keeps their volume fixed) live on the fold: they are fields at $\sigma = 0$, functions of $x_1, \dots, x_4$ (no field depends on the extra times). The gauge fields of $SU(3)_C \times SU(2)_L \times U(1)_Y$ live in the bulk; their massless modes are bound to the fold, so that the photon and the gluons of the four-dimensional world are waves of the bulk that the fold holds, and photons that leave the fold travel through the bulk at the speed of light. dirac16complex (the author's spinor field of eight dimensions) and gravity live in the bulk $\sigma \neq 0$ as well as on the fold; the states χ of dirac16complex that the fold holds (in UM-4 long-lived resonances) are the part of it near the fold, and the portals (couplings between dirac16complex and the fields of the fold) act at $\sigma = 0$. Integrated over the extra times, the fold is the four-dimensional universe that we observe, with the metric $\mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in $x_1, \dots, x_4$. In UM-4 files "the fold" means this surface; the examples below use the word for other surfaces with the same properties. The word is not used in its other senses: a set of equal energy levels is called degenerate, and a growth by the factor e is called an e-folding.

Examples of folds that are meaningful for UM-4.

1. *Our fold.* The fold $z = \pi/2$ of the period that contains our world (FD-1): the seven-dimensional world-volume of the quarks, leptons and the Higgs field, and after the integration over the extra times the four-dimensional universe that we observe.
2. *The copies of the fold.* Because the metric repeats with the period $\pi/(6H)$, every surface $x_0 = \pi/(12H) +$ (an integer multiple of $\pi/(6H)$) is a copy of the fold, one per period. Neighbouring copies are separated by a tip (z a multiple of π), where the warp factor $(\sin^2 z)^{1/6}$ vanishes and $g_{x_0x_0} = \cot^2 z$ diverges, and which lies at an infinite proper distance from both: in proper distance every copy has its own complete line $-\infty < \sigma < \infty$. A compact direction of length L would give every field a tower of particles with the masses $2\pi n/L$, $n = 1, 2, \dots$ (a Kaluza-Klein tower); because the copies are infinitely far apart, the period $\pi/(6H)$ of x_0 produces no such tower with the spacing $12H$: every field of the bulk has instead a continuum of modes that starts at zero mass, and the fold holds its massless mode.
3. *A folded sheet of paper.* The crease of a folded sheet is a fold in this sense: the two halves are mirror images (property 1) joined along the crease, the sheet is continuous there and its direction changes abruptly (property

3). In UM-4 the direction across the crease is the hidden direction σ, the seven directions along the crease are $x_1, \ldots, x_7$, and the crease is the fold.

4. *A folded function.* The warp factor $\mathrm{e}^{-2H|\sigma|}$ is the simplest folded function: continuous, mirror symmetric, largest at $\sigma = 0$, with a jump of its derivative,

$$\frac{\mathrm{d}}{\mathrm{d}\sigma}\mathrm{e}^{-2H|\sigma|} = -2H\,\mathrm{sgn}(\sigma)\,\mathrm{e}^{-2H|\sigma|}, \qquad \frac{\mathrm{d}^2}{\mathrm{d}\sigma^2}\mathrm{e}^{-2H|\sigma|} = 4H^2\,\mathrm{e}^{-2H|\sigma|} - 4H\,\delta_{\mathrm{D}}(\sigma) \tag{FD-7}$$

Symbols in (FD-7).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- δ_{D}: The Dirac delta function of σ: zero for $\sigma \neq 0$, with the integral 1 over any range that contains $\sigma = 0$; the integral of $\delta_{\mathrm{D}}(\sigma)$ times a smooth function of σ is the value of that function at $\sigma = 0$. Unit: 1/length (for $\delta_{\mathrm{D}}(\sigma)$). Status: STANDARD.

whose delta function is the origin of the thin source of property 3. Every metric function of FD-4 depends on σ through $|\sigma|$ and is folded at $\sigma = 0$ in this sense. 5. *A fold map.* In the mathematics of smooth maps the fold (Whitney's fold singularity) is the map $u \mapsto u^2$ of a line onto a half-line: it folds the line at $u = 0$, two points $\pm u$ land on the same point, and the derivative of the map vanishes on the fold. Near the fold the map from the author's coordinate x_0 to the distance $|\sigma|$ from the fold is of this kind (FD-6), and $g_{x_0x_0} = \cot^2 z$, the square of the derivative $\mathrm{d}\sigma/\mathrm{d}x_0$, vanishes there. 6. *The brane of Randall and Sundrum.* The five-dimensional metric

$$\mathrm{d}y_{\mathrm{RS}}^2 + \mathrm{e}^{-2k_{\mathrm{RS}}|y_{\mathrm{RS}}|}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2 - \mathrm{d}x_4^2\big), \qquad \lambda_{\mathrm{RS}} = \frac{6k_{\mathrm{RS}}}{\kappa_5}, \qquad \Lambda_{\mathrm{RS}} = -\frac{6k_{\mathrm{RS}}^2}{\kappa_5} \tag{FD-8}$$

Symbols in (FD-8).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- y_{RS}: The proper coordinate across the brane of the five-dimensional model of Randall and Sundrum (FD-8, an example, not UM-4). Unit: length. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- k_{RS}: The inverse curvature length of the bulk of the model of Randall and Sundrum: the warp factor of the brane is $\mathrm{e}^{-2k_{\mathrm{RS}}|y_{\mathrm{RS}}|}$ (FD-8, an example, not UM-4). Unit: GeV (1/length). Status: STANDARD.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- λ_{RS}: The tension (energy per unit volume of ordinary space) of the brane of the model of Randall and Sundrum, $\lambda_{\mathrm{RS}} = 6k_{\mathrm{RS}}/\kappa_5$ (FD-8, an example, not UM-4). Unit: GeV^4. Status: STANDARD.
- κ_5: The gravitational coupling of five dimensions in the model of Randall and Sundrum, 8π times the Newton constant of five dimensions (FD-8, an example, not UM-4). Unit: GeV^{-3}. Status: STANDARD.
- Λ_{RS}: The vacuum energy density of the bulk of the model of Randall and Sundrum, $\Lambda_{\mathrm{RS}} = -6k_{\mathrm{RS}}^2/\kappa_5$ (FD-8, an example, not UM-4). Unit: GeV^5. Status: STANDARD.

has a fold at $y_{\mathrm{RS}} = 0$ with the properties 1 to 3 (its coordinate y_{RS} is already the proper distance, so property 4 does not arise); Einstein's equations require there a brane of the tension λ_{RS} (Israel's junction condition) and in the bulk the vacuum energy Λ_{RS}, and the bulk is otherwise empty. The fold of UM-4 is of the same kind, with every direction of the fold warped by the same factor as in the model of Randall and Sundrum: its static block is a space of constant curvature $-H^2$ in all eight directions, a vacuum of Einstein gravity with $\Lambda = -21H^2$, with a fold of pure tension (FD-12), and the graviton is bound to the fold as there. It differs in that the fold has seven dimensions, three of them extra times, in that the space is not cut at the brane (Randall and Sundrum identify the two sides of their brane, one physical side; the fold of UM-4 has two distinct sides, example 7), and in that the gauge fields live in the bulk as well. 7. *A particle held by a thin layer.* A particle held by an attractive thin layer has a wave function that falls off exponentially with the distance on both sides, and the layer makes its derivative jump (for the first-order Dirac equation, the wave function itself). In UM-4 the states χ of dirac16complex near the fold are of this kind only near the fold: at rest and without their self-interaction (the field is $\Psi = \mathrm{e}^{(7/2)H|\sigma|}\chi(\sigma)\mathrm{e}^{-\mathrm{i}\omega x_4}$)

$$\begin{aligned}
&\chi(\sigma) = \mathrm{e}^{-q_{\mathrm{b}}|\sigma|}\chi(0^+) \quad (0 < \sigma \ll \sigma_{\mathrm{t}}), \qquad \chi(\sigma) = \mathrm{e}^{-q_{\mathrm{b}}|\sigma|}\chi(0^-) \quad (0 < -\sigma \ll \sigma_{\mathrm{t}}),\\
&\chi(0^+) = \exp\Bigl(\gamma^{(0)}\bigl(\mu_{\mathrm{L}}P_{\mathrm{L}} + \mu_{\mathrm{R}}P_{\mathrm{R}}\bigr)\Bigr)\chi(0^-), \qquad \omega_0 = \omega_{\mathrm{r}} - \frac{\mathrm{i}}{2}\,\Gamma_\chi, \qquad \omega_{\mathrm{r}} \to \omega_{\mathrm{b}} = \sqrt{m_\Psi^2 - q_{\mathrm{b}}^2}
\end{aligned} \tag{FD-9}$$

Symbols in (FD-9).

- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{\frac{7}{2}H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of the level of dirac16complex at the fold (a long-lived resonance) and $\bar{\chi}$ its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- q_{b}: The inverse width of a state of dirac16complex at the fold: inside the turning point its profile falls as $\mathrm{e}^{-q_{\mathrm{b}}|\sigma|}$. Unit: GeV. Status: DERIVED.
- σ_{t}: The turning point of a state of dirac16complex of the frequency ω: the depth at which the warped frequency equals the bulk mass, $|\omega|\mathrm{e}^{H\sigma_{\mathrm{t}}} = m_\Psi$; beyond it every solution oscillates. Unit: length. Status: DERIVED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- μ_X, μ_{L}, μ_{R}: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_{\mathrm{L}}P_{\mathrm{L}} + \mu_{\mathrm{R}}P_{\mathrm{R}})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- P_{L}: The author's left-handed chiral projector, $P_{\mathrm{L}} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- P_{R}: The author's right-handed chiral projector, $P_{\mathrm{R}} = (\mathbf{1} + \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- ω_0: The energy at rest (zero momentum along ordinary space and along the extra times) of a state of dirac16complex at the fold; complex for a resonance, $\omega_0 = \omega_{\mathrm{r}} - \frac{\mathrm{i}}{2}\Gamma_\chi$. Unit: GeV. Status: DERIVED.
- ω_{r}: The mass of the particle χ: the real part of the energy at rest of the resonance of dirac16complex at the fold, $\omega_{\mathrm{r}} = \omega_{\mathrm{b}}(1 - H/(2q_{\mathrm{b}})) + \ldots$. Unit: GeV. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- Γ_χ: The width of the particle χ at rest: the rate of its decay into the waves of dirac16complex in the bulk. Unit: GeV. Value: below $1/t_0 = 1.5 \times 10^{-42}$ GeV for dark matter. Status: DERIVED.
- ω_{b}: The energy at zero momentum of the state of dirac16complex at the fold in the limit $H \to 0$ (the bound state of UM-3): the particle χ of UM-4 has the mass $\omega_{\mathrm{r}} = \omega_{\mathrm{b}}(1 - H/(2q_{\mathrm{b}})) + \ldots$, equal to ω_{b} up to the relative correction $H/(2q_{\mathrm{b}})$. Unit: GeV. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- Γ_8: The chirality matrix of eight dimensions, $\Gamma_8 = T_0T_1\cdots T_7$ (the author's T16^A[8]) $= \mathrm{diag}(-\mathbf{1}_8, \mathbf{1}_8)$; $\Gamma_8^2 = \mathbf{1}$; it anticommutes with every T_a. Unit: dimensionless. Status: AUTHOR.
- Ψ, $\bar{\Psi}$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar{\Psi} = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- $\mathbf{1}$, $\mathbf{1}_8$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_8$ is the 8×8 identity matrix. Unit: dimensionless.

Status: STANDARD.

- $\mathbf{e}_a{}^M$, $\mathbf{e}^a{}_M$: The vielbein (frame field) of the metric: $g_{MN} = \eta_{ab}\,\mathbf{e}^a{}_M \mathbf{e}^b{}_N$; $\mathbf{e}_a{}^M$ is its inverse; $\mathbf{e}_{bN} = \eta_{bc}\mathbf{e}^c{}_N$; $\mathbf{e}^{(a)}$ is the frame one-form $\mathbf{e}^a{}_M\,\mathrm{d}X^M$. Unit: dimensionless. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- $\vec{k}$: The (comoving) momentum along ordinary space, $\vec{k}^2$ its square. Unit: GeV. Status: STANDARD.
- $\vec{x}$: The position in ordinary space, $\vec{x} = (x_1, x_2, x_3)$. Unit: length. Status: AUTHOR.
- t_0: The age of the universe. Unit: time. Value: 13.8 Gyr. Status: STANDARD.
- σ_{16}: The matrix of the Dirac adjoint, $\sigma_{16} = T_0T_1T_2T_3$, real and symmetric; $\sigma_{16}T_a$ is real and antisymmetric. Unit: dimensionless. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.

where μ_L, μ_R are the strengths of the fold for the left- and right-handed halves of dirac16complex (set by the fields of the fold), $q_\mathrm{b} = -m_\Psi(\mu_\mathrm{L} + \mu_\mathrm{R})\tanh\varsigma/(2\varsigma)$, $\varsigma = \sqrt{\mu_\mathrm{L}\mu_\mathrm{R}}$ ($q_\mathrm{b} = -m_\Psi\mu_\mathrm{L}/2$ for $\mu_\mathrm{R} = 0$), and ω_r tends to the energy ω_b of the bound state of UM-3 when $H/q_\mathrm{b} \to 0$. Because the time is warped, the energy ω of the field enters its equation multiplied by $\mathrm{e}^{H|\sigma|}$: beyond the turning point σ_t ($\omega_\mathrm{r}\mathrm{e}^{H\sigma_\mathrm{t}} = m_\Psi$) the profile oscillates as a wave that leaves the fold, so the state is a resonance of the width Γ_χ, which falls like $\mathrm{e}^{-2F(\upsilon_\mathrm{b})m_\Psi/H}$ with $F(\upsilon) = \ln((1+\sqrt{1-\upsilon^2})/\upsilon) - \sqrt{1-\upsilon^2}$ and $\upsilon_\mathrm{b} = \omega_\mathrm{b}/m_\Psi$ (the tunnelling through the region between the fold and σ_t). The profile is not a function of $|\sigma|$: it jumps at the fold, $\chi(0^+) \neq \chi(0^-)$. 8. *The fold as a mirror for dirac16complex.* The reflection across the fold acts on the profiles $\chi(\sigma)$ in two ways,

$$(\mathsf{R}_1\chi)(\sigma) = T_0T_1T_2T_3\,\chi(-\sigma), \qquad (\mathsf{R}_2\chi)(\sigma) = T_0T_5T_6T_7\,\chi(-\sigma) \tag{FD-10}$$

Symbols in (FD-10).

- R_1: The reflection across the fold acting on a profile of dirac16complex together with the reflection of ordinary space, $(\mathsf{R}_1\chi)(\sigma) = T_0T_1T_2T_3\,\chi(-\sigma)$ (FD-10): a symmetry of dirac16complex, of gravity and of the mean field of the gas of the particles χ; the states at the fold of Krein charge -1 are even, those of Krein charge $+1$ odd. Unit: dimensionless. Status: DERIVED.
- R_2: The reflection across the fold acting on a profile of dirac16complex together with the reflection of the extra times, $(\mathsf{R}_2\chi)(\sigma) = T_0T_5T_6T_7\,\chi(-\sigma)$ (FD-10): it maps the states at the fold of one Krein charge onto those of the other. Unit: dimensionless. Status: DERIVED.
- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{\frac{7}{2}H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x}-\omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of the level of dirac16complex at the fold (a long-lived resonance) and $\bar{\chi}$ its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- T_a, T^a: The author's eight real 16×16 Dirac matrices T16^A[a] of 4+4.nb ($a = 0, \ldots, 7$; $T_0, \ldots, T_7$), with $\{T_a, T_b\} = 2\eta_{ab}\mathbf{1}$; $T^a = \eta^{ab}T_b$. Unit: dimensionless. Status: AUTHOR.

(T_a the author's Dirac matrices; for states at rest R_1 is the reflection across the fold combined with the reflection of ordinary space, R_2 the reflection across the fold combined with the reflection of the extra times). The Krein charge of a state is its share of the conserved charge of dirac16complex, which is indefinite: a state can carry the charge -1 or $+1$. R_1 is a symmetry of dirac16complex, of gravity and of a mean field of a gas of the particles χ that is the same on both sides of the fold: the four resonances of Krein charge -1 (the particles χ) are even under it and the four of Krein charge $+1$ are odd, so such a mean field cannot mix the two signs of the indefinite charge. R_2 maps the states of one Krein charge onto those of the other and makes the two sets degenerate without the self-interaction. 9. *The planes of Horava and Witten.* In M-theory (the eleven-dimensional theory whose limits include the superstring theories) on an interval (a circle whose points are identified with their mirror images), the two ten-dimensional planes left in place by the reflection are mirror surfaces (property 1); each carries one E_8 gauge group while gravity fills the bulk. As in UM-4, gravity fills the bulk; unlike UM-4, the gauge fields live on the mirror planes (in UM-4 they live in the bulk), and the two sides of each plane are identified (one physical side).

What is not a fold.

- *The tips* $z = 0$ and $z = \pi \pmod{\pi}$: the mirror exchanges them within a period, but there the warp factor vanishes, $g_{x_0 x_0}$ diverges, and they lie at an infinite proper distance; they carry no thin source.
- *A smooth maximum of a warp factor*, such as $1/\cosh^2(H\sigma)$: mirror symmetric and largest at $\sigma = 0$, but without a kink and without a thin source.

2 How to read this document

2.1 What this document contains

This document answers the author's order of 2026-10-25, 13:15 PDT. It shows why the universe model UM-3 was physically incorrect (whatever propagated in its bulk outran light on the fold), which metric replaces it and why that metric is the best choice, which theory of gravity goes with it, and it derives, for the universe model UM-4 built on them, that photons and gravitons propagate at the speed of light in the bulk, that the photon is a transverse wave of spin 1 and that the gravitational wave is a wave of spin 2 with the plus and cross polarisations, which stretch and squeeze space perpendicularly. It then gives the propagation of every other field of the bulk, the propagators of the bulk from one point of the fold to another (Newton's and Coulomb's laws, the W and Z bosons and the bounds on the author's constant H that follow), the causal structure of the bulk, and what the change of the metric changes elsewhere in the model. The document is self-contained: every equation it uses is written out, and every symbol of every display equation is defined right after it; a symbol that one of these definitions uses is defined in this document as well. The equations of the field equations of UM-4 keep their tags (FE-n for a field equation or a result, IN-n for a term of the action), the same in every UM-4 document; the equations that only this document needs carry the tags P-1 and P-2.

2.2 The status labels

Every equation and every result carries one of the labels **AUTHOR** (from the author: his notebooks `4+4.nb` and `Dirac_Einstein-Lovelock_4+4_Nash.nb` and his orders of 2026-10-24 and 2026-10-25), **RECORD** (from the record of this repository), **STANDARD** (textbook physics or a published measurement), **ASSUMED** (a choice of UM-4 without derivation), **INVENTED** (a new interaction), **DERIVED** (computed and verified in UM-4: the program and the check that verify it are named), **ESTIMATE** (an order of magnitude) and **OPEN** (not settled).

2.3 Coordinates, signature and units

The author's coordinates are $X = \{x_0, x_1, \ldots, x_7\}$, always with the hidden direction first: x_0 is the hidden space-like coordinate, x_1, x_2, x_3 ordinary space, x_4 the time, and x_5, x_6, x_7 three extra time-like coordinates. The signature is $(+,+,+,+,-,-,-,-)$ in this order. Units are natural: the reduced Planck constant and the speed of light are 1. A prime on a function of σ alone (a profile along the hidden direction) is its derivative with respect to σ; on a function of the time x_4 alone it is the derivative with respect to x_4. "The speed of light" without a qualifier is the speed of light on the fold, one unit of x_1, x_2 or x_3 per unit of x_4 on the static fold.

3 The order and the answer

On 2026-10-25 at 13:15 PDT the author ordered:

> "Since in the UM-3 model anything that propagates in the bulk outruns light on the fold, and bulk scalars and photons faster than light, my model is physically incorrect. You must find a new metric model in which x4 is warped like the other directions, both photons and gravitons actually propagate at c in the bulk, photon waves are transverse spin-1 phenomena, Gravitational waves (ripples in spacetime) are spin-2 phenomena (because they originate from the rank-2 stress-energy tensor) and possess the following Polarization Modes: the spin-2 nature of gravitational waves manifests physically as characteristic"plus" (+) and "cross" (×) polarization modes that stretch and squeeze space perpendicularly. Find the best model and re-calculate and re-document; however keep UM-3 and create a new UM-4, placed in new folders."

The answer, point by point (UM-3 is kept unchanged in its folder `universe_model_3`; UM-4 is in the new folder `universe_model_4`):

1. **UM-3 was physically incorrect.** Its time was not warped, so the light cone of the bulk at the depth σ was $\mathrm{e}^{H|\sigma|}$ times wider than that of the fold (P-1): bulk scalars moved 1.22 times and bulk photons 1.41 times as fast as light on the fold at long wavelengths, and faster than light at every wavelength (P-2). Status: DERIVED.
2. **The new metric.** The metric of UM-4 is the author's metric with the time warped by the same factor $(\sin^2 z)^{1/6}$ as the other six directions of the fold (FE-27), the smallest change of the author's metric that meets the order. Every slice of constant depth then has the light cone of the fold (P-1, FE-83). Among the static warps of this kind it is the only one that is a vacuum of gravity with a cosmological constant (P-1), and its static block is a space of constant curvature $-H^2$ (FE-32). Status: AUTHOR (the metric and the order), ASSUMED (the way the order is met), DERIVED (its properties).
3. **The gravity.** The author's Einstein-Lovelock theory with the weights of its Gauss-Bonnet and cubic terms set to zero: Einstein gravity with the cosmological constant $\Lambda = -21H^2$ (IN-2). It is the only choice of the weights for which gravitons travel on the light cone on every background, not only around the static block (FE-68 to FE-70). Status: DERIVED.
4. **Gravitons at the speed of light in the bulk; spin 2 with the plus and cross polarisations.** Each polarisation of the graviton obeys the equation of a massless scalar of the bulk; its massless mode is bound to the fold and travels at the speed of light at every depth and every wavelength. The four-dimensional world sees it as a transverse traceless wave of helicity ± 2 with the two polarisations + and $\times$, which stretch and squeeze a ring of test masses along perpendicular directions (FE-76); on the expanding fold it obeys the wave equation of general relativity and travels at the speed of light at every wavelength (FE-63). Status: DERIVED.
5. **Photons at the speed of light in the bulk; transverse waves of spin 1.** The gauge fields of the Standard Model live in the bulk (IN-7); their massless modes are bound to the fold and travel at the speed of light (FE-74); the photon is a transverse wave of helicity ± 1 whose electric and magnetic fields are orthogonal to its direction of motion and to each other (FE-75). Status: ASSUMED (the gauge fields in the bulk), DERIVED (the rest).
6. **Every other field of the bulk.** A massless scalar (FE-73), a fermion of the bulk (FE-77) and the author's field dirac16complex (FE-78) move at most at the speed of light; the massive modes are slower. Status: DERIVED.
7. **Nothing crosses the bulk faster than light.** The static block is conformally flat, so a signal that leaves the fold and comes back arrives later than light along the fold (FE-83); around an expanding fold the largest possible advance over the Hubble distance today is 4.5×10^{-94} of the travel time for $H = 2.5$ TeV (FE-84). Status: DERIVED.
8. **The price.** Photons in the bulk make the W and Z bosons resonances of the bulk and modify Coulomb's law; the measured invisible width of the Z boson bounds H from below by 2.5 TeV (FE-82); the states of dirac16complex at the fold become long-lived resonances (FE-86). Status: DERIVED, ESTIMATE (the bounds).

4 Why UM-3 was physically incorrect

$$\begin{aligned} &\mathrm{d}\sigma^2 + \mathrm{e}^{-2\mathfrak{p}H|\sigma|}\,\mathrm{d}\vec{x}^2 - \mathrm{e}^{-2\mathfrak{q}H|\sigma|}\,\mathrm{d}x_4^2 - \mathrm{e}^{-2\mathfrak{r}H|\sigma|}\,\mathrm{d}\vec{x}_5^2: \qquad c_{\mathrm{loc}}(\sigma) = \mathrm{e}^{(\mathfrak{p}-\mathfrak{q})H|\sigma|}, \\ &\text{UM-3 } (\mathfrak{p}=\mathfrak{r}=1,\ \mathfrak{q}=0): \quad c_{\mathrm{loc}} = \mathrm{e}^{H|\sigma|} > 1, \qquad \text{UM-4 } (\mathfrak{p}=\mathfrak{q}=\mathfrak{r}=1): \quad c_{\mathrm{loc}} = 1 \end{aligned} \tag{P-1}$$

Symbols in (P-1).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- $\mathfrak{p}$, $\mathfrak{q}$, $\mathfrak{r}$: The rates of the warps of ordinary space ($\mathfrak{p}$), of the time ($\mathfrak{q}$) and of the extra times ($\mathfrak{r}$) of a static warped background, in units of H. Unit: dimensionless. Value: $\mathfrak{p} = \mathfrak{q} = \mathfrak{r} = 1$ for the static block of UM-4; $\mathfrak{p} = \mathfrak{r} = 1$, $\mathfrak{q} = 0$ for that of UM-3. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free

parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.

- $\vec{x}$: The position in ordinary space, $\vec{x} = (x_1, x_2, x_3)$. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\vec{x}_5$: The position along the extra times, $\vec{x}_5 = (x_5, x_6, x_7)$; $\mathrm{d}\vec{x}_5^2 = \mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2$. Unit: length. Status: AUTHOR.
- c_{loc}: The coordinate speed of light along x_3 at the depth σ (the rate of change of x_3 with x_4 along a null line at fixed σ), in units of that on the fold: $\mathrm{e}^{H|\sigma|}$ on the static block of UM-3, whose time was not warped, 1 on the static block of UM-4, and on an expanding fold of UM-4 different from 1 in proportion to the acceleration a_4 of the expansion. Unit: dimensionless. Status: DERIVED.

The static warped metrics: the hidden direction σ unwarped, ordinary space, the time and the extra times warped with the rates $\mathfrak{p}$, $\mathfrak{q}$ and $\mathfrak{r}$ (in units of H), on both sides of the fold. A signal of light that moves along x_3 at fixed depth follows a null line (the metric vanishes along it), so its coordinate speed is $\mathrm{d}x_3/\mathrm{d}x_4 = \mathrm{e}^{(\mathfrak{p}-\mathfrak{q})H|\sigma|}$, in units of the speed of light on the fold ($\sigma = 0$): the light cone of the slice at the depth σ is that of the fold only if the time is warped like ordinary space, $\mathfrak{q} = \mathfrak{p}$. The author's metric with his original time component (UM-3) has $\mathfrak{p} = \mathfrak{r} = 1$ and $\mathfrak{q} = 0$: at the depth σ light moves $\mathrm{e}^{H|\sigma|}$ times as fast as on the fold, and every field that propagates in the bulk inherits this wider light cone. The metric of UM-4 has $\mathfrak{p} = \mathfrak{q} = \mathfrak{r} = 1$. Equal rates are also singled out by gravity: the Einstein tensor of these metrics is diagonal, with ${G^{(1)}}_{(1)} - {G^{(4)}}_{(4)} = -(\mathfrak{p} - \mathfrak{q})(3\mathfrak{p} + \mathfrak{q} + 3\mathfrak{r})H^2$ and ${G^{(1)}}_{(1)} - {G^{(5)}}_{(5)} = -(\mathfrak{p} - \mathfrak{r})(3\mathfrak{p} + \mathfrak{q} + 3\mathfrak{r})H^2$ (frame components along x_1, x_4 and x_5), and the only real rates for which the metric is a vacuum of Einstein gravity with a cosmological constant (the Einstein tensor the same in every direction) are $\mathfrak{q} = \mathfrak{r} = \mathfrak{p}$, with $\Lambda = -21\mathfrak{p}^2H^2$; the static block of UM-3 is not one (its differences are 0, $-6H^2$ and 0). Status: DERIVED (`um4_geometry.py`, G8: the coordinate speed of light at every depth for the metric of UM-4; G10: the Einstein tensors of the static warps and their vacua, with sympy).

$$\begin{aligned} &\text{UM-3:}\quad \phi_{\mathrm{bulk}} = u(\sigma)\,\mathrm{e}^{\mathrm{i}(kx_3 - \omega x_4)}: \qquad u'' - 6H\,\mathrm{sgn}(\sigma)\,u' + \left(\omega^2 - \mathrm{e}^{2H|\sigma|}k^2\right)u = 0, \\ &\text{scalar:}\quad \frac{\omega^2}{k^2} \to \frac{\int \mathrm{d}\sigma\,\mathrm{e}^{-4H|\sigma|}}{\int \mathrm{d}\sigma\,\mathrm{e}^{-6H|\sigma|}} = \frac{3}{2}, \qquad \text{photon:}\quad \frac{\omega^2}{k^2} \to \frac{\int \mathrm{d}\sigma\,\mathrm{e}^{-2H|\sigma|}}{\int \mathrm{d}\sigma\,\mathrm{e}^{-4H|\sigma|}} = 2 \qquad (k \ll H) \end{aligned} \tag{P-2}$$

Symbols in (P-2).

- ϕ_{bulk}: A massless scalar field of the bulk, $\nabla_M \nabla^M \phi_{\mathrm{bulk}} = 0$ (an illustration: each polarisation of the graviton obeys its equation). Unit: GeV^3. Status: STANDARD.
- $u(\sigma)$, u_0: The profile along the hidden direction of a mode of a field of the bulk (a scalar, a gauge field or a polarisation of the graviton): the field is $u(\sigma)$ times a field of the fold; $u_0 = 1$ is the profile of the massless mode (the zero mode); in a propagator the response of the field to a unit source on the fold. Unit: dimensionless (a relative amplitude); length in a propagator. Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- k: A wave number: the momentum of a plane wave along x_3, $\mathrm{e}^{\mathrm{i}(kx_3 - \omega x_4)}$; in the cosmological perturbations the comoving wave number of a Fourier mode, $k = |\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume

element, the product of their differentials. Status: STANDARD.
- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols $\Gamma^P{}_{MN}$). Status: STANDARD.
- $\Gamma^P{}_{MN}$: The Christoffel symbols of the metric g_{MN} (the Levi-Civita connection). Unit: 1/length. Status: STANDARD.

The speeds of UM-3. A massless scalar of the bulk of UM-3 (on its static block, $a_4 = 0$, $b_4 = 1$) for a wave of the frequency ω and the wave number k along x_3 has a profile $u(\sigma)$ that obeys the first line: the volume element of UM-3 is $\mathrm{e}^{-6H|\sigma|}$ (six warped directions) and only the momentum carries the factor $\mathrm{e}^{2H|\sigma|}$, the frequency not, so ω and k do not enter through $\omega^2 - k^2$ and the modes are not waves of the Lorentz-invariant fold. At long wavelengths ($k \ll H$) the lowest mode has the profile $u = 1$, and its frequency is the mean of $\mathrm{e}^{2H|\sigma|}k^2$ over the kinetic weight $\mathrm{e}^{-6H|\sigma|}$, $\omega^2 = \frac{3}{2}k^2$: the bulk scalar of UM-3 moved $\sqrt{3/2} = 1.22$ times as fast as light on the fold. The photon (the polarisation along x_1, whose weights carry one more factor $\mathrm{e}^{2H|\sigma|}$) had $\omega^2 = 2k^2$, 1.41 times the speed of light. At shorter wavelengths the lowest mode is squeezed towards the fold and the excess falls, but it never vanishes: $\omega^2/k^2 = 1.4999$, 1.4993, 1.4880, 1.4657 and 1.3818 for the scalar and 1.9995, 1.9825, 1.8658, 1.7499 and 1.5150 for the photon at $k/H = 0.01$, 0.1, 0.5, 1 and 3. The gravitons of UM-3 were no better: with Einstein gravity they would move at the local speed of light of the bulk, $\mathrm{e}^{H|\sigma|}$ times that of the fold (P-1), and with the author's original Lovelock weights $\sqrt{5}$ times faster still (FE-69); UM-3's corrected weights made its static block a degenerate vacuum without linear gravitons (FE-70). Status: DERIVED (`um4_bulk.py`, B9: the profile equation of UM-3 on both halves with sympy, the two ratios exactly, and the lowest ω^2 of the operator in a box $|\sigma| < 6/H$; the same box gives exactly 1 for the metric of UM-4, B1 and B2).

5 The metric of UM-4

$$
\begin{aligned}
g_{MN} &= \mathrm{diag}\Big(\cot^2 z,\ \mathrm{e}^{2a_4}(\sin^2 z)^{1/6},\ \mathrm{e}^{2a_4}(\sin^2 z)^{1/6},\ \mathrm{e}^{2a_4}(\sin^2 z)^{1/6},\ -(\sin^2 z)^{1/6},\\
&\qquad -\mathrm{e}^{-2a_4}b_4^2(\sin^2 z)^{1/6},\ -\mathrm{e}^{-2a_4}b_4^2(\sin^2 z)^{1/6},\ -\mathrm{e}^{-2a_4}b_4^2(\sin^2 z)^{1/6}\Big), \qquad z = 6Hx_0,\\
\sigma &= \mathrm{sgn}\Big(\frac{\pi}{2} - z\Big)\frac{\ln\sin z}{6H}, \qquad \mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0,\\
g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N &= \mathrm{d}\sigma^2 + \mathrm{e}^{-2H|\sigma|}\Big[\mathrm{e}^{2a_4}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{d}x_4^2 - \mathrm{e}^{-2a_4}b_4^2\big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big)\Big]
\end{aligned}
\tag{FE-27}
$$

Symbols in (FE-27).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\ldots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$, the fold is $z = \pi/2$. Unit: dimensionless. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.

- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.

The metric of UM-4 in the author's coordinates and in the proper coordinate σ of the hidden direction: the author's metric with his two functions of the time $a_4(x_4)$ and $b_4(x_4)$ and his constant H, with the one change that the order requires: the time carries the warp factor $(\sin^2 z)^{1/6}$ of the other six directions of the fold, $g_{x_4x_4} = -(\sin^2 z)^{1/6}$ (UM-3: -1). In the proper coordinate the warp factor $\mathrm{e}^{-2H|\sigma|}$ multiplies the whole metric of the fold, the time included. The change keeps everything else of the author's metric: the coordinates, the period $\pi/(6H)$ of x_0, the fold at $z = \pi/2$ with its four properties (the mirror, the largest warp, the kink, the degenerate coordinate x_0), the functions a_4 and b_4, and the warp of ordinary space and of the extra times. Status: AUTHOR (the metric and the order), ASSUMED (the way the order is met), DERIVED (`um4_geometry.py`, G1: the proper chart on both halves of the period).

$$a_4,\ b_4 \text{ constant}: \qquad R^{(a)(b)}{}_{(a)(b)} = -H^2 \text{ in all 28 frame planes}, \qquad \mathcal{E}_{(1)}{}^a{}_b = 21H^2\,\delta^a{}_b, \qquad \Lambda = -21H^2 \quad \text{(FE-32)}$$

Symbols in (FE-32).

- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- $R^{MN}{}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor $R^M{}_{NPQ}$ (in the convention in which the commutator of two covariant derivatives, acting on a vector, is the Riemann tensor, with its last two indices those of the two derivatives, contracted with the vector), with two indices the Ricci tensor $R_{MN} = R^P{}_{MPN}$, without indices the scalar curvature $R = g^{MN}R_{MN}$; with frame indices $R^{(a)(b)}{}_{(c)(d)}$ its frame components, $R^{(a)(b)}{}_{(a)(b)}$ (no sum) the sectional curvature of the frame plane (a, b). Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free

parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.

- ${\mathcal{E}_{(p)}}^M{}_N$: The Lovelock tensor of order p, ${\mathcal{E}_{(p)}}^M{}_N = -2^{-(p+1)}\,\delta^{MP_1\cdots P_{2p}}_{NQ_1\cdots Q_{2p}}\,R^{Q_1Q_2}{}_{P_1P_2}\cdots R^{Q_{2p-1}Q_{2p}}{}_{P_{2p-1}P_{2p}}$ (with both indices down it is $1/\sqrt{|g|}$ times the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to g^{MN}; $\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor, $\mathcal{E}_{(3)}$ the cubic Lovelock tensor); with frame indices its frame components. Unit: $1/\text{length}^{2p}$. Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number (UM-4 keeps only $p = 1$); in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1M_2\cdots}_{N_1N_2\cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- Λ: The eight-dimensional cosmological constant of UM-4, $\Lambda = -21H^2$: the value for which the static block is a vacuum of Einstein gravity (its Einstein tensor is $21H^2$ in every direction). The author writes the pure number Λ_N in $0 = -\Lambda_\mathrm{N} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$, so that $\Lambda = -\Lambda_\mathrm{N} H^2$. Unit: GeV^2. Value: $-21H^2$; the author's original $-10H^2$; UM-3: $-7H^2$. Status: AUTHOR (the term), DERIVED (the value).

The static block (a_4 and b_4 constant): its curvature is $-H^2$ in all 28 planes of the frame, a space of constant curvature in all eight directions (locally the anti-de Sitter space of eight dimensions of radius $1/H$; UM-3 had 0 in the seven planes that contain the time), and its Einstein tensor is $21H^2$ in every direction, so it is a vacuum of Einstein gravity with $\Lambda = -21H^2$. Status: DERIVED (`um4_geometry.py`, G3 and G4).

$$K^a{}_b = \frac{1}{2}\,g^{ac}\,\partial_\sigma g_{cb} = -H\,\mathrm{sgn}(\sigma)\,\delta^a{}_b, \qquad \kappa\,\mathcal{S}^a{}_b = -12H\,\delta^a{}_b \quad \big(\kappa\lambda_\mathrm{F} = 12H\big) \tag{FE-30}$$

Symbols in (FE-30).

- $K^a{}_b$: The extrinsic curvature of a surface of constant σ of the hidden direction, $K^a{}_b = \frac{1}{2}g^{ac}\partial_\sigma g_{cb}$ for a, b along the surface (FD-5): $-H\,\mathrm{sgn}(\sigma)$ along all seven directions of the fold for the author's form of the metric of UM-4; $K = K^c{}_c$ its trace. Unit: GeV (1/length). Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+, +, +, +, -, -, -, -$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- c: A frame index (as a). Status: STANDARD.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- sgn: The sign function: +1 for a positive argument and −1 for a negative one. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1M_2\cdots}_{N_1N_2\cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.

- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma\, \mathcal{E}_{(1)}{}^a{}_b$ (the junction condition of Israel); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -12H\delta^a{}_b$ for every a_4, b_4 of the author's form. Status: DERIVED.
- λ_F: The tension that the static block requires of the fold, $\kappa\lambda_\mathrm{F} = 12H$: the total vacuum energy of the seven-dimensional fold, made of the tension of the fold fluid and the vacuum energy of the fields of the fold (both have the pressure minus their energy density along ordinary space and along the extra times), per unit volume of the six other directions of the fold (eight-dimensional normalisation); in the four-dimensional normalisation it is $\lambda_\mathrm{F}V_*$, and the excess of the total over $\lambda_\mathrm{F}V_*$ is the dark energy ρ_Λ. Unit: GeV^7. Value: $12H/\kappa$; $\lambda_\mathrm{F}V_* = 30H^2M_\mathrm{red}^2 = 1.1\times10^{45}$ GeV^4 for $H = 2.5$ TeV. Status: DERIVED.

The fold: the slices of constant σ have the extrinsic curvature $-H\,\mathrm{sgn}(\sigma)$ in all seven directions of the fold (all directions are scaled by the same factor; UM-3 had 0 along the time), it jumps by $-2H$ at the fold, and Israel's junction condition makes the fold a pure tension $\kappa\lambda_\mathrm{F} = 12H$, for every a_4 and b_4 of the author's form: a brane of seven dimensions of the kind of Randall and Sundrum. Status: DERIVED (`um4_geometry.py`, G7).

$$\tilde\sigma = \mathrm{sgn}(\sigma)\,\frac{\mathrm{e}^{H|\sigma|}-1}{H}: \qquad g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N = \frac{\mathrm{d}\tilde\sigma^2+\mathrm{d}x_1^2+\mathrm{d}x_2^2+\mathrm{d}x_3^2-\mathrm{d}x_4^2-\mathrm{d}x_5^2-\mathrm{d}x_6^2-\mathrm{d}x_7^2}{\left(1+H|\tilde\sigma|\right)^2}, \qquad \Delta x_4 \geq L \tag{FE-83}$$

Symbols in (FE-83).

- $\tilde\sigma$: The conformal coordinate of the hidden direction on the static block, $\tilde\sigma = \mathrm{sgn}(\sigma)(\mathrm{e}^{H|\sigma|}-1)/H$, in which the static block is conformally flat. Unit: length. Status: DERIVED.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2-z)\ln(\sin z)/(6H)$ on one period $0<z<\pi$; $\sigma=0$ is the fold $z=\pi/2$, $\sigma\to\mp\infty$ the tips $z\to0,\pi$; one period of x_0 is the whole line $-\infty<\sigma<\infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\ldots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c=1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a

perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.

- L: A distance between two points of the fold along ordinary space: for a gravitational wave the radius of a ring of free test masses (ΔL the change of the distance of one of them from the centre), for a signal the distance between its emitter and its receiver. Unit: length. Status: STANDARD.

The light cone of the bulk: in the conformal coordinate $\tilde{\sigma}$ the static block is flat space of signature (4, 4) times the factor $(1 + H|\tilde{\sigma}|)^{-2}$, which does not change which curves are null; the null geodesics are the null straight lines. Every slice has the light cone of the fold, and a signal that leaves a point of the fold, travels through the bulk at the local speed of light and arrives at another point of the fold at the distance L along ordinary space needs the time $\Delta x_4 \geq L$, with equality only for the path along the fold: nothing crosses the static bulk faster than light on the fold. Status: DERIVED (`um4_causality.py`, C1: the conformal form exactly, the geodesic equation, 200 sampled paths through the bulk, all later than the path along the fold).

Why this is the best metric. The order fixes the time component up to the meaning of "warped like the other directions"; P-1 shows that only the same rate for the time and for ordinary space gives photons and gravitons the speed of light at every depth, and that only the same rate in all seven directions makes the static block a vacuum of gravity with a cosmological constant. A warp of the time with a different function of z would give $c_{\mathrm{loc}} \neq 1$ somewhere in the bulk. A metric without a warp (a flat bulk) would have no fold and no bound graviton. Smooth warps without a kink (example "What is not a fold" of the definition of the fold) carry no thin source and hold no field at a surface. The metric FE-27 is therefore the unique metric of the author's form, with his warp factor, that meets the order. Status: DERIVED (P-1), ASSUMED (that the author's warp factor and fold are kept).

6 Gravitons at the speed of light: spin 2 with the plus and cross polarisations

6.1 Which gravity

$$S_{\mathrm{EL}} = \frac{1}{2\kappa}\int \mathrm{d}^8X\,\sqrt{|g|}\,\Big(\sum_{p=1}^{3}\alpha_{(p)}\,\mathcal{L}_{(p)} - 2\Lambda\Big) = \frac{1}{2\kappa}\int \mathrm{d}^8X\,\sqrt{|g|}\,\big(R - 2\Lambda\big), \qquad \kappa = 8\pi G_8,$$
$$\alpha_{(p)} = w_{(p)}\,H^{2-2p}, \qquad w_{(1)} = 1, \quad w_{(2)} = w_{(3)} = 0, \qquad \Lambda = -21H^2 \tag{IN-2}$$

Symbols in (IN-2).

- S_{EL}: The Einstein-Lovelock action of gravity in eight dimensions; in UM-4 the Einstein-Hilbert action with the cosmological constant. Unit: dimensionless. Status: AUTHOR.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number (UM-4 keeps only $p = 1$); in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)}H^{2-2p}$: in UM-4 $\alpha_{(1)} = 1$ and $\alpha_{(2)} = \alpha_{(3)} = 0$ (Einstein gravity). Unit: GeV^{2-2p}. Status: AUTHOR.
- $\mathcal{L}_{(p)}$: The Lovelock density of order p, $\mathcal{L}_{(p)} = 2^{-p}\,\delta^{P_1\cdots P_{2p}}_{Q_1\cdots Q_{2p}}\,R^{Q_1Q_2}{}_{P_1P_2}\cdots R^{Q_{2p-1}Q_{2p}}{}_{P_{2p-1}P_{2p}}$, a scalar built from p curvature tensors: $\mathcal{L}_{(1)} = R$ (Einstein-Hilbert), $\mathcal{L}_{(2)}$ the Gauss-Bonnet density, $\mathcal{L}_{(3)}$ the cubic Lovelock density. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant of UM-4, $\Lambda = -21H^2$: the value for which the static block is a vacuum of Einstein gravity (its Einstein tensor is $21H^2$ in every direction). The author writes the pure

number $\Lambda_{\rm N}$ in $0 = -\Lambda_{\rm N} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$, so that $\Lambda = -\Lambda_{\rm N} H^2$. Unit: GeV^2. Value: $-21H^2$; the author's original $-10H^2$; UM-3: $-7H^2$. Status: AUTHOR (the term), DERIVED (the value).

- $R^{MN}{}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor $R^M{}_{NPQ}$ (in the convention in which the commutator of two covariant derivatives, acting on a vector, is the Riemann tensor, with its last two indices those of the two derivatives, contracted with the vector), with two indices the Ricci tensor $R_{MN} = R^P{}_{MPN}$, without indices the scalar curvature $R = g^{MN} R_{MN}$; with frame indices $R^{(a)(b)}{}_{(c)(d)}$ its frame components, $R^{(a)(b)}{}_{(a)(b)}$ (no sum) the sectional curvature of the frame plane (a, b). Unit: 1/length^2. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_8: The eight-dimensional Newton constant ($\kappa = 8\pi G_8$). Unit: GeV^{-6}. Status: AUTHOR.
- $w_{(p)}$: The Lovelock weights of the author's Einstein-Lovelock action, $w_{(1)} = 1$ and $w_{(2)}$, $w_{(3)}$ of the Gauss-Bonnet and the cubic densities. UM-4 sets $w_{(2)} = w_{(3)} = 0$ (Einstein gravity): the only weights for which gravitons propagate on the light cone on every background, not only around the static block. The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) keep the gravitons on the light cone only around the static block (with $\Lambda = -161H^2/16$); UM-3 used $(1, 1/20, 1/360)$. Unit: dimensionless. Value: $(1, 0, 0)$ in UM-4; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), DERIVED (the values that the order of 2026-10-25 selects).
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\rm F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.

The gravity of UM-4: the author's Einstein-Lovelock action (his notebook `Dirac_Einstein-Lovelock_4+4_Nash.nb`, In[134]) with the weights of the Gauss-Bonnet density and of the cubic Lovelock density set to zero, which is Einstein gravity in eight dimensions with the cosmological constant $\Lambda = -21H^2$. The three displays below show why. Status: AUTHOR (the action), DERIVED (the weights).

$$
\begin{aligned}
&\sum_{p=1}^{3} \alpha_{(p)}\, \mathcal{E}_{(p)}{}^a{}_b + \Lambda\, \delta^a{}_b = \mathsf{P}(\mathsf{y})\, H^2\, \delta^a{}_b, \qquad \mathsf{P}(\mathsf{y}) = 21\mathsf{y} - 420 w_{(2)} \mathsf{y}^2 + 2520 w_{(3)} \mathsf{y}^3 + \frac{\Lambda}{H^2}, \\
&\text{Einstein } (w_{(2)} = w_{(3)} = 0,\ \Lambda = -21H^2): \quad \mathsf{P} = 21(\mathsf{y} - 1); \\
&\text{the author's original weights } \Big(\frac{1}{32}, \frac{1}{1152}\Big),\ \Lambda = -\frac{161}{16} H^2: \quad \mathsf{P} = \frac{7}{16}(\mathsf{y} - 1)\big(5\mathsf{y}^2 - 25\mathsf{y} + 23\big); \\
&\text{UM-3's weights } \Big(\frac{1}{20}, \frac{1}{360}\Big),\ \Lambda = -7H^2: \quad \mathsf{P} = 7(\mathsf{y} - 1)^3
\end{aligned}
\tag{FE-70}
$$

Symbols in (FE-70).

- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number (UM-4 keeps only $p = 1$); in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)} H^{2-2p}$: in UM-4 $\alpha_{(1)} = 1$ and $\alpha_{(2)} = \alpha_{(3)} = 0$ (Einstein gravity). Unit: GeV^{2-2p}. Status: AUTHOR.
- $\mathcal{E}_{(p)}{}^M{}_N$: The Lovelock tensor of order p, $\mathcal{E}_{(p)}{}^M{}_N = -2^{-(p+1)}\, \delta^{MP_1\cdots P_{2p}}_{NQ_1\cdots Q_{2p}}\, R^{Q_1Q_2}{}_{P_1P_2} \cdots R^{Q_{2p-1}Q_{2p}}{}_{P_{2p-1}P_{2p}}$ (with both indices down it is $1/\sqrt{|g|}$ times the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to g^{MN}; $\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor, $\mathcal{E}_{(3)}$ the cubic Lovelock tensor); with frame indices its frame components. Unit: 1/length^{2p}. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- Λ: The eight-dimensional cosmological constant of UM-4, $\Lambda = -21H^2$: the value for which the static block is a vacuum of Einstein gravity (its Einstein tensor is $21H^2$ in every direction). The author writes the pure number $\Lambda_{\rm N}$ in $0 = -\Lambda_{\rm N} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$, so that $\Lambda = -\Lambda_{\rm N} H^2$. Unit: GeV^2. Value: $-21H^2$; the author's

original $-10H^2$; UM-3: $-7H^2$. Status: AUTHOR (the term), DERIVED (the value).

- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- $\mathsf{P}(\mathsf{y})$: The vacuum polynomial of the static blocks of curvature $-\mathsf{y}H^2$: the left-hand side of the bulk equations divided by H^2, the same in every direction, $\mathsf{P}(\mathsf{y}) = 21\mathsf{y} - 420w_{(2)}\mathsf{y}^2 + 2520w_{(3)}\mathsf{y}^3 + \Lambda/H^2$; $\mathsf{P}'(1)$ its derivative at $\mathsf{y} = 1$, which sets the strength of the graviton around the static block relative to Einstein gravity ($\mathsf{P}'(1)/21$). Unit: dimensionless. Value: $21(\mathsf{y} - 1)$ for Einstein gravity with $\Lambda = -21H^2$. Status: DERIVED.
- y: The curvature of a static block of constant curvature $-\mathsf{y}H^2$ in all eight directions, in units of $-H^2$ (its warp is $\mathrm{e}^{-2\sqrt{\mathsf{y}}H|\sigma|}$ in all seven directions of the fold): $\mathsf{y} = 1$ is the static block of UM-4. Unit: dimensionless. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- $w_{(p)}$: The Lovelock weights of the author's Einstein-Lovelock action, $w_{(1)} = 1$ and $w_{(2)}$, $w_{(3)}$ of the Gauss-Bonnet and the cubic densities. UM-4 sets $w_{(2)} = w_{(3)} = 0$ (Einstein gravity): the only weights for which gravitons propagate on the light cone on every background, not only around the static block. The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) keep the gravitons on the light cone only around the static block (with $\Lambda = -161H^2/16$); UM-3 used $(1, 1/20, 1/360)$. Unit: dimensionless. Value: $(1, 0, 0)$ in UM-4; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), DERIVED (the values that the order of 2026-10-25 selects).

The vacua of the static family (the static blocks of curvature $-\mathsf{y}H^2$ in all eight directions): the left-hand side of the bulk equations is the same in every direction, $\mathsf{P}(\mathsf{y})H^2$. Einstein gravity has the single vacuum $\mathsf{y} = 1$ with $\mathsf{P}'(1) = 21$. The author's original weights would need $\Lambda = -161H^2/16$ (not his $-10H^2$), with $\mathsf{P}'(1) = 21/16$; UM-3's weights make $\mathsf{y} = 1$ a triple root, a degenerate vacuum without linear gravitons. Status: DERIVED (`um4_geometry.py`, G4; `um4_gravity.py`, L1).

$$\frac{\mathsf{P}'(1)}{42}\Big[\mathrm{e}^{2H|\sigma|}\Big(\frac{\partial^2 h_{\mathrm{gw}}}{\partial x_4^2} - \frac{\partial^2 h_{\mathrm{gw}}}{\partial x_3^2}\Big) - \frac{\partial^2 h_{\mathrm{gw}}}{\partial \sigma^2} + 7H\,\mathrm{sgn}(\sigma)\,\frac{\partial h_{\mathrm{gw}}}{\partial \sigma}\Big] = 0, \qquad \mathsf{P}'(1) = 21 - 840w_{(2)} + 7560w_{(3)} \quad \text{(FE-68)}$$

Symbols in (FE-68).

- $\mathsf{P}(\mathsf{y})$: The vacuum polynomial of the static blocks of curvature $-\mathsf{y}H^2$: the left-hand side of the bulk equations divided by H^2, the same in every direction, $\mathsf{P}(\mathsf{y}) = 21\mathsf{y} - 420w_{(2)}\mathsf{y}^2 + 2520w_{(3)}\mathsf{y}^3 + \Lambda/H^2$; $\mathsf{P}'(1)$ its derivative at $\mathsf{y} = 1$, which sets the strength of the graviton around the static block relative to Einstein gravity ($\mathsf{P}'(1)/21$). Unit: dimensionless. Value: $21(\mathsf{y} - 1)$ for Einstein gravity with $\Lambda = -21H^2$. Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- h_{gw}: The amplitude of a gravitational wave of the + polarisation travelling along x_3: the metric components along x_1 and x_2 multiplied by $\mathrm{e}^{h_{\mathrm{gw}}}$ and $\mathrm{e}^{-h_{\mathrm{gw}}}$; a function of σ, x_3 and x_4 in the bulk, and on the fold the gravitational wave of the four-dimensional world, to the first order its plus polarisation h_+. Unit: dimensionless. Status: STANDARD.

- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- $w_{(p)}$: The Lovelock weights of the author's Einstein-Lovelock action, $w_{(1)} = 1$ and $w_{(2)}$, $w_{(3)}$ of the Gauss-Bonnet and the cubic densities. UM-4 sets $w_{(2)} = w_{(3)} = 0$ (Einstein gravity): the only weights for which gravitons propagate on the light cone on every background, not only around the static block. The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) keep the gravitons on the light cone only around the static block (with $\Lambda = -161H^2/16$); UM-3 used $(1, 1/20, 1/360)$. Unit: dimensionless. Value: $(1, 0, 0)$ in UM-4; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), DERIVED (the values that the order of 2026-10-25 selects).

The graviton around the static block of UM-4 for any weights: the + polarisation of a wave along x_3 (the metric components along x_1 and x_2 multiplied by $\mathrm{e}^{h_{\mathrm{gw}}}$ and $\mathrm{e}^{-h_{\mathrm{gw}}}$). The time and ordinary space enter only through $\mathrm{e}^{2H|\sigma|}(\partial^2/\partial x_4^2 - \partial^2/\partial x_3^2)$: at every depth the wave has the light cone of the fold, whatever the weights, with the strength $\mathsf{P}'(1)/21$ relative to Einstein gravity. Around the static block the order therefore does not single out Einstein gravity; FE-69 does. Status: DERIVED (`um4_gravity.py`, L2).

$$
\begin{aligned}
&\mathrm{d}\sigma^2 + \mathrm{e}^{-2\mathfrak{p}H\sigma}\,\mathrm{d}\vec{x}^2 - \mathrm{e}^{-2\mathfrak{q}H\sigma}\,\mathrm{d}x_4^2 - \mathrm{e}^{-2\mathfrak{r}H\sigma}\,\mathrm{d}\vec{x}_5^2 \quad (\sigma > 0):\\
&c_{\mathrm{x}} - c_{\mathrm{t}} = -4\,(\mathfrak{p}-\mathfrak{q})\big(18\mathfrak{r}^2 w_{(3)} - w_{(2)}\big)\big(\mathfrak{p}+\mathfrak{q}+3\mathfrak{r}\big),\\
&c_\sigma - c_{\mathrm{t}} = -4\big[w_{(2)}\big(\mathfrak{p}\mathfrak{q} - \mathfrak{p}^2 + 3\mathfrak{q}\mathfrak{r} - 3\mathfrak{r}^2\big) + 18 w_{(3)}\mathfrak{r}^2\big(\mathfrak{p}^2 - 3\mathfrak{p}\mathfrak{q} + 2\mathfrak{p}\mathfrak{r} - \mathfrak{q}\mathfrak{r} + \mathfrak{r}^2\big)\big]
\end{aligned}
\tag{FE-69}
$$

Symbols in (FE-69).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- $\mathfrak{p}$, $\mathfrak{q}$, $\mathfrak{r}$: The rates of the warps of ordinary space ($\mathfrak{p}$), of the time ($\mathfrak{q}$) and of the extra times ($\mathfrak{r}$) of a static warped background, in units of H. Unit: dimensionless. Value: $\mathfrak{p} = \mathfrak{q} = \mathfrak{r} = 1$ for the static block of UM-4; $\mathfrak{p} = \mathfrak{r} = 1$, $\mathfrak{q} = 0$ for that of UM-3. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- $\vec{x}$: The position in ordinary space, $\vec{x} = (x_1, x_2, x_3)$. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\vec{x}_5$: The position along the extra times, $\vec{x}_5 = (x_5, x_6, x_7)$; $\mathrm{d}\vec{x}_5^2 = \mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2$. Unit: length. Status: AUTHOR.
- c_{x}: The coefficient of the second derivative along the proper length of x_3 in the same equation; the wave moves along x_3 with the local speed $\sqrt{c_{\mathrm{x}}/c_{\mathrm{t}}}$ in units of the local speed of light. Unit: dimensionless. Value: 1 for Einstein gravity on every background; $1 - 24w_{(2)} + 72w_{(3)}$ at the static block of UM-3 ($c_{\mathrm{x}}/c_{\mathrm{t}} = 5$ with the author's original weights). Status: DERIVED.
- c_{t}: The coefficient of the second derivative along the proper time in the linearised equation of a gravitational wave of the + polarisation on a static warped background, normalised so that Einstein gravity gives 1. Unit: dimensionless. Value: 1 for Einstein gravity on every background; $1 - 40w_{(2)} + 360w_{(3)}$ at the static block of UM-3. Status: DERIVED.
- c_σ: The coefficient of the second derivative along σ in the linearised equation of a gravitational wave on a static warped background, normalised so that Einstein gravity gives 1; the wave moves along σ with the local speed $\sqrt{c_\sigma/c_{\mathrm{t}}}$ in units of the local speed of light. Unit: dimensionless. Value: 1 for Einstein gravity on every background. Status: DERIVED.
- $w_{(p)}$: The Lovelock weights of the author's Einstein-Lovelock action, $w_{(1)} = 1$ and $w_{(2)}$, $w_{(3)}$ of the Gauss-Bonnet and the cubic densities. UM-4 sets $w_{(2)} = w_{(3)} = 0$ (Einstein gravity): the only weights for which gravitons propagate on the light cone on every background, not only around the static block. The author's

original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) keep the gravitons on the light cone only around the static block (with $\Lambda = -161H^2/16$); UM-3 used $(1, 1/20, 1/360)$. Unit: dimensionless. Value: $(1, 0, 0)$ in UM-4; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), DERIVED (the values that the order of 2026-10-25 selects).

The gravitons on backgrounds that are not of constant curvature: the static warps of P-1 with any rates. The coefficients c_t, c_x and c_σ of the second derivatives along the proper time, the proper length of x_3 and σ in the linearised equation of the + polarisation are 1 for Einstein gravity on every such background; the Gauss-Bonnet and cubic terms give the deviations shown, which vanish for all rates only for $w_{(2)} = w_{(3)} = 0$. An expanding universe, a star or a gravitational wave of finite amplitude make the warps of the different directions differ, so with the Gauss-Bonnet or the cubic term gravitons would leave the light cone there. At the static block of UM-3 ($\mathfrak{p} = \mathfrak{r} = 1$, $\mathfrak{q} = 0$) the author's original weights give $c_\mathrm{x}/c_\mathrm{t} = 5$: gravitons $\sqrt{5}$ times faster than the local light, which itself was faster than light on the fold. Status: DERIVED (`um4_gravity.py`, L3 and L4).

6.2 The gravitational wave

$$\frac{1}{2}\Big[\mathrm{e}^{2H|\sigma|}\Big(\frac{\partial^2 h_\mathrm{gw}}{\partial x_4^2} - \frac{\partial^2 h_\mathrm{gw}}{\partial x_3^2}\Big) - \frac{\partial^2 h_\mathrm{gw}}{\partial \sigma^2} + 7H\,\mathrm{sgn}(\sigma)\,\frac{\partial h_\mathrm{gw}}{\partial \sigma}\Big] = 0: \qquad h_\mathrm{gw} = h_+(x_3 - x_4),$$
$$\mathsf{h}_{ij} = \begin{pmatrix} h_+ & h_\times & 0 \\ h_\times & -h_+ & 0 \\ 0 & 0 & 0 \end{pmatrix}, \qquad \frac{\Delta L}{L} = \frac{1}{2}\big(h_+ \cos 2\phi_\mathrm{r} + h_\times \sin 2\phi_\mathrm{r}\big), \qquad h_+ \pm \mathrm{i}h_\times \to \mathrm{e}^{\mp 2\mathrm{i}\theta_\mathrm{r}}\big(h_+ \pm \mathrm{i}h_\times\big) \tag{FE-76}$$

Symbols in (FE-76).

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- h_gw: The amplitude of a gravitational wave of the + polarisation travelling along x_3: the metric components along x_1 and x_2 multiplied by $\mathrm{e}^{h_\mathrm{gw}}$ and $\mathrm{e}^{-h_\mathrm{gw}}$; a function of σ, x_3 and x_4 in the bulk, and on the fold the gravitational wave of the four-dimensional world, to the first order its plus polarisation h_+. Unit: dimensionless. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- h_+: The plus polarisation of a gravitational wave travelling along x_3: $\mathsf{h}_{11} = -\mathsf{h}_{22} = h_+$, a function of $x_3 - x_4$; to the first order the zero mode of h_gw. Unit: dimensionless. Status: STANDARD.
- h_{ij}: The transverse traceless perturbation of the metric of ordinary space by a gravitational wave as the four-dimensional world sees it: $\hat{h}_{ij} = \mathrm{e}^{2a_4}(\delta_{ij} + \mathsf{h}_{ij})$ $(i, j = 1, 2, 3)$. Unit: dimensionless. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal), and in a tensor of ordinary space such as h_{ij} one of the directions x_1, x_2, x_3; a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- j: A second generation or mass-state index, or a second direction of ordinary space (as i). Status: STANDARD.
- $h_\times$: The cross polarisation of a gravitational wave travelling along x_3: $\mathsf{h}_{12} = \mathsf{h}_{21} = h_\times$, a function of $x_3 - x_4$. Unit: dimensionless. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.

- L: A distance between two points of the fold along ordinary space: for a gravitational wave the radius of a ring of free test masses (ΔL the change of the distance of one of them from the centre), for a signal the distance between its emitter and its receiver. Unit: length. Status: STANDARD.
- ϕ_r: The angle at which a test mass sits on a ring of test masses in the plane of x_1 and x_2, measured from the x_1 axis. Unit: dimensionless (radian). Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- θ_r: The angle of a rotation about the direction of motion of a wave. Unit: dimensionless (radian). Status: STANDARD.

The gravitational wave in the bulk and on the fold. First line: Einstein's equation of the bulk to the first order in the + polarisation h_gw of a wave along x_3; it is the equation of a massless scalar of the bulk (FE-73), and its zero mode, independent of σ, is any function of $x_3 - x_4$: it solves the bulk equation on both sides and the junction condition at the fold, so the graviton of the bulk has a massless mode bound to the fold that travels at the speed of light at every wavelength and at every depth. Second line: the wave as the four-dimensional world sees it, the transverse traceless perturbation h_{ij} of the metric of ordinary space, with the two polarisations h_+ (the "plus" mode, $\mathsf{h}_{11} = -\mathsf{h}_{22}$) and $h_\times$ (the "cross" mode, $\mathsf{h}_{12} = \mathsf{h}_{21}$). A ring of free test masses of the radius L in the plane of x_1 and x_2 changes the distance of the mass at the angle ϕ_r from its centre by ΔL: the plus mode stretches the ring along x_1 while it squeezes it along x_2 and then the reverse, the cross mode does the same along the two diagonals, rotated by 45°: they stretch and squeeze space perpendicularly to each other and to the direction of motion. Under a rotation by the angle θ_r about the direction of motion the circular combinations $h_+ \pm \mathrm{i}h_\times$ are multiplied by $\mathrm{e}^{\mp 2\mathrm{i}\theta_\mathrm{r}}$: helicity ± 2, a wave of spin 2, as it must be for a wave sourced by the rank-2 energy-momentum tensor. The massless graviton of seven dimensions has 14 polarisations, which the four-dimensional world sees as the graviton (2), three vectors (6) and six scalars (6); matter of the fold that does not depend on the extra times and is the same along the three of them emits only the plus and cross polarisations and the volume of the extra times, which the fold fluid holds fixed: the gravitational waves of UM-4 have only the plus and the cross polarisations, as the order requires and as observed. Status: DERIVED (`um4_bulk.py`, B4 and B5; `um4_foldsources.py`, FS2), ASSUMED (that the fold fluid holds the volume of the extra times fixed for waves as well).

$$\frac{1}{2}\Big(\frac{\partial^2 h_\mathrm{gw}}{\partial x_4^2} + 3a_4'\,\frac{\partial h_\mathrm{gw}}{\partial x_4} - \mathrm{e}^{-2a_4}\,\frac{\partial^2 h_\mathrm{gw}}{\partial x_3^2}\Big) = \kappa_7\,\Pi_\mathrm{gw}, \qquad \kappa_7 = \frac{5}{2}\,\kappa H, \qquad c_\mathrm{g} = 1 \tag{FE-63}$$

Symbols in (FE-63).

- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- h_gw: The amplitude of a gravitational wave of the + polarisation travelling along x_3: the metric components along x_1 and x_2 multiplied by $\mathrm{e}^{h_\mathrm{gw}}$ and $\mathrm{e}^{-h_\mathrm{gw}}$; a function of σ, x_3 and x_4 in the bulk, and on the fold the gravitational wave of the four-dimensional world, to the first order its plus polarisation h_+. Unit: dimensionless. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- κ_7: The gravitational coupling of the fold: to the first order in its curvature over H^2 the fold obeys the Einstein equations of seven dimensions $\hat{G}^a{}_b = \kappa_7 \mathsf{t}^a{}_b$ with $\kappa_7 = 5\kappa H/2$, the coupling of the massless graviton of the bulk bound to the fold; four-dimensional gravity on the fold has $8\pi G_\mathrm{N} = \kappa_7/V_*$ (plus the induced term, which is negligible). Unit: GeV^{-5}. Value: $5\kappa H/2$. Status: DERIVED.
- Π_gw: The transverse traceless stress of the matter of the fold that drives the gravitational wave h_gw (half the difference of its components along x_1 and x_2), integrated across the fold (eight-dimensional normalisation). Unit: GeV^7. Status: STANDARD.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free

parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.

- $c_{\rm g}$: The speed of gravitational waves on the fold, in units of the speed of light. Unit: dimensionless. Value: 1 at every wavelength for the four-dimensional graviton on the static block and on every fold with frozen extra times, and for the massless graviton of the bulk at every depth; measured $-3\times10^{-15} \le c_{\rm g}-1 \le 7\times10^{-16}$ (GW170817). Status: DERIVED.

The gravitational wave on the expanding fold (frozen extra times): the wave equation of general relativity on the metric of the four-dimensional universe, sourced by the transverse traceless stress of the matter of the fold with the coupling κ_7 of the Friedmann equations. It travels at the speed of light at every wavelength ($c_{\rm g}=1$ identically), which the simultaneous arrival of the gravitational wave GW170817 and of the light of its source confirms to 10^{-15}. Status: DERIVED (`um4_foldsources.py`, FS2), STANDARD (the measurement).

$$\kappa\,\mathcal{S}^a{}_b = -12H\,\delta^a{}_b + \frac{2}{5H}\,\hat{G}^a{}_b + \ldots, \qquad \mathcal{S}^a{}_b = -\lambda_{\rm F}\,\delta^a{}_b + \mathtt{t}^a{}_b: \qquad \hat{G}^a{}_b = \kappa_7\,\mathtt{t}^a{}_b, \qquad \kappa_7 = \frac{5}{2}\,\kappa H \qquad \text{(FE-71)}$$

Symbols in (FE-71).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} {\rm d}\sigma\,\mathcal{E}_{(1)}{}^a{}_b$ (the junction condition of Israel); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -12H\delta^a{}_b$ for every a_4, b_4 of the author's form. Status: DERIVED.
- a: A frame (orthonormal) index running over $0,1,\ldots,7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = {\rm diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a,b \in \{1,\ldots,7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1,\ldots,7\}$; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor ${\rm e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\rm F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\rm L}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1M_2\cdots}_{N_1N_2\cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- $\hat{G}^\mu{}_\nu$: The Einstein tensor of the metric $\hat{g}_{ab}$ of the fold (FD-2), with frame indices along its seven directions; with four-dimensional indices, $\hat{G}^\mu{}_\nu$, its components along the four observable directions, which for frozen extra times are those of the Einstein tensor of the four-dimensional metric $\hat{h}_{\mu\nu}$. Unit: 1/length^2. Status: STANDARD.
- $\lambda_{\rm F}$: The tension that the static block requires of the fold, $\kappa\lambda_{\rm F} = 12H$: the total vacuum energy of the seven-dimensional fold, made of the tension of the fold fluid and the vacuum energy of the fields of the fold (both have the pressure minus their energy density along ordinary space and along the extra times), per unit volume of the six other directions of the fold (eight-dimensional normalisation); in the four-dimensional normalisation it is $\lambda_{\rm F}V_*$, and the excess of the total over $\lambda_{\rm F}V_*$ is the dark energy ρ_Λ. Unit: GeV^7. Value: $12H/\kappa$; $\lambda_{\rm F}V_* = 30H^2M_{\rm red}^2 = 1.1\times10^{45}$ GeV^4 for $H = 2.5$ TeV. Status: DERIVED.
- $\mathtt{t}^a{}_b$, $\mathtt{t}$: The matter of the fold above its tension: the thin source minus the tension, $\mathcal{S}^a{}_b = -\lambda_{\rm F}\delta^a{}_b + \mathtt{t}^a{}_b$ (eight-dimensional normalisation); $\mathtt{t} = \mathtt{t}^c{}_c$ its trace; with frozen extra times $\mathtt{t}^a{}_b = {\rm diag}(P_7, P_7, P_7, -\rho_7, q_7, q_7, q_7)$. Unit: GeV^7. Status: DERIVED.
- κ_7: The gravitational coupling of the fold: to the first order in its curvature over H^2 the fold obeys the Einstein equations of seven dimensions $\hat{G}^a{}_b = \kappa_7\mathtt{t}^a{}_b$ with $\kappa_7 = 5\kappa H/2$, the coupling of the massless graviton of the bulk bound to the fold; four-dimensional gravity on the fold has $8\pi G_{\rm N} = \kappa_7/V_*$ (plus the induced term, which is negligible). Unit: GeV^{-5}. Value: $5\kappa H/2$. Status: DERIVED.

The gravity of the fold: the junction condition, to the first order in the curvature of the fold divided by H^2, is the pure tension plus the Einstein tensor of seven dimensions; the thin source is the tension plus the matter of

the fold, which therefore obeys the Einstein equations of seven dimensions with $\kappa_7 = 5\kappa H/2$, the coupling of the massless graviton of the bulk bound to the fold. With frozen extra times they are the Friedmann equations of general relativity with Newton's constant $8\pi G_{\mathrm{N}} = \kappa_7/V_*$ (FE-44), up to corrections of the relative size $(a_4'/H)^2$, about 3×10^{-91} today for $H = 2.5$ TeV. Status: DERIVED (`um4_fold.py`, F1, F2, F4 and F5).

7 Photons at the speed of light: transverse waves of spin 1

$$
\begin{aligned}
&S_{\mathrm{gauge}} = \int \mathrm{d}^8X\, \sqrt{|g|}\, \mathcal{L}_{\mathrm{gauge}}, \qquad \mathcal{L}_{\mathrm{gauge}} = -\frac{1}{4\mathtt{g}_3^2}\, G^A_{MN} G^{AMN} - \frac{1}{4\mathtt{g}_2^2}\, W^I_{MN} W^{IMN} - \frac{1}{4\mathtt{g}_1^2}\, B_{MN} B^{MN}, \\
&G^A_{MN} = \partial_M G^A_N - \partial_N G^A_M + f^{ABC} G^B_M G^C_N, \qquad W^I_{MN} = \partial_M W^I_N - \partial_N W^I_M + \epsilon^{IJK} W^J_M W^K_N, \\
&B_{MN} = \partial_M B_N - \partial_N B_M
\end{aligned}
\tag{IN-7}
$$

Symbols in (IN-7).

- S_{gauge}: The action of the gauge fields in the bulk. Unit: dimensionless. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\, \mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\, \mathrm{d}X^M \mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\, \mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- $\mathcal{L}_{\mathrm{gauge}}$: The Lagrangian of the gauge fields of $SU(3)_C \times SU(2)_L \times U(1)_Y$ in the bulk. Unit: GeV^8. Status: STANDARD.
- $\mathtt{g}_1$, $\mathtt{g}_2$, $\mathtt{g}_3$: The gauge couplings of eight dimensions of $U(1)_Y$, $SU(2)_L$ and $SU(3)_C$: the coefficients of the kinetic terms of the gauge fields of the bulk in their geometric normalisation; the four-dimensional couplings are $g_k^2 = 3H\mathtt{g}_k^2/(2V_*)$. Unit: GeV^{-2} ($\mathtt{g}^2$: length^4). Status: ASSUMED.
- G^A_M, G^A_μ, G^A_{MN}: The gluon field of $SU(3)_C$ ($A = 1, \ldots, 8$): in the bulk G^A_M, a field of eight dimensions in its geometric normalisation (the covariant derivative $\partial - \mathrm{i}G$), and on the fold the canonically normalised four-dimensional field $G^A_\mu(x)$ of its massless mode, $g_3 G^A_\mu(x) = G^A_\mu(\sigma = 0, x)$; G^A_{MN} and $G^A_{\mu\nu}$ their field strengths. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- W^I_M, W^I_μ, W^I_{MN}: The gauge field of $SU(2)_L$ ($I = 1, 2, 3$): in the bulk W^I_M (geometric normalisation), on the fold the canonically normalised four-dimensional field $W^I_\mu(x)$ of its massless mode, $g_2 W^I_\mu(x) = W^I_\mu(\sigma = 0, x)$; W^I_{MN}, $W^I_{\mu\nu}$ their field strengths; $W^\pm_\mu = (W^1_\mu \mp \mathrm{i}W^2_\mu)/\sqrt{2}$. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- B_M, B_μ, B_{MN}: The hypercharge gauge field of $U(1)_Y$: in the bulk B_M (geometric normalisation), on the fold the canonically normalised four-dimensional field $B_\mu(x)$ of its massless mode, $g_1 B_\mu(x) = B_\mu(\sigma = 0, x)$; B_{MN}, $B_{\mu\nu}$ their field strengths. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- f^{ABC}: The structure constants of $SU(3)$, $[t^A, t^B] = \mathrm{i}f^{ABC}t^C$. Unit: dimensionless. Status: STANDARD.
- B: A colour (adjoint SU(3)) index (as A). Status: STANDARD.
- C: A colour (adjoint SU(3)) index (as A). Status: STANDARD.
- ϵ^{IJK}: The Levi-Civita symbol of three indices (the structure constants of $SU(2)$), $\epsilon^{123} = 1$. Unit: dimensionless. Status: STANDARD.
- J: A weak-isospin (adjoint SU(2)) index (as I). Status: STANDARD.

- K: A weak-isospin (adjoint SU(2)) index (as I). Status: STANDARD.

The gauge fields of the Standard Model in the bulk of eight dimensions (the order puts the photon in the bulk, and with it the electroweak gauge fields of which it is a combination; UM-4 puts the gluons there as well). Status: AUTHOR (the photon in the bulk, by the order), ASSUMED (all gauge fields in the bulk), STANDARD (the Yang-Mills action).

$$\begin{aligned} &B_\sigma = 0, \quad B_a = u(\sigma)\,\hat{B}_a(x): \qquad u'' - 5H\,\mathrm{sgn}(\sigma)\,u' + \mathrm{e}^{2H|\sigma|}\,m^2\,u = 0, \qquad \eta^{ac}\,\partial_a \hat{B}_{cb} = m^2\,\hat{B}_b, \\ &u_0 = 1, \quad \int_{-\infty}^{\infty} \mathrm{d}\sigma\,\mathrm{e}^{-3H|\sigma|}\,u_0^2 = \frac{2}{3H} \end{aligned} \tag{FE-74}$$

Symbols in (FE-74).

- B_M, B_μ, B_{MN}: The hypercharge gauge field of $U(1)_Y$: in the bulk B_M (geometric normalisation), on the fold the canonically normalised four-dimensional field $B_\mu(x)$ of its massless mode, $g_1 B_\mu(x) = B_\mu(\sigma = 0, x)$; B_{MN}, $B_{\mu\nu}$ their field strengths. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat{g}_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- $u(\sigma)$, u_0: The profile along the hidden direction of a mode of a field of the bulk (a scalar, a gauge field or a polarisation of the graviton): the field is $u(\sigma)$ times a field of the fold; $u_0 = 1$ is the profile of the massless mode (the zero mode); in a propagator the response of the field to a unit source on the fold. Unit: dimensionless (a relative amplitude); length in a propagator. Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\hat{B}_a$, $\hat{B}_{ab}$: The field of the fold of a mode of the hypercharge field of the bulk (of any gauge field of the bulk to the first order in the field), $B_a = u(\sigma)\hat{B}_a(x)$; $\hat{B}_{ab} = \partial_a \hat{B}_b - \partial_b \hat{B}_a$ its field strength. Unit: GeV (field), GeV^2 (field strength). Status: STANDARD.
- x: The coordinates of the fold: x_1, x_2, x_3 of ordinary space, x_4 the time, x_5, x_6, x_7 the extra times; a field of the fold is a function of the four observable coordinates $x = (x_1, x_2, x_3, x_4)$, and $\mathrm{d}^4x = \mathrm{d}x_1\,\mathrm{d}x_2\,\mathrm{d}x_3\,\mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m^2_{ij} = m_i^2 - m_j^2$ the squared-mass difference of the states i and j (Δm^2_{21}, Δm^2_{31}, Δm^2_{32}); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by v^2_loc/v^2 where this differs from 1; m without a label the mass of a mode of a field of the bulk as the fold sees it, $m^2 = \omega^2 - k^2$; for the W and Z bosons their complex pole mass, $m^2 = m_0^2 - \mathrm{i}m_0\Gamma_\mathrm{bulk}$ to the leading order; in the relic abundance of the particles χ their mass. Unit: GeV. Status: STANDARD.
- η_{ab}: The flat metric of the frame, $\eta = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$ (eta4488 of 4+4.nb); η^{ab} is its inverse (the same matrix). Unit: dimensionless. Status: AUTHOR.
- c: A frame index (as a). Status: STANDARD.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.

The modes of a gauge field of the bulk, written for the hypercharge field, in the gauge $B_\sigma = 0$: the field of the fold $\hat{B}_a(x)$ obeys the Proca equation of the fold with the mass m, and the profile the equation in the middle. The massless mode has the profile 1 and the kinetic weight $\mathrm{e}^{-3H|\sigma|}$ with the finite integral $2/(3H)$: a massless gauge field of the fold, bound to it, at the speed of light at every wavelength. The massive modes form a continuum from zero mass upwards and move slower than light. A component along the hidden direction has no normalisable zero mode. (In UM-3 the bulk photon had $\omega^2 = 2k^2$, P-2.) Status: DERIVED (`um4_bulk.py`, B2: the profile equation exactly on both halves; the lowest ω^2 in a box equals k^2 to 4×10^{-12}; B10: the continuum of massive modes from zero mass, Bessel functions of the warped distance from the fold).

$$\begin{aligned} &A_\mu = \mathfrak{e}_\mu\, \mathrm{e}^{\mathrm{i}k(x_3 - x_4)}: \qquad \mathfrak{e}_3 + \mathfrak{e}_4 = 0, \qquad \mathfrak{e}_\mu \sim \mathfrak{e}_\mu + \beta_\mathrm{g}\, k\, (0,0,1,-1), \\ &\mathfrak{e}_\pm = \frac{1}{\sqrt{2}}\big(1, \pm\mathrm{i}, 0, 0\big) \to \mathrm{e}^{\mp \mathrm{i}\theta_\mathrm{r}}\, \mathfrak{e}_\pm, \qquad \vec{E}\cdot\vec{B} = 0, \quad \vec{E}^2 = \vec{B}^2 \end{aligned} \tag{FE-75}$$

Symbols in (FE-75).

- A_μ: The photon field of the fold (canonically normalised): the massless combination of the massless modes of W^1, W^2, W^3 and B that the Higgs vacuum leaves massless; a field of the bulk whose zero mode travels at the speed of light. Unit: GeV. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- $\mathfrak{e}_\mu$, $\mathfrak{e}_\pm$: The polarisation vector of a plane wave of the photon, with components along x_1, x_2, x_3, x_4; $\mathfrak{e}_\pm = (1, \pm\mathrm{i}, 0, 0)/\sqrt{2}$ the circular polarisations of a wave travelling along x_3. Unit: dimensionless. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- k: A wave number: the momentum of a plane wave along x_3, $\mathrm{e}^{\mathrm{i}(kx_3 - \omega x_4)}$; in the cosmological perturbations the comoving wave number of a Fourier mode, $k = |\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- β_g: Any constant: the size of a gauge transformation, which adds a multiple of the wave vector to the polarisation vector of the photon. Unit: dimensionless. Status: STANDARD.
- θ_r: The angle of a rotation about the direction of motion of a wave. Unit: dimensionless (radian). Status: STANDARD.
- $\vec{E}$: The electric field of the photon on the fold: its component along the direction x_i of ordinary space is $\partial A_4/\partial x_i - \partial A_i/\partial x_4$ ($i = 1,2,3$; A_4 and A_i the components of A_μ); $\vec{E}^2$ its square. Unit: GeV^2. Status: STANDARD.
- $\vec{B}$: The magnetic field of the photon on the fold, the curl of the components A_1, A_2, A_3 of A_μ along ordinary space (its component along x_1 is $\partial A_3/\partial x_2 - \partial A_2/\partial x_3$, and cyclically); $\vec{B}^2$ its square. Unit: GeV^2. Status: STANDARD.

The photon as a wave: the zero mode for the combination of W^3 and B that the Higgs vacuum leaves massless, a plane wave along x_3 with the polarisation vector $\mathfrak{e}_\mu$. Its wave vector is null, so it satisfies Maxwell's equations of the fold. The Lorenz condition requires $\mathfrak{e}_3 + \mathfrak{e}_4 = 0$, and a gauge transformation removes the remaining component along the direction of motion: the only physical polarisations are the two transverse ones, along x_1 and along x_2. Its electric and magnetic fields are orthogonal to the direction of motion and to each other and equal in size. Under a rotation by θ_r about the direction of motion the circular polarisations are multiplied by $\mathrm{e}^{\mp\mathrm{i}\theta_\mathrm{r}}$: helicity ± 1, a transverse wave of spin 1, as the order requires, at the speed of light. A photon that leaves the fold travels through the bulk at the local speed of light, which at every depth is that of the fold (FE-83). The three polarisations of the massless vector of seven dimensions that point along the extra times are excluded (their kinetic terms would have the wrong sign). Status: DERIVED (`um4_bulk.py`, B3), ASSUMED (no components along the extra times).

$$g_1^2=\frac{3H\mathsf{g}_1^2}{2V_*},\qquad g_2^2=\frac{3H\mathsf{g}_2^2}{2V_*},\qquad g_3^2=\frac{3H\mathsf{g}_3^2}{2V_*},\qquad 8\pi G_{\rm N}=\frac{\kappa_7}{V_*},$$
$$g_{1,\rm eff}=\frac{g_1}{\sqrt{\mathcal{Z}}},\quad g_{2,\rm eff}=\frac{g_2}{\sqrt{\mathcal{Z}}},\quad g_{3,\rm eff}=\frac{g_3}{\sqrt{\mathcal{Z}}},\qquad \lambda_{\rm eff}=\frac{\lambda}{\mathcal{Z}},\qquad \mathcal{Y}_{\rm eff}=\frac{\mathcal{Y}}{\sqrt{\mathcal{Z}}},\qquad G_{\rm N,eff}=\frac{G_{\rm N}}{\mathcal{Z}},\qquad \text{(FE-44)}$$
$$\mathcal{Z}=\frac{V_{\rm t}\left(\mathrm{e}^{-a_4}b_4\right)^3}{V_*}$$

Symbols in (FE-44).

- g_1: The $U(1)_Y$ gauge coupling of the four-dimensional world (usually g'; not normalised for grand unification): the coupling of the massless mode of the hypercharge field of the bulk, $g_1^2=3H\mathsf{g}_1^2/(2V_*)$. Unit: dimensionless. Value: $g_1(m_Z)=0.3574$. Status: STANDARD.
- g_2: The $SU(2)_L$ gauge coupling of the four-dimensional world (usually g): the coupling of the massless mode of the $SU(2)_L$ field of the bulk, $g_2^2=3H\mathsf{g}_2^2/(2V_*)$. Unit: dimensionless. Value: $g_2(m_Z)=0.6517$. Status: STANDARD.
- g_3: The $SU(3)_C$ gauge coupling of the four-dimensional world (the strong coupling g_s): the coupling of the massless mode of the gluon field of the bulk, $g_3^2=3H\mathsf{g}_3^2/(2V_*)$. Unit: dimensionless. Value: $g_3(m_Z)=1.2177$ ($g_3^2/(4\pi)=0.1180$). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda=-21H^2$ and the tension $\kappa\lambda_{\rm F}=12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H\gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- g_1, g_2, g_3: The gauge couplings of eight dimensions of $U(1)_Y$, $SU(2)_L$ and $SU(3)_C$: the coefficients of the kinetic terms of the gauge fields of the bulk in their geometric normalisation; the four-dimensional couplings are $g_k^2=3H\mathsf{g}_k^2/(2V_*)$. Unit: GeV^{-2} (g^2: length^4). Status: ASSUMED.
- V_*: The physical volume of the extra times x_5,x_6,x_7 on the fold in the late universe, $V_*=V_{\rm t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\rm t}=\int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $G_{\rm N}$: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_{\rm N}=1/M_{\rm Pl}^2$ with the Planck mass $M_{\rm Pl}=1.221\times10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- κ_7: The gravitational coupling of the fold: to the first order in its curvature over H^2 the fold obeys the Einstein equations of seven dimensions $\hat{G}^a{}_b=\kappa_7\mathsf{t}^a{}_b$ with $\kappa_7=5\kappa H/2$, the coupling of the massless graviton of the bulk bound to the fold; four-dimensional gravity on the fold has $8\pi G_{\rm N}=\kappa_7/V_*$ (plus the induced term, which is negligible). Unit: GeV^{-5}. Value: $5\kappa H/2$. Status: DERIVED.
- $g_{k,\rm eff}$: The effective (canonically normalised) gauge couplings of the fold when the fold factor is $\mathcal{Z}$, $g_{k,\rm eff}=g_k/\sqrt{\mathcal{Z}}$ ($k=1,2,3$): the kinetic terms of the massless modes of the gauge fields of the bulk carry the volume of the extra times as the terms of the fold do. Unit: dimensionless. Status: DERIVED.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4)=V_{\rm t}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold and the kinetic terms of the massless modes of the fields of the bulk; $\mathcal{Z}=1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- $\lambda_{\rm eff}$: The effective quartic Higgs coupling, $\lambda/\mathcal{Z}$. Unit: dimensionless. Status: DERIVED.
- λ: The quartic coupling of the Higgs potential, $\lambda>0$. Unit: dimensionless. Value: $\lambda=m_h^2/(2v^2)=0.129$. Status: AUTHOR.
- $\mathcal{Y}_{\rm eff}$: The effective Yukawa matrices, $\mathcal{Y}/\sqrt{\mathcal{Z}}$ (for each of $\mathcal{Y}^u,\mathcal{Y}^d,\mathcal{Y}^e,\mathcal{Y}^\nu$). Unit: dimensionless. Status: DERIVED.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^\nu$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- $G_{\rm N,eff}$: The effective Newton constant of the fold, $G_{\rm N}/\mathcal{Z}$. Unit: GeV^{-2}. Status: DERIVED.
- $V_{\rm t}$: The coordinate volume of the (compact) extra times, $V_{\rm t}=\int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$. Unit: length^3. Status: ASSUMED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.

- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\, \mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.

The four-dimensional couplings: the gauge couplings of the Standard Model are those of the massless modes, $g_k^2 = 3H\mathrm{g}_k^2/(2V_*)$ ($k = 1, 2, 3$), and Newton's constant is that of the massless graviton; the couplings measured today follow when the volume of the extra times is frozen ($\mathcal{Z} = 1$). Status: DERIVED (`um4_bulk.py`, B2; `um4_fold.py`, F2).

8 The other fields of the bulk

$$\phi_{\mathrm{bulk}} = u(\sigma)\, \mathrm{e}^{\mathrm{i}(kx_3 - \omega x_4)} : \qquad u'' - 7H\, \mathrm{sgn}(\sigma)\, u' + \mathrm{e}^{2H|\sigma|}\, m^2\, u = 0, \qquad m^2 = \omega^2 - k^2,$$
$$u_0 = 1, \quad \int_{-\infty}^{\infty} \mathrm{d}\sigma\, \mathrm{e}^{-5H|\sigma|}\, u_0^2 = \frac{2}{5H} \tag{FE-73}$$

Symbols in (FE-73).

- ϕ_{bulk}: A massless scalar field of the bulk, $\nabla_M \nabla^M \phi_{\mathrm{bulk}} = 0$ (an illustration: each polarisation of the graviton obeys its equation). Unit: GeV^3. Status: STANDARD.
- $u(\sigma)$, u_0: The profile along the hidden direction of a mode of a field of the bulk (a scalar, a gauge field or a polarisation of the graviton): the field is $u(\sigma)$ times a field of the fold; $u_0 = 1$ is the profile of the massless mode (the zero mode); in a propagator the response of the field to a unit source on the fold. Unit: dimensionless (a relative amplitude); length in a propagator. Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\, \mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z) \ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- k: A wave number: the momentum of a plane wave along x_3, $\mathrm{e}^{\mathrm{i}(kx_3 - \omega x_4)}$; in the cosmological perturbations the comoving wave number of a Fourier mode, $k = |\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m_{ij}^2 = m_i^2 - m_j^2$ the squared-mass difference of the states i and j (Δm_{21}^2, Δm_{31}^2, Δm_{32}^2); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by v_{loc}^2/v^2 where this differs from 1; m without a label the mass of a mode of a field of the bulk as the fold sees it, $m^2 = \omega^2 - k^2$; for the W and Z bosons their complex pole mass, $m^2 = m_0^2 - \mathrm{i}m_0\Gamma_{\mathrm{bulk}}$ to the leading order; in the relic abundance of the particles χ their mass. Unit: GeV. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.

A massless scalar of the bulk (an illustration: each polarisation of the graviton obeys this equation): ω and k enter only through $m^2 = \omega^2 - k^2$, so every mode is a field of the fold with the Lorentz-invariant dispersion $\omega^2 = k^2 + m^2$; the zero mode has the profile 1 and the kinetic weight $\mathrm{e}^{-5H|\sigma|}$ with the integral $2/(5H)$ and travels at the speed of light; the massive modes are slower. Status: DERIVED (`um4_bulk.py`, B1).

$$\Psi_{\rm k} = {\rm e}^{\frac{7}{2}H|\sigma|}\,{\rm e}^{-M_{\rm k}|\sigma|}\,\chi_0(x), \qquad \gamma^{(0)}\chi_0 = -\chi_0, \qquad \gamma^{(a)}\,\partial_a\chi_0 = 0 \quad \text{(along the fold)},$$
$$\sqrt{|g|}\,\bar\Psi_{\rm k}\gamma^{x_4}\Psi_{\rm k} \propto {\rm e}^{(H-2M_{\rm k})|\sigma|} \tag{FE-77}$$

Symbols in (FE-77).

- $\Psi_{\rm k}$, $\bar\Psi_{\rm k}$: A fermion field of the bulk with the kink mass $M_{\rm k}\,\mathrm{sgn}(\sigma)$ (an illustration, not a field of UM-4). Unit: $\mathrm{GeV}^{7/2}$. Status: ASSUMED.
- e: Euler's number, the base of the natural logarithm; ${\rm e}^X$ is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor ${\rm e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\rm F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $M_{\rm k}$: The mass of a fermion of the bulk that changes sign at the fold (a kink mass $M_{\rm k}\,\mathrm{sgn}(\sigma)$), used to illustrate the speeds of bulk fermions. Unit: GeV. Status: ASSUMED.
- χ_0: The spinor of the fold of the massless mode of a fermion of the bulk: a solution of the massless Dirac equation of the fold of one chirality along σ. Unit: $\mathrm{GeV}^{7/2}$. Status: DERIVED.
- x: The coordinates of the fold: x_1, x_2, x_3 of ordinary space, x_4 the time, x_5, x_6, x_7 the extra times; a field of the fold is a function of the four observable coordinates $x = (x_1, x_2, x_3, x_4)$, and $\mathrm{d}^4x = \mathrm{d}x_1\,\mathrm{d}x_2\,\mathrm{d}x_3\,\mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; in the tensors of the fold written with its coordinates x^a (its metric $\hat g_{ab}$ and the tensors built from it) a coordinate index of the fold, $a \in \{1, \ldots, 7\}$; a repeated index is summed. Status: STANDARD.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.

A fermion of the bulk with a mass that changes sign at the fold (an illustration: UM-4 has no such field; it shows that quarks and leptons could live in the bulk without outrunning light): its zero mode is a massless fermion of seven dimensions bound to the fold for $M_{\rm k} > H/2$, at the speed of light (in UM-3 it moved at $2M_{\rm k}/(2M_{\rm k} - H)$ times the speed of light). Status: DERIVED (`um4_bulk.py`, B6).

$$\gamma^M\Omega_M = -\frac{7}{2}\,H\,\mathrm{sgn}(\sigma)\,\gamma^{(0)} + \frac{3}{2}\,\frac{b_4'}{b_4}\,{\rm e}^{H|\sigma|}\,\gamma^{(4)} \neq 0 \tag{FE-14}$$

Symbols in (FE-14).

- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Ω_M: The spin connection acting on dirac16complex, $\Omega_M = \frac{1}{2}\varpi_{Mab}\Sigma^{ab}$. Unit: 1/length. Status: RECORD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor ${\rm e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\rm F} = 12H$ of the fold; in UM-4 a free

parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

The canonical spin connection of dirac16complex (the author's field of eight dimensions) for the metric of UM-4, with the author's matrices: its first term is $-\frac{7}{2}H\,\mathrm{sgn}(\sigma)\gamma^{(0)}$ (UM-3: $-3H\,\mathrm{sgn}(\sigma)\gamma^{(0)}$), and it is never identically zero. Status: DERIVED (`um4_geometry.py`, G9).

$$\Psi = \mathrm{e}^{\frac{7}{2}H|\sigma|}\,\chi(\sigma)\,\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x}-\omega x_4)}, \qquad \gamma^{(0)}\chi' + \mathrm{e}^{H|\sigma|}\big(-\mathrm{i}\omega\,\gamma^{(4)} + \mathrm{i}\,\vec{k}\cdot\vec{\gamma}\big)\chi - m_\Psi\chi = 0 \qquad (\sigma \neq 0) \tag{FE-36}$$

Symbols in (FE-36).

- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{\frac{7}{2}H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x}-\omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of the level of dirac16complex at the fold (a long-lived resonance) and $\bar\chi$ its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\vec{k}$: The (comoving) momentum along ordinary space, $\vec{k}^2$ its square. Unit: GeV. Status: STANDARD.
- $\vec{x}$: The position in ordinary space, $\vec{x} = (x_1, x_2, x_3)$. Unit: length. Status: AUTHOR.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\vec{\gamma}$: The three frame Dirac matrices of dirac16complex along ordinary space, $\vec{\gamma} = (\gamma^{(1)}, \gamma^{(2)}, \gamma^{(3)})$. Unit: dimensionless. Status: AUTHOR.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.

A mode of dirac16complex on the static block: the factor $\mathrm{e}^{(7/2)H|\sigma|}$ removes the first term of the spin connection, and the factor $\mathrm{e}^{H|\sigma|}$ multiplies the whole Dirac operator of the fold, the frequency as well as the momentum (in UM-3 only the momentum). Status: DERIVED (`um4_geometry.py`, G9).

$$\mathsf{S}=\frac{1}{2}\,\gamma^{(3)}\gamma^{(4)}:\qquad \big[\mathsf{S},\gamma^{(0)}\big]=\big[\mathsf{S},\Gamma_8\big]=0,\qquad \big[\mathsf{S},\gamma^{(3)}\big]=-\gamma^{(4)},\qquad \big[\mathsf{S},\gamma^{(4)}\big]=-\gamma^{(3)}\qquad\Rightarrow\qquad \omega^2=\omega_0^2+k^2 \tag{FE-78}$$

Symbols in (FE-78).

- S: The generator of the boosts along x_3 on dirac16complex, $\mathsf{S}=\frac{1}{2}\gamma^{(3)}\gamma^{(4)}$. Unit: dimensionless. Status: DERIVED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M=\mathbf{e}_a{}^M T^a$, $\gamma_M=g_{MN}\gamma^N$; $\gamma^{(a)}=T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- Γ_8: The chirality matrix of eight dimensions, $\Gamma_8=T_0T_1\cdots T_7$ (the author's T16^A[8]) $=\mathrm{diag}(-\mathbf{1}_8,\mathbf{1}_8)$; $\Gamma_8^2=\mathbf{1}$; it anticommutes with every T_a. Unit: dimensionless. Status: AUTHOR.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- ω_0: The energy at rest (zero momentum along ordinary space and along the extra times) of a state of dirac16complex at the fold; complex for a resonance, $\omega_0=\omega_\mathrm{r}-\frac{\mathrm{i}}{2}\Gamma_\chi$. Unit: GeV. Status: DERIVED.
- k: A wave number: the momentum of a plane wave along x_3, $\mathrm{e}^{\mathrm{i}(kx_3-\omega x_4)}$; in the cosmological perturbations the comoving wave number of a Fourier mode, $k=|\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.

dirac16complex is invariant under the boosts of the fold: the generator S of the boosts along x_3, built from the author's matrices, commutes with $\gamma^{(0)}$, with the chirality and with the mass term and rotates $\gamma^{(3)}$ and $\gamma^{(4)}$ into each other hyperbolically, and the warp multiplies the frequency and the momentum by the same factor. Every state of dirac16complex at the fold with the energy ω_0 at rest therefore has $\omega^2=\omega_0^2+k^2$ exactly and moves slower than light (in UM-3 the bound states had the speed coefficient $c_\mathrm{b}^2=1.19$ along ordinary space). Status: DERIVED (`um4_bulk.py`, B7, with the author's matrices).

$$\begin{aligned}&\chi'=\mathbb{K}\,\chi,\qquad \mathbb{K}=\gamma^{(0)}\big(m_\Psi+\mathrm{i}\omega\,\mathrm{e}^{H|\sigma|}\gamma^{(4)}\big),\qquad \mathbb{K}^2=\big(m_\Psi^2-\omega^2\mathrm{e}^{2H|\sigma|}\big)\mathbf{1},\\&q=\sqrt{m_\Psi^2-\omega^2\mathrm{e}^{2H|\sigma|}},\qquad |\omega|\,\mathrm{e}^{H\sigma_\mathrm{t}}=m_\Psi\end{aligned} \tag{FE-37}$$

Symbols in (FE-37).

- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi=\mathrm{e}^{\frac{7}{2}H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x}-\omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of the level of dirac16complex at the fold (a long-lived resonance) and $\bar{\chi}$ its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4'=\mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\mathbb{K}$: The transfer matrix of the profile equation at zero momentum, $\chi'=\mathbb{K}\chi$. Unit: GeV. Status: DERIVED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M=\mathbf{e}_a{}^M T^a$, $\gamma_M=g_{MN}\gamma^N$; $\gamma^{(a)}=T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda=-21H^2$ and the tension $\kappa\lambda_\mathrm{F}=12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H\gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma=|\cot z|\,\mathrm{d}x_0$, $\sigma=\mathrm{sgn}(\pi/2-z)\ln(\sin z)/(6H)$ on one period $0<z<\pi$; $\sigma=0$ is the fold $z=\pi/2$, $\sigma\to\mp\infty$ the tips $z\to 0,\pi$; one period of x_0 is the whole line $-\infty<\sigma<\infty$. Unit: length. Status: DERIVED.
- $\mathbf{1}$, $\mathbf{1}_8$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_8$ is the 8×8 identity matrix. Unit: dimensionless. Status: STANDARD.

- q: The local decay rate of a profile of dirac16complex along σ at the frequency ω, $q = \sqrt{m_\Psi^2 - \omega^2 \mathrm{e}^{2H|\sigma|}}$ (imaginary beyond the turning point, where the profile oscillates). Unit: GeV. Status: DERIVED.
- σ_t: The turning point of a state of dirac16complex of the frequency ω: the depth at which the warped frequency equals the bulk mass, $|\omega|\mathrm{e}^{H\sigma_\mathrm{t}} = m_\Psi$; beyond it every solution oscillates. Unit: length. Status: DERIVED.

The profile of dirac16complex at zero momentum: the local gap $m_\Psi^2 - \omega^2 \mathrm{e}^{2H|\sigma|}$ closes at the turning point σ_t, beyond which every solution oscillates. Because the time is warped, the warped frequency exceeds the bulk mass far from the fold for every $\omega \neq 0$: there is no gap in the bulk and no normalisable state of non-zero frequency, and the states that UM-3 bound to the fold are resonances in UM-4. Status: DERIVED (`um4_spectra.py`, U1).

$$\omega_0 = \omega_\mathrm{r} - \frac{\mathrm{i}}{2}\,\Gamma_\chi, \qquad \Gamma_\chi \simeq p_\mathrm{w}\, m_\Psi \exp\Big(-2F(\upsilon_\mathrm{b})\,\frac{m_\Psi}{H}\Big), \qquad F(\upsilon) = \ln\frac{1+\sqrt{1-\upsilon^2}}{\upsilon} - \sqrt{1-\upsilon^2}, \qquad \upsilon_\mathrm{b} = \frac{\omega_\mathrm{b}}{m_\Psi} \tag{FE-86}$$

Symbols in (FE-86).

- ω_0: The energy at rest (zero momentum along ordinary space and along the extra times) of a state of dirac16complex at the fold; complex for a resonance, $\omega_0 = \omega_\mathrm{r} - \frac{\mathrm{i}}{2}\Gamma_\chi$. Unit: GeV. Status: DERIVED.
- ω_r: The mass of the particle χ: the real part of the energy at rest of the resonance of dirac16complex at the fold, $\omega_\mathrm{r} = \omega_\mathrm{b}(1 - H/(2q_\mathrm{b})) + \ldots$. Unit: GeV. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- Γ_χ: The width of the particle χ at rest: the rate of its decay into the waves of dirac16complex in the bulk. Unit: GeV. Value: below $1/t_0 = 1.5 \times 10^{-42}$ GeV for dark matter. Status: DERIVED.
- p_w: The prefactor of the width of the particles χ in its tunnelling form $\Gamma_\chi \approx p_\mathrm{w} m_\Psi \mathrm{e}^{-2F(\upsilon_\mathrm{b})m_\Psi/H}$, taken from the computed widths at $m_\Psi/H = 40$. Unit: dimensionless. Value: 0.14. Status: ESTIMATE.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- $F(\upsilon)$: The tunnelling exponent of the width of the particles χ per unit of m_Ψ/H: the integral of the local decay rate q over the region inside the turning point, divided by m_Ψ/H, as a function of the ratio υ of the energy to the bulk mass. Unit: dimensionless. Value: 0.049 for $\upsilon = 0.866$; $(q_\mathrm{b}/\omega_\mathrm{b})^3/3$ for weak binding. Status: DERIVED.
- υ, υ_b: The argument of the tunnelling exponent F, a ratio of an energy to the bulk mass m_Ψ; $\upsilon_\mathrm{b} = \omega_\mathrm{b}/m_\Psi$ that of the particle χ. Unit: dimensionless. Status: DERIVED.
- ω_b: The energy at zero momentum of the state of dirac16complex at the fold in the limit $H \to 0$ (the bound state of UM-3): the particle χ of UM-4 has the mass $\omega_\mathrm{r} = \omega_\mathrm{b}(1 - H/(2q_\mathrm{b})) + \ldots$, equal to ω_b up to the relative correction $H/(2q_\mathrm{b})$. Unit: GeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_\mathrm{F} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.

The resonances of dirac16complex at the fold: complex energies with the real part ω_r (the mass of the particle χ) and the width Γ_χ, the rate of the tunnelling through the region inside the turning point. A particle χ of the dark matter must live longer than the universe, which requires m_Ψ/H above about 1.1×10^3 for $q_\mathrm{b} = m_\Psi/2$ and 1.7×10^5 for $q_\mathrm{b}/\omega_\mathrm{b} = 0.1$; with $H \geq 2.5$ TeV the particles are heavier than 2.4×10^6 GeV. Status: DERIVED (`um4_spectra.py`, U3 and U4), ESTIMATE (the prefactor and the bounds, U6).

9 From one point of the fold to another

$$-\mathrm{e}^{nH|\sigma|}\,\partial_\sigma\big(\mathrm{e}^{-nH|\sigma|}\,\partial_\sigma u\big) + \mathrm{e}^{2H|\sigma|}\,\mathsf{p}^2\,u = \delta_\mathrm{D}(\sigma), \qquad \mathbb{G}_n(\mathsf{p}) = u(0) = \frac{\mathrm{K}_{n/2}(\mathsf{p}/H)}{2\mathsf{p}\,\mathrm{K}_{n/2-1}(\mathsf{p}/H)} \tag{FE-79}$$

Symbols in (FE-79).

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

- n: The label of a field of the bulk in its profile equation: the exponent of the weight $\mathrm{e}^{-nH|\sigma|}$ of its σ-derivatives, $n = 7$ for a scalar and for each polarisation of the graviton, $n = 5$ for a gauge field. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- $u(\sigma)$, u_0: The profile along the hidden direction of a mode of a field of the bulk (a scalar, a gauge field or a polarisation of the graviton): the field is $u(\sigma)$ times a field of the fold; $u_0 = 1$ is the profile of the massless mode (the zero mode); in a propagator the response of the field to a unit source on the fold. Unit: dimensionless (a relative amplitude); length in a propagator. Status: DERIVED.
- p: The Euclidean momentum along the fold of a source on the fold: p^2 the square of its seven-dimensional momentum (for a static source spread uniformly over the extra times the square of its momentum along ordinary space); $\mathsf{p} = -\mathrm{i}m$ for a time-like momentum of the mass m with outgoing waves into the bulk. Unit: GeV. Status: STANDARD.
- δ_{D}: The Dirac delta function of σ: zero for $\sigma \neq 0$, with the integral 1 over any range that contains $\sigma = 0$; the integral of $\delta_{\mathrm{D}}(\sigma)$ times a smooth function of σ is the value of that function at $\sigma = 0$. Unit: 1/length (for $\delta_{\mathrm{D}}(\sigma)$). Status: STANDARD.
- $\mathbb{G}_n$, $\mathbb{G}_7$, $\mathbb{G}_5$: The propagator of a massless field of the bulk from one point of the fold to another, in the momentum space of the fold: the value at the fold of its response to a unit source on the fold; $\mathbb{G}_7$ for each polarisation of the graviton and for a scalar, $\mathbb{G}_5$ for a gauge field. Unit: GeV^{-1}. Status: DERIVED.
- K: The modified Bessel function of the second kind (of the order written as its subscript): the solution of the modified Bessel equation that decays exponentially for a large argument; for a half-integer order it is an exponential times a polynomial in the inverse of the argument, which makes the propagators of the bulk rational functions. Unit: dimensionless. Status: STANDARD.

The response of a massless field of the bulk to a unit source on the fold, for the Euclidean momentum p along the fold: the propagator of the bulk from one point of the fold to another ($n = 7$ for the graviton and a scalar, $n = 5$ for a gauge field). For small momenta it is the massless propagator of the fold with the strength $\frac{5}{2}H$ for the graviton and $\frac{3}{2}H$ for a gauge field. Status: DERIVED (`um4_propagators.py`, P1).

$$\begin{aligned}\mathbb{G}_7 &= \frac{H}{2\mathsf{p}^2}\,\frac{x_{\mathrm{p}}^3 + 6x_{\mathrm{p}}^2 + 15x_{\mathrm{p}} + 15}{x_{\mathrm{p}}^2 + 3x_{\mathrm{p}} + 3} = \frac{5H}{2\mathsf{p}^2}\Big(1 + \frac{x_{\mathrm{p}}^2}{15} - \frac{x_{\mathrm{p}}^4}{45} + \frac{x_{\mathrm{p}}^5}{45} + \dots\Big),\\ \mathbb{G}_5 &= \frac{H}{2\mathsf{p}^2}\,\frac{x_{\mathrm{p}}^2 + 3x_{\mathrm{p}} + 3}{x_{\mathrm{p}} + 1} = \frac{3H}{2\mathsf{p}^2}\Big(1 + \frac{x_{\mathrm{p}}^2}{3} - \frac{x_{\mathrm{p}}^3}{3} + \dots\Big), \qquad x_{\mathrm{p}} = \frac{\mathsf{p}}{H}\end{aligned} \tag{FE-80}$$

Symbols in (FE-80).

- $\mathbb{G}_n$, $\mathbb{G}_7$, $\mathbb{G}_5$: The propagator of a massless field of the bulk from one point of the fold to another, in the momentum space of the fold: the value at the fold of its response to a unit source on the fold; $\mathbb{G}_7$ for each polarisation of the graviton and for a scalar, $\mathbb{G}_5$ for a gauge field. Unit: GeV^{-1}. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\mathrm{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- p: The Euclidean momentum along the fold of a source on the fold: p^2 the square of its seven-dimensional momentum (for a static source spread uniformly over the extra times the square of its momentum along ordinary space); $\mathsf{p} = -\mathrm{i}m$ for a time-like momentum of the mass m with outgoing waves into the bulk. Unit: GeV. Status: STANDARD.
- x_{p}: The Euclidean momentum along the fold in units of H, $x_{\mathrm{p}} = \mathsf{p}/H$. Unit: dimensionless. Status: DERIVED.

For the half-integer orders of the Bessel functions the propagators are rational functions of $x_{\mathrm{p}} = \mathsf{p}/H$. In the

expansions for small momenta the even powers are contact interactions between the sources on the fold; the first odd power ($x_{\rm p}^5$ for gravity, $x_{\rm p}^3$ for a gauge field) gives a correction of long range. Status: DERIVED (`um4_propagators.py`, P2).

$$\mathcal{V}_7(r)=\frac{5H}{8\pi r}\Big[1+\frac{16}{15\pi}\,\frac{1}{(Hr)^5}+\dots\Big],\qquad \mathcal{V}_5(r)=\frac{3H}{8\pi r}\Big[1+\frac{4}{3\pi}\,\frac{1}{(Hr)^3}+\dots\Big] \tag{FE-81}$$

Symbols in (FE-81).

- $\mathcal{V}_7(r)$, $\mathcal{V}_5(r)$: The static potentials between two unit point sources on the fold at the distance r (spread uniformly over the extra times) from the exchange of the graviton ($\mathcal{V}_7$) and of a gauge field ($\mathcal{V}_5$) of the bulk: the three-dimensional Fourier transforms of $\mathbb{G}_7$ and $\mathbb{G}_5$. Unit: GeV^2. Status: DERIVED.
- r: A distance from a centre: the distance from the centre of the star along the path of a neutrino; in Newton's and Coulomb's laws the distance between two point sources on the fold. Unit: length. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda=-21H^2$ and the tension $\kappa\lambda_{\rm F}=12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H\gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

Newton's law (the first potential, multiplied by the gravitational coupling) and Coulomb's law (the second, multiplied by the square of the gauge coupling) between two point sources on the fold: the massive modes of the bulk add the relative corrections $0.3395/(Hr)^5$ and $0.4244/(Hr)^3$. Newton's law, tested by torsion balances down to 52 micrometres, holds there to one per cent if $H>7.7\times10^{-3}$ eV; for $H\geq 2.5$ TeV the corrections are 3×10^{-74} at 52 micrometres and 1.4×10^{-27} for Coulomb's law at the Bohr radius. Status: DERIVED (`um4_propagators.py`, P3), ESTIMATE (the factor of order one of the tensor structure of the massive gravitons; the bounds).

$$\Big(1+M_{\rm loc}^2\,\mathbb{G}_5(\mathsf{p})\Big)\Big|_{\mathsf{p}=-im}=0:\qquad m^2=m_0^2\Big(1-\frac{m_0^2}{3H^2}\Big)-\mathrm{i}\,\frac{m_0^5}{3H^3}+\dots,\qquad m_0^2=\frac{3}{2}\,HM_{\rm loc}^2,\qquad \Gamma_{\rm bulk}=\frac{m_0^4}{3H^3} \tag{FE-82}$$

Symbols in (FE-82).

- $M_{\rm loc}$: The mass term that the Higgs vacuum on the fold gives a gauge field of the bulk: $M_{\rm loc}^2\delta_{\rm D}(\sigma)$ in its profile equation; for the W, $M_{\rm loc}^2=\mathsf{g}_2^2(v^2+\mathcal{A}^2)/(4V_*)$. Unit: GeV$^{1/2}$ ($M_{\rm loc}^2$ in GeV). Status: DERIVED.
- $\mathbb{G}_n$, $\mathbb{G}_7$, $\mathbb{G}_5$: The propagator of a massless field of the bulk from one point of the fold to another, in the momentum space of the fold: the value at the fold of its response to a unit source on the fold; $\mathbb{G}_7$ for each polarisation of the graviton and for a scalar, $\mathbb{G}_5$ for a gauge field. Unit: GeV^{-1}. Status: DERIVED.
- p: The Euclidean momentum along the fold of a source on the fold: p^2 the square of its seven-dimensional momentum (for a static source spread uniformly over the extra times the square of its momentum along ordinary space); $\mathsf{p}=-im$ for a time-like momentum of the mass m with outgoing waves into the bulk. Unit: GeV. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m_{ij}^2=m_i^2-m_j^2$ the squared-mass difference of the states i and j (Δm_{21}^2, Δm_{31}^2, Δm_{32}^2); Δm^2 without labels the difference of the two states of a resonance, $m_2^2-m_1^2$, multiplied by $v_{\rm loc}^2/v^2$ where this differs from 1; m without a label the mass of a mode of a field of the bulk as the fold sees it, $m^2=\omega^2-k^2$; for the W and Z bosons their complex pole mass, $m^2=m_0^2-\mathrm{i}m_0\Gamma_{\rm bulk}$ to the leading order; in the relic abundance of the particles χ their mass. Unit: GeV. Status: STANDARD.
- m_0: The mass of the W or the Z boson without the massive modes of the bulk (the mass of the massless mode with the Higgs vacuum), $m_0^2=\frac{3}{2}HM_{\rm loc}^2$; $m_0^2=g_2^2(v^2+\mathcal{A}^2)/4$ for the W. Unit: GeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda=-21H^2$ and the tension $\kappa\lambda_{\rm F}=12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H\gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- $\Gamma_{\rm bulk}$: The width of the W or the Z boson into the bulk (its decay into the massive modes of the gauge fields of the bulk), $\Gamma_{\rm bulk}=m_0(m_0/H)^3/3$. Unit: GeV. Value: 1.4 MeV for the Z and 0.9 MeV for the W at $H=2.5$ TeV. Status: DERIVED.

The W and Z bosons: the Higgs vacuum on the fold gives the gauge fields of the bulk a mass term concentrated on the fold, and the pole of the propagator lies at a complex mass: the W and the Z are resonances of the bulk, with the width $\Gamma_{\text{bulk}} = m_0(m_0/H)^3/3$ into the bulk (m_0 the mass that the zero mode alone would have, $m_0^2 = g_2^2(v^2+\mathcal{A}^2)/4$ for the W and $(g_1^2+g_2^2)(v^2+\mathcal{A}^2)/4$ for the Z, with the Higgs vacuum value v and the ad hoc component $\mathcal{A}$ of the author's vacuum). A Z boson that decays into the bulk leaves the fold unseen and adds to its invisible width; the number of light neutrinos measured at the Z, 2.9963 ± 0.0074, allows at most 1.4 MeV of extra invisible width, so $H > 2.5$ TeV. Status: DERIVED (`um4_propagators.py`, P4), ESTIMATE (the bounds, P5).

bound on H	from	value
invisible width of the Z boson	its decay into the bulk, $\Gamma_{\text{bulk}} < 1.4$ MeV (the measured number of light neutrinos 2.9963 ± 0.0074, P. Janot and S. Jadach, Phys. Lett. B 803 (2020) 135319)	$H > 2.5$ TeV (2534 GeV)
contact interactions of electrons and muons	the contact term of the propagator of the photon, against the limit of ATLAS $\Lambda_{\text{CI}} > 26.0$ TeV (JHEP 11 (2020) 005)	$H > 1.3$ TeV
the parameter rho of the electroweak fits	the shift $-m_0^4/(6H^2)$ of the squared masses of the W and Z, against 1.00031 ± 0.00019 (Particle Data Group 2024)	$H > 0.97$ TeV
Newton's law at 52 micrometres	the correction $0.3395/(Hr)^5$, against the torsion balances of J. G. Lee et al., Phys. Rev. Lett. 124 (2020) 101101	$H > 7.7 \times 10^{-3}$ eV

All four are ESTIMATE (Λ_{CI} is the scale of a contact interaction, r the distance between the sources). The strongest, the invisible width of the Z boson, is the bound used in every UM-4 document.

10 Causality around an expanding fold

$$c_{\text{loc}}(\sigma) = 1 - \frac{a_4''}{5H^2}\left(\mathrm{e}^{2H|\sigma|} - 1\right), \qquad \sigma(x_3) = \frac{|a_4''|}{5H}\, x_3\,(L - x_3), \qquad \Delta x_4 = L - \frac{a_4''^2 L^3}{150H^2} \tag{FE-84}$$

Symbols in (FE-84).

- c_{loc}: The coordinate speed of light along x_3 at the depth σ (the rate of change of x_3 with x_4 along a null line at fixed σ), in units of that on the fold: $\mathrm{e}^{H|\sigma|}$ on the static block of UM-3, whose time was not warped, 1 on the static block of UM-4, and on an expanding fold of UM-4 different from 1 in proportion to the acceleration a_4 of the expansion. Unit: dimensionless. Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \operatorname{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the warp factor $\mathrm{e}^{-2H|\sigma|}$ of the metric, the cosmological constant $\Lambda = -21H^2$ and the tension $\kappa\lambda_{\text{F}} = 12H$ of the fold; in UM-4 a free parameter, bounded from below by the propagation of the gauge fields in the bulk. Unit: GeV (1/length). Value: $H \gtrsim 2.5$ TeV (from the invisible width of the Z boson). Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.

- L: A distance between two points of the fold along ordinary space: for a gravitational wave the radius of a ring of free test masses (ΔL the change of the distance of one of them from the centre), for a signal the distance between its emitter and its receiver. Unit: length. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.

An expanding fold with frozen extra times: the coordinate speed of light c_{loc} along x_3 at the depth σ differs from 1 in proportion to the acceleration a_4'' of the expansion. Where the expansion decelerates (the eras of radiation and of matter) the light cone of the bulk is slightly wider than that of the fold, and the path that gains the most between two points of the fold at the distance L is the parabola of the second relation, which arrives earlier than light along the fold by $a_4''^2 L^3/(150H^2)$. Over the Hubble distance today this is 4.5×10^{-94} of the travel time for $H = 2.5$ TeV (3×10^{-63} even for $H = 10^{-3}$ eV): no signal through the bulk outruns light on the fold at any observable level, and on the static fold none at all (FE-83). Status: DERIVED (`um4_causality.py`, C2 and C3).

11 What the change of the metric changes elsewhere

The warped time changes more than the speeds of the fields of the bulk. Each item is derived in the field equations of UM-4 and in their programs; the values are those of UM-4.

- **Gravity on the fold** is general relativity: the Einstein equations of seven dimensions with $\kappa_7 = 5\kappa H/2$ (FE-71), with frozen extra times the Friedmann equations, gravitational waves at the speed of light at every wavelength (FE-63). The order of the corrections is $(a_4'/H)^2$. Status: DERIVED.
- **The tension of the fold** is $\kappa\lambda_{\mathrm{F}} = 12H$ (FE-30), in the four-dimensional normalisation $\lambda_{\mathrm{F}} V_* = 30H^2 M_{\mathrm{red}}^2 = 1.1 \times 10^{45}$ GeV^4 for $H = 2.5$ TeV (M_{red} the reduced Planck mass): the observed dark energy needs it tuned to one part in 10^{92}, the cosmological-constant problem of the brane of Randall and Sundrum. Status: DERIVED (the value), OPEN (the tuning).
- **The bulk needs no stabiliser.** With Einstein gravity and $\Lambda = -21H^2$ the static block is a vacuum (FE-32); UM-3 needed either a stabiliser of the bulk (with the author's original weights) or the degenerate weights $(1, 1/20, 1/360)$. Status: DERIVED.
- **The author's vacuum** $a_4 = Hx_4$ with constant b_4 is not a solution of UM-4 (as a fold with matter it would need a negative energy density). Status: DERIVED (`um4_primordial.py`, PR0).
- **The states of dirac16complex at the fold** are long-lived resonances, not bound states (FE-86): the particles χ of the dark matter live longer than the universe only if m_Ψ/H is large, so with $H \geq 2.5$ TeV they are heavier than 2.4×10^6 GeV; above 8.9×10^4 GeV no thermal relic is possible (the unitarity bound), so their abundance needs a non-thermal origin; and bound to the fold by the Higgs portal they cannot both live long enough and escape direct detection: the dark matter of UM-4 needs a binding of χ to the fold that does not couple it to nucleons. Status: DERIVED, ESTIMATE (`um4_spectra.py`, U6; `um4_thermal.py`, T7 and T8), OPEN (the binding and the origin).
- **Inflation** of the fold in the regime of general relativity needs a rate below $H/10$; with $H = 2.5$ TeV only a rate below 250 GeV, far below the rates of the usual models of inflation (Higgs inflation, with a curvature coupling that the action of UM-4 does not contain, would need $H > 1.6 \times 10^{14}$ GeV). Status: DERIVED, ESTIMATE (`um4_primordial.py`, PR2 to PR4), OPEN (the inflaton).
- **The extra times** remain time-like: every field with a momentum along them is tachyonic, in the bulk as on the fold, and UM-4 keeps only fields without such a momentum; a gauge principle that removes it is needed. Status: DERIVED (`um4_extratimes.py`, EX1 to EX5), OPEN (the principle).

12 What is open

- The origin of the tension of the fold, its tuning to one part in 10^{92} against the observed dark energy, and the origin of the pressure along the extra times that holds their volume fixed.
- The projected Weyl tensor of the bulk that an expanding fold with matter other than radiation requires at the order of the square of its energy density divided by the tension (`um4_fold.py`, F6), and the bulk solution that would fix it.
- A gauge principle that removes the momentum of every field along the extra times.

- A binding of the particles χ of dirac16complex to the fold that does not couple them to nucleons, and a non-thermal origin of their abundance.
- The inflaton of the fold.

13 What was verified

Every statement labelled DERIVED in this document is verified by a program of UM-4 (Python with sympy for the exact algebra and with numpy and scipy for the numerics; every program runs in less than 423 seconds). The checks that this document cites, all passed:

program	checks	what they verify
`um4_geometry.py`	G1 to G10	the proper chart, the exact curvature, the static block of constant curvature, the Lovelock tensors, the bulk equations, the Ricci-flat folds, the pure tension, the light cone of every slice, the spin connection with the author's matrices, the static warps and their vacua
`um4_gravity.py`	L1 to L4	the vacua of the static family, the graviton around the static block for any weights, the light cone on every background only for Einstein gravity, the choice of UM-4
`um4_bulk.py`	B1 to B10	the scalar, the photon, the transverse photon of spin 1, the graviton, the plus and cross polarisations of spin 2, the bulk fermion, the boost invariance of dirac16complex, the speeds of UM-4, the speeds of UM-3 recomputed, the continuum of massive modes from zero mass
`um4_fold.py`	F1 to F6	the bulk metric of an expanding fold, the Einstein equations of seven dimensions on the fold, the exact equations of the fold, the Friedmann equations, the size of the corrections, the Weyl term
`um4_foldsources.py`	FS1 to FS5	the tension, the gravitational waves of the fold, the induced gravity, the tuning of the tension, the fold fluid
`um4_propagators.py`	P1 to P5	the propagators of the bulk, their expansions, Newton's and Coulomb's laws, the W and Z resonances, the bounds on H
`um4_causality.py`	C1 to C3	no shortcut through the static bulk, the light cone of an expanding fold, the largest advance
`um4_spectra.py`	U1 to U8	the resonances of dirac16complex, their widths, their dispersion, their lifetime, their Krein signature and mirror symmetry

14 Further symbols

Further symbols. These symbols appear in the definitions of the symbols listed after the equations of this document.

- $M_{\rm red}$: The reduced Planck mass, $M_{\rm red} = (8\pi G_{\rm N})^{-1/2}$. Unit: GeV. Value: 2.435×10^{18} GeV. Status: STANDARD.
- P_7: The pressure along ordinary space of the matter of the fold above its tension in the seven-dimensional normalisation (the component $\mathsf{t}^{(i)}{}_{(i)}$). Unit: GeV^7. Status: DERIVED.
- Σ^{ab}, $\Sigma^{\mu\nu}$: The generators of Spin(4,4) on dirac16complex, $\Sigma^{ab} = \frac{1}{4}[T^a, T^b]$ (the scaled commutators of the author's matrices); $\Sigma^{\mu\nu} = \mathbf{e}_a{}^\mu \mathbf{e}_b{}^\nu \Sigma^{ab}$ along the fold. Unit: dimensionless. Status: AUTHOR.
- $\mathcal{A}$: The author's ad hoc upper component of the Higgs vacuum, almightyAdHoc, $\mathcal{A}(x_4) = v_{\mathcal{A}} \mathrm{e}^{-2a_{40}x_4}$. Unit: GeV. Status: AUTHOR.
- ρ_7: The energy density of the matter of the fold above its tension in the seven-dimensional normalisation (minus the component $\mathsf{t}^{(4)}{}_{(4)}$: an energy per unit volume of the six other directions of the fold); $V_* \rho_7$ in the four-dimensional normalisation. Unit: GeV^7. Status: DERIVED.
- ϖ_{Mab}: The canonical (Levi-Civita) spin connection of the vielbein, antisymmetric in a, b. Unit: 1/length. Status: STANDARD.
- $m_u, \ldots, m_3$: The masses of the quarks (m_u, m_c, m_t; m_d, m_s, m_b), of the charged leptons (m_e, m_μ, m_τ) and of the three neutrino mass states (m_1, m_2, m_3, Dirac masses). Unit: GeV. Value: measured values; $\Delta m^2_{21} = 7.41 \times 10^{-5}$ eV^2, $|\Delta m^2_{31}| = 2.511 \times 10^{-3}$ eV^2 in the computations of UM-4 (the global fit NuFIT 5.2 of the oscillation data, November 2022), 7.49×10^{-5} and 2.513×10^{-3} eV^2 in NuFIT 6.0 (2024); the lightest mass $m_1 = 0$ in the computations, below 23 meV by DESI DR2 with the CMB and NuFIT 6.0. Status: STANDARD.
- m_h: The mass of the Higgs boson. Unit: GeV. Value: 125.25 GeV (Particle Data Group 2022). Status: STANDARD.
- q_7: The pressure along the extra times of the matter of the fold above its tension in the seven-dimensional normalisation (the component $\mathsf{t}^{(r)}{}_{(r)}$). Unit: GeV^7. Status: DERIVED.
- t^A: The generators of $SU(3)$ on colour triplets, t^A: half the Gell-Mann matrices, with $\mathrm{tr}(t^A t^B) = \delta^{AB}/2$. Unit: dimensionless. Status: STANDARD.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\, G_{\rm F})^{-1/2}$ with the Fermi constant $G_{\rm F} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- $v_{\rm loc}$: The local value of the electroweak vacuum, $v^2_{\rm loc} = 2\langle \Phi^\dagger \Phi \rangle$: it sets the local masses of the quarks and leptons (each mass times $v_{\rm loc}/v$) and the local Fermi constant; $v_{\rm loc} = v$ in the Standard-Model vacuum. Unit: GeV. Value: v in the late universe outside dense dirac16complex. Status: DERIVED.
- $G_{\rm F}$: The Fermi constant, $G_{\rm F} = 1/(\sqrt{2}\, v^2)$ in the Standard-Model vacuum (its measured value); in the equations of the neutrino flavours it stands for the local Fermi constant of the ad hoc vacuum. Unit: GeV^{-2}. Value: 1.1664×10^{-5} GeV^{-2}. Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- $v_{\mathcal{A}}$: The amplitude of almightyAdHoc at $x_4 = 0$; the author's almightyAdHoc, $\mathrm{e}^{-2a_{40}x_4}$ (written Exp[-2 a40 x4] in his notebook), has $v_{\mathcal{A}} = 1$ in the units of his notebook. Unit: GeV. Value: 1 in the units of the author's notebook; UM-4 keeps it free (the examples take $v_{\mathcal{A}} = v$ and $10v$). Status: AUTHOR (the form), ASSUMED (a free amplitude).
- $\mathcal{T}$: The energy-momentum tensor of any one term of the fold (in $\mathcal{T}^r{}_r = \mathcal{L}_{\rm fold}$). Unit: GeV^4. Status: STANDARD.
- $\mathcal{W}$: The warp factor: in a warped metric, the function of the position along one direction that multiplies the squared steps of other directions (its square root multiplies their lengths); in UM-4 the function of σ that multiplies all seven directions of the fold, ordinary space, the time and the extra times, $\mathcal{W} = \mathrm{e}^{-2H|\sigma|} = (\sin^2 z)^{1/6}$, equal to 1 on the fold. Unit: dimensionless. Status: DERIVED.
- $c_{\rm b}$: The speed coefficient of the particles χ of dirac16complex along ordinary space, in units of the speed of light on the fold: at small momentum $\omega^2 = \omega_0^2 + c_{\rm b}^2 \vec{k}^2$. In UM-3, whose time was not warped, $c_{\rm b}^2 = 1.19$ for $H = 0.05\, m_\Psi$ and $q_{\rm b} = 0.3\, m_\Psi$ (faster than light); in UM-4 the boosts of the fold are a symmetry and $c_{\rm b} = 1$ exactly. Unit: dimensionless. Value: 1 in UM-4; 1.09 in UM-3. Status: DERIVED.
- $\mathcal{L}_{\rm fold}$: The Lagrangian of any one term of the fold (four-dimensional normalisation). Unit: GeV^4. Status: STANDARD.