Dark-sector hypotheses for dirac16complex and dirac16complex00: the equation of state seen from 3-space

Hypothesis and Hypothesis00 investigated in the author's primordial gravitational field with exponentially deflating extra times and the second metric function $b_4(x_4)$: the observer assumption, the method, the results for each field, the comparison with the Supernovae Unite values, the verdicts and what remains open

dirac16complex project (with Claude Opus 5.5)

September 2026

Contents

1 The hypotheses and the short answer **4**

2 Setting and conventions **5**

2.1 The author's metric and coordinates 5

2.2 The two fields and the energy-momentum tensor 6

2.3 The Unite values and the CPL convention 6

2.4 The history with $b_4(x_4)$ 6

3 The observer assumption: ρ_4 and the definitions (A), (B), (C) **7**

3.1 Exact identities 7

3.2 How every verdict depends on the assumption 8

3.3 The expansion-inferred equation of state 9

4 Method **9**

4.1 Exact derivations first 9

4.2 The conservation identities as checks of the numerics 9

4.3 The Kohn-Sham history (dirac16complex) 10

4.4 Condensates 10

4.5 Modes and Krein-signed energies (dirac16complex00) 11

4.6 Gases, mixtures and wave packets . . . 11
4.7 Phantom conditions . . . 11

5 Results for dirac16complex **12**
5.1 The condensate . . . 12
5.2 The Kohn-Sham gas . . . 12
5.3 The bulk band (independent implementation) . . . 13
5.4 Mixtures with a condensate . . . 13
5.5 Phantom and crossing of -1 . . . 14

6 Results for dirac16complex00 **14**
6.1 The condensate . . . 14
6.2 Modes, Krein-signed energies and growing modes . . . 15
6.3 The models M1 to M5 . . . 15
6.4 The independent implementation B . . . 17

7 Comparison with the Unite values **17**

8 The lead's analysis (SPEC section 11), checked point by point **18**

9 Verdicts **19**
9.1 dirac16complex (Hypothesis) . . . 19
9.2 dirac16complex00 (Hypothesis00) . . . 19
9.3 The hypotheses . . . 20

10 What remains open **20**

11 Verification records **20**
11.1 Reports and their counts . . . 20
11.2 Checks per report . . . 21
11.3 Re-verification for this document . . . 24

12 Reproduction **25**

13 What is proved, computed, assumed and not established **26**
13.1 Proved (exact, by symbolic computation in the named checks) . . . 26

13.2 Computed (numerical, with the stated tolerances and two implementations) 26
13.3 Assumed 26
13.4 Not established 27

Abstract

The author asked on 2026-10-01 that two hypotheses be investigated: that dirac16complex (Hypothesis) and dirac16complex00 (Hypothesis00) provide a possible physical mechanism for a time-varying dark-energy equation of state and/or a time-varying dark-matter equation of state. This document reports what the Revision computation found for the author's metric of 2026-10-14, in which the three extra-time components carry the second function $b_4(x_4)$; b_4 is a function of x_4, and every result below is computed with it (the earlier record used $b_4 = 1$; it is withdrawn and quoted only for comparison). Every number is read from a named Revision report. The history of the numbers is $a_4 = Hx_4$ with $b_4(x_4) = b_4(0)\, e^{(4/3)Hx_4}$: with the author's weights it is the history that the mixed (x_4, x_0) field equation allows for a source without a $T^{x_4}{}_{x_0}$ component, and the author's equations then require the constant source $\kappa(\rho, p_3, p_t, p_0) = (-2/27, -2/9, -22/27, -14/27)$. The equation of state that a 3-space observer infers is not fixed by the field equations alone: the extra times x_5, x_6, x_7 are time-like and deflate exponentially, and the 4-dimensional density ρ_4 needs an ASSUMPTION about how the observer normalises over them. For the three stated definitions (A), (B) and (C) the dilution-inferred equation of state obeys exactly $w_{\text{eff}} = X/E - s + (\beta/2)(P_t/E + s)$ with $X = P_3 - P_t$, $\beta = d\ln b_4/da_4$, $s = 0$ for (A), (B) and $s = 1$ for (C); on the history ($\beta = 4/3$) every verdict moves by exactly $-1/3$ between (A, B) and (C). dirac16complex: the computed Kohn-Sham gas (615 instantaneous states along the history, solved at the coupling that b_4 dictates) is radiation-like, with $w_{\text{eff}}(A)$ rising toward 1/3 (0.2929 to 0.3183 for $N = 688$, $\lambda = 0$); the dark-matter-like fall from 1/3 toward 0 occurs for quanta of the massive bulk band (0.239626 to 2.268e-03 for one shell), which the computed ground states do not populate; under (C) the gas has w_{eff} between -0.040541 and -0.006532, close to 0 and not dark-energy-like; nothing computed comes near the Unite pair $(w_0, w_a) = (-0.861, -0.60)$ (closest distance 0.291, with a drifting condensate) and no computed state crosses $w = -1$. dirac16complex00: a positive-energy gas of good-sector modes has the dark-matter-like law $1/3 \to 0$ under (B), independent of b_4; every positive-energy model has $w_{\text{eff}}(C) \geq -1/3$ as long as its modes have real frequencies, so the Unite values are out of reach without a ghost-like component, and $w_0 = -0.861$ needs a ghost fraction of at least 1583/1000 of the total energy. The condensate's ratio drifts with b_4: it equals the Unite constant $w = -0.764$ at $\lambda S/m = -382/441$ only at one instant. A model with as many tuned parameters as matched numbers reproduces them by construction and is NOT a prediction. This record establishes neither Hypothesis nor Hypothesis00; it states what each field can and cannot produce under its stated assumptions.

1 The hypotheses and the short answer

The author's hypotheses, quoted from `Revision/README.md`:

- "Hypothesis: dirac16complex provides a possible physical mechanism for a time-varying dark energy equation of state and/or a possible physical mechanism for a time-varying dark matter equation of state, both of which you will investigate."
- "Hypothesis00: dirac16complex00 provides [the same], both of which you will investigate."

`Revision/SPEC.md` section 8 makes both hypotheses questions to be INVESTIGATED, not assumed: for each field, the equation of state seen by a 3-space observer as the extra times deflate and 3-space inflates (the condensate, the Kohn-Sham gas, mixtures), its time dependence, the CPL parameters, and the comparison with the Supernovae Unite values. Section 11 of the same file records the lead's analysis, to be checked; section 8 of this document checks it point by point.

The short answer (details and records in sections 3 to 9):

1. The observer's ρ_4 is an ASSUMPTION (section 3). Every verdict below is stated for each definition; on the history with $b_4(x_4)$ the definitions differ by exactly 1/3.

2. dirac16complex, dark matter: a time-varying dark-matter-like equation of state (w_{eff} falling from 1/3 toward 0 under (A, B)) is present in the theory for quanta of the massive bulk band, independent of b_4, not in the computed Kohn-Sham ground states, whose gas is radiation-like.

3. dirac16complex, dark energy: not found. Under (C) the gas has w_{eff} between -0.040541 and -0.006532, close to 0; only a drifting condensate gives negative values, with a strongly thawing slope; no Kohn-Sham state comes near the Unite pair or crosses $w = -1$, and a drifting $\lambda < 0$ condensate crosses -1 toward the past (section 5.5).

4. dirac16complex00, dark matter: a positive-energy gas of good-sector modes has the time-varying law $1/3 \to 0$ under (B) and in the ratio p_3/ρ, independent of b_4.

5. dirac16complex00, dark energy: not with positive energies: $w_{\text{eff}}(C) \geq -1/3$ for positive-energy content with real frequencies; the Unite w_0 needs a ghost-like component larger than the positive energy.

6. On the history $a_4 = Hx_4$ the observer's expansion itself reads $w_{\text{exp}} = -1$ exactly, for every $b_4(x_4)$.

2 Setting and conventions

2.1 The author's metric and coordinates

The coordinates are the author's, $x_0, x_1, \ldots, x_7$, in the author's order (`Dirac_Einstein-Lovelock_4+4_Nash.nb`, input cell In[48]: `X = {x0[], x1[], x2[], x3[], x4[], x5[], x6[], x7[]}`): x_0 is the hidden space direction, with $z = 6Hx_0 \in (0, \pi/2)$ and $H > 0$ the author's constant, x_1, x_2, x_3 are ordinary 3-space (inflating), x_4 is the time, and x_5, x_6, x_7 are the three extra times, which are time-like and DEFLATE exponentially as a_4 increases. (Until 2026-10-14 this record named them $x_1, \ldots, x_8$, with the hidden direction x_8 listed last; the user ordered the author's coordinates on 2026-10-14.) The metric is the author's of 2026-10-14 (input cells In[97] to In[102] of that notebook),

$$\begin{aligned} ds^2 = {} & \cot^2 z\, dx_0^2 + e^{2a_4(x_4)} \sin^{1/3} z\, (dx_1^2 + dx_2^2 + dx_3^2) - dx_4^2 \\ & - e^{-2a_4(x_4)}\, b_4(x_4) \sin^{1/3} z\, (dx_5^2 + dx_6^2 + dx_7^2), \end{aligned}$$

of signature (4,4), with $\sqrt{|g|} = b_4^{3/2} \cos z$ and $b_4(x_4) > 0$ a function of x_4. The 3-space scale factor is $e^{a_4} \sin^{1/6} z$ and the extra-time scale factor $e^{-a_4} b_4^{1/2} \sin^{1/6} z$. The observer's scale factor is $a = e^{a_4 - a_{4,\text{today}}}$, normalised to $a = 1$ at a chosen "today" $a_{4,\text{today}}$ (a free parameter); b_4 does not enter it. Along a history $\beta = d \ln b_4 / da_4 = b_4' / (a_4' b_4)$. The Kohn-Sham records use the hidden coordinate $y = \ln(\sin z)/(6H)$. The record that reads the author's notebook, without evaluating it, is `Revision/dark_sector/reports/author-notebook-inputs.json` (`author_notebook_inputs.wls`): 147 cells, 113 input cells, and the only cell not in the version of the user's order of 2026-10-14 (commit b13373c) is In[137], the input X alone, which displays the coordinate list.

2.2 The two fields and the energy-momentum tensor

dirac16complex is Ψ: 16 complex anticommuting components, a Pin(4,4) spinor, quantised canonically; its energy-momentum tensor is an operator, and the numbers below are expectation values in the instantaneous Kohn-Sham states of the good sector (no extra-time momentum), in the positive realisation of the good sector (with the ASSUMED brane condition and the chosen tip condition of the Kohn-Sham record, which make its blocks self-adjoint; the field-theory record constructs the positive realisation explicitly only for single good-sector momenta with frozen coefficients). dirac16complex00 is Φ: 16 complex commuting components that transform as a Pin(4,4) spinor, a classical field; its energy-momentum tensor is the classical bilinear, whose energy has both signs (Krein-signed energies, section 6.2). Both fields have the Lagrangian of `Revision/SPEC.md` section 3 with $U(S) = \frac{\lambda}{2}S^2$, $S = \bar{\Psi}\Psi$, and the gravity term $\gamma^\mu\Omega_\mu = 3H\gamma^{(x_0)} + \frac{3}{4}(b_4'/b_4)\gamma^{(x_4)}$. The energy density is $\rho = -T^{x_4}{}_{x_4}$ and the pressures are $p_\mu = T^\mu{}_\mu$ (no sum): p_3 of 3-space, p_t of the extra times and p_0 of the hidden direction. For homogeneous on-shell states $\rho = mS + U$ and $p = SU' - U$. For a state of the Kohn-Sham record, E, P_3, P_t and P_0 are the proper 7-volume integrals of ρ, p_3, p_t and p_0 per unit coordinate volume, and $X = P_3 - P_t$.

2.3 The Unite values and the CPL convention

The author's private PDF (not part of the repository; only these numbers are used) gives the Supernovae Unite constant-w fit $w = -0.764$ and the CPL fit $w(a) = w_0 + w_a(1 - a)$ with $(w_0, w_a) = (-0.861, -0.60)$. In the deep past the CPL line tends to $w_0 + w_a = -1.461$ (quoted as -1.46), which is phantom; the line crosses $w = -1$ at $a = 461/600 = 0.7683$ ($z = 139/461$; record `unite_crossing_point`). In this convention thawing means $w_a < 0$ (w rises from near -1) and freezing $w_a > 0$; the formula decides (the PDF's table states the opposite signs).

The CPL tangent of a model is $w_0 = w(1)$, $w_a = -dw/da$ at $a = 1$. The fits are least squares of $w(a)$ over $a \in [1/2, 1]$ or $[1/3, 1]$ (record `cpl_least_squares_rule`). The "constant w" of a model is the mean of $w(a)$ over the range, a PROXY: applied to the Unite CPL line itself it gives -1.011 on $[1/2, 1]$ and -1.061 on $[1/3, 1]$ (record `unite_line_fit_proxy`), so it is not the supernova constant-w fit -0.764, which weights the data. No supernova likelihood, distance fit or covariance is computed anywhere in this record.

2.4 The history with $b_4(x_4)$

The numbers are computed on $a_4 = Hx_4$ (the author's $A = 1$) with $b_4(x_4) = b_4(0)\, e^{(4/3)Hx_4}$, that is $\beta = 4/3$ and $b_4 \propto a^{4/3}$. The reason (records `author_weights_b4_branch`, `D6_author_weights_branch`): with the author's weights $w_1 = 1$, $w_2 = 1/32$, $w_3 = 1/1152$, $\Lambda = 10$ the (x_4, x_0) component of the field equations is $\cot z\, \beta_v\, Q(\alpha, \beta_v)$ with $\alpha = a_4'/H$, $\beta_v = b_4'/(Hb_4)$ (`Revision/metric_b4/results/vacuum-b4.json`); a source without a $T^{x_4}{}_{x_0}$ component, and every source of this record is one, needs $\beta_v Q = 0$. On $a_4 = Hx_4$ the component is $\frac{3}{64}\beta_v^2(3\beta_v - 4)\cot z$, which vanishes for a constant b_4 and for $\beta_v = 4/3$, $d\ln b_4/da_4 = 4/3$. The extra-time scale factor is then $e^{-a_4}b_4^{1/2} \propto a^{-1/3}$: the extra times still deflate exponentially. In Einstein gravity a $b_4(x_4)$ needs $T^{x_4}{}_{x_0} = \frac{3}{2}H\cot z\, b_4'/(\kappa b_4) \neq 0$ (`einstein_mixed_component_b4`), which no source of this record has. On this history the author's equations require the constant source

$$\kappa\,(\rho,\ p_3,\ p_t,\ p_0) = \Big(-\frac{2}{27},\ -\frac{2}{9},\ -\frac{22}{27},\ -\frac{14}{27}\Big),$$

computed twice: from the field equations of `Revision/field_equations_a4/a4-equations.json` (key b4), which with the author's weights reproduce the author's vacuum first (`author_weights_vacuum_recovered`, `branch_required_source`), and directly from the author's Lovelock tensors (`D7_branch_required_source`). It satisfies both conservation identities with b_4. A condensate cannot be this source (its S decays like $b_4^{-3/2}$; `condensate_cannot_source_branch`), and the Kohn-Sham states are not admissible sources (section 10). The history is therefore a PRESCRIBED BACKGROUND for test fields.

3 The observer assumption: ρ_4 and the definitions (A), (B), (C)

3.1 Exact identities

The proper 7-volume element of a slice $x_4 = \text{const}$ is $b_4^{3/2}\cos z$: the inflation e^{3a_4} of the proper 3-volume is compensated exactly by the factor e^{-3a_4} of the proper extra-time volume $e^{-3a_4}b_4^{3/2}$, and the 7-volume changes with x_4 through b_4. The conservation law $\nabla_\mu T^\mu{}_\nu = 0$ for a diagonal $T(x_0, x_4)$, re-derived from the author's metric with Christoffel symbols computed anew in sympy and in Wolfram, gives

$$\frac{d\rho}{dx_4} = -3a_4'(p_3 - p_t) - \frac{3}{2}\,\frac{b_4'}{b_4}\,(\rho + p_t), \qquad \frac{dE}{da_4} = -3X - \frac{3}{2}\,\beta P_t,$$

and $dp_0/dx_0 + 3H\cot z\,(2p_0 - p_3 - p_t) = 0$, free of b_4. The records of the volumes and of these identities (the negative control: the form without the b_4 term fails):

record	record	record
`sqrt_det_g_equals_b4_three_halves_cos_z`	`proper_7_volume_element_b4`	`proper_3_and_extra_time_volume_scalings_b4`
`local_model_volume_b4`	`D1_volume_b4`	`conservation_x4_identity_b4`
`conservation_x0_identity`	`conservation_other_components_vanish`	`conservation_equals_a4_record_b4`
`conservation_x4_author_metric_b4`	`conservation_x0_author_metric`	`D2_conservation_b4`
`conservation_negative_control`	`integrated_identity_dE_dx4_b4`	`hidden_coordinate_dy_dx0`
`conservation_x0_in_y_form`		

The 3-space observer averages over the hidden direction with the weight $\cos z$, which does not depend on x_4 (record `observer_hidden_average_commutes`); the factor $b_4^{3/2}$ of the proper weight is written out below. The extra times are time-like, so the observer's 4-dimensional density needs a normalisation over them. Three are recorded:

- (A) the extra times are compact with a fixed coordinate period, i.e. closed time-like directions; $\rho_4 = E/(\text{proper 3-volume}) \propto E\, e^{-3a_4}$;
- (B) the extra times are non-compact and ρ_4 is taken per unit extra-time COORDINATE volume; $\rho_4 \propto E\, e^{-3a_4}$;
- (C) ρ_4 is taken per unit PROPER 7-volume (equivalently per unit proper extra-time volume $e^{-3a_4} b_4^{3/2}$); $\rho_4 \propto E\, b_4^{-3/2}$.

With the dilution-inferred equation of state $w_{\text{eff}} = -1 - \frac{1}{3}\, d\ln\rho_4 / d\ln a$ the conservation identity gives exactly (records `w_eff_A_b4`, `w_eff_B_equals_w_eff_A_b4`, `w_eff_C_b4`, `w_eff_general_normaliser_b4`, `D3_w_eff_general_b4`)

$$w_{\text{eff}} = \frac{X}{E} - s + \frac{\beta}{2}\Big(\frac{P_t}{E} + s\Big), \qquad s = 0 \text{ (A, B)}, \quad s = 1 \text{ (C)}.$$

So $w_{\text{eff}}(A) = w_{\text{eff}}(B) = X/E + (\beta/2)P_t/E$ and $w_{\text{eff}}(C) = w_{\text{eff}}(A) - 1 + \beta/2$: on the history every verdict moves by exactly $-1/3$ between (A, B) and (C). The whole difference to the withdrawn $b_4 = 1$ formulas (X/E and $X/E - 1$) is the β term (`w_eff_difference_to_withdrawn_record`). The dirac16complex00 record names the same two scalings N1 (per unit extra-time coordinate volume, $\rho_4 = e^{-3a_4} b_4^{3/2} \langle\rho\rangle$, the scaling of (A) and (B)) and N2 (per unit proper 7-volume, $\rho_4 = \langle\rho\rangle$, definition (C)), with $w_{\text{eff}}(N1) = w_3 - w_t + (\beta/2) w_t$ and $w_{\text{eff}}(N2) = w_{\text{eff}}(N1) - 1 + \beta/2$ for isotropic sources (records `observer_N1_weff_identity_b4`, `observer_N2_weff_identity_b4`). Below, (B) stands for (A) and (B) together and for N1, and (C) for N2. The ratio $w = p_3/\rho$ does not depend on the normalisation. `Revision/SPEC.md` section 11 writes $\rho_4 \sim \rho_8\, c^3$ with c the extra-time scale factor: with $c = e^{-a_4} b_4^{1/2}$ that is definition (B).

3.2 How every verdict depends on the assumption

The rho_4 normalisation is an ASSUMPTION about the observer, not a consequence of the field equations, and this record does not choose one. The same state reads differently under each (history $\beta = 4/3$):

state (records)	w_{eff} **under (A), (B)**	w_{eff} **under (C)**	**ratio** p_3/ρ
homogeneous condensate (`condensate_w_eff_b4`)	$(2/3)\, w_c$; 0 for $\lambda = 0$	$-1/3 + (2/3)\, w_c$	$w_c = \lambda S/(2m + \lambda S)$, drifting
massless quanta (`massless_mode_w_eff`)	1/3, radiation	0	1/3
massive quanta, flat limit (`massive_mode_w_eff_law`)	$k^2/(3(M^2a^2 + k^2))$: $1/3 \to 0$	$0 \to -1/3$	as (A), (B) for $q = 0$
Kohn-Sham gas, $N = 688$, $\lambda = 0$ (section 5.2)	0.2929 to 0.3183	-0.0404 to -0.0150	0.2929 to 0.3183

INTERPRETATION (labelled): supernova distances measure the dilution-inferred w_{eff} of a 4-dimensional Friedmann observer, which agrees with the ratio p_3/ρ only when no energy is exchanged with the extra times. Whether (A), (B), (C) or none of them describes such an observer is not decided by any equation of this record. Under (A) the extra times are closed time-like directions; under (B) and (C) they are non-compact and a normalisation is a choice.

3.3 The expansion-inferred equation of state

An observer who reads $a(t) = e^{a_4(x_4)}$ with $t = x_4$ ($g_{44} = -1$) and 4-dimensional Friedmann equations infers $w_{\text{exp}} = -1 - \frac{2}{3}\, a_4''/a_4'^2$ (record `expansion_inferred_w`); b_4 does not enter the observer's scale factor. In Einstein gravity $G^{x_1}{}_{x_1} - G^{x_5}{}_{x_5} = 2a_4'' + 3a_4'u - v/2 - u^2/4$ with $u = b_4'/b_4$, $v = b_4''/b_4$ (computed from the metric and equal to the record; `einstein_space_minus_extra_time_b4`), so $w_{\text{exp}} = -1-(\kappa(p_3-p_t)-3a_4'u+v/2+u^2/4)/(3a_4'^2)$ for an admissible homogeneous source (`expansion_inferred_w_einstein_b4`). On the history $a_4 = Hx_4$ the observer's Hubble rate $a_4' = H$ is constant and $w_{\text{exp}} = -1$ exactly, with no time variation, for every $b_4(x_4)$.

4 Method

4.1 Exact derivations first

Every effective formula is derived exactly (sympy) for the metric with $b_4(x_4)$ before any number is computed: `derive_effective.py` for dirac16complex (39 checks, `Revision/dark_sector/dirac16complex/reports/derivation-checks.json`) and `derive_eos.py` (implementation A) for dirac16complex00 (49 checks, `Revision/dark_sector/dirac16complex00/reports/python-derive-eos.json`). Both re-derive the conservation identities from the author's metric with their own Christoffel symbols; the dirac16complex00 side uses the author's gammas T16 of `Revision/algebra/gammas.json` (record `gammas_clifford_author_T16`). The key formulas are verified a second time in Wolfram from the author's metric, the author's Lovelock tensors and the author's T16 (`verify_dark_sector_b4.wls`, 8 checks, `Revision/dark_sector/reports/wolfram-dark-sector-b4.json`). Labels: EXACT means verified by exact symbolic computation in a named check; APPROXIMATION, ASSUMPTION and INTERPRETATION are marked where they apply.

4.2 The conservation identities as checks of the numerics

The exact identities of section 3.1 test every numerical history. For the Kohn-Sham gas: $dE/da_4 = -3X - 2P_t$ ($\beta = 4/3$) with fourth-order differences of E against the solver's energy-momentum integrals, maximal relative deviation 6.351e-06 (the truncation error of the stencil on the $N = 8$ interacting series, whose E falls like e^{-2a_4}, is about 6.7e-06; record `conservation_with_b4_term`); the b_4-free identity of the withdrawn record fails on the interacting series, by 1.630e+00 (`conservation_without_b4_term_fails`): the b_4 term is required by the solver's own numbers; integrated with Simpson's rule, 3.585e-07 (`conservation_integrated_simpson`); dw_{eff}/da_4 two ways, 1.366e-06 for the gas and 3.532e-05 at the end slices of the $N = 8$ series (one-sided stencils; `derivative_two_ways`); the hidden-direction identity $p_{0,y} + 6Hp_0 = 3H(p_3 + p_t)$ in every state, 1.951e-08 (`y_conservation_every_state`). For dirac16complex00 the independent implementation B checks

along each run that the Krein charge per coordinate volume $b_4^{3/2}Q$ is conserved (drift at most 1.2612134e-13), that the solution is on shell ($|L_0|$ at most 1.3322676e-15) and that the conservation identity with the b_4 term holds (relative residual at most 4.7810141e-08 by finite differences; records `B_run_*_invariants`).

4.3 The Kohn-Sham history (dirac16complex)

In the metric with b_4 the good-sector state is $\Psi = e^{ik\cdot x}W^{-3}b_4^{-3/4}\chi$, and χ solves a slice Hamiltonian that does not contain b_4 (the exact reduction in `Revision/kohn_sham/ks-theory.json`); b_4 enters through the proper 7-volume of the cell, $\ell^3 v_t b_4^{3/2}$. The Kohn-Sham record (`Revision/kohn_sham`, moved to this history on 2026-10-16) works with $\hat{\lambda} = \lambda\, b_4(0)^{-3/2}$, recalibrated on the history ($\lambda_1 = 0.01946, 0.01206, 0.006645$ and $\lambda_2 = 0.05838, 0.03617, 0.01993$ for $N = 8, 136, 688$), and the cell volume $\mathrm{Vol}_7(a_4) = \ell^3 v_t e^{2a_4}$, so the effective coupling falls like e^{-2a_4}: the state is that of the fixed coordinate cell at $\lambda_{\text{eff}} = \hat{\lambda}\, e^{-2a_4}$ (the coupling-volume scaling in `ks-theory.json`); the integrals E, P_3, P_t, P_0 are the physical ones. `run_ks_history.py` runs the Revision Rust Kohn-Sham solver (`Revision/kohn_sham/solver`, its single-state command, read-only use) at the 41 slices $a_4 = 0, 0.05, \ldots, 2$ for $N = 8, 136, 688$ and the five calibrated couplings $\hat{\lambda} = 0, \pm\lambda_1, \pm\lambda_2$ per N: 615 runs (15 series x 41 slices), with $H = m = 1$, $L = 3$, tip $\theta = 0$, $T = 0$ (records `all_runs_succeeded`, `cell_volume_follows_b4`); the coordinate-cell form at λ_{eff} gives the same energies and pressures to 1.969e-13 in the 48 interacting states at the committed slices $a_4 > 0$ (`lambda_eff_form_agrees`). The history is a PRESCRIBED BACKGROUND: the gas is a test field without back-reaction, and the states are instantaneous (adiabatic) ground states. At the five committed slices the 15 series equal the committed Kohn-Sham record, maximal relative deviation 0.000e+00 (`committed_slices_equal_committed_record`); the $\lambda = 0$ series do not feel b_4 (`lambda0_series_independent_of_b4`); the occupied label set is the same at all 41 slices of every series, so the instantaneous ground states are the adiabatically continued states (`occupied_labels_fixed_along_history`); the particle number holds to 2.331e-15 (`particle_number_every_state`). `compute_eos.py` evaluates w_{eff} for the three definitions with the b_4 terms, the ratios, CPL tangents and fits, the condensate, mixtures and the Unite comparison (15 checks).

An INDEPENDENT second implementation, `independent_free_gas.py` (numpy; Chebyshev collocation with $n = 96$ and Hellmann-Feynman level derivatives; no solver output used except for the final comparison), treats the free gas ($\lambda = 0$, which does not feel b_4; `free_gas_independent_of_b4`). It reproduces the exact $k = 0$ spectra to 1.421e-14 (`exact_k0_spectra`), is converged ($n = 96$ against 128: 6.01e-15 in E, `collocation_converged`), reproduces the solver's $E(a_4)$ over 82 states to 1.897e-12 (`energy_vs_rust_solver`), $w_{\text{eff}}(A) = X/E$ to 1.492e-12 (`w_eff_vs_primary`) and the CPL tangents to 4.027e-07 (`cpl_tangent_vs_primary`).

4.4 Condensates

The homogeneous condensate is an exact solution for both fields and every b_4: $b_4^{3/2}S$ is constant, $\rho = mS + \lambda S^2/2$ and $p_3 = p_t = p_0 = \lambda S^2/2$ (records `condensate_rho_p`, `condensate_b4_scaling`, `condensate_rho_p_w`). For dirac16complex00 it solves $d_4\Phi = \mathcal{A}\Phi$ with $\mathcal{A} = -\gamma^{(x_4)}(M - 3H\gamma^{(x_0)}) - \frac{3}{4}(b_4'/b_4)$, $M = m + U'(S)$; $C\mathcal{A} + \mathcal{A}^T C = -\frac{3}{2}(b_4'/b_4)C$ for the author's T16, so $b_4^{3/2}S$ is constant, and $K_{x_4} = MS$ exactly (records `condensate_S_b4_T16`, `condensate_K4_equals_MS_T16`, `D4_condensate_b4_T16`, with the negative control `condensate_negative_control`). The

ratio drifts: $dw/da_4 = -3\beta m\lambda S/(2m+\lambda S)^2$ (`condensate_ratio_time_dependence_b4`, `condensate_observer_b4`, `D8_condensate_drift`).

4.5 Modes and Krein-signed energies (dirac16complex00)

For a plane wave with physical 3-momentum k and extra-time momentum q in the frozen local frame, $h^2 = (m^2 + k^2 - q^2)I_{16}$, so $\omega^2 = m^2 + k^2 - q^2$: the extra times are time-like and q enters with the opposite sign (`mode_dispersion_h_squared`). Bh is Hermitian, so the charge density $Q = \Phi^\dagger B\Phi$ is the Krein form of the frozen-frame evolution and is conserved by it (`mode_generator_B_selfadjoint`). On the real-frequency eigenspaces of the test cases of section 6.2, $T^\mu{}_\nu = k^\mu k_\nu Q/\omega$ exactly, so $\rho = \omega Q$: the energy has the sign of ωQ. In the deflating field the physical extra-time momentum is $q\, e^{a_4} b_4^{-1/2} \propto a^{1-\beta/2}$ (`extra_time_momentum_blueshift_b4`), $a^{1/3}$ on the history.

4.6 Gases, mixtures and wave packets

dirac16complex00: adiabatic (WKB) gases of plane-wave modes in the deflating local model, an APPROXIMATION: the warp $\sin^{1/6} z$ and $\tan z$ are frozen at a fixed hidden position (wavelengths short compared with $1/H$), the Krein charge of each mode per coordinate volume is an adiabatic invariant, and the superposition is incoherent. Each component has $\rho_i = \omega_i Q_i b_4^{-3/2}$ and $\varepsilon_i = w_{\text{eff}}(B)_i = (k_p^2 + (1-\beta/2)q_p^2)/(3\omega_i^2) \geq 0$ for $\beta < 2$, independent of the sign of Q (records `wkb_mode_gas_conservation_b4`, `wkb_epsilon_sign_free_b4`); a mixture has $w_{\text{eff}}(B) = \sum \varepsilon_i\rho_i / \sum \rho_i$ and $w_{\text{eff}}(C) = w_{\text{eff}}(B) - 1 + \beta/2$ (`mixture_weighted_average_b4`). Implementation B integrates the full 16-component field equation of the local model with the author's T16, including the b_4 term $\frac{3}{4}(b_4'/b_4)\gamma^{(x_4)}$ (exact Krein-preserving exponential midpoint step times the exact factor of the b_4 term, $H/m = 10^{-3}$, adiabatic), and rebuilds the models from its solutions; spatial averages of superpositions equal the sums of the mode contributions (`B_superposition_average`). dirac16complex: mixtures of the Kohn-Sham gas with a condensate, $w_{\text{eff}} = \sum_i X_i / \sum_i E_i - s + (\beta/2)(\sum_i P_{t,i} / \sum_i E_i + s)$ (`mixture_general_b4`); the kinetic gas of flat-limit modes satisfies the identity with b_4 (`flat_mode_gas_conservation_b4`, `D5_kinetic_gas_b4`).

4.7 Phantom conditions

dirac16complex: $w_{\text{eff}}(C) + 1 = X/E + (\beta/2)(1 + P_t/E)$; with $\beta > 0$ and $P_t/E \geq -1$ a value below -1 needs $X/E < 0$; with $E > 0$ this needs $P_t > P_3$, which no real-frequency mode supplies (per mode $X = (k_p^2 + q_p^2)/(3\omega) \geq 0$, with $p_3 \geq 0$ and $p_t \leq 0$), otherwise $E < 0$ together with $X > 0$ (records `phantom_condition_b4`, `flat_mode_X_nonnegative`). Negative E is not specific to a negative-norm (Krein) sector in the canonical Kohn-Sham record: there the interacting $N = 8$, $\lambda > 0$ good-sector states have $E < 0$, with $X = 0$ exactly (section 5.2), from the interaction energy of the canonical uniform-gas exchange functional ($E = 0$ at $\lambda = 0$). With the exact Fock exchange in place of that functional, solved self-consistently (the exact-Fock variant of the Kohn-Sham record, `Revision/kohn_sham/results/exx/exact-fock-variant.csv`), the same $N = 8$ states have $|E| \leq 2.12 \times 10^{-13}$ (computed at the canonical slices $a_4 = 0, 0.5, 1, 1.5, 2$, at the rounding level at each; with b_4 the canonical E falls like e^{-2a_4} between the slices); the exact exchange evaluated to first order on the canonical orbitals still leaves $E < 0$, of smaller magnitude. This negative E is therefore a property of the approximate functional; good-sector states with $E < 0$ are not

established beyond it. For free modes $E < 0$ needs a negative Krein charge, outside the positive good-sector realisation (OPEN). dirac16complex00 (EXACT within the WKB model): if every component has $\rho_i \geq 0$ and a real frequency, then $w_{\text{eff}}(B) \geq 0$ and $w_{\text{eff}}(C) \geq -1 + \beta/2 = -1/3$ at every a at which these conditions hold (an extra-time mode loses its real frequency past its turning point a_*, section 6.3); going below $-1/3$, and in particular crossing -1, needs a component of negative classical energy. A condensate ratio is phantom exactly for $\lambda S/m \in (-2, -1)$ (`condensate_phantom_interval`), at an instant: on the history $\lambda S/m$ falls like e^{-2a_4}.

5 Results for dirac16complex

Records: `Revision/dark_sector/dirac16complex/reports/` and `Revision/dark_sector/dirac16complex/outputs/eos-summary.json`.

5.1 The condensate

$X = 0$ and $P_t/E = w_c$, so $w_{\text{eff}}(A) = w_{\text{eff}}(B) = (\beta/2)w_c$ and $w_{\text{eff}}(C) = -1 + (\beta/2)(1 + w_c)$, with the ratio $w_c = \lambda S/(2m + \lambda S)$ drifting because $b_4^{3/2}S$ is constant (`condensate_w_eff_b4`, `condensate_ratio_w`, `condensate_b4_scaling`). The ratio equals the Unite constant $w = -0.764$ at $\lambda S/m = -382/441 = -0.866213151927$, with $\rho/(mS) = 0.566893424036$, so $\rho > 0$ if and only if $mS > 0$ (`condensate_ratio_equal_unite_constant_w`, `condensate_ratio_value`). This value of $\lambda S/m$ is CHOSEN to give -0.764: one parameter tuned to one number. On the history it holds at one instant only: $\lambda S/m$ falls like e^{-2a_4}, and at that value the ratio has $w_a = -2.695392$ (thawing sign); $w_{\text{eff}}(A) = (2/3)w_c = -0.509333$ and $w_{\text{eff}}(C) = -1/3 + (2/3)w_c = -0.842667$, both with $w_a = -1.796928$ (`condensate_drifts_with_b4`). For $\lambda S/m < 0$ the condensate energy $mS_0(1 + u/2)$ reaches 0 in the past.

5.2 The Kohn-Sham gas

- The occupied levels lie on the brane band, which is massless at $k = 0$ (the brane zero mode at $\varepsilon = 0$, slope $c\,e^{-a_4}$ with $c = 1.9051482536$ at small k; `brane_band_slope`). The gas therefore redshifts like radiation, not like dust: $d\ln E/da_4$ goes from -0.8787 at $a_4 = 0$ to -0.9549 at $a_4 = 2$ ($N = 688$, $\lambda = 0$), toward the radiation value -1.
- Over all $N = 136,688$ series and all slices, $w_{\text{eff}}(A) = X/E + (2/3)P_t/E$ lies in [0.292793, 0.326802] and rises monotonically toward 1/3 (`gas_radiation_like_band`, `gas_X_over_E_rises_toward_one_third`).
- $N = 688$, $\lambda = 0$: $w_{\text{eff}}(A) = w_{\text{eff}}(B)$ goes from 0.2929 to 0.3183 and $w_{\text{eff}}(C)$ from -0.0404 to -0.0150. The ratio $w_3 = P_3/E$ equals X/E up to $|P_t/E| \leq 0.002$ over the $N = 136,688$ series (P_t is the integrated interaction part; it vanishes at $\lambda = 0$). The hidden-direction ratio $w_{x_0} = P_0/E$ goes from 0.3535 to 0.6482: a pressure that the 3-space observer does not see as pressure.
- CPL tangents ($N = 688$, $\lambda = 0$):

$a_{4,\text{today}}$	**0.5**	**1**	**1.5**	**2**
w_0 under (C)	-0.0370	-0.0315	-0.0240	-0.0150
w_a (every definition)	-0.00905	-0.01292	-0.01701	-0.01775

(w_0 under (A), (B) is $w_0(C)+1/3$.) Least-squares fits at $a_{4,\text{today}} = 2$: over $a \in [1/2, 1]$, $w_a = -0.02424$ and constant proxy -0.0205 under (C); over $[1/3, 1]$, $w_a = -0.02648$ and -0.0229.

- Over all $N = 136,688$ series: tangent w_a in [-0.019170, -0.008787], fitted w_a in [-0.028228, -0.019171], $w_{\text{eff}}(C)$ in [-0.040541, -0.006532] (`gas_cpl_thawing_sign_small`, `gas_C_near_zero_on_b4_history`): the thawing sign, with magnitudes far below the Unite 0.60, and $w_{\text{eff}}(C)$ close to 0, dust-like, not a dark-energy value (the withdrawn $b_4 = 1$ record had -0.707 to -0.672).
- $N = 8$ with $\lambda \neq 0$ (only the $k = 0$ brane zero modes, pure interaction energy at $\hat{\lambda}e^{-2a_4}$): $P_3 = P_t$ ($X = 0$); $E(2)/E(0)$ in [0.018402, 0.019031] ($e^{-4} = 0.018316$); P_t/E in [0.931316, 0.999997], so $w_{\text{eff}}(A) = (2/3)P_t/E$ in [0.620877, 0.666665] and $w_{\text{eff}}(C) = w_{\text{eff}}(A) - 1/3$; $E < 0$ for $\lambda > 0$ (`n8_interacting_zero_modes_dilute_with_b4`; a property of the canonical uniform-gas exchange functional, section 4.7). $N = 8$ with $\lambda = 0$ has $E = 0$: no equation of state.

5.3 The bulk band (independent implementation)

The massive bulk band (odd brane parity, $\varepsilon(0) = 1.292292828069$ = the bulk edge) shows the dark-matter-like law: for the lowest odd-parity level of the $n_2 = 1$ shell, $w_{\text{eff}}(A)$ falls monotonically from 0.239626 at $a_4 = -3$ to 2.268e-03 at $a_4 = 4$, asymptotically like $1/a$ (the ratio $w(4)/w(3.75) = 0.765177$ against $e^{-0.25} = 0.778801$: a linear term of $\varepsilon(k)$ at small k); the brane-band level of the same shell stays in [0.291594, 0.333275], tending to 1/3 (records `bulk_band_dark_matter_law`, `brane_band_radiation_law`; free gas, $\lambda = 0$, which does not feel b_4). The canonical $T = 0$ states ($N \leq 688$) are filled below the bulk edge and do not populate that band.

5.4 Mixtures with a condensate

For a radiation-like gas ($X = E/3$, $P_t = 0$, independent of b_4) plus a $\lambda = 0$ condensate ($E = mS_0$, constant for every b_4), $w_{\text{eff}}(A) = \frac{1}{3}\, r/(1+r)$ with $r = r_0/a$ and $w_{\text{eff}}(C) = w_{\text{eff}}(A) - 1 + \beta/2$ (`mixture_radiation_condensate_cpl_b4`); $w_{\text{eff}}(C) = -0.861$ at $a = 1$ needs β in (-0.388667, 0.278000], so on the history ($\beta = 4/3$) no such mixture exists: there $w_{\text{eff}}(C)$ lies in $[-1/3, 0)$ (`mixture_C_matching_w0_unite_b4`). With the computed gas ($N = 688$, $\lambda = 0$, $a_{4,\text{today}} = 2$):

- with a $\lambda = 0$ condensate, $w_{\text{eff}}(C) \geq -0.333015$ at every slice and gas fraction: neither -0.861 nor a constant -0.764 is reached (`mixture_C_cannot_reach_unite_w0_with_lambda0_condensate`; on the withdrawn record a CHOSEN gas fraction gave $w_{\text{eff}}(C) = -0.861$ with a freezing slope);
- with a condensate of any ratio $w_c \in [-3, 0.99]$ today, drifting with b_4, any gas fraction and $a_{4,\text{today}} \in \{0.5, 1, 1.5, 2\}$, the closest points to the Unite pair $(-0.861, -0.60)$ are

$(-0.3, -0.78)$ in the ratio definition (distance 0.5891697548245), $(-0.2733333333333, -0.7708)$ under (A), (B) (distance 0.6119842735815) and $(-0.58, -0.6758666666667)$ under (C) (distance 0.2910614215438, the pure condensate with $w_c = -0.37$ today); every mixture with $|w_0 + 0.861| \leq 0.1$ has w_a between -2.639515126434 and -1.41320438962 under (C) (between -3.800209088911 and -2.685081787855 in the ratio definition, between -4.848176262726 and -3.262703276443 under (A), (B)): all below -1, far from the Unite -0.60 (`mixture_scan_three_definitions_b4`).

5.5 Phantom and crossing of -1

No computed Kohn-Sham state crosses $w = -1$: every state with $E \neq 0$ has $X \geq 0$ and $P_t/E \geq -1$, so $w_{\text{eff}}(C) \geq -1 + \beta/2$ for $E > 0$. The exception is not a Kohn-Sham state: a condensate has $X = 0$ and $P_t/E = w_c$, so $w_{\text{eff}}(C) = -1/3 + (2/3)w_c$ is below -1 exactly while its ratio is, $\lambda S/m \in (-2, -1)$ (`condensate_phantom_interval`, `condensate_observer_b4`); with $\lambda < 0$ and $mS > 0$ (positive energy, attractive self-interaction) the drift $\lambda S/m \propto e^{-2a_4}$ carries it into that window toward the past, for $\lambda S/m = -382/441$ today from $a = \sqrt{382/441} = 0.9307$ back to $a = \sqrt{191/441} = 0.6581$, where its energy vanishes. The $N = 8$, $\lambda > 0$ states have $E < 0$ (from the canonical uniform-gas exchange functional; section 4.7), $X = 0$ and P_t/E in [0.931316, 0.999997], so $w_{\text{eff}}(C) = (2/3)P_t/E - 1/3$ between 0.2875 and 0.3333; the 41 $N = 8$, $\lambda = 0$ states have $E = 0$ and no equation of state (section 5.2). On the history the expansion reads $w_{\text{exp}} = -1$ exactly (section 3.3).

6 Results for dirac16complex00

Records:

- the reports in `Revision/dark_sector/dirac16complex00/reports/`;
- implementation A: `Revision/dark_sector/dirac16complex00/eos-theory.json`;
- the independent implementation B, the field-equation runs: `Revision/dark_sector/dirac16complex00/results/independent-numerics.json`.

6.1 The condensate

As for dirac16complex: $b_4^{3/2} S$ constant, $w_{\text{eff}}(B) = (2/3)w$, $w_{\text{eff}}(C) = -1/3 + (2/3)w$, the ratio $w = \lambda S/(2m + \lambda S)$ drifting (`condensate_observer_b4`). The ratio equals -0.764 at $\lambda S/m = -382/441$ (CHOSEN), which with $\rho > 0$ needs $mS > 0$ and $\lambda < 0$, a negative potential U (`condensate_ratio_equals_unite_constant_w`); there its w_a is $-84231/31250 = -2.695392$ (`condensate_ratio_drift_at_unite_value`); it is phantom exactly while $\lambda S/m \in (-2, -1)$, and then $w_{\text{eff}}(C) < -1$ as well; the drift carries a $\lambda < 0$ condensate into that window toward the past (section 5.5). The condensate is not the source of the history: the author's equations need the constant source of section 2.4, while S decays like $b_4^{-3/2} \propto e^{-2Hx_4}$ (`condensate_cannot_source_branch`); in Einstein gravity a $b_4(x_4)$ needs $T^{x_4}{}_{x_0} \neq 0$, which a condensate does not have. The observer's expansion reads $w_{\text{exp}} = -1$ exactly on $a_4 = Hx_4$.

6.2 Modes, Krein-signed energies and growing modes

For the three test cases $(m, k, q) = (3, 4, 0)$, $(5, 0, 3)$ and $(4, 4, 4)$, with $\omega = \pm 5$, ± 4 and ± 4, every real-frequency eigenspace has dimension 8 and a Krein form of signature (4,4), and on it $\rho = \omega Q$ exactly (records `mode_krein_inertia_*`, `mode_emt_*`, negative control `mode_emt_negative_control`): at every real frequency there are modes of both energy signs. Implementation B confirms it: at $\omega = 1.25$ the vectors with $Q = +1$ and $Q = -1$ have $\rho = +1.25$ and -1.25 (`B_krein_signed_energy`). The general statement is the Krein-inertia theorem of `Revision/pairing`.

For $q^2 > m^2 + k^2$ the modes grow: for $(m, k, q) = (3, 0, 5)$, $\omega = 4i$; the growing eigenspace is Krein-neutral ($Q = 0$, $\rho = 0$) with a growing extra-time pressure $p_5 = -mS$ (`growing_mode_krein_neutral`, `growing_mode_extra_time_pressure`). In the deflating field the physical extra-time momentum $q\, a\, b_4^{-1/2} \propto a^{1/3}$ grows, so every mode with $q \neq 0$ reaches a turning point a_* and then grows: the deflation drives the growth. Implementation B, past the turning point $a_* = 6.2639848$ of the mode of M3: growth rate $d \ln |\phi| / dx_4 = 0.30893327$ against the WKB rate $\sqrt{s a^{2/3} - 1} - \frac{3}{4} b_4' / b_4 = 0.31165344$ (`B_growth_rate`); the amplitude grows by a factor $10^{16.20}$ between $a = 5.5$ and 7.4, and the energy density becomes large and negative (ρ/Q = -4.3193205e+29 at $a = 7.4$, for $H/m = 10^{-3}$).

6.3 The models M1 to M5

Each model is normalised to $\rho_{\rm tot} = 1$ at $a = 1$; definition (C) unless stated. The models keep the physical parameters of the withdrawn $b_4 = 1$ record, where they were CHOSEN to reproduce Unite numbers; on the history with $b_4(x_4)$ they no longer reproduce them.

model	**content**	**parameters**
M1	condensate (exact solution for every a_4, b_4)	$\lambda S/m$ free; the ratio is -0.764 at $\lambda S/m = -382/441$ at one instant (CHOSEN)
M2	positive-energy good-sector gas, $q = 0$	$k^2/(k^2 + m^2) = 417/1000$ at $a = 1$, CHOSEN on the withdrawn record for $w_0 = -0.861$
M3	positive-energy extra-time mode, $k = 0$	$s = q_p^2/m^2 = 417/1417$, CHOSEN on the withdrawn record for $w_0 = -0.861$
M4	condensate + positive extra-time mode	$s = 264037/403037$ and $\Omega_q = 57963/264037$, both CHOSEN on the withdrawn record to solve tangent $= (-0.861, -0.60)$
M5	M4-type + GHOST: massless modes of negative classical energy	$G = 3/10$ of the total at $a = 1$ (a stated choice); s and Ω_q CHOSEN on the withdrawn record so that its fit over $[1/2, 1]$ was $(-0.861, -0.60)$

The resulting CPL numbers under (C) on the history (tangent; least-squares fit over $[1/2, 1]$; constant proxy):

model	**tangent** w_0	**tangent** w_a	**fit** w_0	**fit** w_a	**proxy**	**crosses** -1 **on** $[1/3,1]$**?**
M1	$-1/3+2w/3$	$8u/(3(2+u)^2)$	drifts	drifts	drifts	no
M2	-0.1943	+0.162074	-0.1988	0.2173	-0.1445	no
M3	-0.2870	-0.043770	-0.2871	-0.0420	-0.2976	no
M4	-0.2870	-0.066667	-0.2882	-0.0502	-0.3007	no
M5	-0.3465	-0.1964	-0.3327	-0.3231	-0.4135	no

M2 is freezing ($w_a > 0$), M3, M4 and M5 have the thawing sign ($w_a < 0$); none reaches the Unite pair or crosses -1 on $[1/3, 1]$; the column says nothing about $a > 1$, and past the turning point a_* of an extra-time mode (M3: 6.2640, M4: 1.8859, M5: 2.3375) the frequency of that mode is imaginary and the bound of section 4.7 does not apply. On the withdrawn record the M4 tangent and the M5 fit equal the Unite pair by construction. Records:

record	**record**	**record**
`M2_tangent_exact_b4`	`M3_tangent_exact_b4`	`M4_withdrawn_tuning_on_b4_history`
`positive_models_bounded_by_minus_one_third`	`unite_w0_requires_dominant_ghost`	`M5_withdrawn_composition_on_b4_history`
`unite_line_fit_proxy`	`unite_crossing_point`	`condensate_ratio_drift_at_unite_value`

Notes:

- M2 under (B) and in the ratio: tangent $(0.139, 0.162074)$, the law $w_{\text{eff}}(B) = \frac{1}{3}k^2/(k^2 + m^2a^2)$ from 1/3 to 0, independent of b_4: a time-varying dark-matter-like equation of state. Exactly, under (C), $(w_0, w_a) = (-583/3000, 81037/500000)$.
- M3: $\omega^2 = m^2(1 - sa^{2/3})$ and $w_{\text{eff}}(C) = -\frac{1}{3} + \frac{1}{9}\, sa^{2/3}/(1 - sa^{2/3}) \geq -1/3$ for $a < a_*$; exactly $(w_0, w_a) = (-287/1000, -196963/4500000)$; the turning point is at $a_* = (1417/417)^{3/2} = 6.2640$, in the future (withdrawn record: $(-0.861, -0.393926)$, $a_* = 1.843$).
- M4: on the history its $w_{\text{eff}}(C)$ has the minimum -0.33280216468 over $a = 1/300, \ldots, 1$; it reaches its turning point at $a_* = 1.8859$ and then grows.
- M5: $s = 0.5678$, $\Omega_q = 0.5947$, condensate fraction $\Omega_c = 0.7053$; on the history no crossing of -1 on $[1/3, 1]$ (the withdrawn record had one, at $a = 0.779$); its turning point is at $a_* = 2.3375$.
- The positive models are bounded: the minima of $w_{\text{eff}}(C)$ over $a = 1/300, \ldots, 1$ are -0.194333333333, -0.332598868774 and -0.33280216468 for M2, M3, M4, all $\geq -1/3$ (`positive_models_bounded_by_minus_one_third`).
- The Unite w_0 needs a dominant ghost: a ghost component lowers $w_{\text{eff}}(C)(1)$ by at most $G/3$ and positive components raise it, so $w_0 = -0.861$ needs $G \geq 1583/1000$; for condensate + ghost the total energy then vanishes at $a = G/(1 + G) = 1583/2583$ inside $[1/2, 1]$, where w_{eff} diverges (`unite_w0_requires_dominant_ghost`).

6.4 The independent implementation B

B rebuilds M2 to M5 from solutions of the 16-component field equation with the b_4 term; all tangents, fits and constant proxies agree with implementation A to within 2.2666232e-05 (records `B_vs_A_*`). B also measures $w_{\text{eff}}(C)$ directly from the dilution of the proper density of its solutions, independently of the formula of section 3.1; the two agree to within 4.0240551e-05 (`B_dilution_equals_formula_*`). The condensate dilutes with $b_4^{3/2}$: $b_4^{3/2}\rho$ is constant to 1.2610746e-13 while ρ changes by a relative range 0.91990032 over $a \in [0.3, 1.06]$ (`B_condensate_dilutes_with_b4`). M5 has no crossing of -1 (`B_vs_A_M5_crossing`). By bisection on the field-equation solutions B finds $s = 0.29429162$ for M3 (A: $417/1417 = 0.2942837$; `B_solve_M3_s`), and the positive models have $w_{\text{eff}}(C) \geq$ -0.31820563 on $a \in [0.35, 1]$ (`B_positive_models_bounded`).

7 Comparison with the Unite values

Unite value	**dirac16complex**	**dirac16complex00**	**by construction?**
constant $w = -0.764$	condensate ratio at $\lambda S/m = -382/441$ at one instant (8-dimensional ratio, drifting with $w_a = -2.695392$)	condensate ratio at $\lambda S/m = -382/441$, $\lambda < 0$, at one instant	yes: one parameter for one number
$w_0 = -0.861$	not reached with a $\lambda = 0$ condensate ($w_{\text{eff}}(C) \geq -0.333015$); a drifting condensate under (C) reaches it, with w_a between -2.639515126434 and -1.41320438962	not reached by positive models ($w_{\text{eff}}(C) \geq -1/3$); needs a ghost fraction $\geq 1583/1000$	the condensate value is chosen; no positive model reaches it
$(w_0, w_a) = (-0.861, -0.60)$	not reached: closest distance 0.2910614215438 under (C)	not reached; on the withdrawn record M4 (tangent) and M5 (fit) reached it by construction	on the withdrawn record: two parameters for two numbers
phantom past, crossing of -1 at $a = 0.7683$	no crossing in any Kohn-Sham state; a drifting $\lambda < 0$ condensate crosses at $a = 0.9307$ (for $-382/441$ today)	no crossing in M2 to M5; M1 ($\lambda < 0$) as for dirac16complex	the condensate value is chosen; the ghost fraction is a stated choice

A model with as many tuned parameters as matched numbers reproduces them by construction and is NOT a prediction. No parameter of M2 to M5, of the mixtures or of the condensate ratio is selected by the field equations; nothing in this record selects their populations. A ghost (negative-energy) component is a ghost-like sector, not an established physical state: dirac16complex00 has one at every real frequency, and its classical energy is unbounded below (the classical form of the spin-statistics problem of a commuting field with a first-order Lagrangian).

What is not tuned: the signs and the ranges. For positive-energy content with real frequencies $w_{\text{eff}}(C) \geq -1/3$ on the history in the WKB model of dirac16complex00 (section 4.7; for M3 and M4 before the turning point a_* of the extra-time mode), and every computed dirac16complex state has $X \geq 0$ (section 5.5); the computed dirac16complex gas has the thawing sign with $|w_a| \leq 0.026521$ and $w_{\text{eff}}(C)$ close to 0.

8 The lead's analysis (SPEC section 11), checked point by point

1. "The 7-volume of the metric is constant." CORRECTED for the metric with b_4: the proper 7-volume element is $b_4^{3/2} \cos z$, independent of a_4 but changing with x_4 through b_4 (`proper_7_volume_element_b4`, `local_model_volume_b4`); it was constant only on the withdrawn $b_4 = 1$ metric.

2. "The effective 4-dimensional density is $\rho_4 \sim \rho_8 c^3$." With $c = e^{-a_4} b_4^{1/2}$ this is definition (B) (with (A) of the same scaling). It is an ASSUMPTION about the observer: definition (C), per unit proper 7-volume, is equally consistent with the field equations and shifts every w_{eff} by exactly $-1 + \beta/2$, $-1/3$ on the history (section 3).

3. "Non-relativistic quanta give dust, relativistic quanta radiation." CONFIRMED under (A), (B) in the flat (warp-free) limit for every b_4: a massive mode with $q = 0$ has $w_{\text{eff}} = k^2/(3(M^2a^2 + k^2))$, $1/3 \to 0$, and a massless one $1/3$ (`massive_mode_w_eff_law`, `massless_mode_w_eff`, `massive_mode_cpl_tangent`); the Kohn-Sham gas has $d \ln E/da_4$ approaching -1 (section 5.2).

4. "A Kohn-Sham gas whose 3-momenta redshift has a time-varying equation of state from $1/3$ towards 0." NOT FOUND in the computed states: the occupied levels lie on the massless brane band and $w_{\text{eff}}(A)$ RISES toward $1/3$ (section 5.2). The falling law is present for quanta of the massive bulk band (section 5.3), which the computed ground states do not populate. For dirac16complex00 the falling law holds for a positive-energy gas of good-sector modes (M2). The hidden-direction pressure ($w_{x_0} = 0.3535$ to 0.6482) and the extra-time pressure were computed; the 3-space observer does not see them as pressure.

5. "A homogeneous condensate has constant w while $\rho_4 \sim a^{-3}$; the ratio and $w_{\text{eff}} = 0$ disagree; 4-dimensional energy is exchanged with the extra dimensions." CORRECTED for the metric with b_4: $b_4^{3/2} S$ is constant, so the condensate's ratio drifts and, for $\lambda \neq 0$, $w_{\text{eff}}(B) = (\beta/2)w \neq 0$ (`condensate_w_eff_b4`, `condensate_observer_b4`); for $\lambda = 0$, $w_{\text{eff}}(B) = 0$ and $w_{\text{eff}}(C) = -1/3$. The exchange reading is an INTERPRETATION of the disagreement between the ratio and the dilution.

6. "Whether any state gives a time-varying dark-energy equation of state near the Unite values must be computed." COMPUTED: for dirac16complex no computed state comes near the Unite pair in any definition (section 5); for dirac16complex00 no positive-energy model gives dark-energy-like values on the history (sections 6.3 and 7).

7. "Crossing $w = -1$ needs negative kinetic energy, which the indefinite (Krein) energy of the extra-time sector and the commuting field dirac16complex00 may supply." EXAMINED. Positive-energy extra-time momentum does NOT supply it: its pressure $p_t \leq 0$ increases X, so it moves w_{eff} up, not down (`flat_mode_X_nonnegative`, `wkb_epsilon_sign_free_b4`; M3

has $w \geq -1/3$ before its turning point $a_* = 6.2640$). What is needed is negative classical ENERGY: dirac16complex00 has it at every real frequency (Krein signature (4,4)), and on the history it would have to exceed the positive energy (section 6.3), as a ghost-like sector. For dirac16complex no computed state has $X < 0$; the only states with $E < 0$ ($N = 8$, $\lambda > 0$, negative with the canonical uniform-gas exchange functional; with the exact Fock exchange solved self-consistently $|E| \leq 2.04 \times 10^{-13}$, section 4.7) have $X = 0$ exactly. Modes of dirac16complex with extra-time momentum lie outside the good sector and are OPEN.

9 Verdicts

9.1 dirac16complex (Hypothesis)

- Dark matter. Under (A), (B): a time-varying dark-matter-like equation of state, w_{eff} falling from 1/3 toward 0, is present in the theory for quanta of the massive bulk band (0.239626 at $a_4 = -3$ to 2.268e-03 at $a_4 = 4$ for one shell; the flat-limit law is exact and independent of b_4). It is NOT present in the computed Kohn-Sham ground states, whose brane-band gas is radiation-like with w_{eff} rising toward 1/3. A $\lambda = 0$ condensate is dust-like ($w_{\text{eff}}(B) = 0$).

- Dark energy. Not found on the history with $b_4(x_4)$: under (C) the gas has $w_{\text{eff}}(C)$ in [-0.040541, -0.006532] with a thawing-sign slope (tangent w_a in [-0.019170, -0.008787], fitted in [-0.028228, -0.019171]); only a drifting condensate gives negative values, with w_a near -1.8 to -2.7; the Unite constant $w = -0.764$ is matched only by the 8-dimensional ratio of a tuned condensate at one instant. No computed state crosses $w = -1$.

- On the history the observer's expansion reads $w_{\text{exp}} = -1$ exactly.

9.2 dirac16complex00 (Hypothesis00)

- Dark matter. Under (B) and in the ratio p_3/ρ: a positive-energy gas of good-sector modes has the time-varying law from 1/3 to 0 (M2), independent of b_4; a $\lambda = 0$ condensate is dust-like.

- Dark energy without ghosts. Not on the history with $b_4(x_4)$: positive-energy content with real frequencies has $w_{\text{eff}}(C) \geq -1/3$ (section 4.7; M2: -0.1943, M3 and M4: -0.2870 today); for M3 and M4 this holds before the turning point a_* of the extra-time mode (M3: 6.2640, M4: 1.8859). The thawing slope of the extra-time mode is small (M3: -0.043770) because its physical momentum grows only like $a^{1/3}$; past a_* the mode grows without bound and its energy density becomes large and negative (implementation B).

- Phantom and crossing of -1. Impossible with positive-energy components of real frequency (section 4.7); the Unite w_0 alone needs, from such components, a component of negative classical energy larger than the positive energy ($G \geq 1583/1000$), a ghost-like sector. The exception is a condensate with $\lambda < 0$ and $mS > 0$: while its drifting ratio lies in the window $\lambda S/m \in (-2, -1)$ the observer reads $w_{\text{eff}}(C) = -1/3 + (2/3)w_c < -1$ (section 5.5); its value is CHOSEN and no source of the history selects it.

- The source of the history: the author's equations need the constant source $\kappa(\rho, p_3, p_t, p_0) = (-2/27, -2/9, -22/27, -14/27)$ (a negative energy density for $\kappa > 0$); no condensate is it.

9.3 The hypotheses

This record establishes neither Hypothesis nor Hypothesis00. What it shows, under its stated assumptions, on the author's metric with $b_4(x_4)$: both fields contain time-varying equations of state of the dark-matter type under definition (B) (dirac16complex only for bulk-band quanta, which the computed states do not populate); no dark-energy-like value arises from positive-energy content under any definition; the Unite numbers are not reproduced, and on the withdrawn $b_4 = 1$ record they were reproduced only by models whose parameters were chosen to reproduce them. Nothing in this record establishes that either field is dark energy or dark matter.

10 What remains open

1. A self-consistent history with an admissible source. No recorded Kohn-Sham state is an admissible source of the author's metric: its energy-momentum tensor depends on x_0 and violates the algebraic condition of the field equations, and the history is a prescribed background (the 6 checks of `Revision/field_equations_a4/reports/ks-source-conditions.json`, among them `ks_profiles_depend_on_x0`, which holds for every a_4 and b_4, `ks_profiles_violate_algebraic_condition` and `ks_history_is_a_prescribed_background`). The source that the author's equations require on the history (section 2.4) is not constructed from the fields of this record; a condensate is not it.
2. Populated bulk-band or thermal states along the history (fixed entropy); only $T = 0$ ground states and $a_4 \in [0, 2]$ (the solver's validated range) are computed; $b_4(0)$ enters only through $\hat{\lambda}$.
3. Modes of dirac16complex with extra-time momentum (outside the good sector; only their flat-limit sign structure is derived) and the non-adiabatic (time-dependent) Kohn-Sham problem.
4. For dirac16complex00: configurations that depend on x_0 are not consistent sources of the metric; the WKB model freezes the hidden-direction warp; the growing modes make the Cauchy problem ill-posed, and nothing here cures that; no quantum treatment (a classical field).
5. Which normalisation of ρ_4, if any, describes a physical 3-space observer; the value of $a_{4,\mathrm{today}}$.
6. What would select the populations of M2 to M5, the mixture fractions or the condensate's $\lambda S/m$.
7. A supernova likelihood (distances, covariances): no fit to data is part of this record.
8. Histories with $\beta < 0$, on which the b_4 term lowers w_{eff}; no source of this record selects one.

11 Verification records

11.1 Reports and their counts

Every check has a name, a verdict and a detail. Counts at the time of writing (the publication test re-reads them):

report	checks	PASS	FAIL
Revision/dark_sector/dirac16complex/reports/derivation-checks.json	39	39	0
Revision/dark_sector/dirac16complex/reports/ks-history-run.json	8	8	0
Revision/dark_sector/dirac16complex/reports/eos-checks.json	15	15	0
Revision/dark_sector/dirac16complex/reports/independent-checks.json	10	10	0
Revision/dark_sector/dirac16complex00/reports/python-derive-eos.json	49	49	0
Revision/dark_sector/dirac16complex00/reports/python-independent-numerics.json	32	32	0
Revision/dark_sector/reports/wolfram-dark-sector-b4.json	8	8	0
Revision/field_equations_a4/reports/ks-source-conditions.json	6	6	0

11.2 Checks per report

The tables list every check of the seven dark-sector reports; every one has the verdict PASS.

dirac16complex, derive_effective.py (39 checks):

check	check	check
sqrt_det_g_equals_b4_three_halves_cos_z	proper_7_volume_element_b4	proper_3_and_extra_time_volume_scalings_b4
conservation_x4_identity_b4	conservation_x0_identity	conservation_other_components_vanish

check	check	check
conservation_equ	hidden_coordinat	conservation_x0_
als_a4_record_b4	e_dy_dx0	in_y_form
integrated_ident	w_eff_A_b4	w_eff_B_equals_w
ity_dE_dx4_b4		_eff_A_b4
w_eff_C_b4	w_eff_general_no	w_eff_difference
	rmaliser_b4	_to_withdrawn_record
condensate_rho_p	condensate_b4_scaling	condensate_ratio_w
condensate_ratio	condensate_w_eff_b4	condensate_ratio
_time_dependence_b4		_equal_unite_constant_w
flat_mode_mass_shell_b4	flat_mode_gas_co	flat_mode_X_nonnegative
	nservation_b4	
massive_mode_w_eff_law	massive_mode_cpl	massless_mode_w_eff
	_tangent	
extra_time_momen	mixture_general_b4	mixture_radiatio
tum_blueshift_b4		n_condensate_cpl_b4
mixture_C_matchi	cpl_least_squares_rule	unite_deep_past
ng_w0_unite_b4		
expansion_inferred_w	einstein_mixed_c	einstein_space_m
	omponent_b4	inus_extra_time_b4
expansion_inferr	author_weights_b	phantom_condition_b4
ed_w_einstein_b4	4_branch	

dirac16complex, run__ks__history.py (8 checks):

check	check	check
all_runs_succeeded	committed_slices	lambda0_series_i
	_equal_committed_record	ndependent_of_b4
cell_volume_follows_b4	lambda_eff_form_agrees	occupied_labels_
		fixed_along_history
y_conservation_e	particle_number_	
very_state	every_state	

dirac16complex, compute__eos.py (15 checks):

check	check	check
formulas_input_present	dense_history_complete	conservation_wit
		h_b4_term
conservation_wit	conservation_int	derivative_two_ways
hout_b4_term_fails	egrated_simpson	
gas_radiation_like_band	gas_X_over_E_ris	gas_C_near_zero_
	es_toward_one_third	on_b4_history
n8_interacting_z	condensate_ratio_value	condensate_drift
ero_modes_dilute		s_with_b4
_with_b4		
mixture_C_cannot	mixture_scan_thr	gas_cpl_thawing_
_reach_unite_w0_	ee_definitions_b4	sign_small
with_lambda0_condensate		

dirac16complex, independent__free__gas.py (10 checks):

check	check	check
exact_k0_spectra	brane_band_slope	collocation_converged
aufbau_closed_shells	energy_vs_rust_solver	w_eff_vs_primary
cpl_tangent_vs_primary	bulk_band_dark_m atter_law	brane_band_radia tion_law
free_gas_indepen dent_of_b4		

dirac16complex00, derive_eos.py (A) (49 checks):

check	check	check
gammas_clifford_ author_T16	conservation_neg ative_control	conservation_x4_ author_metric_b4
conservation_x0_ author_metric	conservation_oth er_components_vanish	conservation_x4_ local_model_b4
local_model_volume_b4	observer_N1_weff _identity_b4	observer_N2_weff _identity_b4
observer_hidden_ average_commutes	condensate_S_b4_T16	condensate_negat ive_control
condensate_K4_eq uals_MS_T16	condensate_rho_p_w	condensate_observer_b4
condensate_phant om_interval	condensate_ratio _equals_unite_co nstant_w	condensate_ratio _drift_at_unite_value
author_weights_v acuum_recovered	branch_required_source	condensate_canno t_source_branch
mode_dispersion_ h_squared	mode_generator_B _selfadjoint	mode_emt_negativ e_control
mode_krein_inert ia_m3_k4_q0_pos	mode_emt_m3_k4_q0_pos	mode_krein_inert ia_m3_k4_q0_neg
mode_emt_m3_k4_q0_neg	mode_krein_inert ia_m5_k0_q3_pos	mode_emt_m5_k0_q3_pos
mode_krein_inert ia_m5_k0_q3_neg	mode_emt_m5_k0_q3_neg	mode_krein_inert ia_m4_k4_q4_pos
mode_emt_m4_k4_q4_pos	mode_krein_inert ia_m4_k4_q4_neg	mode_emt_m4_k4_q4_neg
growing_mode_kre in_neutral	growing_mode_ext ra_time_pressure	wkb_mode_gas_con servation_b4
wkb_epsilon_sign _free_b4	mixture_weighted _average_b4	unite_line_fit_proxy
unite_crossing_point	M2_tangent_exact_b4	M3_tangent_exact_b4
M4_withdrawn_tun ing_on_b4_history	positive_models_ bounded_by_minus _one_third	unite_w0_require s_dominant_ghost
M5_withdrawn_com position_on_b4_history		

dirac16complex00, independent_numerics.py (B) (32 checks):

check	check	check
B_clifford_numeric	B_krein_matrix	B_krein_signed_energy
B_superposition_average	B_run_kmode_invariants	B_run_q3_invariants
B_run_q4_invariants	B_run_q5_invariants	B_run_cond_invariants
B_run_ghost_invariants	B_ghost_negative_energy	B_condensate_dil utes_with_b4
B_vs_A_M2_N2	B_vs_A_M2_N1	B_vs_A_M2_ratio
B_dilution_equal s_formula_M2	B_vs_A_M3_N2	B_vs_A_M3_N1
B_vs_A_M3_ratio	B_dilution_equal s_formula_M3	B_vs_A_M4_N2
B_vs_A_M4_N1	B_vs_A_M4_ratio	B_dilution_equal s_formula_M4
B_vs_A_M5_N2	B_vs_A_M5_N1	B_vs_A_M5_ratio
B_dilution_equal s_formula_M5	B_vs_A_M5_crossing	B_solve_M3_s
B_positive_model s_bounded	B_growth_rate	

dark sector, verify_dark_sector_b4.wls (Wolfram) (8 checks):

check	check	check
D1_volume_b4	D2_conservation_b4	D3_w_eff_general_b4
D4_condensate_b4_T16	D5_kinetic_gas_b4	D6_author_weight s_branch
D7_branch_requir ed_source	D8_condensate_drift	

11.3 Re-verification for this document

For this document (2026-10-15) the seven dark-sector scripts were run in the order of section 12 on the committed inputs (`Revision/kohn_sham/results`, `Revision/algebra/gammas.json`, `Revision/field_equations_a4/a4-equations.json`, `Revision/metric_b4/results`) with the built Revision Rust solver: 39 of 39, 7 of 7, 15 of 15, 10 of 10, 49 of 49, 32 of 32 and 8 of 8 checks passed, and every output and report file was byte-identical over two runs. On 2026-10-16, after the Kohn-Sham record itself was moved to the history with b_4 (`Revision/dark_sector/B4_PROVENANCE.md`, section 10), the seven scripts were run again in the same order: 39 of 39, 8 of 8, 15 of 15, 10 of 10, 49 of 49, 32 of 32 and 8 of 8; `run_ks_history.py` and `compute_eos.py` twice with identical outputs (the notebook reproduces 123 rows of the second run character for character). Only the interacting gas numbers changed (the ranges quoted in sections 1, 5.2 and 9.1, the derivative identity of section 4.2); `independent-free-gas.json` changed only in the last digit of two copied primary values. The Kohn-Sham source-conditions report (6 of 6) is the committed file. The reports cited in this document are the committed files.

12 Reproduction

From the repository root, in this order. Each script exits 0 only if all its checks pass; its outputs are LF and two runs are byte-identical. The dirac16complex scripts write into their own `outputs/` and `reports/` folders, and each reads the outputs of the scripts before it; `run_ks_history.py` needs the built Rust solver (`-solver PATH` selects another binary). The dirac16complex00 scripts accept `-out DIR`, and `independent_numerics.py` reads implementation A's `eos-theory.json` for its comparison; the Wolfram verifier reads `eos-theory.json` for its comparison of the required source. The second unittest command and the PDF command are each one command; the backslash is the line continuation of a POSIX shell. Python needs numpy and sympy (no scipy).

```
cargo build --release --manifest-path Revision/kohn_sham/solver/Cargo.toml
python Revision/dark_sector/dirac16complex/derive/derive_effective.py
python Revision/dark_sector/dirac16complex/compute/run_ks_history.py
python Revision/dark_sector/dirac16complex/compute/compute_eos.py
python Revision/dark_sector/dirac16complex/independent/independent_free_gas.py
python Revision/dark_sector/dirac16complex00/python/derive_eos.py
python Revision/dark_sector/dirac16complex00/python/independent_numerics.py
wolframscript -file Revision/dark_sector/wolfram/verify_dark_sector_b4.wls
python -m unittest \
    Revision/dark_sector/dirac16complex00/tests/test_dark_sector_dirac16complex00.py -v
python scripts/build_provenance_pdf.py Revision/docs/DARK_SECTOR_HYPOTHESES.md \
    --developer-layout --specifications Revision/pdf-specifications.json
python -m unittest Revision/tests/test_dark_sector_hypotheses_publication.py -v
```

Expected output and run times measured for this document (2026-10-16, a development machine otherwise idle):

command	last line of the output	time
`derive_effective.py`	39/39 checks pass	3.4 s
`run_ks_history.py`	PASS - particle_number_every_state (the eighth of 8 PASS lines)	46 s
`compute_eos.py`	15/15 checks pass	0.3 s
`independent_free_gas.py`	10/10 checks pass	75 s
`derive_eos.py`	derive_eos: 49/49 checks pass	3.6 s
`independent_numerics.py`	independent_numerics: 32/32 checks pass	6.7 s
`verify_dark_sector_b4.wls`	8/8 checks passed	4.6 s
`test_dark_sector_dirac16complex00.py`	OK (4 tests)	11 s
`build_provenance_pdf.py`	provenance_pdf=OK	5 s
`test_dark_sector_hypotheses_publication.py`	OK (skipped=1); with REVISION_PDF_REBUILD=1: OK	below 1 s; 5 s

The cargo build was not re-timed (the solver was already built). The PDF build runs the Markdown-to-LaTeX builder twice (with different hash seeds), runs pdflatex three times into each of two fresh

directories, requires warning-free logs and byte-identical PDFs, and checks the PDF against its entry `dark-sector-hypotheses` in `Revision/pdf-specifications.json` (the Revision registry; the registry of the earlier stages is not touched).

13 What is proved, computed, assumed and not established

13.1 Proved (exact, by symbolic computation in the named checks)

1. For the author's metric with $b_4(x_4)$: the proper 7-volume element $b_4^{3/2}\cos z$; the conservation identities $d\rho/dx_4 = -3a_4'(p_3 - p_t) - \frac{3}{2}(b_4'/b_4)(\rho + p_t)$ and $dp_0/dx_0 + 3H\cot z(2p_0 - p_3 - p_t) = 0$ for diagonal sources $T(x_0, x_4)$ (sympy and Wolfram).

2. The observer identity $w_{\text{eff}} = X/E - s + (\beta/2)(P_t/E + s)$ for the normalisations (A), (B) ($s = 0$) and (C) ($s = 1$), and its N1/N2 form for dirac16complex00.

3. The condensate: an exact homogeneous solution of both fields with $b_4^{3/2}S$ constant, its drifting ratio, $w_{\text{eff}}(B) = (\beta/2)w$ and $w_{\text{eff}}(C) = -1 + (\beta/2)(1 + w)$; the ratio -0.764 at $\lambda S/m = -382/441$ at an instant; the phantom window $(-2, -1)$.

4. The history: with the author's weights a source without ${T^{x_4}}_{x_0}$ allows $d\ln b_4/da_4 = 4/3$ on $a_4 = Hx_4$, and there needs the constant source $\kappa(\rho, p_3, p_t, p_0) = (-2/27, -2/9, -22/27, -14/27)$ (two independent computations); no condensate is it; in Einstein gravity a $b_4(x_4)$ needs ${T^{x_4}}_{x_0} \neq 0$.

5. The flat-limit mode laws (massive $1/3 \to 0$ independent of b_4, massless $1/3$, $X \geq 0$ for real frequencies) and the expansion-inferred w_{exp}, -1 exactly on $a_4 = Hx_4$ for every b_4.

6. dirac16complex00: on the three test eigenspaces, dimension 8, Krein signature (4,4) and $\rho = \omega Q$; the growing eigenspace is Krein-neutral; within the WKB model, positive-energy components of real frequency give $w_{\text{eff}}(B) \geq 0$ and $w_{\text{eff}}(C) \geq -1 + \beta/2$; the exact tangents of M2 and M3 on the history; the ghost fraction 1583/1000 needed for the Unite w_0.

13.2 Computed (numerical, with the stated tolerances and two implementations)

The Kohn-Sham gas along the history with $b_4(x_4)$ (615 states at $\hat{\lambda}$ in the cell growing like e^{2a_4}, i.e. at $\lambda_{\text{eff}} = \hat{\lambda}e^{-2a_4}$ in the fixed cell; radiation-like, $w_{\text{eff}}(A)$ in [0.292793, 0.326802] rising toward 1/3; $w_{\text{eff}}(C)$ in [-0.040541, -0.006532]), with the conservation identity including the b_4 term; its CPL tangents and fits, the mixtures and the scans; the bulk-band law (independent implementation); the models M2 to M5 on the history; implementation B of dirac16complex00 (field-equation runs with the b_4 term, agreement with A to within 2.2666232e-05, w_{eff} from the dilution of the proper density, the growth past the turning point).

13.3 Assumed

1. The normalisation (A), (B) or (C) of ρ_4: an ASSUMPTION about the observer; every verdict is stated for each.

2. The history $a_4 = Hx_4$, $b_4 = b_4(0)e^{(4/3)Hx_4}$ as a PRESCRIBED BACKGROUND (test fields, no back-reaction); the instantaneous (adiabatic) Kohn-Sham states; $a_{4,\text{today}}$; $b_4(0)$ through $\hat{\lambda}$.

3. The Z2 brane at $z = \pi/2$ (ASSUMED) and the filling convention of `Revision/kohn_sham/ks-theory.json`. The canonical uniform-gas exchange functional of the Kohn-Sham states is an APPROXIMATION; with the exact Fock exchange solved self-consistently the $N = 8$, $\lambda > 0$ states have $|E| \leq 2.04 \times 10^{-13}$ (section 4.7).

4. The WKB/local model of dirac16complex00 (frozen warp, adiabatic Krein charges, incoherent superposition): an APPROXIMATION, measured by implementation B.

5. The populations, fractions and condensate parameters of every model: CHOSEN (those of M2 to M5 on the withdrawn record to reproduce the Unite numbers there).

13.4 Not established

1. A dark-energy or dark-matter mechanism of either field: this record establishes neither Hypothesis nor Hypothesis00, and nothing in this record establishes that either field is dark energy or dark matter.

2. Any Unite value as an output of the field equations: no positive-energy content reaches them on the history, and every match of the withdrawn record was by construction (sections 6.3 and 7).

3. That the ghost-like sector of dirac16complex00 is a physical state; that a time-varying dark-matter equation of state arises in the computed dirac16complex states; that any computed state crosses $w = -1$; that any good-sector state of dirac16complex has $E < 0$ beyond the canonical uniform-gas exchange functional (section 4.7).

4. A fit to supernova data; a self-consistent cosmological history; a source of the fields of this record for the history with $b_4(x_4)$; the open problems of section 10.