

Kohn-Sham model of dirac16complex in the author's primordial field with the deflating extra times

The block reduction, the instantaneous ground and first excited states along the prescribed history $a_4 = AHx_4$, Mermin thermodynamics, adiabaticity, the energy-momentum tensor, the rescaling partners and the source conditions, with the verification records and an exact statement of what is computed and what is not

dirac16complex project (with Claude Opus 5.5)

September 2026

Contents

1	The task and the answer	3
2	Setting: the author's field, the good sector and the hidden coordinate	4
3	The exact block reduction	5
4	The functional: Hartree, exact uniform-gas exchange, Mermin	6
5	Boundary conditions, filling convention, numerics and parameters	7
6	Instantaneous ground and first excited states along the prescribed history	9
7	Adiabaticity and Fermi-level crossings	10
8	Mermin thermodynamics	12
9	Energy-momentum tensor profiles	13
10	The rescaling partners	14
11	The recorded states are not admissible sources of the a_4 equations	15
12	The Kohn-Sham-level pairing T3	16

13 Verification records	16
13.1 Reports and their counts	16
13.2 The independent reference and the full cross-check	18
13.3 Determinism and the refinement measurement	18
13.4 Re-verification for this document	19
14 Reproduction	19
15 What is proved, computed, assumed and not established	20
15.1 Proved (exact, two independent verifiers)	20
15.2 Computed (numerical, two independent solvers, cross-checked)	21
15.3 Assumed or chosen	21
15.4 Not established	21

Abstract

This document records the Kohn-Sham (density-functional) model of the fermion field `dirac16complex` in the author's primordial gravitational field, deliverable 4 of `Revision/SPEC.md` section 10, as computed by the code in `Revision/kohn_sham/`. Exact (two independent verifiers, WolframScript and sympy): in the good sector (no extra-time momentum) the ansatz $\Psi = e^{ik \cdot x} W^{-3} \chi(y, x_4)$ removes the spin-connection term $3H\gamma^8$ and reduces the 16-component equation, for every history $a_4(x_4)$, to eight 2×2 blocks $h_j = j[-i\sigma_1 d/dy + M_{\text{eff}}\sigma_2 + \kappa k\sigma_3] + v_v$ in the hidden coordinate y , with $\kappa = e^{-Hy - a_4}$; the deflation of the extra times enters the instantaneous problem only through this redshift of the 3-momenta and through the extra-time volume; the exact local exchange of the uniform good-sector gas gives $M_{\text{eff}} = m + \frac{15}{16}\lambda S$ and $v_v = -\lambda n/16$. Computed (a Rust solver, an independent Python reference with a different discretisation, and a cross-check of the full canonical matrix, 31 of 31 checks PASS over 162691 comparisons): the instantaneous ground and first excited states of 75 parameter sets ($N = 8, 136, 688$; five couplings; five slices $a_{4,0}$ of the prescribed history $a_4 = AHx_4$, $A = 1$), 135 Mermin states at three temperatures, the adiabaticity measure ($Q_{\text{max}} \leq 0.09345324$ over the matrix), the Fermi-level crossings (none in the canonical matrix; a demonstration at $N = 696$), the energy-momentum tensor profiles, the exact rescaling partners and the exact-exchange variant. ASSUMED or chosen: the Z2 brane at the patch end, the regular tip at $y = -3$, the filling convention. Not admissible as sources: no recorded Kohn-Sham state satisfies the source conditions of the a_4 equations, so the history is a prescribed background without back-reaction. Not done: the time-dependent (non-adiabatic) Kohn-Sham problem, correlation beyond exchange, back-reaction.

1 The task and the answer

The author asked (2026-10-01, quoted from `Revision/README.md`): “plan and employ a DFT-motivated approximation similar to the one that you already created [...], with a Kohn-Sham fermion-gas thermodynamic effective potential, that employs the DFT ground and first-excited-state computation”. `Revision/SPEC.md` section 7 makes this precise: a Kohn-Sham fermion-gas model of `dirac16complex` in the good sector with the contact interaction $U = \frac{\lambda}{2}S^2$, the Hartree mean field plus the exact local exchange of the uniform gas, Mermin's finite-temperature functional, the reduction to 2×2 blocks in the hidden coordinate, the boundary conditions, the ground and first excited states, the thermodynamics, and, because a_4 depends on x_4 , the INSTANTANEOUS (adiabatic) Kohn-Sham states along the history with the extra times deflating, with an adiabaticity measure.

The answer of this record:

1. Exact (section 3, section 4): the block reduction, the exchange functional, the boundary conditions, the rescaling identity, the energy-momentum tensor and its conservation laws, the adiabaticity measure. Every formula is in `Revision/kohn_sham/ks-theory.json` and verified twice, in `Revision/kohn_sham/reports/ks-theory-wolfram.json` (46 of 46 PASS) and independently in `Revision/kohn_sham/reports/ks-theory-python.json` (58 of 58 PASS).
2. Computed (sections 5 to 10): the instantaneous Kohn-Sham states of the canonical matrix by the Rust solver, independently by the Python reference, and compared state by state on the full matrix.
3. Not admissible (section 11): the recorded states are not sources of the field equations for a_4 ; the history $a_4 = AHx_4$ is PRESCRIBED.
4. Open (section 15): the time-dependent Kohn-Sham problem, correlation, back-reaction.

Every number below is read from the named Revision report or results file (the publication test re-reads them); units: $H = m = 1$, energies, momenta and temperatures in units of m , as in `Revision/kohn_sham/results/parameters.json`. Interpretations are labelled.

2 Setting: the author's field, the good sector and the hidden coordinate

The coordinates are named as the author names them: x_1, x_2, x_3 are ordinary 3-space, inflating with the scale factor $e^{a_4} \sin^{1/6} z$; x_4 is the time; x_5, x_6, x_7 are the three extra times, which DEFLATE exponentially, with the scale factor $e^{-a_4} \sin^{1/6} z$ and a_4 increasing (they are never treated as static); x_8 is the hidden space direction, $z = 6Hx_8 \in (0, \pi/2)$. The metric of `Revision/SPEC.md` section 1 is

$$ds^2 = e^{2a_4(x_4)} \sin^{1/3} z (dx_1^2 + dx_2^2 + dx_3^2) - dx_4^2 \\ - e^{-2a_4(x_4)} \sin^{1/3} z (dx_5^2 + dx_6^2 + dx_7^2) + \cot^2 z dx_8^2.$$

The Kohn-Sham record uses the hidden coordinate $y = \ln(\sin z)/(6H) \in (-\infty, 0]$, with $dy = \cot z dx_8$; the patch end $z = \pi/2$ is $y = 0$ (the brane) and the tip $z \rightarrow 0$ is $y \rightarrow -\infty$. In it the metric is exactly

$$ds^2 = e^{2Hy} [e^{2a_4}(dx_1^2 + dx_2^2 + dx_3^2) - e^{-2a_4}(dx_5^2 + dx_6^2 + dx_7^2)] - dx_4^2 + dy^2,$$

with the warp $W = e^{Hy}$ and $\sqrt{|g|} = e^{6Hy}$ ($= \cos z$ in the author's chart). The 7-volume is independent of a_4 : the inflation e^{3a_4} of 3-space is compensated exactly by the deflation e^{-3a_4} of the extra times. The canonical spin connection gives $\gamma^\mu \Omega_\mu = 3H\gamma^8$ for every $a_4(x_4)$: the a'_4 pieces of the three inflating directions, $+\frac{3}{2}a'_4\gamma^4$, and of the three deflating directions, $-\frac{3}{2}a'_4\gamma^4$, cancel.

The good sector: no dependence on x_5, x_6, x_7 (zero extra-time momenta), a coordinate 3-torus of side ℓ with the momenta $k \in \Delta k \mathbb{Z}^3$, $\Delta k = 2\pi/\ell$, and the extra-time coordinate volume v_t ; $\text{Vol}_7 = \ell^3 v_t$. The canonical quantisation of `Revision/SPEC.md` section 6 has a positive Fock realisation in this sector, and the expectation-value rule $\langle \Psi^\dagger M \Psi \rangle = \text{Tr}(M\rho)$, $\rho = \sum_{\text{occ}} f uu^\dagger B$, defines the quasi-free (Slater or Mermin) states used here.

item	Wolfram records	sympy records
gammas, C , B , Γ	<code>fixture_input</code> , <code>clifford_relation</code> , <code>C_B_Gamma</code>	<code>fixture_input</code> , <code>fixture_python_equals_wolfram</code> , <code>clifford_relation</code> , <code>C_B_Gamma</code> , <code>BC_and_Cg4</code>
hidden coordinate, $\sqrt{ g }$	<code>geometry_hidden_coordinate</code> , <code>geometry_sqrt_det</code>	<code>geometry_hidden_coordinate</code> , <code>geometry_sqrt_det</code> , <code>geometry_warped_form</code>

item	Wolfram records	sympy records
spin connection	spin_connection_ compatibility, spin_con nection_slash_3H, spin_connection_ time_terms_cancel, spin_connection_ slash_3H_x8_chart	spin_connection_ compatibility, spin_con nection_slash_3H, spin_connection_ time_terms_cancel, spin_connection_ time_term_value, spin_connection_ slash_3H_x8_chart

3 The exact block reduction

Ansatz. $\Psi = e^{ik \cdot x} W^{-3} \chi(y, x_4)$ gives EXACTLY, for a time-dependent a_4 ,

$$\gamma^8 \partial_y \chi + \gamma^4 \partial_{x_4} \chi + i\kappa k \cdot \gamma \chi = (M_{\text{eff}} - iv_v \gamma^4) \chi, \quad \kappa(y, x_4) = e^{-Hy - a_4(x_4)},$$

i.e. $i\partial_{x_4} \chi = h(x_4) \chi$ with h Hermitian in $\int dy \chi^\dagger \chi$. The factor W^{-3} removes the term $3H\gamma^8$; without it the term survives. At a fixed instant the instantaneous (adiabatic) problem is $h(a_{4,0}) \chi = \varepsilon \chi$ with $a_{4,0} = a_4(x_4)$.

Blocks. $J = \gamma^8 \gamma^1 \gamma^4$ ($J^2 = 1$), $K_1 = \gamma^2 \gamma^3$ and $K_2 = \gamma^5 \gamma^6$ ($K^2 = -1$) commute with each other and with γ^8 , $\gamma^8 \gamma^1$, $\gamma^8 \gamma^4$, B and C ; their joint eigenspaces, labelled (j, s_2, s_3) with $j, s_2, s_3 = \pm 1$, are eight rank-2 blocks with an explicit unitary basis V (entries of $2\sqrt{2}V$ in $\{0, \pm 1, \pm i\}$). For k along x_1 (the spinor rotation commutes with γ^8 , γ^4 , B , C , so the spectra depend on $|k|$ only), every block of type j carries

$$h_j = j \left[-i\sigma_1 \frac{d}{dy} + M_{\text{eff}}(y)\sigma_2 + \kappa(y)k\sigma_3 \right] + v_v(y),$$

equivalently $\chi' = N\chi$ with $N = M_{\text{eff}}\sigma_3 - \kappa k\sigma_2 + ij(\varepsilon - v_v)\sigma_1$; in the real form $\chi = (a, ib)$, $a' = M_{\text{eff}}a - (\kappa k + j(\varepsilon - v_v))b$ and $b' = (j(\varepsilon - v_v) - \kappa k)a - M_{\text{eff}}b$. In the block basis $\gamma^8 = \sigma_3$, $B = js_2$ and $C = s_2\sigma_2$.

Symmetries and degeneracies. $h_{-1} - v = -(h_{+1} - v)$ and $\sigma_3 h_j(k) \sigma_3 = h_{-j}(-k)$: the (j, k) and $(-j, -k)$ orbitals are degenerate. On a lattice shell $|k|^2 = \Delta k^2 n_2$ with $r_3(n_2)$ vectors every level of $h_j(|k|)$ is $4r_3(n_2)$ -fold for each j ; at $k = 0$ every level is 4-fold per j . The chirality Γ maps block (j, s_2, s_3) onto $(-j, s_2, s_3)$ with the 2×2 entry $s_2\sigma_2$, and $\sigma_2 h_j(M, k) \sigma_2 = h_{-j}(-M, k)$ with the brane parities exchanged (the ingredient of T3, section 12).

Where the deflation enters (exact). $a_{4,0}$ enters h_j only through $\kappa k = e^{-Hy}(ke^{-a_{4,0}})$: the 3-momenta redshift as ke^{-a_4} . Because the good sector has no extra-time momentum, the extra-time scale factor e^{-a_4} enters the instantaneous problem only through the extra-time volume (section 10).

item	Wolfram records	sympy records
ansatz and Hamiltonian	ansatz_removes_s pin_connection, ansatz_w ithout_W3_term_survives, hamiltonian_16_h ermitian	ansatz_removes_s pin_connection, ansatz_w ithout_W3_term_survives, hamiltonian_16_h ermitian
blocks	blocks_commuting_set, blocks_basis_unitary, blocks_forms, blocks_r elation_to_Gamma, block_hamiltonian, block_ode_equivalent	blocks_commuting_set, blocks_projectors, blocks_basis_unitary, blocks_forms, blocks_n ot_everything_bl ock_diagonal, blocks_r elation_to_Gamma, block_hamiltonian, block_ode_equivalent
symmetries	block_type_relation, block_Gamma_map, rotation_invariance	block_type_relation, block_Gamma_map, rotation_invariance
solver (numerical basis)		block_reduction_numeric, gamma_fixture_numeric (Rust solver report)

4 The functional: Hartree, exact uniform-gas exchange, Mermin

Exchange (exact for the uniform gas). The interaction is $U = \frac{\lambda}{2}S^2$, $S = \bar{\Psi}\Psi$, normal ordered ($\lambda > 0$ repulsive). For a quasi-free state, Wick's theorem gives $\langle : S^2 : \rangle = (\text{Tr } C\rho)^2 - \text{Tr}(C\rho C\rho)$. For every $p \rightarrow -p$ symmetric occupation of the 8-fold good-sector gas, at every temperature, $\rho = (nB + SC)/16$ and

$$e_H = \frac{\lambda}{2}S^2, \quad e_x = -\frac{\lambda}{32}(n^2 + S^2), \quad e_{\text{int}} = \frac{15}{32}\lambda S^2 - \frac{1}{32}\lambda n^2,$$

$$M_{\text{eff}}(y) = m + \frac{15}{16}\lambda S(y), \quad v_v(y) = -\frac{\lambda}{16}n(y),$$

with proper densities; for one filled level at rest $E_x/E_H = -1/8$, and on shell $\langle \mathcal{L} \rangle / \sqrt{|g|} = e_{\text{int}}$. No correlation is included (Hartree plus exchange only). The Rust solver reads the coefficients from `ks-theory.json` and checks them (`theory_input_coefficients`: $M_{\text{eff}} = m + 0.9375\lambda S$, $v_v = -0.0625\lambda n$).

Exact Fock exchange of the slab (diagnostic and variant). For closed shells the exact Fock exchange of the Kohn-Sham determinant is also local, $e_x^{\text{exact}} = -\frac{\lambda}{32}(n^2 + S^2 - Q^2 - Y^2)$ with $Q = \langle \Psi^\dagger B \gamma^8 \Psi \rangle$ and $Y = 0$; the canonical functional is the uniform-gas one, the difference $\Delta E_x = \frac{\lambda}{32} 2\text{Vol}_7 \int e^{6Hy} Q^2 dy$ is reported for every state, and the exact-Fock functional is solved as a variant (60 states, Revision/kohn_sham/results/exx/exact-fock-variant.csv).

Densities, energy, Mermin. Per orbital, with $P(y) = e^{-6Hy}/(\ell^3 v_t)$: $n_o = P\chi^\dagger \chi$, $s_o = P\chi^\dagger j\sigma_2 \chi$, $t_o = P\chi^\dagger j\sigma_3 \chi$, $q_o = P\chi^\dagger \sigma_3 \chi$. Totals are $\sum w g f$ times these, $g = 4r_3(n_2)$, $w = 1/2$ (section 5). The Kohn-Sham energy of the doubled system is $E_{KS} = \sum g f \varepsilon - 2\text{Vol}_7 \int_{-L}^0 e^{6Hy} e_{\text{int}} dy$; Mermin's

occupations are $f = 1/(1 + e^{(\varepsilon - \mu)/T})$ with μ fixed by $\sum gf = N$, the entropy is $S_{\text{ent}} = -\sum g[f \ln f + (1 - f) \ln(1 - f)]$, the grand potential (the thermodynamic effective potential) is $\Omega = -T \sum g \ln(1 + e^{-(\varepsilon - \mu)/T}) - 2\text{Vol}_7 \int_{-L}^0 e^{6Hy} e_{\text{int}} dy$, and $F = \Omega + \mu N$.

Discrepancy recorded, not hidden: the line “mu fixed by sum w_Z2 g f = N” of `ks-theory.json` is inconsistent with its own densities, energy and grand potential; the solver and the reference use $\sum gf = N$, and `Revision/kohn_sham/results/parameters.json` records the discrepancy.

item	Wolfram records	sympy records
uniform gas and exchange	gas_mode_projector, gas_angular_average, gas_densities, exchange _uniform_gas	gas_mode_projector, gas_angular_average, gas_negative_ene rgy_modes, gas_densities, hf_wick_contraction, exchange_uniform_gas, filled_shell_ratio
Kohn-Sham potentials	ks_potentials, ks_onshe ll_lagrangian	ks_potentials, ks_onshe ll_lagrangian, ks_theor y_json_exchange
exact Fock exchange of the slab	exchange_slab_ex act_fock	exchange_slab_ex act_fock

5 Boundary conditions, filling convention, numerics and parameters

Brane (ASSUMED). The Z2 mirror (orbifold) at $y = 0$: $\Psi(-y) = \pm \gamma^8 \Psi(y)$ with the mirror warp $e^{-H|y|}$; continuity gives $(1 \mp \gamma^8) \chi(0) = 0$, i.e. the even parity $\chi_2(0) = 0$ ($b(0) = 0$) and the odd parity $\chi_1(0) = 0$ ($a(0) = 0$). Both conditions kill the y -current and make h_j self-adjoint. The mirror map P_A takes $(M(y), v(y))$ to $(-M(-y), v(-y))$: the doubled problem is P_A -symmetric exactly when the mirror copy carries $(-m, +\lambda)$. Both parities are solved and filled together; N is the particle number of the doubled system (universe and Z2 image), the patch holds $N/2$, and $w = 1/2$ per patch.

Tip (chosen). At $y = -L$: $(1 - Q(\theta)) \chi(-L) = 0$, $Q(\theta) = \cos \theta \sigma_3 + \sin \theta \sigma_2$, canonical $\theta = 0$, cutoff $L = 3$. For $k \neq 0$ an orbital is suppressed toward the tip like $\exp(-|k|(\kappa(-L) - \kappa(y))/H)$.

Exact $k = 0$ spectra. For constant $M > 0$, $v = 0$, $\theta = 0$: even parity $\varepsilon = 0$ with the brane zero mode $\chi = (e^{My}, 0)$, and $\varepsilon = \pm \sqrt{M^2 + (n\pi/L)^2}$; odd parity $\varepsilon = \pm \sqrt{M^2 + p^2}$ with $\tan(pL) = -p/M$. The brane band has the slope $d\varepsilon/dk = jc$ at $k = 0$, $c = e^{-a_{4,0}} \frac{2M}{2M-H} \frac{1 - e^{-(2M-H)L}}{1 - e^{-2ML}}$; for $M = H = 1$, $L = 3$: $c = 2/(1 + e^{-3}) = 1.905148253644866$ (record `brane_band_slope`).

Filling (CONVENTION, justification OPEN). Particles occupy the positive branch (the levels with $\varepsilon > 0$ of the $\lambda = 0$ problem) and the $k = 0$ brane zero modes, each followed continuously in λ ; the negative branch is the normal-ordered sea and contributes nothing to N , n , S or the energy-momentum tensor. The excluded thermal sea holes are reported as a diagnostic (section 8).

Numerics. The Rust solver (`Revision/kohn_sham/solver`, own crate, no external crates) solves each block by shooting with RK4 on 900 steps, Pruefer counting and a root tolerance of 1e-13, Anderson-mixed SCF to 1e-11. The independent Python reference (`Revision/kohn_sham/refer`

ence) uses a staggered finite-difference discretisation on three grids (300, 600, 1200) with Richardson extrapolation and a stated uncertainty, validated on a fourth grid; its convergence order is two (record `free_convergence_order_two`).

Parameters (Revision/kohn_sham/results/parameters.json): $H = m = 1$, $L = 3$, $\Delta k = 0.25$, $v_t = 1$, $\text{Vol}_7 = 15875.21366031351$, tip $\theta = 0$, history $A = 1$, slices $a_{4,0} \in \{0, 0.5, 1, 1.5, 2\}$, temperatures $T \in \{0.01, 0.02, 0.05\}$. Particle numbers: $N = 8$ (the $k = 0$ brane zero modes of both block types), $N = 688$ (the largest closed shell of the free $a_{4,0} = 0$ aufbau whose last level lies below the bulk edge 1.292292828068783) and $N = 136$ (the closed shell nearest 688/4). Couplings: λ_1 and λ_2 keep the first-order mean-field potential below 0.1 and 0.3 (units of m) along the whole history:

N	strength per unit lambda	lambda_1	lambda_2
8	5.138804	0.01946	0.05838
136	107.5483	0.0009298	0.002789
688	541.7126	0.0001846	0.0005538

The canonical matrix holds 75 ground states (3 values of N , the five couplings $0, \pm\lambda_1, \pm\lambda_2$, five slices), each with its first excited state, its a_4 neighbours, its rescaling partner and, for $\lambda \neq 0$, its exact-Fock variant, and 135 Mermin states ($\lambda \in \{0, \pm\lambda_1\}$, three temperatures).

item	records
boundary conditions (Wolfram and sympy)	bc_mirror_map_PA, bc_mirror_parity es_of_densities, bc_brane_parity_ conditions, bc_self_adjoint_boundary _term, bc_tip_family, bc_curre nt_conserved_along_y, bc_exact _k0_spectra, bc_tip_asymptotics, brane_band_slope
free problem (Rust solver)	free_k0_analytic_spectra, free_zer o_mode_exact, free_brane_band_slope, free_block_type_symmetries, free_par ticle_branch_labels, free_tip _angle_insensitivity
free problem and parameters (reference)	free_k0_analytic_spectra, free_con vergence_order_two, free_k0_block_ty pe_symmetry, free_zero_mode, free_bra ne_band_slope, free_particle_branch, parameters_closed_shells, paramete rs_calibration, richardson_asymp totic_ratio, richardson_uncer tainty_validated
parameters (cross-check)	problem_definition_identical, parameters_particle_numbers, parameters_couplings

6 Instantaneous ground and first excited states along the prescribed history

The history is $a_4 = AHx_4$ with $A = 1$ (the exponentially deflating member: the extra-time scale factor is $e^{-Hx_4} \sin^{1/6} z$), PRESCRIBED, not solved for (section 11). At every slice $a_{4,0}$ the ground state is the self-consistent aufbau state of the instantaneous problem; the first excited state is computed three ways: the Kohn-Sham gap $\epsilon_{\text{LUMO}} - \epsilon_{\text{HOMO}}$; the ensemble Δ -SCF energy (one particle moved from the highest occupied degenerate group to the lowest empty group, spread uniformly over each group, self-consistent); and the list of the 24 lowest particle-hole excitations (Revision/kohn_sham/results/excited/particle-hole/). Every ground state of the matrix is closed-shell.

Ground and first excited states for $N = 136$ (Revision/kohn_sham/results/ground/summary.csv and Revision/kohn_sham/results/excited/summary.csv):

lambda	a4,0	E_KS	KS gap	Delta-SCF
0	0	80.28222	0.08267377	0.08267377
0	1	32.38412	0.03498894	0.03498894
0	2	12.44507	0.01445687	0.01445687
0.0009298	0	80.2837	0.08267139	0.08267139
0.0009298	1	32.39295	0.03497394	0.03497395
0.0009298	2	12.44706	0.01447131	0.01447131
-0.0009298	0	80.28074	0.08267616	0.08267615
-0.0009298	1	32.37559	0.03500291	0.0350029
-0.0009298	2	12.44277	0.01444421	0.01444422
0.002789	0	80.28666	0.08266659	0.0826666
0.002789	1	32.41162	0.03494039	0.03494042
0.002789	2	12.449	0.01450788	0.01450787
-0.002789	0	80.27779	0.0826809	0.0826809
-0.002789	1	32.35937	0.03502814	0.03502812
-0.002789	2	12.43783	0.01442269	0.01442269

The free Kohn-Sham gap along the history ($\lambda = 0$, ground/summary.csv):

N	a4,0 = 0	a4,0 = 0.5	a4,0 = 1	a4,0 = 1.5	a4,0 = 2
8	0.4307337	0.2726449	0.1703493	0.1050152	0.06415941
136	0.08267377	0.05383771	0.03498894	0.0226437	0.01445687
688	0.04518029	0.03145504	0.02048367	0.01330982	0.008610228

Reading (labelled interpretation of these numbers): as the extra times deflate, the 3-momenta redshift as $ke^{-a_{4,0}}$, the brane band flattens (its slope at $k = 0$ is c , proportional to $e^{-a_{4,0}}$, section 5), the Kohn-Sham gap falls for every N and E_{KS} falls for $N = 136$ and 688 ; for $N = 8$ (the $k = 0$ zero modes, on which $\kappa k = 0$) E_{KS} does not change with $a_{4,0}$. With $\lambda = 0$ the Δ -SCF energy equals the Kohn-Sham gap (worst difference 2.011e-11 over 15 cases, record `excited_delta_scf_free_equals_gap`); with interaction the two differ slightly, because the Δ -SCF state relaxes the mean field self-consistently.

item	Rust solver records	reference records	cross-check records
ground states	ground_runs_completed, ground_scf_converged, ground_closed_shell, ground_window_complete, ground_N_conservation, ground_energy_two_forms, ground_orbital_boundary_conditions	ground_scf_converged, ground_rid_consistency, ground_closed_shell, ground_window_complete, ground_N_conservation, ground_energy_two_forms	ground_occupations_and_groups, ground_energies, ground_homo_lumo_gap, ground_eigenvalues, ground_profiles
first excited state	excited_delta_scf_free_equals_gap, excited_scf_converged	excited_delta_scf_free_equals_gap, excited_particle_hole_lowest_is_gap	excited_delta_scf, excited_particle_hole_lists
exact-Fock variant	exx_variant_scf_converged, exx_variant_y_conservation	exx_scf_converged, exx_N_conservation, exx_energy_two_forms, exx_empt_energy_integral, exx_y_conservation_integrated	exchange_delta_Ex, exx_variant_scf

7 Adiabaticity and Fermi-level crossings

Exactly (mean field), $i\partial_{x_4}\chi = h(a_4(x_4))\chi$ conserves k , the block and the brane parity; transitions occur only between levels of the same $(k, j, \text{block}, \text{parity})$. The measure along $a_4 = AHx_4$ is

$$Q_{nm} = AH \frac{|\langle n | \partial_a h | m \rangle|}{(\varepsilon_n - \varepsilon_m)^2} = AH \frac{|\langle n | \partial_a m \rangle|}{|\varepsilon_n - \varepsilon_m|}, \quad \partial_a h_j = -j\kappa k\sigma_3 + j(\partial_a M_{\text{eff}})\sigma_2 + \partial_a v_v,$$

for n occupied and m empty in the same sector; the state is adiabatic if $\max Q_{nm} \ll 1$, and Q_{nm}^2 is the leading-order transition probability. The self-consistent derivatives come from the states at $a_{4,0} \pm \delta, \pm 2\delta$ ($\delta = 2 \times 10^{-3}$, Richardson); Hellmann-Feynman, $d\varepsilon_n/da = \langle n | \partial_a h | n \rangle$, is checked in 75 cases (worst 2.609e-9, record `adiabatic_hellmann_feynman`).

Results (`Revision/kohn_sham/results/adiabatic/adiabaticity.csv`): $Q_{\max} = 0$ exactly for $N = 8$ ($\partial_a h$ vanishes on the $k = 0$ sector), and the largest value over the whole matrix is 0.09345324 (state N688_lamm2_a00, pair 11:+1:even:0 -> 11:+1:even:1: two levels of the same shell, block type and parity). For $\lambda = 0$:

N	a4,0	Q_max	delta eps of the pair	dE/da4 (EMT)	dE/da4 (differences)
136	0	0.08514877	2.05293	-71.43976	-71.43976
136	0.5	0.07455997	1.888369	-46.47084	-46.47084
136	1	0.06361841	1.745737	-30.11922	-30.11922
136	1.5	0.05274874	1.616341	-19.33966	-19.33966
136	2	0.03839473	1.524416	-12.18614	-12.18614
688	0	0.09345059	2.253405	-597.9157	-597.9157
688	0.5	0.08526	2.055018	-388.9333	-388.9333
688	1	0.0746905	1.890122	-253.016	-253.016
688	1.5	0.06373822	1.747327	-163.9948	-163.9948
688	2	0.05289328	1.617688	-105.4063	-105.4063

The last two columns are the energy change along the history, from the energy-momentum tensor ($dE/da_4 = -6\text{Vol}_7 \int e^{6Hy}(p_3 - p_t) dy$, section 9) and from finite differences at fixed occupations.

Fermi-level crossings. When levels of different k cross the Fermi level between slices, the instantaneous aufbau ground state differs from the adiabatically continued state, and the solver flags it. No crossing occurs in the canonical matrix: every one of the 15 series of `Revision/kohn_sham/results/adiabatic/history.json` has an empty crossing list, and `adiabatic/fermi-level-crossings.csv` records no change of the occupied set. A demonstration at $\lambda = 0$ and $N = 696$ (the smallest closed shell of the $a_{4,0} = 0$ aufbau that occupies a level off the even $j = +1$ brane band; `adiabatic/crossing-demo.csv`):

a4,0	occupied set as at a4,0 = 0	open shell	E aufbau	E adiabati- cally continued	difference
0	true	false	690.7796	690.7796	0
0.5	false	true	444.1946	447.8513	3.65669
1	false	true	283.7106	289.7632	6.052645
1.5	false	true	179.4593	187.0712	7.611827
2	false	true	112.1016	120.725	8.623384

From $a_{4,0} = 0.5$ on, the $k = 0$ odd bulk level is emptied and the $n_2 = 12$ brane-band shell is filled: the instantaneous ground state is not the adiabatically reached state, which lies above it by the last column.

Interpretation (labelled): with $Q_{\max} \leq 0.09345324$ the instantaneous states are self-consistently adiabatic for the same-sector transitions at the rate $A = 1$ in the canonical matrix; this does not solve the time-dependent problem (OPEN), and where Fermi-level crossings occur (the demonstration) the instantaneous ground state is not the state that the evolution reaches.

Rust solver records	reference records	cross-check records
adiabatic_hellma nn_feynman, adiabati c_crossing_flag_ demonstration	adiabatic_hellma nn_feynman_discrete, crossing_demo_reference	adiabatic_dE_da4, adiabatic_Q_max, crossing_demonstration

8 Mermin thermodynamics

The Mermin states at $T = 0.01, 0.02, 0.05$ solve the self-consistent problem at fixed N with μ the root of $\sum gf = N$ in a well-conditioned form (thermal particles above and holes below a split of the levels, `solver/src/mermin.rs`). The checks: Ω in its two forms, $-dF/dT = S_{\text{ent}}$, $C_V = T dS_{\text{ent}}/dT = dE/dT$ at fixed N , and the chemical potential against 40-digit roots on the solver's own final levels (40-digit and 50-digit roots agree to $7.5\text{e-}35$, record `root_precision_self_check`). For $N = 136$, $\lambda = 0$ (`Revision/kohn_sham/results/thermo/thermodynamics.csv`):

a4,0	T	mu	E	entropy	Omega	sea holes / N
0	0.01	0.8339283	80.34462	7.763517	-33.14726	2.1e-56
0	0.02	0.8250861	80.74027	34.38332	-32.1591	9.49e-29
0	0.05	0.7941344	82.71202	91.45989	-29.86326	4.43e-12
2	0.01	0.1247136	12.99426	104.8261	-5.015053	1.29e-09
2	0.02	0.1150668	14.49922	205.4874	-5.259608	4.25e-05
2	0.05	0.05801333	23.38445	464.1395	-7.712339	0.112

Validity of the particle-only ensemble (diagnostic of the CONVENTION). For every thermal state the number of thermal holes that the excluded sea brane band would carry at the same μ and T is computed. It exceeds 1% of N in 15 of 135 states, all at $a_{4,0} \geq 1$; the largest is 30.976 N (record `thermo_sea_hole_diagnostic_computed`). There the particle-only Mermin ensemble is outside its range of validity and a thermal treatment of the sea (pairs) would be required; in the table above this is the row $a_{4,0} = 2$, $T = 0.05$ (state `N136_lam0_a20_T50`).

Rust solver records	reference records	cross-check records
thermo_runs_completed,	thermo_scf_converged,	thermo_state_functions,
thermo_scf_converged,	thermo_N_conservation,	thermo_derivatives,
thermo_N_conservation,	thermo_grand_pot	thermo_sea_hole_
thermo_grand_pot	ential_two_forms,	diagnostic, thermo_l
ential_two_forms,	thermo_CV_identity,	evels, thermo_mu_high_p
thermo_entropy_identity,	thermo_entropy_identity,	recision, thermo_m
thermo_CV_identity,	thermo_window_cut,	u_rounding_diagnostic,
thermo_window_cut,	thermo_sea_holes_tail	thermo_mu_rust_s
thermo_window_sh		tated_bound
ells_beyond, thermo_m		
u_well_conditioned_root,		
thermo_mu_vs_40d		
igit_roots, thermo_s		
ea_hole_diagnost		
ic_computed		

Mermin-root report	records
Revision/kohn_sham/reports/ks-rust-mermin-roots.json	solver_runs_completed, root_pre cision_self_check, solver_mu_within _rounding_bound, single_reproduce s_committed_matrix, fixture_written

9 Energy-momentum tensor profiles

With the sign convention of Revision/SPEC.md section 4 ($\rho = -T^{x_4}_{x_4}$, $p_\mu = T^\mu_\mu$) and $T^\mu_\nu = -\langle K^\mu_\nu \rangle + \delta^\mu_\nu e_{\text{int}}$, the proper profiles are

$$\begin{aligned}\rho(y) &= \sum wgf \varepsilon n_o - e_{\text{int}}, & p_3(y) &= \sum wgf \frac{\kappa|k|}{3} t_o + e_{\text{int}}, & p_t(y) &= e_{\text{int}}, \\ p_8(y) &= \sum wgf [(\varepsilon - v_v)n_o - M_{\text{eff}}s_o - \kappa|k|t_o] + e_{\text{int}},\end{aligned}$$

with $T^{x_4}_y = 0$ for eigen-orbitals. The extra-time pressure p_t has no kinetic part (good sector). Exact identities, checked for every state: the trace of the kinetic parts is $M_{\text{eff}}S + v_v n$; conservation along the hidden direction, $p'_8 + 6Hp_8 = 3H(p_3 + p_t)$; the energy change $\nabla_\mu T^\mu_{x_4} = -\partial_{x_4}\rho - 3a'_4(p_3 - p_t)$, i.e. $dE/da_4 = -6\text{Vol}_7 \int_{-L}^0 e^{6Hy} (p_3 - p_t) dy$; and $2\text{Vol}_7 \int e^{6Hy} \rho dy = E_{KS}$. Integrated profiles ($2\text{Vol}_7 \int_{-L}^0 e^{6Hy} X dy$, Revision/kohn_sham/results/ground/emt-integrals.csv), $\lambda = 0$:

N	a4,0	int rho	int p3	int p_t	int p8
136	0	80.28222	23.81325	0	35.04349
136	0.5	51.24519	15.49028	0	26.62178
136	1	32.38412	10.03974	0	19.33023
136	1.5	20.20308	6.446555	0	13.173
136	2	12.44507	4.062048	0	8.472703
688	0	680.4412	199.3052	0	240.5598
688	0.5	437.5129	129.6444	0	188.2465
688	1	279.4249	84.33868	0	143.2073
688	1.5	176.7328	54.66494	0	104.3887
688	2	110.3867	35.13543	0	71.5579

The proper profiles on 151 points $y = -3 + 0.02i$ are in Revision/kohn_sham/results/ground/profiles/ (one file per state). The energy density varies along the hidden direction in every state with a nonzero tensor (section 11). Equations of state seen by a 3-space observer are not part of this document; they are the work of Revision/dark_sector/dirac16complex (its Kohn-Sham history report Revision/dark_sector/dirac16complex/reports/ks-history-run.json: 5 of 5 PASS).

item	Wolfram and sympy records	Rust solver records	reference and cross-check records
tensor and identities	emt_orbital_components, emt_trace_identity, emt_y_conservation_orbital, emt_y_conservation_selfconsistent, emt_x4_component	emt_energy_integral, emt_trace_derivative_form, emt_y_conservation_integrated, emt_y_conservation_pointwise, emt_energy_change_dE_da4	emt_energy_integral, emt_y_conservation_integrated, emt_energy_change_dE_da4, emt_integrals, emt_brane_tip_values

10 The rescaling partners

Exact identity. Because $a_{4,0}$ enters only through $\kappa k = e^{-Hy}(ke^{-a_{4,0}})$,

$$KS(a_{4,0}; \Delta k, v_t, \lambda, N, T) = KS(0; \Delta k e^{-a_{4,0}}, v_t e^{-3a_{4,0}}, \lambda, N, T)$$

for the levels, orbitals, proper densities and energy-momentum profiles. Reading (labelled): the slice $a_{4,0}$ of the deflating history is the slice 0 with a 3-torus larger by $e^{a_{4,0}}$ in each direction and an extra-time volume smaller by $e^{-3a_{4,0}}$; the proper 7-volume is the same, and the deflation of the extra times is exactly what keeps it so. The 3-momenta redshift while the proper box stays the same.

Computed. Each solver solves the partner independently (60 partners). The worst level difference is 1.478e-13 and the worst relative difference of E_{KS} and of the profiles 3.318e-13 (records `rescaling_identity_between_slices` and `rescaling_identity_energy_profiles`). For $N = 136$, $\lambda = \lambda_1$ (Revision/kohn_sham/results/rescaling/rescaling.csv):

a4,0	partner Delta k	partner v_t	max delta eps	relative delta E	partner E_KS
0.5	0.1516327	0.2231302	1.478e-13	7.099e-14	51.24916
1	0.09196986	0.04978707	1.101e-13	1.777e-14	32.39295
1.5	0.05578254	0.011109	8.749e-14	0	20.21303
2	0.03383382	0.002478752	7.661e-14	0	12.44706

Wolfram and sympy records	Rust solver records	reference and cross-check records
rescaling_identity	free_rescaling_relation_and_band_monotone, rescaling_identity_between_slices, rescaling_identity_energy_profiles	rescaling_identity_reference, rescaling_partners

11 The recorded states are not admissible sources of the a_4 equations

Revision/SPEC.md section 7 names the Kohn-Sham energy-momentum tensor as the source of the field equations for a_4 . The Einstein-Lovelock equations of the author's metric (Revision/field_equations_a4) require of any source: (C1) every component independent of x_8 ; (C2) $p_3 + p_t = 2p_8$ (the algebraic identity $E^{x_1}_{x_1} + E^{x_5}_{x_5} = 2E^{x_8}_{x_8}$ of the Lovelock tensors); (C3) for the linear member $a_4 = AHx_4 + a_0$: $p_3 = p_t = p_8$ and constant ρ . The record Revision/field_equations_a4/reports/ks-source-conditions.json (5 of 5 PASS) finds:

1. Every one of the 70 nonzero ground-state profiles depends on x_8 : $(\max \rho - \min \rho) / \max |T| \geq 0.0497329$ (record ks_profiles_depend_on_x8).
2. $\max_y |p_3 + p_t - 2p_8| / \max |T|$ lies between 2.09192 and 3.99006 (record ks_profiles_violate_algebraic_condition).
3. Integrated over the patch, $(\int p_3 + \int p_t) / (2 \int p_8)$ differs from 1 for every nonzero state (closest 0.414328); for the history $N = 136$, $\lambda = 0$ it is 0.339767, 0.25969 and 0.239714 at $a_{4,0} = 0, 1, 2$ (record ks_integrals_violate_algebraic_condition): even an x_8 -averaged source violates C2.
4. Along the history ρ changes with $a_{4,0}$ and $p_3 \neq p_t$, so C3 fails: the history is a PRESCRIBED background (record ks_history_is_a_prescribed_background).
5. Five states have an identically vanishing tensor (the free $N = 8$ zero modes), a zero source, for which Einstein gravity has no vacuum solution with $H > 0$ (record ks_zero_source_states_listed).

The full analysis Revision/field_equations_a4/ks_source/reports/ks-source-a4.json (23 of 23 PASS) refines this exactly: the states SATISFY the x_8 conservation identity, and $p_3 + p_t - 2p_8 = p_{8,y} / (3H)$ on shell, so C2 fails exactly because p_8 depends on x_8 ; at least a share 0.583068 of the energy density deviates from its hidden-direction average (L1 measure). Its conclusion: the coupled problem “metric of Revision/SPEC.md section 1 + Kohn-Sham source” has no solution, and nothing about a_4 is derived from it. In a stated approximation (the x_4 and $x_1 - x_5$ moments kept, the others dropped, with dropped residuals of order one) the Kohn-Sham source does not start or select the exponential deflation; it modifies a deflation set by the initial data, e.g. $a'_4(2)/a'_4(0) = 1.95362, 1.1321$ and 0.847549 for the source strengths $\sigma_0 = 10, 1$ and -1 of the series $N = 136$, $\lambda = 0$ in Einstein gravity, and with $\Lambda = 0$ the deflation halts at $a_4 = 0.149623$ (Einstein), 0.0724197 (Einstein-Gauss-Bonnet) and 0.102772 (cubic Lovelock). These are statements of that approximation, not of the field equations.

Consequence: the Kohn-Sham gas of this record is a test field on the prescribed background $a_4 = AHx_4$ without back-reaction; energies, pressures and equations of state along the history are not consequences of the coupled field equations.

report	records
Revision/field_equations_a4/reports/ks-source-conditions.json	ks_profiles_depend_on_x8, ks_profiles_violate_algebraic_condition, ks_integrals_violate_algebraic_condition, ks_history_is_a_prescribed_background, ks_zero_source_states_listed
Revision/field_equations_a4/ks_source/reports/ks-source-a4.json	A1_lovelock_identity_x1_plus_x5_equals_2x8, A6_x8_conservation_in_y, B0_wave1_report_verified, B2_x8_conservation_on_every_profile, B4_hidden_direction_mismatch, C1_averaged_algebraic_condition_fails, D3_lambda_zero_cases_halt, D5_einstein_sign_of_the_effect
Kohn-Sham theory (the record and the label of the history)	adiabatic_offdiagonal_identity, ks_theory_json_written, ks_theory_json_basis, ks_theory_json_slope, ks_theory_json_history_label

12 The Kohn-Sham-level pairing T3

The block map of the chirality Γ , $(\chi, j) \rightarrow (\sigma_2 \chi, -j)$, with the two brane parities exchanged and the tip angle $\theta \rightarrow \pi - \theta$, maps every self-consistent instantaneous Kohn-Sham state with (m, λ, θ) onto one with $(-m, +\lambda, \pi - \theta)$, with equal levels, occupations, Kohn-Sham energies and energy-momentum profiles and $S \rightarrow -S$ (ASSUMED Z2 brane, mean-field level). The theorem, its hypotheses and its proof are in Revision/docs/PAIR_CREATION_PROOFS.md section 8, verified by Revision/pairing/kohn_sham/reports/wolfram-t3.json (10 of 10 PASS) and Revision/pairing/kohn_sham/reports/python-t3.json (13 of 13 PASS); its completion record (Revision/pairing/kohn_sham/t3-completion.json) by Revision/pairing/kohn_sham/reports/wolfram-t3-completion.json (3 of 3 PASS) and Revision/pairing/kohn_sham/reports/python-t3-completion.json (7 of 7 PASS). The Rust solver contains a numerical self-test, labelled in its report as NOT a proof of T3: $(m, \lambda, \theta = 0)$ and $(-m, \lambda, \theta = \pi)$, solved independently, agree with opposite S (worst 9.95e-14, tolerance 1e-9; record t3_block_map_solver_selftest). T3 does not establish a creation process, a rate or an amplitude.

13 Verification records

13.1 Reports and their counts

Every check has a name, a verdict and a detail. Counts at the time of writing (the publication test re-reads them):

report	checks	PASS	FAIL
Revision/kohn_sham/reports/ks-theory-wolfram.json	46	46	0
Revision/kohn_sham/reports/ks-theory-python.json	58	58	0
Revision/kohn_sham/reports/ks-rust-solver.json	42	42	0
Revision/kohn_sham/reports/ks-rust-determinism.json	14	14	0
Revision/kohn_sham/reports/ks-rust-mermin-roots.json	5	5	0
Revision/kohn_sham/reports/ks-reference.json	37	37	0
Revision/kohn_sham/reports/ks-crosscheck.json	31	31	0
Revision/field_equations_a4/reports/ks-source-conditions.json	5	5	0
Revision/field_equations_a4/ks_source/reports/ks-source-a4.json	23	23	0
Revision/field_equations_a4/reports/wolfram-a4-report.json	52	52	0
Revision/field_equations_a4/reports/python-a4-report.json	63	63	0
Revision/pairing/kohn_sham/reports/wolfram-t3.json	10	10	0
Revision/pairing/kohn_sham/reports/python-t3.json	13	13	0

13.2 The independent reference and the full cross-check

The reference shares no code with the Rust solver: different discretisation (staggered finite differences with Richardson extrapolation instead of RK4 shooting), its own SCF, its own parameter derivation (it re-derives the particle numbers and couplings from the stated rules) and its own uncertainty, validated on a fourth grid (record `richardson_uncertainty_validated`). The checker `Revision/kohn_sham/checker/crosscheck_ks.py` compares the two on the FULL canonical matrix: 75 ground states, 135 thermal states, 60 exact-Fock-variant states, 60 rescaling partners, 75 particle-hole lists and the 5 slices of the crossing demonstration, 162691 comparisons in all, under a tolerance rule fixed before the comparison,

$$|x_{\text{Rust}} - x_{\text{ref}}| \leq 3 (U_{\text{ref}} + U_{\text{Rust}}) + 10^{-12} \text{ scale}, \quad U_{\text{Rust}} = \frac{16}{15} |\text{canonical} - \text{refined}|,$$

with U_{ref} the reference's three-grid Richardson uncertainty and U_{Rust} measured for every state of the matrix by the refinement measurement, whose script and record are `Revision/kohn_sham/checker/measure_rust_refinement.py` and `Revision/kohn_sham/checker/rust-refinement.json`. All 31 checks of the cross-check pass:

check	check	check
inputs_all_pass	rust_refinement_applies_to_matrix	rust_refinement_applies_to_matrix_exx
problem_definition_identical	parameters_particle_numbers	parameters_couplings
ground_occupations_and_groups	ground_energies	ground_homo_lumo_gap
ground_eigenvalues	excited_delta_scf	excited_particle_hole_lists
emt_integrals	emt_brane_tip_values	ground_profiles
exchange_delta_E_x	adiabatic_dE_da4	adiabatic_Q_max
exx_variant_scf	rescaling_partners	crossing_demonstration
thermo_state_functions	thermo_derivatives	thermo_sea_hole_diagnostic
thermo_levels	thermo_mu_high_precision	thermo_mu_rounding_diagnostic
thermo_mu_rust_satisfied_bound	reference_output_slf_only	reference_manifest
reference_repeat_byte_identical		

13.3 Determinism and the refinement measurement

The Rust determinism report (14 of 14 PASS): a repeat of the canonical matrix is byte-identical in all 244 result files and in its check report (`repeat_byte_identical`, `repeat_report_byte_identical`, `outputs_slf_only`); a run with refined tolerances (900 to 1800 RK4 steps, root tolerance 1e-14, SCF tolerance 1e-12) agrees with the canonical run to 1.901e-12 in E_{KS} (relative), 2.135e-09 in 23724 levels, 2.800e-09 in the profiles, 9.024e-10 in Q_{max} and dE/da_4 , 2.489e-10 in the thermodynamics and 5.689e-09 in C_V (`refined_same_inputs`, `refined_free_spectra_convergence_order`, `refined_ground_energies`, `refined_ground_homo_lumo_gap`, `refined_eigenvalues`, `refined_delta_sc`

f, refined_profiles, refined_adiabatic_derivatives, refined_thermodynamics, refined_mermin_root_path, refined_heat_capacity).

13.4 Re-verification for this document

For this document (2026-10-08) the committed Rust binary was run on the canonical matrix into a scratch directory (revision_ks_solver all --out <scratch>/rerun -report <scratch>/rerun-report.json): 42 of 42 checks PASS in 128 s of wall time on a machine shared with other jobs, and all 244 result files and the check report were byte-identical to the committed Revision/kohn_sham/results and Revision/kohn_sham/reports/ks-rust-solver.json. The other reports cited are the files of the working tree; their counts are re-read by the publication test.

14 Reproduction

From the repository root, in this order (the Kohn-Sham theory record reads the gammas of Revision/algebra; the Rust solver reads ks-theory.json; the Mermin-root tool runs after the canonical matrix; the cross-checker runs after the reference and the refinement measurement; the source-conditions checker reads the Kohn-Sham results and the Kohn-Sham theory checker reads its record; <scratch> is any empty directory outside the repository; the backslash is the line continuation of a POSIX shell):

```
wolframscript -file Revision/algebra/wolfram/verify_algebra.wls
python Revision/algebra/python/check_algebra.py
wolframscript -file Revision/field_equations_a4/wolfram/verify_field_equations_a4.wls
python Revision/field_equations_a4/python/check_field_equations_a4.py
wolframscript -file Revision/kohn_sham/theory/verify_ks_theory.wls
cargo build --release --manifest-path Revision/kohn_sham/solver/Cargo.toml
cargo test --release --manifest-path Revision/kohn_sham/solver/Cargo.toml
Revision/kohn_sham/solver/target/release/revision_ks_solver all \
    --out Revision/kohn_sham/results \
    --report Revision/kohn_sham/reports/ks-rust-solver.json
python Revision/kohn_sham/solver/tools/mermin_roots_mp.py --work <scratch>/mermin-roots
Revision/kohn_sham/solver/target/release/revision_ks_solver all \
    --out <scratch>/repeat --report <scratch>/repeat-report.json
Revision/kohn_sham/solver/target/release/revision_ks_solver all --refined \
    --out <scratch>/refined --report <scratch>/refined-report.json
python Revision/kohn_sham/solver/tools/compare_runs.py \
    --canonical Revision/kohn_sham/results \
    --canonical-report Revision/kohn_sham/reports/ks-rust-solver.json \
    --repeat <scratch>/repeat \
    --repeat-report <scratch>/repeat-report.json --refined <scratch>/refined \
    --refined-report <scratch>/refined-report.json \
    --report Revision/kohn_sham/reports/ks-rust-determinism.json
python Revision/kohn_sham/reference/run_reference.py --jobs 22 \
    --timing <scratch>/timing.json
python Revision/kohn_sham/checker/measure_rust_refinement.py \
    --work <scratch>/rust-single --jobs 22
python Revision/kohn_sham/checker/crosscheck_ks.py --work <scratch>/cc --jobs 22
python Revision/field_equations_a4/python/check_ks_source_conditions.py
```

```
python Revision/kohn_sham/theory/check_ks_theory.py
python Revision/field_equations_a4/ks_source/ks_source_a4.py
python -m unittest Revision/field_equations_a4/ks_source/test_ks_source_a4.py -v
wolframscript -file Revision/pairing/kohn_sham/wolfram/verify_t3.wls
python Revision/pairing/kohn_sham/python/check_t3.py
python scripts/build_provenance_pdf.py Revision/docs/KOHN_SHAM_DEFLATING_FIELD.md \
    --developer-layout --specifications Revision/pdf-specifications.json
python -m unittest Revision/tests/test_kohn_sham_deflating_field_publication.py -v
```

Expected output: every verifier ends with all of its checks passing and writes the report of section 13.1 with the counts given there (the Wolfram Kohn-Sham verifier prints 46/46 checks passed; the Rust solver prints SUCCESS after 42 PASS lines; the cross-checker writes 31 of 31 PASS); every run is deterministic, and a second run gives byte-identical outputs. Run times recorded in the READMEs of the folders (22 threads or processes; longer when the machine is shared): the Wolfram Kohn-Sham verifier about 40 to 85 s and the sympy checker about 75 to 165 s ([Revision/kohn_sham/theory/WOLFRAMSCRIPT_PROVENANCE.md](#)); the Rust canonical matrix 78.1 s on an idle machine ([Revision/kohn_sham/solver/README.md](#); 128 s in the re-run of section 13.4), the repeat 178.4 s and the refined run 618.6 s on a shared machine; the reference 578.3 s ([Revision/kohn_sham/reference/README.md](#)); the refinement measurement 31.6 s idle and 85.2 s shared; the cross-checker 588.9 s idle and 721.2 s shared, most of it the repeat of the reference that it runs itself ([Revision/kohn_sham/checker/README.md](#)); the Kohn-Sham source analysis 3 s to about 20 s ([Revision/field_equations_a4/ks_source/README.md](#)); the a_4 verifiers 43 to 70 s (Wolfram) and 7 to 28 s (sympy) ([Revision/field_equations_a4/README.md](#)); the T3 verifiers about 3 s and 1 s ([Revision/docs/PAIR_CREATION_PROOFS.md](#) section 9.3).

The PDF build runs the Markdown-to-LaTeX builder twice (with different hash seeds), runs pdflatex three times into each of two fresh directories, requires warning-free logs and byte-identical PDFs, and checks the PDF against its entry kohn-sham-deflating-field in [Revision/pdf-specifications.json](#) (the Revision registry; the registry of the earlier stages is not touched).

15 What is proved, computed, assumed and not established

15.1 Proved (exact, two independent verifiers)

1. The block reduction (section 3): for every history $a_4(x_4)$ the good-sector equation reduces exactly to eight 2×2 blocks h_j in the hidden coordinate; the spin-connection term $3H\gamma^8$ is removed by W^{-3} ; the a'_4 terms of the inflating and the deflating directions cancel.
2. The exchange functional (section 4): the exact local exchange of the uniform 8-fold good-sector gas at every temperature, $M_{\text{eff}} = m + \frac{15}{16}\lambda S$, $v_v = -\lambda n/16$; the exact Fock exchange of the closed-shell slab differs by $+\lambda Q^2/32$.
3. The boundary-value structure (section 5): self-adjointness with current-killing ends, the mirror map and the parities, the exact $k = 0$ spectra, the brane-band slope.
4. The rescaling identity (section 10), the energy-momentum tensor of the Kohn-Sham states with its trace and conservation identities (section 9), and the adiabaticity identity of the measure (section 7).

5. T3 (section 12; proved in `Revision/docs/PAIR_CREATION_PROOFS.md`), under its stated hypotheses.

15.2 Computed (numerical, two independent solvers, cross-checked)

1. The instantaneous ground states, first excited states (gap, Δ -SCF, particle-hole lists), Mermin states, energy-momentum profiles, rescaling partners and exact-Fock variants of the canonical matrix (sections 6 to 10), with the Rust solver and the reference agreeing on the full matrix within the stated uncertainty (31 of 31 checks, 162691 comparisons).
2. The adiabaticity measure ($Q_{\max} \leq 0.09345324$) and the absence of Fermi-level crossings in the canonical matrix; the crossing demonstration at $N = 696$ (section 7).
3. That the recorded states are not admissible sources of the a_4 equations (section 11).

15.3 Assumed or chosen

1. The Z2 brane (orbifold) at the patch end $z = \pi/2$ (ASSUMED); the metric is degenerate there and no junction condition is derived.
2. The regular tip at the cutoff $L = 3$ with $\theta = 0$ (chosen); proper densities near the tip are amplified by $e^{\delta H|y|}$, so the interacting results depend on L .
3. The filling convention (particles on the positive branch and on the $k = 0$ zero modes; the sea not populated thermally), whose justification is OPEN; the sea-hole diagnostic shows where it fails (15 of 135 thermal states).
4. The history $a_4 = AHx_4$, $A = 1$, PRESCRIBED, not solved for.
5. The parameters of section 5 ($N, \lambda, L, \Delta k, v_t, T$).

15.4 Not established

1. The time-dependent (non-adiabatic) Kohn-Sham problem (TDDFT) is OPEN: only instantaneous states are computed; the measure $Q_{\max}$ is a consistency indicator, not a solution of the evolution.
2. Correlation: the functional is Hartree plus exchange only; no correlation functional is derived or used.
3. Back-reaction: the Kohn-Sham states are a test field on a prescribed background; no recorded state is an admissible source of the field equations for a_4 , and no history $a_4(x_4)$ is derived from the Kohn-Sham source.
4. Equations of state and the dark-sector hypothesis are not part of this document (`Revision/dark_sector/dirac16complex`).
5. No creation process, rate or amplitude for universes follows from the Kohn-Sham model or from T3.

6. A thermal treatment of the Dirac sea (pairs) where the particle-only ensemble fails; states beyond the computed range $a_{4,0} \in [0, 2]$; continuum limits $L \rightarrow \infty$ and $\Delta k \rightarrow 0$.