

dirac16complex in the primordial pair-creation gravitational field

Explicit components of the connection, field equations, energy-momentum tensor,
quantization and equations of state

dirac16complex project (with Claude Opus 5.5)

September 2026

Contents

1	Scope and non-claims	4
2	Notation and conventions	5
3	Why this field is a reasonable primordial field	6
4	The metric and the vielbein	7
4.1	The x form	7
4.2	The z, t form and the vielbein	7
4.3	Determinant, volume and signature	8
4.4	The warped zeta form	8
5	Christoffel symbols	9
6	Canonical spin connection and the vielbein postulate	9
7	The spinor connection	10
7.1	Closed forms	10
7.2	Explicit action on the components	11
7.3	Curvature and the Lichnerowicz identity	11
8	Contraction with the gammas and covariant constancy	11
8.1	The contraction with the curved gammas	11
8.2	Covariant constancy of the gammas	12

8.3	The notebook contraction fails	12
9	The Lagrangian in this field	12
9.1	Formula	12
9.2	WolframScript form	13
9.3	Relation to the notebook Lagrangian	14
10	The sixteen component Euler-Lagrange equations	14
10.1	Matrix form	14
10.2	Component form in all eight coordinates	14
10.3	Other presentations of the equations	16
10.4	Evolution form	16
10.5	The (x0, x4) block form and the z, t evolution form	19
11	Comparison with the notebook's cell-1137 equations	19
11.1	Variables and cross reference	19
11.2	The stored equations and the difference	20
11.3	Origin of the q term: the cell-1058 substitution rule	21
11.4	What this means	22
12	The energy-momentum tensor	23
12.1	Definition and form in this field	23
12.2	The 21 nonzero connection terms	23
12.3	Diagonal and off-diagonal structure	24
13	Homogeneous sector and equations of state	24
13.1	Ansatz and bilinears	24
13.2	Energy density, pressures and kinetic/potential splits	25
13.3	Equation of state	25
13.4	Frozen in x4, independent of a4, and the p(0) caveat	26
13.5	Conservation	26
13.6	Exact examples	26
13.7	The state that does not depend on x0	26

14 Modes	27
14.1 Exact zeta reduction	27
14.2 Momenta along 3-space and the extra times	27
14.3 Local WKB dispersion and instability onset	28
15 Einstein tensor and the required source	28
15.1 Curvature	28
15.2 Source required by 8D Einstein gravity	29
15.3 Energy conditions	29
15.4 Linear a4 and the cell-150 alternatives	29
15.5 Which dirac16complex states can supply the source	30
16 The gamma-8 map: a structural plus-minus M pairing	32
17 Canonical quantization in this field	32
17.1 Momentum and anticommutators	32
17.2 The Krein structure	32
17.3 Hamiltonian density and Heisenberg equations	33
17.4 The good sector	33
18 Verification records	34
18.1 WolframScript	34
18.2 Python	35
18.3 Agreement between the implementations	35
18.4 Files and hashes	36
19 Reproduction	36
19.1 PowerShell	36
19.2 Git Bash	37
20 Limitations	37

Abstract

This document repeats, for one fixed background, the calculations that Stage 1 of the `dirac16complex` project carried out in an arbitrary gravitational field. The background is the primordial (pair-creation) field of the author's notebook: the diagonal metric of signature (4,4) called `MatrixMetric44` there, in which ordinary 3-space inflates while three extra times deflate, the 7-volume stays constant, and everything is warped over a hidden space-like coordinate. With coordinates $x_0, \dots, x_7$ and spinor components $\Psi_0, \dots, \Psi_{15}$ we give in closed form the vielbein and $\det g = +\cos^2 z$; the 37 nonzero Christoffel symbols; the 24 nonzero components of the canonical spin connection, for which the vielbein postulate holds in all 512 components; the spinor connection Ω_μ and the contraction $\gamma^\mu \Omega_\mu = 3H\gamma^0$, in which the free function a_4 cancels; the sixteen component Euler-Lagrange equations with all eight derivatives, their evolution form and the (x_0, x_4) block form; the 21 nonzero connection terms of the energy-momentum tensor; the energy density, the seven transverse pressures, two kinetic/potential splits and the equation of state of the homogeneous sector; the modes; the Einstein tensor and the source that 8-dimensional Einstein gravity would need, $\rho_{\text{req}} = -3H^2(7 + a_4'^2)/\kappa < 0$; which `dirac16complex` states can supply that source; and canonical quantization, $\{\Psi, \Psi^\dagger\} = B\delta^7/\cos z$ on a Krein space. For $a_4'' \neq 0$ no `dirac16complex` state that depends on x_0 and x_4 only can supply the source, and no plane wave in the hidden coordinate with real wave number can for any a_4 . For linear a_4 , which includes the notebook's $a_4 = t$, a condensate that does not depend on x_0 is an exact source in all 64 components. Its energy density is negative. The notebook's own block equations (cell 1137) agree term by term with the correct ones except for a term $\pm q y Z_j$. The stored notebook outputs that this document uses are reproduced exactly. That the q term comes from a wrong substitution rule in cell 1058 is a reconstruction. It reproduces the stored cell-1079 output exactly (16/16) if the rule hit only the +1 entries of three curved gammas, whereas a literal re-execution of cell 1058 in Mathematica 15.0.1 applies the rule to all entries and gives no q term. Two independent exact implementations verify the geometric and field-theoretic results: WolframScript (126 of 126 checks true) and Python/sympy (16 of 16 checks true, with 0 mismatches over $320 + 288$ coefficients per printer compared against the Wolfram components). The notebook itself is read by WolframScript alone; the Python checker compares only with transcriptions of three stored outputs. Statements marked as observations are not machine checks.

1 Scope and non-claims

Scope. The object is the Stage-1 field `dirac16complex`: a complex, Grassmann-odd, 16-component spinor Ψ of $\text{Spin}(4,4)$, coupled to gravity through the canonical spin connection, with the Lagrangian of Section 9. The background is the notebook metric of Section 4, with an arbitrary smooth function $a_4(t)$. Every formula below is exact in a_4 , a_4' and a_4'' . The field-independent algebra (the gamma matrices, C , the chirality, and the Krein structure of the matrix B with its subgroups) and four general identities (the Lichnerowicz sign, the curvature of Ω_μ , the tetrad derivation of $T_{\mu\nu}$ and its on-shell trace) come from Stage 1. Where this document only quotes such a result it says "(Stage 1)". Everything else was recomputed in this field by the Stage-2 verifiers of Section 18.

Non-claims.

1. The primordial field is a prescribed background. We do not claim that it solves 8-dimensional Einstein or Einstein-Lovelock gravity with a physically acceptable source. Section 15 shows that in Einstein gravity it needs a negative energy density for every a_4 .
2. For linear a_4 an x_0 -independent `dirac16complex` condensate is an exact source of this field

(Section 15.5), in the c-number reading of the bilinears. Its energy density is negative. We do not claim that such a state is realized in nature or that it is stable. For $a_4'' \neq 0$ no dirac16complex state that depends on x_0 and x_4 only is a source (Section 15.5).

3. The $\pm M$ pairing of Section 16 is a structural property of the equations. We make no claim that universes of masses $\pm M$ are created in pairs.
4. The Einstein-Lovelock terms of order 2 and 3, which the notebook names but does not compute, are not computed here either.
5. No numerical evolution is done here. For $\lambda \neq 0$ the statements about the ζ plane wave are mean-field statements (Section 13.1).
6. The quantization is canonical and formal. The state space is a Krein space, the modes with momentum along the extra times are not Hermitian (Section 14.3), and no interacting Fock space or renormalization is constructed.
7. The stored outputs of cells 1079, 1096, 1111 and 1137 are reproduced exactly. The notebook does not store the value of useT16, so that value, and with it the attribution of the q term to cell 1058, is a reconstruction (Section 11.3).

2 Notation and conventions

- Coordinates $x^\mu = (x_0, \dots, x_7)$, counted from 0. The roles follow the notebook: x_0 is the hidden space, x_1, x_2, x_3 are ordinary 3-space, x_4 is the evolution time, and x_5, x_6, x_7 are the three extra times (the notebook calls them superluminal deflating times).
- $z = 6Hx_0 \in (0, \pi/2)$, $t = Hx_4$, $s = \sin z$. $H > 0$ is the notebook's single inverse length. $a_4 = a_4(t)$ is an arbitrary smooth real function and $a_4' = da_4/dt$.
- $\partial_\mu = \partial/\partial x^\mu$, so $\partial_0 = 6H\partial_z$ and $\partial_4 = H\partial_t$.
- Index groups: $i \in \{1, 2, 3\}$ (3-space) and $j \in \{5, 6, 7\}$ (extra times). "Transverse" means every direction $\mu \neq 4$.
- Tangent metric $\eta = \text{diag}(+1, +1, +1, +1, -1, -1, -1, -1)$. Frame directions 0 to 3 are space-like and 4 to 7 are time-like.
- Frame gamma matrices $\gamma^a = \begin{pmatrix} 0 & \bar{\tau}_a \\ \tau_a & 0 \end{pmatrix}$ ($a = 0, \dots, 7$) in the split-octonion (notebook) basis: real 16 by 16 integer matrices with $\{\gamma^a, \gamma^b\} = 2\eta^{ab}$, symmetric for $a < 4$ and antisymmetric for $a \geq 4$. Every row of every γ^a has exactly one nonzero entry, +1 or -1.
- A gamma matrix with a numeric upper index, $\gamma^0, \dots, \gamma^7$, is always the constant frame matrix. Curved gammas carry a Greek index, $\gamma^\mu = e_a^\mu \gamma^a$; with a numeric label they are written $\gamma^{x\mu}$. Lowered gammas $\gamma_\mu = g_{\mu\nu} \gamma^\nu$ are always curved. In sums with an explicit coefficient, such as $\sum_\mu c_\mu \gamma^\mu \partial_\mu$ with $c_\mu = 1/h_\mu$, γ^μ is the frame matrix with the numeric value of μ .
- $C = \sigma_{16} = \gamma^0 \gamma^1 \gamma^2 \gamma^3 = \text{diag}(-\sigma, \sigma)$ with $\sigma = \begin{pmatrix} 0 & I_4 \\ I_4 & 0 \end{pmatrix}$, so $C^T = C$ and $C^2 = 1$. The Dirac adjoint is $\bar{\Psi} = \Psi^\dagger C$.

- $\gamma^8 = \gamma^0 \gamma^1 \dots \gamma^7 = \text{diag}(-I_8, I_8)$: $\Psi_0, \dots, \Psi_7$ have chirality -1 and $\Psi_8, \dots, \Psi_{15}$ have chirality $+1$.
- $S^{ab} = \frac{1}{4}[\gamma^a, \gamma^b]$ and $\Omega_\mu = \frac{1}{2}\omega_{\mu ab}S^{ab}$, summed over all ordered pairs (a, b) .
- $\Psi = (\Psi_0, \dots, \Psi_{15})^T$ has complex Grassmann-odd components. The notebook's $\text{fl6}[k]$ and $\text{Z}[k]$ are Ψ_k (cross reference in Section 11.1).
- m is the mass parameter. The notebook's M corresponds to $m = -HM$. The self-interaction is $U(S) = \frac{\lambda}{2}S^2$ with $S = \bar{\Psi}\Psi$, and $M_{\text{eff}} = m + U'(S) = m + \lambda S$.
- In the homogeneous-sector formulas (Sections 13 to 15) the amplitude u is a c-number vector. This is the classical, mean-field reading of the bilinears.
- $q = Q_1 \sinh(a_4) a'_4 e^{-a_4}$ always denotes the notebook's spurious term of Section 11. Momenta along x_1 and x_5 are called k_1 and k_5 (Section 14).
- "Cell N " is the N -th cell of the notebook in file order, counting only cells of the styles Input, Code, Text, Section, Subsection, Subsubsection, Title, Chapter, Item, Subtitle, DisplayFormula, ItemNumbered and Program (1279 cells). Output, Print and Message cells are not counted; they belong to the preceding counted cell. This is the enumeration of the function `loadNotebookCells` in `wolfram/Dirac16ComplexPrimordial.wl`. The cited cells have these In and Out labels: 501 In[467] with Out[467]; 583 In[535] with Out[535]; 584 In[536] with Out[536] and Out[537]; 1058 In[1024]; 1060 In[1026] with Out[1026]; 1066 In[1037]; 1075 In[1048]; 1078 In[1051]; 1079 In[1052] with Out[1052]; 1089 In[1059] with Out[1059]; 1096 In[1065] with Out[1065] to Out[1067]; 1111 In[1087] with Out[1087] to Out[1090]; 1137 In[1113] with Out[1115] and Out[1116].
- For a scalar field these sign conventions give $\rho = \frac{1}{2}\dot{\phi}^2 + V$ and $p = \frac{1}{2}\dot{\phi}^2 - V$, the same as the usual treatment in signature $(+, -, -, -)$. The equation of state parameter is $w = p/\rho$.

3 Why this field is a reasonable primordial field

The field is the metric `MatrixMetric44` of the author's notebook

`Pair_Creation_of_Universes_WaveFunctionOfUniverse-4+4-Einstein-Lovelock-Nash.nb`

which is read but never modified; its sha256 is recorded in Section 18.4. The notebook opens with a hypothesis: if superluminal inflation or deflation exists in Einstein (or Einstein-Lovelock) gravity, then at time $x_4 = 0$, before the particles of the standard model exist, a pair of universes with masses $\pm M$ is created. `MatrixMetric44` is the metric in which the notebook expresses this picture, and it has the following features.

1. **Inflation of 3-space.** The three space directions have the scale factor $s^{1/6}e^{a_4(t)}$ and the rate $\partial_4 \ln(s^{1/6}e^{a_4}) = Ha'_4$. For $a_4 = t$ ordinary 3-space expands exponentially, as e^{Hx_4} , at the rate H . This is what the notebook calls superluminal inflation.
2. **Deflation of the three extra times.** The directions x_5, x_6, x_7 have the scale factor $s^{1/6}e^{-a_4(t)}$ and the rate $-Ha'_4$.

3. **Constant 7-volume.** The six transverse scale factors multiply to s , independent of a_4 . Hence $\sqrt{|g|} = \cos z$ does not depend on x_4 : the inflation of 3-space is exactly compensated by the deflation of the extra times.
4. **Hidden-space warp.** In the coordinate ζ of Section 4.4 the six transverse directions carry the common factor $e^{2H\zeta}$ over the hidden coordinate $\zeta \in (-\infty, 0)$, an exponentially warped product.
5. **Control.** One scale H and one free function a_4 . The metric is diagonal and depends on x_0 and x_4 only. Its spin connection enters the Dirac operator only through $\gamma^\mu \Omega_\mu = 3H\gamma^0$, a constant multiple of γ^0 , and its Einstein tensor is diagonal and independent of x_0 .

These properties make it a reasonable primordial background for the Stage-1 field. It is the author's own field, so the notebook's spinor equations can be compared with the correct ones term by term (Section 11). It is exactly tractable. And it contains, in one metric, an inflating 3-space and deflating extra times with an adjustable history a_4 , which is what a primordial scenario of the notebook's type needs.

What it requires as a source. In 8-dimensional Einstein gravity, $G^\mu{}_\nu = \kappa T^\mu{}_\nu$, the field needs the energy density $\rho_{\text{req}} = -3H^2(7 + a_4'^2)/\kappa$. This is negative for every a_4 , and the null energy condition along $e_4 + e_0$ fails for every a_4 (Section 15). It is neither a vacuum solution nor a solution with a cosmological constant: $G^0{}_0 = G^4{}_4$ would require $-3H^2(a_4'^2 - 5) = 3H^2(7 + a_4'^2)$, that is $a_4'^2 = -1$ (derived here from the verified closed forms of Section 15.1). So the notebook's hope that the spinor source tensor is Λg cannot be realized in Einstein gravity. The notebook intends the source of its spinor field to enter the Einstein and/or Einstein-Lovelock field equations, but its curvature code stops at the Einstein tensor (cells 583 and 584, which this document reproduces exactly) and computes no Lovelock terms. This document does not compute them either. Within Einstein gravity `dirac16complex` can supply the source only for linear a_4 . Then a condensate that does not depend on x_0 is an exact source, with negative energy density. For $a_4'' \neq 0$ no `dirac16complex` state that depends on x_0 and x_4 only can be a source (Section 15.5). The field is therefore a reasonable kinematic model of the notebook's primordial picture. For linear a_4 it has an exact `dirac16complex` source of negative energy; what determines a_4 remains open.

4 The metric and the vielbein

4.1 The x form

In the notebook's coordinates the metric is

$$g_{\mu\nu} = \text{diag}(\cot^2(6Hx_0), \sin^{1/3}(6Hx_0) e^{2a_4(Hx_4)} (\times 3), -1, \\ -\sin^{1/3}(6Hx_0) e^{-2a_4(Hx_4)} (\times 3)).$$

4.2 The z, t form and the vielbein

With $z = 6Hx_0$, $t = Hx_4$ and $s = \sin z$ the line element is

$$ds^2 = \cot^2 z \, dx_0^2 + s^{1/3} e^{2a_4} (dx_1^2 + dx_2^2 + dx_3^2) - dx_4^2 \\ - s^{1/3} e^{-2a_4} (dx_5^2 + dx_6^2 + dx_7^2).$$

We use the diagonal vielbein

$$e_\mu^a = \text{diag}(h_\mu), \quad h = (\cot z, s^{1/6} e^{a_4} (\times 3), 1, s^{1/6} e^{-a_4} (\times 3)),$$

with inverse $e_a^\mu = \text{diag}(1/h_\mu)$. The product $e \eta e^T$ equals the metric exactly. The curved gammas are

$$\gamma^{x_0} = \tan z \gamma^0, \quad \gamma^{x_i} = s^{-1/6} e^{-a_4} \gamma^i, \quad \gamma^{x_4} = \gamma^4, \quad \gamma^{x_j} = s^{-1/6} e^{a_4} \gamma^j,$$

and the lowered ones are

$$\gamma_0 = \cot z \gamma^0, \quad \gamma_i = s^{1/6} e^{a_4} \gamma^i, \quad \gamma_4 = -\gamma^4, \quad \gamma_j = -s^{1/6} e^{-a_4} \gamma^j.$$

4.3 Determinant, volume and signature

The diagonal signs are $(+, +, +, +, -, -, -, -)$ for $z \in (0, \pi/2)$ and real a_4 , so the signature is $(4,4)$. The determinant is

$$\det g = \cot^2 z (s^{1/3} e^{2a_4})^3 (-1) (-s^{1/3} e^{-2a_4})^3 = +\cos^2 z, \quad \sqrt{|g|} = \cos z.$$

The sign is $+$ because there is an even number (four) of negative entries; the notebook's cell 1060 shows the same value. (The Stage-2 specification had written $\det g = -\cos^2 z$. That statement is corrected here; $\sqrt{|g|} = \cos z$ is unaffected.) Since $\sqrt{|g|}$ does not depend on x_4 , the 7-volume of a comoving region is constant in x_4 .

4.4 The warped zeta form

Define

$$\zeta = \frac{\ln \sin z}{6H} \in (-\infty, 0), \quad e^{6H\zeta} = s = \sin z, \quad d\zeta = \cot z \, dx_0$$

Then $g_{00} \, dx_0^2 = d\zeta^2$ and $s^{1/3} = e^{2H\zeta}$, and the metric becomes the warped form

$$ds^2 = d\zeta^2 - dx_4^2 + e^{2H\zeta} [e^{2a_4} (dx_1^2 + dx_2^2 + dx_3^2) - e^{-2a_4} (dx_5^2 + dx_6^2 + dx_7^2)]$$

with $\sqrt{|g|} = e^{6H\zeta}$ in the coordinates $(\zeta, x_1, \dots, x_7)$. The x_0 chart covers $\zeta < 0$. At $\zeta = 0$ ($z = \pi/2$) the x_0 coordinate degenerates ($g_{00} = \cot^2 z = 0$ and $\sqrt{|g|} = \cos z = 0$), while the ζ form of the line element is regular there. This is an observation from the verified ζ form, not a separate check.

5 Christoffel symbols

With $\Gamma^\rho_{\mu\nu} = \frac{1}{2}g^{\rho\sigma}(\partial_\mu g_{\nu\sigma} + \partial_\nu g_{\mu\sigma} - \partial_\sigma g_{\mu\nu})$ exactly 37 of the 512 symbols are nonzero. The 25 with $\mu \leq \nu$ are listed below; the others follow from $\Gamma^\rho_{\nu\mu} = \Gamma^\rho_{\mu\nu}$ (12 off-diagonal pairs). Every nonzero symbol has at least one index 0 or 4, because the metric depends on x_0 and x_4 only.

Symbol	Closed form
Γ_{00}^0	$-6H s^{-1} \sec z$
Γ_{11}^0	$-H \tan z s^{1/3} e^{2a_4}$
Γ_{22}^0	$-H \tan z s^{1/3} e^{2a_4}$
Γ_{33}^0	$-H \tan z s^{1/3} e^{2a_4}$
Γ_{55}^0	$H \tan z s^{1/3} e^{-2a_4}$
Γ_{66}^0	$H \tan z s^{1/3} e^{-2a_4}$
Γ_{77}^0	$H \tan z s^{1/3} e^{-2a_4}$
Γ_{01}^1	$H \cot z$
Γ_{14}^1	$H a'_4$
Γ_{02}^2	$H \cot z$
Γ_{24}^2	$H a'_4$
Γ_{03}^3	$H \cot z$
Γ_{34}^3	$H a'_4$
Γ_{11}^4	$H s^{1/3} e^{2a_4} a'_4$
Γ_{22}^4	$H s^{1/3} e^{2a_4} a'_4$
Γ_{33}^4	$H s^{1/3} e^{2a_4} a'_4$
Γ_{55}^4	$H s^{1/3} e^{-2a_4} a'_4$
Γ_{66}^4	$H s^{1/3} e^{-2a_4} a'_4$
Γ_{77}^4	$H s^{1/3} e^{-2a_4} a'_4$
Γ_{05}^5	$H \cot z$
Γ_{45}^5	$-H a'_4$
Γ_{06}^6	$H \cot z$
Γ_{46}^6	$-H a'_4$
Γ_{07}^7	$H \cot z$
Γ_{47}^7	$-H a'_4$

6 Canonical spin connection and the vielbein postulate

The canonical spin connection is fixed by the vielbein postulate, which says that the total covariant derivative of the vielbein is zero:

$$\partial_\mu e_\nu^a - \Gamma^\rho_{\mu\nu} e_\rho^a + \omega_\mu^a{}_b e_\nu^b = 0 \quad \Longleftrightarrow \quad \omega_\mu^a{}_b = e_b^\nu (\Gamma^\rho_{\mu\nu} e_\rho^a - \partial_\mu e_\nu^a),$$

with $\omega_{\mu ab} = \eta_{ac} \omega_\mu^c{}_b = -\omega_{\mu ba}$. All 512 components of the left-hand side vanish identically, and the lowered connection is antisymmetric in (a, b) . Exactly 24 components $\omega_{\mu ab}$ are nonzero; $\omega_{0ab} = \omega_{4ab} = 0$. In closed form, for $i = 1, 2, 3$ and $j = 5, 6, 7$:

$$\begin{aligned}\omega_{i0i} &= -\omega_{i0} = -H s^{1/6} e^{a_4}, & \omega_{i4i} &= -\omega_{i4} = H s^{1/6} e^{a_4} a'_4, \\ \omega_{j0j} &= -\omega_{j0} = H s^{1/6} e^{-a_4}, & \omega_{j4j} &= -\omega_{j4} = H s^{1/6} e^{-a_4} a'_4.\end{aligned}$$

The complete list, each antisymmetric pair in one row:

Component	Value	Component	Value
ω_{101}	$-H s^{1/6} e^{a_4}$	ω_{110}	$H s^{1/6} e^{a_4}$
ω_{114}	$H s^{1/6} e^{a_4} a'_4$	ω_{141}	$-H s^{1/6} e^{a_4} a'_4$
ω_{202}	$-H s^{1/6} e^{a_4}$	ω_{220}	$H s^{1/6} e^{a_4}$
ω_{224}	$H s^{1/6} e^{a_4} a'_4$	ω_{242}	$-H s^{1/6} e^{a_4} a'_4$
ω_{303}	$-H s^{1/6} e^{a_4}$	ω_{330}	$H s^{1/6} e^{a_4}$
ω_{334}	$H s^{1/6} e^{a_4} a'_4$	ω_{343}	$-H s^{1/6} e^{a_4} a'_4$
ω_{505}	$H s^{1/6} e^{-a_4}$	ω_{550}	$-H s^{1/6} e^{-a_4}$
ω_{545}	$-H s^{1/6} e^{-a_4} a'_4$	ω_{554}	$H s^{1/6} e^{-a_4} a'_4$
ω_{606}	$H s^{1/6} e^{-a_4}$	ω_{660}	$-H s^{1/6} e^{-a_4}$
ω_{646}	$-H s^{1/6} e^{-a_4} a'_4$	ω_{664}	$H s^{1/6} e^{-a_4} a'_4$
ω_{707}	$H s^{1/6} e^{-a_4}$	ω_{770}	$-H s^{1/6} e^{-a_4}$
ω_{747}	$-H s^{1/6} e^{-a_4} a'_4$	ω_{774}	$H s^{1/6} e^{-a_4} a'_4$

The mixed components $\omega_\mu{}^a{}_b$ are the notebook's $\omega_{\mu IJ}$; its stored cell-501 output equals them entry by entry. For the 12 components with one space-like and one time-like frame index, the pairs $(i, 4)$ for $\mu = i$ and $(0, j)$ for $\mu = j$, the mixed form is symmetric in (a, b) :

$$\omega_i{}^i{}_4 = \omega_i{}^4{}_i = H s^{1/6} e^{a_4} a'_4, \quad \omega_j{}^0{}_j = \omega_j{}^j{}_0 = H s^{1/6} e^{-a_4}.$$

The notebook contracts the mixed form with S^{ab} (both indices up). The symmetric part then drops out against the antisymmetric S^{ab} , so that contraction silently deletes every boost-type component (Section 8.3).

7 The spinor connection

7.1 Closed forms

The spinor connection $\Omega_\mu = \frac{1}{2}\omega_{\mu ab}S^{ab} = \frac{1}{8}\omega_{\mu ab}[\gamma^a, \gamma^b]$ is

$$\begin{aligned}\Omega_0 &= \Omega_4 = 0, \\ \Omega_i &= -H s^{1/6} e^{a_4} (S^{0i} + a'_4 S^{4i}) \quad (i = 1, 2, 3), \\ \Omega_j &= H s^{1/6} e^{-a_4} (S^{0j} - a'_4 S^{4j}) \quad (j = 5, 6, 7).\end{aligned}$$

Every Ω_μ is block diagonal in chirality, $\Omega_\mu = \text{diag}(\Omega_\mu^{(-)}, \Omega_\mu^{(+)})$ on $\Psi_{0..7}$ and $\Psi_{8..15}$, with the upper block $\frac{1}{2}\omega_{\mu ab}\frac{1}{4}(\bar{\tau}_a\tau_b - \bar{\tau}_b\tau_a)$ and the lower block $\frac{1}{2}\omega_{\mu ab}\frac{1}{4}(\tau_a\bar{\tau}_b - \tau_b\bar{\tau}_a)$, and every Ω_μ commutes with γ^8 .

7.2 Explicit action on the components

Each row of each nonzero Ω_μ has exactly two nonzero entries, one proportional to 1 and one to a'_4 . Write $\alpha_i = H s^{1/6} e^{a_4}$ for $i = 1, 2, 3$ and $\alpha_j = H s^{1/6} e^{-a_4}$ for $j = 5, 6, 7$. Row n of the array below gives $(\Omega_\mu \Psi)_n$ in units of $\frac{1}{2}\alpha_\mu$. For example, the entry $-\Psi_7 + a'_4 \Psi_2$ in row 0, column Ω_1 means $(\Omega_1 \Psi)_0 = \frac{1}{2} H s^{1/6} e^{a_4} (-\Psi_7 + a'_4 \Psi_2)$. The array is generated from the 16 by 16 entries in primordial-components.json.

n	Ω_1	Ω_2	Ω_3	Ω_5	Ω_6	Ω_7
0	$-\Psi_7 + a'_4 \Psi_2$	$\Psi_6 + a'_4 \Psi_3$	$-\Psi_5 - a'_4 \Psi_0$	$-\Psi_6 + a'_4 \Psi_3$	$-\Psi_7 - a'_4 \Psi_2$	$-\Psi_0 + a'_4 \Psi_5$
1	$-\Psi_6 - a'_4 \Psi_3$	$-\Psi_7 + a'_4 \Psi_2$	$\Psi_4 - a'_4 \Psi_1$	$-\Psi_7 - a'_4 \Psi_2$	$\Psi_6 - a'_4 \Psi_3$	$-\Psi_1 - a'_4 \Psi_4$
2	$\Psi_5 + a'_4 \Psi_0$	$-\Psi_4 + a'_4 \Psi_1$	$-\Psi_7 + a'_4 \Psi_2$	$\Psi_4 + a'_4 \Psi_1$	$-\Psi_5 + a'_4 \Psi_0$	$-\Psi_2 - a'_4 \Psi_7$
3	$\Psi_4 - a'_4 \Psi_1$	$\Psi_5 + a'_4 \Psi_0$	$\Psi_6 + a'_4 \Psi_3$	$\Psi_5 - a'_4 \Psi_0$	$\Psi_4 + a'_4 \Psi_1$	$-\Psi_3 + a'_4 \Psi_6$
4	$-\Psi_3 - a'_4 \Psi_6$	$\Psi_2 - a'_4 \Psi_7$	$-\Psi_1 + a'_4 \Psi_4$	$\Psi_2 + a'_4 \Psi_7$	$\Psi_3 - a'_4 \Psi_6$	$\Psi_4 + a'_4 \Psi_1$
5	$-\Psi_2 + a'_4 \Psi_7$	$-\Psi_3 - a'_4 \Psi_6$	$\Psi_0 + a'_4 \Psi_5$	$\Psi_3 - a'_4 \Psi_6$	$-\Psi_2 - a'_4 \Psi_7$	$\Psi_5 - a'_4 \Psi_0$
6	$\Psi_1 - a'_4 \Psi_4$	$-\Psi_0 - a'_4 \Psi_5$	$-\Psi_3 - a'_4 \Psi_6$	$-\Psi_0 + a'_4 \Psi_5$	$\Psi_1 + a'_4 \Psi_4$	$\Psi_6 - a'_4 \Psi_3$
7	$\Psi_0 + a'_4 \Psi_5$	$\Psi_1 - a'_4 \Psi_4$	$\Psi_2 - a'_4 \Psi_7$	$-\Psi_1 - a'_4 \Psi_4$	$-\Psi_0 + a'_4 \Psi_5$	$\Psi_7 + a'_4 \Psi_2$
8	$\Psi_{15} + a'_4 \Psi_{10}$	$-\Psi_{14} + a'_4 \Psi_{11}$	$\Psi_{13} - a'_4 \Psi_8$	$\Psi_{14} + a'_4 \Psi_{11}$	$\Psi_{15} - a'_4 \Psi_{10}$	$\Psi_8 + a'_4 \Psi_{13}$
9	$\Psi_{14} - a'_4 \Psi_{11}$	$\Psi_{15} + a'_4 \Psi_{10}$	$-\Psi_{12} - a'_4 \Psi_9$	$\Psi_{15} - a'_4 \Psi_{10}$	$-\Psi_{14} - a'_4 \Psi_{11}$	$\Psi_9 - a'_4 \Psi_{12}$
10	$-\Psi_{13} + a'_4 \Psi_8$	$\Psi_{12} + a'_4 \Psi_9$	$\Psi_{15} + a'_4 \Psi_{10}$	$-\Psi_{12} + a'_4 \Psi_9$	$\Psi_{13} + a'_4 \Psi_8$	$\Psi_{10} - a'_4 \Psi_{15}$
11	$-\Psi_{12} - a'_4 \Psi_9$	$-\Psi_{13} + a'_4 \Psi_8$	$-\Psi_{14} + a'_4 \Psi_{11}$	$-\Psi_{13} - a'_4 \Psi_8$	$-\Psi_{12} + a'_4 \Psi_9$	$\Psi_{11} + a'_4 \Psi_{14}$
12	$\Psi_{11} - a'_4 \Psi_{14}$	$-\Psi_{10} - a'_4 \Psi_{15}$	$\Psi_9 + a'_4 \Psi_{12}$	$-\Psi_{10} + a'_4 \Psi_{15}$	$-\Psi_{11} - a'_4 \Psi_{14}$	$-\Psi_{12} + a'_4 \Psi_9$
13	$\Psi_{10} + a'_4 \Psi_{15}$	$\Psi_{11} - a'_4 \Psi_{14}$	$-\Psi_8 + a'_4 \Psi_{13}$	$-\Psi_{11} - a'_4 \Psi_{14}$	$\Psi_{10} - a'_4 \Psi_{15}$	$-\Psi_{13} - a'_4 \Psi_8$
14	$-\Psi_9 - a'_4 \Psi_{12}$	$\Psi_8 - a'_4 \Psi_{13}$	$\Psi_{11} - a'_4 \Psi_{14}$	$\Psi_8 + a'_4 \Psi_{13}$	$-\Psi_9 + a'_4 \Psi_{12}$	$-\Psi_{14} - a'_4 \Psi_{11}$
15	$-\Psi_8 + a'_4 \Psi_{13}$	$-\Psi_9 - a'_4 \Psi_{12}$	$-\Psi_{10} - a'_4 \Psi_{15}$	$\Psi_9 - a'_4 \Psi_{12}$	$\Psi_8 + a'_4 \Psi_{13}$	$-\Psi_{15} + a'_4 \Psi_{10}$

7.3 Curvature and the Lichnerowicz identity

The curvature of the spinor connection is $F_{\mu\nu} = \partial_\mu \Omega_\nu - \partial_\nu \Omega_\mu + [\Omega_\mu, \Omega_\nu] = \frac{1}{2} R_{ab\mu\nu} S^{ab}$, with $R_{ab\mu\nu} = \eta_{ac} e_\rho{}^c R^\rho{}_{\sigma\mu\nu} e_b{}^\sigma$ and $R^\rho{}_{\sigma\mu\nu} = \partial_\mu \Gamma^\rho{}_{\nu\sigma} - \partial_\nu \Gamma^\rho{}_{\mu\sigma} + \Gamma^\rho{}_{\mu\lambda} \Gamma^\lambda{}_{\nu\sigma} - \Gamma^\rho{}_{\nu\lambda} \Gamma^\lambda{}_{\mu\sigma}$. The square of the Dirac operator is

$$(\gamma^\mu D_\mu)^2 \Psi = g^{\mu\nu} (D_\mu D_\nu \Psi - \Gamma^\lambda{}_{\mu\nu} D_\lambda \Psi) - \frac{R}{4} \Psi$$

(Stage 1, which also verified both identities at exact points of this field). With $R = 6H^2(a_4'^2 - 7)$ from Section 15.1, the curvature term is $-\frac{R}{4} \Psi = \frac{3}{2} H^2 (7 - a_4'^2) \Psi$, which equals $9H^2 \Psi$ for $a_4 = t$. Since $G^4_4 = 3H^2(7 + a_4'^2)$ never vanishes, the Riemann tensor never vanishes, so Ω_μ is nowhere locally pure gauge. These two statements are consequences drawn here, not separate checks.

8 Contraction with the gammas and covariant constancy

8.1 The contraction with the curved gammas

$$\gamma^\mu \Omega_\mu = 3H\gamma^0, \quad \Omega_\mu \gamma^\mu = -3H\gamma^0, \quad \{\gamma^\mu, \Omega_\mu\} = 0, \quad [\gamma^\mu, \Omega_\mu] = 6H\gamma^0.$$

The free function a_4 cancels. For a diagonal vielbein, $\gamma^\mu \Omega_\mu = \frac{1}{2} \sum_b h_b^{-1} \partial_b \ln(\prod_{c \neq b} h_c) \gamma^b$, and this formula holds exactly here. Only $b = 0$ contributes: $\prod_{c \neq 0} h_c = s$ and $\tan z \partial_0 \ln s = 6H$. For $b = 4$, $\prod_{c \neq 4} h_c = \cos z$ does not depend on x_4 . In other words a_4 cancels because the 7-volume is constant. The divergence identity used in the derivation of the field equations holds:

$$\partial_\mu (\sqrt{|g|} \gamma^\mu) = 6H \cos z \gamma^0 = \sqrt{|g|} [\gamma^\mu, \Omega_\mu].$$

8.2 Covariant constancy of the gammas

With the canonical connection,

$$D_\mu \gamma^\nu := \partial_\mu \gamma^\nu + \Gamma^\nu_{\mu\lambda} \gamma^\lambda + [\Omega_\mu, \gamma^\nu] = 0$$

holds for all 64 pairs (μ, ν) . This is the spinor form of the vielbein postulate.

8.3 The notebook contraction fails

The notebook contraction $\Omega_\mu^{\text{NB}} = \frac{1}{2} \omega_\mu^a{}_b S^{ab}$, with one η missing, violates $D_\mu \gamma^\nu = 0$ in 15 of the 64 pairs, with 288 nonzero matrix entries in total. The values, in frame gammas (both verifiers; check `P_gammaConst_notebookContractionClosedForms`):

Pairs (μ, ν)	$D_\mu \gamma^\nu$ with the notebook contraction
$(i, i), i = 1, 2, 3$	$H a'_4 \gamma^4$
$(i, 4), i = 1, 2, 3$	$H s^{1/6} e^{a_4} a'_4 \gamma^i$
$(j, 0), j = 5, 6, 7$	$H s^{7/6} e^{-a_4} \sec z \gamma^j$
$(j, 4), j = 5, 6, 7$	$2H s^{1/6} e^{-a_4} a'_4 \gamma^j$
$(j, j), j = 5, 6, 7$	$H \gamma^0 - 2H a'_4 \gamma^4$

For example $D_1 \gamma^4 = H s^{1/6} e^{a_4} a'_4 \gamma^1$. With the notebook contraction $\gamma^\mu \Omega_\mu^{\text{NB}} = \frac{3}{2} H (\gamma^0 + a'_4 \gamma^4)$ instead of $3H \gamma^0$, and the divergence identity of Section 8.1 fails.

9 The Lagrangian in this field

9.1 Formula

The Stage-1 Lagrangian is

$$\mathcal{L} = \sqrt{|g|} \left[\frac{1}{2} (\bar{\Psi} \gamma^\mu D_\mu \Psi - (D_\mu \bar{\Psi}) \gamma^\mu \Psi) - m \bar{\Psi} \Psi - U(\bar{\Psi} \Psi) \right],$$

with $D_\mu \Psi = \partial_\mu \Psi + \Omega_\mu \Psi$, $D_\mu \bar{\Psi} = \partial_\mu \bar{\Psi} - \bar{\Psi} \Omega_\mu$ and $U(S) = \frac{\lambda}{2} S^2$. No factor i is needed: $C \gamma^a$ is real and antisymmetric, hence anti-Hermitian, so the symmetrized kinetic term is Hermitian (Stage

1). In this field $\sqrt{|g|} = \cos z$. The connection terms combine to $\frac{1}{2}\bar{\Psi}\{\gamma^\mu, \Omega_\mu\}\Psi$, and $\{\gamma^\mu, \Omega_\mu\} = 0$ (Section 8.1), so they cancel inside $\mathcal{L}$:

$$\mathcal{L} = \cos z \left[\frac{1}{2} \sum_{\mu=0}^7 c_\mu (\bar{\Psi} \gamma^\mu \partial_\mu \Psi - \partial_\mu \bar{\Psi} \gamma^\mu \Psi) - mS - \frac{\lambda}{2} S^2 \right],$$

$$c = (\tan z, s^{-1/6} e^{-a_4} (\times 3), 1, s^{-1/6} e^{a_4} (\times 3)),$$

where the γ^μ in the sum are frame matrices. From $C = \text{diag}(-\sigma, \sigma)$ the scalar bilinear is, in components,

$$S = \bar{\Psi}\Psi = - \sum_{n=0}^3 (\Psi_n^* \Psi_{n+4} + \Psi_{n+4}^* \Psi_n) + \sum_{n=8}^{11} (\Psi_n^* \Psi_{n+4} + \Psi_{n+4}^* \Psi_n).$$

The cancellation inside $\mathcal{L}$ and the component form of S are consequences of verified identities (the first of $\{\gamma^\mu, \Omega_\mu\} = 0$, the second of the committed C). The connection does not disappear from the physics. It returns in the field equations through $\partial_\mu(\sqrt{|g|}\gamma^\mu) = \sqrt{|g|}[\gamma^\mu, \Omega_\mu]$, as the term $3H\gamma^0\Psi$. The Euler-Lagrange equations derived from the full $\mathcal{L}$, with $\Psi^\dagger$ and Ψ varied independently, are those of Section 10 (checks P_EL_fromLagrangianPsiDagger and P_EL_fromLagrangianPsi).

9.2 WolframScript form

In wolfram/Dirac16ComplexPrimordial.wl the field, the connection and the Lagrangian are built as follows. G[a] is γ^a , C16 is C , Sab[a, b] is S^{ab} , omLF is $\omega_{\mu ab}$ of Section 6 (all lists are 1-based), mm is m and lam is λ . Ps[n] and Pb[n] are the components of Ψ and $\Psi^\dagger$, treated as independent formal symbols in a fixed order ($\Psi^\dagger$ on the left, Ψ on the right).

```
Yv = {z, x1, x2, x3, t, x5, x6, x7}; (* z = 6 H x0, t = H x4 *)
jac = {6 H, 1, 1, 1, H, 1, 1, 1};
d1[k_Integer, f_] := jac[[k]] D[f, Yv[[k]]]; (* partial w.r.t. x^(k-1) *)
sF = Sin[z];
hhF = {Cot[z], sF^(1/6) E^a4[t], sF^(1/6) E^a4[t], sF^(1/6) E^a4[t], 1,
  sF^(1/6) E^(-a4[t]), sF^(1/6) E^(-a4[t]), sF^(1/6) E^(-a4[t])};
eiF = DiagonalMatrix[1/hhF]; (* e_a^mu *)
sqrtgF = Cos[z]; (* sqrt|g| *)
gUF = Table[Sum[eiF[[a, m]] G[a - 1], {a, 8}], {m, 8}]; (* gamma^mu *)
OmF = Table[(1/2) Sum[omLF[[m, a, b]] Sab[a - 1, b - 1], {a, 8}, {b, 8}], {m, 8}];
psi = Table[Ps[n] @@ Yv, {n, 0, 15}];
psib = Table[Pb[n] @@ Yv, {n, 0, 15}];
S0 = psib.C16.psi; (* S = Psibar Psi *)
Dpsi = Table[d1[mu, psi] + OmF[[mu]].psi, {mu, 8}];
Dpsib = Table[d1[mu, psib.C16] - psib.C16.OmF[[mu]], {mu, 8}];
Lag = sqrtgF ((1/2) Sum[psib.C16.gUF[[mu]].Dpsi[[mu]]
  - Dpsib[[mu]].gUF[[mu]].psi, {mu, 8}] - mm S0 - (lam/2) S0^2);
ELb = Table[D[Lag, Pb[a] @@ Yv]
  - Sum[D[D[Lag, D[Pb[a] @@ Yv, Yv[[k]]]], Yv[[k]]], {k, 8}], {a, 0, 15}];
(* check P_EL_fromLagrangianPsiDagger: ELb == cos z C.(gamma^mu D_mu psi - M_eff psi) *)
```

9.3 Relation to the notebook Lagrangian

For a Grassmann-odd Ψ the notebook's Lagrangian $\text{Lg}[]$ is a pure divergence: its mass term vanishes identically because σ_{16} is symmetric, and its kinetic and connection terms cancel up to a total derivative (Stage 1, also checked at exact points of this field). The notebook treats $\text{f16}[k]$ as commuting functions, which is why its equations are not empty. Section 11 compares them with the correct ones.

10 The sixteen component Euler-Lagrange equations

10.1 Matrix form

The field equations are

$$\gamma^\mu D_\mu \Psi = (m + U'(S))\Psi, \quad (D_\mu \bar{\Psi})\gamma^\mu = -(m + U'(S))\bar{\Psi}.$$

In this field, with $i = 1, 2, 3$ and $j = 5, 6, 7$,

$$\begin{aligned} \tan z \gamma^0 \partial_0 \Psi + s^{-1/6} e^{-a_4} \gamma^i \partial_i \Psi + \gamma^4 \partial_4 \Psi + s^{-1/6} e^{a_4} \gamma^j \partial_j \Psi \\ + 3H \gamma^0 \Psi = (m + \lambda S) \Psi. \end{aligned}$$

10.2 Component form in all eight coordinates

Equation (En) is row n of the matrix equation. Because every row of every γ^a has exactly one entry ± 1 , each equation has exactly ten terms: one derivative term for each of the eight coordinates, the term $3H(\gamma^0)_{nk}\Psi_k$ and the mass term. The coefficients are $\tan z$ for ∂_0 , $s^{-1/6}e^{-a_4}$ for $\partial_1, \partial_2, \partial_3$, 1 for ∂_4 , $s^{-1/6}e^{a_4}$ for $\partial_5, \partial_6, \partial_7$, then $3H$ and $m + \lambda S$. Here $\partial_\mu = \partial/\partial x^\mu$, $z = 6Hx_0$, $s = \sin z$ and $a_4 = a_4(Hx_4)$. The TeX of every equation below is taken verbatim from `primordial-components.json`; only the line breaks are chosen here.

$$\begin{aligned} (\tan z \partial_0 + 3H) \Psi_8 - \partial_4 \Psi_{13} \\ + s^{-1/6} e^{-a_4} (-\partial_1 \Psi_{15} + \partial_2 \Psi_{14} - \partial_3 \Psi_{13}) \\ + s^{-1/6} e^{a_4} (\partial_5 \Psi_{14} + \partial_6 \Psi_{15} + \partial_7 \Psi_8) = (m + \lambda S) \Psi_0 \end{aligned} \tag{E0}$$

$$\begin{aligned} (\tan z \partial_0 + 3H) \Psi_9 + \partial_4 \Psi_{12} \\ + s^{-1/6} e^{-a_4} (-\partial_1 \Psi_{14} - \partial_2 \Psi_{15} + \partial_3 \Psi_{12}) \\ + s^{-1/6} e^{a_4} (\partial_5 \Psi_{15} - \partial_6 \Psi_{14} + \partial_7 \Psi_9) = (m + \lambda S) \Psi_1 \end{aligned} \tag{E1}$$

$$\begin{aligned} (\tan z \partial_0 + 3H) \Psi_{10} + \partial_4 \Psi_{15} \\ + s^{-1/6} e^{-a_4} (\partial_1 \Psi_{13} - \partial_2 \Psi_{12} - \partial_3 \Psi_{15}) \\ + s^{-1/6} e^{a_4} (-\partial_5 \Psi_{12} + \partial_6 \Psi_{13} + \partial_7 \Psi_{10}) = (m + \lambda S) \Psi_2 \end{aligned} \tag{E2}$$

$$\begin{aligned}
& (\tan z \partial_0 + 3H) \Psi_{11} - \partial_4 \Psi_{14} \\
& + s^{-1/6} e^{-a_4} (\partial_1 \Psi_{12} + \partial_2 \Psi_{13} + \partial_3 \Psi_{14}) \\
& + s^{-1/6} e^{a_4} (-\partial_5 \Psi_{13} - \partial_6 \Psi_{12} + \partial_7 \Psi_{11}) = (m + \lambda S) \Psi_3
\end{aligned} \tag{E3}$$

$$\begin{aligned}
& (\tan z \partial_0 + 3H) \Psi_{12} + \partial_4 \Psi_9 \\
& + s^{-1/6} e^{-a_4} (-\partial_1 \Psi_{11} + \partial_2 \Psi_{10} - \partial_3 \Psi_9) \\
& + s^{-1/6} e^{a_4} (-\partial_5 \Psi_{10} - \partial_6 \Psi_{11} - \partial_7 \Psi_{12}) = (m + \lambda S) \Psi_4
\end{aligned} \tag{E4}$$

$$\begin{aligned}
& (\tan z \partial_0 + 3H) \Psi_{13} - \partial_4 \Psi_8 \\
& + s^{-1/6} e^{-a_4} (-\partial_1 \Psi_{10} - \partial_2 \Psi_{11} + \partial_3 \Psi_8) \\
& + s^{-1/6} e^{a_4} (-\partial_5 \Psi_{11} + \partial_6 \Psi_{10} - \partial_7 \Psi_{13}) = (m + \lambda S) \Psi_5
\end{aligned} \tag{E5}$$

$$\begin{aligned}
& (\tan z \partial_0 + 3H) \Psi_{14} - \partial_4 \Psi_{11} \\
& + s^{-1/6} e^{-a_4} (\partial_1 \Psi_9 - \partial_2 \Psi_8 - \partial_3 \Psi_{11}) \\
& + s^{-1/6} e^{a_4} (\partial_5 \Psi_8 - \partial_6 \Psi_9 - \partial_7 \Psi_{14}) = (m + \lambda S) \Psi_6
\end{aligned} \tag{E6}$$

$$\begin{aligned}
& (\tan z \partial_0 + 3H) \Psi_{15} + \partial_4 \Psi_{10} \\
& + s^{-1/6} e^{-a_4} (\partial_1 \Psi_8 + \partial_2 \Psi_9 + \partial_3 \Psi_{10}) \\
& + s^{-1/6} e^{a_4} (\partial_5 \Psi_9 + \partial_6 \Psi_8 - \partial_7 \Psi_{15}) = (m + \lambda S) \Psi_7
\end{aligned} \tag{E7}$$

$$\begin{aligned}
& (\tan z \partial_0 + 3H) \Psi_0 + \partial_4 \Psi_5 \\
& + s^{-1/6} e^{-a_4} (\partial_1 \Psi_7 - \partial_2 \Psi_6 + \partial_3 \Psi_5) \\
& + s^{-1/6} e^{a_4} (-\partial_5 \Psi_6 - \partial_6 \Psi_7 - \partial_7 \Psi_0) = (m + \lambda S) \Psi_8
\end{aligned} \tag{E8}$$

$$\begin{aligned}
& (\tan z \partial_0 + 3H) \Psi_1 - \partial_4 \Psi_4 \\
& + s^{-1/6} e^{-a_4} (\partial_1 \Psi_6 + \partial_2 \Psi_7 - \partial_3 \Psi_4) \\
& + s^{-1/6} e^{a_4} (-\partial_5 \Psi_7 + \partial_6 \Psi_6 - \partial_7 \Psi_1) = (m + \lambda S) \Psi_9
\end{aligned} \tag{E9}$$

$$\begin{aligned}
& (\tan z \partial_0 + 3H) \Psi_2 - \partial_4 \Psi_7 \\
& + s^{-1/6} e^{-a_4} (-\partial_1 \Psi_5 + \partial_2 \Psi_4 + \partial_3 \Psi_7) \\
& + s^{-1/6} e^{a_4} (\partial_5 \Psi_4 - \partial_6 \Psi_5 - \partial_7 \Psi_2) = (m + \lambda S) \Psi_{10}
\end{aligned} \tag{E10}$$

$$\begin{aligned}
& (\tan z \partial_0 + 3H) \Psi_3 + \partial_4 \Psi_6 \\
& + s^{-1/6} e^{-a_4} (-\partial_1 \Psi_4 - \partial_2 \Psi_5 - \partial_3 \Psi_6) \\
& + s^{-1/6} e^{a_4} (\partial_5 \Psi_5 + \partial_6 \Psi_4 - \partial_7 \Psi_3) = (m + \lambda S) \Psi_{11}
\end{aligned} \tag{E11}$$

$$\begin{aligned}
& (\tan z \partial_0 + 3H) \Psi_4 - \partial_4 \Psi_1 \\
& + s^{-1/6} e^{-a_4} (\partial_1 \Psi_3 - \partial_2 \Psi_2 + \partial_3 \Psi_1) \\
& + s^{-1/6} e^{a_4} (\partial_5 \Psi_2 + \partial_6 \Psi_3 + \partial_7 \Psi_4) = (m + \lambda S) \Psi_{12}
\end{aligned} \tag{E12}$$

$$\begin{aligned}
& (\tan z \partial_0 + 3H) \Psi_5 + \partial_4 \Psi_0 \\
& + s^{-1/6} e^{-a_4} (\partial_1 \Psi_2 + \partial_2 \Psi_3 - \partial_3 \Psi_0) \\
& + s^{-1/6} e^{a_4} (\partial_5 \Psi_3 - \partial_6 \Psi_2 + \partial_7 \Psi_5) = (m + \lambda S) \Psi_{13}
\end{aligned} \tag{E13}$$

$$\begin{aligned}
& (\tan z \partial_0 + 3H) \Psi_6 + \partial_4 \Psi_3 \\
& + s^{-1/6} e^{-a_4} (-\partial_1 \Psi_1 + \partial_2 \Psi_0 + \partial_3 \Psi_3) \\
& + s^{-1/6} e^{a_4} (-\partial_5 \Psi_0 + \partial_6 \Psi_1 + \partial_7 \Psi_6) = (m + \lambda S) \Psi_{14}
\end{aligned} \tag{E14}$$

$$\begin{aligned}
& (\tan z \partial_0 + 3H) \Psi_7 - \partial_4 \Psi_2 \\
& + s^{-1/6} e^{-a_4} (-\partial_1 \Psi_0 - \partial_2 \Psi_1 - \partial_3 \Psi_2) \\
& + s^{-1/6} e^{a_4} (-\partial_5 \Psi_1 - \partial_6 \Psi_0 + \partial_7 \Psi_7) = (m + \lambda S) \Psi_{15}
\end{aligned} \tag{E15}$$

10.3 Other presentations of the equations

The same equations in explicit x notation and in the ζ form follow by the substitutions $\tan z \partial_0 = \partial_\zeta$ and $s^{-1/6} = e^{-H\zeta}$; the component file contains all four presentations (x form, main form, ζ form, z, t form) for each row. Row 0 as an example:

$$\begin{aligned}
& (\tan(6Hx_0) \partial_{x_0} + 3H) \Psi_8 - \partial_{x_4} \Psi_{13} \\
& + \sin^{-1/6}(6Hx_0) e^{-a_4(Hx_4)} (-\partial_{x_1} \Psi_{15} + \partial_{x_2} \Psi_{14} - \partial_{x_3} \Psi_{13}) \\
& + \sin^{-1/6}(6Hx_0) e^{a_4(Hx_4)} (\partial_{x_5} \Psi_{14} + \partial_{x_6} \Psi_{15} + \partial_{x_7} \Psi_8) = (m + \lambda S) \Psi_0
\end{aligned} \tag{E0: x form}$$

$$\begin{aligned}
& (\partial_\zeta + 3H) \Psi_8 - \partial_4 \Psi_{13} \\
& + e^{-H\zeta - a_4} (-\partial_1 \Psi_{15} + \partial_2 \Psi_{14} - \partial_3 \Psi_{13}) \\
& + e^{-H\zeta + a_4} (\partial_5 \Psi_{14} + \partial_6 \Psi_{15} + \partial_7 \Psi_8) = (m + \lambda S) \Psi_0
\end{aligned} \tag{E0: zeta form}$$

$$\begin{aligned}
& (6H \tan z \partial_z + 3H) \Psi_8 - H \partial_t \Psi_{13} \\
& + s^{-1/6} e^{-a_4} (-\partial_1 \Psi_{15} + \partial_2 \Psi_{14} - \partial_3 \Psi_{13}) \\
& + s^{-1/6} e^{a_4} (\partial_5 \Psi_{14} + \partial_6 \Psi_{15} + \partial_7 \Psi_8) = (m + \lambda S) \Psi_0
\end{aligned} \tag{E0: z, t form}$$

10.4 Evolution form

Multiplying by $-\gamma^4$ (with $\gamma^4 \gamma^4 = -1$) solves for the x_4 derivative:

$$\partial_4 \Psi = -(m + \lambda S) \gamma^4 \Psi + 3H \gamma^4 \gamma^0 \Psi + \sum_{\mu \neq 4} h_\mu^{-1} \gamma^4 \gamma^\mu \partial_\mu \Psi.$$

In components, each equation (Vn) has nine terms:

$$\begin{aligned}
\partial_4 \Psi_0 &= -\tan z \partial_0 \Psi_5 \\
&- s^{-1/6} e^{-a_4} \partial_1 \Psi_2 - s^{-1/6} e^{-a_4} \partial_2 \Psi_3 + s^{-1/6} e^{-a_4} \partial_3 \Psi_0 \\
&- s^{-1/6} e^{a_4} \partial_5 \Psi_3 + s^{-1/6} e^{a_4} \partial_6 \Psi_2 - s^{-1/6} e^{a_4} \partial_7 \Psi_5 \\
&- 3H \Psi_5 + (m + \lambda S) \Psi_{13}
\end{aligned} \tag{V0}$$

$$\begin{aligned}
\partial_4 \Psi_1 &= \tan z \partial_0 \Psi_4 \\
&+ s^{-1/6} e^{-a_4} \partial_1 \Psi_3 - s^{-1/6} e^{-a_4} \partial_2 \Psi_2 + s^{-1/6} e^{-a_4} \partial_3 \Psi_1 \\
&+ s^{-1/6} e^{a_4} \partial_5 \Psi_2 + s^{-1/6} e^{a_4} \partial_6 \Psi_3 + s^{-1/6} e^{a_4} \partial_7 \Psi_4 \\
&+ 3H \Psi_4 - (m + \lambda S) \Psi_{12}
\end{aligned} \tag{V1}$$

$$\begin{aligned}
\partial_4 \Psi_2 &= \tan z \partial_0 \Psi_7 \\
&- s^{-1/6} e^{-a_4} \partial_1 \Psi_0 - s^{-1/6} e^{-a_4} \partial_2 \Psi_1 - s^{-1/6} e^{-a_4} \partial_3 \Psi_2 \\
&- s^{-1/6} e^{a_4} \partial_5 \Psi_1 - s^{-1/6} e^{a_4} \partial_6 \Psi_0 + s^{-1/6} e^{a_4} \partial_7 \Psi_7 \\
&+ 3H \Psi_7 - (m + \lambda S) \Psi_{15}
\end{aligned} \tag{V2}$$

$$\begin{aligned}
\partial_4 \Psi_3 &= -\tan z \partial_0 \Psi_6 \\
&+ s^{-1/6} e^{-a_4} \partial_1 \Psi_1 - s^{-1/6} e^{-a_4} \partial_2 \Psi_0 - s^{-1/6} e^{-a_4} \partial_3 \Psi_3 \\
&+ s^{-1/6} e^{a_4} \partial_5 \Psi_0 - s^{-1/6} e^{a_4} \partial_6 \Psi_1 - s^{-1/6} e^{a_4} \partial_7 \Psi_6 \\
&- 3H \Psi_6 + (m + \lambda S) \Psi_{14}
\end{aligned} \tag{V3}$$

$$\begin{aligned}
\partial_4 \Psi_4 &= \tan z \partial_0 \Psi_1 \\
&+ s^{-1/6} e^{-a_4} \partial_1 \Psi_6 + s^{-1/6} e^{-a_4} \partial_2 \Psi_7 - s^{-1/6} e^{-a_4} \partial_3 \Psi_4 \\
&- s^{-1/6} e^{a_4} \partial_5 \Psi_7 + s^{-1/6} e^{a_4} \partial_6 \Psi_6 - s^{-1/6} e^{a_4} \partial_7 \Psi_1 \\
&+ 3H \Psi_1 - (m + \lambda S) \Psi_9
\end{aligned} \tag{V4}$$

$$\begin{aligned}
\partial_4 \Psi_5 &= -\tan z \partial_0 \Psi_0 \\
&- s^{-1/6} e^{-a_4} \partial_1 \Psi_7 + s^{-1/6} e^{-a_4} \partial_2 \Psi_6 - s^{-1/6} e^{-a_4} \partial_3 \Psi_5 \\
&+ s^{-1/6} e^{a_4} \partial_5 \Psi_6 + s^{-1/6} e^{a_4} \partial_6 \Psi_7 + s^{-1/6} e^{a_4} \partial_7 \Psi_0 \\
&- 3H \Psi_0 + (m + \lambda S) \Psi_8
\end{aligned} \tag{V5}$$

$$\begin{aligned}
\partial_4 \Psi_6 &= -\tan z \partial_0 \Psi_3 \\
&+ s^{-1/6} e^{-a_4} \partial_1 \Psi_4 + s^{-1/6} e^{-a_4} \partial_2 \Psi_5 + s^{-1/6} e^{-a_4} \partial_3 \Psi_6 \\
&- s^{-1/6} e^{a_4} \partial_5 \Psi_5 - s^{-1/6} e^{a_4} \partial_6 \Psi_4 + s^{-1/6} e^{a_4} \partial_7 \Psi_3 \\
&- 3H \Psi_3 + (m + \lambda S) \Psi_{11}
\end{aligned} \tag{V6}$$

$$\begin{aligned}
\partial_4 \Psi_7 &= \tan z \partial_0 \Psi_2 \\
&- s^{-1/6} e^{-a_4} \partial_1 \Psi_5 + s^{-1/6} e^{-a_4} \partial_2 \Psi_4 + s^{-1/6} e^{-a_4} \partial_3 \Psi_7 \\
&+ s^{-1/6} e^{a_4} \partial_5 \Psi_4 - s^{-1/6} e^{a_4} \partial_6 \Psi_5 - s^{-1/6} e^{a_4} \partial_7 \Psi_2 \\
&+ 3H \Psi_2 - (m + \lambda S) \Psi_{10}
\end{aligned} \tag{V7}$$

$$\begin{aligned}
\partial_4 \Psi_8 &= \tan z \partial_0 \Psi_{13} \\
&- s^{-1/6} e^{-a_4} \partial_1 \Psi_{10} - s^{-1/6} e^{-a_4} \partial_2 \Psi_{11} + s^{-1/6} e^{-a_4} \partial_3 \Psi_8 \\
&- s^{-1/6} e^{a_4} \partial_5 \Psi_{11} + s^{-1/6} e^{a_4} \partial_6 \Psi_{10} - s^{-1/6} e^{a_4} \partial_7 \Psi_{13} \\
&+ 3H \Psi_{13} - (m + \lambda S) \Psi_5
\end{aligned} \tag{V8}$$

$$\begin{aligned}
\partial_4 \Psi_9 &= -\tan z \partial_0 \Psi_{12} \\
&+ s^{-1/6} e^{-a_4} \partial_1 \Psi_{11} - s^{-1/6} e^{-a_4} \partial_2 \Psi_{10} + s^{-1/6} e^{-a_4} \partial_3 \Psi_9 \\
&+ s^{-1/6} e^{a_4} \partial_5 \Psi_{10} + s^{-1/6} e^{a_4} \partial_6 \Psi_{11} + s^{-1/6} e^{a_4} \partial_7 \Psi_{12} \\
&- 3H \Psi_{12} + (m + \lambda S) \Psi_4
\end{aligned} \tag{V9}$$

$$\begin{aligned}
\partial_4 \Psi_{10} &= -\tan z \partial_0 \Psi_{15} \\
&- s^{-1/6} e^{-a_4} \partial_1 \Psi_8 - s^{-1/6} e^{-a_4} \partial_2 \Psi_9 - s^{-1/6} e^{-a_4} \partial_3 \Psi_{10} \\
&- s^{-1/6} e^{a_4} \partial_5 \Psi_9 - s^{-1/6} e^{a_4} \partial_6 \Psi_8 + s^{-1/6} e^{a_4} \partial_7 \Psi_{15} \\
&- 3H \Psi_{15} + (m + \lambda S) \Psi_7
\end{aligned} \tag{V10}$$

$$\begin{aligned}
\partial_4 \Psi_{11} &= \tan z \partial_0 \Psi_{14} \\
&+ s^{-1/6} e^{-a_4} \partial_1 \Psi_9 - s^{-1/6} e^{-a_4} \partial_2 \Psi_8 - s^{-1/6} e^{-a_4} \partial_3 \Psi_{11} \\
&+ s^{-1/6} e^{a_4} \partial_5 \Psi_8 - s^{-1/6} e^{a_4} \partial_6 \Psi_9 - s^{-1/6} e^{a_4} \partial_7 \Psi_{14} \\
&+ 3H \Psi_{14} - (m + \lambda S) \Psi_6
\end{aligned} \tag{V11}$$

$$\begin{aligned}
\partial_4 \Psi_{12} &= -\tan z \partial_0 \Psi_9 \\
&+ s^{-1/6} e^{-a_4} \partial_1 \Psi_{14} + s^{-1/6} e^{-a_4} \partial_2 \Psi_{15} - s^{-1/6} e^{-a_4} \partial_3 \Psi_{12} \\
&- s^{-1/6} e^{a_4} \partial_5 \Psi_{15} + s^{-1/6} e^{a_4} \partial_6 \Psi_{14} - s^{-1/6} e^{a_4} \partial_7 \Psi_9 \\
&- 3H \Psi_9 + (m + \lambda S) \Psi_1
\end{aligned} \tag{V12}$$

$$\begin{aligned}
\partial_4 \Psi_{13} &= \tan z \partial_0 \Psi_8 \\
&- s^{-1/6} e^{-a_4} \partial_1 \Psi_{15} + s^{-1/6} e^{-a_4} \partial_2 \Psi_{14} - s^{-1/6} e^{-a_4} \partial_3 \Psi_{13} \\
&+ s^{-1/6} e^{a_4} \partial_5 \Psi_{14} + s^{-1/6} e^{a_4} \partial_6 \Psi_{15} + s^{-1/6} e^{a_4} \partial_7 \Psi_8 \\
&+ 3H \Psi_8 - (m + \lambda S) \Psi_0
\end{aligned} \tag{V13}$$

$$\begin{aligned}
\partial_4 \Psi_{14} &= \tan z \partial_0 \Psi_{11} \\
&+ s^{-1/6} e^{-a_4} \partial_1 \Psi_{12} + s^{-1/6} e^{-a_4} \partial_2 \Psi_{13} + s^{-1/6} e^{-a_4} \partial_3 \Psi_{14} \\
&- s^{-1/6} e^{a_4} \partial_5 \Psi_{13} - s^{-1/6} e^{a_4} \partial_6 \Psi_{12} + s^{-1/6} e^{a_4} \partial_7 \Psi_{11} \\
&+ 3H \Psi_{11} - (m + \lambda S) \Psi_3
\end{aligned} \tag{V14}$$

$$\begin{aligned}
\partial_4 \Psi_{15} &= -\tan z \partial_0 \Psi_{10} \\
&- s^{-1/6} e^{-a_4} \partial_1 \Psi_{13} + s^{-1/6} e^{-a_4} \partial_2 \Psi_{12} + s^{-1/6} e^{-a_4} \partial_3 \Psi_{15} \\
&+ s^{-1/6} e^{a_4} \partial_5 \Psi_{12} - s^{-1/6} e^{a_4} \partial_6 \Psi_{13} - s^{-1/6} e^{a_4} \partial_7 \Psi_{10} \\
&- 3H \Psi_{10} + (m + \lambda S) \Psi_2
\end{aligned} \tag{V15}$$

10.5 The (x0, x4) block form and the z, t evolution form

For fields that depend on x_0 and x_4 only, the equations reduce to

$$\tan z \gamma^0 \partial_0 \Psi + \gamma^4 \partial_4 \Psi + 3H \gamma^0 \Psi = (m + \lambda S) \Psi.$$

No a_4 appears: $\gamma^\mu \Omega_\mu = 3H \gamma^0$ is independent of a_4 , and the coefficients that contain a_4 multiply $\partial_1, \dots, \partial_3$ and $\partial_5, \dots, \partial_7$ only. The coupling graph of γ^0 and γ^4 splits the 16 components into four blocks of four,

$$\{0, 5, 8, 13\}, \quad \{1, 4, 9, 12\}, \quad \{2, 7, 10, 15\}, \quad \{3, 6, 11, 14\},$$

each with two components of chirality -1 and two of chirality $+1$. These are exactly the notebook's coupling sets (cell 1089). For $\lambda \neq 0$ the blocks are coupled only through the scalar S , a bilinear in all 16 components; the linear operator is block diagonal. In the variables z, t ($\partial_0 = 6H \partial_z$, $\partial_4 = H \partial_t$) the evolution form of the four blocks is:

$$\begin{aligned} H \partial_t \Psi_0 &= -(6H \tan z \partial_z + 3H) \Psi_5 + (m + \lambda S) \Psi_{13} \\ H \partial_t \Psi_5 &= -(6H \tan z \partial_z + 3H) \Psi_0 + (m + \lambda S) \Psi_8 \\ H \partial_t \Psi_8 &= (6H \tan z \partial_z + 3H) \Psi_{13} - (m + \lambda S) \Psi_5 \\ H \partial_t \Psi_{13} &= (6H \tan z \partial_z + 3H) \Psi_8 - (m + \lambda S) \Psi_0 \end{aligned} \quad (\text{block } 0,5,8,13)$$

$$\begin{aligned} H \partial_t \Psi_1 &= (6H \tan z \partial_z + 3H) \Psi_4 - (m + \lambda S) \Psi_{12} \\ H \partial_t \Psi_4 &= (6H \tan z \partial_z + 3H) \Psi_1 - (m + \lambda S) \Psi_9 \\ H \partial_t \Psi_9 &= -(6H \tan z \partial_z + 3H) \Psi_{12} + (m + \lambda S) \Psi_4 \\ H \partial_t \Psi_{12} &= -(6H \tan z \partial_z + 3H) \Psi_9 + (m + \lambda S) \Psi_1 \end{aligned} \quad (\text{block } 1,4,9,12)$$

$$\begin{aligned} H \partial_t \Psi_2 &= (6H \tan z \partial_z + 3H) \Psi_7 - (m + \lambda S) \Psi_{15} \\ H \partial_t \Psi_7 &= (6H \tan z \partial_z + 3H) \Psi_2 - (m + \lambda S) \Psi_{10} \\ H \partial_t \Psi_{10} &= -(6H \tan z \partial_z + 3H) \Psi_{15} + (m + \lambda S) \Psi_7 \\ H \partial_t \Psi_{15} &= -(6H \tan z \partial_z + 3H) \Psi_{10} + (m + \lambda S) \Psi_2 \end{aligned} \quad (\text{block } 2,7,10,15)$$

$$\begin{aligned} H \partial_t \Psi_3 &= -(6H \tan z \partial_z + 3H) \Psi_6 + (m + \lambda S) \Psi_{14} \\ H \partial_t \Psi_6 &= -(6H \tan z \partial_z + 3H) \Psi_3 + (m + \lambda S) \Psi_{11} \\ H \partial_t \Psi_{11} &= (6H \tan z \partial_z + 3H) \Psi_{14} - (m + \lambda S) \Psi_6 \\ H \partial_t \Psi_{14} &= (6H \tan z \partial_z + 3H) \Psi_{11} - (m + \lambda S) \Psi_3 \end{aligned} \quad (\text{block } 3,6,11,14)$$

11 Comparison with the notebook's cell-1137 equations

11.1 Variables and cross reference

The notebook's production Lagrangian $\text{La}[]$ (cell 1066) is, for commuting fields $\text{f16}[k](x_0, x_4)$,

$$\text{La} = \cos(6Hx_0) \left[\Psi_{16}^T \sigma_{16} \sum_{\alpha} \text{useT16}_{\alpha} \left(\partial_{\alpha} \Psi_{16} + \frac{Q_1}{2} \sum_{A,B} \omega_{\alpha}{}^A{}_B S^{AB} \Psi_{16} \right) + HM \Psi_{16}^T \sigma_{16} \Psi_{16} \right],$$

where useT16 are the curved gammas built in cell 1058, $\omega_{\alpha}{}^A{}_B$ is the mixed connection of cell 501, and Q_1 is the notebook's spin-connection switch. Its Euler-Lagrange expressions eL[] (cell 1075) give the stored eLa (cell 1079). Cell 1096 applies $(1/(2H))$, $\text{f16}[k](x_0, x_4) = Z[k](z, t)$ and $x_0 = z/(6H)$, $x_4 = t/H$. Cell 1111 relabels the components block by block into yZ, and cell 1137 stores the equations solved for ∂_t . The correct equations of this document (Grassmann fields, $U = 0$) are compared in the same variables with $M_{\text{notebook}} = -m/H$. Both use

$$q = Q_1 \sinh(a_4) a'_4 e^{-a_4} = \frac{1}{2} Q_1 a'_4 (1 - e^{-2a_4}), \quad a'_4 = \frac{da_4}{dt}$$

The cross reference between Ψ_k (the notebook's f16[k] and Z[k]), the yZ index, the block and the difference "stored minus correct" in the ∂_t equation:

yZ	Ψ_k	Block	Difference
yZ ₀	Ψ_0	{0, 5, 8, 13}	+q yZ ₀
yZ ₁	Ψ_5	{0, 5, 8, 13}	-q yZ ₁
yZ ₂	Ψ_8	{0, 5, 8, 13}	-q yZ ₂
yZ ₃	Ψ_{13}	{0, 5, 8, 13}	+q yZ ₃
yZ ₄	Ψ_1	{1, 4, 9, 12}	+q yZ ₄
yZ ₅	Ψ_4	{1, 4, 9, 12}	-q yZ ₅
yZ ₆	Ψ_9	{1, 4, 9, 12}	-q yZ ₆
yZ ₇	Ψ_{12}	{1, 4, 9, 12}	+q yZ ₇
yZ ₈	Ψ_2	{2, 7, 10, 15}	none
yZ ₉	Ψ_7	{2, 7, 10, 15}	none
yZ ₁₀	Ψ_{10}	{2, 7, 10, 15}	none
yZ ₁₁	Ψ_{15}	{2, 7, 10, 15}	none
yZ ₁₂	Ψ_3	{3, 6, 11, 14}	none
yZ ₁₃	Ψ_6	{3, 6, 11, 14}	none
yZ ₁₄	Ψ_{11}	{3, 6, 11, 14}	none
yZ ₁₅	Ψ_{14}	{3, 6, 11, 14}	none

11.2 The stored equations and the difference

The notebook's stored cell-1137 equations, reproduced exactly from the rebuilt chain for all 16 rows, are (M is the notebook's mass parameter and q is defined in Section 11.1):

$$\begin{aligned} \partial_t yZ_0 &= q yZ_0 - 3 yZ_1 - M yZ_3 - 6 \tan z \partial_z yZ_1 \\ \partial_t yZ_1 &= -3 yZ_0 - q yZ_1 - M yZ_2 - 6 \tan z \partial_z yZ_0 \\ \partial_t yZ_2 &= M yZ_1 - q yZ_2 + 3 yZ_3 + 6 \tan z \partial_z yZ_3 \\ \partial_t yZ_3 &= M yZ_0 + 3 yZ_2 + q yZ_3 + 6 \tan z \partial_z yZ_2 \end{aligned} \quad (\text{stored, block 0,5,8,13})$$

$$\begin{aligned}
\partial_t y Z_4 &= q y Z_4 + 3 y Z_5 + M y Z_7 + 6 \tan z \partial_z y Z_5 \\
\partial_t y Z_5 &= 3 y Z_4 - q y Z_5 + M y Z_6 + 6 \tan z \partial_z y Z_4 \\
\partial_t y Z_6 &= -M y Z_5 - q y Z_6 - 3 y Z_7 - 6 \tan z \partial_z y Z_7 \\
\partial_t y Z_7 &= -M y Z_4 - 3 y Z_6 + q y Z_7 - 6 \tan z \partial_z y Z_6
\end{aligned}
\tag{stored, block 1,4,9,12}$$

$$\begin{aligned}
\partial_t y Z_8 &= 3 y Z_9 + M y Z_{11} + 6 \tan z \partial_z y Z_9 \\
\partial_t y Z_9 &= 3 y Z_8 + M y Z_{10} + 6 \tan z \partial_z y Z_8 \\
\partial_t y Z_{10} &= -M y Z_9 - 3 y Z_{11} - 6 \tan z \partial_z y Z_{11} \\
\partial_t y Z_{11} &= -M y Z_8 - 3 y Z_{10} - 6 \tan z \partial_z y Z_{10}
\end{aligned}
\tag{stored, block 2,7,10,15}$$

$$\begin{aligned}
\partial_t y Z_{12} &= -3 y Z_{13} - M y Z_{15} - 6 \tan z \partial_z y Z_{13} \\
\partial_t y Z_{13} &= -3 y Z_{12} - M y Z_{14} - 6 \tan z \partial_z y Z_{12} \\
\partial_t y Z_{14} &= M y Z_{13} + 3 y Z_{15} + 6 \tan z \partial_z y Z_{15} \\
\partial_t y Z_{15} &= M y Z_{12} + 3 y Z_{14} + 6 \tan z \partial_z y Z_{14}
\end{aligned}
\tag{stored, block 3,6,11,14}$$

The correct equations are the same with every q removed. Every other term agrees exactly. In the correct theory no a_4 dependence at all survives for fields of (x_0, x_4) , in agreement with Section 10.5. The blocks $\{2, 7, 10, 15\}$ and $\{3, 6, 11, 14\}$ ($y Z_8, \dots, y Z_{15}$) contain no q and are identical.

11.3 Origin of the q term: the cell-1058 substitution rule

Three facts are established by exact computation.

1. With Clifford-consistent curved gammas $\gamma^\mu = e_a^\mu \gamma^a$, every Q_1 term drops out of the commuting-field Euler-Lagrange equations, both for the notebook contraction and for the correct Ω_μ . The reason is that $\sigma_{16} \gamma^\mu \Omega_\mu$ has no symmetric part for a diagonal vielbein. For commuting fields the gravitational term of the notebook's equations comes from $\partial_\mu(\sqrt{|g|} \gamma^\mu)$, not from the spin connection.
2. The stored eLa (cell 1079) contains Q_1 terms in exactly the 8 rows of the coupling sets $\{0, 5, 8, 13\}$ and $\{1, 4, 9, 12\}$.
3. These terms are reproduced exactly, 16/16 rows with zero residual, by the reconstruction in which the curved gammas $\gamma^{x_5}, \gamma^{x_6}, \gamma^{x_7}$ of useT16 carry the factor $e^{-a_4} s^{-1/6}$ on their +1 entries and $e^{+a_4} s^{-1/6}$ on their -1 entries:

$$\gamma'^{x_j} = s^{-1/6} (\cosh(a_4) \gamma^j - \sinh(a_4) |\gamma^j|) \quad (j = 5, 6, 7),$$

where $|\gamma^j|$ is the entrywise absolute value. The correct curved gamma is $e^{+a_4} s^{-1/6} \gamma^j$. The same reconstruction, carried through the notebook's own chain, reproduces the stored cell-1096 output and the stored cell-1137 equations exactly (16/16), and the coupling sets (cell 1089) and the relabelling (cell 1111) match.

The rule. Cell 1058 contains the substitution rule

$$1/\text{Sqrt}[\text{Sin}[6*H*x0]^{(1/3)}/E^{(2*a4[H*x4])}] \rightarrow 1/(E^{a4[H*x4]}*\text{Sin}[6*H*x0]^{(1/6)})$$

which is mathematically wrong. The left-hand side equals $e^{a_4} s^{-1/6}$, so the correct rule is

$$1/\text{Sqrt}[\text{Sin}[6*H*x0]^{(1/3)}/E^{(2*a4[H*x4])}] \rightarrow E^{a4[H*x4]}/\text{Sin}[6*H*x0]^{(1/6)}$$

The stored eLa is consistent with this rule having hit only the +1 entries of $\gamma^{x_5}, \gamma^{x_6}, \gamma^{x_7}$ in the stored session. The notebook does not store the value of useT16. The exact 16/16 reproduction of the stored cell-1079 eLa confirms the reconstructed useT16 of fact 3; that the rule of cell 1058 produced it by acting on the +1 entries only is a reconstruction. When cell 1058 is re-executed literally in this kernel (Mathematica 15.0.1), the wrong factor e^{-a_4} is applied uniformly to all entries of $\gamma^{x_5}, \gamma^{x_6}, \gamma^{x_7}$ (the other five curved gammas come out correct). The resulting equations then contain no q term, and they differ from the stored eLa by exactly the q terms.

The notebook's metadata bear on this difference. The In labels of the chain increase monotonically, from In[1024] for cell 1058 to In[1113] for cell 1137 (Section 2), which indicates that the chain ran in one kernel session. Cell 1096 stores the date of that evaluation, 2026-01-30, and the last recorded change of cell 1058 is dated 2025-12-04. That session therefore ran a kernel build older than the verifier's, which is 15.0.1 of July 2, 2026. The form that FullSimplify gives the -1 entries, and hence whether the rule matches them, can differ between versions. This explanation is a reconstruction as well.

Mechanism. The reconstructed γ'^{x_j} is not in the Clifford algebra: $(\gamma'^{x_5})^2 = -s^{-1/3} \cdot 1$, whereas the Clifford relation requires $g^{55} = -s^{-1/3} e^{2a_4}$, and γ'^{x_5} anticommutes neither with γ'^{x_6} nor with γ^{x_0} . Consequently $\sigma_{16} \gamma'^{x_j} S^{4j}$ acquires a symmetric part, which commuting fields do not annihilate. Call the q vector the difference between the stored eLa and the Clifford-consistent rebuild, in the normalisation of cell 1079. It equals exactly

$$Q_1(\sigma_{16}Y + (\sigma_{16}Y)^T)\Psi, \quad Y = \sum_{j=5}^7 (\gamma'^{x_j} - \gamma^{x_j}) \Omega_j^{\text{NB}},$$

with $\Omega_j^{\text{NB}} = \frac{1}{2} \omega_j^{a_b} S^{ab}$ as in Section 8.3; equivalently $\frac{Q_1}{2}(\dots)$ with Y built from $\omega_j^{a_b} S^{ab}$ (check P_notebookCompare_qTermFromNonCliffordExtraTimeGammas). Its size is the difference of the two factors, $s^{-1/6}(e^{a_4} - e^{-a_4})$, times $\omega_j^{4_j} = H a'_4 s^{1/6} e^{-a_4}$, times Q_1 . That is $2Hq$, with $q = Q_1 \sinh(a_4) a'_4 e^{-a_4}$, and it becomes q after the factor $1/(2H)$ of cell 1096: each of the 8 nonzero rows of the q vector is $\pm 2Hq \Psi_k$ for one component k (check P_notebookCompare_qVectorIs2HqAtCell1079).

11.4 What this means

The notebook's block equations are correct in every term except q , once the Grassmann-correct Lagrangian is used and $M_{\text{notebook}} = -m/H$. The q term, and with it every a_4 dependence of the (x_0, x_4) equations, is an artifact of the non-Clifford curved gammas $\gamma'^{x_5}, \gamma'^{x_6}, \gamma'^{x_7}$; by the reconstruction of Section 11.3 these come from the wrong rule in cell 1058. Later steps of the notebook that remove or use the q term (a rescaling of the yZ and an equation for a_4 built from the same coefficient) therefore have no counterpart in the correct equations. This last remark is a reading of the notebook text, not a machine check.

12 The energy-momentum tensor

12.1 Definition and form in this field

The energy-momentum tensor of Stage 1 is

$$T_{\mu\nu} = -\frac{1}{4}[\bar{\Psi}\gamma_\mu D_\nu \Psi + \bar{\Psi}\gamma_\nu D_\mu \Psi - (D_\mu \bar{\Psi})\gamma_\nu \Psi - (D_\nu \bar{\Psi})\gamma_\mu \Psi] + g_{\mu\nu}\mathcal{L}_s, \quad \mathcal{L}_s = \mathcal{L}/\sqrt{|g|}.$$

Stage 1 derived it as $T_{\mu\nu} = -(2/\sqrt{|g|})\delta I/\delta g^{\mu\nu}$, with the action $I = \int \mathcal{L} d^8x$, by tetrad variation and showed that it is symmetric, Hermitian and covariantly conserved on shell, with $\mathcal{L}_s = SU'(S) - U(S)$ and trace $T^\mu{}_\mu = -mS + 7SU' - 8U$ on shell. The sign is fixed so that $T_{44} = \rho$. In this field, for Ψ depending on all eight coordinates and off shell,

$$T_{\mu\nu} = -\frac{1}{4}[\bar{\Psi}\gamma_\mu \partial_\nu \Psi + \bar{\Psi}\gamma_\nu \partial_\mu \Psi - \partial_\mu \bar{\Psi}\gamma_\nu \Psi - \partial_\nu \bar{\Psi}\gamma_\mu \Psi] - \frac{1}{4}\bar{\Psi}A_{\mu\nu}\Psi + g_{\mu\nu}\mathcal{L}_s,$$

with the lowered gammas of Section 4.2 and $A_{\mu\nu} = \{\gamma_\mu, \Omega_\nu\} + \{\gamma_\nu, \Omega_\mu\}$. The tensor is symmetric.

12.2 The 21 nonzero connection terms

Exactly 21 of the 36 matrices $A_{\mu\nu}$ with $\mu \leq \nu$ are nonzero, and each is a single product of three frame gammas:

(μ, ν)	Product	Coefficient of $A_{\mu\nu}$
(0, 1)	$\gamma^0\gamma^1\gamma^4$	$H \cot z s^{1/6} e^{a_4} a'_4$
(0, 2)	$\gamma^0\gamma^2\gamma^4$	$H \cot z s^{1/6} e^{a_4} a'_4$
(0, 3)	$\gamma^0\gamma^3\gamma^4$	$H \cot z s^{1/6} e^{a_4} a'_4$
(0, 5)	$\gamma^0\gamma^4\gamma^5$	$-H \cot z s^{1/6} e^{-a_4} a'_4$
(0, 6)	$\gamma^0\gamma^4\gamma^6$	$-H \cot z s^{1/6} e^{-a_4} a'_4$
(0, 7)	$\gamma^0\gamma^4\gamma^7$	$-H \cot z s^{1/6} e^{-a_4} a'_4$
(1, 4)	$\gamma^0\gamma^1\gamma^4$	$H s^{1/6} e^{a_4}$
(1, 5)	$\gamma^1\gamma^4\gamma^5$	$-2H s^{1/3} a'_4$
(1, 6)	$\gamma^1\gamma^4\gamma^6$	$-2H s^{1/3} a'_4$
(1, 7)	$\gamma^1\gamma^4\gamma^7$	$-2H s^{1/3} a'_4$
(2, 4)	$\gamma^0\gamma^2\gamma^4$	$H s^{1/6} e^{a_4}$
(2, 5)	$\gamma^2\gamma^4\gamma^5$	$-2H s^{1/3} a'_4$
(2, 6)	$\gamma^2\gamma^4\gamma^6$	$-2H s^{1/3} a'_4$
(2, 7)	$\gamma^2\gamma^4\gamma^7$	$-2H s^{1/3} a'_4$
(3, 4)	$\gamma^0\gamma^3\gamma^4$	$H s^{1/6} e^{a_4}$
(3, 5)	$\gamma^3\gamma^4\gamma^5$	$-2H s^{1/3} a'_4$
(3, 6)	$\gamma^3\gamma^4\gamma^6$	$-2H s^{1/3} a'_4$
(3, 7)	$\gamma^3\gamma^4\gamma^7$	$-2H s^{1/3} a'_4$
(4, 5)	$\gamma^0\gamma^4\gamma^5$	$H s^{1/6} e^{-a_4}$
(4, 6)	$\gamma^0\gamma^4\gamma^6$	$H s^{1/6} e^{-a_4}$

(μ, ν)	Product	Coefficient of $A_{\mu\nu}$
$(4, 7)$	$\gamma^0 \gamma^4 \gamma^7$	$H s^{1/6} e^{-a_4}$

All diagonal $A_{\mu\mu}$ vanish, as do A_{04} and $A_{ii'}$, $A_{jj'}$ within the groups $\{1, 2, 3\}$ and $\{5, 6, 7\}$.

12.3 Diagonal and off-diagonal structure

- **Transverse diagonal.** For every $\Psi(x_0, x_4)$, off shell and for every U , $T^i_i = T^j_j = \mathcal{L}_s$ for all six directions 1, 2, 3, 5, 6, 7 (check `P_source_transversePressuresEqualForEveryXOX4State`). On shell this is $SU'(S) - U(S)$.
- **Transverse off-diagonal.** For $\Psi(x_0, x_4)$ and $i \in \{1, 2, 3\}$, $j \in \{5, 6, 7\}$ the derivative terms vanish and $T_{ij} = -\frac{1}{4}\bar{\Psi}A_{ij}\Psi = \frac{1}{2}H s^{1/3}a'_4\bar{\Psi}\gamma^i\gamma^4\gamma^j\Psi$. This agrees with the Stage-1 formula $T_{ij} = \frac{1}{4}\epsilon_i\epsilon_j h_i h_j (H_i - H_j)\bar{\Psi}\gamma^i\gamma^j\gamma^4\Psi$ with $\epsilon_i = +1$, $\epsilon_j = -1$, $h_i h_j = s^{1/3}$ and the rates $H_i = H a'_4$, $H_j = -H a'_4$. Within a group the rates coincide and $T_{ii'} = T_{jj'} = 0$.
- **Components with 0 or 4.** T_{0i} , T_{0j} , T_{4i} , T_{4j} and T_{04} contain vector-type derivative bilinears and, except for T_{04} , connection terms: A_{0i} and A_{0j} come from the x_4 dependence (a'_4), and A_{i4} and A_{j4} come from the x_0 dependence (the hidden-space warp). The Stage-1 statement that T_{4i} vanishes on shell for homogeneous states in diagonal backgrounds assumes homogeneity in all transverse directions. Here the background depends on x_0 , and $T_{4i} \neq 0$ in general, even for the state of Section 13.7, which does not depend on x_0 .
- **Consistency with a diagonal metric.** The Einstein tensor of this field is diagonal (Section 15.1). A `dirac16complex` state can be a consistent source of it only if the expectation values of all these off-diagonal bilinears vanish. For the ζ plane wave of Section 13 with real K all 22 off-diagonal pairs $(0, 1), \dots, (0, 7)$, $(1, 4), \dots, (1, 7)$, $(2, 4), \dots, (2, 7)$, $(3, 4), \dots, (3, 7)$, $(4, 5), (4, 6), (4, 7)$ are nonzero for generic amplitudes. Of these only T_{04} , which is proportional to $K \cos z u^\dagger B u / \sin^2 z$, is constant in x_4 . For the x_0 -independent state of Section 13.7, $T_{04} = 0$ and the other 21 off-diagonal components are multiples of 15 three-gamma bilinears, which can all vanish (Section 15.5).

13 Homogeneous sector and equations of state

13.1 Ansatz and bilinears

The homogeneous sector has no dependence on $x_1, x_2, x_3, x_5, x_6, x_7$. Sections 13.1 to 13.6 treat the plane wave in ζ with real wave number K :

$$\Psi = e^{-3H\zeta} e^{iK\zeta} u(x_4) = s^{-1/2+iK/(6H)} u(x_4).$$

The bilinears are $S = \bar{\Psi}\Psi = s^{-1}u^\dagger C u$, $K_0 = iK s^{-1}u^\dagger C \gamma^0 u$ and $K_4 = \frac{1}{2}(\bar{\Psi}\gamma^4\partial_4\Psi - \partial_4\bar{\Psi}\gamma^4\Psi)$. The ansatz solves the Dirac equation exactly if and only if M_{eff} does not depend on ζ . That is exact for $U = 0$ ($M_{\text{eff}} = m$). For $\lambda \neq 0$ the condensate $S = u^\dagger C u / \sin z$ makes $M_{\text{eff}} = m + \lambda S$ depend on z , and the reduction $\gamma^4 \dot{u} = (M_{\text{eff}} - iK\gamma^0)u$ holds only pointwise, with M_{eff} treated as given (mean

field). The complex value $K = -3iH$ gives the state of Section 13.7, which does not depend on x_0 and is exact for every λ .

13.2 Energy density, pressures and kinetic/potential splits

Two splits of ρ are used. The L split takes half of the time-derivative part of the kinetic term: $\text{KE}_L = \frac{1}{2}K_4$ and $\text{PE}_L = \rho - \text{KE}_L$. The H split divides the Hamiltonian density (Section 17.3) into its gradient part KE_H and its mass-plus-potential part PE_H . The pressures are $p_{(\mu)} = T^\mu{}_\mu$ (no sum) for every transverse μ .

Off shell, for every $u(x_4)$:

$$\begin{aligned}\rho &= T_{44} = mS + U - K_0, & \mathcal{L}_s &= K_4 + K_0 - mS - U, \\ p_{(0)} &= T^0{}_0 = \mathcal{L}_s - K_0, & p_{(i)} &= T^i{}_i = \mathcal{L}_s \quad (i = 1, 2, 3, 5, 6, 7), \\ \text{KE}_L &= \frac{1}{2}K_4, & \text{PE}_L &= \rho - \text{KE}_L, \\ \text{KE}_H &= -\frac{1}{2} \sum_{\mu \neq 4} (\bar{\Psi} \gamma^\mu D_\mu \Psi - D_\mu \bar{\Psi} \gamma^\mu \Psi) = -K_0, & \text{PE}_H &= mS + U.\end{aligned}$$

On shell, $\gamma^4 \dot{u} = (M_{\text{eff}} - iK\gamma^0)u$ with $\dot{u} = du/dx_4$, so $K_4 + K_0 = M_{\text{eff}}S$ and

$$\begin{aligned}\mathcal{L}_s &= SU' - U, & \rho &= mS + U - K_0, \\ p_{(0)} &= SU' - U - K_0, & p_{(1,2,3,5,6,7)} &= SU' - U, \\ \text{KE}_L &= \frac{1}{2}(S(m + U') - K_0), & \text{PE}_L &= \frac{1}{2}(mS + 2U - SU') - \frac{1}{2}K_0, \\ \text{KE}_H &= -K_0, & \text{PE}_H &= mS + U.\end{aligned}$$

For $K = 0$ the zero mode gives $T_{44} = mS + U$.

13.3 Equation of state

- **$K = 0$, isotropic.** $\rho = mS + U$ and $p = SU' - U$ in all seven transverse directions, so the equation of state is $w = \frac{SU' - U}{mS + U} = \frac{\text{KE}_L - \text{PE}_L}{\text{KE}_L + \text{PE}_L}$.
- **Free massive** ($U = 0$, $K = 0$). $\text{KE}_L = \text{PE}_L = \frac{1}{2}mS$ and $p = 0$: $w = 0$, dust.
- **Default potential** ($U = \frac{\lambda}{2}S^2$, $K = 0$). $\rho = mS + \frac{\lambda}{2}S^2$, $p = \frac{\lambda}{2}S^2$, $w = \frac{\lambda S}{2m + \lambda S}$, $\text{KE}_L = \frac{1}{2}S(m + \lambda S)$ and $\text{PE}_L = \frac{1}{2}mS$. For $\rho > 0$ the state is phantom ($w < -1$) exactly when $\text{KE}_L < 0$, that is when $S(m + U') < 0$.
- **$K \neq 0$, anisotropic.** $w_{(0)} = p_{(0)}/\rho = (SU' - U - K_0)/(mS + U - K_0)$ and $w_{(i)} = (SU' - U)/(mS + U - K_0)$ for $i \neq 0, 4$. The pressure is isotropic only in the six directions 1, 2, 3, 5, 6, 7; the hidden direction carries the extra $-K_0$. With the mean transverse pressure $\bar{p} = \frac{1}{7} \sum_{\mu \neq 4} p_{(\mu)}$ and $U = 0$ this gives $\bar{w} = \bar{p}/\rho = (-K_0/7)/(mS - K_0)$.

The bilinear $S = u^\dagger C u / \sin z$ is not positive, because C has eigenvalues $+1$ and -1 (8 each). So $\rho = mS$ can be negative even for the free field; see the examples of Section 13.6.

13.4 Frozen in x_4 , independent of a_4 , and the $p(0)$ caveat

- No a_4 appears in ρ , $\mathcal{L}_s$, K_0 , K_4 , S or KE_H .
- For $U = 0$ and any K , $\partial_4 \rho = 0$ on shell. For $K = 0$ and any M_{eff} , $\partial_4 S = 0$, so ρ and all pressures are frozen.
- **Caveat for $p_{(0)}$.** For $K \neq 0$ (and $U = 0$) the energy density $\rho = mS - K_0$ and the six transverse pressures (zero on shell) are frozen, but $p_{(0)} = -K_0$ is not. Its rate is $\partial_4 p_{(0)} = -2iKm u^\dagger C \gamma^4 \gamma^0 u / \sin z$, the commutator $i[h, Q_{p_0}]$ with the single-particle operator h of Section 14.1. It oscillates at the frequency $2E$ with $E = \sqrt{m^2 + K^2}$, a Zitterbewegung between the $+E$ and $-E$ eigenspaces. $p_{(0)}$ is frozen only for $K = 0$ or on eigenstates of h . The Stage-2 specification's phrase "frozen in x_4 " is corrected accordingly.

13.5 Conservation

$\nabla_\mu T^\mu{}_\nu = 0$ holds on shell ($\lambda = 0$) for all eight ν , both in the homogeneous sector and for general $\Psi(x_0, x_4)$ (then on the solutions of the evolution equation of Section 10.5). As negative controls, the same divergence is nonzero off shell in both sectors.

13.6 Exact examples

Two exact sample points with $\lambda = 0$. P1 has $\sin z = 3/5$, $H = 1$, $m = 5/7$, $K = 3/2$, and P2 has $\sin z = 5/13$, $H = 3/4$, $m = -1/3$, $K = 2/5$. The amplitudes are $u_a = e_0 + e_4$ and $u_b = e_0 + ie_1 + e_4 + ie_{12}$, where e_n are the unit vectors of $\mathbb{C}^{16}$ (0-based). The transverse pressures $p_{(1,2,3,5,6,7)}$ vanish in all four cases. Since $\lambda = 0$ we have $\text{KE}_L = \text{PE}_L = \rho/2$ on shell, and $\bar{w} = \bar{p}/\rho$.

	S	ρ	$p_{(0)}$	$\text{KE}_L = \text{PE}_L$	KE_H	PE_H	$\partial_4 p_{(0)}$	$\bar{w}$
P1, u_a	-10/3	-50/21	0	-25/21	0	-50/21	0	0
P1, u_b	-10/3	-155/21	-5	-155/42	-5	-50/21	-50/7	3/31
P2, u_a	-26/5	26/15	0	13/15	0	26/15	0	0
P2, u_b	-26/5	-26/75	-52/25	-13/75	-52/25	26/15	104/75	6/7

For u_b at P1 this gives $S = -10/3$, $\rho = -155/21$ and $p_{(0)} = -5$. The nonzero $\partial_4 p_{(0)}$ of u_b is the caveat of Section 13.4.

13.7 The state that does not depend on x_0

For $K = -3iH$ the exponent $-\frac{1}{2} + iK/(6H)$ of the ansatz of Section 13.1 vanishes, and the state is

$$\Psi = u(x_4), \quad \gamma^4 \partial_4 u = (M_{\text{eff}} - 3H\gamma^0)u, \quad \partial_4 u = Au, \quad A = -\gamma^4(M_{\text{eff}} - 3H\gamma^0).$$

It depends on x_4 only. Its bilinears do not depend on z , and $S = u^\dagger C u$ is conserved, $\partial_4 S = u^\dagger (A^\dagger C + CA)u = 0$. Hence $M_{\text{eff}} = m + \lambda S$ is a constant, and unlike the real- K plane wave this state solves the Dirac equation exactly for every λ (check `P_source_x0IndependentStateSolvesDiracExactly`). The identity $h^2 = (M_{\text{eff}}^2 + K^2) \cdot 1$ of Section 14.1 at $K = -3iH$, with $h = iA$, gives $A^2 = -(M_{\text{eff}}^2 - 9H^2) \cdot 1$.

For $M_{\text{eff}}^2 > 9H^2$ the solutions oscillate with $\omega = \sqrt{M_{\text{eff}}^2 - 9H^2}$. A solution $u = e^{i\omega x_4} u_0$ with $Au_0 = i\omega u_0$ is stationary: every bilinear $u^\dagger X u$ is constant in x_4 .

On shell the diagonal components are (check P_source_x0IndependentDiagonalOnShell)

$$T^4_4 = -(mS + U) = -\rho, \quad T^\mu_\mu = SU' - U \quad (\text{all seven } \mu \neq 4),$$

so $\rho = mS + U$ and the pressure is isotropic, $p = SU' - U$, as for $K = 0$ in Section 13.3, but with a z -independent S . The component T_{04} vanishes, and so do $T_{ii'}$ and $T_{jj'}$ within the groups $\{1, 2, 3\}$ and $\{5, 6, 7\}$. Each of the other 21 off-diagonal components is a single three-gamma bilinear (check P_source_x0IndependentOffDiagonalAre15Bilinears), with $i = 1, 2, 3$ and $j = 5, 6, 7$:

(μ, ν)	Bilinear	Coefficient of $T_{\mu\nu}$
$(0, i)$	$\bar{\Psi} \gamma^0 \gamma^i \gamma^4 \Psi$	$-\frac{1}{4} H \cot z s^{1/6} e^{a_4} a'_4$
$(0, j)$	$\bar{\Psi} \gamma^0 \gamma^4 \gamma^j \Psi$	$\frac{1}{4} H \cot z s^{1/6} e^{-a_4} a'_4$
$(i, 4)$	$\bar{\Psi} \gamma^0 \gamma^i \gamma^4 \Psi$	$-\frac{7}{4} H s^{1/6} e^{a_4}$
$(4, j)$	$\bar{\Psi} \gamma^0 \gamma^4 \gamma^j \Psi$	$-\frac{7}{4} H s^{1/6} e^{-a_4}$
(i, j)	$\bar{\Psi} \gamma^i \gamma^4 \gamma^j \Psi$	$\frac{1}{2} H s^{1/3} a'_4$

So the off-diagonal part is governed by 15 bilinears: $V_a = \bar{\Psi} \gamma^0 \gamma^a \gamma^4 \Psi$ for $a = 1, 2, 3, 5, 6, 7$ and $W_{ij} = \bar{\Psi} \gamma^i \gamma^4 \gamma^j \Psi$. For a stationary state they are constant. In the canonical measure $\cos z dx_0$ of Section 17.2 this state is normalizable in x_0 , since $\int_0^{\pi/2} \cos z dz = 1$. The real- K plane wave is not: its $\cos z |\Psi|^2$ is proportional to $\cot z$, which is not integrable at $z \rightarrow 0$ (check P_source_x0IndependentNormalizableRealKNot).

14 Modes

14.1 Exact zeta reduction

For $\Psi = e^{-3H\zeta} e^{iK\zeta} u(x_4)$ the Dirac equation reduces exactly (for $\lambda = 0$, or with M_{eff} given) to

$$\begin{aligned} \gamma^4 \dot{u} &= (M_{\text{eff}} - iK\gamma^0)u \quad \Longleftrightarrow \quad i\dot{u} = hu, \quad h = -iM_{\text{eff}}\gamma^4 - K\gamma^4\gamma^0, \\ h &= h^\dagger, \quad h^2 = (M_{\text{eff}}^2 + K^2) \cdot 1, \quad E = \pm \sqrt{M_{\text{eff}}^2 + K^2} \quad (8 \text{ each}). \end{aligned}$$

The operator h is traceless. The Wolfram check evaluates the spectrum for $M_{\text{eff}} = 3$ and $K = 4$: the eigenvalues are $+5$ and -5 , eight each. The identity $h^2 = (M_{\text{eff}}^2 + K^2) \cdot 1$ is polynomial in K and therefore also holds at $K = -3iH$ (Section 13.7), where h is no longer Hermitian.

14.2 Momenta along 3-space and the extra times

For $\Psi = e^{-3H\zeta} e^{ik_1 x_1 + ik_5 x_5} \phi(\zeta, x_4)$, with momentum k_1 along x_1 and k_5 along x_5 , the equation becomes

$$\gamma^0 \partial_\zeta \phi + \gamma^4 \partial_4 \phi + i(k_1 e^{-H\zeta - a_4(t)} \gamma^1 + k_5 e^{-H\zeta + a_4(t)} \gamma^5) \phi = M_{\text{eff}} \phi.$$

The coefficients $k_1 e^{-H\zeta - a_4(t)}$ and $k_5 e^{-H\zeta + a_4(t)}$ are products of a nonconstant function of ζ and a nonconstant function of t . They multiply γ^1 and γ^5 , which anticommute with γ^0 and γ^4 . So the reduced equation keeps an explicit ζ dependence, and for $k_1 \neq 0$ or $k_5 \neq 0$ no product ansatz separates the ζ and t dependence.

14.3 Local WKB dispersion and instability onset

With frozen coefficients (a local, WKB-type statement) the single-particle operator $h_{\text{loc}} = -iM_{\text{eff}}\gamma^4 - \gamma^4(K\gamma^0 + k_{1,\text{eff}}\gamma^1 + k_{5,\text{eff}}\gamma^5)$, with $k_{1,\text{eff}} = k_1 e^{-H\zeta - a_4}$ and $k_{5,\text{eff}} = k_5 e^{-H\zeta + a_4}$, satisfies $h_{\text{loc}}^2 = E^2 \cdot 1$ with

$$E^2 = M_{\text{eff}}^2 + K^2 + (k_1 e^{-H\zeta - a_4})^2 - (k_5 e^{-H\zeta + a_4})^2.$$

h_{loc} is Hermitian if and only if $k_5 = 0$. Since $\gamma^4\gamma^5$ is anti-Hermitian, $h_{\text{loc}} - h_{\text{loc}}^\dagger = -2k_{5,\text{eff}}\gamma^4\gamma^5$, and the anti-Hermitian part of h_{loc} is the term $-k_{5,\text{eff}}\gamma^4\gamma^5$ itself. For $k_1 = 0$ and $k_5 \neq 0$, $E^2 < 0$ once

$$a_4(t) > H\zeta + \ln(\sqrt{M_{\text{eff}}^2 + K^2}/|k_5|),$$

and for $a_4 = t$ the onset time is $t_* = H\zeta + \ln(\sqrt{M_{\text{eff}}^2 + K^2}/|k_5|)$. The onset formula is verified at $M_{\text{eff}} = 3$, $K = 4$, $k_5 = 1/2$, where it gives $a_4 = H\zeta + \ln 10$. An unboundedly growing a_4 , for example $a_4 = t$ (3-space inflation, extra-time deflation), reaches the onset at every ζ . A bounded a_4 need not, although at fixed t the condition holds for sufficiently negative ζ (read off from the formula). This is the ultrahyperbolic ill-posedness of momenta along the extra times: as the extra times deflate, the physical momentum $k_5 e^{-H\zeta + a_4}$ grows.

15 Einstein tensor and the required source

15.1 Curvature

The Ricci scalar and the mixed Einstein tensor are

$$\begin{aligned} R &= 6H^2(a_4'^2 - 7), \\ G^0_0 &= -3H^2(a_4'^2 - 5), \\ G^i_i &= H^2(15 - 3a_4'^2 + a_4'') \quad (i = 1, 2, 3), \\ G^4_4 &= 3H^2(7 + a_4'^2), \\ G^j_j &= H^2(15 - 3a_4'^2 - a_4'') \quad (j = 5, 6, 7). \end{aligned}$$

All off-diagonal components of G^μ_ν and $R_{\mu\nu}$ vanish, and $R_{44} = -6H^2a_4'^2$. The Ricci scalar equals the notebook's cell-583 output, and the covariant $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R$ equals its cell-584 output entry by entry. The contracted Bianchi identity holds. Nothing here depends on x_0 .

15.2 Source required by 8D Einstein gravity

If $G^\mu{}_\nu = \kappa T^\mu{}_\nu$ with $\rho = -T^4{}_4$ and $p_{(\mu)} = T^\mu{}_\mu$, the required source is

$$\begin{aligned}\rho_{\text{req}} &= -3H^2(7 + a_4'^2)/\kappa, & p_{(0)} &= -3H^2(a_4'^2 - 5)/\kappa, \\ p_{(1,2,3)} &= H^2(15 - 3a_4'^2 + a_4'')/\kappa, & p_{(5,6,7)} &= H^2(15 - 3a_4'^2 - a_4'')/\kappa.\end{aligned}$$

In particular $\rho_{\text{req}} \leq -21H^2/\kappa < 0$: a negative energy density for every a_4 .

15.3 Energy conditions

Evaluated in the orthonormal frame for the observer $u = e_4$ (the evolution time), with the null vectors formed from e_4 or a time-like e_j and a space-like e_0 or e_i . Each quantity must be nonnegative for the condition to hold.

Condition	Quantity	Status
WEC	$\rho = -\frac{3H^2}{\kappa}(7 + a_4'^2)$	violated for every a_4
NEC, null vector $e_4 + e_0$	$\rho + p_{(0)} = -\frac{6H^2}{\kappa}(1 + a_4'^2)$	violated for every a_4
NEC, null vector $e_4 + e_i$	$\rho + p_{(i)} = -\frac{H^2}{\kappa}(6 + 6a_4'^2 - a_4'')$	violated unless $a_4'' \geq 6(1 + a_4'^2)$
NEC, null vector $e_j + e_0$	$p_{(0)} - p_{(j)} = \frac{H^2}{\kappa}a_4''$	holds if and only if $a_4'' \geq 0$
NEC, null vector $e_j + e_i$	$p_{(i)} - p_{(j)} = \frac{2H^2}{\kappa}a_4''$	holds if and only if $a_4'' \geq 0$
SEC (time-like convergence)	$R_{44} = -6H^2 a_4'^2$	violated whenever $a_4' \neq 0$
DEC	requires $\rho \geq 0$	violated

The 8-dimensional strong energy condition $5\rho + \sum_{\mu \neq 4} p_{(\mu)} = 6R_{44}/\kappa = -36H^2 a_4'^2/\kappa$ fails whenever $a_4' \neq 0$. In signature (4,4) there are four time-like directions, so these are the conditions for the chosen observer and null vectors, not a complete classification.

15.4 Linear a4 and the cell-150 alternatives

For $a_4'' = 0$, that is $a_4 = ct$ up to a constant, all seven transverse pressures coincide:

$$R = 6H^2(c^2 - 7), \quad \rho_{\text{req}} = -3H^2(7 + c^2)/\kappa, \quad p_{(\mu)} = H^2(15 - 3c^2)/\kappa \quad (\mu \neq 4), \quad w = \frac{c^2 - 5}{c^2 + 7}.$$

The equation of state parameter $w = (c^2 - 5)/(c^2 + 7)$ increases from $-5/7$ at $c = 0$ towards 1 and is negative for $c^2 < 5$ (read off from the formula). With $\rho < 0$ and $p > 0$ the usual reading of w (for example $w < -1/3$ as acceleration) does not carry over. For $a_4 = t$:

$$\begin{aligned}R &= -36H^2, & G^\mu{}_\nu &= \text{diag}(12, 12, 12, 12, 24, 12, 12, 12) H^2, \\ \rho_{\text{req}} &= -24H^2/\kappa, & p_{(\mu)} &= 12H^2/\kappa \quad (\text{all seven transverse directions}), & w &= -\frac{1}{2}.\end{aligned}$$

and the notebook's q becomes $q = Q_1 \sinh t e^{-t} = \frac{1}{2}Q_1(1 - e^{-2t})$. Cell 150 of the notebook lists, without derivation, the slopes $a'_4 = \frac{2}{3}(M - 1)$ and $a'_4 = \frac{2}{3}(M + 1)$ with $a''_4 = 0$. Writing both as $a'_4 = \frac{2}{3}(M \mp 1)$:

$$\begin{aligned} R &= \frac{2}{3}H^2(4M^2 \mp 8M - 59), & \kappa\rho_{\text{req}} &= -\frac{1}{3}H^2(4M^2 \mp 8M + 67), \\ \kappa p_{(\mu)} &= -\frac{1}{3}H^2(4M^2 \mp 8M - 41) \quad (\mu \neq 4), & w &= \frac{4M^2 \mp 8M - 41}{4M^2 \mp 8M + 67}. \end{aligned}$$

$\rho_{\text{req}} < 0$ for every real M .

15.5 Which dirac16complex states can supply the source

The bilinears are read as c-numbers, as in Sections 13 and 14. There are three results.

Proposition 1 ($a''_4 \neq 0$). No dirac16complex state that depends on x_0 and x_4 only satisfies $G^\mu{}_\nu = \kappa T^\mu{}_\nu$ on an open set where $a''_4 \neq 0$.

Proof. For every such state, off shell and for every U , $T^i{}_i = T^j{}_j = \mathcal{L}_s$ for $i = 1, 2, 3$ and $j = 5, 6, 7$. The reason is that $\partial_i \Psi = \partial_j \Psi = 0$ and $A_{ii} = A_{jj} = 0$ (Section 12.3; check P_source_transversePressuresEqualForEveryX0X4State). But $G^i{}_i - G^j{}_j = 2H^2 a''_4$ (check P_source_einsteinTransverseDifferenceIs2H2a4pp), so $\kappa(T^i{}_i - T^j{}_j) = 0 \neq 2H^2 a''_4$. $\square$

Proposition 2 (real K). No plane wave $\Psi = s^{-1/2+iK/(6H)}u(x_4)$ with real K satisfies $G^\mu{}_\nu = \kappa T^\mu{}_\nu$ on an open range of z , for any a_4 .

Proof. For real K every bilinear is a function of x_4 divided by $\sin z$. Hence every diagonal component $T^\mu{}_\mu$, in particular $T^4{}_4 = -\rho$, has the form $c_1(t)/\sin z + c_2(t)/\sin^2 z$, with $c_2 \propto \lambda$ from $U = \frac{\lambda}{2}S^2$ (check P_source_realKDiagonalIsC1OverSPplusC2OverS2). The off-diagonal mixed components do not have this form; for example $T^0{}_4 = g^{00}T_{04}$ is proportional to $1/\cos z$. For the energy density

$$\rho = \frac{m u^\dagger C u - i K u^\dagger C \gamma^0 u}{\sin z} + \frac{\lambda (u^\dagger C u)^2}{2 \sin^2 z}.$$

The required source is independent of z and nonzero: $G^4{}_4 = 3H^2(7 + a_4'^2) > 0$. The functions 1 , $1/\sin z$ and $1/\sin^2 z$ are linearly independent on $(0, \pi/2)$; their Wronskian is

$$W(1, \frac{1}{\sin z}, \frac{1}{\sin^2 z}) = -2 \cot^3 z \csc^3 z \neq 0.$$

Comparing the coefficients of 1 in $G^4{}_4 = \kappa T^4{}_4$ gives $3H^2(7 + a_4'^2) = 0$, which is impossible (check P_source_realKPlaneWaveCannotSource). $\square$

The Python checker confirms this for $\lambda = 0$ from the other side: $\rho \sin z$ is independent of x_0 , so $\rho = \rho_{\text{req}}$ for all z would force $\rho = 0 \neq \rho_{\text{req}}$. For complex K the modulus of the prefactor is $|s^{-1/2+iK/(6H)}|^2 = s^{-1-\text{Im}K/(3H)}$. For $\lambda = 0$ the argument therefore excludes every K with $\text{Im} K \neq -3H$ as well. On the line $\text{Im} K = -3H$ the bilinears do not depend on z . Its member $K = -3iH$ is the state of Section 13.7; the members with $\text{Re} K \neq 0$ are not analysed here. These two remarks are derived here, not separate checks.

Proposition 3 (the state that does not depend on x_0). Let $a_4 = ct$ up to a constant, so $a_4'' = 0$. The state $\Psi = u(x_4)$ of Section 13.7 satisfies $G^\mu{}_\nu = \kappa T^\mu{}_\nu$ in all 64 components for all x_4 if and only if

1. the six bilinears V_a vanish and, if $c \neq 0$, the nine bilinears W_{ij} vanish as well;
2. $mS = -36H^2/\kappa$;
3. $\lambda S^2 = 2H^2(15 - 3c^2)/\kappa$.

Then $M_{\text{eff}} = m + \lambda S = -6H^2(1 + c^2)/(\kappa S)$. For $\kappa > 0$ the solutions oscillate, $M_{\text{eff}}^2 > 9H^2$, if and only if $\kappa|S| < 2H(1 + c^2)$ (derived here from the formula for M_{eff}). For $a_4'' \neq 0$ the state is excluded by Proposition 1.

Proof. The off-diagonal components are the bilinears of Section 13.7 times coefficients; the coefficients of the V_a never vanish, and those of the W_{ij} vanish only for $c = 0$. With $U = \frac{\lambda}{2}S^2$ the diagonal components are $T^4{}_4 = -(mS + \frac{\lambda}{2}S^2)$ and $T^\mu{}_\mu = \frac{\lambda}{2}S^2$ for $\mu \neq 4$. The two equations for $G^4{}_4$ and $G^0{}_0$ have the unique solution 2 and 3, and with it the other six diagonal equations hold if and only if $a_4'' = 0$ (check P_source_x0IndependentSourceConditions). $\square$

The energy density of such a source is $\rho = mS + \frac{\lambda}{2}S^2 = -3H^2(7 + c^2)/\kappa < 0$. A negative ρ is possible because C is indefinite, so S can have either sign. Condition 2 fixes S for a given m , condition 3 then fixes λ , and condition 1 restricts the amplitude u .

Existence. For $M_{\text{eff}} = \pm 5H$ the $i\omega$ eigenspace of A meets the singlets of the diagonal $\text{Spin}(3)$ generated by $S^{23} - S^{67}$, $S^{31} - S^{75}$ and $S^{12} - S^{56}$ (simultaneous rotations of x_1, x_2, x_3 and x_5, x_6, x_7) in a 2-dimensional space. On it 12 of the 15 bilinears vanish identically and the other three coincide, $W_{15} = W_{26} = W_{37}$, so that a single real condition remains (check P_source_x0IndependentConstruction). Two exact solutions with $H = \kappa = 1$, both of the form $\Psi = e^{4ix_4}u_0$ with $Au_0 = 4iu_0$:

Case A. $a_4 = \sqrt{5}t$, $\lambda = 0$, $m = M_{\text{eff}} = 5$ and $S = -36/5$, with

$$u_0 = \sqrt{9/40} (3, -i, 0, 0, 3, i, 0, 0, 1, -3i, 0, 0, 1, 3i, 0, 0).$$

Here $\rho = -36$ and $p = 0$: negative-energy dust, $w = 0$, in agreement with $w = (c^2 - 5)/(c^2 + 7)$ of Section 15.4.

Case B, the notebook's $a_4 = t$: $m = -15$, $\lambda = 25/6$, $M_{\text{eff}} = -5$ and $S = 12/5$, with

$$u_0 = \sqrt{3/40} (1, 3i, 0, 0, 1, -3i, 0, 0, -3, -i, 0, 0, -3, i, 0, 0).$$

Here $\rho = -24$ and $p = 12$ in all seven transverse directions, $w = -\frac{1}{2}$, which is the required source of Section 15.4.

In both cases Ψ solves the Dirac equation exactly, all 15 bilinears vanish, and $G^\mu{}_\nu - \kappa T^\mu{}_\nu = 0$ in all 64 components. As a negative control the same state fails for the slope $c + 1$ (check P_source_x0IndependentExactExamples; the Python checker verifies the same examples independently).

16 The gamma-8 map: a structural plus-minus M pairing

The chirality matrix $\gamma^8 = \text{diag}(-I_8, I_8)$ anticommutes with every γ^μ , commutes with every Ω_μ and leaves $S = \bar{\Psi}\Psi$ invariant, while the kinetic bilinear changes sign. Hence, in this field,

$$\begin{aligned}\mathcal{L}_{m,\lambda}[\gamma^8\Psi] &= -\mathcal{L}_{-m,-\lambda}[\Psi], \\ \gamma^\mu D_\mu\Psi &= (-m - \lambda S)\Psi \iff \gamma^\mu D_\mu(\gamma^8\Psi) = (m + \lambda S)\gamma^8\Psi.\end{aligned}$$

The map flips the sign of the eight upper components $\Psi_0, \dots, \Psi_7$ and keeps $\Psi_8, \dots, \Psi_{15}$. For $U = 0$ it relates the theory of mass $+m$ to the theory of mass $-m$, that is $+M$ to $-M$ in the notebook's normalization. With $U = \frac{\lambda}{2}S^2$ the self-coupling flips too, so $\mathcal{L}_m \rightarrow -\mathcal{L}_{-m}$ holds exactly only for $U = 0$. Stage 1 showed the same relation for the energy-momentum tensor: $T_{\mu\nu}^{(m,\lambda)}[\gamma^8\Psi] = -T_{\mu\nu}^{(-m,-\lambda)}[\Psi]$. This pairing of solutions echoes the notebook's hypothesis of universes created in pairs with masses $\pm M$. It is **structural only**: a field redefinition that maps one theory onto another. Nothing about a creation process at $x_4 = 0$, about a pair of universes, or about a physical preference between the two signs follows from it.

17 Canonical quantization in this field

17.1 Momentum and anticommutators

Adding the total derivative $\frac{1}{2}\partial_4(\sqrt{|g|}\bar{\Psi}\gamma^4\Psi)$ to $\mathcal{L}$ cancels the term with $\partial_4\bar{\Psi}$; no other term appears, because $\partial_4(\sqrt{|g|}\gamma^4) = 0$ here. The momentum conjugate to Ψ is then

$$\Pi = \frac{\partial\mathcal{L}}{\partial(\partial_4\Psi)} = \sqrt{|g|}\Psi^\dagger C\gamma^4 = \cos z \Psi^\dagger C\gamma^4.$$

The constraints $\Pi - \cos z \Psi^\dagger C\gamma^4 \approx 0$ are second class. The Dirac brackets give the equal- x_4 anticommutators

$$\{\Psi_a(x), \Psi_b^\dagger(y)\}_{x^4=y^4} = i[(C\gamma^4)^{-1}]_{ab} \frac{\delta^7(x-y)}{\cos z} = B_{ab} \frac{\delta^7(x-y)}{\cos z}, \quad B = -iC\gamma^4,$$

and all other anticommutators vanish. The general curved-space form $i[\gamma^4 C]_{ab} \delta^7/(g^{44}\sqrt{|g|})$ of Stage 1 reduces to this because $g^{44} = -1$ and $\gamma^{x_4} = \gamma^4$ ($h_4 = 1$).

17.2 The Krein structure

$$B = B^\dagger, \quad B^2 = 1, \quad \text{spec } B = \{+1^{(8)}, -1^{(8)}\}, \quad [C, B] = 0, \quad BC = -i\gamma^4.$$

The form $(\Psi, \Phi)_B = \int \cos z \Psi^\dagger B \Phi d^7x$ defined by the anticommutator is therefore indefinite, with signature (8,8) per mode. The canonical state space is a Krein space with fundamental symmetry $J = B$, and the positive Hilbert product is $\int \cos z \Psi^\dagger \Phi d^7x$. The conserved current $J^\mu = -i\Psi\gamma^\mu\Psi$ (Stage 1) has $J^4 = \Psi^\dagger B \Psi$ here (from $C\gamma^4 = iB$), so the charge $\int \cos z \Psi^\dagger B \Psi d^7x$ is indefinite. Stage

1 showed that this indefiniteness is intrinsic to (4,4): every $\text{Spin}(4,4)$ -invariant Hermitian form is block diagonal in chirality, while γ^4 is off-diagonal. Stage 1 also classified the frame generators S^{ab} (these are field-independent facts about B):

- 13 of the 28 commute with B : the 9 with a, b both in $\{0, 1, 2, 3\}$ or both in $\{5, 6, 7\}$, plus the 4 boosts S^{a4} with $a \in \{0, 1, 2, 3\}$.
- The 9 compact ones are anti-Hermitian and generate the unitarily implemented $\text{Spin}(4) \times \text{Spin}(3)$. The 4 boosts commute with B but are Hermitian, not unitary.
- The 21 generators with $a, b \neq 4$ are Krein-unitary ($\text{Spin}(4,3)$); the 7 generators S^{4b} fail the Krein condition.

In this field the rotations of (x_1, x_2, x_3) and of (x_5, x_6, x_7) are isometries of the metric. They are generated by 6 of those 9 compact generators and are therefore unitarily implemented, together with the translations in these six directions (an observation, not a separate check).

17.3 Hamiltonian density and Heisenberg equations

$$\mathcal{H} = \cos z \left[-\frac{1}{2} \sum_{\mu \neq 4} h_\mu^{-1} (\bar{\Psi} \gamma^\mu \partial_\mu \Psi - \partial_\mu \bar{\Psi} \gamma^\mu \Psi) + m \bar{\Psi} \Psi + \frac{\lambda}{2} (\bar{\Psi} \Psi)^2 \right], \quad h = (\cot z, s^{1/6} e^{a_4} (\times 3), 1, s^{1/6} e^{-a_4} (\times 3)),$$

$\mathcal{H}$ contains no x_4 derivatives, and up to a total derivative $\mathcal{H} = \sqrt{|g|} T_{44}$. The Heisenberg equation

$$i \partial_4 \Psi = B \frac{1}{\cos z} \frac{\delta H}{\delta \Psi^\dagger} \implies \partial_4 \Psi = -\gamma^4 \left[(m + \lambda S) \Psi - \sum_{\mu \neq 4} \gamma^\mu D_\mu \Psi \right], \quad H = \int \mathcal{H} d^7 x,$$

reproduces the Euler-Lagrange evolution of Section 10.4.

17.4 The good sector

In the sector without dependence on x_5, x_6, x_7 (no momentum k_5 along the extra times),

$$i \partial_4 \Psi = h \Psi, \quad h = -im \gamma^4 + 3iH \gamma^4 \gamma^0 + i \tan z \gamma^4 \gamma^0 \partial_0 + i s^{-1/6} e^{-a_4} \gamma^4 \gamma^i \partial_i,$$

and h is Hermitian with respect to $\int \cos z \Psi^\dagger \Phi d^7 x$. The term $3iH \gamma^4 \gamma^0$ is exactly what the weight $\cos z$ requires; in the ζ form the ζ operator is symmetric for the measure $e^{6H\zeta}$. The matrices $\gamma^4 \gamma^i$ ($i = 1, 2, 3$) are Hermitian, while the extra-time terms $i s^{-1/6} e^{a_4} \gamma^4 \gamma^j \partial_j$ are anti-Hermitian ($\gamma^4 \gamma^j$ is anti-Hermitian). So h is Hermitian if and only if there is no dependence on x_5, x_6, x_7 , in agreement with Section 14.3. In the ζ reduction h has the spectrum $\pm \sqrt{M_{\text{eff}}^2 + K^2}$, eight each (Section 14.1). For flat-space modes of the same form Stage 1 found that the B form has signature (4,4) on each of the two energy eigenspaces, so the Krein structure of Section 17.2 persists in the good sector.

18 Verification records

18.1 WolframScript

WolframScript 1.14 with Mathematica 15.0.1 runs the Wolfram implementation, a driver script and a module (file names in Section 18.4), in about 39 s. It writes the report wolfram-primordial-report.json and the component file primordial-components.json. The tables and displays of Sections 5, 6, 7.2, 10, 11.1, 11.2, 12.2, 13.7 (the coefficients of the off-diagonal components), 15.1 and 15.3 take their TeX from the component file, and tests/test_d16c_primordial_publication.py compares them with it. The values of Sections 8.3 and 13.6 are those recorded in python-primordial-report.json. The other displays are written by hand from the verified identities. All 126 checks are true:

Group	Checks	Group	Checks
fixture	4	notebookCompare	16
metric	6	EMT	18
zeta	4	modes	7
christoffel	3	einstein	10
spinconn	6	source	11
Omega	8	quant	11
gammaConst	7	a4linear	3
EL	7	internal	2
blocks	3		

Exactness method: every scalar is mapped to the ring $\mathbb{Q}(\text{parameters})[s^{1/6}, \cos z, e^{a_4}, a'_4, a''_4, \dots]$ modulo $\cos^2 z + s^2 - 1$. This is a complete zero test because a_4 is arbitrary, so e^{a_4} and its derivatives are independent. In addition there are exact point checks at $\sin z = 3/5$ and $\sin z = 5/13$ with rational jets of a_4 ($e^{1/6}$ treated as transcendental, coefficients RootReduce'd). Stored notebook outputs, among them those of cells 501, 583, 584, 1060, 1079, 1089, 1096, 1111 and 1137, are parsed from the committed notebook. The 15 source cells 106, 111, 386, 431, 475, 499, 501, 1058, 1061, 1066, 1075, 1078, 1096, 1111 and 1137 are checked to contain the code that is rebuilt. The check names are:

```

fixture: gammasMatchCommittedFixture CAndChiralityMatch cliffordAndC
        gammaRowsSignedPermutations
metric: vielbeinProduct signature44 detG_equals_plus_cos2z sqrtAbsDetG_cosz
        notebookCell11060Det pointCheck
zeta: gZetaZetaIsOne warpFactor range warpedMetric
christoffel: closedForms512 count pointCheck
spinconn: vielbeinPostulate512 antisymmetry closedForms count24
          notebookCell1501OmegaMuIJEqualsMixedOmega pointCheck
Omega: closedForms zeroForXOX4 blockDiagonalChirality gammaSlash3Hgamma0
        OmegaGammaAndAnticommutator slashA4Independent contractDiagonalFormula pointCheck
gammaConst: DmuGammaNuZero64 notebookContractionFails
            notebookContractionClosedForms divergenceIdentity
            divergenceIdentityFailsNotebookContraction divergenceLhsValue pointCheck
EL: componentsMatchOperator tenTermsPerEquation pointCheck evolutionFormEquivalent
    fromLagrangianPsiDagger fromLagrangianPsi zetaFormChainRule
blocks: fourBlocksOfFour closedUnderCoupling matchNotebookSets

```

```

notebookCompare: sourceCellsAsRebuilt spinCoefficientsCell499EqualsOmegaMuIJ storedEla16
commutingQ1DropsOutCliffordGammas reconstructionReproducesStoredEla
literalRebuildResidualIsExactlyQTerms qTermFromNonCliffordExtraTimeGammas
qVectorIs2HqAtCell1079 reconstructedGamma5NotClifford
reconstructedGammaForm eLaztCell1096 couplingSetsCell1089 relabelCell1111
cell11137Reproduced16of16 correctVsStoredDifferOnlyByQ qOnlyInYZ0to7
EMT: anticommutatorsAreThreeGammaProducts OmegaPartIsMinusQuarterPsi symmetric
homogeneous_rhoOffShell homogeneous_LsOffShell homogeneous_pressuresOffShell
homogeneous_KEH_equals_minusK0 T44equals_mSplusU_zeroMode
homogeneous_onShell_K4plusK0_equals_MeffS reductionSolvesDiracEquation
homogeneous_frozenInX4 homogeneous_a4Independent homogeneous_conservationOnShell
homogeneous_conservationFailsOffShell x0x4EvolutionEquationIsEL
conservationOnShellX0X4Sector conservationFailsOffShellX0X4Sector
homogeneous_offDiagonalTransverseFromA
modes: dispersion spectrumPlusMinusE8each exactReduction kqNonzeroNotSeparable
localWKBDispersion localHermitianIffQZero instabilityOnset
einstein: ricciScalar GmixedClosedForms offDiagonalZero pointCheck notebookCell584
notebookCell583 R44 requiredSource rhoRequiredNegative energyConditionForms
source: transversePressuresEqualForEveryX0X4State
einsteinTransverseDifferenceIs2H2a4pp realKDiagonalIsC10overSPlusC20overS2
realKPlaneWaveCannotSource x0IndependentStateSolvesDiracExactly
x0IndependentDiagonalOnShell x0IndependentOffDiagonalAre15Bilinears
x0IndependentSourceConditions x0IndependentConstruction
x0IndependentExactExamples x0IndependentNormalizableRealKNot
quant: Bproperties anticommutatorMatrix momentum hamiltonianHasNoTimeDerivatives
hamiltonianDensityClosedForm heisenbergReproducesEL goodSectorHermitian
extraTimeMomentaNonHermitian singleParticleOperatorMatchesEvolution
gamma8MapsMtoMinusM gamma8LagrangianSign
a4linear: values isotropicWhenA4ppZero cell150AlternativesRhoNegative
internal: zeroTestSanity noException

```

18.2 Python

The independent Python checker (sympy and the standard library only; file name in Section 18.4) writes the report python-primordial-report.json. Nothing produced by Wolfram is used as truth. The gamma matrices are rebuilt from the split-octonion recipe and only compared with the committed fixture, and the geometry is recomputed from the metric. The 16 by 16 identities are proved in the Clifford algebra $Cl(4,4)$ in the blade basis; the notebook matrices are verified to be a faithful representation (all 65536 blade products). Every matrix identity is re-checked with the actual matrices at the two exact sample points P1 and P2, with $s^{1/6}$ kept as an exact radical and e^{a_4} as a transcendental. The Python checker has no notebook reader: for the notebook it compares only with transcriptions of the stored outputs of cells 583, 584 and 1137. All 16 checks are true:

```

P_algebraSetup P_metric P_zeta P_christoffel P_spinconn P_Omega P_gammaConst P_EL
P_blocks P_EMT P_modes P_einstein P_source P_quant P_a4linear P_EL_agreesWithWolfram

```

18.3 Agreement between the implementations

`P_EL_agreesWithWolfram` parses the Wolfram component file and compares every coefficient of the 16 component equations and of the 16 evolution equations with the Python derivation. Both

printers of the file (py and wl) are compared, exactly, at the two sample points, with $\lambda = 1/2$ and $S = 7/3$ inserted so that the λS term is compared too. That is 320 coefficients for the equations (16 rows, 10 terms, 2 points) and 288 for the evolution form (16 rows, 9 terms, 2 points), for each printer, with 0 mismatches. The Python report records wolframAgreement = compared and the sha256 of the component file it compared. Three Wolfram runs produced a byte-identical component file.

Where the Stage-2 specification disagreed with the measured results, the checks test the measured truth and record the correction: $\det g = +\cos^2 z$ (Section 4.3); $p_{(0)}$ frozen only for $K = 0$ or on eigenstates (Section 13.4); isotropy only in the six directions 1, 2, 3, 5, 6, 7 (Section 13.3); the ζ plane wave exact only for $\lambda = 0$ (Section 13.1); the off-diagonal $T_{0\mu}$, $T_{4\mu}$ and T_{ij} nonzero for generic plane waves (Section 12.3). The binding contract of the project states that a homogeneous condensate cannot source this field because $\bar{\Psi}\Psi \propto 1/\sin z$. That holds for the real- K plane waves only. The state that does not depend on x_0 has a constant $\bar{\Psi}\Psi$ and is an exact source for $a_4'' = 0$ (Section 15.5); the checks of the group source test the corrected statement.

18.4 Files and hashes

sha256 of the inputs and producers, as recorded in the two reports for this edition:

```
Pair_Creation_of_Universes_WaveFunctionOfUniverse-4+4-Einstein-Lovelock-Nash.nb
5ee5cb2a95146136ee65636a4aef174f303b6c41da130d2c57e4684f9c8ff69f
artifacts/dirac16complex/arbitrary-field/algebra-fixture.json
8b4f15462ca4d61ef6ec0e72f04c8a77d23ce8bcabc9b6ce19f921c90e02653b
wolfram/Dirac16ComplexPrimordial.wl
0b95682601ace6343a89ad8eaf0f4896d071015041f15cccbeb3beac193d81a1
scripts/verify_dirac16complex_primordial.wls
f49606ba044ace6eb0ebc5e7661a7823f826416f234fce756acd9a2ccab958f9
scripts/check_dirac16complex_primordial.py
4e86497e8fee662fc43117a5a55341c1450e6644dbe7629e14c6b96a9798793f
artifacts/dirac16complex/primordial-field/primordial-components.json
a5d8caa8cb2260824c29dcdb1ed0e62ec22079c297682939a277a6b98dfa3e52
```

19 Reproduction

Run every command from the repository root. The Wolfram report path must be passed positionally, not after a double-hyphen argument: WolframScript 1.14 drops a double hyphen and every argument after it (see the header of the gate scripts).

19.1 PowerShell

The Stage-2 gate runs the four steps below and audits both reports:

```
pwsh -NoProfile -File scripts/verify_stage2_primordial_field.ps1
```

Its last line is `stage2_primordial_field_verification=OK`.

The individual steps:

```
$r = "artifacts/dirac16complex/primordial-field/wolfram-primordial-report.json"
wolframscript -file scripts/verify_dirac16complex_primordial.wls $r
python scripts/check_dirac16complex_primordial.py
python -m unittest discover -s tests -p "test_d16c_primordial*.py" -v
python scripts/build_provenance_pdf.py provenance/DIRAC16COMPLEX_PRIMORDIAL_FIELD.md
```

19.2 Git Bash

```
bash scripts/verify_stage2_primordial_field.sh
```

The individual steps:

```
wolframscript -file scripts/verify_dirac16complex_primordial.wls \
    artifacts/dirac16complex/primordial-field/wolfram-primordial-report.json
python scripts/check_dirac16complex_primordial.py
python -m unittest discover -s tests -p "test_d16c_primordial*.py" -v
python scripts/build_provenance_pdf.py provenance/DIRAC16COMPLEX_PRIMORDIAL_FIELD.md
```

The Wolfram step prints `check_count=126` and `failed_check_count=0`. The Python step prints `check_count=16`, `failed_check_count=0` and `measurement_wolframAgreement=compared`. The PDF step rebuilds this document twice (two builder runs, three pdflatex passes each), requires warning-free logs and byte-identical PDFs, compares them with the registered edition `dirac16complex-primordial-field` in `provenance/pdf-specifications.json`, and prints `provenance_pdf=OK`. After an edit of this document the edition is registered once more with

```
python scripts/build_provenance_pdf.py --register \
    provenance/DIRAC16COMPLEX_PRIMORDIAL_FIELD.md
```

(in PowerShell the two lines are written as one line)

and the sha256 pins in `tests/test_d16c_primordial_publication.py` are updated.

20 Limitations

1. Only 8-dimensional Einstein gravity is used for the source. The Einstein-Lovelock terms of order 2 and 3, which the notebook names but does not compute, are not computed here either.
2. The source analysis of Section 15.5 covers states that depend on x_0 and x_4 only, with the bilinears read as c-numbers. States that depend on the transverse coordinates are not analysed; for them the off-diagonal components of Section 12.3 have to vanish as well. The plane waves with $\text{Im } K = -3H$ and $\text{Re } K \neq 0$ are not analysed. The exact source found for linear a_4 has negative energy density, and neither its stability nor its quantum (Krein-space) status is examined.

3. For $\lambda \neq 0$ the ζ plane wave is not an exact solution. The homogeneous-sector formulas then hold pointwise, with M_{eff} treated as given.
4. The bilinears are evaluated on c-number amplitudes. The operator expectation values in a Krein-space state, with normal ordering, are not computed.
5. Modes with momentum along the extra times are non-Hermitian, and for an unboundedly growing a_4 they become unstable (Section 14.3). Only the good sector has a standard positive single-particle structure.
6. The dynamics of a_4 is not determined: a_4 is a free function, and the notebook's slopes of cell 150 are tabulated, not derived.
7. The attribution of the q term to cell 1058 is a reconstruction. The reconstructed useT16 reproduces the stored outputs exactly, but the notebook does not store useT16, and a literal re-execution in Mathematica 15.0.1 behaves differently (Section 11.3).
8. The x_0 chart covers only $\zeta < 0$. Continuation beyond $\zeta = 0$ is not discussed.