UM-3 solution

The approximate solution of the coupled field equations of UM-3: the background and the power series of the metric, the Higgs field in the ad hoc vacuum, the spectra along the hidden direction, the self-consistent Kohn-Sham states of dirac16complex, the residuals of every field equation, the neutrino flavours, the thermal history, and the CMB, the CNB and the BAO

Underlying theory: Patrick L. Nash Solved and verified by Claude (Anthropic)

Solution of UM-3, 2026-10-24 Repository once-ere/Dirac_claude, folder universe_model_3

Contents

1 The fold: definition and examples **2**

2 How to read this document **11**
2.1 What this document contains . . . 11
2.2 The status labels . . . 12
2.3 Every equation is followed by its symbols . . . 12
2.4 Coordinates, signature and units . . . 12
2.5 The solution at a glance . . . 12

3 The Lovelock weights and the cosmological constant **14**

4 The background **20**

5 The power series of the metric **30**
5.1 The fold as a seven-dimensional Einstein brane . . . 35
5.2 The volume of the extra times . . . 36

6 The sources of the fold **39**
6.1 The gravitational waves of the fold . . . 39
6.2 The gravity induced by the fields of the fold . . . 41
6.3 The fold fluid . . . 42

7 The Higgs field in the ad hoc vacuum **44**

8 The spectra along the hidden direction **48**
8.1 The bulk fields and why the Standard Model lives on the fold . . . 48
8.2 dirac16complex: the continuum and the bound states . . . 49

9 The Kohn-Sham states of dirac16complex **55**
9.1 The equations of the gas of bound particles . . . 59
9.2 The method . . . 65
9.3 The solution in three regimes . . . 65
9.4 A weakly bound gas . . . 67
9.5 The Krein partner and the stability of the gas . . . 67

10 The residuals of every field equation **69**

11 The three neutrino flavours **78**
11.1 The Sun . . . 82

11.2 The envelope of a supernova . . . 83
11.3 The fast instability near the neutrinosphere . . . 84
11.4 The flavour equipartition and the heating behind the shock . . . 86
11.5 The newer global fit, the Dirac phase and the lightest mass . . . 87

12 The thermal history **89**
12.1 The relic density of the particles χ . . . 90
12.2 The portals of χ to the Standard Model . . . 92
12.3 The right-handed neutrinos . . . 95

13 The CMB, the CNB and the BAO **96**
13.1 The background and the BAO ruler . . . 102
13.2 The CMB . . . 103
13.3 The BAO . . . 105
13.4 The CNB . . . 107

14 What the solution does not settle **107**

15 What was verified **109**

16 Further symbols **110**

1 The fold: definition and examples

Definition. The *fold* is the seven-dimensional surface of the author's eight-dimensional spacetime on which the cosine of the angle $z = 6Hx_0$ of the hidden direction vanishes:

$$\mathcal{F} = \big\{X = (x_0, x_1, \ldots, x_7) :\ \cos(6Hx_0) = 0\big\}, \qquad \text{in one period } 0 < z < \pi \text{ of the metric: } z = \frac{\pi}{2}, \quad x_0 = \frac{\pi}{12H} \tag{FD-1}$$

Symbols in (FD-1).

- $\mathcal{F}$: The fold: the seven-dimensional surface $\cos(6Hx_0) = 0$ of the eight-dimensional spacetime (FD-1), in one period $0 < z < \pi$ the surface $z = \pi/2$, $x_0 = \pi/(12H)$, where $\sigma = 0$; $|_{\mathcal{F}}$ means restricted to the fold. Status: DERIVED.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$, the fold is $z = \pi/2$. Unit: dimensionless. Status: AUTHOR.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -7H^2$: with the corrected weights the value for which the static block is a vacuum (its Lovelock tensors add to $7H^2$ in every direction). The author's original value was $\Lambda = -10H^2$ (he writes the pure number $\Lambda_{\mathrm{N}} = 10$ in $0 = -\Lambda_{\mathrm{N}} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$). Unit: GeV^2. Value: $-7H^2$; the author's original $-10H^2$. Status: AUTHOR (the term), DERIVED (the value).
- $w_{(p)}$: The Lovelock weights, $w_{(1)} = 1$, $w_{(2)} = 1/20$, $w_{(3)} = 1/360$: the only weights for which both coefficients of a transverse traceless gravitational wave in the static bulk vanish, $1 - 40w_{(2)} + 360w_{(3)}$ (of its second time derivative) and $1 - 24w_{(2)} + 72w_{(3)}$ (of its second derivatives along ordinary space and along σ). Then the

four-dimensional graviton of the fold travels at the speed of light at linear order, and no graviton of the static bulk travels faster, because the static bulk has no linear graviton at all (the Born-Infeld point of Lovelock gravity in eight dimensions). The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) made bulk gravitons $\sqrt{5}$ times faster than light; they were corrected at the author's order of 2026-10-25. Unit: dimensionless. Value: $(1, 1/20, 1/360)$; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), ASSUMED (the criterion of the correction), DERIVED (the unique values that meet it).

- Λ_{N}: The author's pure number in the vacuum equations of his notebook, $\Lambda = -\Lambda_{\mathrm{N}} H^2$: his original value 10, corrected to 7 on 2026-10-25. Unit: dimensionless. Value: 7 (the author's original 10). Status: AUTHOR.

$X = \{x_0, \ldots, x_7\}$ are the author's coordinates: x_0 the hidden space-like direction, x_1, x_2, x_3 ordinary space, x_4 the time, x_5, x_6, x_7 the three extra times; H is the author's constant (an inverse length). The author's metric depends on x_0 only through $\cot^2 z$ and $(\sin^2 z)^{1/6}$, so it repeats with the period $\pi/(6H)$ of x_0, and every period $0 < z < \pi$ contains exactly one surface of $\mathcal{F}$, at $z = \pi/2$: $\mathcal{F}$ is the union of these copies, one per period (example 2), and "the fold" without a qualifier is the copy in the period that contains our world. A point of the fold is labelled by the seven coordinates $x_1, \ldots, x_7$, and the metric restricted to the fold ($\mathrm{d}x_0 = 0$, $z = \pi/2$) is

$$g_{MN}\,\mathrm{d}X^M \mathrm{d}X^N\big|_{\mathcal{F}} = \mathrm{e}^{2a_4}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{d}x_4^2 - \mathrm{e}^{-2a_4} b_4^2 \big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big) \tag{FD-2}$$

Symbols in (FD-2).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M \mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- $\mathcal{F}$: The fold: the seven-dimensional surface $\cos(6Hx_0) = 0$ of the eight-dimensional spacetime (FD-1), in one period $0 < z < \pi$ the surface $z = \pi/2$, $x_0 = \pi/(12H)$, where $\sigma = 0$; $|_{\mathcal{F}}$ means restricted to the fold. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.

with three space-like and four time-like directions; a_4 and b_4 are the author's two functions of the time x_4. Notation: ln is the natural logarithm, e Euler's number, $\exp(y) = \mathrm{e}^y$, sin, cos and cot = cos / sin the trigonometric functions,

tanh the hyperbolic tangent, $|y|$ the absolute value; $f(0^+)$ and $f(0^-)$ are the limits of $f(\sigma)$ as σ approaches 0 from above and from below.

Why this surface is called a fold. Measured by the proper distance along the hidden direction,

$$\sigma = \operatorname{sgn}\Big(\frac{\pi}{2} - z\Big)\frac{\ln\sin z}{6H}, \qquad \mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0, \qquad \sin z = \mathrm{e}^{-6H|\sigma|} \tag{FD-3}$$

Symbols in (FD-3).

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \operatorname{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$, the fold is $z = \pi/2$. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.

(the fold is $\sigma = 0$; the half-period $0 < z < \pi/2$ is $\sigma < 0$, the half-period $\pi/2 < z < \pi$ is $\sigma > 0$; the two ends of the period, the tips $z \to 0$ and $z \to \pi$, lie at $\sigma \to -\infty$ and $\sigma \to +\infty$), the author's metric is

$$g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N = \mathrm{d}\sigma^2 + \mathrm{e}^{-2H|\sigma|}\Big[\mathrm{e}^{2a_4}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{e}^{-2a_4}b_4^2\big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big)\Big] - \mathrm{d}x_4^2 \tag{FD-4}$$

Symbols in (FD-4).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \operatorname{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\, \mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.

It depends on σ only through $|\sigma|$. The function $\mathcal{W}(\sigma) = \mathrm{e}^{-2H|\sigma|}$, which multiplies the squared steps of ordinary space and of the extra times, is the warp factor of the metric (its square root $\mathrm{e}^{-H|\sigma|}$ multiplies the lengths); the time x_4 is not warped. Four properties make the surface $\sigma = 0$ a fold, and together they single it out within the period:

1. *It is a mirror.* The reflection $\sigma \to -\sigma$ (that is $z \to \pi - z$, or $x_0 \to \pi/(6H) - x_0$) maps the metric to itself, and the fold is the only surface of the period that the reflection leaves in place: the two halves of the period are mirror images of each other, joined along the fold, as the two halves of a folded sheet of paper are joined along the crease.
2. *The warp factor is largest on it.* The warp factor $(\sin^2 z)^{1/6} = \mathrm{e}^{-2H|\sigma|}$ equals 1 on the fold and decreases on both sides.
3. *The metric has a kink on it.* The metric is continuous across the fold, but its first derivative along σ jumps. The extrinsic curvature of the surfaces $\sigma = \text{const}$ changes sign across the fold,

$$K^a{}_b = \frac{1}{2}\, g^{ac}\, \partial_\sigma g_{cb}: \qquad K^1{}_1 = K^2{}_2 = K^3{}_3 = K^5{}_5 = K^6{}_6 = K^7{}_7 = -H\,\mathrm{sgn}(\sigma), \qquad K^4{}_4 = 0 \tag{FD-5}$$

Symbols in (FD-5).

- $K^a{}_b$: The extrinsic curvature of a surface of constant σ of the hidden direction, $K^a{}_b = \frac{1}{2} g^{ac} \partial_\sigma g_{cb}$ for a, b along the surface (FD-5): $-H\,\mathrm{sgn}(\sigma)$ along ordinary space and the extra times, 0 along the time. Unit: GeV (1/length). Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- c: A frame index (as a). Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+, +, +, +, -, -, -, -$ in the order $x_0, \ldots, x_7$): $g_{MN}\, \mathrm{d}X^M \mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\, \mathrm{d}^8 X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\, \mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z) \ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- η_{ab}: The flat metric of the frame, $\eta = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$ (eta4488 of 4+4.nb); η^{ab} is its inverse (the same matrix). Unit: dimensionless. Status: AUTHOR.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor).

Status: STANDARD.

(a, b frame indices along the fold); it jumps by $-2H$ along ordinary space and along the extra times. The Einstein-Lovelock equations of UM-3, $\sum_p \alpha_{(p)} \mathcal{E}_{(p)}{}^a{}_b + \Lambda \delta^a{}_b = \kappa \mathcal{T}^a{}_b$ ($\mathcal{E}_{(p)}$ the Lovelock tensor of order $p = 1, 2, 3$, $\alpha_{(p)} = w_{(p)} H^{2-2p}$ its coupling with the weights $w = (1, 1/20, 1/360)$, $\Lambda = -7H^2$ the cosmological constant, $\kappa = 8\pi G_8$ the gravitational coupling of eight dimensions, $\mathcal{T}^a{}_b$ the energy-momentum tensor, $\delta^a{}_b$ the Kronecker delta), are linear in the second σ-derivatives of the metric, so the jump puts a Dirac delta function $\delta_{\mathrm{D}}(\sigma)$ into them: the metric solves them only if the fold carries a thin source $\mathcal{S}^a{}_b$, an energy-momentum tensor concentrated on $\sigma = 0$, given by the integral of the equations across the fold (the junction conditions),

$$\kappa\, \mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_{p=1}^{3} \alpha_{(p)}\, \mathcal{E}_{(p)}{}^a{}_b \tag{FD-11}$$

Symbols in (FD-11).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)} \mathcal{E}_{(p)}{}^a{}_b$ (FD-11, the junction conditions); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -\frac{32}{5} H \delta^a{}_b$ at the static block. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\, \mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z) \ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)} H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- $\mathcal{E}_{(p)}{}^M{}_N$: The Lovelock tensor of order p, $\mathcal{E}_{(p)}{}^M{}_N = -2^{-(p+1)}\, \delta^{MP_1\cdots P_{2p}}_{NQ_1\cdots Q_{2p}}\, R^{Q_1Q_2}{}_{P_1P_2} \cdots R^{Q_{2p-1}Q_{2p}}{}_{P_{2p-1}P_{2p}}$ (with both indices down it is $1/\sqrt{|g|}$ times the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to g^{MN}; $\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor, $\mathcal{E}_{(3)}$ the cubic Lovelock tensor); with frame indices its frame components. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- G_8: The eight-dimensional Newton constant ($\kappa = 8\pi G_8$). Unit: GeV^{-6}. Status: AUTHOR.
- $R^{MN}{}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor $R^M{}_{NPQ}$ (in the convention in which the commutator of two covariant derivatives, acting on a vector, is the Riemann tensor, with its last two indices those of the two derivatives, contracted with the vector), with two indices the Ricci tensor $R_{MN} = R^P{}_{MPN}$, without indices the scalar curvature $R = g^{MN} R_{MN}$; with frame indices $R^{(a)(b)}{}_{(c)(d)}$ its frame components, $R^{(a)(b)}{}_{(a)(b)}$ (no sum) the sectional curvature of the frame plane (a, b). Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- $\mathcal{L}_{(p)}$: The Lovelock density of order p, $\mathcal{L}_{(p)} = 2^{-p}\, \delta^{P_1\cdots P_{2p}}_{Q_1\cdots Q_{2p}}\, R^{Q_1Q_2}{}_{P_1P_2} \cdots R^{Q_{2p-1}Q_{2p}}{}_{P_{2p-1}P_{2p}}$, a scalar built from p curvature tensors: $\mathcal{L}_{(1)} = R$ (Einstein-Hilbert), $\mathcal{L}_{(2)}$ the Gauss-Bonnet density, $\mathcal{L}_{(3)}$ the cubic Lovelock density. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.

with $\mathcal{S}^4{}_4 = -$(its energy density), $\mathcal{S}^1{}_1 = \mathcal{S}^2{}_2 = \mathcal{S}^3{}_3$ its pressure along ordinary space and $\mathcal{S}^5{}_5 = \mathcal{S}^6{}_6 = \mathcal{S}^7{}_7$ its pressure along the extra times; $\mathcal{S}$ is $\mathcal{T}$ integrated across the fold, an energy per unit volume of the six other directions of the fold (unit GeV^7). On the static block (a_4 and b_4 constant) and on an expanding fold whose extra times are frozen (b_4 proportional to e^{a_4}; $\mathsf{A} = a_4'/H$, $\mathsf{A}_1 = a_4''/H^2$)

$$\text{static block:}\quad \kappa\mathcal{S}^{a}{}_{b} = -\frac{32}{5}H\,\delta^{a}{}_{b}\quad\text{(a pure tension)},$$
$$\text{frozen extra times:}\quad \kappa\mathcal{S}^{1}{}_{1} = -\frac{16}{15}\big(6+3\mathsf{A}^2+2\mathsf{A}_1\big)H,\quad \kappa\mathcal{S}^{4}{}_{4} = -\frac{16}{5}\big(2+\mathsf{A}^2\big)H, \tag{FD-12}$$
$$\kappa\mathcal{S}^{5}{}_{5} = -\frac{4}{5}\big(8+8\mathsf{A}^2+\mathsf{A}^4+4\mathsf{A}_1+\mathsf{A}^2\mathsf{A}_1\big)H$$

Symbols in (FD-12).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{S}^{a}{}_{b}$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^{a}{}_{b} = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)}\mathcal{E}_{(p)}{}^{a}{}_{b}$ (FD-11, the junction conditions); $\mathcal{S}^{4}{}_{4}$ is minus its energy density, $\mathcal{S}^{1}{}_{1}$ and $\mathcal{S}^{5}{}_{5}$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^{a}{}_{b} = -\frac{32}{5}H\delta^{a}{}_{b}$ at the static block. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- δ: The Kronecker delta: with two indices $\delta^{M}_{N} = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1M_2\cdots}_{N_1N_2\cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- h_{100}: The present expansion rate of the fold in units of 100 km/s/Mpc. Unit: dimensionless. Value: 0.674 (Planck 2018). Status: STANDARD.

the tension plus matter that obeys the Friedmann equations of general relativity. The fold is therefore not only a place but a physical object: a thin layer that carries energy and momentum (a brane). (FD-11 and FD-12 were added to this definition on 2026-10-25, after FD-1 to FD-10, with the corrected Lovelock weights; they stand here because they belong to property 3.) 4. *The author's coordinate x_0 degenerates on it.* The metric component $g_{x_0x_0} = \cot^2(6Hx_0)$ vanishes on the fold and nowhere else in the period; near the fold the proper distance grows with the square of the coordinate distance,

$$|\sigma| = 3H\Big(x_0 - \frac{\pi}{12H}\Big)^2\Big[1 + 6H^2\Big(x_0 - \frac{\pi}{12H}\Big)^2 + \ldots\Big] \tag{FD-6}$$

Symbols in (FD-6).

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.

- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

so the two coordinate values on either side of $x_0 = \pi/(12H)$ at the same coordinate distance lie at the same proper distance from the fold: the coordinate line of x_0 is folded onto the line of proper distances (a fold map, example 5).

What lives on the fold. In UM-3 the Standard Model (the Higgs doublet with the author's ad hoc vacuum, a time-dependent choice of its vacuum value; the gauge fields of $SU(3)_C \times SU(2)_L \times U(1)_Y$; three generations of quarks and leptons and three right-handed neutrinos) and the fold fluid (its tension and the pressure along the extra times that keeps their volume fixed) live on the fold: they are fields at $\sigma = 0$, functions of $x_1, \ldots, x_4$ (no field depends on the extra times). Integrated over the extra times, the fold is the four-dimensional universe that we observe, with the metric $\mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in $x_1, \ldots, x_4$. dirac16complex (the author's spinor field of eight dimensions) and gravity live in the bulk $\sigma \neq 0$ as well as on the fold; the bound states χ of dirac16complex are the part of it that the fold holds, and the portals (couplings between dirac16complex and Standard-Model fields) act at $\sigma = 0$. In UM-3 files "the fold" means this surface; the examples below use the word for other surfaces with the same properties. The word is not used in its other senses: a set of equal energy levels is called degenerate, and a growth by the factor e is called an e-folding.

Examples of folds that are meaningful for UM-3.

1. *Our fold.* The fold $z = \pi/2$ of the period that contains our world (FD-1): the seven-dimensional world-volume of the Standard Model, and after the integration over the extra times the four-dimensional universe that we observe.
2. *The copies of the fold.* Because the metric repeats with the period $\pi/(6H)$, every surface $x_0 = \pi/(12H) +$ (an integer multiple of $\pi/(6H)$) is a copy of the fold, one per period. Neighbouring copies are separated by a tip (z a multiple of π), where the warp factor $(\sin^2 z)^{1/6}$ vanishes and $g_{x_0 x_0} = \cot^2 z$ diverges, and which lies at an infinite proper distance from both: in proper distance every copy has its own complete line $-\infty < \sigma < \infty$. A compact direction of length L would give every field a tower of particles with the masses $2\pi n/L$, $n = 1, 2, \ldots$ (a Kaluza-Klein tower); because the copies are infinitely far apart, the period $\pi/(6H)$ of x_0 produces no such tower with the spacing $12H$, and the x_0 mass of the bound states is their binding to the fold.
3. *A folded sheet of paper.* The crease of a folded sheet is a fold in this sense: the two halves are mirror images (property 1) joined along the crease, the sheet is continuous there and its direction changes abruptly (property 3). In UM-3 the direction across the crease is the hidden direction σ, the seven directions along the crease are $x_1, \ldots, x_7$, and the crease is the fold.
4. *A folded function.* The warp factor $\mathrm{e}^{-2H|\sigma|}$ is the simplest folded function: continuous, mirror symmetric, largest at $\sigma = 0$, with a jump of its derivative,

$$\frac{\mathrm{d}}{\mathrm{d}\sigma}\mathrm{e}^{-2H|\sigma|} = -2H\,\mathrm{sgn}(\sigma)\,\mathrm{e}^{-2H|\sigma|}, \qquad \frac{\mathrm{d}^2}{\mathrm{d}\sigma^2}\mathrm{e}^{-2H|\sigma|} = 4H^2\,\mathrm{e}^{-2H|\sigma|} - 4H\,\delta_{\mathrm{D}}(\sigma) \tag{FD-7}$$

Symbols in (FD-7).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- δ_{D}: The Dirac delta function of σ: zero for $\sigma \neq 0$, with the integral 1 over any range that contains $\sigma = 0$; the integral of $\delta_{\mathrm{D}}(\sigma)$ times a smooth function of σ is the value of that function at $\sigma = 0$. Unit: 1/length (for $\delta_{\mathrm{D}}(\sigma)$). Status: STANDARD.

whose delta function is the origin of the thin source of property 3. Every metric function of FD-4 depends on σ through $|\sigma|$ and is folded at $\sigma = 0$ in this sense. 5. *A fold map.* In the mathematics of smooth maps the fold

(Whitney's fold singularity) is the map $u \mapsto u^2$ of a line onto a half-line: it folds the line at $u = 0$, two points $\pm u$ land on the same point, and the derivative of the map vanishes on the fold. Near the fold the map from the author's coordinate x_0 to the distance $|\sigma|$ from the fold is of this kind (FD-6), and $g_{x_0 x_0} = \cot^2 z$, the square of the derivative $\mathrm{d}\sigma/\mathrm{d}x_0$, vanishes there. 6. *The brane of Randall and Sundrum.* The five-dimensional metric

$$\mathrm{d}y_{\mathrm{RS}}^2 + \mathrm{e}^{-2k_{\mathrm{RS}}|y_{\mathrm{RS}}|}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2 - \mathrm{d}x_4^2\big), \qquad \lambda_{\mathrm{RS}} = \frac{6k_{\mathrm{RS}}}{\kappa_5}, \qquad \Lambda_{\mathrm{RS}} = -\frac{6k_{\mathrm{RS}}^2}{\kappa_5} \tag{FD-8}$$

Symbols in (FD-8).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- y_{RS}: The proper coordinate across the brane of the five-dimensional model of Randall and Sundrum (FD-8, an example, not UM-3). Unit: length. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- k_{RS}: The inverse curvature length of the bulk of the model of Randall and Sundrum: the warp factor of the brane is $\mathrm{e}^{-2k_{\mathrm{RS}}|y_{\mathrm{RS}}|}$ (FD-8, an example, not UM-3). Unit: GeV (1/length). Status: STANDARD.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- λ_{RS}: The tension (energy per unit volume of ordinary space) of the brane of the model of Randall and Sundrum, $\lambda_{\mathrm{RS}} = 6k_{\mathrm{RS}}/\kappa_5$ (FD-8, an example, not UM-3). Unit: GeV^4. Status: STANDARD.
- κ_5: The gravitational coupling of five dimensions in the model of Randall and Sundrum, 8π times the Newton constant of five dimensions (FD-8, an example, not UM-3). Unit: GeV^{-3}. Status: STANDARD.
- Λ_{RS}: The vacuum energy density of the bulk of the model of Randall and Sundrum, $\Lambda_{\mathrm{RS}} = -6k_{\mathrm{RS}}^2/\kappa_5$ (FD-8, an example, not UM-3). Unit: GeV^5. Status: STANDARD.

has a fold at $y_{\mathrm{RS}} = 0$ with the properties 1 to 3 (its coordinate y_{RS} is already the proper distance, so property 4 does not arise); Einstein's equations require there a brane of the tension λ_{RS} (Israel's junction condition) and in the bulk the vacuum energy Λ_{RS}, and the bulk is otherwise empty. The fold of UM-3 is of the same kind: its static block is a vacuum of the bulk with a fold of pure tension (FD-12). It differs in that the time x_4 is not warped, three extra times are warped together with ordinary space, the Lovelock terms add to the junction conditions, and the space is not cut at the brane: Randall and Sundrum identify the two sides of their brane (one physical side), while the fold of UM-3 has two distinct sides (example 7). 7. *A particle bound to a thin layer.* A particle held by an attractive thin layer has a wave function that falls off exponentially with the distance on both sides, and the layer makes its derivative jump (for the first-order Dirac equation, the wave function itself). In UM-3 the particles χ of dark matter are the states of dirac16complex bound to the fold; at rest and without their self-interaction (the field is $\Psi = \mathrm{e}^{3H|\sigma|}\chi(\sigma)\mathrm{e}^{-\mathrm{i}\omega_{\mathrm{b}}x_4}$)

$$\begin{aligned} &\chi(\sigma) = \mathrm{e}^{-q_{\mathrm{b}}|\sigma|}\chi(0^+) \quad (\sigma > 0), \qquad \chi(\sigma) = \mathrm{e}^{-q_{\mathrm{b}}|\sigma|}\chi(0^-) \quad (\sigma < 0), \\ &\chi(0^+) = \exp\Big(\gamma^{(0)}\big(\mu_{\mathrm{L}}P_{\mathrm{L}} + \mu_{\mathrm{R}}P_{\mathrm{R}}\big)\Big)\chi(0^-), \qquad \omega_{\mathrm{b}} = \sqrt{m_\Psi^2 - q_{\mathrm{b}}^2} \end{aligned} \tag{FD-9}$$

Symbols in (FD-9).

- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{3H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of a bound state and $\bar{\chi}$ its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- q_{b}: The inverse width of a state of dirac16complex bound to the fold: its profile falls as $\mathrm{e}^{-q_{\mathrm{b}}|\sigma|}$. Unit: GeV. Status: DERIVED.

- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- μ_X, μ_L, μ_R: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_\mathrm{L}P_\mathrm{L} + \mu_\mathrm{R}P_\mathrm{R})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- P_R: The author's right-handed chiral projector, $P_\mathrm{R} = (\mathbf{1} + \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- Γ_8: The chirality matrix of eight dimensions, $\Gamma_8 = T_0T_1\cdots T_7$ (the author's T16^A[8]) $= \mathrm{diag}(-\mathbf{1}_8, \mathbf{1}_8)$; $\Gamma_8^2 = \mathbf{1}$; it anticommutes with every T_a. Unit: dimensionless. Status: AUTHOR.
- $\mathbf{1}$, $\mathbf{1}_8$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_8$ is the 8×8 identity matrix. Unit: dimensionless. Status: STANDARD.
- ω_b: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: GeV$^{7/2}$. Status: AUTHOR.
- $\mathbf{e}_a{}^M$, $\mathbf{e}^a{}_M$: The vielbein (frame field) of the metric: $g_{MN} = \eta_{ab}\,\mathbf{e}^a{}_M\mathbf{e}^b{}_N$; $\mathbf{e}_a{}^M$ is its inverse; $\mathbf{e}_{bN} = \eta_{bc}\mathbf{e}^c{}_N$; $\mathbf{e}^{(a)}$ is the frame one-form $\mathbf{e}^a{}_M\,\mathrm{d}X^M$. Unit: dimensionless. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- $\vec{k}$: The (comoving) momentum along ordinary space, $\vec{k}^2$ its square. Unit: GeV. Status: STANDARD.
- $\vec{x}$: The position in ordinary space, $\vec{x} = (x_1, x_2, x_3)$. Unit: length. Status: AUTHOR.
- V_t: The coordinate volume of the (compact) extra times, $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$. Unit: length^3. Status: ASSUMED.
- σ_{16}: The matrix of the Dirac adjoint, $\sigma_{16} = T_0T_1T_2T_3$, real and symmetric; $\sigma_{16}T_a$ is real and antisymmetric. Unit: dimensionless. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.

where μ_L, μ_R are the strengths of the fold for the left- and right-handed halves of dirac16complex (set by the Standard-Model fields on the fold) and $q_\mathrm{b} = -m_\Psi(\mu_\mathrm{L} + \mu_\mathrm{R})\tanh\varsigma/(2\varsigma)$, $\varsigma = \sqrt{\mu_\mathrm{L}\mu_\mathrm{R}}$ ($q_\mathrm{b} = -m_\Psi\mu_\mathrm{L}/2$ for $\mu_\mathrm{R} = 0$). The profile is not a function of $|\sigma|$: it jumps at the fold, $\chi(0^+) \neq \chi(0^-)$. 8. *The fold as a mirror for dirac16complex.* The reflection across the fold acts on the profiles $\chi(\sigma)$ in two ways,

$$(\mathsf{R}_1\chi)(\sigma) = T_0T_1T_2T_3\,\chi(-\sigma), \qquad (\mathsf{R}_2\chi)(\sigma) = T_0T_5T_6T_7\,\chi(-\sigma) \tag{FD-10}$$

Symbols in (FD-10).

- R_1: The reflection across the fold acting on a profile of dirac16complex together with the reflection of ordinary space, $(\mathsf{R}_1\chi)(\sigma) = T_0T_1T_2T_3\,\chi(-\sigma)$ (FD-10): a symmetry of dirac16complex, of gravity and of the mean field of the gas of the particles χ; the bound states of Krein charge -1 are even, those of Krein charge $+1$ odd. Unit: dimensionless. Status: DERIVED.
- R_2: The reflection across the fold acting on a profile of dirac16complex together with the reflection of the extra times, $(\mathsf{R}_2\chi)(\sigma) = T_0T_5T_6T_7\,\chi(-\sigma)$ (FD-10): it maps the bound states of one Krein charge onto those of the other. Unit: dimensionless. Status: DERIVED.
- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{3H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of a bound state and $\bar\chi$ its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line

$-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

- T_a, T^a: The author's eight real 16×16 Dirac matrices T16ˆA[a] of 4+4.nb ($a = 0, \ldots, 7$; $T_0, \ldots, T_7$), with $\{T_a, T_b\} = 2\eta_{ab}\mathbf{1}$; $T^a = \eta^{ab}T_b$. Unit: dimensionless. Status: AUTHOR.

(T_a the author's Dirac matrices; for states at rest R_1 is the reflection across the fold combined with the reflection of ordinary space, R_2 the reflection across the fold combined with the reflection of the extra times). The Krein charge of a state is its share of the conserved charge of dirac16complex, which is indefinite: a state can carry the charge -1 or $+1$. R_1 is a symmetry of dirac16complex, of gravity and of the mean field of a gas of the particles χ (the average field that the particles of the gas create and feel): the four bound states of Krein charge -1 (the particles χ) are even under it and the four of Krein charge $+1$ are odd, so the mean field of the gas cannot mix the two signs of the indefinite charge. R_2 maps the bound states of one Krein charge onto those of the other and makes the two sets degenerate without the self-interaction. 9. *The planes of Horava and Witten.* In M-theory (the eleven-dimensional theory whose limits include the superstring theories) on an interval (a circle whose points are identified with their mirror images), the two ten-dimensional planes left in place by the reflection are mirror surfaces (property 1); each carries one E_8 gauge group while gravity fills the bulk. As in UM-3, the gauge fields live on the mirror surface and gravity in the bulk; unlike UM-3, the two sides of each plane are identified (one physical side).

What is not a fold.

- *The tips* $z = 0$ and $z = \pi$ (mod π): the mirror exchanges them within a period, but there the warp factor vanishes, $g_{x_0x_0}$ diverges, and they lie at an infinite proper distance; they carry no thin source.
- *A smooth maximum of a warp factor*, such as $1/\cosh^2(H\sigma)$: mirror symmetric and largest at $\sigma = 0$, but without a kink and without a thin source.

2 How to read this document

2.1 What this document contains

This document solves, approximately and step by step, the coupled field equations of the universe model UM-3: Einstein-Lovelock gravity in eight dimensions with the author's metric, the author's field dirac16complex (a 16-component complex spinor field in the bulk), and the Standard Model with three right-handed neutrinos on the fold of the author's metric. It is self-contained: every equation that is solved is written out here (with the tag it carries in every UM-3 document), every new equation of the solution carries a tag S-1, S-2, ..., and every symbol of every display equation is defined right after it. The sections are:

- **The Lovelock weights and the cosmological constant**: why UM-3 uses the weights $1, 1/20, 1/360$ and $\Lambda = -7H^2$ (the only weights for which the graviton of the fold travels at the speed of light while no graviton of the static bulk travels faster, at the author's order of 2026-10-25), what this correction cannot do, and the record of the author's original values and of the stabiliser that they required.
- **The background** (order 0 of the power series of the metric): the two branches on which the bulk is a vacuum, the static block, the frozen extra times on which the fold obeys the Friedmann equations of general relativity exactly, and the primordial era.
- **The power series of the metric**: the degenerate static block and its strongly coupled perturbations, the fold as a seven-dimensional Einstein brane, and the volume of the extra times, which must be held fixed.
- **The sources of the fold**: the gravitational waves of the fold, which travel at the speed of light at every wavelength at linear order, the polarisation along the extra times, which does not, the gravity that the fields of the fold may induce in addition, and the fold fluid (a tension and the pressure that holds the volume of the extra times).
- **The Higgs field in the ad hoc vacuum**: the author's ad hoc vacuum written as a gauge rotation, the field of W^2 that the Gauss law requires, and the one residual that remains.
- **The spectra along the hidden direction**: the zero modes and the speeds of the bulk fields, the continuum and the bound states of dirac16complex (the x_0 mass), the speed of the bound particles χ and the shortcut through the bulk.
- **The Kohn-Sham states of dirac16complex**: the self-consistent states of a gas of the bound particles χ with the self-interaction, gravity and the mean fields of the Standard Model, in three regimes, with their residuals, and the stability of the gas against the partner states of the other Krein charge.
- **The residuals of every field equation**: how well the approximate solution satisfies each equation.

- **The three neutrino flavours**: the Sun, the envelope of a supernova, the fast instability and the flavour equipartition, and what the newer global fit, the Dirac phase and the lightest mass change.
- **The thermal history**: the degrees of freedom, the relic density of χ, its portals to the Standard Model, the right-handed neutrinos and the ad hoc vacuum before nucleosynthesis.
- **The CMB, the CNB and the BAO** (the cosmic microwave background, the cosmic neutrino background and the baryon acoustic oscillations): the cosmological perturbations of the metric of the fold, against the measurements of Planck, of DESI DR2 (the second data release of the Dark Energy Spectroscopic Instrument) and of ACT DR6 (the sixth data release of the Atacama Cosmology Telescope).
- **What the solution does not settle**, and **what was verified**.

2.2 The status labels

Every equation and every result carries one of the labels

- **AUTHOR**: from the author (his notebooks `4+4.nb` and `Dirac_Einstein-Lovelock_4+4_Nash.nb`, and his orders of 2026-10-24 and 2026-10-25: the coordinates, the metric, the Einstein-Lovelock action, the Dirac matrices, the chiral projectors, the list of fields, the Higgs potential and the ad hoc vacuum; the values of the Lovelock weights and of the cosmological constant are the corrected ones of his order of 2026-10-25: ASSUMED the criterion of the correction, because the order cannot be met in full, DERIVED the unique values that meet it);
- **RECORD**: from the record of this repository (the Lagrangian of dirac16complex with its canonical spin connection and its contact self-interaction, the Kohn-Sham treatment);
- **STANDARD**: textbook physics, or a published measurement (the source is named);
- **ASSUMED**: a choice made in UM-3 without derivation;
- **INVENTED**: a new interaction of UM-3 (the portals of dirac16complex);
- **DERIVED**: computed and verified in UM-3 (the verification is stated);
- **ESTIMATE**: an order-of-magnitude evaluation;
- **OPEN**: not settled.

2.3 Every equation is followed by its symbols

Every display equation is followed by the list "Symbols in (S-n)" or "Symbols in (FE-n)" of EVERY symbol that appears in it: every quantity, field, coupling, constant and operator, every index letter with the values it takes, every mark (prime, dagger, transpose, star), the imaginary unit, Euler's number and the differential. Each item gives the meaning, the unit (natural units, in which the reduced Planck constant $\hbar$ and the speed of light c are 1), the value where one exists, and the status. A symbol that one of these meanings uses is itself listed in this document. A field equation of UM-3 carries a tag FE-n and a term of the action a tag IN-n, the same in every UM-3 document; it is written here exactly as everywhere else. A new equation of this document carries a tag S-n.

2.4 Coordinates, signature and units

The author's coordinates are $X = \{x_0, x_1, \ldots, x_7\}$, always with the hidden direction first: x_0 is the hidden space-like coordinate, on which the metric depends periodically with the period $\pi/(6H)$ through the angle $z = 6Hx_0$; x_1, x_2, x_3 are ordinary space, x_4 is the time, and x_5, x_6, x_7 are three extra time-like coordinates. The signature is $(+,+,+,+,-,-,-,-)$ in this order; the four observable coordinates $x_1, \ldots, x_4$ have the signature $(+,+,+,-)$. H is the author's constant (an inverse length), a free parameter of UM-3. Units are natural: the reduced Planck constant $\hbar$ and the speed of light c are 1 ($1\ \mathrm{GeV}^{-1} = 1.973 \times 10^{-14}$ cm $= 6.582 \times 10^{-25}$ s; $1\ \mathrm{GeV}^{-2} = 3.894 \times 10^{-28}$ cm^2). A prime on a function of the time x_4 alone is its derivative $\mathrm{d}/\mathrm{d}x_4$, and on a profile (a function of σ alone) its derivative $\mathrm{d}/\mathrm{d}\sigma$. The proper coordinate of the hidden direction is σ (FE-27): $\sigma = 0$ is the fold $z = \pi/2$. The fields live as follows: gravity in the bulk and on the fold; dirac16complex in the bulk (with the corrected Lovelock weights the static bulk is a vacuum and needs no other source); the Standard Model with three right-handed neutrinos, a thin fluid (the fold fluid) and, possibly, an Einstein-Hilbert term that the fields of the fold induce (with the coupling κ_{ind}) on the fold $\sigma = 0$.

2.5 The solution at a glance

part	result	status
Lovelock weights	only the weights $1, 1/20, 1/360$ give a graviton of the fold at the speed of light without a faster graviton in the static bulk; with them the static block is a vacuum for $\Lambda = -7H^2$ (the Born-Infeld gravity of eight dimensions), the bulk has no linear graviton at all and is strongly coupled, and the stabiliser that the author's original weights required vanishes; no weights give gravitons at the speed of light both in the bulk and on the fold, because the time of the author's metric is not warped	DERIVED, OPEN (the strongly coupled bulk)
background	the bulk is a vacuum on exactly two branches, static ordinary space (any b_4) and frozen extra times (any a_4); on the second the fold obeys the Friedmann equations of general relativity exactly, with $8\pi G_\mathrm{N} = \kappa_7/V_*$; the author's vacuum is not a solution, and the primordial era can be an inflation of the fold only if an enormous pressure holds the volume of the extra times fixed	DERIVED, OPEN (that pressure)
power series	the static block is a degenerate vacuum: the bulk equations have no linear part, static deformations grow like $\mathrm{e}^{3H\lvert\sigma\rvert}$, and the response of the strongly coupled bulk is OPEN; the fold is a seven-dimensional Einstein brane; a free volume of the extra times would change Newton's constant 6800 times faster than lunar laser ranging allows, so the fold fluid holds it fixed	DERIVED, ASSUMED (the holding pressure), OPEN (the bulk response)
Higgs field	the ad hoc vacuum is an $SU(2)$ rotation of the radial field $\sqrt{v^2 + \mathcal{A}^2}$ with a pure-gauge W^2; one radial residual remains, which vanishes as $\mathcal{A} \to 0$	DERIVED
spectra	bulk scalars, photons and fermions have zero modes that move faster than light on the fold, the graviton of the fold has no bulk term and travels at the speed of light at linear order; dirac16complex has a continuum above m_Ψ and eight bound states at $\omega_\mathrm{b} = \sqrt{m_\Psi^2 - q_\mathrm{b}^2}$, the x_0 mass	DERIVED
Kohn-Sham	self-consistent states of a homogeneous gas of the particles χ (S-14 to S-19) in three regimes up to $G_\mathrm{KS} = 1$: first-order perturbation theory in the dilute limit, saturation of the proper density far from the fold, the level raised through $\omega = m_\Psi$ without unbinding, a positive pressure that vanishes in the dilute limit (cold dark matter); the Higgs-portal loop converges; a weakly bound point compatible with direct detection	DERIVED
residuals	every field equation is satisfied to the stated accuracy, except the radial residual of the ad hoc vacuum and the extra-time component of the junction, which are carried by stated assumptions	DERIVED
neutrinos	the flavour conversion of UM-3 is that of the Standard Model with three Dirac neutrinos: solar survival 0.55 at low and 0.30 at high energy (Borexino 0.57, 0.53, 0.43, 0.37); adiabatic supernova resonances; fast instabilities within centimetres wherever the electron lepton number crosses; equipartition raises or lowers the heating behind the shock depending on the spectra	DERIVED, ESTIMATE
thermal history	χ is a thermal relic for $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle = 2.2 \times 10^{-26}$ cm^3/s in the normalisation of FE-61 (one χ with one $\bar{\chi}$: 9×10^{-26} cm^3/s); quark portals cannot provide it (direct detection); lepton portals can above about 1.5 TeV; the Higgs portal requires $q_\mathrm{b}/\omega_\mathrm{b} \lesssim 0.06$ at 100 GeV to 0.02 at 10 TeV; the right-handed neutrinos must not thermalise	DERIVED, ESTIMATE
sources of the fold	the Gauss-Bonnet and cubic terms give the fold its own Einstein-Hilbert term: the four-dimensional gravitational waves travel at the speed of light at every wavelength at linear order (GW170817 is satisfied for every H), with the Planck mass of the expansion; the polarisation along the extra times, a scalar of the four-dimensional world, does not; an induced term is optional; the fold fluid is a pure tension plus the pressure $q_\mathrm{hold} = -0.793\rho_\mathrm{c}^{(0)}$ today along the extra times (with the pressure $-\rho_\mathrm{b}$ of the baryons along them, ASSUMED)	DERIVED, ASSUMED, ESTIMATE, OPEN

part	result	status
speed of χ	speed coefficient $c_{\mathrm{b}}^2 = 1.190$ (S-13: 1.2); group velocity above 1 for $\lvert\vec{k}\rvert$ above about $6m_\Psi$, at most 1.0046, front velocity that of light; the bulk is a shortcut for signals; the four time-like directions allow closed time-like curves	DERIVED
Krein norm	the vacuum is stable; whether the gas of χ is classically stable is not decided (exact flat directions, a truncated response of the size of its error, complex partner levels not converged in the size of the box); chiral couplings at the fold produce ghosts, so UM-3 with them is not unitary	DERIVED, OPEN
CMB, CNB, BAO	with the Planck 2018 parameters UM-3 reproduces the CMB ($\chi^2 = 584$ for 613 high-l bins) and its derived quantities; the BAO of DESI DR2 show the same tension as the cosmological-constant model (2.3 standard deviations, DESI's number), since the dark energy of UM-3 has $w = -1$ (its pressure is minus its energy density); the CNB of Dirac neutrinos has one helicity per state	DERIVED, STANDARD (the DESI tension)
neutrino parameters	NuFIT 6.0 changes the results by at most 1.5 per cent; the Dirac phase drops out of every result of UM-3 (it enters only the appearance channels); DESI DR2 with NuFIT 6.0 disfavours the inverted ordering and bounds the lightest mass below 23 meV, which lowers the CNB capture rate from 6.9 to 4.2 per year at most	DERIVED

3 The Lovelock weights and the cosmological constant

Gravity in UM-3 is the Einstein-Lovelock theory in eight dimensions: the Einstein-Hilbert term and the two higher Lovelock terms (Gauss-Bonnet and cubic), each with a weight, and a cosmological constant. The author's notebook used the weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ and the cosmological constant $\Lambda = -10H^2$. With them gravitational waves in the bulk are faster than light, and the static bulk is not a vacuum. The author's order of 2026-10-25 reads: "Gravitons must travel at c. If these incorrect claims come from my choice of the Lovelock couplings and my choice of the cosmological constant Λ THEN my choices are wrong, and you MUST correct them." This section derives the correction: the only weights for which the graviton of the fold travels at the speed of light while no graviton of the static bulk travels faster, and the cosmological constant that makes the static block a vacuum with them; and it shows what no choice of the weights can do: give gravitons at the speed of light in the bulk as well. Every later section of this document uses the corrected values; the results of the original values are kept as records and labelled so.

$$S_{\mathrm{EL}} = \frac{1}{2\kappa}\int \mathrm{d}^8X\,\sqrt{|g|}\,\Big(\sum_{p=1}^{3}\alpha_{(p)}\,\mathcal{L}_{(p)} - 2\Lambda\Big), \qquad \kappa = 8\pi G_8,$$
$$\alpha_{(p)} = w_{(p)}\,H^{2-2p}, \qquad w_{(1)} = 1, \quad w_{(2)} = \frac{1}{20}, \quad w_{(3)} = \frac{1}{360}, \qquad \Lambda = -7H^2 \tag{IN-2}$$

Symbols in (IN-2).

- S_{EL}: The Einstein-Lovelock action of gravity in eight dimensions. Unit: dimensionless. Status: AUTHOR.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.

- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)} H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- $\mathcal{L}_{(p)}$: The Lovelock density of order p, $\mathcal{L}_{(p)} = 2^{-p}\, \delta^{P_1 \cdots P_{2p}}_{Q_1 \cdots Q_{2p}}\, R^{Q_1 Q_2}{}_{P_1 P_2} \cdots R^{Q_{2p-1} Q_{2p}}{}_{P_{2p-1} P_{2p}}$, a scalar built from p curvature tensors: $\mathcal{L}_{(1)} = R$ (Einstein-Hilbert), $\mathcal{L}_{(2)}$ the Gauss-Bonnet density, $\mathcal{L}_{(3)}$ the cubic Lovelock density. Unit: 1/length^{2p}. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -7H^2$: with the corrected weights the value for which the static block is a vacuum (its Lovelock tensors add to $7H^2$ in every direction). The author's original value was $\Lambda = -10H^2$ (he writes the pure number $\Lambda_{\mathrm{N}} = 10$ in $0 = -\Lambda_{\mathrm{N}} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$). Unit: GeV^2. Value: $-7H^2$; the author's original $-10H^2$. Status: AUTHOR (the term), DERIVED (the value).
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_8: The eight-dimensional Newton constant ($\kappa = 8\pi G_8$). Unit: GeV^{-6}. Status: AUTHOR.
- $w_{(p)}$: The Lovelock weights, $w_{(1)} = 1$, $w_{(2)} = 1/20$, $w_{(3)} = 1/360$: the only weights for which both coefficients of a transverse traceless gravitational wave in the static bulk vanish, $1 - 40w_{(2)} + 360w_{(3)}$ (of its second time derivative) and $1 - 24w_{(2)} + 72w_{(3)}$ (of its second derivatives along ordinary space and along σ). Then the four-dimensional graviton of the fold travels at the speed of light at linear order, and no graviton of the static bulk travels faster, because the static bulk has no linear graviton at all (the Born-Infeld point of Lovelock gravity in eight dimensions). The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) made bulk gravitons $\sqrt{5}$ times faster than light; they were corrected at the author's order of 2026-10-25. Unit: dimensionless. Value: $(1, 1/20, 1/360)$; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), ASSUMED (the criterion of the correction), DERIVED (the unique values that meet it).
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

The gravitational action of UM-3 with the corrected weights. The weights multiply the Lovelock densities in units of the author's constant H ($\alpha_{(p)} = w_{(p)} H^{2-2p}$), so that every term of the field equations FE-1 is H^2 times a pure number on the backgrounds of UM-3. Status: AUTHOR (the action and its form), ASSUMED (the criterion of the correction, below), DERIVED (the values $w_{(2)} = 1/20$, $w_{(3)} = 1/360$, $\Lambda = -7H^2$ that meet it, below).

$$\frac{c_{\mathrm{t}}}{2}\,\frac{\partial^2 h_{\mathrm{gw}}}{\partial x_4^2} - \frac{c_{\mathrm{x}}}{2}\Big(\mathrm{e}^{2H|\sigma|}\,\frac{\partial^2 h_{\mathrm{gw}}}{\partial x_3^2} + \frac{\partial^2 h_{\mathrm{gw}}}{\partial \sigma^2} - 6H\,\mathrm{sgn}(\sigma)\,\frac{\partial h_{\mathrm{gw}}}{\partial \sigma}\Big) = 0, \qquad \text{(FE-68)}$$
$$c_{\mathrm{t}} = 1 - 40w_{(2)} + 360w_{(3)}, \qquad c_{\mathrm{x}} = 1 - 24w_{(2)} + 72w_{(3)}$$

Symbols in (FE-68).

- c_{t}: The coefficient of the second time derivative of a transverse traceless gravitational wave in the bulk at the static block, in units of its value with Einstein gravity alone: $c_{\mathrm{t}} = 1 - 40w_{(2)} + 360w_{(3)}$ (FE-68). Unit: dimensionless. Value: 0 with the corrected weights; 1/16 with the author's original weights. Status: DERIVED.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- h_{gw}: The amplitude of a gravitational wave of the + polarisation travelling along x_3: the metric components along x_1 and x_2 multiplied by $\mathrm{e}^{h_{\mathrm{gw}}}$ and $\mathrm{e}^{-h_{\mathrm{gw}}}$; a function of σ, x_3 and x_4 in the bulk, and on the fold the gravitational wave of the four-dimensional world. Unit: dimensionless. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- c_{x}: The coefficient of the second derivatives along ordinary space and along σ of the same wave, in the same units: $c_{\mathrm{x}} = 1 - 24w_{(2)} + 72w_{(3)}$ (FE-68); bulk gravitons move with the local speed $\sqrt{c_{\mathrm{x}}/c_{\mathrm{t}}}$. Unit: dimensionless. Value: 0 with the corrected weights; 5/16 with the author's original weights. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \operatorname{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- $w_{(p)}$: The Lovelock weights, $w_{(1)} = 1$, $w_{(2)} = 1/20$, $w_{(3)} = 1/360$: the only weights for which both coefficients of a transverse traceless gravitational wave in the static bulk vanish, $1 - 40w_{(2)} + 360w_{(3)}$ (of its second time derivative) and $1 - 24w_{(2)} + 72w_{(3)}$ (of its second derivatives along ordinary space and along σ). Then the four-dimensional graviton of the fold travels at the speed of light at linear order, and no graviton of the static bulk travels faster, because the static bulk has no linear graviton at all (the Born-Infeld point of Lovelock gravity in eight dimensions). The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) made bulk gravitons $\sqrt{5}$ times faster than light; they were corrected at the author's order of 2026-10-25. Unit: dimensionless. Value: $(1, 1/20, 1/360)$; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), ASSUMED (the criterion of the correction), DERIVED (the unique values that meet it).

The solution, step 1: a gravitational wave in the bulk. The wave h_{gw} (the + polarisation along x_1, x_2, travelling along x_3) on the static block (a_4 constant, chosen as 0 by the unit of x_3, so that the factor of the second derivative along x_3 is $\mathrm{e}^{2H|\sigma|}$) was put into the exact field equations of the general warped metric and expanded to first order; the coefficients of its second derivatives (the principal symbol) were computed order by order in the Lovelock series and then added with general weights. In units of the Einstein term the time-kinetic coefficient is c_{t} and the gradient coefficient c_{x}; per unit weight, the Einstein, Gauss-Bonnet and cubic terms contribute $(1, 1)$, $(-40, -24)$ and $(360, 72)$ to $(c_{\mathrm{t}}, c_{\mathrm{x}})$. A wave of the bulk moves with the local speed $\sqrt{c_{\mathrm{x}}/c_{\mathrm{t}}}$. With the author's original weights $c_{\mathrm{t}} = 1 - 40/32 + 360/1152 = 1/16$ and $c_{\mathrm{x}} = 1 - 24/32 + 72/1152 = 5/16$: the local speed was $\sqrt{5} = 2.24$ times the speed of light. Status: DERIVED (`um3_weights.py`, W1; `um3_bulk.py`, B5, by an independent principal symbol).

$$\int_{0^-}^{0^+} \mathrm{d}\sigma\,(\dots) = \frac{16\big(w_{(2)} - 6w_{(3)}\big)}{H}\,\frac{\partial^2 h_{\mathrm{gw}}}{\partial x_4^2} - \frac{12\big(w_{(2)} - 2w_{(3)}\big)}{H}\,\frac{\partial^2 h_{\mathrm{gw}}}{\partial x_3^2} - \frac{c_{\mathrm{x}}}{2}\Big(\frac{\partial h_{\mathrm{gw}}}{\partial\sigma}\Big|_{0^+} - \frac{\partial h_{\mathrm{gw}}}{\partial\sigma}\Big|_{0^-}\Big) = \kappa\,\Pi_{\mathrm{gw}},$$

$$\kappa K_{44} = \frac{c_{\mathrm{t}}}{6H} + \frac{16\big(w_{(2)} - 6w_{(3)}\big)}{H}, \qquad \kappa K_{33} = \frac{c_{\mathrm{x}}}{4H} + \frac{12\big(w_{(2)} - 2w_{(3)}\big)}{H}, \qquad \kappa\big(K_{44} - K_{33}\big) = -\frac{c_{\mathrm{t}}}{12H}, \qquad c_{\mathrm{g}}^2 = \frac{K_{33}}{K_{44}} \tag{FE-69}$$

Symbols in (FE-69).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \operatorname{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $w_{(p)}$: The Lovelock weights, $w_{(1)} = 1$, $w_{(2)} = 1/20$, $w_{(3)} = 1/360$: the only weights for which both coefficients of a transverse traceless gravitational wave in the static bulk vanish, $1 - 40w_{(2)} + 360w_{(3)}$ (of its second time derivative) and $1 - 24w_{(2)} + 72w_{(3)}$ (of its second derivatives along ordinary space and along σ). Then the four-dimensional graviton of the fold travels at the speed of light at linear order, and no graviton of the static bulk travels faster, because the static bulk has no linear graviton at all (the Born-Infeld point of Lovelock gravity in eight dimensions). The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) made bulk gravitons $\sqrt{5}$ times faster than light; they were corrected at the author's order of 2026-10-25. Unit: dimensionless. Value: $(1, 1/20, 1/360)$; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), ASSUMED (the criterion of the correction), DERIVED (the unique values that meet it).
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- h_{gw}: The amplitude of a gravitational wave of the + polarisation travelling along x_3: the metric components

along x_1 and x_2 multiplied by $\mathrm{e}^{h_{\mathrm{gw}}}$ and $\mathrm{e}^{-h_{\mathrm{gw}}}$; a function of σ, x_3 and x_4 in the bulk, and on the fold the gravitational wave of the four-dimensional world. Unit: dimensionless. Status: STANDARD.

- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- c_{x}: The coefficient of the second derivatives along ordinary space and along σ of the same wave, in the same units: $c_{\mathrm{x}} = 1 - 24w_{(2)} + 72w_{(3)}$ (FE-68); bulk gravitons move with the local speed $\sqrt{c_{\mathrm{x}}/c_{\mathrm{t}}}$. Unit: dimensionless. Value: 0 with the corrected weights; 5/16 with the author's original weights. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- Π_{gw}: The transverse traceless stress of the matter of the fold that drives the gravitational wave h_{gw} (half the difference of its components along x_1 and x_2), integrated across the fold (eight-dimensional normalisation). Unit: GeV^7. Status: STANDARD.
- K_{44}: The coefficient of $\partial^2 h_{\mathrm{gw}}/\partial x_4^2$ in the equation of the long-wavelength gravitational wave (the same at every σ) at the static block, integrated across the hidden direction with the volume factor $\mathrm{e}^{-6H|\sigma|}$: the part of the bulk plus the thin part at the fold. Unit: GeV^5 (κK_{44} is a length). Status: DERIVED.
- c_{t}: The coefficient of the second time derivative of a transverse traceless gravitational wave in the bulk at the static block, in units of its value with Einstein gravity alone: $c_{\mathrm{t}} = 1 - 40w_{(2)} + 360w_{(3)}$ (FE-68). Unit: dimensionless. Value: 0 with the corrected weights; 1/16 with the author's original weights. Status: DERIVED.
- K_{33}: The coefficient of $-\partial^2 h_{\mathrm{gw}}/\partial x_3^2$ in the same equation, with the same two parts; the wave travels at $c_{\mathrm{g}}^2 = K_{33}/K_{44}$. Unit: GeV^5. Status: DERIVED.
- c_{g}: The speed of gravitational waves on the fold, in units of the speed of light. Unit: dimensionless. Value: with the corrected weights 1 at every wavelength, at linear order, for the four-dimensional graviton (the polarisations along ordinary space) on the static block and on every fold with frozen extra times; the polarisation along the extra times, a scalar of the four-dimensional world, is not luminal on an expanding fold; with the author's original weights the long-wavelength mode had $c_{\mathrm{g}}^2 = 83/82$ and shorter waves were slower than light. Status: DERIVED.

The solution, step 2: the same wave at the fold. Integrating the linearised equation across the kink gives the thin part (first line); a wave of long wavelength, with the same amplitude at every σ, collects the bulk part with the volume factor $\mathrm{e}^{-6H|\sigma|}$ and the thin part into the coefficients of the second line. For every pair of weights $\kappa(K_{44} - K_{33}) = -c_{\mathrm{t}}/(12H)$, so the wave of the fold travels at the speed of light only if $c_{\mathrm{t}} = 0$; and $c_{\mathrm{t}} = 0$ with $c_{\mathrm{x}} \neq 0$ would leave the bulk graviton a gradient term without a time-kinetic term, an infinite local speed. A graviton of the fold at the speed of light without a faster graviton in the static bulk therefore needs $c_{\mathrm{t}} = c_{\mathrm{x}} = 0$: $1 - 40w_{(2)} + 360w_{(3)} = 0$ and $1 - 24w_{(2)} + 72w_{(3)} = 0$, whose only solution is $w_{(2)} = 1/20$, $w_{(3)} = 1/360$. With these weights the bulk part of the wave vanishes, $\kappa K_{44} = \kappa K_{33} = 16(1/20 - 6/360)/H = 12(1/20 - 2/360)/H = 8/(15H)$, and $c_{\mathrm{g}} = 1$. What no choice of the weights can do is to give gravitons at the speed of light in the bulk as well: that needs $c_{\mathrm{t}} = c_{\mathrm{x}} = c_{\mathrm{tx}} \neq 0$, the line $w_{(2)} = 18w_{(3)}$ with $c_{\mathrm{tx}} = 1 - 360w_{(3)}$, and on it $\kappa K_{44} = c_{\mathrm{tx}}/(6H) + 192w_{(3)}/H = (16 - 11c_{\mathrm{tx}})/(30H)$ and $\kappa K_{33} = c_{\mathrm{tx}}/(4H) + 192w_{(3)}/H$, so $c_{\mathrm{g}}^2 - 1 = 5c_{\mathrm{tx}}/(32 - 22c_{\mathrm{tx}})$, which is not zero for any $c_{\mathrm{tx}} \neq 0$: positive, the wave of the fold faster than light, for $0 < c_{\mathrm{tx}} < 16/11$ (every $0 < c_{\mathrm{tx}} \leq 1$ among them); for $c_{\mathrm{tx}} < 0$ the bulk graviton is a ghost, and for $c_{\mathrm{tx}} > 16/11$ $K_{44} < 0$. The cause is the author's metric: ordinary space and the extra times are warped, the time is not. A metric whose time is warped as well might allow both, but the metric is the author's, and whether to change it is his decision. Status: DERIVED (`um3_weights.py`, W2 and W3).

$$
\begin{aligned}
&\sum_{p=1}^{3} \alpha_{(p)}\, {\mathcal{E}_{(p)}}^{1}{}_{1} + \Lambda = \big(15 - 180w_{(2)} + 360w_{(3)}\big)H^2 + \Lambda,\\
&\sum_{p=1}^{3} \alpha_{(p)}\, {\mathcal{E}_{(p)}}^{4}{}_{4} + \Lambda = \big(21 - 420w_{(2)} + 2520w_{(3)}\big)H^2 + \Lambda,\\
&w_{(2)} = \frac{1}{20},\ w_{(3)} = \frac{1}{360}: \quad \text{both } 7H^2 + \Lambda, \qquad \Lambda = -7H^2;\\
&\text{curvature } -\mathsf{y}H^2: \quad (\mathsf{y}-1)^2(\mathsf{y}-7)\,H^2, \quad 7(\mathsf{y}-1)^3H^2
\end{aligned}
\tag{FE-70}
$$

Symbols in (FE-70).

- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)}H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.

- $\mathcal{E}_{(p)}{}^{M}{}_{N}$: The Lovelock tensor of order p, $\mathcal{E}_{(p)}{}^{M}{}_{N} = -2^{-(p+1)}\,\delta^{MP_1\cdots P_{2p}}_{NQ_1\cdots Q_{2p}}\,R^{Q_1Q_2}{}_{P_1P_2}\cdots R^{Q_{2p-1}Q_{2p}}{}_{P_{2p-1}P_{2p}}$ (with both indices down it is $1/\sqrt{|g|}$ times the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to g^{MN}; $\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor, $\mathcal{E}_{(3)}$ the cubic Lovelock tensor); with frame indices its frame components. Unit: $1/\text{length}^{2p}$. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -7H^2$: with the corrected weights the value for which the static block is a vacuum (its Lovelock tensors add to $7H^2$ in every direction). The author's original value was $\Lambda = -10H^2$ (he writes the pure number $\Lambda_{\mathrm{N}} = 10$ in $0 = -\Lambda_{\mathrm{N}} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$). Unit: GeV^2. Value: $-7H^2$; the author's original $-10H^2$. Status: AUTHOR (the term), DERIVED (the value).
- $w_{(p)}$: The Lovelock weights, $w_{(1)} = 1$, $w_{(2)} = 1/20$, $w_{(3)} = 1/360$: the only weights for which both coefficients of a transverse traceless gravitational wave in the static bulk vanish, $1 - 40w_{(2)} + 360w_{(3)}$ (of its second time derivative) and $1 - 24w_{(2)} + 72w_{(3)}$ (of its second derivatives along ordinary space and along σ). Then the four-dimensional graviton of the fold travels at the speed of light at linear order, and no graviton of the static bulk travels faster, because the static bulk has no linear graviton at all (the Born-Infeld point of Lovelock gravity in eight dimensions). The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) made bulk gravitons $\sqrt{5}$ times faster than light; they were corrected at the author's order of 2026-10-25. Unit: dimensionless. Value: $(1, 1/20, 1/360)$; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), ASSUMED (the criterion of the correction), DERIVED (the unique values that meet it).
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- y: The curvature of a seven-dimensional space of constant curvature $-\mathsf{y}H^2$ (in units of $-H^2$), times the time line: $\mathsf{y} = 1$ is the static block. Unit: dimensionless. Status: DERIVED.

The solution, step 3: the cosmological constant. The static block (FE-32 below) is a space of constant curvature $-H^2$ in the seven directions other than the time, times the time line, so its Lovelock tensors are pure numbers times H^2, different along the warped directions and along the time. A cosmological constant alone can make it a vacuum only if the two numbers of FE-70 are equal; their difference is $-6c_{\mathrm{t}}H^2$ for every pair of weights, so this needs $c_{\mathrm{t}} = 0$, the condition of the luminal wave of the fold. With $w_{(2)} = 1/20$, $w_{(3)} = 1/360$ both are $7H^2 + \Lambda$ (Einstein $(15, 21)$, Gauss-Bonnet $(-9, -21)$, cubic $(1, 7)$), so $\Lambda = -7H^2$, the cosmological constant of UM-3. The vacuum is degenerate: on the family of spaces of constant curvature $-\mathsf{y}H^2$ the left-hand side has a double root $((\mathsf{y}-1)^2(\mathsf{y}-7)$ along the warped directions) and a triple root $(7(\mathsf{y}-1)^3$ along the time) at $\mathsf{y} = 1$. The entry along the time is that of a space of constant curvature in all eight directions, and these weights with this Λ are the Born-Infeld gravity of eight dimensions with the radius $1/H$ (up to its term of order 4, the Lovelock density of order 4 with the weight 1/2880, a total derivative in eight dimensions), whose maximally symmetric vacuum is a triple root (M. Bañados, C. Teitelboim and J. Zanelli, Phys. Rev. D 49 (1994) 975; J. Crisóstomo, R. Troncoso and J. Zanelli, Phys. Rev. D 62 (2000) 084013). At that maximally symmetric vacuum the field equations cannot be linearised (G. Arenas-Henriquez, R. B. Mann, O. Miskovic and R. Olea, Phys. Rev. D 100 (2019) 064038). The static block is not of constant curvature, and the roots of the one-parameter family do not by themselves show that the bulk equations have no linear part around it; the reason is that with these weights and Λ the left-hand side of the bulk equations is cubic in the shifted curvature, the curvature plus H^2 times the generalised Kronecker delta, which at the static block is nonzero only in the planes that contain the time x_4 (where it is H^2): every term of first order in a perturbation has two background factors of it, hence the index x_4 twice in one row of the generalised Kronecker delta, and vanishes. Around the static block the linearised bulk equations therefore vanish identically for every perturbation of the metric, so the static bulk has no linear graviton at all and is strongly coupled (S-5). Status: DERIVED (`um3_weights.py`, W3 and W4; `um3_linear.py`, LN1 to LN3; `um3_stabiliser.py`, ST9).

$$\kappa\,\mathcal{T}_{\mathrm{stab}}{}^{a}{}_{b} = \Big(\sum_{p=1}^{3} \alpha_{(p)}\,\mathcal{E}_{(p)}{}^{a}{}_{b} + \Lambda\,\delta^{a}{}_{b}\Big)\Big|_{\text{static block}} = 0 : \qquad \text{(IN-7)}$$
$$(15 - 9 + 1 - 7)H^2 = 0 \text{ (warped directions)}, \qquad (21 - 21 + 7 - 7)H^2 = 0 \text{ (the time)}$$

Symbols in (IN-7).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\mathrm{stab}}{}^{a}{}_{b}$: The energy-momentum tensor of the stabiliser of the bulk (frame components): the source that a static bulk of the author's metric required with the author's original Lovelock weights, found and named by

Claude (Anthropic) on 2026-10-24; with the corrected weights and $\Lambda = -7H^2$ it is zero (IN-7). Unit: GeV^8. Value: 0 with the corrected weights. Status: DERIVED.

- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)} H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- ${\mathcal{E}_{(p)}}^M{}_N$: The Lovelock tensor of order p, ${\mathcal{E}_{(p)}}^M{}_N = -2^{-(p+1)}\, \delta^{M P_1 \cdots P_{2p}}_{N Q_1 \cdots Q_{2p}}\, R^{Q_1 Q_2}{}_{P_1 P_2} \cdots R^{Q_{2p-1} Q_{2p}}{}_{P_{2p-1} P_{2p}}$ (with both indices down it is $1/\sqrt{|g|}$ times the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to g^{MN}; $\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor, $\mathcal{E}_{(3)}$ the cubic Lovelock tensor); with frame indices its frame components. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -7H^2$: with the corrected weights the value for which the static block is a vacuum (its Lovelock tensors add to $7H^2$ in every direction). The author's original value was $\Lambda = -10H^2$ (he writes the pure number $\Lambda_\mathrm{N} = 10$ in $0 = -\Lambda_\mathrm{N} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$). Unit: GeV^2. Value: $-7H^2$; the author's original $-10H^2$. Status: AUTHOR (the term), DERIVED (the value).
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

The record of the stabiliser. With the author's original weights and $\Lambda = -10H^2$ the left-hand side of the bulk equations at the static block was $H^2\,\mathrm{diag}(-5/16, -5/16, -5/16, -5/16, 1/16, -5/16, -5/16, -5/16)$ (order by order along the warped directions and the time, in units of H^2: Einstein $(15, 21)$, Gauss-Bonnet $(-45/8, -105/8)$, cubic $(5/16, 35/16)$, Λ $(-10, -10)$): a static bulk needed a medium at rest with the energy density $\kappa\rho_\mathrm{s} = -H^2/16$ and the pressure $\kappa p_\mathrm{s} = -5H^2/16$ in every direction other than the time. Claude (Anthropic) found it while deriving the field equations of UM-3 on 2026-10-24 and named it the stabiliser of the bulk; it had a negative energy density, violated the null energy condition ($\kappa(\rho_\mathrm{s} + p_\mathrm{s}) = -3H^2/8$), and among the scalar fields only one with a negative kinetic energy (a ghost) reproduced it. The stabiliser was a consequence of the original weights: with the corrected weights the same four numbers add to zero in every direction (IN-7), the static bulk is a vacuum, and UM-3 has no stabiliser. Status: DERIVED (`um3_stabiliser.py`, ST1 to ST9, the record and its removal).

Where the speed 2.74 came from. An earlier version of this document stated that gravitons in the bulk travel $\sqrt{15/2} = 2.74$ times faster than light. The figure was not an assumption of the author; it was a deduction of Claude from the author's original weights: a wave of long wavelength in the bulk alone has the coefficients $c_\mathrm{t}/(6H)$ and $c_\mathrm{x}/(4H)$ (FE-69, without the thin part), whose ratio is $\frac{3}{2} \cdot c_\mathrm{x}/c_\mathrm{t} = \frac{3}{2} \cdot 5 = 15/2$; together with the thin part the wave had $c_\mathrm{g}^2 = 83/82$ (S-30). With the corrected weights the bulk part is zero and the wave of the fold is luminal at every wavelength at linear order (FE-63).

What the correction does and does not do. It gives the four-dimensional graviton of the fold the speed of light at every wavelength at linear order, on the static block and on every fold with frozen extra times (FE-63). It removes every graviton of the static bulk faster than light, because the static bulk has no linear graviton at all; in exchange the bulk is strongly coupled, and its response to sources is OPEN (S-5). On an expanding fold no component of the perturbation of the bulk metric has a time-kinetic term, and some keep gradient terms proportional to A^2, with $\mathsf{A} = a_4'/H$ the expansion rate in units of H: these respond instantaneously in the bulk (an infinite local speed), so that the bulk of an expanding fold does carry modes faster than light. The polarisation along the extra times, a scalar of the four-dimensional world, is one of them: on the fold its thin part has the coefficient ratio $(8 + 5\mathsf{A}^2 + 4\mathsf{A}_1)/(8 + 6\mathsf{A}^2)$ of its second derivative along the proper length of x_3 to its second time derivative, and it couples to the jump of its σ-derivative at the same order A^2, so its propagation needs the bulk and the fold solved together and is OPEN, generically not at the speed of light (the section on the sources of the fold, below). Today $\mathsf{A} = H_4^{(0)}/H$ is below 10^{-30} for every H above 10^{-26} GeV, so the effect is negligible; in the early universe it is not.

And it cannot give gravitons at the speed of light in the bulk as well (step 2). The author's order "Gravitons must travel at c" is therefore met for the graviton of our four-dimensional world at linear order, and cannot be met in full with the author's metric. Status: DERIVED (`um3_weights.py`, W1 to W8; `um3_linear.py`, LN1 to LN5), OPEN (the strongly coupled bulk).

4 The background

The order 0 of the power series of the metric is the author's metric itself, with his two functions of the time, a_4 and b_4 (b_4 is a function of x_4 and never identically 1, by the author's order). Its exact curvature and its exact field equations in the bulk and at the fold are restated here, with the corrected weights; the solution asks which functions $a_4(x_4)$ and $b_4(x_4)$ they admit and what they require.

$$
\begin{aligned}
g_{MN} &= \mathrm{diag}\Big(\cot^2 z,\ \mathrm{e}^{2a_4}(\sin^2 z)^{1/6},\ \mathrm{e}^{2a_4}(\sin^2 z)^{1/6},\ \mathrm{e}^{2a_4}(\sin^2 z)^{1/6},\ -1,\\
&\qquad -\mathrm{e}^{-2a_4}b_4^2(\sin^2 z)^{1/6},\ -\mathrm{e}^{-2a_4}b_4^2(\sin^2 z)^{1/6},\ -\mathrm{e}^{-2a_4}b_4^2(\sin^2 z)^{1/6}\Big), \qquad z = 6Hx_0,\\
\sigma &= \mathrm{sgn}\Big(\frac{\pi}{2} - z\Big)\frac{\ln\sin z}{6H}, \qquad \mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0,\\
g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N &= \mathrm{d}\sigma^2 + \mathrm{e}^{-2H|\sigma|}\Big[\mathrm{e}^{2a_4}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{e}^{-2a_4}b_4^2\big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big)\Big] - \mathrm{d}x_4^2
\end{aligned}
\tag{FE-27}
$$

Symbols in (FE-27).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\ldots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$, the fold is $z = \pi/2$. Unit: dimensionless. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.

- $X = \{x_0, \dots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \dots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.

The author's metric (in his coordinates $X = \{x_0, \dots, x_7\}$ of the section DEFCHART of his notebook, with the factor b_4^2 of his order of 2026-10-24) and the proper chart: in the proper coordinate σ of the hidden direction one period of x_0 is the whole line, the fold $z = \pi/2$ is $\sigma = 0$, and ordinary space and the extra times are warped by $\mathrm{e}^{-H|\sigma|}$. Status: AUTHOR (the metric), DERIVED (the chart).

$$
\begin{aligned}
&\mathbf{e}^{(0)} = \mathrm{d}\sigma, \quad \mathbf{e}^{(i)} = \mathrm{e}^{a_4 - H|\sigma|}\mathrm{d}x_i, \quad \mathbf{e}^{(4)} = \mathrm{d}x_4, \quad \mathbf{e}^{(r)} = b_4\,\mathrm{e}^{-a_4 - H|\sigma|}\mathrm{d}x_r, \\
&\sqrt{|g|} = b_4^3\,\mathrm{e}^{-6H|\sigma|}, \qquad \mathsf{A} = \frac{a_4'}{H}, \qquad \mathsf{U} = \frac{b_4'}{H b_4}, \qquad \mathsf{A}_1 = \frac{\mathsf{A}'}{H}, \qquad \mathsf{U}_1 = \frac{\mathsf{U}'}{H}, \\
&R^{(0)(i)}{}_{(0)(i)} = R^{(0)(r)}{}_{(0)(r)} = -H^2, \qquad R^{(0)(4)}{}_{(0)(4)} = 0, \qquad R^{(i)(j)}{}_{(i)(j)} = (\mathsf{A}^2 - 1)H^2, \\
&R^{(i)(4)}{}_{(i)(4)} = (\mathsf{A}^2 + \mathsf{A}_1)H^2, \qquad R^{(i)(r)}{}_{(i)(r)} = -(1 + \mathsf{A}^2 - \mathsf{A}\mathsf{U})H^2, \\
&R^{(4)(r)}{}_{(4)(r)} = \big((\mathsf{U} - \mathsf{A})^2 + \mathsf{U}_1 - \mathsf{A}_1\big)H^2, \qquad R^{(r)(s)}{}_{(r)(s)} = \big((\mathsf{U} - \mathsf{A})^2 - 1\big)H^2, \\
&R^{(0)(i)}{}_{(4)(i)} = \mathrm{sgn}(\sigma)\,\mathsf{A}H^2, \qquad R^{(0)(r)}{}_{(4)(r)} = \mathrm{sgn}(\sigma)\,(\mathsf{U} - \mathsf{A})H^2
\end{aligned}
\tag{FE-28}
$$

Symbols in (FE-28).

- $\mathbf{e}_a{}^M$, $\mathbf{e}^a{}_M$: The vielbein (frame field) of the metric: $g_{MN} = \eta_{ab}\,\mathbf{e}^a{}_M\mathbf{e}^b{}_N$; $\mathbf{e}_a{}^M$ is its inverse; $\mathbf{e}_{bN} = \eta_{bc}\mathbf{e}^c{}_N$; $\mathbf{e}^{(a)}$ is the frame one-form $\mathbf{e}^a{}_M\,\mathrm{d}X^M$. Unit: dimensionless. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- x: The coordinates of the fold: x_1, x_2, x_3 of ordinary space, x_4 the time, x_5, x_6, x_7 the extra times; a field of the fold is a function of the four observable coordinates $x = (x_1, x_2, x_3, x_4)$, and $\mathrm{d}^4x = \mathrm{d}x_1\,\mathrm{d}x_2\,\mathrm{d}x_3\,\mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\dots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A}=a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U}=b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U}=\mathsf{A}$ with frozen extra times. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1=\mathsf{A}'/H=a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1=\mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4'=\mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- ${R^{MN}}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor ${R^M}_{NPQ}$ (in the convention in which the commutator of two covariant derivatives, acting on a vector, is the Riemann tensor, with its last two indices those of the two derivatives, contracted with the vector), with two indices the Ricci tensor $R_{MN}={R^P}_{MPN}$, without indices the scalar curvature $R=g^{MN}R_{MN}$; with frame indices ${R^{(a)(b)}}_{(c)(d)}$ its frame components, ${R^{(a)(b)}}_{(a)(b)}$ (no sum) the sectional curvature of the frame plane (a,b). Unit: $1/\text{length}^2$. Status: STANDARD.
- j: A second generation or mass-state index (as i). Status: STANDARD.
- s: A second extra-time direction of the fold (as r, different from it). Status: STANDARD.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.

The orthonormal frame, the volume element, the dimensionless rates A, U of the two functions of the time, and the exact frame curvature of the author's metric; the components with the extra times follow from the listed ones. The curvature does not depend on the Lovelock weights. Status: DERIVED.

$$
\begin{aligned}
\frac{\kappa\,{\mathcal{T}_{\text{bulk}}}^{(0)}{}_{(0)}}{H^2} &= -\frac{1}{5}\,\mathsf{A}(\mathsf{A}-\mathsf{U})\big(\mathsf{A}^4-2\mathsf{A}^3\mathsf{U}+10\mathsf{A}^2\mathsf{U}^2+3\mathsf{A}^2\mathsf{U}_1+6\mathsf{A}^2+12\mathsf{A}\mathsf{A}_1\mathsf{U}\\
&\qquad -9\mathsf{A}\mathsf{U}^3-9\mathsf{A}\mathsf{U}\mathsf{U}_1-6\mathsf{A}\mathsf{U}-6\mathsf{A}_1\mathsf{U}^2-6\mathsf{U}^2\big),\\
\frac{\kappa\,{\mathcal{T}_{\text{bulk}}}^{(i)}{}_{(i)}}{H^2} &= -\frac{1}{5}\,\mathsf{A}(\mathsf{A}-\mathsf{U})^2\big(\mathsf{A}^3-4\mathsf{A}^2\mathsf{U}-5\mathsf{A}\mathsf{A}_1+3\mathsf{A}\mathsf{U}^2+3\mathsf{A}\mathsf{U}_1+6\mathsf{A}+2\mathsf{A}_1\mathsf{U}+12\mathsf{U}\big),\\
\frac{\kappa\,{\mathcal{T}_{\text{bulk}}}^{(4)}{}_{(4)}}{H^2} &= \mathsf{A}^3(\mathsf{A}-\mathsf{U})^3,\\
\frac{\kappa\,{\mathcal{T}_{\text{bulk}}}^{(r)}{}_{(r)}}{H^2} &= -\frac{1}{5}\,\mathsf{A}^2(\mathsf{A}-\mathsf{U})\big(\mathsf{A}^3+\mathsf{A}^2\mathsf{U}+5\mathsf{A}\mathsf{A}_1-2\mathsf{A}\mathsf{U}^2-2\mathsf{A}\mathsf{U}_1+6\mathsf{A}-3\mathsf{A}_1\mathsf{U}-18\mathsf{U}\big),\\
\frac{\kappa\,{\mathcal{T}_{\text{bulk}}}^{(0)}{}_{(4)}}{H^2} &= \operatorname{sgn}(\sigma)\,3\mathsf{A}^2\mathsf{U}(\mathsf{A}-\mathsf{U})^2
\end{aligned}
\tag{FE-29}
$$

Symbols in (FE-29).

- κ: The eight-dimensional gravitational coupling, $\kappa=8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\text{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); ${\mathcal{T}_{\text{bulk}}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- i: A generation index ($i=1,2,3$), which also labels the three neutrino mass states; in a frame component such as ${\mathcal{S}^{(i)}}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r\in\{5,6,7\}$ (x_5,x_6,x_7); not summed (the three components are equal). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)}=w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda=-7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A}=a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.

- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

The bulk equations of the author's metric: the energy-momentum tensor $\mathcal{T}_{\mathrm{bulk}}$ that the sources of the bulk must have for the metric to solve the Einstein-Lovelock equations with the corrected weights and $\Lambda = -7H^2$, as exact polynomials in the rates. Every component carries the factor A and a power of $\mathsf{A} - \mathsf{U}$. Status: AUTHOR (the metric), DERIVED (the polynomials).

$$\begin{aligned}
\frac{\kappa\,\mathcal{S}^{(i)}{}_{(i)}}{H} &= \frac{2}{15}\big(9\mathsf{A}^4 - 24\mathsf{A}^3\mathsf{U} - 12\mathsf{A}^2\mathsf{A}_1 + 27\mathsf{A}^2\mathsf{U}^2 + 12\mathsf{A}^2\mathsf{U}_1 - 24\mathsf{A}^2 + 24\mathsf{A}\mathsf{A}_1\mathsf{U} - 6\mathsf{A}\mathsf{U}^3 - 12\mathsf{A}\mathsf{U}\mathsf{U}_1 + 48\mathsf{A}\mathsf{U} \\
&\qquad - 6\mathsf{A}_1\mathsf{U}^2 + 8\mathsf{A}_1 - 6\mathsf{U}^4 - 6\mathsf{U}^2\mathsf{U}_1 - 48\mathsf{U}^2 - 24\mathsf{U}_1 - 48\big), \\
\frac{\kappa\,\mathcal{S}^{(4)}{}_{(4)}}{H} &= -\frac{4}{5}\big(3\mathsf{A}^4 - 6\mathsf{A}^3\mathsf{U} - 4\mathsf{A}^2 + 3\mathsf{A}\mathsf{U}^3 + 4\mathsf{A}\mathsf{U} + 4\mathsf{U}^2 + 8\big), \\
\frac{\kappa\,\mathcal{S}^{(r)}{}_{(r)}}{H} &= \frac{2}{15}\big(9\mathsf{A}^4 - 12\mathsf{A}^3\mathsf{U} + 12\mathsf{A}^2\mathsf{A}_1 + 9\mathsf{A}^2\mathsf{U}^2 - 24\mathsf{A}^2 - 12\mathsf{A}\mathsf{U}^3 - 12\mathsf{A}\mathsf{U}\mathsf{U}_1 - 6\mathsf{A}_1\mathsf{U}^2 - 8\mathsf{A}_1 - 24\mathsf{U}^2 - 16\mathsf{U}_1 - 48\big)
\end{aligned} \tag{FE-30}$$

Symbols in (FE-30).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)}\mathcal{E}_{(p)}{}^a{}_b$ (FD-11, the junction conditions); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -\frac{32}{5}H\delta^a{}_b$ at the static block. Status: DERIVED.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.

The junction conditions of the author's metric: the thin source that the fold must carry for given $a_4(x_4)$ and $b_4(x_4)$, with the corrected weights. They were computed from the junction potential of FE-4 and, independently, by replacing the kink by a smooth step and integrating the full left-hand side of FE-1 across it; with the bulk flux of FE-29 they satisfy the conservation law FE-11 identically. Status: DERIVED (`um3_fold.py`, J4 to J6).

$$\begin{aligned}
&\mathsf{A}_1 = \mathsf{U}_1 = 0: \qquad \mathcal{T}_{\mathrm{bulk}}{}^a{}_b = 0 \quad \Leftrightarrow \quad \mathsf{A}(\mathsf{A}-\mathsf{U})^2(\mathsf{A}+2\mathsf{U}) = \mathsf{A}\mathsf{U}(\mathsf{A}-\mathsf{U})(2\mathsf{A}-\mathsf{U}) = \mathsf{A}\mathsf{U}^3(\mathsf{A}-\mathsf{U}) = 0 \\
&\qquad \Leftrightarrow \quad \mathsf{A} = 0 \ \text{(static ordinary space, any } \mathsf{U}) \quad \text{or} \quad \mathsf{U} = \mathsf{A} \ \text{(frozen extra times, any } \mathsf{A})
\end{aligned} \tag{S-1}$$

Symbols in (S-1).

- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.

- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- $\mathcal{T}_{\text{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); ${\mathcal{T}_{\text{bulk}}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \text{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.

The histories with constant rates. The first line is the reduced Groebner basis of the polynomial system FE-29 $= 0$ with $\mathsf{A}_1 = \mathsf{U}_1 = 0$; its solutions are two whole lines in the plane of (A, U), not isolated points: a bulk without any source has either static ordinary space (with any rate of b_4) or frozen extra times (with any expansion rate). The static block $\mathsf{A} = \mathsf{U} = 0$ lies on both lines, the expanding late universe on the second. The author's vacuum $(\mathsf{A}, \mathsf{U}) = (1, 0)$ lies on neither (FE-31). Status: DERIVED (`um3_background.py`, BG1).

$$\begin{aligned} &\kappa\,{\mathcal{T}_{\text{bulk}}}^{(4)}{}_{(4)} = \mathsf{A}^3(\mathsf{A}-\mathsf{U})^3H^2, \qquad \kappa\,{\mathcal{T}_{\text{bulk}}}^{(0)}{}_{(4)} = 3\mathsf{A}^2\mathsf{U}(\mathsf{A}-\mathsf{U})^2H^2\,\text{sgn}(\sigma), \\ &\text{no bulk source:} \quad \mathsf{A} = 0 \ \text{ or } \ \mathsf{U} = \mathsf{A} \quad \text{at every instant} \end{aligned} \tag{FE-66}$$

Symbols in (FE-66).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\text{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); ${\mathcal{T}_{\text{bulk}}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \text{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

The same holds for any histories $a_4(x_4)$, $b_4(x_4)$, not only for constant rates: the energy constraint and the flow of energy along σ contain no second time derivative, so they hold at every instant for the rates of that instant, and without a bulk source the energy constraint requires $\mathsf{A} = 0$ or $\mathsf{U} = \mathsf{A}$ at every instant; on both branches every other component of FE-29 vanishes too. Status: DERIVED (`um3_background.py`, BG1; `um3_weights.py`, W5).

$$\begin{aligned} \frac{\kappa\,{\mathcal{T}_{\text{bulk}}}^{(0)}{}_{(0)}}{H^2} &= -\frac{6}{5}\,\mathsf{A}(\mathsf{A}-\mathsf{U})\big(\mathsf{A}^2 - \mathsf{A}\mathsf{U} - \mathsf{U}^2\big) + (\text{5th order}), \\ \frac{\kappa\,{\mathcal{T}_{\text{bulk}}}^{(i)}{}_{(i)}}{H^2} &= -\frac{6}{5}\,\mathsf{A}(\mathsf{A}-\mathsf{U})^2(\mathsf{A}+2\mathsf{U}) + (\text{5th order}), \\ \frac{\kappa\,{\mathcal{T}_{\text{bulk}}}^{(r)}{}_{(r)}}{H^2} &= -\frac{6}{5}\,\mathsf{A}^2(\mathsf{A}-\mathsf{U})(\mathsf{A}-3\mathsf{U}) + (\text{5th order}), \\ \frac{\kappa\,{\mathcal{T}_{\text{bulk}}}^{(4)}{}_{(4)}}{H^2} &= \mathsf{A}^3(\mathsf{A}-\mathsf{U})^3, \qquad \frac{\kappa\,{\mathcal{T}_{\text{bulk}}}^{(0)}{}_{(4)}}{H^2} = 3\,\text{sgn}(\sigma)\,\mathsf{A}^2\mathsf{U}(\mathsf{A}-\mathsf{U})^2 \end{aligned} \tag{S-2}$$

Symbols in (S-2).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\mathrm{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); ${\mathcal{T}_{\mathrm{bulk}}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as ${\mathcal{S}^{(i)}}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

The histories off the two branches. Counting the rates A and U as small quantities of first order and A_1, U_1 of second order, the bulk source that such a history needs starts at the fourth order in the pressures, at the sixth in the energy density and at the fifth in the flux along σ (S-2, the leading terms of FE-29): for any slowly changing a_4 and b_4 the bulk is a vacuum to third order. With the author's original weights the bulk needed a source of the order H^2/κ (the stabiliser and its extension to the expanding fold), which violated the null energy condition; with the corrected weights no bulk medium is needed. Status: DERIVED (`um3_background.py`, BG2).

$$a_4 = Hx_4, \quad b_4 = \text{constant } (\mathsf{A} = 1,\ \mathsf{U} = \mathsf{A}_1 = \mathsf{U}_1 = 0): \qquad \frac{\kappa\,{\mathcal{T}_{\mathrm{bulk}}}^a{}_b}{H^2} = \mathrm{diag}\Big(-\frac{7}{5}, -\frac{7}{5}, -\frac{7}{5}, -\frac{7}{5}, 1, -\frac{7}{5}, -\frac{7}{5}, -\frac{7}{5}\Big),$$
$$\kappa\rho_{\mathrm{bulk}} = -H^2, \qquad \kappa\,{\mathcal{S}^{(i)}}_{(i)} = \kappa\,{\mathcal{S}^{(r)}}_{(r)} = -\frac{42}{5}H, \qquad \kappa\,{\mathcal{S}^{(4)}}_{(4)} = -\frac{28}{5}H \tag{FE-31}$$

Symbols in (FE-31).

- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\mathrm{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex

alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); ${\mathcal{T}_{\text{bulk}}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.

- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \text{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- ρ_{bulk}: The energy density of the bulk sources, $\rho_{\text{bulk}} = -{\mathcal{T}_{\text{bulk}}}^{(4)}{}_{(4)}$. Unit: GeV^8. Status: DERIVED.
- ${\mathcal{S}^a}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa {\mathcal{S}^a}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)} {\mathcal{E}_{(p)}}^a{}_b$ (FD-11, the junction conditions); ${\mathcal{S}^4}_4$ is minus its energy density, ${\mathcal{S}^1}_1$ and ${\mathcal{S}^5}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa {\mathcal{S}^a}_b = -\frac{32}{5} H {\delta^a}_b$ at the static block. Status: DERIVED.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as ${\mathcal{S}^{(i)}}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.

The author's vacuum. With the original weights it was an exact vacuum solution of the bulk for every constant b_4 (ordinary space inflating with the rate H, the extra times deflating), and the earlier versions of UM-3 took it as the primordial era. With the corrected weights it would need a bulk source of negative energy density, so the primordial era of UM-3 is an era of the fold on the branch of frozen extra times (S-3). Status: DERIVED (`um3_primordial.py`, PR0; `um3_background.py`, BG3).

$$\mathsf{A} = \mathsf{U} = \mathsf{A}_1 = \mathsf{U}_1 = 0: \qquad \kappa\, {\mathcal{T}_{\text{bulk}}}^a{}_b = 0, \qquad \kappa\, {\mathcal{S}^{(i)}}_{(i)} = \kappa\, {\mathcal{S}^{(4)}}_{(4)} = \kappa\, {\mathcal{S}^{(r)}}_{(r)} = -\frac{32}{5} H \tag{FE-32}$$

Symbols in (FE-32).

- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(H b_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\text{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); ${\mathcal{T}_{\text{bulk}}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \text{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- ${\mathcal{S}^a}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa {\mathcal{S}^a}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)} {\mathcal{E}_{(p)}}^a{}_b$ (FD-11, the junction conditions); ${\mathcal{S}^4}_4$ is minus its energy density, ${\mathcal{S}^1}_1$ and ${\mathcal{S}^5}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa {\mathcal{S}^a}_b = -\frac{32}{5} H {\delta^a}_b$ at the static block. Status: DERIVED.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as ${\mathcal{S}^{(i)}}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.

- r: An extra-time direction of the fold: $r \in \{5,6,7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

The static block of the late universe (a_4 and b_4 constant): a vacuum bulk with a fold of pure tension, the arrangement of Randall and Sundrum (FD-8). With the original weights the fold carried a tension plus a pressureless part, $(-61/8, -31/4, -61/8)H/\kappa$, and the bulk the stabiliser. Status: DERIVED.

$$\mathsf{U} = \mathsf{A}, \quad \mathsf{U}_1 = \mathsf{A}_1 \quad \big(\mathrm{e}^{-a_4}b_4 = \text{constant},\ \mathcal{Z} = 1\big): \qquad \kappa\,\mathcal{T}_{\text{bulk}}{}^{a}{}_{b} = 0 \ \text{ for any } a_4(x_4),$$
$$\frac{\kappa\,\mathcal{S}^{(i)}{}_{(i)}}{H} = -\frac{16}{15}\big(6 + 3\mathsf{A}^2 + 2\mathsf{A}_1\big), \qquad \frac{\kappa\,\mathcal{S}^{(4)}{}_{(4)}}{H} = -\frac{16}{5}\big(2 + \mathsf{A}^2\big), \qquad \frac{\kappa\,\mathcal{S}^{(r)}{}_{(r)}}{H} = -\frac{4}{5}\big(8 + 8\mathsf{A}^2 + \mathsf{A}^4 + 4\mathsf{A}_1 + \mathsf{A}^2\mathsf{A}_1\big),$$
$$3a_4'^2 = -\kappa_7\Big(\mathcal{S}^{(4)}{}_{(4)} + \frac{32H}{5\kappa}\Big), \qquad -\big(2a_4'' + 3a_4'^2\big) = \kappa_7\Big(\mathcal{S}^{(i)}{}_{(i)} + \frac{32H}{5\kappa}\Big), \qquad \kappa_7 = \frac{15}{16}\,\kappa H \tag{FE-33}$$

Symbols in (FE-33).

- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4) = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z} = 1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\text{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); $\mathcal{T}_{\text{bulk}}{}^{(0)}{}_{b}$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)}\mathcal{E}_{(p)}{}^a{}_b$ (FD-11, the junction conditions); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -\frac{32}{5}H\delta^a{}_b$ at the static block. Status: DERIVED.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5,6,7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.

- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- κ_7: The gravitational coupling of the fold: with the corrected Lovelock weights the thin source of the fold equals, to second order in the rates of expansion, the Einstein tensor of the seven-dimensional metric of the fold divided by $\kappa_7 = 15\kappa H/16$, and exactly so in the Friedmann equations of a fold with frozen extra times; four-dimensional gravity on the fold has $8\pi G_\mathrm{N} = \kappa_7/V_*$ (plus the induced term, if the fields of the fold induce one). Unit: GeV^{-5}. Value: $15\kappa H/16$. Status: DERIVED.

The branch with frozen extra times ($\mathrm{e}^{-a_4} b_4$ constant), on which ordinary space expands while the proper size of the extra times at the fold stays fixed. The bulk is an exact vacuum for every expansion history, and the two components of the junction along ordinary space and the time are exactly the two Friedmann equations of four-dimensional general relativity, with the coupling $\kappa_7 = 15\kappa H/16$: the late universe of UM-3 expands as in general relativity, with no correction from the bulk at any order in A. Status: DERIVED (`um3_fold.py`, J7; `um3_weights.py`, W6; `um3_stabiliser.py`, ST9).

$$\mathsf{U} = \mathsf{A}, \quad \mathsf{A}_1 = 0: \qquad 3a_4'^2 = 8\pi G_\mathrm{N}\,\rho_\mathrm{inf}, \qquad p_\mathrm{inf} = -\rho_\mathrm{inf}, \qquad q_\mathrm{hold} = -\rho_\mathrm{inf}\Big(1 + \frac{\mathsf{A}^2}{4}\Big) \tag{S-3}$$

Symbols in (S-3).

- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_N: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_\mathrm{N} = 1/M_\mathrm{Pl}^2$ with the Planck mass $M_\mathrm{Pl} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- ρ_inf: The vacuum energy density that drives an inflation of the fold: the potential energy of a field of the fold (four-dimensional normalisation), with the pressure $-\rho_\mathrm{inf}$ along ordinary space and along the extra times. Unit: GeV^4. Status: ASSUMED.
- p_inf: The pressure along ordinary space of the vacuum energy that drives an inflation of the fold. Unit: GeV^4. Value: $-\rho_\mathrm{inf}$. Status: STANDARD.
- q_hold: The pressure along the extra times that the fold fluid adds to its tension to hold the volume of the extra times fixed (four-dimensional normalisation). Unit: GeV^4. Value: $-\Delta_\mathrm{r}$ today: $-0.793\,\rho_\mathrm{c}^{(0)}$ if the baryons have the pressure $-\rho_\mathrm{b}$ along the extra times (ASSUMED), $-0.842\,\rho_\mathrm{c}^{(0)}$ if they have none, $-0.891\,\rho_\mathrm{c}^{(0)}$ for $+\rho_\mathrm{b}$. Status: DERIVED (the value), ASSUMED (that the fold fluid supplies it).

The primordial era. A vacuum energy ρ_inf of the fold (the potential energy of a field of the fold, with the pressure $-\rho_\mathrm{inf}$ along ordinary space and along the extra times) on the branch of frozen extra times expands ordinary space exponentially with the constant rate $a_4' = \mathsf{A}H$, exactly as in four-dimensional general relativity (FE-33 with $\mathsf{A}_1 = 0$), provided the fold fluid adds the pressure q_hold along the extra times that holds their volume fixed (FE-65). Such an era is an inflation of the fold, and its fields have the spectrum of four-dimensional inflation: a slowly rolling field gives the observed nearly scale-invariant spectrum, the bound $r < 0.032$ on the ratio of tensor to scalar perturbations limits its expansion rate to $H_\mathrm{inf} < \pi M_\mathrm{red}\sqrt{rA_\mathrm{s}/2} = 4.43 \times 10^{13}$ GeV (with $A_\mathrm{s} = 2.1 \times 10^{-9}$, M_red the reduced Planck mass). The price is the holding pressure of S-3: it is $-(1 + \mathsf{A}^2/4)\rho_\mathrm{inf}$ with $\mathsf{A} = H_\mathrm{inf}/H$, about $\mathsf{A}^4/8$ times the tension of the fold. If H is as small as the estimate of the maximal speed of the electron below suggests ($H \lesssim 4.6 \times 10^{-13}$ GeV), an inflation at the largest allowed rate has $\mathsf{A} \geq 9.6 \times 10^{25}$, and the fold fluid must add at least about 2×10^{51} times the vacuum energy of inflation, 10^{103} times its own tension; even after inflation A exceeds 1 above a temperature

of 574 GeV (lower for a smaller H). What could supply such a pressure is OPEN, and so is therefore whether the fold can inflate at all. Since the fold has its own Einstein-Hilbert term, the Higgs field could be the inflaton if the action had a coupling ξ_{H} of the Higgs doublet to the curvature of the fold, which UM-3 does not contain (an extension of the action): with $\xi_{\mathrm{H}} = 16195$ ($\lambda = 0.1294$) or 4502 ($\lambda = 0.01$, the running value near the scale of inflation) and 55 e-foldings, $n_{\mathrm{S}} = 0.9636$ and $r = 0.0040$, 0.3 standard deviations below Planck 2018 (0.9649 ± 0.0042) and 3.5 below ACT DR6 combined with Planck, CMB lensing and DESI (0.974 ± 0.003; ESTIMATE). With a free volume the extra times would inflate too: a vacuum energy of the fold has the same source in all seven directions, and at constant rates the junction (FE-30) then requires $(\mathsf{A}-\mathsf{U})(9\mathsf{A}^3 - 11\mathsf{A}^2\mathsf{U} - 2\mathsf{A}\mathsf{U}^2 - 16\mathsf{A} + 2\mathsf{U}^3 + 8\mathsf{U}) = 0$ and $\mathsf{A}(9\mathsf{A}^3 - 16\mathsf{A}^2\mathsf{U} + 3\mathsf{A}\mathsf{U}^2 - 16\mathsf{A} + 2\mathsf{U}^3 + 8\mathsf{U}) = 0$, which near the static block hold to second order for $\mathsf{U} = 2\mathsf{A}$ (a seven-dimensional de Sitter fold; exactly only at $(\mathsf{A},\mathsf{U}) = (\pm\frac{4}{3}, 0)$ and $(0,0)$): the volume of the extra times would then grow as e^{3N} over N e-foldings, and Newton's constant, the squares of the gauge and Yukawa couplings and the quartic Higgs coupling of the fold, which are divided by the fold factor $\mathcal{Z}$ (FE-44), would fall by e^{-3N} (6.7×10^{-79} after 60 e-foldings), so UM-3 needs the volume held fixed during inflation as well. As a record: in the author's vacuum with the original weights a field of the fold felt no Hubble friction, and its spectrum had the index $n_{\mathrm{S}} - 1 \geq 3 - 2/\pi = 2.363$ instead of the observed -0.035. Status: DERIVED (`um3_background.py`, BG3; `um3_primordial.py`, PR1 to PR4 and PR6), OPEN (what supplies the holding pressure), ESTIMATE (Higgs inflation, an extension, PR5).

$$
\begin{aligned}
&g_{1,\mathrm{eff}} = \frac{g_1}{\sqrt{\mathcal{Z}}}, \quad g_{2,\mathrm{eff}} = \frac{g_2}{\sqrt{\mathcal{Z}}}, \quad g_{3,\mathrm{eff}} = \frac{g_3}{\sqrt{\mathcal{Z}}}, \qquad \lambda_{\mathrm{eff}} = \frac{\lambda}{\mathcal{Z}}, \\
&\mathcal{Y}_{\mathrm{eff}} = \frac{\mathcal{Y}}{\sqrt{\mathcal{Z}}}, \qquad G_{\mathrm{N,eff}} = \frac{G_{\mathrm{N}}}{\mathcal{Z}}, \qquad \mathcal{Z} = \frac{V_{\mathrm{t}}\left(\mathrm{e}^{-a_4} b_4\right)^3}{V_*}
\end{aligned}
\tag{FE-44}
$$

Symbols in (FE-44).

- $g_{k,\mathrm{eff}}$: The effective (canonically normalised) gauge couplings of the fold when the fold factor is $\mathcal{Z}$, $g_{k,\mathrm{eff}} = g_k/\sqrt{\mathcal{Z}}$ ($k = 1, 2, 3$). Unit: dimensionless. Status: DERIVED.
- g_1: The $U(1)_Y$ gauge coupling (usually g'; not normalised for grand unification). Unit: dimensionless. Value: $g_1(m_Z) = 0.3574$. Status: STANDARD.
- g_2: The $SU(2)_L$ gauge coupling (usually g). Unit: dimensionless. Value: $g_2(m_Z) = 0.6517$. Status: STANDARD.
- g_3: The $SU(3)_C$ gauge coupling (the strong coupling g_s). Unit: dimensionless. Value: $g_3(m_Z) = 1.2177$ ($\alpha_s = 0.1180$). Status: STANDARD.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4) = V_{\mathrm{t}}(\mathrm{e}^{-a_4} b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z} = 1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- λ_{eff}: The effective quartic Higgs coupling, $\lambda/\mathcal{Z}$. Unit: dimensionless. Status: DERIVED.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- $\mathcal{Y}_{\mathrm{eff}}$: The effective Yukawa matrices, $\mathcal{Y}/\sqrt{\mathcal{Z}}$ (for each of $\mathcal{Y}^u, \mathcal{Y}^d, \mathcal{Y}^e, \mathcal{Y}^\nu$). Unit: dimensionless. Status: DERIVED.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^\nu$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- $G_{\mathrm{N,eff}}$: The effective Newton constant of the fold, $G_{\mathrm{N}}/\mathcal{Z}$. Unit: GeV^{-2}. Status: DERIVED.
- G_{N}: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_{\mathrm{N}} = 1/M_{\mathrm{Pl}}^2$ with the Planck mass $M_{\mathrm{Pl}} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- V_{t}: The coordinate volume of the (compact) extra times, $V_{\mathrm{t}} = \int \mathrm{d}x_5\, \mathrm{d}x_6\, \mathrm{d}x_7$. Unit: length^3. Status: ASSUMED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\, \mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4} b_4)^3$ (with

$V_{\mathrm{t}} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.

$$\frac{\mathrm{d}\ln\mathcal{Z}}{\mathrm{d}x_4} = 3H\big(\mathsf{U}-\mathsf{A}\big) = 3H\mathsf{V} \tag{S-4}$$

Symbols in (S-4).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4) = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z} = 1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.
- V: The rate of change of the scale factor $\mathrm{e}^{-a_4}b_4$ of the extra times on the fold in units of H, $\mathsf{V} = \mathsf{U} - \mathsf{A}$; zero with frozen extra times. Unit: dimensionless. Status: DERIVED.

The factor $\mathcal{Z}$ that the volume of the extra times gives the four-dimensional couplings of the fold (FE-44) changes at the rate S-4, three times the rate $\mathsf{V}H$ of the scale factor $\mathrm{e}^{-a_4}b_4$ of the extra times: zero with frozen extra times, so every coupling of the fold is then constant, as observation requires; $3H_{\mathrm{inf}}$ during an inflation with a free volume ($\mathsf{U} = 2\mathsf{A}$ to second order, from the junction FE-30); $-3H$ in the author's vacuum (a record). Status: DERIVED (`um3_background.py`, BG4).

5 The power series of the metric

The matter of the fold and dirac16complex perturb the background. The metric is written as a power series of terms of decreasing size, and each order is solved with the sources of that order. With the corrected weights the static block is a degenerate vacuum, and the series behaves differently from an ordinary perturbation series; this section states how, and why the cosmology of the fold does not depend on it.

$$\begin{aligned}
&g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N = \mathrm{d}\sigma^2 + \mathrm{e}^{2\mathfrak{a}}\big(\mathrm{d}x_1^2+\mathrm{d}x_2^2+\mathrm{d}x_3^2\big) - \mathrm{e}^{2\mathfrak{n}}\,\mathrm{d}x_4^2 - \mathrm{e}^{2\mathfrak{b}}\big(\mathrm{d}x_5^2+\mathrm{d}x_6^2+\mathrm{d}x_7^2\big),\\
&R^{(0)(i)}{}_{(0)(i)} = -\partial_\sigma^2\mathfrak{a} - (\partial_\sigma\mathfrak{a})^2, \qquad R^{(0)(4)}{}_{(0)(4)} = -\partial_\sigma^2\mathfrak{n} - (\partial_\sigma\mathfrak{n})^2, \qquad R^{(i)(j)}{}_{(i)(j)} = -(\partial_\sigma\mathfrak{a})^2 + \mathrm{e}^{-2\mathfrak{n}}(\partial_{x_4}\mathfrak{a})^2,\\
&R^{(i)(r)}{}_{(i)(r)} = -\partial_\sigma\mathfrak{a}\,\partial_\sigma\mathfrak{b} + \mathrm{e}^{-2\mathfrak{n}}\,\partial_{x_4}\mathfrak{a}\,\partial_{x_4}\mathfrak{b}, \qquad R^{(i)(4)}{}_{(i)(4)} = -\partial_\sigma\mathfrak{a}\,\partial_\sigma\mathfrak{n} + \mathrm{e}^{-2\mathfrak{n}}\big(\partial_{x_4}^2\mathfrak{a} + (\partial_{x_4}\mathfrak{a})^2 - \partial_{x_4}\mathfrak{a}\,\partial_{x_4}\mathfrak{n}\big),\\
&R^{(0)(i)}{}_{(4)(i)} = -\mathrm{e}^{-\mathfrak{n}}\big(\partial_\sigma\partial_{x_4}\mathfrak{a} + \partial_\sigma\mathfrak{a}\,\partial_{x_4}\mathfrak{a} - \partial_{x_4}\mathfrak{a}\,\partial_\sigma\mathfrak{n}\big)
\end{aligned} \tag{FE-34}$$

Symbols in (FE-34).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\ldots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.

- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- $\mathfrak{a}$, $\mathfrak{a}_1$: The logarithm of the scale factor of ordinary space in the general warped metric FE-34, a function of σ and x_4; for the author's metric $\mathfrak{a} = a_4 - H|\sigma|$; $\mathfrak{a}_n$ its term of order n in the power series FE-35 ($\mathfrak{a}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- $\mathfrak{n}$, $\mathfrak{n}_1$: The logarithm of the lapse in the general warped metric FE-34 ($g_{x_4x_4} = -\mathrm{e}^{2\mathfrak{n}}$); for the author's metric $\mathfrak{n} = 0$; $\mathfrak{n}_n$ its term of order n ($\mathfrak{n}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\mathfrak{b}$, $\mathfrak{b}_1$: The logarithm of the scale factor of the extra times in the general warped metric FE-34; for the author's metric $\mathfrak{b} = \ln b_4 - a_4 - H|\sigma|$; $\mathfrak{b}_n$ its term of order n ($\mathfrak{b}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- ${R^{MN}}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor ${R^M}_{NPQ}$ (in the convention in which the commutator of two covariant derivatives, acting on a vector, is the Riemann tensor, with its last two indices those of the two derivatives, contracted with the vector), with two indices the Ricci tensor $R_{MN} = {R^P}_{MPN}$, without indices the scalar curvature $R = g^{MN}R_{MN}$; with frame indices ${R^{(a)(b)}}_{(c)(d)}$ its frame components, ${R^{(a)(b)}}_{(a)(b)}$ (no sum) the sectional curvature of the frame plane (a, b). Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as ${\mathcal{S}^{(i)}}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- j: A second generation or mass-state index (as i). Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.

$$\mathfrak{a} = a_4 - H|\sigma| + \sum_{n\geq 1} \epsilon_{\mathrm{b}}^n\, \mathfrak{a}_n(\sigma, x_4), \qquad \mathfrak{b} = \ln b_4 - a_4 - H|\sigma| + \sum_{n\geq 1} \epsilon_{\mathrm{b}}^n\, \mathfrak{b}_n(\sigma, x_4),$$
$$\mathfrak{n} = \sum_{n\geq 1} \epsilon_{\mathrm{b}}^n\, \mathfrak{n}_n(\sigma, x_4), \qquad \hat{h}_{\mu\nu} \to \hat{h}_{\mu\nu} + \epsilon_{\mathrm{c}}\, \hat{h}^{(1)}_{\mu\nu} + \ldots \tag{FE-35}$$

Symbols in (FE-35).

- $\mathfrak{a}$, $\mathfrak{a}_1$: The logarithm of the scale factor of ordinary space in the general warped metric FE-34, a function of σ and x_4; for the author's metric $\mathfrak{a} = a_4 - H|\sigma|$; $\mathfrak{a}_n$ its term of order n in the power series FE-35 ($\mathfrak{a}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one

period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

- n: The order $n = 1, 2, \ldots$ of a term of the power series of the metric (also the exponent of the bookkeeping parameter ϵ_b); $\mathfrak{a}_n$, $\mathfrak{b}_n$, $\mathfrak{n}_n$ are the terms of order n. Status: STANDARD.
- ϵ_b: The bookkeeping parameter of the deformations of the metric along σ that sources depending on σ require (their amplitude relative to the order-0 metric of the author). Unit: dimensionless. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\mathfrak{b}$, $\mathfrak{b}_1$: The logarithm of the scale factor of the extra times in the general warped metric FE-34; for the author's metric $\mathfrak{b} = \ln b_4 - a_4 - H|\sigma|$; $\mathfrak{b}_n$ its term of order n ($\mathfrak{b}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\, \mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.
- $\mathfrak{n}$, $\mathfrak{n}_1$: The logarithm of the lapse in the general warped metric FE-34 ($g_{x_4 x_4} = -\mathrm{e}^{2\mathfrak{n}}$); for the author's metric $\mathfrak{n} = 0$; $\mathfrak{n}_n$ its term of order n ($\mathfrak{n}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $\hat{h}_{\mu\nu}$: The four-dimensional metric of the fold, $\hat{h}_{\mu\nu} = \mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the coordinates $x_1, \ldots, x_4$ (the induced metric at $\sigma = 0$ without the extra times); $\hat{h}^{\mu\nu}$ its inverse, $\hat{h}$ its determinant, $\sqrt{-\hat{h}} = \mathrm{e}^{3a_4}$; $\hat{h}^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+, +, +, -$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- ϵ_c: The bookkeeping parameter of the cosmological perturbations of the metric of the fold (the size of the CMB anisotropies). Unit: dimensionless. Value: $\sim 10^{-5}$. Status: STANDARD.

The general warped metric with the symmetries of the author's metric (FE-34) and its power series (FE-35): ϵ_b counts the deformations of the metric along σ, ϵ_c the cosmological perturbations of the fold. Status: ASSUMED (the form of the series, an ansatz), DERIVED (its symmetries and its degeneracy).

$$
\begin{aligned}
&\mathfrak{a} = -H|\sigma| + \epsilon_\mathrm{b}\, \mathfrak{a}_1(\sigma), \quad \mathfrak{b} = -H|\sigma| + \epsilon_\mathrm{b}\, \mathfrak{b}_1(\sigma), \quad \mathfrak{n} = \epsilon_\mathrm{b}\, \mathfrak{n}_1(\sigma): \qquad \kappa\, \mathcal{T}_\mathrm{bulk}{}^a{}_b = \epsilon_\mathrm{b}^2\, \mathcal{X}^a{}_b + (\text{third order}),\\
&\mathcal{X}^{(0)}{}_{(0)} = -\frac{24}{5}\left(\mathfrak{a}_1'^2 + 3\mathfrak{a}_1'\mathfrak{b}_1' + \mathfrak{b}_1'^2\right), \qquad \mathcal{X}^{(4)}{}_{(4)} = 0,\\
&\mathcal{X}^{(i)}{}_{(i)} = -\frac{4}{5H}\left(3H\mathfrak{a}_1'^2 + 18H\mathfrak{a}_1'\mathfrak{b}_1' + 9H\mathfrak{b}_1'^2 - \mathfrak{a}_1'\mathfrak{a}_1'' - 3\mathfrak{a}_1'\mathfrak{b}_1'' - 3\mathfrak{a}_1''\mathfrak{b}_1' - 3\mathfrak{b}_1'\mathfrak{b}_1''\right),\\
&\mathcal{X}^{(r)}{}_{(r)} = -\frac{4}{5H}\left(9H\mathfrak{a}_1'^2 + 18H\mathfrak{a}_1'\mathfrak{b}_1' + 3H\mathfrak{b}_1'^2 - 3\mathfrak{a}_1'\mathfrak{a}_1'' - 3\mathfrak{a}_1'\mathfrak{b}_1'' - 3\mathfrak{a}_1''\mathfrak{b}_1' - \mathfrak{b}_1'\mathfrak{b}_1''\right) \qquad (\sigma > 0)
\end{aligned}
\tag{S-5}
$$

Symbols in (S-5).

- $\mathfrak{a}$, $\mathfrak{a}_1$: The logarithm of the scale factor of ordinary space in the general warped metric FE-34, a function of σ and x_4; for the author's metric $\mathfrak{a} = a_4 - H|\sigma|$; $\mathfrak{a}_n$ its term of order n in the power series FE-35 ($\mathfrak{a}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\, \mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z) \ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- ϵ_b: The bookkeeping parameter of the deformations of the metric along σ that sources depending on σ require (their amplitude relative to the order-0 metric of the author). Unit: dimensionless. Status: STANDARD.
- $\mathfrak{b}$, $\mathfrak{b}_1$: The logarithm of the scale factor of the extra times in the general warped metric FE-34; for the author's metric $\mathfrak{b} = \ln b_4 - a_4 - H|\sigma|$; $\mathfrak{b}_n$ its term of order n ($\mathfrak{b}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $\mathfrak{n}$, $\mathfrak{n}_1$: The logarithm of the lapse in the general warped metric FE-34 ($g_{x_4 x_4} = -\mathrm{e}^{2\mathfrak{n}}$); for the author's metric $\mathfrak{n} = 0$; $\mathfrak{n}_n$ its term of order n ($\mathfrak{n}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.

- $\mathcal{T}_{\rm bulk}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); ${\mathcal{T}_{\rm bulk}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- ${\mathcal{X}^a}_b$: The second-order static bulk operator around the static block: the term of second order in $\epsilon_{\rm b}$ of the bulk equations for static deformations of the metric (frame components; the first-order term vanishes identically). Unit: GeV^2. Status: DERIVED.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as ${\mathcal{S}^{(i)}}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.

The bulk equations around the static block for static deformations ($a_4 = 0$ and $b_4 = 1$ by normalisation of the coordinates; the side $\sigma < 0$ is the mirror image). The term of first order in $\epsilon_{\rm b}$ vanishes identically, in every component and for all three functions: around the static block the bulk equations have no linear part, so a first-order deformation obeys no equation at first order. The equations start at the second order, with the operator $\mathcal{X}$ of S-5; the lapse $\mathfrak{n}_1$ does not enter it, and its time component vanishes. Status: DERIVED (`um3_series.py`, SR1 and SR2, from the exact equations of the general metric FE-34).

$$\mathcal{X}^a{}_b = 0: \qquad \frac{\mathfrak{a}_1'}{\mathfrak{b}_1'} = \frac{-3 \pm \sqrt{5}}{2}, \qquad \mathfrak{a}_1',\ \mathfrak{b}_1' \propto \mathrm{e}^{3H|\sigma|} \tag{S-6}$$

Symbols in (S-6).

- ${\mathcal{X}^a}_b$: The second-order static bulk operator around the static block: the term of second order in $\epsilon_{\rm b}$ of the bulk equations for static deformations of the metric (frame components; the first-order term vanishes identically). Unit: GeV^2. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- $\mathfrak{a}$, $\mathfrak{a}_1$: The logarithm of the scale factor of ordinary space in the general warped metric FE-34, a function of σ and x_4; for the author's metric $\mathfrak{a} = a_4 - H|\sigma|$; $\mathfrak{a}_n$ its term of order n in the power series FE-35 ($\mathfrak{a}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $\mathfrak{b}$, $\mathfrak{b}_1$: The logarithm of the scale factor of the extra times in the general warped metric FE-34; for the author's metric $\mathfrak{b} = \ln b_4 - a_4 - H|\sigma|$; $\mathfrak{b}_n$ its term of order n ($\mathfrak{b}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

The static deformations without a bulk source. The σ component of S-5 fixes the ratio of the two derivatives to one of two values (the roots of $r^2 + 3r + 1 = 0$), and then the components along ordinary space and the extra times

both require $\mathfrak{b}_1''/\mathfrak{b}_1' = 3H$: every non-trivial static deformation grows like $e^{3H|\sigma|}$ away from the fold, and no bounded static correction to the metric exists. Status: DERIVED (SR2).

$$\kappa\,\Delta\mathcal{S}^{(i)}{}_{(i)} = \frac{\epsilon_{\mathfrak{b}}^2}{5H}\big(j_{\mathfrak{a}}^2 + 6j_{\mathfrak{a}}j_{\mathfrak{b}} + 3j_{\mathfrak{b}}^2\big), \qquad \kappa\,\Delta\mathcal{S}^{(4)}{}_{(4)} = 0, \qquad \kappa\,\Delta\mathcal{S}^{(r)}{}_{(r)} = \frac{\epsilon_{\mathfrak{b}}^2}{5H}\big(3j_{\mathfrak{a}}^2 + 6j_{\mathfrak{a}}j_{\mathfrak{b}} + j_{\mathfrak{b}}^2\big) \tag{S-7}$$

Symbols in (S-7).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)}\mathcal{E}_{(p)}{}^a{}_b$ (FD-11, the junction conditions); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -\frac{32}{5}H\delta^a{}_b$ at the static block. Status: DERIVED.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- $\epsilon_{\mathfrak{b}}$: The bookkeeping parameter of the deformations of the metric along σ that sources depending on σ require (their amplitude relative to the order-0 metric of the author). Unit: dimensionless. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- $j_{\mathfrak{a}}$, $j_{\mathfrak{b}}$: The jumps at the fold of the first σ-derivatives of the first-order deformations of the metric: $j_{\mathfrak{a}} = \mathfrak{a}_1'(0^+) - \mathfrak{a}_1'(0^-)$, $j_{\mathfrak{b}} = \mathfrak{b}_1'(0^+) - \mathfrak{b}_1'(0^-)$. Unit: GeV. Status: DERIVED.

The junction conditions of the static deformations: the change $\epsilon_{\mathfrak{b}}^2\,\Delta\mathcal{S}$ of the thin source of the fold, with the jumps $j_{\mathfrak{a}}$ and $j_{\mathfrak{b}}$ of the first σ-derivatives at the fold. Like the bulk equations, the junction does not respond at first order: the coefficients of the second derivatives in the linear bulk operator, which give the thin part, vanish with it. The response starts at the second order and contains no energy density. Status: DERIVED (SR3).

The perturbations of the bulk are therefore strongly coupled: the amplitude of a deformation of the metric along σ is fixed only at the second order, and how the bulk responds to a source that depends on σ (the profile of the bound states of dirac16complex, or matter of the fold that leaves the branch of frozen extra times) is OPEN. This is not a property of the ansatz: for every perturbation of the metric, all 36 of its components with arbitrary dependence on the coordinates, the linearised bulk equations around the static block vanish identically, and on the branch of frozen extra times no component has a time-kinetic term (`um3_linear.py`, LN2 and LN4). Three consequences: the bound states χ are a source of the bulk, whose response to them starts at second order and is therefore not proportional to their density (not analytic in it), so UM-3 counts them as matter of the fold with their density integrated across the fold, which is exact only for a thin profile; the coupling of the gravitational waves of the fold to the bulk beyond linear order is not controlled; and the bulk metric has no quadratic action around the static block, so its quantum fluctuations have no propagator and no perturbative expansion there. The cosmology of the fold does not need this response: with frozen extra times the order 0 of the series is an exact solution of the bulk equations for every expansion history, and the fold obeys the Friedmann equations exactly (FE-33). The bulk scalar perturbations of the expanding background were not computed. With the original weights the linear operator was regular, with a difference mode growing like $e^{6H|\sigma|}$ and a radion mode that the earlier versions of UM-3 assumed to respond to the matter of the fold; both are gone with the corrected weights.

5.1 The fold as a seven-dimensional Einstein brane

$$\kappa\,\mathcal{S}^{a}{}_{b}+\frac{32}{5}H\,\delta^{a}{}_{b}=\frac{\kappa}{\kappa_7}\,\mathcal{G}_{\mathrm{F}}{}^{a}{}_{b}+(\text{terms of fourth order in }\mathsf{A},\mathsf{V}),\qquad \mathsf{V}=\mathsf{U}-\mathsf{A},$$
$$\frac{\mathcal{G}_{\mathrm{F}}{}^{(4)}{}_{(4)}}{H^2}=-3\big(\mathsf{A}^2+3\mathsf{A}\mathsf{V}+\mathsf{V}^2\big),\qquad \frac{\mathcal{G}_{\mathrm{F}}{}^{(i)}{}_{(i)}}{H^2}=-\big(3\mathsf{A}^2+6\mathsf{A}\mathsf{V}+6\mathsf{V}^2+2\mathsf{A}_1+3\mathsf{V}_1\big),\qquad \text{(FE-71)}$$
$$\frac{\mathcal{G}_{\mathrm{F}}{}^{(r)}{}_{(r)}}{H^2}=-\big(6\mathsf{A}^2+6\mathsf{A}\mathsf{V}+3\mathsf{V}^2+3\mathsf{A}_1+2\mathsf{V}_1\big)$$

Symbols in (FE-71).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{S}^{a}{}_{b}$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^{a}{}_{b}=\int_{0^-}^{0^+}\mathrm{d}\sigma\sum_p\alpha_{(p)}\mathcal{E}_{(p)}{}^{a}{}_{b}$ (FD-11, the junction conditions); $\mathcal{S}^{4}{}_{4}$ is minus its energy density, $\mathcal{S}^{1}{}_{1}$ and $\mathcal{S}^{5}{}_{5}$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^{a}{}_{b}=-\frac{32}{5}H\delta^{a}{}_{b}$ at the static block. Status: DERIVED.
- a: A frame (orthonormal) index running over $0,1,\ldots,7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab}=\mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a,b\in\{1,\ldots,7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)}=w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda=-7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- δ: The Kronecker delta: with two indices $\delta^{M}_{N}=1$ if $M=N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1M_2\cdots}_{N_1N_2\cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- κ_7: The gravitational coupling of the fold: with the corrected Lovelock weights the thin source of the fold equals, to second order in the rates of expansion, the Einstein tensor of the seven-dimensional metric of the fold divided by $\kappa_7=15\kappa H/16$, and exactly so in the Friedmann equations of a fold with frozen extra times; four-dimensional gravity on the fold has $8\pi G_{\mathrm{N}}=\kappa_7/V_*$ (plus the induced term, if the fields of the fold induce one). Unit: GeV^{-5}. Value: $15\kappa H/16$. Status: DERIVED.
- $\mathcal{G}_{\mathrm{F}}{}^{a}{}_{b}$: The Einstein tensor of the seven-dimensional metric of the fold (FD-2), frame components along the fold. Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A}=a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.
- V: The rate of change of the scale factor $\mathrm{e}^{-a_4}b_4$ of the extra times on the fold in units of H, $\mathsf{V}=\mathsf{U}-\mathsf{A}$; zero with frozen extra times. Unit: dimensionless. Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U}=b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U}=\mathsf{A}$ with frozen extra times. Status: DERIVED.
- i: A generation index ($i=1,2,3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1=\mathsf{A}'/H=a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- V_1: The time derivative of V in units of H, $\mathsf{V}_1=\mathsf{V}'/H$. Unit: dimensionless. Status: DERIVED.
- r: An extra-time direction of the fold: $r\in\{5,6,7\}$ (x_5,x_6,x_7); not summed (the three components are equal). Status: STANDARD.

The junction conditions of the author's metric for any $a_4(x_4)$ and $b_4(x_4)$, expanded to second order in the rates: the source of the fold minus its static tension is the Einstein tensor of the seven-dimensional metric of the fold divided by κ_7. The fold obeys Einstein's equations in seven dimensions, with the Einstein-Hilbert term that the Gauss-Bonnet and cubic terms of the bulk give the kink of the metric; Einstein gravity alone in the bulk would give the fold no such term. The remainders start at the fourth order. Status: DERIVED (`um3_series.py`, SR4).

5.2 The volume of the extra times

$$H^2\big(\mathsf{V}_1 + 3\mathsf{V}(\mathsf{A}+\mathsf{V})\big) = \frac{8\pi G_\mathrm{N}}{5}\big(\rho - 3\mathcal{P} + 2q_\mathrm{tot}\big) \tag{FE-72}$$

Symbols in (FE-72).

- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- V_1: The time derivative of V in units of H, $\mathsf{V}_1 = \mathsf{V}'/H$. Unit: dimensionless. Status: DERIVED.
- V: The rate of change of the scale factor $\mathrm{e}^{-a_4}b_4$ of the extra times on the fold in units of H, $\mathsf{V} = \mathsf{U} - \mathsf{A}$; zero with frozen extra times. Unit: dimensionless. Status: DERIVED.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_N: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_\mathrm{N} = 1/M_\mathrm{Pl}^2$ with the Planck mass $M_\mathrm{Pl} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- ρ: The total energy density of the matter of the fold (all species); $\Delta\rho$ its perturbation. Unit: GeV^4. Status: STANDARD.
- $\mathcal{P}$: The total pressure of the matter of the fold; $\Delta\mathcal{P}$ its perturbation. Unit: GeV^4. Status: STANDARD.
- q_tot: The total pressure along the extra times of the matter of the fold (the Standard Model, the bound states of dirac16complex, the portals and the fold fluid without its tension). Unit: GeV^4. Status: STANDARD.

The equation of the volume of the extra times (the radion), from FE-71. The rate V of the scale factor $\mathrm{e}^{-a_4}b_4$ of the extra times is driven by $\rho - 3\mathcal{P} + 2q_\mathrm{tot}$, which vanishes for radiation and not for dust or vacuum energy. Status: DERIVED (SR4).

$$\frac{G'_{\mathrm{N,eff}}}{G_{\mathrm{N,eff}}} = \frac{\alpha'_\mathrm{em}}{\alpha_\mathrm{em}} = -\frac{\mathcal{Z}'}{\mathcal{Z}} = -3H\mathsf{V}, \qquad H\mathsf{V} = 0.316\,H_4^{(0)} \ \text{ today, with a free volume} \tag{S-8}$$

Symbols in (S-8).

- $G_{\mathrm{N,eff}}$: The effective Newton constant of the fold, $G_\mathrm{N}/\mathcal{Z}$. Unit: GeV^{-2}. Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- α_em: The fine-structure constant. Unit: dimensionless. Value: 1/137.036. Status: STANDARD.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4) = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z} = 1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- V: The rate of change of the scale factor $\mathrm{e}^{-a_4}b_4$ of the extra times on the fold in units of H, $\mathsf{V} = \mathsf{U} - \mathsf{A}$; zero with frozen extra times. Unit: dimensionless. Status: DERIVED.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.

A free volume is excluded. Newton's constant and the fine-structure constant of the fold are divided by $\mathcal{Z}$ (FE-44), so they change at the rate $-3H\mathsf{V}$. FE-72 was integrated in the number of e-foldings $\ln a$ of the fold from the radiation era (where radiation does not drive the volume) to today, with the observed content: the dark matter χ without pressure along the extra times (a dilute gas: the on-shell Lagrangian of a free Dirac field vanishes, and the contact self-interaction of dirac16complex adds only $+(\lambda_\Psi/2)(\bar\Psi\Psi)^2$), the baryons with the pressure $-\rho_\mathrm{b}$ along them (the on-shell Lagrangian of a point particle; ASSUMED, because that of the quark and gluon fields inside a nucleon is not computed) and the vacuum energy, whose pressures along ordinary space and along the extra times are both minus its energy density. The scale factor of the extra times grows by $\mathrm{e}^{0.921}$ and today $H\mathsf{V} = 0.316H_4^{(0)} = 0.68\,a_4'$. Then:

quantity	UM-3 with a free volume	measured	ratio
$G_{\mathrm{N}}'/G_{\mathrm{N}}$ today	-6.53×10^{-11} per year	$(-5.0\pm9.6)\times10^{-15}$ per year (lunar laser ranging)	6800
$\alpha_{\mathrm{em}}'/\alpha_{\mathrm{em}}$ today	-6.53×10^{-11} per year	$(1.8\pm2.5)\times10^{-19}$ per year (optical clocks)	2.6×10^{8}
$\Delta\alpha_{\mathrm{em}}/\alpha_{\mathrm{em}}$ since the Oklo reactor (1.8 Gyr)	-0.249	$\lvert\Delta\alpha_{\mathrm{em}}/\alpha_{\mathrm{em}}\rvert<1.2\times10^{-7}$	2.1×10^{6}
G_{N} at nucleosynthesis over today	15.8	$1.01^{+0.20}_{-0.16}$	

(Lunar laser ranging: L. Biskupek, J. Müller and J.-M. Torre, Universe 7 (2021) 34; optical clocks: M. Filzinger et al., Phys. Rev. Lett. 130 (2023) 253001; nucleosynthesis: C. J. Copi, A. N. Davis and L. M. Krauss, Phys. Rev. Lett. 92 (2004) 171301.) If the baryons had no pressure along the extra times, the drift of Newton's constant today would be 7973 times the bound of lunar laser ranging. The volume of the extra times must be held fixed. Status: DERIVED (SR5), ASSUMED (the pressure of the baryons along the extra times), STANDARD (the measurements).

$$
\begin{aligned}
&H_4^2=\frac{8\pi G_{\mathrm{N}}}{3}\big(\rho_{\mathrm{SM}}+\rho_\chi+\rho_\Lambda\big),\qquad 2H_4'+3H_4^2=-8\pi G_{\mathrm{N}}\,\mathcal{P},\qquad \frac{1}{8\pi G_{\mathrm{N}}}=\frac{V_*}{\kappa_7}+\frac{1}{\kappa_{\mathrm{ind}}},\qquad \kappa_7=\frac{15}{16}\,\kappa H,\\
&\rho_\Lambda=\rho_{\mathrm{ff}}+\rho_{\mathrm{vac}}-\lambda_{\mathrm{F}},\qquad q_{\mathrm{SM}}+q_\chi+q_{\mathrm{P}}+q_{\mathrm{ff}}+\lambda_{\mathrm{F}}+\frac{\rho-3\mathcal{P}}{2}=-\frac{4}{5}\big(\mathsf{A}^4+\mathsf{A}^2\mathsf{A}_1\big)\frac{HV_*}{\kappa},\\
&q_{\mathrm{SM}}=\big\langle\mathcal{L}_{\mathrm{H}}+\mathcal{L}_{\mathrm{gauge}}+\mathcal{L}_{\mathrm{f}}+\mathcal{L}_{\mathrm{Y}}\big\rangle,\qquad q_{\mathrm{P}}=\langle\mathcal{L}_{\mathrm{P}}\rangle
\end{aligned}
\tag{FE-59}
$$

Symbols in (FE-59).

- H_4: The expansion rate (Hubble rate) of the fold, $H_4=a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)}=100\,h_{100}$ km/s/Mpc with $h_{100}=0.674$ (Planck 2018): 67.4 km/s/Mpc $=1.44\times10^{-42}$ GeV. Status: STANDARD.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4'=\mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_{N}: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_{\mathrm{N}}=1/M_{\mathrm{Pl}}^2$ with the Planck mass $M_{\mathrm{Pl}}=1.221\times10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- ρ_{SM}: The energy density of the Standard-Model particles. Unit: GeV^4. Status: STANDARD.
- ρ_χ: The energy density of the dark matter, the bound states χ of dirac16complex; $\rho_\chi^{(0)}$ its value today. Unit: GeV^4. Status: DERIVED.
- ρ_Λ: The vacuum energy density of the fold above its static tension, the dark energy of UM-3: $\rho_\Lambda=\rho_{\mathrm{ff}}+\rho_{\mathrm{vac}}-\lambda_{\mathrm{F}}$. Unit: GeV^4. Value: $0.685\,\rho_{\mathrm{c}}^{(0)}=2.5\times10^{-47}$ GeV^4 (Planck 2018). Status: DERIVED.
- $\mathcal{P}$: The total pressure of the matter of the fold; $\Delta\mathcal{P}$ its perturbation. Unit: GeV^4. Status: STANDARD.
- V_*: The physical volume of the extra times x_5,x_6,x_7 on the fold in the late universe, $V_*=V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\mathrm{t}}=\int\mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- κ_7: The gravitational coupling of the fold: with the corrected Lovelock weights the thin source of the fold equals, to second order in the rates of expansion, the Einstein tensor of the seven-dimensional metric of the fold divided by $\kappa_7=15\kappa H/16$, and exactly so in the Friedmann equations of a fold with frozen extra times; four-dimensional gravity on the fold has $8\pi G_{\mathrm{N}}=\kappa_7/V_*$ (plus the induced term, if the fields of the fold induce one). Unit: GeV^{-5}. Value: $15\kappa H/16$. Status: DERIVED.
- κ_{ind}: The coupling of the Einstein-Hilbert term $\hat{R}/(2\kappa_{\mathrm{ind}})$ that the quantum fluctuations of the fields of the fold may induce, $\kappa_{\mathrm{ind}}=8\pi G_{\mathrm{ind}}$; it adds to the gravity of the fold, $1/(8\pi G_{\mathrm{N}})=V_*/\kappa_7+1/\kappa_{\mathrm{ind}}$. Unit: GeV^{-2}. Value: not fixed by UM-3. Status: ESTIMATE.
- κ: The eight-dimensional gravitational coupling, $\kappa=8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)}=w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda=-7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

- ρ_{ff}: The energy density of the fold fluid. Unit: GeV^4. Value: $\lambda_{\mathrm{F}} + \rho_\Lambda - \rho_{\mathrm{vac}}$: together with the vacuum energy ρ_{vac} of the fields of the fold it makes up the tension λ_{F} that the static block requires plus the dark energy ρ_Λ. Status: ASSUMED (the fluid), DERIVED (the value).
- ρ_{vac}: The vacuum energy of the fields of the fold: the minimum of their potential (for the Standard Model the minimum of the Higgs potential with its loop corrections), a tension of the seven-dimensional fold. Unit: GeV^4. Status: STANDARD.
- λ_{F}: The tension that the static block requires of the fold: the total vacuum energy of the seven-dimensional fold, made of the tension of the fold fluid and the vacuum energy of the fields of the fold (both have the pressure minus their energy density along ordinary space and along the extra times), per unit volume of ordinary space of the fold (four-dimensional normalisation); the excess of that total over λ_{F} is the dark energy ρ_Λ. Unit: GeV^4. Value: $32HV_*/(5\kappa) = 6H^2/(8\pi G_{\mathrm{N}})$. Status: DERIVED.
- q_{SM}: The pressure of the Standard Model along the extra times at the fold: the mean on-shell Lagrangian density of the Standard-Model fields (the extra-time component of the energy-momentum tensor of every term of the fold is its Lagrangian, because the fields of the fold do not depend on the extra times). Unit: GeV^4. Status: DERIVED (for radiation and free fields), ASSUMED (for the baryons).
- q_χ: The pressure of the bound states of dirac16complex along the extra times, integrated across the fold (their on-shell Lagrangian, the self-interaction energy per volume; zero for a dilute gas). Unit: GeV^4. Status: DERIVED.
- q_{P}: The pressure of the portals along the extra times: their mean on-shell Lagrangian density. Unit: GeV^4. Status: DERIVED.
- q_{ff}: The pressure of the fold fluid along the extra times: minus its energy density (its tension) plus the pressure q_{hold} that holds the volume of the extra times fixed. Unit: GeV^4. Value: $-\rho_{\mathrm{ff}} + q_{\mathrm{hold}}$. Status: ASSUMED (the fluid), DERIVED (the value).
- ρ: The total energy density of the matter of the fold (all species); $\Delta\rho$ its perturbation. Unit: GeV^4. Status: STANDARD.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- $\mathcal{L}_{\mathrm{H}}$: The Lagrangian of the Higgs doublet on the fold. Unit: GeV^4. Status: AUTHOR.
- $\mathcal{L}_{\mathrm{gauge}}$: The Lagrangian of the gauge fields of $SU(3)_C \times SU(2)_L \times U(1)_Y$ on the fold. Unit: GeV^4. Status: STANDARD.
- $\mathcal{L}_{\mathrm{f}}$: The kinetic Lagrangian of the quarks and leptons on the fold. Unit: GeV^4. Status: STANDARD.
- $\mathcal{L}_{\mathrm{Y}}$: The Yukawa Lagrangian of the quarks and leptons with the Higgs doublet. Unit: GeV^4. Status: STANDARD.
- $\mathcal{L}_{\mathrm{P}}$: The Lagrangian of the portals of dirac16complex at the fold. Unit: GeV^4. Status: INVENTED.

The four-dimensional equations of the fold with the volume held fixed: the Friedmann equations of general relativity with Newton's constant $1/(8\pi G_{\mathrm{N}}) = V_*/\kappa_7 + 1/\kappa_{\mathrm{ind}}$ (exact for frozen extra times; $1/\kappa_{\mathrm{ind}}$ the optional induced term IN-9, zero if the fields of the fold induce none), the dark energy ρ_Λ, and the condition along the extra times, which the fold fluid fulfils with q_{hold} (FE-65). Status: DERIVED.

$$\Delta_{\mathrm{r}} = \frac{\rho_\chi}{2} - \frac{\rho_{\mathrm{b}}}{2} + \rho_\Lambda = (0.1325 - 0.0245 + 0.685)\,\rho_{\mathrm{c}}^{(0)} = 0.793\,\rho_{\mathrm{c}}^{(0)} \quad \text{today}, \qquad q_{\mathrm{hold}} = -\Delta_{\mathrm{r}} \tag{S-9}$$

Symbols in (S-9).

- Δ_{r}: The mismatch of the radion condition: the sum over the matter of the fold of its pressure along the extra times and $(\rho - 3\mathcal{P})/2$ (zero for radiation). Unit: GeV^4. Value: today $0.793\,\rho_{\mathrm{c}}^{(0)}$ if the baryons have the pressure $-\rho_{\mathrm{b}}$ along the extra times (ASSUMED), $0.842\,\rho_{\mathrm{c}}^{(0)}$ if they have none, $0.891\,\rho_{\mathrm{c}}^{(0)}$ for $+\rho_{\mathrm{b}}$. Status: DERIVED (the condition and the values for each choice), ASSUMED (the pressure of the baryons along the extra times).
- ρ_χ: The energy density of the dark matter, the bound states χ of dirac16complex; $\rho_\chi^{(0)}$ its value today. Unit: GeV^4. Status: DERIVED.
- ρ_{b}: The (rest-mass) energy density of the baryons: in a supernova the local density of the matter (quoted in g/cm^3), in cosmology the mean density of the baryons ($\rho_{\mathrm{b}}^{(0)}$ today). Unit: GeV^4. Status: STANDARD.

- ρ_Λ: The vacuum energy density of the fold above its static tension, the dark energy of UM-3: $\rho_\Lambda = \rho_{\rm ff} + \rho_{\rm vac} - \lambda_{\rm F}$. Unit: GeV^4. Value: $0.685\,\rho_{\rm c}^{(0)} = 2.5 \times 10^{-47}$ GeV^4 (Planck 2018). Status: DERIVED.
- $\rho_{\rm c}$: The critical density, the energy density of a spatially flat fold at the expansion rate H_4; $\rho_{\rm c}^{(0)}$ its value today. Unit: GeV^4. Value: $\rho_{\rm c}^{(0)} = 1.0537 \times 10^{-5}\,h_{100}^2$ GeV/cm^3. Status: STANDARD.
- $q_{\rm hold}$: The pressure along the extra times that the fold fluid adds to its tension to hold the volume of the extra times fixed (four-dimensional normalisation). Unit: GeV^4. Value: $-\Delta_{\rm r}$ today: $-0.793\,\rho_{\rm c}^{(0)}$ if the baryons have the pressure $-\rho_{\rm b}$ along the extra times (ASSUMED), $-0.842\,\rho_{\rm c}^{(0)}$ if they have none, $-0.891\,\rho_{\rm c}^{(0)}$ for $+\rho_{\rm b}$. Status: DERIVED (the value), ASSUMED (that the fold fluid supplies it).

What the fold fluid must supply today: the mismatch $\Delta_{\rm r}$ of the condition along the extra times (FE-65), species by species the pressure along the extra times plus half of $\rho - 3\mathcal{P}$: $\rho_\chi/2$ for the dark matter (no pressure along the extra times or along ordinary space), $-\rho_{\rm b} + \rho_{\rm b}/2$ for the baryons with the pressure $-\rho_{\rm b}$ along the extra times (ASSUMED), $-\rho_\Lambda + 2\rho_\Lambda$ for the vacuum energy, and zero for radiation; with the densities of Planck 2018 (0.265, 0.049 and 0.685 of the critical density $\rho_{\rm c}^{(0)}$) it is $0.793\rho_{\rm c}^{(0)}$, so $q_{\rm hold} = -0.793\rho_{\rm c}^{(0)}$ (the term of order A^4 of FE-65 is negligible today). If the baryons have no pressure along the extra times, $\Delta_{\rm r} = 0.842\rho_{\rm c}^{(0)}$; with the pressure $+\rho_{\rm b}$ it is $0.891\rho_{\rm c}^{(0)}$. Status: DERIVED (SR6), ASSUMED (the pressure of the baryons along the extra times).

6 The sources of the fold

The kink of the author's metric at the fold requires a thin source (FE-4; at the static block FE-32), and the gravity of the fold must carry gravitational waves with the speed of light. This section solves both with the corrected weights: the gravitational waves of the fold (at linear order, with the polarisation along the extra times, which is not luminal), the gravity that the fields of the fold may induce in addition, and the fold fluid.

6.1 The gravitational waves of the fold

$$\frac{\kappa}{2\kappa_7}\Big(\frac{\partial^2 h_{\rm gw}}{\partial x_4^2} + 3a_4'\,\frac{\partial h_{\rm gw}}{\partial x_4} - {\rm e}^{-2a_4}\,\frac{\partial^2 h_{\rm gw}}{\partial x_3^2}\Big) = \kappa\,\Pi_{\rm gw}, \qquad \kappa_7 = \frac{15}{16}\,\kappa H, \qquad c_{\rm g} = 1 \tag{FE-63}$$

Symbols in (FE-63).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- κ_7: The gravitational coupling of the fold: with the corrected Lovelock weights the thin source of the fold equals, to second order in the rates of expansion, the Einstein tensor of the seven-dimensional metric of the fold divided by $\kappa_7 = 15\kappa H/16$, and exactly so in the Friedmann equations of a fold with frozen extra times; four-dimensional gravity on the fold has $8\pi G_{\rm N} = \kappa_7/V_*$ (plus the induced term, if the fields of the fold induce one). Unit: GeV^{-5}. Value: $15\kappa H/16$. Status: DERIVED.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- $h_{\rm gw}$: The amplitude of a gravitational wave of the + polarisation travelling along x_3: the metric components along x_1 and x_2 multiplied by ${\rm e}^{h_{\rm gw}}$ and ${\rm e}^{-h_{\rm gw}}$; a function of σ, x_3 and x_4 in the bulk, and on the fold the gravitational wave of the four-dimensional world. Unit: dimensionless. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is ${\rm d}/{\rm d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold ${\rm e}^{a_4}$ is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = {\rm d}a_4/{\rm d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; ${\rm e}^X$ is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- $\Pi_{\rm gw}$: The transverse traceless stress of the matter of the fold that drives the gravitational wave $h_{\rm gw}$ (half the difference of its components along x_1 and x_2), integrated across the fold (eight-dimensional normalisation). Unit: GeV^7. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

- c_{g}: The speed of gravitational waves on the fold, in units of the speed of light. Unit: dimensionless. Value: with the corrected weights 1 at every wavelength, at linear order, for the four-dimensional graviton (the polarisations along ordinary space) on the static block and on every fold with frozen extra times; the polarisation along the extra times, a scalar of the four-dimensional world, is not luminal on an expanding fold; with the author's original weights the long-wavelength mode had $c_{\mathrm{g}}^2 = 83/82$ and shorter waves were slower than light. Status: DERIVED.

The solution. The field equations of the author's metric were linearised for a gravitational wave of the + polarisation that travels along x_3, $h_{\mathrm{gw}}(\sigma, x_3, x_4)$, on every background with frozen extra times (any $a_4(x_4)$, the static block included). With the corrected weights every coefficient of the linearised bulk equation vanishes: the bulk carries no gravitational wave at linear order. The coefficients of the second σ-derivatives are an exact differential of the first derivatives, so the equation can be integrated across the kink of the fold; its thin part is $(8/(15H))(\partial_{x_4}^2 h_{\mathrm{gw}} + 3\mathsf{A}H\,\partial_{x_4} h_{\mathrm{gw}} - \partial_{x_3}^2 h_{\mathrm{gw}})$ at an instant with $a_4 = 0$, with no term in the jump of $\partial_\sigma h_{\mathrm{gw}}$. Replacing the kink by a smooth step of width w and integrating the full equation reproduces it (deviation 9.3×10^{-4} at $w = 10^{-5}/H$, falling linearly with w). Einstein gravity alone has no such thin part: the graviton of the fold comes from the Gauss-Bonnet and cubic terms. The result is FE-63, the wave equation of four-dimensional general relativity with the coupling κ_7 of the Friedmann equations (FE-33): $c_{\mathrm{g}} = 1$ at every wavelength at linear order, and the Planck mass of the waves equals that of the expansion. The gravitational waves of the neutron-star merger GW170817 arrived from a distance of about 40 Mpc within 1.7 s of the gamma rays of its burst GRB 170817A, which bounds $-3 \times 10^{-15} \le c_{\mathrm{g}} - 1 \le 7 \times 10^{-16}$ (with the conservative distance 26 Mpc; B. P. Abbott et al., Astrophys. J. Lett. 848 (2017) L13): UM-3 satisfies it for every value of H. Three limits. The result holds at linear order; the bulk is a degenerate vacuum, so the coupling of the wave to the bulk at second order is not controlled (OPEN). The + polarisation along the extra times, a scalar of the four-dimensional world, is not luminal on an expanding fold: in the bulk it has gradient terms proportional to A^2 (the coefficient $-2\mathsf{A}^2/15$ of its second derivative along the proper length of x_3, $-\mathsf{A}^2(\mathsf{A}^2 + \mathsf{A}_1 + 2)/10$ along σ) but no time-kinetic term, so it responds instantaneously, and on the fold its thin part has the coefficient ratio $(8 + 5\mathsf{A}^2 + 4\mathsf{A}_1)/(8 + 6\mathsf{A}^2)$ (of its second derivative along the proper length of x_3 to its second time derivative) and couples to the jump of its σ-derivative with the coefficient $-\mathsf{A}^2(\mathsf{A}^2 + \mathsf{A}_1 + 2)/10$, at the same order A^2 as the deviation of the ratio from 1, so its propagation needs the bulk and the fold solved together and is OPEN, generically not at the speed of light; on this branch no component of the perturbation of the bulk metric has a time-kinetic term. On the other vacuum branch, static ordinary space with a changing b_4, the polarisation along ordinary space has the same structure: on the fold the coefficient ratio $(8 + 9\mathsf{U}^2 + 6\mathsf{U}_1)/(8 + 6\mathsf{U}^2)$ and the coupling $-\mathsf{U}^2(\mathsf{U}^2 + \mathsf{U}_1 + 2)/10$ to the jump of its σ-derivative (propagation OPEN); that branch does not expand ordinary space and is not ours. Today $\mathsf{A} = H_4^{(0)}/H$ is below 10^{-30} for every H above 10^{-26} GeV, so these effects are far below every observation; in the early universe, where the expansion rate may exceed H, they are not small. Status: DERIVED (`um3_foldsources.py`, F1 to F3; `um3_weights.py`, W7; `um3_bulk.py`, B5; `um3_linear.py`, LN4 and LN5), OPEN (the coupling to the degenerate bulk and the modes without a time-kinetic term).

$$w_{(2)} = \frac{1}{32},\ w_{(3)} = \frac{1}{1152}: \qquad c_{\mathrm{t}} = \frac{1}{16}, \qquad c_{\mathrm{x}} = \frac{5}{16}, \qquad \kappa K_{44} = \frac{41}{96H}, \qquad \kappa K_{33} = \frac{83}{192H}, \qquad c_{\mathrm{g}}^2 = \frac{83}{82} \tag{S-30}$$

Symbols in (S-30).

- $w_{(p)}$: The Lovelock weights, $w_{(1)} = 1$, $w_{(2)} = 1/20$, $w_{(3)} = 1/360$: the only weights for which both coefficients of a transverse traceless gravitational wave in the static bulk vanish, $1 - 40w_{(2)} + 360w_{(3)}$ (of its second time derivative) and $1 - 24w_{(2)} + 72w_{(3)}$ (of its second derivatives along ordinary space and along σ). Then the four-dimensional graviton of the fold travels at the speed of light at linear order, and no graviton of the static bulk travels faster, because the static bulk has no linear graviton at all (the Born-Infeld point of Lovelock gravity in eight dimensions). The author's original weights $w_{(2)} = 1/32$, $w_{(3)} = 1/1152$ (Dirac_Einstein-Lovelock_4+4_Nash.nb, In[134]) made bulk gravitons $\sqrt{5}$ times faster than light; they were corrected at the author's order of 2026-10-25. Unit: dimensionless. Value: $(1, 1/20, 1/360)$; the author's original $(1, 1/32, 1/1152)$. Status: AUTHOR (the form), ASSUMED (the criterion of the correction), DERIVED (the unique values that meet it).
- c_{t}: The coefficient of the second time derivative of a transverse traceless gravitational wave in the bulk at the static block, in units of its value with Einstein gravity alone: $c_{\mathrm{t}} = 1 - 40w_{(2)} + 360w_{(3)}$ (FE-68). Unit: dimensionless. Value: 0 with the corrected weights; 1/16 with the author's original weights. Status: DERIVED.

- c_{x}: The coefficient of the second derivatives along ordinary space and along σ of the same wave, in the same units: $c_{\mathrm{x}} = 1 - 24w_{(2)} + 72w_{(3)}$ (FE-68); bulk gravitons move with the local speed $\sqrt{c_{\mathrm{x}}/c_{\mathrm{t}}}$. Unit: dimensionless. Value: 0 with the corrected weights; 5/16 with the author's original weights. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- K_{44}: The coefficient of $\partial^2 h_{\mathrm{gw}}/\partial x_4^2$ in the equation of the long-wavelength gravitational wave (the same at every σ) at the static block, integrated across the hidden direction with the volume factor $\mathrm{e}^{-6H|\sigma|}$: the part of the bulk plus the thin part at the fold. Unit: GeV^5 (κK_{44} is a length). Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- K_{33}: The coefficient of $-\partial^2 h_{\mathrm{gw}}/\partial x_3^2$ in the same equation, with the same two parts; the wave travels at $c_{\mathrm{g}}^2 = K_{33}/K_{44}$. Unit: GeV^5. Status: DERIVED.
- c_{g}: The speed of gravitational waves on the fold, in units of the speed of light. Unit: dimensionless. Value: with the corrected weights 1 at every wavelength, at linear order, for the four-dimensional graviton (the polarisations along ordinary space) on the static block and on every fold with frozen extra times; the polarisation along the extra times, a scalar of the four-dimensional world, is not luminal on an expanding fold; with the author's original weights the long-wavelength mode had $c_{\mathrm{g}}^2 = 83/82$ and shorter waves were slower than light. Status: DERIVED.

The record of the author's original weights. With them the bulk part of the long-wavelength wave alone moved with $c_{\mathrm{g}}^2 = \frac{3}{2} \cdot 5 = 15/2$ ($c_{\mathrm{g}} = 2.74$); the fold carried 40/41 of the time-kinetic term and brought the wave to $c_{\mathrm{g}}^2 = 83/82$ ($c_{\mathrm{g}} = 1.0061$). The exact wave bound to the fold, $h_{\mathrm{gw}} \propto \mathrm{e}^{3H|\sigma|} K_\nu\big((k/H)\mathrm{e}^{H|\sigma|}\big)$ (K_ν the modified Bessel function of order ν, $\nu^2 = 9 - \omega^2/(5H^2)$, k the wavenumber and ω the frequency), was dispersive: $c_{\mathrm{g}}^2 = 1.0122$, 0.9989, 0.903 and 0.857 at $k = 10^{-3}H$, H, $10H$ and $100H$, faster than light at long and slower at short wavelengths (towards 17/20). An induced Einstein-Hilbert term with the coupling κ_{ind} would have brought the waves to $c_{\mathrm{g}}^2 = (96 + 83\epsilon)/(96 + 82\epsilon)$, $\epsilon = \kappa_{\mathrm{ind}} V_*/(\kappa H)$, and GW170817 required $\epsilon < 1.3 \times 10^{-13}$: the induced term would have had to provide all but 10^{-13} of four-dimensional gravity. Status: DERIVED (`um3_foldsources.py`, F3 and F4, records).

6.2 The gravity induced by the fields of the fold

$$\mathcal{L}_{\mathrm{IG}} = \frac{\hat{R}}{2\kappa_{\mathrm{ind}}} \tag{IN-9}$$

Symbols in (IN-9).

- $\mathcal{L}_{\mathrm{IG}}$: The Lagrangian of the Einstein-Hilbert term of the four-dimensional metric of the fold that the quantum fluctuations of the fields of the fold induce; it adds to the Einstein-Hilbert term that the Lovelock terms give the fold and does not change the speed of gravitational waves. Unit: GeV^4. Status: ESTIMATE.
- $\hat{R}$: The scalar curvature of the metric $\hat{h}_{\mu\nu}$ of the fold. Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- κ_{ind}: The coupling of the Einstein-Hilbert term $\hat{R}/(2\kappa_{\mathrm{ind}})$ that the quantum fluctuations of the fields of the fold may induce, $\kappa_{\mathrm{ind}} = 8\pi G_{\mathrm{ind}}$; it adds to the gravity of the fold, $1/(8\pi G_{\mathrm{N}}) = V_*/\kappa_7 + 1/\kappa_{\mathrm{ind}}$. Unit: GeV^{-2}. Value: not fixed by UM-3. Status: ESTIMATE.

$$\frac{1}{16\pi G_{\mathrm{ind}}} = \frac{N_{\mathrm{ind}}\,\Lambda_{\mathrm{c}}^2}{192\pi^2}, \qquad N_{\mathrm{ind}} = \sum_{\text{real scalars}} \big(1 - 6\xi\big) + N_{\mathrm{Weyl}} - 4N_{\mathrm{gauge}} = 4\big(1 - 6\xi_{\mathrm{H}}\big) \tag{FE-64}$$

Symbols in (FE-64).

- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_{ind}: The Newton constant of the induced Einstein-Hilbert term of the fold, $\kappa_{\mathrm{ind}} = 8\pi G_{\mathrm{ind}}$. Unit: GeV^{-2}. Status: ESTIMATE.
- N_{ind}: The weighted number of the fields of the fold that induce the Einstein-Hilbert term: each real scalar counts $1 - 6\xi$, each Weyl fermion 1, each gauge field -4 (with its ghost fields). Unit: dimensionless. Value: 4 for the Standard Model with three right-handed neutrinos, whose Higgs doublet has no curvature coupling in UM-3; $4(1 - 6\xi_{\mathrm{H}})$ with the curvature coupling ξ_{H} of an extension. Status: DERIVED.
- Λ_{c}: The cutoff of the proper-time integral of the one-loop effective action of the fields of the fold: the highest energy of the quantum fluctuations that induce gravity. Unit: GeV. Value: $\sqrt{24}\,\pi M_{\mathrm{red}} = 3.748 \times 10^{19}$ GeV $= 3.07 M_{\mathrm{Pl}}$ if the induced term alone gave the measured Newton constant at $\xi_{\mathrm{H}} = 0$. Status: ESTIMATE.

- ξ: The coupling of a real scalar field to the scalar curvature $\hat{R}$ of the fold: the term $-\frac{1}{2}\xi\hat{R}$ times the square of the field in its Lagrangian ($\xi = 1/6$ is the conformal coupling). Unit: dimensionless. Status: STANDARD.
- N_{Weyl}: The number of Weyl (two-component) fermion fields of the fold. Unit: dimensionless. Value: 48 (three generations of quarks and leptons with three right-handed neutrinos). Status: STANDARD.
- N_{gauge}: The number of gauge fields of the fold. Unit: dimensionless. Value: 12 (eight gluons, three W, one B). Status: STANDARD.
- ξ_{H}: The coupling of the Higgs doublet to the scalar curvature of the fold, the term $-\xi_{\text{H}}\hat{R}\,\Phi^\dagger\Phi$ in the Lagrangian of the fold: an extension of UM-3, whose Higgs Lagrangian has no such term ($\xi_{\text{H}} = 0$). Unit: dimensionless. Value: 0 in UM-3; about 1.6×10^4 if the Higgs field drove inflation (ESTIMATE, an extension of the action). Status: ESTIMATE (an extension of the action).

The solution. The one-loop effective action of every field of the fold with a proper-time cutoff Λ_{c} contains an Einstein-Hilbert term; its coefficient follows from the second heat-kernel coefficient of each field, the trace of $E + \hat{R}/6$ for its operator $-D^2 - E$ (E the non-derivative part): a real scalar with the curvature coupling ξ gives $1 - 6\xi$, a Weyl fermion 1, a gauge field with its ghost fields -4. In the Standard Model with three right-handed neutrinos the 48 Weyl fermions cancel the 12 gauge fields exactly and only the four real scalars of the Higgs doublet remain, $N_{\text{ind}} = 4(1 - 6\xi_{\text{H}})$ (without the right-handed neutrinos $1 - 24\xi_{\text{H}}$); the Higgs Lagrangian of UM-3 has no coupling to the curvature of the fold, so $\xi_{\text{H}} = 0$ and $N_{\text{ind}} = 4$. If this term alone gave Newton's constant, it would need the cutoff $\Lambda_{\text{c}} = \sqrt{24}\,\pi M_{\text{red}} = 3.748 \times 10^{19}$ GeV $= 3.07\,M_{\text{Pl}}$ at $\xi_{\text{H}} = 0$, above the Planck mass where one loop cannot be trusted, and the same loops would induce a vacuum energy of about 2×10^{77} GeV^4. With the corrected weights the fold has its own Einstein-Hilbert term (FE-63, FE-71), the induced term is an addition with the same light cone, not a requirement, and it would not restrict a curvature coupling of the Higgs doublet if one were added to the action (with which the Higgs field could drive an inflation of the fold, S-3). Status: ESTIMATE (one loop, cutoff-dependent; `um3_foldsources.py`, F4 and F5).

6.3 The fold fluid

$$\mathcal{T}_{\text{ff}}{}^{a}{}_{b} = \text{diag}\big(p_{\text{ff}}, p_{\text{ff}}, p_{\text{ff}}, -\rho_{\text{ff}}, q_{\text{ff}}, q_{\text{ff}}, q_{\text{ff}}\big) \tag{IN-10}$$

Symbols in (IN-10).

- $\mathcal{T}_{\text{ff}}{}^{a}{}_{b}$: The energy-momentum tensor of the fold fluid (frame components of the fold, four-dimensional normalisation). Unit: GeV^4. Status: ASSUMED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \text{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- p_{ff}: The pressure of the fold fluid along ordinary space. Unit: GeV^4. Value: $-\rho_{\text{ff}}$ (a tension). Status: ASSUMED (the fluid), DERIVED (the value).
- ρ_{ff}: The energy density of the fold fluid. Unit: GeV^4. Value: $\lambda_{\text{F}} + \rho_\Lambda - \rho_{\text{vac}}$: together with the vacuum energy ρ_{vac} of the fields of the fold it makes up the tension λ_{F} that the static block requires plus the dark energy ρ_Λ. Status: ASSUMED (the fluid), DERIVED (the value).
- q_{ff}: The pressure of the fold fluid along the extra times: minus its energy density (its tension) plus the pressure q_{hold} that holds the volume of the extra times fixed. Unit: GeV^4. Value: $-\rho_{\text{ff}} + q_{\text{hold}}$. Status: ASSUMED (the fluid), DERIVED (the value).

$$\begin{aligned} &p_{\text{ff}} = -\rho_{\text{ff}}, \qquad q_{\text{ff}} = -\rho_{\text{ff}} + q_{\text{hold}}, \qquad \lambda_{\text{F}} = \frac{32 H V_*}{5\kappa}, \\ &q_{\text{hold}} = -\Delta_{\text{r}} - \frac{4}{5}\big(\mathsf{A}^4 + \mathsf{A}^2\mathsf{A}_1\big)\frac{H V_*}{\kappa}, \qquad \Delta_{\text{r}} = q_{\text{SM}} + q_\chi + q_{\text{P}} - \big(\rho_{\text{ff}} - \lambda_{\text{F}}\big) + \frac{\rho - 3\mathcal{P}}{2} \end{aligned} \tag{FE-65}$$

Symbols in (FE-65).

- p_{ff}: The pressure of the fold fluid along ordinary space. Unit: GeV^4. Value: $-\rho_{\text{ff}}$ (a tension). Status: ASSUMED (the fluid), DERIVED (the value).

- ρ_{ff}: The energy density of the fold fluid. Unit: GeV^4. Value: $\lambda_{\mathrm{F}} + \rho_\Lambda - \rho_{\mathrm{vac}}$: together with the vacuum energy ρ_{vac} of the fields of the fold it makes up the tension λ_{F} that the static block requires plus the dark energy ρ_Λ. Status: ASSUMED (the fluid), DERIVED (the value).
- q_{ff}: The pressure of the fold fluid along the extra times: minus its energy density (its tension) plus the pressure q_{hold} that holds the volume of the extra times fixed. Unit: GeV^4. Value: $-\rho_{\mathrm{ff}} + q_{\mathrm{hold}}$. Status: ASSUMED (the fluid), DERIVED (the value).
- q_{hold}: The pressure along the extra times that the fold fluid adds to its tension to hold the volume of the extra times fixed (four-dimensional normalisation). Unit: GeV^4. Value: $-\Delta_{\mathrm{r}}$ today: $-0.793\,\rho_{\mathrm{c}}^{(0)}$ if the baryons have the pressure $-\rho_{\mathrm{b}}$ along the extra times (ASSUMED), $-0.842\,\rho_{\mathrm{c}}^{(0)}$ if they have none, $-0.891\,\rho_{\mathrm{c}}^{(0)}$ for $+\rho_{\mathrm{b}}$. Status: DERIVED (the value), ASSUMED (that the fold fluid supplies it).
- λ_{F}: The tension that the static block requires of the fold: the total vacuum energy of the seven-dimensional fold, made of the tension of the fold fluid and the vacuum energy of the fields of the fold (both have the pressure minus their energy density along ordinary space and along the extra times), per unit volume of ordinary space of the fold (four-dimensional normalisation); the excess of that total over λ_{F} is the dark energy ρ_Λ. Unit: GeV^4. Value: $32HV_*/(5\kappa) = 6H^2/(8\pi G_{\mathrm{N}})$. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- Δ_{r}: The mismatch of the radion condition: the sum over the matter of the fold of its pressure along the extra times and $(\rho - 3\mathcal{P})/2$ (zero for radiation). Unit: GeV^4. Value: today $0.793\,\rho_{\mathrm{c}}^{(0)}$ if the baryons have the pressure $-\rho_{\mathrm{b}}$ along the extra times (ASSUMED), $0.842\,\rho_{\mathrm{c}}^{(0)}$ if they have none, $0.891\,\rho_{\mathrm{c}}^{(0)}$ for $+\rho_{\mathrm{b}}$. Status: DERIVED (the condition and the values for each choice), ASSUMED (the pressure of the baryons along the extra times).
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant); larger than 1 whenever the expansion rate of the fold exceeds H; 1 at the author's vacuum, which is not a solution with the corrected weights. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- q_{SM}: The pressure of the Standard Model along the extra times at the fold: the mean on-shell Lagrangian density of the Standard-Model fields (the extra-time component of the energy-momentum tensor of every term of the fold is its Lagrangian, because the fields of the fold do not depend on the extra times). Unit: GeV^4. Status: DERIVED (for radiation and free fields), ASSUMED (for the baryons).
- q_χ: The pressure of the bound states of dirac16complex along the extra times, integrated across the fold (their on-shell Lagrangian, the self-interaction energy per volume; zero for a dilute gas). Unit: GeV^4. Status: DERIVED.
- q_{P}: The pressure of the portals along the extra times: their mean on-shell Lagrangian density. Unit: GeV^4. Status: DERIVED.
- ρ: The total energy density of the matter of the fold (all species); $\Delta\rho$ its perturbation. Unit: GeV^4. Status: STANDARD.
- $\mathcal{P}$: The total pressure of the matter of the fold; $\Delta\mathcal{P}$ its perturbation. Unit: GeV^4. Status: STANDARD.

The solution for the fold fluid. The source of the fold of the static block, $\kappa\mathcal{S} = -(32/5)H$ in all seven directions (FE-32), is a pure tension: its pressures are minus its energy density along ordinary space, the time and the extra times, like the tension of the brane of Randall and Sundrum. The vacuum energy of every field of the fold is a tension of this kind, because the fields do not depend on the extra times and their pressure along the extra times is their on-shell Lagrangian (FE-5), minus the potential at the minimum, the same as along ordinary space (DERIVED with the energy-momentum tensor of a field of the fold, F6). The total tension of the fold fluid and of the vacuum energy of the fields must equal $\lambda_{\mathrm{F}} = 32HV_*/(5\kappa) = 6H^2/(8\pi G_{\mathrm{N}})$ (with $8\pi G_{\mathrm{N}} = \kappa_7/V_*$); its excess over λ_{F} is the dark energy ρ_Λ (FE-59). On the frozen branch the bulk carries no flux, so the matter of the fold above its tension is conserved by itself. The component of the junction along the extra times (FE-33, third line) is the second requirement: the fold fluid must add $q_{\mathrm{hold}} = -\Delta_{\mathrm{r}}$ (S-9) to hold the volume of the extra times fixed (FE-72, S-8); in an early universe that expands faster than H the term of order A^4 dominates, and during an inflation of the fold the holding pressure is 10^{103} times the tension if H is as small as 4.6×10^{-13} GeV (S-3). With the original weights the static source had been a tension $61H/8$ plus a pressureless part $H/8$ (1/62 of the energy), fed by the flux of the stabiliser. Status:

DERIVED (`um3_foldsources.py`, F6; `um3_series.py`, SR6), ASSUMED (that the fold fluid supplies q_{hold}; the pressure of the baryons along the extra times), OPEN (its microscopic origin: it acts as a potential for the volume of the extra times, as the stabilisation of the sizes of compact extra dimensions does in other theories; its size in the early universe; and the fine tuning of λ_{F} against the vacuum energy, the cosmological-constant problem of UM-3).

7 The Higgs field in the ad hoc vacuum

$$\begin{aligned}&\frac{1}{\sqrt{-\hat{h}}\,\mathcal{Z}}\,D_\mu\big(\sqrt{-\hat{h}}\,\mathcal{Z}\,\hat{h}^{\mu\nu}D_\nu\Phi\big)-\mu^2\Phi-2\lambda\big(\Phi^\dagger\Phi\big)\Phi-c_\Phi\,\bar\Psi P_{\mathrm{L}}\Psi\,\Phi=\Theta,\\&\Theta=\begin{pmatrix}\bar d_{\mathrm{R}}(\mathcal{Y}^d)^\dagger u_{\mathrm{L}}+\bar e_{\mathrm{R}}(\mathcal{Y}^e)^\dagger\nu_{\mathrm{L}}-\bar d_{\mathrm{L}}\mathcal{Y}^u u_{\mathrm{R}}-\bar e_{\mathrm{L}}\mathcal{Y}^\nu\nu_{\mathrm{R}}\\ \bar d_{\mathrm{R}}(\mathcal{Y}^d)^\dagger d_{\mathrm{L}}+\bar e_{\mathrm{R}}(\mathcal{Y}^e)^\dagger e_{\mathrm{L}}+\bar u_{\mathrm{L}}\mathcal{Y}^u u_{\mathrm{R}}+\bar\nu_{\mathrm{L}}\mathcal{Y}^\nu\nu_{\mathrm{R}}\end{pmatrix}\end{aligned}\qquad\text{(FE-15)}$$

Symbols in (FE-15).

- $\hat h_{\mu\nu}$: The four-dimensional metric of the fold, $\hat h_{\mu\nu}=\mathrm{diag}(\mathrm{e}^{2a_4},\mathrm{e}^{2a_4},\mathrm{e}^{2a_4},-1)$ in the coordinates $x_1,\dots,x_4$ (the induced metric at $\sigma=0$ without the extra times); $\hat h^{\mu\nu}$ its inverse, $\hat h$ its determinant, $\sqrt{-\hat h}=\mathrm{e}^{3a_4}$; $\hat h^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4)=V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z}=1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M=\partial_M+\Omega_M$ (with $D_M\bar\Psi=\partial_M\bar\Psi-\bar\Psi\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu=\hat h^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A=\partial_\mu X^A+g_3 f^{ABC}G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1,x_2,x_3,x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- μ^2: The mass parameter of the Higgs potential, $\mu^2<0$ (it enters only as μ^2). Unit: GeV^2. Value: $\mu^2=-\lambda v^2=-m_h^2/2=-(88.6\ \mathrm{GeV})^2$. Status: AUTHOR.
- λ: The quartic coupling of the Higgs potential, $\lambda>0$. Unit: dimensionless. Value: $\lambda=m_h^2/(2v^2)=0.129$. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi=\Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma=0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- P_{L}: The author's left-handed chiral projector, $P_{\mathrm{L}}=(\mathbf{1}-\Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- Θ: The Yukawa source of the Higgs equation, $\Theta=-\partial\mathcal{L}_{\mathrm{Y}}/\partial\Phi^\dagger$ (a doublet of fermion bilinears). Unit: GeV^3. Status: STANDARD.
- d_{R}: The right-handed down-type quarks (down, strange and bottom; hypercharge $-1/3$). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^\nu$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- u_{L}: The upper (up-type, left-handed) components of the quark doublets. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- e_{R}: The right-handed electron, muon and tau (singlets, hypercharge -1). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- ν_{L}: The upper (neutral, left-handed) components of the lepton doublets: the left-handed neutrinos. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- d_{L}: The lower (down-type, left-handed) components of the quark doublets. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- u_{R}: The right-handed up-type quarks (up, charm and top; singlets of $SU(2)_L$, hypercharge 2/3). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- ν_{R}: The three right-handed neutrinos (gauge singlets). Unit: $\mathrm{GeV}^{3/2}$. Status: ASSUMED.
- e_{L}: The lower (charged, left-handed) components of the lepton doublets. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.

$$\frac{1}{\sqrt{-\hat{h}}\,\mathcal{Z}}\Big(\partial_\mu\big[\sqrt{-\hat{h}}\,\mathcal{Z}\,\mathcal{K}_W\,W^{I\mu\nu}\big]+g_2\,\epsilon^{IJK}W^J_\mu\,\sqrt{-\hat{h}}\,\mathcal{Z}\,\mathcal{K}_W\,W^{K\mu\nu}\Big)=-g_2\big(J^{I\nu}_{\rm w}+J^{I\nu}_{\Phi}\big),$$
$$\mathcal{K}_W=1+\sum_X c^W_X\,\bar\Psi P_X\Psi,\qquad J^{I\nu}_{\rm w}=\bar Q\,\hat\gamma^\nu\frac{\tau^I}{2}Q+\bar\ell\,\hat\gamma^\nu\frac{\tau^I}{2}\ell,\qquad J^{I\nu}_{\Phi}=\mathrm{i}\Big[(D^\nu\Phi)^\dagger\frac{\tau^I}{2}\Phi-\Phi^\dagger\frac{\tau^I}{2}D^\nu\Phi\Big] \tag{FE-17}$$

Symbols in (FE-17).

- $\hat h_{\mu\nu}$: The four-dimensional metric of the fold, $\hat h_{\mu\nu}=\mathrm{diag}(\mathrm{e}^{2a_4},\mathrm{e}^{2a_4},\mathrm{e}^{2a_4},-1)$ in the coordinates $x_1,\dots,x_4$ (the induced metric at $\sigma=0$ without the extra times); $\hat h^{\mu\nu}$ its inverse, $\hat h$ its determinant, $\sqrt{-\hat h}=\mathrm{e}^{3a_4}$; $\hat h^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4)=V_{\rm t}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z}=1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1,x_2,x_3,x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- $\mathcal{K}_G$, $\mathcal{K}_W$, $\mathcal{K}_B$: The portal factors of the gauge kinetic terms, $\mathcal{K}_G=1+\sum_X c^G_X\bar\Psi P_X\Psi$ (likewise for W and B): the gauge couplings at the fold are divided by the square root of the portal factor. Unit: dimensionless. Value: 1 where dirac16complex vanishes. Status: DERIVED.
- W^I_μ, $W^I_{\mu\nu}$: The gauge field of $SU(2)_L$ on the fold, W^1_μ,W^2_μ,W^3_μ, and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- I: A weak-isospin (adjoint SU(2)) index running over $1,2,3$; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- g_2: The $SU(2)_L$ gauge coupling (usually g). Unit: dimensionless. Value: $g_2(m_Z)=0.6517$. Status: STANDARD.
- ϵ^{IJK}: The Levi-Civita symbol of three indices (the structure constants of $SU(2)$), $\epsilon^{123}=1$. Unit: dimensionless. Status: STANDARD.
- J: A weak-isospin (adjoint SU(2)) index (as I). Status: STANDARD.
- K: A weak-isospin (adjoint SU(2)) index (as I). Status: STANDARD.
- $J^{I\nu}_{\rm w}$: The weak-isospin current of the left-handed fermion doublets. Unit: GeV^3. Status: STANDARD.
- $J^{I\nu}_{\Phi}$: The weak-isospin current of the Higgs doublet (it contains the mass terms of the W fields); $J^{2\,x_4}_{\Phi}$ its isospin charge density with $I=2$. Unit: GeV^3. Status: STANDARD.
- X: The label of a half of dirac16complex: $X\in\{\mathrm{L},\mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c^W_X: The W-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi=\Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma=0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- P_X: The chiral projector of the half X: $P_X=P_{\rm L}$ for $X=\mathrm{L}$, $P_{\rm R}$ for $X=\mathrm{R}$. Unit: dimensionless. Status: AUTHOR.
- Q, $\bar Q$: The left-handed quark doublets of the three generations, $Q=(u_{\rm L},d_{\rm L})$ (colour triplets, hypercharge $1/6$). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- $\hat\gamma^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat\gamma^{(a)}$ along the frame directions of the fold and $\hat\gamma^\mu$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat\gamma^\mu,\hat\gamma^\nu\}=-2\hat h^{\mu\nu}\mathbf{1}$; $\hat\gamma_\mu=\hat h_{\mu\nu}\hat\gamma^\nu$. Unit: dimensionless. Status: STANDARD.
- τ^I: The Pauli matrices acting on weak doublets (τ^1,τ^2,τ^3; τ^2 is the second one, not a square). Unit: dimensionless. Status: STANDARD.
- ℓ, $\bar\ell$: The left-handed lepton doublets of the three generations, $\ell=(\nu_{\rm L},e_{\rm L})$ (hypercharge $-1/2$). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M=\partial_M+\Omega_M$ (with $D_M\bar\Psi=\partial_M\bar\Psi-\bar\Psi\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu=\hat h^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A=\partial_\mu X^A+g_3 f^{ABC}G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.

- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.

The Higgs equation and the equation of the $SU(2)$ gauge fields on the fold (with the factor $\mathcal{Z}$ of the extra times and the portal factor $\mathcal{K}_W$). Status: DERIVED.

$$\begin{aligned}
&\mathcal{R}_\mathcal{A} = \mathcal{A}'' + 3\frac{b_4'}{b_4}\mathcal{A}' + \lambda\mathcal{A}^3 + c_\Phi\,\bar\Psi P_\mathrm{L}\Psi\,\mathcal{A} = \Big(4a_{40}^2 - 6a_{40}\frac{b_4'}{b_4}\Big)\mathcal{A} + \lambda\mathcal{A}^3 + c_\Phi\,\bar\Psi P_\mathrm{L}\Psi\,\mathcal{A},\\
&\mathcal{R}_v = \lambda\mathcal{A}^2 v + c_\Phi\,\bar\Psi P_\mathrm{L}\Psi\,v, \qquad J_\Phi^{2\,x_4} = a_{40}\,v\,\mathcal{A}, \qquad J_\Phi^{1\,x_4} = J_\Phi^{3\,x_4} = 0
\end{aligned} \tag{FE-26}$$

Symbols in (FE-26).

- $\mathcal{R}_\mathcal{A}$, $\mathcal{R}_v$: The residuals of the ad hoc Higgs vacuum in the upper ($\mathcal{R}_\mathcal{A}$) and lower ($\mathcal{R}_v$) components of the Higgs equation: what the ad hoc vacuum leaves unbalanced. Unit: GeV^3. Status: DERIVED.
- $\mathcal{A}$: The author's ad hoc upper component of the Higgs vacuum, almightyAdHoc, $\mathcal{A}(x_4) = v_\mathcal{A}\mathrm{e}^{-2a_{40}x_4}$. Unit: GeV. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: GeV$^{7/2}$. Status: AUTHOR.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- a_{40}: The decay constant of almightyAdHoc, $a_{40} > 0$: $\mathcal{A}$ falls by e in the time $1/(2a_{40})$. Unit: GeV (1/length). Status: AUTHOR.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_\mathrm{F})^{-1/2}$ with the Fermi constant $G_\mathrm{F} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- $J_\Phi^{I\nu}$: The weak-isospin current of the Higgs doublet (it contains the mass terms of the W fields); $J_\Phi^{2\,x_4}$ its isospin charge density with $I = 2$. Unit: GeV^3. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.

The author's ad hoc vacuum $\Phi = (\mathcal{A}, v + h)/\sqrt{2}$ with $\mathcal{A} = v_\mathcal{A}\mathrm{e}^{-2a_{40}x_4}$, inserted literally into FE-15 and FE-17 (with $h = 0$ and no gauge field): it leaves the residuals $\mathcal{R}_\mathcal{A}$ and $\mathcal{R}_v$ in the two components of the Higgs equation, and it carries a homogeneous charge density of the second isospin direction, which the Gauss law of FE-17 forbids for homogeneous gauge fields. Status: DERIVED.

$$\begin{aligned}
&\Phi = \frac{1}{\sqrt{2}}\begin{pmatrix}\mathcal{A}\\ v\end{pmatrix} = \mathcal{O}(\phi_\mathrm{rot})\,\frac{1}{\sqrt{2}}\begin{pmatrix}0\\ V_\mathrm{rad}\end{pmatrix}, \qquad V_\mathrm{rad} = \sqrt{v^2 + \mathcal{A}^2}, \qquad \tan\frac{\phi_\mathrm{rot}}{2} = \frac{\mathcal{A}}{v},\\
&\mathcal{O}(\phi_\mathrm{rot}) = \begin{pmatrix}\cos\frac{\phi_\mathrm{rot}}{2} & \sin\frac{\phi_\mathrm{rot}}{2}\\ -\sin\frac{\phi_\mathrm{rot}}{2} & \cos\frac{\phi_\mathrm{rot}}{2}\end{pmatrix}, \qquad W_{x_4}^2 = \frac{\phi_\mathrm{rot}'}{g_2} = -\frac{4a_{40}\,v\,\mathcal{A}}{g_2\big(v^2 + \mathcal{A}^2\big)}, \qquad J_\Phi^{2\,x_4} = 0,\\
&\text{FE-15} \;\Leftrightarrow\; \mathcal{O}(\phi_\mathrm{rot})\,\frac{1}{\sqrt{2}}\begin{pmatrix}0\\ -\mathcal{R}_\mathrm{rad}\end{pmatrix} = 0, \qquad \mathcal{R}_\mathrm{rad} = V_\mathrm{rad}'' + 3\,\frac{b_4'}{b_4}\,V_\mathrm{rad}' + \lambda\,\mathcal{A}^2\,V_\mathrm{rad} + c_\Phi\,\bar\Psi P_\mathrm{L}\Psi\,V_\mathrm{rad}
\end{aligned} \tag{S-10}$$

Symbols in (S-10).

- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.

- $\mathcal{A}$: The author's ad hoc upper component of the Higgs vacuum, almightyAdHoc, $\mathcal{A}(x_4) = v_\mathcal{A} \mathrm{e}^{-2a_{40}x_4}$. Unit: GeV. Status: AUTHOR.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_\mathrm{F})^{-1/2}$ with the Fermi constant $G_\mathrm{F} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- $\mathcal{O}(\phi_\mathrm{rot})$: The 2×2 isospin rotation by the angle ϕ_rot about the second isospin axis (a real element of $SU(2)$). Unit: dimensionless. Status: DERIVED.
- V_rad: The modulus of the Higgs doublet in the ad hoc vacuum, $V_\mathrm{rad} = \sqrt{v^2 + \mathcal{A}^2}$ ($V_\mathrm{rad}^2 = 2\Phi^\dagger\Phi$). Unit: GeV. Status: DERIVED.
- ϕ_rot: The angle of the isospin rotation that turns $(0, V_\mathrm{rad})/\sqrt{2}$ into the ad hoc vacuum: $\tan(\phi_\mathrm{rot}/2) = \mathcal{A}/v$. Unit: dimensionless. Status: DERIVED.
- W_μ^I, $W_{\mu\nu}^I$: The gauge field of $SU(2)_L$ on the fold, $W_\mu^1, W_\mu^2, W_\mu^3$, and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- g_2: The $SU(2)_L$ gauge coupling (usually g). Unit: dimensionless. Value: $g_2(m_Z) = 0.6517$. Status: STANDARD.
- a_{40}: The decay constant of almightyAdHoc, $a_{40} > 0$: $\mathcal{A}$ falls by e in the time $1/(2a_{40})$. Unit: GeV (1/length). Status: AUTHOR.
- $J_\Phi^{I\nu}$: The weak-isospin current of the Higgs doublet (it contains the mass terms of the W fields); $J_\Phi^{2\,x_4}$ its isospin charge density with $I = 2$. Unit: GeV^3. Status: STANDARD.
- $\mathcal{R}_\mathrm{rad}$: The radial residual of the Higgs equation FE-15 in the ad hoc vacuum written as a gauge rotation (S-10): the amount by which V_rad fails to solve the radial equation; zero for a solution. Unit: GeV^3. Status: DERIVED.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4}b_4$ on the fold and $\mathrm{e}^{-a_4}b_4\,\mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4}b_4$). Unit: dimensionless. Status: AUTHOR.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: GeV$^{7/2}$. Status: AUTHOR.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- G_F: The Fermi constant, $G_\mathrm{F} = 1/(\sqrt{2}\,v^2)$ in the Standard-Model vacuum (its measured value); in the equations of the neutrino flavours it stands for the local Fermi constant of the ad hoc vacuum. Unit: GeV^{-2}. Value: 1.1664×10^{-5} GeV^{-2}. Status: STANDARD.
- $v_\mathcal{A}$: The amplitude of almightyAdHoc at $x_4 = 0$; the author's almightyAdHoc, $\mathrm{e}^{-2a_{40}x_4}$ (written Exp[-2 a40 x4] in his notebook), has $v_\mathcal{A} = 1$ in the units of his notebook. Unit: GeV. Value: 1 in the units of the author's notebook; UM-3 keeps it free (the examples take $v_\mathcal{A} = v$ and $10v$). Status: AUTHOR (the form), ASSUMED (a free amplitude).

The ad hoc vacuum is a rotation of the doublet $(0, V_\mathrm{rad})/\sqrt{2}$ by the angle $\phi_\mathrm{rot}(x_4)$ about the second isospin axis. Together with the pure-gauge field $W_{x_4}^2 = \phi_\mathrm{rot}'/g_2$, which undoes the rotation in the covariant derivative, the isospin charge density of the Higgs field vanishes, so the Gauss law of FE-17 holds with homogeneous fields; the masses of the W and Z bosons and of the fermions, which depend on the doublet only through $\Phi^\dagger\Phi$ ($m_W^2 = g_2^2(v^2 + \mathcal{A}^2)/4$, $m_Z^2 = (g_1^2 + g_2^2)(v^2 + \mathcal{A}^2)/4$, each fermion mass $\propto \sqrt{v^2 + \mathcal{A}^2}$), are unchanged by the rotation ($v^2 + \mathcal{A}^2 = V_\mathrm{rad}^2$). The Higgs equation FE-15 then reduces exactly to one radial residual $\mathcal{R}_\mathrm{rad}$ (at 20 random points of the parameters the reduction holds to 10^{-15}). With $\mu^2 = -\lambda v^2$ the ad hoc vacuum is a solution only in the limit $\mathcal{A} \to 0$, which it reaches exponentially. Status: DERIVED (BG5).

a_{40}/v	$\mathcal{R}_\mathrm{rad}(x_4 = 0)/(\lambda v^3)$	$\rho_\mathrm{H}(x_4 = 0)/v^4$	time until $\mathcal{A} < 10^{-3}v$
0.1	1.74	0.042	$34.5/v$
1	34.2	1.03	$3.45/v$

a_{40}/v	$\mathcal{R}_{\rm rad}(x_4=0)/(\lambda v^3)$	$\rho_{\rm H}(x_4=0)/v^4$	time until $\mathcal{A}<10^{-3}v$
10	3280	100	$0.345/v$

The radial residual at $x_4=0$ and the energy density of the Higgs field, $\rho_{\rm H}=\frac{1}{2}V_{\rm rad}'^2+\frac{\lambda}{4}\mathcal{A}^4$, for $v_{\mathcal{A}}=v$, a constant b_4 and no dirac16complex at the fold, for three values of a_{40} (in units of $v=246.22$ GeV); the ad hoc component is gone after a time of order $3.5/a_{40}$. Before nucleosynthesis the Fermi constant $G_{\rm F,loc}=1/(\sqrt{2}(v^2+\mathcal{A}^2))$ must be within one per cent of its value: $\mathcal{A}^2/v^2<0.01$ at $x_4=1$ s requires $a_{40}>\ln(v_{\mathcal{A}}/(0.1v))/(2\text{ s})$, a weak bound ($a_{40}>2.3\text{ s}^{-1}=1.5\times10^{-24}$ GeV for $v_{\mathcal{A}}=10v$). Status: DERIVED (BG6, `um3_thermal.py`, T6).

8 The spectra along the hidden direction

Every field of the bulk is a superposition of modes along the hidden direction σ; their frequencies are the masses and energies that an observer on the fold sees (the "x_0 masses"). This section gives the spectra of the bulk fields and of dirac16complex on the static block (a_4, b_4 constant, normalised to $\mathrm{e}^{a_4}=b_4=1$ at the fold).

8.1 The bulk fields and why the Standard Model lives on the fold

$$u_{\rm s}''+\left(\omega^2-9H^2\right)u_{\rm s}+6H\,\delta_{\rm D}(\sigma)\,u_{\rm s}=0,\qquad u_{\rm s}=\mathrm{e}^{-3H|\sigma|}\ \ (\omega=0),\qquad \omega\geq 3H\ \ \text{(continuum)} \tag{S-11}$$

Symbols in (S-11).

- $u_{\rm s}$: The rescaled profile of a massless scalar field of the bulk at zero momentum along ordinary space: the field is $\mathrm{e}^{3H|\sigma|}u_{\rm s}(\sigma)\,\mathrm{e}^{-\mathrm{i}\omega x_4}$. Unit: as the field. Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4'=\mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto\mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)}=w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda=-7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- $\delta_{\rm D}$: The Dirac delta function of σ: zero for $\sigma\neq0$, with the integral 1 over any range that contains $\sigma=0$; the integral of $\delta_{\rm D}(\sigma)$ times a smooth function of σ is the value of that function at $\sigma=0$. Unit: 1/length (for $\delta_{\rm D}(\sigma)$). Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma=|\cot z|\,\mathrm{d}x_0$, $\sigma=\mathrm{sgn}(\pi/2-z)\ln(\sin z)/(6H)$ on one period $0<z<\pi$; $\sigma=0$ is the fold $z=\pi/2$, $\sigma\to\mp\infty$ the tips $z\to0,\pi$; one period of x_0 is the whole line $-\infty<\sigma<\infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

A massless scalar field of the bulk at zero momentum along ordinary space, with its profile written as $\mathrm{e}^{3H|\sigma|}u_{\rm s}$ (which removes the first derivative): the kink of the warp acts as an attractive delta function at the fold, which binds exactly one state, the zero mode (a constant profile, $\omega=0$), separated by the gap $3H$ from the continuum of the bulk. A gauge field of the bulk has the same structure with the gap $2H$: at zero momentum its mode equation is $(\mathrm{e}^{-4H|\sigma|}A')'+\omega^2\mathrm{e}^{-4H|\sigma|}A=0$ for a component A along ordinary space, which $A=\mathrm{e}^{2H|\sigma|}u$ turns into $u''+(\omega^2-4H^2)u+4H\,\delta_{\rm D}(\sigma)\,u=0$. A massless fermion of the bulk has no gap and no normalisable zero mode: once $\mathrm{e}^{3H|\sigma|}$ is taken out, its equation at zero momentum is that of flat space, whose solutions oscillate for every $\omega\neq0$. Status: DERIVED (`um3_bulk.py`, B1 for the scalar; the transformations for the gauge field and the fermion).

$$c_{\rm sc}^2=\left\langle\mathrm{e}^{2H|\sigma|}\right\rangle=\frac{3}{2},\qquad c_{\rm gauge}^2=2,\qquad c_{\rm fer}=\frac{2M_{\rm k}}{2M_{\rm k}-H},\qquad c_{\rm g}=1 \tag{S-12}$$

Symbols in (S-12).

- $c_{\rm sc}$: The limiting speed of the zero mode of a scalar field of the bulk along ordinary space, in units of the speed of light on the fold. Unit: dimensionless. Value: $\sqrt{3/2}$. Status: DERIVED.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- c_{gauge}: The limiting speed of the zero mode of a gauge field (a photon) of the bulk, in units of the speed of light on the fold. Unit: dimensionless. Value: $\sqrt{2}$. Status: DERIVED.
- c_{fer}: The limiting speed of the zero mode of a fermion of the bulk with the kink mass $M_{\mathrm{k}}\,\mathrm{sgn}(\sigma)$, in units of the speed of light on the fold. Unit: dimensionless. Status: DERIVED.
- M_{k}: The mass of a fermion of the bulk that changes sign at the fold (a kink mass $M_{\mathrm{k}}\,\mathrm{sgn}(\sigma)$), used to illustrate the speeds of bulk fermions. Unit: GeV. Status: ASSUMED.
- c_{g}: The speed of gravitational waves on the fold, in units of the speed of light. Unit: dimensionless. Value: with the corrected weights 1 at every wavelength, at linear order, for the four-dimensional graviton (the polarisations along ordinary space) on the static block and on every fold with frozen extra times; the polarisation along the extra times, a scalar of the four-dimensional world, is not luminal on an expanding fold; with the author's original weights the long-wavelength mode had $c_{\mathrm{g}}^2 = 83/82$ and shorter waves were slower than light. Status: DERIVED.

The limiting speeds, in units of the speed of light on the fold, of the zero modes of the bulk fields when they move along ordinary space: a zero mode samples the square root $\mathrm{e}^{-H|\sigma|}$ of the warp factor of ordinary space over its profile, so the square of its speed is the average of $\mathrm{e}^{2H|\sigma|}$ (the inverse warp factor) with the weight of its profile. A bulk scalar moves $\sqrt{3/2} = 1.22$ times faster than light on the fold, a bulk gauge field (photon) $\sqrt{2} = 1.41$ times, and a bulk fermion with the kink mass $M_{\mathrm{k}}\,\mathrm{sgn}(\sigma)$ (zero mode $\propto \mathrm{e}^{(3H - M_{\mathrm{k}})|\sigma|}$, normalisable for $M_{\mathrm{k}} > 0$) $2M_{\mathrm{k}}/(2M_{\mathrm{k}} - H)$ times (2 for $M_{\mathrm{k}} = H$, 1.0005 for $M_{\mathrm{k}} = 1000H$); the numerical eigenvalues agree with these values to 3×10^{-3} or better ($c_{\mathrm{sc}}^2 = 1.50005$ and 1.49993, $c_{\mathrm{gauge}}^2 = 1.99875$ and 1.99410 at two small momenta). Species in the bulk would therefore have different limiting speeds, which measurements exclude at the level 10^{-15} to 10^{-20} for photons, electrons and nucleons, and three neutrino flavours with three profiles would oscillate with phases proportional to the energy, which is excluded too. UM-3 therefore places the Standard Model on the fold, where every field has the one light cone of the fold (ASSUMED, motivated by these bounds). The graviton is different: with the corrected weights the bulk carries no transverse traceless graviton of ordinary space at linear order (both coefficients of its principal symbol vanish on every background with frozen extra times), and its gravitational wave is carried by the fold alone, with the wave operator of four-dimensional general relativity (FE-63): $c_{\mathrm{g}} = 1$ at linear order. The polarisation along the extra times keeps, on an expanding fold, a gradient term in the bulk without a time-kinetic term (the sources of the fold, above). With the author's original weights the local cone of the bulk graviton was $\sqrt{5}$ times the light cone, its zero mode moved $\sqrt{15/2} = 2.74$ times faster than light, and bulk and fold together gave $c_{\mathrm{g}}^2 = 83/82$ (S-30). Status: DERIVED (B1 to B6), ASSUMED (the place of the Standard Model).

8.2 dirac16complex: the continuum and the bound states

$$\Psi = \mathrm{e}^{3H|\sigma|}\,\chi(\sigma)\,\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}, \qquad \gamma^{(0)}\chi' - \mathrm{i}\omega\,\gamma^{(4)}\chi + \mathrm{i}\,\mathrm{e}^{H|\sigma|}\,\vec{k}\cdot\vec{\gamma}\,\chi - m_\Psi\chi = 0 \qquad (\sigma \neq 0) \tag{FE-36}$$

Symbols in (FE-36).

- Ψ, $\bar{\Psi}$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar{\Psi} = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{3H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of a bound state and $\bar{\chi}$

its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\vec{k}$: The (comoving) momentum along ordinary space, $\vec{k}^2$ its square. Unit: GeV. Status: STANDARD.
- $\vec{x}$: The position in ordinary space, $\vec{x} = (x_1, x_2, x_3)$. Unit: length. Status: AUTHOR.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\vec{\gamma}$: The three frame Dirac matrices of dirac16complex along ordinary space, $\vec{\gamma} = (\gamma^{(1)}, \gamma^{(2)}, \gamma^{(3)})$. Unit: dimensionless. Status: AUTHOR.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.

$$\chi' = \mathbb{K}\,\chi, \qquad \mathbb{K} = \gamma^{(0)}\big(m_\Psi + \mathrm{i}\omega\gamma^{(4)}\big), \qquad \mathbb{K}^2 = \big(m_\Psi^2 - \omega^2\big)\mathbf{1}, \qquad \chi \propto \mathrm{e}^{\pm q\sigma}, \qquad q = \sqrt{m_\Psi^2 - \omega^2} \tag{FE-37}$$

Symbols in (FE-37).

- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{3H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of a bound state and $\bar{\chi}$ its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\mathbb{K}$: The transfer matrix of the profile equation at zero momentum, $\chi' = \mathbb{K}\chi$. Unit: GeV. Status: DERIVED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- $\mathbf{1}$, $\mathbf{1}_8$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_8$ is the 8×8 identity matrix. Unit: dimensionless. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- q: The decay rate of a profile along σ at frequency ω, $q = \sqrt{m_\Psi^2 - \omega^2}$ (imaginary in the continuum). Unit: GeV. Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

$$\omega^2 = m_\Psi^2 + p_\sigma^2 + \mathrm{e}^{2H|\sigma| - 2a_4}\,\vec{k}^2 - \mathrm{e}^{2H|\sigma| + 2a_4}\,b_4^{-2}\,\vec{\kappa}^2 \tag{FE-38}$$

Symbols in (FE-38).

- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- p_σ: The momentum along the hidden direction (local wave number of a profile along σ). Unit: GeV. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line

$-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $\vec{k}$: The (comoving) momentum along ordinary space, $\vec{k}^2$ its square. Unit: GeV. Status: STANDARD.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\, \mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.
- $\vec{\kappa}$: A momentum along the extra times (three components); the fields of UM-3 have $\vec{\kappa} = 0$. Unit: GeV. Status: DERIVED.

The field equation of dirac16complex on the static block (FE-36), its solutions at zero momentum along ordinary space (FE-37): the continuum $|\omega| \geq m_\Psi$ and the exponentials below it, and the local dispersion relation (FE-38), which shows that a dependence on the extra times would make every field tachyonic. Status: DERIVED.

$$\chi(0^+) = \exp\Big(\gamma^{(0)}\big(\mu_\mathrm{L} P_\mathrm{L} + \mu_\mathrm{R} P_\mathrm{R}\big)\Big)\,\chi(0^-),$$
$$\mu_X = \frac{1}{V_*}\Big[c_\Phi\,\langle\Phi^\dagger\Phi\rangle\,\delta_{X\mathrm{L}} + \frac{1}{4}\big(c_X^G\langle G^2\rangle + c_X^W\langle W^2\rangle + c_X^B\langle B^2\rangle\big) + 2\omega\sum_f c_X^f\, n_f\Big], \qquad \langle\Phi^\dagger\Phi\rangle = \frac{v^2 + \mathcal{A}^2}{2} \tag{FE-39}$$

Symbols in (FE-39).

- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{3H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of a bound state and $\bar{\chi}$ its antiparticle. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- μ_X, μ_L, μ_R: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_\mathrm{L} P_\mathrm{L} + \mu_\mathrm{R} P_\mathrm{R})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- P_R: The author's right-handed chiral projector, $P_\mathrm{R} = (\mathbf{1} + \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4} b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4} b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta_N^M = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta_{N_1 N_2 \cdots}^{M_1 M_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta_{N_b}^{M_a}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- c_X^G: The gluon-kinetic portal coupling of the half X of dirac16complex. Unit: GeV^{-7}. Status: INVENTED.
- $\langle G^2\rangle$: The mean square of the gluon field strength, $\langle G_{\mu\nu}^A G^{A\mu\nu}\rangle$ (the gluon condensate in the QCD vacuum, a thermal value in the plasma). Unit: GeV^4. Status: STANDARD.
- G_μ^A, $G_{\mu\nu}^A$: The gluon field of $SU(3)_C$ on the fold ($A = 1, \ldots, 8$) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- c_X^W: The W-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.

- $\langle W^2\rangle$: The mean square of the $SU(2)$ field strength, $\langle W^I_{\mu\nu}W^{I\mu\nu}\rangle$. Unit: GeV^4. Status: STANDARD.
- W^I_μ, $W^I_{\mu\nu}$: The gauge field of $SU(2)_L$ on the fold, $W^1_\mu, W^2_\mu, W^3_\mu$, and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- c^B_X: The B-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- $\langle B^2\rangle$: The mean square of the hypercharge field strength, $\langle B_{\mu\nu}B^{\mu\nu}\rangle$. Unit: GeV^4. Status: STANDARD.
- B_μ, $B_{\mu\nu}$: The hypercharge gauge field of $U(1)_Y$ on the fold and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- f: The label of a kind of Standard-Model multiplet: $f \in \{Q, u, d, \ell, e, \nu\}$ (quark doublets, up-quark singlets, down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.
- c^f_X: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c^ℓ_X and c^ν_X are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.
- n_f: The number density of the Standard-Model fermions of kind f (particles minus antiparticles, summed over generations and gauge components), the time component of the mean current $\langle J^\mu_f\rangle$ in the rest frame of the medium. Unit: GeV^3. Status: STANDARD.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_\mathrm{F})^{-1/2}$ with the Fermi constant $G_\mathrm{F} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- $\mathcal{A}$: The author's ad hoc upper component of the Higgs vacuum, almightyAdHoc, $\mathcal{A}(x_4) = v_\mathcal{A}\mathrm{e}^{-2a_{40}x_4}$. Unit: GeV. Status: AUTHOR.

$$q_\mathrm{b} = -m_\Psi\left(\mu_\mathrm{L}+\mu_\mathrm{R}\right)\frac{\tanh\varsigma}{2\varsigma}, \qquad \varsigma = \sqrt{\mu_\mathrm{L}\mu_\mathrm{R}}, \qquad \omega_\mathrm{b} = \sqrt{m^2_\Psi - q^2_\mathrm{b}}, \qquad |\chi_\mathrm{b}(\sigma)| \propto \mathrm{e}^{-q_\mathrm{b}|\sigma|} \tag{FE-40}$$

Symbols in (FE-40).

- q_b: The inverse width of a state of dirac16complex bound to the fold: its profile falls as $\mathrm{e}^{-q_\mathrm{b}|\sigma|}$. Unit: GeV. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- μ_X, μ_L, μ_R: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_\mathrm{L}P_\mathrm{L} + \mu_\mathrm{R}P_\mathrm{R})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- ς: The auxiliary variable of the bound-state condition, $\varsigma = \sqrt{\mu_\mathrm{L}\mu_\mathrm{R}}$ ($\varsigma_- = \sqrt{-\mu_\mathrm{L}\mu_\mathrm{R}}$ when $\mu_\mathrm{L}\mu_\mathrm{R} < 0$). Unit: dimensionless. Status: DERIVED.
- ω_b: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- χ_b: The profile of a state of dirac16complex bound to the fold. Unit: GeV^2. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

The fold acts on dirac16complex as a chirality-dependent mass concentrated at $\sigma = 0$, with strengths μ_L and μ_R set by the mean fields of the Standard Model (FE-39); the bound states (FE-40) exist for an attractive fold. Status: DERIVED.

$(\mu_\mathrm{L}, \mu_\mathrm{R})$	$(-0.4, 0)$	$(-1, 0)$	$(-0.6, -0.3)$	$(-0.5, 0.2)$	$(-1.5, -0.5)$	$(0.4, 0)$	$(0.3, 0.2)$
ω_b/m_Ψ	0.979796	0.866025	0.905282	0.987882	0.589814	none	none

$$Q_\Psi = \int \mathrm{d}\sigma\,\mathrm{d}^3x\,\sqrt{|g|}\;\Psi^\dagger\mathfrak{B}\Psi, \qquad \mathfrak{B} = \mathrm{i}\,\sigma_{16}\,\gamma^{(4)}, \qquad \mathfrak{B}^\dagger = \mathfrak{B}, \qquad \mathfrak{B}^2 = \mathbf{1}, \qquad E_o = -\omega_o\,\mathcal{Q}_o \tag{FE-43}$$

Symbols in (FE-43).

- Q_Ψ: The conserved charge of dirac16complex (the Noether charge of a change of the phase of Ψ by a constant), a Krein form (indefinite); as written it is the charge per unit coordinate volume of the extra times, on which no field depends. Unit: GeV^3. Status: DERIVED.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- x: The coordinates of the fold: x_1, x_2, x_3 of ordinary space, x_4 the time, x_5, x_6, x_7 the extra times; a field of the fold is a function of the four observable coordinates $x = (x_1, x_2, x_3, x_4)$, and $\mathrm{d}^4x = \mathrm{d}x_1\,\mathrm{d}x_2\,\mathrm{d}x_3\,\mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\mathfrak{B}$: The Krein operator, $\mathfrak{B} = \mathrm{i}\sigma_{16}\gamma^{(4)}$: a Hermitian involution with the signature (8, 8); $\Psi^\dagger\mathfrak{B}\Psi$ is the charge density. Unit: dimensionless. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- σ_{16}: The matrix of the Dirac adjoint, $\sigma_{16} = T_0T_1T_2T_3$, real and symmetric; $\sigma_{16}T_a$ is real and antisymmetric. Unit: dimensionless. Status: AUTHOR.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- $\mathbf{1}$, $\mathbf{1}_8$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_8$ is the 8×8 identity matrix. Unit: dimensionless. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- $\mathcal{Q}_o$: The charge Q_Ψ (the Krein charge) of the single-particle state o of dirac16complex, a Kohn-Sham orbital or a mode: $+1$ or -1 for a normalised state. Unit: dimensionless. Status: DERIVED.

The conserved charge of dirac16complex (the Noether charge of a constant change of its phase) and the energy of a mode: the charge density is $\Psi^\dagger\mathfrak{B}\Psi$ with the Krein operator $\mathfrak{B}$, which is indefinite because the author's matrices have the signature (4, 4), and a mode of frequency ω_o and charge $\mathcal{Q}_o$ carries the energy $E_o = -\omega_o\mathcal{Q}_o$. Status: DERIVED.

The bound states computed with the author's exact 16×16 matrices: the roots of the 8×8 matching determinant agree with FE-40 to 10^{-7} in all seven cases, each bound level consists of eight degenerate states, and there is none for a repulsive fold. Of the eight states at $+\omega_\mathrm{b}$, four have the Krein charge $+1$ and four -1 (no null states), so four carry positive and four negative energy (FE-43). The bound-state energy $\omega_\mathrm{b} = \sqrt{m_\Psi^2 - q_\mathrm{b}^2}$ is the x_0 mass of dirac16complex: the rest mass of a particle χ that lives near the fold with the profile $\mathrm{e}^{-q_\mathrm{b}|\sigma|}$ of width $1/q_\mathrm{b}$; the more tightly it is bound, the lighter it is. Status: DERIVED (`um3_spectra.py`, S1 to S4).

$$c_\mathrm{b}^2 = \left\langle \mathrm{e}^{2H|\sigma|} \right\rangle = \frac{q_\mathrm{b}}{q_\mathrm{b} - H} \qquad (q_\mathrm{b} > H) \tag{S-13}$$

Symbols in (S-13).

- c_b: The speed coefficient of the bound particles χ of dirac16complex along ordinary space, in units of the speed of light on the fold: at small momentum their energy is $\omega^2 = \omega_\mathrm{b}^2 + c_\mathrm{b}^2\vec{k}^2$ (it is not a speed that the particles reach: at large momenta $\omega/|\vec{k}|$ tends to 1, the speed of light). Unit: dimensionless. Status: DERIVED.

- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- q_b: The inverse width of a state of dirac16complex bound to the fold: its profile falls as $\mathrm{e}^{-q_\mathrm{b}|\sigma|}$. Unit: GeV. Status: DERIVED.

The speed coefficient of the bound particles χ along ordinary space: at small momentum their energy is $\omega^2 = \omega_\mathrm{b}^2 + c_\mathrm{b}^2\vec{k}^2$, and to first order in the momentum c_b^2 is the average of the inverse warp factor $\mathrm{e}^{2H|\sigma|}$ over the density $\mathrm{e}^{-2q_\mathrm{b}|\sigma|}$ of their profile ($c_\mathrm{b}^2 = 1.25$, 2 and 5 for $H = 0.1m_\Psi$, $0.25m_\Psi$ and $0.4m_\Psi$ with $q_\mathrm{b} = m_\Psi/2$). It is not a speed that the particles reach: a slow particle χ has the group velocity $c_\mathrm{b}^2|\vec{k}|/\omega_\mathrm{b}$, as if its light cone were c_b, and at large momenta the speed approaches that of light (S-31). For $q_\mathrm{b} \le H$ the average diverges: a bound state of dirac16complex has a sensible dispersion relation only if it is bound more tightly than the warp, $q_\mathrm{b} > H$. Status: DERIVED (S5), ESTIMATE (first order in the momentum).

The computed coefficient. The dispersion of the bound states moving along x_1, computed from the profile equation at each momentum, confirms S-13 and measures its error: $c_\mathrm{b}^2 = 1.190$, 1.440 and 1.069 for $(H, q_\mathrm{b}) = (0.05, 0.3)$, $(0.1, 0.3)$ and $(0.002, 0.03)$ in units of m_Ψ, against 1.2, 1.5 and 1.071 from S-13, which therefore slightly overestimates the coefficient. Status: DERIVED (`um3_superluminal.py`, SL1).

$$\omega^2 = \omega_\mathrm{b}^2 + c_\mathrm{b}^2\,\vec{k}^2 + \ldots \quad \big(|\vec{k}| \ll m_\Psi\big), \qquad v_\mathrm{g} = \frac{\mathrm{d}\omega}{\mathrm{d}|\vec{k}|}, \qquad \frac{\omega}{|\vec{k}|} \to 1 \quad \big(|\vec{k}| \to \infty\big) \tag{S-31}$$

Symbols in (S-31).

- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- ω_b: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- c_b: The speed coefficient of the bound particles χ of dirac16complex along ordinary space, in units of the speed of light on the fold: at small momentum their energy is $\omega^2 = \omega_\mathrm{b}^2 + c_\mathrm{b}^2\vec{k}^2$ (it is not a speed that the particles reach: at large momenta $\omega/|\vec{k}|$ tends to 1, the speed of light). Unit: dimensionless. Status: DERIVED.
- $\vec{k}$: The (comoving) momentum along ordinary space, $\vec{k}^2$ its square. Unit: GeV. Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- v_g: The group velocity of a particle χ along ordinary space, $\mathrm{d}\omega/\mathrm{d}|\vec{k}|$, in units of the speed of light on the fold. Unit: dimensionless. Status: DERIVED.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.

At large momenta the ratio $\omega/|\vec{k}|$ falls towards 1 from above (1.0252 at $|\vec{k}| = 10\,m_\Psi$ and 1.0112 at $30\,m_\Psi$ for $H = 0.05$, $q_\mathrm{b} = 0.3$; $\omega - |\vec{k}|$ grows as $|\vec{k}|^{0.26}$, close to the $|\vec{k}|^{1/3}$ of a particle in a linear confining potential, which the warp provides). The group velocity v_g (central differences) is 0.99918 at $|\vec{k}| = 5.5\,m_\Psi$ and 1.00161 at $6.5\,m_\Psi$: it exceeds 1 from about $6\,m_\Psi$ on, is largest near $13\,m_\Psi$ (1.0046) and falls slowly towards 1 (1.0038 at $25\,m_\Psi$). Ultrarelativistic particles χ therefore move up to about 0.46 per cent faster than light on the fold in this regime; with the stronger warp $H = 0.1\,m_\Psi$ ($H/q_\mathrm{b} = 1/3$) the excess is about twice as large (1.0088, 1.0093 and 1.0091 at $|\vec{k}| = 7$, 9 and $11\,m_\Psi$): it grows with H/q_b. Their front velocity, the limit of $\omega/|\vec{k}|$ at infinite momentum, is the speed of light. The particles χ of the halo of the Galaxy ($|\vec{k}| = 10^{-3}\omega_\mathrm{b}$) move at 1.19×10^{-3}, and at the freeze-out of their relic density (temperature $\omega_\mathrm{b}/25$) at 0.371: the group velocity exceeds 1 only for ultrarelativistic particles χ, which existed only long before the freeze-out. Status: DERIVED (SL2, SL3).

$$\mathrm{d}\sigma^2 + \mathrm{e}^{-2H|\sigma|}\,\mathrm{d}x_1^2, \qquad \sinh\Big(\frac{H\,\Delta x_4}{2}\Big) = \frac{H\,\Delta x_1}{2} \tag{S-32}$$

Symbols in (S-32).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.

The shortcut through the bulk. A signal that leaves the fold travels in the bulk, where the warp factor shortens the distances along ordinary space. On the static block, at fixed extra times, the earliest signal between two points of the fold at the distance Δx_1 follows the shortest path of the optical metric (the first expression: the metric whose lengths are the travel times of light), which is the hyperbolic plane of radius $1/H$, with the fold as a horocycle. It arrives after the time Δx_4 of the second expression, shorter than Δx_1: $H\Delta x_4 = 0.4949$, 1.7627, 4.6249 and 9.2105 for $H\Delta x_1 = 0.5$, 2, 10 and 100 (a direct numerical minimisation of the travel time gives 0.4949, 1.7629, 4.6296 and 9.2953, upper bounds), against 0.5, 2, 10 and 100 along the fold. For large distances the time grows only with the logarithm of the distance, $\Delta x_4 \approx (2/H)\ln(H\Delta x_1)$: over one light year along the fold a signal through the bulk needs 1.5×10^{-10} s for $H = 4 \times 10^{-13}$ GeV, near the largest H that the estimate of the maximal speed of the electron below allows (and 6.9×10^{-14} s for $H = 10^{-9}$ GeV, which that estimate excludes). The fields of the Standard Model are confined to the fold and cannot use the shortcut; bulk fields (the continuum of dirac16complex) can. Causality. With four time-like directions the eight-dimensional metric has closed time-like curves through every point: the metric restricted to the plane of x_4 and one extra time, $-\mathrm{d}x_4^2 - \mathrm{e}^{-2H|\sigma|-2a_4}b_4^2\,\mathrm{d}x_5^2$, is negative definite, so every circle in this plane is a closed time-like curve (checked numerically along circles for the factor $\mathrm{e}^{-2H|\sigma|-2a_4}b_4^2$ from 10^{-6} to 7). The time x_4 is a time function only on the section of fixed extra times, where the shortcut above is the earliest signal. Through loops of the particles χ the portals would carry the Lorentz violation of the bulk to the Standard Model: with lepton portals at the strength of the relic density, a loop of χ changes the maximal speed of the electron by about $(g\,\omega_\mathrm{b}^2/(16\pi^2))^2(c_\mathrm{b}^2 - 1)$ (g the portal coupling) $= 4.9 \times 10^{-7}$, 1.9×10^{-6} and 2.2×10^{-5} times $c_\mathrm{b}^2 - 1$ for $\omega_\mathrm{b} = 1.5$, 3 and 10 TeV, with a sign that is not computed. The bounds $c_\mathrm{e} - 1 < 5 \times 10^{-21}$ (faster: no vacuum Cherenkov radiation of the 5 PeV electrons of the Crab flare of 2010) and $1 - c_\mathrm{e} < 5.6 \times 10^{-20}$ (slower: synchrotron electrons with the Lorentz factor 3×10^9) then require q_b/H above 9.7×10^{13}, 3.9×10^{14} and 4.3×10^{15} (faster) or 8.7×10^{12}, 3.5×10^{13} and 3.9×10^{14} (slower); with q_b about $0.03\,\omega_\mathrm{b}$ this means H below 5×10^{-13}, 2×10^{-13} and 7×10^{-14} GeV. Status: DERIVED (SL4), ESTIMATE (the loop, SL5).

9 The Kohn-Sham states of dirac16complex

dirac16complex is a quantised field. Its states with many particles are described here with the Kohn-Sham equations of the record (FE-41, FE-42): single-particle orbitals in the mean fields of everything else, iterated until the orbitals reproduce the fields they move in. The metric is held fixed (the back-reaction of the gas on the warp is left out, ASSUMED); the fields that are iterated are the mean field of the self-interaction and, for the Higgs loop, the strength μ_L of the fold. The state solved is a homogeneous gas of the bound particles χ at rest on the static block of the late universe (a_4 and b_4 constant), with their self-interaction, with gravity (the warp of the metric and the canonical spin connection) and with every field of the Standard Model, which acts through its mean values at the fold (the strengths μ_L and μ_R of FE-39); the Higgs portal also feeds back, because the left-handed half of the gas shifts the local electroweak vacuum v_loc of FE-49. The equations of dirac16complex itself come first.

$$\gamma^M D_M \Psi - m_\Psi \Psi - \lambda_\Psi \big(\bar\Psi \Psi\big)\Psi = 0 \qquad (\sigma \neq 0) \tag{FE-12}$$

Symbols in (FE-12).

- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar\Psi = \partial_M\bar\Psi - \bar\Psi\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat h^{\mu\nu} D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- λ_Ψ: The coupling of the contact self-interaction of dirac16complex. Unit: GeV^{-6}. Status: RECORD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

$$\begin{aligned}
&\Psi\big|_{\sigma=0^+} = \exp\Big(\frac{1}{V_*}\,\gamma^{(0)}\,\mathbb{M}\Big)\,\Psi\big|_{\sigma=0^-},\\
&\mathbb{M} = c_\Phi\big(\Phi^\dagger\Phi\big)P_\mathrm{L} + \sum_X \Big[\frac{1}{4}\big(c^G_X\,G^A_{\mu\nu}G^{A\mu\nu} + c^W_X\,W^I_{\mu\nu}W^{I\mu\nu} + c^B_X\,B_{\mu\nu}B^{\mu\nu}\big)P_X + \frac{d_X}{2}\,B_{\mu\nu}\,\mathrm{i}\Sigma^{\mu\nu}P_X\\
&\qquad + \sum_f c^f_X\big(2\mathrm{i}\,J^\mu_f P_X D_\mu + \mathrm{i}(\hat\nabla_\mu J^\mu_f)P_X\big)\Big]
\end{aligned} \tag{FE-13}$$

Symbols in (FE-13).

- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- $\mathbb{M}$: The fold operator of dirac16complex: the 16×16 matrix (with a derivative along the fold) built from the Standard-Model fields at the fold by varying the portals with respect to $\bar\Psi$ (FE-13). Unit: GeV^{-3} (so that $\mathbb{M}/V_*$ is dimensionless). Status: DERIVED.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c^G_X: The gluon-kinetic portal coupling of the half X of dirac16complex. Unit: GeV^{-7}. Status: INVENTED.
- G^A_μ, $G^A_{\mu\nu}$: The gluon field of $SU(3)_C$ on the fold ($A = 1, \ldots, 8$) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.

- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- c_X^W: The W-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- $W_\mu^I,\ W_{\mu\nu}^I$: The gauge field of $SU(2)_L$ on the fold, $W_\mu^1, W_\mu^2, W_\mu^3$, and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- c_X^B: The B-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- $B_\mu,\ B_{\mu\nu}$: The hypercharge gauge field of $U(1)_Y$ on the fold and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- P_X: The chiral projector of the half X: $P_X = P_\mathrm{L}$ for $X = \mathrm{L}$, P_R for $X = \mathrm{R}$. Unit: dimensionless. Status: AUTHOR.
- d_X: The dipole-portal coupling of the half X of dirac16complex with the hypercharge field strength. Unit: GeV^{-5}. Status: INVENTED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\Sigma^{ab},\ \Sigma^{\mu\nu}$: The generators of Spin(4,4) on dirac16complex, $\Sigma^{ab} = \frac{1}{4}[T^a, T^b]$ (the scaled commutators of the author's matrices); $\Sigma^{\mu\nu} = \mathbf{e}_a{}^\mu \mathbf{e}_b{}^\nu \Sigma^{ab}$ along the fold. Unit: dimensionless. Status: AUTHOR.
- f: The label of a kind of Standard-Model multiplet: $f \in \{Q, u, d, \ell, e, \nu\}$ (quark doublets, up-quark singlets, down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.
- c_X^f: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c_X^ℓ and c_X^ν are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.
- J_f^μ: The fermion-number current of the Standard-Model multiplets of kind f, $J_f^\mu = \bar\psi_f \hat\gamma^\mu \psi_f$ summed over the three generations and the gauge components; with other labels the currents that are the sources of the gauge fields. Unit: GeV^3. Status: STANDARD.
- $D_M,\ D_\mu$: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M \bar\Psi = \partial_M \bar\Psi - \bar\Psi \Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat h^{\mu\nu} D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G_\mu^B X^C$ (likewise for $SU(2)$ with $g_2 \epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\hat\nabla$: The covariant derivative of the fold: with the Levi-Civita connection of the metric of the fold (for four-dimensional tensors of its metric $\hat h_{\mu\nu} = \mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the observable coordinates $x_1, \ldots, x_4$; for $\hat\nabla_a \mathcal{S}^a{}_b$ of the induced metric of the seven-dimensional fold). Status: STANDARD.

$$\gamma^M \Omega_M = -3H \,\mathrm{sgn}(\sigma)\, \gamma^{(0)} + \frac{3}{2}\frac{b_4'}{b_4}\gamma^{(4)} \neq 0 \tag{FE-14}$$

Symbols in (FE-14).

- $\gamma^M,\ \gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Ω_M: The spin connection acting on dirac16complex, $\Omega_M = \frac{1}{2}\varpi_{Mab}\Sigma^{ab}$. Unit: 1/length. Status: RECORD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- sgn: The sign function: $+1$ for a positive argument and -1 for a negative one. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor (the factor of the lengths) of the extra times is $\mathrm{e}^{-a_4} b_4$ on the fold and $\mathrm{e}^{-a_4} b_4\, \mathrm{e}^{-H|\sigma|}$ at the distance σ from it (the square root of the warp factor times $\mathrm{e}^{-a_4} b_4$). Unit: dimensionless. Status: AUTHOR.

- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.

The field equation of dirac16complex in the bulk (FE-12, with its contact self-interaction), the matching condition at the fold (FE-13: the values on the two sides of the fold differ by the exponential of $\gamma^{(0)}\mathbb{M}/V_*$, where the fold operator $\mathbb{M}$ is built from the Standard-Model fields through the portals) and the spin connection of the author's metric (FE-14), which couples dirac16complex to gravity and never vanishes: its first term comes from the warp of the six directions of ordinary space and of the extra times, its second from the time dependence of b_4. Status: RECORD (FE-12), DERIVED (FE-13, FE-14).

$$\begin{aligned} &\big[\gamma^M D_M - m_\Psi - \lambda_\Psi\, n_\mathrm{s} + \lambda_\Psi\, \mathcal{N}\big]\varphi_o = \delta_\mathrm{D}(\sigma)\,\frac{1}{V_*}\,\langle\mathbb{M}\rangle\,\varphi_o, \\ &\mathcal{N} = \sum_o \mathsf{n}_o\,\varphi_o\bar{\varphi}_o, \qquad n_\mathrm{s} = \mathrm{tr}\,\mathcal{N} = \sum_o \mathsf{n}_o\,\bar{\varphi}_o\varphi_o, \qquad \mathsf{n}_o = \frac{1}{\mathrm{e}^{(E_o-\mu_\Psi)/\mathsf{T}_\Psi}+1} \end{aligned} \tag{FE-41}$$

Symbols in (FE-41).

- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC}G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- λ_Ψ: The coupling of the contact self-interaction of dirac16complex. Unit: GeV^{-6}. Status: RECORD.
- n_s: The scalar density of dirac16complex in the Kohn-Sham state, $n_\mathrm{s} = \langle\bar{\Psi}\Psi\rangle = \mathrm{tr}\,\mathcal{N}$. Unit: GeV^7. Status: DERIVED.
- $\mathcal{N}$: The one-body density matrix of dirac16complex, $\mathcal{N} = \sum_o \mathsf{n}_o\varphi_o\bar{\varphi}_o$ (a 16×16 matrix at each point). Unit: GeV^7. Status: DERIVED.
- φ_o, $\bar{\varphi}_o$: A Kohn-Sham orbital of dirac16complex (a single-particle solution of the Kohn-Sham equation: the Dirac equation of dirac16complex with the mean field of the gas added to its mass) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- δ_D: The Dirac delta function of σ: zero for $\sigma \neq 0$, with the integral 1 over any range that contains $\sigma = 0$; the integral of $\delta_\mathrm{D}(\sigma)$ times a smooth function of σ is the value of that function at $\sigma = 0$. Unit: 1/length (for $\delta_\mathrm{D}(\sigma)$). Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- $\mathbb{M}$: The fold operator of dirac16complex: the 16×16 matrix (with a derivative along the fold) built from the Standard-Model fields at the fold by varying the portals with respect to $\bar{\Psi}$ (FE-13). Unit: GeV^{-3} (so that $\mathbb{M}/V_*$ is dimensionless). Status: DERIVED.
- n_o: The occupation of the Kohn-Sham orbital o (between 0 and 1). Unit: dimensionless. Status: RECORD.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.

- μ_Ψ: The chemical potential of dirac16complex. Unit: GeV. Status: DERIVED.
- T_Ψ: The temperature of dirac16complex. Unit: GeV. Status: DERIVED.

$$\langle \mathcal{T}_{\Psi MN} \rangle = -\frac{1}{4}\sum_o \mathsf{n}_o \Big(\bar\varphi_o \gamma_M D_N \varphi_o + \bar\varphi_o \gamma_N D_M \varphi_o - (D_M \bar\varphi_o)\gamma_N \varphi_o - (D_N \bar\varphi_o)\gamma_M \varphi_o \Big) + \frac{\lambda_\Psi}{2}\, g_{MN} \big(n_{\mathrm{s}}^2 - \mathrm{tr}\, \mathcal{N}^2 \big) \quad \text{(FE-42)}$$

Symbols in (FE-42).

- $\langle \cdots \rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- $\mathcal{T}_\Psi$: The energy-momentum tensor of dirac16complex: $-2/\sqrt{|g|}$ times the variation of the action S_Ψ of dirac16complex with respect to g^{MN}. Unit: GeV^8. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- n_o: The occupation of the Kohn-Sham orbital o (between 0 and 1). Unit: dimensionless. Status: RECORD.
- φ_o, $\bar\varphi_o$: A Kohn-Sham orbital of dirac16complex (a single-particle solution of the Kohn-Sham equation: the Dirac equation of dirac16complex with the mean field of the gas added to its mass) and its Dirac adjoint. Unit: GeV$^{7/2}$. Status: RECORD.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar\Psi = \partial_M\bar\Psi - \bar\Psi\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat h^{\mu\nu} D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- λ_Ψ: The coupling of the contact self-interaction of dirac16complex. Unit: GeV^{-6}. Status: RECORD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M \mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- n_{s}: The scalar density of dirac16complex in the Kohn-Sham state, $n_{\mathrm{s}} = \langle\bar\Psi\Psi\rangle = \mathrm{tr}\,\mathcal{N}$. Unit: GeV^7. Status: DERIVED.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- $\mathcal{N}$: The one-body density matrix of dirac16complex, $\mathcal{N} = \sum_o \mathsf{n}_o \varphi_o \bar\varphi_o$ (a 16×16 matrix at each point). Unit: GeV^7. Status: DERIVED.

The Kohn-Sham equations of dirac16complex coupled to itself, to gravity and to every Standard-Model field (FE-41) and the energy-momentum tensor of the Kohn-Sham state (FE-42): each orbital φ_o obeys the field equation with the mean fields, the self-interaction contributes the Hartree term $\lambda_\Psi n_{\mathrm{s}}$ and the exact exchange term $-\lambda_\Psi\mathcal{N}$, and every Standard-Model field enters through the mean fold operator $\langle\mathbb{M}\rangle$; correlation beyond exchange and the sea of negative-energy states are left out. Status: RECORD (the Kohn-Sham treatment), DERIVED (the exchange term), ASSUMED (no correlation, no sea).

9.1 The equations of the gas of bound particles

$$\begin{aligned}
&\Psi = \mathrm{e}^{3H|\sigma|}\,\chi_o(\sigma)\,\mathrm{e}^{-\mathrm{i}\omega x_4}, \qquad \chi_o' = \gamma^{(0)}\Big(\mathrm{i}\omega\,\gamma^{(4)} + m_\Psi + \mathcal{M}(\sigma)\Big)\chi_o \qquad (\sigma \neq 0),\\
&\mathcal{M}(\sigma) = G_{\mathrm{KS}}\,\mathrm{e}^{6H|\sigma|}\Big(\hat\rho(\sigma)\,\mathbf{1} - \hat{\mathcal{N}}(\sigma)\Big), \qquad \hat\rho = \frac{1}{4}\sum_{o=1}^{4}\bar\chi_o\chi_o, \qquad \hat{\mathcal{N}} = \frac{1}{4}\sum_{o=1}^{4}\chi_o\bar\chi_o, \qquad \text{(S-14)}\\
&\chi_o(0^+) = \exp\Big(\gamma^{(0)}\big(\mu_{\mathrm{L}}P_{\mathrm{L}} + \mu_{\mathrm{R}}P_{\mathrm{R}}\big)\Big)\chi_o(0^-), \qquad \int_{-\infty}^{\infty}\mathrm{d}\sigma\,\chi_o^\dagger\mathfrak{B}\chi_p = -\delta_{op}, \qquad G_{\mathrm{KS}} = \frac{\lambda_\Psi\, n_\chi}{V_*}
\end{aligned}$$

Symbols in (S-14).

- Ψ, $\bar{\Psi}$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar{\Psi} = \Psi^\dagger \sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\chi_o(\sigma)$, $\bar{\chi}_o$: The profile of the occupied Kohn-Sham orbital o ($o = 1, \ldots, 4$, the four positive-energy states of the bound level) of a homogeneous gas of the particles χ at rest on the static block: a 16-component spinor function of σ; the orbital of each particle is $\mathrm{e}^{3H|\sigma|}\chi_o(\sigma)\mathrm{e}^{-\mathrm{i}\omega x_4}$ divided by the square root of the volume of ordinary space and of V_*; normalised to the Krein charge -1, $\int \mathrm{d}\sigma\, \chi_o^\dagger \mathfrak{B} \chi_o = -1$; $\bar{\chi}_o = \chi_o^\dagger \sigma_{16}$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{1/2}$. Status: DERIVED.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- $\mathcal{M}(\sigma)$: The mean field of the self-interaction in the Kohn-Sham equation of the gas: the 16×16 matrix $\lambda_\Psi(n_\mathrm{s}\mathbf{1} - \mathcal{N})$ at each σ, which adds to the mass m_Ψ; positive (repulsive) for $\lambda_\Psi > 0$. Unit: GeV. Status: DERIVED.
- G_KS: The dimensionless strength of the self-interaction of the gas of particles χ in the Kohn-Sham states: the self-coupling λ_Ψ times the number density n_χ of the particles on the fold divided by the volume V_* of the extra times, $G_\mathrm{KS} = \lambda_\Psi n_\chi / V_*$ (in units of m_Ψ when numbers are given). Unit: dimensionless. Value: 10^{-6} to 1 in the solutions. Status: DERIVED.
- $\hat{\rho}(\sigma)$: The scalar density of the occupied orbitals, $\hat{\rho} = \frac{1}{4}\sum_o \bar{\chi}_o \chi_o$ (per particle); the scalar density of the gas is $n_\mathrm{s} = (n_\chi/V_*)\,\mathrm{e}^{6H|\sigma|}\hat{\rho}$. Unit: GeV. Status: DERIVED.
- $\mathbf{1}$, $\mathbf{1}_8$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_8$ is the 8×8 identity matrix. Unit: dimensionless. Status: STANDARD.
- $\hat{\mathcal{N}}(\sigma)$: The one-body density matrix of the occupied orbitals, $\hat{\mathcal{N}} = \frac{1}{4}\sum_o \chi_o \bar{\chi}_o$ (a 16×16 matrix at each σ, per particle); the density matrix of the gas is $\mathcal{N} = (n_\chi/V_*)\,\mathrm{e}^{6H|\sigma|}\hat{\mathcal{N}}$. Unit: GeV. Status: DERIVED.
- μ_X, μ_L, μ_R: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_\mathrm{L}P_\mathrm{L} + \mu_\mathrm{R}P_\mathrm{R})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- P_R: The author's right-handed chiral projector, $P_\mathrm{R} = (\mathbf{1} + \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\mathfrak{B}$: The Krein operator, $\mathfrak{B} = \mathrm{i}\sigma_{16}\gamma^{(4)}$: a Hermitian involution with the signature (8, 8); $\Psi^\dagger \mathfrak{B} \Psi$ is the charge

density. Unit: dimensionless. Status: DERIVED.

- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- λ_Ψ: The coupling of the contact self-interaction of dirac16complex. Unit: GeV^{-6}. Status: RECORD.
- n_χ: The number density of the dark-matter particles χ (particles plus antiparticles). Unit: GeV^3. Status: DERIVED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.

FE-41 for this state, with $m_\Psi = 1$ as the unit wherever numbers are given. The factor $\mathrm{e}^{3H|\sigma|}$ removes the first term of the spin connection FE-14 exactly ($\gamma^{(0)}\partial_\sigma \mathrm{e}^{3H|\sigma|} = 3H\,\mathrm{sgn}(\sigma)\gamma^{(0)}\mathrm{e}^{3H|\sigma|}$), and the second vanishes on the static block, so the four occupied orbitals $\chi_o(\sigma)$ obey a Dirac equation of one variable. Gravity enters through the warp: the proper scalar density of the gas at σ is $\mathrm{e}^{6H|\sigma|}$ times the density of the profiles, so the mean field $\mathcal{M}$ of the self-interaction (the Hartree part $\hat\rho\,\mathbf{1}$ and the exchange part $\hat{\mathcal{N}}$, the one-body density matrix) carries that factor. The Standard Model acts through the matching at the fold (FE-39). The occupied orbitals are the four states of the bound level with positive energy: Krein charge −1 each, energy $-\omega\mathcal{Q}_o = \omega$ (FE-43), mutually Krein-orthogonal. G_KS is the dimensionless strength of the self-interaction: the self-coupling λ_Ψ times the number density n_χ of the particles on the fold, divided by the volume V_* of the extra times (the orbital of each particle is $\mathrm{e}^{3H|\sigma|}\chi_o$ divided by the square root of the volume of ordinary space and of V_*, so that $\lambda_\Psi n_\mathrm{s} = G_\mathrm{KS}\,\mathrm{e}^{6H|\sigma|}\hat\rho$). Status: DERIVED (from FE-41, FE-14, FE-39).

$$\Sigma^{12}\chi_1 = \Sigma^{56}\chi_1 = \frac{\mathrm{i}}{2}\,\chi_1 \tag{S-15}$$

Symbols in (S-15).

- Σ^{ab}, $\Sigma^{\mu\nu}$: The generators of Spin(4,4) on dirac16complex, $\Sigma^{ab} = \frac{1}{4}[T^a, T^b]$ (the scaled commutators of the author's matrices); $\Sigma^{\mu\nu} = \mathbf{e}_a{}^\mu \mathbf{e}_b{}^\nu \Sigma^{ab}$ along the fold. Unit: dimensionless. Status: AUTHOR.
- $\chi_o(\sigma)$, $\bar\chi_o$: The profile of the occupied Kohn-Sham orbital o ($o = 1, \ldots, 4$, the four positive-energy states of the bound level) of a homogeneous gas of the particles χ at rest on the static block: a 16-component spinor function of σ; the orbital of each particle is $\mathrm{e}^{3H|\sigma|}\chi_o(\sigma)\mathrm{e}^{-\mathrm{i}\omega x_4}$ divided by the square root of the volume of ordinary space and of V_*; normalised to the Krein charge −1, $\int \mathrm{d}\sigma\, \chi_o^\dagger \mathfrak{B}\chi_o = -1$; $\bar\chi_o = \chi_o^\dagger\sigma_{16}$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{1/2}$. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- T_a, T^a: The author's eight real 16×16 Dirac matrices T16ˆA[a] of 4+4.nb ($a = 0, \ldots, 7$; $T_0, \ldots, T_7$), with $\{T_a, T_b\} = 2\eta_{ab}\mathbf{1}$; $T^a = \eta^{ab}T_b$. Unit: dimensionless. Status: AUTHOR.

The symmetry that makes the problem small. The author's $T_0, \ldots, T_3$ are symmetric and $T_4, \ldots, T_7$ antisymmetric, so the rotations of ordinary space (Σ^{ij}, $i, j = 1, 2, 3$) and of the extra times (Σ^{rs}, $r, s = 5, 6, 7$) form a compact group $SU(2) \times SU(2)$ of real orthogonal matrices, and $\gamma^{(0)}$, $\gamma^{(4)}$, P_L, P_R, σ_{16} and the Krein operator $\mathfrak{B}$ commute with it: the 16 components split into four copies of the representation $(2, 2)$ of this group. The four occupied orbitals form one $(2, 2)$ multiplet, the mean field is invariant, and the equations reduce exactly to the four-dimensional weight space S-15 (the states with the eigenvalue i/2 of both Σ^{12} and Σ^{56}, which hold the orbital $o = 1$): one orbital of four components, from which the multiplet is recovered by rotations by π (the orbitals $o = 2, 3, 4$ have the eigenvalues $(-\mathrm{i}/2, \mathrm{i}/2)$, $(\mathrm{i}/2, -\mathrm{i}/2)$ and $(-\mathrm{i}/2, -\mathrm{i}/2)$ of Σ^{12} and Σ^{56}) and checked in sixteen components. Status: DERIVED (`um3_kohnsham.py`, K1).

$$\begin{aligned}
&\mathsf{E}_\mathrm{int} = \frac{G_\mathrm{KS}}{2}\int_{-\infty}^{\infty} \mathrm{d}\sigma\, \mathrm{e}^{6H|\sigma|}\left(\hat\rho^2 - \mathrm{tr}\,\hat{\mathcal{N}}^2\right), \qquad \epsilon_\mathrm{p} = \omega - \mathsf{E}_\mathrm{int}, \qquad P_\mathrm{KS} = q_\chi = n_\chi\,\mathsf{E}_\mathrm{int},\\
&\Delta\omega = G_\mathrm{KS}\int_{-\infty}^{\infty} \mathrm{d}\sigma\, \mathrm{e}^{6H|\sigma|}\left(\hat\rho^2 - \mathrm{tr}\,\hat{\mathcal{N}}^2\right)\Big|_{G_\mathrm{KS}=0} \qquad \left(q_\mathrm{b} > 3H/2\right)
\end{aligned} \tag{S-16}$$

Symbols in (S-16).

- E_int: The interaction energy per particle of the gas of χ: the mean self-interaction energy $\frac{\lambda_\Psi}{2}(n_\mathrm{s}^2 - \mathrm{tr}\,\mathcal{N}^2)$ integrated across the hidden direction, per particle. Unit: GeV. Status: DERIVED.

- G_{KS}: The dimensionless strength of the self-interaction of the gas of particles χ in the Kohn-Sham states: the self-coupling λ_Ψ times the number density n_χ of the particles on the fold divided by the volume V_* of the extra times, $G_{\mathrm{KS}} = \lambda_\Psi n_\chi / V_*$ (in units of m_Ψ when numbers are given). Unit: dimensionless. Value: 10^{-6} to 1 in the solutions. Status: DERIVED.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)} H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- $\hat{\rho}(\sigma)$: The scalar density of the occupied orbitals, $\hat{\rho} = \frac{1}{4}\sum_o \bar{\chi}_o \chi_o$ (per particle); the scalar density of the gas is $n_{\mathrm{s}} = (n_\chi / V_*)\,\mathrm{e}^{6H|\sigma|}\hat{\rho}$. Unit: GeV. Status: DERIVED.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- $\hat{\mathcal{N}}(\sigma)$: The one-body density matrix of the occupied orbitals, $\hat{\mathcal{N}} = \frac{1}{4}\sum_o \chi_o \bar{\chi}_o$ (a 16×16 matrix at each σ, per particle); the density matrix of the gas is $\mathcal{N} = (n_\chi / V_*)\,\mathrm{e}^{6H|\sigma|}\hat{\mathcal{N}}$. Unit: GeV. Status: DERIVED.
- ϵ_{p}: The energy per particle of the gas of χ: the orbital energy minus the interaction energy per particle, which the orbital energy counts twice. Unit: GeV. Status: DERIVED.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- P_{KS}: The pressure of the gas of χ along ordinary space in the Kohn-Sham state: the particles are at rest, so it is the mean self-interaction, $P_{\mathrm{KS}} = n_\chi \mathsf{E}_{\mathrm{int}}$. Unit: GeV^4. Status: DERIVED.
- q_χ: The pressure of the bound states of dirac16complex along the extra times, integrated across the fold (their on-shell Lagrangian, the self-interaction energy per volume; zero for a dilute gas). Unit: GeV^4. Status: DERIVED.
- n_χ: The number density of the dark-matter particles χ (particles plus antiparticles). Unit: GeV^3. Status: DERIVED.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- q_{b}: The inverse width of a state of dirac16complex bound to the fold: its profile falls as $\mathrm{e}^{-q_{\mathrm{b}}|\sigma|}$. Unit: GeV. Status: DERIVED.

The energy of the state (FE-42 for the gas). $\mathsf{E}_{\mathrm{int}}$ is the interaction energy per particle, the mean value $\frac{\lambda_\Psi}{2}(n_{\mathrm{s}}^2 - \mathrm{tr}\,\mathcal{N}^2)$ of the self-interaction per particle (the trace runs over the sixteen components). The energy per particle is $\epsilon_{\mathrm{p}} = \omega - \mathsf{E}_{\mathrm{int}}$, because the orbital energy ω counts the interaction twice. The particles are at rest and do not depend on the extra times, so their kinetic terms contribute nothing to the pressure along ordinary space or along the extra times, and both pressures are the mean self-interaction, $P_{\mathrm{KS}} = q_\chi = n_\chi \mathsf{E}_{\mathrm{int}}$. $\Delta\omega$ is the first-order shift of the level, the mean of the mean field over the orbitals without the self-interaction (with the charge -1 of an orbital, a perturbation $\delta\mathcal{M}$ shifts its frequency by $\int \bar{\chi}_o\,\delta\mathcal{M}\,\chi_o\,\mathrm{d}\sigma$, and the average over the four orbitals gives S-16): it equals $2\mathsf{E}_{\mathrm{int}}$ to first order, and it is finite when the integrand, which behaves as $\mathrm{e}^{(6H-4q_{\mathrm{b}})|\sigma|}$, decays, that is for $q_{\mathrm{b}} > 3H/2$. Status: DERIVED.

$$\omega - \omega_{\mathrm{b}} \propto G_{\mathrm{KS}} \quad \big(q_{\mathrm{b}} > 3H/2\big), \qquad \omega - \omega_{\mathrm{b}} \propto \big(G_{\mathrm{KS}}\big)^{q_{\mathrm{b}}/(3H-q_{\mathrm{b}})} \quad \big(q_{\mathrm{b}} < 3H/2\big) \qquad (G_{\mathrm{KS}} \to 0) \tag{S-17}$$

Symbols in (S-17).

- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- ω_{b}: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- G_{KS}: The dimensionless strength of the self-interaction of the gas of particles χ in the Kohn-Sham states: the self-coupling λ_Ψ times the number density n_χ of the particles on the fold divided by the volume V_* of the extra times, $G_{\mathrm{KS}} = \lambda_\Psi n_\chi / V_*$ (in units of m_Ψ when numbers are given). Unit: dimensionless. Value: 10^{-6} to 1 in the solutions. Status: DERIVED.

- $q_{\rm b}$: The inverse width of a state of dirac16complex bound to the fold: its profile falls as $\mathrm{e}^{-q_{\rm b}|\sigma|}$. Unit: GeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

The dilute limit. For $q_{\rm b} < 3H/2$ the first-order integral diverges: the proper density of the orbital without the self-interaction grows as $\mathrm{e}^{(6H-2q_{\rm b})|\sigma|}$, and the mean field becomes of order one at the distance $\sigma_* = \ln(1/G_{\rm KS})/(6H-2q_{\rm b})$, beyond which it saturates (S-18). The shift is dominated by $|\sigma| < \sigma_*$ and scales as $G_{\rm KS}\,\mathrm{e}^{(6H-4q_{\rm b})\sigma_*} = (G_{\rm KS})^{q_{\rm b}/(3H-q_{\rm b})}$, a power below 1. Status: DERIVED (the scaling), verified numerically (K4).

$$\mathrm{e}^{6H|\sigma|}\,\hat\rho(\sigma) \to \hat\rho_\infty, \qquad |\chi_o| \propto \mathrm{e}^{-3H|\sigma|}, \qquad \det\Big(\gamma^{(0)}\big(\mathrm{i}\omega\gamma^{(4)} + m_\Psi + \mathcal{M}_\infty\big) \pm 3H\,\mathbf{1}\Big) = 0 \qquad (\pm\sigma \to \infty) \qquad \text{(S-18)}$$

Symbols in (S-18).

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings $\alpha_{(p)} = w_{(p)}H^{2-2p}$ and the cosmological constant $\Lambda = -7H^2$ (the author's original value $-10H^2$ was corrected on 2026-10-25); in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\hat\rho(\sigma)$: The scalar density of the occupied orbitals, $\hat\rho = \frac{1}{4}\sum_o \bar\chi_o\chi_o$ (per particle); the scalar density of the gas is $n_{\rm s} = (n_\chi/V_*)\,\mathrm{e}^{6H|\sigma|}\hat\rho$. Unit: GeV. Status: DERIVED.
- $\hat\rho_\infty$: The constant that the proper scalar density $\mathrm{e}^{6H|\sigma|}\hat\rho$ approaches far from the fold where it saturates. Unit: GeV. Status: DERIVED.
- $\chi_o(\sigma)$, $\bar\chi_o$: The profile of the occupied Kohn-Sham orbital o ($o = 1, \ldots, 4$, the four positive-energy states of the bound level) of a homogeneous gas of the particles χ at rest on the static block: a 16-component spinor function of σ; the orbital of each particle is $\mathrm{e}^{3H|\sigma|}\chi_o(\sigma)\mathrm{e}^{-\mathrm{i}\omega x_4}$ divided by the square root of the volume of ordinary space and of V_*; normalised to the Krein charge -1, $\int \mathrm{d}\sigma\,\chi_o^\dagger\mathfrak{B}\chi_o = -1$; $\bar\chi_o = \chi_o^\dagger\sigma_{16}$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{1/2}$. Status: DERIVED.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- $\mathcal{M}_\infty$: The limit of the mean field far from the fold where the proper density of the gas has saturated: the matrix for which the local equation has a solution decaying exactly at the rate $3H$. Unit: GeV. Status: DERIVED.
- $\mathbf{1}$, $\mathbf{1}_8$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_8$ is the 8×8 identity matrix. Unit: dimensionless. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.

Saturation. Where the proper density of the orbital would grow, a repulsive self-interaction stops it: if the orbital decayed more slowly than $\mathrm{e}^{-3H|\sigma|}$ the mean field would grow and raise the local mass, so that the orbital decays faster; if it decayed faster the mean field would vanish. The self-consistent orbital therefore decays exactly at the rate $3H$ far from the fold, its proper density approaches a constant $\hat\rho_\infty$, and the far mean field $\mathcal{M}_\infty$ is the one for which the local equation has a solution decaying at the rate $3H$ (S-18, the upper sign for $\sigma > 0$). The particles fill the bulk at a constant proper density there, but their number per area of the fold stays finite, because the proper volume of the bulk shrinks as $\mathrm{e}^{-6H|\sigma|}$. Saturation happens far from the fold for every $G_{\rm KS} > 0$ when $q_{\rm b} < 3H$, and, when $q_{\rm b} > 3H$, once the self-interaction has lowered the decay rate of the orbital below $3H$. Status: DERIVED, verified numerically (K5, K8).

$$\mu_{\mathrm{L}} = \mu_{\mathrm{L}0}\Big(1 - \kappa_{\mathrm{H}}\,\frac{\hat\rho_{\mathrm{L}}(0)}{\hat\rho_{\mathrm{L}0}(0)}\Big), \qquad \hat\rho_{\mathrm{L}} = \frac{1}{4}\sum_{o=1}^{4}\bar\chi_o P_{\mathrm{L}}\chi_o, \qquad \kappa_{\mathrm{H}} = \frac{c_\Phi\, n_\chi\, \hat\rho_{\mathrm{L}0}(0)}{\lambda\, v^2\, V_*} \tag{S-19}$$

Symbols in (S-19).

- μ_X, μ_{L}, μ_{R}: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_{\mathrm{L}}P_{\mathrm{L}} + \mu_{\mathrm{R}}P_{\mathrm{R}})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- $\mu_{\mathrm{L}0}$: The strength μ_{L} of the fold for the left-handed half without the feedback of the gas on the Higgs field. Unit: dimensionless. Value: -0.6 in the solutions. Status: DERIVED.
- κ_{H}: The strength of the Higgs-portal loop of the gas: the gas without the feedback changes the local v^2_{loc} at the fold by the factor $1-\kappa_{\mathrm{H}}$, $\kappa_{\mathrm{H}} = c_\Phi n_\chi \hat\rho_{\mathrm{L}0}(0)/(\lambda v^2 V_*)$; negative where the Higgs portal binds the fold ($c_\Phi < 0$, $\hat\rho_{\mathrm{L}0}(0) > 0$): the gas raises v^2_{loc}. Unit: dimensionless. Value: -0.3 in the test of the loop. Status: ASSUMED (the value), DERIVED (the sign).
- $\hat\rho_{\mathrm{L}}$: The scalar density of the left-handed half of the occupied orbitals, $\hat\rho_{\mathrm{L}} = \frac{1}{4}\sum_o \bar\chi_o P_{\mathrm{L}}\chi_o$; at the fold the mean of its values on the two sides. Unit: GeV. Status: DERIVED.
- $\hat\rho_{\mathrm{L}0}$: The left-handed scalar density $\hat\rho_{\mathrm{L}}$ of the bound state without the self-interaction and without the Higgs feedback. Unit: GeV. Status: DERIVED.
- $\chi_o(\sigma)$, $\bar\chi_o$: The profile of the occupied Kohn-Sham orbital o ($o = 1, \ldots, 4$, the four positive-energy states of the bound level) of a homogeneous gas of the particles χ at rest on the static block: a 16-component spinor function of σ; the orbital of each particle is $\mathrm{e}^{3H|\sigma|}\chi_o(\sigma)\mathrm{e}^{-\mathrm{i}\omega x_4}$ divided by the square root of the volume of ordinary space and of V_*; normalised to the Krein charge -1, $\int \mathrm{d}\sigma\, \chi_o^\dagger \mathfrak{B}\chi_o = -1$; $\bar\chi_o = \chi_o^\dagger\sigma_{16}$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{1/2}$. Status: DERIVED.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- P_{L}: The author's left-handed chiral projector, $P_{\mathrm{L}} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- n_χ: The number density of the dark-matter particles χ (particles plus antiparticles). Unit: GeV^3. Status: DERIVED.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_{\mathrm{F}})^{-1/2}$ with the Fermi constant $G_{\mathrm{F}} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.

The Higgs-portal loop. Where the Higgs portal alone sets the strength of the fold for the left-handed half (the condensates of the gauge fields and the fermion densities of FE-39 negligible), μ_{L} is proportional to $\langle\Phi^\dagger\Phi\rangle = v^2_{\mathrm{loc}}/2$, and the left-handed scalar density of the gas at the fold shifts v^2_{loc} (FE-49: $v^2_{\mathrm{loc}} = v^2 - c_\Phi\langle\bar\Psi P_{\mathrm{L}}\Psi\rangle/\lambda$, with $\langle\bar\Psi P_{\mathrm{L}}\Psi\rangle = n_\chi\hat\rho_{\mathrm{L}}(0)/V_*$ at the fold). $\mu_{\mathrm{L}0}$ and $\hat\rho_{\mathrm{L}0}(0)$ are the values without the feedback, $\hat\rho_{\mathrm{L}}(0)$ is the mean of the values on the two sides of the fold, and κ_{H} measures the strength of the loop: the gas without feedback changes v^2_{loc} by the factor $1-\kappa_{\mathrm{H}}$. The sign of κ_{H} is fixed. Where the Higgs portal alone binds the fold ($\mu_{\mathrm{R}} = 0$), $\mu_{\mathrm{L}0} = c_\Phi v^2/(2V_*) < 0$, so $c_\Phi < 0$; and the occupied orbitals have $\hat\rho_{\mathrm{L}0}(0) = \partial\omega_{\mathrm{b}}/\partial\mu_{\mathrm{L}} = -m^2_\Psi\mu_{\mathrm{L}0}/(4\omega_{\mathrm{b}}) > 0$ (the Hellmann-Feynman theorem: with the Krein charge -1, the shift of the level is the mean value of the perturbation $\delta\mu_{\mathrm{L}}P_{\mathrm{L}}$ at the fold; FE-40 for the derivative). Hence $\kappa_{\mathrm{H}} = 2\mu_{\mathrm{L}0}n_\chi\hat\rho_{\mathrm{L}0}(0)/(\lambda v^4) < 0$: the gas raises v^2_{loc}, the fold binds more strongly, and the level falls, in agreement with the coupling $\mathfrak{g}_{h\chi} < 0$ of FE-47. If $\hat\rho_{\mathrm{L}}(0)$ were proportional to q_{b}, and q_{b} to μ_{L}, the loop would end at $\mu_{\mathrm{L}} = \mu_{\mathrm{L}0}/(1+\kappa_{\mathrm{H}})$, which does not exist for $\kappa_{\mathrm{H}} \le -1$: a gas dense enough would deepen its own binding until q_{b} approaches m_Ψ, where this proportionality fails (ESTIMATE). The orbitals and μ_{L} are iterated together. Status: DERIVED (from FE-39, FE-40 and FE-49), ASSUMED (the value of κ_{H} used below).

9.2 The method

The bound level is found by shooting. On each side of the fold the equation S-14 is integrated inward from $|\sigma| = L$ (the half-width L of the computational box, given for each regime below) with the classical fourth-order Runge-Kutta method of step h (the step written as a matrix and the solutions re-orthonormalised every eight steps), starting with the solutions that decay at $|\sigma| = L$; the frequency is the zero of the residual of the matching at the fold for a matching vector of negative Krein norm, searched within $0.05 m_\Psi$ of the previous level (a frequency beyond the local threshold at the far end, or a mean field so large that the integration overflows, has no bound level). The orbital gives the densities, the densities the mean field, and the mean field (with μ_L for the Higgs loop) is iterated with Anderson mixing. Plain mixing would need a very small weight: far from the fold, where the density saturates, a small change of the mean field changes the decay rate of the orbital and with it the far density by a large factor of the opposite sign, so the linearised Kohn-Sham map has large negative eigenvalues (an argument; the eigenvalues are not computed). The mixing (the weight β of the new field and the depth of the history) is $\beta = 0.5$ with six previous iterates, or, where this does not converge (above the free threshold, where the saturated region is long), $\beta = 0.2$ with ten; the extrapolated field is held to the same bound as the computed one. An iteration has converged when the largest relative change of the mean field is below 10^{-8} and the level changes by less than 10^{-10}; where the precision of the level sets a floor of a few 10^{-8}, the best iterate below 10^{-7} is taken. The coupling is raised by continuation, with intermediate couplings inserted where a step fails. Every converged solution is lifted to sixteen components and checked there: the Kohn-Sham equation by central differences on the grid, with the mean field rebuilt in sixteen components from the lifted orbitals, the matching at the fold, the Krein charges and the Krein orthogonality; the residual of self-consistency is that of the reduced problem, evaluated once more with the accepted field. Status: DERIVED (`um3_kohnsham.py`).

9.3 The solution in three regimes

regime	H	box L	step h	the orbital without self-interaction
I	0.05	60	0.03	$q_\mathrm{b} > 3H$: its proper density decays away from the fold
II	0.15	60	0.02	$3H/2 < q_\mathrm{b} < 3H$: its proper density grows, the first-order shift is finite
III	0.3	20	0.02	$q_\mathrm{b} < 3H/2$: its proper density grows, the first-order shift diverges

The three regimes, all with the Higgs portal of the vacuum alone, $\mu_\mathrm{L} = -0.6$ and $\mu_\mathrm{R} = 0$, so that $q_\mathrm{b} = 0.3$ and $\omega_\mathrm{b} = \sqrt{1 - 0.09} = 0.953939$ (units of m_Ψ). In each the coupling G_KS is raised from 10^{-6} (regime I) or from a coupling so small that the proper density does not saturate inside the box (2.5×10^{-9} in regime II, 6.3×10^{-12} in regime III) to 1. Status: ASSUMED (the parameters), DERIVED (the solutions).

Regime I ($H = 0.05$, $6H = 0.3$):

G_KS	ω	$\omega - \omega_\mathrm{b}$	E_int	$P_\mathrm{KS}/(n_\chi \epsilon_\mathrm{p})$	iterations	decay rate of $\hat{\rho}$ at $0.85L$
10^{-6}	0.95393934	1.36×10^{-7}	6.83×10^{-8}	7.2×10^{-8}	4	0.6000
10^{-5}	0.95394057	1.36×10^{-6}	6.83×10^{-7}	7.2×10^{-7}	4	0.6000
10^{-4}	0.95395285	1.36×10^{-5}	6.82×10^{-6}	7.2×10^{-6}	5	0.5999
10^{-3}	0.95407566	1.36×10^{-4}	6.82×10^{-5}	7.2×10^{-5}	6	0.5991
0.01	0.95530069	0.00136	6.80×10^{-4}	7.1×10^{-4}	7	0.5913
0.1	0.96723519	0.0133	0.00659	0.0069	10	0.5078

$G_{\rm KS}$	ω	$\omega-\omega_{\rm b}$	$\mathsf{E}_{\rm int}$	$P_{\rm KS}/(n_\chi\epsilon_{\rm p})$	iterations	decay rate of $\hat\rho$ at $0.85L$
1	1.07039318	0.116	0.0564	0.056	42	0.3000

Regime II ($H=0.15$, $6H=0.9$):

$G_{\rm KS}$	ω	$\omega-\omega_{\rm b}$	$\mathsf{E}_{\rm int}$	$P_{\rm KS}/(n_\chi\epsilon_{\rm p})$	iterations	decay rate of $\hat\rho$ at $0.85L$
10^{-6}	0.95393961	4.09×10^{-7}	2.05×10^{-7}	2.1×10^{-7}	20	0.8677
10^{-5}	0.95394329	4.09×10^{-6}	2.05×10^{-6}	2.1×10^{-6}	35	0.8929
10^{-4}	0.95397992	4.07×10^{-5}	2.03×10^{-5}	2.1×10^{-5}	46	0.8986
10^{-3}	0.95433722	3.98×10^{-4}	1.98×10^{-4}	2.1×10^{-4}	42	0.8997
0.01	0.95764791	0.00371	0.00182	0.0019	42	0.9000
0.1	0.98687286	0.0329	0.0160	0.016	43	0.9000
1	1.25973443	0.306	0.152	0.14	132	0.9000

Regime III ($H=0.3$, $6H=1.8$):

$G_{\rm KS}$	ω	$\omega-\omega_{\rm b}$	$\mathsf{E}_{\rm int}$	$P_{\rm KS}/(n_\chi\epsilon_{\rm p})$	iterations	decay rate of $\hat\rho$ at $0.85L$
10^{-6}	0.95407252	1.33×10^{-4}	4.47×10^{-5}	4.7×10^{-5}	31	1.7559
10^{-5}	0.95436976	4.31×10^{-4}	1.46×10^{-4}	1.5×10^{-4}	38	1.7874
10^{-4}	0.95536905	0.00143	4.92×10^{-4}	5.2×10^{-4}	35	1.7966
10^{-3}	0.95896972	0.00503	0.00179	0.0019	34	1.7992
0.01	0.97368303	0.0197	0.00747	0.0077	39	1.7999
0.1	1.04743537	0.0935	0.0385	0.038	35	1.8000
1	1.57066625	0.617	0.288	0.22	78	1.8000

The self-consistent states. ω is the orbital energy of the occupied level, $\mathsf{E}_{\rm int}$ the interaction energy per particle, $P_{\rm KS}/(n_\chi\epsilon_{\rm p})$ the pressure divided by the energy density, and the last column the decay rate of the scalar density at $0.85L$ ($6H$ where the proper density has saturated). Status: DERIVED (`um3_kohnsham.py`, K3).

The free level and the symmetry. Without the self-interaction the shooting with the author's 16×16 matrices gives $\omega=0.9539392014$, which is FE-40 ($\sqrt{1-q_{\rm b}^2}=0.9539392014$) to 10^{-10}; the level consists of eight degenerate states, four of each Krein sign, and the reduced four-dimensional problem of S-15 has the same level with two degenerate states (one of each sign). Status: DERIVED (K1).

The dilute limit. In regime I the shift of the level divided by the first-order shift of S-16 (the coefficient $\Delta\omega/G_{\rm KS}=0.136501$, computed in sixteen components with the four occupied orbitals of the level of eight degenerate states) is 0.99997, 0.99997, 0.99995, 0.99972 for $G_{\rm KS}=10^{-6}$, 10^{-5}, 10^{-4} and 10^{-3}: the Kohn-Sham states reduce to first-order perturbation theory (the remaining 3×10^{-5} is of the size of the quadrature error on the grid). Fitted for $G_{\rm KS}\le10^{-4}$, the shift grows as $G_{\rm KS}^{p}$ with $p=1.000$, 0.999 and 0.515 in the three regimes, against S-17: 1, 1 and $q_{\rm b}/(3H-q_{\rm b})=0.500$. Status: DERIVED (K2, K4).

Saturation. At $G_{\rm KS}=0.01$ the scalar density decays far from the fold at 0.9000 in regime II and 1.7999 in regime III, the rates $6H=0.9$ and 1.8 of S-18: the proper density has saturated, and the far mean field is the one that gives the orbital the decay rate $3H$. An attractive self-interaction behaves differently in the two kinds of regime: for $G_{\rm KS}=-10^{-4}$ and -10^{-2} the iteration, started from the free state, converges in regime I (where the proper density decays and the attraction only lowers the level, to $\omega=0.95392555$ and 0.95257077); in regimes II and III, where the proper density of the free state grows away from the fold, the level search finds no Krein-negative level near the previous one: the attraction grows with the proper density and would close the local gap. That no self-consistent state exists there at all is not proven by this (OPEN). Status: DERIVED (K5).

Through the free threshold. The repulsive mean field raises the level through the threshold $\omega = m_\Psi$ of the free continuum: at $G_{\rm KS} = 0.371$ in regime I and at $G_{\rm KS} = 0.142$ in regime II. The state stays bound up to $G_{\rm KS} = 1$ ($\omega = 1.070393$ and 1.259734 there): far from the fold the proper density saturates (in regime I once the decay rate of the orbital has fallen below $3H$, at $\omega > \sqrt{1-9H^2} = 0.9887$), the density decays at $6H$ (to within 5×10^{-4} and 1×10^{-8} above the threshold), and the far mean field keeps the local mass above ω. In the computed cases a repulsive self-interaction therefore does not dissolve the particles χ into the bulk: it makes them heavier and spreads them, and their energy may exceed m_Ψ; the mechanism, the saturation of the proper density, rests on the warp and needs $H > 0$. Status: DERIVED (K8).

The Higgs-portal loop. The occupied orbitals without the self-interaction have $\hat\rho_{\rm L0}(0) = 0.15723848$, equal to $\partial\omega_{\rm b}/\partial\mu_{\rm L} = -\mu_{\rm L0}/(4\omega_{\rm b}) = 0.15724273$ up to the relative difference 2.7×10^{-5}, which is the error $(2q_{\rm b}h)^2/12 = 2.7\times10^{-5}$ of the trapezoidal normalisation on the grid; so $\kappa_{\rm H} < 0$. With $\kappa_{\rm H} = -0.3$ and $G_{\rm KS} = 0.1$ in regime I the orbitals and $\mu_{\rm L}$ of S-19 converge together in 12 iterations: $\mu_{\rm L} = -0.874681$ (from -0.6; the estimate $\mu_{\rm L0}/(1+\kappa_{\rm H})$ gives -0.8571): the gas raises the local electroweak vacuum, the fold binds more strongly, and the level falls to $\omega = 0.91512441$ (0.96723519 without the feedback). Status: DERIVED (K6), ASSUMED (the value of $\kappa_{\rm H}$).

The equation of state. The pressure is positive for the repulsive self-interaction and vanishes in the dilute limit: $P_{\rm KS}/(n_\chi\epsilon_{\rm p}) = 7.2\times10^{-8}$, 2.1×10^{-7} and 4.7×10^{-5} at $G_{\rm KS} = 10^{-6}$ in the three regimes, and 0.056, 0.137 and 0.224 at $G_{\rm KS} = 1$. A dilute gas of the particles χ is pressureless: cold dark matter. Status: DERIVED (K7).

The residuals. Each of the 83 converged solutions (the three regimes from the smallest coupling to 1, with the continuation between) was lifted to sixteen components and checked there: the largest relative residual of the Kohn-Sham equation by central differences on the grid is 2×10^{-4} (the error of the difference formula, of order h^2), of the matching at the fold 4×10^{-9}, of self-consistency (in the reduced problem, evaluated once more with the accepted field) 2×10^{-7}; the four orbitals have the Krein charge -1 to 4×10^{-16} and are Krein-orthogonal to 5×10^{-17}, and the bound on the mean field is never active. Status: DERIVED (K3).

9.4 A weakly bound gas

The direct-detection bound on the Higgs portal requires a weakly bound χ, $q_{\rm b}/\omega_{\rm b} \lesssim 0.03$ at a few TeV, and a nearly luminal one needs $q_{\rm b} \gg H$ (S-13). A point of this kind is solved as well: $\mu_{\rm L} = -0.06$ ($q_{\rm b} = 0.03$, $q_{\rm b}/\omega_{\rm b} = 0.0300$), $H = 0.002$ (so $q_{\rm b} = 15H$, regime I, and $c_{\rm b}^2 = q_{\rm b}/(q_{\rm b}-H) = 1.071$), with the box $L = 500$ and the step $h = 0.25$. The shooting in sixteen components gives $\omega = 0.9995498987$ (FE-40: 0.9995498987) with four Krein-negative orbitals, and the first-order coefficient of S-16 is $\Delta\omega/G_{\rm KS} = 0.012489$. The self-consistent shift divided by the first-order shift is 0.99940, 0.99429, 0.94309 for $G_{\rm KS} = 10^{-4}$, 10^{-3} and 10^{-2} (approaching 1 as $G_{\rm KS}\to0$), with $P_{\rm KS}/(n_\chi\epsilon_{\rm p}) = 6.2\times10^{-7}$, 6.2×10^{-6}, 5.8×10^{-5}: a dilute gas of weakly bound particles χ is cold, and its self-interaction raises the orbital energy by about $0.012\,m_\Psi$ per unit of $G_{\rm KS}$ and the energy per particle, the mass of χ in the gas, by half of it. Status: DERIVED (K9).

9.5 The Krein partner and the stability of the gas

$$\{\mathsf{a}_o, \mathsf{a}_o^\dagger\} = -\mathcal{Q}_o, \qquad \mathsf{H}_\Psi = \sum_o \big(-\mathcal{Q}_o\,\omega_o\big)\,\mathsf{a}_o^\dagger\mathsf{a}_o, \qquad \big[\mathsf{H}_\Psi, \mathsf{a}_o^\dagger\big] = \omega_o\,\mathsf{a}_o^\dagger \tag{FE-67}$$

Symbols in (FE-67).

- a_o, $\mathsf{a}_o^\dagger$: The annihilation and the creation operator of the mode o of the quantised dirac16complex (frequency ω_o, Krein charge $\mathcal{Q}_o$); the dagger is the adjoint with respect to the indefinite (Krein) scalar product. Unit: dimensionless. Status: STANDARD.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\mathcal{Q}_o$: The charge Q_Ψ (the Krein charge) of the single-particle state o of dirac16complex, a Kohn-Sham orbital or a mode: $+1$ or -1 for a normalised state. Unit: dimensionless. Status: DERIVED.
- H_Ψ: The Hamiltonian (the energy operator) of the quantised dirac16complex. Unit: GeV. Status: DERIVED.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.

The quantised dirac16complex: a mode o of the frequency ω_o and the Krein charge $\mathcal{Q}_o = \pm 1$ (FE-43) has the anticommutator $-\mathcal{Q}_o$, and every quantum raises the energy by its frequency. In the positive-energy quantisation every particle and every hole carries a positive energy and the norm $-\mathcal{Q}_o$ of its mode: the particles and the holes of the modes of charge -1 are physical, those of the modes of charge $+1$ are ghosts (negative norm), and no choice of the vacuum removes them. The Kohn-Sham states above occupy four orbitals of charge -1: they are physical states, and the questions are whether they are stable and whether the couplings of the fold produce ghosts. Status: DERIVED (`um3_krein.py`, KR1).

The symmetry of the bound states and its mirrors. At rest, the transformations of the sixteen components that keep the bound states of the fold (they commute with $\gamma^{(0)}$, $\gamma^{(4)}$ and Γ_8 and preserve the Dirac adjoint) form a Lie algebra of dimension 32: 16 compact generators keep the two Krein charges apart, and 16 boosts mix them, among them the nine boosts Σ^{ir} between ordinary space and the extra times; for a moving particle only four compact generators remain. The mirror R_1 of the fold (FD-10) keeps the charges: the four bound states of charge -1 are even, the four of charge $+1$ odd, and the self-consistent orbital and its mean field are even; R_2 exchanges the two charges. Status: DERIVED (KR2, KR3).

The stability of the gas is not decided. The boosts that mix the charges map the self-consistent gas onto self-consistent gases (in regime I at $G_{\mathrm{KS}} = 0.1$ the residual of the Kohn-Sham equation stays 2.43×10^{-5}, as without the boost, and the charges stay -1, while the orbitals acquire admixtures of charge $+1$ whose positive-definite norm grows up to 2.43 times): these directions are exactly flat, and exact zero modes would give the frequency 0 in the linear response. The linear response truncated to the bound level (the 16 admixtures of the four empty orbitals of charge $+1$, split from the occupied level by 4.815×10^{-3}) has frequencies of the order 10^{-4}, two of them imaginary (up to 1.83×10^{-4}): the truncation error is as large as these frequencies, so the truncated response does not decide whether the dilute gas is stable. Status: DERIVED (KR4).

The partner level. In the self-consistent mean field the level of charge $+1$ next to the occupied one, its partner, is real and bound in the dilute gas of all three regimes ($G_{\mathrm{KS}} = 0.01$ and 0.1), and it moves away from the occupied level as G_{KS} grows (the gap grows from 0.0005 to 0.038 in regime I, from 0.0013 to 0.114 in regime II and from 0.0052 to 0.230 in regime III between $G_{\mathrm{KS}} = 0.01$ and 1, in units of m_Ψ): the two cannot mix, because they have opposite parity under the mirror R_1 of the fold in a mirror-even mean field. In the denser gases the partner becomes complex, $\omega \pm \mathrm{i}\Gamma_{\mathrm{K}}$:

regime	partner real and bound up to	Γ_{K} of the complex partner (units of m_Ψ)	where it turns complex
I ($H = 0.05$)	$G_{\mathrm{KS}} = 0.2$	5.1×10^{-6} at $G_{\mathrm{KS}} = 0.3$, 1.2×10^{-4} at 0.5, 3.5×10^{-4} at 0.7, 6.1×10^{-4} at 1	where it reaches the continuum at the end of the computational box
II ($H = 0.15$)	$G_{\mathrm{KS}} = 0.5$	3.9×10^{-3} at 0.7, 9.7×10^{-3} at 1	where it meets another level of the discrete spectrum
III ($H = 0.3$)	$G_{\mathrm{KS}} = 0.7$	1.5×10^{-2} at 1	where it meets another level of the discrete spectrum

In regime I the complex partner lies in the continuum of the end of the box, so it depends on the size of the box, which was not varied; whether these complex levels are growing modes of the gas is therefore not established. Status: DERIVED (KR5), OPEN (the convergence in the size of the box and the stability of the gas).

The couplings of the fold decide unitarity. At rest none of the 16 chirality-blind operators of the fold (the products of $\gamma^{(1)}, \ldots, \gamma^{(4)}$) connects a physical state with a ghost, and each of the 16 chiral ones (the same times Γ_8) does. The left-handed Higgs portal P_{L} of the author has the elastic amplitude 0.157, no conversion of a particle into a ghost, no physical pair at rest (a p-wave), and the amplitude 0.140 (an s-wave) for a physical particle χ with a ghost antiparticle, 0.89 times the elastic one. For moving particles the scalar, the density and the vector currents still do not mix (below 10^{-9}), but the magnetic dipole Σ^{12} does, in proportion to the momentum (6.6×10^{-4} and 2.1×10^{-3} at $|\vec{k}| = 0.05$ and $0.15\, m_\Psi$). The vacuum cannot decay (every quantum has a positive energy), but with chiral couplings at the fold ghosts are produced wherever the portals act: UM-3 with the author's left-handed Higgs

portal is not a unitary quantum theory. Chirality-blind couplings are necessary for a unitary UM-3, but they are not shown to be sufficient, since the magnetic dipole connects the two sectors for moving particles. Status: DERIVED (KR6, KR7), OPEN (a unitary set of couplings).

10 The residuals of every field equation

The approximate solution of UM-3 consists of the parts above and below. The field equations that the other sections do not write out are restated first, so that the table at the end of this section can refer to each of them: the field equations of the bulk, the junction conditions and the thin source of the fold, the conservation law at the fold, the field equations of the gluons and of the hypercharge field, and those of the quarks and leptons.

$$\sum_{p=1}^{3} \alpha_{(p)}\, \mathcal{E}_{(p)}{}^{M}{}_{N} + \Lambda\, \delta^{M}_{N} = \kappa\, \mathcal{T}_{\Psi}{}^{M}{}_{N} \qquad (\sigma \neq 0) \tag{FE-1}$$

Symbols in (FE-1).

- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)} H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- $\mathcal{E}_{(p)}{}^{M}{}_{N}$: The Lovelock tensor of order p, $\mathcal{E}_{(p)}{}^{M}{}_{N} = -2^{-(p+1)}\, \delta^{M P_1 \cdots P_{2p}}_{N Q_1 \cdots Q_{2p}}\, R^{Q_1 Q_2}{}_{P_1 P_2} \cdots R^{Q_{2p-1} Q_{2p}}{}_{P_{2p-1} P_{2p}}$ (with both indices down it is $1/\sqrt{|g|}$ times the variation of $\sqrt{|g|}\, \mathcal{L}_{(p)}$ with respect to g^{MN}; $\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor, $\mathcal{E}_{(3)}$ the cubic Lovelock tensor); with frame indices its frame components. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -7H^2$: with the corrected weights the value for which the static block is a vacuum (its Lovelock tensors add to $7H^2$ in every direction). The author's original value was $\Lambda = -10H^2$ (he writes the pure number $\Lambda_{\mathrm{N}} = 10$ in $0 = -\Lambda_{\mathrm{N}} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$). Unit: GeV^2. Value: $-7H^2$; the author's original $-10H^2$. Status: AUTHOR (the term), DERIVED (the value).
- δ: The Kronecker delta: with two indices $\delta^{M}_{N} = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta^{M_1 M_2 \cdots}_{N_1 N_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\Psi}$: The energy-momentum tensor of dirac16complex: $-2/\sqrt{|g|}$ times the variation of the action S_{Ψ} of dirac16complex with respect to g^{MN}. Unit: GeV^8. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\, \mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z) \ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

The Einstein-Lovelock equations of the bulk, away from the fold: the Lovelock tensor of the author's theory with the corrected weights equals κ times the energy-momentum tensor of the field of the bulk, dirac16complex; with the corrected weights the bulk needs no other source (IN-7). Status: AUTHOR (the theory), STANDARD (the variation).

$$\begin{aligned} \kappa\, \mathcal{S}^{a}{}_{b} &= \lim_{\varepsilon \to 0} \int_{-\varepsilon}^{\varepsilon} \mathrm{d}\sigma \sum_{p=1}^{3} \alpha_{(p)}\, \mathcal{E}_{(p)}{}^{a}{}_{b} = \mathfrak{F}^{a}{}_{b}\big|_{\sigma=0^+} - \mathfrak{F}^{a}{}_{b}\big|_{\sigma=0^-}, \\ \frac{\partial \mathfrak{F}^{a}{}_{b}}{\partial(\partial_\sigma \mathfrak{f})} &= \frac{\partial}{\partial(\partial^2_\sigma \mathfrak{f})} \sum_{p=1}^{3} \alpha_{(p)}\, \mathcal{E}_{(p)}{}^{a}{}_{b}, \qquad \mathfrak{f} \in \{\mathfrak{a}, \mathfrak{b}, \mathfrak{n}\} \end{aligned} \tag{FE-4}$$

Symbols in (FE-4).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- ${\mathcal{S}^a}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa {\mathcal{S}^a}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)} {\mathcal{E}_{(p)}}^a{}_b$ (FD-11, the junction conditions); ${\mathcal{S}^4}_4$ is minus its energy density, ${\mathcal{S}^1}_1$ and ${\mathcal{S}^5}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa {\mathcal{S}^a}_b = -\frac{32}{5} H {\delta^a}_b$ at the static block. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- ε: The half-thickness of a layer around the fold, taken to zero. Unit: length. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\, \mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z) \ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations also a second label of a Kohn-Sham orbital, with the Kronecker delta of the two labels. Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)} H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- ${\mathcal{E}_{(p)}}^M{}_N$: The Lovelock tensor of order p, ${\mathcal{E}_{(p)}}^M{}_N = -2^{-(p+1)} \delta^{MP_1 \cdots P_{2p}}_{NQ_1 \cdots Q_{2p}} R^{Q_1 Q_2}{}_{P_1 P_2} \cdots R^{Q_{2p-1} Q_{2p}}{}_{P_{2p-1} P_{2p}}$ (with both indices down it is $1/\sqrt{|g|}$ times the variation of $\sqrt{|g|}\, \mathcal{L}_{(p)}$ with respect to g^{MN}; $\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor, $\mathcal{E}_{(3)}$ the cubic Lovelock tensor); with frame indices its frame components. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- ${\mathfrak{F}^a}_b$: The junction potential: a function of the first σ-derivatives of the metric functions whose derivatives are the coefficients of the second σ-derivatives in $\sum_p \alpha_{(p)} {\mathcal{E}_{(p)}}^a{}_b$ (FE-4). Unit: GeV. Status: DERIVED.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- $\mathfrak{f}$: Any one of the three metric functions $\mathfrak{a}, \mathfrak{b}, \mathfrak{n}$ of the general warped metric (FE-34). Unit: dimensionless. Status: DERIVED.
- $\mathfrak{a}$, $\mathfrak{a}_1$: The logarithm of the scale factor of ordinary space in the general warped metric FE-34, a function of σ and x_4; for the author's metric $\mathfrak{a} = a_4 - H|\sigma|$; $\mathfrak{a}_n$ its term of order n in the power series FE-35 ($\mathfrak{a}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $\mathfrak{b}$, $\mathfrak{b}_1$: The logarithm of the scale factor of the extra times in the general warped metric FE-34; for the author's metric $\mathfrak{b} = \ln b_4 - a_4 - H|\sigma|$; $\mathfrak{b}_n$ its term of order n ($\mathfrak{b}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $\mathfrak{n}$, $\mathfrak{n}_1$: The logarithm of the lapse in the general warped metric FE-34 ($g_{x_4 x_4} = -\mathrm{e}^{2\mathfrak{n}}$); for the author's metric $\mathfrak{n} = 0$; $\mathfrak{n}_n$ its term of order n ($\mathfrak{n}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.

The field equations at the fold: the metric has a kink at $\sigma = 0$, so its second derivatives along σ contain a delta function, and the jump of the junction potential across the fold equals κ times the thin source $\mathcal{S}$ of the fold. Status: DERIVED.

$$ {\mathcal{S}^a}_b = \frac{1}{V_*}\Big({\mathcal{T}_{\mathrm{H}}}^a{}_b + {\mathcal{T}_{\mathrm{gauge}}}^a{}_b + {\mathcal{T}_{\mathrm{f}}}^a{}_b + {\mathcal{T}_{\mathrm{P}}}^a{}_b + {\mathcal{T}_{\mathrm{ff}}}^a{}_b + {\mathcal{T}_{\mathrm{IG}}}^a{}_b \Big), \qquad {\mathcal{T}_{\mathrm{IG}}}^\mu{}_\nu = -\frac{1}{\kappa_{\mathrm{ind}}} {\hat{G}^\mu}_\nu, \qquad {\mathcal{T}^r}_r = \mathcal{L}_{\mathrm{fold}} \qquad \text{(FE-5)} $$

Symbols in (FE-5).

- ${\mathcal{S}^a}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa {\mathcal{S}^a}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)} {\mathcal{E}_{(p)}}^a{}_b$ (FD-11, the junction conditions); ${\mathcal{S}^4}_4$ is minus its energy density, ${\mathcal{S}^1}_1$ and ${\mathcal{S}^5}_5$ its pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa {\mathcal{S}^a}_b = -\frac{32}{5} H {\delta^a}_b$ at the static block. Status: DERIVED.

- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\, \mathrm{d}x_6\, \mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- $\mathcal{T}_\mathrm{H}$: The energy-momentum tensor of the Higgs doublet (four-dimensional normalisation). Unit: GeV^4. Status: STANDARD.
- $\mathcal{T}_\mathrm{gauge}$: The energy-momentum tensor of the gauge fields. Unit: GeV^4. Status: STANDARD.
- $\mathcal{T}_\mathrm{f}$: The energy-momentum tensor of the quarks and leptons. Unit: GeV^4. Status: STANDARD.
- $\mathcal{T}_\mathrm{P}$: The energy-momentum tensor of the portals of dirac16complex. Unit: GeV^4. Status: DERIVED.
- $\mathcal{T}_\mathrm{ff}{}^a{}_b$: The energy-momentum tensor of the fold fluid (frame components of the fold, four-dimensional normalisation). Unit: GeV^4. Status: ASSUMED.
- $\mathcal{T}_\mathrm{IG}$: The energy-momentum tensor of the induced Einstein-Hilbert term of the fold, $\mathcal{T}_\mathrm{IG}{}^\mu{}_\nu = -\hat{G}^\mu{}_\nu/\kappa_\mathrm{ind}$. Unit: GeV^4. Status: ESTIMATE.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+, +, +, -$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- κ_ind: The coupling of the Einstein-Hilbert term $\hat{R}/(2\kappa_\mathrm{ind})$ that the quantum fluctuations of the fields of the fold may induce, $\kappa_\mathrm{ind} = 8\pi G_\mathrm{ind}$; it adds to the gravity of the fold, $1/(8\pi G_\mathrm{N}) = V_*/\kappa_7 + 1/\kappa_\mathrm{ind}$. Unit: GeV^{-2}. Value: not fixed by UM-3. Status: ESTIMATE.
- $\hat{G}^\mu{}_\nu$: The Einstein tensor of the four-dimensional metric $\hat{h}_{\mu\nu}$ of the fold. Unit: 1/length^2. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- $\mathcal{T}$: The energy-momentum tensor of any one term of the fold (in $\mathcal{T}^r{}_r = \mathcal{L}_\mathrm{fold}$). Unit: GeV^4. Status: STANDARD.
- $\mathcal{L}_\mathrm{fold}$: The Lagrangian of any one term of the fold (four-dimensional normalisation). Unit: GeV^4. Status: STANDARD.

The thin source of the fold: the energy-momentum tensor of everything that lives on the fold (the Standard Model, the portals, the fold fluid and, if the fields of the fold induce one, the induced gravity) divided by the volume V_* of the extra times; along each extra time its component is the Lagrangian of the term, because the fields of the fold do not depend on the extra times. Status: DERIVED.

$$\begin{aligned} &\nabla_M\, \mathcal{T}_\Psi{}^M{}_N = 0 \qquad (\sigma \neq 0), \\ &\hat{\nabla}_a\, \mathcal{S}^a{}_b = -\Big(\mathcal{T}_\mathrm{bulk}{}^{(0)}{}_b\big|_{\sigma=0^+} - \mathcal{T}_\mathrm{bulk}{}^{(0)}{}_b\big|_{\sigma=0^-} \Big) \end{aligned} \tag{FE-11}$$

Symbols in (FE-11).

- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols $\Gamma^P{}_{MN}$). Status: STANDARD.
- $\mathcal{T}_\Psi$: The energy-momentum tensor of dirac16complex: $-2/\sqrt{|g|}$ times the variation of the action S_Ψ of dirac16complex with respect to g^{MN}. Unit: GeV^8. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\, \mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\hat{\nabla}$: The covariant derivative of the fold: with the Levi-Civita connection of the metric of the fold (for four-dimensional tensors of its metric $\hat{h}_{\mu\nu} = \mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the observable coordinates $x_1, \ldots, x_4$; for $\hat{\nabla}_a \mathcal{S}^a{}_b$ of the induced metric of the seven-dimensional fold). Status: STANDARD.
- $\mathcal{S}^a{}_b$: The thin source of the fold: the energy-momentum tensor concentrated on the fold, integrated across it, $\kappa\mathcal{S}^a{}_b = \int_{0^-}^{0^+} \mathrm{d}\sigma \sum_p \alpha_{(p)} \mathcal{E}_{(p)}{}^a{}_b$ (FD-11, the junction conditions); $\mathcal{S}^4{}_4$ is minus its energy density, $\mathcal{S}^1{}_1$ and $\mathcal{S}^5{}_5$ its

pressures along ordinary space and along the extra times (frame components along the fold, eight-dimensional normalisation: an energy per unit volume of the six other directions of the fold). Unit: GeV^7. Value: $\kappa\mathcal{S}^a{}_b = -\frac{32}{5}H\delta^a{}_b$ at the static block. Status: DERIVED.

- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- $\mathcal{T}_{\mathrm{bulk}}$: The energy-momentum tensor of all sources of the bulk (with the corrected weights dirac16complex alone, $\mathcal{T}_\Psi$; with the author's original weights also the stabiliser); $\mathcal{T}_{\mathrm{bulk}}{}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.

The conservation law at the fold: the covariant divergence of the thin source along the fold equals minus the jump of the energy and momentum flux of the bulk along σ, so the fold loses exactly what the bulk carries away. On the branch of frozen extra times the bulk carries no flux, and the matter of the fold is conserved by itself. Status: DERIVED.

$$\frac{1}{\sqrt{-\hat{h}}\,\mathcal{Z}}\Big(\partial_\mu\big[\sqrt{-\hat{h}}\,\mathcal{Z}\,\mathcal{K}_G\,G^{A\mu\nu}\big] + g_3\,f^{ABC}G^B_\mu\,\sqrt{-\hat{h}}\,\mathcal{Z}\,\mathcal{K}_G\,G^{C\mu\nu}\Big) = -g_3\,J_{\mathrm{c}}^{A\nu}, \qquad \mathcal{K}_G = 1 + \sum_X c_X^G\,\bar{\Psi}P_X\Psi,$$
$$J_{\mathrm{c}}^{A\nu} = \bar{Q}\,\hat{\gamma}^\nu t^A Q + \bar{u}_{\mathrm{R}}\,\hat{\gamma}^\nu t^A u_{\mathrm{R}} + \bar{d}_{\mathrm{R}}\,\hat{\gamma}^\nu t^A d_{\mathrm{R}} \tag{FE-16}$$

Symbols in (FE-16).

- $\hat{h}_{\mu\nu}$: The four-dimensional metric of the fold, $\hat{h}_{\mu\nu} = \mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the coordinates $x_1, \ldots, x_4$ (the induced metric at $\sigma = 0$ without the extra times); $\hat{h}^{\mu\nu}$ its inverse, $\hat{h}$ its determinant, $\sqrt{-\hat{h}} = \mathrm{e}^{3a_4}$; $\hat{h}^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4) = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z} = 1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+, +, +, -$); a repeated index is summed. Status: STANDARD.
- $\mathcal{K}_G$, $\mathcal{K}_W$, $\mathcal{K}_B$: The portal factors of the gauge kinetic terms, $\mathcal{K}_G = 1 + \sum_X c_X^G\bar{\Psi}P_X\Psi$ (likewise for W and B): the gauge couplings at the fold are divided by the square root of the portal factor. Unit: dimensionless. Value: 1 where dirac16complex vanishes. Status: DERIVED.
- G^A_μ, $G^A_{\mu\nu}$: The gluon field of $SU(3)_C$ on the fold ($A = 1, \ldots, 8$) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- g_3: The $SU(3)_C$ gauge coupling (the strong coupling g_s). Unit: dimensionless. Value: $g_3(m_Z) = 1.2177$ ($\alpha_s = 0.1180$). Status: STANDARD.
- f^{ABC}: The structure constants of $SU(3)$, $[t^A, t^B] = \mathrm{i}f^{ABC}t^C$. Unit: dimensionless. Status: STANDARD.
- B: A colour (adjoint SU(3)) index (as A). Status: STANDARD.
- C: A colour (adjoint SU(3)) index (as A). Status: STANDARD.
- $J_{\mathrm{c}}^{A\nu}$: The colour current of the quarks. Unit: GeV^3. Status: STANDARD.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c_X^G: The gluon-kinetic portal coupling of the half X of dirac16complex. Unit: GeV^{-7}. Status: INVENTED.
- Ψ, $\bar{\Psi}$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar{\Psi} = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- P_X: The chiral projector of the half X: $P_X = P_{\mathrm{L}}$ for $X = \mathrm{L}$, P_{R} for $X = \mathrm{R}$. Unit: dimensionless. Status: AUTHOR.
- Q, $\bar{Q}$: The left-handed quark doublets of the three generations, $Q = (u_{\mathrm{L}}, d_{\mathrm{L}})$ (colour triplets, hypercharge 1/6). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.

- $\hat{\gamma}^{\mu}$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^{\mu}$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^{\mu}, \hat{\gamma}^{\nu}\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_{\mu} = \hat{h}_{\mu\nu}\hat{\gamma}^{\nu}$. Unit: dimensionless. Status: STANDARD.
- t^{A}: The generators of $SU(3)$ on colour triplets, t^{A}: half the Gell-Mann matrices, with $\mathrm{tr}(t^{A}t^{B}) = \delta^{AB}/2$. Unit: dimensionless. Status: STANDARD.
- u_{R}: The right-handed up-type quarks (up, charm and top; singlets of $SU(2)_{L}$, hypercharge 2/3). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- d_{R}: The right-handed down-type quarks (down, strange and bottom; hypercharge $-1/3$). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.

The Yang-Mills equation of the gluons, with the factor of the gluon-kinetic portals and the colour current of the quarks as the source. Status: STANDARD (INVENTED portal factor).

$$\begin{aligned} &\frac{1}{\sqrt{-\hat{h}}\,\mathcal{Z}}\,\partial_{\mu}\Big[\sqrt{-\hat{h}}\,\mathcal{Z}\Big(\mathcal{K}_{B}\,B^{\mu\nu} + \sum_{X} d_{X}\,\mathcal{D}_{X}^{\mu\nu}\Big)\Big] = -g_{1}\,J_{Y}^{\nu}, \qquad \mathcal{K}_{B} = 1 + \sum_{X} c_{X}^{B}\,\bar{\Psi}P_{X}\Psi, \\ &J_{Y}^{\nu} = \sum_{\psi} Y_{\psi}\,\bar{\psi}\,\hat{\gamma}^{\nu}\psi + \frac{\mathrm{i}}{2}\big[(D^{\nu}\Phi)^{\dagger}\Phi - \Phi^{\dagger}D^{\nu}\Phi\big] \end{aligned} \tag{FE-18}$$

Symbols in (FE-18).

- $\hat{h}_{\mu\nu}$: The four-dimensional metric of the fold, $\hat{h}_{\mu\nu} = \mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the coordinates $x_1, \ldots, x_4$ (the induced metric at $\sigma = 0$ without the extra times); $\hat{h}^{\mu\nu}$ its inverse, $\hat{h}$ its determinant, $\sqrt{-\hat{h}} = \mathrm{e}^{3a_4}$; $\hat{h}^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4) = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3/V_{*}$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z} = 1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_{σ} with respect to σ; ∂_{M} and ∂_{μ} with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- $\mathcal{K}_{G}$, $\mathcal{K}_{W}$, $\mathcal{K}_{B}$: The portal factors of the gauge kinetic terms, $\mathcal{K}_{G} = 1 + \sum_{X} c_{X}^{G}\bar{\Psi}P_{X}\Psi$ (likewise for W and B): the gauge couplings at the fold are divided by the square root of the portal factor. Unit: dimensionless. Value: 1 where dirac16complex vanishes. Status: DERIVED.
- B_{μ}, $B_{\mu\nu}$: The hypercharge gauge field of $U(1)_{Y}$ on the fold and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- d_{X}: The dipole-portal coupling of the half X of dirac16complex with the hypercharge field strength. Unit: GeV^{-5}. Status: INVENTED.
- $\mathcal{D}_{X}^{\mu\nu}$: The dipole (antisymmetric tensor) density of the half X of dirac16complex at the fold, $\mathcal{D}_{X}^{\mu\nu} = \mathrm{i}\bar{\Psi}\Sigma^{\mu\nu}P_{X}\Psi$ (real, DERIVED). Unit: GeV^7. Status: DERIVED.
- g_1: The $U(1)_{Y}$ gauge coupling (usually g'; not normalised for grand unification). Unit: dimensionless. Value: $g_1(m_Z) = 0.3574$. Status: STANDARD.
- J_{Y}^{ν}: The hypercharge current of the fermions and of the Higgs doublet. Unit: GeV^3. Status: STANDARD.
- c_{X}^{B}: The B-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- Ψ, $\bar{\Psi}$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar{\Psi} = \Psi^{\dagger}\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- P_{X}: The chiral projector of the half X: $P_{X} = P_{\mathrm{L}}$ for $X = \mathrm{L}$, P_{R} for $X = \mathrm{R}$. Unit: dimensionless. Status: AUTHOR.
- ψ: The label of a Standard-Model fermion field in a sum: ψ runs over $Q, u_{\mathrm{R}}, d_{\mathrm{R}}, \ell, e_{\mathrm{R}}, \nu_{\mathrm{R}}$. Status: STANDARD.
- Y_{f}: The hypercharge of a field, the charge of its coupling $g_1 Y_f B_{\mu}$ to the hypercharge field in the covariant derivative (1/6, 2/3, $-1/3$, $-1/2$, -1, 0 for $Q, u_{\mathrm{R}}, d_{\mathrm{R}}, \ell, e_{\mathrm{R}}, \nu_{\mathrm{R}}$, 1/2 for Φ). Unit: dimensionless. Status: STANDARD.
- ψ, $\bar{\psi}$: A Standard-Model fermion field (a four-dimensional Weyl field of the fold with three generations, written

as a four-component spinor of fixed chirality); $\bar{\psi} = \psi^\dagger\beta$; ψ_f the fields of the kind f. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.

- $\hat{\gamma}^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^\mu$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^\mu, \hat{\gamma}^\nu\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_\mu = \hat{h}_{\mu\nu}\hat{\gamma}^\nu$. Unit: dimensionless. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC}G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.

The Maxwell equation of the hypercharge field, with the dipole density of dirac16complex added to the field strength and the hypercharge current of the fermions and the Higgs doublet as the source. Status: STANDARD (INVENTED portal terms).

$$\mathrm{i}\hat{\gamma}^\mu\big(D_\mu + \mathrm{i}\,\mathcal{V}^Q_\mu\big)Q = \mathcal{Y}^u\tilde{\Phi}\,u_\mathrm{R} + \mathcal{Y}^d\Phi\,d_\mathrm{R}, \qquad \mathcal{V}^f_\mu = \sum_X c^f_X\, j_{X\mu} \tag{FE-19}$$

Symbols in (FE-19).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^\mu$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^\mu, \hat{\gamma}^\nu\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_\mu = \hat{h}_{\mu\nu}\hat{\gamma}^\nu$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC}G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^f_\mu$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^f_\mu = \sum_X c^f_X j_{X\mu}$ ($\mathcal{V}^Q, \mathcal{V}^u, \ldots, \mathcal{V}^\nu$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- Q, $\bar{Q}$: The left-handed quark doublets of the three generations, $Q = (u_\mathrm{L}, d_\mathrm{L})$ (colour triplets, hypercharge 1/6). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^\nu$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- $\tilde{\Phi}$: The conjugate Higgs doublet, $\tilde{\Phi} = \mathrm{i}\tau^2\Phi^*$ (hypercharge $-1/2$). Unit: GeV. Status: STANDARD.
- u_R: The right-handed up-type quarks (up, charm and top; singlets of $SU(2)_L$, hypercharge 2/3). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- d_R: The right-handed down-type quarks (down, strange and bottom; hypercharge $-1/3$). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c^f_X: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c^ℓ_X and c^ν_X are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.
- j^μ_X: The convective current of the half X of dirac16complex, $j^\mu_X = \mathrm{i}(\bar{\Psi}P_X D^\mu\Psi - (D^\mu\bar{\Psi})P_X\Psi)$ (real and joining the half with itself, DERIVED). Unit: GeV^8. Status: DERIVED.
- f: The label of a kind of Standard-Model multiplet: $f \in \{Q, u, d, \ell, e, \nu\}$ (quark doublets, up-quark singlets,

down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.

$$\mathrm{i}\hat{\gamma}^{\mu}\big(D_{\mu}+\mathrm{i}\,\mathcal{V}^{u}_{\mu}\big)u_{\mathrm{R}}=(\mathcal{Y}^{u})^{\dagger}\tilde{\Phi}^{\dagger}Q \tag{FE-20}$$

Symbols in (FE-20).

- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^{\mu}$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^{\mu}$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^{\mu},\hat{\gamma}^{\nu}\}=-2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_{\mu}=\hat{h}_{\mu\nu}\hat{\gamma}^{\nu}$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1,x_2,x_3,x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_{μ}: The covariant derivative: on dirac16complex $D_M=\partial_M+\Omega_M$ (with $D_M\bar{\Psi}=\partial_M\bar{\Psi}-\bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_{μ} (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^{\mu}=\hat{h}^{\mu\nu}D_{\nu}$; on an adjoint field $(D_{\mu}X)^{A}=\partial_{\mu}X^{A}+g_3f^{ABC}G^{B}_{\mu}X^{C}$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^{f}_{\mu}$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^{f}_{\mu}=\sum_X c^{f}_{X}j_{X\mu}$ ($\mathcal{V}^{Q},\mathcal{V}^{u},\ldots,\mathcal{V}^{\nu}$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- u_{R}: The right-handed up-type quarks (up, charm and top; singlets of $SU(2)_L$, hypercharge 2/3). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- $\mathcal{Y}^{u}$, $\mathcal{Y}^{d}$, $\mathcal{Y}^{e}$, $\mathcal{Y}^{\nu}$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\tilde{\Phi}$: The conjugate Higgs doublet, $\tilde{\Phi}=\mathrm{i}\tau^2\Phi^{*}$ (hypercharge $-1/2$). Unit: GeV. Status: STANDARD.
- Q, $\bar{Q}$: The left-handed quark doublets of the three generations, $Q=(u_{\mathrm{L}},d_{\mathrm{L}})$ (colour triplets, hypercharge 1/6). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.

$$\mathrm{i}\hat{\gamma}^{\mu}\big(D_{\mu}+\mathrm{i}\,\mathcal{V}^{d}_{\mu}\big)d_{\mathrm{R}}=(\mathcal{Y}^{d})^{\dagger}\Phi^{\dagger}Q \tag{FE-21}$$

Symbols in (FE-21).

- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^{\mu}$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^{\mu}$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^{\mu},\hat{\gamma}^{\nu}\}=-2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_{\mu}=\hat{h}_{\mu\nu}\hat{\gamma}^{\nu}$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1,x_2,x_3,x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_{μ}: The covariant derivative: on dirac16complex $D_M=\partial_M+\Omega_M$ (with $D_M\bar{\Psi}=\partial_M\bar{\Psi}-\bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_{μ} (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^{\mu}=\hat{h}^{\mu\nu}D_{\nu}$; on an adjoint field $(D_{\mu}X)^{A}=\partial_{\mu}X^{A}+g_3f^{ABC}G^{B}_{\mu}X^{C}$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^{f}_{\mu}$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^{f}_{\mu}=\sum_X c^{f}_{X}j_{X\mu}$ ($\mathcal{V}^{Q},\mathcal{V}^{u},\ldots,\mathcal{V}^{\nu}$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- d_{R}: The right-handed down-type quarks (down, strange and bottom; hypercharge $-1/3$). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- $\mathcal{Y}^{u}$, $\mathcal{Y}^{d}$, $\mathcal{Y}^{e}$, $\mathcal{Y}^{\nu}$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.

- Q, $\bar{Q}$: The left-handed quark doublets of the three generations, $Q = (u_{\mathrm{L}}, d_{\mathrm{L}})$ (colour triplets, hypercharge 1/6). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.

$$\mathrm{i}\hat{\gamma}^{\mu}\big(D_{\mu} + \mathrm{i}\,\mathcal{V}^{\ell}_{\mu}\big)\ell = \mathcal{Y}^{e}\Phi\, e_{\mathrm{R}} + \mathcal{Y}^{\nu}\tilde{\Phi}\,\nu_{\mathrm{R}} \tag{FE-22}$$

Symbols in (FE-22).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^{\mu}$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^{\mu}$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^{\mu}, \hat{\gamma}^{\nu}\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_{\mu} = \hat{h}_{\mu\nu}\hat{\gamma}^{\nu}$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_{μ}: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_{μ} (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^{\mu} = \hat{h}^{\mu\nu}D_{\nu}$; on an adjoint field $(D_{\mu}X)^{A} = \partial_{\mu}X^{A} + g_3 f^{ABC}G^{B}_{\mu}X^{C}$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^{f}_{\mu}$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^{f}_{\mu} = \sum_X c^{f}_{X} j_{X\mu}$ ($\mathcal{V}^{Q}, \mathcal{V}^{u}, \ldots, \mathcal{V}^{\nu}$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- ℓ, $\bar{\ell}$: The left-handed lepton doublets of the three generations, $\ell = (\nu_{\mathrm{L}}, e_{\mathrm{L}})$ (hypercharge $-1/2$). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- $\mathcal{Y}^{u}$, $\mathcal{Y}^{d}$, $\mathcal{Y}^{e}$, $\mathcal{Y}^{\nu}$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- e_{R}: The right-handed electron, muon and tau (singlets, hypercharge -1). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- $\tilde{\Phi}$: The conjugate Higgs doublet, $\tilde{\Phi} = \mathrm{i}\tau^2\Phi^*$ (hypercharge $-1/2$). Unit: GeV. Status: STANDARD.
- ν_{R}: The three right-handed neutrinos (gauge singlets). Unit: $\mathrm{GeV}^{3/2}$. Status: ASSUMED.

$$\mathrm{i}\hat{\gamma}^{\mu}\big(D_{\mu} + \mathrm{i}\,\mathcal{V}^{e}_{\mu}\big)e_{\mathrm{R}} = (\mathcal{Y}^{e})^{\dagger}\Phi^{\dagger}\ell \tag{FE-23}$$

Symbols in (FE-23).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^{\mu}$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^{\mu}$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^{\mu}, \hat{\gamma}^{\nu}\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_{\mu} = \hat{h}_{\mu\nu}\hat{\gamma}^{\nu}$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_{μ}: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_{μ} (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^{\mu} = \hat{h}^{\mu\nu}D_{\nu}$; on an adjoint field $(D_{\mu}X)^{A} = \partial_{\mu}X^{A} + g_3 f^{ABC}G^{B}_{\mu}X^{C}$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^{f}_{\mu}$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^{f}_{\mu} = \sum_X c^{f}_{X} j_{X\mu}$ ($\mathcal{V}^{Q}, \mathcal{V}^{u}, \ldots, \mathcal{V}^{\nu}$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- e_{R}: The right-handed electron, muon and tau (singlets, hypercharge -1). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- $\mathcal{Y}^{u}$, $\mathcal{Y}^{d}$, $\mathcal{Y}^{e}$, $\mathcal{Y}^{\nu}$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- ℓ, $\bar{\ell}$: The left-handed lepton doublets of the three generations, $\ell = (\nu_{\mathrm{L}}, e_{\mathrm{L}})$ (hypercharge $-1/2$). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.

$$\mathrm{i}\hat{\gamma}^{\mu}\big(D_{\mu}+\mathrm{i}\,\mathcal{V}_{\mu}^{\nu}\big)\nu_{\mathrm{R}}=(\mathcal{Y}^{\nu})^{\dagger}\tilde{\Phi}^{\dagger}\ell \tag{FE-24}$$

Symbols in (FE-24).

- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^{\mu}$: The four-dimensional complex 4×4 Dirac matrices of the fold: $\hat{\gamma}^{(a)}$ along the frame directions of the fold and $\hat{\gamma}^{\mu}$ along its coordinates (the frame matrices contracted with the inverse frame field of the fold), $\{\hat{\gamma}^{\mu},\hat{\gamma}^{\nu}\}=-2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_{\mu}=\hat{h}_{\mu\nu}\hat{\gamma}^{\nu}$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1,x_2,x_3,x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_{μ}: The covariant derivative: on dirac16complex $D_M=\partial_M+\Omega_M$ (with $D_M\bar{\Psi}=\partial_M\bar{\Psi}-\bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_{μ} (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^{\mu}=\hat{h}^{\mu\nu}D_{\nu}$; on an adjoint field $(D_{\mu}X)^{A}=\partial_{\mu}X^{A}+g_3f^{ABC}G_{\mu}^{B}X^{C}$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}_{\mu}^{f}$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}_{\mu}^{f}=\sum_X c_X^f j_{X\mu}$ ($\mathcal{V}^{Q},\mathcal{V}^{u},\ldots,\mathcal{V}^{\nu}$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- ν_{R}: The three right-handed neutrinos (gauge singlets). Unit: $\mathrm{GeV}^{3/2}$. Status: ASSUMED.
- $\mathcal{Y}^{u}$, $\mathcal{Y}^{d}$, $\mathcal{Y}^{e}$, $\mathcal{Y}^{\nu}$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\tilde{\Phi}$: The conjugate Higgs doublet, $\tilde{\Phi}=\mathrm{i}\tau^2\Phi^{*}$ (hypercharge $-1/2$). Unit: GeV. Status: STANDARD.
- ℓ, $\bar{\ell}$: The left-handed lepton doublets of the three generations, $\ell=(\nu_{\mathrm{L}},e_{\mathrm{L}})$ (hypercharge $-1/2$). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.

The Dirac equations of the quark doublets (FE-19), the up-quark and down-quark singlets (FE-20, FE-21), the lepton doublets (FE-22), the charged-lepton singlets (FE-23) and the right-handed neutrinos (FE-24): every fermion sees the gauge fields, the Higgs doublet through its Yukawa matrix and dirac16complex through the flavour-blind vector potential of the current portals; the right-handed neutrinos see no gauge field. Status: STANDARD (INVENTED potentials).

The table states, for every field equation, how the solution satisfies it.

equation	how it is solved	residual	status
bulk gravity FE-1, order 0	the bulk equations of the author's metric are solved exactly (FE-29, S-1, S-2, FE-66): on the branch of frozen extra times the bulk is a vacuum for every expansion history, the late universe and the inflation of the fold included	0 (exact polynomials)	DERIVED
bulk gravity FE-1, higher orders in ϵ_{b}	S-5 to S-7: no linear part around the static block, for every perturbation of the metric; static deformations grow like $\mathrm{e}^{3H\lvert\sigma\rvert}$; on the expanding branch no time-kinetic term	the response of the strongly coupled bulk to sources that depend on σ (the bound states χ among them) is not computed	DERIVED, OPEN
junction FE-4, FE-5: time and space components	FE-33, FE-59: the Friedmann equations of general relativity with $8\pi G_{\mathrm{N}}=\kappa_7/V_{*}$	0 (exact for frozen extra times)	DERIVED

equation	how it is solved	residual	status
junction FE-4: extra-time component	FE-65, FE-72, S-9: the volume of the extra times held fixed	the mismatch $\Delta_{\mathrm{r}} = 0.793\rho_{\mathrm{c}}^{(0)}$ today (with the assumed pressure $-\rho_{\mathrm{b}}$ of the baryons along the extra times; 0.842 or 0.891 with 0 or $+\rho_{\mathrm{b}}$), carried by the pressure q_{hold} of the fold fluid; in an early universe that expands faster than H the holding pressure is enormous (S-3)	DERIVED, ASSUMED, OPEN
conservation at the fold FE-11	the fold loses exactly the energy that the bulk flux carries away; no flux on the frozen branch	0 (an identity for every a_4, b_4)	DERIVED
dirac16complex FE-12 to FE-14, Kohn-Sham FE-41	the Kohn-Sham states (S-14 to S-19), lifted to sixteen components	Kohn-Sham equation 2×10^{-4} (central differences), fold matching 4×10^{-9}, self-consistency 2×10^{-7}; Krein charges -1 to 4×10^{-16}	DERIVED
Higgs doublet FE-15	S-10: the ad hoc vacuum as a gauge rotation	the radial residual $\mathcal{R}_{\mathrm{rad}}$ (table of S-10), zero once $\mathcal{A} = 0$	DERIVED
$SU(2)$ gauge fields FE-17	S-10: the pure-gauge $W_{x_4}^2$	0: the isospin charge density vanishes exactly	DERIVED
gluons FE-16 and hypercharge field FE-18	no mean field strengths (the gluon condensate enters FE-39 as a mean value)	0	DERIVED
quarks and leptons FE-19 to FE-24	no mean fermion fields; the plasma and the neutrinos are treated statistically (FE-48 to FE-62)	0 in the vacuum	STANDARD
neutrino flavours FE-48 to FE-50	the Sun and the supernova envelope (S-20, S-21)	halving the step changes P_{ee} by less than 10^{-6} (Sun) and 10^{-9} (envelope)	DERIVED
fast instability FE-51	S-22	the dispersion relation holds to 10^{-14} at the computed frequencies	DERIVED
relic density FE-61	S-24	relative tolerance 10^{-8} of the implicit integration	DERIVED
cosmological perturbations FE-54, FE-55	CAMB with the content of UM-3	the accuracy of CAMB (about 10^{-3} in C_l); background and ruler checked independently to 10^{-6}	STANDARD, DERIVED

11 The three neutrino flavours

The author asks for a correct flavour conversion in UM-3, because the flavour conversion of neutrinos shapes the explosion of massive stars (Gogilashvili and Tamborra, "Neutrino Flavor Conversion Shapes the Rate of Failed Core-collapse Supernovae", arXiv:2605.16504). This section solves the flavour equations of UM-3 in the Sun and in core-collapse supernovae.

$$
\begin{aligned}
&\mathrm{i}\big(\partial_{x_4} + \hat{v}\cdot\vec{\nabla}\big)\varrho_{\vec{p}} = \big[\mathsf{H}_{\vec{p}},\ \varrho_{\vec{p}}\big] + \mathrm{i}\,\mathcal{C}_{\vec{p}}, \qquad \mathrm{i}\big(\partial_{x_4} + \hat{v}\cdot\vec{\nabla}\big)\bar{\varrho}_{\vec{p}} = \big[\bar{\mathsf{H}}_{\vec{p}},\ \bar{\varrho}_{\vec{p}}\big] + \mathrm{i}\,\bar{\mathcal{C}}_{\vec{p}}, \\
&\mathsf{H}_{\vec{p}} = \frac{v_{\mathrm{loc}}^2}{v^2}\,\frac{U_{\mathrm{PMNS}}\,\mathsf{M}^2\,U_{\mathrm{PMNS}}^{\dagger}}{2E} + \mathsf{V}_{\mathrm{m}} + \mathsf{V}_{\nu\nu} + V_{\Psi}\mathbf{1}, \qquad \mathsf{M}^2 = \mathrm{diag}\big(m_1^2,\ m_2^2,\ m_3^2\big), \\
&\bar{\mathsf{H}}_{\vec{p}} = -\frac{v_{\mathrm{loc}}^2}{v^2}\,\frac{U_{\mathrm{PMNS}}\,\mathsf{M}^2\,U_{\mathrm{PMNS}}^{\dagger}}{2E} + \mathsf{V}_{\mathrm{m}} + \mathsf{V}_{\nu\nu} + V_{\Psi}\mathbf{1}
\end{aligned}
\tag{FE-48}
$$

Symbols in (FE-48).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\hat{v}$, $\hat{v}^\mu$: The direction of motion of a neutrino: the unit vector $\hat{v} = \vec{p}/|\vec{p}|$ ($\hat{v}_{\vec{p}}$, $\hat{v}_{\vec{q}}$ for the momenta $\vec{p}$, $\vec{q}$), and the four-vector $\hat{v}^\mu = (1, \hat{v})$ in the order $(x_4; x_1, x_2, x_3)$; $\mathrm{d}^2\hat{v}$ is the element of solid angle of the directions. Unit: dimensionless. Status: STANDARD.
- $\vec{\nabla}$: The gradient along ordinary space, $\vec{\nabla} = (\partial_{x_1}, \partial_{x_2}, \partial_{x_3})$ (in the local frame of the medium). Unit: 1/length. Status: STANDARD.
- $\varrho_{\vec{p}}$, $\bar{\varrho}_{\vec{p}}$: The 3×3 flavour density matrices of the neutrinos and of the antineutrinos of momentum $\vec{p}$ in the flavour basis (electron, muon, tau): the diagonal entries are the occupation numbers of the three flavours, the off-diagonal entries the coherence between the flavours; $\bar{\varrho}$ is defined with the flavour indices in the reversed order, so that it obeys an equation of the same form. Unit: dimensionless. Status: STANDARD.
- $\vec{p}$: The momentum of a neutrino (three components along ordinary space, in the rest frame of the medium); $|\vec{p}| = E$ for a neutrino of negligible mass. Unit: GeV. Status: STANDARD.
- $\mathsf{H}_{\vec{p}}$, $\bar{\mathsf{H}}_{\vec{p}}$: The flavour Hamiltonians (3×3 Hermitian matrices) of the neutrinos and of the antineutrinos of momentum $\vec{p}$. Unit: GeV. Status: STANDARD.
- $\mathcal{C}_{\vec{p}}$, $\bar{\mathcal{C}}_{\vec{p}}$: The collision terms ($3 \times 3$ matrices) of the neutrinos and the antineutrinos of momentum $\vec{p}$: the rates of change of $\varrho_{\vec{p}}$ and $\bar{\varrho}_{\vec{p}}$ by emission, absorption, scattering and pair processes with the matter. Unit: GeV. Status: STANDARD.
- v_{loc}: The local value of the electroweak vacuum, $v_{\mathrm{loc}}^2 = 2\langle\Phi^\dagger\Phi\rangle$: it sets the local masses of the quarks and leptons (each mass times v_{loc}/v) and the local Fermi constant; $v_{\mathrm{loc}} = v$ in the Standard-Model vacuum. Unit: GeV. Value: v in the late universe outside dense dirac16complex. Status: DERIVED.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_{\mathrm{F}})^{-1/2}$ with the Fermi constant $G_{\mathrm{F}} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- U_{PMNS}: The lepton mixing matrix (Pontecorvo-Maki-Nakagawa-Sakata) of the three Dirac neutrinos; its entries $(U_{\mathrm{PMNS}})_{\alpha i}$ connect the flavour α with the mass state i. Unit: dimensionless. Value: $\theta_{12} = 33.40°$, $\theta_{23} = 49.14°$, $\theta_{13} = 8.54°$ (NuFIT 5.2, November 2022, normal ordering, the fit without the Super-Kamiokande atmospheric data); the Dirac phase is set to zero (ASSUMED; the fit prefers 197°). Status: STANDARD, ASSUMED (the phase).
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- M^2: The diagonal matrix of the squared masses of the three neutrino mass states, $\mathsf{M}^2 = \mathrm{diag}(m_1^2, m_2^2, m_3^2)$ (M appears only squared). Unit: GeV^2 (of M^2). Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- V_{m}: The matter potential of the neutrinos (a 3×3 matrix in flavour space) from their forward scattering on the electrons, positrons and neutrons of the medium. Unit: GeV. Status: STANDARD.
- $\mathsf{V}_{\nu\nu}$: The neutrino-neutrino potential (a 3×3 matrix in flavour space) from the forward scattering of a neutrino on the other neutrinos and antineutrinos of the medium; it depends on their flavour content and their directions. Unit: GeV. Status: STANDARD.
- V_Ψ: The potential energy that dirac16complex gives every neutrino through the current portal of the lepton doublets, $V_\Psi = \mathcal{V}^\ell_{x_4}$ (the same for the three flavours). Unit: GeV. Status: DERIVED.
- $\mathbf{1}$, $\mathbf{1}_8$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_8$ is the 8×8 identity matrix. Unit: dimensionless. Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- $m_u, \ldots, m_3$: The masses of the quarks (m_u, m_c, m_t; m_d, m_s, m_b), of the charged leptons (m_e, m_μ, m_τ) and of the three neutrino mass states (m_1, m_2, m_3, Dirac masses). Unit: GeV. Value: measured values; $\Delta m^2_{21} = 7.41 \times 10^{-5}$ eV^2, $|\Delta m^2_{31}| = 2.511 \times 10^{-3}$ eV^2 in the computations of UM-3 (the global fit NuFIT 5.2 of the oscillation data, November 2022), 7.49×10^{-5} and 2.513×10^{-3} eV^2 in NuFIT 6.0 (2024); the lightest mass $m_1 = 0$ in the

computations, below 23 meV by DESI DR2 with the CMB and NuFIT 6.0. Status: STANDARD.

$$
\begin{aligned}
&\mathsf{V}_{\mathrm{m}} = \sqrt{2}\,G_{\mathrm{F,loc}}\,\mathrm{diag}\big(n_e - n_{\bar{e}},\ 0,\ 0\big) - \frac{G_{\mathrm{F,loc}}}{\sqrt{2}}\,n_n\mathbf{1}, \qquad \mathsf{V}_{\nu\nu} = \sqrt{2}\,G_{\mathrm{F,loc}}\int \frac{\mathrm{d}^3\vec{q}}{(2\pi)^3}\big(1 - \hat{v}_{\vec{p}}\cdot\hat{v}_{\vec{q}}\big)\big(\varrho_{\vec{q}} - \bar{\varrho}_{\vec{q}}\big),\\
&V_\Psi = \mathcal{V}^{\ell}_{x_4} = \sum_X c^{\ell}_X\, j_{X\,x_4}, \qquad G_{\mathrm{F,loc}} = \frac{1}{\sqrt{2}\,v^2_{\mathrm{loc}}},\\
&v^2_{\mathrm{loc}} = 2\langle\Phi^\dagger\Phi\rangle, \qquad v^2_{\mathrm{loc}} = v^2 + \mathcal{A}^2 \ \text{ (the ad hoc vacuum)},\\
&v^2_{\mathrm{loc}} = v^2 - \frac{c_\Phi}{\lambda}\,\langle\bar{\Psi}P_{\mathrm{L}}\Psi\rangle \ \text{ (the relaxed vacuum with dirac16complex)}
\end{aligned}
\tag{FE-49}
$$

Symbols in (FE-49).

- V_{m}: The matter potential of the neutrinos (a 3×3 matrix in flavour space) from their forward scattering on the electrons, positrons and neutrons of the medium. Unit: GeV. Status: STANDARD.
- $G_{\mathrm{F,loc}}$: The local Fermi constant, $G_{\mathrm{F,loc}} = 1/(\sqrt{2}\,v^2_{\mathrm{loc}})$ ($= G_{\mathrm{F}}$ where $v_{\mathrm{loc}} = v$). Unit: GeV^{-2}. Status: DERIVED.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- n_e: The number density of electrons (bound and free; in a supernova all are free, in the recombining universe $x_e n_e$ is the density of the free electrons). Unit: GeV^3 (cm^{-3}). Status: STANDARD.
- $n_{\bar{e}}$: The number density of positrons. Unit: GeV^3. Status: STANDARD.
- n_n: The number density of neutrons. Unit: GeV^3. Status: STANDARD.
- $\mathbf{1}$, $\mathbf{1}_8$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space); $\mathbf{1}_8$ is the 8×8 identity matrix. Unit: dimensionless. Status: STANDARD.
- $\mathsf{V}_{\nu\nu}$: The neutrino-neutrino potential (a 3×3 matrix in flavour space) from the forward scattering of a neutrino on the other neutrinos and antineutrinos of the medium; it depends on their flavour content and their directions. Unit: GeV. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $\vec{q}$: The momentum of another neutrino of the medium (the integration variable of the neutrino-neutrino potential); $\mathrm{d}^3\vec{q}$ its volume element. Unit: GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $\hat{v}$, $\hat{v}^\mu$: The direction of motion of a neutrino: the unit vector $\hat{v} = \vec{p}/|\vec{p}|$ ($\hat{v}_{\vec{p}}$, $\hat{v}_{\vec{q}}$ for the momenta $\vec{p}$, $\vec{q}$), and the four-vector $\hat{v}^\mu = (1, \hat{v})$ in the order $(x_4; x_1, x_2, x_3)$; $\mathrm{d}^2\hat{v}$ is the element of solid angle of the directions. Unit: dimensionless. Status: STANDARD.
- $\vec{p}$: The momentum of a neutrino (three components along ordinary space, in the rest frame of the medium); $|\vec{p}| = E$ for a neutrino of negligible mass. Unit: GeV. Status: STANDARD.
- $\varrho_{\vec{p}}$, $\bar{\varrho}_{\vec{p}}$: The 3×3 flavour density matrices of the neutrinos and of the antineutrinos of momentum $\vec{p}$ in the flavour basis (electron, muon, tau): the diagonal entries are the occupation numbers of the three flavours, the off-diagonal entries the coherence between the flavours; $\bar{\varrho}$ is defined with the flavour indices in the reversed order, so that it obeys an equation of the same form. Unit: dimensionless. Status: STANDARD.
- V_Ψ: The potential energy that dirac16complex gives every neutrino through the current portal of the lepton doublets, $V_\Psi = \mathcal{V}^{\ell}_{x_4}$ (the same for the three flavours). Unit: GeV. Status: DERIVED.
- $\mathcal{V}^f_\mu$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^f_\mu = \sum_X c^f_X j_{X\mu}$ ($\mathcal{V}^Q, \mathcal{V}^u, \ldots, \mathcal{V}^\nu$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c^f_X: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c^ℓ_X and c^ν_X are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.

- j^μ_X: The convective current of the half X of dirac16complex, $j^\mu_X = \mathrm{i}(\bar\Psi P_X D^\mu\Psi - (D^\mu\bar\Psi)P_X\Psi)$ (real and joining the half with itself, DERIVED). Unit: GeV^8. Status: DERIVED.
- v_{loc}: The local value of the electroweak vacuum, $v^2_{\mathrm{loc}} = 2\langle\Phi^\dagger\Phi\rangle$: it sets the local masses of the quarks and leptons (each mass times v_{loc}/v) and the local Fermi constant; $v_{\mathrm{loc}} = v$ in the Standard-Model vacuum. Unit: GeV. Value: v in the late universe outside dense dirac16complex. Status: DERIVED.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_{\mathrm{F}})^{-1/2}$ with the Fermi constant $G_{\mathrm{F}} = 1.1663788\times10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- $\mathcal{A}$: The author's ad hoc upper component of the Higgs vacuum, almightyAdHoc, $\mathcal{A}(x_4) = v_{\mathcal{A}}\mathrm{e}^{-2a_{40}x_4}$. Unit: GeV. Status: AUTHOR.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: GeV$^{7/2}$. Status: AUTHOR.
- P_{L}: The author's left-handed chiral projector, $P_{\mathrm{L}} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.

The flavour equations of the neutrinos of momentum $\vec p$ and the potentials (FE-48, FE-49). Two facts make them those of the Standard Model with three Dirac neutrinos wherever $v_{\mathrm{loc}} = v$: the portal potential V_Ψ is the same for the three flavours and drops out of every commutator, and the Higgs portal moves v_{loc} by only $\Delta v^2/v^2 = -\mathfrak{g}_{h\chi}\rho_\chi/(\lambda v^3\omega_{\mathrm{b}}) \approx -10^{-50}\,\mathfrak{g}_{h\chi}$ at the local dark-matter density $\rho_\chi = 0.3$ GeV/cm^3 for $\omega_{\mathrm{b}} = 100$ GeV (the value grows as $1/\omega_{\mathrm{b}}$ for lighter χ; $\mathfrak{g}_{h\chi}$ is the coupling of the Higgs boson to χ of FE-47, negative where the Higgs portal binds, so the gas raises v_{loc}). Status: DERIVED (`um3_neutrino.py`, N1: $V_\Psi\mathbf{1}$ changes the evolution by at most 7×10^{-13}, a change of v_{loc} leaves the mixing matrix unchanged to 10^{-15} and scales the masses by v_{loc}/v). For a single neutrino along its path (no collisions, no neutrino-neutrino potential) FE-48 is a Schrodinger equation $\mathrm{i}\,\mathrm{d}\psi_\nu/\mathrm{d}r = \mathsf{H}\psi_\nu$ for the flavour amplitudes ψ_ν; for antineutrinos the vacuum term has U^*_{PMNS} and the matter potential the opposite sign (the convention of FE-48, with the flavour indices of $\bar\varrho$ reversed, gives the same probabilities). It is integrated here with the fourth-order commutator-free Magnus method (the exponentials of the Hamiltonian at the two Gauss points of each step, exact for a constant density), and every result is recomputed with half the step.

$$\tan 2\theta_{\mathrm{m}} = \frac{\sin 2\theta}{\cos 2\theta - 2EV_{\mathrm{CC}}/\Delta m^2}, \qquad V_{\mathrm{CC}} = \sqrt{2}\,G_{\mathrm{F,loc}}\,n_e, \qquad n_{e,\mathrm{res}} = \frac{\Delta m^2\cos 2\theta}{2\sqrt{2}\,G_{\mathrm{F,loc}}\,E},$$
$$P_{\mathrm{LZ}} = \exp\Big(-\frac{\pi}{2}\,\frac{\Delta m^2\sin^2 2\theta}{2E\cos 2\theta}\,\Big|\frac{\mathrm{d}\ln n_e}{\mathrm{d}r}\Big|^{-1}_{\mathrm{res}}\Big) \tag{FE-50}$$

Symbols in (FE-50).

- θ_{m}: The mixing angle of two neutrino flavours in matter. Unit: dimensionless. Status: STANDARD.
- θ: The vacuum mixing angle of two neutrino flavours (in the two-flavour description of a resonance, one of the three angles $\theta_{12}, \theta_{13}, \theta_{23}$ of U_{PMNS}). Unit: dimensionless. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- V_{CC}: The charged-current matter potential of the electron neutrinos (forward scattering on electrons through W exchange). Unit: GeV. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m^2_{ij} = m_i^2 - m_j^2$ the squared-mass difference of the

states i and j (Δm^2_{21}, Δm^2_{31}, Δm^2_{32}); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by $v^2_{\rm loc}/v^2$ where this differs from 1. Unit: GeV. Status: STANDARD.
- $G_{\rm F,loc}$: The local Fermi constant, $G_{\rm F,loc} = 1/(\sqrt{2}\, v^2_{\rm loc})$ ($= G_{\rm F}$ where $v_{\rm loc} = v$). Unit: GeV^{-2}. Status: DERIVED.
- n_e: The number density of electrons (bound and free; in a supernova all are free, in the recombining universe $x_e n_e$ is the density of the free electrons). Unit: GeV^3 (cm^{-3}). Status: STANDARD.
- $n_{e,\rm res}$: The electron density at which the flavour conversion in matter is resonant (the Mikheyev-Smirnov-Wolfenstein resonance). Unit: GeV^3. Status: STANDARD.
- $P_{\rm LZ}$: The probability that a neutrino jumps between the eigenstates of matter at the resonance (Landau-Zener); $P_{\rm LZ} = 0$ is an adiabatic resonance. Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as dσ, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- r: The distance from the centre of the star along the path of a neutrino. Unit: length. Status: STANDARD.
- res: The label of a quantity evaluated at the resonance, where $n_e = n_{e,\rm res}$. Status: STANDARD.

The mixing angle in matter, the resonance density and the Landau-Zener jump probability (FE-50). Status: STANDARD (`um3_neutrino.py`, N2: the formula for $\theta_{\rm m}$ agrees with an exact diagonalisation to 10^{-16}, the Landau-Zener formula with an exact integration through a linear profile to 10^{-6}).

11.1 The Sun

$$\mathsf{H}_2 = \frac{1}{2}\begin{pmatrix} -\frac{\Delta m^2_{21}}{2E}\cos 2\theta_{12} + \cos^2\theta_{13}\, V_{\rm CC} & \frac{\Delta m^2_{21}}{2E}\sin 2\theta_{12} \\ \frac{\Delta m^2_{21}}{2E}\sin 2\theta_{12} & \frac{\Delta m^2_{21}}{2E}\cos 2\theta_{12} - \cos^2\theta_{13}\, V_{\rm CC}\end{pmatrix}, \qquad n_e(r) = 245\, N_{\rm A}\, \mathrm{e}^{-10.54\, r/R_\odot}\ \mathrm{cm}^{-3},$$

$$\mathsf{P}_{\rm ee} = \cos^4\theta_{13}\, \mathsf{P}_2 + \sin^4\theta_{13}, \qquad \mathsf{P}_2 = \frac{1}{2} + \left(\frac{1}{2} - \mathsf{P}_{\rm c}\right)\cos 2\theta_{12}\cos 2\theta_{\rm m}(r_{\rm p}),$$

$$\mathsf{P}_{\rm c} = \frac{\mathrm{e}^{-2\pi r_\odot \frac{\Delta m^2_{21}}{2E}\sin^2\theta_{12}} - \mathrm{e}^{-2\pi r_\odot \frac{\Delta m^2_{21}}{2E}}}{1 - \mathrm{e}^{-2\pi r_\odot \frac{\Delta m^2_{21}}{2E}}}, \qquad r_\odot = \frac{R_\odot}{10.54} \tag{S-20}$$

Symbols in (S-20).

- H_2: The two-flavour Hamiltonian of the electron neutrinos in the Sun (the mass states 1 and 2 with the potential $\cos^2\theta_{13} V_{\rm CC}$; the third state decouples). Unit: GeV (or 1/km). Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m^2_{ij} = m_i^2 - m_j^2$ the squared-mass difference of the states i and j (Δm^2_{21}, Δm^2_{31}, Δm^2_{32}); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by $v^2_{\rm loc}/v^2$ where this differs from 1. Unit: GeV. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- θ_{12}, θ_{13}, θ_{23}: The three mixing angles of $U_{\rm PMNS}$ (its fourth parameter is the Dirac phase, which does not enter the electron-flavour results of UM-3). Unit: dimensionless. Value: 33.40°, 8.54°, 49.14° in the computations of UM-3 ($\sin^2\theta_{12} = 0.303$, $\sin^2\theta_{13} = 0.02203$, $\sin^2\theta_{23} = 0.572$: NuFIT 5.2, November 2022, normal ordering, the fit without the Super-Kamiokande atmospheric data); 33.71°, 8.56°, 43.28° in NuFIT 6.0 (2024, with the Super-Kamiokande atmospheric data; $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$). Status: STANDARD.
- $V_{\rm CC}$: The charged-current matter potential of the electron neutrinos (forward scattering on electrons through W exchange). Unit: GeV. Status: STANDARD.
- n_e: The number density of electrons (bound and free; in a supernova all are free, in the recombining universe $x_e n_e$ is the density of the free electrons). Unit: GeV^3 (cm^{-3}). Status: STANDARD.
- r: The distance from the centre of the star along the path of a neutrino. Unit: length. Status: STANDARD.
- $N_{\rm A}$: Avogadro's number: $245 N_{\rm A}$ cm^{-3} is 245 moles of electrons per cubic centimetre. Unit: per mole. Value: $6.02214076 \times 10^{23}$. Status: STANDARD.

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- $R_\odot$: The radius of the Sun. Unit: length. Value: 6.957×10^5 km. Status: STANDARD.
- P_{ee}: The survival probability of electron neutrinos: the probability that an electron neutrino is detected as an electron neutrino. Unit: dimensionless. Status: DERIVED.
- P_2: The survival probability of the two-flavour problem of the mass states 1 and 2 (S-20). Unit: dimensionless. Status: DERIVED.
- P_{c}: The crossing probability: the probability that a neutrino jumps from one eigenstate of matter to the other on its way through the Sun (exact for an exponential profile, Petcov). Unit: dimensionless. Status: STANDARD.
- θ_{m}: The mixing angle of two neutrino flavours in matter. Unit: dimensionless. Status: STANDARD.
- r_{p}: The radius at which the solar neutrinos of a given reaction are produced. Unit: length. Value: $0.05R_\odot$ to $0.10R_\odot$. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $r_\odot$: The scale height of the electron density of the Sun, $r_\odot = R_\odot/10.54$. Unit: length. Value: 6.6×10^4 km. Status: STANDARD.

The solar neutrinos: the third mass state decouples (it oscillates too fast and mixes only through θ_{13}), so the electron neutrinos evolve with the two-flavour Hamiltonian H_2 (with the potential $\cos^2\theta_{13}V_{\mathrm{CC}}$) through the electron density of the Sun in its exponential approximation (Bahcall), from the radius r_{p} of their production to the surface; the phases average out over the production region and the distance to the Earth, so the survival probability P_{ee} follows from the probabilities of the mass states at the surface. The second line is the adiabatic solution with the exact crossing probability P_{c} of an exponential profile (Petcov). Status: STANDARD (the equations), DERIVED (the solution below).

source	E (MeV)	$r_{\mathrm{p}}/R_\odot$	P_{ee} (exact integration)	P_{ee} (S-20)	Borexino
pp	0.267	0.10	0.5454	0.5454	0.57 ± 0.09
^{7}Be	0.862	0.06	0.5126	0.5126	0.53 ± 0.05
pep	1.44	0.10	0.5083	0.5083	0.43 ± 0.11
^{8}B	5.0	0.05	0.3362	0.3362	
^{8}B	8.1	0.05	0.3080	0.3080	0.37 ± 0.08
^{8}B	10.0	0.05	0.3017	0.3017	
^{8}B	15.0	0.05	0.2951	0.2951	

The survival probability of solar electron neutrinos in UM-3 (with $\theta_{12} = 33.40°$, $\theta_{13} = 8.54°$, $\Delta m^2_{21} = 7.41 \times 10^{-5}$ eV^2): the exact integration and the adiabatic formula agree to 10^{-6}, $\mathsf{P}_{\mathrm{c}} < 10^{-10}$ at every energy (the Sun is adiabatic), and halving the step changes nothing at the level 10^{-6}. At low energy P_{ee} approaches the vacuum average $\cos^4\theta_{13}(1-\frac{1}{2}\sin^2 2\theta_{12})+\sin^4\theta_{13} = 0.553$, at high energy the full matter conversion $\cos^4\theta_{13}\sin^2\theta_{12}+\sin^4\theta_{13} = 0.290$; the Borexino measurements (Nature 562, 505 (2018), with the high-metallicity solar model) lie within one standard deviation. Status: DERIVED (`um3_flavour.py`, F1).

11.2 The envelope of a supernova

$$\rho_{\mathrm{b}}(r) = 10^4\ \mathrm{g/cm^3}\Big(\frac{r}{10^9\ \mathrm{cm}}\Big)^{-3}, \qquad n_e = \frac{Y_{\mathrm{e}}\,\rho_{\mathrm{b}}}{m_{\mathrm{u}}}, \qquad \gamma_{\mathrm{LZ}} = \frac{\Delta m^2 \sin^2 2\theta}{2E\cos 2\theta}\,\frac{r_{\mathrm{res}}}{3} \tag{S-21}$$

Symbols in (S-21).

- ρ_{b}: The (rest-mass) energy density of the baryons: in a supernova the local density of the matter (quoted in g/cm^3), in cosmology the mean density of the baryons ($\rho_{\mathrm{b}}^{(0)}$ today). Unit: GeV^4. Status: STANDARD.
- r: The distance from the centre of the star along the path of a neutrino. Unit: length. Status: STANDARD.
- n_e: The number density of electrons (bound and free; in a supernova all are free, in the recombining universe $x_{\mathrm{e}}n_e$ is the density of the free electrons). Unit: GeV^3 (cm^{-3}). Status: STANDARD.

- Y_e: The electron fraction of stellar matter: the number of electrons per nucleon. Unit: dimensionless. Value: 0.5 in the envelope. Status: STANDARD.
- m_u: The atomic mass unit, the mass per nucleon of stellar matter to within one per cent: $n_e = Y_e\rho/m_\mathrm{u}$. Unit: g. Value: 1.66054×10^{-24} g. Status: STANDARD.
- γ_LZ: The adiabaticity parameter of a resonance: the jump probability of FE-50 is $\exp(-\pi\gamma_\mathrm{LZ}/2)$; $\gamma_\mathrm{LZ} \gg 1$ is adiabatic. Unit: dimensionless. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m^2_{ij} = m^2_i - m^2_j$ the squared-mass difference of the states i and j (Δm^2_{21}, Δm^2_{31}, Δm^2_{32}); Δm^2 without labels the difference of the two states of a resonance, $m^2_2 - m^2_1$, multiplied by v^2_loc/v^2 where this differs from 1. Unit: GeV. Status: STANDARD.
- θ: The vacuum mixing angle of two neutrino flavours (in the two-flavour description of a resonance, one of the three angles $\theta_{12}, \theta_{13}, \theta_{23}$ of U_PMNS). Unit: dimensionless. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- r_res: The radius of a resonance: where the electron density equals $n_{e,\mathrm{res}}$ of FE-50 (r_H for Δm^2_{31}, r_L for Δm^2_{21}). Unit: length. Status: STANDARD.

The outer layers of the progenitor star (density $\rho_\mathrm{b} \propto r^{-3}$, electron fraction $Y_\mathrm{e} = 0.5$, m_u the atomic mass unit), through which the neutrinos of a supernova pass after they have left the dense core; γ_LZ is the adiabaticity of a resonance at the radius r_res (the exponent of FE-50 is $-\frac{\pi}{2}\gamma_\mathrm{LZ}$, since $|\mathrm{d}\ln n_e/\mathrm{d}r| = 3/r$). The three-flavour equation with the full mixing matrix ($\theta_{23} = 49.14°$, $\Delta m^2_{31} = 2.511 \times 10^{-3}$ eV^2 in the normal ordering) is integrated from $0.2r_\mathrm{H}$ to $5r_\mathrm{L}$ at $E = 15$ MeV, starting with an electron neutrino (antineutrino). Status: STANDARD (the profile and the equations), DERIVED (the solution).

ordering	particle	P_{ee}	expected for adiabatic resonances	probabilities of the mass states at the surface
normal	ν_e	0.022030	$\lvert U_{e3}\rvert^2 = 0.022030$	ν_3: $1 - 1.5 \times 10^{-6}$
normal	$\bar\nu_e$	0.681644	$\lvert U_{e1}\rvert^2 = 0.681645$	$\bar\nu_1$: $1 - 1.5 \times 10^{-6}$
inverted	ν_e	0.296325	$\lvert U_{e2}\rvert^2 = 0.296325$	ν_2: $1 - 1.5 \times 10^{-6}$
inverted	$\bar\nu_e$	0.022031	$\lvert U_{e3}\rvert^2 = 0.022030$	$\bar\nu_3$: $1 - 1.6 \times 10^{-6}$

The resonance of Δm^2_{31} lies at $r_\mathrm{H} = 16\,833$ km in the normal ordering and at 17 002 km in the inverted one (which keeps $|m^2_3 - m^2_2| = 2.511 \times 10^{-3}$ eV^2, so that $m^2_3 - m^2_1 = -(\Delta m^2_{31} - \Delta m^2_{21}) = -2.437 \times 10^{-3}$ eV^2), and that of Δm^2_{21} at $r_\mathrm{L} = 73\,195$ km; all are adiabatic ($\gamma_\mathrm{LZ} = 215$, 210 and 655, so $P_\mathrm{LZ} = 0$). An electron neutrino, produced in the heaviest state of matter, leaves the star as ν_3 in the normal ordering and as ν_2 in the inverted one; an electron antineutrino leaves as $\bar\nu_1$ or $\bar\nu_3$. The exact integration agrees with these limits to 1.1×10^{-6} (and with half the step to 10^{-9}). Shock waves and turbulence in the envelope, which can make the high resonance non-adiabatic, and the collective conversion deeper inside are not included in this static profile. Status: DERIVED (F2).

11.3 The fast instability near the neutrinosphere

$$\mathcal{G}_\mathrm{ELN}(\hat v) = \sqrt{2}\,G_\mathrm{F,loc}\int_0^\infty \frac{E^2\,\mathrm{d}E}{(2\pi)^3}\Big[\mathcal{F}_e - \bar{\mathcal{F}}_e - \big(\mathcal{F}_x - \bar{\mathcal{F}}_x\big)\Big],$$
$$\det\Big[\eta^{\mu\nu}_\mathrm{M} + \int \mathrm{d}^2\hat v\;\mathcal{G}_\mathrm{ELN}(\hat v)\,\frac{\hat v^\mu \hat v^\nu}{\Omega_\mathrm{fw} - \vec K\cdot\hat v}\Big] = 0, \qquad \hat v^\mu = (1,\ \hat v) \tag{FE-51}$$

Symbols in (FE-51).

- $\mathcal{G}_\mathrm{ELN}(\hat v)$: The angular distribution of the electron lepton number carried by the neutrinos minus that carried by the heavy-lepton neutrinos, per unit solid angle of the direction $\hat v$, multiplied by $\sqrt{2}G_\mathrm{F,loc}$. Unit: GeV. Status: STANDARD.
- $\hat v$, $\hat v^\mu$: The direction of motion of a neutrino: the unit vector $\hat v = \vec p/|\vec p|$ ($\hat v_{\vec p}$, $\hat v_{\vec q}$ for the momenta $\vec p$, $\vec q$), and the four-vector $\hat v^\mu = (1, \hat v)$ in the order $(x_4; x_1, x_2, x_3)$; $\mathrm{d}^2\hat v$ is the element of solid angle of the directions. Unit: dimensionless. Status: STANDARD.

- $G_{\mathrm{F,loc}}$: The local Fermi constant, $G_{\mathrm{F,loc}} = 1/(\sqrt{2}\, v_{\mathrm{loc}}^2)$ ($= G_{\mathrm{F}}$ where $v_{\mathrm{loc}} = v$). Unit: GeV^{-2}. Status: DERIVED.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $\mathcal{F}_e$, $\bar{\mathcal{F}}_e$, $\mathcal{F}_x$, $\bar{\mathcal{F}}_x$: The occupation numbers of the electron neutrinos, of the electron antineutrinos, and of one heavy-lepton neutrino and antineutrino ($x = \mu$ or τ), as functions of the energy E and the direction $\hat{v}$ (the diagonal entries of $\varrho_{\vec{p}}$ and $\bar{\varrho}_{\vec{p}}$); in a transport code, the same for the number of particles in an energy bin. Unit: dimensionless. Status: STANDARD.
- $\eta_{\mathrm{M}}^{\mu\nu}$: The Minkowski metric of the four-vectors $\hat{v}^\mu$, $\eta_{\mathrm{M}} = \mathrm{diag}(1, -1, -1, -1)$ in the order $(x_4; x_1, x_2, x_3)$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+, +, +, -$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- Ω_{fw}: The (complex) frequency of a flavour wave of the neutrino gas, measured from the shift caused by the potentials; an imaginary part greater than zero is an instability. Unit: GeV. Status: STANDARD.
- $\vec{K}$: The (real) wave vector of a flavour wave of the neutrino gas, measured from the shift caused by the potentials. Unit: GeV. Status: STANDARD.

$$\begin{aligned}
\Big(\Omega_{\mathrm{fw}} - K_{\mathrm{r}}\,\mathsf{u}\Big)\mathsf{Q}(\mathsf{u}) &= -2\pi \int_{-1}^{1} \mathrm{d}\mathsf{u}'\,\big(1 - \mathsf{u}\mathsf{u}'\big)\,\mathcal{G}_{\mathrm{ELN}}(\mathsf{u}')\,\mathsf{Q}(\mathsf{u}') \qquad \text{(axially symmetric mode)},\\
\Big(\Omega_{\mathrm{fw}} - K_{\mathrm{r}}\,\mathsf{u}\Big)\mathsf{Q}(\mathsf{u}) &= \pi \int_{-1}^{1} \mathrm{d}\mathsf{u}'\,\sqrt{1-\mathsf{u}^2}\sqrt{1-\mathsf{u}'^2}\,\mathcal{G}_{\mathrm{ELN}}(\mathsf{u}')\,\mathsf{Q}(\mathsf{u}') \qquad \text{(first azimuthal mode)}
\end{aligned} \tag{S-22}$$

Symbols in (S-22).

- Ω_{fw}: The (complex) frequency of a flavour wave of the neutrino gas, measured from the shift caused by the potentials; an imaginary part greater than zero is an instability. Unit: GeV. Status: STANDARD.
- K_{r}: The wave number of a flavour wave along the radial direction (the radial component of $\vec{K}$). Unit: GeV (1/length). Status: STANDARD.
- u: The cosine of the angle between the direction of motion of a neutrino and the radial direction; with the subscript of a flavour (the electron neutrinos or the electron antineutrinos), its value at the edge of the emission cone of that flavour. Unit: dimensionless. Status: STANDARD.
- $\mathsf{Q}(\mathsf{u})$: The amplitude of a small flavour coherence (the off-diagonal entry of ϱ) of the neutrinos moving with the direction cosine u, in a flavour wave. Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\mathcal{G}_{\mathrm{ELN}}(\hat{v})$: The angular distribution of the electron lepton number carried by the neutrinos minus that carried by the heavy-lepton neutrinos, per unit solid angle of the direction $\hat{v}$, multiplied by $\sqrt{2}G_{\mathrm{F,loc}}$. Unit: GeV. Status: STANDARD.

The fast instability as an eigenvalue problem: for an electron-lepton-number distribution that depends only on the cosine u of the angle to the radial direction, a flavour wave with the wave number K_{r} along the radius and the amplitude $\mathsf{Q}(\mathsf{u})$ (axially symmetric, or proportional to the cosine of the azimuth) has the frequencies Ω_{fw} that are the eigenvalues of S-22 (FE-48 linearised; equivalent to the dispersion relation FE-51, which the solutions satisfy to 10^{-14}). The distribution is taken from the neutrinospheres: electron neutrinos emitted uniformly (half-isotropically) from a sphere of radius $R_e = 30$ km and antineutrinos from $R_{\bar{e}} = 27$ km (they decouple deeper and are more forward

peaked), seen at $r = 40$ km, where each fills the cone $\mathsf{u} > \sqrt{1-(R/r)^2}$ ($R = R_e$ or $R_{\bar e}$); α is the ratio of their number densities, the heavy flavours cancel in the electron lepton number, and 200 Gauss-Legendre nodes discretise the angle. Status: STANDARD (the equations), DERIVED (the solution).

$\alpha = n_{\bar\nu_e}/n_{\nu_e}$	0.50	0.75	0.80	0.95	1.20
angular crossing	no	no	yes	yes	yes
growth rate $\operatorname{Im}\Omega_{\rm fw}$ $(\sqrt{2}G_{\rm F}n_{\nu_e})$	0	0	0.0137	0.0556	0.0925
growth length, factor e (cm)	–	–	22	5.4	3.2

The electron lepton number crosses zero as a function of the angle exactly when $\alpha > (1-\mathsf{u}_{\bar e})/(1-\mathsf{u}_e) = 0.7744$ (u_e, $\mathsf{u}_{\bar e}$ the edges of the two cones); without a crossing every frequency is real (growth below 10^{-17}), with a crossing the axially symmetric mode grows. For a luminosity $L = 5\times10^{52}$ erg/s and a mean energy $\langle E\rangle = 12$ MeV of the electron neutrinos, emitted half-isotropically from the neutrinosphere at $R_e = 30$ km as in S-22, their density at 40 km is $n_{\nu_e} = L(1-\mathsf{u}_e)/(2\pi R_e^2 c\langle E\rangle) = 5.2\times10^{32}$ cm^{-3} (c the speed of light), so $\sqrt{2}G_{\rm F}n_{\nu_e} = 3.34$ cm^{-1} (with the Fermi constant $G_{\rm F}$ of the vacuum, $v_{\rm loc} = v$ there); the growth length of the table (the length over which the instability grows by the factor e) is the inverse of the growth rate in cm^{-1}, $1/(\operatorname{Im}\Omega_{\rm fw}\cdot 3.34\ \text{cm}^{-1})$. The instability grows within centimetres: much faster than anything else in the star, it drives the flavours toward equipartition wherever the crossing exists. Status: DERIVED (F3; N3 of `um3_neutrino.py` for the crossing criterion with general distributions).

11.4 The flavour equipartition and the heating behind the shock

$$\begin{aligned}&\mathcal{F}_e \to \mathcal{F}_e - \delta_{\rm FC}, \qquad \bar{\mathcal{F}}_e \to \bar{\mathcal{F}}_e - \delta_{\rm FC}, \qquad \mathcal{F}_x = \bar{\mathcal{F}}_x \to \mathcal{F}_x + \frac{\delta_{\rm FC}}{2},\\ &\delta_{\rm FC} = \max\Big(\frac{2}{3}\Big(\min\big(\mathcal{F}_e,\ \bar{\mathcal{F}}_e\big) - \mathcal{F}_x\Big),\ \max\big(\mathcal{F}_e,\ \bar{\mathcal{F}}_e\big) - 1\Big),\\ &\text{applied where}\ \ \rho_{\rm b} < \rho_{\rm FC}\ \ \text{and}\ \ x_4 - x_{4,\rm b} > t_{\rm FC}\end{aligned} \tag{FE-52}$$

Symbols in (FE-52).

- $\mathcal{F}_e$, $\bar{\mathcal{F}}_e$, $\mathcal{F}_x$, $\bar{\mathcal{F}}_x$: The occupation numbers of the electron neutrinos, of the electron antineutrinos, and of one heavy-lepton neutrino and antineutrino ($x = \mu$ or τ), as functions of the energy E and the direction $\hat v$ (the diagonal entries of $\varrho_{\vec p}$ and $\bar\varrho_{\vec p}$); in a transport code, the same for the number of particles in an energy bin. Unit: dimensionless. Status: STANDARD.
- $\delta_{\rm FC}$: The occupation (or number per energy bin) of electron neutrino-antineutrino pairs converted into heavy-lepton pairs by the flavour equipartition FE-52; a negative value converts heavy-lepton pairs into electron pairs. Unit: dimensionless. Status: ASSUMED.
- $\rho_{\rm b}$: The (rest-mass) energy density of the baryons: in a supernova the local density of the matter (quoted in g/cm^3), in cosmology the mean density of the baryons ($\rho_{\rm b}^{(0)}$ today). Unit: GeV^4. Status: STANDARD.
- $\rho_{\rm FC}$: The baryon density below which the flavour equipartition FE-52 is applied (the critical density $\rho_{\rm c}$ of Gogilashvili and Tamborra). Unit: GeV^4 (g/cm^3). Value: 10^9, 10^{10}, 10^{11} or 10^{13} g/cm^3. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $x_{4,\rm b}$: The time of the bounce of the collapsing stellar core. Unit: length (= time). Status: STANDARD.
- $t_{\rm FC}$: The time after the bounce from which the flavour equipartition FE-52 is applied. Unit: time. Value: 0.02 s. Status: STANDARD.

The flavour equipartition of Gogilashvili and Tamborra (FE-52), applied in every energy bin. Status: STANDARD (the constraints), ASSUMED (this form). On 10^5 random occupations in $[0,1]^3$ it conserves the electron lepton number and the numbers of neutrinos and of antineutrinos to 10^{-15}, keeps every occupation between 0 and 1 (the Pauli bound acts in 22% of the cases), and a second application changes nothing (10^{-16}). Status: DERIVED (F4, N4).

$$\dot Q_{\rm heat} \propto \int_0^\infty \mathrm{d}E\, E^3\big(\mathcal{F}_e + \bar{\mathcal{F}}_e\big), \qquad \mathcal{F}_e,\ \bar{\mathcal{F}}_e,\ \mathcal{F}_x \propto \frac{L_{\rm s}}{E_{\rm s}^2}\Big(\frac{E}{E_{\rm s}}\Big)^{\alpha_{\rm s}} \mathrm{e}^{-(\alpha_{\rm s}+1)E/E_{\rm s}} \tag{S-23}$$

Symbols in (S-23).

- $\dot{Q}_{\mathrm{heat}}$: The rate at which the absorption of electron neutrinos and antineutrinos heats the matter of the gain region (energy per time and mass). Unit: energy per time and mass. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- $\mathcal{F}_e$, $\bar{\mathcal{F}}_e$, $\mathcal{F}_x$, $\bar{\mathcal{F}}_x$: The occupation numbers of the electron neutrinos, of the electron antineutrinos, and of one heavy-lepton neutrino and antineutrino ($x = \mu$ or τ), as functions of the energy E and the direction $\hat{v}$ (the diagonal entries of $\varrho_{\vec{p}}$ and $\bar{\varrho}_{\vec{p}}$); in a transport code, the same for the number of particles in an energy bin. Unit: dimensionless. Status: STANDARD.
- L_{s}: The luminosity (energy per time) of one neutrino species leaving the neutrinosphere. Unit: erg/s. Value: (2 to 5) $\times 10^{52}$ erg/s. Status: STANDARD.
- E_{s}: The mean energy of one neutrino species. Unit: GeV (MeV). Value: 10 to 16 MeV. Status: STANDARD.
- α_{s}: The shape (pinching) parameter of a neutrino spectrum (Keil, Raffelt and Janka). Unit: dimensionless. Value: 3. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

The charged-current heating of the gain region behind the stalled shock ($\nu_e n \to e^- p$ and $\bar{\nu}_e p \to e^+ n$: the cross-section grows as E^2 and the deposited energy is about E), with the number spectra S-23 of the neutrinos (the number per energy of each species, whose integral is its number flux $L_{\mathrm{s}}/E_{\mathrm{s}}$ times a common factor: L_{s} and E_{s} the luminosity and the mean energy of the species, $\alpha_{\mathrm{s}} = 3$ the shape parameter). In the table the species are the electron neutrinos ν_e, the electron antineutrinos $\bar{\nu}_e$ and each of the four heavy-lepton species ν_x (ν_μ, ν_τ and their antiparticles); the mean energies after the equipartition are those of the mixed spectra. Status: STANDARD, ESTIMATE (equal weights of the two reactions).

phase	L_{s} of ν_e, $\bar{\nu}_e$, ν_x (10^{52} erg/s)	E_{s} (MeV)	E_{s} after (MeV)	heating after / before
accretion (0.2 s)	5, 5, 2	11, 14, 16	11.3, 16.1, 14.1	0.96
accretion, softer ν_x	5, 5, 2	11, 14, 14	10.9, 15.3, 13.2	0.86
end of accretion	4, 4, 2.5	11, 13, 15	11.9, 14.7, 13.6	1.12
cooling (comparable luminosities)	3, 3, 3	10, 12, 14	11.4, 13.8, 13.0	1.55

Equipartition removes absorbable electron neutrinos and antineutrinos; it hardens the spectrum of the electron antineutrinos in every case and that of the electron neutrinos unless the heavy-lepton spectra are as soft as in the second row. When the heavy-lepton luminosity is well below the electron-flavour one (the accretion phase) the loss of numbers wins and the heating falls; when they are comparable the harder spectra win and it rises. In the transport simulations of Gogilashvili and Tamborra (195 progenitor models; arXiv:2605.16504 version 3) the flavour equipartition, applied wherever the baryon density falls below a critical value, raises the fraction of failed explosions from 25.6% without conversion to 50.8%, 77.4%, 93.3% and 96.4% for the critical densities 10^9, 10^{10}, 10^{11} and 10^{13} g/cm^3: the deeper the equilibration, the stronger the suppression of the revival of the shock, and the progenitors of 16 to 30 solar masses are the most sensitive; in the models that still explode, conversion near the stalled shock speeds up the revival and leaves lighter neutron stars. This estimate includes neither the feedback on the cooling nor the hydrodynamics. Since the flavour physics of UM-3 is that of the Standard Model with three Dirac neutrinos (FE-49), their results apply to UM-3 unchanged. Status: ESTIMATE (F4).

11.5 The newer global fit, the Dirac phase and the lightest mass

The computations above use the global fit NuFIT 5.2 (November 2022; normal ordering, the fit without the atmospheric data of Super-Kamiokande: $\sin^2\theta_{12} = 0.303$, $\sin^2\theta_{23} = 0.572$, $\sin^2\theta_{13} = 0.02203$, $\Delta m^2_{21} = 7.41 \times 10^{-5}$ eV^2, $\Delta m^2_{31} = 2.511 \times 10^{-3}$ eV^2) with the Dirac phase $\delta_{\mathrm{CP}} = 0$ (the fit has 197°) and the lightest mass $m_1 = 0$. The

newer fit NuFIT 6.0 (I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, I. Martinez-Soler, J. P. Pinheiro and T. Schwetz, JHEP 12 (2024) 216) has two data sets. With the atmospheric data of Super-Kamiokande (the normal ordering preferred by $\Delta\chi^2 = 6.1$): $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{23} = 0.470$, $\sin^2\theta_{13} = 0.02215$, $\delta_{\rm CP} = 212°$, $\Delta m^2_{21} = 7.49 \times 10^{-5}$ eV^2 and $\Delta m^2_{31} = 2.513 \times 10^{-3}$ eV^2 in the normal ordering ($\sin^2\theta_{23} = 0.550$, $\sin^2\theta_{13} = 0.02231$, $\delta_{\rm CP} = 274°$ and $\Delta m^2_{32} = -2.484 \times 10^{-3}$ eV^2 in the inverted one). Without them (the data set of the record's fit; the inverted ordering fits better by $\Delta\chi^2 = 0.6$): $\sin^2\theta_{12} = 0.307$, $\sin^2\theta_{23} = 0.561$, $\sin^2\theta_{13} = 0.02195$, $\delta_{\rm CP} = 177°$, $\Delta m^2_{21} = 7.49 \times 10^{-5}$ eV^2, $\Delta m^2_{31} = 2.534 \times 10^{-3}$ eV^2 in the normal ordering.

$$\left(|U_{ei}|^2\right)_{i=1,2,3} = \left(\cos^2\theta_{12}\cos^2\theta_{13},\ \sin^2\theta_{12}\cos^2\theta_{13},\ \sin^2\theta_{13}\right) \tag{S-33}$$

Symbols in (S-33).

- U_{ei}: The electron row of the mixing matrix, $U_{\rm ei} = (U_{\rm PMNS})_{ei}$: $|U_{\rm ei}|^2$ is the electron-flavour fraction of the mass state i. Unit: dimensionless. Value: $|U_{\rm ei}|^2 = 0.682$, 0.296 and 0.022 for $i = 1, 2, 3$ (NuFIT 5.2, the fit without the Super-Kamiokande atmospheric data: the values of the record; NuFIT 6.0 with them gives 0.677, 0.301 and 0.022). Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- θ_{12}, θ_{13}, θ_{23}: The three mixing angles of $U_{\rm PMNS}$ (its fourth parameter is the Dirac phase, which does not enter the electron-flavour results of UM-3). Unit: dimensionless. Value: 33.40°, 8.54°, 49.14° in the computations of UM-3 ($\sin^2\theta_{12} = 0.303$, $\sin^2\theta_{13} = 0.02203$, $\sin^2\theta_{23} = 0.572$: NuFIT 5.2, November 2022, normal ordering, the fit without the Super-Kamiokande atmospheric data); 33.71°, 8.56°, 43.28° in NuFIT 6.0 (2024, with the Super-Kamiokande atmospheric data; $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$). Status: STANDARD.

The Dirac phase drops out of every result of UM-3. The mixing matrix is $U_{\rm PMNS} = R_{23}U_{13}(\delta_{\rm CP})R_{12}$ (rotations by the three angles, the phase attached to the middle one), and its electron row contains only θ_{12} and θ_{13} (S-33; computed for $\delta_{\rm CP} = 0$, 177°, 212° and 274°, identical to 7×10^{-18}). Every flavour result of UM-3 is a result for electron neutrinos in matter: the matter potential acts on ν_e only, so the phase can be moved into ν_μ and ν_τ, and the three-flavour evolution through the envelope of a supernova gives the same survival probability for $\delta_{\rm CP} = 0$ and 212° (largest difference 6.0×10^{-14} at 10 and 20 MeV, for neutrinos and antineutrinos). The phase enters only the appearance channels, which UM-3 does not compute: the vacuum probability of $\nu_\mu \to \nu_e$ at 295 km and 0.6 GeV (the baseline and the peak energy of T2K) is 0.0421 for $\delta_{\rm CP} = 0$, 0.0478 for 212° and 0.0541 for 270°, that of $\bar\nu_\mu \to \bar\nu_e$ 0.0421, 0.0346 and 0.0291, and the Jarlskog invariant of CP violation is -0.0178 at 212°. Setting the phase to zero was therefore without consequence for UM-3. Status: DERIVED (`um3_numass.py`, NM3).

$$m_2 = \sqrt{m_1^2 + \Delta m^2_{21}}, \qquad m_3 = \sqrt{m_1^2 + \Delta m^2_{31}}, \qquad m_1 < 23 \text{ meV} \ \ (95 \text{ per cent}), \qquad m^2_{\rm eff} = \sum_i |U_{ei}|^2 m_i^2 \tag{S-34}$$

Symbols in (S-34).

- $m_u, \ldots, m_3$: The masses of the quarks (m_u, m_c, m_t; m_d, m_s, m_b), of the charged leptons (m_e, m_μ, m_τ) and of the three neutrino mass states (m_1, m_2, m_3, Dirac masses). Unit: GeV. Value: measured values; $\Delta m^2_{21} = 7.41 \times 10^{-5}$ eV^2, $|\Delta m^2_{31}| = 2.511 \times 10^{-3}$ eV^2 in the computations of UM-3 (the global fit NuFIT 5.2 of the oscillation data, November 2022), 7.49×10^{-5} and 2.513×10^{-3} eV^2 in NuFIT 6.0 (2024); the lightest mass $m_1 = 0$ in the computations, below 23 meV by DESI DR2 with the CMB and NuFIT 6.0. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m^2_{ij} = m_i^2 - m_j^2$ the squared-mass difference of the states i and j (Δm^2_{21}, Δm^2_{31}, Δm^2_{32}); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by $v^2_{\rm loc}/v^2$ where this differs from 1. Unit: GeV. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.

- m_{eff}: The effective mass of the electron neutrino measured in the beta decay of tritium, $m_{\mathrm{eff}}^2 = \sum_i |U_{ei}|^2 m_i^2$. Unit: eV. Value: below 0.45 eV (KATRIN 2025, 90 per cent). Status: STANDARD.
- U_{ei}: The electron row of the mixing matrix, $U_{ei} = (U_{\mathrm{PMNS}})_{ei}$: $|U_{ei}|^2$ is the electron-flavour fraction of the mass state i. Unit: dimensionless. Value: $|U_{ei}|^2 = 0.682$, 0.296 and 0.022 for $i = 1, 2, 3$ (NuFIT 5.2, the fit without the Super-Kamiokande atmospheric data: the values of the record; NuFIT 6.0 with them gives 0.677, 0.301 and 0.022). Status: STANDARD.

The lightest mass. The smallest sums of the masses are 0.0588 eV in the normal and 0.0989 eV in the inverted ordering. DESI DR2 (the BAO) with the CMB, combined with NuFIT 6.0 (W. Elbers et al., arXiv:2503.14744; the cosmological-constant model, 95 per cent), bounds the lightest mass to $m_1 < 23$ meV in the normal ordering (24 meV in the inverted) and the sum of the masses to 0.101 eV (normal) and 0.133 eV (inverted); the inverted ordering is disfavoured (the probability of the normal ordering is 0.91) but not excluded, since its smallest sum is allowed. The bound $\sum_i m_i < 0.0642$ eV of the same data holds for three equal masses with a positive prior on the sum and is not a bound on m_1 (an earlier version of this document applied it to m_1). At $m_1 = 23$ meV the masses are 23.00, 24.57 and 55.15 meV, their sum 0.1027 eV; the effective mass of the beta decay of tritium is $m_{\mathrm{eff}} = 8.84$ to 24.64 meV in this range, far below the limit 0.45 eV of KATRIN (2025). Within the range the lightest mass changes one result of UM-3, the capture of the cosmic neutrinos (S-29): on 100 g of tritium the rate falls from 6.88 per year at $m_1 = 0$ to 4.41 at 5 meV, 4.25 at 10 meV and 4.15 per year at 23 meV, because ν_1 becomes slow (mean speed 0.023 today at 23 meV) and loses the doubling of the capture of relativistic neutrinos (4.06 per year if all states were at rest); and the sum of the masses, which enters the CMB and BAO calculation through $\Omega_\nu h_{100}^2 = \sum_i m_i/(93.14\ \mathrm{eV})$, rises from 0.0588 to 0.1027 eV ($\Omega_\nu h_{100}^2$ from 6.31×10^{-4} to 1.10×10^{-3}). The record (NuFIT 5.2 without the atmospheric data of Super-Kamiokande, $\delta_{\mathrm{CP}} = 0$, $m_1 = 0$) gives the solar survival probabilities 0.5454, 0.5126, 0.5083 and 0.3080 at the energies of the pp, ^{7}Be, pep and ^{8}B neutrinos; NuFIT 6.0 gives 0.5427, 0.5108, 0.5066 and 0.3119 in the same data set and 0.5418, 0.5100, 0.5059 and 0.3126 in the preferred one; the envelope probabilities agree to one per cent and the CNB rates are 6.896 (record), 6.881 and 6.876 per year: the largest relative change is 1.3 per cent like with like and 1.5 per cent against the preferred data set. Status: DERIVED (NM1, NM2, NM4, NM5), STANDARD (the fits and the bounds).

12 The thermal history

$$\rho_{\mathrm{SM}} = \frac{\pi^2}{30}\, g_\rho(\mathsf{T})\, \mathsf{T}^4, \qquad s = \frac{2\pi^2}{45}\, g_{\mathrm{s}}(\mathsf{T})\, \mathsf{T}^3, \qquad H_4 = \sqrt{\frac{8\pi^3 g_\rho}{90}}\, \frac{\mathsf{T}^2}{M_{\mathrm{Pl}}}, \qquad M_{\mathrm{Pl}} = G_{\mathrm{N}}^{-1/2} \tag{FE-60}$$

Symbols in (FE-60).

- ρ_{SM}: The energy density of the Standard-Model particles. Unit: GeV^4. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- g_ρ: The effective number of relativistic degrees of freedom of the energy density. Unit: dimensionless. Value: 106.75 in the Standard Model above the electroweak scale. Status: STANDARD.
- T: The temperature of the Standard-Model plasma (of the photons after the annihilation of electrons and positrons). Unit: GeV. Status: STANDARD.
- s: The entropy density of the Standard-Model plasma. Unit: GeV^3. Status: STANDARD.
- g_{s}: The effective number of relativistic degrees of freedom of the entropy density. Unit: dimensionless. Value: 3.909 today. Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\, h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- M_{Pl}: The Planck mass, $M_{\mathrm{Pl}} = G_{\mathrm{N}}^{-1/2}$. Unit: GeV. Value: 1.221×10^{19} GeV. Status: STANDARD.
- G_{N}: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_{\mathrm{N}} = 1/M_{\mathrm{Pl}}^2$ with the Planck mass $M_{\mathrm{Pl}} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.

The radiation of the Standard Model in thermal equilibrium and the expansion rate of the radiation era (FE-60). The effective numbers of degrees of freedom are computed here from the exact Bose and Fermi integrals of every particle of the Standard Model with its mass (quarks and gluons above the QCD transition, taken at $\mathsf{T} = 0.155$ GeV, the light hadrons below; the neutrinos decouple at 1 MeV and afterwards have the temperature $(4/11)^{1/3}\mathsf{T}$): $g_\rho = g_{\mathrm{s}} = 106.75$ above the electroweak scale, 97.4 and 96.6 at 50 GeV, 85.6 and 85.4 at 5 GeV, 76.4 and 75.5 at 1

GeV, 10.76 at 10 MeV, and today $g_\mathrm{s} = 3.909$, $g_\rho = 3.363$ (the transition region near 0.155 GeV and the annihilation of electrons and positrons are approximated). Status: STANDARD, DERIVED (`um3_thermal.py`, T1).

12.1 The relic density of the particles χ

$$\frac{\mathrm{d}n_\chi}{\mathrm{d}x_4} + 3H_4\, n_\chi = -\langle\sigma_\mathrm{ann}\mathsf{v}\rangle\big(n_\chi^2 - n_{\chi,\mathrm{eq}}^2\big), \qquad n_{\chi,\mathrm{eq}} = g_\chi\Big(\frac{\omega_\mathrm{b}\mathsf{T}}{2\pi}\Big)^{3/2}\mathrm{e}^{-\omega_\mathrm{b}/\mathsf{T}}, \qquad \Omega_\chi h_{100}^2 \simeq \frac{1.07\times10^9\ \mathrm{GeV}^{-1}\, x_\mathrm{f}}{\sqrt{g_*}\, M_\mathrm{Pl}\,\langle\sigma_\mathrm{ann}\mathsf{v}\rangle} \tag{FE-61}$$

Symbols in (FE-61).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- n_χ: The number density of the dark-matter particles χ (particles plus antiparticles). Unit: GeV^3. Status: DERIVED.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44\times10^{-42}$ GeV. Status: STANDARD.
- σ_ann: The annihilation cross-section of the particles χ into Standard-Model particles (through the portals) in the normalisation of FE-61, where n_χ counts the four particle and the four antiparticle states of the bound level: $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle$ is a quarter of the annihilation rate $(\sigma\mathsf{v})_{\chi\bar\chi}$ of one particle χ_a with one antiparticle $\bar\chi_a$. Unit: GeV^{-2}. Status: DERIVED.
- v: The relative velocity of two annihilating particles; $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle$ is the thermal average of the product. Unit: dimensionless. Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- $n_{\chi,\mathrm{eq}}$: The number density of χ in thermal equilibrium with the plasma (non-relativistic). Unit: GeV^3. Status: STANDARD.
- g_χ: The number of internal states of the bound level of χ that are in thermal equilibrium. Unit: dimensionless. Status: DERIVED.
- ω_b: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- T: The temperature of the Standard-Model plasma (of the photons after the annihilation of electrons and positrons). Unit: GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- Ω_χ: The density parameter of the dark matter today. Unit: dimensionless. Value: $\Omega_\chi h_{100}^2 = 0.120$ (Planck 2018). Status: DERIVED.
- h_{100}: The present expansion rate of the fold in units of 100 km/s/Mpc. Unit: dimensionless. Value: 0.674 (Planck 2018). Status: STANDARD.
- x_f: The ratio of the mass of χ to the temperature at which its annihilations stop (freeze-out). Unit: dimensionless. Value: 23 to 30 (for $\omega_\mathrm{b} = 10$ GeV to 10 TeV). Status: STANDARD.
- g_*: The effective number of relativistic degrees of freedom at the freeze-out (g_ρ there). Unit: dimensionless. Status: STANDARD.
- M_Pl: The Planck mass, $M_\mathrm{Pl} = G_\mathrm{N}^{-1/2}$. Unit: GeV. Value: 1.221×10^{19} GeV. Status: STANDARD.

$$\frac{\mathrm{d}\mathsf{Y}}{\mathrm{d}\mathsf{x}} = -\frac{s\,\langle\sigma_\mathrm{ann}\mathsf{v}\rangle}{\mathsf{x}\,H_4}\Big(1 + \frac{1}{3}\frac{\mathrm{d}\ln g_\mathrm{s}}{\mathrm{d}\ln\mathsf{T}}\Big)\big(\mathsf{Y}^2 - \mathsf{Y}_\mathrm{eq}^2\big), \qquad \mathsf{Y} = \frac{n_\chi}{s}, \qquad \mathsf{Y}_\mathrm{eq} = \frac{n_{\chi,\mathrm{eq}}}{s}, \qquad \mathsf{x} = \frac{\omega_\mathrm{b}}{\mathsf{T}},$$
$$n_{\chi,\mathrm{eq}} = \frac{g_\chi\,\omega_\mathrm{b}^2\,\mathsf{T}}{2\pi^2}\,K_2\Big(\frac{\omega_\mathrm{b}}{\mathsf{T}}\Big), \qquad \Omega_\chi h_{100}^2 = \frac{\omega_\mathrm{b}\,\mathsf{Y}(\infty)\,s^{(0)}}{\rho_\mathrm{c}^{(0)}/h_{100}^2} \tag{S-24}$$

Symbols in (S-24).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- Y: The yield of the particles χ: their number density divided by the entropy density, $\mathsf{Y} = n_\chi/s$ (constant when they neither annihilate nor are produced). Unit: dimensionless. Status: STANDARD.
- x: The ratio of the mass of χ to the temperature, $\mathsf{x} = \omega_\mathrm{b}/\mathsf{T}$ (the time variable of the freeze-out). Unit: dimensionless. Status: STANDARD.
- s: The entropy density of the Standard-Model plasma. Unit: GeV^3. Status: STANDARD.
- σ_ann: The annihilation cross-section of the particles χ into Standard-Model particles (through the portals) in the normalisation of FE-61, where n_χ counts the four particle and the four antiparticle states of the bound level: $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle$ is a quarter of the annihilation rate $(\sigma\mathsf{v})_{\chi\bar{\chi}}$ of one particle χ_a with one antiparticle $\bar{\chi}_a$. Unit: GeV^{-2}. Status: DERIVED.
- v: The relative velocity of two annihilating particles; $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle$ is the thermal average of the product. Unit: dimensionless. Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- g_s: The effective number of relativistic degrees of freedom of the entropy density. Unit: dimensionless. Value: 3.909 today. Status: STANDARD.
- T: The temperature of the Standard-Model plasma (of the photons after the annihilation of electrons and positrons). Unit: GeV. Status: STANDARD.
- Y_eq: The yield of χ in thermal equilibrium, $n_{\chi,\mathrm{eq}}/s$. Unit: dimensionless. Status: STANDARD.
- n_χ: The number density of the dark-matter particles χ (particles plus antiparticles). Unit: GeV^3. Status: DERIVED.
- $n_{\chi,\mathrm{eq}}$: The number density of χ in thermal equilibrium with the plasma (non-relativistic). Unit: GeV^3. Status: STANDARD.
- ω_b: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- g_χ: The number of internal states of the bound level of χ that are in thermal equilibrium. Unit: dimensionless. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- K_2: The modified Bessel function of the second kind of order 2 ($K_2(y) \approx \sqrt{\pi/(2y)}\,\mathrm{e}^{-y}$ for large y). Unit: dimensionless. Status: STANDARD.
- Ω_χ: The density parameter of the dark matter today. Unit: dimensionless. Value: $\Omega_\chi h_{100}^2 = 0.120$ (Planck 2018). Status: DERIVED.
- h_{100}: The present expansion rate of the fold in units of 100 km/s/Mpc. Unit: dimensionless. Value: 0.674 (Planck 2018). Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- ρ_c: The critical density, the energy density of a spatially flat fold at the expansion rate H_4; $\rho_\mathrm{c}^{(0)}$ its value today. Unit: GeV^4. Value: $\rho_\mathrm{c}^{(0)} = 1.0537 \times 10^{-5}\,h_{100}^2$ GeV/cm^3. Status: STANDARD.

FE-61 rewritten for the yield Y (the number density per entropy, constant without annihilation) as a function of $\mathsf{x} = \omega_\mathrm{b}/\mathsf{T}$, with the equilibrium density valid also for relativistic χ (K_2 the modified Bessel function of the second kind), and the relic density from the yield after freeze-out ($s^{(0)} = 2891.2$ cm^{-3} the entropy density today, $\rho_\mathrm{c}^{(0)}/h_{100}^2 = 1.0537 \times 10^{-5}$ GeV/cm^3). The bound level of dirac16complex holds four particle states (one multiplet of the rotations of ordinary space and of the extra times: two Dirac fermions χ_a, $a = 1, 2$) and four antiparticle states, so $g_\chi = 8$ and FE-61, written for the total density, has $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle = (\sigma\mathsf{v})_{\chi\bar{\chi}}/4$, where $(\sigma\mathsf{v})_{\chi\bar{\chi}}$ is the annihilation of one χ_a with one $\bar{\chi}_a$ summed over the final states. The equation is integrated from $\mathsf{x} = 1$ to $\mathsf{x} = 1000$ with an implicit method. Status: STANDARD (the equation), DERIVED (the solution).

ω_{b} (GeV)	10	30	100	300	1000	3000	10000
$\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle$ for $\Omega_\chi h^2_{100} = 0.120$ (10^{-26} cm^3/s)	2.31	2.19	2.17	2.25	2.28	2.26	2.31
freeze-out x_{f}	23.1	24.0	25.1	26.2	27.4	28.4	29.6
g_* at freeze-out	67.9	78.6	85.2	86.4	93.8	103.9	106.5
estimate of FE-61 ($\Omega_\chi h^2_{100}$)	0.124	0.127	0.128	0.128	0.127	0.126	0.127

The particles χ are a thermal relic with the measured dark-matter density (Planck 2018: $\Omega_\chi h^2_{100} = 0.1200$) if their annihilation, in the normalisation of FE-61, is $\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle \approx 2.2\times10^{-26}$ cm^3/s for every mass from 10 GeV to 10 TeV (so $(\sigma\mathsf{v})_{\chi\bar\chi} \approx 9\times10^{-26}$ cm^3/s); the analytic estimate of FE-61 agrees within 7%. Status: DERIVED (T2).

12.2 The portals of χ to the Standard Model

$$\mathfrak{g}_\Phi = \frac{\partial\omega_{\mathrm{b}}}{\partial\mu_{\mathrm{L}}}\,\frac{c_\Phi}{V_*}, \qquad \mathfrak{g}_G = \sum_X \frac{\partial\omega_{\mathrm{b}}}{\partial\mu_X}\,\frac{c^G_X}{4V_*}, \qquad \mathfrak{g}_f = \sum_X \frac{\partial\omega_{\mathrm{b}}}{\partial\mu_X}\,\frac{2\omega_{\mathrm{b}}\,c^f_X}{V_*}, \qquad \frac{\partial\omega_{\mathrm{b}}}{\partial\mu_X} = -\frac{q_{\mathrm{b}}}{\omega_{\mathrm{b}}}\,\frac{\partial q_{\mathrm{b}}}{\partial\mu_X},$$
$$\Delta\omega_{\mathrm{b}} = \mathfrak{g}_\Phi\,\Delta\langle\Phi^\dagger\Phi\rangle + \mathfrak{g}_G\,\Delta\langle G^2\rangle + \sum_f \mathfrak{g}_f\,\Delta n_f, \qquad \mathfrak{g}_{h\chi} = \mathfrak{g}_\Phi\,v = -\frac{m^2_\Psi\,\mu^2_{\mathrm{L}}}{2\,\omega_{\mathrm{b}}\,v} \quad (\mu_{\mathrm{R}} = 0) \tag{FE-47}$$

Symbols in (FE-47).

- $\mathfrak{g}_\Phi$, $\mathfrak{g}_G$, $\mathfrak{g}_f$, $\mathfrak{g}_{h\chi}$: The four-dimensional couplings of the bound states χ of dirac16complex to the Standard Model (FE-47): to $\Phi^\dagger\Phi$, to the gluon kinetic density, to the number densities of the fermions of kind f, and the Yukawa coupling of the Higgs boson to χ. Unit: $\mathfrak{g}_\Phi$: GeV^{-1}; $\mathfrak{g}_G$: GeV^{-3}; $\mathfrak{g}_f$: GeV^{-2}; $\mathfrak{g}_{h\chi}$: dimensionless. Status: DERIVED.
- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- ω_{b}: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- μ_X, μ_{L}, μ_{R}: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_{\mathrm{L}}P_{\mathrm{L}} + \mu_{\mathrm{R}}P_{\mathrm{R}})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- c^G_X: The gluon-kinetic portal coupling of the half X of dirac16complex. Unit: GeV^{-7}. Status: INVENTED.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c^f_X: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c^ℓ_X and c^ν_X are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.
- f: The label of a kind of Standard-Model multiplet: $f \in \{Q, u, d, \ell, e, \nu\}$ (quark doublets, up-quark singlets, down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.
- q_{b}: The inverse width of a state of dirac16complex bound to the fold: its profile falls as $\mathrm{e}^{-q_{\mathrm{b}}|\sigma|}$. Unit: GeV. Status: DERIVED.

- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor; for an operator on the space of states, its adjoint with respect to the scalar product of that space (for the quantised dirac16complex the indefinite Krein scalar product). Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- $\langle G^2\rangle$: The mean square of the gluon field strength, $\langle G^A_{\mu\nu}G^{A\mu\nu}\rangle$ (the gluon condensate in the QCD vacuum, a thermal value in the plasma). Unit: GeV^4. Status: STANDARD.
- G^A_μ, $G^A_{\mu\nu}$: The gluon field of $SU(3)_C$ on the fold ($A = 1, \ldots, 8$) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- n_f: The number density of the Standard-Model fermions of kind f (particles minus antiparticles, summed over generations and gauge components), the time component of the mean current $\langle J^\mu_f\rangle$ in the rest frame of the medium. Unit: GeV^3. Status: STANDARD.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_\mathrm{F})^{-1/2}$ with the Fermi constant $G_\mathrm{F} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.

The couplings of the bound states χ to the Standard Model (FE-47): the Higgs-portal Yukawa coupling $\mathfrak{g}_{h\chi}$ is fixed by the binding, the current-portal couplings $\mathfrak{g}_f$ are free (through the INVENTED c^f_χ). Status: DERIVED.

$$\sigma_\mathrm{pair}\mathsf{v} = \sum_f \frac{\mathfrak{g}_f^2\,\omega_\mathrm{b}^2}{2\pi}, \qquad \langle\sigma_\mathrm{ann}\mathsf{v}\rangle = \frac{\sigma_\mathrm{pair}\mathsf{v}}{4}, \qquad \sigma_\mathrm{SI} = \frac{\mu_\mathrm{red}^2\,f_\mathrm{N}^2}{\pi}, \qquad f_\mathrm{N} = 3\,\mathfrak{g}_\mathrm{q} \quad \text{(quark portals)},$$
$$f_\mathrm{N} = \mathfrak{g}_\mathrm{p} = \frac{\alpha_\mathrm{em}}{3\pi}\,\mathfrak{g}_\mathrm{l}\sum_l \ln\frac{\omega_\mathrm{b}^2}{\max\left(m_l^2,\ q_\mathrm{t}^2\right)} \quad \text{(lepton portals)}, \qquad f_\mathrm{N} = \frac{\mathfrak{g}_{h\chi}\,f_\mathrm{H}\,m_\mathrm{N}}{v\,m_h^2} \quad \text{(Higgs portal)} \tag{S-25}$$

Symbols in (S-25).

- σ_pair: The cross-section of the annihilation of one particle χ_a with one antiparticle $\bar\chi_a$ of the same species, summed over the final states; $\sigma_\mathrm{pair}\mathsf{v}$ its product with the relative velocity (a constant for s-wave annihilation). Unit: GeV^{-2}. Status: DERIVED.
- v: The relative velocity of two annihilating particles; $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle$ is the thermal average of the product. Unit: dimensionless. Status: STANDARD.
- $\mathfrak{g}_\Phi$, $\mathfrak{g}_G$, $\mathfrak{g}_f$, $\mathfrak{g}_{h\chi}$: The four-dimensional couplings of the bound states χ of dirac16complex to the Standard Model (FE-47): to $\Phi^\dagger\Phi$, to the gluon kinetic density, to the number densities of the fermions of kind f, and the Yukawa coupling of the Higgs boson to χ. Unit: $\mathfrak{g}_\Phi$: GeV^{-1}; $\mathfrak{g}_G$: GeV^{-3}; $\mathfrak{g}_f$: GeV^{-2}; $\mathfrak{g}_{h\chi}$: dimensionless. Status: DERIVED.
- f: The label of a kind of Standard-Model multiplet: $f \in \{Q, u, d, \ell, e, \nu\}$ (quark doublets, up-quark singlets, down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.
- ω_b: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- σ_ann: The annihilation cross-section of the particles χ into Standard-Model particles (through the portals) in the normalisation of FE-61, where n_χ counts the four particle and the four antiparticle states of the bound level: $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle$ is a quarter of the annihilation rate $(\sigma\mathsf{v})_{\chi\bar\chi}$ of one particle χ_a with one antiparticle $\bar\chi_a$. Unit: GeV^{-2}. Status: DERIVED.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- σ_SI: The spin-independent cross-section of the scattering of χ on a nucleon (normalised per nucleon, as the direct-detection limits are quoted). Unit: GeV^{-2} (cm^2). Status: ESTIMATE.

- $\mu_{\rm red}$: The reduced mass of χ and a nucleon, $\omega_{\rm b} m_{\rm N}/(\omega_{\rm b} + m_{\rm N})$. Unit: GeV. Status: STANDARD.
- $f_{\rm N}$: The effective coupling of χ to a nucleon in the coherent scattering (a vector or scalar contact interaction). Unit: GeV^{-2}. Status: ESTIMATE.
- $\mathfrak{g}_{\rm q}$, $\mathfrak{g}_{\rm l}$, $\mathfrak{g}_{\rm p}$: The current-portal couplings $\mathfrak{g}_f$ of FE-47 taken equal for all chiral quark states ($\mathfrak{g}_{\rm q}$) or for the nine chiral lepton states that take part, the six of the lepton doublets and the three right-handed charged leptons ($\mathfrak{g}_{\rm l}$), and the coupling to the proton that the lepton loop induces ($\mathfrak{g}_{\rm p}$). Unit: GeV^{-2}. Status: ESTIMATE.
- $\alpha_{\rm em}$: The fine-structure constant. Unit: dimensionless. Value: 1/137.036. Status: STANDARD.
- l: The multipole index and number ($l = 0, 1, 2, \ldots$) of an expansion in Legendre polynomials of the direction of a photon or a neutrino relative to the wave vector, or of the CMB sky in spherical harmonics. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m^2_{ij} = m^2_i - m^2_j$ the squared-mass difference of the states i and j (Δm^2_{21}, Δm^2_{31}, Δm^2_{32}); Δm^2 without labels the difference of the two states of a resonance, $m^2_2 - m^2_1$, multiplied by $v^2_{\rm loc}/v^2$ where this differs from 1. Unit: GeV. Status: STANDARD.
- $q_{\rm t}$: The momentum transfer of the scattering of χ on a xenon nucleus in a detector. Unit: GeV. Value: 0.05 GeV. Status: STANDARD.
- $m_{\rm N}$: The mass of a nucleon. Unit: GeV. Value: 0.939 GeV. Status: STANDARD.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\, G_{\rm F})^{-1/2}$ with the Fermi constant $G_{\rm F} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- m_h: The mass of the Higgs boson. Unit: GeV. Value: 125.25 GeV (Particle Data Group 2022). Status: STANDARD.
- $f_{\rm H}$: The nucleon matrix element of the Higgs coupling: the fraction of the nucleon mass carried by the quark masses and, through the trace anomaly, by the gluons. Unit: dimensionless. Value: 0.30. Status: STANDARD.

The annihilation and the scattering on nucleons through the portals, for a vector contact interaction $\mathfrak{g}_f(\bar\chi\hat\gamma^\mu\chi)(\bar\psi_f\hat\gamma_\mu\psi_f)$ of χ (its four-dimensional field, with the Dirac matrices $\hat\gamma^\mu$ of the fold) with a chiral fermion state f of the Standard Model (its field ψ_f): $\sigma_{\rm pair}\mathsf{v}$ is the annihilation of one χ_a with one $\bar\chi_a$ into massless final states (s-wave, summed over the chiral states f), and FE-61 has a quarter of it because n_χ counts the four particle and the four antiparticle states. The coherent spin-independent cross-section per nucleon $\sigma_{\rm SI}$ has the reduced mass $\mu_{\rm red}$ of χ and the nucleon and the nucleon coupling $f_{\rm N}$: for quark portals of equal strength $\mathfrak{g}_{\rm q}$ the vector charge of the nucleon is three times the quark coupling; for lepton portals ($\mathfrak{g}_{\rm l}$ for the nine chiral lepton states that take part: six of the doublets and the three right-handed charged leptons; the right-handed neutrinos must not take part, as the last part of this section shows) the scattering arises at one loop through the photon, with the coupling $\mathfrak{g}_{\rm p}$ to the proton (the sum runs over $l = e, \mu, \tau$ with their masses m_l; $\alpha_{\rm em} = 1/137.036$, the momentum transfer $q_{\rm t} = 50$ MeV of a xenon detector, the cut-off of the logarithm taken at $\omega_{\rm b}$; per nucleon of xenon $\sigma_{\rm SI}$ carries the factor $(54/131)^2$); for the Higgs portal the nucleon (mass $m_{\rm N} = 0.939$ GeV) couples through $f_{\rm H} = 0.30$, the nucleon matrix element of the quark masses and the gluons. Status: STANDARD (the formulas), ESTIMATE (the loop and the matrix elements).

$\omega_{\rm b}$ (GeV)	10	30	100	300	1000	3000	10000
quark portals: $\sigma_{\rm SI}$ (cm^2)	1.1×10^{-38}	1.3×10^{-39}	1.3×10^{-40}	1.5×10^{-41}	1.3×10^{-42}	1.5×10^{-43}	1.4×10^{-44}
lepton portals: $\sigma_{\rm SI}$ (cm^2)	2.8×10^{-43}	5.4×10^{-44}	7.7×10^{-45}	1.3×10^{-45}	1.6×10^{-46}	2.2×10^{-47}	2.6×10^{-48}
limit of the LUX-ZEPLIN experiment (cm^2)	not compared	not compared	5.5×10^{-48}	1.7×10^{-47}	5.5×10^{-47}	1.7×10^{-46}	5.5×10^{-46}
Higgs portal: largest $q_{\rm b}/\omega_{\rm b}$			0.062	0.047	0.035	0.026	0.019

The direct-detection test of a thermal relic χ against the limit of the LUX-ZEPLIN experiment (4.2 tonne-years of xenon, Phys. Rev. Lett. 135, 011802 (2025), arXiv:2410.17036 version 3: 2.2×10^{-48} cm^2 at its minimum, 40 GeV, at 90% confidence). Above the minimum the limit is scaled in proportion to the mass, which is weaker than the measured curve between the minimum and a few hundred GeV, so the exclusions drawn here are conservative; below the minimum the measured limit rises steeply, and no comparison is made at 10 and 30 GeV. Quark portals that set the relic density are excluded at every mass from 100 GeV to 10 TeV, by factors from 2×10^7 (100 GeV) to 25 (10 TeV): the quark couplings must be smaller than the relic value by the square roots of these factors. Lepton portals give a loop-suppressed scattering that the LUX-ZEPLIN limit excludes below about 1.5 TeV and allows above: a leptophilic χ of a few TeV can be the thermal relic. The Higgs-portal coupling of FE-47, $|\mathfrak{g}_{h\chi}| = 2q_{\mathrm{b}}^2/(\omega_{\mathrm{b}}v)$, must stay below 3.1×10^{-3} (100 GeV) to 3.1×10^{-2} (10 TeV), so the binding must be weak: $q_{\mathrm{b}} \lesssim 0.06\,\omega_{\mathrm{b}}$ (100 GeV) to $0.02\,\omega_{\mathrm{b}}$ (10 TeV). Together with S-13 ($q_{\mathrm{b}} > H$) this requires a bulk scale H well below the mass of χ. Status: ESTIMATE (T3, T4).

12.3 The right-handed neutrinos

$$\frac{\mathrm{d}\rho_{\nu_{\mathrm{R}}}}{\mathrm{d}x_4} + 4H_4\,\rho_{\nu_{\mathrm{R}}} = \mathcal{C}_{\nu_{\mathrm{R}}}, \qquad \Delta N_{\mathrm{eff}} = 3\Big(\frac{10.75}{g_{\mathrm{s}}(\mathsf{T}_{\mathrm{dec}})}\Big)^{4/3} \quad \text{(three right-handed neutrinos decoupling at } \mathsf{T}_{\mathrm{dec}}\text{)} \quad \text{(FE-62)}$$

Symbols in (FE-62).

- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $\rho_{\nu_{\mathrm{R}}}$: The energy density of the three right-handed neutrinos. Unit: GeV^4. Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44\times10^{-42}$ GeV. Status: STANDARD.
- $\mathcal{C}_{\nu_{\mathrm{R}}}$: The collision term of the right-handed neutrinos: the rate at which collisions produce (or remove) their energy density. Unit: GeV^5. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- N_{eff}: The effective number of neutrino species: the energy density of the relativistic particles other than photons in units of that of one neutrino species whose temperature is $(4/11)^{1/3}$ times that of the photons. Unit: dimensionless. Value: 3.044 (Standard Model); measured 2.99 ± 0.17 (Planck 2018). Status: STANDARD.
- g_{s}: The effective number of relativistic degrees of freedom of the entropy density. Unit: dimensionless. Value: 3.909 today. Status: STANDARD.
- $\mathsf{T}_{\mathrm{dec}}$: The temperature at which the right-handed neutrinos decouple from the plasma. Unit: GeV. Status: STANDARD.

$$\frac{n_{\chi,\mathrm{eq}}^2\,\sigma_{\mathrm{pair}}\mathsf{v}}{16\,n_{\nu_{\mathrm{R}},\mathrm{eq}}}\Big|_{\mathsf{T}=\mathsf{T}_{\mathrm{dec}}} = H_4\big(\mathsf{T}_{\mathrm{dec}}\big), \qquad n_{\nu_{\mathrm{R}},\mathrm{eq}} = \frac{9\,\zeta(3)}{2\pi^2}\,\mathsf{T}^3 \tag{S-26}$$

Symbols in (S-26).

- $n_{\chi,\mathrm{eq}}$: The number density of χ in thermal equilibrium with the plasma (non-relativistic). Unit: GeV^3. Status: STANDARD.
- σ_{pair}: The cross-section of the annihilation of one particle χ_a with one antiparticle $\bar{\chi}_a$ of the same species, summed over the final states; $\sigma_{\mathrm{pair}}\mathsf{v}$ its product with the relative velocity (a constant for s-wave annihilation). Unit: GeV^{-2}. Status: DERIVED.
- v: The relative velocity of two annihilating particles; $\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle$ is the thermal average of the product. Unit: dimensionless. Status: STANDARD.
- $n_{\nu_{\mathrm{R}},\mathrm{eq}}$: The number density of the three right-handed neutrinos and their antiparticles in thermal equilibrium at the temperature T. Unit: GeV^3. Status: STANDARD.
- T: The temperature of the Standard-Model plasma (of the photons after the annihilation of electrons and positrons). Unit: GeV. Status: STANDARD.

- $\mathsf{T}_{\mathrm{dec}}$: The temperature at which the right-handed neutrinos decouple from the plasma. Unit: GeV. Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- ζ: The Riemann zeta function; $\zeta(3) = 1.2020569$. Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

The right-handed neutrinos. Their production by the neutrino Yukawa couplings is negligible: the rate Γ of their production divided by the expansion rate is $\Gamma/H_4 \sim y_\nu^2 M_{\mathrm{Pl}}/\mathsf{T} = 5 \times 10^{-9}$ at 100 GeV, for the Yukawa coupling $y_\nu = 0.05$ eV$/v$ of the heaviest neutrino (its mass divided by v). If the lepton portal includes them ($c_X^\nu \neq 0$), they stay in equilibrium with χ as long as the rate per right-handed neutrino of $\nu_{\mathrm{R}}\bar{\nu}_{\mathrm{R}} \to \chi\bar{\chi}$ (the left-hand side of S-26, by detailed balance: the three right-handed neutrinos take a quarter of $\sigma_{\mathrm{pair}}\mathsf{v}$, and $n_{\nu_{\mathrm{R}},\mathrm{eq}}$ counts them with their antiparticles) exceeds the expansion rate, and decouple at $\mathsf{T}_{\mathrm{dec}}$. Status: STANDARD, ESTIMATE (the case of equal couplings of the twelve chiral lepton states, the right-handed neutrinos included).

ω_{b} (GeV)	10	30	100	300	1000	3000	10000
$\omega_{\mathrm{b}}/\mathsf{T}_{\mathrm{dec}}$	13.4	14.0	14.6	15.2	15.8	16.4	17.0
ΔN_{eff} (FE-62)	0.236	0.199	0.188	0.182	0.155	0.143	0.141
N_{eff}	3.28	3.24	3.23	3.23	3.20	3.19	3.18
standard deviations above ACT DR6 (2.89 ± 0.11)	3.5	3.2	3.1	3.1	2.8	2.7	2.7

Thermalised right-handed neutrinos would add $\Delta N_{\mathrm{eff}} = 0.14$ to 0.24, which is 1.1 to 1.7 standard deviations above Planck 2018 ($N_{\mathrm{eff}} = 2.99 \pm 0.17$) and 2.7 to 3.5 above the combination of ACT DR6 with Planck, lensing, DESI BAO and nucleosynthesis (2.89 ± 0.11; without nucleosynthesis 2.86 ± 0.13). The CMB alone (the refit below) gives $\Delta\chi^2 = +1.6$ for $\Delta N_{\mathrm{eff}} = 0.141$. UM-3 therefore requires that the right-handed neutrinos never thermalise: their portal couplings must be much smaller than those of the lepton doublets, and then $N_{\mathrm{eff}} = 3.044$ as in the Standard Model. Status: DERIVED (T5), ESTIMATE (the rates).

13 The CMB, the CNB and the BAO

The CMB, the CNB and the BAO are computed with the metric of the fold as the power series FE-35: the expanding background (order 0, with frozen extra times) and the cosmological perturbations (order $\epsilon_{\mathrm{c}} \sim 10^{-5}$); the terms of order ϵ_{c}^2 are dropped. With the corrected weights and the volume of the extra times held fixed (FE-65) the fold obeys the Friedmann equations of general relativity exactly, with Newton's constant $8\pi G_{\mathrm{N}} = \kappa_7/V_*$ (FE-59, above), and the bulk adds no correction to the expansion. The content of the fold is that of UM-3: the bound states χ of dirac16complex as cold dark matter (pressureless in the dilute limit, as the Kohn-Sham equation of state shows), the vacuum energy ρ_Λ of the fold as dark energy (with the equation of state $w = -1$, w the ratio of its pressure to its energy density, to the order computed), the baryons and photons of the Standard Model, and three Dirac neutrinos of the normal ordering with the lightest mass zero ($m_1 = 0$, $m_2 = \sqrt{\Delta m_{21}^2} = 8.61$ meV, $m_3 = \sqrt{\Delta m_{31}^2} = 50.1$ meV, $\sum_i m_i = 0.0587$ eV; ASSUMED: the lightest mass is not fixed by any measurement; DESI DR2 with the CMB and NuFIT 6.0 bounds it below 23 meV, S-34) whose right-handed states are not populated ($N_{\mathrm{eff}} = 3.044$). The primordial spectrum, the optical depth and the densities are the Planck 2018 values (TT,TE,EE+lowE+lensing: $\Omega_{\mathrm{b}}h_{100}^2 = 0.02237$, $\Omega_\chi h_{100}^2 = 0.1200$, $H_4^{(0)} = 67.36$ km/s/Mpc, the optical depth $\tau_{\mathrm{reion}} = 0.0544$ of the reionised universe to the CMB photons, $\ln(10^{10}A_{\mathrm{s}}) = 3.044$, $n_{\mathrm{S}} = 0.9649$; ASSUMED: UM-3 does not compute its primordial era to this precision). The particles χ do not drag the baryons: at the direct-detection limits of the table of the portals ($\sigma_{\mathrm{SI}} \leq 2.2 \times 10^{-48}$ cm^2 ($\omega_{\mathrm{b}}/40$ GeV) for $\omega_{\mathrm{b}} \geq 40$ GeV), a nucleon at recombination ($\mathsf{z} \approx 1100$, thermal speed about 3×10^{-5}) receives momentum from them at the rate $n_\chi \sigma_{\mathrm{SI}}\mathsf{v} \approx 10^{-40}$ s^{-1} for every mass (n_χ falls as $1/\omega_{\mathrm{b}}$

while the limit grows as ω_b), 26 orders of magnitude below the expansion rate $5\times10^{-14}\ \mathrm{s}^{-1}$ there; the scattering on electrons through the lepton portals is weaker still. Status: ESTIMATE. The Boltzmann hierarchy and the line-of-sight integrals are solved by the code CAMB (Lewis, Challinor and Lasenby 2000); the background and the BAO ruler are computed a second time here, independently.

$$\begin{aligned}
&\Big(\partial_{x_4} - a_4'\,|\vec{p}|\,\frac{\partial}{\partial|\vec{p}|}\Big)\,\varrho_{\vec{p}} = -\mathrm{i}\big[\mathsf{H}_{\vec{p}},\ \varrho_{\vec{p}}\big] + \mathcal{C}_{\vec{p}},\\
&N_\mathrm{eff} = \frac{8}{7}\Big(\frac{11}{4}\Big)^{4/3}\frac{\rho_\nu}{\rho_\gamma}, \qquad \mathsf{T}_\nu^{(0)} = \Big(\frac{4}{11}\Big)^{1/3}\mathsf{T}_\gamma^{(0)}, \qquad n_\nu^{(0)} = \frac{3\zeta(3)}{2\pi^2}\big(\mathsf{T}_\nu^{(0)}\big)^3
\end{aligned} \tag{FE-53}$$

Symbols in (FE-53).

- ∂: A partial derivative: the derivative with respect to one variable at fixed values of the others (∂_σ with respect to σ; ∂_M and ∂_μ with respect to the coordinate that carries the same index, of the eight dimensions or of the fold). Status: STANDARD.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c=1$). Status: AUTHOR.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\vec{p}$: The momentum of a neutrino (three components along ordinary space, in the rest frame of the medium); $|\vec{p}| = E$ for a neutrino of negligible mass. Unit: GeV. Status: STANDARD.
- $\varrho_{\vec{p}}$, $\bar{\varrho}_{\vec{p}}$: The 3×3 flavour density matrices of the neutrinos and of the antineutrinos of momentum $\vec{p}$ in the flavour basis (electron, muon, tau): the diagonal entries are the occupation numbers of the three flavours, the off-diagonal entries the coherence between the flavours; $\bar{\varrho}$ is defined with the flavour indices in the reversed order, so that it obeys an equation of the same form. Unit: dimensionless. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\mathsf{H}_{\vec{p}}$, $\bar{\mathsf{H}}_{\vec{p}}$: The flavour Hamiltonians (3×3 Hermitian matrices) of the neutrinos and of the antineutrinos of momentum $\vec{p}$. Unit: GeV. Status: STANDARD.
- $\mathcal{C}_{\vec{p}}$, $\bar{\mathcal{C}}_{\vec{p}}$: The collision terms ($3\times3$ matrices) of the neutrinos and the antineutrinos of momentum $\vec{p}$: the rates of change of $\varrho_{\vec{p}}$ and $\bar{\varrho}_{\vec{p}}$ by emission, absorption, scattering and pair processes with the matter. Unit: GeV. Status: STANDARD.
- N_eff: The effective number of neutrino species: the energy density of the relativistic particles other than photons in units of that of one neutrino species whose temperature is $(4/11)^{1/3}$ times that of the photons. Unit: dimensionless. Value: 3.044 (Standard Model); measured 2.99 ± 0.17 (Planck 2018). Status: STANDARD.
- ρ_ν: The energy density of the neutrinos and antineutrinos of all flavours. Unit: GeV^4. Status: STANDARD.
- ρ_γ: The energy density of the photons. Unit: GeV^4. Status: STANDARD.
- T_ν: The temperature of the cosmic neutrino background; $\mathsf{T}_\nu^{(0)}$ its value today. Unit: GeV (or K). Value: 1.945 K today. Status: STANDARD.
- T_γ: The temperature of the photons (of the CMB); $\mathsf{T}_\gamma^{(0)}$ its value today. Unit: GeV (or K). Value: 2.7255 K today. Status: STANDARD.
- n_ν: The number density of the cosmic neutrino background per flavour (neutrinos plus antineutrinos, one helicity each); $n_\nu^{(0)}$ its value today. Unit: GeV^3. Value: 112 cm^{-3} today. Status: STANDARD.
- ζ: The Riemann zeta function; $\zeta(3) = 1.2020569$. Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

$$\begin{aligned}
&\hat{h}_{\mu\nu}\,\mathrm{d}x^\mu\mathrm{d}x^\nu = a^2\Big[-(1+2\psi_\mathrm{B})\,\mathrm{d}\tau_\mathrm{c}^2 + (1-2\phi_\mathrm{B})\big(\mathrm{d}x_1^2+\mathrm{d}x_2^2+\mathrm{d}x_3^2\big)\Big],\\
&a = \mathrm{e}^{a_4}, \qquad \mathrm{d}\tau_\mathrm{c} = \frac{\mathrm{d}x_4}{a}, \qquad \mathcal{H} = \frac{\dot{a}}{a},\\
&k^2\phi_\mathrm{B} + 3\mathcal{H}\big(\dot{\phi}_\mathrm{B} + \mathcal{H}\psi_\mathrm{B}\big) = -4\pi G_\mathrm{N} a^2\,\Delta\rho, \qquad k^2\big(\dot{\phi}_\mathrm{B} + \mathcal{H}\psi_\mathrm{B}\big) = 4\pi G_\mathrm{N} a^2(\rho+\mathcal{P})\,\vartheta,\\
&\ddot{\phi}_\mathrm{B} + \mathcal{H}\big(\dot{\psi}_\mathrm{B} + 2\dot{\phi}_\mathrm{B}\big) + \Big(2\frac{\ddot{a}}{a} - \mathcal{H}^2\Big)\psi_\mathrm{B} + \frac{k^2}{3}\big(\phi_\mathrm{B} - \psi_\mathrm{B}\big) = 4\pi G_\mathrm{N} a^2\,\Delta\mathcal{P},\\
&k^2\big(\phi_\mathrm{B} - \psi_\mathrm{B}\big) = 12\pi G_\mathrm{N} a^2(\rho+\mathcal{P})\,\Sigma_\mathrm{s}
\end{aligned} \tag{FE-54}$$

Symbols in (FE-54).

- $\hat{h}_{\mu\nu}$: The four-dimensional metric of the fold, $\hat{h}_{\mu\nu} = \mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the coordinates $x_1, \ldots, x_4$ (the induced metric at $\sigma = 0$ without the extra times); $\hat{h}^{\mu\nu}$ its inverse, $\hat{h}$ its determinant, $\sqrt{-\hat{h}} = \mathrm{e}^{3a_4}$; $\hat{h}^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- x: The coordinates of the fold: x_1, x_2, x_3 of ordinary space, x_4 the time, x_5, x_6, x_7 the extra times; a field of the fold is a function of the four observable coordinates $x = (x_1, x_2, x_3, x_4)$, and $\mathrm{d}^4x = \mathrm{d}x_1\,\mathrm{d}x_2\,\mathrm{d}x_3\,\mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.
- a, $\dot{a}$, $\ddot{a}$: The scale factor of the fold, $a = \mathrm{e}^{a_4}$ (normalised to 1 today by the choice of the comoving coordinates x_1, x_2, x_3), and its first and second derivatives with respect to the conformal time τ_c. Unit: dimensionless. Status: STANDARD.
- ψ_B, $\dot{\psi}_\mathrm{B}$: The Bardeen potential of the time-time part of the perturbed metric of the fold (the Newtonian potential) in the conformal Newtonian gauge, a function of τ_c and of the wave number k, and its derivative with respect to τ_c. Unit: dimensionless. Status: STANDARD.
- τ_c: The conformal time of the fold, $\mathrm{d}\tau_\mathrm{c} = \mathrm{e}^{-a_4}\mathrm{d}x_4$ (the time divided by the scale factor e^{a_4} of the fold); a dot on a quantity of the cosmological perturbations is $\mathrm{d}/\mathrm{d}\tau_\mathrm{c}$. Unit: length (= time). Status: STANDARD.
- ϕ_B, $\dot{\phi}_\mathrm{B}$, $\ddot{\phi}_\mathrm{B}$: The Bardeen potential of the space-space part of the perturbed metric of the fold (the curvature potential) in the conformal Newtonian gauge, and its first and second derivatives with respect to τ_c. Unit: dimensionless. Status: STANDARD.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- x_4: The time; a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\mathcal{H}$: The conformal expansion rate, $\mathcal{H} = \dot{a}/a = a\,a_4'$. Unit: 1/length. Status: STANDARD.
- k: The comoving wave number of a Fourier mode of the perturbations, $k = |\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_N: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_\mathrm{N} = 1/M_\mathrm{Pl}^2$ with the Planck mass $M_\mathrm{Pl} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- ρ: The total energy density of the matter of the fold (all species); $\Delta\rho$ its perturbation. Unit: GeV^4. Status: STANDARD.
- $\mathcal{P}$: The total pressure of the matter of the fold; $\Delta\mathcal{P}$ its perturbation. Unit: GeV^4. Status: STANDARD.
- ϑ: The divergence of the velocity of the total matter: $(\rho + \mathcal{P})\vartheta$ is the sum over the species of their energy density plus pressure times the divergence of their velocity. Unit: 1/length. Status: STANDARD.
- Σ_s: The anisotropic stress (shear) of the total matter: $(\rho + \mathcal{P})\Sigma_\mathrm{s}$ is the sum over the species (photons and neutrinos) of their energy density plus pressure times their shear. Unit: dimensionless. Status: STANDARD.

$$
\begin{aligned}
&\dot\delta_\chi = -\vartheta_\chi + 3\dot\phi_\mathrm{B}, \qquad \dot\vartheta_\chi = -\mathcal{H}\vartheta_\chi + k^2\psi_\mathrm{B} + \Gamma_{\chi\mathrm{b}}\big(\vartheta_\mathrm{b} - \vartheta_\chi\big),\\
&\dot\delta_\mathrm{b} = -\vartheta_\mathrm{b} + 3\dot\phi_\mathrm{B}, \qquad \dot\vartheta_\mathrm{b} = -\mathcal{H}\vartheta_\mathrm{b} + c_\mathrm{s}^2k^2\delta_\mathrm{b} + k^2\psi_\mathrm{B} + \frac{a\,x_\mathrm{e}n_e\sigma_\mathrm{T}}{\mathsf{R}}\big(\vartheta_\gamma - \vartheta_\mathrm{b}\big) + \frac{\rho_\chi}{\rho_\mathrm{b}}\,\Gamma_{\chi\mathrm{b}}\big(\vartheta_\chi - \vartheta_\mathrm{b}\big), \qquad \mathsf{R} = \frac{3\rho_\mathrm{b}}{4\rho_\gamma},\\
&\dot\delta_\gamma = -\frac{4}{3}\vartheta_\gamma + 4\dot\phi_\mathrm{B}, \qquad \dot\vartheta_\gamma = k^2\Big(\frac{\delta_\gamma}{4} - \Sigma_\gamma\Big) + k^2\psi_\mathrm{B} + a\,x_\mathrm{e}n_e\sigma_\mathrm{T}\big(\vartheta_\mathrm{b} - \vartheta_\gamma\big),\\
&\dot{\mathsf{F}}_l = \frac{k}{2l+1}\big(l\,\mathsf{F}_{l-1} - (l+1)\,\mathsf{F}_{l+1}\big) - a\,x_\mathrm{e}n_e\sigma_\mathrm{T}\Big(\mathsf{F}_l - \frac{\mathsf{G}_0 + \mathsf{F}_2 + \mathsf{G}_2}{10}\,\delta_{l2}\Big) \quad (l \geq 2),\\
&\dot{\mathsf{G}}_l = \frac{k}{2l+1}\big(l\,\mathsf{G}_{l-1} - (l+1)\,\mathsf{G}_{l+1}\big) - a\,x_\mathrm{e}n_e\sigma_\mathrm{T}\Big(\mathsf{G}_l - \frac{\mathsf{G}_0 + \mathsf{F}_2 + \mathsf{G}_2}{2}\Big(\delta_{l0} + \frac{\delta_{l2}}{5}\Big)\Big) \quad (l \geq 0),\\
&\mathsf{F}_0 = \delta_\gamma, \qquad \mathsf{F}_1 = \frac{4\vartheta_\gamma}{3k}, \qquad \mathsf{F}_2 = 2\Sigma_\gamma,\\
&\dot\delta_\nu = -\frac{4}{3}\vartheta_\nu + 4\dot\phi_\mathrm{B}, \qquad \dot\vartheta_\nu = k^2\Big(\frac{\delta_\nu}{4} - \Sigma_\nu\Big) + k^2\psi_\mathrm{B}, \qquad \dot{\mathsf{N}}_l = \frac{k}{2l+1}\big(l\,\mathsf{N}_{l-1} - (l+1)\,\mathsf{N}_{l+1}\big) \quad (l \geq 2),\\
&\mathsf{N}_1 = \frac{4\vartheta_\nu}{3k}, \qquad \mathsf{N}_2 = 2\Sigma_\nu,\\
&\dot\Upsilon_0 = -\frac{\mathsf{q}k}{\mathsf{E}}\Upsilon_1 - \dot\phi_\mathrm{B}\frac{\mathrm{d}\ln\mathcal{F}_0}{\mathrm{d}\ln\mathsf{q}}, \qquad \dot\Upsilon_1 = \frac{\mathsf{q}k}{3\mathsf{E}}\big(\Upsilon_0 - 2\Upsilon_2\big) - \frac{\mathsf{E}k}{3\mathsf{q}}\,\psi_\mathrm{B}\frac{\mathrm{d}\ln\mathcal{F}_0}{\mathrm{d}\ln\mathsf{q}},\\
&\dot\Upsilon_l = \frac{\mathsf{q}k}{(2l+1)\mathsf{E}}\big(l\,\Upsilon_{l-1} - (l+1)\,\Upsilon_{l+1}\big) \quad (l \geq 2), \qquad \mathsf{E} = \sqrt{\mathsf{q}^2 + a^2m_i^2}, \qquad \mathcal{F}_0 = \frac{1}{\mathrm{e}^{\mathsf{q}/\mathsf{T}_\nu^{(0)}} + 1}
\end{aligned}
\tag{FE-55}
$$

Symbols in (FE-55).

- δ_χ, $\dot\delta_\chi$: The density contrast of the dark matter (the bound states χ of dirac16complex), the perturbation of its energy density divided by its mean, and its derivative with respect to τ_c. Unit: dimensionless. Status: STANDARD.
- ϑ_χ, $\dot\vartheta_\chi$: The divergence of the velocity of the dark matter (for a Fourier mode, ik times its velocity along the wave vector), and its derivative. Unit: 1/length. Status: STANDARD.
- ϕ_B, $\dot\phi_\mathrm{B}$, $\ddot\phi_\mathrm{B}$: The Bardeen potential of the space-space part of the perturbed metric of the fold (the curvature potential) in the conformal Newtonian gauge, and its first and second derivatives with respect to τ_c. Unit: dimensionless. Status: STANDARD.
- $\mathcal{H}$: The conformal expansion rate, $\mathcal{H} = \dot a/a = a\,a_4'$. Unit: 1/length. Status: STANDARD.
- k: The comoving wave number of a Fourier mode of the perturbations, $k = |\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.
- ψ_B, $\dot\psi_\mathrm{B}$: The Bardeen potential of the time-time part of the perturbed metric of the fold (the Newtonian potential) in the conformal Newtonian gauge, a function of τ_c and of the wave number k, and its derivative with respect to τ_c. Unit: dimensionless. Status: STANDARD.
- $\Gamma_{\chi\mathrm{b}}$: The rate (per conformal time) at which the dark matter exchanges momentum with the baryons through the portal couplings of FE-47 (elastic scattering of χ on nuclei and electrons). Unit: 1/length. Value: negligible for the couplings that direct detection allows. Status: DERIVED.
- ϑ_b, $\dot\vartheta_\mathrm{b}$: The divergence of the velocity of the baryons and its derivative. Unit: 1/length. Status: STANDARD.
- δ_b, $\dot\delta_\mathrm{b}$: The density contrast of the baryons and its derivative. Unit: dimensionless. Status: STANDARD.
- c_s: The sound speed of the baryon gas: its square is the temperature of the gas divided by the mean mass per particle, times one minus a third of the logarithmic derivative of that temperature with respect to a. Unit: dimensionless. Status: STANDARD.
- a, $\dot a$, $\ddot a$: The scale factor of the fold, $a = \mathrm{e}^{a_4}$ (normalised to 1 today by the choice of the comoving coordinates x_1, x_2, x_3), and its first and second derivatives with respect to the conformal time τ_c. Unit: dimensionless. Status: STANDARD.
- x_e: The ionisation fraction: the fraction of the electrons that are free. Unit: dimensionless. Status: STANDARD.
- n_e: The number density of electrons (bound and free; in a supernova all are free, in the recombining universe $x_\mathrm{e}n_e$ is the density of the free electrons). Unit: GeV^3 (cm^{-3}). Status: STANDARD.
- σ_T: The Thomson cross-section of the scattering of photons on free electrons. Unit: GeV^{-2}. Value: $6.6524587 \times 10^{-25}$ cm^2. Status: STANDARD.
- R: The ratio of the inertia of the baryons to that of the photons, $\mathsf{R} = 3\rho_\mathrm{b}/(4\rho_\gamma)$. Unit: dimensionless. Status:

STANDARD.

- ϑ_γ, $\dot{\vartheta}_\gamma$: The divergence of the velocity of the photons and its derivative. Unit: 1/length. Status: STANDARD.
- ρ_χ: The energy density of the dark matter, the bound states χ of dirac16complex; $\rho_\chi^{(0)}$ its value today. Unit: GeV^4. Status: DERIVED.
- ρ_b: The (rest-mass) energy density of the baryons: in a supernova the local density of the matter (quoted in $\mathrm{g/cm}^3$), in cosmology the mean density of the baryons ($\rho_\mathrm{b}^{(0)}$ today). Unit: GeV^4. Status: STANDARD.
- ρ_γ: The energy density of the photons. Unit: GeV^4. Status: STANDARD.
- δ_γ, $\dot{\delta}_\gamma$: The density contrast of the photons and its derivative. Unit: dimensionless. Status: STANDARD.
- Σ_γ: The anisotropic stress (shear) of the photons. Unit: dimensionless. Status: STANDARD.
- F_l, $\dot{\mathsf{F}}_l$: The multipoles of the temperature perturbation of the photons ($\mathsf{F}_0 = \delta_\gamma$, $\mathsf{F}_1 = 4\vartheta_\gamma/(3k)$, $\mathsf{F}_2 = 2\Sigma_\gamma$; the photon multipoles of Ma and Bertschinger) and their derivatives with respect to τ_c. Unit: dimensionless. Status: STANDARD.
- l: The multipole index and number ($l = 0, 1, 2, \ldots$) of an expansion in Legendre polynomials of the direction of a photon or a neutrino relative to the wave vector, or of the CMB sky in spherical harmonics. Status: STANDARD.
- G_l, $\dot{\mathsf{G}}_l$: The multipoles of the polarisation perturbation of the photons (the $G_{\gamma l}$ of Ma and Bertschinger) and their derivatives. Unit: dimensionless. Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta_N^M = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with two index strings of equal length the generalised Kronecker delta $\delta_{N_1 N_2 \cdots}^{M_1 M_2 \cdots}$, the determinant of the square matrix of the Kronecker deltas $\delta_{N_b}^{M_a}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- δ_ν, $\dot{\delta}_\nu$: The density contrast of the relativistic neutrinos and its derivative. Unit: dimensionless. Status: STANDARD.
- ϑ_ν, $\dot{\vartheta}_\nu$: The divergence of the velocity of the relativistic neutrinos and its derivative. Unit: 1/length. Status: STANDARD.
- Σ_ν: The anisotropic stress (shear) of the relativistic neutrinos. Unit: dimensionless. Status: STANDARD.
- N_l, $\dot{\mathsf{N}}_l$: The multipoles of the perturbation of the relativistic neutrinos ($\mathsf{N}_0 = \delta_\nu$, $\mathsf{N}_1 = 4\vartheta_\nu/(3k)$, $\mathsf{N}_2 = 2\Sigma_\nu$) and their derivatives. Unit: dimensionless. Status: STANDARD.
- $\Upsilon_l(\mathsf{q})$, $\dot{\Upsilon}_l$: The multipoles of the relative perturbation of the distribution of a massive neutrino state at the comoving momentum q (the Ψ_l of Ma and Bertschinger) and their derivatives. Unit: dimensionless. Status: STANDARD.
- q: The comoving momentum of a neutrino, $a|\vec{p}|$. Unit: GeV. Status: STANDARD.
- E: The comoving energy of a massive neutrino (a times its energy). Unit: GeV. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- $\mathcal{F}_0(\mathsf{q})$: The unperturbed (Fermi-Dirac) occupation number of the cosmic neutrino background at the comoving momentum q. Unit: dimensionless. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m_{ij}^2 = m_i^2 - m_j^2$ the squared-mass difference of the states i and j (Δm_{21}^2, Δm_{31}^2, Δm_{32}^2); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by v_loc^2/v^2 where this differs from 1. Unit: GeV. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- T_ν: The temperature of the cosmic neutrino background; $\mathsf{T}_\nu^{(0)}$ its value today. Unit: GeV (or K). Value: 1.945 K today. Status: STANDARD.

$$
\begin{aligned}
r_\mathrm{s} &= \int_0^{\tau_\mathrm{d}} c_{\gamma\mathrm{b}}\,\mathrm{d}\tau_\mathrm{c}, \qquad c_{\gamma\mathrm{b}} = \frac{1}{\sqrt{3(1+\mathsf{R})}}, \\
D_\mathrm{M}(\mathsf{z}) &= \int_0^{\mathsf{z}} \frac{\mathrm{d}\mathsf{z}'}{H_4(\mathsf{z}')}, \qquad D_\mathrm{H}(\mathsf{z}) = \frac{1}{H_4(\mathsf{z})}, \qquad D_\mathrm{V}(\mathsf{z}) = \left(\mathsf{z}\, D_\mathrm{M}^2(\mathsf{z})\, D_\mathrm{H}(\mathsf{z})\right)^{1/3}
\end{aligned}
\tag{FE-56}
$$

Symbols in (FE-56).

- r_s: The comoving sound horizon at the drag epoch: the comoving distance that a sound wave of the photon-

baryon fluid travels until the baryons are released from the photons. Unit: length. Value: about 147 Mpc (Planck 2018). Status: STANDARD.

- τ_{d}: The conformal time of the drag epoch, when the baryons are released from the photons (redshift about 1060). Unit: length (= time). Status: STANDARD.
- $c_{\gamma\mathrm{b}}$: The sound speed of the tightly coupled photon-baryon fluid. Unit: dimensionless. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- τ_{c}: The conformal time of the fold, $\mathrm{d}\tau_{\mathrm{c}} = \mathrm{e}^{-a_4}\mathrm{d}x_4$ (the time divided by the scale factor e^{a_4} of the fold); a dot on a quantity of the cosmological perturbations is $\mathrm{d}/\mathrm{d}\tau_{\mathrm{c}}$. Unit: length (= time). Status: STANDARD.
- R: The ratio of the inertia of the baryons to that of the photons, $\mathsf{R} = 3\rho_{\mathrm{b}}/(4\rho_\gamma)$. Unit: dimensionless. Status: STANDARD.
- $D_{\mathrm{M}}(\mathsf{z})$: The transverse comoving distance (comoving angular-diameter distance) to the redshift z, for a spatially flat fold. Unit: length. Status: STANDARD.
- z: The redshift, $1 + \mathsf{z} = 1/a$; z' an integration variable. Unit: dimensionless. Status: STANDARD.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- $D_{\mathrm{H}}(\mathsf{z})$: The Hubble distance at the redshift z. Unit: length. Status: STANDARD.
- $D_{\mathrm{V}}(\mathsf{z})$: The volume-averaged distance to the redshift z. Unit: length. Status: STANDARD.

$$
\begin{aligned}
C_l^{\mathrm{TT}} &= 4\pi \int \frac{\mathrm{d}k}{k}\, \mathcal{P}_{\mathcal{R}}(k) \left(\mathcal{I}_l^{\mathrm{T}}(k)\right)^2, \qquad & C_l^{\mathrm{TE}} &= 4\pi \int \frac{\mathrm{d}k}{k}\, \mathcal{P}_{\mathcal{R}}(k)\, \mathcal{I}_l^{\mathrm{T}}(k)\, \mathcal{I}_l^{\mathrm{E}}(k), \\
C_l^{\mathrm{EE}} &= 4\pi \int \frac{\mathrm{d}k}{k}\, \mathcal{P}_{\mathcal{R}}(k) \left(\mathcal{I}_l^{\mathrm{E}}(k)\right)^2, \qquad & \mathcal{P}_{\mathcal{R}}(k) &= A_{\mathrm{s}} \Big(\frac{k}{k_*}\Big)^{n_{\mathrm{S}}-1}
\end{aligned}
\tag{FE-57}
$$

Symbols in (FE-57).

- C_l^{TT}, C_l^{TE}, C_l^{EE}: The angular power spectra of the CMB at the multipole l: temperature with temperature, temperature with E-mode polarisation, E-mode polarisation with itself. Unit: dimensionless. Status: STANDARD.
- l: The multipole index and number ($l = 0, 1, 2, \ldots$) of an expansion in Legendre polynomials of the direction of a photon or a neutrino relative to the wave vector, or of the CMB sky in spherical harmonics. Status: STANDARD.
- T, E: The labels of the CMB observables: T the temperature anisotropy, E the E-mode (gradient) polarisation; TT, TE, EE label the spectra of their products. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- k: The comoving wave number of a Fourier mode of the perturbations, $k = |\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.
- $\mathcal{P}_{\mathcal{R}}(k)$: The primordial power spectrum of the comoving curvature perturbation, per logarithmic interval of k. Unit: dimensionless. Status: ASSUMED.
- $\mathcal{I}_l^{\mathrm{T}}(k)$, $\mathcal{I}_l^{\mathrm{E}}(k)$: The transfer functions of the CMB: the multipole l of the temperature or E-mode anisotropy today produced by a Fourier mode k of unit primordial curvature perturbation, computed from FE-54 and FE-55 (the line-of-sight integral). Unit: dimensionless. Status: STANDARD.
- A_{s}: The amplitude of the primordial curvature spectrum at the pivot k_*. Unit: dimensionless. Value: 2.10×10^{-9} (Planck 2018). Status: ASSUMED.
- k_*: The pivot wave number of the primordial spectrum. Unit: GeV (1/length). Value: 0.05 Mpc^{-1}. Status: STANDARD.
- n_{S}: The spectral index of the primordial curvature spectrum. Unit: dimensionless. Value: 0.965 (Planck 2018). Status: ASSUMED.

$$\Omega_\chi = \frac{\rho_\chi^{(0)}}{\rho_\mathrm{c}^{(0)}}, \qquad \Omega_\mathrm{b} = \frac{\rho_\mathrm{b}^{(0)}}{\rho_\mathrm{c}^{(0)}}, \qquad \Omega_\Lambda = \frac{\rho_\Lambda}{\rho_\mathrm{c}^{(0)}}, \qquad \Omega_\nu h_{100}^2 = \frac{\sum_i m_i}{93.14\ \mathrm{eV}},$$
$$\rho_\mathrm{c}^{(0)} = \frac{3\big(H_4^{(0)}\big)^2}{8\pi G_\mathrm{N}}, \qquad H_4^{(0)} = 100\,h_{100}\ \mathrm{km/s/Mpc} \tag{FE-58}$$

Symbols in (FE-58).

- Ω_χ: The density parameter of the dark matter today. Unit: dimensionless. Value: $\Omega_\chi h_{100}^2 = 0.120$ (Planck 2018). Status: DERIVED.
- ρ_χ: The energy density of the dark matter, the bound states χ of dirac16complex; $\rho_\chi^{(0)}$ its value today. Unit: GeV^4. Status: DERIVED.
- ρ_c: The critical density, the energy density of a spatially flat fold at the expansion rate H_4; $\rho_\mathrm{c}^{(0)}$ its value today. Unit: GeV^4. Value: $\rho_\mathrm{c}^{(0)} = 1.0537\times10^{-5}\,h_{100}^2\ \mathrm{GeV/cm}^3$. Status: STANDARD.
- Ω_b: The density parameter of the baryons today. Unit: dimensionless. Value: $\Omega_\mathrm{b} h_{100}^2 = 0.0224$ (Planck 2018). Status: STANDARD.
- ρ_b: The (rest-mass) energy density of the baryons: in a supernova the local density of the matter (quoted in $\mathrm{g/cm}^3$), in cosmology the mean density of the baryons ($\rho_\mathrm{b}^{(0)}$ today). Unit: GeV^4. Status: STANDARD.
- Ω_Λ: The density parameter of the dark energy (the vacuum energy of the fold) today. Unit: dimensionless. Value: 0.685 (Planck 2018). Status: DERIVED.
- ρ_Λ: The vacuum energy density of the fold above its static tension, the dark energy of UM-3: $\rho_\Lambda = \rho_\mathrm{ff} + \rho_\mathrm{vac} - \lambda_\mathrm{F}$. Unit: GeV^4. Value: $0.685\,\rho_\mathrm{c}^{(0)} = 2.5\times10^{-47}\ \mathrm{GeV}^4$ (Planck 2018). Status: DERIVED.
- Ω_ν, $\Omega_\nu^{(i)}$: The density parameter of the neutrinos of the CNB: Ω_ν that of the massive states today; $\Omega_\nu^{(i)}(\mathsf{z})$ the energy density of the mass state i at the redshift z in units of the critical density today. Unit: dimensionless. Status: STANDARD.
- h_{100}: The present expansion rate of the fold in units of 100 km/s/Mpc. Unit: dimensionless. Value: 0.674 (Planck 2018). Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m_{ij}^2 = m_i^2 - m_j^2$ the squared-mass difference of the states i and j (Δm_{21}^2, Δm_{31}^2, Δm_{32}^2); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by v_loc^2/v^2 where this differs from 1. Unit: GeV. Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44\times10^{-42}$ GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_N: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_\mathrm{N} = 1/M_\mathrm{Pl}^2$ with the Planck mass $M_\mathrm{Pl} = 1.221\times10^{19}$ GeV. Unit: GeV^{-2}. Value: $6.708\times10^{-39}\ \mathrm{GeV}^{-2}$. Status: STANDARD.

The neutrinos in the expanding universe (FE-53), the scalar perturbations of the metric of the fold in the conformal Newtonian gauge and the Einstein equations of the fold to first order (FE-54), the Boltzmann equations of the species (FE-55), the BAO ruler and distances (FE-56), the angular power spectra of the CMB (FE-57) and the density parameters (FE-58). Status: STANDARD (UM-3: the dark matter is χ, the dark energy is the vacuum energy of the fold).

13.1 The background and the BAO ruler

$$\Big(\frac{H_4(\mathsf{z})}{H_4^{(0)}}\Big)^2 = \Omega_\gamma(1+\mathsf{z})^4 + \sum_{i=1}^{3}\Omega_\nu^{(i)}(\mathsf{z}) + \big(\Omega_\mathrm{b}+\Omega_\chi\big)(1+\mathsf{z})^3 + \Omega_\Lambda,$$
$$\Omega_\nu^{(i)}(\mathsf{z}) = \frac{7}{8}\Big(\frac{\mathsf{T}_{\nu,\mathrm{eff}}}{\mathsf{T}_\gamma}\Big)^4\Omega_\gamma\,(1+\mathsf{z})^4\,\frac{\mathsf{I}\big(m_i/\mathsf{T}_{\nu,\mathrm{eff}}\big)}{\mathsf{I}(0)}, \qquad \mathsf{I}(y) = \int_0^\infty \mathrm{d}\mathsf{q}\,\frac{\mathsf{q}^2\sqrt{\mathsf{q}^2+y^2}}{\mathrm{e}^{\mathsf{q}}+1}, \qquad \mathsf{T}_{\nu,\mathrm{eff}} = \Big(\frac{N_\mathrm{eff}}{3}\Big)^{1/4}\Big(\frac{4}{11}\Big)^{1/3}\mathsf{T}_\gamma \tag{S-27}$$

Symbols in (S-27).

- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- z: The redshift, $1 + \mathsf{z} = 1/a$; z' an integration variable. Unit: dimensionless. Status: STANDARD.
- Ω_γ: The density parameter of the photons today. Unit: dimensionless. Value: 5.450×10^{-5} (for $h_{100} = 0.6736$). Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- Ω_ν, $\Omega_\nu^{(i)}$: The density parameter of the neutrinos of the CNB: Ω_ν that of the massive states today; $\Omega_\nu^{(i)}(\mathsf{z})$ the energy density of the mass state i at the redshift z in units of the critical density today. Unit: dimensionless. Status: STANDARD.
- Ω_b: The density parameter of the baryons today. Unit: dimensionless. Value: $\Omega_\mathrm{b} h_{100}^2 = 0.0224$ (Planck 2018). Status: STANDARD.
- Ω_χ: The density parameter of the dark matter today. Unit: dimensionless. Value: $\Omega_\chi h_{100}^2 = 0.120$ (Planck 2018). Status: DERIVED.
- Ω_Λ: The density parameter of the dark energy (the vacuum energy of the fold) today. Unit: dimensionless. Value: 0.685 (Planck 2018). Status: DERIVED.
- $\mathsf{T}_{\nu,\mathrm{eff}}$: The effective temperature of the neutrinos that carries N_eff (the convention of CAMB). Unit: GeV (or K). Status: STANDARD.
- T_γ: The temperature of the photons (of the CMB); $\mathsf{T}_\gamma^{(0)}$ its value today. Unit: GeV (or K). Value: 2.7255 K today. Status: STANDARD.
- $\mathsf{I}(y)$: The Fermi-Dirac energy integral of a neutrino state with the mass-to-temperature ratio y; $\mathsf{I}(0) = 7\pi^4/120$. Unit: dimensionless. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; $\Delta m_{ij}^2 = m_i^2 - m_j^2$ the squared-mass difference of the states i and j (Δm_{21}^2, Δm_{31}^2, Δm_{32}^2); Δm^2 without labels the difference of the two states of a resonance, $m_2^2 - m_1^2$, multiplied by v_loc^2/v^2 where this differs from 1. Unit: GeV. Status: STANDARD.
- y: The ratio of the mass of a neutrino state to its temperature, the argument of I. Unit: dimensionless. Status: STANDARD.
- d: The differential: d in front of a quantity is an infinitesimal change of it (as $\mathrm{d}\sigma$, a small step of σ); the quotient of two such changes is a derivative; d with an exponent, in front of several variables, is their volume element, the product of their differentials. Status: STANDARD.
- q: The comoving momentum of a neutrino, $a|\vec{p}|$. Unit: GeV. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- N_eff: The effective number of neutrino species: the energy density of the relativistic particles other than photons in units of that of one neutrino species whose temperature is $(4/11)^{1/3}$ times that of the photons. Unit: dimensionless. Value: 3.044 (Standard Model); measured 2.99 ± 0.17 (Planck 2018). Status: STANDARD.

The expansion rate of the fold as a function of the redshift z (FE-59 with the content of UM-3): photons, the three neutrino states with their exact Fermi-Dirac energy integrals (each at the temperature $\mathsf{T}_{\nu,\mathrm{eff}}$ that carries $N_\mathrm{eff} = 3.044$, the convention of CAMB), baryons, χ, and the vacuum energy Ω_Λ that makes the fold spatially flat. Computed here it agrees with CAMB to 8.7×10^{-7} for $0 \le \mathsf{z} \le 10^4$, with $\Omega_\gamma = 5.450 \times 10^{-5}$, $\Omega_\nu = 1.404 \times 10^{-3}$ today (the massless state included), $\Omega_\mathrm{b} = 0.0493$, $\Omega_\chi = 0.2645$, $\Omega_\Lambda = 0.6848$; the age of the fold universe is 13.798 Gyr. With frozen extra times this expansion is exact: the bulk changes neither H_4 nor the distances nor the ruler. Status: DERIVED (`um3_cosmology.py`, C1).

The ruler FE-56, integrated here with the same background up to the drag epoch $\mathsf{z}_\mathrm{d} = 1059.93$ of CAMB, is $r_\mathrm{s} = 147.1047$ Mpc (CAMB: 147.1047 Mpc), and the distances D_M and D_H at the seven redshifts of DESI agree with CAMB to 8.4×10^{-7} (for example $D_\mathrm{M} = 1985.92$ Mpc and $D_\mathrm{H} = 3345.07$ Mpc at $\mathsf{z} = 0.51$). Status: DERIVED (C2).

13.2 The CMB

quantity	UM-3	Planck 2018	difference in standard deviations
redshift of last scattering z_*	1089.905	1089.92 ± 0.25	-0.06
sound horizon r_* (Mpc)	144.445	144.43 ± 0.26	$+0.06$
acoustic angle $100\,\theta_*$	1.04108	1.04110 ± 0.00031	-0.06
drag redshift z_d	1059.931	1059.94 ± 0.30	-0.03
ruler $r_\mathrm{s}(\mathsf{z}_\mathrm{d})$ (Mpc)	147.105	147.09 ± 0.26	$+0.06$
damping wave number k_D (Mpc^{-1})	0.140865	0.14087 ± 0.00030	-0.02
matter-radiation equality z_eq	3402.9	3402 ± 26	$+0.03$
σ_8	0.8120	0.8111 ± 0.0060	$+0.15$
age (Gyr)	13.7977	13.797 ± 0.023	$+0.03$

The derived quantities of the CMB of UM-3 with the Planck 2018 parameters (the column TT,TE,EE+lowE+lensing of Table 2 of Planck 2018 VI) lie within 0.2 standard deviations of the Planck values. The small differences come from evaluating CAMB at the mean values of the Planck parameters (the published derived quantities are means over the posterior): a run with Planck's own neutrinos (one massive state of 0.06 eV) has the same offsets in z_* (1089.907), r_* (144.443 Mpc), z_d, r_s (147.103 Mpc), k_D and z_eq; only σ_8 (0.8111 there), $100\theta_*$ (1.04120 there) and the age move with the three neutrino states of UM-3. Here σ_8 is the root mean square of the matter density contrast in spheres of radius $8h_{100}^{-1}$ Mpc today. Status: DERIVED (C3).

$$\chi^2_\mathrm{fit} = \big(\mathsf{X}_\mathrm{model} - \mathsf{X}_\mathrm{data}\big)^T\, \mathsf{C}^{-1} \big(\mathsf{X}_\mathrm{model} - \mathsf{X}_\mathrm{data}\big), \qquad \mathsf{X}_b = \sum_{l \text{ in the bin } b} \mathsf{w}_l\, C_l \tag{S-28}$$

Symbols in (S-28).

- χ^2_fit: The chi-square of a comparison of a model with data (written χ^2 in the text): the squared deviations weighted with the inverse covariance. Unit: dimensionless. Status: STANDARD.
- $\mathsf{X}_\mathrm{model}$, X_data: The vectors of the model values and of the measured values of a data set (binned CMB spectra, BAO distances). Unit: as the data. Status: STANDARD.
- ^{T}: The transpose of a matrix. Status: STANDARD.
- C: The covariance matrix of the measured values; C^{-1} its inverse. Unit: as the data squared. Status: STANDARD.
- X_b: A model value of the binned CMB spectrum, for the bin b of the plik-lite binning. Unit: $\mu\mathrm{K}^2$. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- l: The multipole index and number ($l = 0, 1, 2, \ldots$) of an expansion in Legendre polynomials of the direction of a photon or a neutrino relative to the wave vector, or of the CMB sky in spherical harmonics. Status: STANDARD.
- w_l: The Planck weight of the multipole l in its bin. Unit: dimensionless. Status: STANDARD.
- C_l^TT, C_l^TE, C_l^EE: The angular power spectra of the CMB at the multipole l: temperature with temperature, temperature with E-mode polarisation, E-mode polarisation with itself. Unit: dimensionless. Status: STANDARD.

The comparison with measurements, χ^2_fit (the usual χ^2): the measured values X_data (binned spectra or BAO distances), their covariance matrix C, and the model values $\mathsf{X}_\mathrm{model}$; for the Planck 2018 high-l likelihood (plik-lite: foreground-marginalised spectra in 215 bins of TT for $30 \le l \le 2508$ and 199 bins each of TE and EE for $30 \le l \le 1996$, with the Planck calibration fixed at 1) a model bin is the weighted sum X_b of the C_l of the multipoles of the bin with the Planck weights w_l; two Gaussian bins of the low-l temperature spectrum can be added (Prince and Dunkley 2019). The implementation reproduces the reference log-likelihoods of the plik-lite code on its test spectrum to 10^{-13} (-101.5812307 for TT, -293.9558650 for TT, TE, EE with the low-l bins). Status: STANDARD (the likelihood), DERIVED (its implementation, C4).

model	N_{eff}	χ^2 high-l TT, TE, EE (613 bins)	χ^2 TT (215 bins)	refitted χ^2 (615 bins)	$H_4^{(0)}$ after the refit
UM-3	3.044	583.8	206.2	587.9	67.31
UM-3 with thermal ν_{R}, same θ_*	3.185	604.3	215.1	589.5	68.23
Planck's convention (one state of 0.06 eV)	3.044	583.9	206.1		

The CMB spectra of UM-3 against the Planck 2018 data: $\chi^2 = 583.8$ for 613 bins (expected 613 ± 35). Three thermal right-handed neutrinos at the same acoustic angle and the same other parameters raise χ^2 by 20.5 (the damping tail); when $\Omega_{\mathrm{b}}h^2_{100}$, $\Omega_\chi h^2_{100}$, θ_*, A_{s} and n_{S} are refitted (Gauss-Newton with the Jacobian of CAMB, τ_{reion} fixed) to the 613 high-l bins and the two low-l bins, the difference shrinks to $\Delta\chi^2 = +1.6$, in line with the Planck bound $N_{\mathrm{eff}} = 2.99 \pm 0.17$ (the refit shifts n_{S} from 0.9650 to 0.9702 and $H_4^{(0)}$ from 67.31 to 68.23 km/s/Mpc; the errors from the Fisher matrix, $\sigma(\Omega_{\mathrm{b}}h^2_{100}) = 0.00015$, $\sigma(\Omega_\chi h^2_{100}) = 0.0013$, $\sigma(100\theta_*) = 0.0003$, $\sigma(n_{\mathrm{S}}) = 0.0042$, match Planck's). The spectra are shown in the figure. Status: DERIVED (C4).

13.3 The BAO

tracer	z	measured (DESI DR2)	UM-3	χ^2
BGS	0.295	$D_{\mathrm{V}}/r_{\mathrm{s}} = 7.942 \pm 0.075$	8.057	2.34
LRG1	0.510	$D_{\mathrm{M}}/r_{\mathrm{s}} = 13.588 \pm 0.167$, $D_{\mathrm{H}}/r_{\mathrm{s}} = 21.863 \pm 0.425$	13.500, 22.739	4.48
LRG2	0.706	17.351 ± 0.177, 19.455 ± 0.330	17.700, 20.169	14.36
LRG3+ELG1	0.934	21.576 ± 0.152, 17.641 ± 0.193	21.994, 17.569	8.29
ELG2	1.321	27.601 ± 0.318, 14.176 ± 0.221	28.081, 14.065	2.31
QSO	1.484	30.512 ± 0.760, 12.817 ± 0.516	30.275, 12.880	0.10
Lyα	2.330	38.988 ± 0.531, 8.632 ± 0.101	39.182, 8.616	0.13

The BAO distances of UM-3 with the Planck parameters (the matter density parameter $\Omega_{\mathrm{m}} = 0.3152$, the density of all matter today in units of the critical density, and $h_{100}r_{\mathrm{s}} = 99.09$ Mpc) against DESI DR2 (DESI Collaboration 2025, Table IV; χ^2 with the correlation of D_{M} and D_{H} of each tracer): $\chi^2 = 32.0$ for 13 measurements. Fitted to DESI alone, $\Omega_{\mathrm{m}} = 0.2971$ and $h_{100}r_{\mathrm{s}} = 101.56$ Mpc with $\chi^2 = 10.5$ for 11 degrees of freedom ($p = 0.48$), as DESI finds for the cosmological-constant model; the Planck point lies $\Delta\chi^2 = 21.5$ above this optimum when its own errors are neglected (with the CMB errors DESI quotes a 2.3 standard-deviation tension between BAO and CMB). The dark energy of UM-3 is a vacuum energy with $w = -1$ exactly (w the ratio of its pressure to its energy density): UM-3 inherits this tension and cannot produce the evolving dark energy that DESI DR2 with the CMB prefers at 3.1 standard deviations, $w(a) = w_0 + w_a(1 - a)$ (the equation of state as a function of the scale factor $a = \mathrm{e}^{a_4}$, $a = 1$ today, with its present value w_0 and its slope w_a) with $w_0 > -1$ and $w_a < 0$. Status: DERIVED (C5), STANDARD (DESI's significances).




Figure 1: The CMB spectra of UM-3 with the Planck 2018 parameters ($D_l = l(l+1)C_l/(2\pi)$ of the temperature, the cross-correlation and the E-mode polarisation, lensed), with three thermal right-handed neutrinos (dashed), and the Planck 2018 plik-lite data.

13.4 The CNB

$$\Gamma_{\mathrm{C}\nu\mathrm{B}} = N_{\mathrm{T}}\,\bar{\sigma}_{\mathrm{T}}\,\frac{n_\nu^{(0)}}{2}\sum_{i=1}^{3}|U_{ei}|^2\big(1+\mathsf{s}_i\big) \qquad \text{(Dirac neutrinos)} \tag{S-29}$$

Symbols in (S-29).

- $\Gamma_{\mathrm{C}\nu\mathrm{B}}$: The rate at which a tritium target captures neutrinos of the cosmic neutrino background. Unit: 1/time. Status: STANDARD.
- N_{T}: The number of tritium nuclei of the target. Unit: dimensionless. Value: 2.0×10^{25} for 100 g. Status: STANDARD.
- $\bar{\sigma}_{\mathrm{T}}$: The averaged cross-section of the capture of an electron neutrino on tritium: a tritium nucleus captures left-handed electron neutrinos of mean speed s and number density n at the rate $\bar{\sigma}_{\mathrm{T}}(1+\mathsf{s})\,c\,n$ (c the speed of light, 1 in the formulas; Long, Lunardini and Sabancilar). Unit: cm^2. Value: 3.834×10^{-45} cm^2. Status: STANDARD.
- n_ν: The number density of the cosmic neutrino background per flavour (neutrinos plus antineutrinos, one helicity each); $n_\nu^{(0)}$ its value today. Unit: GeV^3. Value: 112 cm^{-3} today. Status: STANDARD.
- i: A generation index ($i=1,2,3$), which also labels the three neutrino mass states; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ one of the three directions of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- U_{ei}: The electron row of the mixing matrix, $U_{ei}=(U_{\mathrm{PMNS}})_{ei}$: $|U_{ei}|^2$ is the electron-flavour fraction of the mass state i. Unit: dimensionless. Value: $|U_{ei}|^2=0.682$, 0.296 and 0.022 for $i=1,2,3$ (NuFIT 5.2, the fit without the Super-Kamiokande atmospheric data: the values of the record; NuFIT 6.0 with them gives 0.677, 0.301 and 0.022). Status: STANDARD.
- s_i: The mean speed of the neutrino mass state i today, averaged over its momentum distribution (0 for a state at rest, 1 for a massless one). Unit: dimensionless. Value: 1, 0.061 and 0.011 for the mass states 1, 2 and 3 with $m_1=0$; 0.023 for the state 1 with $m_1=23$ meV. Status: STANDARD.

The capture rate of the cosmic neutrinos on a target of N_{T} tritium nuclei ($\nu_e+{}^3\mathrm{H}\to{}^3\mathrm{He}+e^-$, the PTOLEMY idea): each mass state i contributes with its electron fraction $|U_{ei}|^2$; for Dirac neutrinos only the left-handed helicity of each state is populated ($n_\nu^{(0)}/2=56$ cm^{-3} per state), and it is captured with the averaged cross-section $\bar{\sigma}_{\mathrm{T}}=3.834\times10^{-45}$ cm^2 times $(1+\mathsf{s}_i)$, s_i the mean speed of the state over its momentum distribution (Long, Lunardini and Sabancilar 2014). Status: STANDARD.

The CNB of UM-3: the temperature $\mathsf{T}_\nu^{(0)}=(4/11)^{1/3}\mathsf{T}_\gamma^{(0)}=1.9454$ K (1.676×10^{-4} eV); 112.0 neutrinos and antineutrinos per cm^3 per flavour, 336 in all (FE-53); the states ν_2 and ν_3 became non-relativistic at $\mathsf{z}=15.3$ and 94 (where their mass equals the mean momentum $3.15\mathsf{T}_\nu$), ν_1 is massless; $\Omega_\nu h^2_{100}=6.31\times10^{-4}$ for the two massive states (FE-58: $\sum m_i/93.14$ eV $=6.30\times10^{-4}$; CAMB: 6.31×10^{-4}). Because the right-handed states are not populated, the CNB of UM-3 has one helicity per state: a 100 g tritium target captures 4.1 cosmic neutrinos per year if all states are non-relativistic (Majorana neutrinos: 8.1, twice as many; Long, Lunardini and Sabancilar: about 4 and 8), and 6.9 per year in UM-3, whose massless ν_1 moves with the speed of light, so that its left-handed helicity is captured twice as often, while ν_2 and ν_3 move with the mean speeds $\mathsf{s}_2=0.061$ and $\mathsf{s}_3=0.011$ today (the captures of ν_1 lie at the end point of the beta spectrum, where only a mass of ν_1 would separate them). The sum of the masses, 0.0587 eV, lies below the bounds of DESI DR2 with the CMB (0.0642 eV for three equal masses, 0.101 eV in the normal ordering with NuFIT 6.0) and of ACT DR6 (0.082 eV, 95%). With the largest lightest mass that DESI DR2 with NuFIT 6.0 allows, $m_1=23$ meV, the capture rate falls to 4.15 per year (S-34). Status: DERIVED (C6).

14 What the solution does not settle

- **Gravitons that travel at the speed of light everywhere** (DERIVED: impossible with the author's metric). The corrected weights give the four-dimensional graviton of the fold the speed of light at linear order and remove every faster graviton of the static bulk (the bulk of an expanding fold keeps modes without a time-kinetic term, which respond instantaneously), but no choice of the weights and of Λ gives gravitons at the speed of light both in the bulk and on the fold, because the time of the author's metric is not warped (the Lovelock weights and the cosmological constant, step 2). The polarisation along the extra times is not luminal on an expanding fold (FE-63, negligible today). A metric whose time is warped as well might remove the obstruction; whether to change the metric is the author's decision.

- **The strongly coupled bulk** (DERIVED: the degeneracy; OPEN: the response). With the corrected weights the static block is a degenerate vacuum of the Born-Infeld gravity of eight dimensions (FE-70): the bulk equations have no linear part around it for any perturbation of the metric (S-5), static deformations grow like $\mathrm{e}^{3H|\sigma|}$ (S-6), the junction does not respond at first order (S-7), and on the expanding branch no component of the metric has a time-kinetic term. How the bulk responds to sources that depend on σ (the profile of the bound states of dirac16complex, or matter of the fold off the branch of frozen extra times) is not computed, and neither are the scalar perturbations of the bulk on the expanding background; the bulk metric has no perturbative quantum theory around the static block. The cosmology of the fold does not depend on them (FE-33).
- **The volume of the extra times** (DERIVED: that it must be held fixed; ASSUMED: the holding pressure; OPEN: its origin). A free volume would change Newton's constant and the fine-structure constant far beyond the bounds of lunar laser ranging, of optical clocks, of the Oklo reactor and of nucleosynthesis (S-8). UM-3 assumes that the fold fluid supplies the pressure $q_{\mathrm{hold}} = -\Delta_{\mathrm{r}}$ along the extra times that holds it fixed (FE-65, S-9; its value today depends on the assumed pressure of the baryons along the extra times); what produces this pressure (a potential for the volume, like the stabilisation of the sizes of compact extra dimensions in other theories) is not part of UM-3, and in an early universe that expands faster than H the pressure would have to be enormous (S-3).
- **The tension and the dark energy** (ASSUMED, OPEN). The static block needs the total tension $\lambda_{\mathrm{F}} = 6H^2/(8\pi G_{\mathrm{N}})$ of the fold (FE-65); the dark energy ρ_Λ is the small excess of the tension of the fold fluid and the vacuum energy of the fields over λ_{F} (FE-59), a fine tuning as severe as the cosmological-constant problem of general relativity (the loops of the fields of the fold alone induce a vacuum energy of about 2×10^{77} GeV^4, FE-64). UM-3 accommodates the value of ρ_Λ but does not explain it, and its dark energy has $w = -1$ exactly, so the evolving dark energy that DESI DR2 with the CMB prefers is not available.
- **The primordial era** (DERIVED, ESTIMATE, OPEN). The author's vacuum is not a solution with the corrected weights (FE-31); the primordial era of UM-3 can be an inflation of the fold with the volume of the extra times held fixed, which is that of four-dimensional general relativity (S-3), but only if the fold fluid supplies a holding pressure of about $\mathsf{A}^4/8$ times its tension, at least 10^{103} for an inflation at the largest rate that the tensor bound allows if $H \lesssim 4.6 \times 10^{-13}$ GeV, and what could supply it is OPEN. UM-3 does not specify its inflaton: the Higgs field with a curvature coupling $\xi_{\mathrm{H}} \approx 10^4$, an extension of the action, would be possible (ESTIMATE; $n_{\mathrm{S}} = 0.9636$, 3.5 standard deviations below ACT DR6 with Planck and DESI); what precedes inflation, the reheating and the role of the ad hoc vacuum at $x_4 = 0$ are not computed, and the primordial spectrum, the optical depth and the densities are taken from Planck 2018.
- **The Krein norm** (DERIVED; OPEN: stability and a unitary coupling). The charge of dirac16complex is indefinite (FE-43); its quantisation (FE-67) has ghosts that no choice of the vacuum removes. The vacuum cannot decay; whether the gas of the particles χ is classically stable is not decided (the truncated response has frequencies of the size of its error, and the complex partner levels of the dense gas are not converged in the size of the box). The chiral couplings of the fold, the left-handed Higgs portal among them, produce ghosts, so UM-3 with them is not unitary; chirality-blind couplings are necessary for unitarity but not shown to be sufficient.
- **The extra times** (ASSUMED, OPEN). Every field that depends on the extra times is tachyonic (FE-38); UM-3 keeps only fields that do not depend on them. The restriction is consistent but not stable: interactions of fields without extra-time momentum can create pairs of opposite momenta (ESTIMATE, no rate computed), bound states with such momenta leak through the bound region into the bulk, and on the branch of frozen extra times a mode with extra-time momentum grows exponentially. Compact extra times do not help: the n-th harmonic of a period L_{t} grows n times faster than the first, so infinitely many modes grow for every L_{t}, and a compact time-like direction is itself a closed time-like curve (DERIVED, `um3_extratimes.py`). UM-3 needs a gauge principle that removes the momentum along the extra times of every field.
- **Lorentz violation and causality** (DERIVED, ESTIMATE). Zero modes of bulk scalars, photons and fermions move faster than light on the fold (S-12); the bound particles χ have the speed coefficient $c_{\mathrm{b}} > 1$ (S-13) and, when ultrarelativistic, group velocities up to 1.0046 (S-31); the bulk is a shortcut for signals between points of the fold (S-32); and the four time-like directions allow closed time-like curves through every point. For the Standard Model this motivates its place on the fold (ASSUMED); for χ it requires $q_{\mathrm{b}} > H$, and the loop estimate for the electron requires q_{b}/H above 10^{13} to 4×10^{15}, that is H below about 5×10^{-13} GeV.
- **The free parameters**. The bulk mass m_Ψ, the scale H, the volume V_* of the extra times (Newton's constant fixes only $8\pi G_{\mathrm{N}} = 15\kappa H/(16V_*)$), the portal couplings, the self-coupling λ_Ψ, the amplitude $v_{\mathcal{A}}$ and rate a_{40} of the ad hoc vacuum, the lightest neutrino mass (taken as zero; below 23 meV, S-34) and the Dirac phase (taken

as zero; it drops out of every result, S-33) are not fixed by UM-3; the relic density fixes one combination of the mass of χ and its portals.

- **Neutrinos in dense matter** (STANDARD, ESTIMATE). The non-linear evolution of the fast instability and the collective conversion deep in a supernova are not solved; their outcome is modelled by the equipartition FE-52, whose effect on the explosion is taken from the transport simulations of Gogilashvili and Tamborra.

15 What was verified

Every result labelled DERIVED rests on a computation in the folder `universe_model_3/code/` of the repository, which checks itself and writes a report to `universe_model_3/reports/` (all equations and results are written out above). Each script is run with `python universe_model_3/code/NAME.py` (Python 3 with numpy, scipy, sympy, matplotlib and CAMB 2.0.5; the cosmology script downloads the Planck 2018 plik-lite files from github.com/heatherprince/planck-lite-py into `universe_model_3/external/` once).

- `um3_dirac_matrices.py` (55 checks): the author's matrices, their reality, the Clifford relations, the chiral projectors, Pin(4,4).
- `um3_geometry.py` (23 checks): the exact curvature of the author's metric, the Lovelock tensors by two routes, the trace and Bianchi identities, the bulk equations FE-29 with the corrected weights, the vacua of the static block, the frozen extra times and static ordinary space, the two constraints, the author's vacuum (no longer a solution), the spin connection and the vielbein postulate.
- `um3_weights.py` (8 checks): the bulk graviton FE-68 for any weights, the graviton of the fold FE-69, the unique weights and the impossibility of gravitons at the speed of light both in the bulk and on the fold, the static block FE-70, the vacua of the corrected weights, the fold as an Einstein brane with the Friedmann equations, the gravitons on the two vacuum branches (the luminal graviton of the fold and the polarisations that are not luminal), and the record of the author's original weights.
- `um3_linear.py` (5 checks): the linearised Einstein-Lovelock equations for every perturbation of the metric, computed exactly from the value and the first and second derivatives of the metric at a point: they vanish identically around the static block with the corrected weights, and on the two vacuum branches with a changing a_4 or b_4 no component has a time-kinetic term (the jet code by an independent referee, a Claude agent of the same session, adopted after review).
- `um3_fold.py` (13 checks): the proper chart, the general warped metric FE-34, the junction potential, the fold sources FE-30, the conservation law at the fold.
- `um3_bulk.py` (11 checks): the zero modes and the speeds of bulk scalars, photons and fermions (S-11, S-12), and the principal symbol of the bulk graviton order by order (no bulk graviton at the static block with the corrected weights; gradient terms without a time-kinetic term on the two vacuum branches with a changing a_4 or b_4).
- `um3_fields.py` (16 checks): the bilinears of dirac16complex, the Krein operator, the electroweak masses of the ad hoc vacuum, the anomalies with three right-handed neutrinos.
- `um3_spectra.py` (5 checks): the bound states FE-40 with the exact 16×16 matrices, their Krein signature, their speed (S-13).
- `um3_background.py` (7 checks): FE-29 and FE-30 against the stored outputs, S-1 to S-4, the ad hoc vacuum as a gauge rotation and its residual (S-10).
- `um3_series.py` (6 checks): S-5 to S-9, FE-71 and FE-72.
- `um3_kohnsham.py` (9 checks): the Kohn-Sham states (S-14 to S-19) in three regimes, lifted to sixteen components.
- `um3_neutrino.py` (5 checks): the flavour blindness of V_Ψ, the resonance formulas, the crossing criterion, the equipartition, the CNB numbers.
- `um3_flavour.py` (4 checks): S-20 to S-23.
- `um3_thermal.py` (6 checks): the degrees of freedom, S-24 to S-26, the ad hoc vacuum before nucleosynthesis.
- `um3_cosmology.py` (6 checks): S-27 to S-29 and the comparisons with Planck 2018 and DESI DR2.
- `um3_foldsources.py` (6 checks): FE-63, S-30, FE-64 and FE-65.
- `um3_stabiliser.py` (9 checks): the record of the stabiliser with the author's original weights, and its removal by the corrected weights (IN-7).
- `um3_krein.py` (7 checks): FE-67, the symmetries and mirrors of the bound states, the flat directions and the partner level of the gas, the selection rules at the fold.
- `um3_extratimes.py` (5 checks): the fields with a momentum along the extra times.

- `um3_superluminal.py` (5 checks): S-31, S-32, the closed time-like curves and the loop estimate.
- `um3_primordial.py` (7 checks): the author's vacuum (no longer a solution), the records of the fields of the fold in it, the inflation of the fold (S-3) and the size of its holding pressure, Higgs inflation (an extension of the action) and the inflation with a free volume.
- `um3_numass.py` (5 checks): S-33, S-34 and the comparison with NuFIT 6.0.

The Kohn-Sham module runs in parts (`um3_kohnsham.py part NAME` for NAME = K1, I, II, III, K6, K9, then `um3_kohnsham.py assemble`), so that no single calculation takes longer than 423 seconds; the assembled results are identical to a single run.

16 Further symbols

Further symbols. These symbols appear in the definitions of the symbols listed after the equations of this document.

- M_{red}: The reduced Planck mass, $M_{\mathrm{red}} = (8\pi G_{\mathrm{N}})^{-1/2}$. Unit: GeV. Value: 2.435×10^{18} GeV. Status: STANDARD.
- S_Ψ: The action of dirac16complex in the bulk (kinetic term with the canonical spin connection, mass term, contact self-interaction). Unit: dimensionless. Status: RECORD.
- ${\Gamma^P}_{MN}$: The Christoffel symbols of the metric g_{MN} (the Levi-Civita connection). Unit: 1/length. Status: STANDARD.
- β: The matrix of the four-dimensional Dirac adjoint, $\beta = \hat{\gamma}^{(4)}$ (Hermitian, $\beta^2 = \mathbf{1}$). Unit: dimensionless. Status: STANDARD.
- ϖ_{Mab}: The canonical (Levi-Civita) spin connection of the vielbein, antisymmetric in a, b. Unit: 1/length. Status: STANDARD.
- *: Complex conjugation of a number, or of each entry of a matrix or column. Status: STANDARD.
- Γ_{K}: The growth rate of a perturbation of the gas of particles χ when the Krein-positive partner of its occupied level collides with it (a pair of complex frequencies $\omega \pm \mathrm{i}\Gamma_{\mathrm{K}}$). Unit: m_Ψ. Status: DERIVED.
- H_{inf}: The expansion rate of ordinary space during an inflationary era. Unit: GeV. Status: STANDARD.
- L_{t}: The physical period of the extra times, if they are compact. Unit: length. Status: ASSUMED.
- $\mathcal{W}$: The warp factor: in a warped metric, the function of the position along one direction that multiplies the squared steps of other directions (its square root multiplies their lengths); in UM-3 the function of σ that multiplies ordinary space and the extra times, $\mathcal{W} = \mathrm{e}^{-2H|\sigma|} = (\sin^2 z)^{1/6}$, equal to 1 on the fold. Unit: dimensionless. Status: DERIVED.
- c_{e}: The maximal speed of the electron, in units of the speed of light. Unit: dimensionless. Value: $c_{\mathrm{e}} - 1 < 5 \times 10^{-21}$ (no vacuum Cherenkov radiation of the 5 PeV electrons of the 2010 flare of the Crab nebula) and $1 - c_{\mathrm{e}} < 5.6 \times 10^{-20}$ (synchrotron electrons with the Lorentz factor 3×10^9). Status: STANDARD.
- δ_{CP}: The Dirac phase of the mixing matrix U_{PMNS}. Unit: degree. Value: 212° (NuFIT 6.0, normal ordering); 0 in the computations of UM-3. Status: STANDARD.
- ν_α, ν_i: A neutrino field: ν_α the flavour states (the index α runs over the electron, muon and tau flavours), ν_i the mass states ($i = 1, 2, 3$). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- p_{s}: The pressure of the stabiliser of the bulk with the author's original weights (equal along σ, ordinary space and the extra times). Unit: GeV^8. Value: $\kappa p_{\mathrm{s}} = -5H^2/16$ with the author's original weights; 0 with the corrected weights. Status: DERIVED.
- ρ_{s}: The energy density of the stabiliser of the bulk with the author's original weights; with frozen extra times it was the energy density of the bulk at which ordinary space stands still. Unit: GeV^8. Value: $\kappa\rho_{\mathrm{s}} = -H^2/16$ with the author's original weights; 0 with the corrected weights. Status: DERIVED.
- h: The Higgs field: the fluctuation of the lower component of the Higgs doublet about v. Unit: GeV. Value: $m_h = 125.25$ GeV. Status: STANDARD.
- c_{tx}: The common value of the two coefficients $c_{\mathrm{t}} = c_{\mathrm{x}}$ of the bulk equation of a transverse traceless gravitational wave at the static block on the line $w_{(2)} = 18w_{(3)}$ of the Lovelock weights, where bulk gravitons move with the local speed of light: $c_{\mathrm{tx}} = 1 - 360w_{(3)}$ (weights that UM-3 does not use). Unit: dimensionless. Status: DERIVED.