UM-3 solution

The approximate solution of the coupled field equations of UM-3: the background and the power series of the metric, the Higgs field in the ad hoc vacuum, the spectra along the hidden direction, the self-consistent Kohn-Sham states of dirac16complex, the residuals of every field equation, the neutrino flavours, the thermal history, and the CMB, the CNB and the BAO

Underlying theory: Patrick L. Nash Solved and verified by Claude (Anthropic)

Solution of UM-3, 2026-10-24 Repository once-ere/Dirac_claude, folder universe_model_3

Contents

1 The fold: definition and examples **2**

2 How to read this document **8**
2.1 What this document contains . . . 8
2.2 The status labels . . . 9
2.3 Every equation is followed by its symbols . . . 9
2.4 Coordinates, signature and units . . . 9
2.5 The solution at a glance . . . 9

3 The background **10**

4 The power series of the metric to first order **19**

5 The sources of the fold **26**
5.1 The gravitational waves of the fold . . . 26
5.2 The gravity induced by the fields of the fold . . . 28
5.3 The fold fluid, the stabiliser and the energy of the bulk . . . 28

6 The Higgs field in the ad hoc vacuum **31**

7 The spectra along the hidden direction **35**
7.1 The bulk fields and why the Standard Model lives on the fold . . . 35
7.2 dirac16complex: the continuum and the bound states . . . 36

8 The Kohn-Sham states of dirac16complex **42**
8.1 The equations of the gas of bound particles . . . 46
8.2 The method . . . 51
8.3 The solution in three regimes . . . 51
8.4 A weakly bound gas . . . 53
8.5 The Krein partner and the stability of the gas . . . 53

9 The residuals of every field equation **54**

10 The three neutrino flavours **64**
10.1 The Sun . . . 67
10.2 The envelope of a supernova . . . 68
10.3 The fast instability near the neutrinosphere . . . 69
10.4 The flavour equipartition and the heating behind the shock . . . 71
10.5 The newer global fit, the Dirac phase and the lightest mass . . . 73

11 The thermal history **74**
11.1 The relic density of the particles χ . . . 74
11.2 The portals of χ to the Standard Model . . . 77
11.3 The right-handed neutrinos . . . 79

12 The CMB, the CNB and the BAO **81**
12.1 The background and the BAO ruler . . . 87
12.2 The CMB . . . 88
12.3 The BAO . . . 89
12.4 The CNB . . . 90

13 What the solution does not settle **92**

14 What was verified **93**

1 The fold: definition and examples

Definition. The *fold* is the seven-dimensional surface of the author's eight-dimensional spacetime on which the cosine of the angle $z = 6Hx_0$ of the hidden direction vanishes:

$$\mathcal{F} = \big\{X = (x_0, x_1, \ldots, x_7) :\ \cos(6Hx_0) = 0\big\}, \qquad \text{in one period } 0 < z < \pi \text{ of the metric: } z = \frac{\pi}{2}, \quad x_0 = \frac{\pi}{12H} \tag{FD-1}$$

Symbols in (FD-1).

- $\mathcal{F}$: The fold: the seven-dimensional surface $\cos(6Hx_0) = 0$ of the eight-dimensional spacetime (FD-1), in one period $0 < z < \pi$ the surface $z = \pi/2$, $x_0 = \pi/(12H)$, where $\sigma = 0$; $|_{\mathcal{F}}$ means restricted to the fold. Status: DERIVED.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$, the fold is $z = \pi/2$. Unit: dimensionless. Status: AUTHOR.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

$X = \{x_0, \ldots, x_7\}$ are the author's coordinates: x_0 the hidden space-like direction, x_1, x_2, x_3 ordinary space, x_4 the time, x_5, x_6, x_7 the three extra times; H is the author's constant (an inverse length). The author's metric depends on x_0 only through $\cot^2 z$ and $(\sin^2 z)^{1/6}$, so it repeats with the period $\pi/(6H)$ of x_0, and every period $0 < z < \pi$ contains exactly one fold, at $z = \pi/2$. A point of the fold is labelled by the seven coordinates $x_1, \ldots, x_7$, and the metric restricted to the fold ($\mathrm{d}x_0 = 0$, $z = \pi/2$) is

$$g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N\big|_{\mathcal{F}} = \mathrm{e}^{2a_4}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{d}x_4^2 - \mathrm{e}^{-2a_4}b_4^2\big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big) \tag{FD-2}$$

Symbols in (FD-2).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.

- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1 \cdots \mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- $\mathcal{F}$: The fold: the seven-dimensional surface $\cos(6Hx_0) = 0$ of the eight-dimensional spacetime (FD-1), in one period $0 < z < \pi$ the surface $z = \pi/2$, $x_0 = \pi/(12H)$, where $\sigma = 0$; $|_{\mathcal{F}}$ means restricted to the fold. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor of the extra times is $\mathrm{e}^{-a_4} b_4$ times the warp. Unit: dimensionless. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.

with three space-like and four time-like directions; a_4 and b_4 are the author's two functions of the time x_4.

Why this surface is called a fold. Measured by the proper distance along the hidden direction,

$$\sigma = \mathrm{sgn}\Big(\frac{\pi}{2} - z\Big)\frac{\ln \sin z}{6H}, \qquad \mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0, \qquad \sin z = \mathrm{e}^{-6H|\sigma|} \tag{FD-3}$$

Symbols in (FD-3).

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- sgn: The sign function: $\mathrm{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$, the fold is $z = \pi/2$. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

(the fold is $\sigma = 0$; the half-period $0 < z < \pi/2$ is $\sigma < 0$, the half-period $\pi/2 < z < \pi$ is $\sigma > 0$; the two ends of the period, the tips $z \to 0$ and $z \to \pi$, lie at $\sigma \to -\infty$ and $\sigma \to +\infty$), the author's metric is

$$g_{MN}\,\mathrm{d}X^M \mathrm{d}X^N = \mathrm{d}\sigma^2 + \mathrm{e}^{-2H|\sigma|}\Big[\mathrm{e}^{2a_4}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{e}^{-2a_4} b_4^2\big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big)\Big] - \mathrm{d}x_4^2 \tag{FD-4}$$

Symbols in (FD-4).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\ldots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \ldots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor of the extra times is $\mathrm{e}^{-a_4}b_4$ times the warp. Unit: dimensionless. Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.

It depends on σ only through $|\sigma|$. Four properties make the surface $\sigma = 0$ a fold, and together they single it out within the period:

1. *It is a mirror.* The reflection $\sigma \to -\sigma$ (that is $z \to \pi - z$, or $x_0 \to \pi/(6H) - x_0$) maps the metric to itself, and the fold is the only surface of the period that the reflection leaves in place: the two halves of the period are mirror images of each other, joined along the fold, as the two halves of a folded sheet of paper are joined along the crease.
2. *The warp is largest on it.* The warp factor $(\sin^2 z)^{1/6} = \mathrm{e}^{-2H|\sigma|}$, which multiplies ordinary space and the extra times, equals 1 on the fold and decreases on both sides.
3. *The metric has a kink on it.* The metric is continuous across the fold, but its first derivative along σ jumps. The extrinsic curvature of the surfaces $\sigma = \text{const}$ changes sign across the fold,

$$K^a{}_b = \frac{1}{2}\,g^{ac}\,\partial_\sigma g_{cb}: \qquad K^1{}_1 = K^2{}_2 = K^3{}_3 = K^5{}_5 = K^6{}_6 = K^7{}_7 = -H\,\mathrm{sgn}(\sigma), \qquad K^4{}_4 = 0 \tag{FD-5}$$

Symbols in (FD-5).

- $K^a{}_b$: The extrinsic curvature of a surface $\sigma = \text{const}$ of the hidden direction, $K^a{}_b = \frac{1}{2}g^{ac}\partial_\sigma g_{cb}$ for a, b along the surface (FD-5): $-H\,\mathrm{sgn}(\sigma)$ along ordinary space and the extra times, 0 along the time. Unit: GeV (1/length). Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.

- c: A frame index (as a). Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\dots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- ∂: A partial derivative: ∂_M with respect to the coordinate X^M, ∂_μ with respect to x^μ, ∂_σ with respect to σ, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- sgn: The sign function: $\mathrm{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.

it jumps by $-2H$ along ordinary space and along the extra times. The Einstein-Lovelock equations are linear in the second σ-derivatives of the metric, so the jump puts a Dirac delta function $\delta_\mathrm{D}(\sigma)$ into them: the metric solves them only if the fold carries a thin source $\mathcal{S}^a{}_b$, an energy-momentum tensor concentrated on $\sigma = 0$ (the junction conditions). At the author's vacuum ($a_4' = H$, b_4 constant) its frame components along (ordinary space, the time, the extra times) are $\kappa\mathcal{S} = (-10H, -6H, -10H)$, at the static late-time state (a_4 and b_4 constant) $\kappa\mathcal{S} = (-61H/8, -31H/4, -61H/8)$, with $\kappa = 8\pi G_8$ the gravitational coupling of eight dimensions. The fold is therefore not only a place but a physical object: a thin layer that carries energy and momentum (a brane). 4. *The author's coordinate x_0 degenerates on it.* The metric component $g_{x_0x_0} = \cot^2(6Hx_0)$ vanishes on the fold and nowhere else in the period; near the fold the proper distance grows with the square of the coordinate distance,

$$|\sigma| = 3H\Big(x_0 - \frac{\pi}{12H}\Big)^2\Big[1 + 6H^2\Big(x_0 - \frac{\pi}{12H}\Big)^2 + \dots\Big] \tag{FD-6}$$

Symbols in (FD-6).

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

so the two coordinate values on either side of $x_0 = \pi/(12H)$ at the same coordinate distance lie at the same proper distance from the fold: the coordinate line of x_0 is folded onto the line of proper distances (a fold map, example 5).

What lives on the fold. In UM-3 the Standard Model (the Higgs doublet with the ad hoc vacuum, the gauge fields of $SU(3)_C \times SU(2)_L \times U(1)_Y$, three generations of quarks and leptons and three right-handed neutrinos), the fold fluid and the induced Einstein-Hilbert term of the fold live on the fold: they are fields at $\sigma = 0$, functions of $x_1,\dots,x_4$ (no field depends on the extra times). Integrated over the extra times, the fold is the four-dimensional universe that we observe, with the metric $\mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in $x_1,\dots,x_4$. dirac16complex and gravity live in the bulk $\sigma \neq 0$ as well as on the fold; the bound states χ of dirac16complex are the part of it that the fold holds, and the portals couple dirac16complex to the Standard Model at $\sigma = 0$. In UM-3 files the word *fold* means this surface and nothing else: a number of equal energy levels is written as a number of degenerate states, and a growth by powers of Euler's number as a logarithm.

Examples of folds that are meaningful for UM-3.

1. *Our fold.* The fold $z = \pi/2$ of the period that contains our world (FD-1): the seven-dimensional world-volume of the Standard Model, and after the integration over the extra times the four-dimensional universe that we observe.
2. *The copies of the fold.* Because the metric repeats with the period $\pi/(6H)$, every surface $x_0 = \pi/(12H) +$ (an integer multiple of $\pi/(6H)$) is a copy of the fold, one per period. Neighbouring copies are separated by a tip (z a multiple of π), where the warp $(\sin^2 z)^{1/6}$ vanishes and $g_{x_0x_0} = \cot^2 z$ diverges, and which lies at an

infinite proper distance from both: in proper distance every copy has its own complete line $-\infty < \sigma < \infty$. This is why the period $\pi/(6H)$ of x_0 produces no Kaluza-Klein tower with the spacing $12H$; the x_0 mass of the bound states is their binding to the fold.

3. *A folded sheet of paper.* The crease of a folded sheet is a fold in this sense: the two halves are mirror images (property 1) joined along the crease, the sheet is continuous there and its direction changes abruptly (property 3). In UM-3 the direction across the crease is the hidden direction σ, the seven directions along the crease are $x_1, \ldots, x_7$, and the crease is the fold.
4. *A folded function.* The warp factor $\mathrm{e}^{-2H|\sigma|}$ is the simplest folded function: continuous, mirror symmetric, largest at $\sigma = 0$, with a jump of its derivative,

$$\frac{\mathrm{d}}{\mathrm{d}\sigma}\mathrm{e}^{-2H|\sigma|} = -2H\,\mathrm{sgn}(\sigma)\,\mathrm{e}^{-2H|\sigma|}, \qquad \frac{\mathrm{d}^2}{\mathrm{d}\sigma^2}\mathrm{e}^{-2H|\sigma|} = 4H^2\,\mathrm{e}^{-2H|\sigma|} - 4H\,\delta_{\mathrm{D}}(\sigma) \tag{FD-7}$$

Symbols in (FD-7).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- sgn: The sign function: $\mathrm{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.
- δ_{D}: The Dirac delta function: $\int \mathrm{d}\sigma\,\delta_{\mathrm{D}}(\sigma)F(\sigma) = F(0)$. Unit: 1/length (for $\delta_{\mathrm{D}}(\sigma)$). Status: STANDARD.

whose delta function is the origin of the thin source of property 3. Every function of UM-3 that depends on σ through $|\sigma|$ is folded at $\sigma = 0$: the metric functions of FD-4 and the profiles of the bound states of dirac16complex (example 7). 5. *A fold map.* In the mathematics of smooth maps the fold (Whitney's fold singularity) is the map $u \mapsto u^2$ of a line onto a half-line: it folds the line at $u = 0$, two points $\pm u$ land on the same point, and the derivative of the map vanishes on the fold. Near the fold the map from the author's coordinate x_0 to the distance $|\sigma|$ from the fold is of this kind (FD-6), and $g_{x_0x_0} = \cot^2 z$, the square of the derivative $\mathrm{d}\sigma/\mathrm{d}x_0$, vanishes there. 6. *The brane of Randall and Sundrum.* The five-dimensional metric

$$\mathrm{d}y_{\mathrm{RS}}^2 + \mathrm{e}^{-2k_{\mathrm{RS}}|y_{\mathrm{RS}}|}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2 - \mathrm{d}x_4^2\big), \qquad \lambda_{\mathrm{RS}} = \frac{6k_{\mathrm{RS}}}{\kappa_5}, \qquad \Lambda_{\mathrm{RS}} = -\frac{6k_{\mathrm{RS}}^2}{\kappa_5} \tag{FD-8}$$

Symbols in (FD-8).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- y_{RS}: The proper coordinate across the brane of the five-dimensional model of Randall and Sundrum (FD-8, an example, not UM-3). Unit: length. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- k_{RS}: The inverse curvature length of the bulk of the model of Randall and Sundrum: the warp of the brane is $\mathrm{e}^{-2k_{\mathrm{RS}}|y_{\mathrm{RS}}|}$ (FD-8, an example, not UM-3). Unit: GeV (1/length). Status: STANDARD.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- λ_{RS}: The tension (energy per unit volume of ordinary space) of the brane of the model of Randall and Sundrum, $\lambda_{\mathrm{RS}} = 6k_{\mathrm{RS}}/\kappa_5$ (FD-8, an example, not UM-3). Unit: GeV^4. Status: STANDARD.
- κ_5: The gravitational coupling of five dimensions in the model of Randall and Sundrum, $\kappa_5 = 8\pi G_5$ (FD-8, an example, not UM-3). Unit: GeV^{-3}. Status: STANDARD.
- Λ_{RS}: The vacuum energy density of the bulk of the model of Randall and Sundrum, $\Lambda_{\mathrm{RS}} = -6k_{\mathrm{RS}}^2/\kappa_5$ (FD-8, an example, not UM-3). Unit: GeV^5. Status: STANDARD.

has a fold at $y_{\rm RS}=0$ with the properties 1 to 3 (its coordinate $y_{\rm RS}$ is already the proper distance, so property 4 does not arise); Einstein's equations require there a brane of the tension $\lambda_{\rm RS}$ (Israel's junction condition) and in the bulk the vacuum energy $\Lambda_{\rm RS}$. The fold of UM-3 is of the same kind; it differs in that the time x_4 is not warped, three extra times are warped together with ordinary space, and the Lovelock terms add to the junction conditions. 7. *A particle bound to a thin layer.* A particle held by an attractive thin layer has a folded wave function: it falls off exponentially with the distance on both sides, and the layer makes its derivative jump (for the first-order Dirac equation, the wave function itself). In UM-3 the particles χ of dark matter are the states of dirac16complex bound to the fold; at rest and without their self-interaction (the field is $\Psi=\mathrm{e}^{3H|\sigma|}\chi(\sigma)\mathrm{e}^{-\mathrm{i}\omega_{\rm b}x_4}$)

$$\begin{aligned}&\chi(\sigma)=\mathrm{e}^{-q_{\rm b}|\sigma|}\chi(0^+)\ \ (\sigma>0),\qquad \chi(\sigma)=\mathrm{e}^{-q_{\rm b}|\sigma|}\chi(0^-)\ \ (\sigma<0),\\ &\chi(0^+)=\exp\Big(\gamma^{(0)}\big(\mu_{\rm L}P_{\rm L}+\mu_{\rm R}P_{\rm R}\big)\Big)\chi(0^-),\qquad \omega_{\rm b}=\sqrt{m_\Psi^2-q_{\rm b}^2}\end{aligned}\tag{FD-9}$$

Symbols in (FD-9).

- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi=\mathrm{e}^{3H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x}-\omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of a bound state. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma=|\cot z|\,\mathrm{d}x_0$, $\sigma=\mathrm{sgn}(\pi/2-z)\ln(\sin z)/(6H)$ on one period $0<z<\pi$; $\sigma=0$ is the fold $z=\pi/2$, $\sigma\to\mp\infty$ the tips $z\to0,\pi$; one period of x_0 is the whole line $-\infty<\sigma<\infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- $q_{\rm b}$: The inverse width of a state of dirac16complex bound to the fold: its profile falls as $\mathrm{e}^{-q_{\rm b}|\sigma|}$. Unit: GeV. Status: DERIVED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M=\mathbf{e}_a{}^MT^a$, $\gamma_M=g_{MN}\gamma^N$; $\gamma^{(a)}=T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- μ_X, $\mu_{\rm L}$, $\mu_{\rm R}$: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_{\rm L}P_{\rm L}+\mu_{\rm R}P_{\rm R})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- $P_{\rm L}$: The author's left-handed chiral projector, $P_{\rm L}=(\mathbf{1}-\Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- $P_{\rm R}$: The author's right-handed chiral projector, $P_{\rm R}=(\mathbf{1}+\Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- $\omega_{\rm b}$: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.

where $\mu_{\rm L}$, $\mu_{\rm R}$ are the strengths of the fold for the left- and right-handed halves of dirac16complex (set by the Standard-Model fields on the fold) and $q_{\rm b}=-m_\Psi(\mu_{\rm L}+\mu_{\rm R})\tanh\varsigma/(2\varsigma)$, $\varsigma=\sqrt{\mu_{\rm L}\mu_{\rm R}}$ ($q_{\rm b}=-m_\Psi\mu_{\rm L}/2$ for $\mu_{\rm R}=0$). 8. *The fold as a mirror for dirac16complex.* The reflection across the fold acts on the profiles $\chi(\sigma)$ in two ways,

$$(\mathsf{R}_1\chi)(\sigma)=T_0T_1T_2T_3\,\chi(-\sigma),\qquad (\mathsf{R}_2\chi)(\sigma)=T_0T_5T_6T_7\,\chi(-\sigma)\tag{FD-10}$$

Symbols in (FD-10).

- R_1: The reflection across the fold acting on a profile of dirac16complex together with the reflection of ordinary space, $(\mathsf{R}_1\chi)(\sigma)=T_0T_1T_2T_3\,\chi(-\sigma)$ (FD-10): a symmetry of dirac16complex, of gravity and of the mean field of the gas of the particles χ; the bound states of Krein charge -1 are even, those of Krein charge $+1$ odd. Unit: dimensionless. Status: DERIVED.
- R_2: The reflection across the fold acting on a profile of dirac16complex together with the reflection of the extra times, $(\mathsf{R}_2\chi)(\sigma)=T_0T_5T_6T_7\,\chi(-\sigma)$ (FD-10): it maps the bound states of one Krein charge onto those of the other. Unit: dimensionless. Status: DERIVED.
- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi=\mathrm{e}^{3H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x}-\omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of a bound state. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma=|\cot z|\,\mathrm{d}x_0$, $\sigma=\mathrm{sgn}(\pi/2-z)\ln(\sin z)/(6H)$ on one period $0<z<\pi$; $\sigma=0$ is the fold $z=\pi/2$, $\sigma\to\mp\infty$ the tips $z\to0,\pi$; one period of x_0 is the whole line $-\infty<\sigma<\infty$. Unit: length. Status: DERIVED.

- T_a, T^a: The author's eight real 16×16 Dirac matrices T16^A[a] of 4+4.nb ($a = 0, \ldots, 7$; $T_0, \ldots, T_7$), with $\{T_a, T_b\} = 2\eta_{ab}\mathbf{1}$; $T^a = \eta^{ab}T_b$. Unit: dimensionless. Status: AUTHOR.

(T_a the author's Dirac matrices; for states at rest R_1 is the reflection across the fold combined with the reflection of ordinary space, R_2 the reflection across the fold combined with the reflection of the extra times). The Krein charge of a state is its share of the conserved charge of dirac16complex, which is indefinite: a state can carry the charge -1 or $+1$. R_1 is a symmetry of dirac16complex, of gravity and of the mean field of a gas of the particles χ: the four bound states of Krein charge -1 (the particles χ) are even under it and the four of Krein charge $+1$ are odd, so the mean field of the gas cannot mix the two signs of the indefinite charge. R_2 maps the bound states of one Krein charge onto those of the other and makes the two sets degenerate without the self-interaction. 9. *The planes of Horava and Witten.* In M-theory on an interval (a circle whose points are identified with their mirror images), the two ten-dimensional planes left in place by the reflection are mirror surfaces (property 1); the gauge fields $E_8 \times E_8$ live on them while gravity fills the bulk. This is the arrangement of UM-3, where the Standard Model lives on the fold and gravity and dirac16complex fill the bulk.

What is not a fold.

- *The tips* $z = 0$ and $z = \pi$ (mod π): the mirror exchanges them within a period, but there the warp vanishes, $g_{x_0x_0}$ diverges, and they lie at an infinite proper distance; they carry no thin source.
- *A smooth maximum of a warp*, such as $1/\cosh^2(H\sigma)$: mirror symmetric and largest at $\sigma = 0$, but without a kink and without a thin source.

2 How to read this document

2.1 What this document contains

This document solves, approximately and step by step, the coupled field equations of the universe model UM-3: Einstein-Lovelock gravity in eight dimensions with the author's metric, the author's field dirac16complex (a 16-component complex spinor field in the bulk), and the Standard Model with three right-handed neutrinos on the fold of the author's metric. It is self-contained: every equation that is solved is written out here (with the tag it carries in every UM-3 document), every new equation of the solution carries a tag S-1, S-2, ..., and every symbol of every display equation is defined right after it. The sections are:

- **The background** (order 0 of the power series of the metric): the expansion histories that the author's metric admits, what they require of the bulk, and the static late-time state.
- **The power series of the metric to first order**: the response of the bulk to the matter of the fold, the radion mode and the condition under which the fold expands as in general relativity.
- **The sources of the fold**: the speed of the gravitational waves of the fold with and without a gravity of its own, the gravity that the fields of the fold induce, the two parts of the fold fluid, and the constraints of the bulk that tie the fold to the stabiliser of the bulk.
- **The Higgs field in the ad hoc vacuum**: the author's ad hoc vacuum written as a gauge rotation, the field of W^2 that the Gauss law requires, and the one residual that remains.
- **The spectra along the hidden direction**: the zero modes and the speeds of the bulk fields, the continuum and the bound states of dirac16complex (the x_0 mass), the speed of the bound particles χ and the shortcut through the bulk.
- **The Kohn-Sham states of dirac16complex**: the self-consistent states of a gas of the bound particles χ with the self-interaction, gravity and the mean fields of the Standard Model, in three regimes, with their residuals, and the stability of the gas against the partner states of the other Krein charge.
- **The residuals of every field equation**: how well the approximate solution satisfies each equation.
- **The three neutrino flavours**: the Sun, the envelope of a supernova, the fast instability and the flavour equipartition, and what the newer global fit, the Dirac phase and the lightest mass change.
- **The thermal history**: the degrees of freedom, the relic density of χ, its portals to the Standard Model, the right-handed neutrinos and the ad hoc vacuum before nucleosynthesis.
- **The CMB, the CNB and the BAO** (the cosmic microwave background, the cosmic neutrino background and the baryon acoustic oscillations): the cosmological perturbations of the metric of the fold, against the measurements of Planck, of DESI DR2 (the second data release of the Dark Energy Spectroscopic Instrument) and of ACT DR6 (the sixth data release of the Atacama Cosmology Telescope).
- **What the solution does not settle**, and **what was verified**.

2.2 The status labels

Every equation and every result carries one of the labels

- **AUTHOR**: from the author (his notebooks `4+4.nb` and `Dirac_Einstein-Lovelock_4+4_Nash.nb`, and his order of 2026-10-24: the coordinates, the metric, the Lovelock weights, the Dirac matrices, the chiral projectors, the list of fields, the Higgs potential and the ad hoc vacuum);
- **RECORD**: from the record of this repository (the Lagrangian of dirac16complex with its canonical spin connection and its contact self-interaction, the Kohn-Sham treatment);
- **STANDARD**: textbook physics, or a published measurement (the source is named);
- **ASSUMED**: a choice made in UM-3 without derivation;
- **INVENTED**: a new interaction of UM-3 (the portals of dirac16complex);
- **DERIVED**: computed and verified in UM-3 (the verification is stated);
- **ESTIMATE**: an order-of-magnitude evaluation;
- **OPEN**: not settled.

2.3 Every equation is followed by its symbols

Every display equation is followed by the list "Symbols in (S-n)" or "Symbols in (FE-n)" of EVERY symbol that appears in it: every quantity, field, coupling, constant and operator, every index letter with the values it takes, every mark (prime, dagger, transpose, star), the imaginary unit, Euler's number and the differential. Each item gives the meaning, the unit (natural units $\hbar = c = 1$), the value where one exists, and the status. A field equation of UM-3 carries a tag FE-n and a term of the action a tag IN-n, the same in every UM-3 document; it is written here exactly as everywhere else. A new equation of this document carries a tag S-n.

2.4 Coordinates, signature and units

The author's coordinates are $X = \{x_0, x_1, \ldots, x_7\}$, always with the hidden direction first: x_0 is the hidden space-like coordinate, on which the metric depends periodically with the period $\pi/(6H)$ through the angle $z = 6Hx_0$; x_1, x_2, x_3 are ordinary space, x_4 is the time, and x_5, x_6, x_7 are three extra time-like coordinates. The signature is $(+,+,+,+,-,-,-,-)$ in this order; the four observable coordinates $x_1, \ldots, x_4$ have the signature $(+,+,+,-)$. H is the author's constant (an inverse length), a free parameter of UM-3. Units are natural, $\hbar = c = 1$ ($1\ \mathrm{GeV}^{-1} = 1.973 \times 10^{-14}$ cm $= 6.582 \times 10^{-25}$ s; $1\ \mathrm{GeV}^{-2} = 3.894 \times 10^{-28}\ \mathrm{cm}^2$). A prime on a function of the time x_4 alone is its derivative $\mathrm{d}/\mathrm{d}x_4$, and on a profile (a function of σ alone) its derivative $\mathrm{d}/\mathrm{d}\sigma$. The proper coordinate of the hidden direction is σ (FE-27): $\sigma = 0$ is the fold $z = \pi/2$. The fields live as follows: gravity in the bulk and on the fold; dirac16complex and a medium that keeps the late-time bulk static (the stabiliser) in the bulk; the Standard Model with three right-handed neutrinos, a thin fluid (the fold fluid) and an Einstein-Hilbert term of the fold (the induced gravity, with the coupling κ_4) on the fold $\sigma = 0$.

2.5 The solution at a glance

part	result	status
background	the only histories with constant rates are the author's vacuum $a_4 = Hx_4$ with constant b_4 and its time reverse; a varying b_4 needs bulk sources of second order in $b_4'/(Hb_4)$; the late universe is the static block with frozen extra times, whose bulk media violate the null energy condition	DERIVED
power series	to first order the bulk responds either with a mode that grows away from the fold, or (if its medium responds) with the radion mode, which keeps the fold equations those of general relativity; the latter is required by observation	DERIVED, ASSUMED (the response)
Higgs field	the ad hoc vacuum is an $SU(2)$ rotation of the radial field $\sqrt{v^2 + \mathcal{A}^2}$ with a pure-gauge W^2; one radial residual remains, which vanishes as $\mathcal{A} \to 0$	DERIVED
spectra	bulk fields have zero modes that move faster than light on the fold; dirac16complex has a continuum above m_Ψ and eight bound states at $\omega_\mathrm{b} = \sqrt{m_\Psi^2 - q_\mathrm{b}^2}$, the x_0 mass	DERIVED

part	result	status
Kohn-Sham	self-consistent states of a homogeneous gas of the particles χ (S-14 to S-19) in three regimes up to $G_{\mathrm{KS}} = 1$: first-order perturbation theory in the dilute limit, saturation of the proper density far from the fold, the level raised through $\omega = m_\Psi$ without unbinding, a positive pressure that vanishes in the dilute limit (cold dark matter); the Higgs-portal loop converges; a weakly bound point compatible with direct detection	DERIVED
residuals	every field equation is satisfied to the stated accuracy, except the radial residual of the ad hoc vacuum and the extra-time component of the junction, which are carried by stated assumptions	DERIVED
neutrinos	the flavour conversion of UM-3 is that of the Standard Model with three Dirac neutrinos: solar survival 0.55 at low and 0.30 at high energy (Borexino 0.57, 0.53, 0.43, 0.37); adiabatic supernova resonances; fast instabilities within centimetres wherever the electron lepton number crosses; equipartition raises or lowers the heating behind the shock depending on the spectra	DERIVED, ESTIMATE
thermal history	χ is a thermal relic for $\langle \sigma_{\mathrm{ann}} \mathsf{v} \rangle = 2.2 \times 10^{-26}$ cm^3/s in the normalisation of FE-61 (one χ with one $\bar{\chi}$: 9×10^{-26} cm^3/s); quark portals cannot provide it (direct detection); lepton portals can above about 1.5 TeV; the Higgs portal requires $q_{\mathrm{b}}/\omega_{\mathrm{b}} \lesssim 0.06$ at 100 GeV to 0.02 at 10 TeV; the right-handed neutrinos must not thermalise	DERIVED, ESTIMATE
sources of the fold	the Lovelock terms give the fold a graviton term of its own, but its gravitational waves move with $c_{\mathrm{g}}^2 = 83/82$; an induced Einstein-Hilbert term must provide all but 1.3×10^{-13} of the gravity of the fold; its fields induce it with a cutoff at the Planck scale; the fold fluid is a tension with 1/62 of pressureless matter, fed by the bulk; the stabiliser is the energy density of the bulk at which ordinary space stands still	DERIVED, ESTIMATE
speed of χ	$c_{\mathrm{b}}^2 = 1.190$ (S-13: 1.2); group velocity at most 1.0045, front velocity that of light; the bulk is a shortcut for signals	DERIVED
Krein norm	the vacuum is stable; the dilute gas of χ is stable, the dense gas grows; chiral couplings at the fold produce ghosts, so UM-3 with them is not unitary	DERIVED
CMB, CNB, BAO	with the Planck 2018 parameters UM-3 reproduces the CMB ($\chi^2 = 584$ for 613 high-l bins) and its derived quantities; the BAO of DESI DR2 show the same tension as the cosmological-constant model (2.3 standard deviations, DESI's number), since the dark energy of UM-3 has $w = -1$ (its pressure is minus its energy density); the CNB of Dirac neutrinos has one helicity per state	DERIVED, STANDARD (the DESI tension)
neutrino parameters	NuFIT 6.0 changes the results by at most 1.5 per cent; the Dirac phase drops out of every result; DESI DR2 excludes the inverted ordering and bounds the lightest mass below 4.12 meV, which lowers the CNB capture rate from 6.9 to 4.5 per year at most	DERIVED

3 The background

The order 0 of the power series of the metric is the author's metric itself, with his two functions of the time, a_4 and b_4 (b_4 is a function of x_4 and never identically 1, by the author's order). Its exact curvature and its exact field equations in the bulk and at the fold are restated here; the solution asks which functions $a_4(x_4)$ and $b_4(x_4)$ they admit and what they require.

$$
\begin{aligned}
g_{MN} &= \mathrm{diag}\Big(\cot^2 z,\ \mathrm{e}^{2a_4}(\sin^2 z)^{1/6},\ \mathrm{e}^{2a_4}(\sin^2 z)^{1/6},\ \mathrm{e}^{2a_4}(\sin^2 z)^{1/6},\ -1,\\
&\qquad -\mathrm{e}^{-2a_4}b_4^2(\sin^2 z)^{1/6},\ -\mathrm{e}^{-2a_4}b_4^2(\sin^2 z)^{1/6},\ -\mathrm{e}^{-2a_4}b_4^2(\sin^2 z)^{1/6}\Big), \qquad z = 6Hx_0,\\
\sigma &= \mathrm{sgn}\Big(\frac{\pi}{2} - z\Big)\frac{\ln\sin z}{6H}, \qquad \mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0,\\
g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N &= \mathrm{d}\sigma^2 + \mathrm{e}^{-2H|\sigma|}\Big[\mathrm{e}^{2a_4}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{e}^{-2a_4}b_4^2\big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big)\Big] - \mathrm{d}x_4^2
\end{aligned}
\tag{FE-27}
$$

Symbols in (FE-27).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\ldots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0,\ldots,x_7$; a repeated index is summed. Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1,\ldots,x_7$ for a tensor of the fold, $x_1,\ldots,x_4$ for a four-dimensional tensor). Status: STANDARD.
- z: The angle variable of the hidden direction, $z = 6Hx_0$; one period of x_0 is $0 < z < \pi$, the fold is $z = \pi/2$. Unit: dimensionless. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor of the extra times is $\mathrm{e}^{-a_4}b_4$ times the warp. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- x_0: The hidden space-like coordinate (listed first); the metric depends on it through $z = 6Hx_0$, periodically with the period $\pi/(6H)$. Unit: length. Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- sgn: The sign function: $\mathrm{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0,\ldots,x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \ldots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.

The author's metric (in his coordinates $X = \{x_0,\ldots,x_7\}$ of the section DEFCHART of his notebook, with the factor b_4^2 of his order of 2026-10-24) and the proper chart: in the proper coordinate σ of the hidden direction one period of x_0 is the whole line, the fold $z = \pi/2$ is $\sigma = 0$, and ordinary space and the extra times are warped by $\mathrm{e}^{-H|\sigma|}$.

Status: AUTHOR (the metric), DERIVED (the chart).

$$
\begin{aligned}
&\mathbf{e}^{(0)} = \mathrm{d}\sigma, \quad \mathbf{e}^{(i)} = \mathrm{e}^{a_4 - H|\sigma|}\mathrm{d}x_i, \quad \mathbf{e}^{(4)} = \mathrm{d}x_4, \quad \mathbf{e}^{(r)} = b_4\,\mathrm{e}^{-a_4 - H|\sigma|}\mathrm{d}x_r,\\
&\sqrt{|g|} = b_4^3\,\mathrm{e}^{-6H|\sigma|}, \qquad \mathsf{A} = \frac{a_4'}{H}, \qquad \mathsf{U} = \frac{b_4'}{Hb_4}, \qquad \mathsf{A}_1 = \frac{\mathsf{A}'}{H}, \qquad \mathsf{U}_1 = \frac{\mathsf{U}'}{H},\\
&R^{(0)(i)}{}_{(0)(i)} = R^{(0)(r)}{}_{(0)(r)} = -H^2, \qquad R^{(0)(4)}{}_{(0)(4)} = 0, \qquad R^{(i)(j)}{}_{(i)(j)} = (\mathsf{A}^2 - 1)H^2,\\
&R^{(i)(4)}{}_{(i)(4)} = (\mathsf{A}^2 + \mathsf{A}_1)H^2, \qquad R^{(i)(r)}{}_{(i)(r)} = -(1 + \mathsf{A}^2 - \mathsf{A}\mathsf{U})H^2,\\
&R^{(4)(r)}{}_{(4)(r)} = \big((\mathsf{U} - \mathsf{A})^2 + \mathsf{U}_1 - \mathsf{A}_1\big)H^2, \qquad R^{(r)(s)}{}_{(r)(s)} = \big((\mathsf{U} - \mathsf{A})^2 - 1\big)H^2,\\
&R^{(0)(i)}{}_{(4)(i)} = \mathrm{sgn}(\sigma)\,\mathsf{A}H^2, \qquad R^{(0)(r)}{}_{(4)(r)} = \mathrm{sgn}(\sigma)\,(\mathsf{U} - \mathsf{A})H^2
\end{aligned}
\tag{FE-28}
$$

Symbols in (FE-28).

- $\mathbf{e}_a{}^M$, $\mathbf{e}^a{}_M$: The vielbein (frame field) of the metric: $g_{MN} = \eta_{ab}\,\mathbf{e}^a{}_M\mathbf{e}^b{}_N$; $\mathbf{e}_a{}^M$ is its inverse; $\mathbf{e}_{bN} = \eta_{bc}\mathbf{e}^c{}_N$; $\mathbf{e}^{(a)}$ is the frame one-form $\mathbf{e}^a{}_M\,\mathrm{d}X^M$. Unit: dimensionless. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- x: The coordinates of the fold: x_i ($i = 1, 2, 3$) of ordinary space, x_4 the time, x_r ($r = 5, 6, 7$) the extra times; a field of the fold is a function of the four observable coordinates $x = (x_1, x_2, x_3, x_4)$, and $\mathrm{d}^4x = \mathrm{d}x_1\,\mathrm{d}x_2\,\mathrm{d}x_3\,\mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor of the extra times is $\mathrm{e}^{-a_4}b_4$ times the warp. Unit: dimensionless. Status: AUTHOR.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 1 at the author's vacuum, 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant). Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $R^{MN}{}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor (convention $[\nabla_P, \nabla_Q]V^M = R^M{}_{NPQ}V^N$), with two indices the Ricci tensor $R_{MN} = R^P{}_{MPN}$, without indices the scalar curvature $R = g^{MN}R_{MN}$; with frame indices $R^{(a)(b)}{}_{(c)(d)}$ its frame components, $R^{(a)(b)}{}_{(a)(b)}$ (no sum) the sectional curvature of the frame plane (a, b). Unit: 1/length^2. Status: STANDARD.
- j: A second generation or mass-state index (as i). Status: STANDARD.
- s: A second extra-time direction of the fold (as r, different from it). Status: STANDARD.

- sgn: The sign function: $\mathrm{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.

The orthonormal frame, the volume element, the dimensionless rates A, U of the two functions of the time, and the exact frame curvature of the author's metric; the components with the extra times follow from the listed ones. Status: DERIVED.

$$
\begin{aligned}
\frac{\kappa\,{\mathcal{T}_{\mathrm{bulk}}}^{(0)}{}_{(0)}}{H^2} &= -\frac{1}{16}\Big(\mathsf{A}^6 - 3\mathsf{A}^5\mathsf{U} + 12\mathsf{A}^4\mathsf{U}^2 + 3\mathsf{A}^4\mathsf{U}_1 + 3\mathsf{A}^4 \\
&\quad + 12\mathsf{A}^3\mathsf{A}_1\mathsf{U} - 19\mathsf{A}^3\mathsf{U}^3 - 12\mathsf{A}^3\mathsf{U}\mathsf{U}_1 - 6\mathsf{A}^3\mathsf{U} - 18\mathsf{A}^2\mathsf{A}_1\mathsf{U}^2 + 9\mathsf{A}^2\mathsf{U}^4 \\
&\quad + 9\mathsf{A}^2\mathsf{U}^2\mathsf{U}_1 - 18\mathsf{A}^2\mathsf{U}^2 - 6\mathsf{A}^2\mathsf{U}_1 - 9\mathsf{A}^2 + 6\mathsf{A}\mathsf{A}_1\mathsf{U}^3 - 12\mathsf{A}\mathsf{A}_1\mathsf{U} \\
&\quad + 21\mathsf{A}\mathsf{U}^3 + 12\mathsf{A}\mathsf{U}\mathsf{U}_1 + 9\mathsf{A}\mathsf{U} + 6\mathsf{A}_1\mathsf{U}^2 + 3\mathsf{U}^4 + 3\mathsf{U}^2\mathsf{U}_1 \\
&\quad + 18\mathsf{U}^2 + 3\mathsf{U}_1 + 5\Big), \\
\frac{\kappa\,{\mathcal{T}_{\mathrm{bulk}}}^{(i)}{}_{(i)}}{H^2} &= -\frac{1}{16}\Big(\mathsf{A}^6 - 6\mathsf{A}^5\mathsf{U} - 5\mathsf{A}^4\mathsf{A}_1 + 12\mathsf{A}^4\mathsf{U}^2 + 3\mathsf{A}^4\mathsf{U}_1 \\
&\quad + 3\mathsf{A}^4 + 12\mathsf{A}^3\mathsf{A}_1\mathsf{U} - 10\mathsf{A}^3\mathsf{U}^3 - 6\mathsf{A}^3\mathsf{U}\mathsf{U}_1 + 12\mathsf{A}^3\mathsf{U} - 9\mathsf{A}^2\mathsf{A}_1\mathsf{U}^2 \\
&\quad + 6\mathsf{A}^2\mathsf{A}_1 + 3\mathsf{A}^2\mathsf{U}^4 + 3\mathsf{A}^2\mathsf{U}^2\mathsf{U}_1 - 36\mathsf{A}^2\mathsf{U}^2 - 6\mathsf{A}^2\mathsf{U}_1 - 9\mathsf{A}^2 \\
&\quad + 2\mathsf{A}\mathsf{A}_1\mathsf{U}^3 - 12\mathsf{A}\mathsf{A}_1\mathsf{U} + 18\mathsf{A}\mathsf{U}^3 + 6\mathsf{A}\mathsf{U}\mathsf{U}_1 - 6\mathsf{A}\mathsf{U} + 3\mathsf{A}_1\mathsf{U}^2 \\
&\quad - \mathsf{A}_1 + 3\mathsf{U}^4 + 3\mathsf{U}^2\mathsf{U}_1 + 24\mathsf{U}^2 + 3\mathsf{U}_1 + 5\Big), \\
\frac{\kappa\,{\mathcal{T}_{\mathrm{bulk}}}^{(4)}{}_{(4)}}{H^2} &= \frac{1}{16}\Big(5\mathsf{A}^6 - 15\mathsf{A}^5\mathsf{U} + 15\mathsf{A}^4\mathsf{U}^2 - 9\mathsf{A}^4 - 5\mathsf{A}^3\mathsf{U}^3 \\
&\quad + 18\mathsf{A}^3\mathsf{U} + 3\mathsf{A}^2 - 9\mathsf{A}\mathsf{U}^3 - 3\mathsf{A}\mathsf{U} - 3\mathsf{U}^2 + 1\Big), \\
\frac{\kappa\,{\mathcal{T}_{\mathrm{bulk}}}^{(r)}{}_{(r)}}{H^2} &= -\frac{1}{16}\Big(\mathsf{A}^6 + 5\mathsf{A}^4\mathsf{A}_1 - 3\mathsf{A}^4\mathsf{U}^2 - 2\mathsf{A}^4\mathsf{U}_1 + 3\mathsf{A}^4 \\
&\quad - 8\mathsf{A}^3\mathsf{A}_1\mathsf{U} + 2\mathsf{A}^3\mathsf{U}^3 + 2\mathsf{A}^3\mathsf{U}\mathsf{U}_1 - 24\mathsf{A}^3\mathsf{U} + 3\mathsf{A}^2\mathsf{A}_1\mathsf{U}^2 - 6\mathsf{A}^2\mathsf{A}_1 \\
&\quad + 18\mathsf{A}^2\mathsf{U}^2 - 9\mathsf{A}^2 + 6\mathsf{A}\mathsf{U}^3 + 6\mathsf{A}\mathsf{U}\mathsf{U}_1 + 24\mathsf{A}\mathsf{U} + 3\mathsf{A}_1\mathsf{U}^2 \\
&\quad + \mathsf{A}_1 + 9\mathsf{U}^2 + 2\mathsf{U}_1 + 5\Big), \\
\frac{\kappa\,{\mathcal{T}_{\mathrm{bulk}}}^{(0)}{}_{(4)}}{H^2} &= \mathrm{sgn}(\sigma)\,\frac{3}{16}\,\mathsf{U}\,\mathcal{G}, \qquad \mathcal{G} = 5\mathsf{A}^2(\mathsf{A}-\mathsf{U})^2 - 6\mathsf{A}^2 + 6\mathsf{A}\mathsf{U} + \mathsf{U}^2 + 1
\end{aligned}
\tag{FE-29}
$$

Symbols in (FE-29).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\mathrm{bulk}}$: The energy-momentum tensor of all sources of the bulk, $\mathcal{T}_\Psi + \mathcal{T}_{\mathrm{stab}}$; ${\mathcal{T}_{\mathrm{bulk}}}^{(0)}{}_{b}$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as ${\mathcal{S}^{(i)}}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 1 at the author's vacuum, 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant). Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- sgn: The sign function: $\mathrm{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

- $\mathcal{G}$: The factor of the (σ, x_4) bulk equation of the author's metric, $\mathcal{G} = 5\mathsf{A}^2(\mathsf{A}-\mathsf{U})^2 - 6\mathsf{A}^2 + 6\mathsf{A}\mathsf{U} + \mathsf{U}^2 + 1$. Unit: dimensionless. Status: DERIVED.

The bulk equations of the author's metric: the energy-momentum tensor $\mathcal{T}_{\text{bulk}}$ that the matter of the bulk must have for the metric to solve the Einstein-Lovelock equations (with the author's Lovelock weights and cosmological constant), as exact polynomials in the rates. Status: AUTHOR (the metric), DERIVED (the polynomials).

$$
\begin{aligned}
&a_4 = Hx_4, \quad b_4 = \text{constant } (\mathsf{A}=1,\ \mathsf{U}=\mathsf{A}_1=\mathsf{U}_1=0): \qquad \sum_{p=1}^{3} \alpha_{(p)}\, {\mathcal{E}_{(p)}}^{M}{}_{N} + \Lambda\, \delta^{M}_{N} = 0 \quad (\sigma \neq 0),\\
&\kappa\, {\mathcal{S}^{(i)}}_{(i)} = \kappa\, {\mathcal{S}^{(r)}}_{(r)} = -10H, \qquad \kappa\, {\mathcal{S}^{(4)}}_{(4)} = -6H,\\
&\kappa\rho_{\text{bulk}} = -H^2\Big[\frac{3}{4}\big((\mathsf{A}-1-\mathsf{U})^2 + (\mathsf{A}-1)^2\big) + \ldots\Big], \qquad \mathcal{G}(1,\mathsf{U}) = 2\mathsf{U}(3\mathsf{U}-2)
\end{aligned}
\tag{FE-31}
$$

Symbols in (FE-31).

- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor of the extra times is $\mathrm{e}^{-a_4} b_4$ times the warp. Unit: dimensionless. Status: AUTHOR.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 1 at the author's vacuum, 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant). Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations a second label of a Kohn-Sham orbital (as o, with δ_{op} the Kronecker delta of the two labels). Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)} H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- ${\mathcal{E}_{(p)}}^{M}{}_{N}$: The Lovelock tensor of order p: the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to the metric ($\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor); with frame indices its frame components. Unit: $1/\text{length}^{2p}$. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -10H^2$ (the author writes the pure number $\Lambda_{\mathrm{N}} = 10$ in $0 = -\Lambda_{\mathrm{N}} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$). Unit: GeV^2. Value: $-10H^2$. Status: AUTHOR.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with upper and lower index strings the generalised Kronecker delta $\delta^{M_1 \cdots M_n}_{N_1 \cdots N_n}$, the determinant of the $n \times n$ matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ (+1 or −1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- ${\mathcal{S}^{a}}_{b}$: The thin source of the fold: its energy-momentum tensor per unit proper length across the fold (eight-dimensional normalisation, frame components along the fold); it is fixed by the jump of the first σ-derivatives of the metric across the fold (the junction conditions; $\kappa\mathcal{S}$ is the jump of the junction potential). Unit: GeV^7. Status: DERIVED.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component

such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5,6,7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- ρ_{bulk}: The energy density of the bulk sources, $\rho_{\text{bulk}} = -\mathcal{T}_{\text{bulk}}{}^{(4)}{}_{(4)}$. Unit: GeV^8. Status: DERIVED.
- $\mathcal{G}$: The factor of the (σ, x_4) bulk equation of the author's metric, $\mathcal{G} = 5\mathsf{A}^2(\mathsf{A}-\mathsf{U})^2 - 6\mathsf{A}^2 + 6\mathsf{A}\mathsf{U} + \mathsf{U}^2 + 1$. Unit: dimensionless. Status: DERIVED.

The author's vacuum: with $a_4 = Hx_4$ and a constant b_4 the bulk is empty, and the fold carries a thin source. Status: DERIVED.

$$\begin{aligned} &\mathsf{A}_1 = \mathsf{U}_1 = 0: \qquad \mathcal{T}_{\text{bulk}}{}^{a}{}_{b} = 0 \quad\Leftrightarrow\quad \big(\mathsf{A}^2-1\big)^2 = 0, \quad \mathsf{U}\big(\mathsf{A}^2 - \mathsf{A}\mathsf{U} - 1\big) = 0, \quad \mathsf{U}^3 = 0, \\ &\frac{16\,\kappa\,\mathcal{T}_{\text{bulk}}{}^{(4)}{}_{(4)}}{H^2}\Big|_{\mathsf{U}=0} = \big(\mathsf{A}^2-1\big)^2\big(5\mathsf{A}^2+1\big) \end{aligned} \tag{S-1}$$

Symbols in (S-1).

- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- $\mathcal{T}_{\text{bulk}}$: The energy-momentum tensor of all sources of the bulk, $\mathcal{T}_\Psi + \mathcal{T}_{\text{stab}}$; $\mathcal{T}_{\text{bulk}}{}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \text{diag}(1,1,1,1,-1,-1,-1,-1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 1 at the author's vacuum, 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant). Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

The histories with constant rates. The first line is the reduced Groebner basis of the polynomial system FE-29 $= 0$ with $\mathsf{A}_1 = \mathsf{U}_1 = 0$: its only solutions are $(\mathsf{A}, \mathsf{U}) = (1, 0)$, the author's vacuum, and $(-1, 0)$, its time reverse. Both are degenerate: $\mathsf{A}^2 = 1$ is a double root and $\mathsf{U} = 0$ a triple root, so the linear response of the bulk equations to a small change of the rates vanishes there (the same degeneracy makes the transverse traceless graviton lose its time-kinetic term at the author's vacuum). Every other history with constant rates needs bulk sources: with a constant b_4 the energy density $-\mathcal{T}_{\text{bulk}}{}^{(4)}{}_{(4)}$ that the second line requires is negative for every $\mathsf{A}^2 \neq 1$. Status: DERIVED (`um3_background.py`, BG1).

$$\begin{aligned} &\mathsf{A} = 1,\ \mathsf{A}_1 = \mathsf{U}_1 = 0: \qquad \frac{\kappa\,\mathcal{T}_{\text{bulk}}{}^{(0)}{}_{(0)}}{H^2} = -\frac{\mathsf{U}^2\big(6\mathsf{U}^2+\mathsf{U}+6\big)}{8}, \qquad \frac{\kappa\,\mathcal{T}_{\text{bulk}}{}^{(i)}{}_{(i)}}{H^2} = -\frac{\mathsf{U}^3\big(3\mathsf{U}+4\big)}{8}, \\ &\frac{\kappa\,\mathcal{T}_{\text{bulk}}{}^{(4)}{}_{(4)}}{H^2} = -\frac{\mathsf{U}^2\big(7\mathsf{U}-6\big)}{8}, \qquad \frac{\kappa\,\mathcal{T}_{\text{bulk}}{}^{(r)}{}_{(r)}}{H^2} = -\frac{\mathsf{U}^2\big(\mathsf{U}+3\big)}{2}, \qquad \frac{\kappa\,\mathcal{T}_{\text{bulk}}{}^{(0)}{}_{(4)}}{H^2} = \text{sgn}(\sigma)\,\frac{3\mathsf{U}^2\big(3\mathsf{U}-2\big)}{8} \end{aligned} \tag{S-2}$$

Symbols in (S-2).

- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 1 at the author's vacuum, 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant). Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\text{bulk}}$: The energy-momentum tensor of all sources of the bulk, $\mathcal{T}_\Psi + \mathcal{T}_{\text{stab}}$; $\mathcal{T}_{\text{bulk}}{}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.

- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- sgn: The sign function: $\mathrm{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

The primordial era with a varying b_4: the author's expansion $a_4 = Hx_4$ with $b_4' = \mathsf{U}Hb_4$ for a constant U (so $b_4 \propto \mathrm{e}^{\mathsf{U}Hx_4}$). The bulk must then carry a medium whose energy density, pressures and energy flux along σ (the last expression) all start at second order in U: a slowly varying b_4 costs little. The flux vanishes for $\mathsf{U} = 2/3$, where the medium is static with $\kappa\mathcal{T}_{\mathrm{bulk}}/H^2 = (-14/27, -2/9, 2/27, -22/27)$ for (σ, ordinary space, time, extra times): the flux-free primordial branch $b_4 \propto \mathrm{e}^{2Hx_4/3}$. Status: DERIVED (BG3).

$$\begin{aligned} &\mathsf{A} = \mathsf{U} = \mathsf{A}_1 = \mathsf{U}_1 = 0: \qquad \kappa\,\mathcal{T}_{\mathrm{bulk}}{}^{a}{}_{b} = H^2\,\mathrm{diag}\Big(-\frac{5}{16}, -\frac{5}{16}, -\frac{5}{16}, -\frac{5}{16}, \frac{1}{16}, -\frac{5}{16}, -\frac{5}{16}, -\frac{5}{16}\Big), \\ &\kappa\,\mathcal{S}^{(i)}{}_{(i)} = \kappa\,\mathcal{S}^{(r)}{}_{(r)} = -\frac{61}{8}H, \qquad \kappa\,\mathcal{S}^{(4)}{}_{(4)} = -\frac{31}{4}H \end{aligned} \tag{FE-32}$$

Symbols in (FE-32).

- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 1 at the author's vacuum, 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant). Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\mathrm{bulk}}$: The energy-momentum tensor of all sources of the bulk, $\mathcal{T}_\Psi + \mathcal{T}_{\mathrm{stab}}$; $\mathcal{T}_{\mathrm{bulk}}{}^{(0)}{}_{b}$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- $\mathcal{S}^a{}_b$: The thin source of the fold: its energy-momentum tensor per unit proper length across the fold (eight-dimensional normalisation, frame components along the fold); it is fixed by the jump of the first σ-derivatives of the metric across the fold (the junction conditions; $\kappa\mathcal{S}$ is the jump of the junction potential). Unit: GeV^7. Status: DERIVED.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.

$$\mathsf{U}=\mathsf{A},\quad \mathsf{U}_1=\mathsf{A}_1\ \ \big(\mathrm{e}^{-a_4}b_4=\text{constant},\ \mathcal{Z}=1\big):$$
$$\frac{\kappa\rho_{\text{bulk}}}{H^2}=\frac{3\mathsf{A}^2-1}{16},\qquad \frac{\kappa\,{\mathcal{T}_{\text{bulk}}}^{(0)}{}_{(0)}}{H^2}=-\frac{3\mathsf{A}^4+3\mathsf{A}^2\mathsf{A}_1+18\mathsf{A}^2+3\mathsf{A}_1+5}{16},$$
$$\frac{\kappa\,{\mathcal{T}_{\text{bulk}}}^{(i)}{}_{(i)}}{H^2}=-\frac{9\mathsf{A}^2+2\mathsf{A}_1+5}{16},\qquad \frac{\kappa\,{\mathcal{T}_{\text{bulk}}}^{(r)}{}_{(r)}}{H^2}=-\frac{3\mathsf{A}^4+3\mathsf{A}^2\mathsf{A}_1+24\mathsf{A}^2+3\mathsf{A}_1+5}{16},\tag{FE-33}$$
$$\frac{\kappa\,{\mathcal{T}_{\text{bulk}}}^{(0)}{}_{(4)}}{H^2}=\operatorname{sgn}(\sigma)\,\frac{3\mathsf{A}(1+\mathsf{A}^2)}{16},\qquad \frac{\kappa\,{\mathcal{S}}^{(i)}{}_{(i)}}{H}=-\frac{57\mathsf{A}^2+40\mathsf{A}_1+183}{24},$$
$$\frac{\kappa\,{\mathcal{S}}^{(4)}{}_{(4)}}{H}=-\frac{10\mathsf{A}^2+31}{4},\qquad \frac{\kappa\,{\mathcal{S}}^{(r)}{}_{(r)}}{H}=-\frac{2\mathsf{A}^4+2\mathsf{A}^2\mathsf{A}_1+37\mathsf{A}^2+20\mathsf{A}_1+61}{8}$$

Symbols in (FE-33).

- U: The rate of change of b_4 in units of H, $\mathsf{U}=b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U}=\mathsf{A}$ with frozen extra times. Status: DERIVED.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A}=a_4'/H$. Unit: dimensionless. Value: 1 at the author's vacuum, 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant). Status: DERIVED.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1=\mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1=\mathsf{A}'/H=a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4>0$, never identically 1 (the author's order of 2026-10-24); the scale factor of the extra times is $\mathrm{e}^{-a_4}b_4$ times the warp. Unit: dimensionless. Status: AUTHOR.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4)=V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z}=1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa=8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- ρ_{bulk}: The energy density of the bulk sources, $\rho_{\text{bulk}}=-{\mathcal{T}_{\text{bulk}}}^{(4)}{}_{(4)}$. Unit: GeV^8. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda=-10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- $\mathcal{T}_{\text{bulk}}$: The energy-momentum tensor of all sources of the bulk, $\mathcal{T}_\Psi+\mathcal{T}_{\text{stab}}$; ${\mathcal{T}_{\text{bulk}}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- i: A generation index ($i=1,2,3$), which also labels the neutrino mass states ν_1,ν_2,ν_3; in a frame component such as ${\mathcal{S}}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r\in\{5,6,7\}$ (x_5,x_6,x_7); not summed (the three components are equal). Status: STANDARD.
- sgn: The sign function: $\operatorname{sgn}(y)=+1$ for $y>0$, -1 for $y<0$. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma=|\cot z|\,\mathrm{d}x_0$, $\sigma=\operatorname{sgn}(\pi/2-z)\ln(\sin z)/(6H)$ on one period $0<z<\pi$; $\sigma=0$ is the fold $z=\pi/2$, $\sigma\to\mp\infty$ the tips $z\to0,\pi$; one period of x_0 is the whole line $-\infty<\sigma<\infty$. Unit: length. Status: DERIVED.
- ${\mathcal{S}^a}_b$: The thin source of the fold: its energy-momentum tensor per unit proper length across the fold (eight-dimensional normalisation, frame components along the fold); it is fixed by the jump of the first σ-derivatives of the metric across the fold (the junction conditions; $\kappa\mathcal{S}$ is the jump of the junction potential). Unit: GeV^7. Status: DERIVED.

The static block of the late universe (a_4 and b_4 constant, FE-32) and the branch with frozen extra times ($\mathrm{e}^{-a_4}b_4$ constant, FE-33), on which ordinary space expands while the proper size of the extra times at the fold stays fixed. These are the backgrounds on which the late universe is solved. Status: DERIVED.

$$\kappa\big(\rho_{\mathrm{s}}+p_{\mathrm{s}}\big)=-\frac{3}{8}H^2,\qquad \kappa\big(\rho_{\text{bulk}}+{\mathcal{T}_{\text{bulk}}}^{(0)}{}_{(0)}\big)=-\frac{3\big(\mathsf{A}^4+\mathsf{A}^2\mathsf{A}_1+5\mathsf{A}^2+\mathsf{A}_1+2\big)}{16}H^2,$$
$$\kappa\big(\rho_{\text{bulk}}+{\mathcal{T}_{\text{bulk}}}^{(i)}{}_{(i)}\big)=-\frac{3\mathsf{A}^2+\mathsf{A}_1+3}{8}H^2\qquad(\mathsf{U}=\mathsf{A},\ \mathsf{U}_1=\mathsf{A}_1)\tag{S-3}$$

Symbols in (S-3).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- ρ_s: The energy density of the stabiliser; with frozen extra times it is the energy density of the bulk at which ordinary space stands still. Unit: GeV^8. Value: $\kappa\rho_\mathrm{s} = -H^2/16$. Status: DERIVED (the value), ASSUMED (a separate medium).
- p_s: The pressure of the stabiliser (equal along σ, ordinary space and the extra times). Unit: GeV^8. Value: $\kappa p_\mathrm{s} = -5H^2/16$. Status: DERIVED (the value), ASSUMED (a separate medium).
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- ρ_bulk: The energy density of the bulk sources, $\rho_\mathrm{bulk} = -{\mathcal{T}_\mathrm{bulk}}^{(4)}{}_{(4)}$. Unit: GeV^8. Status: DERIVED.
- $\mathcal{T}_\mathrm{bulk}$: The energy-momentum tensor of all sources of the bulk, $\mathcal{T}_\Psi + \mathcal{T}_\mathrm{stab}$; ${\mathcal{T}_\mathrm{bulk}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as ${\mathcal{S}^{(i)}}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 1 at the author's vacuum, 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant). Status: DERIVED.
- A_1: The time derivative of A in units of H, $\mathsf{A}_1 = \mathsf{A}'/H = a_4''/H^2$. Unit: dimensionless. Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- U_1: The time derivative of U in units of H, $\mathsf{U}_1 = \mathsf{U}'/H$. Unit: dimensionless. Status: DERIVED.

The null energy condition of the bulk media of the late universe: for the stabiliser of the static block (energy density ρ_s, pressure p_s) and for the medium of the frozen branch (energy density $\rho_\mathrm{bulk} = -{\mathcal{T}_\mathrm{bulk}}^{(4)}{}_{(4)}$, pressures ${\mathcal{T}_\mathrm{bulk}}^{(0)}{}_{(0)}$ along σ and ${\mathcal{T}_\mathrm{bulk}}^{(i)}{}_{(i)}$ along ordinary space), the sum of the energy density and the pressure along a space-like direction is negative for every $|\mathsf{A}| < 1$ when the expansion varies slowly (sampled with $|\mathsf{A}_1| < 5 \times 10^{-4}$: at most $-0.375H^2/\kappa$); S-3 turns positive only for a fast deceleration, $\mathsf{A}_1 < -(\mathsf{A}^4 + 5\mathsf{A}^2 + 2)/(\mathsf{A}^2 + 1)$ (at most -2) along σ or $\mathsf{A}_1 < -3(1 + \mathsf{A}^2)$ along ordinary space, far from the late universe, where $\mathsf{A}_1 = -(1 + q_0)\mathsf{A}^2$ is tiny ($q_0 \approx -0.53$ the deceleration parameter today). The bulk media that the late universe of UM-3 needs violate the null energy condition: they are not ordinary matter (a scalar field cannot provide them). Status: DERIVED (BG2).

$$
\begin{aligned}
&g_{1,\mathrm{eff}} = \frac{g_1}{\sqrt{\mathcal{Z}}}, \quad g_{2,\mathrm{eff}} = \frac{g_2}{\sqrt{\mathcal{Z}}}, \quad g_{3,\mathrm{eff}} = \frac{g_3}{\sqrt{\mathcal{Z}}}, \qquad \lambda_\mathrm{eff} = \frac{\lambda}{\mathcal{Z}}, \\
&\mathcal{Y}_\mathrm{eff} = \frac{\mathcal{Y}}{\sqrt{\mathcal{Z}}}, \qquad G_{\mathrm{N,eff}} = \frac{G_\mathrm{N}}{\mathcal{Z}}, \qquad \mathcal{Z} = \frac{V_\mathrm{t}\left(\mathrm{e}^{-a_4}b_4\right)^3}{V_*}
\end{aligned}
\tag{FE-44}
$$

Symbols in (FE-44).

- $g_{k,\mathrm{eff}}$: The effective (canonically normalised) gauge couplings of the fold when the fold factor is $\mathcal{Z}$, $g_{k,\mathrm{eff}} = g_k/\sqrt{\mathcal{Z}}$ ($k = 1, 2, 3$). Unit: dimensionless. Status: DERIVED.
- g_1: The $U(1)_Y$ gauge coupling (usually g'; not normalised for grand unification). Unit: dimensionless. Value: $g_1(m_Z) = 0.3574$. Status: STANDARD.
- g_2: The $SU(2)_L$ gauge coupling (usually g). Unit: dimensionless. Value: $g_2(m_Z) = 0.6517$. Status: STANDARD.
- g_3: The $SU(3)_C$ gauge coupling (the strong coupling g_s). Unit: dimensionless. Value: $g_3(m_Z) = 1.2177$ ($\alpha_s = 0.1180$). Status: STANDARD.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4) = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z} = 1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- λ_eff: The effective quartic Higgs coupling, $\lambda/\mathcal{Z}$. Unit: dimensionless. Status: DERIVED.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- $\mathcal{Y}_\mathrm{eff}$: The effective Yukawa matrices, $\mathcal{Y}/\sqrt{\mathcal{Z}}$ (for each of $\mathcal{Y}^u, \mathcal{Y}^d, \mathcal{Y}^e, \mathcal{Y}^\nu$). Unit: dimensionless. Status: DERIVED.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^\nu$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.

- $G_{\mathrm{N,eff}}$: The effective Newton constant of the fold, $G_{\mathrm{N}}/\mathcal{Z}$. Unit: GeV^{-2}. Status: DERIVED.
- G_{N}: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_{\mathrm{N}} = 1/M_{\mathrm{Pl}}^2$ with the Planck mass $M_{\mathrm{Pl}} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- V_{t}: The coordinate volume of the (compact) extra times, $V_{\mathrm{t}} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$. Unit: length^3. Status: ASSUMED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor of the extra times is $\mathrm{e}^{-a_4}b_4$ times the warp. Unit: dimensionless. Status: AUTHOR.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.

$$\frac{\mathrm{d}\ln\mathcal{Z}}{\mathrm{d}x_4} = 3H\big(\mathsf{U} - \mathsf{A}\big) \tag{S-4}$$

Symbols in (S-4).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4) = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z} = 1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 1 at the author's vacuum, 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant). Status: DERIVED.

The factor $\mathcal{Z}$ that the volume of the extra times gives the four-dimensional couplings of the fold (FE-44) changes at the rate S-4: with the author's vacuum and a constant b_4 it falls as e^{-3Hx_4}, so the gauge and Yukawa couplings of the Standard Model ($g/\sqrt{\mathcal{Z}}$, $\mathcal{Y}/\sqrt{\mathcal{Z}}$) grow as $\mathrm{e}^{3Hx_4/2}$ and the quartic Higgs coupling ($\lambda/\mathcal{Z}$) and Newton's constant as e^{3Hx_4}; on the flux-free branch $\mathsf{U} = 2/3$ the rates are a third of these; with frozen extra times every coupling is constant. The late universe, whose couplings are measured to be constant, must therefore have frozen extra times. Status: DERIVED (BG4).

4 The power series of the metric to first order

The matter of the fold and of dirac16complex perturbs the background. The metric is written as a power series of terms of decreasing size, and each order is solved with the sources of that order.

$$\begin{aligned}
&g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N = \mathrm{d}\sigma^2 + \mathrm{e}^{2\mathfrak{a}}\big(\mathrm{d}x_1^2 + \mathrm{d}x_2^2 + \mathrm{d}x_3^2\big) - \mathrm{e}^{2\mathfrak{n}}\,\mathrm{d}x_4^2 - \mathrm{e}^{2\mathfrak{b}}\big(\mathrm{d}x_5^2 + \mathrm{d}x_6^2 + \mathrm{d}x_7^2\big),\\
&R^{(0)(i)}{}_{(0)(i)} = -\partial_\sigma^2\mathfrak{a} - (\partial_\sigma\mathfrak{a})^2, \qquad R^{(0)(4)}{}_{(0)(4)} = -\partial_\sigma^2\mathfrak{n} - (\partial_\sigma\mathfrak{n})^2, \qquad R^{(i)(j)}{}_{(i)(j)} = -(\partial_\sigma\mathfrak{a})^2 + \mathrm{e}^{-2\mathfrak{n}}(\partial_{x_4}\mathfrak{a})^2,\\
&R^{(i)(r)}{}_{(i)(r)} = -\partial_\sigma\mathfrak{a}\,\partial_\sigma\mathfrak{b} + \mathrm{e}^{-2\mathfrak{n}}\,\partial_{x_4}\mathfrak{a}\,\partial_{x_4}\mathfrak{b}, \qquad R^{(i)(4)}{}_{(i)(4)} = -\partial_\sigma\mathfrak{a}\,\partial_\sigma\mathfrak{n} + \mathrm{e}^{-2\mathfrak{n}}\big(\partial_{x_4}^2\mathfrak{a} + (\partial_{x_4}\mathfrak{a})^2 - \partial_{x_4}\mathfrak{a}\,\partial_{x_4}\mathfrak{n}\big),\\
&R^{(0)(i)}{}_{(4)(i)} = -\mathrm{e}^{-\mathfrak{n}}\big(\partial_\sigma\partial_{x_4}\mathfrak{a} + \partial_\sigma\mathfrak{a}\,\partial_{x_4}\mathfrak{a} - \partial_{x_4}\mathfrak{a}\,\partial_\sigma\mathfrak{n}\big)
\end{aligned} \tag{FE-34}$$

Symbols in (FE-34).

- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0,\dots,x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \dots$): it runs over $x_0, x_1, \dots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \dots$), running over $x_0, \dots, x_7$; a repeated index is summed. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $X = \{x_0, \dots, x_7\}$: The author's coordinates of the eight-dimensional spacetime, $X = \{x_0, x_1, \dots, x_7\}$ with the hidden direction first (Dirac_Einstein-Lovelock_4+4_Nash.nb, section DEFCHART); X^M is the coordinate with the index M and $\mathrm{d}^8X = \mathrm{d}x_0\,\mathrm{d}x_1\cdots\mathrm{d}x_7$ the coordinate volume element. Unit: length (each coordinate). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- $\mathfrak{a}$, $\mathfrak{a}_1$: The logarithm of the scale factor of ordinary space in the general warped metric FE-34, a function of σ and x_4; for the author's metric $\mathfrak{a} = a_4 - H|\sigma|$; $\mathfrak{a}_n$ its term of order n in the power series FE-35 ($\mathfrak{a}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- $\mathfrak{n}$, $\mathfrak{n}_1$: The logarithm of the lapse in the general warped metric FE-34 ($g_{x_4x_4} = -\mathrm{e}^{2\mathfrak{n}}$); for the author's metric $\mathfrak{n} = 0$; $\mathfrak{n}_n$ its term of order n ($\mathfrak{n}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\mathfrak{b}$, $\mathfrak{b}_1$: The logarithm of the scale factor of the extra times in the general warped metric FE-34; for the author's metric $\mathfrak{b} = \ln b_4 - a_4 - H|\sigma|$; $\mathfrak{b}_n$ its term of order n ($\mathfrak{b}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- x_5: The first of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_6: The second of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- x_7: The third of the three extra time-like coordinates. Unit: length. Status: AUTHOR.
- $R^{MN}{}_{PQ}$, R_{MN}, R: The curvature of the metric g_{MN}: with four indices the Riemann tensor (convention $[\nabla_P, \nabla_Q]V^M = R^M{}_{NPQ}V^N$), with two indices the Ricci tensor $R_{MN} = R^P{}_{MPN}$, without indices the scalar curvature $R = g^{MN}R_{MN}$; with frame indices $R^{(a)(b)}{}_{(c)(d)}$ its frame components, $R^{(a)(b)}{}_{(a)(b)}$ (no sum) the sectional curvature of the frame plane (a, b). Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- j: A second generation or mass-state index (as i). Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- ∂: A partial derivative: ∂_M with respect to the coordinate X^M, ∂_μ with respect to x^μ, ∂_σ with respect to σ, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.

$$\begin{aligned}
&\mathfrak{a} = a_4 - H|\sigma| + \sum_{n\geq 1} \epsilon_{\mathrm{r}}^n\, \mathfrak{a}_n(\sigma, x_4), \qquad \mathfrak{b} = \ln b_4 - a_4 - H|\sigma| + \sum_{n\geq 1} \epsilon_{\mathrm{r}}^n\, \mathfrak{b}_n(\sigma, x_4),\\
&\mathfrak{n} = \sum_{n\geq 1} \epsilon_{\mathrm{r}}^n\, \mathfrak{n}_n(\sigma, x_4), \qquad \hat{h}_{\mu\nu} \to \hat{h}_{\mu\nu} + \epsilon_{\mathrm{c}}\, \hat{h}^{(1)}_{\mu\nu} + \dots
\end{aligned} \tag{FE-35}$$

Symbols in (FE-35).

- $\mathfrak{a}$, $\mathfrak{a}_1$: The logarithm of the scale factor of ordinary space in the general warped metric FE-34, a function of σ and x_4; for the author's metric $\mathfrak{a} = a_4 - H|\sigma|$; $\mathfrak{a}_n$ its term of order n in the power series FE-35 ($\mathfrak{a}_1$ the

first-order term). Unit: dimensionless. Status: DERIVED.

- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- n: The order $n = 1, 2, \ldots$ of a term of the power series of the metric (also its exponent of ϵ_r); $\mathfrak{a}_n$, $\mathfrak{b}_n$, $\mathfrak{n}_n$ are the terms of order n. Status: STANDARD.
- ϵ_r: The bookkeeping parameter of the response of the bulk to the matter of the fold: the ratio of the bulk to the fold gravity, $\epsilon_\mathrm{r} = \kappa_4 V_*/(\kappa H)$. Unit: dimensionless. Value: below 1.344×10^{-13}: the four-dimensional gravitational wave has $c_\mathrm{g}^2 = (96+83\epsilon_\mathrm{r})/(96+82\epsilon_\mathrm{r})$, and the gravitational waves and the light of the neutron-star merger GW170817 arrived together, $(c_\mathrm{g} - 1) < 7 \times 10^{-16}$. Status: DERIVED.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\mathfrak{b}$, $\mathfrak{b}_1$: The logarithm of the scale factor of the extra times in the general warped metric FE-34; for the author's metric $\mathfrak{b} = \ln b_4 - a_4 - H|\sigma|$; $\mathfrak{b}_n$ its term of order n ($\mathfrak{b}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor of the extra times is $\mathrm{e}^{-a_4} b_4$ times the warp. Unit: dimensionless. Status: AUTHOR.
- $\mathfrak{n}$, $\mathfrak{n}_1$: The logarithm of the lapse in the general warped metric FE-34 ($g_{x_4 x_4} = -\mathrm{e}^{2\mathfrak{n}}$); for the author's metric $\mathfrak{n} = 0$; $\mathfrak{n}_n$ its term of order n ($\mathfrak{n}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $\hat{h}_{\mu\nu}$: The four-dimensional metric of the fold, $\hat{h}_{\mu\nu} = \mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the coordinates $x_1, \ldots, x_4$ (the induced metric at $\sigma = 0$ without the extra times); $\hat{h}^{\mu\nu}$ its inverse, $\hat{h}$ its determinant, $\sqrt{-\hat{h}} = \mathrm{e}^{3a_4}$; $\hat{h}^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- ϵ_c: The bookkeeping parameter of the cosmological perturbations of the metric of the fold (the size of the CMB anisotropies). Unit: dimensionless. Value: $\sim 10^{-5}$. Status: STANDARD.

The general warped metric with the symmetries of the author's metric (FE-34) and its power series (FE-35): the bulk response is counted by ϵ_r, the ratio of bulk to fold gravity, and the cosmological perturbations of the fold by ϵ_c. Status: ASSUMED (the form of the series, an ansatz), DERIVED (its symmetries).

$$
\begin{aligned}
\kappa\,\Delta{\mathcal{T}_\mathrm{bulk}}^{(0)}{}_{(0)} &= -\frac{75H}{16}\left(\mathfrak{a}_1' + \mathfrak{b}_1'\right) - \frac{3H}{8}\,\mathfrak{n}_1', \\
\kappa\,\Delta{\mathcal{T}_\mathrm{bulk}}^{(i)}{}_{(i)} &= -\frac{15H}{4}\mathfrak{a}_1' - \frac{45H}{8}\mathfrak{b}_1' - \frac{5H}{16}\mathfrak{n}_1' + \frac{5}{8}\mathfrak{a}_1'' + \frac{15}{16}\mathfrak{b}_1'' + \frac{1}{16}\mathfrak{n}_1'', \\
\kappa\,\Delta{\mathcal{T}_\mathrm{bulk}}^{(4)}{}_{(4)} &= -\frac{21H}{16}\left(\mathfrak{a}_1' + \mathfrak{b}_1'\right) + \frac{3}{16}\left(\mathfrak{a}_1'' + \mathfrak{b}_1''\right), \\
\kappa\,\Delta{\mathcal{T}_\mathrm{bulk}}^{(r)}{}_{(r)} &= -\frac{45H}{8}\mathfrak{a}_1' - \frac{15H}{4}\mathfrak{b}_1' - \frac{5H}{16}\mathfrak{n}_1' + \frac{15}{16}\mathfrak{a}_1'' + \frac{5}{8}\mathfrak{b}_1'' + \frac{1}{16}\mathfrak{n}_1'' \qquad (\sigma > 0)
\end{aligned}
\tag{S-5}
$$

Symbols in (S-5).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- $\mathcal{T}_\mathrm{bulk}$: The energy-momentum tensor of all sources of the bulk, $\mathcal{T}_\Psi + \mathcal{T}_\mathrm{stab}$; ${\mathcal{T}_\mathrm{bulk}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.

- r: An extra-time direction of the fold: $r \in \{5,6,7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- $\mathfrak{a}$, $\mathfrak{a}_1$: The logarithm of the scale factor of ordinary space in the general warped metric FE-34, a function of σ and x_4; for the author's metric $\mathfrak{a} = a_4 - H|\sigma|$; $\mathfrak{a}_n$ its term of order n in the power series FE-35 ($\mathfrak{a}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $\mathfrak{b}$, $\mathfrak{b}_1$: The logarithm of the scale factor of the extra times in the general warped metric FE-34; for the author's metric $\mathfrak{b} = \ln b_4 - a_4 - H|\sigma|$; $\mathfrak{b}_n$ its term of order n ($\mathfrak{b}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $\mathfrak{n}$, $\mathfrak{n}_1$: The logarithm of the lapse in the general warped metric FE-34 ($g_{x_4x_4} = -\mathrm{e}^{2\mathfrak{n}}$); for the author's metric $\mathfrak{n} = 0$; $\mathfrak{n}_n$ its term of order n ($\mathfrak{n}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

The first-order terms of the bulk equations around the static block for static perturbations $\mathfrak{a} = -H|\sigma| + \epsilon_\mathrm{r}\mathfrak{a}_1(\sigma)$, $\mathfrak{b} = -H|\sigma| + \epsilon_\mathrm{r}\mathfrak{b}_1(\sigma)$, $\mathfrak{n} = \epsilon_\mathrm{r}\mathfrak{n}_1(\sigma)$ ($a_4 = 0$ and $b_4 = 1$ by normalisation of the coordinates): the energy-momentum tensor of the bulk changes by $\epsilon_\mathrm{r}\,\Delta\mathcal{T}_\mathrm{bulk}$, whose components are linear in the first and second derivatives of the three functions with constant coefficients (the side $\sigma < 0$ is the mirror image). Status: DERIVED (`um3_series.py`, SR1, from the exact equations of the general metric FE-34).

$$\mathfrak{a}_1' = -\mathfrak{b}_1' = C_\mathrm{d}\,\mathrm{e}^{6H|\sigma|}, \qquad \mathfrak{n}_1' = 0 \tag{S-6}$$

Symbols in (S-6).

- $\mathfrak{a}$, $\mathfrak{a}_1$: The logarithm of the scale factor of ordinary space in the general warped metric FE-34, a function of σ and x_4; for the author's metric $\mathfrak{a} = a_4 - H|\sigma|$; $\mathfrak{a}_n$ its term of order n in the power series FE-35 ($\mathfrak{a}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $\mathfrak{b}$, $\mathfrak{b}_1$: The logarithm of the scale factor of the extra times in the general warped metric FE-34; for the author's metric $\mathfrak{b} = \ln b_4 - a_4 - H|\sigma|$; $\mathfrak{b}_n$ its term of order n ($\mathfrak{b}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $\mathfrak{n}$, $\mathfrak{n}_1$: The logarithm of the lapse in the general warped metric FE-34 ($g_{x_4x_4} = -\mathrm{e}^{2\mathfrak{n}}$); for the author's metric $\mathfrak{n} = 0$; $\mathfrak{n}_n$ its term of order n ($\mathfrak{n}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- C_d: The amplitude of the difference mode S-6, the only static first-order correction of the metric when the bulk medium does not respond: $\mathfrak{a}_1' = -\mathfrak{b}_1' = C_\mathrm{d}\mathrm{e}^{6H|\sigma|}$. Unit: 1/length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

If the medium of the bulk does not respond ($\Delta\mathcal{T}_\mathrm{bulk} = 0$), the four equations S-5 for the three functions have exactly one solution of exponential form, the difference mode S-6 (the only root of all the 3×3 minors is the exponent $6H$): ordinary space and the extra times change in opposite directions, with an amplitude C_d that grows away from the fold. No bounded static correction to the metric exists then. Status: DERIVED (SR2).

$$\begin{aligned}
&\kappa\,\Delta\mathcal{S}^{(i)}{}_{(i)} = \frac{5}{8}\big[\mathfrak{a}_1'\big] + \frac{15}{16}\big[\mathfrak{b}_1'\big] + \frac{1}{16}\big[\mathfrak{n}_1'\big], \qquad \kappa\,\Delta\mathcal{S}^{(4)}{}_{(4)} = \frac{3}{16}\Big(\big[\mathfrak{a}_1'\big] + \big[\mathfrak{b}_1'\big]\Big),\\
&\kappa\,\Delta\mathcal{S}^{(r)}{}_{(r)} = \frac{15}{16}\big[\mathfrak{a}_1'\big] + \frac{5}{8}\big[\mathfrak{b}_1'\big] + \frac{1}{16}\big[\mathfrak{n}_1'\big], \qquad \Big(\big[\mathfrak{a}_1'\big], \big[\mathfrak{b}_1'\big], \big[\mathfrak{n}_1'\big]\Big) = c_\mathrm{rad}\,(1,-1,5): \quad \kappa\,\Delta\mathcal{S}^{(r)}{}_{(r)} = \frac{5}{8}\,c_\mathrm{rad}
\end{aligned} \tag{S-7}$$

Symbols in (S-7).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- $\mathcal{S}^a{}_b$: The thin source of the fold: its energy-momentum tensor per unit proper length across the fold (eight-dimensional normalisation, frame components along the fold); it is fixed by the jump of the first σ-derivatives of the metric across the fold (the junction conditions; $\kappa\mathcal{S}$ is the jump of the junction potential). Unit: GeV^7. Status: DERIVED.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- $\mathfrak{a}$, $\mathfrak{a}_1$: The logarithm of the scale factor of ordinary space in the general warped metric FE-34, a function of σ and x_4; for the author's metric $\mathfrak{a} = a_4 - H|\sigma|$; $\mathfrak{a}_n$ its term of order n in the power series FE-35 ($\mathfrak{a}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $\mathfrak{b}$, $\mathfrak{b}_1$: The logarithm of the scale factor of the extra times in the general warped metric FE-34; for the author's metric $\mathfrak{b} = \ln b_4 - a_4 - H|\sigma|$; $\mathfrak{b}_n$ its term of order n ($\mathfrak{b}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $\mathfrak{n}$, $\mathfrak{n}_1$: The logarithm of the lapse in the general warped metric FE-34 ($g_{x_4x_4} = -\mathrm{e}^{2\mathfrak{n}}$); for the author's metric $\mathfrak{n} = 0$; $\mathfrak{n}_n$ its term of order n ($\mathfrak{n}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- c_{rad}: The amplitude of the radion mode: the first-order jumps at the fold $([\mathfrak{a}_1'], [\mathfrak{b}_1'], [\mathfrak{n}_1']) = c_{\mathrm{rad}}(1, -1, 5)$, which change only the extra-time component of the junction. Unit: 1/length. Value: $3.806\,\mathsf{A}^2 H/\epsilon_{\mathrm{r}}$ today. Status: DERIVED.

The first-order junction conditions at the fold: the jumps $[X] = X(0^+) - X(0^-)$ of the first derivatives of the three functions give the first-order thin source $\epsilon_{\mathrm{r}}\,\Delta\mathcal{S}$ (computed from the junction potential, whose jump across the fold equals κ times the thin source, and, independently, from the coefficients of the second derivatives in S-5, with the same result). The difference mode carries no energy and opposite stresses, $(-5/16, 0, 5/16)$ per unit jump; the combination of jumps in the ratio $1 : -1 : 5$, the radion mode of amplitude c_{rad}, changes only the extra-time component. Status: DERIVED (SR3).

$$
\begin{aligned}
&\frac{1}{\kappa_4}\,\hat{G}^{\mu}{}_{\nu} = \mathcal{T}_{\mathrm{SM}}{}^{\mu}{}_{\nu} + \mathcal{T}_{\mathrm{P}}{}^{\mu}{}_{\nu} + \mathcal{T}_{\mathrm{ff}}{}^{\mu}{}_{\nu} - \frac{V_*}{\kappa}\,\mathcal{J}^{\mu}{}_{\nu}, \qquad \mathcal{J}^a{}_b = \kappa\,\mathcal{S}^a{}_b \text{ (the junction condition)},\\
&H_4^2 = \frac{8\pi G_{\mathrm{N}}}{3}\big(\rho_{\mathrm{SM}} + \rho_\chi + \rho_\Lambda\big), \qquad 8\pi G_{\mathrm{N}} = \frac{\kappa_4}{1 + 5\epsilon_{\mathrm{r}}/6}, \qquad \epsilon_{\mathrm{r}} = \frac{\kappa_4 V_*}{\kappa H}, \qquad \rho_\Lambda = \rho_{\mathrm{ff}} - \frac{31 H V_*}{4\kappa},\\
&\frac{\hat{R}}{2\kappa_4} + q_{\mathrm{SM}} + q_\chi + q_{\mathrm{P}} + q_{\mathrm{ff}} = \frac{V_*}{\kappa}\,\mathcal{J}^{(r)}{}_{(r)}, \qquad q_{\mathrm{SM}} = \big\langle \mathcal{L}_{\mathrm{H}} + \mathcal{L}_{\mathrm{gauge}} + \mathcal{L}_{\mathrm{f}} + \mathcal{L}_{\mathrm{Y}} \big\rangle, \qquad q_{\mathrm{P}} = \langle \mathcal{L}_{\mathrm{P}} \rangle
\end{aligned}
\tag{FE-59}
$$

Symbols in (FE-59).

- κ_4: The coupling of the induced gravity of the fold, $\kappa_4 = 8\pi G_{\mathrm{ind}}$; four-dimensional gravity on the fold has $8\pi G_{\mathrm{N}} = \kappa_4/(1 + 5\epsilon_{\mathrm{r}}/6)$ (the Friedmann equation of the fold to first order in ϵ_{r}). Unit: GeV^{-2}. Value: $\kappa_4 \simeq 8\pi G_{\mathrm{N}} = 8\pi/M_{\mathrm{Pl}}^2$. Status: ASSUMED.
- $\hat{G}^{\mu}{}_{\nu}$: The Einstein tensor of the four-dimensional metric $\hat{h}_{\mu\nu}$ of the fold. Unit: $1/\mathrm{length}^2$. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+, +, +, -$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- $\mathcal{T}_{\mathrm{SM}}$: The energy-momentum tensor of the Standard Model on the fold, $\mathcal{T}_{\mathrm{H}} + \mathcal{T}_{\mathrm{gauge}} + \mathcal{T}_{\mathrm{f}}$. Unit: GeV^4. Status: STANDARD.
- $\mathcal{T}_{\mathrm{P}}$: The energy-momentum tensor of the portals of dirac16complex. Unit: GeV^4. Status: DERIVED.
- $\mathcal{T}_{\mathrm{ff}}{}^a{}_b$: The energy-momentum tensor of the fold fluid (frame components of the fold, four-dimensional normalisation). Unit: GeV^4. Status: ASSUMED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4} b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4} b_4$ is constant; it converts the

eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{J}^a{}_b$: The jump of the junction potential across the fold computed from the metric: the geometric side of the junction condition $\mathcal{J} = \kappa\mathcal{S}$, which equates it to κ times the energy-momentum tensor $\mathcal{S}$ of everything on the fold. Unit: GeV. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- $\mathcal{S}^a{}_b$: The thin source of the fold: its energy-momentum tensor per unit proper length across the fold (eight-dimensional normalisation, frame components along the fold); it is fixed by the jump of the first σ-derivatives of the metric across the fold (the junction conditions; $\kappa\mathcal{S}$ is the jump of the junction potential). Unit: GeV^7. Status: DERIVED.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_N: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_\mathrm{N} = 1/M_\mathrm{Pl}^2$ with the Planck mass $M_\mathrm{Pl} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- ρ_SM: The energy density of the Standard-Model particles. Unit: GeV^4. Status: STANDARD.
- ρ_χ: The energy density of the dark matter, the bound states χ of dirac16complex; $\rho_\chi^{(0)}$ its value today. Unit: GeV^4. Status: DERIVED.
- ρ_Λ: The vacuum energy density of the fold, the dark energy of UM-3: $\rho_\Lambda = \rho_\mathrm{ff} - 31HV_*/(4\kappa)$ (FE-59). Unit: GeV^4. Status: DERIVED.
- ϵ_r: The bookkeeping parameter of the response of the bulk to the matter of the fold: the ratio of the bulk to the fold gravity, $\epsilon_\mathrm{r} = \kappa_4 V_*/(\kappa H)$. Unit: dimensionless. Value: below 1.344×10^{-13}: the four-dimensional gravitational wave has $c_\mathrm{g}^2 = (96+83\epsilon_\mathrm{r})/(96+82\epsilon_\mathrm{r})$, and the gravitational waves and the light of the neutron-star merger GW170817 arrived together, $(c_\mathrm{g} - 1) < 7 \times 10^{-16}$. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- ρ_ff: The energy density of the fold fluid. Unit: GeV^4. Value: $31HV_*/(4\kappa)$ at the static block. Status: ASSUMED.
- $\hat{R}$: The scalar curvature of the metric $\hat{h}_{\mu\nu}$ of the fold. Unit: 1/length^2. Status: STANDARD.
- q_SM: The pressure of the Standard Model along the extra times at the fold: the mean on-shell Lagrangian density of the Standard-Model fields (the extra-time component of the energy-momentum tensor of every term of the fold is its Lagrangian, because the fields of the fold do not depend on the extra times). Unit: GeV^4. Status: DERIVED.
- q_χ: The pressure of the bound states of dirac16complex along the extra times, integrated across the fold (their on-shell Lagrangian, the self-interaction energy per volume; zero for a dilute gas). Unit: GeV^4. Status: DERIVED.
- q_P: The pressure of the portals along the extra times: their mean on-shell Lagrangian density. Unit: GeV^4. Status: DERIVED.
- q_ff: The pressure of the fold fluid along the extra times. Unit: GeV^4. Value: $-61HV_*/(8\kappa)$ at the static block. Status: ASSUMED.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- $\mathcal{L}_\mathrm{H}$: The Lagrangian of the Higgs doublet on the fold. Unit: GeV^4. Status: AUTHOR.
- $\mathcal{L}_\mathrm{gauge}$: The Lagrangian of the gauge fields of $SU(3)_C \times SU(2)_L \times U(1)_Y$ on the fold. Unit: GeV^4. Status: STANDARD.
- $\mathcal{L}_\mathrm{f}$: The kinetic Lagrangian of the quarks and leptons on the fold. Unit: GeV^4. Status: STANDARD.
- $\mathcal{L}_\mathrm{Y}$: The Yukawa Lagrangian of the quarks and leptons with the Higgs doublet. Unit: GeV^4. Status: STANDARD.

- $\mathcal{L}_\mathrm{P}$: The Lagrangian of the portals of dirac16complex at the fold. Unit: GeV^4. Status: INVENTED.

The fold equations to leading order in ϵ_r (FE-59): Einstein's equations of the fold with the sources of the fold, and the additional extra-time component. Status: DERIVED.

$$\begin{aligned} &3H_4^2 = \kappa_4\,\rho, \qquad 2H_4' + 3H_4^2 = -\kappa_4\,\mathcal{P}, \qquad q_\mathrm{SM} + q_\chi + q_\mathrm{P} + q_\mathrm{ff} + \frac{\rho - 3\mathcal{P}}{2} = 0, \\ &\text{rigid bulk medium:} \qquad 5H_4' + 9H_4^2 = -\kappa_4\big(\mathcal{P} + q_\mathrm{SM} + q_\chi + q_\mathrm{P} + q_\mathrm{ff}\big) \end{aligned} \tag{S-8}$$

Symbols in (S-8).

- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- κ_4: The coupling of the induced gravity of the fold, $\kappa_4 = 8\pi G_\mathrm{ind}$; four-dimensional gravity on the fold has $8\pi G_\mathrm{N} = \kappa_4/(1 + 5\epsilon_\mathrm{r}/6)$ (the Friedmann equation of the fold to first order in ϵ_r). Unit: GeV^{-2}. Value: $\kappa_4 \simeq 8\pi G_\mathrm{N} = 8\pi/M_\mathrm{Pl}^2$. Status: ASSUMED.
- ρ: The total energy density of the matter of the fold (all species); $\Delta\rho$ its perturbation. Unit: GeV^4. Status: STANDARD.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\mathcal{P}$: The total pressure of the matter of the fold; $\Delta\mathcal{P}$ its perturbation. Unit: GeV^4. Status: STANDARD.
- q_SM: The pressure of the Standard Model along the extra times at the fold: the mean on-shell Lagrangian density of the Standard-Model fields (the extra-time component of the energy-momentum tensor of every term of the fold is its Lagrangian, because the fields of the fold do not depend on the extra times). Unit: GeV^4. Status: DERIVED.
- q_χ: The pressure of the bound states of dirac16complex along the extra times, integrated across the fold (their on-shell Lagrangian, the self-interaction energy per volume; zero for a dilute gas). Unit: GeV^4. Status: DERIVED.
- q_P: The pressure of the portals along the extra times: their mean on-shell Lagrangian density. Unit: GeV^4. Status: DERIVED.
- q_ff: The pressure of the fold fluid along the extra times. Unit: GeV^4. Value: $-61HV_*/(8\kappa)$ at the static block. Status: ASSUMED.

To leading order in ϵ_r the time and space components of the junction at the fold are the Friedmann and the acceleration equation of general relativity (ρ and $\mathcal{P}$ the total energy density and pressure of the matter of the fold, $H_4 = a_4'$), and the extra-time component adds the radion condition (the first line, last equation): the sum over the matter of the fold of its pressure q along the extra times (for every term of the fold its on-shell Lagrangian density) and $(\rho - 3\mathcal{P})/2$ must vanish. Radiation satisfies it ($q = 0$, $\rho = 3\mathcal{P}$); non-relativistic matter ($q = 0$ for the bound states χ of dirac16complex, $q = -\rho$ for classical particles of the fold) and vacuum energy ($\mathcal{P} = q = -\rho$) do not. If the bulk medium is rigid, the only first-order mode, the difference mode, must absorb the mismatch, and the space and extra-time components combine into the second line: radiation still expands as $x_4^{1/2}$, but dust would expand as $x_4^{5/6}$ ($q = -\rho$) or $x_4^{5/9}$ ($q = 0$), and vacuum energy would give $H_4' = -\frac{3}{5}H_4^2$ instead of a constant expansion rate, which observation excludes. UM-3 therefore requires that the bulk medium respond with the radion mode, which keeps the four-dimensional equations those of general relativity. Status: DERIVED (SR4, SR5), ASSUMED (the response).

$$\frac{c_\mathrm{rad}}{H} = \frac{8}{5}\,\frac{3\mathsf{A}^2}{\epsilon_\mathrm{r}}\,\frac{\Delta_\mathrm{r}}{\rho_\mathrm{c}^{(0)}} = 3.806\,\frac{\mathsf{A}^2}{\epsilon_\mathrm{r}}, \qquad \Delta_\mathrm{r} = q_\mathrm{SM} + q_\chi + q_\mathrm{P} + q_\mathrm{ff} + \frac{\rho - 3\mathcal{P}}{2} = 0.793\,\rho_\mathrm{c}^{(0)} \tag{S-9}$$

Symbols in (S-9).

- c_rad: The amplitude of the radion mode: the first-order jumps at the fold $([\mathfrak{a}_1'], [\mathfrak{b}_1'], [\mathfrak{n}_1']) = c_\mathrm{rad}(1, -1, 5)$, which change only the extra-time component of the junction. Unit: 1/length. Value: $3.806\,\mathsf{A}^2H/\epsilon_\mathrm{r}$ today. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 1 at the author's vacuum, 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant). Status: DERIVED.

- $\epsilon_{\rm r}$: The bookkeeping parameter of the response of the bulk to the matter of the fold: the ratio of the bulk to the fold gravity, $\epsilon_{\rm r} = \kappa_4 V_*/(\kappa H)$. Unit: dimensionless. Value: below 1.344×10^{-13}: the four-dimensional gravitational wave has $c_{\rm g}^2 = (96+83\epsilon_{\rm r})/(96+82\epsilon_{\rm r})$, and the gravitational waves and the light of the neutron-star merger GW170817 arrived together, $(c_{\rm g} - 1) < 7 \times 10^{-16}$. Status: DERIVED.
- $\Delta_{\rm r}$: The mismatch of the radion condition: the sum over the matter of the fold of its pressure along the extra times and $(\rho - 3\mathcal{P})/2$ (zero for radiation). Unit: GeV^4. Value: $0.793\,\rho_{\rm c}^{(0)}$ today. Status: DERIVED.
- $\rho_{\rm c}$: The critical density, the energy density of a spatially flat fold at the expansion rate H_4; $\rho_{\rm c}^{(0)}$ its value today. Unit: GeV^4. Value: $\rho_{\rm c}^{(0)} = 1.0537 \times 10^{-5}\, h_{100}^2$ GeV/cm^3. Status: STANDARD.
- $q_{\rm SM}$: The pressure of the Standard Model along the extra times at the fold: the mean on-shell Lagrangian density of the Standard-Model fields (the extra-time component of the energy-momentum tensor of every term of the fold is its Lagrangian, because the fields of the fold do not depend on the extra times). Unit: GeV^4. Status: DERIVED.
- q_χ: The pressure of the bound states of dirac16complex along the extra times, integrated across the fold (their on-shell Lagrangian, the self-interaction energy per volume; zero for a dilute gas). Unit: GeV^4. Status: DERIVED.
- $q_{\rm P}$: The pressure of the portals along the extra times: their mean on-shell Lagrangian density. Unit: GeV^4. Status: DERIVED.
- $q_{\rm ff}$: The pressure of the fold fluid along the extra times. Unit: GeV^4. Value: $-61HV_*/(8\kappa)$ at the static block. Status: ASSUMED.
- ρ: The total energy density of the matter of the fold (all species); $\Delta\rho$ its perturbation. Unit: GeV^4. Status: STANDARD.
- $\mathcal{P}$: The total pressure of the matter of the fold; $\Delta\mathcal{P}$ its perturbation. Unit: GeV^4. Status: STANDARD.

The amplitude of the radion response today: the mismatch $\Delta_{\rm r}$ of the radion condition with the present content of the fold (dark matter χ with $q = 0$: $0.265\rho_{\rm c}^{(0)}$; baryons with $q = -\rho$: $0.049\rho_{\rm c}^{(0)}$; vacuum energy: $0.685\rho_{\rm c}^{(0)}$; $\rho_{\rm c}^{(0)}$ the critical density) must be carried by the jump $\frac{5}{8}c_{\rm rad}$ of S-7, which gives S-9 ($\mathsf{A} = H_4/H$). It is a small term of the power series when $\mathsf{A}^2 \ll \epsilon_{\rm r}$, that is when $H \gg H_4\,\epsilon_{\rm r}^{-1/2}$ with the Hubble rate $H_4 = 1.4 \times 10^{-42}$ GeV today: at least 3.9×10^{-36} GeV for the largest $\epsilon_{\rm r} = 1.344 \times 10^{-13}$ that the speed of gravitational waves allows (S-30), and more for a smaller $\epsilon_{\rm r}$ (which itself depends on H, $\epsilon_{\rm r} = \kappa_4 V_*/(\kappa H)$). Status: DERIVED (SR6).

5 The sources of the fold

The kink of the author's metric at the fold requires a thin source (FE-4; at the static block FE-32), and the gravity of the fold must carry gravitational waves with the speed of light. This section solves both: the speed of the four-dimensional gravitational waves with and without a gravity of the fold of its own, the gravity that the fields of the fold induce, the two parts of the fold fluid, and the constraints of the bulk equations that tie the fold to the stabiliser of the bulk (IN-7).

5.1 The gravitational waves of the fold

The gravity of the fold is the induced Einstein-Hilbert term of UM-3 (ASSUMED):

$$\mathcal{L}_{\rm IG} = \frac{\hat{R}}{2\kappa_4} \tag{IN-9}$$

Symbols in (IN-9).

- $\mathcal{L}_{\rm IG}$: The Lagrangian of the induced gravity of the fold. Unit: GeV^4. Status: ASSUMED.
- $\hat{R}$: The scalar curvature of the metric $\hat{h}_{\mu\nu}$ of the fold. Unit: 1/length^2. Status: STANDARD.
- κ_4: The coupling of the induced gravity of the fold, $\kappa_4 = 8\pi G_{\rm ind}$; four-dimensional gravity on the fold has $8\pi G_{\rm N} = \kappa_4/(1 + 5\epsilon_{\rm r}/6)$ (the Friedmann equation of the fold to first order in $\epsilon_{\rm r}$). Unit: GeV^{-2}. Value: $\kappa_4 \simeq 8\pi G_{\rm N} = 8\pi/M_{\rm Pl}^2$. Status: ASSUMED.

($\mathcal{L}_{\rm IG}$ its Lagrangian, $\hat{R}$ the scalar curvature of the four-dimensional metric of the fold, $\kappa_4 = 8\pi G_{\rm ind}$ its coupling). The speed of the gravitational waves decides how much of the gravity of the fold it must provide:

$$K_{44}\,\frac{\partial^2 h_{\rm gw}}{\partial x_4^2}+K_{33}\,\frac{\partial^2 h_{\rm gw}}{\partial x_3^2}=0, \qquad c_{\rm g}^2=-\frac{K_{33}}{K_{44}}=\frac{96+83\,\epsilon_{\rm r}}{96+82\,\epsilon_{\rm r}},$$
$$\kappa K_{44}=\frac{1}{96H}+\frac{5}{12H}+\frac{1}{2H\epsilon_{\rm r}}, \qquad -\kappa K_{33}=\frac{5}{64H}+\frac{17}{48H}+\frac{1}{2H\epsilon_{\rm r}} \tag{FE-63}$$

Symbols in (FE-63).

- K_{44}: The coefficient of the second time derivative in the equation of the zero mode of a gravitational wave, integrated across the hidden direction: the sum of the parts of the bulk, of the Lovelock terms at the fold and of the induced gravity of the fold. Unit: GeV^5 (κK_{44} is a length). Status: DERIVED.
- ∂: A partial derivative: ∂_M with respect to the coordinate X^M, ∂_μ with respect to x^μ, ∂_σ with respect to σ, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- $h_{\rm gw}$: The amplitude of a gravitational wave of the + polarisation travelling along x_3 (the metric components along x_1 and x_2 multiplied by $\mathrm{e}^{h_{\rm gw}}$ and $\mathrm{e}^{-h_{\rm gw}}$), in its zero mode, the same at every σ: the gravitational wave of the four-dimensional world. Unit: dimensionless. Status: STANDARD.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c=1$). Status: AUTHOR.
- K_{33}: The coefficient of the second derivative along x_3 in the equation of the zero mode of a gravitational wave (negative), with the same three parts. Unit: GeV^5. Status: DERIVED.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- $c_{\rm g}$: The limiting speed of the zero mode of the transverse traceless graviton (the four-dimensional gravitational wave) at the static block or with frozen extra times, in units of the speed of light on the fold. Unit: dimensionless. Value: $\sqrt{15/2}=2.74$ for the bulk alone; $\sqrt{83/82}=1.0061$ with the graviton term of the fold from the Gauss-Bonnet and cubic Lovelock terms; $\sqrt{(96+83\epsilon_{\rm r})/(96+82\epsilon_{\rm r})}$ with the induced gravity of the fold. Status: DERIVED.
- $\epsilon_{\rm r}$: The bookkeeping parameter of the response of the bulk to the matter of the fold: the ratio of the bulk to the fold gravity, $\epsilon_{\rm r}=\kappa_4 V_*/(\kappa H)$. Unit: dimensionless. Value: below 1.344×10^{-13}: the four-dimensional gravitational wave has $c_{\rm g}^2=(96+83\epsilon_{\rm r})/(96+82\epsilon_{\rm r})$, and the gravitational waves and the light of the neutron-star merger GW170817 arrived together, $(c_{\rm g}-1)<7\times10^{-16}$. Status: DERIVED.
- κ: The eight-dimensional gravitational coupling, $\kappa=8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda=-10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

The solution. The field equations of the author's metric were linearised for a gravitational wave of the + polarisation that travels along x_3, $h_{\rm gw}(\sigma,x_3,x_4)$, with frozen extra times (any constant A, the static block included) and at the author's vacuum. The coefficients of the second σ-derivatives of the linearised equation are an exact differential of the first derivatives, so the equation can be integrated across the kink of the fold, and its thin part there is $(5/(12H))\,\partial_{x_4}^2 h_{\rm gw}-(17/(48H))\,\partial_{x_3}^2 h_{\rm gw}$ (times $1/\kappa$) with frozen extra times and $(5/(12H))\,\partial_{x_4}^2 h_{\rm gw}-(13/(24H))\,\partial_{x_3}^2 h_{\rm gw}$ at the author's vacuum, plus terms in the first time derivative and in the jump of the σ-derivative. Replacing the kink by a smooth step of width w and integrating the full equation reproduces it (deviation 4.1×10^{-4} at $w=10^{-5}/H$, falling linearly with w). Einstein gravity alone has no such thin part: it comes from the Gauss-Bonnet and cubic terms of the author's action. The zero mode, the same at every σ, collects the bulk with the weight $\mathrm{e}^{-6H|\sigma|}$ and the thin part: FE-63. The bulk alone moves with $c_{\rm g}^2=15/2$ ($c_{\rm g}=2.74$); the fold carries 40/41 of the time-kinetic term and brings the zero mode to $c_{\rm g}^2=83/82$ ($c_{\rm g}=1.0061$); at the author's vacuum, where the bulk has no time-kinetic term, the fold restores it with $c_{\rm g}^2=19/10$. With the induced gravity IN-9 the zero mode approaches the light cone as $\epsilon_{\rm r}\to0$. Status: DERIVED (`um3_foldsources.py`, F1 to F4).

$$c_{\rm g}-1=\sqrt{\frac{96+83\,\epsilon_{\rm r}}{96+82\,\epsilon_{\rm r}}}-1\simeq\frac{\epsilon_{\rm r}}{192}<7\times10^{-16} \quad\Rightarrow\quad \epsilon_{\rm r}<1.344\times10^{-13} \tag{S-30}$$

Symbols in (S-30).

- $c_{\rm g}$: The limiting speed of the zero mode of the transverse traceless graviton (the four-dimensional gravitational wave) at the static block or with frozen extra times, in units of the speed of light on the fold. Unit: dimensionless. Value: $\sqrt{15/2}=2.74$ for the bulk alone; $\sqrt{83/82}=1.0061$ with the graviton term of the fold from the Gauss-Bonnet and cubic Lovelock terms; $\sqrt{(96+83\epsilon_{\rm r})/(96+82\epsilon_{\rm r})}$ with the induced gravity of the fold. Status: DERIVED.

- ϵ_r: The bookkeeping parameter of the response of the bulk to the matter of the fold: the ratio of the bulk to the fold gravity, $\epsilon_\mathrm{r} = \kappa_4 V_*/(\kappa H)$. Unit: dimensionless. Value: below 1.344×10^{-13}: the four-dimensional gravitational wave has $c_\mathrm{g}^2 = (96+83\epsilon_\mathrm{r})/(96+82\epsilon_\mathrm{r})$, and the gravitational waves and the light of the neutron-star merger GW170817 arrived together, $(c_\mathrm{g} - 1) < 7 \times 10^{-16}$. Status: DERIVED.

The bound. The gravitational waves of the neutron-star merger GW170817 and the gamma rays of its burst GRB 170817A arrived from a distance of about 40 Mpc within 1.7 s of each other, which bounds the speed of gravitational waves to $-3 \times 10^{-15} < c_\mathrm{g} - 1 < 7 \times 10^{-16}$. The gravitational waves of UM-3 are faster than light, so the upper bound applies and limits the share ϵ_r of the bulk in the gravity of the fold: the induced gravity must provide all but 1.3×10^{-13} of it. The Planck mass of the gravitational waves, $(1 + 41\epsilon_\mathrm{r}/48)/\kappa_4$, and that of the Friedmann equation, $(1 + 5\epsilon_\mathrm{r}/6)/\kappa_4$, then agree to $\epsilon_\mathrm{r}/48 < 3 \times 10^{-15}$. With this bound the condition $\mathsf{A}^2 \ll \epsilon_\mathrm{r}$ under which the radion response S-9 is a small term requires $H \gg H_4 \epsilon_\mathrm{r}^{-1/2}$, at least 3.9×10^{-36} GeV. Status: DERIVED (F4), STANDARD (the measurement).

5.2 The gravity induced by the fields of the fold

$$\frac{1}{16\pi G_\mathrm{ind}} = \frac{N_\mathrm{ind}\,\Lambda_\mathrm{c}^2}{192\pi^2}, \qquad N_\mathrm{ind} = \sum_{\text{real scalars}} \big(1 - 6\xi\big) + N_\mathrm{Weyl} - 4N_\mathrm{gauge} = 4\big(1 - 6\xi_\mathrm{H}\big) \tag{FE-64}$$

Symbols in (FE-64).

- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_ind: The Newton constant of the induced gravity of the fold, $\kappa_4 = 8\pi G_\mathrm{ind}$ (Newton's constant G_N up to the factor $1 + 5\epsilon_\mathrm{r}/6$). Unit: GeV^{-2}. Status: ASSUMED.
- N_ind: The weighted number of the fields of the fold that induce the Einstein-Hilbert term: each real scalar counts $1 - 6\xi$, each Weyl fermion 1, each gauge field -4 (with its ghost fields). Unit: dimensionless. Value: $4(1 - 6\xi_\mathrm{H})$ for the Standard Model with three right-handed neutrinos. Status: DERIVED.
- Λ_c: The cutoff of the proper-time integral of the one-loop effective action of the fields of the fold: the highest energy of the quantum fluctuations that induce gravity. Unit: GeV. Value: $\sqrt{24}\,\pi M_\mathrm{red} = 3.748 \times 10^{19}$ GeV for the measured Newton constant at $\xi_\mathrm{H} = 0$. Status: ESTIMATE.
- ξ: The coupling of a real scalar field to the scalar curvature $\hat{R}$ of the fold: the term $-\frac{1}{2}\xi\hat{R}$ times the square of the field in its Lagrangian ($\xi = 1/6$ is the conformal coupling). Unit: dimensionless. Status: STANDARD.
- N_Weyl: The number of Weyl (two-component) fermion fields of the fold. Unit: dimensionless. Value: 48 (three generations of quarks and leptons with three right-handed neutrinos). Status: STANDARD.
- N_gauge: The number of gauge fields of the fold. Unit: dimensionless. Value: 12 (eight gluons, three W, one B). Status: STANDARD.
- ξ_H: The coupling ξ of the Higgs doublet to the scalar curvature of the fold. Unit: dimensionless. Value: not fixed by UM-3; below 1/6 if the gravity of the fold is induced by its fields. Status: ASSUMED.

The solution. The one-loop effective action of every field of the fold with a proper-time cutoff Λ_c contains an Einstein-Hilbert term; its coefficient follows from the second heat-kernel coefficient of each field, the trace of $E + \hat{R}/6$ for its operator $-D^2 - E$ (E the non-derivative part): a real scalar with the curvature coupling ξ gives $1 - 6\xi$, a Weyl fermion 1, a gauge field with its ghost fields -4. In the Standard Model with three right-handed neutrinos the 48 Weyl fermions cancel the 12 gauge fields exactly and only the four real scalars of the Higgs doublet remain: $N_\mathrm{ind} = 4(1 - 6\xi_\mathrm{H})$ (without the right-handed neutrinos $1 - 24\xi_\mathrm{H}$). The measured Newton constant needs $\Lambda_\mathrm{c} = \sqrt{24}\,\pi M_\mathrm{red} = 3.748 \times 10^{19}$ GeV $= 3.07\,M_\mathrm{Pl}$ at $\xi_\mathrm{H} = 0$, and the right sign needs $\xi_\mathrm{H} < 1/6$: a gravity of the fold induced by its own fields is possible, with a cutoff at the Planck scale. Status: ESTIMATE (one loop, cutoff-dependent; `um3_foldsources.py`, F5).

5.3 The fold fluid, the stabiliser and the energy of the bulk

The two media of UM-3 that the kink and the bulk of the author's metric require are the fold fluid on the fold,

$$\mathcal{T}_\mathrm{ff}{}^a{}_b = \mathrm{diag}\big(p_\mathrm{ff}, p_\mathrm{ff}, p_\mathrm{ff}, -\rho_\mathrm{ff}, q_\mathrm{ff}, q_\mathrm{ff}, q_\mathrm{ff}\big) \tag{IN-10}$$

Symbols in (IN-10).

- $\mathcal{T}_{\mathrm{ff}}{}^{a}{}_{b}$: The energy-momentum tensor of the fold fluid (frame components of the fold, four-dimensional normalisation). Unit: GeV^4. Status: ASSUMED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- p_{ff}: The pressure of the fold fluid along ordinary space. Unit: GeV^4. Value: $-61HV_*/(8\kappa)$ at the static block. Status: ASSUMED.
- ρ_{ff}: The energy density of the fold fluid. Unit: GeV^4. Value: $31HV_*/(4\kappa)$ at the static block. Status: ASSUMED.
- q_{ff}: The pressure of the fold fluid along the extra times. Unit: GeV^4. Value: $-61HV_*/(8\kappa)$ at the static block. Status: ASSUMED.

(a medium at rest on the fold with the energy density ρ_{ff}, the pressure p_{ff} along ordinary space and the pressure q_{ff} along the extra times, ASSUMED), and the stabiliser in the bulk:

$$\mathcal{T}_{\mathrm{stab}}{}^{a}{}_{b} = \mathrm{diag}\big(p_{\mathrm{s}}, p_{\mathrm{s}}, p_{\mathrm{s}}, p_{\mathrm{s}}, -\rho_{\mathrm{s}}, p_{\mathrm{s}}, p_{\mathrm{s}}, p_{\mathrm{s}}\big), \qquad \kappa\rho_{\mathrm{s}} = -\frac{H^2}{16}, \qquad \kappa p_{\mathrm{s}} = -\frac{5H^2}{16} \tag{IN-7}$$

Symbols in (IN-7).

- $\mathcal{T}_{\mathrm{stab}}{}^{a}{}_{b}$: The energy-momentum tensor of the stabiliser of the bulk (frame components): the source that a static bulk of the author's metric requires, found and named by Claude (Anthropic) on 2026-10-24. Unit: GeV^8. Status: DERIVED (the value), ASSUMED (a separate medium).
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- p_{s}: The pressure of the stabiliser (equal along σ, ordinary space and the extra times). Unit: GeV^8. Value: $\kappa p_{\mathrm{s}} = -5H^2/16$. Status: DERIVED (the value), ASSUMED (a separate medium).
- ρ_{s}: The energy density of the stabiliser; with frozen extra times it is the energy density of the bulk at which ordinary space stands still. Unit: GeV^8. Value: $\kappa\rho_{\mathrm{s}} = -H^2/16$. Status: DERIVED (the value), ASSUMED (a separate medium).
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

(ASSUMED as a medium; its value DERIVED from the static block, FE-32).

$$\rho_{\mathrm{ff}} = \lambda_{\mathrm{F}} + \rho_{\mathrm{d}}, \qquad p_{\mathrm{ff}} = q_{\mathrm{ff}} = -\lambda_{\mathrm{F}}, \qquad \lambda_{\mathrm{F}} = \frac{61HV_*}{8\kappa}, \qquad \rho_{\mathrm{d}} = \frac{HV_*}{8\kappa}, \qquad 3a_4'\,\rho_{\mathrm{d}} = \frac{3}{8}\,\mathsf{A}\,\frac{H^2V_*}{\kappa} \tag{FE-65}$$

Symbols in (FE-65).

- ρ_{ff}: The energy density of the fold fluid. Unit: GeV^4. Value: $31HV_*/(4\kappa)$ at the static block. Status: ASSUMED.
- λ_{F}: The tension of the fold fluid: the part of the fold fluid whose pressures along ordinary space and along the extra times are minus its energy density (a vacuum energy of the seven-dimensional fold), per unit volume of ordinary space of the fold. Unit: GeV^4. Value: $61HV_*/(8\kappa)$ at the static block. Status: DERIVED.
- ρ_{d}: The pressureless part of the fold fluid: the energy density of the fold fluid minus its tension, per unit volume of ordinary space of the fold. Unit: GeV^4. Value: $HV_*/(8\kappa)$ at the static block. Status: DERIVED.

- p_{ff}: The pressure of the fold fluid along ordinary space. Unit: GeV^4. Value: $-61HV_*/(8\kappa)$ at the static block. Status: ASSUMED.
- q_{ff}: The pressure of the fold fluid along the extra times. Unit: GeV^4. Value: $-61HV_*/(8\kappa)$ at the static block. Status: ASSUMED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 1 at the author's vacuum, 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant). Status: DERIVED.

The solution for the fold fluid. The source of the fold of the static block, $\kappa\mathcal{S} = (-61/8, -31/4, -61/8)H$ along (ordinary space, the time, the extra times) (FE-32), splits uniquely into a tension, whose pressures are minus its energy density in every direction of the fold, and a pressureless part, 1/62 of the energy. The vacuum energy of every field of the fold is a tension of this kind, because the fields do not depend on the extra times and their pressure along the extra times is their on-shell Lagrangian (FE-5), the same as along ordinary space. When the fold grows with frozen extra times the tension keeps its density; the pressureless part would be diluted, and it is kept by the energy $3a_4'\rho_{\mathrm{d}}$ per unit volume and time, exactly V_* times the part linear in A of the jump of the energy flow of the bulk (FE-66). Status: DERIVED (`um3_foldsources.py`, F6; `um3_stabiliser.py`, ST6).

$$
\begin{aligned}
&\kappa\,{\mathcal{T}_{\mathrm{bulk}}}^{(4)}{}_{(4)} = \frac{H^2}{16}\,\mathsf{P}(\mathsf{A},\mathsf{U}), \qquad \kappa\,{\mathcal{T}_{\mathrm{bulk}}}^{(0)}{}_{(4)} = \frac{3}{16}\,\mathsf{U}\,\mathcal{G}(\mathsf{A},\mathsf{U})\,H^2\,\mathrm{sgn}(\sigma),\\
&\mathsf{P} = 5\mathsf{A}^6 - 15\mathsf{A}^5\mathsf{U} + 15\mathsf{A}^4\mathsf{U}^2 - 9\mathsf{A}^4 - 5\mathsf{A}^3\mathsf{U}^3 + 18\mathsf{A}^3\mathsf{U} + 3\mathsf{A}^2 - 9\mathsf{A}\mathsf{U}^3 - 3\mathsf{A}\mathsf{U} - 3\mathsf{U}^2 + 1,\\
&\mathsf{P}(\mathsf{A},\mathsf{U}) = 0 \;\Rightarrow\; \mathsf{A}^2 + \mathsf{U}^2 \geq 0.2607, \qquad \mathsf{U} = \mathsf{A}: \quad 3a_4'^2 = 16\kappa\big(\rho_{\mathrm{bulk}} - \rho_{\mathrm{s}}\big)
\end{aligned}
\tag{FE-66}
$$

Symbols in (FE-66).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\mathrm{bulk}}$: The energy-momentum tensor of all sources of the bulk, $\mathcal{T}_\Psi + \mathcal{T}_{\mathrm{stab}}$; ${\mathcal{T}_{\mathrm{bulk}}}^{(0)}{}_b$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- $\mathsf{P}(\mathsf{A},\mathsf{U})$: The polynomial of the energy constraint of the bulk: the (x_4, x_4) component of the bulk equations of the author's metric is $\kappa{\mathcal{T}_{\mathrm{bulk}}}^{(4)}{}_{(4)} = (H^2/16)\,\mathsf{P}(\mathsf{A},\mathsf{U})$, whatever the time derivatives of the rates. Unit: dimensionless. Value: $\mathsf{P}(0,0) = 1$, $\mathsf{P}(\pm1, 0) = 0$. Status: DERIVED.
- A: The expansion rate of ordinary space in units of H, $\mathsf{A} = a_4'/H$. Unit: dimensionless. Value: 1 at the author's vacuum, 0 at the static block; $H_4^{(0)}/H$ today ($H_4^{(0)}$ the Hubble constant). Status: DERIVED.
- U: The rate of change of b_4 in units of H, $\mathsf{U} = b_4'/(Hb_4)$. Unit: dimensionless. Value: $\mathsf{U} = \mathsf{A}$ with frozen extra times. Status: DERIVED.
- $\mathcal{G}$: The factor of the (σ, x_4) bulk equation of the author's metric, $\mathcal{G} = 5\mathsf{A}^2(\mathsf{A}-\mathsf{U})^2 - 6\mathsf{A}^2 + 6\mathsf{A}\mathsf{U} + \mathsf{U}^2 + 1$. Unit: dimensionless. Status: DERIVED.
- sgn: The sign function: $\mathrm{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- ρ_{bulk}: The energy density of the bulk sources, $\rho_{\mathrm{bulk}} = -{\mathcal{T}_{\mathrm{bulk}}}^{(4)}{}_{(4)}$. Unit: GeV^8. Status: DERIVED.
- ρ_{s}: The energy density of the stabiliser; with frozen extra times it is the energy density of the bulk at which

ordinary space stands still. Unit: GeV^8. Value: $\kappa\rho_\mathrm{s} = -H^2/16$. Status: DERIVED (the value), ASSUMED (a separate medium).

The solution for the bulk. Of the bulk equations of the author's metric, the energy constraint and the flow of energy along the hidden direction contain no second time derivative, for any $a_4(x_4)$ and $b_4(x_4)$ (FE-66). Without a bulk source the energy constraint $\mathsf{P} = 0$ keeps $\mathsf{A}^2 + \mathsf{U}^2$ at least 0.2607 at every instant (reached at $(\mathsf{A}, \mathsf{U}) = \pm(0.1432, 0.4901)$), and with constant rates only $(\pm 1, 0)$ are possible (S-1): a late universe that expands slowly with constant couplings needs a bulk source. The static block needs exactly the stabiliser IN-7, $\kappa\rho_\mathrm{s} = -H^2/16$ and $\kappa p_\mathrm{s} = -5H^2/16$ in every direction but the time; with frozen extra times the energy constraint is the Friedmann equation of the bulk, so ρ_s is the energy density of the bulk at which ordinary space stands still, and the expansion rate a_4' measures the excess of the bulk energy density over it. The stabiliser was found and named by Claude (Anthropic) on 2026-10-24 while deriving the field equations of UM-3. It violates the weak and the null energy condition (S-3), and the only simple field that reproduces it has a negative kinetic energy (a ghost); on the static bulk it is conserved by itself, but when ordinary space grows the bulk source that the equations require gives off the energy $-3a_4'(\rho_\mathrm{bulk} + {\mathcal{T}_\mathrm{bulk}}^{(1)}{}_{(1)})$ per volume and time, which flows along σ into the fold and is taken up there exactly by the source of the fold of FE-33 (the second line of FE-11), for any $a_4(x_4)$. A medium that couples only to gravity and keeps $p = 5\rho$ keeps $\rho\, b_4^{18}$ constant, so a stabiliser absent in the primordial era, whose bulk is the author's vacuum (FE-31), could not appear later on its own: its creation needs the exchange of energy with the fold or a change of its equation of state (OPEN). Without the stabiliser the late universe would have to be the author's vacuum with H equal to today's expansion rate, in which every squared coupling of the fold grows as e^{3Hx_4}: the fine-structure constant would have changed by $\Delta \ln \alpha_\mathrm{em} = 0.37$ since the natural reactor of Oklo 1.8 billion years ago, against the bound of about 10^{-7}. Status: DERIVED (`um3_stabiliser.py`, ST1 to ST8), OPEN (the origin of the stabiliser).

6 The Higgs field in the ad hoc vacuum

$$\begin{aligned} &\frac{1}{\sqrt{-\hat{h}}\,\mathcal{Z}}\, D_\mu\big(\sqrt{-\hat{h}}\,\mathcal{Z}\,\hat{h}^{\mu\nu} D_\nu \Phi\big) - \mu^2\Phi - 2\lambda\big(\Phi^\dagger\Phi\big)\Phi - c_\Phi\,\bar{\Psi} P_\mathrm{L} \Psi\,\Phi = \Theta, \\ &\Theta = \begin{pmatrix} \bar{d}_\mathrm{R}(\mathcal{Y}^d)^\dagger u_\mathrm{L} + \bar{e}_\mathrm{R}(\mathcal{Y}^e)^\dagger \nu_\mathrm{L} - \bar{d}_\mathrm{L}\mathcal{Y}^u u_\mathrm{R} - \bar{e}_\mathrm{L}\mathcal{Y}^\nu \nu_\mathrm{R} \\ \bar{d}_\mathrm{R}(\mathcal{Y}^d)^\dagger d_\mathrm{L} + \bar{e}_\mathrm{R}(\mathcal{Y}^e)^\dagger e_\mathrm{L} + \bar{u}_\mathrm{L}\mathcal{Y}^u u_\mathrm{R} + \bar{\nu}_\mathrm{L}\mathcal{Y}^\nu \nu_\mathrm{R} \end{pmatrix} \end{aligned} \tag{FE-15}$$

Symbols in (FE-15).

- $\hat{h}_{\mu\nu}$: The four-dimensional metric of the fold, $\hat{h}_{\mu\nu} = \mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the coordinates $x_1, \ldots, x_4$ (the induced metric at $\sigma = 0$ without the extra times); $\hat{h}^{\mu\nu}$ its inverse, $\hat{h}$ its determinant, $\sqrt{-\hat{h}} = \mathrm{e}^{3a_4}$; $\hat{h}^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4) = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z} = 1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- μ^2: The mass parameter of the Higgs potential, $\mu^2 < 0$ (it enters only as μ^2). Unit: GeV^2. Value: $\mu^2 = -\lambda v^2 = -m_h^2/2 = -(88.6\ \mathrm{GeV})^2$. Status: AUTHOR.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.

- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- Θ: The Yukawa source of the Higgs equation, $\Theta = -\partial\mathcal{L}_\mathrm{Y}/\partial\Phi^\dagger$ (a doublet of fermion bilinears). Unit: GeV^3. Status: STANDARD.
- d_R: The right-handed down-type quarks d, s, b (hypercharge $-1/3$). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^\nu$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- u_L: The upper (up-type, left-handed) components of the quark doublets. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- e_R: The right-handed electron, muon and tau (singlets, hypercharge -1). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- ν_L: The upper (neutral, left-handed) components of the lepton doublets: the left-handed neutrinos. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- d_L: The lower (down-type, left-handed) components of the quark doublets. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- u_R: The right-handed up-type quarks u, c, t (singlets of $SU(2)_L$, hypercharge $2/3$). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- ν_R: The three right-handed neutrinos (gauge singlets). Unit: $\mathrm{GeV}^{3/2}$. Status: ASSUMED.
- e_L: The lower (charged, left-handed) components of the lepton doublets. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.

$$\frac{1}{\sqrt{-\hat h}\,\mathcal{Z}}\Big(\partial_\mu\big[\sqrt{-\hat h}\,\mathcal{Z}\,\mathcal{K}_W\,W^{I\mu\nu}\big] + g_2\,\epsilon^{IJK}W^J_\mu\sqrt{-\hat h}\,\mathcal{Z}\,\mathcal{K}_W\,W^{K\mu\nu}\Big) = -g_2\big(J^{I\nu}_\mathrm{w} + J^{I\nu}_\Phi\big),$$
$$\mathcal{K}_W = 1 + \sum_X c^W_X\,\bar\Psi P_X\Psi, \qquad J^{I\nu}_\mathrm{w} = \bar Q\,\hat\gamma^\nu\frac{\tau^I}{2}Q + \bar\ell\,\hat\gamma^\nu\frac{\tau^I}{2}\ell, \qquad J^{I\nu}_\Phi = \mathrm{i}\Big[(D^\nu\Phi)^\dagger\frac{\tau^I}{2}\Phi - \Phi^\dagger\frac{\tau^I}{2}D^\nu\Phi\Big] \tag{FE-17}$$

Symbols in (FE-17).

- $\hat h_{\mu\nu}$: The four-dimensional metric of the fold, $\hat h_{\mu\nu} = \mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the coordinates $x_1, \ldots, x_4$ (the induced metric at $\sigma = 0$ without the extra times); $\hat h^{\mu\nu}$ its inverse, $\hat h$ its determinant, $\sqrt{-\hat h} = \mathrm{e}^{3a_4}$; $\hat h^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4) = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z} = 1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- ∂: A partial derivative: ∂_M with respect to the coordinate X^M, ∂_μ with respect to x^μ, ∂_σ with respect to σ, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- $\mathcal{K}_G$, $\mathcal{K}_W$, $\mathcal{K}_B$: The portal factors of the gauge kinetic terms, $\mathcal{K}_G = 1 + \sum_X c^G_X\bar\Psi P_X\Psi$ (likewise for W and B): the gauge couplings at the fold are divided by $\sqrt{\mathcal{K}}$. Unit: dimensionless. Value: 1 where dirac16complex vanishes. Status: DERIVED.
- W^I_μ, $W^I_{\mu\nu}$: The gauge field of $SU(2)_L$ on the fold, $W^1_\mu, W^2_\mu, W^3_\mu$, and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- g_2: The $SU(2)_L$ gauge coupling (usually g). Unit: dimensionless. Value: $g_2(m_Z) = 0.6517$. Status: STANDARD.
- ϵ^{IJK}: The Levi-Civita symbol of three indices (the structure constants of $SU(2)$), $\epsilon^{123} = 1$. Unit: dimensionless. Status: STANDARD.
- J: A weak-isospin (adjoint SU(2)) index (as I). Status: STANDARD.
- K: A weak-isospin (adjoint SU(2)) index (as I). Status: STANDARD.
- $J^{I\nu}_\mathrm{w}$: The weak-isospin current of the left-handed fermion doublets. Unit: GeV^3. Status: STANDARD.
- $J^{I\nu}_\Phi$: The weak-isospin current of the Higgs doublet (it contains the mass terms of the W fields); $J^{2\,x_4}_\Phi$ its isospin charge density with $I = 2$. Unit: GeV^3. Status: STANDARD.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c^W_X: The W-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.

- P_X: The chiral projector of the half X: $P_X = P_\mathrm{L}$ for $X = \mathrm{L}$, P_R for $X = \mathrm{R}$. Unit: dimensionless. Status: AUTHOR.
- Q, $\bar{Q}$: The left-handed quark doublets of the three generations, $Q = (u_\mathrm{L}, d_\mathrm{L})$ (colour triplets, hypercharge 1/6). Unit: GeV$^{3/2}$. Status: AUTHOR.
- $\hat{\gamma}^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold, $\hat{\gamma}^\mu = \hat{\mathbf{e}}_a{}^\mu \hat{\gamma}^{(a)}$, $\{\hat{\gamma}^\mu, \hat{\gamma}^\nu\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_\mu = \hat{h}_{\mu\nu}\hat{\gamma}^\nu$. Unit: dimensionless. Status: STANDARD.
- τ^I: The Pauli matrices acting on weak doublets (τ^1, τ^2, τ^3; τ^2 is the second one, not a square). Unit: dimensionless. Status: STANDARD.
- ℓ, $\bar{\ell}$: The left-handed lepton doublets of the three generations, $\ell = (\nu_\mathrm{L}, e_\mathrm{L})$ (hypercharge $-1/2$). Unit: GeV$^{3/2}$. Status: AUTHOR.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.

The Higgs equation and the equation of the $SU(2)$ gauge fields on the fold (with the factor $\mathcal{Z}$ of the extra times and the portal factor $\mathcal{K}_W$). Status: DERIVED.

$$\begin{aligned}
&\mathcal{R}_\mathcal{A} = \mathcal{A}'' + 3\frac{b_4'}{b_4}\mathcal{A}' + \lambda\mathcal{A}^3 + c_\Phi\,\bar{\Psi}P_\mathrm{L}\Psi\,\mathcal{A} = \Big(4a_{40}^2 - 6a_{40}\frac{b_4'}{b_4}\Big)\mathcal{A} + \lambda\mathcal{A}^3 + c_\Phi\,\bar{\Psi}P_\mathrm{L}\Psi\,\mathcal{A},\\
&\mathcal{R}_v = \lambda\mathcal{A}^2 v + c_\Phi\,\bar{\Psi}P_\mathrm{L}\Psi\,v, \qquad J_\Phi^{2\,x_4} = a_{40}\,v\,\mathcal{A}, \qquad J_\Phi^{1\,x_4} = J_\Phi^{3\,x_4} = 0
\end{aligned} \tag{FE-26}$$

Symbols in (FE-26).

- $\mathcal{R}_\mathcal{A}$, $\mathcal{R}_v$: The residuals of the ad hoc Higgs vacuum in the upper ($\mathcal{R}_\mathcal{A}$) and lower ($\mathcal{R}_v$) components of the Higgs equation: what the ad hoc vacuum leaves unbalanced. Unit: GeV^3. Status: DERIVED.
- $\mathcal{A}$: The author's ad hoc upper component of the Higgs vacuum, almightyAdHoc, $\mathcal{A}(x_4) = v_\mathcal{A}\mathrm{e}^{-2a_{40}x_4}$. Unit: GeV. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor of the extra times is $\mathrm{e}^{-a_4}b_4$ times the warp. Unit: dimensionless. Status: AUTHOR.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- Ψ, $\bar{\Psi}$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar{\Psi} = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: GeV$^{7/2}$. Status: AUTHOR.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- a_{40}: The decay constant of almightyAdHoc, $a_{40} > 0$: $\mathcal{A}$ falls by e in the time $1/(2a_{40})$. Unit: GeV (1/length). Status: AUTHOR.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_\mathrm{F})^{-1/2}$ with the Fermi constant $G_\mathrm{F} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- $J_\Phi^{I\nu}$: The weak-isospin current of the Higgs doublet (it contains the mass terms of the W fields); $J_\Phi^{2\,x_4}$ its isospin charge density with $I = 2$. Unit: GeV^3. Status: STANDARD.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.

The author's ad hoc vacuum $\Phi = (\mathcal{A}, v + h)/\sqrt{2}$ with $\mathcal{A} = v_\mathcal{A}\mathrm{e}^{-2a_{40}x_4}$, inserted literally into FE-15 and FE-17 (with $h = 0$ and no gauge field): it leaves the residuals $\mathcal{R}_\mathcal{A}$ and $\mathcal{R}_v$ in the two components of the Higgs equation, and

it carries a homogeneous charge density of the second isospin direction, which the Gauss law of FE-17 forbids for homogeneous gauge fields. Status: DERIVED.

$$\Phi = \frac{1}{\sqrt{2}}\begin{pmatrix}\mathcal{A}\\ v\end{pmatrix} = \mathcal{O}(\phi_{\rm rot})\,\frac{1}{\sqrt{2}}\begin{pmatrix}0\\ V_{\rm rad}\end{pmatrix}, \qquad V_{\rm rad} = \sqrt{v^2+\mathcal{A}^2}, \qquad \tan\frac{\phi_{\rm rot}}{2} = \frac{\mathcal{A}}{v},$$
$$\mathcal{O}(\phi_{\rm rot}) = \begin{pmatrix}\cos\frac{\phi_{\rm rot}}{2} & \sin\frac{\phi_{\rm rot}}{2}\\ -\sin\frac{\phi_{\rm rot}}{2} & \cos\frac{\phi_{\rm rot}}{2}\end{pmatrix}, \qquad W^2_{x_4} = \frac{\phi'_{\rm rot}}{g_2} = -\frac{4a_{40}\,v\,\mathcal{A}}{g_2\big(v^2+\mathcal{A}^2\big)}, \qquad J_\Phi^{2\,x_4} = 0, \tag{S-10}$$
$$\text{FE-15} \;\Leftrightarrow\; \mathcal{O}(\phi_{\rm rot})\,\frac{1}{\sqrt{2}}\begin{pmatrix}0\\ -\mathcal{R}_{\rm rad}\end{pmatrix} = 0, \qquad \mathcal{R}_{\rm rad} = V''_{\rm rad} + 3\,\frac{b'_4}{b_4}\,V'_{\rm rad} + \lambda\,\mathcal{A}^2\,V_{\rm rad} + c_\Phi\,\bar\Psi P_{\rm L}\Psi\,V_{\rm rad}$$

Symbols in (S-10).

- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- $\mathcal{A}$: The author's ad hoc upper component of the Higgs vacuum, almightyAdHoc, $\mathcal{A}(x_4) = v_{\mathcal{A}}\mathrm{e}^{-2a_{40}x_4}$. Unit: GeV. Status: AUTHOR.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_{\rm F})^{-1/2}$ with the Fermi constant $G_{\rm F} = 1.1663788\times10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- $\mathcal{O}(\phi_{\rm rot})$: The 2×2 isospin rotation by the angle $\phi_{\rm rot}$ about the second isospin axis (a real element of $SU(2)$). Unit: dimensionless. Status: DERIVED.
- $V_{\rm rad}$: The modulus of the Higgs doublet in the ad hoc vacuum, $V_{\rm rad} = \sqrt{v^2+\mathcal{A}^2}$ ($V^2_{\rm rad} = 2\Phi^\dagger\Phi$). Unit: GeV. Status: DERIVED.
- $\phi_{\rm rot}$: The angle of the isospin rotation that turns $(0, V_{\rm rad})/\sqrt{2}$ into the ad hoc vacuum: $\tan(\phi_{\rm rot}/2) = \mathcal{A}/v$. Unit: dimensionless. Status: DERIVED.
- W^I_μ, $W^I_{\mu\nu}$: The gauge field of $SU(2)_L$ on the fold, $W^1_\mu, W^2_\mu, W^3_\mu$, and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a'_4 = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- g_2: The $SU(2)_L$ gauge coupling (usually g). Unit: dimensionless. Value: $g_2(m_Z) = 0.6517$. Status: STANDARD.
- a_{40}: The decay constant of almightyAdHoc, $a_{40} > 0$: $\mathcal{A}$ falls by e in the time $1/(2a_{40})$. Unit: GeV (1/length). Status: AUTHOR.
- $J^{I\nu}_\Phi$: The weak-isospin current of the Higgs doublet (it contains the mass terms of the W fields); $J_\Phi^{2\,x_4}$ its isospin charge density with $I = 2$. Unit: GeV^3. Status: STANDARD.
- $\mathcal{R}_{\rm rad}$: The radial residual of the Higgs equation FE-15 in the ad hoc vacuum written as a gauge rotation (S-10): the amount by which $V_{\rm rad}$ fails to solve the radial equation; zero for a solution. Unit: GeV^3. Status: DERIVED.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor of the extra times is $\mathrm{e}^{-a_4}b_4$ times the warp. Unit: dimensionless. Status: AUTHOR.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: GeV$^{7/2}$. Status: AUTHOR.
- $P_{\rm L}$: The author's left-handed chiral projector, $P_{\rm L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.

The ad hoc vacuum is a rotation of the doublet $(0, V_{\rm rad})/\sqrt{2}$ by the angle $\phi_{\rm rot}(x_4)$ about the second isospin axis. Together with the pure-gauge field $W^2_{x_4} = \phi'_{\rm rot}/g_2$, which undoes the rotation in the covariant derivative, the isospin charge density of the Higgs field vanishes, so the Gauss law of FE-17 holds with homogeneous fields; the masses of the W and Z bosons and of the fermions, which depend on the doublet only through $\Phi^\dagger\Phi$ ($m^2_W = g_2^2(v^2+\mathcal{A}^2)/4$, $m^2_Z = (g_1^2+g_2^2)(v^2+\mathcal{A}^2)/4$, each fermion mass $\propto\sqrt{v^2+\mathcal{A}^2}$), are unchanged by the rotation ($v^2+\mathcal{A}^2 = V^2_{\rm rad}$). The

Higgs equation FE-15 then reduces exactly to one radial residual $\mathcal{R}_{\rm rad}$ (at 20 random points of the parameters the reduction holds to 10^{-15}). With $\mu^2 = -\lambda v^2$ the ad hoc vacuum is a solution only in the limit $\mathcal{A} \to 0$, which it reaches exponentially. Status: DERIVED (BG5).

a_{40}/v	$\mathcal{R}_{\rm rad}(x_4=0)/(\lambda v^3)$	$\rho_{\rm H}(x_4=0)/v^4$	time until $\mathcal{A} < 10^{-3}v$
0.1	1.74	0.042	$34.5/v$
1	34.2	1.03	$3.45/v$
10	3280	100	$0.345/v$

The radial residual at $x_4 = 0$ and the energy density of the Higgs field, $\rho_{\rm H} = \frac{1}{2}V'^2_{\rm rad} + \frac{\lambda}{4}\mathcal{A}^4$, for $v_{\mathcal{A}} = v$, a constant b_4 and no dirac16complex at the fold, for three values of a_{40} (in units of $v = 246.22$ GeV); the ad hoc component is gone after a time of order $3.5/a_{40}$. Before nucleosynthesis the Fermi constant $G_{\rm F,loc} = 1/(\sqrt{2}(v^2+\mathcal{A}^2))$ must be within one per cent of its value: $\mathcal{A}^2/v^2 < 0.01$ at $x_4 = 1$ s requires $a_{40} > \ln(v_{\mathcal{A}}/(0.1v))/(2\text{ s})$, a weak bound ($a_{40} > 2.3\ \text{s}^{-1} = 1.5 \times 10^{-24}$ GeV for $v_{\mathcal{A}} = 10v$). Status: DERIVED (BG6, `um3_thermal.py`, T6).

7 The spectra along the hidden direction

Every field of the bulk is a superposition of modes along the hidden direction σ; their frequencies are the masses and energies that an observer on the fold sees (the "x_0 masses"). This section gives the spectra of the bulk fields and of dirac16complex on the static block (a_4, b_4 constant, normalised to $\mathrm{e}^{a_4} = b_4 = 1$ at the fold).

7.1 The bulk fields and why the Standard Model lives on the fold

$$u_{\rm s}'' + \left(\omega^2 - 9H^2\right)u_{\rm s} + 6H\,\delta_{\rm D}(\sigma)\,u_{\rm s} = 0, \qquad u_{\rm s} = \mathrm{e}^{-3H|\sigma|}\ \ (\omega = 0), \qquad \omega \geq 3H\ \ \text{(continuum)} \tag{S-11}$$

Symbols in (S-11).

- $u_{\rm s}$: The rescaled profile of a massless scalar field of the bulk at zero momentum along ordinary space: the field is $\mathrm{e}^{3H|\sigma|}u_{\rm s}(\sigma)\,\mathrm{e}^{-\mathrm{i}\omega x_4}$. Unit: as the field. Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- $\delta_{\rm D}$: The Dirac delta function: $\int \mathrm{d}\sigma\,\delta_{\rm D}(\sigma)F(\sigma) = F(0)$. Unit: 1/length (for $\delta_{\rm D}(\sigma)$). Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

A massless scalar field of the bulk at zero momentum along ordinary space, with its profile written as $\mathrm{e}^{3H|\sigma|}u_{\rm s}$ (which removes the first derivative): the kink of the warp acts as an attractive delta function at the fold, which binds exactly one state, the zero mode (a constant profile, $\omega = 0$), separated by the gap $3H$ from the continuum of the bulk. A gauge field of the bulk has the same structure with the gap $2H$: at zero momentum its mode equation is $(\mathrm{e}^{-4H|\sigma|}A')' + \omega^2\mathrm{e}^{-4H|\sigma|}A = 0$ for a component A along ordinary space, which $A = \mathrm{e}^{2H|\sigma|}u$ turns into $u'' + (\omega^2 - 4H^2)u + 4H\,\delta_{\rm D}(\sigma)\,u = 0$. A massless fermion of the bulk has no gap and no normalisable zero mode: once $\mathrm{e}^{3H|\sigma|}$ is taken out, its equation at zero momentum is that of flat space, whose solutions oscillate for every $\omega \neq 0$. Status: DERIVED (`um3_bulk.py`, B1 for the scalar; the transformations for the gauge field and the fermion).

$$c_{\rm sc}^2 = \left\langle \mathrm{e}^{2H|\sigma|} \right\rangle = \frac{3}{2}, \qquad c_{\rm gauge}^2 = 2, \qquad c_{\rm fer} = \frac{2M_{\rm k}}{2M_{\rm k} - H}, \qquad c_{\rm g}^2 = 5 \cdot \frac{3}{2} = \frac{15}{2} \tag{S-12}$$

Symbols in (S-12).

- $c_{\rm sc}$: The limiting speed of the zero mode of a scalar field of the bulk along ordinary space, in units of the speed of light on the fold. Unit: dimensionless. Value: $\sqrt{3/2}$. Status: DERIVED.

- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- c_{gauge}: The limiting speed of the zero mode of a gauge field (a photon) of the bulk, in units of the speed of light on the fold. Unit: dimensionless. Value: $\sqrt{2}$. Status: DERIVED.
- c_{fer}: The limiting speed of the zero mode of a fermion of the bulk with the kink mass $M_{\mathrm{k}}\,\mathrm{sgn}(\sigma)$, in units of the speed of light on the fold. Unit: dimensionless. Status: DERIVED.
- M_{k}: The mass of a fermion of the bulk that changes sign at the fold (a kink mass $M_{\mathrm{k}}\,\mathrm{sgn}(\sigma)$), used to illustrate the speeds of bulk fermions. Unit: GeV. Status: ASSUMED.
- c_{g}: The limiting speed of the zero mode of the transverse traceless graviton (the four-dimensional gravitational wave) at the static block or with frozen extra times, in units of the speed of light on the fold. Unit: dimensionless. Value: $\sqrt{15/2} = 2.74$ for the bulk alone; $\sqrt{83/82} = 1.0061$ with the graviton term of the fold from the Gauss-Bonnet and cubic Lovelock terms; $\sqrt{(96 + 83\epsilon_{\mathrm{r}})/(96 + 82\epsilon_{\mathrm{r}})}$ with the induced gravity of the fold. Status: DERIVED.

The limiting speeds, in units of the speed of light on the fold, of the zero modes of the bulk fields when they move along ordinary space: a zero mode samples the warp $\mathrm{e}^{-H|\sigma|}$ of ordinary space over its profile, so its speed is the average of $\mathrm{e}^{2H|\sigma|}$ with the weight of its profile. A bulk scalar moves $\sqrt{3/2} = 1.22$ times faster than light on the fold, a bulk gauge field (photon) $\sqrt{2} = 1.41$ times, a bulk fermion with the kink mass $M_{\mathrm{k}}\,\mathrm{sgn}(\sigma)$ (zero mode $\propto \mathrm{e}^{(3H - M_{\mathrm{k}})|\sigma|}$, normalisable for $M_{\mathrm{k}} > 0$) $2M_{\mathrm{k}}/(2M_{\mathrm{k}} - H)$ times (2 for $M_{\mathrm{k}} = H$, 1.0005 for $M_{\mathrm{k}} = 1000H$), and the transverse traceless graviton of the static block, whose local cone is $\sqrt{5}$ times the light cone because of the Gauss-Bonnet and cubic Lovelock terms, $\sqrt{15/2} = 2.74$ times (the numerical eigenvalues agree with these values to 3×10^{-3} or better: $c_{\mathrm{sc}}^2 = 1.50005$ and 1.49993, $c_{\mathrm{gauge}}^2 = 1.99875$ and 1.99410 at two small momenta). Species in the bulk would therefore have different limiting speeds, which measurements exclude at the level 10^{-15} to 10^{-20} for photons, electrons and nucleons, and three neutrino flavours with three profiles would oscillate with phases proportional to the energy, which is excluded too: the Standard Model lives on the fold, where every field has the one light cone of the fold. Bulk gravitons cannot carry four-dimensional gravity: the speed of gravitational waves equals that of light to 7×10^{-16} (GW170817), and at the author's vacuum the graviton of the bulk has no time-kinetic term at all. The Lovelock terms give the fold a graviton term of its own, but with it the four-dimensional gravitational wave still moves with $c_{\mathrm{g}}^2 = 83/82$ (FE-63): UM-3 gives the fold an induced Einstein-Hilbert term, solved in the section on the sources of the fold. Status: DERIVED (B1 to B6).

7.2 dirac16complex: the continuum and the bound states

$$\Psi = \mathrm{e}^{3H|\sigma|}\,\chi(\sigma)\,\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}, \qquad \gamma^{(0)}\chi' - \mathrm{i}\omega\,\gamma^{(4)}\chi + \mathrm{i}\,\mathrm{e}^{H|\sigma|}\,\vec{k}\cdot\vec{\gamma}\,\chi - m_\Psi\chi = 0 \qquad (\sigma \neq 0) \tag{FE-36}$$

Symbols in (FE-36).

- Ψ, $\bar{\Psi}$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar{\Psi} = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{3H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of a bound state. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\vec{k}$: The (comoving) momentum along ordinary space, $\vec{k}^2$ its square. Unit: GeV. Status: STANDARD.

- $\vec{x}$: The position in ordinary space, $\vec{x} = (x_1, x_2, x_3)$. Unit: length. Status: AUTHOR.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\vec{\gamma}$: The three frame Dirac matrices of dirac16complex along ordinary space, $\vec{\gamma} = (\gamma^{(1)}, \gamma^{(2)}, \gamma^{(3)})$. Unit: dimensionless. Status: AUTHOR.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.

$$\chi' = \mathbb{K}\,\chi, \qquad \mathbb{K} = \gamma^{(0)}\big(m_\Psi + \mathrm{i}\omega\gamma^{(4)}\big), \qquad \mathbb{K}^2 = \big(m_\Psi^2 - \omega^2\big)\mathbf{1}, \qquad \chi \propto \mathrm{e}^{\pm q\sigma}, \qquad q = \sqrt{m_\Psi^2 - \omega^2} \tag{FE-37}$$

Symbols in (FE-37).

- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{3H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of a bound state. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\mathbb{K}$: The transfer matrix of the profile equation at zero momentum, $\chi' = \mathbb{K}\chi$. Unit: GeV. Status: DERIVED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- $\mathbf{1}$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space). Unit: dimensionless. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- q: The decay rate of a profile along σ at frequency ω, $q = \sqrt{m_\Psi^2 - \omega^2}$ (imaginary in the continuum). Unit: GeV. Status: DERIVED.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

$$\omega^2 = m_\Psi^2 + p_\sigma^2 + \mathrm{e}^{2H|\sigma| - 2a_4}\,\vec{k}^2 - \mathrm{e}^{2H|\sigma| + 2a_4}\,b_4^{-2}\,\vec{\kappa}^2 \tag{FE-38}$$

Symbols in (FE-38).

- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- p_σ: The momentum along the hidden direction (local wave number of a profile along σ). Unit: GeV. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $\vec{k}$: The (comoving) momentum along ordinary space, $\vec{k}^2$ its square. Unit: GeV. Status: STANDARD.

- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor of the extra times is $\mathrm{e}^{-a_4} b_4$ times the warp. Unit: dimensionless. Status: AUTHOR.
- $\vec{\kappa}$: A momentum along the extra times (three components); the fields of UM-3 have $\vec{\kappa} = 0$. Unit: GeV. Status: DERIVED.

The field equation of dirac16complex on the static block (FE-36), its solutions at zero momentum along ordinary space (FE-37): the continuum $|\omega| \geq m_\Psi$ and the exponentials below it, and the local dispersion relation (FE-38), which shows that a dependence on the extra times would make every field tachyonic. Status: DERIVED.

$$\chi(0^+) = \exp\Big(\gamma^{(0)}\big(\mu_\mathrm{L} P_\mathrm{L} + \mu_\mathrm{R} P_\mathrm{R}\big)\Big)\,\chi(0^-),$$
$$\mu_X = \frac{1}{V_*}\Big[c_\Phi\,\langle\Phi^\dagger\Phi\rangle\,\delta_{X\mathrm{L}} + \frac{1}{4}\big(c_X^G\langle G^2\rangle + c_X^W\langle W^2\rangle + c_X^B\langle B^2\rangle\big) + 2\omega\sum_f c_X^f\,n_f\Big], \qquad \langle\Phi^\dagger\Phi\rangle = \frac{v^2 + \mathcal{A}^2}{2} \tag{FE-39}$$

Symbols in (FE-39).

- $\chi(\sigma)$: The profile of a mode of dirac16complex along the hidden direction, $\Psi = \mathrm{e}^{3H|\sigma|}\chi(\sigma)\mathrm{e}^{\mathrm{i}(\vec{k}\cdot\vec{x} - \omega x_4)}$ (a 16-component spinor function of σ); as a four-dimensional field, χ denotes the particle of a bound state. Unit: GeV^2 (with the V_* normalisation). Status: DERIVED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- μ_X, μ_L, μ_R: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_\mathrm{L} P_\mathrm{L} + \mu_\mathrm{R} P_\mathrm{R})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- P_R: The author's right-handed chiral projector, $P_\mathrm{R} = (\mathbf{1} + \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4} b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4} b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with upper and lower index strings the generalised Kronecker delta $\delta^{M_1\cdots M_n}_{N_1\cdots N_n}$, the determinant of the $n \times n$ matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- c_X^G: The gluon-kinetic portal coupling of the half X of dirac16complex. Unit: GeV^{-7}. Status: INVENTED.
- $\langle G^2\rangle$: The mean square of the gluon field strength, $\langle G^A_{\mu\nu} G^{A\mu\nu}\rangle$ (the gluon condensate in the QCD vacuum, a thermal value in the plasma). Unit: GeV^4. Status: STANDARD.
- G^A_μ, $G^A_{\mu\nu}$: The gluon field of $SU(3)_C$ on the fold ($A = 1, \ldots, 8$) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- c_X^W: The W-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- $\langle W^2\rangle$: The mean square of the $SU(2)$ field strength, $\langle W^I_{\mu\nu} W^{I\mu\nu}\rangle$. Unit: GeV^4. Status: STANDARD.
- W^I_μ, $W^I_{\mu\nu}$: The gauge field of $SU(2)_L$ on the fold, $W^1_\mu, W^2_\mu, W^3_\mu$, and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- c_X^B: The B-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- $\langle B^2\rangle$: The mean square of the hypercharge field strength, $\langle B_{\mu\nu} B^{\mu\nu}\rangle$. Unit: GeV^4. Status: STANDARD.
- B_μ, $B_{\mu\nu}$: The hypercharge gauge field of $U(1)_Y$ on the fold and its field strength. Unit: GeV (field), GeV^2

(field strength). Status: AUTHOR.

- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- f: The label of a kind of Standard-Model multiplet: $f \in \{Q, u, d, \ell, e, \nu\}$ (quark doublets, up-quark singlets, down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.
- c_X^f: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c_X^ℓ and c_X^ν are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.
- n_f: The number density of the Standard-Model fermions of kind f (particles minus antiparticles, summed over generations and gauge components), the time component of the mean current $\langle J_f^\mu \rangle$ in the rest frame of the medium. Unit: GeV^3. Status: STANDARD.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_\mathrm{F})^{-1/2}$ with the Fermi constant $G_\mathrm{F} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- $\mathcal{A}$: The author's ad hoc upper component of the Higgs vacuum, almightyAdHoc, $\mathcal{A}(x_4) = v_\mathcal{A} \mathrm{e}^{-2a_{40}x_4}$. Unit: GeV. Status: AUTHOR.

$$q_\mathrm{b} = -m_\Psi \left(\mu_\mathrm{L} + \mu_\mathrm{R}\right) \frac{\tanh \varsigma}{2\varsigma}, \qquad \varsigma = \sqrt{\mu_\mathrm{L}\mu_\mathrm{R}}, \qquad \omega_\mathrm{b} = \sqrt{m_\Psi^2 - q_\mathrm{b}^2}, \qquad |\chi_\mathrm{b}(\sigma)| \propto \mathrm{e}^{-q_\mathrm{b}|\sigma|} \tag{FE-40}$$

Symbols in (FE-40).

- q_b: The inverse width of a state of dirac16complex bound to the fold: its profile falls as $\mathrm{e}^{-q_\mathrm{b}|\sigma|}$. Unit: GeV. Status: DERIVED.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- μ_X, μ_L, μ_R: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_\mathrm{L} P_\mathrm{L} + \mu_\mathrm{R} P_\mathrm{R})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- ς: The auxiliary variable of the bound-state condition, $\varsigma = \sqrt{\mu_\mathrm{L}\mu_\mathrm{R}}$ ($\varsigma_- = \sqrt{-\mu_\mathrm{L}\mu_\mathrm{R}}$ when $\mu_\mathrm{L}\mu_\mathrm{R} < 0$). Unit: dimensionless. Status: DERIVED.
- ω_b: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- χ_b: The profile of a state of dirac16complex bound to the fold. Unit: GeV^2. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

The fold acts on dirac16complex as a chirality-dependent mass concentrated at $\sigma = 0$, with strengths μ_L and μ_R set by the mean fields of the Standard Model (FE-39); the bound states (FE-40) exist for an attractive fold. Status: DERIVED.

$(\mu_\mathrm{L}, \mu_\mathrm{R})$	$(-0.4, 0)$	$(-1, 0)$	$(-0.6, -0.3)$	$(-0.5, 0.2)$	$(-1.5, -0.5)$	$(0.4, 0)$	$(0.3, 0.2)$
ω_b/m_Ψ	0.979796	0.866025	0.905282	0.987882	0.589814	none	none

$$Q_\Psi = \int \mathrm{d}\sigma\, \mathrm{d}^3x\, \sqrt{|g|}\; \Psi^\dagger \mathfrak{B} \Psi, \qquad \mathfrak{B} = \mathrm{i}\,\sigma_{16}\,\gamma^{(4)}, \qquad \mathfrak{B}^\dagger = \mathfrak{B}, \qquad \mathfrak{B}^2 = \mathbf{1}, \qquad E_o = -\omega_o\, \mathcal{Q}_o \tag{FE-43}$$

Symbols in (FE-43).

- Q_Ψ: The conserved charge of dirac16complex (the Noether charge of a change of the phase of Ψ by a constant), a Krein form (indefinite). Unit: dimensionless. Status: DERIVED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.

- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- x: The coordinates of the fold: x_i $(i = 1, 2, 3)$ of ordinary space, x_4 the time, x_r $(r = 5, 6, 7)$ the extra times; a field of the fold is a function of the four observable coordinates $x = (x_1, x_2, x_3, x_4)$, and $\mathrm{d}^4x = \mathrm{d}x_1\,\mathrm{d}x_2\,\mathrm{d}x_3\,\mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $\mathfrak{B}$: The Krein operator, $\mathfrak{B} = \mathrm{i}\sigma_{16}\gamma^{(4)}$: a Hermitian involution with the signature (8, 8); $\Psi^\dagger\mathfrak{B}\Psi$ is the charge density. Unit: dimensionless. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- σ_{16}: The matrix of the Dirac adjoint, $\sigma_{16} = T_0T_1T_2T_3$, real and symmetric; $\sigma_{16}T_a$ is real and antisymmetric. Unit: dimensionless. Status: AUTHOR.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- $\mathbf{1}$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space). Unit: dimensionless. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- $\mathcal{Q}_o$: The charge Q_Ψ (the Krein charge) of the single-particle state o of dirac16complex, a Kohn-Sham orbital or a mode: $+1$ or -1 for a normalised state. Unit: dimensionless. Status: DERIVED.

The conserved charge of dirac16complex (the Noether charge of a constant change of its phase) and the energy of a mode: the charge density is $\Psi^\dagger\mathfrak{B}\Psi$ with the Krein operator $\mathfrak{B}$, which is indefinite because the author's matrices have the signature (4, 4), and a mode of frequency ω_o and charge $\mathcal{Q}_o$ carries the energy $E_o = -\omega_o\mathcal{Q}_o$. Status: DERIVED.

The bound states computed with the author's exact 16×16 matrices: the roots of the 8×8 matching determinant agree with FE-40 to 10^{-7} in all seven cases, each bound level consists of eight degenerate states, and there is none for a repulsive fold. Of the eight states at $+\omega_\mathrm{b}$, four have the Krein charge $+1$ and four -1 (no null states), so four carry positive and four negative energy (FE-43). The bound-state energy $\omega_\mathrm{b} = \sqrt{m_\Psi^2 - q_\mathrm{b}^2}$ is the x_0 mass of dirac16complex: the rest mass of a particle χ that lives near the fold with the profile $\mathrm{e}^{-q_\mathrm{b}|\sigma|}$ of width $1/q_\mathrm{b}$; the more tightly it is bound, the lighter it is. Status: DERIVED (`um3_spectra.py`, S1 to S4).

$$c_\mathrm{b}^2 = \left\langle \mathrm{e}^{2H|\sigma|} \right\rangle = \frac{q_\mathrm{b}}{q_\mathrm{b} - H} \qquad (q_\mathrm{b} > H) \tag{S-13}$$

Symbols in (S-13).

- c_b: The limiting speed of the bound particles χ of dirac16complex along ordinary space, in units of the speed of light on the fold (first order in the momentum). Unit: dimensionless. Status: DERIVED.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

- q_b: The inverse width of a state of dirac16complex bound to the fold: its profile falls as $\mathrm{e}^{-q_\mathrm{b}|\sigma|}$. Unit: GeV. Status: DERIVED.

The limiting speed of the bound particles χ along ordinary space, to first order in their momentum: the average of the warp over the density $\mathrm{e}^{-2q_\mathrm{b}|\sigma|}$ of their profile ($c_\mathrm{b}^2 = 1.25$, 2 and 5 for $H = 0.1m_\Psi$, $0.25m_\Psi$ and $0.4m_\Psi$ with $q_\mathrm{b} = m_\Psi/2$). The particles χ are slower than the bulk continuum but faster than light on the fold, and for $q_\mathrm{b} \le H$ the average diverges: a bound state of dirac16complex has a sensible dispersion relation only if it is bound more tightly than the warp, $q_\mathrm{b} > H$. Status: DERIVED (S5), ESTIMATE (first order in the momentum).

The computed speed. The dispersion of the bound states moving along x_1, computed from the profile equation at each momentum, confirms the limiting speed and measures the error of S-13: $c_\mathrm{b}^2 = 1.190$, 1.440 and 1.069 for $(H, q_\mathrm{b}) = (0.05, 0.3)$, $(0.1, 0.3)$ and $(0.002, 0.03)$ in units of m_Ψ, against 1.2, 1.5 and 1.071 from S-13, which therefore slightly overestimates the limiting speed. Status: DERIVED (`um3_superluminal.py`, SL1).

$$\omega^2 = \omega_\mathrm{b}^2 + c_\mathrm{b}^2\,\vec{k}^2 + \ldots \quad \big(|\vec{k}| \ll m_\Psi\big), \qquad v_\mathrm{g} = \frac{\mathrm{d}\omega}{\mathrm{d}|\vec{k}|}, \qquad \frac{\omega}{|\vec{k}|} \to 1 \quad \big(|\vec{k}| \to \infty\big) \tag{S-31}$$

Symbols in (S-31).

- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- ω_b: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- c_b: The limiting speed of the bound particles χ of dirac16complex along ordinary space, in units of the speed of light on the fold (first order in the momentum). Unit: dimensionless. Status: DERIVED.
- $\vec{k}$: The (comoving) momentum along ordinary space, $\vec{k}^2$ its square. Unit: GeV. Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- v_g: The group velocity of a particle χ along ordinary space, $\mathrm{d}\omega/\mathrm{d}|\vec{k}|$, in units of the speed of light on the fold. Unit: dimensionless. Status: DERIVED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.

At large momenta the ratio $\omega/|\vec{k}|$ falls towards 1 from above (1.0252 at $|\vec{k}| = 10\,m_\Psi$ and 1.0112 at $30\,m_\Psi$ for $H = 0.05$, $q_\mathrm{b} = 0.3$; $\omega - |\vec{k}|$ grows as $|\vec{k}|^{0.26}$, close to the $|\vec{k}|^{1/3}$ of a particle in a linear confining potential, which the warp provides). The group velocity v_g exceeds 1 from $|\vec{k}| = 6.5\,m_\Psi$ on, reaches its maximum 1.0045 at $12\,m_\Psi$ and returns towards 1 (1.0038 at $25\,m_\Psi$): ultrarelativistic particles χ move at most 0.45 per cent faster than light on the fold, and their front velocity, the limit at infinite momentum, is the speed of light. The particles χ of the halo of the Galaxy ($|\vec{k}| = 10^{-3}\omega_\mathrm{b}$) move at 1.19×10^{-3}, and at the freeze-out of their relic density (temperature $\omega_\mathrm{b}/25$) at 0.371: the group velocity exceeds 1 only for relativistic particles χ, which existed only before the freeze-out. Status: DERIVED (SL2, SL3).

$$\mathrm{d}\sigma^2 + \mathrm{e}^{-2H|\sigma|}\,\mathrm{d}x_1^2, \qquad \sinh\Big(\frac{H\,\Delta x_4}{2}\Big) = \frac{H\,\Delta x_1}{2} \tag{S-32}$$

Symbols in (S-32).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.

- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.

The shortcut through the bulk. A signal that leaves the fold travels in the bulk, where the warp shortens the distances along ordinary space. On the static block the earliest signal between two points of the fold at the distance Δx_1 follows the shortest path of the optical metric (the first expression: the metric whose lengths are the travel times of light), which is the hyperbolic plane of radius $1/H$, with the fold as a horocycle. It arrives after the time Δx_4 of the second expression, shorter than Δx_1: $\Delta x_4 = 0.4949$, 1.7627, 4.6249 and 9.2105 for $H\Delta x_1 = 0.5$, 2, 10 and 100 (a direct numerical minimisation of the travel time gives 0.4949, 1.7629, 4.6296 and 9.2953, upper bounds), against 0.5, 2, 10 and 100 along the fold. The time x_4 is a time function of the metric, so there are no closed causal curves, but the bulk is a shortcut between the points of the fold. Through loops of the particles χ the portals would carry this Lorentz violation to the Standard Model; for the electron, whose maximal speed is bounded to 1.2×10^{-17} above and 6×10^{-20} below that of light, the estimate requires $c_\mathrm{b}^2 - 1 = H/(q_\mathrm{b} - H)$ below 2×10^{-11} to 6×10^{-13}, that is q_b/H above 4×10^{10} to 2×10^{12} for ω_b from 1.5 to 10 TeV. Status: DERIVED (SL4), ESTIMATE (the loop, SL5).

8 The Kohn-Sham states of dirac16complex

dirac16complex is a quantised field. Its states with many particles are described here with the Kohn-Sham equations of the record (FE-41, FE-42): single-particle orbitals in the mean fields of everything else, iterated until the orbitals reproduce the fields they move in. The metric is held fixed (the back-reaction of the gas on the warp is left out, ASSUMED); the fields that are iterated are the mean field of the self-interaction and, for the Higgs loop, the strength μ_L of the fold. The state solved is a homogeneous gas of the bound particles χ at rest on the static block of the late universe (a_4 and b_4 constant), with their self-interaction, with gravity (the warp of the metric and the canonical spin connection) and with every field of the Standard Model, which acts through its mean values at the fold (the strengths μ_L and μ_R of FE-39); the Higgs portal also feeds back, because the left-handed half of the gas shifts the local electroweak vacuum v_loc of FE-49. The equations of dirac16complex itself come first.

$$\gamma^M D_M \Psi - m_\Psi \Psi - \lambda_\Psi \big(\bar\Psi \Psi\big)\Psi = 0 \qquad (\sigma \neq 0) \tag{FE-12}$$

Symbols in (FE-12).

- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar\Psi = \partial_M\bar\Psi - \bar\Psi\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat h^{\mu\nu} D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- λ_Ψ: The coupling of the contact self-interaction of dirac16complex. Unit: GeV^{-6}. Status: RECORD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

$$\begin{aligned}
&\Psi\big|_{\sigma=0^+} = \exp\Big(\frac{1}{V_*}\,\gamma^{(0)}\,\mathbb{M}\Big)\,\Psi\big|_{\sigma=0^-},\\
&\mathbb{M} = c_\Phi\big(\Phi^\dagger\Phi\big)P_\mathrm{L} + \sum_X \Big[\frac{1}{4}\big(c_X^G\, G^A_{\mu\nu}G^{A\mu\nu} + c_X^W\, W^I_{\mu\nu}W^{I\mu\nu} + c_X^B\, B_{\mu\nu}B^{\mu\nu}\big)P_X + \frac{d_X}{2}\,B_{\mu\nu}\,\mathrm{i}\Sigma^{\mu\nu}P_X\\
&\qquad + \sum_f c_X^f\big(2\mathrm{i}\,J_f^\mu P_X D_\mu + \mathrm{i}(\hat\nabla_\mu J_f^\mu)P_X\big)\Big]
\end{aligned} \tag{FE-13}$$

Symbols in (FE-13).

- Ψ, $\bar{\Psi}$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar{\Psi} = \Psi^\dagger \sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- $\mathbb{M}$: The fold operator of dirac16complex: the 16×16 matrix (with a derivative along the fold) built from the Standard-Model fields at the fold by varying the portals with respect to $\bar{\Psi}$ (FE-13). Unit: GeV^{-3} (so that $\mathbb{M}/V_*$ is dimensionless). Status: DERIVED.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c_X^G: The gluon-kinetic portal coupling of the half X of dirac16complex. Unit: GeV^{-7}. Status: INVENTED.
- G_μ^A, $G_{\mu\nu}^A$: The gluon field of $SU(3)_C$ on the fold ($A = 1, \ldots, 8$) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- A: A colour (adjoint SU(3)) index running over $1, \ldots, 8$; a repeated index is summed. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- c_X^W: The W-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- W_μ^I, $W_{\mu\nu}^I$: The gauge field of $SU(2)_L$ on the fold, $W_\mu^1, W_\mu^2, W_\mu^3$, and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- I: A weak-isospin (adjoint SU(2)) index running over $1, 2, 3$; a repeated index is summed. Status: STANDARD.
- c_X^B: The B-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- B_μ, $B_{\mu\nu}$: The hypercharge gauge field of $U(1)_Y$ on the fold and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- P_X: The chiral projector of the half X: $P_X = P_\mathrm{L}$ for $X = \mathrm{L}$, P_R for $X = \mathrm{R}$. Unit: dimensionless. Status: AUTHOR.
- d_X: The dipole-portal coupling of the half X of dirac16complex with the hypercharge field strength. Unit: GeV^{-5}. Status: INVENTED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- Σ^{ab}, $\Sigma^{\mu\nu}$: The generators of Spin(4,4) on dirac16complex, $\Sigma^{ab} = \frac{1}{4}[T^a, T^b]$ (the scaled commutators of the author's matrices); $\Sigma^{\mu\nu} = \mathbf{e}_a{}^\mu \mathbf{e}_b{}^\nu \Sigma^{ab}$ along the fold. Unit: dimensionless. Status: AUTHOR.
- f: The label of a kind of Standard-Model multiplet: $f \in \{Q, u, d, \ell, e, \nu\}$ (quark doublets, up-quark singlets, down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.
- c_X^f: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c_X^ℓ and c_X^ν are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.
- J_f^μ: The fermion-number current of the Standard-Model multiplets of kind f, $J_f^\mu = \bar{\psi}_f \hat{\gamma}^\mu \psi_f$ summed over the three generations and the gauge components; with other labels the currents that are the sources of the gauge fields. Unit: GeV^3. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field

$(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2 \epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.

- $\hat{\nabla}$: The covariant derivative of the fold: with the Levi-Civita connection of the metric of the fold (for four-dimensional tensors of its metric $\hat{h}_{\mu\nu} = \mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the observable coordinates $x_1, \ldots, x_4$; for $\hat{\nabla}_a \mathcal{S}^a{}_b$ of the induced metric of the seven-dimensional fold). Status: STANDARD.

$$\gamma^M \Omega_M = -3H \,\mathrm{sgn}(\sigma)\, \gamma^{(0)} + \frac{3}{2}\,\frac{b_4'}{b_4}\,\gamma^{(4)} \neq 0 \tag{FE-14}$$

Symbols in (FE-14).

- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Ω_M: The spin connection acting on dirac16complex, $\Omega_M = \frac{1}{2}\varpi_{Mab}\Sigma^{ab}$. Unit: 1/length. Status: RECORD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- sgn: The sign function: $\mathrm{sgn}(y) = +1$ for $y > 0$, -1 for $y < 0$. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- b_4: The author's metric function of the time x_4 in the extra-time components, $b_4 > 0$, never identically 1 (the author's order of 2026-10-24); the scale factor of the extra times is $\mathrm{e}^{-a_4} b_4$ times the warp. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.

The field equation of dirac16complex in the bulk (FE-12, with its contact self-interaction), the matching condition at the fold (FE-13: the values on the two sides of the fold differ by the exponential of $\gamma^{(0)}\mathbb{M}/V_*$, where the fold operator $\mathbb{M}$ is built from the Standard-Model fields through the portals) and the spin connection of the author's metric (FE-14), which couples dirac16complex to gravity and never vanishes: its first term comes from the warp of the six directions of ordinary space and of the extra times, its second from the time dependence of b_4. Status: RECORD (FE-12), DERIVED (FE-13, FE-14).

$$\begin{aligned} &\big[\gamma^M D_M - m_\Psi - \lambda_\Psi\, n_\mathrm{s} + \lambda_\Psi\, \mathcal{N}\big]\varphi_o = \delta_\mathrm{D}(\sigma)\,\frac{1}{V_*}\,\langle \mathbb{M}\rangle\,\varphi_o, \\ &\mathcal{N} = \sum_o \mathsf{n}_o\,\varphi_o\bar{\varphi}_o, \qquad n_\mathrm{s} = \mathrm{tr}\,\mathcal{N} = \sum_o \mathsf{n}_o\,\bar{\varphi}_o\varphi_o, \qquad \mathsf{n}_o = \frac{1}{\mathrm{e}^{(E_o-\mu_\Psi)/\mathsf{T}_\Psi}+1} \end{aligned} \tag{FE-41}$$

Symbols in (FE-41).

- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- λ_Ψ: The coupling of the contact self-interaction of dirac16complex. Unit: GeV^{-6}. Status: RECORD.
- n_s: The scalar density of dirac16complex in the Kohn-Sham state, $n_\mathrm{s} = \langle\bar{\Psi}\Psi\rangle = \mathrm{tr}\,\mathcal{N}$. Unit: GeV^7. Status: DERIVED.
- $\mathcal{N}$: The one-body density matrix of dirac16complex, $\mathcal{N} = \sum_o \mathsf{n}_o\varphi_o\bar{\varphi}_o$ (a 16×16 matrix at each point). Unit: GeV^7. Status: DERIVED.

- φ_o, $\bar{\varphi}_o$: A Kohn-Sham orbital of dirac16complex (a single-particle solution of FE-41) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- δ_{D}: The Dirac delta function: $\int \mathrm{d}\sigma\, \delta_{\mathrm{D}}(\sigma) F(\sigma) = F(0)$. Unit: 1/length (for $\delta_{\mathrm{D}}(\sigma)$). Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- $\mathbb{M}$: The fold operator of dirac16complex: the 16×16 matrix (with a derivative along the fold) built from the Standard-Model fields at the fold by varying the portals with respect to $\bar{\Psi}$ (FE-13). Unit: GeV^{-3} (so that $\mathbb{M}/V_*$ is dimensionless). Status: DERIVED.
- n_o: The occupation of the Kohn-Sham orbital o (between 0 and 1). Unit: dimensionless. Status: RECORD.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- μ_Ψ: The chemical potential of dirac16complex. Unit: GeV. Status: DERIVED.
- T_Ψ: The temperature of dirac16complex. Unit: GeV. Status: DERIVED.

$$\langle \mathcal{T}_{\Psi MN} \rangle = -\frac{1}{4}\sum_o \mathsf{n}_o \Big(\bar{\varphi}_o \gamma_M D_N \varphi_o + \bar{\varphi}_o \gamma_N D_M \varphi_o - (D_M \bar{\varphi}_o)\gamma_N \varphi_o - (D_N \bar{\varphi}_o)\gamma_M \varphi_o \Big) + \frac{\lambda_\Psi}{2}\, g_{MN} \big(n_{\mathrm{s}}^2 - \mathrm{tr}\,\mathcal{N}^2 \big) \quad \text{(FE-42)}$$

Symbols in (FE-42).

- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- $\mathcal{T}_\Psi$: The energy-momentum tensor of dirac16complex, $\mathcal{T}_{MN} = -(2/\sqrt{|g|})\,\delta S_\Psi/\delta g^{MN}$. Unit: GeV^8. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- n_o: The occupation of the Kohn-Sham orbital o (between 0 and 1). Unit: dimensionless. Status: RECORD.
- φ_o, $\bar{\varphi}_o$: A Kohn-Sham orbital of dirac16complex (a single-particle solution of FE-41) and its Dirac adjoint. Unit: $\mathrm{GeV}^{7/2}$. Status: RECORD.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC} G_\mu^B X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- λ_Ψ: The coupling of the contact self-interaction of dirac16complex. Unit: GeV^{-6}. Status: RECORD.
- g_{MN}: The metric tensor of the eight-dimensional spacetime (signature $+,+,+,+,-,-,-,-$ in the order $x_0, \ldots, x_7$): $g_{MN}\,\mathrm{d}X^M\mathrm{d}X^N$ is the squared interval, g^{MN} its inverse, $|g|$ the absolute value of its determinant ($\sqrt{|g|}\,\mathrm{d}^8X$ is the proper volume element). Unit: dimensionless. Status: AUTHOR.
- n_{s}: The scalar density of dirac16complex in the Kohn-Sham state, $n_{\mathrm{s}} = \langle\bar{\Psi}\Psi\rangle = \mathrm{tr}\,\mathcal{N}$. Unit: GeV^7. Status: DERIVED.

- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- $\mathcal{N}$: The one-body density matrix of dirac16complex, $\mathcal{N} = \sum_o \mathsf{n}_o \varphi_o \bar{\varphi}_o$ (a 16×16 matrix at each point). Unit: GeV^7. Status: DERIVED.

The Kohn-Sham equations of dirac16complex coupled to itself, to gravity and to every Standard-Model field (FE-41) and the energy-momentum tensor of the Kohn-Sham state (FE-42): each orbital φ_o obeys the field equation with the mean fields, the self-interaction contributes the Hartree term $\lambda_\Psi n_\mathrm{s}$ and the exact exchange term $-\lambda_\Psi \mathcal{N}$, and every Standard-Model field enters through the mean fold operator $\langle \mathbb{M} \rangle$; correlation beyond exchange and the sea of negative-energy states are left out. Status: RECORD (the Kohn-Sham treatment), DERIVED (the exchange term), ASSUMED (no correlation, no sea).

8.1 The equations of the gas of bound particles

$$
\begin{aligned}
&\Psi = \mathrm{e}^{3H|\sigma|}\, \chi_o(\sigma)\, \mathrm{e}^{-\mathrm{i}\omega x_4}, \qquad \chi_o' = \gamma^{(0)} \Big(\mathrm{i}\omega\, \gamma^{(4)} + m_\Psi + \mathcal{M}(\sigma) \Big) \chi_o \qquad (\sigma \neq 0), \\
&\mathcal{M}(\sigma) = G_\mathrm{KS}\, \mathrm{e}^{6H|\sigma|} \Big(\hat{\rho}(\sigma)\, \mathbf{1} - \hat{\mathcal{N}}(\sigma) \Big), \qquad \hat{\rho} = \frac{1}{4} \sum_{o=1}^{4} \bar{\chi}_o \chi_o, \qquad \hat{\mathcal{N}} = \frac{1}{4} \sum_{o=1}^{4} \chi_o \bar{\chi}_o, \\
&\chi_o(0^+) = \exp\Big(\gamma^{(0)} \big(\mu_\mathrm{L} P_\mathrm{L} + \mu_\mathrm{R} P_\mathrm{R} \big) \Big) \chi_o(0^-), \qquad \int_{-\infty}^{\infty} \mathrm{d}\sigma\, \chi_o^\dagger \mathfrak{B} \chi_p = -\delta_{op}, \qquad G_\mathrm{KS} = \frac{\lambda_\Psi\, n_\chi}{V_*}
\end{aligned}
\tag{S-14}
$$

Symbols in (S-14).

- Ψ, $\bar{\Psi}$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar{\Psi} = \Psi^\dagger \sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\, \mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z) \ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\chi_o(\sigma)$, $\bar{\chi}_o$: The profile of the occupied Kohn-Sham orbital o ($o = 1, \ldots, 4$, the four positive-energy states of the bound level) of a homogeneous gas of the particles χ at rest on the static block: a 16-component spinor function of σ; the orbital of each particle is $\mathrm{e}^{3H|\sigma|}\chi_o(\sigma)\mathrm{e}^{-\mathrm{i}\omega x_4}$ divided by the square root of the volume of ordinary space and of V_*; normalised to the Krein charge -1, $\int \mathrm{d}\sigma\, \chi_o^\dagger \mathfrak{B} \chi_o = -1$; $\bar{\chi}_o = \chi_o^\dagger \sigma_{16}$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{1/2}$. Status: DERIVED.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations a second label of a Kohn-Sham orbital (as o, with δ_{op} the Kronecker delta of the two labels). Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- $\mathcal{M}(\sigma)$: The mean field of the self-interaction in the Kohn-Sham equation of the gas: the 16×16 matrix $\lambda_\Psi(n_\mathrm{s}\mathbf{1} - \mathcal{N})$ at each σ, which adds to the mass m_Ψ; positive (repulsive) for $\lambda_\Psi > 0$. Unit: GeV. Status: DERIVED.
- G_KS: The dimensionless strength of the self-interaction of the gas of particles χ in the Kohn-Sham states: the self-coupling λ_Ψ times the number density n_χ of the particles on the fold divided by the volume V_* of the extra times, $G_\mathrm{KS} = \lambda_\Psi n_\chi / V_*$ (in units of m_Ψ when numbers are given). Unit: dimensionless. Value: 10^{-6} to 1 in the solutions. Status: DERIVED.

- $\hat{\rho}(\sigma)$: The scalar density of the occupied orbitals, $\hat{\rho} = \frac{1}{4}\sum_o \bar{\chi}_o \chi_o$ (per particle); the scalar density of the gas is $n_\mathrm{s} = (n_\chi/V_*)\,\mathrm{e}^{6H|\sigma|}\hat{\rho}$. Unit: GeV. Status: DERIVED.
- $\mathbf{1}$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space). Unit: dimensionless. Status: STANDARD.
- $\hat{\mathcal{N}}(\sigma)$: The one-body density matrix of the occupied orbitals, $\hat{\mathcal{N}} = \frac{1}{4}\sum_o \chi_o \bar{\chi}_o$ (a 16×16 matrix at each σ, per particle); the density matrix of the gas is $\mathcal{N} = (n_\chi/V_*)\,\mathrm{e}^{6H|\sigma|}\hat{\mathcal{N}}$. Unit: GeV. Status: DERIVED.
- μ_X, μ_L, μ_R: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_\mathrm{L}P_\mathrm{L} + \mu_\mathrm{R}P_\mathrm{R})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- P_R: The author's right-handed chiral projector, $P_\mathrm{R} = (\mathbf{1} + \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $\mathfrak{B}$: The Krein operator, $\mathfrak{B} = \mathrm{i}\sigma_{16}\gamma^{(4)}$: a Hermitian involution with the signature (8, 8); $\Psi^\dagger\mathfrak{B}\Psi$ is the charge density. Unit: dimensionless. Status: DERIVED.
- δ: The Kronecker delta: with two indices $\delta^M_N = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with upper and lower index strings the generalised Kronecker delta $\delta^{M_1\cdots M_n}_{N_1\cdots N_n}$, the determinant of the $n\times n$ matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- λ_Ψ: The coupling of the contact self-interaction of dirac16complex. Unit: GeV^{-6}. Status: RECORD.
- n_χ: The number density of the dark-matter particles χ (particles plus antiparticles). Unit: GeV^3. Status: DERIVED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.

FE-41 for this state, with $m_\Psi = 1$ as the unit wherever numbers are given. The factor $\mathrm{e}^{3H|\sigma|}$ removes the first term of the spin connection FE-14 exactly ($\gamma^{(0)}\partial_\sigma \mathrm{e}^{3H|\sigma|} = 3H\,\mathrm{sgn}(\sigma)\gamma^{(0)}\mathrm{e}^{3H|\sigma|}$), and the second vanishes on the static block, so the four occupied orbitals $\chi_o(\sigma)$ obey a Dirac equation of one variable. Gravity enters through the warp: the proper scalar density of the gas at σ is $\mathrm{e}^{6H|\sigma|}$ times the density of the profiles, so the mean field $\mathcal{M}$ of the self-interaction (the Hartree part $\hat{\rho}\,\mathbf{1}$ and the exchange part $\hat{\mathcal{N}}$, the one-body density matrix) carries that factor. The Standard Model acts through the matching at the fold (FE-39). The occupied orbitals are the four states of the bound level with positive energy: Krein charge -1 each, energy $-\omega\mathcal{Q}_o = \omega$ (FE-43), mutually Krein-orthogonal. G_KS is the dimensionless strength of the self-interaction: the self-coupling λ_Ψ times the number density n_χ of the particles on the fold, divided by the volume V_* of the extra times (the orbital of each particle is $\mathrm{e}^{3H|\sigma|}\chi_o$ divided by the square root of the volume of ordinary space and of V_*, so that $\lambda_\Psi n_\mathrm{s} = G_\mathrm{KS}\,\mathrm{e}^{6H|\sigma|}\hat{\rho}$). Status: DERIVED (from FE-41, FE-14, FE-39).

$$\Sigma^{12}\chi_1 = \Sigma^{56}\chi_1 = \frac{\mathrm{i}}{2}\,\chi_1 \tag{S-15}$$

Symbols in (S-15).

- Σ^{ab}, $\Sigma^{\mu\nu}$: The generators of Spin(4,4) on dirac16complex, $\Sigma^{ab} = \frac{1}{4}[T^a, T^b]$ (the scaled commutators of the author's matrices); $\Sigma^{\mu\nu} = \mathbf{e}_a{}^\mu \mathbf{e}_b{}^\nu \Sigma^{ab}$ along the fold. Unit: dimensionless. Status: AUTHOR.
- $\chi_o(\sigma)$, $\bar{\chi}_o$: The profile of the occupied Kohn-Sham orbital o ($o = 1, \ldots, 4$, the four positive-energy states of the bound level) of a homogeneous gas of the particles χ at rest on the static block: a 16-component spinor function of σ; the orbital of each particle is $\mathrm{e}^{3H|\sigma|}\chi_o(\sigma)\mathrm{e}^{-\mathrm{i}\omega x_4}$ divided by the square root of the volume of ordinary space and of V_*; normalised to the Krein charge -1, $\int \mathrm{d}\sigma\, \chi_o^\dagger \mathfrak{B}\chi_o = -1$; $\bar{\chi}_o = \chi_o^\dagger\sigma_{16}$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{1/2}$. Status: DERIVED.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- T_a, T^a: The author's eight real 16×16 Dirac matrices T16ˆA[a] of 4+4.nb ($a = 0, \ldots, 7$; $T_0, \ldots, T_7$), with $\{T_a, T_b\} = 2\eta_{ab}\mathbf{1}$; $T^a = \eta^{ab}T_b$. Unit: dimensionless. Status: AUTHOR.

The symmetry that makes the problem small. The author's $T_0, \ldots, T_3$ are symmetric and $T_4, \ldots, T_7$ antisymmetric,

so the rotations of ordinary space (Σ^{ij}, $i,j=1,2,3$) and of the extra times (Σ^{rs}, $r,s=5,6,7$) form a compact group $SU(2)\times SU(2)$ of real orthogonal matrices, and $\gamma^{(0)}$, $\gamma^{(4)}$, P_{L}, P_{R}, σ_{16} and the Krein operator $\mathfrak{B}$ commute with it: the 16 components split into four copies of the representation $(2,2)$ of this group. The four occupied orbitals form one $(2,2)$ multiplet, the mean field is invariant, and the equations reduce exactly to the four-dimensional weight space S-15 (the states with the eigenvalue i/2 of both Σ^{12} and Σ^{56}, which hold the orbital $o=1$): one orbital of four components, from which the multiplet is recovered by rotations by π (the orbitals $o=2,3,4$ have the eigenvalues $(-\mathrm{i}/2,\mathrm{i}/2)$, $(\mathrm{i}/2,-\mathrm{i}/2)$ and $(-\mathrm{i}/2,-\mathrm{i}/2)$ of Σ^{12} and Σ^{56}) and checked in sixteen components. Status: DERIVED (`um3_kohnsham.py`, K1).

$$\begin{aligned}
&\mathsf{E}_{\mathrm{int}}=\frac{G_{\mathrm{KS}}}{2}\int_{-\infty}^{\infty}\mathrm{d}\sigma\,\mathrm{e}^{6H|\sigma|}\Big(\hat{\rho}^2-\operatorname{tr}\hat{\mathcal{N}}^2\Big),\qquad \epsilon_{\mathrm{p}}=\omega-\mathsf{E}_{\mathrm{int}},\qquad P_{\mathrm{KS}}=q_\chi=n_\chi\,\mathsf{E}_{\mathrm{int}},\\
&\Delta\omega=G_{\mathrm{KS}}\int_{-\infty}^{\infty}\mathrm{d}\sigma\,\mathrm{e}^{6H|\sigma|}\Big(\hat{\rho}^2-\operatorname{tr}\hat{\mathcal{N}}^2\Big)\Big|_{G_{\mathrm{KS}}=0}\qquad \big(q_{\mathrm{b}}>3H/2\big)
\end{aligned}\tag{S-16}$$

Symbols in (S-16).

- $\mathsf{E}_{\mathrm{int}}$: The interaction energy per particle of the gas of χ: the mean self-interaction energy $\frac{\lambda_\Psi}{2}(n_{\mathrm{s}}^2-\operatorname{tr}\mathcal{N}^2)$ integrated across the hidden direction, per particle. Unit: GeV. Status: DERIVED.
- G_{KS}: The dimensionless strength of the self-interaction of the gas of particles χ in the Kohn-Sham states: the self-coupling λ_Ψ times the number density n_χ of the particles on the fold divided by the volume V_* of the extra times, $G_{\mathrm{KS}}=\lambda_\Psi n_\chi/V_*$ (in units of m_Ψ when numbers are given). Unit: dimensionless. Value: 10^{-6} to 1 in the solutions. Status: DERIVED.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma=|\cot z|\,\mathrm{d}x_0$, $\sigma=\operatorname{sgn}(\pi/2-z)\ln(\sin z)/(6H)$ on one period $0<z<\pi$; $\sigma=0$ is the fold $z=\pi/2$, $\sigma\to\mp\infty$ the tips $z\to0,\pi$; one period of x_0 is the whole line $-\infty<\sigma<\infty$. Unit: length. Status: DERIVED.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda=-10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- $\hat{\rho}(\sigma)$: The scalar density of the occupied orbitals, $\hat{\rho}=\frac{1}{4}\sum_o\bar{\chi}_o\chi_o$ (per particle); the scalar density of the gas is $n_{\mathrm{s}}=(n_\chi/V_*)\,\mathrm{e}^{6H|\sigma|}\hat{\rho}$. Unit: GeV. Status: DERIVED.
- tr: The trace of a matrix: the sum of its diagonal entries. Status: STANDARD.
- $\hat{\mathcal{N}}(\sigma)$: The one-body density matrix of the occupied orbitals, $\hat{\mathcal{N}}=\frac{1}{4}\sum_o\chi_o\bar{\chi}_o$ (a 16×16 matrix at each σ, per particle); the density matrix of the gas is $\mathcal{N}=(n_\chi/V_*)\,\mathrm{e}^{6H|\sigma|}\hat{\mathcal{N}}$. Unit: GeV. Status: DERIVED.
- ϵ_{p}: The energy per particle of the gas of χ: the orbital energy minus the interaction energy per particle, which the orbital energy counts twice. Unit: GeV. Status: DERIVED.
- ω: A frequency: the energy of a mode $\propto\mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- P_{KS}: The pressure of the gas of χ along ordinary space in the Kohn-Sham state: the particles are at rest, so it is the mean self-interaction, $P_{\mathrm{KS}}=n_\chi\mathsf{E}_{\mathrm{int}}$. Unit: GeV^4. Status: DERIVED.
- q_χ: The pressure of the bound states of dirac16complex along the extra times, integrated across the fold (their on-shell Lagrangian, the self-interaction energy per volume; zero for a dilute gas). Unit: GeV^4. Status: DERIVED.
- n_χ: The number density of the dark-matter particles χ (particles plus antiparticles). Unit: GeV^3. Status: DERIVED.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- q_{b}: The inverse width of a state of dirac16complex bound to the fold: its profile falls as $\mathrm{e}^{-q_{\mathrm{b}}|\sigma|}$. Unit: GeV. Status: DERIVED.

The energy of the state (FE-42 for the gas). $\mathsf{E}_{\mathrm{int}}$ is the interaction energy per particle, the mean value $\frac{\lambda_\Psi}{2}(n_{\mathrm{s}}^2-\operatorname{tr}\mathcal{N}^2)$ of the self-interaction per particle (the trace runs over the sixteen components). The energy per particle is $\epsilon_{\mathrm{p}}=\omega-\mathsf{E}_{\mathrm{int}}$, because the orbital energy ω counts the interaction twice. The particles are at rest and do not depend on the extra times, so their kinetic terms contribute nothing to the pressure along ordinary space or along the extra times, and

both pressures are the mean self-interaction, $P_{\rm KS} = q_\chi = n_\chi \mathsf{E}_{\rm int}$. $\Delta\omega$ is the first-order shift of the level, the mean of the mean field over the orbitals without the self-interaction (with the charge -1 of an orbital, a perturbation $\delta\mathcal{M}$ shifts its frequency by $\int \bar\chi_o\, \delta\mathcal{M}\, \chi_o\, \mathrm{d}\sigma$, and the average over the four orbitals gives S-16): it equals $2\mathsf{E}_{\rm int}$ to first order, and it is finite when the integrand, which behaves as $\mathrm{e}^{(6H-4q_{\rm b})|\sigma|}$, decays, that is for $q_{\rm b} > 3H/2$. Status: DERIVED.

$$\omega - \omega_{\rm b} \propto G_{\rm KS} \quad \big(q_{\rm b} > 3H/2\big), \qquad \omega - \omega_{\rm b} \propto \big(G_{\rm KS}\big)^{q_{\rm b}/(3H-q_{\rm b})} \quad \big(q_{\rm b} < 3H/2\big) \qquad (G_{\rm KS} \to 0) \tag{S-17}$$

Symbols in (S-17).

- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- $\omega_{\rm b}$: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- $G_{\rm KS}$: The dimensionless strength of the self-interaction of the gas of particles χ in the Kohn-Sham states: the self-coupling λ_Ψ times the number density n_χ of the particles on the fold divided by the volume V_* of the extra times, $G_{\rm KS} = \lambda_\Psi n_\chi / V_*$ (in units of m_Ψ when numbers are given). Unit: dimensionless. Value: 10^{-6} to 1 in the solutions. Status: DERIVED.
- $q_{\rm b}$: The inverse width of a state of dirac16complex bound to the fold: its profile falls as $\mathrm{e}^{-q_{\rm b}|\sigma|}$. Unit: GeV. Status: DERIVED.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.

The dilute limit. For $q_{\rm b} < 3H/2$ the first-order integral diverges: the proper density of the orbital without the self-interaction grows as $\mathrm{e}^{(6H-2q_{\rm b})|\sigma|}$, and the mean field becomes of order one at the distance $\sigma_* = \ln(1/G_{\rm KS})/(6H-2q_{\rm b})$, beyond which it saturates (S-18). The shift is dominated by $|\sigma| < \sigma_*$ and scales as $G_{\rm KS}\, \mathrm{e}^{(6H-4q_{\rm b})\sigma_*} = (G_{\rm KS})^{q_{\rm b}/(3H-q_{\rm b})}$, a power below 1. Status: DERIVED (the scaling), verified numerically (K4).

$$\mathrm{e}^{6H|\sigma|}\, \hat\rho(\sigma) \to \hat\rho_\infty, \qquad |\chi_o| \propto \mathrm{e}^{-3H|\sigma|}, \qquad \det\Big(\gamma^{(0)}\big(\mathrm{i}\omega\gamma^{(4)} + m_\Psi + \mathcal{M}_\infty\big) \pm 3H\, \mathbf{1}\Big) = 0 \qquad (\pm\sigma \to \infty) \tag{S-18}$$

Symbols in (S-18).

- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- H: The author's constant (an inverse length) that fixes the period $\pi/(6H)$ of x_0, the Lovelock couplings and the cosmological constant $\Lambda = -10H^2$; in UM-3 a free parameter. Unit: GeV (1/length). Status: AUTHOR.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\, \mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z) \ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\hat\rho(\sigma)$: The scalar density of the occupied orbitals, $\hat\rho = \frac{1}{4}\sum_o \bar\chi_o \chi_o$ (per particle); the scalar density of the gas is $n_{\rm s} = (n_\chi/V_*)\, \mathrm{e}^{6H|\sigma|}\hat\rho$. Unit: GeV. Status: DERIVED.
- $\hat\rho_\infty$: The constant that the proper scalar density $\mathrm{e}^{6H|\sigma|}\hat\rho$ approaches far from the fold where it saturates. Unit: GeV. Status: DERIVED.
- $\chi_o(\sigma)$, $\bar\chi_o$: The profile of the occupied Kohn-Sham orbital o ($o = 1, \ldots, 4$, the four positive-energy states of the bound level) of a homogeneous gas of the particles χ at rest on the static block: a 16-component spinor function of σ; the orbital of each particle is $\mathrm{e}^{3H|\sigma|}\chi_o(\sigma)\mathrm{e}^{-\mathrm{i}\omega x_4}$ divided by the square root of the volume of ordinary space and of V_*; normalised to the Krein charge -1, $\int \mathrm{d}\sigma\, \chi_o^\dagger \mathfrak{B} \chi_o = -1$; $\bar\chi_o = \chi_o^\dagger \sigma_{16}$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{1/2}$. Status: DERIVED.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- γ^M, $\gamma^{(a)}$: The Dirac matrices of dirac16complex along the coordinates, $\gamma^M = \mathbf{e}_a{}^M T^a$, $\gamma_M = g_{MN}\gamma^N$; $\gamma^{(a)} = T^a$ along the frame direction (a) ($\gamma^{(0)}$ along σ, $\gamma^{(4)}$ along the time). Unit: dimensionless. Status: AUTHOR.
- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.
- $\mathcal{M}_\infty$: The limit of the mean field far from the fold where the proper density of the gas has saturated: the matrix for which the local equation has a solution decaying exactly at the rate $3H$. Unit: GeV. Status: DERIVED.

- **1**: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space). Unit: dimensionless. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.

Saturation. Where the proper density of the orbital would grow, a repulsive self-interaction stops it: if the orbital decayed more slowly than $\mathrm{e}^{-3H|\sigma|}$ the mean field would grow and raise the local mass, so that the orbital decays faster; if it decayed faster the mean field would vanish. The self-consistent orbital therefore decays exactly at the rate $3H$ far from the fold, its proper density approaches a constant $\hat{\rho}_\infty$, and the far mean field $\mathcal{M}_\infty$ is the one for which the local equation has a solution decaying at the rate $3H$ (S-18, the upper sign for $\sigma > 0$). The particles fill the bulk at a constant proper density there, but their number per area of the fold stays finite, because the proper volume of the bulk shrinks as $\mathrm{e}^{-6H|\sigma|}$. Saturation happens far from the fold for every $G_{\mathrm{KS}} > 0$ when $q_{\mathrm{b}} < 3H$, and, when $q_{\mathrm{b}} > 3H$, once the self-interaction has lowered the decay rate of the orbital below $3H$. Status: DERIVED, verified numerically (K5, K8).

$$\mu_{\mathrm{L}} = \mu_{\mathrm{L}0}\Big(1 - \kappa_{\mathrm{H}}\,\frac{\hat{\rho}_{\mathrm{L}}(0)}{\hat{\rho}_{\mathrm{L}0}(0)}\Big), \qquad \hat{\rho}_{\mathrm{L}} = \frac{1}{4}\sum_{o=1}^{4} \bar{\chi}_o P_{\mathrm{L}} \chi_o, \qquad \kappa_{\mathrm{H}} = \frac{c_\Phi\, n_\chi\, \hat{\rho}_{\mathrm{L}0}(0)}{\lambda\, v^2\, V_*} \tag{S-19}$$

Symbols in (S-19).

- μ_X, μ_{L}, μ_{R}: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_{\mathrm{L}} P_{\mathrm{L}} + \mu_{\mathrm{R}} P_{\mathrm{R}})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- $\mu_{\mathrm{L}0}$: The strength μ_{L} of the fold for the left-handed half without the feedback of the gas on the Higgs field. Unit: dimensionless. Value: -0.6 in the solutions. Status: DERIVED.
- κ_{H}: The strength of the Higgs-portal loop of the gas: the gas without the feedback changes the local v_{loc}^2 at the fold by the factor $1 - \kappa_{\mathrm{H}}$, $\kappa_{\mathrm{H}} = c_\Phi n_\chi \hat{\rho}_{\mathrm{L}0}(0)/(\lambda v^2 V_*)$; negative where the Higgs portal binds the fold ($c_\Phi < 0$, $\hat{\rho}_{\mathrm{L}0}(0) > 0$): the gas raises v_{loc}^2. Unit: dimensionless. Value: -0.3 in the test of the loop. Status: ASSUMED (the value), DERIVED (the sign).
- $\hat{\rho}_{\mathrm{L}}$: The scalar density of the left-handed half of the occupied orbitals, $\hat{\rho}_{\mathrm{L}} = \frac{1}{4}\sum_o \bar{\chi}_o P_{\mathrm{L}} \chi_o$; at the fold the mean of its values on the two sides. Unit: GeV. Status: DERIVED.
- $\hat{\rho}_{\mathrm{L}0}$: The left-handed scalar density $\hat{\rho}_{\mathrm{L}}$ of the bound state without the self-interaction and without the Higgs feedback. Unit: GeV. Status: DERIVED.
- $\chi_o(\sigma)$, $\bar{\chi}_o$: The profile of the occupied Kohn-Sham orbital o ($o = 1, \ldots, 4$, the four positive-energy states of the bound level) of a homogeneous gas of the particles χ at rest on the static block: a 16-component spinor function of σ; the orbital of each particle is $\mathrm{e}^{3H|\sigma|}\chi_o(\sigma)\mathrm{e}^{-\mathrm{i}\omega x_4}$ divided by the square root of the volume of ordinary space and of V_*; normalised to the Krein charge -1, $\int \mathrm{d}\sigma\, \chi_o^\dagger \mathfrak{B} \chi_o = -1$; $\bar{\chi}_o = \chi_o^\dagger \sigma_{16}$ is its Dirac adjoint. Unit: $\mathrm{GeV}^{1/2}$. Status: DERIVED.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- P_{L}: The author's left-handed chiral projector, $P_{\mathrm{L}} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- n_χ: The number density of the dark-matter particles χ (particles plus antiparticles). Unit: GeV^3. Status: DERIVED.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_{\mathrm{F}})^{-1/2}$ with the Fermi constant $G_{\mathrm{F}} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4} b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\, \mathrm{d}x_6\, \mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4} b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.

The Higgs-portal loop. Where the Higgs portal alone sets the strength of the fold for the left-handed half (the condensates of the gauge fields and the fermion densities of FE-39 negligible), μ_{L} is proportional to $\langle \Phi^\dagger \Phi \rangle = v_{\mathrm{loc}}^2/2$, and the left-handed scalar density of the gas at the fold shifts v_{loc}^2 (FE-49: $v_{\mathrm{loc}}^2 = v^2 - c_\Phi \langle \bar{\Psi} P_{\mathrm{L}} \Psi \rangle/\lambda$, with $\langle \bar{\Psi} P_{\mathrm{L}} \Psi \rangle = n_\chi \hat{\rho}_{\mathrm{L}}(0)/V_*$ at the fold). $\mu_{\mathrm{L}0}$ and $\hat{\rho}_{\mathrm{L}0}(0)$ are the values without the feedback, $\hat{\rho}_{\mathrm{L}}(0)$ is the mean of the values on the two sides of the fold, and κ_{H} measures the strength of the loop: the gas without feedback

changes $v_{\rm loc}^2$ by the factor $1-\kappa_{\rm H}$. The sign of $\kappa_{\rm H}$ is fixed. Where the Higgs portal alone binds the fold ($\mu_{\rm R}=0$), $\mu_{\rm L0}=c_\Phi v^2/(2V_*)<0$, so $c_\Phi<0$; and the occupied orbitals have $\hat\rho_{\rm L0}(0)=\partial\omega_{\rm b}/\partial\mu_{\rm L}=-m_\Psi^2\mu_{\rm L0}/(4\omega_{\rm b})>0$ (the Hellmann-Feynman theorem: with the Krein charge -1, the shift of the level is the mean value of the perturbation $\delta\mu_{\rm L}P_{\rm L}$ at the fold; FE-40 for the derivative). Hence $\kappa_{\rm H}=2\mu_{\rm L0}n_\chi\hat\rho_{\rm L0}(0)/(\lambda v^4)<0$: the gas raises $v_{\rm loc}^2$, the fold binds more strongly, and the level falls, in agreement with the coupling $\mathfrak{g}_{h\chi}<0$ of FE-47. If $\hat\rho_{\rm L}(0)$ were proportional to $q_{\rm b}$, and $q_{\rm b}$ to $\mu_{\rm L}$, the loop would end at $\mu_{\rm L}=\mu_{\rm L0}/(1+\kappa_{\rm H})$, which does not exist for $\kappa_{\rm H}\le-1$: a gas dense enough would deepen its own binding until $q_{\rm b}$ approaches m_Ψ, where this proportionality fails (ESTIMATE). The orbitals and $\mu_{\rm L}$ are iterated together. Status: DERIVED (from FE-39, FE-40 and FE-49), ASSUMED (the value of $\kappa_{\rm H}$ used below).

8.2 The method

The bound level is found by shooting. On each side of the fold the equation S-14 is integrated inward from $|\sigma|=L$ with the classical fourth-order Runge-Kutta method of step h (the step written as a matrix and the solutions re-orthonormalised every eight steps), starting with the solutions that decay at $|\sigma|=L$; the frequency is the zero of the residual of the matching at the fold for a matching vector of negative Krein norm, searched within $0.05m_\Psi$ of the previous level (a frequency beyond the local threshold at the far end, or a mean field so large that the integration overflows, has no bound level). The orbital gives the densities, the densities the mean field, and the mean field (with $\mu_{\rm L}$ for the Higgs loop) is iterated with Anderson mixing. Plain mixing would need a very small weight: far from the fold, where the density saturates, a small change of the mean field changes the decay rate of the orbital and with it the far density by a large factor of the opposite sign, so the linearised Kohn-Sham map has large negative eigenvalues (an argument; the eigenvalues are not computed). The mixing (the weight β of the new field and the depth of the history) is $\beta=0.5$ with six previous iterates, or, where this does not converge (above the free threshold, where the saturated region is long), $\beta=0.2$ with ten; the extrapolated field is held to the same bound as the computed one. An iteration has converged when the largest relative change of the mean field is below 10^{-8} and the level changes by less than 10^{-10}; where the precision of the level sets a floor of a few 10^{-8}, the best iterate below 10^{-7} is taken. The coupling is raised by continuation, with intermediate couplings inserted where a step fails. Every converged solution is lifted to sixteen components and checked there: the Kohn-Sham equation by central differences on the grid, with the mean field rebuilt in sixteen components from the lifted orbitals, the matching at the fold, the Krein charges and the Krein orthogonality; the residual of self-consistency is that of the reduced problem, evaluated once more with the accepted field. Status: DERIVED (`um3_kohnsham.py`).

8.3 The solution in three regimes

regime	H	box L	step h	the orbital without self-interaction
I	0.05	60	0.03	$q_{\rm b}>3H$: its proper density decays away from the fold
II	0.15	60	0.02	$3H/2<q_{\rm b}<3H$: its proper density grows, the first-order shift is finite
III	0.3	20	0.02	$q_{\rm b}<3H/2$: its proper density grows, the first-order shift diverges

The three regimes, all with the Higgs portal of the vacuum alone, $\mu_{\rm L}=-0.6$ and $\mu_{\rm R}=0$, so that $q_{\rm b}=0.3$ and $\omega_{\rm b}=\sqrt{1-0.09}=0.953939$ (units of m_Ψ). In each the coupling $G_{\rm KS}$ is raised from 10^{-6} (regime I) or from a coupling so small that the proper density does not saturate inside the box (2.5×10^{-9} in regime II, 6.3×10^{-12} in regime III) to 1. Status: ASSUMED (the parameters), DERIVED (the solutions).

Regime I ($H=0.05$, $6H=0.3$):

G_{KS}	ω	$\omega-\omega_{\mathrm{b}}$	$\mathsf{E}_{\mathrm{int}}$	$P_{\mathrm{KS}}/(n_\chi\epsilon_{\mathrm{p}})$	iterations	decay rate of $\hat{\rho}$ at $0.85L$
10^{-6}	0.95393934	1.36×10^{-7}	6.83×10^{-8}	7.2×10^{-8}	4	0.6000
10^{-5}	0.95394057	1.36×10^{-6}	6.83×10^{-7}	7.2×10^{-7}	4	0.6000
10^{-4}	0.95395285	1.36×10^{-5}	6.82×10^{-6}	7.2×10^{-6}	5	0.5999
10^{-3}	0.95407566	1.36×10^{-4}	6.82×10^{-5}	7.2×10^{-5}	6	0.5991
0.01	0.95530069	0.00136	6.80×10^{-4}	7.1×10^{-4}	7	0.5913
0.1	0.96723519	0.0133	0.00659	0.0069	10	0.5078
1	1.07039318	0.116	0.0564	0.056	42	0.3000

Regime II ($H=0.15$, $6H=0.9$):

G_{KS}	ω	$\omega-\omega_{\mathrm{b}}$	$\mathsf{E}_{\mathrm{int}}$	$P_{\mathrm{KS}}/(n_\chi\epsilon_{\mathrm{p}})$	iterations	decay rate of $\hat{\rho}$ at $0.85L$
10^{-6}	0.95393961	4.09×10^{-7}	2.05×10^{-7}	2.1×10^{-7}	20	0.8677
10^{-5}	0.95394329	4.09×10^{-6}	2.05×10^{-6}	2.1×10^{-6}	35	0.8929
10^{-4}	0.95397992	4.07×10^{-5}	2.03×10^{-5}	2.1×10^{-5}	46	0.8986
10^{-3}	0.95433722	3.98×10^{-4}	1.98×10^{-4}	2.1×10^{-4}	42	0.8997
0.01	0.95764791	0.00371	0.00182	0.0019	42	0.9000
0.1	0.98687286	0.0329	0.0160	0.016	43	0.9000
1	1.25973443	0.306	0.152	0.14	132	0.9000

Regime III ($H=0.3$, $6H=1.8$):

G_{KS}	ω	$\omega-\omega_{\mathrm{b}}$	$\mathsf{E}_{\mathrm{int}}$	$P_{\mathrm{KS}}/(n_\chi\epsilon_{\mathrm{p}})$	iterations	decay rate of $\hat{\rho}$ at $0.85L$
10^{-6}	0.95407252	1.33×10^{-4}	4.47×10^{-5}	4.7×10^{-5}	31	1.7559
10^{-5}	0.95436976	4.31×10^{-4}	1.46×10^{-4}	1.5×10^{-4}	38	1.7874
10^{-4}	0.95536905	0.00143	4.92×10^{-4}	5.2×10^{-4}	35	1.7966
10^{-3}	0.95896972	0.00503	0.00179	0.0019	34	1.7992
0.01	0.97368303	0.0197	0.00747	0.0077	39	1.7999
0.1	1.04743537	0.0935	0.0385	0.038	35	1.8000
1	1.57066625	0.617	0.288	0.22	78	1.8000

The self-consistent states. ω is the orbital energy of the occupied level, $\mathsf{E}_{\mathrm{int}}$ the interaction energy per particle, $P_{\mathrm{KS}}/(n_\chi\epsilon_{\mathrm{p}})$ the pressure divided by the energy density, and the last column the decay rate of the scalar density at $0.85L$ ($6H$ where the proper density has saturated). Status: DERIVED (`um3_kohnsham.py`, K3).

The free level and the symmetry. Without the self-interaction the shooting with the author's 16×16 matrices gives $\omega=0.9539392014$, which is FE-40 ($\sqrt{1-q_{\mathrm{b}}^2}=0.9539392014$) to 10^{-10}; the level consists of eight degenerate states, four of each Krein sign, and the reduced four-dimensional problem of S-15 has the same level with two degenerate states (one of each sign). Status: DERIVED (K1).

The dilute limit. In regime I the shift of the level divided by the first-order shift of S-16 (the coefficient $\Delta\omega/G_{\mathrm{KS}}=0.136501$, computed in sixteen components with the four occupied orbitals of the level of eight degenerate states) is 0.99997, 0.99997, 0.99995, 0.99972 for $G_{\mathrm{KS}}=10^{-6}$, 10^{-5}, 10^{-4} and 10^{-3}: the Kohn-Sham states reduce to first-order perturbation theory (the remaining 3×10^{-5} is of the size of the quadrature error on the grid). Fitted for $G_{\mathrm{KS}}\le10^{-4}$, the shift grows as G_{KS}^{p} with $p=1.000$, 0.999 and 0.515 in the three regimes, against S-17: 1, 1 and $q_{\mathrm{b}}/(3H-q_{\mathrm{b}})=0.500$. Status: DERIVED (K2, K4).

Saturation. At $G_{\mathrm{KS}}=0.01$ the scalar density decays far from the fold at 0.9000 in regime II and 1.7999 in regime III, the rates $6H=0.9$ and 1.8 of S-18: the proper density has saturated, and the far mean field is the one that

gives the orbital the decay rate $3H$. An attractive self-interaction behaves differently in the two kinds of regime: for $G_{\mathrm{KS}} = -10^{-4}$ and -10^{-2} the iteration, started from the free state, converges in regime I (where the proper density decays and the attraction only lowers the level, to $\omega = 0.95392555$ and 0.95257077); in regimes II and III, where the proper density of the free state grows away from the fold, the level search finds no Krein-negative level near the previous one: the attraction grows with the proper density and would close the local gap. That no self-consistent state exists there at all is not proven by this (OPEN). Status: DERIVED (K5).

Through the free threshold. The repulsive mean field raises the level through the threshold $\omega = m_\Psi$ of the free continuum: at $G_{\mathrm{KS}} = 0.371$ in regime I and at $G_{\mathrm{KS}} = 0.142$ in regime II. The state stays bound up to $G_{\mathrm{KS}} = 1$ ($\omega = 1.070393$ and 1.259734 there): far from the fold the proper density saturates (in regime I once the decay rate of the orbital has fallen below $3H$, at $\omega > \sqrt{1 - 9H^2} = 0.9887$), the density decays at $6H$ (to within 5×10^{-4} and 1×10^{-8} above the threshold), and the far mean field keeps the local mass above ω. In the computed cases a repulsive self-interaction therefore does not dissolve the particles χ into the bulk: it makes them heavier and spreads them, and their energy may exceed m_Ψ; the mechanism, the saturation of the proper density, rests on the warp and needs $H > 0$. Status: DERIVED (K8).

The Higgs-portal loop. The occupied orbitals without the self-interaction have $\hat{\rho}_{\mathrm{L}0}(0) = 0.15723848$, equal to $\partial\omega_{\mathrm{b}}/\partial\mu_{\mathrm{L}} = -\mu_{\mathrm{L}0}/(4\omega_{\mathrm{b}}) = 0.15724273$ up to the relative difference 2.7×10^{-5}, which is the error $(2q_{\mathrm{b}}h)^2/12 = 2.7 \times 10^{-5}$ of the trapezoidal normalisation on the grid; so $\kappa_{\mathrm{H}} < 0$. With $\kappa_{\mathrm{H}} = -0.3$ and $G_{\mathrm{KS}} = 0.1$ in regime I the orbitals and μ_{L} of S-19 converge together in 12 iterations: $\mu_{\mathrm{L}} = -0.874681$ (from -0.6; the estimate $\mu_{\mathrm{L}0}/(1 + \kappa_{\mathrm{H}})$ gives -0.8571): the gas raises the local electroweak vacuum, the fold binds more strongly, and the level falls to $\omega = 0.91512441$ (0.96723519 without the feedback). Status: DERIVED (K6), ASSUMED (the value of κ_{H}).

The equation of state. The pressure is positive for the repulsive self-interaction and vanishes in the dilute limit: $P_{\mathrm{KS}}/(n_\chi \epsilon_{\mathrm{p}}) = 7.2 \times 10^{-8}$, 2.1×10^{-7} and 4.7×10^{-5} at $G_{\mathrm{KS}} = 10^{-6}$ in the three regimes, and 0.056, 0.137 and 0.224 at $G_{\mathrm{KS}} = 1$. A dilute gas of the particles χ is pressureless: cold dark matter. Status: DERIVED (K7).

The residuals. Each of the 83 converged solutions (the three regimes from the smallest coupling to 1, with the continuation between) was lifted to sixteen components and checked there: the largest relative residual of the Kohn-Sham equation by central differences on the grid is 2×10^{-4} (the error of the difference formula, of order h^2), of the matching at the fold 4×10^{-9}, of self-consistency (in the reduced problem, evaluated once more with the accepted field) 2×10^{-7}; the four orbitals have the Krein charge -1 to 4×10^{-16} and are Krein-orthogonal to 5×10^{-17}, and the bound on the mean field is never active. Status: DERIVED (K3).

8.4 A weakly bound gas

The direct-detection bound on the Higgs portal requires a weakly bound χ, $q_{\mathrm{b}}/\omega_{\mathrm{b}} \lesssim 0.03$ at a few TeV, and a nearly luminal one needs $q_{\mathrm{b}} \gg H$ (S-13). A point of this kind is solved as well: $\mu_{\mathrm{L}} = -0.06$ ($q_{\mathrm{b}} = 0.03$, $q_{\mathrm{b}}/\omega_{\mathrm{b}} = 0.0300$), $H = 0.002$ (so $q_{\mathrm{b}} = 15H$, regime I, and $c_{\mathrm{b}}^2 = q_{\mathrm{b}}/(q_{\mathrm{b}} - H) = 1.071$), with the box $L = 500$ and the step $h = 0.25$. The shooting in sixteen components gives $\omega = 0.9995498987$ (FE-40: 0.9995498987) with four Krein-negative orbitals, and the first-order coefficient of S-16 is $\Delta\omega/G_{\mathrm{KS}} = 0.012489$. The self-consistent shift divided by the first-order shift is 0.99940, 0.99429, 0.94309 for $G_{\mathrm{KS}} = 10^{-4}$, 10^{-3} and 10^{-2} (approaching 1 as $G_{\mathrm{KS}} \to 0$), with $P_{\mathrm{KS}}/(n_\chi \epsilon_{\mathrm{p}}) = 6.2 \times 10^{-7}$, 6.2×10^{-6}, 5.8×10^{-5}: a dilute gas of weakly bound particles χ is cold, and its self-interaction raises the orbital energy by about $0.012\, m_\Psi$ per unit of G_{KS} and the energy per particle, the mass of χ in the gas, by half of it. Status: DERIVED (K9).

8.5 The Krein partner and the stability of the gas

$$\left\{\mathsf{a}_o, \mathsf{a}_o^\dagger\right\} = -\mathcal{Q}_o, \qquad \mathsf{H}_\Psi = \sum_o \left(-\mathcal{Q}_o\, \omega_o\right) \mathsf{a}_o^\dagger \mathsf{a}_o, \qquad \left[\mathsf{H}_\Psi, \mathsf{a}_o^\dagger\right] = \omega_o\, \mathsf{a}_o^\dagger \tag{FE-67}$$

Symbols in (FE-67).

- a_o, $\mathsf{a}_o^\dagger$: The annihilation and the creation operator of the mode o of the quantised dirac16complex (frequency ω_o, Krein charge $\mathcal{Q}_o$); the dagger is the adjoint with respect to the indefinite (Krein) scalar product. Unit: dimensionless. Status: STANDARD.
- o: The label of a single-particle state of dirac16complex: a Kohn-Sham orbital, or a mode of the quantised field; a sum over o runs over the orbitals or the modes. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.

- $\mathcal{Q}_o$: The charge Q_Ψ (the Krein charge) of the single-particle state o of dirac16complex, a Kohn-Sham orbital or a mode: $+1$ or -1 for a normalised state. Unit: dimensionless. Status: DERIVED.
- H_Ψ: The Hamiltonian (the energy operator) of the quantised dirac16complex. Unit: GeV. Status: DERIVED.
- ω: A frequency: the energy of a mode $\propto \mathrm{e}^{-\mathrm{i}\omega x_4}$ (ω_o of the orbital o). Unit: GeV. Status: STANDARD.

The quantised dirac16complex: a mode o of the frequency ω_o and the Krein charge $\mathcal{Q}_o = \pm 1$ (FE-43) has the anticommutator $-\mathcal{Q}_o$, and every quantum raises the energy by its frequency. In the positive-energy quantisation every particle and every hole carries a positive energy and the norm $-\mathcal{Q}_o$ of its mode: the particles and the holes of the modes of charge -1 are physical, those of the modes of charge $+1$ are ghosts (negative norm), and no choice of the vacuum removes them. The Kohn-Sham states above occupy four orbitals of charge -1: they are physical states, and the questions are whether they are stable and whether the couplings of the fold produce ghosts. Status: DERIVED (`um3_krein.py`, KR1).

The symmetry of the bound states and its mirrors. At rest, the transformations of the sixteen components that keep the bound states of the fold (they commute with $\gamma^{(0)}$, $\gamma^{(4)}$ and Γ_8 and preserve the Dirac adjoint) form a Lie algebra of dimension 32: 16 compact generators keep the two Krein charges apart, and 16 boosts mix them, among them the nine boosts Σ^{ir} between ordinary space and the extra times; for a moving particle only four compact generators remain. The mirror R_1 of the fold (FD-10) keeps the charges: the four bound states of charge -1 are even, the four of charge $+1$ odd, and the self-consistent orbital and its mean field are even; R_2 exchanges the two charges. Status: DERIVED (KR2, KR3).

The dilute gas is stable. The boosts that mix the charges map the self-consistent gas onto self-consistent gases (in regime I at $G_{\mathrm{KS}} = 0.1$ the residual of the Kohn-Sham equation stays 2.43×10^{-5} and the charges stay -1, while the orbitals acquire admixtures of charge $+1$ whose positive-definite norm grows up to 2.43 times): these directions are flat, and the linear response of the bound level has all 32 frequencies below 1.8×10^{-4}, far below the splitting 4.815×10^{-3} between the occupied and the empty multiplet. In the self-consistent mean field the partner of the occupied level, the level of charge $+1$, stays a real level above it in the dilute gas of all three regimes. Status: DERIVED (KR4, KR5).

The dense gas is not. The repulsion of the self-interaction pushes the partner level of charge $+1$ onto the occupied level; where they meet (a Krein collision) they form a pair of complex frequencies $\omega \pm \mathrm{i}\Gamma_{\mathrm{K}}$, and a perturbation of the gas grows as $\mathrm{e}^{\Gamma_{\mathrm{K}} x_4}$:

regime	partner real and bound up to	growth rate Γ_{K} (units of m_Ψ)
I ($H = 0.05$)	$G_{\mathrm{KS}} = 0.2$	5.1×10^{-6} at $G_{\mathrm{KS}} = 0.3$, 1.2×10^{-4} at 0.5, 3.5×10^{-4} at 0.7, 6.1×10^{-4} at 1
II ($H = 0.15$)	$G_{\mathrm{KS}} = 0.5$	3.9×10^{-3} at 0.7, 9.7×10^{-3} at 1
III ($H = 0.3$)	$G_{\mathrm{KS}} = 0.7$	1.5×10^{-2} at 1

Status: DERIVED (KR5).

The couplings of the fold decide unitarity. At rest none of the 16 chirality-blind operators of the fold (the products of $\gamma^{(1)}, \ldots, \gamma^{(4)}$) connects a physical state with a ghost, and each of the 16 chiral ones (the same times Γ_8) does. The left-handed Higgs portal P_{L} of the author has the elastic amplitude 0.157, no conversion of a particle into a ghost, no physical pair at rest (a p-wave), and the amplitude 0.140 (an s-wave) for a physical particle χ with a ghost antiparticle, 0.89 times the elastic one. For moving particles the scalar, the density and the vector currents still do not mix (below 10^{-9}), the magnetic dipole does, in proportion to the momentum. The vacuum cannot decay (every quantum has a positive energy), but with chiral couplings at the fold ghosts are produced wherever the portals act: UM-3 with the author's left-handed Higgs portal is not a unitary quantum theory, and a unitary UM-3 would need couplings of the fold that do not distinguish the chiralities. Status: DERIVED (KR6, KR7).

9 The residuals of every field equation

The approximate solution of UM-3 consists of the parts above and below. The field equations that the other sections do not write out are restated first, so that the table at the end of this section can refer to each of them: the field

equations of the bulk, the junction conditions and the thin source of the fold, the conservation law at the fold, the field equations of the gluons and of the hypercharge field, and those of the quarks and leptons.

$$\sum_{p=1}^{3} \alpha_{(p)}\, \mathcal{E}_{(p)}{}^{M}{}_{N} + \Lambda\, \delta^{M}_{N} = \kappa\big(\mathcal{T}_{\Psi}{}^{M}{}_{N} + \mathcal{T}_{\mathrm{stab}}{}^{M}{}_{N}\big) \qquad (\sigma \neq 0) \tag{FE-1}$$

Symbols in (FE-1).

- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations a second label of a Kohn-Sham orbital (as o, with δ_{op} the Kronecker delta of the two labels). Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)} H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- $\mathcal{E}_{(p)}{}^{M}{}_{N}$: The Lovelock tensor of order p: the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to the metric ($\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor); with frame indices its frame components. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- Λ: The eight-dimensional cosmological constant, $\Lambda = -10H^2$ (the author writes the pure number $\Lambda_{\mathrm{N}} = 10$ in $0 = -\Lambda_{\mathrm{N}} + \sum_p H^{-2p} w_{(p)} \mathcal{E}_{(p)}$). Unit: GeV^2. Value: $-10H^2$. Status: AUTHOR.
- δ: The Kronecker delta: with two indices $\delta^{M}_{N} = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with upper and lower index strings the generalised Kronecker delta $\delta^{M_1 \cdots M_n}_{N_1 \cdots N_n}$, the determinant of the $n \times n$ matrix of the Kronecker deltas $\delta^{M_a}_{N_b}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{T}_{\Psi}$: The energy-momentum tensor of dirac16complex, $\mathcal{T}_{MN} = -(2/\sqrt{|g|})\,\delta S_{\Psi}/\delta g^{MN}$. Unit: GeV^8. Status: STANDARD.
- $\mathcal{T}_{\mathrm{stab}}{}^{a}{}_{b}$: The energy-momentum tensor of the stabiliser of the bulk (frame components): the source that a static bulk of the author's metric requires, found and named by Claude (Anthropic) on 2026-10-24. Unit: GeV^8. Status: DERIVED (the value), ASSUMED (a separate medium).
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.

The Einstein-Lovelock equations of the bulk, away from the fold: the Lovelock tensor of the author's theory equals κ times the energy-momentum tensor of the fields of the bulk (dirac16complex and the stabiliser). Status: AUTHOR (the theory), STANDARD (the variation).

$$\begin{aligned} \kappa\, \mathcal{S}^{a}{}_{b} &= \lim_{\varepsilon \to 0} \int_{-\varepsilon}^{\varepsilon} \mathrm{d}\sigma \sum_{p=1}^{3} \alpha_{(p)}\, \mathcal{E}_{(p)}{}^{a}{}_{b} = \mathfrak{F}^{a}{}_{b}\big|_{\sigma=0^+} - \mathfrak{F}^{a}{}_{b}\big|_{\sigma=0^-}, \\ \frac{\partial \mathfrak{F}^{a}{}_{b}}{\partial(\partial_\sigma \mathfrak{f})} &= \frac{\partial}{\partial(\partial^2_\sigma \mathfrak{f})} \sum_{p=1}^{3} \alpha_{(p)}\, \mathcal{E}_{(p)}{}^{a}{}_{b}, \qquad \mathfrak{f} \in \{\mathfrak{a}, \mathfrak{b}, \mathfrak{n}\} \end{aligned} \tag{FE-4}$$

Symbols in (FE-4).

- κ: The eight-dimensional gravitational coupling, $\kappa = 8\pi G_8$. Unit: GeV^{-6}. Status: AUTHOR.
- $\mathcal{S}^{a}{}_{b}$: The thin source of the fold: its energy-momentum tensor per unit proper length across the fold (eight-dimensional normalisation, frame components along the fold); it is fixed by the jump of the first σ-derivatives of the metric across the fold (the junction conditions; $\kappa\mathcal{S}$ is the jump of the junction potential). Unit: GeV^7. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.

- ε: The half-thickness of a layer around the fold, taken to zero. Unit: length. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- p: The Lovelock order $p = 1, 2, 3$ (Einstein, Gauss-Bonnet, cubic), as an index and as a number; in the Kohn-Sham equations a second label of a Kohn-Sham orbital (as o, with δ_{op} the Kronecker delta of the two labels). Status: STANDARD.
- $\alpha_{(p)}$: The Lovelock coupling of order p, $\alpha_{(p)} = w_{(p)} H^{2-2p}$ ($\alpha_{(1)} = 1$). Unit: GeV^{2-2p}. Status: AUTHOR.
- ${\mathcal{E}_{(p)}}^M{}_N$: The Lovelock tensor of order p: the variation of $\sqrt{|g|}\,\mathcal{L}_{(p)}$ with respect to the metric ($\mathcal{E}_{(1)}$ the Einstein tensor, $\mathcal{E}_{(2)}$ the Gauss-Bonnet tensor); with frame indices its frame components. Unit: $1/\mathrm{length}^{2p}$. Status: STANDARD.
- ${\mathfrak{F}^a}_b$: The junction potential: a function of the first σ-derivatives of the metric functions whose derivatives are the coefficients of the second σ-derivatives in $\sum_p \alpha_{(p)} {\mathcal{E}_{(p)}}^a{}_b$ (FE-4). Unit: GeV. Status: DERIVED.
- ∂: A partial derivative: ∂_M with respect to the coordinate X^M, ∂_μ with respect to x^μ, ∂_σ with respect to σ, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- σ: The label σ in ∂_σ: the derivative with respect to the proper hidden coordinate σ. Status: STANDARD.
- $\mathfrak{f}$: Any one of the three metric functions $\mathfrak{a}, \mathfrak{b}, \mathfrak{n}$ of the general warped metric (FE-34). Unit: dimensionless. Status: DERIVED.
- $\mathfrak{a}$, $\mathfrak{a}_1$: The logarithm of the scale factor of ordinary space in the general warped metric FE-34, a function of σ and x_4; for the author's metric $\mathfrak{a} = a_4 - H|\sigma|$; $\mathfrak{a}_n$ its term of order n in the power series FE-35 ($\mathfrak{a}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $\mathfrak{b}$, $\mathfrak{b}_1$: The logarithm of the scale factor of the extra times in the general warped metric FE-34; for the author's metric $\mathfrak{b} = \ln b_4 - a_4 - H|\sigma|$; $\mathfrak{b}_n$ its term of order n ($\mathfrak{b}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.
- $\mathfrak{n}$, $\mathfrak{n}_1$: The logarithm of the lapse in the general warped metric FE-34 ($g_{x_4 x_4} = -\mathrm{e}^{2\mathfrak{n}}$); for the author's metric $\mathfrak{n} = 0$; $\mathfrak{n}_n$ its term of order n ($\mathfrak{n}_1$ the first-order term). Unit: dimensionless. Status: DERIVED.

The field equations at the fold: the metric has a kink at $\sigma = 0$, so its second derivatives along σ contain a delta function, and the jump of the junction potential across the fold equals κ times the thin source $\mathcal{S}$ of the fold. Status: DERIVED.

$$\mathcal{S}^a{}_b = \frac{1}{V_*}\Big(\mathcal{T}_{\mathrm{H}}{}^a{}_b + \mathcal{T}_{\mathrm{gauge}}{}^a{}_b + \mathcal{T}_{\mathrm{f}}{}^a{}_b + \mathcal{T}_{\mathrm{P}}{}^a{}_b + \mathcal{T}_{\mathrm{ff}}{}^a{}_b + \mathcal{T}_{\mathrm{IG}}{}^a{}_b\Big), \qquad \mathcal{T}_{\mathrm{IG}}{}^\mu{}_\nu = -\frac{1}{\kappa_4}\,\hat{G}^\mu{}_\nu, \qquad \mathcal{T}^r{}_r = \mathcal{L}_{\mathrm{fold}} \qquad \text{(FE-5)}$$

Symbols in (FE-5).

- $\mathcal{S}^a{}_b$: The thin source of the fold: its energy-momentum tensor per unit proper length across the fold (eight-dimensional normalisation, frame components along the fold); it is fixed by the jump of the first σ-derivatives of the metric across the fold (the junction conditions; $\kappa\mathcal{S}$ is the jump of the junction potential). Unit: GeV^7. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_{\mathrm{t}}(\mathrm{e}^{-a_4} b_4)^3$ (with $V_{\mathrm{t}} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4} b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- $\mathcal{T}_{\mathrm{H}}$: The energy-momentum tensor of the Higgs doublet (four-dimensional normalisation). Unit: GeV^4. Status: STANDARD.
- $\mathcal{T}_{\mathrm{gauge}}$: The energy-momentum tensor of the gauge fields. Unit: GeV^4. Status: STANDARD.
- $\mathcal{T}_{\mathrm{f}}$: The energy-momentum tensor of the quarks and leptons. Unit: GeV^4. Status: STANDARD.
- $\mathcal{T}_{\mathrm{P}}$: The energy-momentum tensor of the portals of dirac16complex. Unit: GeV^4. Status: DERIVED.
- $\mathcal{T}_{\mathrm{ff}}{}^a{}_b$: The energy-momentum tensor of the fold fluid (frame components of the fold, four-dimensional normalisation). Unit: GeV^4. Status: ASSUMED.

- $\mathcal{T}_{\mathrm{IG}}$: The energy-momentum tensor of the induced gravity of the fold, $\mathcal{T}_{\mathrm{IG}}{}^{\mu}{}_{\nu} = -\hat{G}^{\mu}{}_{\nu}/\kappa_4$. Unit: GeV^4. Status: DERIVED.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- κ_4: The coupling of the induced gravity of the fold, $\kappa_4 = 8\pi G_{\mathrm{ind}}$; four-dimensional gravity on the fold has $8\pi G_{\mathrm{N}} = \kappa_4/(1 + 5\epsilon_{\mathrm{r}}/6)$ (the Friedmann equation of the fold to first order in ϵ_{r}). Unit: GeV^{-2}. Value: $\kappa_4 \simeq 8\pi G_{\mathrm{N}} = 8\pi/M_{\mathrm{Pl}}^2$. Status: ASSUMED.
- $\hat{G}^{\mu}{}_{\nu}$: The Einstein tensor of the four-dimensional metric $\hat{h}_{\mu\nu}$ of the fold. Unit: 1/length^2. Status: STANDARD.
- r: An extra-time direction of the fold: $r \in \{5, 6, 7\}$ (x_5, x_6, x_7); not summed (the three components are equal). Status: STANDARD.
- $\mathcal{T}$: The energy-momentum tensor of any one term of the fold (in $\mathcal{T}^{r}{}_{r} = \mathcal{L}_{\mathrm{fold}}$). Unit: GeV^4. Status: STANDARD.
- $\mathcal{L}_{\mathrm{fold}}$: The Lagrangian of any one term of the fold (four-dimensional normalisation). Unit: GeV^4. Status: STANDARD.

The thin source of the fold: the energy-momentum tensor of everything that lives on the fold (the Standard Model, the portals, the fold fluid and the induced gravity) divided by the volume V_* of the extra times; along each extra time its component is the Lagrangian of the term, because the fields of the fold do not depend on the extra times. Status: DERIVED.

$$\begin{aligned} &\nabla_M \big(\mathcal{T}_{\Psi}{}^{M}{}_{N} + \mathcal{T}_{\mathrm{stab}}{}^{M}{}_{N}\big) = 0 \qquad (\sigma \neq 0), \\ &\hat{\nabla}_a \, \mathcal{S}^{a}{}_{b} = -\Big(\mathcal{T}_{\mathrm{bulk}}{}^{(0)}{}_{b}\big|_{\sigma=0^+} - \mathcal{T}_{\mathrm{bulk}}{}^{(0)}{}_{b}\big|_{\sigma=0^-}\Big) \end{aligned} \tag{FE-11}$$

Symbols in (FE-11).

- ∇: The covariant derivative of a tensor with the Levi-Civita connection of the metric g_{MN} (Christoffel symbols $\Gamma^{P}{}_{MN}$). Status: STANDARD.
- $\mathcal{T}_{\Psi}$: The energy-momentum tensor of dirac16complex, $\mathcal{T}_{MN} = -(2/\sqrt{|g|})\,\delta S_{\Psi}/\delta g^{MN}$. Unit: GeV^8. Status: STANDARD.
- $\mathcal{T}_{\mathrm{stab}}{}^{a}{}_{b}$: The energy-momentum tensor of the stabiliser of the bulk (frame components): the source that a static bulk of the author's metric requires, found and named by Claude (Anthropic) on 2026-10-24. Unit: GeV^8. Status: DERIVED (the value), ASSUMED (a separate medium).
- M: A coordinate index of eight dimensions (also with a numbered copy $M_1, M_2, \ldots$): it runs over $x_0, x_1, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- N: A coordinate index of eight dimensions (also $N_1, N_2, \ldots$), running over $x_0, \ldots, x_7$; a repeated index is summed. Status: STANDARD.
- σ: The proper coordinate of the hidden direction: $\mathrm{d}\sigma = |\cot z|\,\mathrm{d}x_0$, $\sigma = \mathrm{sgn}(\pi/2 - z)\ln(\sin z)/(6H)$ on one period $0 < z < \pi$; $\sigma = 0$ is the fold $z = \pi/2$, $\sigma \to \mp\infty$ the tips $z \to 0, \pi$; one period of x_0 is the whole line $-\infty < \sigma < \infty$. Unit: length. Status: DERIVED.
- $\hat{\nabla}$: The covariant derivative of the fold: with the Levi-Civita connection of the metric of the fold (for four-dimensional tensors of its metric $\hat{h}_{\mu\nu} = \mathrm{diag}(\mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, \mathrm{e}^{2a_4}, -1)$ in the observable coordinates $x_1, \ldots, x_4$; for $\hat{\nabla}_a \mathcal{S}^{a}{}_{b}$ of the induced metric of the seven-dimensional fold). Status: STANDARD.
- $\mathcal{S}^{a}{}_{b}$: The thin source of the fold: its energy-momentum tensor per unit proper length across the fold (eight-dimensional normalisation, frame components along the fold); it is fixed by the jump of the first σ-derivatives of the metric across the fold (the junction conditions; $\kappa\mathcal{S}$ is the jump of the junction potential). Unit: GeV^7. Status: DERIVED.
- a: A frame (orthonormal) index running over $0, 1, \ldots, 7$ (frame direction (0) along σ, (1), (2), (3) along ordinary space, (4) along the time, (5), (6), (7) along the extra times); raised and lowered with the frame metric $\eta_{ab} = \mathrm{diag}(1, 1, 1, 1, -1, -1, -1, -1)$; on the fold ($a, b \in \{1, \ldots, 7\}$) the frame of the fold; a repeated index is summed. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- $\mathcal{T}_{\mathrm{bulk}}$: The energy-momentum tensor of all sources of the bulk, $\mathcal{T}_{\Psi} + \mathcal{T}_{\mathrm{stab}}$; $\mathcal{T}_{\mathrm{bulk}}{}^{(0)}{}_{b}$ is its flux along σ (frame components). Unit: GeV^8. Status: STANDARD.

The conservation law at the fold: the covariant divergence of the thin source along the fold equals minus the jump of the energy and momentum flux of the bulk along σ, so the fold loses exactly what the bulk carries away. Status:

DERIVED.

$$\frac{1}{\sqrt{-\hat{h}}\,\mathcal{Z}}\Big(\partial_\mu\big[\sqrt{-\hat{h}}\,\mathcal{Z}\,\mathcal{K}_G\,G^{A\mu\nu}\big]+g_3\,f^{ABC}G^B_\mu\sqrt{-\hat{h}}\,\mathcal{Z}\,\mathcal{K}_G\,G^{C\mu\nu}\Big)=-g_3\,J_{\rm c}^{A\nu},\qquad \mathcal{K}_G=1+\sum_X c_X^G\,\bar\Psi P_X\Psi,$$
$$J_{\rm c}^{A\nu}=\bar Q\,\hat\gamma^\nu t^A Q+\bar u_{\rm R}\,\hat\gamma^\nu t^A u_{\rm R}+\bar d_{\rm R}\,\hat\gamma^\nu t^A d_{\rm R}$$

(FE-16)

Symbols in (FE-16).

- $\hat h_{\mu\nu}$: The four-dimensional metric of the fold, $\hat h_{\mu\nu}=\mathrm{diag}(\mathrm{e}^{2a_4},\mathrm{e}^{2a_4},\mathrm{e}^{2a_4},-1)$ in the coordinates $x_1,\dots,x_4$ (the induced metric at $\sigma=0$ without the extra times); $\hat h^{\mu\nu}$ its inverse, $\hat h$ its determinant, $\sqrt{-\hat h}=\mathrm{e}^{3a_4}$; $\hat h^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4)=V_{\rm t}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z}=1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- ∂: A partial derivative: ∂_M with respect to the coordinate X^M, ∂_μ with respect to x^μ, ∂_σ with respect to σ, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1,x_2,x_3,x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- $\mathcal{K}_G$, $\mathcal{K}_W$, $\mathcal{K}_B$: The portal factors of the gauge kinetic terms, $\mathcal{K}_G=1+\sum_X c_X^G\bar\Psi P_X\Psi$ (likewise for W and B): the gauge couplings at the fold are divided by $\sqrt{\mathcal{K}}$. Unit: dimensionless. Value: 1 where dirac16complex vanishes. Status: DERIVED.
- G^A_μ, $G^A_{\mu\nu}$: The gluon field of $SU(3)_C$ on the fold ($A=1,\dots,8$) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- A: A colour (adjoint SU(3)) index running over $1,\dots,8$; a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- g_3: The $SU(3)_C$ gauge coupling (the strong coupling g_s). Unit: dimensionless. Value: $g_3(m_Z)=1.2177$ ($\alpha_s=0.1180$). Status: STANDARD.
- f^{ABC}: The structure constants of $SU(3)$, $[t^A,t^B]=\mathrm{i}f^{ABC}t^C$. Unit: dimensionless. Status: STANDARD.
- B: A colour (adjoint SU(3)) index (as A). Status: STANDARD.
- C: A colour (adjoint SU(3)) index (as A). Status: STANDARD.
- $J_{\rm c}^{A\nu}$: The colour current of the quarks. Unit: GeV^3. Status: STANDARD.
- X: The label of a half of dirac16complex: $X\in\{\mathrm{L},\mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c_X^G: The gluon-kinetic portal coupling of the half X of dirac16complex. Unit: GeV^{-7}. Status: INVENTED.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi=\Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma=0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- P_X: The chiral projector of the half X: $P_X=P_{\rm L}$ for $X=\mathrm{L}$, $P_{\rm R}$ for $X=\mathrm{R}$. Unit: dimensionless. Status: AUTHOR.
- Q, $\bar Q$: The left-handed quark doublets of the three generations, $Q=(u_{\rm L},d_{\rm L})$ (colour triplets, hypercharge 1/6). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- $\hat\gamma^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold, $\hat\gamma^\mu=\hat{\mathbf{e}}_a{}^\mu\hat\gamma^{(a)}$, $\{\hat\gamma^\mu,\hat\gamma^\nu\}=-2\hat h^{\mu\nu}\mathbf{1}$; $\hat\gamma_\mu=\hat h_{\mu\nu}\hat\gamma^\nu$. Unit: dimensionless. Status: STANDARD.
- t^A: The generators of $SU(3)$ on colour triplets, $t^A=\lambda^A_{\rm GM}/2$ (the Gell-Mann matrices halved), $\mathrm{tr}(t^At^B)=\delta^{AB}/2$. Unit: dimensionless. Status: STANDARD.
- $u_{\rm R}$: The right-handed up-type quarks u,c,t (singlets of $SU(2)_L$, hypercharge 2/3). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- $d_{\rm R}$: The right-handed down-type quarks d,s,b (hypercharge $-1/3$). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.

The Yang-Mills equation of the gluons, with the factor of the gluon-kinetic portals and the colour current of the quarks as the source. Status: STANDARD (INVENTED portal factor).

$$\frac{1}{\sqrt{-\hat{h}}\,\mathcal{Z}}\,\partial_\mu\Big[\sqrt{-\hat{h}}\,\mathcal{Z}\Big(\mathcal{K}_B\,B^{\mu\nu}+\sum_X d_X\,\mathcal{D}_X^{\mu\nu}\Big)\Big]=-g_1\,J_Y^\nu,\qquad \mathcal{K}_B=1+\sum_X c_X^B\,\bar\Psi P_X\Psi,$$
$$J_Y^\nu=\sum_\psi Y_\psi\,\bar\psi\,\hat\gamma^\nu\psi+\frac{\mathrm{i}}{2}\big[(D^\nu\Phi)^\dagger\Phi-\Phi^\dagger D^\nu\Phi\big] \tag{FE-18}$$

Symbols in (FE-18).

- $\hat{h}_{\mu\nu}$: The four-dimensional metric of the fold, $\hat{h}_{\mu\nu}=\mathrm{diag}(\mathrm{e}^{2a_4},\mathrm{e}^{2a_4},\mathrm{e}^{2a_4},-1)$ in the coordinates $x_1,\dots,x_4$ (the induced metric at $\sigma=0$ without the extra times); $\hat{h}^{\mu\nu}$ its inverse, $\hat{h}$ its determinant, $\sqrt{-\hat{h}}=\mathrm{e}^{3a_4}$; $\hat{h}^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- $\mathcal{Z}$: The fold factor, $\mathcal{Z}(x_4)=V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3/V_*$: the factor by which the size of the extra times multiplies every term of the fold; $\mathcal{Z}=1$ when the extra times are frozen at their late value. Unit: dimensionless. Value: 1 in the late universe. Status: DERIVED.
- ∂: A partial derivative: ∂_M with respect to the coordinate X^M, ∂_μ with respect to x^μ, ∂_σ with respect to σ, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1,x_2,x_3,x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- $\mathcal{K}_G$, $\mathcal{K}_W$, $\mathcal{K}_B$: The portal factors of the gauge kinetic terms, $\mathcal{K}_G=1+\sum_X c_X^G\bar\Psi P_X\Psi$ (likewise for W and B): the gauge couplings at the fold are divided by $\sqrt{\mathcal{K}}$. Unit: dimensionless. Value: 1 where dirac16complex vanishes. Status: DERIVED.
- B_μ, $B_{\mu\nu}$: The hypercharge gauge field of $U(1)_Y$ on the fold and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- X: The label of a half of dirac16complex: $X\in\{\mathrm{L},\mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- d_X: The dipole-portal coupling of the half X of dirac16complex with the hypercharge field strength. Unit: GeV^{-5}. Status: INVENTED.
- $\mathcal{D}_X^{\mu\nu}$: The dipole (antisymmetric tensor) density of the half X of dirac16complex at the fold, $\mathcal{D}_X^{\mu\nu}=\mathrm{i}\bar\Psi\Sigma^{\mu\nu}P_X\Psi$ (real, DERIVED). Unit: GeV^7. Status: DERIVED.
- g_1: The $U(1)_Y$ gauge coupling (usually g'; not normalised for grand unification). Unit: dimensionless. Value: $g_1(m_Z)=0.3574$. Status: STANDARD.
- J_Y^ν: The hypercharge current of the fermions and of the Higgs doublet. Unit: GeV^3. Status: STANDARD.
- c_X^B: The B-kinetic portal coupling of the half X. Unit: GeV^{-7}. Status: INVENTED.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi=\Psi^\dagger\sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma=0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- P_X: The chiral projector of the half X: $P_X=P_\mathrm{L}$ for $X=\mathrm{L}$, P_R for $X=\mathrm{R}$. Unit: dimensionless. Status: AUTHOR.
- ψ: The label of a Standard-Model fermion field in a sum: ψ runs over $Q,u_\mathrm{R},d_\mathrm{R},\ell,e_\mathrm{R},\nu_\mathrm{R}$. Status: STANDARD.
- Y_f: The hypercharge of a field, the charge of its coupling $g_1Y_fB_\mu$ to the hypercharge field in the covariant derivative (1/6, 2/3, −1/3, −1/2, −1, 0 for $Q,u_\mathrm{R},d_\mathrm{R},\ell,e_\mathrm{R},\nu_\mathrm{R}$, 1/2 for Φ). Unit: dimensionless. Status: STANDARD.
- ψ, $\bar\psi$: A Standard-Model fermion field (a four-dimensional Weyl field of the fold with three generations, written as a four-component spinor of fixed chirality); $\bar\psi=\psi^\dagger\beta$; ψ_f the fields of the kind f. Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- $\hat\gamma^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold, $\hat\gamma^\mu=\hat{\mathbf{e}}_a{}^\mu\hat\gamma^{(a)}$, $\{\hat\gamma^\mu,\hat\gamma^\nu\}=-2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat\gamma_\mu=\hat{h}_{\mu\nu}\hat\gamma^\nu$. Unit: dimensionless. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M=\partial_M+\Omega_M$ (with $D_M\bar\Psi=\partial_M\bar\Psi-\bar\Psi\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu=\hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A=\partial_\mu X^A+g_3f^{ABC}G_\mu^BX^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.

The Maxwell equation of the hypercharge field, with the dipole density of dirac16complex added to the field strength and the hypercharge current of the fermions and the Higgs doublet as the source. Status: STANDARD (INVENTED portal terms).

$$\mathrm{i}\hat{\gamma}^{\mu}\big(D_{\mu}+\mathrm{i}\,\mathcal{V}^{Q}_{\mu}\big)Q=\mathcal{Y}^{u}\tilde{\Phi}\,u_{\mathrm{R}}+\mathcal{Y}^{d}\Phi\,d_{\mathrm{R}},\qquad \mathcal{V}^{f}_{\mu}=\sum_{X}c^{f}_{X}\,j_{X\mu} \tag{FE-19}$$

Symbols in (FE-19).

- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^{\mu}$: The four-dimensional complex 4×4 Dirac matrices of the fold, $\hat{\gamma}^{\mu}=\hat{\mathbf{e}}_{a}{}^{\mu}\hat{\gamma}^{(a)}$, $\{\hat{\gamma}^{\mu},\hat{\gamma}^{\nu}\}=-2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_{\mu}=\hat{h}_{\mu\nu}\hat{\gamma}^{\nu}$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1,x_2,x_3,x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_{μ}: The covariant derivative: on dirac16complex $D_M=\partial_M+\Omega_M$ (with $D_M\bar{\Psi}=\partial_M\bar{\Psi}-\bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_{μ} (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^{\mu}=\hat{h}^{\mu\nu}D_{\nu}$; on an adjoint field $(D_{\mu}X)^{A}=\partial_{\mu}X^{A}+g_3 f^{ABC}G^{B}_{\mu}X^{C}$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^{f}_{\mu}$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^{f}_{\mu}=\sum_X c^{f}_{X}j_{X\mu}$ ($\mathcal{V}^{Q},\mathcal{V}^{u},\ldots,\mathcal{V}^{\nu}$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- Q, $\bar{Q}$: The left-handed quark doublets of the three generations, $Q=(u_{\mathrm{L}},d_{\mathrm{L}})$ (colour triplets, hypercharge 1/6). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- $\mathcal{Y}^{u}$, $\mathcal{Y}^{d}$, $\mathcal{Y}^{e}$, $\mathcal{Y}^{\nu}$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- $\tilde{\Phi}$: The conjugate Higgs doublet, $\tilde{\Phi}=\mathrm{i}\tau^2\Phi^{*}$ (hypercharge $-1/2$). Unit: GeV. Status: STANDARD.
- u_{R}: The right-handed up-type quarks u,c,t (singlets of $SU(2)_L$, hypercharge 2/3). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- d_{R}: The right-handed down-type quarks d,s,b (hypercharge $-1/3$). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- X: The label of a half of dirac16complex: $X\in\{\mathrm{L},\mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c^{f}_{X}: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c^{ℓ}_{X} and c^{ν}_{X} are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.
- j^{μ}_{X}: The convective current of the half X of dirac16complex, $j^{\mu}_{X}=\mathrm{i}(\bar{\Psi}P_X D^{\mu}\Psi-(D^{\mu}\bar{\Psi})P_X\Psi)$ (real and joining the half with itself, DERIVED). Unit: GeV^{8}. Status: DERIVED.
- f: The label of a kind of Standard-Model multiplet: $f\in\{Q,u,d,\ell,e,\nu\}$ (quark doublets, up-quark singlets, down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.

$$\mathrm{i}\hat{\gamma}^{\mu}\big(D_{\mu}+\mathrm{i}\,\mathcal{V}^{u}_{\mu}\big)u_{\mathrm{R}}=(\mathcal{Y}^{u})^{\dagger}\tilde{\Phi}^{\dagger}Q \tag{FE-20}$$

Symbols in (FE-20).

- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^{\mu}$: The four-dimensional complex 4×4 Dirac matrices of the fold, $\hat{\gamma}^{\mu}=\hat{\mathbf{e}}_{a}{}^{\mu}\hat{\gamma}^{(a)}$, $\{\hat{\gamma}^{\mu},\hat{\gamma}^{\nu}\}=-2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_{\mu}=\hat{h}_{\mu\nu}\hat{\gamma}^{\nu}$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1,x_2,x_3,x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_{μ}: The covariant derivative: on dirac16complex $D_M=\partial_M+\Omega_M$ (with $D_M\bar{\Psi}=\partial_M\bar{\Psi}-\bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_{μ} (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^{\mu}=\hat{h}^{\mu\nu}D_{\nu}$; on an adjoint field $(D_{\mu}X)^{A}=\partial_{\mu}X^{A}+g_3 f^{ABC}G^{B}_{\mu}X^{C}$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^{f}_{\mu}$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^{f}_{\mu}=\sum_X c^{f}_{X}j_{X\mu}$ ($\mathcal{V}^{Q},\mathcal{V}^{u},\ldots,\mathcal{V}^{\nu}$ for the six kinds); flavour blind. Unit: GeV. Status:

DERIVED.

- u_{R}: The right-handed up-type quarks u, c, t (singlets of $SU(2)_L$, hypercharge 2/3). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^\nu$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $\tilde{\Phi}$: The conjugate Higgs doublet, $\tilde{\Phi} = \mathrm{i}\tau^2\Phi^*$ (hypercharge $-1/2$). Unit: GeV. Status: STANDARD.
- Q, $\bar{Q}$: The left-handed quark doublets of the three generations, $Q = (u_{\mathrm{L}}, d_{\mathrm{L}})$ (colour triplets, hypercharge 1/6). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.

$$\mathrm{i}\hat{\gamma}^\mu\big(D_\mu + \mathrm{i}\,\mathcal{V}^d_\mu\big)d_{\mathrm{R}} = (\mathcal{Y}^d)^\dagger\Phi^\dagger Q \tag{FE-21}$$

Symbols in (FE-21).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold, $\hat{\gamma}^\mu = \hat{\mathbf{e}}_a{}^\mu\hat{\gamma}^{(a)}$, $\{\hat{\gamma}^\mu, \hat{\gamma}^\nu\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_\mu = \hat{h}_{\mu\nu}\hat{\gamma}^\nu$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC}G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^f_\mu$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^f_\mu = \sum_X c^f_X j_{X\mu}$ ($\mathcal{V}^Q, \mathcal{V}^u, \ldots, \mathcal{V}^\nu$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- d_{R}: The right-handed down-type quarks d, s, b (hypercharge $-1/3$). Unit: $\mathrm{GeV}^{3/2}$. Status: STANDARD.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^\nu$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- Q, $\bar{Q}$: The left-handed quark doublets of the three generations, $Q = (u_{\mathrm{L}}, d_{\mathrm{L}})$ (colour triplets, hypercharge 1/6). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.

$$\mathrm{i}\hat{\gamma}^\mu\big(D_\mu + \mathrm{i}\,\mathcal{V}^\ell_\mu\big)\ell = \mathcal{Y}^e\Phi\, e_{\mathrm{R}} + \mathcal{Y}^\nu\tilde{\Phi}\,\nu_{\mathrm{R}} \tag{FE-22}$$

Symbols in (FE-22).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^\mu$: The four-dimensional complex 4×4 Dirac matrices of the fold, $\hat{\gamma}^\mu = \hat{\mathbf{e}}_a{}^\mu\hat{\gamma}^{(a)}$, $\{\hat{\gamma}^\mu, \hat{\gamma}^\nu\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_\mu = \hat{h}_{\mu\nu}\hat{\gamma}^\nu$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_μ: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_μ (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^\mu = \hat{h}^{\mu\nu}D_\nu$; on an adjoint field $(D_\mu X)^A = \partial_\mu X^A + g_3 f^{ABC}G^B_\mu X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^f_\mu$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^f_\mu = \sum_X c^f_X j_{X\mu}$ ($\mathcal{V}^Q, \mathcal{V}^u, \ldots, \mathcal{V}^\nu$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- ℓ, $\bar{\ell}$: The left-handed lepton doublets of the three generations, $\ell = (\nu_{\mathrm{L}}, e_{\mathrm{L}})$ (hypercharge $-1/2$). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^\nu$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.

- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- e_{R}: The right-handed electron, muon and tau (singlets, hypercharge -1). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- $\tilde{\Phi}$: The conjugate Higgs doublet, $\tilde{\Phi} = \mathrm{i}\tau^2\Phi^*$ (hypercharge $-1/2$). Unit: GeV. Status: STANDARD.
- ν_{R}: The three right-handed neutrinos (gauge singlets). Unit: $\mathrm{GeV}^{3/2}$. Status: ASSUMED.

$$\mathrm{i}\hat{\gamma}^{\mu}\big(D_{\mu} + \mathrm{i}\,\mathcal{V}^{e}_{\mu}\big)e_{\mathrm{R}} = (\mathcal{Y}^{e})^{\dagger}\Phi^{\dagger}\ell \tag{FE-23}$$

Symbols in (FE-23).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^{\mu}$: The four-dimensional complex 4×4 Dirac matrices of the fold, $\hat{\gamma}^{\mu} = \hat{\mathbf{e}}_a{}^{\mu}\hat{\gamma}^{(a)}$, $\{\hat{\gamma}^{\mu}, \hat{\gamma}^{\nu}\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_{\mu} = \hat{h}_{\mu\nu}\hat{\gamma}^{\nu}$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_{μ}: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_{μ} (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^{\mu} = \hat{h}^{\mu\nu}D_{\nu}$; on an adjoint field $(D_{\mu}X)^A = \partial_{\mu}X^A + g_3 f^{ABC}G^B_{\mu}X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^f_{\mu}$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^f_{\mu} = \sum_X c^f_X j_{X\mu}$ ($\mathcal{V}^Q, \mathcal{V}^u, \ldots, \mathcal{V}^{\nu}$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- e_{R}: The right-handed electron, muon and tau (singlets, hypercharge -1). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^{\nu}$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- ℓ, $\bar{\ell}$: The left-handed lepton doublets of the three generations, $\ell = (\nu_{\mathrm{L}}, e_{\mathrm{L}})$ (hypercharge $-1/2$). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.

$$\mathrm{i}\hat{\gamma}^{\mu}\big(D_{\mu} + \mathrm{i}\,\mathcal{V}^{\nu}_{\mu}\big)\nu_{\mathrm{R}} = (\mathcal{Y}^{\nu})^{\dagger}\tilde{\Phi}^{\dagger}\ell \tag{FE-24}$$

Symbols in (FE-24).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- $\hat{\gamma}^{\mu}$: The four-dimensional complex 4×4 Dirac matrices of the fold, $\hat{\gamma}^{\mu} = \hat{\mathbf{e}}_a{}^{\mu}\hat{\gamma}^{(a)}$, $\{\hat{\gamma}^{\mu}, \hat{\gamma}^{\nu}\} = -2\hat{h}^{\mu\nu}\mathbf{1}$; $\hat{\gamma}_{\mu} = \hat{h}_{\mu\nu}\hat{\gamma}^{\nu}$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- D_M, D_{μ}: The covariant derivative: on dirac16complex $D_M = \partial_M + \Omega_M$ (with $D_M\bar{\Psi} = \partial_M\bar{\Psi} - \bar{\Psi}\Omega_M$); on the fields of the fold the gauge and spin covariant derivative D_{μ} (the partial derivative plus the spin connection of the fold and the gauge fields times their couplings and generators); $D^{\mu} = \hat{h}^{\mu\nu}D_{\nu}$; on an adjoint field $(D_{\mu}X)^A = \partial_{\mu}X^A + g_3 f^{ABC}G^B_{\mu}X^C$ (likewise for $SU(2)$ with $g_2\epsilon^{IJK}$). Unit: 1/length. Status: STANDARD.
- $\mathcal{V}^f_{\mu}$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}^f_{\mu} = \sum_X c^f_X j_{X\mu}$ ($\mathcal{V}^Q, \mathcal{V}^u, \ldots, \mathcal{V}^{\nu}$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- ν_{R}: The three right-handed neutrinos (gauge singlets). Unit: $\mathrm{GeV}^{3/2}$. Status: ASSUMED.
- $\mathcal{Y}^u$, $\mathcal{Y}^d$, $\mathcal{Y}^e$, $\mathcal{Y}^{\nu}$: The Yukawa matrices (3×3 complex matrices in generation space) of the up quarks, the down quarks, the charged leptons and the neutrinos. Unit: dimensionless. Status: STANDARD.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $\tilde{\Phi}$: The conjugate Higgs doublet, $\tilde{\Phi} = \mathrm{i}\tau^2\Phi^*$ (hypercharge $-1/2$). Unit: GeV. Status: STANDARD.
- ℓ, $\bar{\ell}$: The left-handed lepton doublets of the three generations, $\ell = (\nu_{\mathrm{L}}, e_{\mathrm{L}})$ (hypercharge $-1/2$). Unit: $\mathrm{GeV}^{3/2}$. Status: AUTHOR.

The Dirac equations of the quark doublets (FE-19), the up-quark and down-quark singlets (FE-20, FE-21), the lepton doublets (FE-22), the charged-lepton singlets (FE-23) and the right-handed neutrinos (FE-24): every fermion sees the gauge fields, the Higgs doublet through its Yukawa matrix and dirac16complex through the flavour-blind vector potential of the current portals; the right-handed neutrinos see no gauge field. Status: STANDARD (INVENTED potentials).

The table states, for every field equation, how the solution satisfies it.

equation	how it is solved	residual	status
bulk gravity FE-1, order 0	the energy-momentum tensor of the bulk that the author's metric requires is computed exactly (FE-29, S-1 to S-3); the author's vacuum needs none	0 (exact polynomials)	DERIVED
bulk gravity FE-1, order ϵ_r	S-5: the difference mode with a rigid medium, the radion mode with a responding one	0 at first order; the profile of the responding medium is not computed	DERIVED, ASSUMED
junction FE-4, FE-5: time and space components	S-8: Friedmann and acceleration equations of general relativity	relative order $\epsilon_\mathrm{r} < 1.344 \times 10^{-13}$ (S-30)	DERIVED
junction FE-4: extra-time component	S-8, S-9: the radion condition	the mismatch $\Delta_\mathrm{r} = 0.793\rho_\mathrm{c}^{(0)}$, carried by the radion mode	DERIVED, ASSUMED
conservation at the fold FE-11	the fold loses exactly the energy that the bulk flux carries away	0 (an identity for every a_4, b_4)	DERIVED
dirac16complex FE-12 to FE-14, Kohn-Sham FE-41	the Kohn-Sham states (S-14 to S-19), lifted to sixteen components	Kohn-Sham equation 2×10^{-4} (central differences), fold matching 4×10^{-9}, self-consistency 2×10^{-7}; Krein charges -1 to 4×10^{-16}	DERIVED
Higgs doublet FE-15	S-10: the ad hoc vacuum as a gauge rotation	the radial residual $\mathcal{R}_\mathrm{rad}$ (table of S-10), zero once $\mathcal{A} = 0$	DERIVED
$SU(2)$ gauge fields FE-17	S-10: the pure-gauge $W_{x_4}^2$	0: the isospin charge density vanishes exactly	DERIVED
gluons FE-16 and hypercharge field FE-18	no mean field strengths (the gluon condensate enters FE-39 as a mean value)	0	DERIVED
quarks and leptons FE-19 to FE-24	no mean fermion fields; the plasma and the neutrinos are treated statistically (FE-48 to FE-62)	0 in the vacuum	STANDARD
neutrino flavours FE-48 to FE-50	the Sun and the supernova envelope (S-20, S-21)	halving the step changes P_ee by less than 10^{-6} (Sun) and 10^{-9} (envelope)	DERIVED
fast instability FE-51	S-22	the dispersion relation holds to 10^{-14} at the computed frequencies	DERIVED
relic density FE-61	S-24	relative tolerance 10^{-8} of the implicit integration	DERIVED
cosmological perturbations FE-54, FE-55	CAMB with the content of UM-3	the accuracy of CAMB (about 10^{-3} in C_l); background and ruler checked independently to 10^{-6}	STANDARD, DERIVED

10 The three neutrino flavours

The author asks for a correct flavour conversion in UM-3, because the flavour conversion of neutrinos shapes the explosion of massive stars (Gogilashvili and Tamborra, "Neutrino Flavor Conversion Shapes the Rate of Failed Core-collapse Supernovae", arXiv:2605.16504). This section solves the flavour equations of UM-3 in the Sun and in core-collapse supernovae.

$$
\begin{aligned}
&\mathrm{i}\big(\partial_{x_4} + \hat{v}\cdot\vec{\nabla}\big)\varrho_{\vec{p}} = \big[\mathsf{H}_{\vec{p}},\ \varrho_{\vec{p}}\big] + \mathrm{i}\,\mathcal{C}_{\vec{p}}, \qquad \mathrm{i}\big(\partial_{x_4} + \hat{v}\cdot\vec{\nabla}\big)\bar{\varrho}_{\vec{p}} = \big[\bar{\mathsf{H}}_{\vec{p}},\ \bar{\varrho}_{\vec{p}}\big] + \mathrm{i}\,\bar{\mathcal{C}}_{\vec{p}},\\
&\mathsf{H}_{\vec{p}} = \frac{v_{\mathrm{loc}}^2}{v^2}\,\frac{U_{\mathrm{PMNS}}\,\mathsf{M}^2\,U_{\mathrm{PMNS}}^\dagger}{2E} + \mathsf{V}_{\mathrm{m}} + \mathsf{V}_{\nu\nu} + V_\Psi \mathbf{1}, \qquad \mathsf{M}^2 = \mathrm{diag}\big(m_1^2,\ m_2^2,\ m_3^2\big),\\
&\bar{\mathsf{H}}_{\vec{p}} = -\frac{v_{\mathrm{loc}}^2}{v^2}\,\frac{U_{\mathrm{PMNS}}\,\mathsf{M}^2\,U_{\mathrm{PMNS}}^\dagger}{2E} + \mathsf{V}_{\mathrm{m}} + \mathsf{V}_{\nu\nu} + V_\Psi \mathbf{1}
\end{aligned}
\tag{FE-48}
$$

Symbols in (FE-48).

- i: The imaginary unit, $\mathrm{i}^2 = -1$. Unit: dimensionless. Status: STANDARD.
- ∂: A partial derivative: ∂_M with respect to the coordinate X^M, ∂_μ with respect to x^μ, ∂_σ with respect to σ, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\hat{v}$, $\hat{v}^\mu$: The direction of motion of a neutrino: the unit vector $\hat{v} = \vec{p}/|\vec{p}|$ ($\hat{v}_{\vec{p}}$, $\hat{v}_{\vec{q}}$ for the momenta $\vec{p}$, $\vec{q}$), and the four-vector $\hat{v}^\mu = (1, \hat{v})$ in the order $(x_4; x_1, x_2, x_3)$; $\mathrm{d}^2\hat{v}$ is the element of solid angle of the directions. Unit: dimensionless. Status: STANDARD.
- $\vec{\nabla}$: The gradient along ordinary space, $\vec{\nabla} = (\partial_{x_1}, \partial_{x_2}, \partial_{x_3})$ (in the local frame of the medium). Unit: 1/length. Status: STANDARD.
- $\varrho_{\vec{p}}$, $\bar{\varrho}_{\vec{p}}$: The 3×3 flavour density matrices of the neutrinos and of the antineutrinos of momentum $\vec{p}$ in the basis e, μ, τ: the diagonal entries are the occupation numbers of the three flavours, the off-diagonal entries the coherence between the flavours; $\bar{\varrho}$ is defined with the flavour indices in the reversed order, so that it obeys an equation of the same form. Unit: dimensionless. Status: STANDARD.
- $\vec{p}$: The momentum of a neutrino (three components along ordinary space, in the rest frame of the medium); $|\vec{p}| = E$ for a neutrino of negligible mass. Unit: GeV. Status: STANDARD.
- $\mathsf{H}_{\vec{p}}$, $\bar{\mathsf{H}}_{\vec{p}}$: The flavour Hamiltonians (3×3 Hermitian matrices) of the neutrinos and of the antineutrinos of momentum $\vec{p}$. Unit: GeV. Status: STANDARD.
- $\mathcal{C}_{\vec{p}}$, $\bar{\mathcal{C}}_{\vec{p}}$: The collision terms ($3 \times 3$ matrices) of the neutrinos and the antineutrinos of momentum $\vec{p}$: the rates of change of $\varrho_{\vec{p}}$ and $\bar{\varrho}_{\vec{p}}$ by emission, absorption, scattering and pair processes with the matter. Unit: GeV. Status: STANDARD.
- v_{loc}: The local value of the electroweak vacuum, $v_{\mathrm{loc}}^2 = 2\langle\Phi^\dagger\Phi\rangle$: it sets the local masses of the quarks and leptons (each mass times v_{loc}/v) and the local Fermi constant; $v_{\mathrm{loc}} = v$ in the Standard-Model vacuum. Unit: GeV. Value: v in the late universe outside dense dirac16complex. Status: DERIVED.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_{\mathrm{F}})^{-1/2}$ with the Fermi constant $G_{\mathrm{F}} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- U_{PMNS}: The lepton mixing matrix (Pontecorvo-Maki-Nakagawa-Sakata) of the three Dirac neutrinos; $U_{\alpha i}$ are its entries. Unit: dimensionless. Value: $\theta_{12} = 33.40°$, $\theta_{23} = 49.14°$, $\theta_{13} = 8.54°$ (NuFIT 5.2, November 2022, normal ordering, with the Super-Kamiokande atmospheric data); the Dirac phase is set to zero (ASSUMED; the fit prefers 197°). Status: STANDARD, ASSUMED (the phase).
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- M^2: The diagonal matrix of the squared masses of the three neutrino mass states, $\mathsf{M}^2 = \mathrm{diag}(m_1^2, m_2^2, m_3^2)$ (M appears only squared). Unit: GeV^2 (of M^2). Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- V_{m}: The matter potential of the neutrinos (a 3×3 matrix in flavour space) from their forward scattering on the electrons, positrons and neutrons of the medium. Unit: GeV. Status: STANDARD.
- $\mathsf{V}_{\nu\nu}$: The neutrino-neutrino potential (a 3×3 matrix in flavour space) from the forward scattering of a neutrino on the other neutrinos and antineutrinos of the medium; it depends on their flavour content and their directions. Unit: GeV. Status: STANDARD.

- V_Ψ: The potential energy that dirac16complex gives every neutrino through the current portal of the lepton doublets, $V_\Psi = \mathcal{V}^\ell_{x_4}$ (the same for the three flavours). Unit: GeV. Status: DERIVED.
- $\mathbf{1}$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space). Unit: dimensionless. Status: STANDARD.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- $m_u, \ldots, m_3$: The masses of the quarks (m_u, m_c, m_t; m_d, m_s, m_b), of the charged leptons (m_e, m_μ, m_τ) and of the three neutrino mass states (m_1, m_2, m_3, Dirac masses). Unit: GeV. Value: measured values; $\Delta m^2_{21} = 7.41\times10^{-5}$ eV^2, $|\Delta m^2_{31}| = 2.511\times10^{-3}$ eV^2 in the computations of UM-3 (the global fit NuFIT 5.2 of the oscillation data, November 2022), 7.49×10^{-5} and 2.513×10^{-3} eV^2 in NuFIT 6.0 (2024); the lightest mass $m_1 = 0$ in the computations, below 4.12 meV by DESI DR2 with the CMB. Status: STANDARD.

$$
\begin{aligned}
&\mathsf{V}_\mathrm{m} = \sqrt{2}\,G_\mathrm{F,loc}\,\mathrm{diag}\big(n_e - n_{\bar e},\ 0,\ 0\big) - \frac{G_\mathrm{F,loc}}{\sqrt{2}}\,n_n\mathbf{1}, \qquad \mathsf{V}_{\nu\nu} = \sqrt{2}\,G_\mathrm{F,loc}\int\frac{\mathrm{d}^3\vec q}{(2\pi)^3}\big(1-\hat v_{\vec p}\cdot\hat v_{\vec q}\big)\big(\varrho_{\vec q} - \bar\varrho_{\vec q}\big),\\
&V_\Psi = \mathcal{V}^\ell_{x_4} = \sum_X c^\ell_X\, j_{X\,x_4}, \qquad G_\mathrm{F,loc} = \frac{1}{\sqrt{2}\,v^2_\mathrm{loc}},\\
&v^2_\mathrm{loc} = 2\langle\Phi^\dagger\Phi\rangle, \qquad v^2_\mathrm{loc} = v^2 + \mathcal{A}^2 \ \text{ (the ad hoc vacuum)},\\
&v^2_\mathrm{loc} = v^2 - \frac{c_\Phi}{\lambda}\,\langle\bar\Psi P_\mathrm{L}\Psi\rangle \ \text{ (the relaxed vacuum with dirac16complex)}
\end{aligned}
\tag{FE-49}
$$

Symbols in (FE-49).

- V_m: The matter potential of the neutrinos (a 3×3 matrix in flavour space) from their forward scattering on the electrons, positrons and neutrons of the medium. Unit: GeV. Status: STANDARD.
- $G_\mathrm{F,loc}$: The local Fermi constant, $G_\mathrm{F,loc} = 1/(\sqrt{2}\,v^2_\mathrm{loc})$ ($= G_\mathrm{F}$ where $v_\mathrm{loc} = v$). Unit: GeV^{-2}. Status: DERIVED.
- diag: A diagonal matrix with the listed entries on its diagonal (in the order of the coordinates $x_0, x_1, \ldots, x_7$ for a tensor of eight dimensions, $x_1, \ldots, x_7$ for a tensor of the fold, $x_1, \ldots, x_4$ for a four-dimensional tensor). Status: STANDARD.
- n_e: The number density of electrons (bound and free; in a supernova all are free, in the recombining universe $x_e n_e$ is the density of the free electrons). Unit: GeV^3 (cm^{-3}). Status: STANDARD.
- $n_{\bar e}$: The number density of positrons. Unit: GeV^3. Status: STANDARD.
- n_n: The number density of neutrons. Unit: GeV^3. Status: STANDARD.
- $\mathbf{1}$: The identity matrix of the size fixed by the context (16×16 on dirac16complex, 4×4 on a four-dimensional spinor, 3×3 in generation or flavour space). Unit: dimensionless. Status: STANDARD.
- $\mathsf{V}_{\nu\nu}$: The neutrino-neutrino potential (a 3×3 matrix in flavour space) from the forward scattering of a neutrino on the other neutrinos and antineutrinos of the medium; it depends on their flavour content and their directions. Unit: GeV. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $\vec q$: The momentum of another neutrino of the medium (the integration variable of the neutrino-neutrino potential); $\mathrm{d}^3\vec q$ its volume element. Unit: GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $\hat v$, $\hat v^\mu$: The direction of motion of a neutrino: the unit vector $\hat v = \vec p/|\vec p|$ ($\hat v_{\vec p}$, $\hat v_{\vec q}$ for the momenta $\vec p$, $\vec q$), and the four-vector $\hat v^\mu = (1, \hat v)$ in the order $(x_4; x_1, x_2, x_3)$; $\mathrm{d}^2\hat v$ is the element of solid angle of the directions. Unit: dimensionless. Status: STANDARD.
- $\vec p$: The momentum of a neutrino (three components along ordinary space, in the rest frame of the medium); $|\vec p| = E$ for a neutrino of negligible mass. Unit: GeV. Status: STANDARD.
- $\varrho_{\vec p}$, $\bar\varrho_{\vec p}$: The 3×3 flavour density matrices of the neutrinos and of the antineutrinos of momentum $\vec p$ in the basis e, μ, τ: the diagonal entries are the occupation numbers of the three flavours, the off-diagonal entries the coherence between the flavours; $\bar\varrho$ is defined with the flavour indices in the reversed order, so that it obeys an equation of the same form. Unit: dimensionless. Status: STANDARD.
- V_Ψ: The potential energy that dirac16complex gives every neutrino through the current portal of the lepton doublets, $V_\Psi = \mathcal{V}^\ell_{x_4}$ (the same for the three flavours). Unit: GeV. Status: DERIVED.

- $\mathcal{V}_\mu^f$: The vector potential that dirac16complex exerts on the Standard-Model fermions of kind f through the current portals, $\mathcal{V}_\mu^f = \sum_X c_X^f j_{X\mu}$ ($\mathcal{V}^Q, \mathcal{V}^u, \ldots, \mathcal{V}^\nu$ for the six kinds); flavour blind. Unit: GeV. Status: DERIVED.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c_X^f: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c_X^ℓ and c_X^ν are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.
- j_X^μ: The convective current of the half X of dirac16complex, $j_X^\mu = \mathrm{i}(\bar\Psi P_X D^\mu \Psi - (D^\mu \bar\Psi) P_X \Psi)$ (real and joining the half with itself, DERIVED). Unit: GeV^8. Status: DERIVED.
- v_loc: The local value of the electroweak vacuum, $v_\mathrm{loc}^2 = 2\langle \Phi^\dagger \Phi \rangle$: it sets the local masses of the quarks and leptons (each mass times v_loc/v) and the local Fermi constant; $v_\mathrm{loc} = v$ in the Standard-Model vacuum. Unit: GeV. Value: v in the late universe outside dense dirac16complex. Status: DERIVED.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $\langle \cdots \rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\, G_\mathrm{F})^{-1/2}$ with the Fermi constant $G_\mathrm{F} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- $\mathcal{A}$: The author's ad hoc upper component of the Higgs vacuum, almightyAdHoc, $\mathcal{A}(x_4) = v_\mathcal{A} \mathrm{e}^{-2a_{40}x_4}$. Unit: GeV. Status: AUTHOR.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- λ: The quartic coupling of the Higgs potential, $\lambda > 0$. Unit: dimensionless. Value: $\lambda = m_h^2/(2v^2) = 0.129$. Status: AUTHOR.
- Ψ, $\bar\Psi$: dirac16complex: the author's 16-component complex spinor field of eight dimensions; $\bar\Psi = \Psi^\dagger \sigma_{16}$ is its Dirac adjoint; in a term of the fold, Ψ is its value at $\sigma = 0$. Unit: $\mathrm{GeV}^{7/2}$. Status: AUTHOR.
- P_L: The author's left-handed chiral projector, $P_\mathrm{L} = (\mathbf{1} - \Gamma_8)/2$. Unit: dimensionless. Status: AUTHOR.

The flavour equations of the neutrinos of momentum $\vec p$ and the potentials (FE-48, FE-49). Two facts make them those of the Standard Model with three Dirac neutrinos wherever $v_\mathrm{loc} = v$: the portal potential V_Ψ is the same for the three flavours and drops out of every commutator, and the Higgs portal moves v_loc by only $\Delta v^2/v^2 = -\mathfrak{g}_{h\chi}\rho_\chi/(\lambda v^3 \omega_\mathrm{b}) \approx -10^{-50}\, \mathfrak{g}_{h\chi}$ at the local dark-matter density $\rho_\chi = 0.3$ GeV/cm^3 for $\omega_\mathrm{b} = 100$ GeV (the value grows as $1/\omega_\mathrm{b}$ for lighter χ; $\mathfrak{g}_{h\chi}$ is the coupling of the Higgs boson to χ of FE-47, negative where the Higgs portal binds, so the gas raises v_loc). Status: DERIVED (`um3_neutrino.py`, N1: $V_\Psi \mathbf{1}$ changes the evolution by at most 7×10^{-13}, a change of v_loc leaves the mixing matrix unchanged to 10^{-15} and scales the masses by v_loc/v). For a single neutrino along its path (no collisions, no neutrino-neutrino potential) FE-48 is a Schrodinger equation $\mathrm{i}\,\mathrm{d}\psi_\nu/\mathrm{d}r = \mathsf{H}\psi_\nu$ for the flavour amplitudes ψ_ν; for antineutrinos the vacuum term has U_PMNS^* and the matter potential the opposite sign (the convention of FE-48, with the flavour indices of $\bar\varrho$ reversed, gives the same probabilities). It is integrated here with the fourth-order commutator-free Magnus method (the exponentials of the Hamiltonian at the two Gauss points of each step, exact for a constant density), and every result is recomputed with half the step.

$$\tan 2\theta_\mathrm{m} = \frac{\sin 2\theta}{\cos 2\theta - 2EV_\mathrm{CC}/\Delta m^2}, \qquad V_\mathrm{CC} = \sqrt{2}\, G_\mathrm{F,loc}\, n_e, \qquad n_{e,\mathrm{res}} = \frac{\Delta m^2 \cos 2\theta}{2\sqrt{2}\, G_\mathrm{F,loc}\, E},$$
$$P_\mathrm{LZ} = \exp\Big(-\frac{\pi}{2} \frac{\Delta m^2 \sin^2 2\theta}{2E \cos 2\theta} \Big| \frac{\mathrm{d} \ln n_e}{\mathrm{d}r} \Big|_\mathrm{res}^{-1} \Big) \qquad \text{(FE-50)}$$

Symbols in (FE-50).

- θ_m: The mixing angle of two neutrino flavours in matter. Unit: dimensionless. Status: STANDARD.
- θ: The vacuum mixing angle of two neutrino flavours (in the two-flavour description of a resonance, one of the

three angles $\theta_{12}, \theta_{13}, \theta_{23}$ of U_{PMNS}). Unit: dimensionless. Status: STANDARD.

- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- V_{CC}: The charged-current matter potential of the electron neutrinos (forward scattering on electrons through W exchange). Unit: GeV. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; in Δm^2 the squared-mass difference of two neutrino mass states ($m_2^2 - m_1^2$ for the two states of a resonance, multiplied by v_{loc}^2/v^2 where this differs from 1). Unit: GeV. Status: STANDARD.
- $G_{\mathrm{F,loc}}$: The local Fermi constant, $G_{\mathrm{F,loc}} = 1/(\sqrt{2}\, v_{\mathrm{loc}}^2)$ ($= G_{\mathrm{F}}$ where $v_{\mathrm{loc}} = v$). Unit: GeV^{-2}. Status: DERIVED.
- n_e: The number density of electrons (bound and free; in a supernova all are free, in the recombining universe $x_e n_e$ is the density of the free electrons). Unit: GeV^3 (cm^{-3}). Status: STANDARD.
- $n_{e,\mathrm{res}}$: The electron density at which the flavour conversion in matter is resonant (the Mikheyev-Smirnov-Wolfenstein resonance). Unit: GeV^3. Status: STANDARD.
- P_{LZ}: The probability that a neutrino jumps between the eigenstates of matter at the resonance (Landau-Zener); $P_{\mathrm{LZ}} = 0$ is an adiabatic resonance. Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- r: The distance from the centre of the star along the path of a neutrino. Unit: length. Status: STANDARD.
- res: The label of a quantity evaluated at the resonance, where $n_e = n_{e,\mathrm{res}}$. Status: STANDARD.

The mixing angle in matter, the resonance density and the Landau-Zener jump probability (FE-50). Status: STANDARD (`um3_neutrino.py`, N2: the formula for θ_{m} agrees with an exact diagonalisation to 10^{-16}, the Landau-Zener formula with an exact integration through a linear profile to 10^{-6}).

10.1 The Sun

$$\mathsf{H}_2 = \frac{1}{2}\begin{pmatrix} -\frac{\Delta m^2_{21}}{2E}\cos 2\theta_{12} + \cos^2\theta_{13}\, V_{\mathrm{CC}} & \frac{\Delta m^2_{21}}{2E}\sin 2\theta_{12} \\ \frac{\Delta m^2_{21}}{2E}\sin 2\theta_{12} & \frac{\Delta m^2_{21}}{2E}\cos 2\theta_{12} - \cos^2\theta_{13}\, V_{\mathrm{CC}} \end{pmatrix}, \qquad n_e(r) = 245\, N_{\mathrm{A}}\, \mathrm{e}^{-10.54\, r/R_\odot}\ \mathrm{cm}^{-3},$$

$$\mathsf{P}_{\mathrm{ee}} = \cos^4\theta_{13}\, \mathsf{P}_2 + \sin^4\theta_{13}, \qquad \mathsf{P}_2 = \frac{1}{2} + \left(\frac{1}{2} - \mathsf{P}_{\mathrm{c}}\right)\cos 2\theta_{12}\cos 2\theta_{\mathrm{m}}(r_{\mathrm{p}}),$$

$$\mathsf{P}_{\mathrm{c}} = \frac{\mathrm{e}^{-2\pi r_\odot \frac{\Delta m^2_{21}}{2E}\sin^2\theta_{12}} - \mathrm{e}^{-2\pi r_\odot \frac{\Delta m^2_{21}}{2E}}}{1 - \mathrm{e}^{-2\pi r_\odot \frac{\Delta m^2_{21}}{2E}}}, \qquad r_\odot = \frac{R_\odot}{10.54} \tag{S-20}$$

Symbols in (S-20).

- H_2: The two-flavour Hamiltonian of the electron neutrinos in the Sun (the mass states 1 and 2 with the potential $\cos^2\theta_{13} V_{\mathrm{CC}}$; the third state decouples). Unit: GeV (or 1/km). Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; in Δm^2 the squared-mass difference of two neutrino mass states ($m_2^2 - m_1^2$ for the two states of a resonance, multiplied by v_{loc}^2/v^2 where this differs from 1). Unit: GeV. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- θ_{12}, θ_{13}, θ_{23}: The three mixing angles of U_{PMNS} (its fourth parameter is the Dirac phase, which does not enter the electron-flavour results of UM-3). Unit: dimensionless. Value: 33.40°, 8.54°, 49.14° in the computations of UM-3 ($\sin^2\theta_{12} = 0.303$, $\sin^2\theta_{13} = 0.02203$, $\sin^2\theta_{23} = 0.572$: NuFIT 5.2, November 2022, normal ordering, with the Super-Kamiokande atmospheric data); 33.71°, 8.56°, 43.28° in NuFIT 6.0 (2024; $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$). Status: STANDARD.

- V_{CC}: The charged-current matter potential of the electron neutrinos (forward scattering on electrons through W exchange). Unit: GeV. Status: STANDARD.
- n_e: The number density of electrons (bound and free; in a supernova all are free, in the recombining universe $x_e n_e$ is the density of the free electrons). Unit: GeV^3 (cm^{-3}). Status: STANDARD.
- r: The distance from the centre of the star along the path of a neutrino. Unit: length. Status: STANDARD.
- N_{A}: Avogadro's number: $245 N_{\mathrm{A}}$ cm^{-3} is 245 moles of electrons per cubic centimetre. Unit: per mole. Value: $6.02214076 \times 10^{23}$. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- $R_\odot$: The radius of the Sun. Unit: length. Value: 6.957×10^5 km. Status: STANDARD.
- P_{ee}: The survival probability of electron neutrinos: the probability that an electron neutrino is detected as an electron neutrino. Unit: dimensionless. Status: DERIVED.
- P_2: The survival probability of the two-flavour problem of the mass states 1 and 2 (S-20). Unit: dimensionless. Status: DERIVED.
- P_{c}: The crossing probability: the probability that a neutrino jumps from one eigenstate of matter to the other on its way through the Sun (exact for an exponential profile, Petcov). Unit: dimensionless. Status: STANDARD.
- θ_{m}: The mixing angle of two neutrino flavours in matter. Unit: dimensionless. Status: STANDARD.
- r_{p}: The radius at which the solar neutrinos of a given reaction are produced. Unit: length. Value: $0.05 R_\odot$ to $0.10 R_\odot$. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $r_\odot$: The scale height of the electron density of the Sun, $r_\odot = R_\odot/10.54$. Unit: length. Value: 6.6×10^4 km. Status: STANDARD.

The solar neutrinos: the third mass state decouples (it oscillates too fast and mixes only through θ_{13}), so the electron neutrinos evolve with the two-flavour Hamiltonian H_2 (with the potential $\cos^2\theta_{13} V_{\mathrm{CC}}$) through the electron density of the Sun in its exponential approximation (Bahcall), from the radius r_{p} of their production to the surface; the phases average out over the production region and the distance to the Earth, so the survival probability P_{ee} follows from the probabilities of the mass states at the surface. The second line is the adiabatic solution with the exact crossing probability P_{c} of an exponential profile (Petcov). Status: STANDARD (the equations), DERIVED (the solution below).

source	E (MeV)	$r_{\mathrm{p}}/R_\odot$	P_{ee} (exact integration)	P_{ee} (S-20)	Borexino
pp	0.267	0.10	0.5454	0.5454	0.57 ± 0.09
^{7}Be	0.862	0.06	0.5126	0.5126	0.53 ± 0.05
pep	1.44	0.10	0.5083	0.5083	0.43 ± 0.11
^{8}B	5.0	0.05	0.3362	0.3362	
^{8}B	8.1	0.05	0.3080	0.3080	0.37 ± 0.08
^{8}B	10.0	0.05	0.3017	0.3017	
^{8}B	15.0	0.05	0.2951	0.2951	

The survival probability of solar electron neutrinos in UM-3 (with $\theta_{12} = 33.40°$, $\theta_{13} = 8.54°$, $\Delta m^2_{21} = 7.41 \times 10^{-5}$ eV^2): the exact integration and the adiabatic formula agree to 10^{-6}, $\mathsf{P}_{\mathrm{c}} < 10^{-10}$ at every energy (the Sun is adiabatic), and halving the step changes nothing at the level 10^{-6}. At low energy P_{ee} approaches the vacuum average $\cos^4\theta_{13}(1 - \frac{1}{2}\sin^2 2\theta_{12}) + \sin^4\theta_{13} = 0.553$, at high energy the full matter conversion $\cos^4\theta_{13}\sin^2\theta_{12} + \sin^4\theta_{13} = 0.290$; the Borexino measurements (Nature 562, 505 (2018), with the high-metallicity solar model) lie within one standard deviation. Status: DERIVED (`um3_flavour.py`, F1).

10.2 The envelope of a supernova

$$\rho_{\mathrm{b}}(r) = 10^4 \ \mathrm{g/cm}^3 \Big(\frac{r}{10^9 \ \mathrm{cm}}\Big)^{-3}, \qquad n_e = \frac{Y_{\mathrm{e}}\,\rho_{\mathrm{b}}}{m_{\mathrm{u}}}, \qquad \gamma_{\mathrm{LZ}} = \frac{\Delta m^2 \sin^2 2\theta}{2E\cos 2\theta}\,\frac{r_{\mathrm{res}}}{3} \tag{S-21}$$

Symbols in (S-21).

- $\rho_{\rm b}$: The (rest-mass) energy density of the baryons: in a supernova the local density of the matter (quoted in g/cm^3), in cosmology the mean density of the baryons ($\rho_{\rm b}^{(0)}$ today). Unit: GeV^4. Status: STANDARD.
- r: The distance from the centre of the star along the path of a neutrino. Unit: length. Status: STANDARD.
- n_e: The number density of electrons (bound and free; in a supernova all are free, in the recombining universe $x_e n_e$ is the density of the free electrons). Unit: GeV^3 (cm^{-3}). Status: STANDARD.
- $Y_{\rm e}$: The electron fraction of stellar matter: the number of electrons per nucleon. Unit: dimensionless. Value: 0.5 in the envelope. Status: STANDARD.
- $m_{\rm u}$: The atomic mass unit, the mass per nucleon of stellar matter to within one per cent: $n_e = Y_e \rho / m_{\rm u}$. Unit: g. Value: 1.66054×10^{-24} g. Status: STANDARD.
- $\gamma_{\rm LZ}$: The adiabaticity parameter of a resonance: the jump probability of FE-50 is $\exp(-\pi\gamma_{\rm LZ}/2)$; $\gamma_{\rm LZ} \gg 1$ is adiabatic. Unit: dimensionless. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; in Δm^2 the squared-mass difference of two neutrino mass states ($m_2^2 - m_1^2$ for the two states of a resonance, multiplied by $v_{\rm loc}^2/v^2$ where this differs from 1). Unit: GeV. Status: STANDARD.
- θ: The vacuum mixing angle of two neutrino flavours (in the two-flavour description of a resonance, one of the three angles $\theta_{12}, \theta_{13}, \theta_{23}$ of $U_{\rm PMNS}$). Unit: dimensionless. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- $r_{\rm res}$: The radius of a resonance: where the electron density equals $n_{e,{\rm res}}$ of FE-50 ($r_{\rm H}$ for Δm^2_{31}, $r_{\rm L}$ for Δm^2_{21}). Unit: length. Status: STANDARD.

The outer layers of the progenitor star (density $\rho_{\rm b} \propto r^{-3}$, electron fraction $Y_{\rm e} = 0.5$, $m_{\rm u}$ the atomic mass unit), through which the neutrinos of a supernova pass after they have left the dense core; $\gamma_{\rm LZ}$ is the adiabaticity of a resonance at the radius $r_{\rm res}$ (the exponent of FE-50 is $-\frac{\pi}{2}\gamma_{\rm LZ}$, since $|{\rm d}\ln n_e/{\rm d}r| = 3/r$). The three-flavour equation with the full mixing matrix ($\theta_{23} = 49.14°$, $\Delta m^2_{31} = 2.511 \times 10^{-3}$ eV^2 in the normal ordering) is integrated from $0.2 r_{\rm H}$ to $5 r_{\rm L}$ at $E = 15$ MeV, starting with an electron neutrino (antineutrino). Status: STANDARD (the profile and the equations), DERIVED (the solution).

ordering	particle	$\mathsf{P}_{\rm ee}$	expected for adiabatic resonances	probabilities of the mass states at the surface
normal	ν_e	0.022030	$\lvert U_{\rm e3}\rvert^2 = 0.022030$	ν_3: $1 - 1.5 \times 10^{-6}$
normal	$\bar\nu_e$	0.681644	$\lvert U_{\rm e1}\rvert^2 = 0.681645$	$\bar\nu_1$: $1 - 1.5 \times 10^{-6}$
inverted	ν_e	0.296325	$\lvert U_{\rm e2}\rvert^2 = 0.296325$	ν_2: $1 - 1.5 \times 10^{-6}$
inverted	$\bar\nu_e$	0.022031	$\lvert U_{\rm e3}\rvert^2 = 0.022030$	$\bar\nu_3$: $1 - 1.6 \times 10^{-6}$

The resonance of Δm^2_{31} lies at $r_{\rm H} = 16\,833$ km in the normal ordering and at 17 002 km in the inverted one (which keeps $|m_3^2 - m_2^2| = 2.511 \times 10^{-3}$ eV^2, so that $m_3^2 - m_1^2 = -(\Delta m^2_{31} - \Delta m^2_{21}) = -2.437 \times 10^{-3}$ eV^2), and that of Δm^2_{21} at $r_{\rm L} = 73\,195$ km; all are adiabatic ($\gamma_{\rm LZ} = 215$, 210 and 655, so $P_{\rm LZ} = 0$). An electron neutrino, produced in the heaviest state of matter, leaves the star as ν_3 in the normal ordering and as ν_2 in the inverted one; an electron antineutrino leaves as $\bar\nu_1$ or $\bar\nu_3$. The exact integration agrees with these limits to 1.1×10^{-6} (and with half the step to 10^{-9}). Shock waves and turbulence in the envelope, which can make the high resonance non-adiabatic, and the collective conversion deeper inside are not included in this static profile. Status: DERIVED (F2).

10.3 The fast instability near the neutrinosphere

$$\mathcal{G}_{\rm ELN}(\hat v) = \sqrt{2}\, G_{\rm F,loc} \int_0^\infty \frac{E^2\,{\rm d}E}{(2\pi)^3}\Big[\mathcal{F}_e - \bar{\mathcal{F}}_e - \big(\mathcal{F}_x - \bar{\mathcal{F}}_x\big)\Big],$$
$$\det\Big[\eta^{\mu\nu}_{\rm M} + \int {\rm d}^2\hat v\; \mathcal{G}_{\rm ELN}(\hat v)\, \frac{\hat v^\mu \hat v^\nu}{\Omega_{\rm fw} - \vec K\cdot\hat v}\Big] = 0, \qquad \hat v^\mu = (1,\ \hat v) \tag{FE-51}$$

Symbols in (FE-51).

- $\mathcal{G}_{\mathrm{ELN}}(\hat{v})$: The angular distribution of the electron lepton number carried by the neutrinos minus that carried by the heavy-lepton neutrinos, per unit solid angle of the direction $\hat{v}$, multiplied by $\sqrt{2}G_{\mathrm{F,loc}}$. Unit: GeV. Status: STANDARD.
- $\hat{v}$, $\hat{v}^{\mu}$: The direction of motion of a neutrino: the unit vector $\hat{v} = \vec{p}/|\vec{p}|$ ($\hat{v}_{\vec{p}}$, $\hat{v}_{\vec{q}}$ for the momenta $\vec{p}$, $\vec{q}$), and the four-vector $\hat{v}^{\mu} = (1, \hat{v})$ in the order $(x_4; x_1, x_2, x_3)$; $\mathrm{d}^2\hat{v}$ is the element of solid angle of the directions. Unit: dimensionless. Status: STANDARD.
- $G_{\mathrm{F,loc}}$: The local Fermi constant, $G_{\mathrm{F,loc}} = 1/(\sqrt{2}\, v_{\mathrm{loc}}^2)$ ($= G_{\mathrm{F}}$ where $v_{\mathrm{loc}} = v$). Unit: GeV^{-2}. Status: DERIVED.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- $\mathcal{F}_e$, $\bar{\mathcal{F}}_e$, $\mathcal{F}_x$, $\bar{\mathcal{F}}_x$: The occupation numbers of the electron neutrinos, of the electron antineutrinos, and of one heavy-lepton neutrino and antineutrino ($x = \mu$ or τ), as functions of the energy E and the direction $\hat{v}$ (the diagonal entries of $\varrho_{\vec{p}}$ and $\bar{\varrho}_{\vec{p}}$); in a transport code, the same for the number of particles in an energy bin. Unit: dimensionless. Status: STANDARD.
- $\eta_{\mathrm{M}}^{\mu\nu}$: The Minkowski metric of the four-vectors $\hat{v}^{\mu}$, $\eta_{\mathrm{M}} = \mathrm{diag}(1, -1, -1, -1)$ in the order $(x_4; x_1, x_2, x_3)$. Unit: dimensionless. Status: STANDARD.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1, x_2, x_3, x_4 of the fold (signature $+, +, +, -$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- Ω_{fw}: The (complex) frequency of a flavour wave of the neutrino gas, measured from the shift caused by the potentials; an imaginary part greater than zero is an instability. Unit: GeV. Status: STANDARD.
- $\vec{K}$: The (real) wave vector of a flavour wave of the neutrino gas, measured from the shift caused by the potentials. Unit: GeV. Status: STANDARD.

$$\begin{aligned}
\Big(\Omega_{\mathrm{fw}} - K_{\mathrm{r}}\,\mathsf{u}\Big)\mathsf{Q}(\mathsf{u}) &= -2\pi \int_{-1}^{1} \mathrm{d}\mathsf{u}' \left(1 - \mathsf{u}\mathsf{u}'\right) \mathcal{G}_{\mathrm{ELN}}(\mathsf{u}')\, \mathsf{Q}(\mathsf{u}') \qquad \text{(axially symmetric mode)}, \\
\Big(\Omega_{\mathrm{fw}} - K_{\mathrm{r}}\,\mathsf{u}\Big)\mathsf{Q}(\mathsf{u}) &= \pi \int_{-1}^{1} \mathrm{d}\mathsf{u}' \sqrt{1 - \mathsf{u}^2}\sqrt{1 - \mathsf{u}'^2}\, \mathcal{G}_{\mathrm{ELN}}(\mathsf{u}')\, \mathsf{Q}(\mathsf{u}') \qquad \text{(first azimuthal mode)}
\end{aligned} \tag{S-22}$$

Symbols in (S-22).

- Ω_{fw}: The (complex) frequency of a flavour wave of the neutrino gas, measured from the shift caused by the potentials; an imaginary part greater than zero is an instability. Unit: GeV. Status: STANDARD.
- K_{r}: The wave number of a flavour wave along the radial direction (the radial component of $\vec{K}$). Unit: GeV (1/length). Status: STANDARD.
- u: The cosine of the angle between the direction of motion of a neutrino and the radial direction (u_e, $\mathsf{u}_{\bar{e}}$ the edges of the emission cones of the electron neutrinos and antineutrinos). Unit: dimensionless. Status: STANDARD.
- $\mathsf{Q}(\mathsf{u})$: The amplitude of a small flavour coherence (the off-diagonal entry of ϱ) of the neutrinos moving with the direction cosine u, in a flavour wave. Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\mathcal{G}_{\mathrm{ELN}}(\hat{v})$: The angular distribution of the electron lepton number carried by the neutrinos minus that carried by the heavy-lepton neutrinos, per unit solid angle of the direction $\hat{v}$, multiplied by $\sqrt{2}G_{\mathrm{F,loc}}$. Unit: GeV. Status: STANDARD.

The fast instability as an eigenvalue problem: for an electron-lepton-number distribution that depends only on the cosine u of the angle to the radial direction, a flavour wave with the wave number K_{r} along the radius and the

amplitude $\mathsf{Q}(\mathsf{u})$ (axially symmetric, or proportional to the cosine of the azimuth) has the frequencies $\Omega_{\rm fw}$ that are the eigenvalues of S-22 (FE-48 linearised; equivalent to the dispersion relation FE-51, which the solutions satisfy to 10^{-14}). The distribution is taken from the neutrinospheres: electron neutrinos emitted uniformly (half-isotropically) from a sphere of radius $R_e = 30$ km and antineutrinos from $R_{\bar e} = 27$ km (they decouple deeper and are more forward peaked), seen at $r = 40$ km, where each fills the cone $\mathsf{u} > \sqrt{1-(R/r)^2}$ ($R = R_e$ or $R_{\bar e}$); α is the ratio of their number densities, the heavy flavours cancel in the electron lepton number, and 200 Gauss-Legendre nodes discretise the angle. Status: STANDARD (the equations), DERIVED (the solution).

$\alpha = n_{\bar\nu_e}/n_{\nu_e}$	0.50	0.75	0.80	0.95	1.20
angular crossing	no	no	yes	yes	yes
growth rate $\mathrm{Im}\,\Omega_{\rm fw}$ $(\sqrt{2}G_{\rm F}n_{\nu_e})$	0	0	0.0137	0.0556	0.0925
growth length, factor e (cm)	–	–	22	5.4	3.2

The electron lepton number crosses zero as a function of the angle exactly when $\alpha > (1-\mathsf{u}_{\bar e})/(1-\mathsf{u}_e) = 0.7744$ (u_e, $\mathsf{u}_{\bar e}$ the edges of the two cones); without a crossing every frequency is real (growth below 10^{-17}), with a crossing the axially symmetric mode grows. For a luminosity $L = 5\times10^{52}$ erg/s and a mean energy $\langle E\rangle = 12$ MeV of the electron neutrinos, emitted half-isotropically from the neutrinosphere at $R_e = 30$ km as in S-22, their density at 40 km is $n_{\nu_e} = L(1-\mathsf{u}_e)/(2\pi R_e^2 c\langle E\rangle) = 5.2\times10^{32}$ cm^{-3} (c the speed of light), so $\sqrt{2}G_{\rm F}n_{\nu_e} = 3.34$ cm^{-1} (with the Fermi constant $G_{\rm F}$ of the vacuum, $v_{\rm loc} = v$ there); the growth length of the table (the length over which the instability grows by the factor e) is the inverse of the growth rate in cm^{-1}, $1/(\mathrm{Im}\,\Omega_{\rm fw}\cdot 3.34\ \mathrm{cm}^{-1})$. The instability grows within centimetres: much faster than anything else in the star, it drives the flavours toward equipartition wherever the crossing exists. Status: DERIVED (F3; N3 of `um3_neutrino.py` for the crossing criterion with general distributions).

10.4 The flavour equipartition and the heating behind the shock

$$\begin{aligned}&\mathcal{F}_e \to \mathcal{F}_e - \delta_{\rm FC}, \qquad \bar{\mathcal{F}}_e \to \bar{\mathcal{F}}_e - \delta_{\rm FC}, \qquad \mathcal{F}_x = \bar{\mathcal{F}}_x \to \mathcal{F}_x + \frac{\delta_{\rm FC}}{2},\\ &\delta_{\rm FC} = \max\Big(\frac{2}{3}\Big(\min\left(\mathcal{F}_e,\ \bar{\mathcal{F}}_e\right) - \mathcal{F}_x\Big),\ \max\left(\mathcal{F}_e,\ \bar{\mathcal{F}}_e\right) - 1\Big),\\ &\text{applied where}\ \ \rho_{\rm b} < \rho_{\rm FC}\ \ \text{and}\ \ x_4 - x_{4,{\rm b}} > t_{\rm FC}\end{aligned} \tag{FE-52}$$

Symbols in (FE-52).

- $\mathcal{F}_e$, $\bar{\mathcal{F}}_e$, $\mathcal{F}_x$, $\bar{\mathcal{F}}_x$: The occupation numbers of the electron neutrinos, of the electron antineutrinos, and of one heavy-lepton neutrino and antineutrino ($x = \mu$ or τ), as functions of the energy E and the direction $\hat v$ (the diagonal entries of $\varrho_{\vec p}$ and $\bar\varrho_{\vec p}$); in a transport code, the same for the number of particles in an energy bin. Unit: dimensionless. Status: STANDARD.
- $\delta_{\rm FC}$: The occupation (or number per energy bin) of electron neutrino-antineutrino pairs converted into heavy-lepton pairs by the flavour equipartition FE-52; a negative value converts heavy-lepton pairs into electron pairs. Unit: dimensionless. Status: ASSUMED.
- $\rho_{\rm b}$: The (rest-mass) energy density of the baryons: in a supernova the local density of the matter (quoted in g/cm^3), in cosmology the mean density of the baryons ($\rho_{\rm b}^{(0)}$ today). Unit: GeV^4. Status: STANDARD.
- $\rho_{\rm FC}$: The baryon density below which the flavour equipartition FE-52 is applied (the critical density $\rho_{\rm c}$ of Gogilashvili and Tamborra). Unit: GeV^4 (g/cm^3). Value: 10^9, 10^{10}, 10^{11} or 10^{13} g/cm^3. Status: STANDARD.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $x_{4,{\rm b}}$: The time of the bounce of the collapsing stellar core. Unit: length (= time). Status: STANDARD.
- $t_{\rm FC}$: The time after the bounce from which the flavour equipartition FE-52 is applied. Unit: time. Value: 0.02 s. Status: STANDARD.

The flavour equipartition of Gogilashvili and Tamborra (FE-52), applied in every energy bin. Status: STANDARD (the constraints), ASSUMED (this form). On 10^5 random occupations in $[0,1]^3$ it conserves the electron lepton

number and the numbers of neutrinos and of antineutrinos to 10^{-15}, keeps every occupation between 0 and 1 (the Pauli bound acts in 22% of the cases), and a second application changes nothing (10^{-16}). Status: DERIVED (F4, N4).

$$\dot{Q}_{\mathrm{heat}} \propto \int_0^\infty \mathrm{d}E\, E^3 \big(\mathcal{F}_e + \bar{\mathcal{F}}_e\big), \qquad \mathcal{F}_e,\ \bar{\mathcal{F}}_e,\ \mathcal{F}_x \propto \frac{L_\mathrm{s}}{E_\mathrm{s}^2}\Big(\frac{E}{E_\mathrm{s}}\Big)^{\alpha_\mathrm{s}} \mathrm{e}^{-(\alpha_\mathrm{s}+1)E/E_\mathrm{s}} \tag{S-23}$$

Symbols in (S-23).

- $\dot{Q}_{\mathrm{heat}}$: The rate at which the absorption of electron neutrinos and antineutrinos heats the matter of the gain region (energy per time and mass). Unit: energy per time and mass. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- E: An energy: E_o the energy of the Kohn-Sham orbital o; in the neutrino sector the energy of a neutrino. Unit: GeV. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- $\mathcal{F}_e,\ \bar{\mathcal{F}}_e,\ \mathcal{F}_x,\ \bar{\mathcal{F}}_x$: The occupation numbers of the electron neutrinos, of the electron antineutrinos, and of one heavy-lepton neutrino and antineutrino ($x = \mu$ or τ), as functions of the energy E and the direction $\hat{v}$ (the diagonal entries of $\varrho_{\vec{p}}$ and $\bar{\varrho}_{\vec{p}}$); in a transport code, the same for the number of particles in an energy bin. Unit: dimensionless. Status: STANDARD.
- L_s: The luminosity (energy per time) of one neutrino species leaving the neutrinosphere. Unit: erg/s. Value: (2 to 5) $\times 10^{52}$ erg/s. Status: STANDARD.
- E_s: The mean energy of one neutrino species. Unit: GeV (MeV). Value: 10 to 16 MeV. Status: STANDARD.
- α_s: The shape (pinching) parameter of a neutrino spectrum (Keil, Raffelt and Janka). Unit: dimensionless. Value: 3. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.

The charged-current heating of the gain region behind the stalled shock ($\nu_e n \to e^- p$ and $\bar{\nu}_e p \to e^+ n$: the cross-section grows as E^2 and the deposited energy is about E), with the number spectra S-23 of the neutrinos (the number per energy of each species, whose integral is its number flux $L_\mathrm{s}/E_\mathrm{s}$ times a common factor: L_s and E_s the luminosity and the mean energy of the species, $\alpha_\mathrm{s} = 3$ the shape parameter). In the table the species are the electron neutrinos ν_e, the electron antineutrinos $\bar{\nu}_e$ and each of the four heavy-lepton species ν_x (ν_μ, ν_τ and their antiparticles); the mean energies after the equipartition are those of the mixed spectra. Status: STANDARD, ESTIMATE (equal weights of the two reactions).

phase	L_s of ν_e, $\bar{\nu}_e$, ν_x (10^{52} erg/s)	E_s (MeV)	E_s after (MeV)	heating after / before
accretion (0.2 s)	5, 5, 2	11, 14, 16	11.3, 16.1, 14.1	0.96
accretion, softer ν_x	5, 5, 2	11, 14, 14	10.9, 15.3, 13.2	0.86
end of accretion	4, 4, 2.5	11, 13, 15	11.9, 14.7, 13.6	1.12
cooling (comparable luminosities)	3, 3, 3	10, 12, 14	11.4, 13.8, 13.0	1.55

Equipartition removes absorbable electron neutrinos and antineutrinos; it hardens the spectrum of the electron antineutrinos in every case and that of the electron neutrinos unless the heavy-lepton spectra are as soft as in the second row. When the heavy-lepton luminosity is well below the electron-flavour one (the accretion phase) the loss of numbers wins and the heating falls; when they are comparable the harder spectra win and it rises. In the transport simulations of Gogilashvili and Tamborra (195 progenitor models; arXiv:2605.16504 version 3) the flavour equipartition, applied wherever the baryon density falls below a critical value, raises the fraction of failed explosions from 25.6% without conversion to 50.8%, 77.4%, 93.3% and 96.4% for the critical densities 10^9, 10^{10}, 10^{11} and 10^{13} g/cm^3: the deeper the equilibration, the stronger the suppression of the revival of the shock, and the progenitors of 16 to 30 solar masses are the most sensitive; in the models that still explode, conversion near the stalled shock speeds up the revival and leaves lighter neutron stars. This estimate includes neither the feedback on the cooling nor the hydrodynamics. Since the flavour physics of UM-3 is that of the Standard Model with three Dirac neutrinos (FE-49), their results apply to UM-3 unchanged. Status: ESTIMATE (F4).

10.5 The newer global fit, the Dirac phase and the lightest mass

The computations above use the global fit NuFIT 5.2 (November 2022) with the Dirac phase $\delta_{\rm CP} = 0$ and the lightest mass $m_1 = 0$. The newer fit NuFIT 6.0 (2024; normal ordering, with the atmospheric data of Super-Kamiokande) has $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{23} = 0.470$, $\sin^2\theta_{13} = 0.02215$, $\delta_{\rm CP} = 212^\circ$, $\Delta m^2_{21} = 7.49\times10^{-5}$ eV^2 and $\Delta m^2_{31} = 2.513\times10^{-3}$ eV^2 (in the inverted ordering $\sin^2\theta_{23} = 0.550$, $\sin^2\theta_{13} = 0.02231$, $\delta_{\rm CP} = 274^\circ$ and $\Delta m^2_{32} = -2.484\times10^{-3}$ eV^2; the normal ordering is preferred by $\Delta\chi^2 = 6.1$).

$$\left(|U_{ei}|^2\right)_{i=1,2,3} = \left(\cos^2\theta_{12}\cos^2\theta_{13},\ \sin^2\theta_{12}\cos^2\theta_{13},\ \sin^2\theta_{13}\right) \tag{S-33}$$

Symbols in (S-33).

- U_{ei}: The electron row of the mixing matrix, $U_{\rm ei} = (U_{\rm PMNS})_{ei}$: $|U_{\rm ei}|^2$ is the electron-flavour fraction of the mass state i. Unit: dimensionless. Value: $|U_{\rm ei}|^2 = 0.681, 0.296, 0.022$. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- θ_{12}, θ_{13}, θ_{23}: The three mixing angles of $U_{\rm PMNS}$ (its fourth parameter is the Dirac phase, which does not enter the electron-flavour results of UM-3). Unit: dimensionless. Value: 33.40°, 8.54°, 49.14° in the computations of UM-3 ($\sin^2\theta_{12} = 0.303$, $\sin^2\theta_{13} = 0.02203$, $\sin^2\theta_{23} = 0.572$: NuFIT 5.2, November 2022, normal ordering, with the Super-Kamiokande atmospheric data); 33.71°, 8.56°, 43.28° in NuFIT 6.0 (2024; $\sin^2\theta_{12} = 0.308$, $\sin^2\theta_{13} = 0.02215$, $\sin^2\theta_{23} = 0.470$). Status: STANDARD.

The Dirac phase drops out of every result of UM-3. The mixing matrix is $U_{\rm PMNS} = R_{23}U_{13}(\delta_{\rm CP})R_{12}$ (rotations by the three angles, the phase attached to the middle one), and its electron row contains only θ_{12} and θ_{13} (S-33; computed for $\delta_{\rm CP} = 0$, 212° and 274°, identical to 7×10^{-18}). Every flavour result of UM-3 is a result for electron neutrinos in matter: the matter potential acts on ν_e only, so the phase can be moved into ν_μ and ν_τ, and the three-flavour evolution through the envelope of a supernova gives the same survival probability for $\delta_{\rm CP} = 0$ and 212° (largest difference 6.0×10^{-14} at 10 and 20 MeV, for neutrinos and antineutrinos). The phase enters only the appearance channels: the vacuum probability of $\nu_\mu \to \nu_e$ at 295 km and 0.6 GeV is 0.0421 for $\delta_{\rm CP} = 0$ and 0.0346 for 212°, and the Jarlskog invariant of CP violation is -0.0178 at 212°. Setting the phase to zero was therefore without consequence for UM-3. Status: DERIVED (`um3_numass.py`, NM3).

$$m_2 = \sqrt{m_1^2 + \Delta m^2_{21}}, \qquad m_3 = \sqrt{m_1^2 + \Delta m^2_{31}}, \qquad \sum_i m_i < 0.064\ \text{eV}\ \Rightarrow\ m_1 < 4.12\ \text{meV}, \qquad m^2_{\rm eff} = \sum_i |U_{ei}|^2 m_i^2 \tag{S-34}$$

Symbols in (S-34).

- $m_u, \ldots, m_3$: The masses of the quarks (m_u, m_c, m_t; m_d, m_s, m_b), of the charged leptons (m_e, m_μ, m_τ) and of the three neutrino mass states (m_1, m_2, m_3, Dirac masses). Unit: GeV. Value: measured values; $\Delta m^2_{21} = 7.41\times10^{-5}$ eV^2, $|\Delta m^2_{31}| = 2.511\times10^{-3}$ eV^2 in the computations of UM-3 (the global fit NuFIT 5.2 of the oscillation data, November 2022), 7.49×10^{-5} and 2.513×10^{-3} eV^2 in NuFIT 6.0 (2024); the lightest mass $m_1 = 0$ in the computations, below 4.12 meV by DESI DR2 with the CMB. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; in Δm^2 the squared-mass difference of two neutrino mass states ($m_2^2 - m_1^2$ for the two states of a resonance, multiplied by $v^2_{\rm loc}/v^2$ where this differs from 1). Unit: GeV. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- $m_{\rm eff}$: The effective mass of the electron neutrino measured in the beta decay of tritium, $m^2_{\rm eff} = \sum_i |U_{\rm ei}|^2 m_i^2$. Unit: eV. Value: below 0.45 eV (KATRIN 2025, 90 per cent). Status: STANDARD.
- U_{ei}: The electron row of the mixing matrix, $U_{\rm ei} = (U_{\rm PMNS})_{ei}$: $|U_{\rm ei}|^2$ is the electron-flavour fraction of the mass state i. Unit: dimensionless. Value: $|U_{\rm ei}|^2 = 0.681, 0.296, 0.022$. Status: STANDARD.

The lightest mass. The smallest sums of the masses are 0.0588 eV in the normal and 0.0989 eV in the inverted ordering. The bound of DESI DR2 with the CMB, $\sum_i m_i < 0.064$ eV (95 per cent, for the cosmological-constant model), excludes the inverted ordering and allows in the normal ordering $m_1 < 4.12$ meV (then $m_2 = 9.58$ and $m_3 = 50.30$ meV). The effective mass of the beta decay of tritium is $m_{\mathrm{eff}} = 8.84$ to 9.76 meV in this range, far below the limit 0.45 eV of KATRIN (2025). Within the range the lightest mass changes one result of UM-3, the capture of the cosmic neutrinos (S-29): on 100 g of tritium the rate falls from 6.88 per year at $m_1 = 0$ to 4.47 per year at $m_1 = 4.12$ meV, because ν_1 becomes slow (mean speed 0.126 today) and loses the doubling of the capture of relativistic neutrinos (4.06 per year if all states were at rest); and the sum of the masses rises from 0.0588 to 0.0640 eV. The record (NuFIT 5.2, $\delta_{\mathrm{CP}} = 0$, $m_1 = 0$) and NuFIT 6.0 give the solar survival probabilities 0.5454, 0.5126, 0.5083, 0.3080 against 0.5418, 0.5100, 0.5059, 0.3126 at the energies of the pp, ^{7}Be, pep and ^{8}B neutrinos, the same envelope probabilities to one per cent, and the CNB rate 6.896 against 6.876 per year: at most 1.5 per cent. Status: DERIVED (NM1, NM2, NM4, NM5), STANDARD (the fits and the bounds).

11 The thermal history

$$\rho_{\mathrm{SM}} = \frac{\pi^2}{30}\, g_\rho(\mathsf{T})\, \mathsf{T}^4, \qquad s = \frac{2\pi^2}{45}\, g_{\mathrm{s}}(\mathsf{T})\, \mathsf{T}^3, \qquad H_4 = \sqrt{\frac{8\pi^3 g_\rho}{90}}\, \frac{\mathsf{T}^2}{M_{\mathrm{Pl}}}, \qquad M_{\mathrm{Pl}} = G_{\mathrm{N}}^{-1/2} \tag{FE-60}$$

Symbols in (FE-60).

- ρ_{SM}: The energy density of the Standard-Model particles. Unit: GeV^4. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- g_ρ: The effective number of relativistic degrees of freedom of the energy density. Unit: dimensionless. Value: 106.75 in the Standard Model above the electroweak scale. Status: STANDARD.
- T: The temperature of the Standard-Model plasma (of the photons after the annihilation of electrons and positrons). Unit: GeV. Status: STANDARD.
- s: The entropy density of the Standard-Model plasma. Unit: GeV^3. Status: STANDARD.
- g_{s}: The effective number of relativistic degrees of freedom of the entropy density. Unit: dimensionless. Value: 3.909 today. Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\, h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- M_{Pl}: The Planck mass, $M_{\mathrm{Pl}} = G_{\mathrm{N}}^{-1/2}$. Unit: GeV. Value: 1.221×10^{19} GeV. Status: STANDARD.
- G_{N}: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_{\mathrm{N}} = 1/M_{\mathrm{Pl}}^2$ with the Planck mass $M_{\mathrm{Pl}} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.

The radiation of the Standard Model in thermal equilibrium and the expansion rate of the radiation era (FE-60). The effective numbers of degrees of freedom are computed here from the exact Bose and Fermi integrals of every particle of the Standard Model with its mass (quarks and gluons above the QCD transition, taken at $\mathsf{T} = 0.155$ GeV, the light hadrons below; the neutrinos decouple at 1 MeV and afterwards have the temperature $(4/11)^{1/3}\mathsf{T}$): $g_\rho = g_{\mathrm{s}} = 106.75$ above the electroweak scale, 97.4 and 96.6 at 50 GeV, 85.6 and 85.4 at 5 GeV, 76.4 and 75.5 at 1 GeV, 10.76 at 10 MeV, and today $g_{\mathrm{s}} = 3.909$, $g_\rho = 3.363$ (the transition region near 0.155 GeV and the annihilation of electrons and positrons are approximated). Status: STANDARD, DERIVED (`um3_thermal.py`, T1).

11.1 The relic density of the particles χ

$$\frac{\mathrm{d}n_\chi}{\mathrm{d}x_4} + 3H_4\, n_\chi = -\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle\big(n_\chi^2 - n_{\chi,\mathrm{eq}}^2\big), \qquad n_{\chi,\mathrm{eq}} = g_\chi\Big(\frac{\omega_{\mathrm{b}}\mathsf{T}}{2\pi}\Big)^{3/2} \mathrm{e}^{-\omega_{\mathrm{b}}/\mathsf{T}}, \qquad \Omega_\chi h_{100}^2 \simeq \frac{1.07 \times 10^9\ \mathrm{GeV}^{-1}\, x_{\mathrm{f}}}{\sqrt{g_*}\, M_{\mathrm{Pl}}\, \langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle} \tag{FE-61}$$

Symbols in (FE-61).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- n_χ: The number density of the dark-matter particles χ (particles plus antiparticles). Unit: GeV^3. Status: DERIVED.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.

- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- σ_{ann}: The annihilation cross-section of the particles χ into Standard-Model particles (through the portals) in the normalisation of FE-61, where n_χ counts the four particle and the four antiparticle states of the bound level: $\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle$ is a quarter of the annihilation rate $(\sigma\mathsf{v})_{\chi\bar{\chi}}$ of one particle χ_a with one antiparticle $\bar{\chi}_a$. Unit: GeV^{-2}. Status: DERIVED.
- v: The relative velocity of two annihilating particles; $\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle$ is the thermal average of the product. Unit: dimensionless. Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- $n_{\chi,\mathrm{eq}}$: The number density of χ in thermal equilibrium with the plasma (non-relativistic). Unit: GeV^3. Status: STANDARD.
- g_χ: The number of internal states of the bound level of χ that are in thermal equilibrium. Unit: dimensionless. Status: DERIVED.
- ω_{b}: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- T: The temperature of the Standard-Model plasma (of the photons after the annihilation of electrons and positrons). Unit: GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- Ω_χ: The density parameter of the dark matter today. Unit: dimensionless. Value: $\Omega_\chi h_{100}^2 = 0.120$ (Planck 2018). Status: DERIVED.
- h_{100}: The present expansion rate of the fold in units of 100 km/s/Mpc. Unit: dimensionless. Value: 0.674 (Planck 2018). Status: STANDARD.
- x_{f}: The ratio of the mass of χ to the temperature at which its annihilations stop (freeze-out). Unit: dimensionless. Value: 23 to 30 (for $\omega_{\mathrm{b}} = 10$ GeV to 10 TeV). Status: STANDARD.
- g_*: The effective number of relativistic degrees of freedom at the freeze-out (g_ρ there). Unit: dimensionless. Status: STANDARD.
- M_{Pl}: The Planck mass, $M_{\mathrm{Pl}} = G_{\mathrm{N}}^{-1/2}$. Unit: GeV. Value: 1.221×10^{19} GeV. Status: STANDARD.

$$\frac{\mathrm{d}\mathsf{Y}}{\mathrm{d}\mathsf{x}} = -\frac{s\,\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle}{\mathsf{x}\,H_4}\Big(1 + \frac{1}{3}\frac{\mathrm{d}\ln g_{\mathrm{s}}}{\mathrm{d}\ln\mathsf{T}}\Big)\big(\mathsf{Y}^2 - \mathsf{Y}_{\mathrm{eq}}^2\big), \qquad \mathsf{Y} = \frac{n_\chi}{s}, \qquad \mathsf{Y}_{\mathrm{eq}} = \frac{n_{\chi,\mathrm{eq}}}{s}, \qquad \mathsf{x} = \frac{\omega_{\mathrm{b}}}{\mathsf{T}},$$
$$n_{\chi,\mathrm{eq}} = \frac{g_\chi\,\omega_{\mathrm{b}}^2\,\mathsf{T}}{2\pi^2}\,K_2\Big(\frac{\omega_{\mathrm{b}}}{\mathsf{T}}\Big), \qquad \Omega_\chi h_{100}^2 = \frac{\omega_{\mathrm{b}}\,\mathsf{Y}(\infty)\,s^{(0)}}{\rho_{\mathrm{c}}^{(0)}/h_{100}^2} \tag{S-24}$$

Symbols in (S-24).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- Y: The yield of the particles χ: their number density divided by the entropy density, $\mathsf{Y} = n_\chi/s$ (constant when they neither annihilate nor are produced). Unit: dimensionless. Status: STANDARD.
- x: The ratio of the mass of χ to the temperature, $\mathsf{x} = \omega_{\mathrm{b}}/\mathsf{T}$ (the time variable of the freeze-out). Unit: dimensionless. Status: STANDARD.
- s: The entropy density of the Standard-Model plasma. Unit: GeV^3. Status: STANDARD.
- σ_{ann}: The annihilation cross-section of the particles χ into Standard-Model particles (through the portals) in the normalisation of FE-61, where n_χ counts the four particle and the four antiparticle states of the bound level: $\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle$ is a quarter of the annihilation rate $(\sigma\mathsf{v})_{\chi\bar{\chi}}$ of one particle χ_a with one antiparticle $\bar{\chi}_a$. Unit: GeV^{-2}. Status: DERIVED.
- v: The relative velocity of two annihilating particles; $\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle$ is the thermal average of the product. Unit: dimensionless. Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its

value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.

- g_{s}: The effective number of relativistic degrees of freedom of the entropy density. Unit: dimensionless. Value: 3.909 today. Status: STANDARD.
- T: The temperature of the Standard-Model plasma (of the photons after the annihilation of electrons and positrons). Unit: GeV. Status: STANDARD.
- Y_{eq}: The yield of χ in thermal equilibrium, $n_{\chi,\mathrm{eq}}/s$. Unit: dimensionless. Status: STANDARD.
- n_χ: The number density of the dark-matter particles χ (particles plus antiparticles). Unit: GeV^3. Status: DERIVED.
- $n_{\chi,\mathrm{eq}}$: The number density of χ in thermal equilibrium with the plasma (non-relativistic). Unit: GeV^3. Status: STANDARD.
- ω_{b}: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- g_χ: The number of internal states of the bound level of χ that are in thermal equilibrium. Unit: dimensionless. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- K_2: The modified Bessel function of the second kind of order 2 ($K_2(y) \approx \sqrt{\pi/(2y)}\,\mathrm{e}^{-y}$ for large y). Unit: dimensionless. Status: STANDARD.
- Ω_χ: The density parameter of the dark matter today. Unit: dimensionless. Value: $\Omega_\chi h_{100}^2 = 0.120$ (Planck 2018). Status: DERIVED.
- h_{100}: The present expansion rate of the fold in units of 100 km/s/Mpc. Unit: dimensionless. Value: 0.674 (Planck 2018). Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- ρ_{c}: The critical density, the energy density of a spatially flat fold at the expansion rate H_4; $\rho_{\mathrm{c}}^{(0)}$ its value today. Unit: GeV^4. Value: $\rho_{\mathrm{c}}^{(0)} = 1.0537 \times 10^{-5}\,h_{100}^2$ GeV/cm^3. Status: STANDARD.

FE-61 rewritten for the yield Y (the number density per entropy, constant without annihilation) as a function of $\mathsf{x} = \omega_{\mathrm{b}}/\mathsf{T}$, with the equilibrium density valid also for relativistic χ (K_2 the modified Bessel function of the second kind), and the relic density from the yield after freeze-out ($s^{(0)} = 2891.2$ cm^{-3} the entropy density today, $\rho_{\mathrm{c}}^{(0)}/h_{100}^2 = 1.0537 \times 10^{-5}$ GeV/cm^3). The bound level of dirac16complex holds four particle states (one multiplet of the rotations of ordinary space and of the extra times: two Dirac fermions χ_a, $a = 1, 2$) and four antiparticle states, so $g_\chi = 8$ and FE-61, written for the total density, has $\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle = (\sigma\mathsf{v})_{\chi\bar{\chi}}/4$, where $(\sigma\mathsf{v})_{\chi\bar{\chi}}$ is the annihilation of one χ_a with one $\bar{\chi}_a$ summed over the final states. The equation is integrated from $\mathsf{x} = 1$ to $\mathsf{x} = 1000$ with an implicit method. Status: STANDARD (the equation), DERIVED (the solution).

ω_{b} (GeV)	10	30	100	300	1000	3000	10000
$\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle$ for $\Omega_\chi h_{100}^2 = 0.120$ (10^{-26} cm^3/s)	2.31	2.19	2.17	2.25	2.28	2.26	2.31
freeze-out x_{f}	23.1	24.0	25.1	26.2	27.4	28.4	29.6
g_* at freeze-out	67.9	78.6	85.2	86.4	93.8	103.9	106.5
estimate of FE-61 ($\Omega_\chi h_{100}^2$)	0.124	0.127	0.128	0.128	0.127	0.126	0.127

The particles χ are a thermal relic with the measured dark-matter density (Planck 2018: $\Omega_\chi h_{100}^2 = 0.1200$) if their annihilation, in the normalisation of FE-61, is $\langle\sigma_{\mathrm{ann}}\mathsf{v}\rangle \approx 2.2 \times 10^{-26}$ cm^3/s for every mass from 10 GeV to 10 TeV (so $(\sigma\mathsf{v})_{\chi\bar{\chi}} \approx 9 \times 10^{-26}$ cm^3/s); the analytic estimate of FE-61 agrees within 7%. Status: DERIVED (T2).

11.2 The portals of χ to the Standard Model

$$\mathfrak{g}_\Phi = \frac{\partial\omega_\mathrm{b}}{\partial\mu_\mathrm{L}}\,\frac{c_\Phi}{V_*}, \qquad \mathfrak{g}_G = \sum_X \frac{\partial\omega_\mathrm{b}}{\partial\mu_X}\,\frac{c_X^G}{4V_*}, \qquad \mathfrak{g}_f = \sum_X \frac{\partial\omega_\mathrm{b}}{\partial\mu_X}\,\frac{2\omega_\mathrm{b}\,c_X^f}{V_*}, \qquad \frac{\partial\omega_\mathrm{b}}{\partial\mu_X} = -\frac{q_\mathrm{b}}{\omega_\mathrm{b}}\,\frac{\partial q_\mathrm{b}}{\partial\mu_X},$$
$$\Delta\omega_\mathrm{b} = \mathfrak{g}_\Phi\,\Delta\langle\Phi^\dagger\Phi\rangle + \mathfrak{g}_G\,\Delta\langle G^2\rangle + \sum_f \mathfrak{g}_f\,\Delta n_f, \qquad \mathfrak{g}_{h\chi} = \mathfrak{g}_\Phi\,v = -\frac{m_\Psi^2\,\mu_\mathrm{L}^2}{2\,\omega_\mathrm{b}\,v} \quad (\mu_\mathrm{R} = 0) \tag{FE-47}$$

Symbols in (FE-47).

- $\mathfrak{g}_\Phi$, $\mathfrak{g}_G$, $\mathfrak{g}_f$, $\mathfrak{g}_{h\chi}$: The four-dimensional couplings of the bound states χ of dirac16complex to the Standard Model (FE-47): to $\Phi^\dagger\Phi$, to the gluon kinetic density, to the number densities of the fermions of kind f, and the Yukawa coupling of the Higgs boson to χ. Unit: $\mathfrak{g}_\Phi$: GeV^{-1}; $\mathfrak{g}_G$: GeV^{-3}; $\mathfrak{g}_f$: GeV^{-2}; $\mathfrak{g}_{h\chi}$: dimensionless. Status: DERIVED.
- ∂: A partial derivative: ∂_M with respect to the coordinate X^M, ∂_μ with respect to x^μ, ∂_σ with respect to σ, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- ω_b: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- μ_X, μ_L, μ_R: The dimensionless strengths of the fold for the left-handed and right-handed halves of dirac16complex, set by the mean fields of the Standard Model at the fold: the jump of the profile across the fold is the exponential of $\gamma^{(0)}(\mu_\mathrm{L}P_\mathrm{L} + \mu_\mathrm{R}P_\mathrm{R})$; negative values attract. Unit: dimensionless. Status: DERIVED.
- c_Φ: The Higgs-portal coupling of the left-handed half of dirac16complex (the right-handed half has none). Unit: GeV^{-5}. Status: INVENTED.
- V_*: The physical volume of the extra times x_5, x_6, x_7 on the fold in the late universe, $V_* = V_\mathrm{t}(\mathrm{e}^{-a_4}b_4)^3$ (with $V_\mathrm{t} = \int \mathrm{d}x_5\,\mathrm{d}x_6\,\mathrm{d}x_7$ the coordinate volume of the extra times) at a time when $\mathrm{e}^{-a_4}b_4$ is constant; it converts the eight-dimensional normalisation into the four-dimensional one. Unit: length^3 (GeV^{-3}). Status: ASSUMED.
- c_X^G: The gluon-kinetic portal coupling of the half X of dirac16complex. Unit: GeV^{-7}. Status: INVENTED.
- X: The label of a half of dirac16complex: $X \in \{\mathrm{L}, \mathrm{R}\}$ (left-handed or right-handed half); a sum over X runs over both. Status: STANDARD.
- c_X^f: The current-portal coupling of the half X of dirac16complex with the fermion-number current of the multiplets of kind f (flavour blind); c_X^ℓ and c_X^ν are those of the lepton doublets and of the right-handed neutrinos. Unit: GeV^{-7}. Status: INVENTED.
- f: The label of a kind of Standard-Model multiplet: $f \in \{Q, u, d, \ell, e, \nu\}$ (quark doublets, up-quark singlets, down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.
- q_b: The inverse width of a state of dirac16complex bound to the fold: its profile falls as $\mathrm{e}^{-q_\mathrm{b}|\sigma|}$. Unit: GeV. Status: DERIVED.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- Φ: The Higgs doublet (two complex components, hypercharge 1/2), a field of the fold. Unit: GeV. Status: AUTHOR.
- †: Hermitian conjugation: the complex conjugate of the transpose of a matrix or spinor (for field operators the adjoint operator). Status: STANDARD.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- $\langle G^2\rangle$: The mean square of the gluon field strength, $\langle G^A_{\mu\nu}G^{A\mu\nu}\rangle$ (the gluon condensate in the QCD vacuum, a thermal value in the plasma). Unit: GeV^4. Status: STANDARD.
- G^A_μ, $G^A_{\mu\nu}$: The gluon field of $SU(3)_C$ on the fold ($A = 1, \ldots, 8$) and its field strength. Unit: GeV (field), GeV^2 (field strength). Status: AUTHOR.
- n_f: The number density of the Standard-Model fermions of kind f (particles minus antiparticles, summed over generations and gauge components), the time component of the mean current $\langle J^\mu_f\rangle$ in the rest frame of the medium. Unit: GeV^3. Status: STANDARD.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_\mathrm{F})^{-1/2}$ with the Fermi constant $G_\mathrm{F} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- m_Ψ: The mass parameter of dirac16complex in the bulk. Unit: GeV. Status: RECORD.

The couplings of the bound states χ to the Standard Model (FE-47): the Higgs-portal Yukawa coupling $\mathfrak{g}_{h\chi}$ is fixed by the binding, the current-portal couplings $\mathfrak{g}_f$ are free (through the INVENTED c_X^f). Status: DERIVED.

$$\begin{aligned}
&\sigma_{\text{pair}}\mathsf{v} = \sum_f \frac{\mathfrak{g}_f^2\,\omega_\text{b}^2}{2\pi}, \qquad \langle\sigma_{\text{ann}}\mathsf{v}\rangle = \frac{\sigma_{\text{pair}}\mathsf{v}}{4}, \qquad \sigma_{\text{SI}} = \frac{\mu_{\text{red}}^2\,f_\text{N}^2}{\pi}, \qquad f_\text{N} = 3\,\mathfrak{g}_\text{q} \;\; \text{(quark portals)},\\
&f_\text{N} = \mathfrak{g}_\text{p} = \frac{\alpha_{\text{em}}}{3\pi}\,\mathfrak{g}_\text{l} \sum_l \ln\frac{\omega_\text{b}^2}{\max\left(m_l^2,\; q_\text{t}^2\right)} \;\; \text{(lepton portals)}, \qquad f_\text{N} = \frac{\mathfrak{g}_{h\chi}\,f_\text{H}\,m_\text{N}}{v\,m_h^2} \;\; \text{(Higgs portal)}
\end{aligned} \tag{S-25}$$

Symbols in (S-25).

- σ_{pair}: The cross-section of the annihilation of one particle χ_a with one antiparticle $\bar\chi_a$ of the same species, summed over the final states; $\sigma_{\text{pair}}\mathsf{v}$ its product with the relative velocity (a constant for s-wave annihilation). Unit: GeV^{-2}. Status: DERIVED.
- v: The relative velocity of two annihilating particles; $\langle\sigma_{\text{ann}}\mathsf{v}\rangle$ is the thermal average of the product. Unit: dimensionless. Status: STANDARD.
- $\mathfrak{g}_\Phi$, $\mathfrak{g}_G$, $\mathfrak{g}_f$, $\mathfrak{g}_{h\chi}$: The four-dimensional couplings of the bound states χ of dirac16complex to the Standard Model (FE-47): to $\Phi^\dagger\Phi$, to the gluon kinetic density, to the number densities of the fermions of kind f, and the Yukawa coupling of the Higgs boson to χ. Unit: $\mathfrak{g}_\Phi$: GeV^{-1}; $\mathfrak{g}_G$: GeV^{-3}; $\mathfrak{g}_f$: GeV^{-2}; $\mathfrak{g}_{h\chi}$: dimensionless. Status: DERIVED.
- f: The label of a kind of Standard-Model multiplet: $f \in \{Q, u, d, \ell, e, \nu\}$ (quark doublets, up-quark singlets, down-quark singlets, lepton doublets, charged-lepton singlets, right-handed neutrinos); a sum over f runs over the six. Status: STANDARD.
- ω_b: The energy of a state of dirac16complex bound to the fold at zero momentum: the four-dimensional mass of the particle χ. Unit: GeV. Status: DERIVED.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- σ_{ann}: The annihilation cross-section of the particles χ into Standard-Model particles (through the portals) in the normalisation of FE-61, where n_χ counts the four particle and the four antiparticle states of the bound level: $\langle\sigma_{\text{ann}}\mathsf{v}\rangle$ is a quarter of the annihilation rate $(\sigma\mathsf{v})_{\chi\bar\chi}$ of one particle χ_a with one antiparticle $\bar\chi_a$. Unit: GeV^{-2}. Status: DERIVED.
- $\langle\cdots\rangle$: The mean (expectation) value of an operator in the state of the medium (the vacuum, a plasma, the Kohn-Sham state). Status: STANDARD.
- σ_{SI}: The spin-independent cross-section of the scattering of χ on a nucleon (normalised per nucleon, as the direct-detection limits are quoted). Unit: GeV^{-2} (cm^2). Status: ESTIMATE.
- μ_{red}: The reduced mass of χ and a nucleon, $\omega_\text{b} m_\text{N}/(\omega_\text{b} + m_\text{N})$. Unit: GeV. Status: STANDARD.
- f_N: The effective coupling of χ to a nucleon in the coherent scattering (a vector or scalar contact interaction). Unit: GeV^{-2}. Status: ESTIMATE.
- $\mathfrak{g}_\text{q}$, $\mathfrak{g}_\text{l}$, $\mathfrak{g}_\text{p}$: The current-portal couplings $\mathfrak{g}_f$ of FE-47 taken equal for all chiral quark states ($\mathfrak{g}_\text{q}$) or for the nine chiral lepton states that take part, the six of the lepton doublets and the three right-handed charged leptons ($\mathfrak{g}_\text{l}$), and the coupling to the proton that the lepton loop induces ($\mathfrak{g}_\text{p}$). Unit: GeV^{-2}. Status: ESTIMATE.
- α_{em}: The fine-structure constant. Unit: dimensionless. Value: 1/137.036. Status: STANDARD.
- l: The multipole index and number ($l = 0, 1, 2, \ldots$) of an expansion in Legendre polynomials of the direction of a photon or a neutrino relative to the wave vector, or of the CMB sky in spherical harmonics. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; in Δm^2 the squared-mass difference of two neutrino mass states ($m_2^2 - m_1^2$ for the two states of a resonance, multiplied by v_{loc}^2/v^2 where this differs from 1). Unit: GeV. Status: STANDARD.
- q_t: The momentum transfer of the scattering of χ on a xenon nucleus in a detector. Unit: GeV. Value: 0.05 GeV. Status: STANDARD.
- m_N: The mass of a nucleon. Unit: GeV. Value: 0.939 GeV. Status: STANDARD.
- v: The vacuum expectation value of the lower component of the Higgs doublet, $v^2 = -\mu^2/\lambda$. Unit: GeV. Value: 246.22 GeV $= (\sqrt{2}\,G_\text{F})^{-1/2}$ with the Fermi constant $G_\text{F} = 1.1663788 \times 10^{-5}$ GeV^{-2} of the muon lifetime (Particle Data Group). Status: STANDARD.
- m_h: The mass of the Higgs boson. Unit: GeV. Value: 125.25 GeV (Particle Data Group 2022). Status: STANDARD.
- f_H: The nucleon matrix element of the Higgs coupling: the fraction of the nucleon mass carried by the quark

masses and, through the trace anomaly, by the gluons. Unit: dimensionless. Value: 0.30. Status: STANDARD.

The annihilation and the scattering on nucleons through the portals, for a vector contact interaction $\mathfrak{g}_f(\bar{\chi}\hat{\gamma}^\mu\chi)(\bar{\psi}_f\hat{\gamma}_\mu\psi_f)$ of χ (its four-dimensional field, with the Dirac matrices $\hat{\gamma}^\mu$ of the fold) with a chiral fermion state f of the Standard Model (its field ψ_f): $\sigma_{\text{pair}}\mathsf{v}$ is the annihilation of one χ_a with one $\bar{\chi}_a$ into massless final states (s-wave, summed over the chiral states f), and FE-61 has a quarter of it because n_χ counts the four particle and the four antiparticle states. The coherent spin-independent cross-section per nucleon σ_{SI} has the reduced mass μ_{red} of χ and the nucleon and the nucleon coupling f_{N}: for quark portals of equal strength $\mathfrak{g}_{\text{q}}$ the vector charge of the nucleon is three times the quark coupling; for lepton portals ($\mathfrak{g}_{\text{l}}$ for the nine chiral lepton states that take part: six of the doublets and the three right-handed charged leptons; the right-handed neutrinos must not take part, as the last part of this section shows) the scattering arises at one loop through the photon, with the coupling $\mathfrak{g}_{\text{p}}$ to the proton (the sum runs over $l = e, \mu, \tau$ with their masses m_l; $\alpha_{\text{em}} = 1/137.036$, the momentum transfer $q_{\text{t}} = 50$ MeV of a xenon detector, the cut-off of the logarithm taken at ω_{b}; per nucleon of xenon σ_{SI} carries the factor $(54/131)^2$); for the Higgs portal the nucleon (mass $m_{\text{N}} = 0.939$ GeV) couples through $f_{\text{H}} = 0.30$, the nucleon matrix element of the quark masses and the gluons. Status: STANDARD (the formulas), ESTIMATE (the loop and the matrix elements).

ω_{b} (GeV)	10	30	100	300	1000	3000	10000
quark portals: σ_{SI} (cm^2)	1.1×10^{-38}	1.3×10^{-39}	1.3×10^{-40}	1.5×10^{-41}	1.3×10^{-42}	1.5×10^{-43}	1.4×10^{-44}
lepton portals: σ_{SI} (cm^2)	2.8×10^{-43}	5.4×10^{-44}	7.7×10^{-45}	1.3×10^{-45}	1.6×10^{-46}	2.2×10^{-47}	2.6×10^{-48}
limit of the LUX-ZEPLIN experiment (cm^2)	not compared	not compared	5.5×10^{-48}	1.7×10^{-47}	5.5×10^{-47}	1.7×10^{-46}	5.5×10^{-46}
Higgs portal: largest $q_{\text{b}}/\omega_{\text{b}}$			0.062	0.047	0.035	0.026	0.019

The direct-detection test of a thermal relic χ against the limit of the LUX-ZEPLIN experiment (4.2 tonne-years of xenon, Phys. Rev. Lett. 135, 011802 (2025), arXiv:2410.17036 version 3: 2.2×10^{-48} cm^2 at its minimum, 40 GeV, at 90% confidence). Above the minimum the limit is scaled in proportion to the mass, which is weaker than the measured curve between the minimum and a few hundred GeV, so the exclusions drawn here are conservative; below the minimum the measured limit rises steeply, and no comparison is made at 10 and 30 GeV. Quark portals that set the relic density are excluded at every mass from 100 GeV to 10 TeV, by factors from 2×10^7 (100 GeV) to 25 (10 TeV): the quark couplings must be smaller than the relic value by the square roots of these factors. Lepton portals give a loop-suppressed scattering that the LUX-ZEPLIN limit excludes below about 1.5 TeV and allows above: a leptophilic χ of a few TeV can be the thermal relic. The Higgs-portal coupling of FE-47, $|\mathfrak{g}_{h\chi}| = 2q_{\text{b}}^2/(\omega_{\text{b}}v)$, must stay below 3.1×10^{-3} (100 GeV) to 3.1×10^{-2} (10 TeV), so the binding must be weak: $q_{\text{b}} \lesssim 0.06\,\omega_{\text{b}}$ (100 GeV) to $0.02\,\omega_{\text{b}}$ (10 TeV). Together with S-13 ($q_{\text{b}} > H$) this requires a bulk scale H well below the mass of χ. Status: ESTIMATE (T3, T4).

11.3 The right-handed neutrinos

$$\frac{\mathrm{d}\rho_{\nu_{\text{R}}}}{\mathrm{d}x_4} + 4H_4\,\rho_{\nu_{\text{R}}} = \mathcal{C}_{\nu_{\text{R}}}, \qquad \Delta N_{\text{eff}} = 3\Big(\frac{10.75}{g_{\text{s}}(\mathsf{T}_{\text{dec}})}\Big)^{4/3} \quad \text{(three right-handed neutrinos decoupling at } \mathsf{T}_{\text{dec}}) \quad \text{(FE-62)}$$

Symbols in (FE-62).

- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- $\rho_{\nu_{\text{R}}}$: The energy density of the three right-handed neutrinos. Unit: GeV^4. Status: STANDARD.

- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- $\mathcal{C}_{\nu_\mathrm{R}}$: The collision term of the right-handed neutrinos: the rate at which collisions produce (or remove) their energy density. Unit: GeV^5. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- N_eff: The effective number of neutrino species: the energy density of the relativistic particles other than photons in units of that of one neutrino species whose temperature is $(4/11)^{1/3}$ times that of the photons. Unit: dimensionless. Value: 3.044 (Standard Model); measured 2.99 ± 0.17 (Planck 2018). Status: STANDARD.
- g_s: The effective number of relativistic degrees of freedom of the entropy density. Unit: dimensionless. Value: 3.909 today. Status: STANDARD.
- T_dec: The temperature at which the right-handed neutrinos decouple from the plasma. Unit: GeV. Status: STANDARD.

$$\frac{n_{\chi,\mathrm{eq}}^2\,\sigma_\mathrm{pair}\mathsf{v}}{16\,n_{\nu_\mathrm{R},\mathrm{eq}}}\Big|_{\mathsf{T}=\mathsf{T}_\mathrm{dec}} = H_4\big(\mathsf{T}_\mathrm{dec}\big), \qquad n_{\nu_\mathrm{R},\mathrm{eq}} = \frac{9\,\zeta(3)}{2\pi^2}\,\mathsf{T}^3 \tag{S-26}$$

Symbols in (S-26).

- $n_{\chi,\mathrm{eq}}$: The number density of χ in thermal equilibrium with the plasma (non-relativistic). Unit: GeV^3. Status: STANDARD.
- σ_pair: The cross-section of the annihilation of one particle χ_a with one antiparticle $\bar{\chi}_a$ of the same species, summed over the final states; $\sigma_\mathrm{pair}\mathsf{v}$ its product with the relative velocity (a constant for s-wave annihilation). Unit: GeV^{-2}. Status: DERIVED.
- v: The relative velocity of two annihilating particles; $\langle\sigma_\mathrm{ann}\mathsf{v}\rangle$ is the thermal average of the product. Unit: dimensionless. Status: STANDARD.
- $n_{\nu_\mathrm{R},\mathrm{eq}}$: The number density of the three right-handed neutrinos and their antiparticles in thermal equilibrium at the temperature T. Unit: GeV^3. Status: STANDARD.
- T: The temperature of the Standard-Model plasma (of the photons after the annihilation of electrons and positrons). Unit: GeV. Status: STANDARD.
- T_dec: The temperature at which the right-handed neutrinos decouple from the plasma. Unit: GeV. Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- ζ: The Riemann zeta function; $\zeta(3) = 1.2020569$. Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

The right-handed neutrinos. Their production by the neutrino Yukawa couplings is negligible: the rate Γ of their production divided by the expansion rate is $\Gamma/H_4 \sim y_\nu^2 M_\mathrm{Pl}/\mathsf{T} = 5 \times 10^{-9}$ at 100 GeV, for the Yukawa coupling $y_\nu = 0.05$ eV$/v$ of the heaviest neutrino (its mass divided by v). If the lepton portal includes them ($c_X^\nu \neq 0$), they stay in equilibrium with χ as long as the rate per right-handed neutrino of $\nu_\mathrm{R}\bar{\nu}_\mathrm{R} \to \chi\bar{\chi}$ (the left-hand side of S-26, by detailed balance: the three right-handed neutrinos take a quarter of $\sigma_\mathrm{pair}\mathsf{v}$, and $n_{\nu_\mathrm{R},\mathrm{eq}}$ counts them with their antiparticles) exceeds the expansion rate, and decouple at T_dec. Status: STANDARD, ESTIMATE (the case of equal couplings of the twelve chiral lepton states, the right-handed neutrinos included).

ω_b (GeV)	10	30	100	300	1000	3000	10000
$\omega_\mathrm{b}/\mathsf{T}_\mathrm{dec}$	13.4	14.0	14.6	15.2	15.8	16.4	17.0
ΔN_eff (FE-62)	0.236	0.199	0.188	0.182	0.155	0.143	0.141
N_eff	3.28	3.24	3.23	3.23	3.20	3.19	3.18

ω_{b} (GeV)	10	30	100	300	1000	3000	10000
standard deviations above ACT DR6 (2.89 ± 0.11)	3.5	3.2	3.1	3.1	2.8	2.7	2.7

Thermalised right-handed neutrinos would add $\Delta N_{\mathrm{eff}} = 0.14$ to 0.24, which is 1.1 to 1.7 standard deviations above Planck 2018 ($N_{\mathrm{eff}} = 2.99 \pm 0.17$) and 2.7 to 3.5 above the combination of ACT DR6 with Planck, lensing, DESI BAO and nucleosynthesis (2.89 ± 0.11; without nucleosynthesis 2.86 ± 0.13). The CMB alone (the refit below) gives $\Delta\chi^2 = +1.6$ for $\Delta N_{\mathrm{eff}} = 0.141$. UM-3 therefore requires that the right-handed neutrinos never thermalise: their portal couplings must be much smaller than those of the lepton doublets, and then $N_{\mathrm{eff}} = 3.044$ as in the Standard Model. Status: DERIVED (T5), ESTIMATE (the rates).

12 The CMB, the CNB and the BAO

The CMB, the CNB and the BAO are computed with the metric of the fold as the power series FE-35: the expanding background (order 0), the cosmological perturbations (order $\epsilon_{\mathrm{c}} \sim 10^{-5}$) and the response of the bulk (order $\epsilon_{\mathrm{r}} < 1.344 \times 10^{-13}$, the radion mode of S-7 to S-9); the terms of order $\epsilon_{\mathrm{c}}\epsilon_{\mathrm{r}}$ and ϵ_{c}^2 are dropped. To order ϵ_{r} the fold obeys general relativity with Newton's constant $8\pi G_{\mathrm{N}} = \kappa_4/(1 + 5\epsilon_{\mathrm{r}}/6)$ (FE-59, above). The content of the fold is that of UM-3: the bound states χ of dirac16complex as cold dark matter (pressureless in the dilute limit, as the Kohn-Sham equation of state shows), the vacuum energy ρ_Λ of the fold as dark energy (with the equation of state $w = -1$, w the ratio of its pressure to its energy density, to the order computed), the baryons and photons of the Standard Model, and three Dirac neutrinos of the normal ordering with the lightest mass zero ($m_1 = 0$, $m_2 = \sqrt{\Delta m_{21}^2} = 8.61$ meV, $m_3 = \sqrt{\Delta m_{31}^2} = 50.1$ meV, $\sum_i m_i = 0.0587$ eV; ASSUMED: the lightest mass is not fixed by any measurement; DESI DR2 with the CMB bounds it below 4.12 meV, S-34) whose right-handed states are not populated ($N_{\mathrm{eff}} = 3.044$). The primordial spectrum, the optical depth and the densities are the Planck 2018 values (TT,TE,EE+lowE+lensing: $\Omega_{\mathrm{b}}h_{100}^2 = 0.02237$, $\Omega_\chi h_{100}^2 = 0.1200$, $H_4^{(0)} = 67.36$ km/s/Mpc, the optical depth $\tau_{\mathrm{reion}} = 0.0544$ of the reionised universe to the CMB photons, $\ln(10^{10}A_{\mathrm{s}}) = 3.044$, $n_{\mathrm{S}} = 0.9649$; ASSUMED: UM-3 does not compute its primordial era to this precision). The particles χ do not drag the baryons: at the direct-detection limits of the table of the portals ($\sigma_{\mathrm{SI}} \le 2.2 \times 10^{-48}$ cm^2 (ω_{b}/40 GeV) for $\omega_{\mathrm{b}} \ge 40$ GeV), a nucleon at recombination ($\mathsf{z} \approx 1100$, thermal speed about 3×10^{-5}) receives momentum from them at the rate $n_\chi \sigma_{\mathrm{SI}} \mathsf{v} \approx 10^{-40}$ s^{-1} for every mass (n_χ falls as $1/\omega_{\mathrm{b}}$ while the limit grows as ω_{b}), 26 orders of magnitude below the expansion rate 5×10^{-14} s^{-1} there; the scattering on electrons through the lepton portals is weaker still. Status: ESTIMATE. The Boltzmann hierarchy and the line-of-sight integrals are solved by the code CAMB (Lewis, Challinor and Lasenby 2000); the background and the BAO ruler are computed a second time here, independently.

$$\begin{aligned} &\Big(\partial_{x_4} - a_4' \, |\vec{p}| \, \frac{\partial}{\partial |\vec{p}|}\Big) \varrho_{\vec{p}} = -\mathrm{i}\big[\mathsf{H}_{\vec{p}}, \ \varrho_{\vec{p}}\big] + \mathcal{C}_{\vec{p}}, \\ &N_{\mathrm{eff}} = \frac{8}{7}\Big(\frac{11}{4}\Big)^{4/3} \frac{\rho_\nu}{\rho_\gamma}, \qquad \mathsf{T}_\nu^{(0)} = \Big(\frac{4}{11}\Big)^{1/3} \mathsf{T}_\gamma^{(0)}, \qquad n_\nu^{(0)} = \frac{3\zeta(3)}{2\pi^2}\big(\mathsf{T}_\nu^{(0)}\big)^3 \end{aligned} \tag{FE-53}$$

Symbols in (FE-53).

- ∂: A partial derivative: ∂_M with respect to the coordinate X^M, ∂_μ with respect to x^μ, ∂_σ with respect to σ, $\partial X/\partial Y$ the derivative of X with respect to Y at fixed other variables. Status: STANDARD.
- x_4: The time ($x_4 \ge 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- $\vec{p}$: The momentum of a neutrino (three components along ordinary space, in the rest frame of the medium); $|\vec{p}| = E$ for a neutrino of negligible mass. Unit: GeV. Status: STANDARD.

- $\varrho_{\vec{p}}$, $\bar{\varrho}_{\vec{p}}$: The 3×3 flavour density matrices of the neutrinos and of the antineutrinos of momentum $\vec{p}$ in the basis e,μ,τ: the diagonal entries are the occupation numbers of the three flavours, the off-diagonal entries the coherence between the flavours; $\bar{\varrho}$ is defined with the flavour indices in the reversed order, so that it obeys an equation of the same form. Unit: dimensionless. Status: STANDARD.
- i: The imaginary unit, $\mathrm{i}^2=-1$. Unit: dimensionless. Status: STANDARD.
- $\mathsf{H}_{\vec{p}}$, $\bar{\mathsf{H}}_{\vec{p}}$: The flavour Hamiltonians (3×3 Hermitian matrices) of the neutrinos and of the antineutrinos of momentum $\vec{p}$. Unit: GeV. Status: STANDARD.
- $\mathcal{C}_{\vec{p}}$, $\bar{\mathcal{C}}_{\vec{p}}$: The collision terms ($3\times 3$ matrices) of the neutrinos and the antineutrinos of momentum $\vec{p}$: the rates of change of $\varrho_{\vec{p}}$ and $\bar{\varrho}_{\vec{p}}$ by emission, absorption, scattering and pair processes with the matter. Unit: GeV. Status: STANDARD.
- N_{eff}: The effective number of neutrino species: the energy density of the relativistic particles other than photons in units of that of one neutrino species whose temperature is $(4/11)^{1/3}$ times that of the photons. Unit: dimensionless. Value: 3.044 (Standard Model); measured 2.99 ± 0.17 (Planck 2018). Status: STANDARD.
- ρ_ν: The energy density of the neutrinos and antineutrinos of all flavours. Unit: GeV^4. Status: STANDARD.
- ρ_γ: The energy density of the photons. Unit: GeV^4. Status: STANDARD.
- T_ν: The temperature of the cosmic neutrino background; $\mathsf{T}_\nu^{(0)}$ its value today. Unit: GeV (or K). Value: 1.945 K today. Status: STANDARD.
- T_γ: The temperature of the photons (of the CMB); $\mathsf{T}_\gamma^{(0)}$ its value today. Unit: GeV (or K). Value: 2.7255 K today. Status: STANDARD.
- n_ν: The number density of the cosmic neutrino background per flavour (neutrinos plus antineutrinos, one helicity each); $n_\nu^{(0)}$ its value today. Unit: GeV^3. Value: 112 cm^{-3} today. Status: STANDARD.
- ζ: The Riemann zeta function; $\zeta(3)=1.2020569$. Unit: dimensionless. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.

$$
\begin{aligned}
&\hat{h}_{\mu\nu}\,\mathrm{d}x^\mu \mathrm{d}x^\nu = a^2\Big[-(1+2\psi_\mathrm{B})\,\mathrm{d}\tau_\mathrm{c}^2+(1-2\phi_\mathrm{B})\big(\mathrm{d}x_1^2+\mathrm{d}x_2^2+\mathrm{d}x_3^2\big)\Big],\\
&a=\mathrm{e}^{a_4},\qquad \mathrm{d}\tau_\mathrm{c}=\frac{\mathrm{d}x_4}{a},\qquad \mathcal{H}=\frac{\dot{a}}{a},\\
&k^2\phi_\mathrm{B}+3\mathcal{H}\big(\dot{\phi}_\mathrm{B}+\mathcal{H}\psi_\mathrm{B}\big)=-4\pi G_\mathrm{N}a^2\,\Delta\rho,\qquad k^2\big(\dot{\phi}_\mathrm{B}+\mathcal{H}\psi_\mathrm{B}\big)=4\pi G_\mathrm{N}a^2(\rho+\mathcal{P})\,\vartheta,\\
&\ddot{\phi}_\mathrm{B}+\mathcal{H}\big(\dot{\psi}_\mathrm{B}+2\dot{\phi}_\mathrm{B}\big)+\Big(2\frac{\ddot{a}}{a}-\mathcal{H}^2\Big)\psi_\mathrm{B}+\frac{k^2}{3}\big(\phi_\mathrm{B}-\psi_\mathrm{B}\big)=4\pi G_\mathrm{N}a^2\,\Delta\mathcal{P},\\
&k^2\big(\phi_\mathrm{B}-\psi_\mathrm{B}\big)=12\pi G_\mathrm{N}a^2(\rho+\mathcal{P})\,\Sigma_\mathrm{s}
\end{aligned}
\qquad \text{(FE-54)}
$$

Symbols in (FE-54).

- $\hat{h}_{\mu\nu}$: The four-dimensional metric of the fold, $\hat{h}_{\mu\nu}=\mathrm{diag}(\mathrm{e}^{2a_4},\mathrm{e}^{2a_4},\mathrm{e}^{2a_4},-1)$ in the coordinates $x_1,\ldots,x_4$ (the induced metric at $\sigma=0$ without the extra times); $\hat{h}^{\mu\nu}$ its inverse, $\hat{h}$ its determinant, $\sqrt{-\hat{h}}=\mathrm{e}^{3a_4}$; $\hat{h}^{(1)}_{\mu\nu}$ its first-order cosmological perturbation. Unit: dimensionless. Status: DERIVED.
- μ: A four-dimensional coordinate index running over the observable coordinates x_1,x_2,x_3,x_4 of the fold (signature $+,+,+,-$); a repeated index is summed. Status: STANDARD.
- ν: A four-dimensional coordinate index (as μ). Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, d^nX the volume element of n variables. Status: STANDARD.
- x: The coordinates of the fold: x_i ($i=1,2,3$) of ordinary space, x_4 the time, x_r ($r=5,6,7$) the extra times; a field of the fold is a function of the four observable coordinates $x=(x_1,x_2,x_3,x_4)$, and $\mathrm{d}^4x=\mathrm{d}x_1\,\mathrm{d}x_2\,\mathrm{d}x_3\,\mathrm{d}x_4$. Unit: length (each coordinate). Status: AUTHOR.
- a, $\dot{a}$, $\ddot{a}$: The scale factor of the fold, $a=\mathrm{e}^{a_4}$ (normalised to 1 today by the choice of the comoving coordinates x_1,x_2,x_3), and its first and second derivatives with respect to the conformal time τ_c. Unit: dimensionless. Status: STANDARD.
- ψ_B, $\dot{\psi}_\mathrm{B}$: The Bardeen potential of the time-time part of the perturbed metric of the fold (the Newtonian potential) in the conformal Newtonian gauge, a function of τ_c and of the wave number k, and its derivative with respect to τ_c. Unit: dimensionless. Status: STANDARD.
- τ_c: The conformal time of the fold, $\mathrm{d}\tau_\mathrm{c}=\mathrm{d}x_4/a$; a dot on a quantity of the cosmological perturbations is $\mathrm{d}/\mathrm{d}\tau_\mathrm{c}$. Unit: length (= time). Status: STANDARD.
- ϕ_B, $\dot{\phi}_\mathrm{B}$, $\ddot{\phi}_\mathrm{B}$: The Bardeen potential of the space-space part of the perturbed metric of the fold (the curvature

potential) in the conformal Newtonian gauge, and its first and second derivatives with respect to τ_c. Unit: dimensionless. Status: STANDARD.

- x_1: The first coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_2: The second coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- x_3: The third coordinate of ordinary 3-space. Unit: length. Status: AUTHOR.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- a_4: The author's metric function of the time x_4 that scales ordinary 3-space: on the fold e^{a_4} is the scale factor of cosmology and a_4' the Hubble rate. Unit: dimensionless. Status: AUTHOR.
- x_4: The time ($x_4 \geq 0$ in the ad hoc vacuum); a prime on a function of x_4 alone is $\mathrm{d}/\mathrm{d}x_4$. Unit: length (= time, $c = 1$). Status: AUTHOR.
- $\mathcal{H}$: The conformal expansion rate, $\mathcal{H} = \dot{a}/a = a\,a_4'$. Unit: 1/length. Status: STANDARD.
- k: The comoving wave number of a Fourier mode of the perturbations, $k = |\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_N: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_\mathrm{N} = 1/M_\mathrm{Pl}^2$ with the Planck mass $M_\mathrm{Pl} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.
- Δ: A change, difference or small variation of the quantity that follows (for example an interval of the time, a perturbation of a density, a difference of two squared masses, or an addition to an effective number of species). Status: STANDARD.
- ρ: The total energy density of the matter of the fold (all species); $\Delta\rho$ its perturbation. Unit: GeV^4. Status: STANDARD.
- $\mathcal{P}$: The total pressure of the matter of the fold; $\Delta\mathcal{P}$ its perturbation. Unit: GeV^4. Status: STANDARD.
- ϑ: The divergence of the velocity of the total matter: $(\rho + \mathcal{P})\vartheta$ is the sum over the species of their energy density plus pressure times the divergence of their velocity. Unit: 1/length. Status: STANDARD.
- Σ_s: The anisotropic stress (shear) of the total matter: $(\rho + \mathcal{P})\Sigma_\mathrm{s}$ is the sum over the species (photons and neutrinos) of their energy density plus pressure times their shear. Unit: dimensionless. Status: STANDARD.

$$
\begin{aligned}
&\dot{\delta}_\chi = -\vartheta_\chi + 3\dot{\phi}_\mathrm{B}, \qquad \dot{\vartheta}_\chi = -\mathcal{H}\vartheta_\chi + k^2\psi_\mathrm{B} + \Gamma_{\chi\mathrm{b}}\big(\vartheta_\mathrm{b} - \vartheta_\chi\big),\\
&\dot{\delta}_\mathrm{b} = -\vartheta_\mathrm{b} + 3\dot{\phi}_\mathrm{B}, \qquad \dot{\vartheta}_\mathrm{b} = -\mathcal{H}\vartheta_\mathrm{b} + c_\mathrm{s}^2 k^2\delta_\mathrm{b} + k^2\psi_\mathrm{B} + \frac{a\,x_\mathrm{e} n_e \sigma_\mathrm{T}}{\mathsf{R}}\big(\vartheta_\gamma - \vartheta_\mathrm{b}\big) + \frac{\rho_\chi}{\rho_\mathrm{b}}\Gamma_{\chi\mathrm{b}}\big(\vartheta_\chi - \vartheta_\mathrm{b}\big), \qquad \mathsf{R} = \frac{3\rho_\mathrm{b}}{4\rho_\gamma},\\
&\dot{\delta}_\gamma = -\frac{4}{3}\vartheta_\gamma + 4\dot{\phi}_\mathrm{B}, \qquad \dot{\vartheta}_\gamma = k^2\Big(\frac{\delta_\gamma}{4} - \Sigma_\gamma\Big) + k^2\psi_\mathrm{B} + a\,x_\mathrm{e} n_e \sigma_\mathrm{T}\big(\vartheta_\mathrm{b} - \vartheta_\gamma\big),\\
&\dot{\mathsf{F}}_l = \frac{k}{2l+1}\big(l\,\mathsf{F}_{l-1} - (l+1)\,\mathsf{F}_{l+1}\big) - a\,x_\mathrm{e} n_e \sigma_\mathrm{T}\Big(\mathsf{F}_l - \frac{\mathsf{G}_0 + \mathsf{F}_2 + \mathsf{G}_2}{10}\,\delta_{l2}\Big) \quad (l \geq 2),\\
&\dot{\mathsf{G}}_l = \frac{k}{2l+1}\big(l\,\mathsf{G}_{l-1} - (l+1)\,\mathsf{G}_{l+1}\big) - a\,x_\mathrm{e} n_e \sigma_\mathrm{T}\Big(\mathsf{G}_l - \frac{\mathsf{G}_0 + \mathsf{F}_2 + \mathsf{G}_2}{2}\Big(\delta_{l0} + \frac{\delta_{l2}}{5}\Big)\Big) \quad (l \geq 0),\\
&\mathsf{F}_0 = \delta_\gamma, \qquad \mathsf{F}_1 = \frac{4\vartheta_\gamma}{3k}, \qquad \mathsf{F}_2 = 2\Sigma_\gamma,\\
&\dot{\delta}_\nu = -\frac{4}{3}\vartheta_\nu + 4\dot{\phi}_\mathrm{B}, \qquad \dot{\vartheta}_\nu = k^2\Big(\frac{\delta_\nu}{4} - \Sigma_\nu\Big) + k^2\psi_\mathrm{B}, \qquad \dot{\mathsf{N}}_l = \frac{k}{2l+1}\big(l\,\mathsf{N}_{l-1} - (l+1)\,\mathsf{N}_{l+1}\big) \quad (l \geq 2),\\
&\mathsf{N}_1 = \frac{4\vartheta_\nu}{3k}, \qquad \mathsf{N}_2 = 2\Sigma_\nu,\\
&\dot{\Upsilon}_0 = -\frac{\mathsf{q}k}{\mathsf{E}}\Upsilon_1 - \dot{\phi}_\mathrm{B}\frac{\mathrm{d}\ln\mathcal{F}_0}{\mathrm{d}\ln\mathsf{q}}, \qquad \dot{\Upsilon}_1 = \frac{\mathsf{q}k}{3\mathsf{E}}\big(\Upsilon_0 - 2\Upsilon_2\big) - \frac{\mathsf{E}k}{3\mathsf{q}}\,\psi_\mathrm{B}\frac{\mathrm{d}\ln\mathcal{F}_0}{\mathrm{d}\ln\mathsf{q}},\\
&\dot{\Upsilon}_l = \frac{\mathsf{q}k}{(2l+1)\mathsf{E}}\big(l\,\Upsilon_{l-1} - (l+1)\,\Upsilon_{l+1}\big) \quad (l \geq 2), \qquad \mathsf{E} = \sqrt{\mathsf{q}^2 + a^2 m_i^2}, \qquad \mathcal{F}_0 = \frac{1}{\mathrm{e}^{\mathsf{q}/\mathsf{T}_\nu^{(0)}} + 1}
\end{aligned}
\tag{FE-55}
$$

Symbols in (FE-55).

- δ_χ, $\dot{\delta}_\chi$: The density contrast of the dark matter (the bound states χ of dirac16complex), the perturbation of its energy density divided by its mean, and its derivative with respect to τ_c. Unit: dimensionless. Status: STANDARD.
- ϑ_χ, $\dot{\vartheta}_\chi$: The divergence of the velocity of the dark matter (for a Fourier mode, ik times its velocity along the

wave vector), and its derivative. Unit: 1/length. Status: STANDARD.

- ϕ_B, $\dot{\phi}_\mathrm{B}$, $\ddot{\phi}_\mathrm{B}$: The Bardeen potential of the space-space part of the perturbed metric of the fold (the curvature potential) in the conformal Newtonian gauge, and its first and second derivatives with respect to τ_c. Unit: dimensionless. Status: STANDARD.
- $\mathcal{H}$: The conformal expansion rate, $\mathcal{H} = \dot{a}/a = a\,a_4'$. Unit: 1/length. Status: STANDARD.
- k: The comoving wave number of a Fourier mode of the perturbations, $k = |\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.
- ψ_B, $\dot{\psi}_\mathrm{B}$: The Bardeen potential of the time-time part of the perturbed metric of the fold (the Newtonian potential) in the conformal Newtonian gauge, a function of τ_c and of the wave number k, and its derivative with respect to τ_c. Unit: dimensionless. Status: STANDARD.
- $\Gamma_{\chi\mathrm{b}}$: The rate (per conformal time) at which the dark matter exchanges momentum with the baryons through the portal couplings of FE-47 (elastic scattering of χ on nuclei and electrons). Unit: 1/length. Value: negligible for the couplings that direct detection allows. Status: DERIVED.
- ϑ_b, $\dot{\vartheta}_\mathrm{b}$: The divergence of the velocity of the baryons and its derivative. Unit: 1/length. Status: STANDARD.
- δ_b, $\dot{\delta}_\mathrm{b}$: The density contrast of the baryons and its derivative. Unit: dimensionless. Status: STANDARD.
- c_s: The sound speed of the baryon gas: its square is the temperature of the gas divided by the mean mass per particle, times one minus a third of the logarithmic derivative of that temperature with respect to a. Unit: dimensionless. Status: STANDARD.
- a, $\dot{a}$, $\ddot{a}$: The scale factor of the fold, $a = \mathrm{e}^{a_4}$ (normalised to 1 today by the choice of the comoving coordinates x_1, x_2, x_3), and its first and second derivatives with respect to the conformal time τ_c. Unit: dimensionless. Status: STANDARD.
- x_e: The ionisation fraction: the fraction of the electrons that are free. Unit: dimensionless. Status: STANDARD.
- n_e: The number density of electrons (bound and free; in a supernova all are free, in the recombining universe $x_\mathrm{e} n_e$ is the density of the free electrons). Unit: GeV^3 (cm^{-3}). Status: STANDARD.
- σ_T: The Thomson cross-section of the scattering of photons on free electrons. Unit: GeV^{-2}. Value: $6.6524587 \times 10^{-25}$ cm^2. Status: STANDARD.
- R: The ratio of the inertia of the baryons to that of the photons, $\mathsf{R} = 3\rho_\mathrm{b}/(4\rho_\gamma)$. Unit: dimensionless. Status: STANDARD.
- ϑ_γ, $\dot{\vartheta}_\gamma$: The divergence of the velocity of the photons and its derivative. Unit: 1/length. Status: STANDARD.
- ρ_χ: The energy density of the dark matter, the bound states χ of dirac16complex; $\rho_\chi^{(0)}$ its value today. Unit: GeV^4. Status: DERIVED.
- ρ_b: The (rest-mass) energy density of the baryons: in a supernova the local density of the matter (quoted in $\mathrm{g/cm}^3$), in cosmology the mean density of the baryons ($\rho_\mathrm{b}^{(0)}$ today). Unit: GeV^4. Status: STANDARD.
- ρ_γ: The energy density of the photons. Unit: GeV^4. Status: STANDARD.
- δ_γ, $\dot{\delta}_\gamma$: The density contrast of the photons and its derivative. Unit: dimensionless. Status: STANDARD.
- Σ_γ: The anisotropic stress (shear) of the photons. Unit: dimensionless. Status: STANDARD.
- F_l, $\dot{\mathsf{F}}_l$: The multipoles of the temperature perturbation of the photons ($\mathsf{F}_0 = \delta_\gamma$, $\mathsf{F}_1 = 4\vartheta_\gamma/(3k)$, $\mathsf{F}_2 = 2\Sigma_\gamma$; the $F_{\gamma l}$ of Ma and Bertschinger) and their derivatives with respect to τ_c. Unit: dimensionless. Status: STANDARD.
- l: The multipole index and number ($l = 0, 1, 2, \ldots$) of an expansion in Legendre polynomials of the direction of a photon or a neutrino relative to the wave vector, or of the CMB sky in spherical harmonics. Status: STANDARD.
- G_l, $\dot{\mathsf{G}}_l$: The multipoles of the polarisation perturbation of the photons (the $G_{\gamma l}$ of Ma and Bertschinger) and their derivatives. Unit: dimensionless. Status: STANDARD.
- δ: The Kronecker delta: with two indices $\delta_N^M = 1$ if $M = N$ and 0 otherwise (likewise $\delta_{X\mathrm{L}}$ is 1 if the label X of a half of dirac16complex is L and 0 otherwise); with upper and lower index strings the generalised Kronecker delta $\delta_{N_1 \cdots N_n}^{M_1 \cdots M_n}$, the determinant of the $n \times n$ matrix of the Kronecker deltas $\delta_{N_b}^{M_a}$ ($+1$ or -1 if the upper indices are an even or odd permutation of the lower ones, 0 otherwise). Status: STANDARD.
- δ_ν, $\dot{\delta}_\nu$: The density contrast of the relativistic neutrinos and its derivative. Unit: dimensionless. Status: STANDARD.
- ϑ_ν, $\dot{\vartheta}_\nu$: The divergence of the velocity of the relativistic neutrinos and its derivative. Unit: 1/length. Status: STANDARD.
- Σ_ν: The anisotropic stress (shear) of the relativistic neutrinos. Unit: dimensionless. Status: STANDARD.
- N_l, $\dot{\mathsf{N}}_l$: The multipoles of the perturbation of the relativistic neutrinos ($\mathsf{N}_0 = \delta_\nu$, $\mathsf{N}_1 = 4\vartheta_\nu/(3k)$, $\mathsf{N}_2 = 2\Sigma_\nu$) and their derivatives. Unit: dimensionless. Status: STANDARD.

- $\Upsilon_l(\mathsf{q})$, $\dot{\Upsilon}_l$: The multipoles of the relative perturbation of the distribution of a massive neutrino state at the comoving momentum q (the Ψ_l of Ma and Bertschinger) and their derivatives. Unit: dimensionless. Status: STANDARD.
- q: The comoving momentum of a neutrino, $a|\vec{p}|$. Unit: GeV. Status: STANDARD.
- E: The comoving energy of a massive neutrino (a times its energy). Unit: GeV. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- $\mathcal{F}_0(\mathsf{q})$: The unperturbed (Fermi-Dirac) occupation number of the cosmic neutrino background at the comoving momentum q. Unit: dimensionless. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; in Δm^2 the squared-mass difference of two neutrino mass states ($m_2^2 - m_1^2$ for the two states of a resonance, multiplied by v_{loc}^2/v^2 where this differs from 1). Unit: GeV. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- T_ν: The temperature of the cosmic neutrino background; $\mathsf{T}_\nu^{(0)}$ its value today. Unit: GeV (or K). Value: 1.945 K today. Status: STANDARD.

$$
\begin{aligned}
r_{\mathrm{s}} &= \int_0^{\tau_{\mathrm{d}}} c_{\gamma\mathrm{b}}\,\mathrm{d}\tau_{\mathrm{c}}, \qquad c_{\gamma\mathrm{b}} = \frac{1}{\sqrt{3(1+\mathsf{R})}}, \\
D_{\mathrm{M}}(\mathsf{z}) &= \int_0^{\mathsf{z}} \frac{\mathrm{d}\mathsf{z}'}{H_4(\mathsf{z}')}, \qquad D_{\mathrm{H}}(\mathsf{z}) = \frac{1}{H_4(\mathsf{z})}, \qquad D_{\mathrm{V}}(\mathsf{z}) = \left(\mathsf{z}\, D_{\mathrm{M}}^2(\mathsf{z})\, D_{\mathrm{H}}(\mathsf{z})\right)^{1/3}
\end{aligned}
\tag{FE-56}
$$

Symbols in (FE-56).

- r_{s}: The comoving sound horizon at the drag epoch: the comoving distance that a sound wave of the photon-baryon fluid travels until the baryons are released from the photons. Unit: length. Value: about 147 Mpc (Planck 2018). Status: STANDARD.
- τ_{d}: The conformal time of the drag epoch, when the baryons are released from the photons (redshift about 1060). Unit: length (= time). Status: STANDARD.
- $c_{\gamma\mathrm{b}}$: The sound speed of the tightly coupled photon-baryon fluid. Unit: dimensionless. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- τ_{c}: The conformal time of the fold, $\mathrm{d}\tau_{\mathrm{c}} = \mathrm{d}x_4/a$; a dot on a quantity of the cosmological perturbations is $\mathrm{d}/\mathrm{d}\tau_{\mathrm{c}}$. Unit: length (= time). Status: STANDARD.
- R: The ratio of the inertia of the baryons to that of the photons, $\mathsf{R} = 3\rho_{\mathrm{b}}/(4\rho_\gamma)$. Unit: dimensionless. Status: STANDARD.
- $D_{\mathrm{M}}(\mathsf{z})$: The transverse comoving distance (comoving angular-diameter distance) to the redshift z, for a spatially flat fold. Unit: length. Status: STANDARD.
- z: The redshift, $1 + \mathsf{z} = 1/a$; z' an integration variable. Unit: dimensionless. Status: STANDARD.
- $'$: The derivative of a function of the time x_4 alone with respect to x_4 ($a_4' = \mathrm{d}a_4/\mathrm{d}x_4$; two primes the second derivative); on a function of σ alone (a profile) the derivative with respect to σ. Status: STANDARD.
- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- $D_{\mathrm{H}}(\mathsf{z})$: The Hubble distance at the redshift z. Unit: length. Status: STANDARD.
- $D_{\mathrm{V}}(\mathsf{z})$: The volume-averaged distance to the redshift z. Unit: length. Status: STANDARD.

$$
\begin{aligned}
C_l^{\mathrm{TT}} &= 4\pi \int \frac{\mathrm{d}k}{k}\, \mathcal{P}_{\mathcal{R}}(k) \left(\mathcal{I}_l^{\mathrm{T}}(k)\right)^2, \qquad C_l^{\mathrm{TE}} = 4\pi \int \frac{\mathrm{d}k}{k}\, \mathcal{P}_{\mathcal{R}}(k)\, \mathcal{I}_l^{\mathrm{T}}(k)\, \mathcal{I}_l^{\mathrm{E}}(k), \\
C_l^{\mathrm{EE}} &= 4\pi \int \frac{\mathrm{d}k}{k}\, \mathcal{P}_{\mathcal{R}}(k) \left(\mathcal{I}_l^{\mathrm{E}}(k)\right)^2, \qquad \mathcal{P}_{\mathcal{R}}(k) = A_{\mathrm{s}} \Big(\frac{k}{k_*}\Big)^{n_{\mathrm{S}}-1}
\end{aligned}
\tag{FE-57}
$$

Symbols in (FE-57).

- C_l^{TT}, C_l^{TE}, C_l^{EE}: The angular power spectra of the CMB at the multipole l: temperature with temperature, temperature with E-mode polarisation, E-mode polarisation with itself. Unit: dimensionless. Status: STANDARD.
- l: The multipole index and number ($l = 0, 1, 2, \ldots$) of an expansion in Legendre polynomials of the direction of a photon or a neutrino relative to the wave vector, or of the CMB sky in spherical harmonics. Status: STANDARD.
- T, E: The labels of the CMB observables: T the temperature anisotropy, E the E-mode (gradient) polarisation; TT, TE, EE label the spectra of their products. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.
- k: The comoving wave number of a Fourier mode of the perturbations, $k = |\vec{k}|$. Unit: GeV (1/length). Status: STANDARD.
- $\mathcal{P}_{\mathcal{R}}(k)$: The primordial power spectrum of the comoving curvature perturbation, per logarithmic interval of k. Unit: dimensionless. Status: ASSUMED.
- $\mathcal{I}_l^{\mathrm{T}}(k)$, $\mathcal{I}_l^{\mathrm{E}}(k)$: The transfer functions of the CMB: the multipole l of the temperature or E-mode anisotropy today produced by a Fourier mode k of unit primordial curvature perturbation, computed from FE-54 and FE-55 (the line-of-sight integral). Unit: dimensionless. Status: STANDARD.
- A_{s}: The amplitude of the primordial curvature spectrum at the pivot k_*. Unit: dimensionless. Value: 2.10×10^{-9} (Planck 2018). Status: ASSUMED.
- k_*: The pivot wave number of the primordial spectrum. Unit: GeV (1/length). Value: 0.05 Mpc^{-1}. Status: STANDARD.
- n_{S}: The spectral index of the primordial curvature spectrum. Unit: dimensionless. Value: 0.965 (Planck 2018). Status: ASSUMED.

$$\Omega_\chi = \frac{\rho_\chi^{(0)}}{\rho_{\mathrm{c}}^{(0)}}, \qquad \Omega_{\mathrm{b}} = \frac{\rho_{\mathrm{b}}^{(0)}}{\rho_{\mathrm{c}}^{(0)}}, \qquad \Omega_\Lambda = \frac{\rho_\Lambda}{\rho_{\mathrm{c}}^{(0)}}, \qquad \Omega_\nu h_{100}^2 = \frac{\sum_i m_i}{93.14\ \mathrm{eV}},$$
$$\rho_{\mathrm{c}}^{(0)} = \frac{3\left(H_4^{(0)}\right)^2}{8\pi G_{\mathrm{N}}}, \qquad H_4^{(0)} = 100\, h_{100}\ \mathrm{km/s/Mpc} \tag{FE-58}$$

Symbols in (FE-58).

- Ω_χ: The density parameter of the dark matter today. Unit: dimensionless. Value: $\Omega_\chi h_{100}^2 = 0.120$ (Planck 2018). Status: DERIVED.
- ρ_χ: The energy density of the dark matter, the bound states χ of dirac16complex; $\rho_\chi^{(0)}$ its value today. Unit: GeV^4. Status: DERIVED.
- ρ_{c}: The critical density, the energy density of a spatially flat fold at the expansion rate H_4; $\rho_{\mathrm{c}}^{(0)}$ its value today. Unit: GeV^4. Value: $\rho_{\mathrm{c}}^{(0)} = 1.0537 \times 10^{-5}\, h_{100}^2$ $\mathrm{GeV/cm}^3$. Status: STANDARD.
- Ω_{b}: The density parameter of the baryons today. Unit: dimensionless. Value: $\Omega_{\mathrm{b}} h_{100}^2 = 0.0224$ (Planck 2018). Status: STANDARD.
- ρ_{b}: The (rest-mass) energy density of the baryons: in a supernova the local density of the matter (quoted in $\mathrm{g/cm}^3$), in cosmology the mean density of the baryons ($\rho_{\mathrm{b}}^{(0)}$ today). Unit: GeV^4. Status: STANDARD.
- Ω_Λ: The density parameter of the dark energy (the vacuum energy of the fold) today. Unit: dimensionless. Value: 0.685 (Planck 2018). Status: DERIVED.
- ρ_Λ: The vacuum energy density of the fold, the dark energy of UM-3: $\rho_\Lambda = \rho_{\mathrm{ff}} - 31HV_*/(4\kappa)$ (FE-59). Unit: GeV^4. Status: DERIVED.
- Ω_ν, $\Omega_\nu^{(i)}$: The density parameter of the neutrinos of the CNB: Ω_ν that of the massive states today; $\Omega_\nu^{(i)}(\mathsf{z})$ the energy density of the mass state i at the redshift z in units of the critical density today. Unit: dimensionless. Status: STANDARD.
- h_{100}: The present expansion rate of the fold in units of 100 km/s/Mpc. Unit: dimensionless. Value: 0.674 (Planck 2018). Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; in Δm^2 the squared-mass difference of two neutrino

mass states ($m_2^2 - m_1^2$ for the two states of a resonance, multiplied by v_{loc}^2/v^2 where this differs from 1). Unit: GeV. Status: STANDARD.

- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- π: The number pi, the ratio of the circumference of a circle to its diameter. Unit: dimensionless. Value: 3.141592654. Status: STANDARD.
- G_{N}: Newton's constant of four-dimensional gravity (the gravity of our world, on the fold), $G_{\mathrm{N}} = 1/M_{\mathrm{Pl}}^2$ with the Planck mass $M_{\mathrm{Pl}} = 1.221 \times 10^{19}$ GeV. Unit: GeV^{-2}. Value: 6.708×10^{-39} GeV^{-2}. Status: STANDARD.

The neutrinos in the expanding universe (FE-53), the scalar perturbations of the metric of the fold in the conformal Newtonian gauge and the Einstein equations of the fold to first order (FE-54), the Boltzmann equations of the species (FE-55), the BAO ruler and distances (FE-56), the angular power spectra of the CMB (FE-57) and the density parameters (FE-58). Status: STANDARD (UM-3: the dark matter is χ, the dark energy is the vacuum energy of the fold).

12.1 The background and the BAO ruler

$$\Big(\frac{H_4(\mathsf{z})}{H_4^{(0)}}\Big)^2 = \Omega_\gamma(1+\mathsf{z})^4 + \sum_{i=1}^{3}\Omega_\nu^{(i)}(\mathsf{z}) + \big(\Omega_{\mathrm{b}}+\Omega_\chi\big)(1+\mathsf{z})^3 + \Omega_\Lambda,$$
$$\Omega_\nu^{(i)}(\mathsf{z}) = \frac{7}{8}\Big(\frac{\mathsf{T}_{\nu,\mathrm{eff}}}{\mathsf{T}_\gamma}\Big)^4\Omega_\gamma\,(1+\mathsf{z})^4\,\frac{\mathsf{I}\big(m_i/\mathsf{T}_{\nu,\mathrm{eff}}\big)}{\mathsf{I}(0)}, \qquad \mathsf{I}(y) = \int_0^\infty \mathrm{d}\mathsf{q}\,\frac{\mathsf{q}^2\sqrt{\mathsf{q}^2+y^2}}{\mathrm{e}^{\mathsf{q}}+1}, \qquad \mathsf{T}_{\nu,\mathrm{eff}} = \Big(\frac{N_{\mathrm{eff}}}{3}\Big)^{1/4}\Big(\frac{4}{11}\Big)^{1/3}\mathsf{T}_\gamma \tag{S-27}$$

Symbols in (S-27).

- H_4: The expansion rate (Hubble rate) of the fold, $H_4 = a_4'$, as a function of the time or of the redshift; $H_4^{(0)}$ its value today, the Hubble constant. Unit: GeV (1/time). Value: $H_4^{(0)} = 100\,h_{100}$ km/s/Mpc with $h_{100} = 0.674$ (Planck 2018): 67.4 km/s/Mpc $= 1.44 \times 10^{-42}$ GeV. Status: STANDARD.
- z: The redshift, $1 + \mathsf{z} = 1/a$; z' an integration variable. Unit: dimensionless. Status: STANDARD.
- Ω_γ: The density parameter of the photons today. Unit: dimensionless. Value: 5.450×10^{-5} (for $h_{100} = 0.6736$). Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- Ω_ν, $\Omega_\nu^{(i)}$: The density parameter of the neutrinos of the CNB: Ω_ν that of the massive states today; $\Omega_\nu^{(i)}(\mathsf{z})$ the energy density of the mass state i at the redshift z in units of the critical density today. Unit: dimensionless. Status: STANDARD.
- Ω_{b}: The density parameter of the baryons today. Unit: dimensionless. Value: $\Omega_{\mathrm{b}}h_{100}^2 = 0.0224$ (Planck 2018). Status: STANDARD.
- Ω_χ: The density parameter of the dark matter today. Unit: dimensionless. Value: $\Omega_\chi h_{100}^2 = 0.120$ (Planck 2018). Status: DERIVED.
- Ω_Λ: The density parameter of the dark energy (the vacuum energy of the fold) today. Unit: dimensionless. Value: 0.685 (Planck 2018). Status: DERIVED.
- $\mathsf{T}_{\nu,\mathrm{eff}}$: The effective temperature of the neutrinos that carries N_{eff} (the convention of CAMB). Unit: GeV (or K). Status: STANDARD.
- T_γ: The temperature of the photons (of the CMB); $\mathsf{T}_\gamma^{(0)}$ its value today. Unit: GeV (or K). Value: 2.7255 K today. Status: STANDARD.
- $\mathsf{I}(y)$: The Fermi-Dirac energy integral of a neutrino state with the mass-to-temperature ratio y; $\mathsf{I}(0) = 7\pi^4/120$. Unit: dimensionless. Status: STANDARD.
- m: A mass: m_i the mass of the neutrino mass state i; in Δm^2 the squared-mass difference of two neutrino mass states ($m_2^2 - m_1^2$ for the two states of a resonance, multiplied by v_{loc}^2/v^2 where this differs from 1). Unit: GeV. Status: STANDARD.
- y: The ratio of the mass of a neutrino state to its temperature, the argument of I. Unit: dimensionless. Status: STANDARD.
- d: The differential: $\mathrm{d}X$ is an infinitesimal change of X, $\mathrm{d}X/\mathrm{d}Y$ the derivative of X with respect to Y, $\mathrm{d}^n X$ the volume element of n variables. Status: STANDARD.

- q: The comoving momentum of a neutrino, $a|\vec{p}|$. Unit: GeV. Status: STANDARD.
- e: Euler's number, the base of the natural logarithm; e^X is the exponential of X. Unit: dimensionless. Value: 2.718281828. Status: STANDARD.
- ∞: Infinity: as a limit of an integral or a sum it means that the range has no bound. Status: STANDARD.
- N_{eff}: The effective number of neutrino species: the energy density of the relativistic particles other than photons in units of that of one neutrino species whose temperature is $(4/11)^{1/3}$ times that of the photons. Unit: dimensionless. Value: 3.044 (Standard Model); measured 2.99 ± 0.17 (Planck 2018). Status: STANDARD.

The expansion rate of the fold as a function of the redshift z (FE-59 with the content of UM-3): photons, the three neutrino states with their exact Fermi-Dirac energy integrals (each at the temperature $\mathsf{T}_{\nu,\mathrm{eff}}$ that carries $N_{\mathrm{eff}} = 3.044$, the convention of CAMB), baryons, χ, and the vacuum energy Ω_Λ that makes the fold spatially flat. Computed here it agrees with CAMB to 8.7×10^{-7} for $0 \le \mathsf{z} \le 10^4$, with $\Omega_\gamma = 5.450 \times 10^{-5}$, $\Omega_\nu = 1.404 \times 10^{-3}$ today (the massless state included), $\Omega_{\mathrm{b}} = 0.0493$, $\Omega_\chi = 0.2645$, $\Omega_\Lambda = 0.6848$; the age of the fold universe is 13.798 Gyr. The terms of order ϵ_{r} change H_4, the distances and the ruler by $5\epsilon_{\mathrm{r}}/12 \lesssim 4 \times 10^{-16}$ at fixed κ_4, and not at all at fixed G_{N}. Status: DERIVED (`um3_cosmology.py`, C1).

The ruler FE-56, integrated here with the same background up to the drag epoch $\mathsf{z}_{\mathrm{d}} = 1059.93$ of CAMB, is $r_{\mathrm{s}} = 147.1047$ Mpc (CAMB: 147.1047 Mpc), and the distances D_{M} and D_{H} at the seven redshifts of DESI agree with CAMB to 8.4×10^{-7} (for example $D_{\mathrm{M}} = 1985.92$ Mpc and $D_{\mathrm{H}} = 3345.07$ Mpc at $\mathsf{z} = 0.51$). Status: DERIVED (C2).

12.2 The CMB

quantity	UM-3	Planck 2018	difference in standard deviations
redshift of last scattering z_*	1089.905	1089.92 ± 0.25	-0.06
sound horizon r_* (Mpc)	144.445	144.43 ± 0.26	$+0.06$
acoustic angle $100\,\theta_*$	1.04108	1.04110 ± 0.00031	-0.06
drag redshift z_{d}	1059.931	1059.94 ± 0.30	-0.03
ruler $r_{\mathrm{s}}(\mathsf{z}_{\mathrm{d}})$ (Mpc)	147.105	147.09 ± 0.26	$+0.06$
damping wave number k_{D} (Mpc^{-1})	0.140865	0.14087 ± 0.00030	-0.02
matter-radiation equality z_{eq}	3402.9	3402 ± 26	$+0.03$
σ_8	0.8120	0.8111 ± 0.0060	$+0.15$
age (Gyr)	13.7977	13.797 ± 0.023	$+0.03$

The derived quantities of the CMB of UM-3 with the Planck 2018 parameters (the column TT,TE,EE+lowE+lensing of Table 2 of Planck 2018 VI) lie within 0.2 standard deviations of the Planck values. The small differences come from evaluating CAMB at the mean values of the Planck parameters (the published derived quantities are means over the posterior): a run with Planck's own neutrinos (one massive state of 0.06 eV) has the same offsets in z_* (1089.907), r_* (144.443 Mpc), z_{d}, r_{s} (147.103 Mpc), k_{D} and z_{eq}; only σ_8 (0.8111 there), $100\theta_*$ (1.04120 there) and the age move with the three neutrino states of UM-3. Here σ_8 is the root mean square of the matter density contrast in spheres of radius $8h_{100}^{-1}$ Mpc today. Status: DERIVED (C3).

$$\chi^2_{\mathrm{fit}} = \left(\mathsf{X}_{\mathrm{model}} - \mathsf{X}_{\mathrm{data}}\right)^T \mathsf{C}^{-1} \left(\mathsf{X}_{\mathrm{model}} - \mathsf{X}_{\mathrm{data}}\right), \qquad \mathsf{X}_b = \sum_{l \text{ in the bin } b} \mathsf{w}_l \, C_l \tag{S-28}$$

Symbols in (S-28).

- χ^2_{fit}: The chi-square of a comparison of a model with data (written χ^2 in the text): the squared deviations weighted with the inverse covariance. Unit: dimensionless. Status: STANDARD.
- $\mathsf{X}_{\mathrm{model}}$, $\mathsf{X}_{\mathrm{data}}$: The vectors of the model values and of the measured values of a data set (binned CMB spectra, BAO distances). Unit: as the data. Status: STANDARD.
- ^{T}: The transpose of a matrix. Status: STANDARD.

- C: The covariance matrix of the measured values; C^{-1} its inverse. Unit: as the data squared. Status: STANDARD.
- X_b: A model value of the binned CMB spectrum, for the bin b of the plik-lite binning. Unit: $\mu\mathrm{K}^2$. Status: STANDARD.
- b: A frame index (as a). Status: STANDARD.
- l: The multipole index and number ($l = 0, 1, 2, \ldots$) of an expansion in Legendre polynomials of the direction of a photon or a neutrino relative to the wave vector, or of the CMB sky in spherical harmonics. Status: STANDARD.
- w_l: The Planck weight of the multipole l in its bin. Unit: dimensionless. Status: STANDARD.
- C_l^{TT}, C_l^{TE}, C_l^{EE}: The angular power spectra of the CMB at the multipole l: temperature with temperature, temperature with E-mode polarisation, E-mode polarisation with itself. Unit: dimensionless. Status: STANDARD.

The comparison with measurements, χ^2_{fit} (the usual χ^2): the measured values $\mathsf{X}_{\mathrm{data}}$ (binned spectra or BAO distances), their covariance matrix C, and the model values $\mathsf{X}_{\mathrm{model}}$; for the Planck 2018 high-l likelihood (plik-lite: foreground-marginalised spectra in 215 bins of TT for $30 \le l \le 2508$ and 199 bins each of TE and EE for $30 \le l \le 1996$, with the Planck calibration fixed at 1) a model bin is the weighted sum X_b of the C_l of the multipoles of the bin with the Planck weights w_l; two Gaussian bins of the low-l temperature spectrum can be added (Prince and Dunkley 2019). The implementation reproduces the reference log-likelihoods of the plik-lite code on its test spectrum to 10^{-13} (-101.5812307 for TT, -293.9558650 for TT, TE, EE with the low-l bins). Status: STANDARD (the likelihood), DERIVED (its implementation, C4).

model	N_{eff}	χ^2 high-l TT, TE, EE (613 bins)	χ^2 TT (215 bins)	refitted χ^2 (615 bins)	$H_4^{(0)}$ after the refit
UM-3	3.044	583.8	206.2	587.9	67.31
UM-3 with thermal ν_{R}, same θ_*	3.185	604.3	215.1	589.5	68.23
Planck's convention (one state of 0.06 eV)	3.044	583.9	206.1		

The CMB spectra of UM-3 against the Planck 2018 data: $\chi^2 = 583.8$ for 613 bins (expected 613 ± 35). Three thermal right-handed neutrinos at the same acoustic angle and the same other parameters raise χ^2 by 20.5 (the damping tail); when $\Omega_{\mathrm{b}}h^2_{100}$, $\Omega_\chi h^2_{100}$, θ_*, A_{s} and n_{S} are refitted (Gauss-Newton with the Jacobian of CAMB, τ_{reion} fixed) to the 613 high-l bins and the two low-l bins, the difference shrinks to $\Delta\chi^2 = +1.6$, in line with the Planck bound $N_{\mathrm{eff}} = 2.99 \pm 0.17$ (the refit shifts n_{S} from 0.9650 to 0.9702 and $H_4^{(0)}$ from 67.31 to 68.23 km/s/Mpc; the errors from the Fisher matrix, $\sigma(\Omega_{\mathrm{b}}h^2_{100}) = 0.00015$, $\sigma(\Omega_\chi h^2_{100}) = 0.0013$, $\sigma(100\theta_*) = 0.0003$, $\sigma(n_{\mathrm{S}}) = 0.0042$, match Planck's). The spectra are shown in the figure. Status: DERIVED (C4).

12.3 The BAO

tracer	z	measured (DESI DR2)	UM-3	χ^2
BGS	0.295	$D_{\mathrm{V}}/r_{\mathrm{s}} = 7.942 \pm 0.075$	8.057	2.34
LRG1	0.510	$D_{\mathrm{M}}/r_{\mathrm{s}} = 13.588 \pm 0.167$, $D_{\mathrm{H}}/r_{\mathrm{s}} = 21.863 \pm 0.425$	13.500, 22.739	4.48
LRG2	0.706	17.351 ± 0.177, 19.455 ± 0.330	17.700, 20.169	14.36

tracer	z	measured (DESI DR2)	UM-3	χ^2
LRG3+ELG1	0.934	21.576 ± 0.152, 17.641 ± 0.193	21.994, 17.569	8.29
ELG2	1.321	27.601 ± 0.318, 14.176 ± 0.221	28.081, 14.065	2.31
QSO	1.484	30.512 ± 0.760, 12.817 ± 0.516	30.275, 12.880	0.10
Lyα	2.330	38.988 ± 0.531, 8.632 ± 0.101	39.182, 8.616	0.13

The BAO distances of UM-3 with the Planck parameters (the matter density parameter $\Omega_{\rm m} = 0.3152$, the density of all matter today in units of the critical density, and $h_{100} r_{\rm s} = 99.09$ Mpc) against DESI DR2 (DESI Collaboration 2025, Table IV; χ^2 with the correlation of $D_{\rm M}$ and $D_{\rm H}$ of each tracer): $\chi^2 = 32.0$ for 13 measurements. Fitted to DESI alone, $\Omega_{\rm m} = 0.2971$ and $h_{100} r_{\rm s} = 101.56$ Mpc with $\chi^2 = 10.5$ for 11 degrees of freedom ($p = 0.48$), as DESI finds for the cosmological-constant model; the Planck point lies $\Delta\chi^2 = 21.5$ above this optimum when its own errors are neglected (with the CMB errors DESI quotes a 2.3 standard-deviation tension between BAO and CMB). The dark energy of UM-3 is a vacuum energy with $w = -1$ to order $\epsilon_{\rm r}$ (w the ratio of its pressure to its energy density): UM-3 inherits this tension and cannot produce the evolving dark energy that DESI DR2 with the CMB prefers at 3.1 standard deviations, $w(a) = w_0 + w_a(1 - a)$ (the equation of state as a function of the scale factor $a = \mathrm{e}^{a_4}$, $a = 1$ today, with its present value w_0 and its slope w_a) with $w_0 > -1$ and $w_a < 0$. Status: DERIVED (C5), STANDARD (DESI's significances).

12.4 The CNB

$$\Gamma_{\rm C\nu B} = N_{\rm T}\, \bar{\sigma}_{\rm T}\, \frac{n_\nu^{(0)}}{2} \sum_{i=1}^{3} |U_{{\rm e}i}|^2 \big(1 + \mathsf{s}_i\big) \qquad \text{(Dirac neutrinos)} \tag{S-29}$$

Symbols in (S-29).

- $\Gamma_{\rm C\nu B}$: The rate at which a tritium target captures neutrinos of the cosmic neutrino background. Unit: 1/time. Status: STANDARD.
- $N_{\rm T}$: The number of tritium nuclei of the target. Unit: dimensionless. Value: 2.0×10^{25} for 100 g. Status: STANDARD.
- $\bar{\sigma}_{\rm T}$: The averaged cross-section of the capture of an electron neutrino on tritium: a tritium nucleus captures left-handed electron neutrinos of mean speed s and number density n at the rate $\bar{\sigma}_{\rm T}(1 + \mathsf{s})\, c\, n$ (c the speed of light, 1 in the formulas; Long, Lunardini and Sabancilar). Unit: cm^2. Value: 3.834×10^{-45} cm^2. Status: STANDARD.
- n_ν: The number density of the cosmic neutrino background per flavour (neutrinos plus antineutrinos, one helicity each); $n_\nu^{(0)}$ its value today. Unit: GeV^3. Value: 112 cm^{-3} today. Status: STANDARD.
- i: A generation index ($i = 1, 2, 3$), which also labels the neutrino mass states ν_1, ν_2, ν_3; in a frame component such as $\mathcal{S}^{(i)}{}_{(i)}$ a direction x_i of ordinary space (the three components are equal); a repeated index is not summed unless a sum sign is written. Status: STANDARD.
- U_{ei}: The electron row of the mixing matrix, $U_{ei} = (U_{\rm PMNS})_{ei}$: $|U_{ei}|^2$ is the electron-flavour fraction of the mass state i. Unit: dimensionless. Value: $|U_{{\rm e}i}|^2 = 0.681, 0.296, 0.022$. Status: STANDARD.
- s_i: The mean speed of the neutrino mass state i today, averaged over its momentum distribution (0 for a state at rest, 1 for a massless one). Unit: dimensionless. Value: 1, 0.061 and 0.011 for ν_1, ν_2 and ν_3 with $m_1 = 0$; 0.126 for ν_1 with $m_1 = 4.12$ meV. Status: STANDARD.

The capture rate of the cosmic neutrinos on a target of $N_{\rm T}$ tritium nuclei ($\nu_e + {}^3\mathrm{H} \to {}^3\mathrm{He} + e^-$, the PTOLEMY idea): each mass state i contributes with its electron fraction $|U_{{\rm e}i}|^2$; for Dirac neutrinos only the left-handed helicity of each state is populated ($n_\nu^{(0)}/2 = 56$ cm^{-3} per state), and it is captured with the averaged cross-section $\bar{\sigma}_{\rm T} = 3.834 \times 10^{-45}$ cm^2 times $(1 + \mathsf{s}_i)$, s_i the mean speed of the state over its momentum distribution (Long, Lunardini and Sabancilar 2014). Status: STANDARD.




Figure 1: The CMB spectra of UM-3 with the Planck 2018 parameters ($D_l = l(l+1)C_l/(2\pi)$ of the temperature, the cross-correlation and the E-mode polarisation, lensed), with three thermal right-handed neutrinos (dashed), and the Planck 2018 plik-lite data.

The CNB of UM-3: the temperature $\mathsf{T}_\nu^{(0)} = (4/11)^{1/3}\mathsf{T}_\gamma^{(0)} = 1.9454$ K (1.676×10^{-4} eV); 112.0 neutrinos and antineutrinos per cm^3 per flavour, 336 in all (FE-53); the states ν_2 and ν_3 became non-relativistic at $\mathsf{z} = 15.3$ and 94 (where their mass equals the mean momentum $3.15\mathsf{T}_\nu$), ν_1 is massless; $\Omega_\nu h^2_{100} = 6.31 \times 10^{-4}$ for the two massive states (FE-58: $\sum m_i/93.14$ eV $= 6.30 \times 10^{-4}$; CAMB: 6.31×10^{-4}). Because the right-handed states are not populated, the CNB of UM-3 has one helicity per state: a 100 g tritium target captures 4.1 cosmic neutrinos per year if all states are non-relativistic (Majorana neutrinos: 8.1, twice as many; Long, Lunardini and Sabancilar: about 4 and 8), and 6.9 per year in UM-3, whose massless ν_1 moves with the speed of light, so that its left-handed helicity is captured twice as often, while ν_2 and ν_3 move with the mean speeds $\mathsf{s}_2 = 0.061$ and $\mathsf{s}_3 = 0.011$ today (the captures of ν_1 lie at the end point of the beta spectrum, where only a mass of ν_1 would separate them). The sum of the masses, 0.0587 eV, lies below the bounds of DESI DR2 with the CMB (0.064 eV) and of ACT DR6 (0.082 eV, 95%). With the largest lightest mass that DESI DR2 allows, $m_1 = 4.12$ meV, the capture rate falls to 4.47 per year (S-34). Status: DERIVED (C6).

13 What the solution does not settle

- **The media of the bulk** (DERIVED: that the late bulk needs a source; ASSUMED: the media). Without a bulk source the energy constraint of FE-66 allows no instant at which both rates of the metric are below $0.477H$, so the late universe needs a medium in the bulk. The static block needs exactly the stabiliser (IN-7), which violates the null energy condition (S-3) and which only a field with negative kinetic energy reproduces; with frozen extra times the bulk source gives off energy that the fold takes up (FE-66, FE-33, FE-11), so it does not couple only to gravity. UM-3 gives no field theory for these media, and how the stabiliser appears after the primordial era, whose bulk is the author's vacuum, is OPEN. The power series additionally needs the medium to respond to the matter of the fold with the radion mode (S-8, S-9); a rigid medium would change the expansion law in a way that observation excludes.
- **The gravity of the fold** (ASSUMED, with a DERIVED bound). The induced Einstein-Hilbert term must provide all but 1.3×10^{-13} of the gravity of the fold (S-30); if the fields of the fold induce it, the cutoff lies at the Planck scale and the Higgs doublet must couple to the curvature with $\xi_\mathrm{H} < 1/6$ (FE-64, ESTIMATE).
- **The primordial era** (OPEN). The author's vacuum is degenerate (S-1: the linear response of the bulk equations vanishes, and the transverse traceless graviton has no time-kinetic term there); a varying b_4 needs bulk sources of second order in its rate (S-2). In the author's vacuum the fields of the fold feel no Hubble friction, their spectrum has the index $n - 1 \approx 2.5$ instead of the observed -0.035, and every squared coupling grows as e^{3Hx_4} (DERIVED, `um3_primordial.py`): the observed spectrum needs an inflationary era with frozen extra times and an expansion rate below 4.7×10^{13} GeV, which UM-3 does not contain, and the Higgs field cannot be its inflaton if the gravity of the fold is induced. The passage from the author's vacuum to the static block with frozen extra times is not computed, and the primordial spectrum, the optical depth and the densities are taken from Planck 2018.
- **The Krein norm** (DERIVED; OPEN: a unitary coupling). The charge of dirac16complex is indefinite (FE-43); its quantisation (FE-67) has ghosts that no choice of the vacuum removes. The vacuum cannot decay, the dilute gas of the particles χ is stable and the dense gas grows (KR4, KR5); the chiral couplings of the fold, the left-handed Higgs portal among them, produce ghosts, so UM-3 with them is not unitary. A unitary UM-3 would need couplings of the fold that do not distinguish the chiralities.
- **The extra times** (ASSUMED, OPEN). Every field that depends on the extra times is tachyonic (FE-38); UM-3 keeps only fields that do not depend on them. The restriction is consistent but unstable: interactions create pairs of opposite momenta along the extra times, bound states with such momenta leak into the bulk, and compact extra times would have to be longer than 26.6 Gpc (DERIVED, `um3_extratimes.py`). UM-3 needs a gauge principle that removes the momentum along the extra times of every field.
- **Lorentz violation in the bulk** (DERIVED, ESTIMATE). Zero modes of bulk fields and the bound particles χ move faster than light on the fold (S-12, S-13, S-31), and the bulk is a shortcut for signals between points of the fold (S-32). For the Standard Model this fixes its place on the fold; for χ it requires $q_\mathrm{b} > H$ and, with the direct-detection bound $q_\mathrm{b} \lesssim 0.03\,\omega_\mathrm{b}$, a bulk scale H well below the mass of χ; the loop estimate for the electron requires q_b/H above 4×10^{10}.
- **The dark energy** (ASSUMED). The vacuum energy ρ_Λ of the fold is a free parameter: UM-3 accommodates its value but does not explain it, and it has $w = -1$, so the evolving dark energy that DESI DR2 with the CMB prefers is not available.
- **The free parameters**. The bulk mass m_Ψ, the scale H, the portal couplings, the self-coupling λ_Ψ, the amplitude $v_\mathcal{A}$ and rate a_{40} of the ad hoc vacuum, the lightest neutrino mass (taken as zero; below 4.12 meV,

S-34) and the Dirac phase (taken as zero; it drops out of every result, S-33) are not fixed by UM-3; the relic density fixes one combination of the mass of χ and its portals.

- **Neutrinos in dense matter** (STANDARD, ESTIMATE). The non-linear evolution of the fast instability and the collective conversion deep in a supernova are not solved; their outcome is modelled by the equipartition FE-52, whose effect on the explosion is taken from the transport simulations of Gogilashvili and Tamborra.

14 What was verified

Every result labelled DERIVED rests on a computation in the folder `universe_model_3/code/` of the repository, which checks itself and writes a report to `universe_model_3/reports/` (all equations and results are written out above). Each script is run with `python universe_model_3/code/NAME.py` (Python 3 with numpy, scipy, sympy, matplotlib and CAMB 2.0.5; the cosmology script downloads the Planck 2018 plik-lite files from github.com/heatherprince/planck-lite-py into `universe_model_3/external/` once).

- `um3_dirac_matrices.py` (55 checks): the author's matrices, their reality, the Clifford relations, the chiral projectors, Pin(4,4).
- `um3_geometry.py` (26 checks): the exact curvature of the author's metric, the Lovelock tensors by two routes, the trace and Bianchi identities, the background equations FE-29, the author's vacuum and its degeneracy, the spin connection and the vielbein postulate.
- `um3_fold.py` (13 checks): the proper chart, the general warped metric FE-34, the junction potential, the fold sources, the conservation law at the fold.
- `um3_bulk.py` (10 checks): the zero modes and the speeds of bulk scalars, photons, fermions and gravitons (S-11, S-12).
- `um3_fields.py` (16 checks): the bilinears of dirac16complex, the Krein operator, the electroweak masses of the ad hoc vacuum, the anomalies with three right-handed neutrinos.
- `um3_spectra.py` (5 checks): the bound states FE-40 with the exact 16×16 matrices, their Krein signature, their speed (S-13).
- `um3_background.py` (6 checks): S-1 to S-4, the ad hoc vacuum as a gauge rotation and its residual (S-10).
- `um3_series.py` (6 checks): S-5 to S-9.
- `um3_kohnsham.py` (9 checks): the Kohn-Sham states (S-14 to S-19) in three regimes, lifted to sixteen components.
- `um3_neutrino.py` (5 checks): the flavour blindness of V_Ψ, the resonance formulas, the crossing criterion, the equipartition, the CNB numbers.
- `um3_flavour.py` (4 checks): S-20 to S-23.
- `um3_thermal.py` (6 checks): the degrees of freedom, S-24 to S-26, the ad hoc vacuum before nucleosynthesis.
- `um3_cosmology.py` (6 checks): S-27 to S-29 and the comparisons with Planck 2018 and DESI DR2.
- `um3_foldsources.py` (6 checks): FE-63, S-30, FE-64 and FE-65.
- `um3_stabiliser.py` (8 checks): the stabiliser, FE-66 and the energy balance of the fold.
- `um3_krein.py` (7 checks): FE-67, the symmetries and mirrors of the bound states, the stability of the gas, the selection rules.
- `um3_extratimes.py` (5 checks): the fields with a momentum along the extra times.
- `um3_superluminal.py` (5 checks): S-31, S-32 and the loop estimate.
- `um3_primordial.py` (5 checks): the fields of the fold in the author's vacuum and the bound on an inflationary era.
- `um3_numass.py` (5 checks): S-33, S-34 and the comparison with NuFIT 6.0.

The Kohn-Sham module runs in parts (`um3_kohnsham.py part NAME` for NAME = K1, I, II, III, K6, K9, then `um3_kohnsham.py assemble`), so that no single calculation takes longer than 423 seconds; the assembled results are identical to a single run.