

The generalized Kronecker delta and the Lovelock tensors of the author's primordial metric

GKD, the pure-Rust generalized Kronecker delta, and the exact Lovelock tensors of order $k = 1, 2, 3$ of Lovelock's equation (4.38) for the 4+4 metric with deflating extra times, with their independent verifications and an exact statement of what they establish

dirac16complex project (with Claude Opus 5.5)

September 2026

Contents

1	The task and the result	3
2	Setting and conventions	4
2.1	The author's metric and coordinates	4
2.2	Curvature conventions	4
3	The generalized Kronecker delta	4
3.1	The author's definition	4
3.2	GKD and the proof that it equals the definition	5
3.3	The self-test of GKD	5
4	The Lovelock tensors	6
4.1	Lovelock's equation (4.38)	6
4.2	Definitions and index conventions of this record	6
4.3	Normalisations	6
4.4	Why the series stops at $k = 3$	7
5	How the code computes them	7
5.1	The crate	7
5.2	The enumeration	9

6	Results	9
6.1	Christoffel symbols and curvature	9
6.2	The Lovelock scalars	10
6.3	The Lovelock tensors	10
6.4	The checks of the Rust program	11
7	Independent verifications	12
7.1	The Wolfram check (29 checks)	12
7.2	The sympy checker (49 checks)	13
8	The comparison with the author's notebook	15
9	How the field equations for a_4 use the Lovelock tensors	15
10	Verification records	17
11	Reproduction	17
11.1	Commands	17
11.2	Expected output	18
11.3	Run times	19
11.4	The PDF	19
12	What is established and what is not	19
12.1	Established	19
12.2	Not established	20

Abstract

This document records the computation of `Revision/gkd_lovelock`: the generalized Kronecker delta of the author's notebook, re-implemented in pure Rust as the function `GKD`, and the three nonzero Lovelock tensors $P_{(1)}$, $P_{(2)}$, $P_{(3)}$ of Lovelock's equation (4.38) for the author's primordial metric of signature (4,4), in which the three extra times x_5, x_6, x_7 deflate as $e^{-a_4(x_4)}$ while 3-space inflates as $e^{a_4(x_4)}$. Every component of $P_{(k)}$ is an exact polynomial in H , a'_4 and a''_4 ; the tensors are diagonal, free of the warp factor $\sin^{1/3} z$, of e^{a_4} and of x_8 , with equal components in the three 3-space directions and in the three extra-time directions, and $P_{(4)} = 0$ identically in eight dimensions. The Rust program checks 19 identities (19 of 19 pass); `GKD` equals the author's determinant on all 16,777,216 pairs of index lists of length 4 and on 200,000 pseudo-random pairs of each length 5 to 9. Two independent verifications agree with every component: a Wolfram Language check that evaluates the author's definition verbatim (29 of 29 checks pass) and a sympy checker that shares no code with the Rust program (49 of 49 checks pass). The field equations for a_4 use these tensors with the normalisation $E_{(k)} = -P_{(k)}/2^{k+1}$, $E_{(1)} = G$. Not established here: any solution of the field equations, anything at or beyond the brane $z = \pi/2$, and a comparison with the answers in the output cells of the author's notebook, which were never read.

1 The task and the result

The author's test (2026-10-01, `Revision/README.md` and `Revision/SPEC.md` sections 5 and 10): refactor the author's Wolfram Language definition of the generalized Kronecker delta into a pure-Rust function `GKD`, use it to compute, with provenance, the three nonzero Lovelock tensors of Lovelock's equation (4.38) for the author's 8 x 8 metric, write every component, and do the work without reading the author's answers. The result, in one paragraph each:

1. `GKD` (section 3). For index lists of equal length, `GKD` returns the determinant of the author's 0/1 matrix without computing a determinant: it is the sign of the permutation that maps the lower list onto the upper list, or 0. The equality with the author's definition is a theorem (section 3.2), confirmed exhaustively and on random samples by three programs.
2. The Lovelock tensors (sections 4 to 6). $P_{(k)}{}^h{}_j$, $A_{(k)}{}^{lh}$ and $L_{(k)}$ for $k = 1, 2, 3$ are computed exactly from the metric alone, with every term of the sums weighted by an explicit call of `GKD`. Their nonzero components are listed in section 6.3.
3. The verifications (sections 7 and 10). The Rust program, the Wolfram check and the sympy checker agree exactly on every one of the 64 components of each tensor; every check of the three reports has the verdict PASS.
4. The use (section 9). The field equations for a_4 of `Revision/field_equations_a4` recompute the three tensors in both of their verifiers and compare them with the file of this record.

Every formula and number below is taken from the Revision files named with it; nothing is taken from the earlier stages of the repository. Interpretations are labelled as such.

2 Setting and conventions

2.1 The author's metric and coordinates

The coordinates are named as the author names them; arrays index them $0, \dots, 7$. x_1, x_2, x_3 are ordinary 3-space, which inflates with the scale factor $e^{a_4} \sin^{1/6} z$; x_4 is the time; x_5, x_6, x_7 are the three extra times, time-like, which DEFLATE exponentially with the scale factor $e^{-a_4} \sin^{1/6} z$ as $a_4(x_4)$ increases (they are never treated as static); x_8 is the hidden space direction with $z = 6Hx_8 \in (0, \pi/2)$; $H > 0$ is the author's constant and $a'_4 = da_4/dx_4$. The metric of `Revision/SPEC.md` section 1, as the Rust program stores it (`metricDiagonal` of `curvature.json`), is

$$g = \text{diag}(e^{2a_4} \sin^{1/3} z, e^{2a_4} \sin^{1/3} z, e^{2a_4} \sin^{1/3} z, -1, \\ -e^{-2a_4} \sin^{1/3} z, -e^{-2a_4} \sin^{1/3} z, -e^{-2a_4} \sin^{1/3} z, \cot^2 z),$$

of signature (4,4): space-like x_1, x_2, x_3, x_8 , time-like x_4, x_5, x_6, x_7 . Its determinant is $\det g = \cos^2 z$, so $\sqrt{|g|} = \cos z$ on $0 < z < \pi/2$; the Rust program writes it as $\sin z \cot z$ (`sqrtAbsDetG`). The 7-volume factor $e^{3a_4} e^{-3a_4} = 1$ shows that the inflation of 3-space is compensated by the deflation of the extra times.

2.2 Curvature conventions

Stated in `Revision/gkd_lovelock/code/src/geometry.rs` and used by all three programs:

- Christoffel symbols $\Gamma^a_{bc} = \frac{1}{2} g^{ad} (\partial_b g_{dc} + \partial_c g_{db} - \partial_d g_{bc})$.
- Riemann tensor (Misner-Thorne-Wheeler) $R^a_{bcd} = \partial_c \Gamma^a_{bd} - \partial_d \Gamma^a_{bc} + \Gamma^a_{ce} \Gamma^e_{bd} - \Gamma^a_{de} \Gamma^e_{bc}$; a round 2-sphere of radius r has $R^{12}_{12} = +1/r^2$.
- $R^{ab}_{cd} = g^{be} R^a_{ecd}$; Ricci $R_{bd} = R^a_{bad}$, so $R^h_j = R^{ha}_{ja}$; $R = R^{ab}_{ab}$; Einstein $G^h_j = R^h_j - \frac{1}{2} \delta^h_j R$.

3 The generalized Kronecker delta

3.1 The author's definition

The cell stored as `In[87]` in the author's notebook `Generalized_Kronecker_Delta_4+4.nb` (the author's `In[54]`) defines

```
kδ[lower_, upper_] /; Length[lower] == Length[upper] := Det[Outer[delta, lower, upper]]
```

For lists $l = (l_1, \dots, l_p)$ and $u = (u_1, \dots, u_p)$ the matrix `Outer[delta, lower, upper]` has the entries $M_{ij} = \delta(l_i, u_j)$ (1 if $l_i = u_j$, otherwise 0), and

$$\delta^{u_1 \dots u_p}_{l_1 \dots l_p} = \det [\delta(l_i, u_j)]_{i,j=1}^p.$$

For lists of different length the definition does not apply and the expression stays unevaluated. Because $\det M = \det M^T$, exchanging the two lists does not change the value.

3.2 GKD and the proof that it equals the definition

Theorem (GKD). For lists of equal length p , $\det M = 0$ if two lower indices are equal, if two upper indices are equal, or if some lower index is not among the upper indices; otherwise $\det M = \text{sign}(\sigma)$, where σ is the permutation with $l_i = u_{\sigma(i)}$.

Proof (the comment of `Revision/gkd_lovelock/code/src/gkd.rs`). (a) Two equal lower indices give two equal rows of M , so $\det M = 0$. (b) Two equal upper indices give two equal columns, so $\det M = 0$. (c) A lower index l_i that is not among the upper indices gives a zero row i , so $\det M = 0$. (d) Otherwise both lists consist of p distinct indices and every l_i equals exactly one $u_{\sigma(i)}$; M has a single 1 in each row i , in column $\sigma(i)$, so it is the permutation matrix of σ , and in the Leibniz formula only the term of σ survives: $\det M = \text{sign}(\sigma)$. QED.

`gkd.rs::GKD` implements cases (a) to (d) with a position table and counts the inversions of σ : $O(p^2)$ integer operations ($p \leq 64$) instead of a $p \times p$ determinant. With more than eight indices over eight values two indices of a list are always equal (pigeonhole), so GKD is 0. `gkd.rs::kdelta_det` is a literal transcription of the definition (the Leibniz determinant of the 0/1 matrix, permutations by Heap's algorithm) used only to test GKD.

3.3 The self-test of GKD

`lovelock_gkd gkd-selftest -exhaustive-max 4` compares GKD with `gkd.rs::kdelta_det` on every pair of index lists of length 1 to 4 over the eight labels and on 200,000 pseudo-random pairs (xorshift64*, half of them permutations of each other) of each length 5 to 9. The report `Revision/gkd_lovelock/results/gkd-selftest.json` has the verdict **SUCCESS**:

length p	mode	pairs	nonzero values	mismatches
1	exhaustive	64		0
2	exhaustive	4096		0
3	exhaustive	262144		0
4	exhaustive	16777216		0
5	random	200000	20538	0
6	random	200000	7812	0
7	random	200000	1949	0
8	random	200000	244	0
9	random	200000	0	0

At length 9 (nine indices over eight values) no pair has a nonzero value, as the pigeonhole argument requires. The Lovelock program records two values of GKD itself (`gkdSelfCheck` of `lovelock-report.json`): +1 for a 3-cycle (`length3Pair`) and -1 for a transposition (`transposition`).

4 The Lovelock tensors

4.1 Lovelock's equation (4.38)

The cell stored as `In[101]` in the author's notebook (the author's `In[68]`) is a picture of Lovelock's equation (4.38) (Lovelock and Rund), rendered to `Revision/gkd_lovelock/results/notebook-in68-image.png` and read from that picture. In n dimensions, with arbitrary constants $\alpha_{(k)}$ and λ and $m = n/2$ for even n ,

$$A^{lh} = \sqrt{g} \sum_{k=1}^{m-1} \alpha_{(k)} g^{jl} \delta_{jj_1 \dots j_{2k}}^{hh_1 \dots h_{2k}} R^{j_1 j_2}_{h_1 h_2} \dots R^{j_{2k-1} j_{2k}}_{h_{2k-1} h_{2k}} + \lambda \sqrt{g} g^{lh}.$$

For the author's metric $n = 8$ and $m = 4$, so the sum runs over $k = 1, 2, 3$; $\sqrt{g}$ means $\sqrt{|\det g|} = \cos z$ (section 2.1).

4.2 Definitions and index conventions of this record

For $k = 1, 2, 3$ (`lovelock-tensors.json`, key `definition`):

$$\begin{aligned} P_{(k)}^h{}_j &= \delta_{jj_1 \dots j_{2k}}^{hh_1 \dots h_{2k}} R^{j_1 j_2}_{h_1 h_2} \dots R^{j_{2k-1} j_{2k}}_{h_{2k-1} h_{2k}} \\ &= \sum \text{GKD}([j, j_1, \dots, j_{2k}], [h, h_1, \dots, h_{2k}]) \prod_{i=1}^k R^{j_{2i-1} j_{2i}}_{h_{2i-1} h_{2i}}, \\ L_{(k)} &= \delta_{jj_1 \dots j_{2k}}^{hh_1 \dots h_{2k}} R^{j_1 j_2}_{h_1 h_2} \dots R^{j_{2k-1} j_{2k}}_{h_{2k-1} h_{2k}}, \\ A_{(k)}^{lh} &= \sqrt{|g|} g^{jl} P_{(k)}^h{}_j, \end{aligned}$$

summed over all repeated indices. Conventions, stated once:

- The lower list of GKD is $[j, j_1, \dots, j_{2k}]$ (the lower indices of δ , which carry the UPPER index pairs of the curvature factors) and the upper list is $[h, h_1, \dots, h_{2k}]$; in the code (`lovelock.rs::dfs`) the upper pair a, b of a factor R^{ab}_{cd} joins the lower row of δ and the lower pair c, d joins the upper row. The field-equations branch writes the lists in the other order; by section 3.1 the value is the same.
- $P_{(k)}$ is stored with its first index up and its second down (`P1_mixed_up_h_down_j`, `P2_mixed_up_h_down_j`, `P3_mixed_up_h_down_j`); $A_{(k)}$ with both indices up (`A1_contravariant_1_h`, `A2_contravariant_1_h`, `A3_contravariant_1_h`). For the diagonal metric $A_{(k)}^{lh} = \cos z g^{ll} P_{(k)}^h{}_l$.
- In this notation equation (4.38) reads $A^{lh} = \sum_{k=1}^3 \alpha_{(k)} A_{(k)}^{lh} + \lambda \sqrt{|g|} g^{lh}$.

4.3 Normalisations

$P_{(k)}$ carries the normalisation of the δ sum. The usual normalised Lovelock tensors are (`lovelock-tensors.json`, key `normalisationNote`; `lovelock.rs::usual_normalisation`)

$$E_{(k)}{}^h{}_j = -\frac{P_{(k)}{}^h{}_j}{2^{k+1}}, \quad E_{(1)} = G,$$

with the following exact relations, derived in the docstring of `Revision/gkd_lovelock/verification/check_lovelock_gkd.py` by the Laplace expansion of the determinant, $P_{(k)}{}^h{}_j = \delta_j^h L_{(k)} - 2k \delta_{jj_2 \dots j_{2k}}^{h_1 \dots h_{2k}} R^{hj_2}{}_{h_1 h_2} \dots$, and verified for this metric:

- $k = 1$: $P_{(1)} = 2R\delta - 4\text{Ric} = -4G$ and $L_{(1)} = 2R$ (`k1_equals_minus_4_einstein`, `L1_equals_2R`).
- $k = 2$: $P_{(2)} = -8\mathcal{H}$, with the Gauss-Bonnet (Lanczos) tensor $\mathcal{H}^h{}_j = 2(RR^h{}_j - 2R^h{}_a R^a{}_j - 2R^{ha}{}_{jb} R^b{}_a + R^{habc} R_{jabc}) - \frac{1}{2}\delta_j^h (R^2 - 4R_{ab}R^{ab} + R_{abcd}R^{abcd})$, and $L_{(2)} = 4(R^2 - 4R^a{}_b R^b{}_a + R^{ab}{}_{cd} R^{cd}{}_{ab})$ (`k2_equals_minus_8_gauss_bonnet`, `L2_equals_4_gauss_bonnet`, `gauss_bonnet_index_placement`).
- $k = 3$: $L_{(3)} = 8(2T_1 + 8T_2 + 24T_3 + 3T_4 + 24T_5 + 16T_6 - 12T_7 + T_8)$ with the cubic invariants $T_1 = R^{ab}{}_{cd} R^{cd}{}_{ef} R^{ef}{}_{ab}$, $T_2 = R^{ab}{}_{cd} R^{ce}{}_{bf} R^{df}{}_{ae}$, $T_3 = R^{ab}{}_{cd} R^{cd}{}_{be} R^e{}_a$, $T_4 = R R^{ab}{}_{cd} R^{cd}{}_{ab}$, $T_5 = R^{ab}{}_{cd} R^c{}_a R^d{}_b$, $T_6 = R^a{}_b R^b{}_c R^c{}_a$, $T_7 = R R^a{}_b R^b{}_a$ and $T_8 = R^3$ (`L3_equals_8_cubic_lovelock_density`).
- The constants are also derived, not only assumed: from literal δ sums on random algebraic curvature tensors, $P_{(1)}/G$ takes the single value -4 in $d = 4$, $P_{(2)}/\mathcal{H}$ the single value -8 in $d = 5$, and in $d = 6$ the 9×8 system of nine random tensors has rank 8 with the unique solution `c = [16, 64, 192, 24, 192, 128, -96, 8] = 8 x [2, 8, 24, 3, 24, 16, -12, 1]` (`normalisation_P1_derived_minus_4`, `normalisation_P2_derived_minus_8`, `normalisation_L3_cubic_derived`).
- Trace identity: $\sum_h P_{(k)}{}^h{}_h = (n - 2k)L_{(k)} = (8 - 2k)L_{(k)}$.

4.4 Why the series stops at $k = 3$

$P_{(4)}$ would need δ with $2 \cdot 4 + 1 = 9$ upper and 9 lower indices over eight values: two indices of each list are equal, so every term vanishes and $P_{(4)} = 0$ identically (`k4_tensor_vanishes`, `k4_terms_vanish_literally`, `gkd_nine_indices_in_eight_dimensions_vanish`; `lovelock-tensors.json` key `k4`: “identically zero (GKD of 9 indices in 8 dimensions)”). Equation (4.38) stops at $k = m - 1 = 3$, so $k = 1, 2, 3$ is the complete series in eight dimensions.

5 How the code computes them

5.1 The crate

`Revision/gkd_lovelock/code` is the Rust crate `lovelock_gkd` (edition 2021, no dependencies, its own empty workspace). Its modules:

file	function or type	what it does
gkd.rs	gkd.rs::GKD	the generalized Kronecker delta by cases (a) to (d) of section 3.2
gkd.rs	gkd.rs::kdelta_det, gkd.rs::compare_exhaustive, gkd.rs::compare_random	the literal Leibniz determinant and the two self-test loops
rational.rs	rational.rs::Rational	exact rationals on i128; every operation that would overflow stops with a message
poly.rs	poly.rs::Poly, poly.rs::d	exact Laurent polynomials with rational coefficients in H , a'_4 to a''''_4 , $E = e^{a_4}$, $S = \sin^{1/3} z$, $C = \cot z$, with the exact derivatives in x_4 and x_8
geometry.rs	geometry.rs::metric_diagonal, geometry.rs::Geometry	the metric, its inverse, $\sqrt{ g } = S^3 C$, the Christoffel symbols, $R^a{}_{bcd}$, $R^{ab}{}_{cd}$, Ricci and R
geometry.rs	geometry.rs::nonzero_mixed, geometry.rs::einstein_mixed	the list of nonzero $R^{ab}{}_{cd}$; the Einstein tensor from the Ricci tensor
lovelock.rs	lovelock.rs::lovelock_mixed, lovelock.rs::lovelock_scalar, lovelock.rs::dfs	$P_{(k)}{}^h{}_j$ and $L_{(k)}$ by a depth-first search over the nonzero curvature entries
lovelock.rs	lovelock.rs::lovelock_contravariant, lovelock.rs::divergence	$A_{(k)}{}^{lh}$; the covariant divergence $\nabla_h P^h{}_j$
lovelock.rs	lovelock.rs::brute_force_numeric, lovelock.rs::usual_normalisation	the unpruned literal sum at a numerical point; the factor $-1/2^{k+1}$
output.rs	output.rs::tensor_json, output.rs::monomials_json, output.rs::tensor_markdown	deterministic JSON and Markdown writers
main.rs	main.rs::gkd_selftest, main.rs::run_lovelock, main.rs::print_config	the command line <code>gkd-selftest</code> , <code>lovelock</code> , <code>print-config</code>

The symbols S and C satisfy $S^6(1 + C^2) = 1$, so the representation of `poly.rs` is not canonical in general; S cancels identically in every mixed tensor the program prints, every zero test is done on mixed tensors, and the independent checkers use a canonical form (section 7.2).

5.2 The enumeration

For each pair (h, j) , `lovelock.rs::dfs` chooses k ordered nonzero entries R^{ab}_{cd} one after another and appends a, b to the lower list and c, d to the upper list. A branch is cut when an index would repeat within a list (GKD is 0 by cases (a) and (b)); at a leaf the two index SETS must be equal (otherwise GKD is 0 by case (c)), and only then is GKD called; a nonzero value multiplies the product of the chosen entries. A vanishing curvature factor gives a vanishing term, so only the nonzero entries are enumerated. The counters of `lovelock-report.json`:

k	leaves	GKD calls	nonzero GKD values	GKD calls for $L_{(k)}$	nonzero components of $P_{(k)}$
1	5616	696	696	108	8
2	176640	32640	32640	7200	8
3	1128960	495360	495360	213120	8
4	0	0	0	1105920	0

The pruning is checked, not assumed: `lovelock.rs::brute_force_numeric` evaluates the completely unpruned literal sum, every index list sent to GKD, at the point $H = 0.23$, $a_4 = 0.17$, $a'_4 = 0.61$, $a''_4 = -0.37$, $x_8 = 0.41$: 262144 index lists for $k = 1$ (maximal relative deviation from the exact $P_{(1)}$ 1.40e-15) and 1073741824 index lists for $k = 2$ (maximal relative deviation from the exact $P_{(2)}$ 4.41e-14) (`k1_brute_force_numeric`, `k2_brute_force_numeric`). The two independent checkers repeat the unpruned sums exactly (section 7).

6 Results

6.1 Christoffel symbols and curvature

`curvature.json` lists 25 nonzero Christoffel symbols Γ^a_{bc} with $b \leq c$. With $i = 1, 2, 3$ (3-space) and $t = 5, 6, 7$ (extra times):

$$\begin{aligned}
\Gamma^{x_i}_{x_i x_4} &= a'_4, & \Gamma^{x_t}_{x_4 x_t} &= -a'_4, & \Gamma^{x_i}_{x_i x_8} &= \Gamma^{x_t}_{x_t x_8} = H \cot z, \\
\Gamma^{x_4}_{x_i x_i} &= a'_4 e^{2a_4} \sin^{1/3} z, & \Gamma^{x_4}_{x_t x_t} &= a'_4 e^{-2a_4} \sin^{1/3} z, \\
\Gamma^{x_8}_{x_i x_i} &= -H e^{2a_4} \sin^{1/3} z \tan z, & \Gamma^{x_8}_{x_t x_t} &= H e^{-2a_4} \sin^{1/3} z \tan z, \\
\Gamma^{x_8}_{x_8 x_8} &= -6H \tan z - 6H \cot z.
\end{aligned}$$

The mixed Riemann tensor R^{ab}_{cd} has 156 nonzero entries. Up to the antisymmetry in each index pair they are, with $i \neq i'$ in 3-space and $t \neq t'$ among the extra times:

entry	value
$R^{x_i x_{i'}}_{x_i x_{i'}}$ and $R^{x_t x_{t'}}_{x_t x_{t'}}$	$(a'_4)^2 - H^2$
$R^{x_i x_t}_{x_i x_t}$	$-(a'_4)^2 - H^2$
$R^{x_i x_4}_{x_i x_4}$	$a''_4 + (a'_4)^2$
$R^{x_4 x_t}_{x_4 x_t}$	$-a''_4 + (a'_4)^2$

entry	value
$R^{x_i x_8}_{x_i x_8}$ and $R^{x_t x_8}_{x_t x_8}$	$-H^2$
$R^{x_i x_4}_{x_i x_8}$ and $R^{x_4 x_t}_{x_t x_8}$	$H a'_4 \cot z$
$R^{x_i x_8}_{x_i x_4}$ and $R^{x_t x_8}_{x_4 x_t}$	$-H a'_4 \tan z$

$R^{x_4 x_8}_{x_4 x_8} = 0$. Every R^{ab}_{cd} is free of $\sin^{1/3} z$ and of e^{a_4} (`mixed_riemann_free_of_sin_third`, `mixed_riemann_free_of_warp_and_exponential`). The deflation enters through a'_4 and a''_4 with the opposite sign of the inflating directions: the sectional curvature of the plane (x_i, x_4) is $a''_4 + (a'_4)^2$ and that of (x_4, x_t) is $-a''_4 + (a'_4)^2$.

The mixed Ricci tensor is diagonal, $R^{x_i}_{x_i} = a''_4 - 6H^2$, $R^{x_4}_{x_4} = 6(a'_4)^2$, $R^{x_t}_{x_t} = -a''_4 - 6H^2$, $R^{x_8}_{x_8} = -6H^2$; the Ricci scalar is $R = 6(a'_4)^2 - 42H^2$ (`ricciScalar`), and the Einstein tensor is

$$\begin{aligned} G^{x_i}_{x_i} &= a''_4 - 3(a'_4)^2 + 15H^2, & G^{x_4}_{x_4} &= 3(a'_4)^2 + 21H^2, \\ G^{x_t}_{x_t} &= -a''_4 - 3(a'_4)^2 + 15H^2, & G^{x_8}_{x_8} &= -3(a'_4)^2 + 15H^2. \end{aligned}$$

6.2 The Lovelock scalars

From `lovelock-tensors.json` (keys L1, L2, L3):

$$\begin{aligned} L_{(1)} &= 12(a'_4)^2 - 84H^2, \\ L_{(2)} &= -96(a'_4)^4 - 2112H^2(a'_4)^2 + 3360H^4, \\ L_{(3)} &= 1152(a'_4)^6 + 31104H^2(a'_4)^4 + 100224H^4(a'_4)^2 - 40320H^6. \end{aligned}$$

6.3 The Lovelock tensors

Every $P_{(k)}^{h_j}$ is diagonal: all 56 off-diagonal components of each tensor vanish, including x_4 - x_8 . The diagonal satisfies $x_1 = x_2 = x_3$ and $x_5 = x_6 = x_7$, so four components per order are independent. The formulas below are the LaTeX texts of `lovelock-tensors.json`, unchanged.

$k = 1$:

$$\begin{aligned} P_{(1)}^{x_1}_{x_1} &= -4a''_4 + 12(a'_4)^2 - 60H^2 \\ P_{(1)}^{x_4}_{x_4} &= -12(a'_4)^2 - 84H^2 \\ P_{(1)}^{x_5}_{x_5} &= 4a''_4 + 12(a'_4)^2 - 60H^2 \\ P_{(1)}^{x_8}_{x_8} &= 12(a'_4)^2 - 60H^2 \end{aligned}$$

$k = 2$:

$$\begin{aligned}
P_{(2)}^{x_1 x_1} &= 192 (a'_4)^2 a''_4 - 96 (a'_4)^4 + 320 H^2 a''_4 \\
&\quad - 1344 H^2 (a'_4)^2 + 1440 H^4 \\
P_{(2)}^{x_4 x_4} &= 288 (a'_4)^4 + 960 H^2 (a'_4)^2 + 3360 H^4 \\
P_{(2)}^{x_5 x_5} &= -192 (a'_4)^2 a''_4 - 96 (a'_4)^4 - 320 H^2 a''_4 \\
&\quad - 1344 H^2 (a'_4)^2 + 1440 H^4 \\
P_{(2)}^{x_8 x_8} &= -96 (a'_4)^4 - 1344 H^2 (a'_4)^2 + 1440 H^4
\end{aligned}$$

$k = 3$:

$$\begin{aligned}
P_{(3)}^{x_1 x_1} &= -5760 (a'_4)^4 a''_4 + 1152 (a'_4)^6 - 6912 H^2 (a'_4)^2 a''_4 + 10368 H^2 (a'_4)^4 \\
&\quad - 5760 H^4 a''_4 + 31104 H^4 (a'_4)^2 - 5760 H^6 \\
P_{(3)}^{x_4 x_4} &= -5760 (a'_4)^6 - 10368 H^2 (a'_4)^4 - 17280 H^4 (a'_4)^2 - 40320 H^6 \\
P_{(3)}^{x_5 x_5} &= 5760 (a'_4)^4 a''_4 + 1152 (a'_4)^6 + 6912 H^2 (a'_4)^2 a''_4 + 10368 H^2 (a'_4)^4 \\
&\quad + 5760 H^4 a''_4 + 31104 H^4 (a'_4)^2 - 5760 H^6 \\
P_{(3)}^{x_8 x_8} &= 1152 (a'_4)^6 + 10368 H^2 (a'_4)^4 + 31104 H^4 (a'_4)^2 - 5760 H^6
\end{aligned}$$

Remarks, each an exact property of the listed components:

- $P_{(1)} = -4G$ component by component (compare section 6.1).
- Of the metric function only the derivatives a'_4 and a''_4 occur, not a_4 itself; a''_4 occurs only in the 3-space and extra-time components, with opposite signs (the inflating and the deflating directions), and never in the x_4 and x_8 components.
- $P_{(k)}^{x_1 x_1} + P_{(k)}^{x_5 x_5} = 2P_{(k)}^{x_8 x_8}$ for $k = 1, 2, 3$ (read off the lists above).
- The contravariant densities are $A_{(k)}^{ll} = \cos z g^{ll} P_{(k)}^l{}_l$, with $g^{x_i x_i} = e^{-2a_4} \sin^{-1/3} z$, $g^{x_4 x_4} = -1$, $g^{x_t x_t} = -e^{2a_4} \sin^{-1/3} z$ and $g^{x_8 x_8} = \tan^2 z$; for example the file gives $A_{(1)}^{x_4 x_4} = 12 (a'_4)^2 \sin(6Hx_8) \cot(6Hx_8) + 84 H^2 \sin(6Hx_8) \cot(6Hx_8)$. All 64 components of each $A_{(k)}$, each with its exact monomial list, are in `lovelock-tensors.json`; all 384 components of $P_{(k)}$ and $A_{(k)}$ are also written out in `lovelock-components.md`.

6.4 The checks of the Rust program

`lovelock_gkd lovelock -brute-force-k2` writes `curvature.json`, `lovelock-tensors.json`, `lovelock-components.md` and `lovelock-report.json`. The report gives "checkCount": 19, "failedCheckCount": 0 and "verdict": "SUCCESS"; every one of its 19 checks has "passed": true:

check	check	check
riemann_antisymmetry	riemann_first_bianchi	mixed_riemann_frr ee_of_sin_third

check	check	check
k1_equals_minus_4_einstein	k1_trace_identity	k1_divergence_free
k1_symmetric	k1_free_of_sin_third	k2_trace_identity
k2_divergence_free	k2_symmetric	k2_free_of_sin_third
k3_trace_identity	k3_divergence_free	k3_symmetric
k3_free_of_sin_third	k4_tensor_vanishes	k1_brute_force_numeric
k2_brute_force_numeric		

They are mathematical identities, not comparisons with the author's numbers: the antisymmetries and the first Bianchi identity of R^{ab}_{cd} for all 4096 index lists, $P_{(1)} = -4G$ by an independent route, the trace identities, $\nabla_h P_{(k)}^h{}_j = 0$ for every j , the symmetry of $g_{hh} P_{(k)}^h{}_j$ (so $A_{(k)}^{lh} = A_{(k)}^{hl}$), the absence of the warp factor, $P_{(4)} = 0$ and the unpruned numerical sums. Two runs give byte-identical files (Revision/gkd_lovelock/results/PROVENANCE_OF_THE_COMPUTATION.md).

7 Independent verifications

7.1 The Wolfram check (29 checks)

Revision/gkd_lovelock/verification/verify_lovelock_gkd.wls with the package Lovelock GKDCheck.wl writes wolfram-gkd-report.json: "checkCount": 29, "expectedCheckCount": 29, "failedCheckCount": 0, "verdict": "SUCCESS", producer Wolfram Language 15.0.1. What it does:

- The author's definition, re-typed verbatim, is the kernel's definition of `kδ` (compared with the text recorded in PROVENANCE_OF_THE_COMPUTATION.md), with `delta[a_Integer, b_Integer] := KroneckerDelta[a, b]`; unequal lengths stay unevaluated.
- GKD is NOT re-implemented in Wolfram Language: a small exporter compiled with cargo against the unchanged crate writes the values of the Rust function GKD for 346304 pairs (all 64, 4096 and 262144 pairs of lengths 1, 2, 3 and 20000 pseudo-random pairs of each length 4 to 7; values file of 3161984 bytes), and the verbatim `kδ` is evaluated on every pair: 0 mismatches. The negative control against `-GKD` reports a mismatch at exactly every pair where GKD is nonzero (8, 112, 2016, 10022, 10005, 10002, 10000 pairs for lengths 1 to 7).
- The curvature is recomputed from the metric with `D`, `Inverse` and `Simplify` (156 nonzero R^{ab}_{cd} , the same as `curvature.json`), and $\det g = \cos^2 z$.
- The Lovelock tensors are recomputed with the verbatim `kδ`: for $k = 1$ and $k = 2$ every k -tuple of the 156 nonzero entries for each of the 64 components (9984 and 1557504 calls), for $k = 3$ every list left after skipping lists with a repeated label in a row (1128960 calls). `FullSimplify` of the difference with the Rust tensors is 0 for all 64 components of each $P_{(k)}$ and $A_{(k)}$. The numbers of nonzero `kδ` terms, 696, 32640 and 495360, equal the Rust counters. A tampered $P_{(2)}$ component and a tampered $A_{(3)}$ component are flagged, and only those.

check	check
definition_is_the_authors_verbatim	definition_unequal_lengths_stay_unevaluated
gkd_values_file_layout	gkd_equals_kdelta_exhaustive_len_gth_1
gkd_equals_kdelta_exhaustive_len_gth_2	gkd_equals_kdelta_exhaustive_len_gth_3
gkd_equals_kdelta_random_length_4	gkd_equals_kdelta_random_length_5
gkd_equals_kdelta_random_length_6	gkd_equals_kdelta_random_length_7
negative_control_gkd_comparison_detects_a_sign_flip	riemann_mixed_equals_rust_curvature
sqrt_det_g_is_cos	lovelock_kdelta_values_are_integers
k1_P_equals_rust_all_64_components	k1_A_equals_rust_all_64_components
k2_P_equals_rust_all_64_components	k2_A_equals_rust_all_64_components
k3_P_equals_rust_all_64_components	k3_A_equals_rust_all_64_components
negative_control_tensor_comparison_detects_a_change	k1_equals_minus_4_einstein
k1_trace_equals_6_L1	k2_trace_equals_4_L2
k3_trace_equals_2_L3	k1_nonzero_kdelta_terms_equal_rust_counter
k2_nonzero_kdelta_terms_equal_rust_counter	k3_nonzero_kdelta_terms_equal_rust_counter
rust_json_text_equals_its_monomials	

Scope. The Wolfram check is independent of the Rust code in its definition of δ (the author's verbatim text), its curvature and its Lovelock sums; it uses the Rust program for the GKD values it compares and for the tensors it compares with. Its default run skips, for $k = 3$, lists with a repeated label in a row (their determinant is 0 because two rows or columns are equal); an optional mode recomputes the eight diagonal $k = 3$ components with no skip at all, and its single recorded run passed 30 of 30 checks (Revision/gkd_lovelock/verification/WOLFRAMSCRIPT_PROVENANCE.md, part 4.5); that report is not committed. Only Wolfram 15.0.1 was used.

7.2 The sympy checker (49 checks)

The sympy checker is the file `check_lovelock_gkd.py` in the folder `Revision/gkd_lovelock/verification/`. Its report `python-lovelock-report.json` gives "checkCount": 49, "failedCheckCount": 0 and "verdict": "SUCCESS". It shares no code with the Rust crate: the geometry is sympy's own differentiation, δ is the author's definition evaluated with sympy's determinant (memoised by the 0/1 matrix), and the sums are organised breadth-first (all k -fold products with pairwise different lower and pairwise different upper indices, then completed by every admissible h and j , one literal call of the determinant per term). Its counters:

k	products	determinant calls	nonzero values	distinct multisets
1	156	5772	804	156
2	11040	187680	39840	5520
3	282240	1411200	708480	47040

with 4096401 determinant calls in all and 29030 distinct 0/1 matrices determined by sympy. The skipped terms are checked as well: the completely unpruned literal sums for $k = 1$ (10140 calls) and $k = 2$ (1581840 calls) agree exactly with the pruned sums; 20000 random terms of the unpruned $k = 3$ sum, 19900 of them with a repeated index, give 0 whenever skipped; 300 random 9-index lists give 0. The comparison with the Rust output is exact three times over: by an exact canonical form (`canon`, which uses $s^6(1 + c^2) = 1$ to reach a unique form, so a zero difference is a proof of equality), by plain sympy simplification, and numerically with 60 digits at 5 random rational points.

check	check	check
gkd_literal_equa ls_cofactor_expansion	gkd_examples	gkd_nine_indices _in_eight_dims ions_vanish
normalisation_P1 _derived_minus_4 metric_inverse riemann_first_bianchi	normalisation_P2 _derived_minus_8 sqrt_abs_det_g mixed_riemann_fr ee_of_warp_and_e xponential	normalisation_L3 _cubic_derived christoffel_symmetric riemann_antisymm etry_and_pair_symmetry
k1_unpruned_lite ral_sum_agrees k4_terms_vanish_ literally k2_equals_minus_ 8_gauss_bonnet L3_equals_8_cubi c_lovelock_density k3_trace_identity k1_free_of_warp k2_free_of_warp k3_free_of_warp	k2_unpruned_lite ral_sum_agrees k1_equals_minus_ 4_einstein L2_equals_4_gaus s_bonnet k1_trace_identity k1_divergence_free k2_divergence_free k3_divergence_free rust_k1_mixed_co mponents_agree rust_k2_mixed_co mponents_agree rust_k3_mixed_co mponents_agree negative_control s_detected rust_riemann_agrees	k3_pruned_terms_ vanish_literally L1_equals_2R gauss_bonnet_ind ex_placement k2_trace_identity k1_symmetric k2_symmetric k3_symmetric rust_k1_contrava riant_components_agree rust_k2_contrava riant_components_agree rust_k3_contrava riant_components_agree rust_metric_and_ sqrt_g_agree rust_ricci_einst ein_scalar_agree
rust_L1_agrees rust_L2_agrees rust_L3_agrees rust_christoffels_agree rust_text_matche s_monomials		

Scope. The sympy checker reads only the metric (typed in from the task text) and, for the comparison, `curvature.json` and `lovelock-tensors.json`; it never reads the author's notebook. Its literal δ is compared with an independent cofactor expansion on all 266304 pairs of lengths 1 to 3 and 7500 random pairs of lengths 4 to 8. The negative controls (a coefficient increased by 1, a wrong power of $\sin^{1/3} z$, a wrong power of e^{a_4} , an added $10^{-30}H^4$) are each detected by each of its three comparison methods.

8 The comparison with the author's notebook

What was read is recorded in `PROVENANCE_OF_THE_COMPUTATION.md`, written before any comparison, and in the provenance file `WOLFRAMSCRIPT_PROVENANCE.md` of the folder `Revision/gkd_lovelock/notebook_reading/`: only the 58 INPUT cells of `Generalized_Kronecker_Delta_4+4.nb` (sha256 23bb4e0c70943e766d9b081a3a399ef29664889aad088041329ce2e065b80afb), converted to text without evaluation; the committed digest `notebook-input-cells.txt` gives, for each, its label, its length and its first 160 characters. The definition of $k\delta$ was taken from the cell `In[87]` and Lovelock's equation from the picture in the cell `In[101]` (1372 x 435 pixels). No output cell, no file `Generalized_Kronecker_Delta.txt` and no other Mathematica file of the author was opened; the metric was taken from the author's message.

What the notebook computes, as far as its input cells show (a reading of the digest, labelled as such): it sets up the `xAct` packages, an 8-dimensional manifold and a metric of signature (4,4) (`In[29]`); it builds the generalized delta in a second way, as contracted products of two Levi-Civita tensors divided by $(8-p)!$ (`delta11`, `delta22`, `delta33`, `delta55`); it defines $k\delta$ (`In[87]`) and evaluates it for 3, 5 and 7 index pairs (`kdelta33`, `kdelta55`, `kdelta77`); it compares the two constructions (`In[102]`, `In[105]`, `In[108]`); and it contains the picture of (4.38). Its answers are in the output cells, which were not read.

The comparison that was made:

- The definition: `definition_is_the_authors_verbatim` shows that the definition evaluated by the Wolfram check is the author's text, character for character.
- The function: `GKD` equals the author's $k\delta$ on every pair of lists of length 1 to 3, on 20000 random pairs of each length 4 to 7 (Wolfram), and equals the literal determinant on every pair of length 1 to 4 and 200000 random pairs of each length 5 to 9 (Rust self-test), as the theorem of section 3.2 requires.
- The equation: the tensors are those of (4.38) as shown in the notebook's picture, with the conventions of section 4.2.

The comparison that was NOT made: `PROVENANCE_OF_THE_COMPUTATION.md` announces a comparison with the author's own answers in a later commit; no file under `Revision/` records it. Whether the Lovelock tensors of this record agree with any tensors computed in the author's notebooks is therefore not established (section 12).

9 How the field equations for a_4 use the Lovelock tensors

`Revision/field_equations_a4` writes the Einstein-Lovelock equations of `Revision/SPEC.md` section 5 for the author's metric,

$$\sum_{k=1}^3 \alpha_k E_{(k)}{}^\mu{}_\nu + \Lambda \delta^\mu_\nu = \kappa T^\mu{}_\nu, \quad E_{(k)} = -\frac{P_{(k)}}{2^{k+1}}, \quad E_{(1)} = G,$$

with $\alpha_1 = 1$ for Einstein gravity ($\alpha_2 = \alpha_3 = 0$). Both of its verifiers recompute the three tensors with their own GKD (Wolfram: as the determinant of the delta matrix; sympy: as the permutation sign) and compare all 64 components of each with the exact monomial lists of `Revision/gkd_lovelock/results/lovelock-tensors.json`:

step	Wolfram records (<code>wolfram-a4-report.json</code>)	sympy records (<code>python-a4-report.json</code>)
$P_{(k)}$ equal to this record	P1_direct_equals _gkd_branch_monomials, P2_direct_equals _gkd_branch_monomials, P3_direct_equals _gkd_branch_monomials	P1_equals_gkd_branch_monomials, P2_equals_gkd_branch_monomials, P3_equals_gkd_branch_monomials
$L_{(k)}$ equal to this record	L1_direct_equals _gkd_branch, L2_direct_equals_gkd_branch, L3_direct_equals _gkd_branch	L1_equals_gkd_branch, L2_equals_gkd_branch, L3_equals_gkd_branch
$P_{(1)} = -4G$, traces	P1_direct_equals _minus_4_Einstein, P1_trace_identity, P2_trace_identity, P3_trace_identity	P1_equals_minus_4_Einstein, P1_trace_identity, P2_trace_identity, P3_trace_identity
structure, $P_{(4)} = 0$	P1_structure, P2_structure, P3_structure, independent_components, P4_vanishes_pigeonhole	P1_structure, P2_structure, P3_structure, other_components_vanish
divergence of $E_{(k)}$; the $E_{(k)}$ of <code>a4-equations.json</code>	E1_divergence_free, E2_divergence_free, E3_divergence_free	E1_divergence_free, E2_divergence_free, E3_divergence_free, json_lovelock_components

What follows from the tensors of this record (taken from those reports, not derived here): the independent components are $x_1 = x_2 = x_3$, x_4 , $x_5 = x_6 = x_7$, x_8 and the pair x_4 - x_8 , whose left-hand side is 0; the evolution equation, the 3-space minus extra-time combination, factorises as $a_4'' F(a_4') = \kappa(p_3 - p_t)$ with

$$F(a_4') = 2\alpha_1 - 48\alpha_2(a_4')^2 - 80\alpha_2 H^2 + 720\alpha_3(a_4')^4 + 864\alpha_3(a_4')^2 H^2 + 720\alpha_3 H^4$$

(Wolfram record `evolution_factorises_a4pp_times_F`, sympy record `evolution_factorises`). The identity $P_{(k)x_1}^{x_1} + P_{(k)x_5}^{x_5} = 2P_{(k)x_8}^{x_8}$ of section 6.3 becomes the source condition $p_3 + p_t = 2p_8$ (`algebraic_identity_x1_plus_x5_minus_2x8`, `algebraic_identity`). The solutions of these equations and their sources are the subject of the field-theory documents `Revision/docs/DIRAC16COMPLEX_FIELD_THEORY.md` and `Revision/docs/DIRAC16COMPLEX00_FIELD_THEORY.md`, not of this one.

10 Verification records

Every check has a name, a verdict and a detail. Counts at the time of writing, as the JSON files give them (the publication test re-reads them):

report	checks	PASS	FAIL
Revision/gkd_lovelock/results/lovelock-report.json	19	19	0
Revision/gkd_lovelock/results/wolfram-gkd-report.json	29	29	0
Revision/gkd_lovelock/results/python-lovelock-report.json	49	49	0
Revision/field_equations_a4/reports/wolfram-a4-report.json	52	52	0
Revision/field_equations_a4/reports/python-a4-report.json	63	63	0

The Rust report stores each check with `"passed": true` instead of a verdict; its 19 checks are counted as PASS. The GKD self-test `Revision/gkd_lovelock/results/gkd-selftest.json` has no list of checks; its nine comparisons (section 3.3) all have 0 mismatches and its verdict is `SUCCESS`. The test file `Revision/tests/test_gkd_lovelock.py` pins the sha256 of the seven result files of the computation and its verifications (`curvature.json`, `lovelock-tensors.json`, `lovelock-components.md`, `lovelock-report.json`, `gkd-selftest.json`, `python-lovelock-report.json`, `wolfram-gkd-report.json`), requires every check to pass, and, in its slow tests, re-runs the Rust program, the sympy checker and the Wolfram check and requires byte-identical outputs.

Re-verification for this document (2026-10-08, on the shared machine of the provenance files; written to a scratch directory, the committed files untouched): `lovelock_gkd lovelock -brute-force-k2` gave 19 of 19 checks in 18.9 s and four files byte-identical to the committed ones; the sympy checker gave 49 of 49 checks in 40.3 s and a report byte-identical to the committed one. These two run times are measurements made while writing this document; they are not recorded in a report.

11 Reproduction

11.1 Commands

From the repository root (Rust 1.87 or newer, Python 3 with sympy and numpy, Wolfram Language with WolframScript). A backslash at the end of a line is the line continuation of a POSIX shell; in

PowerShell write such a command on one line:

```
cargo build --release --manifest-path Revision/gkd_lovelock/code/Cargo.toml
Revision/gkd_lovelock/code/target/release/lovelock_gkd gkd-selftest \
  --exhaustive-max 4 --output Revision/gkd_lovelock/results
Revision/gkd_lovelock/code/target/release/lovelock_gkd lovelock \
  --output Revision/gkd_lovelock/results --brute-force-k2
python Revision/gkd_lovelock/verification/check_lovelock_gkd.py
wolframscript -file Revision/gkd_lovelock/verification/verify_lovelock_gkd.wls \
  Revision/gkd_lovelock/results/wolfram-gkd-report.json
wolframscript -file \
  Revision/gkd_lovelock/notebook_reading/lovelock_extract_nb_inputs.wls
python Revision/gkd_lovelock/notebook_reading/lovelock_digest_nb_inputs.py
wolframscript -file \
  Revision/gkd_lovelock/notebook_reading/lovelock_export_nb_image.wls
python -m unittest Revision/tests/test_gkd_lovelock.py -v
python scripts/build_provenance_pdf.py Revision/docs/LOVELOCK_GKD.md \
  --developer-layout --specifications Revision/pdf-specifications.json
python -m unittest Revision/tests/test_lovelock_gkd_publication.py -v
```

On Windows the binary is `lovelock_gkd.exe`. Each of the commands that write into `Revision/gkd_lovelock/results` overwrites a committed file with a byte-identical one; to leave the committed files untouched, give another output directory or report path and compare. The Wolfram check builds its exporter in `<$TemporaryDirectory>/revision_gkd_export` (or in `$LOVELOCK_GKD_EXPORT_DIR`); run only one copy at a time with the default directory. The notebook-reading commands need the author's notebook in the repository root and write the uncommitted text `build/lovelock_nb_inputs.txt`.

11.2 Expected output

- `gkd-selftest`: nine lines starting with `PASS - GKD vs literal`, then `gkd-selftest: SUCCESS (...)`; the file `gkd-selftest.json` with the table of section 3.3.
- `lovelock`: the 19 check lines starting with `PASS -`, three lines starting with `info -` (the scalar $L_{(4)}$ and the times of the two unpruned sums), the counter lines of section 5.2, then `check_count=19, failed_check_count=0` and `lovelock: SUCCESS (...)`; exit code 0.
- The sympy checker: 49 lines `[PASS] <name>` in six groups, then `total ... s; 49 checks, 0 failed; verdict SUCCESS`; exit code 0.
- The Wolfram check: ten `time_<step>=<seconds>` lines, the 29 lines `check_<name>=true`, then `check_count=29, failed_check_count=0, time_total=<seconds>` and `report=<path>`; exit code 0 (1 if a check fails, 2 on a load, build or input/output error).
- The notebook reading: `input cells written: 58, 58 input cells, {1372, 435}` and `28229 bytes (117 bytes of text/time chunks removed)`, with both committed outputs reproduced byte for byte.

11.3 Run times

From the provenance files (shared 24-core machine, Windows 11):

- The Wolfram check, from part 4.4 of the provenance file `WOLFRAMSCRIPT_PROVENANCE.md` in `Revision/gkd_lovelock/verification/`: `time_total` 68.4 s in the run of 2026-10-08 that regenerated the committed report, 68-135 s over seven runs; the $k = 2$ sum took 26-43 s and the $k = 3$ sum 35-77 s over eight runs; peak memory of the kernel about 486 MB working set. The optional no-skip mode for $k = 3$ took 1637.7 s (part 4.5).
- The notebook reading, from the provenance file `WOLFRAMSCRIPT_PROVENANCE.md` in `Revision/gkd_lovelock/notebook_reading/`: the whole set takes about 10 to 25 seconds, and up to about 40 seconds on a fully loaded machine.
- The GKD self-test with `-exhaustive-max 4`: several minutes (`Revision/tests/test_gkd_lovelock.py`, which runs it only when `LOVELOCK_GKD_SELFTEST=1`).
- The Rust `lovelock` run and the sympy checker: no run time is recorded in a provenance file; see the measurement of section 10 (18.9 s and 40.3 s).

11.4 The PDF

The PDF build runs the Markdown-to-LaTeX builder twice (with different hash seeds), runs `pdflatex` three times into each of two fresh directories, requires warning-free logs and byte-identical PDFs, and checks the PDF against its entry `lovelock-gkd` in `Revision/pdf-specifications.json` (the Revision registry; the registry of the earlier stages is not touched).

12 What is established and what is not

12.1 Established

1. GKD equals the author's definition $k\delta$ for all index lists of equal length (theorem of section 3.2, with its proof); confirmed exhaustively up to length 4 (Rust) and length 3 (Wolfram, sympy) and on pseudo-random samples up to length 9.
2. For the author's metric on $0 < z < \pi/2$, with the conventions of sections 2.2 and 4.2: the 25 nonzero Christoffel symbols, the 156 nonzero R^{ab}_{cd} , the Ricci and Einstein tensors and the Lovelock tensors $P_{(k)}$, $A_{(k)}$ and scalars $L_{(k)}$, $k = 1, 2, 3$, exactly as listed in section 6 and in `lovelock-tensors.json`; three programs that share no code for the sums agree on every component.
3. $P_{(1)} = -4G$, $P_{(2)} = -8\mathcal{H}$ (Gauss-Bonnet), $L_{(1)} = 2R$, $L_{(2)} = 4GB$, $L_{(3)} = 8(2T_1 + \dots + T_8)$; the trace identities; $\nabla_h P_{(k)}^h{}_j = 0$; the symmetry of $A_{(k)}$; $P_{(4)} = 0$ in eight dimensions.
4. The structure: diagonal, free of $\sin^{1/3} z$, e^{a_4} and x_8 , with $x_1 = x_2 = x_3$ and $x_5 = x_6 = x_7$, and with a_4'' entering the 3-space and extra-time components with opposite signs.

12.2 Not established

1. No solution: this record gives the left-hand sides of the field equations only. Which a_4 and which sources solve them is not part of it (section 9).
2. The domain: only the patch $0 < z < \pi/2$, where $\sqrt{|g|} = \cos z$. Nothing is computed at the brane $z = \pi/2$, where $g_{88} = \cot^2 z = 0$ and $\det g = 0$, or on a mirror patch.
3. No comparison with the author's answers: the output cells of the author's notebook were never read, and the announced comparison is not recorded in **Revision/** (section 8). Agreement with the author's own Lovelock tensors is therefore not established.
4. For $k = 3$ the tensor $P_{(3)}$ is verified by three independent recomputations, the trace, the divergence and the symmetry, and the scalar $L_{(3)}$ against the classical cubic density; it is not compared with an independent closed formula for the third-order Lovelock tensor. $E_{(k)} = -P_{(k)}/2^{k+1}$ is a normalisation convention, which equals the Einstein tensor for $k = 1$ and the Gauss-Bonnet tensor for $k = 2$.
5. The unpruned sums are exact for $k = 1, 2$ (sympy) and numerical at one point (Rust); for $k = 3$ the skipped terms are confirmed by a random sample (sympy) and by the theorem of section 3.2; the Wolfram no-skip run of the diagonal $k = 3$ components is recorded in its provenance file only, and its report is not committed.
6. Only Wolfram Language 15.0.1 and the toolchains named in the provenance files were used.
7. No physical interpretation is attached to the components here; in particular nothing is said about the sign of an energy or about the stability of a solution.