

dirac16complex: a complex Grassmann spinor of $\text{Pin}(4,4)$ in an arbitrary gravitational field

Lagrangian, covariant field equations, energy-momentum tensor, canonical quantization and equations of state

dirac16complex project (with Claude Opus 5.5)

September 2026

Contents

1	Scope, claims and non-claims	5
1.1	What is proved	5
1.2	What is conventional	6
1.3	What is not claimed	6
2	Counting and notation dictionary	7
2.1	Counting and complex vector spaces	7
2.2	This document, the notebook and dirac-main	7
2.3	The scalar-field PDF	8
3	The Clifford module in the split-octonion and Clifford pictures	9
3.1	The split-octonion (notebook) construction	9
3.2	Charge matrix, expression [1] and chirality	10
3.3	The Clifford picture of dirac-main and the intertwiners	11
3.4	Verification record	12
4	Definition of dirac16complex	13
4.1	The field	13
4.2	The action of $\text{Pin}(4,4)$ and the two lifts	13
4.3	Irreducibility under $\text{Pin}(4,4)$	13
4.4	Two inequivalent 8-dimensional modules under $\text{Spin}(4,4)$	14

4.5	The spinor bundle	14
5	Canonical spin connection and covariant derivative	14
5.1	Vielbein, metric and Christoffel symbols	14
5.2	The vielbein postulate and the canonical spin connection	15
5.3	Spinor connection and covariant derivatives	15
5.4	Covariant constancy of the gammas and the divergence identity	15
5.5	Local spin transformations and curvature	16
5.6	The notebook contraction and why it is wrong	17
5.7	Test geometries	17
6	Why the notebook Lagrangian $\text{Lg}[]$ is empty for a Grassmann field	18
6.1	The notebook Lagrangian, verbatim	18
6.2	Lemmas	19
6.3	Theorem	19
6.4	Machine verification	19
6.5	Remark on commuting fields	20
7	The dirac16complex Lagrangian	20
7.1	Formula	20
7.2	WolframScript form in the notebook's variable names	21
7.3	Relation to $\text{Lg}[]$, term by term	21
7.4	Hermiticity	22
7.5	Symmetries	22
7.6	The potential: polynomials only	22
7.7	Discrete maps: the chirality map and the Pin characters	23
8	Euler-Lagrange equations in an arbitrary gravitational field	23
8.1	Derivation	23
8.2	The covariant field equations	24
8.3	Component form	24
8.4	Non-triviality in an arbitrary gravitational field	25
8.5	Diagonal vielbeins	25

9	Energy-momentum tensor operator, energies, pressure and equation of state	26
9.1	Derivation from the metric variation	26
9.2	Formula	26
9.3	Properties	27
9.4	Observer decomposition	27
9.5	Kinetic and potential energy: two splits	27
9.6	Equation of state	28
9.7	Homogeneous diagonal backgrounds	29
9.8	Side-by-side with the scalar field of the PDF	29
10	Canonical quantization in 4+4 dimensions	30
10.1	Slicing and momenta	30
10.2	Constraints and Dirac brackets	30
10.3	Equal-x4 anticommutators in curved space	31
10.4	Hamiltonian and Heisenberg equations	31
10.5	Krein structure	32
10.6	Flat-space modes and the good sector	32
10.7	Fock space, Dirac sea and normal ordering	33
10.8	Unitary and Krein-unitary symmetries	33
10.9	The U(1) current and charge	34
10.10	The extra-time sector	34
10.11	Expectation values	34
11	Verification records	35
11.1	WolframScript, algebra	35
11.2	WolframScript, geometry	35
11.3	Python, algebra	36
11.4	Python, geometry	36
11.5	Python, Grassmann demonstrations	37
11.6	Agreement between the implementations	37
11.7	Files and hashes	38
12	Reproduction	39

12.1 PowerShell	39
12.2 Git Bash	39
13 Limitations	40

Abstract

This document defines `dirac16complex`, a new 16-component fermion field on an 8-dimensional manifold whose metric has signature (4,4), and derives its physics in an arbitrary gravitational field. The components $\Psi_0, \dots, \Psi_{15}$ are complex-valued and anticommuting (complex Grassmann-odd). They carry the irreducible 16-dimensional complex representation of $\text{Pin}(4,4)$, and under $\text{Spin}(4,4)$ the same space splits into two inequivalent irreducible 8-dimensional modules. The split-octonion gamma matrices of the author's notebook and the tensor-product gamma matrices of `dirac-main` describe one Clifford module; exact integer intertwiners, unique up to scale, connect the two pictures. The field couples to gravity only through the canonical spin connection, which is fixed by the vielbein postulate (the total covariant derivative of the vielbein vanishes). We prove that the notebook's Lagrangian `Lg[]` is a pure divergence for a Grassmann field, so its Euler-Lagrange equations are empty, and we replace it by a Hermitian first-order Lagrangian. Its covariant field equations $\gamma^\mu D_\mu \Psi = (m + U'(S))\Psi$, $S = \bar{\Psi}\Psi$, contain the spin connection in every curved field: the connection is pure gauge exactly when the Riemann tensor vanishes, and the squared Dirac operator carries the curvature term cR with the measured constant $c = -1/4$. The energy-momentum tensor operator follows from the metric variation of the action; it is symmetric, Hermitian and conserved on shell, and its on-shell trace is $-mS + 7SU' - 8U$. In homogeneous diagonal backgrounds the energy density is $\rho = mS + U$ and the pressure is $p = SU' - U$ in all seven transverse directions. We define the kinetic and potential energies in two ways, give the equation of state $w = p/\rho$, and compare everything term by term with the scalar-field formulas $\rho_\phi = \frac{1}{2}\dot{\phi}^2 + V$, $P_\phi = \frac{1}{2}\dot{\phi}^2 - V$ of the reference PDF. Canonical quantization with respect to the time x_4 gives the equal-time anticommutator $\{\Psi, \Psi^\dagger\} = i(\sqrt{|g|} C \gamma^{x_4})^{-1} \delta^7$ and a Krein structure with fundamental symmetry $J = B = -iC\gamma^4$ of signature (8,8). The sector without momenta along the three extra times has an ordinary Fock space; momenta along the extra times give complex frequencies. Two independent exact implementations, one in `WolframScript` and one in `Python`, pass 153 of 153 checks in five reports. Their cross-comparison of 47 algebra and 84 geometry measurements finds no disagreement.

1 Scope, claims and non-claims

1.1 What is proved

Every numbered result below is backed by named checks in the five Stage-1 reports:

```
artifacts/dirac16complex/arbitrary-field/wolfram-algebra-report.json
artifacts/dirac16complex/arbitrary-field/wolfram-geometry-report.json
artifacts/dirac16complex/arbitrary-field/python-algebra-report.json
artifacts/dirac16complex/arbitrary-field/python-geometry-report.json
artifacts/dirac16complex/arbitrary-field/grassmann-demo-report.json
```

The check names are quoted in a "checks" block after each result, and Section 11 lists all of them. Numbers are copied from these reports. A statement marked "derived" follows from the checked results by the argument given, but is not itself a machine check.

- **Algebra (exact).** The Clifford relations, the charge matrix, expression [1], the chirality, the representation theory (commutants, intertwiners, invariant forms), the Krein signatures and the subgroup counts are exact integer or Gaussian-rational computations with the actual 16 by 16 matrices. Where a statement depends on parameters (masses and momenta in Section 10), the text says whether it was proved for symbolic values or checked at exact sample values.

- **Geometry (general proof, exact test points).** The statements about an arbitrary gravitational field are proved in this document by general arguments. They are then checked with exact arithmetic in three test geometries (Section 5.7): G1, a generic non-diagonal vielbein at three rational points; G2, the primordial family of Stage 2 (checked by Wolfram at three exact points and by Python at a fully symbolic point); and G3, homogeneous diagonal (Bianchi-I type) frames at three times.
- **Grassmann statements** are checked in genuine Grassmann algebras with up to 1440 odd generators (grassmann-demo-report.json), and the Euler-Lagrange checks of the Wolfram geometry verifier also use a genuine Grassmann algebra.

1.2 What is conventional

The following are choices, not results:

- counting from 0;
- the tangent metric $\eta = \text{diag}(+1, +1, +1, +1, -1, -1, -1, -1)$, with x_4 as the evolution time;
- the split-octonion basis of the notebook as the primary basis;
- the Dirac adjoint $\bar{\Psi} = \Psi^\dagger C$ with $C = \sigma_{16}$;
- the symmetrized form of the kinetic term and the default potential $U(S) = \frac{\lambda}{2} S^2$;
- the sign of $T_{\mu\nu}$, fixed so that T_{44} is the energy density;
- the two definitions of kinetic and potential energy in Section 9.5;
- units with $\hbar = 1$.

1.3 What is not claimed

1. No observational claim is made. The signature (4,4) is not the signature of observed spacetime, and nothing here is fitted to data.
2. The dark-energy and dark-matter questions are not answered in this document. Section 9 supplies the exact energy density, pressures and equation of state that such a study needs; the numerical investigation is separate work.
3. Perturbative stability of the classical field equations is not studied.
4. The quantization is canonical and formal. The state space carries an indefinite (Krein) metric (Section 10.5), the modes with momentum along the extra times have complex frequencies (Section 10.10), and no interacting Fock space or renormalization is constructed.
5. The homogeneous-sector formulas of Section 9.7 treat the bilinears as classical (mean-field) quantities.
6. The pairing of the masses $\pm m$ by the chirality map (Section 7.7) is a structural property of the equations, not a claim that universes of masses $\pm M$ are created in pairs.

2 Counting and notation dictionary

2.1 Counting and complex vector spaces

As the task prescribes, everything is counted from 0. A complex vector space is a set whose elements can be added and multiplied by complex numbers while satisfying the usual distributive rules. A basis $(e_0, \dots, e_7)$ of an 8-dimensional complex vector space is an ordered list in which every vector v has one and only one expansion $v = v_0 e_0 + v_1 e_1 + \dots + v_7 e_7$; the coordinate column of v is $(v_0, \dots, v_7)^T$. The dimension counts basis vectors, not the number of vectors: an 8-dimensional complex space contains infinitely many vectors. The spinor space $\mathbb{C}^{16}$ of `dirac16complex` is used in the same way, with the basis $(e_0, \dots, e_{15})$ and coordinate columns $(\Psi_0, \dots, \Psi_{15})^T$.

- Coordinates $x = \{x_0, x_1, \dots, x_7\}$; curved indices $\mu, \nu, \rho, \sigma, \lambda = 0, \dots, 7$; frame indices $a, b, c = 0, \dots, 7$; spinor indices $0, \dots, 15$.
- Roles of the coordinates (from the notebook): x_0 is a hidden space direction, x_1, x_2, x_3 are ordinary 3-space, x_4 is the evolution time, and x_5, x_6, x_7 are three extra times. Frame directions 0 to 3 are space-like ($\eta_{aa} = +1$) and 4 to 7 are time-like ($\eta_{aa} = -1$).
- A gamma matrix with a numeric upper index, $\gamma^0, \dots, \gamma^7$, is always the constant frame matrix. Curved gammas carry a Greek index, $\gamma^\mu = e_a^\mu \gamma^a$; with a numeric value they are written γ^{x_4} and so on. Lowered gammas $\gamma_\mu = g_{\mu\nu} \gamma^\nu$ are curved.
- Mathematica lists are 1-based. A notebook entry such as `SAB[[a,b]]` with $a, b = 1, \dots, 8$ is $S^{a-1 b-1}$ in this document.
- The only other 1-based indices are those of the notebook's 4 by 4 building blocks in Section 3.1, which are labelled as such.

2.2 This document, the notebook and dirac-main

The notebook is the file

`Pair_Creation_of_Universes_WaveFunctionOfUniverse-4+4-Einstein-Lovelock-Nash.nb`

in the repository root (read only, never modified). Cell numbers refer to the cell numbering of the project's text dump of the notebook. `dirac-main` is the reference implementation of the Clifford and split-octonion pictures (read only).

This document	Notebook (WolframScript)	dirac-main
$x_0, \dots, x_7$	X (cell 61)	$x^0, \dots, x^7$
η_{ab}	η_{4488}	η_{ab}
frame gamma γ^a	$(T16^{\wedge}A)[a]$	γ^a (list below)
$\gamma^\mu = e_a^\mu \gamma^a$	$(T16^{\wedge}\alpha)[\mu]$	$\gamma^\mu = e_a^\mu \gamma^a$
$C = \sigma_{16}$	σ_{16}	C (tensor picture)
$S^{ab} = \frac{1}{4}[\gamma^a, \gamma^b]$	$SAB[[a+1,b+1]]$	$\Sigma_{ab} = \frac{1}{4}[\gamma_a, \gamma_b]$

This document	Notebook (WolframScript)	dirac-main
$\omega_\mu^a{}_b$	$\omega\text{mat}[[\mu+1,\text{a}+1,\text{b}+1]]$	$\omega_\mu^a{}_b$
$\omega_{\mu ab} = \eta_{ac}\omega_\mu^c{}_b$	not formed	$\omega_{\mu ab}$
$\Omega_\mu = \frac{1}{2}\omega_{\mu ab}S^{ab}$	(Q1/2) ωmat SAB, see 5.6	$\frac{1}{8}\omega_{\mu ab}[\gamma^a, \gamma^b]$
$D_\mu\Psi = \partial_\mu\Psi + \Omega_\mu\Psi$	$\text{D}[\Psi16, \text{X}[[\mu+1]]] + \dots$	$D_\mu\psi$
Ψ , complex Grassmann	$\Psi16$, commuting	ψ , commuting
$\bar{\Psi} = \Psi^\dagger C$	$\text{Transpose}[\Psi16].\sigma16$	$\bar{\psi} = \psi^T C$
$\sqrt{ g }$	$\text{Sqrt}[\text{detgg}]$	$\sqrt{ g }$
mass m	$-HM$	none
$\Psi_0, \dots, \Psi_7$	$\Psi16\text{upper}$	image of P_-
$\Psi_8, \dots, \Psi_{15}$	$\Psi16\text{lower}$	image of P_+

Further notebook names used in `Lg[]` (Section 6.1): Q1 is the notebook's book-keeping switch for the spin connection (1 switches it on); sg is the list of rules that turns the metric symbols into diagonal functions of (x_0, x_4) (cell 269); detgg is `Det[g4488 /. sg]` (cell 275); constraintVars is the list of positivity assumptions used by Simplify (cell 73); H and M are the notebook's inverse length and mass.

The notebook's $\Psi16$ is the column of the 16 functions `f16[k]`, which the notebook treats as real commuting functions of (x_0, x_4) . The dirac-main gammas form the ordered list $(\gamma_1^+, \gamma_2^+, \gamma_3^+, \gamma_4^+, \gamma_1^-, \gamma_2^-, \gamma_3^-, \gamma_4^-)$ of Section 3.3, with 1-based labels: γ_k^+ is the frame matrix γ^{k-1} and γ_k^- is γ^{k+3} ($k = 1, \dots, 4$), so the evolution-time generator γ^4 is γ_1^- .

2.3 The scalar-field PDF

The reference on the equation of state is the PDF file whose name begins with "Gmail - w = equation of state parameter" (repository root, read only). It treats a canonical scalar field ϕ in four dimensions with metric signature $(+, -, -, -)$ and $\mathcal{L}_\phi = \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - V(\phi)$.

PDF	This document
signature $(+, -, -, -)$, time first	signature (4,4), time x_4 ; the PDF metric is $-g$ on the coordinates x_1, x_2, x_3, x_4
$\mathcal{L}_\phi = \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - V$	the same function: $\mathcal{L}_\phi = -\frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi - V$
$T^\mu{}_\nu = \partial^\mu\phi\partial_\nu\phi - \delta^\mu{}_\nu\mathcal{L}_\phi$	$T_{\mu\nu} = \partial_\mu\phi\partial_\nu\phi + g_{\mu\nu}\mathcal{L}_\phi$ (same covariant components)
$\rho_\phi = \frac{1}{2}\dot{\phi}^2 + V$	$\rho = T_{44}$
$P_\phi = \frac{1}{2}\dot{\phi}^2 - V$	$p_{(i)} = T^i{}_i$ (no sum), $i \neq 4$
$w_\phi = P_\phi/\rho_\phi$	$w = p/\rho$
$\text{KE} = \frac{1}{2}\dot{\phi}^2$, $\text{PE} = V$	KE_L , PE_L and KE_H , PE_H (Section 9.5)
$\dot{\phi} = d\phi/dt$	$\partial_4 = \partial/\partial x_4$
CPL: $w(a) = w_0 + w_a(1 - a)$	not used; a is the PDF's scale factor

In both conventions a homogeneous scalar field has $\rho = \frac{1}{2}\dot{\phi}^2 + V$ and $p = \frac{1}{2}\dot{\phi}^2 - V$. The two metrics differ by an overall sign on the 4-dimensional subspace; the covariant tensor $T_{\mu\nu}$ is the same in both,

and the mixed tensor $T^\mu{}_\nu$ changes sign. (These scalar-field formulas are standard and are quoted for comparison; they are not part of the machine checks.)

Sign hazard in the PDF. The PDF writes $w(a) = w_0 + w_a(1 - a)$, so $dw/da = -w_a$. Since a grows with time, " w increases with time" means $w_a < 0$, and " w decreases towards -1 " means $w_a > 0$. The PDF's table of thawing and freezing models nevertheless labels thawing (" w starts near -1 and increases over time") with $w_a > 0$ and freezing (" w starts less negative and approaches -1 ") with $w_a < 0$. That is the reverse of what its own formula implies. Wherever thawing or freezing is meant in this project, the formula-consistent signs are used: thawing has $w_a < 0$ and freezing has $w_a > 0$.

3 The Clifford module in the split-octonion and Clifford pictures

3.1 The split-octonion (notebook) construction

The notebook builds its gamma matrices from 4 by 4 blocks. With 1-based $p, q \in \{1, 2, 3, 4\}$ and $h \in \{1, 2, 3\}$,

$$\begin{aligned} Q_a[h]_{pq} &= \epsilon_{hpq4}, & Q_b[h]_{pq} &= \delta_{p4}\delta_{qh} - \delta_{ph}\delta_{q4}, \\ s_4[h] &= Q_a[h] - Q_b[h], & t_4[h] &= Q_a[h] + Q_b[h], \\ \tau_0 &= I_8, & \tau_h &= \begin{pmatrix} 0 & s_4[h] \\ s_4[h] & 0 \end{pmatrix}, & \tau_{7-h} &= \begin{pmatrix} 0 & t_4[h] \\ -t_4[h] & 0 \end{pmatrix}, \\ \tau_7 &= \tau_1\tau_2\tau_3\tau_4\tau_5\tau_6, & \sigma &= \begin{pmatrix} 0 & I_4 \\ I_4 & 0 \end{pmatrix}, & \bar{\tau}_0 &= I_8, & \bar{\tau}_a &= \sigma \tau_a^T \sigma \quad (a = 1, \dots, 7), \\ \gamma^a &= \begin{pmatrix} 0 & \bar{\tau}_a \\ \tau_a & 0 \end{pmatrix} & (a = 0, \dots, 7), \end{aligned}$$

where ϵ_{hpq4} is Mathematica's `Signature[{h,p,q,4}]`. In the notebook γ^a is `(T16^A)[a]` and $\bar{\tau}_a$ is `OverBar[tau][a]`.

Result 3.1. The γ^a are integer matrices with $\{\gamma^a, \gamma^b\} = 2\eta^{ab}I_{16}$ for all 64 ordered pairs; $(\gamma^a)^2 = +1$ for $a < 4$ and -1 for $a \geq 4$. The matrix γ^a is symmetric for $a < 4$ and antisymmetric for $a \geq 4$.

checks

```
ALG_clifford, ALG_gammaTransposeSymmetry: wolfram-algebra, python-algebra
ALG_cliffordRelations: wolfram-geometry, python-geometry
```

Result 3.2 (faithfulness). The 256 ordered monomials $\gamma^{a_1} \dots \gamma^{a_k}$ ($a_1 < \dots < a_k$) are linearly independent (rank 256), and the 128 even ones have rank 128. The gammas therefore generate the full matrix algebra, $\text{Cl}(4, 4) \cong \text{Mat}_{16}(\mathbb{R})$.

checks

```
ALG_faithful: wolfram-algebra, python-algebra
```

3.2 Charge matrix, expression [1] and chirality

The charge (Dirac-adjoint) matrix is the notebook's σ_{16} :

$$C := \sigma_{16} = \gamma^0 \gamma^1 \gamma^2 \gamma^3 = \begin{pmatrix} -\sigma & 0 \\ 0 & \sigma \end{pmatrix}, \quad C^T = C, \quad C^2 = I_{16}.$$

Result 3.3. $C = \gamma^0 \gamma^1 \gamma^2 \gamma^3 = \text{diag}(-\sigma, \sigma)$, C is symmetric with $C^2 = 1$, its characteristic polynomial is $(x-1)^8(x+1)^8$, and its signature is (8,8).

checks

```
ALG_chargeMatrix: wolfram-algebra, python-algebra, wolfram-geometry
ALG_sigma16EqualsGamma0123, ALG_CSymmetricInvolution: python-geometry
```

Expression [1] of the task, verbatim in WolframScript:

```
\[Sigma]16.(T16^A)[#]==-Transpose[\[Sigma]16.(T16^A)[#]]&/@Range[0,7]
```

It states $(C\gamma^a)^T = -C\gamma^a$ for $a = 0, \dots, 7$.

Result 3.4 (expression [1]). Expression [1] evaluates to {True, True, True, True, True, True, True, True}. The Wolfram verifier evaluates it twice: with the package's matrices, and literally, from the 11 verbatim notebook definition cells evaluated in an isolated context (the notebook's Symbolize'd names $T16^A$ and $\text{OverBar}[\tau]$ are reproduced in two independent ways, and both reproduce the package's η , σ , τ , $\bar{\tau}$, γ^a and σ_{16} exactly).

checks

```
ALG_expression1: wolfram-algebra, python-algebra
ALG_expression1CgammaAntisymmetric: python-geometry
```

Proof. $(C\gamma^a)^T = (\gamma^a)^T C$. For $a < 4$, γ^a is symmetric and anticommutes with three of the four factors of $C = \gamma^0 \gamma^1 \gamma^2 \gamma^3$, so $\gamma^a C = -C\gamma^a$. For $a \geq 4$, γ^a is antisymmetric and anticommutes with all four factors, so $(\gamma^a)^T C = -\gamma^a C = -C\gamma^a$. In both cases $(C\gamma^a)^T = -C\gamma^a$.

Result 3.5. With $S^{ab} = \frac{1}{4}[\gamma^a, \gamma^b]$: $(CS^{ab})^T = -CS^{ab}$ for all 64 pairs; $C\{\gamma^c, S^{ab}\}$ is symmetric and $C[\gamma^c, S^{ab}]$ is antisymmetric for all 512 triples; $[S^{ab}, \gamma^c] = \eta^{bc}\gamma^a - \eta^{ac}\gamma^b$. The Python algebra checker counts 476 verified identities of this kind.

checks

```
ALG_spinTransposeProperties: wolfram-algebra, python-algebra
ALG_CSabAntisymmetric, ALG_CAnticommutatorGammaSSymmetric,
ALG_CCommutatorGammaSAntisymmetric, ALG_SabGammaCommutator: python-geometry
```

Result 3.6 (chirality). $\gamma^8 := \gamma^0 \gamma^1 \dots \gamma^7 = \text{diag}(-I_8, I_8)$, $(\gamma^8)^2 = 1$, γ^8 anticommutes with every γ^a and commutes with every S^{ab} and with C . The upper components $\Psi_0, \dots, \Psi_7$ (the notebook's $\Psi_{16\text{upper}}$) have chirality -1 and the lower components $\Psi_8, \dots, \Psi_{15}$ ($\Psi_{16\text{lower}}$) have chirality $+1$. The chiral projectors are $P_{\mp} = \frac{1}{2}(1 \mp \gamma^8)$.

checks

ALG_chirality: wolfram-algebra, python-algebra

ALG_chiralityDiag: python-geometry

3.3 The Clifford picture of dirac-main and the intertwiners

dirac-main builds its gammas from the 2 by 2 matrices

$$P = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad N = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad G = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

$$\hat{\gamma}_k^+ = G^{\otimes(k-1)} \otimes P \otimes I_2^{\otimes(4-k)}, \quad \hat{\gamma}_k^- = G^{\otimes(k-1)} \otimes N \otimes I_2^{\otimes(4-k)} \quad (k = 1, 2, 3, 4),$$

ordered $(\hat{\gamma}_1^+, \dots, \hat{\gamma}_4^+, \hat{\gamma}_1^-, \dots, \hat{\gamma}_4^-) = (\hat{\gamma}^0, \dots, \hat{\gamma}^7)$, with charge matrix $C_{\text{dm}} = \hat{\gamma}^0 \hat{\gamma}^1 \hat{\gamma}^2 \hat{\gamma}^3$ and volume element $G \otimes G \otimes G \otimes G$. Its split-octonion picture uses the Zorn vector-matrix model: a vector $x = (x_0, \dots, x_7)$ is the Zorn matrix (a, u, v, b) with $a = x_0 + x_4$, $u_i = x_i + x_{i+4}$, $v_i = -x_i + x_{i+4}$ ($i = 1, 2, 3$) and $b = x_0 - x_4$, and

$$(a, u, v, b)(c, r, w, d) = (ac + u \cdot w, ar + du - v \times w, cv + bw + u \times r, v \cdot r + bd),$$

$$\Gamma(x) = \begin{pmatrix} 0 & L_{\bar{x}} \\ L_x & 0 \end{pmatrix}, \quad \Gamma^a = \Gamma(e_a),$$

where $L_x(y) = xy$ and $\bar{x} = (x_0, -x_1, \dots, -x_7)$.

Result 3.7 (Clifford picture). The tensor matrices satisfy the Clifford relations with the same η and equal dirac-main's committed cl44-seed generators. The intertwiner equation $\hat{\gamma}^a K = K \gamma^a$ ($a = 0, \dots, 7$) has a solution space of dimension exactly 1. Its primitive integer generator K_{clifford} (first nonzero entry positive, coprime entries) has rank 16, so it is invertible, and it satisfies $KCK^{-1} = C_{\text{dm}}$ and $K\gamma^8 K^{-1} = +G \otimes G \otimes G \otimes G$. The notebook's chirality -1 block $\Psi_0, \dots, \Psi_7$ maps onto the dirac-main basis indices $\{1, 2, 4, 7, 8, 11, 13, 14\}$ and the $+1$ block onto $\{0, 3, 5, 6, 9, 10, 12, 15\}$. K_{clifford} is not a signed permutation: its entries are $0, \pm 1$ with four nonzero entries per row.

checks

ALG_cliffordPictureIntertwiner: wolfram-algebra, python-algebra

K_clifford =

```
[ 0  0  0  0  0  0  0  0  0  1 -1  0  0  1  1  0  0 ]
[-1  1  0  0  1  1  0  0  0  0  0  0  0  0  0  0 ]
[ 0  0 -1  1  0  0 -1 -1  0  0  0  0  0  0  0  0 ]
[ 0  0  0  0  0  0  0  0  0  0  0  1 -1  0  0 -1 -1 ]
[ 0  0  1  1  0  0 -1  1  0  0  0  0  0  0  0  0  0 ]
[ 0  0  0  0  0  0  0  0  0  0  0 -1 -1  0  0 -1  1 ]
[ 0  0  0  0  0  0  0  0 -1 -1  0  0  1 -1  0  0  0 ]
[ 1  1  0  0  1 -1  0  0  0  0  0  0  0  0  0  0  0 ]
[ 1 -1  0  0  1  1  0  0  0  0  0  0  0  0  0  0  0 ]
[ 0  0  0  0  0  0  0  0 -1  1  0  0  1  1  0  0  0 ]
[ 0  0  0  0  0  0  0  0  0  0 -1  1  0  0 -1 -1  0 ]
```

```
[ 0  0  1 -1  0  0 -1 -1  0  0  0  0  0  0  0  0 ]
[ 0  0  0  0  0  0  0  0  0  0  1  1  0  0 -1  1 ]
[ 0  0 -1 -1  0  0 -1  1  0  0  0  0  0  0  0  0 ]
[-1 -1  0  0  1 -1  0  0  0  0  0  0  0  0  0  0 ]
[ 0  0  0  0  0  0  0  0  1  1  0  0  1 -1  0  0 ]
```

Result 3.8 (split-octonion picture). The Zorn product has the unit e_0 , satisfies $N(xy) = N(x)N(y)$ for the norm $N(x) = \eta(x, x)$, has conjugation reversing products, and is not associative (the associator of e_1, e_2, e_4 is $2e_7$). It has 64 nonzero structure constants and equals dirac-main's split-octonion.json multiplication tensor, and the Γ^a equal dirac-main's triality44.json octonion Clifford generators. The intertwiner equation $\gamma^a K = K\Gamma^a$ has a solution space of dimension exactly 1; its primitive generator $K_octonion$ has rank 16 and is block diagonal, $K_{octonion} = \text{diag}(Q, Q)$ with the matrix Q below, $Q^T Q = 2I_8$ and $Q^T \sigma Q = 2\eta$. Hence $\tau_a = QL_{e_a}Q^{-1}$ and $\bar{\tau}_a = QL_{\bar{e}_a}Q^{-1}$ for every a . The notebook's τ_a equal the octonion multiplications L_{e_a} only for $a = 0$, and no signed permutation relates them. Since $Q/\sqrt{2}$ is orthogonal, the notebook's 8-space basis is a null (light-cone) basis of the split octonions, in which the octonion norm η appears as $\sigma/2$. Finally, the product $K_clifford K_octonion$, made primitive, equals dirac-main's canonicalCliffordIntertwiner.

checks

ALG_octonionPictureIntertwiner: wolfram-algebra, python-algebra

```
Q =
[ 1  0  0  0  0  0  0  0 -1 ]
[ 0  0  0 -1 -1  0  0  0  0 ]
[ 0  0  1  0  0  1  0  0  0 ]
[ 0 -1  0  0  0  0  1  0  0 ]
[ 1  0  0  0  0  0  0  0  1 ]
[ 0  0  0 -1  1  0  0  0  0 ]
[ 0  0  1  0  0 -1  0  0  0 ]
[ 0 -1  0  0  0  0 -1  0  0 ]
```

Compatibility. The three sets of matrices, γ^a (notebook), $\hat{\gamma}^a$ (tensor) and Γ^a (octonion), are one Clifford module written in three bases: $\hat{\gamma}^a = K_{clifford}\gamma^a K_{clifford}^{-1}$ and $\Gamma^a = K_{octonion}^{-1}\gamma^a K_{octonion}$. Each intertwiner is unique up to a scalar, as Schur's lemma requires for an irreducible module (Section 4.3), and the notebook-to-tensor map sends C to C_{dm} and the chirality blocks to the chirality blocks. The intertwiners change coordinates only; they do not make the octonion product associative.

3.4 Verification record

Both algebra implementations build the matrices independently: the Wolfram package wolfram/Dirac16ComplexAlgebra.wl and the Python module scripts/d16c_exact.py (exact rationals and Gaussian rationals, standard library only). Python writes the exact fixture algebra-fixture.json (η , σ , τ_a , $\bar{\tau}_a$, the gammas of all three pictures, C , the chirality, the 28 S^{ab} , B and both intertwiners), and the Wolfram verifier compares every fixture matrix with its own construction: 75 of 75 matrices agree by name, none disagree and none is unmatched (ALG_fixtureAgreement in wolfram-algebra-report.json). In the other direction, the Python checker reads the Wolfram report and requires the same $K_clifford$, $K_octonion$ and chirality, the same verdict for all 20 check names both reports contain, and agreement of 47 shared measurements; all 47 agree (ALG_wolframAgreement in python-algebra-report.json).

4 Definition of dirac16complex

4.1 The field

Definition. dirac16complex is the field

$$\Psi = (\Psi_0, \Psi_1, \dots, \Psi_{15})^T,$$

whose components are complex-valued Grassmann-odd (anticommuting) fields on an 8-dimensional manifold M with a metric of signature (4,4):

$$\begin{aligned} \Psi_a \Psi_b &= -\Psi_b \Psi_a, & \Psi_a \Psi_b^* &= -\Psi_b^* \Psi_a, & \Psi_a^* \Psi_b^* &= -\Psi_b^* \Psi_a^*, \\ (\theta_1 \theta_2)^* &= \theta_2^* \theta_1^*. \end{aligned}$$

Complex conjugation is an involution of the Grassmann algebra that reverses the order of products. We write $\Psi^\dagger = (\Psi^*)^T$ and $\bar{\Psi} = \Psi^\dagger C$. The components transform under $\text{Pin}(4,4)$ through the Clifford module of Section 3 (Section 4.2); for the determinant-1 transformations, $\text{Spin}(4,4)$, the module splits into the two chirality blocks (Section 4.4). The conjugation rules, including the product rule and the involution property, are checked in an explicit Grassmann algebra (`GR_conjugationRules`).

4.2 The action of $\text{Pin}(4,4)$ and the two lifts

$\text{Pin}(4,4)$ is the group generated, inside $\text{Cl}(4,4)$, by the unit vectors $u = u_a \gamma^a$ with $n(u) := \eta(u, u) = \pm 1$, so that $u^2 = n(u)$. It acts on dirac16complex by Clifford multiplication, $\rho(u)\Psi = u\Psi$. $\text{Spin}(4,4)$ is the subgroup of products of an even number of unit vectors. $\text{Pin}(4,4)$ is the double cover of the orthogonal group $\text{O}(4,4)$ and $\text{Spin}(4,4)$ is the double cover of $\text{SO}(4,4)$. Vectors transform by the twisted adjoint action $v \mapsto \alpha(g)vg^{-1}$, where α is the parity automorphism; a unit vector u then acts as the reflection in the hyperplane orthogonal to u , and the kernel is $\{\pm 1\}$ (standard facts, not machine-checked). The untwisted adjoint action $v \mapsto gvg^{-1}$ gives, for a unit vector, minus that reflection. Because -1_8 lies in the identity component $\text{SO}_0(4,4)$ (it is the product of rotations by π in the planes (0,1), (2,3), (4,5) and (6,7)), the untwisted action also maps $\text{Pin}(4,4)$ onto $\text{O}(4,4)$. For every tested unit vector the untwisted matrix is exactly in $\text{O}(4,4)$ (`ALG_pinLiftCharacter`). The choice of lift matters for the sign characters of Section 7.7.

4.3 Irreducibility under $\text{Pin}(4,4)$

Theorem 4.1. $\mathbb{C}^{16}$, with $\text{Pin}(4,4)$ acting through the gamma matrices, is an irreducible complex representation of $\text{Pin}(4,4)$.

Proof. Every monomial $\gamma^{a_1} \dots \gamma^{a_k}$ is, up to sign, a product of unit vectors, so the linear span of $\text{Pin}(4,4)$ is the span of the 256 monomials. By Result 3.2 these are linearly independent, so they span $\text{Mat}_{16}(\mathbb{R})$, and over $\mathbb{C}$ they span $\text{Mat}_{16}(\mathbb{C})$. A subspace of $\mathbb{C}^{16}$ invariant under $\text{Pin}(4,4)$ is invariant under all of $\text{Mat}_{16}(\mathbb{C})$, hence it is 0 or $\mathbb{C}^{16}$. Equivalently (Schur's lemma), the commutant $\{X : X\gamma^a = \gamma^a X \ \forall a\}$ consists of the scalars.

Result 4.1. The commutant of the eight gammas on $\mathbb{C}^{16}$ has dimension exactly 1 and consists of the scalar matrices. The dimension is computed over $\mathbb{Q}$; the coefficient matrix of the equations is rational, so its rank, and hence the solution dimension, is the same over $\mathbb{C}$.

checks

ALG_pinIrreducibleComplex, ALG_faithful: wolfram-algebra, python-algebra

4.4 Two inequivalent 8-dimensional modules under $\text{Spin}(4,4)$

Theorem 4.2. Under $\text{Spin}(4,4)$, $\mathbb{C}^{16} = \mathbb{C}_-^8 \oplus \mathbb{C}_+^8$ (the images of P_- and P_+), and the two summands are irreducible and inequivalent.

Proof. $\text{Spin}(4,4)$ is generated by even elements, and every even element commutes with γ^8 , so the two chirality blocks are invariant. The even monomials restricted to each block span an algebra of rank $64 = \dim \text{End}(\mathbb{C}^8)$, so each block is irreducible, and each block's commutant is one-dimensional. If the two blocks were equivalent, the commutant on $\mathbb{C}^{16}$ would be $\text{Mat}_2(\mathbb{C})$, of dimension 4. It has dimension 2, spanned by P_- and P_+ , and the intertwiners between the blocks form the zero space in both directions, so the blocks are inequivalent.

Result 4.2. The commutant of the 28 spin generators S^{ab} has dimension 2 and is spanned by P_- and P_+ ; every S^{ab} and every even monomial is block diagonal; the block commutants have dimensions (1, 1); the even-algebra ranks on the blocks are (64, 64); the cross-intertwiner dimensions are (0, 0).

checks

ALG_spinDecomposition: wolfram-algebra, python-algebra

The same statements hold for the identity component, which is generated by the $\exp(\theta S^{ab})$, and for the full group, whose elements lie in the even algebra.

4.5 The spinor bundle

On a manifold with a spin structure $P_{\text{Spin}}(M)$ for the (4,4) frame bundle (assumed to exist), dirac16complex is a section, with values in the odd part of a complex Grassmann algebra, of the bundle

$$S_{\mathbb{C}} = P_{\text{Spin}}(M) \times_{\rho} (\mathbb{C} \otimes \Delta_{\mathbb{R}}), \quad \Delta_{\mathbb{R}} = \Delta_- \oplus \Delta_+,$$

where $\Delta_{\mathbb{R}} \cong \mathbb{R}^{16}$ is the real module on which the real gammas act and $\Delta_{\mp}$ are its chirality halves. The spin connection of Section 5 preserves Δ_- and Δ_+ because every Ω_{μ} is even.

5 Canonical spin connection and covariant derivative

5.1 Vielbein, metric and Christoffel symbols

The gravitational field is a vielbein $e_{\mu}^a(x)$ (row index μ) with inverse e_a^{μ} :

$$g_{\mu\nu} = e_\mu^a \eta_{ab} e_\nu^b, \quad \gamma^\mu = e_a^\mu \gamma^a, \quad \{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu},$$

$$\Gamma^\rho_{\mu\nu} = \frac{1}{2}g^{\rho\sigma}(\partial_\mu g_{\nu\sigma} + \partial_\nu g_{\mu\sigma} - \partial_\sigma g_{\mu\nu}).$$

The Christoffel symbols are those of the torsion-free metric-compatible (Levi-Civita) connection.

5.2 The vielbein postulate and the canonical spin connection

The canonical spin connection is defined by the vielbein postulate: the total covariant derivative of the vielbein, built with the Christoffel symbols on the curved index and the spin connection on the frame index, is zero,

$$\nabla_\mu e_\nu^a := \partial_\mu e_\nu^a - \Gamma^\rho_{\mu\nu} e_\rho^a + \omega_\mu^a{}_b e_\nu^b = 0.$$

Multiplying by the inverse vielbein solves it uniquely:

$$\omega_\mu^a{}_b = e_b^\nu (\Gamma^\rho_{\mu\nu} e_\rho^a - \partial_\mu e_\nu^a), \quad \omega_{\mu ab} := \eta_{ac} \omega_\mu^c{}_b = -\omega_{\mu ba}.$$

The antisymmetry in (a, b) follows from metric compatibility.

Result 5.1. In every test geometry all $8 \cdot 8 \cdot 8 = 512$ components of the vielbein postulate vanish together with their 4096 first derivatives, and $\omega_{\mu ab} = -\omega_{\mu ba}$. The Python checker computes ω by two routes, from the Christoffel symbols and from the anholonomy coefficients, and they agree. The number of nonzero $\omega_{\mu ab}$ is 448 at each G1 point and 24 in G2.

checks

```
GEO_vielbeinPostulate_G1, GEO_vielbeinPostulate_G2,
GEO_omegaAntisymmetry_G1, GEO_omegaAntisymmetry_G2: wolfram-geometry, python-geometry
```

5.3 Spinor connection and covariant derivatives

$$\Omega_\mu = \frac{1}{2}\omega_{\mu ab}S^{ab} = \frac{1}{8}\omega_{\mu ab}[\gamma^a, \gamma^b], \quad D_\mu \Psi = \partial_\mu \Psi + \Omega_\mu \Psi, \quad D_\mu \bar{\Psi} = \partial_\mu \bar{\Psi} - \bar{\Psi} \Omega_\mu.$$

The sums run over all ordered pairs (a, b) ; with a sum over $a < b$ only, the coefficient of $[\gamma^a, \gamma^b]$ is $\frac{1}{4}$. The rule for $\bar{\Psi}$ follows from $\Psi^\dagger \Omega_\mu^\dagger C = -\bar{\Psi} \Omega_\mu$, which holds because S^{ab} is real and $(S^{ab})^T C = -C S^{ab}$ (Result 3.5).

5.4 Covariant constancy of the gammas and the divergence identity

Result 5.2. The curved gammas are covariantly constant,

$$D_\mu \gamma^\nu := \partial_\mu \gamma^\nu + \Gamma^\nu_{\mu\lambda} \gamma^\lambda + [\Omega_\mu, \gamma^\nu] = 0,$$

in all 64 pairs (μ, ν) of every test geometry. This identity is equivalent to the vielbein postulate, since $[\Omega_\mu, \gamma^b] = -\omega_\mu^b{}_c \gamma^c$ by Result 3.5.

checks

GEO_gammaCovariantConstancy_G1,
GEO_gammaCovariantConstancy_G2: wolfram-geometry, python-geometry

Result 5.3 (divergence identity). Contracting $D_\mu \gamma^\mu = 0$ and using $\partial_\mu \sqrt{|g|} = \sqrt{|g|} \Gamma^\rho_{\rho\mu}$ gives

$$\partial_\mu (\sqrt{|g|} \gamma^\mu) = \sqrt{|g|} [\gamma^\mu, \Omega_\mu],$$

which holds exactly in every test geometry; the Wolfram check also verifies $\partial_\mu \sqrt{|g|} = \sqrt{|g|} \Gamma^\rho_{\rho\mu}$. This identity is used twice: in the proof that $\text{Lg}[]$ is empty (Section 6) and in the derivation of the field equations (Section 8).

checks

GEO_divergenceIdentity_G1, GEO_divergenceIdentity_G2: wolfram-geometry, python-geometry

5.5 Local spin transformations and curvature

Under a local spin transformation $R(x)$,

$$\Psi \rightarrow R\Psi, \quad \Omega_\mu \rightarrow R\Omega_\mu R^{-1} - (\partial_\mu R)R^{-1}, \quad D_\mu \Psi \rightarrow R D_\mu \Psi.$$

Result 5.4. For a finite local transformation $R = \gamma(n)\gamma(m)$, $\gamma(n) := n_a \gamma^a$, built from exact rational unit vector fields $n(x)$ and $m(x)$, recomputing Ω from the rotated frame gives exactly $\Omega' = R\Omega R^{-1} - (\partial R)R^{-1}$, $\gamma'^\mu = R\gamma^\mu R^{-1}$ and an unchanged metric; for $\eta(n, n)\eta(m, m) = +1$ the Lagrangian of Section 7 is unchanged (value and first derivatives). A first-order check with exact dual numbers ($\kappa^2 = 0$) gives zero first-order change of the Lagrangian.

checks

LAG_localSpinInvariance_G1, LAG_localSpinInvariance_G2: wolfram-geometry

The curvature of the spinor connection is

$$F_{\mu\nu} = \partial_\mu \Omega_\nu - \partial_\nu \Omega_\mu + [\Omega_\mu, \Omega_\nu] = +\frac{1}{2} R_{ab\mu\nu} S^{ab},$$

$$R_{ab\mu\nu} := \eta_{ac} e_\rho{}^c R^\rho{}_{\sigma\mu\nu} e_b{}^\sigma,$$

with $R^\rho{}_{\sigma\mu\nu} = \partial_\mu \Gamma^\rho_{\nu\sigma} - \partial_\nu \Gamma^\rho_{\mu\sigma} + \Gamma^\rho_{\mu\lambda} \Gamma^\lambda_{\nu\sigma} - \Gamma^\rho_{\nu\lambda} \Gamma^\lambda_{\mu\sigma}$, Ricci tensor $R_{\sigma\nu} = R^\rho{}_{\sigma\rho\nu}$ and scalar curvature $R = g^{\sigma\nu} R_{\sigma\nu}$.

Result 5.5. The form with $+\frac{1}{2}$ holds at all six Wolfram points and at the Python points; the form with $-\frac{1}{2}$ and the form without η (mixed indices) fail. The frame curvature equals the Riemann tensor and the Ricci tensor is symmetric.

checks

GEO_curvature_G1, GEO_curvature_G2: wolfram-geometry, python-geometry

5.6 The notebook contraction and why it is wrong

The notebook's $\omega_{\text{mat}}[[\mu, a, b]]$ is the mixed connection $\omega_{\mu}^a{}_b$, and $\text{Lg}[]$ contracts it with $\text{SAB}[[a, b]] = S^{ab}$ (both indices up):

$$\Omega_{\mu}^{\text{nb}} = \frac{1}{2} \omega_{\mu}^a{}_b S^{ab} \quad (\text{one } \eta \text{ is missing}).$$

Since $\omega_{\mu}^a{}_b = \eta^{aa} \omega_{\mu ab}$ (no sum), the mixed components are symmetric in (a, b) whenever $\eta^{aa} = -\eta^{bb}$, that is for every space-time (boost) pair, and the antisymmetric S^{ab} deletes them. For time-time pairs $(a, b \geq 4)$ the sign is reversed. So Ω_{μ}^{nb} equals the space-space part of Ω_{μ} minus its time-time part, and has no boost part (derived from the index placement).

Result 5.6. With the notebook contraction the gammas are not covariantly constant: $D_{\mu} \gamma^{\nu} \neq 0$ in all 64 pairs (μ, ν) at every G1 point, with 4096 nonzero matrix entries, and in 15 pairs with 288 nonzero entries in G2. The largest violation $\max |D_{\mu} \gamma^{\nu}|$ is $2/3$, $1/5$ and $9/5$ at the three G2 points and approximately 1.46822, 1.52979 and 1.68937 at the three G1 points. The symmetric part of $\omega_{\mu}^a{}_b$ that the contraction deletes has 256 nonzero entries at each G1 point and 12 in G2. In the primordial field the notebook contraction gives $\gamma^{\mu} \Omega_{\mu}^{\text{nb}} = \frac{3H}{2}(\gamma^0 + a'_4 \gamma^4)$ instead of the correct $3H\gamma^0$, and it breaks local spin invariance already at first order. Replacing Ω by Ω^{nb} makes all eleven connection-sensitive G2 checks fail.

checks

```
GEO_notebookContractionFails_G1,
GEO_notebookContractionFails_G2: wolfram-geometry, python-geometry
NEG_notebookConnectionDetected_G2: wolfram-geometry
```

5.7 Test geometries

G1 is a generic non-diagonal vielbein,

$$e_{\mu}^a = \delta_{\mu}^a + P_{\mu}^a(x), \quad P_{\mu}^a = \frac{1}{10}((\mu + 1)x_a - (a + 1)x_{\mu}) + \frac{1}{20}x_{\mu}x_a + \frac{1}{30}x_{(\mu+a) \bmod 8}^2 \quad (\mu \neq a),$$

$$P_{\mu}^{\mu} = \frac{1}{10}x_{\mu}^2 + \frac{1}{40}x_{(\mu+1) \bmod 8},$$

evaluated at three rational points:

```
p1 = (1/7, -2/9, 1/5, 3/11, -1/13, 2/17, -3/19, 1/23)
p2 = (-1/3, 1/4, 2/7, -1/5, 1/6, -2/11, 1/9, 3/13)
p3 = (2/9, 1/8, -1/7, 1/10, -3/14, 1/12, 2/15, -1/16)
```

At each point $\det e$ is an exact positive rational (approximately 1.49738, 1.85501 and 1.37123), the metric has inertia $(4, 4, 0)$, $g^{44} \neq 0$, and the scalar curvature is approximately -2.42693 , 13.2081 and 36.6612 . The exact rationals are in the reports, and the two implementations agree on them exactly (Section 11.6).

checks

```
GEO_frameNondegenerate_G1: wolfram-geometry, python-geometry
GEO_sqrtgSquaredEqualsDet_G1: python-geometry
```

G2 is the primordial field of Stage 2 with an arbitrary function a_4 :

$$e_\mu^a = \text{diag}(\cot z, s^{1/6} e^{a_4} (\times 3), 1, s^{1/6} e^{-a_4} (\times 3)), \quad z = 6Hx_0, \quad t = Hx_4, \quad s = \sin z.$$

Wolfram evaluates it at three exact points; Python keeps $\sin^{1/6} z$, $\cos z$, e^{a_4} , a_4' , a_4'' , a_4''' and H as symbols. The results are $\sqrt{|g|} = \cos z$, $R = 6H^2(a_4'^2 - 7)$ (at the three Wolfram points $R = -2672/147$, $-1102/675$ and $-834/49$) and $\gamma^\mu \Omega_\mu = 3H\gamma^0$.

checks

```
GEO_primordialInvariants_G2, GEO_diagonalSlashFormula_G2: wolfram-geometry,
python-geometry
GEO_symbolicJetAgreement_G2, GEO_frameNondegenerate_G2: wolfram-geometry
GEO_sqrtgSquaredEqualsDetg_G2: python-geometry
```

G3 is a homogeneous diagonal frame, $ds^2 = -N^2 dx_4^2 + \sum_{i \neq 4} \eta_{ii} h_i^2 dx_i^2$, evaluated at $N = 1$, $h_i = 1 + x_4^2/(i + 2)$ and $x_4 \in \{1/3, -2/5, 3/7\}$. Fully symbolic diagonal frames are used for the metric variation (Section 9.1).

Field data at each point (the values of Ψ , $\Psi^\dagger$ and their first and second derivatives) are independent exact random rationals from recorded seeds: a documented 64-bit linear congruential generator in Wolfram, recorded seeds in Python. The Wolfram parameters are $m \in \{3/7, -2/5, 5/9\}$ and $\lambda \in \{5/11, 7/13, -3/8\}$ at p1, p2, p3.

6 Why the notebook Lagrangian Lg[] is empty for a Grassmann field

6.1 The notebook Lagrangian, verbatim

The task quotes the notebook's Lagrangian as expression [2]. Verbatim in WolframScript (the line breaks are inserted only for the page width; joining the lines without any separator gives the expression character for character):

```
Lg[]:=Sqrt[detgg] *( Transpose[\[CapitalPsi]16].\[Sigma]16.
Sum[FullSimplify[(T16^\[Alpha])\[Alpha]1-1]/.sg),constraintVars].
(D[ \[CapitalPsi]16,X[[\[Alpha]1]]]+(Q1/2)*Sum[\[Omega]mat[[\[Alpha]1,a,b]]*SAB[[a,
b]].\[CapitalPsi]16,{a,1,8},{b,1,8}]],{\[Alpha]1,1,Length[X]}]+
(H*M)*Transpose[\[CapitalPsi]16].\[Sigma]16.\[CapitalPsi]16)//Simplify[#,
constraintVars]&
```

This is the definition in notebook cell 1064 (In[1034]); its InputForm there reads $\text{Lg[]} := (\text{Simplify}[\#1, \text{constraintVars}] \&)[\text{Sqrt}[\text{detgg}] * (...)]$, and both forms parse to the identical held expression in WolframScript 1.14 (a one-off comparison made while writing this document, not one of the recorded checks). In the notation of this document, with $Q1 = 1$ and $m = -HM$,

$$\text{Lg} = \sqrt{|g|} \left[\Psi^T C \gamma^\mu (\partial_\mu \Psi + \Omega_\mu^{\text{nb}} \Psi) - m \Psi^T C \Psi \right],$$

where the notebook's Ψ_{16} is a real 16-component column and $(T16\hat{\alpha})[\mu]$ is the curved gamma γ^μ . The notebook treats the components $f_{16}[k]$ as commuting functions. For a fermion they must anticommute, and then $\text{Lg}[]$ carries no dynamics at all.

6.2 Lemmas

Let Ψ be a real 16-component column of Grassmann-odd fields.

Lemma 6.1. For every matrix M , $\Psi^T M \Psi = \Psi^T M_A \Psi$ with $M_A = \frac{1}{2}(M - M^T)$. In particular $\Psi^T C \Psi \equiv 0$, because C is symmetric.

Proof. $\Psi^T M \Psi = \sum_{ab} \Psi_a M_{ab} \Psi_b = -\sum_{ab} \Psi_b M_{ab} \Psi_a = -\Psi^T M^T \Psi$, so the symmetric part drops out.

Lemma 6.2. For an antisymmetric matrix function $A(x)$, $\Psi^T A \partial_\mu \Psi = \frac{1}{2} \partial_\mu (\Psi^T A \Psi) - \frac{1}{2} \Psi^T (\partial_\mu A) \Psi$.

Proof. $\partial_\mu \Psi^T A \Psi = -\Psi^T A^T \partial_\mu \Psi = \Psi^T A \partial_\mu \Psi$ by the anticommutation of the components and $A^T = -A$; the product rule gives the claim.

Lemma 6.3. $C \gamma^c S^{ab} = \frac{1}{2} C \{\gamma^c, S^{ab}\} + \frac{1}{2} C [\gamma^c, S^{ab}]$, where the first term is symmetric and the second antisymmetric (Result 3.5). Hence $\Psi^T C \gamma^\mu \Omega_\mu \Psi = \frac{1}{2} \Psi^T C [\gamma^\mu, \Omega_\mu] \Psi$.

6.3 Theorem

Theorem 6.1. For a real Grassmann-odd Ψ and the canonical spin connection, $\text{Lg}[]$ (which already contains the factor $\text{Sqrt}[\text{det}g]$) is a pure divergence,

$$\text{Lg} = \partial_\mu V^\mu, \quad V^\mu = \frac{1}{2} \sqrt{|g|} \Psi^T C \gamma^\mu \Psi,$$

so its Euler-Lagrange operator vanishes identically: the field equations are $0 = 0$.

Proof. The mass term vanishes by Lemma 6.1. With $A^\mu = \sqrt{|g|} C \gamma^\mu$, which is antisymmetric by expression [1], Lemma 6.2 gives $\sqrt{|g|} \Psi^T C \gamma^\mu \partial_\mu \Psi = \frac{1}{2} \partial_\mu (\Psi^T A^\mu \Psi) - \frac{1}{2} \Psi^T C \partial_\mu (\sqrt{|g|} \gamma^\mu) \Psi$. By Lemma 6.3 and the divergence identity (Result 5.3) the connection term is $\sqrt{|g|} \Psi^T C \gamma^\mu \Omega_\mu \Psi = \frac{1}{2} \Psi^T C \partial_\mu (\sqrt{|g|} \gamma^\mu) \Psi$, which cancels the second term exactly. The Euler-Lagrange operator of a total divergence is identically zero.

With the notebook's own contraction Ω^{nb} in place of Ω the cancellation fails, but what remains has no derivatives: the Euler-Lagrange equations become the algebraic equations $X \Psi = 0$ with $X = \sqrt{|g|} C \sum_\alpha [\gamma^\alpha, \Omega_\alpha^{\text{nb}} - \Omega_\alpha]$. Either way $\text{Lg}[]$ gives no wave equation for a Grassmann field.

6.4 Machine verification

Result 6.1. In a genuine Grassmann algebra (real Ψ and its jets as odd generators; 720 generators in the Python jet space):

1. $\Psi^T C \Psi$ has 0 monomials, while each $\Psi^T C \gamma^a \Psi$ has 8 monomials with coefficients ± 2 ; for a random integer matrix only the antisymmetric part survives.

2. $\Psi^T A \partial_\mu \Psi = \frac{1}{2} \partial_\mu (\Psi^T A \Psi)$ for each constant $C\gamma^a$ and a random antisymmetric matrix, with identically vanishing Euler-Lagrange expression; with the symmetric C instead, the kinetic term does not reduce and its Euler-Lagrange expression is $2(C\partial_0\Psi)_0 \neq 0$.
3. For `Lg[]` itself in curved space (G1 at p1 and the symbolic G2): with the canonical Ω , 0 of the 16 Euler-Lagrange components are nonzero, and `Lg[]` equals $\partial_\mu V^\mu$ exactly. With the notebook contraction all 16 components are nonzero, contain no derivative, and equal $X\Psi$ with the X above.
4. The Wolfram geometry verifier repeats the computation at all six G1 and G2 points in its own Grassmann algebra: the Euler-Lagrange operator vanishes identically with the canonical Ω ; with the notebook contraction it is $X\Psi$ with X of rank 16, $\max |X|$ approximately 9.927, 12.36 and 10.72 at the G1 points and $3\sqrt{455}/32 \approx 1.99976$ at the first G2 point.

checks

```
GR_massTermVanishesReal, GR_bilinearOnlyAntisymmetricPartSurvives: grassmann-demo
ALG_grassmannLemmas: wolfram-geometry
GR_kineticTotalDerivativeReal, GR_kineticSymmetricMatrixContrast: grassmann-demo
GR_notebookLgELTrivial, GR_notebookLgPureDivergence: grassmann-demo
LAG_notebookLgGrassmannTrivial_G1, LAG_notebookLgGrassmannTrivial_G2: wolfram-geometry
```

Since X has rank 16 at every tested point, the equations $X\Psi = 0$ admit only $\Psi = 0$ there.

6.5 Remark on commuting fields

For commuting components (the notebook's actual use) Lemma 6.1 is reversed: only the symmetric part of a bilinear survives. The mass term $\Psi^T C \Psi$ then survives, and from the connection term only $\frac{1}{2} C \{\gamma^\mu, \Omega_\mu\}$ survives; it contains only the totally antisymmetric part $\omega_{[cab]}$ of the connection, which vanishes for every diagonal vielbein (for G2, $\sum_\mu \{\gamma^\mu, \Omega_\mu\} = 0$ for both contractions: `GEO_anticommutatorGammaOmegaVanishesDiagonal_G2`). In the notebook's diagonal fields the gravitational term in its Euler-Lagrange equations therefore comes from $\partial_\mu(\sqrt{|g|}\gamma^\mu)$, not from the spin connection. This remark is derived from the lemmas; it is not a separate machine check.

7 The dirac16complex Lagrangian

7.1 Formula

The Lagrangian density of `dirac16complex` in an arbitrary gravitational field is

$$\mathcal{L} = \sqrt{|g|} \left[\frac{1}{2} (\bar{\Psi} \gamma^\mu D_\mu \Psi - (D_\mu \bar{\Psi}) \gamma^\mu \Psi) - m \bar{\Psi} \Psi - U(\bar{\Psi} \Psi) \right],$$

$$\bar{\Psi} = \Psi^\dagger C, \quad U(S) = \frac{\lambda}{2} S^2.$$

We write $S = \bar{\Psi} \Psi$, $\mathcal{L}_s = \mathcal{L}/\sqrt{|g|}$ and $M_{\text{eff}} = m + U'(S)$, and $K = \frac{1}{2} (\bar{\Psi} \gamma^\mu D_\mu \Psi - (D_\mu \bar{\Psi}) \gamma^\mu \Psi)$ for the kinetic term.

7.2 WolframScript form in the notebook's variable names

The same Lagrangian, written with the notebook's variable names (1-based lists, as in Lg[]):

```
(* dirac16complex Lagrangian density in the notebook's variable names. *)
(* Lists are 1-based as in the notebook: SAB[[a,b]] is S^(a-1,b-1). *)
\[CapitalPsi]16 = (f16[#] @@ X &) /@ Range[0, 15]; (* all eight coordinates *)
(* lowered spin connection omega_{mu ab} = eta_{ac} omega_mu^c_b *)
\[Omega]low = Table[Sum[\[Eta]4488[[a, c]] \[Omega]mat[[\[Alpha]1, c, b]], {c, 1, 8}],
  {\[Alpha]1, 1, 8}, {a, 1, 8}, {b, 1, 8}];
\[CapitalOmega]16[[\[Alpha]1_] := (1/2) Sum[\[Omega]low[[\[Alpha]1, a, b]] SAB[[a, b]],
  {a, 1, 8}, {b, 1, 8}];
\[CapitalPsi]16bar = ConjugateTranspose[\[CapitalPsi]16].\[Sigma]16;
D\[CapitalPsi]16[[\[Alpha]1_] := D[\[CapitalPsi]16, X[[\[Alpha]1]]] +
  \[CapitalOmega]16[[\[Alpha]1]].\[CapitalPsi]16;
D\[CapitalPsi]16bar[[\[Alpha]1_] := D[\[CapitalPsi]16bar, X[[\[Alpha]1]]] -
  \[CapitalPsi]16bar.\[CapitalOmega]16[[\[Alpha]1];
S16 = \[CapitalPsi]16bar.\[CapitalPsi]16; (* S = Psibar Psi *)
U[s_] := (\[Lambda]/2) s^2; (* default potential *)
m = -H*M; (* mass, m = -H M *)
Ldirac16complex[] := Sqrt[Abs[detgg]]*(
  (1/2) Sum[
    \[CapitalPsi]16bar.(T16^\[Alpha])[[\[Alpha]1 - 1].D\[CapitalPsi]16[[\[Alpha]1] -
    D\[CapitalPsi]16bar[[\[Alpha]1].(T16^\[Alpha])[[\[Alpha]1 - 1].\[CapitalPsi]16,
    {\[Alpha]1, 1, Length[X]}] -
    m S16 - U[S16])
```

The field depends on all eight coordinates, so Ψ_{16} is redefined accordingly, and the curved gammas $(T_{16}^\alpha)_{[\mu]}$ are those of the arbitrary vielbein (no /.sg specialization). Mathematica's Times and Dot treat these symbols as commuting, so this code is a transcription that keeps every $\bar{\Psi}$ factor to the left of every Ψ factor; it is not executed as a Grassmann computation. The verified implementation is wolfram/Dirac16ComplexGeometry.wl, which carries its own exact Grassmann algebra (convention.grassmann in wolfram-geometry-report.json).

7.3 Relation to Lg[], term by term

Lg[]	dirac16complex
Transpose[Ψ ₁₆]	ConjugateTranspose[Ψ ₁₆]
real commuting Ψ	complex Grassmann Ψ
Transpose[Ψ ₁₆].σ ₁₆	$\bar{\Psi} = \Psi^\dagger C$, the same $C = \sigma_{16}$
unsymmetrized $\Psi^T C \gamma^\mu D_\mu \Psi$	symmetrized $\frac{1}{2}(\bar{\Psi} \gamma^\mu D_\mu \Psi - (D_\mu \bar{\Psi}) \gamma^\mu \Psi)$
ω _{mat} , mixed ω _μ ^a _b	ω _{low} , lowered ω _{μab} = η _{ac} ω _μ ^c _b
(Q1/2) with a switch Q1	the fixed factor $\frac{1}{2}$ of Ω _μ = $\frac{1}{2} \omega_{\mu ab} S^{ab}$
Sqrt[detgg]	Sqrt[Abs[detgg]]
+(H M) Transpose[Ψ ₁₆].σ ₁₆ .Ψ ₁₆	-mΨΨ with m = -HM (same sign)
no self-interaction	-U(ΨΨ), U = $\frac{\lambda}{2} S^2$

7.4 Hermiticity

No factor i is needed. C is real symmetric and $C\gamma^a$ is real antisymmetric (expression [1]), hence anti-Hermitian, and with the conjugation rule $(\theta_1\theta_2)^* = \theta_2^*\theta_1^*$ the symmetrized kinetic term is Hermitian. $S = \bar{\Psi}\Psi$ is Hermitian.

Result 7.1. $(C\gamma^a)^\dagger = -C\gamma^a$ and $C^\dagger = C$; at the coefficient level the Lagrangian has the form $\bar{\Psi}K_0\Psi + \bar{\Psi}K^\mu\partial_\mu\Psi + \partial_\mu\bar{\Psi}L^\mu\Psi$ with $K_0 = K_0^\dagger$ and $L^\mu = (K^\mu)^\dagger$. In the Grassmann algebra $\mathcal{L}^* = \mathcal{L}$ exactly, and S is Hermitian with 16 monomials. As a negative control, the unsymmetrized kinetic term $\bar{\Psi}\gamma^\mu D_\mu\Psi$ alone is not Hermitian.

checks

```
LAG_hermiticity_G1, LAG_hermiticity_G2: wolfram-geometry
GR_lagrangianHermitian, GR_scalarBilinearHermitian,
GR_unsymmetrizedKineticNotHermitian: grassmann-demo
```

7.5 Symmetries

- **Local spin transformations.** $\mathcal{L}$ is invariant under every local $R(x)$ of spinor norm $+1$, in particular under the identity component $\text{Spin}_0(4,4)$ (derived from $R^T C R = C$ and the transformation law of Ω_μ ; tested in Result 5.4 for finite products of two unit vector fields and infinitesimally). Elements of spinor norm -1 are treated in Section 7.7.
- **Diffeomorphisms.** $\mathcal{L}$ is a scalar density by construction (not a separate check).
- **U(1).** $\Psi \rightarrow e^{i\alpha}\Psi$ leaves $\mathcal{L}$ invariant. The Noether current $J^\mu = -i\bar{\Psi}\gamma^\mu\Psi$ is Hermitian (GR_currentHermitian; QNT_currentHermiticity) and is conserved on shell (derived from the field equations of Section 8).

7.6 The potential: polynomials only

$S = \bar{\Psi}\Psi$ is even, of Grassmann degree 2. At a point the algebra is generated by 32 odd elements (Ψ_a and Ψ_a^*), so $S^{17} = 0$ and every function of S is a polynomial of degree at most 16. We require $U(0) = U'(0) = 0$: no constant term (no cosmological constant) and no second mass term. The default $U = \frac{\lambda}{2}S^2$ is a four-fermion contact interaction.

Result 7.2. $(\bar{\Psi}\Psi)^k$ has $\binom{16}{k}$ nonzero monomials with coefficients of absolute value $k!$ for $k = 1, \dots, 16$, namely 16, 120, 560, 1820, 4368, 8008, 11440, 12870, 11440, 8008, 4368, 1820, 560, 120, 16 and 1, and $(\bar{\Psi}\Psi)^{17} = 0$. The Grassmann derivatives of the quartic term are $\partial(\frac{\lambda}{2}S^2)/\partial\Psi_a^\dagger = \lambda S(C\Psi)_a$ (left derivative) and $\lambda S\bar{\Psi}_a$ (right derivative with respect to Ψ_a), and S commutes with Ψ .

checks

```
GR_quarticTermPolynomial: grassmann-demo
ALG_grassmannLemmas: wolfram-geometry
```

7.7 Discrete maps: the chirality map and the Pin characters

Chirality map. Under $\Psi \rightarrow \gamma^8 \Psi$ one has $\bar{\Psi} \rightarrow \bar{\Psi} \gamma^8$ (because $(\gamma^8)^T C = C \gamma^8$), γ^8 commutes with D_μ , every kinetic matrix changes sign and S is unchanged. Hence

$$\begin{aligned}\mathcal{L}_{m,U}[\gamma^8 \Psi] &= -\mathcal{L}_{-m,-U}[\Psi], \\ \mathcal{L}_{m,\lambda}[\gamma^8 \Psi] &= -\mathcal{L}_{-m,-\lambda}[\Psi] \quad \text{for } U = \frac{\lambda}{2} S^2.\end{aligned}$$

The simpler form $\mathcal{L}_m \rightarrow -\mathcal{L}_{-m}$ holds only for $U = 0$. The energy-momentum tensor of Section 9 is built from the same bilinears and maps the same way, $T_{\mu\nu}^{(m,\lambda)}[\gamma^8 \Psi] = -T_{\mu\nu}^{(-m,-\lambda)}[\Psi]$ (derived). If Ψ solves the field equations with (m, λ) , then $\gamma^8 \Psi$ solves them with $(-m, -\lambda)$. This pairing of $\pm m$ is a structural property of the equations and not a physical claim. The measured signs are -1 for every kinetic matrix and $+1$ for the mass and potential terms.

checks

ALG_gamma8Map: wolfram-algebra, python-algebra

Pin characters. For a unit vector u with $u^2 = n(u)$, $\Psi \rightarrow u\Psi$ gives $\bar{\Psi}\Psi \rightarrow -n(u)\bar{\Psi}\Psi$. With the untwisted lift the kinetic term picks up the same factor $-n(u)$, so the free Lagrangian goes to $-n(u)\mathcal{L}$ and the free field equations are $\text{Pin}(4,4)$ -covariant; with the twisted lift the kinetic factor is $+n(u)$, opposite to the mass factor. For a product $u_1 u_2$ the factor is the spinor norm $n(u_1)n(u_2)$; the 16 products $e_a e_b$ with $a < 4 \leq b$ have spinor norm -1 and change the sign of both terms. The even potential does not change sign: for a transformation with character -1 ,

$$\mathcal{L}_{m,\lambda} \rightarrow -\mathcal{L}_{m,-\lambda},$$

and the interacting field equation is mapped to the one with $\lambda \rightarrow -\lambda$. Exact covariance of the interacting theory under character -1 holds only for $U = 0$ or an odd U ; elements with character $+1$, in particular all of $\text{Spin}_0(4,4)$, are exact symmetries. In the geometry verifier, a local $R = \gamma(n)\gamma(m)$ with $\eta(n,n)\eta(m,m) = -1$ maps $\mathcal{L}[m,\lambda]$ exactly to $-\mathcal{L}[m,-\lambda]$, and the naive $-\mathcal{L}[m,\lambda]$ fails for $\lambda \neq 0$.

checks

ALG_pinLiftCharacter: wolfram-algebra, python-algebra

LAG_localSpinInvariance_G1, LAG_localSpinInvariance_G2: wolfram-geometry

8 Euler-Lagrange equations in an arbitrary gravitational field

8.1 Derivation

Vary $\Psi^\dagger$ (left derivative) and Ψ (right derivative) independently. The terms of $\mathcal{L}$ that contain $\Psi^\dagger$ are $\frac{1}{2}\sqrt{|g|}\Psi^\dagger C\gamma^\mu D_\mu \Psi$, $-\frac{1}{2}\sqrt{|g|}(\partial_\mu \Psi^\dagger C - \Psi^\dagger C\Omega_\mu)\gamma^\mu \Psi$ and $-\sqrt{|g|}(mS + U)$. Integrating the derivative term by parts and using the divergence identity $\partial_\mu(\sqrt{|g|}\gamma^\mu) = \sqrt{|g|}[\gamma^\mu, \Omega_\mu]$,

$$\begin{aligned}
E &:= \frac{\partial_L \mathcal{L}}{\partial \Psi^\dagger} - \partial_\mu \frac{\partial_L \mathcal{L}}{\partial (\partial_\mu \Psi^\dagger)} \\
&= \sqrt{|g|} C \left[\frac{1}{2} \gamma^\mu D_\mu \Psi + \frac{1}{2} \Omega_\mu \gamma^\mu \Psi - (m + U'(S)) \Psi \right] + \frac{1}{2} \partial_\mu (\sqrt{|g|} C \gamma^\mu \Psi) \\
&= \sqrt{|g|} C \left[\frac{1}{2} \gamma^\mu D_\mu \Psi + \frac{1}{2} \Omega_\mu \gamma^\mu \Psi + \frac{1}{2} \gamma^\mu \partial_\mu \Psi + \frac{1}{2} [\gamma^\mu, \Omega_\mu] \Psi - (m + U'(S)) \Psi \right] \\
&= \sqrt{|g|} C [\gamma^\mu D_\mu \Psi - (m + U'(S)) \Psi].
\end{aligned}$$

C is invertible, so E vanishes exactly when the Dirac equation below holds. The quartic term contributes $\lambda S C \Psi$ (Result 7.2). The variation with respect to Ψ gives the conjugate equation in the same way.

8.2 The covariant field equations

$$\gamma^\mu D_\mu \Psi = (m + U'(\bar{\Psi} \Psi)) \Psi, \quad (D_\mu \bar{\Psi}) \gamma^\mu = -(m + U'(\bar{\Psi} \Psi)) \bar{\Psi},$$

with $U'(S) = \lambda S$ for the default potential. The second equation is the Dirac conjugate of the first: $(\gamma^\mu D_\mu \Psi)^\dagger C = -(D_\mu \bar{\Psi}) \gamma^\mu$ by expression [1] and Result 3.5.

Result 8.1. At every G1 and G2 point, with $\lambda \neq 0$, the Euler-Lagrange expression with respect to $\Psi_a^\dagger$ equals $\sqrt{|g|} (C(\gamma^\mu D_\mu \Psi - (m + \lambda S) \Psi))_a$, and the one with respect to Ψ_a equals $\sqrt{|g|} ((D_\mu \bar{\Psi}) \gamma^\mu + (m + \lambda S) \bar{\Psi})_a$. The Wolfram verifier works in a genuine Grassmann algebra and computes each Euler-Lagrange operator in two independent ways (direct total derivative and a jet-commutator identity), which agree. The Python geometry checker evaluates the same expressions with commuting exact numbers, which is exact here because the Lagrangian is bilinear with every $\bar{\Psi}$ factor on the left (measurement LAG_commutingProxyJustification), and the Grassmann demonstration repeats them in a genuine Grassmann algebra at G1 p1 and in the symbolic G2.

checks

```

LAG_eulerLagrangePsibar_G1,
LAG_eulerLagrangePsibar_G2, LAG_eulerLagrangePsi_G1,
LAG_eulerLagrangePsi_G2: wolfram-geometry, python-geometry
GR_complexLagrangianNonTrivial, GR_complexQuarticEL,
GR_complexPsiEquation: grassmann-demo

```

8.3 Component form

$$\begin{aligned}
\gamma^\mu D_\mu \Psi &= e_a^\mu \gamma^a \left(\partial_\mu + \frac{1}{8} \omega_{\mu bc} [\gamma^b, \gamma^c] \right) \Psi, \\
\omega_{\mu bc} &= \eta_{bd} e_c^\nu (\Gamma^\rho_{\mu\nu} e_\rho^d - \partial_\mu e_\nu^d).
\end{aligned}$$

Written out, the 16 component equations are

$$\sum_{\mu=0}^7 \sum_{a=0}^7 \sum_{k=0}^{15} e_a^\mu (\gamma^a)_{jk} \left(\partial_\mu \Psi_k + \sum_{l=0}^{15} (\Omega_\mu)_{kl} \Psi_l \right) = (m + \lambda S) \Psi_j \quad (j = 0, \dots, 15),$$

$$S = \sum_{k,l=0}^{15} \Psi_k^* C_{kl} \Psi_l.$$

8.4 Non-triviality in an arbitrary gravitational field

Theorem 8.1. (a) The spin connection is locally pure gauge, $\Omega_\mu = -(\partial_\mu R)R^{-1}$ for some local spin transformation R , if and only if the Riemann tensor vanishes. (b) In a curved field no local spin transformation removes the spin connection from the field equations, and the squared Dirac operator contains the scalar curvature:

$$(\gamma^\mu D_\mu)^2 \Psi = g^{\mu\nu} (D_\mu D_\nu \Psi - \Gamma_{\mu\nu}^\lambda D_\lambda \Psi) + c R \Psi, \quad c = -\frac{1}{4}.$$

Proof. (a) A pure-gauge connection has $F_{\mu\nu} = 0$ (direct computation). Conversely $F_{\mu\nu} = \frac{1}{2} R_{ab\mu\nu} S^{ab}$ (Result 5.5), and the 28 matrices S^{ab} , $a < b$, are linearly independent (Result 3.2), so $F = 0$ forces the Riemann tensor to vanish; a flat connection is locally pure gauge. (b) If Ω could be gauged away the curvature would vanish, contradicting (a) in a curved field. The Lichnerowicz form follows from $\gamma^\mu \gamma^\nu = g^{\mu\nu} + \frac{1}{2} [\gamma^\mu, \gamma^\nu]$, $D_\mu \gamma^\nu = 0$ and $[D_\mu, D_\nu] \Psi = F_{\mu\nu} \Psi$; the constant c is fixed by the measurement below. Consequently, for $U = 0$, every solution satisfies $g^{\mu\nu} (D_\mu D_\nu \Psi - \Gamma_{\mu\nu}^\lambda D_\lambda \Psi) - \frac{1}{4} R \Psi = m^2 \Psi$: the curvature enters the second-order equation explicitly.

Result 8.2. $c = -1/4$ exactly at G1 p1, p2, p3 and at the three Wolfram G2 points and the symbolic Python G2 point, and $c = +1/4$ fails. In curved space $\gamma^\mu \Omega_\mu \neq 0$: it has 128 nonzero entries at G1 p1 and equals $3H\gamma^0$ in G2, and the spin-connection term of the $\Psi^\dagger$ equation is nonzero in all 16 components at every G1 point and G2 sample.

checks

```
GEO_lichnerowicz_G1, GEO_lichnerowicz_G2: wolfram-geometry, python-geometry
GEO_lichnerowiczConstantSameG1G2: python-geometry
LAG_eulerLagrangePsibar_spinConnectionTermNonzeroComponents:
python-geometry measurements
```

8.5 Diagonal vielbeins

For a diagonal vielbein $e_\mu^a = \text{diag}(h_0, \dots, h_7)(x)$ the anticommutator part vanishes, $\{\gamma^\mu, \Omega_\mu\} = 0$, and

$$\gamma^\mu \Omega_\mu = \frac{1}{2\sqrt{|g|}} \partial_\mu (\sqrt{|g|} \gamma^\mu) = \frac{1}{2} \sum_{b=0}^7 \frac{1}{h_b} \partial_b \ln \left(\prod_{c \neq b} h_c \right) \gamma^b.$$

In the primordial field this is $3H\gamma^0$, independent of a_4 .

checks

GEO_diagonalSlashFormula_G2: wolfram-geometry, python-geometry
 GEO_diagonalSlashFormula_G3,
 GEO_anticommutatorGammaOmegaVanishesDiagonal_G2: python-geometry

9 Energy-momentum tensor operator, energies, pressure and equation of state

9.1 Derivation from the metric variation

The energy-momentum tensor is defined by $T_{\mu\nu} = -\frac{2}{\sqrt{|g|}} \frac{\delta \mathcal{S}}{\delta g^{\mu\nu}}$ with $\mathcal{S} = \int d^8x \mathcal{L}$. For a spinor the metric is varied through the vielbein, $\delta e_a{}^\mu = \frac{1}{2} e_{a\nu} \delta g^{\mu\nu}$ (the antisymmetric part of δe is an infinitesimal local frame rotation, under which $\mathcal{S}$ is invariant). The explicit $e_a{}^\mu$ in $\gamma^\mu = e_a{}^\mu \gamma^a$ contributes $\frac{1}{2}(\bar{\Psi} \gamma^a D_\mu \Psi - (D_\mu \bar{\Psi}) \gamma^a \Psi) \delta e_a{}^\mu$ and $\delta \sqrt{|g|} = -\frac{1}{2} \sqrt{|g|} g_{\mu\nu} \delta g^{\mu\nu}$; the variation of Ω_μ enters the symmetrized kinetic term only through the totally antisymmetric part of the connection and, by the standard argument, does not contribute to the symmetric tensor (for diagonal metrics the reduced Lagrangian contains no metric derivatives at all, Result 9.1). With $e_{a\nu} \gamma^a = \gamma_\nu$ this gives the formula below.

Result 9.1. For diagonal metrics the variation is carried out explicitly and reproduces the formula off shell and on shell: in the minisuperspace $h_\mu(x_4)$ with lapse $N = h_4$, $\rho = -\frac{N}{\sqrt{|g|}} \frac{\partial \mathcal{L}}{\partial N}$ and $T^i{}_i = \frac{h_i}{\sqrt{|g|}} \frac{\partial \mathcal{L}}{\partial h_i}$ (no sum), and Python repeats it for a local diagonal frame with symbolic h_μ and all $\partial_\nu h_\mu$. The reduced Lagrangian does not depend on the metric velocities. The off-diagonal components are not varied explicitly; they are tested through symmetry, Hermiticity and on-shell conservation in G1 (Limitations).

checks

EMT_variation_G3 (and measurement emt.signConvention): wolfram-geometry
 EMT_variation: python-geometry

9.2 Formula

$$T_{\mu\nu} = -\frac{1}{4} \left[\bar{\Psi} \gamma_\mu D_\nu \Psi + \bar{\Psi} \gamma_\nu D_\mu \Psi - (D_\mu \bar{\Psi}) \gamma_\nu \Psi - (D_\nu \bar{\Psi}) \gamma_\mu \Psi \right] + g_{\mu\nu} \mathcal{L}_s,$$

$$\mathcal{L}_s = K - mS - U(S).$$

On shell $K = M_{\text{eff}} S$ (contract the two field equations with $\bar{\Psi}$ and Ψ), hence $\mathcal{L}_s = SU'(S) - U(S)$ on shell. The sign is fixed so that the energy density is T_{44} ; for the homogeneous zero mode this is $mS + U$ (Section 9.7).

checks

EMT_onshellLagrangianSUprimeMinusU_G1,
 EMT_onshellLagrangianSUprimeMinusU_G2: python-geometry

9.3 Properties

Result 9.2.

1. **Symmetric:** $T_{\mu\nu} = T_{\nu\mu}$.
2. **Hermitian:** at the coefficient level $T_{\mu\nu} = \bar{\Psi}K_{\mu\nu}\Psi + \dots$ with Hermitian $K_{\mu\nu}$ and mutually adjoint derivative coefficients, and $T_{\mu\nu}^* = T_{\mu\nu}$ in the Grassmann algebra.
3. **Conserved on shell:** $\nabla^\mu T_{\mu\nu} = 0$ when the field equations and their first derivatives hold, solved exactly at the point for the x_4 -derivatives; as a negative control the off-shell divergence is nonzero.
4. **Trace:** off shell $T^\mu{}_\mu = 7K - 8(mS + U)$; on shell

$$T^\mu{}_\mu = -mS + 7SU'(S) - 8U(S) = -mS + 3\lambda S^2 \quad \text{for } U = \frac{\lambda}{2}S^2$$

The off-shell trace follows from $\gamma^\mu\gamma_\mu = 8$: $T^\mu{}_\mu = -K + 8\mathcal{L}_s$ (measurement `emt.trace` records the on-shell formula).

checks

```
EMT_symmetric_G1, EMT_symmetric_G2: python-geometry
EMT_symmetricHermitian_G1, EMT_symmetricHermitian_G2: wolfram-geometry
GR_emtHermitian: grassmann-demo
EMT_conservation_G1, EMT_conservation_G2: wolfram-geometry, python-geometry
EMT_onshellJetSolve_G1, EMT_onshellJetSolve_G2: python-geometry
EMT_traceOffShellIdentity_G1, EMT_traceOffShellIdentity_G2: python-geometry
EMT_trace_G1, EMT_trace_G2: wolfram-geometry, python-geometry
```

9.4 Observer decomposition

Take Gaussian normal time, $g_{44} = -1$, $g_{4i} = 0$, and the observer $u = \partial_4$. Then

$$\rho = T_{\mu\nu}u^\mu u^\nu = T_{44}, \quad p_{(i)} = T^i{}_i \quad (\text{no sum, } i \neq 4), \quad \bar{p} = \frac{1}{7} \sum_{i \neq 4} p_{(i)}.$$

There are seven transverse directions: four space-like ($i = 0, 1, 2, 3$) and three time-like ($i = 5, 6, 7$). The equation of state is $w = p/\rho$ when the transverse pressures are equal, and $w_{(i)} = p_{(i)}/\rho$ or $\bar{w} = \bar{p}/\rho$ otherwise.

9.5 Kinetic and potential energy: two splits

Split the kinetic term into its x_4 part and the rest, $K = K_4 + K_\perp$, with

$$K_4 = \frac{1}{2}(\bar{\Psi}\gamma^{x_4}D_4\Psi - (D_4\bar{\Psi})\gamma^{x_4}\Psi), \quad K_\perp = \frac{1}{2}\sum_{\mu \neq 4}(\bar{\Psi}\gamma^\mu D_\mu\Psi - (D_\mu\bar{\Psi})\gamma^\mu\Psi).$$

- **(A) Lagrangian split.** $\text{KE}_L := \frac{1}{2}K_4$ (half of the time-derivative part of the kinetic term) and $\text{PE}_L := \rho - \text{KE}_L$. This is the exact analogue of the scalar field's $\frac{1}{2}\dot{\phi}^2$ and V : $\rho = \text{KE}_L + \text{PE}_L$ by definition, and for a homogeneous isotropic state on shell also $p = \text{KE}_L - \text{PE}_L$ (Section 9.6).
- **(B) Hamiltonian split.** $\text{KE}_H := -K_\perp$ (gradient and transverse-connection energy) and $\text{PE}_H := mS + U$ (mass and self-interaction energy). In Gaussian normal time, $\gamma_4 = -\gamma^{x_4}$ and $g_{44} = -1$ turn the formula of Section 9.2 into $T_{44} = K_4 - \mathcal{L}_s = -K_\perp + mS + U$, so

$$\rho = T_{44} = \text{KE}_H + \text{PE}_H \quad \text{identically (off shell).}$$

This identity is derived here from the verified formula; it is the energy density as the Hamiltonian density of Section 10.4. The time derivatives do not appear in split (B): for a first-order field they are not an energy.

9.6 Equation of state

For a homogeneous isotropic state on shell (Section 9.7):

$$\begin{aligned} \rho &= mS + U, & p &= SU' - U, \\ w &= \frac{p}{\rho} = \frac{SU' - U}{mS + U} = \frac{\text{KE}_L - \text{PE}_L}{\text{KE}_L + \text{PE}_L}, \\ \text{KE}_L &= \frac{1}{2}S(m + U'), & \text{PE}_L &= \frac{1}{2}(mS + 2U - SU'), & \text{KE}_H &= 0, & \text{PE}_H &= \rho. \end{aligned}$$

- **Free massive field** ($U = 0$). $\text{KE}_L = \text{PE}_L = \frac{1}{2}mS$, $p = 0$, $w = 0$: dust.
- **Default potential.** $\rho = mS + \frac{\lambda}{2}S^2$, $p = \frac{\lambda}{2}S^2$, $w = \frac{\lambda S}{2m + \lambda S}$, $\text{KE}_L = \frac{1}{2}S(m + \lambda S)$, $\text{PE}_L = \frac{1}{2}mS$.
- **Limits.** $w \rightarrow -1$ when $\text{KE}_L/\text{PE}_L \rightarrow 0$ and $w \rightarrow +1$ when $\text{PE}_L/\text{KE}_L \rightarrow 0$, as for the scalar field.
- **Phantom criterion.** $\rho + p = 2\text{KE}_L = S(m + U')$. For $\rho > 0$ the state is phantom ($w < -1$) exactly when $\text{KE}_L < 0$, that is when $S(m + U') < 0$. Unlike $\frac{1}{2}\dot{\phi}^2$, KE_L has no fixed sign, because $S = \Psi^\dagger C \Psi$ is built from C of signature (8,8) (Result 3.3).

Consequence (derived from the verified formulas, not machine-checked). Section 9.7 gives $\partial_4 \rho + \sum_{i \neq 4} H_i(\rho + p_i) = 0$; with $\rho + p = S(m + U')$ this becomes $(m + U')(\partial_4 S + S \sum_i H_i) = 0$, so $S\sqrt{|g|}$ is constant wherever $m + U' \neq 0$: S dilutes as the inverse 7-volume. For the default potential $w = \lambda S/(2m + \lambda S)$ therefore changes as the 7-volume changes. For example, with $m > 0$, $\lambda < 0$ and $x = |\lambda|S/m$: $w = -x/(2 - x)$, the state is phantom with $\rho > 0$ for $1 < x < 2$, $w = -1$ at $x = 1$, and $w \rightarrow 0$ as $x \rightarrow 0$. Whether such states are physically admissible (energy conditions, stability, the quantum Krein structure) is not decided here.

9.7 Homogeneous diagonal backgrounds

For a diagonal Gaussian-normal frame $h_i(x_4)$, $h_4 = N = 1$, with Hubble rates $H_i = \partial_4 h_i / h_i$, and a homogeneous $\Psi(x_4)$:

$$\begin{aligned} \rho &= mS + U \quad (\text{off shell}), & p_{(i)} &= \mathcal{L}_s \quad (\text{off shell}) = SU' - U \quad (\text{on shell, all seven } i \neq 4), \\ T_{ij} &= \frac{1}{4}(H_i - H_j) \bar{\Psi} \gamma_i \gamma_j \gamma^{x_4} \Psi \quad (i \neq j, i, j \neq 4), & \gamma_i &= g_{ii} \gamma^{x_i}, \\ T_{4i} &= -\frac{1}{4}(\bar{\Psi} \gamma_i \partial_4 \Psi - \partial_4 \bar{\Psi} \gamma_i \Psi) \quad (\text{off shell}), & T_{4i} &= 0 \quad (\text{on shell}), \\ \partial_4 \rho + \sum_{i \neq 4} H_i (\rho + p_{(i)}) &= 0, \\ \partial_4 (\bar{\Psi} \gamma^i \gamma^j \gamma^4 \Psi) &= -\left(\sum_k H_k\right) \bar{\Psi} \gamma^i \gamma^j \gamma^4 \Psi \quad (\text{frame gammas, on shell}). \end{aligned}$$

The pressure is isotropic in all seven transverse directions although the h_i differ. The off-diagonal T_{4i} vanish on shell because $\{\gamma_i, \gamma^{x_4}\} = 0$. The T_{ij} are proportional to a tensor bilinear that decays like the inverse 7-volume, so zero initial data stay zero, and a diagonal metric is consistent with states for which this bilinear vanishes. In the flat-gamma notation of the Python checker the same component reads $T_{ij} = \frac{1}{4} \eta_{ii} \eta_{jj} h_i h_j (H_i - H_j) \bar{\Psi} \gamma^i \gamma^j \gamma^4 \Psi$; the two forms are identical because $\gamma_i = \eta_{ii} h_i \gamma^i$ (notation conversion recorded in stage1-summary.json).

Result 9.3. The Wolfram verifier checks all of these at the three G3 times, off and on shell as indicated. The Python checker independently checks, with its own random on-shell data at three G3 times, the energy density, the seven equal pressures, KE_L , PE_L , $\rho = \text{KE}_L + \text{PE}_L$, $p = \text{KE}_L - \text{PE}_L$ and the off-diagonal forms; the off-shell forms $\rho = mS + U$ and $p_{(i)} = \mathcal{L}_s$ are part of its metric-variation check. For generic states the T_{ij} are nonzero in 42 of 42 ordered pairs and the T_{4i} vanish on shell. A sample: at $x_4 = 1/3$ with $m = 4$, $\lambda = 7/6$ and $S = -75899/7560$ the Python report gives $\rho = 260864863/13996800$, $\text{KE}_L = 3793356121/97977600$, $\text{PE}_L = -75899/3780$ and $w = 75899/24059$.

checks

```
EMT_homogeneousReduction_G3: wolfram-geometry
EMT_homogeneousReduction, EMT_homogeneousTimeSpaceVanishesOnShell_G3: python-geometry
EMT_homogeneous_G3_q1 (measurement): python-geometry
```

9.8 Side-by-side with the scalar field of the PDF

Quantity	Scalar field (PDF)	dirac16complex (homogeneous, on shell)
Lagrangian	$\frac{1}{2} \dot{\phi}^2 - V(\phi)$	$\mathcal{L}_s = K - mS - U(S)$
energy density	$\rho_\phi = \frac{1}{2} \dot{\phi}^2 + V$	$\rho = mS + U$
pressure	$P_\phi = \frac{1}{2} \dot{\phi}^2 - V = \mathcal{L}_\phi$	$p = SU' - U = \mathcal{L}_s$
kinetic energy	$\frac{1}{2} \dot{\phi}^2 \geq 0$	$\text{KE}_L = \frac{1}{2} S(m+U')$, either sign
potential energy	V	$\text{PE}_L = \frac{1}{2} (mS + 2U - SU')$
$\rho = \text{KE} + \text{PE}$, $p = \text{KE} - \text{PE}$	yes	yes (split A)
Hamiltonian split	$\frac{1}{2} \dot{\phi}^2 + V$	$\text{KE}_H = 0$, $\text{PE}_H = mS + U$

Quantity	Scalar field (PDF)	dirac16complex (homogeneous, on shell)
equation of state	$w_\phi = \frac{\frac{1}{2}\dot{\phi}^2 - V}{\frac{1}{2}\dot{\phi}^2 + V}$	$w = \frac{SU' - U}{mS + U}$
$w \rightarrow -1$	$\dot{\phi}^2 \ll V$	$ \text{KE}_L \ll \text{PE}_L$
$w \rightarrow +1$	$\dot{\phi}^2 \gg V$ (kination)	$\text{PE}_L \ll \text{KE}_L$
phantom $w < -1$	impossible for a canonical field	possible: $\text{KE}_L < 0$ with $\rho > 0$
equation of motion	$\ddot{\phi} + 3H\dot{\phi} + V'(\phi) = 0$	$\gamma^\mu D_\mu \Psi = (m + U')\Psi$
dilution	depends on V	$S \propto 1/\sqrt{ g }$ (derived, 9.6)

Exact analogy. In both cases the energy density is the Hamiltonian density (T_{44}) and, for a homogeneous state, the pressure equals the Lagrangian density; the Lagrangian split gives $\rho = \text{KE} + \text{PE}$ and $p = \text{KE} - \text{PE}$, hence the same formula for w . **Difference.** The fermion Lagrangian is first order in time. Its time-derivative term is not a positive kinetic energy: in the Hamiltonian split it contributes nothing to ρ , and in the Lagrangian split $\text{KE}_L = \frac{1}{2}S(m + U')$ can be negative. The canonical scalar field satisfies $w \geq -1$ whenever $\rho > 0$; dirac16complex can cross $w = -1$ at the level of these classical (mean-field) formulas.

10 Canonical quantization in 4+4 dimensions

10.1 Slicing and momenta

We quantize with respect to x_4 on the 7-dimensional slices $x_4 = \text{const}$, with coordinates $(x_0, x_1, x_2, x_3, x_5, x_6, x_7)$. This requires $g^{44} \neq 0$ (the slices are non-characteristic). Split $\mathcal{L}$ into its x_4 part and the rest and add the total derivative $\frac{1}{2}\partial_4(\sqrt{|g|}\bar{\Psi}\gamma^{x_4}\Psi)$:

$$\mathcal{L}' = \mathcal{L} + \frac{1}{2}\partial_4(\sqrt{|g|}\bar{\Psi}\gamma^{x_4}\Psi) = \sqrt{|g|}\Psi^\dagger C\gamma^{x_4}\partial_4\Psi + (\text{terms without } \partial_4\Psi \text{ or } \partial_4\Psi^\dagger).$$

The momentum conjugate to Ψ (right derivative) is $\Pi = \sqrt{|g|}\Psi^\dagger C\gamma^{x_4}$, and $\mathcal{L}'$ does not depend on $\partial_4\Psi^\dagger$.

Result 10.1. In a genuine Grassmann algebra, $\Pi_a = \partial_R\mathcal{L}'/\partial(\partial_4\Psi_a) = \sqrt{|g|}(\Psi^\dagger C\gamma^{x_4})_a$ and $\partial_L\mathcal{L}'/\partial(\partial_4\Psi_a^\dagger) = 0$ at every G1 and G2 point.

checks

QNT_canonicalMomentum_G1, QNT_canonicalMomentum_G2: wolfram-geometry

10.2 Constraints and Dirac brackets

The momenta do not contain velocities, so there are primary constraints $\chi_a = \Pi_a - \sqrt{|g|}(\Psi^\dagger C\gamma^{x_4})_a \approx 0$ and $\bar{\chi}_a = \Pi_{\Psi^\dagger, a} \approx 0$. Their graded Poisson brackets form, up to sign and transposition, the matrix $\mathcal{K} = \sqrt{|g|}C\gamma^{x_4}$. It is invertible exactly when $g^{44} \neq 0$, because

$$(C\gamma^{x_4})(\gamma^{x_4}C) = (\gamma^{x_4}C)(C\gamma^{x_4}) = g^{44} I_{16}, \quad (C\gamma^{x_4})^{-1} = \frac{\gamma^{x_4}C}{g^{44}},$$

so the constraints are second class. The Dirac bracket eliminates them, and the quantization rule $\{\cdot, \cdot\}_+ = i\{\cdot, \cdot\}_D$ gives the anticommutators below. The matrix identity is verified symbolically for an arbitrary e_a^4 and at the G1 points. The graded constraint-matrix computation itself is not a separate machine check; the verifiers check the result as the matrix identity $\{\Psi, \Psi^\dagger\} = i\mathcal{K}^{-1}$ that follows from the first-order form of $\mathcal{L}'$.

checks

QNT_curvedAnticommutatorMatrix: wolfram-algebra

10.3 Equal- x_4 anticommutators in curved space

For x and y on the same slice ($\hbar = 1$):

$$\begin{aligned} \{\Psi_a(x), \Psi_b^\dagger(y)\} &= i[(C\gamma^{x_4})^{-1}]_{ab} \frac{\delta^7(x-y)}{\sqrt{|g|}} = i \frac{[\gamma^{x_4}C]_{ab}}{g^{44}} \frac{\delta^7(x-y)}{\sqrt{|g|}}, \\ \{\Psi_a(x), \Psi_b(y)\} &= \{\Psi_a^\dagger(x), \Psi_b^\dagger(y)\} = 0. \end{aligned}$$

The matrix $i(\sqrt{|g|}C\gamma^{x_4})^{-1}$ is Hermitian. In Gaussian normal gauge ($g^{44} = -1$ and $\gamma^{x_4} = \gamma^4$, as in the primordial field G2) this becomes

$$\{\Psi_a(x), \Psi_b^\dagger(y)\} = B_{ab} \frac{\delta^7(x-y)}{\sqrt{|g|}}, \quad B := -iC\gamma^4.$$

Result 10.2. B is Hermitian, $B^2 = 1$, its eigenvalues are $+1$ and -1 with multiplicity 8 each (characteristic polynomial $(x-1)^8(x+1)^8$), $[C, B] = 0$ and $BC = -i\gamma^4$. The Gaussian-normal reduction of the anticommutator to $B/\sqrt{|g|}$ is checked in the primordial field.

checks

ALG_chargeFormB: wolfram-algebra, python-algebra

QNT_canonicalMomentum_G1, QNT_canonicalMomentum_G2: wolfram-geometry

10.4 Hamiltonian and Heisenberg equations

The canonical Hamiltonian density is $\mathcal{H} = \Pi \partial_4 \Psi - \mathcal{L}'$. Using the split of Section 9.5 (derived here, not a separate machine check):

$$\begin{aligned} \mathcal{H} &= \sqrt{|g|} (\text{KE}_H + \text{PE}_H) \\ &\quad - \frac{1}{2} \sqrt{|g|} \bar{\Psi} \{\gamma^{x_4}, \Omega_4\} \Psi - \frac{1}{2} \bar{\Psi} \partial_4 (\sqrt{|g|} \gamma^{x_4}) \Psi. \end{aligned}$$

The last two terms come from the connection along x_4 and from the explicit x_4 -dependence of the added total derivative. In flat space in Cartesian coordinates both vanish and $\mathcal{H} = T_{44} = \rho$.

There, with $\{\Psi_a, \Psi_b^\dagger\} = B_{ab}\delta^7$ and $U = 0$, $H = \int d^7x \Psi^\dagger(mC - C\gamma^j\partial_j)\Psi$ (sum over $j \neq 4$), and the Heisenberg equation gives

$$\partial_4\Psi = i[H, \Psi] = -iB(mC - C\gamma^j\partial_j)\Psi = -m\gamma^4\Psi + \gamma^4\gamma^j\partial_j\Psi,$$

which is the field equation $\gamma^\mu\partial_\mu\Psi = m\Psi$ multiplied by γ^4 (using $BC = -i\gamma^4$ and $(\gamma^4)^2 = -1$).

10.5 Krein structure

Theorem 10.1. Every Spin(4,4)-invariant Hermitian form on $\mathbb{C}^{16}$ is even in chirality, whereas γ^4 is odd. Consequently every charge density of the form $\text{Herm}(cH\gamma^4)$ built from an invariant form H is indefinite, of signature (8,8), or vanishes. The canonical state space is therefore a Krein space; this is intrinsic to signature (4,4) and not a consequence of the particular choice B .

Result 10.3. The Spin(4,4)-invariant bilinear forms form a 2-dimensional space spanned by CP_- and CP_+ (both symmetric and block diagonal in chirality); an invariant Hermitian form is $H = \alpha CP_- + \beta CP_+$ with real α, β ; $H\gamma^4$ is block off-diagonal, traceless and similar to its negative, with $(H\gamma^4)^2 = -\alpha\beta I_{16}$; and for every constant c the Hermitian part of $cH\gamma^4$ is traceless with square $|k|^2 I_{16}$, $k = \frac{1}{2}(\bar{c}\beta - c\alpha)$, hence has signature (8,8) unless it vanishes.

checks

ALG_invariantForms: wolfram-algebra, python-algebra

The canonical anticommutator defines the Krein form $[f, h] = f^\dagger B h$ on the one-particle space. $J = B$ is a fundamental symmetry ($J = J^\dagger = J^{-1}$), and the associated positive Hilbert product is $[f, Jh] = f^\dagger h$, the standard one. In a Hilbert-space representation the Hilbert adjoint of Ψ is $\chi := \Psi^\dagger B$, with $\{\Psi_a, \chi_b\} = \delta_{ab}\delta^7$ (Gaussian normal gauge, flat): the canonical $\Psi^\dagger = \chi B$ is the Krein adjoint of Ψ , not its Hilbert adjoint.

Result 10.4. At rest the positive- and negative-frequency spaces (the eigenspaces of $-i\gamma^4$, since $h_0 = m(-i\gamma^4)$) are 8-dimensional, B preserves them, and the B -form on each has signature (4,4). The same (4,4) holds for moving states in the good sector: at $m = 1$, $k = (1, 1, 2, 3)$, $E = 4$ for both signs of the energy, and at $m = 3$, $k = (1, 1, 1, 2)$, $E = 4$ for positive energy. The Krein form is thus indefinite even among positive-energy states.

checks

QNT_kreinSignature: wolfram-algebra, python-algebra

10.6 Flat-space modes and the good sector

In flat space the field equation takes the Schrödinger form $i\partial_4\Psi = h\Psi$ with, for a plane wave $e^{ik_j x^j}$ ($j \in \{0, 1, 2, 3, 5, 6, 7\}$),

$$\begin{aligned} h_k &= -im\gamma^4 - \gamma^4 \sum_{j \neq 4} k_j \gamma^j, & [h_k, B] &= 2i \sum_{j=5}^7 k_j C \gamma^j, \\ h_k^2 &= (m^2 + k_0^2 + k_1^2 + k_2^2 + k_3^2 - k_5^2 - k_6^2 - k_7^2) I_{16}. \end{aligned}$$

The anti-Hermitian part of h_k is $-\gamma^4(k_5\gamma^5 + k_6\gamma^6 + k_7\gamma^7)$, whose square is $-(k_5^2 + k_6^2 + k_7^2)I_{16}$. Hence h_k is Hermitian, and commutes with B , if and only if $k_5 = k_6 = k_7 = 0$.

Result 10.5. These identities hold exactly for symbolic m and k (Wolfram) and by a structural proof for all real m, k plus 10 exact samples (Python); 2 of the samples with extra-time momentum have $E^2 < 0$.

checks

QNT_flatModeHamiltonian: wolfram-algebra, python-algebra

The good sector is the sector of fields that do not depend on x_5, x_6, x_7 ($k_5 = k_6 = k_7 = 0$). There h_k is Hermitian and traceless with $h_k^2 = E_k^2$, $E_k = \sqrt{m^2 + k_0^2 + k_1^2 + k_2^2 + k_3^2}$, so its eigenvalues are $+E_k$ and $-E_k$ with multiplicity 8 each: 8 particle and 8 antiparticle states for every momentum $k = (k_0, k_1, k_2, k_3)$.

10.7 Fock space, Dirac sea and normal ordering

In the good sector (flat space, Gaussian normal time) let $u_s(k)$ and $v_s(k)$, $s = 0, \dots, 7$, be orthonormal (for $f^\dagger h$) eigenvectors of h_k with eigenvalues $+E_k$ and $-E_k$. The mode expansion

$$\Psi(x) = \int \frac{d^4 k}{(2\pi)^4} \sum_{s=0}^7 \left[b_s(k) u_s(k) e^{-iE_k x_4} + d_s(-k)^* v_s(k) e^{iE_k x_4} \right] e^{ik \cdot x},$$

$$\{b_s(k), b_{s'}(k')^*\} = \{d_s(k), d_{s'}(k')^*\} = \delta_{ss'} (2\pi)^4 \delta^4(k - k'),$$

with the Hilbert adjoints $*$ of the positive structure $J = B$, realizes $\{\Psi_a, \chi_b\} = \delta_{ab} \delta^4$ on a positive-definite Fock space. The vacuum $|0\rangle$ is annihilated by all $b_s(k)$ and $d_s(k)$: the negative-energy states are filled (Dirac sea), and d^* creates antiparticles. After normal ordering

$$H = \int \frac{d^4 k}{(2\pi)^4} E_k \sum_{s=0}^7 (b_s(k)^* b_s(k) + d_s(k)^* d_s(k)) \geq 0.$$

This construction is derived here from Results 10.2, 10.4 and 10.5; it is not a separate machine check.

10.8 Unitary and Krein-unitary symmetries

A spin transformation $\Psi \rightarrow R\Psi$ preserves the canonical anticommutator iff $RBR^\dagger = B$, equivalently $R^\dagger BR = B$ (Krein-unitary). It is implemented unitarily on the positive Fock space iff in addition R is unitary for $f^\dagger h$ and commutes with $J = B$ (derived). For $R = \exp(\theta S^{ab})$ the conditions concern the generators.

Result 10.6.

1. Exactly 13 of the 28 generators S^{ab} commute with B : the 6 of $\mathfrak{so}(4)$ ($a, b \in \{0, 1, 2, 3\}$), the 3 of $\mathfrak{so}(3)$ ($a, b \in \{5, 6, 7\}$) and the 4 boosts S^{a4} , $a \in \{0, 1, 2, 3\}$. Since $B \propto \gamma^0 \gamma^1 \gamma^2 \gamma^3 \gamma^4$, the boosts

S^{a4} anticommute with C and with γ^4 and therefore commute with B ; they are Hermitian, so $\exp(\theta S^{a4})$ is neither unitary nor Krein-unitary.

2. Exactly 9 generators are anti-Hermitian (unitary for $\Psi^\dagger \Psi$) and commute with B . They generate the unitarily implemented group $\text{Spin}(4) \times \text{Spin}(3)$. The literal claim "exactly 9 of the 28 S^{ab} commute with B " is false and is recorded as false in both reports.
3. Exactly 21 generators, those with $a, b \neq 4$, satisfy the Krein condition $(S^{ab})^\dagger B + B S^{ab} = 0$. They generate the Krein-unitary slice group $\text{Spin}(4,3)$. The Krein condition fails for all 7 generators S^{4b} .
4. Exactly 12 generators are anti-Hermitian for $\Psi^\dagger \Psi$ ($\text{so}(4)$ and $\text{so}(4)$ on the directions 4 to 7).

Both algebra reports use this meaning of the check, as recorded in `stage1-summary.json`. Time translations in the good sector are unitary (h_k Hermitian), and the $U(1)$ phase is unitary.

checks

`QNT_unitaryAndKreinSubgroups: wolfram-algebra, python-algebra`

10.9 The $U(1)$ current and charge

$J^\mu = -i\bar{\Psi}\gamma^\mu\Psi$ is Hermitian, and in Gaussian normal gauge $J^4 = \Psi^\dagger B\Psi$: all eight current matrices are Hermitian and the matrix of J^4 is B . The conserved charge is $Q = \int d^7x \sqrt{|g|} J^4$. As a quadratic form in classical spinors it is indefinite (signature (8,8)); in the Hilbert representation of Section 10.5, $\Psi^\dagger B\Psi = \chi\Psi$, and after normal ordering $Q = \sum(b^*b - d^*d)$: particles have charge +1 and antiparticles -1 (derived).

checks

`QNT_currentHermiticity: wolfram-algebra, python-algebra`

`GR_currentHermitian: grassmann-demo`

10.10 The extra-time sector

For momenta along x_5, x_6, x_7 the operator h_k is not Hermitian and does not commute with B (Result 10.5), and $E^2 = m^2 + k_0^2 + \dots + k_3^2 - k_5^2 - k_6^2 - k_7^2$ can be negative: two of the Python samples have $E^2 = -271/196$ and $E^2 = -3$. The corresponding frequencies are imaginary and the modes grow exponentially in x_4 . This is the ill-posedness of the initial-value problem for ultrahyperbolic equations: the slice $x_4 = \text{const}$ contains three time-like directions. The positive Fock space of Section 10.7 exists only in the good sector.

10.11 Expectation values

In the Hilbert representation the canonical bilinear is $\Psi^\dagger M\Psi = \chi B M\Psi$. For a normalized one-particle state with wave function u ($u^\dagger u = 1$, positive energy, normal ordering),

$$\langle \Psi^\dagger M\Psi \rangle = u^\dagger B M u.$$

Examples (derived): the charge ($M = B$) gives $u^\dagger u = 1$; the energy ($M = Bh$) gives $u^\dagger h u = E$; the scalar density ($M = C$) gives $u^\dagger B C u = u^\dagger (-i\gamma^4) u$, which is +1 for a positive-energy state at rest with $m > 0$, so $\langle \rho \rangle = m \langle S \rangle = m = E$. The naive density $\Psi^\dagger \Psi$ gives $u^\dagger B u$, which is indefinite even on positive-energy states (Result 10.4); it is not the physical density.

11 Verification records

All five reports have schemaVersion 1, and every check in them is true. The counts below are those of stage1-summary.json, which re-reads the five reports, re-compares the key measurements and records an empty list of disagreements:

report	producer	checks true
wolfram-algebra-report.json	scripts/verify_dirac16complex_algebra.wls	21 of 21
wolfram-geometry-report.json	scripts/verify_dirac16complex_geometry.wls	43 of 43
python-algebra-report.json	scripts/check_dirac16complex_algebra.py	21 of 21
python-geometry-report.json	scripts/check_dirac16complex_geometry.py	52 of 52
grassmann-demo-report.json	scripts/demo_grassmann_lagrangians.py	16 of 16
stage1-summary.json	scripts/build_stage1_summary.py	153 of 153

The producers record Wolfram Language 15.0.1 and Python 3.14.5, sympy 1.14.0, numpy 2.4.6.

11.1 WolframScript, algebra

The Wolfram algebra verifier uses exact integer, rational and Gaussian-rational arithmetic. Commutants, intertwiners and invariant forms are exact null spaces of integer coefficient matrices, and signatures come from exact characteristic polynomials. Files and checks:

```
wolfram/Dirac16ComplexAlgebra.wl
scripts/verify_dirac16complex_algebra.wls
```

```
wolfram-algebra: ALG_clifford ALG_gammaTransposeSymmetry ALG_chargeMatrix ALG_expression1
  ALG_spinTransposeProperties ALG_chirality ALG_faithful ALG_pinIrreducibleComplex
  ALG_spinDecomposition ALG_cliffordPictureIntertwiner ALG_octonionPictureIntertwiner
  ALG_chargeFormB ALG_invariantForms ALG_gamma8Map ALG_pinLiftCharacter
  QNT_curvedAnticommutatorMatrix QNT_flatModeHamiltonian QNT_kreinSignature
  QNT_unitaryAndKreinSubgroups QNT_currentHermiticity ALG_fixtureAgreement
```

11.2 WolframScript, geometry

Every derived object of the Wolfram geometry verifier is an exact order-1 jet (value and eight first derivatives) at the test point, and the G2 arithmetic is exact in $\mathbb{Q}(\cos z)$. The Euler-Lagrange, canonical-momentum and Lg[] checks use a genuine Grassmann algebra; the energy-momentum, Lichnerowicz and invariance checks use commuting exact numbers, which is exact for bilinears with $\Psi^\dagger$ on the left and Ψ on the right (measurement convention.commutingEvaluation). Files and checks:

```
wolfram/Dirac16ComplexGeometry.wl
scripts/verify_dirac16complex_geometry.wls
```

```
wolfram-geometry: ALG_chargeMatrix ALG_cliffordRelations ALG_grassmannLemmas
  EMT_conservation_G1 EMT_conservation_G2 EMT_homogeneousReduction_G3
  EMT_symmetricHermitian_G1 EMT_symmetricHermitian_G2 EMT_trace_G1 EMT_trace_G2
  EMT_variation_G3 GEO_curvature_G1 GEO_curvature_G2 GEO_diagonalSlashFormula_G2
  GEO_divergenceIdentity_G1 GEO_divergenceIdentity_G2 GEO_frameNondegenerate_G1
  GEO_frameNondegenerate_G2 GEO_gammaCovariantConstancy_G1
  GEO_gammaCovariantConstancy_G2 GEO_lichnerowicz_G1 GEO_lichnerowicz_G2
  GEO_notebookContractionFails_G1 GEO_notebookContractionFails_G2
  GEO_omegaAntisymmetry_G1 GEO_omegaAntisymmetry_G2 GEO_primordialInvariants_G2
  GEO_symbolicJetAgreement_G2 GEO_vielbeinPostulate_G1 GEO_vielbeinPostulate_G2
  LAG_eulerLagrangePsibar_G1 LAG_eulerLagrangePsibar_G2 LAG_eulerLagrangePsi_G1
  LAG_eulerLagrangePsi_G2 LAG_hermiticity_G1 LAG_hermiticity_G2
  LAG_localSpinInvariance_G1 LAG_localSpinInvariance_G2
  LAG_notebookLgGrassmannTrivial_G1 LAG_notebookLgGrassmannTrivial_G2
  NEG_notebookConnectionDetected_G2 QNT_canonicalMomentum_G1 QNT_canonicalMomentum_G2
```

11.3 Python, algebra

The Python algebra checker uses the standard library only (exact rationals, no floating point). Files and checks:

```
scripts/d16c_exact.py
scripts/build_dirac16complex_fixture.py
scripts/check_dirac16complex_algebra.py
```

```
python-algebra: ALG_clifford ALG_gammaTransposeSymmetry ALG_chargeMatrix ALG_expression1
  ALG_spinTransposeProperties ALG_chirality ALG_faithful ALG_pinIrreducibleComplex
  ALG_spinDecomposition ALG_cliffordPictureIntertwiner ALG_octonionPictureIntertwiner
  ALG_chargeFormB ALG_invariantForms ALG_gamma8Map ALG_pinLiftCharacter
  QNT_flatModeHamiltonian QNT_kreinSignature QNT_unitaryAndKreinSubgroups
  QNT_currentHermiticity ALG_fixtureAgreement ALG_wolframAgreement
```

11.4 Python, geometry

The Python geometry checker uses sympy with exact rationals: G1 by exact truncated Taylor jets, and G2 at a fully symbolic point in $\mathbb{Q}(w, c, E, A_1, A_2, A_3, H)$ with $w = \sin^{1/6} z$, $c = \cos z$, $E = e^{a_4}$ and A_k the derivatives of a_4 . The relation $w^{12} + c^2 = 1$ was needed in 0 of its zero tests. Files and checks:

```
scripts/d16c_geometry_sympy.py
scripts/check_dirac16complex_geometry.py
```

```
python-geometry: ALG_cliffordRelations ALG_sigma16EqualsGamma0123
  ALG_CSymmetricInvolution ALG_expression1CgammaAntisymmetric ALG_gammaSymmetryPattern
  ALG_CSabAntisymmetric ALG_chiralityDiag ALG_CAnticommutatorGammaSSymmetric
```

```

ALG_CCommutatorGammaSAntisymmetric ALG_SabGammaCommutator GEO_frameNondegenerate_G1
GEO_vielbeinPostulate_G1 GEO_omegaAntisymmetry_G1 GEO_gammaCovariantConstancy_G1
GEO_notebookContractionFails_G1 GEO_divergenceIdentity_G1
GEO_sqrtgSquaredEqualsDetg_G1 GEO_curvature_G1 GEO_lichnerowicz_G1
LAG_eulerLagrangePsibar_G1 LAG_eulerLagrangePsi_G1 EMT_onshellJetSolve_G1
EMT_symmetric_G1 EMT_trace_G1 EMT_traceOffShellIdentity_G1
EMT_onshellLagrangianSUprimeMinusU_G1 EMT_conservation_G1 GEO_vielbeinPostulate_G2
GEO_omegaAntisymmetry_G2 GEO_gammaCovariantConstancy_G2
GEO_notebookContractionFails_G2 GEO_divergenceIdentity_G2
GEO_sqrtgSquaredEqualsDetg_G2 GEO_curvature_G2
GEO_anticommutatorGammaOmegaVanishesDiagonal_G2 GEO_primordialInvariants_G2
GEO_diagonalSlashFormula_G2 GEO_lichnerowicz_G2 LAG_eulerLagrangePsibar_G2
LAG_eulerLagrangePsi_G2 EMT_onshellJetSolve_G2 EMT_symmetric_G2 EMT_trace_G2
EMT_traceOffShellIdentity_G2 EMT_onshellLagrangianSUprimeMinusU_G2
EMT_conservation_G2 GEO_lichnerowiczConstantSameG1G2 EMT_homogeneousReduction
GEO_diagonalSlashFormula_G3 EMT_homogeneousTimeSpaceVanishesOnShell_G3 EMT_variation
GEO_wolframAgreement

```

11.5 Python, Grassmann demonstrations

The demonstrations work in exact Grassmann algebras with 720 generators for real Ψ jets and 1440 for complex (Ψ, Ψ^*) jets. Files and checks:

```

scripts/grassmann_algebra.py
scripts/demo_grassmann_lagrangians.py

```

```

grassmann-demo: GR_conjugationRules GR_massTermVanishesReal
GR_bilinearOnlyAntisymmetricPartSurvives GR_kineticTotalDerivativeReal
GR_kineticSymmetricMatrixContrast GR_quarticTermPolynomial GR_scalarBilinearHermitian
GR_currentHermitian GR_notebookLgELTrivial GR_notebookLgPureDivergence
GR_complexLagrangianNonTrivial GR_complexQuarticEL GR_complexPsiEquation
GR_lagrangianHermitian GR_emtHermitian GR_unsymmetrizedKineticNotHermitian

```

11.6 Agreement between the implementations

The cross-implementation checks:

```

ALG_fixtureAgreement (wolfram-algebra):  Wolfram reads the Python fixture;
75 of 75 matrices agree by name
ALG_wolframAgreement (python-algebra):  Python reads the Wolfram algebra report;
K_clifford, K_octonion and the chirality agree; 20 shared check names
with equal verdicts; 47 of 47 shared measurements agree
GEO_wolframAgreement (python-geometry):  Python reads the Wolfram geometry report;
26 shared check names with equal verdicts;
84 of 84 shared measurements agree

```

The geometry comparison covers the G1 point data, the nonzero counts, the Lichnerowicz constant, the curvature sign, the Python G2 formulas evaluated at the exact Wolfram points, and the energy-momentum sign and trace conventions. The field-dependent values (traces, ρ , p , KE and PE at the

test points) are not compared between the implementations, because each draws its own random field data and parameters; they are compared through the formulas that both verify. stage1-summary.json records 30 agreed key measurements, the hash links between the reports, and the equivalences that hold after a notation conversion.

11.7 Files and hashes

sha256 of the reports, of the fixture and of the producers, as recorded in the reports and in stage1-summary.json:

```
artifacts/dirac16complex/arbitrary-field/wolfram-algebra-report.json
d43adeeba580bdd8cec56fac5fa74906b7578c06c8589e52dbe3da1404cd49e7
artifacts/dirac16complex/arbitrary-field/wolfram-geometry-report.json
cec9ee0d577c4e7f0e82efd404d412a503140ccb1547118ab48a78a3d773f0ac
artifacts/dirac16complex/arbitrary-field/python-algebra-report.json
3275b2af1a2f56f9bd779ad359e7ed078cb2c64c744863732f789f410bb50c97
artifacts/dirac16complex/arbitrary-field/python-geometry-report.json
deec576c1613bb06ce8ac4ad9f89a7b008239bd9d4b624bea14728fa77b2a197
artifacts/dirac16complex/arbitrary-field/grassmann-demo-report.json
b83997857ccfc9fb71f05efbcaebe16f29bf36687cc08c16e0823a26a6369f09
artifacts/dirac16complex/arbitrary-field/stage1-summary.json
a65972b5a52f947d9bee9f67e09f7151fff617c03f496ba313e07db168c10770
artifacts/dirac16complex/arbitrary-field/algebra-fixture.json
8b4f15462ca4d61ef6ec0e72f04c8a77d23ce8bcabc9b6ce19f921c90e02653b
wolfram/Dirac16ComplexAlgebra.wl
ba0d00c818fa3707f04f7b2a7478b0c635fbf9fdaaa08b2e036d2704180f2262
scripts/verify_dirac16complex_algebra.wls
86904ac07e1237cf0dec14431a3707d7bf24f951e12def504705d331c934949a
wolfram/Dirac16ComplexGeometry.wl
f5b674665eee4000750161e6ab6312c38bfac9da7a17450a2b3bdd88ef292af2
scripts/verify_dirac16complex_geometry.wls
81b1b0febf1a5d6956f042179dff4eed949718cc15955e0753c71606f3f09953
scripts/d16c_exact.py
eb2fcbe498b6ae927df61f73abcee4e58c319984fb8f6e1b28defcd0ca13a3f5
scripts/build_dirac16complex_fixture.py
585051a6b2e01ab1d8ae92ae5e1949c4d0c23e37815332c44dc5af525a36d776
scripts/check_dirac16complex_algebra.py
3d2fd639b655491897b68f8719db3618d638248b2b1176f22668723e273bea12
scripts/d16c_geometry_sympy.py
4f7eca4cf247354650ed9beb0b9c46530d14c9965f095979ea5ee24651af9f25
scripts/check_dirac16complex_geometry.py
ac26111ce6758cebdf473e943d1cd4f1e14f87e4961ee694623d888742824653
scripts/grassmann_algebra.py
dfce851d3dc7ab82202a62d414cc259644a64a101f08e9ba80ed747faa8bc24c
scripts/demo_grassmann_lagrangians.py
3053c7011037e29cd9efa7c2067aafd23ebc67c6156eb581fef184810ef1515a
dirac-main/artifacts/exact/cl44-seed.json
52e7a73b14467435d7986b33bf29547510b321853ffbd76121896d70b796d133
dirac-main/artifacts/exact/split-octonion.json
cbc28db6709ec1fc35ccfcc26353d1952b42bb516f6c50d6a52e2d6d355be710
dirac-main/artifacts/exact/triality44.json
a8c31c07809fc884ae0495d91f4105fac0909c954bb185c643306c57699cd395
```

12 Reproduction

Run every command from the repository root. The Wolfram report path is a plain positional argument: WolframScript 1.14 drops "-" and every argument after it when it is combined with -file. The gates set PYTHONUTF8=1 for their own process; set it yourself for single steps, because Python output contains Greek letters.

12.1 PowerShell

The Stage-1 gate runs all eleven steps through the logging wrapper scripts/run_logged.ps1, with the logs in build/logs/, and then audits the five reports and the cross-implementation checks:

```
pwsh -NoProfile -File scripts/verify_stage1_arbitrary_field.ps1
```

Its last line is

```
stage1_arbitrary_field_verification=OK
```

The individual steps:

```
$env:PYTHONUTF8 = "1"
$a = "artifacts/dirac16complex/arbitrary-field"
python scripts/build_dirac16complex_fixture.py
python scripts/check_dirac16complex_algebra.py --wolfram-report=
wolframscript -file scripts/verify_dirac16complex_algebra.wls '
    "$a/wolfram-algebra-report.json"
wolframscript -file scripts/verify_dirac16complex_geometry.wls '
    "$a/wolfram-geometry-report.json"
python scripts/check_dirac16complex_geometry.py --wolfram-report '
    "$a/wolfram-geometry-report.json"
python scripts/demo_grassmann_lagrangians.py
python scripts/check_dirac16complex_algebra.py --wolfram-report '
    "$a/wolfram-algebra-report.json"
python -m unittest discover -s tests -p "test_d16c_[ag]*.py" -v
python -m unittest discover -s tests -p "test_publication_tooling.py" -v
python scripts/build_stage1_summary.py --output "$a/stage1-summary.json"
python scripts/build_provenance_pdf.py provenance/DIRAC16COMPLEX_ARBITRARY_FIELD.md
```

12.2 Git Bash

The Git Bash twin runs the same steps through scripts/run_logged.sh and ends with the same line:

```
bash scripts/verify_stage1_arbitrary_field.sh
```

The individual steps:

```

export PYTHONUTF8=1
a=artifacts/dirac16complex/arbitrary-field
python scripts/build_dirac16complex_fixture.py
python scripts/check_dirac16complex_algebra.py --wolfram-report=
wolframscript -file scripts/verify_dirac16complex_algebra.wls \
    "$a/wolfram-algebra-report.json"
wolframscript -file scripts/verify_dirac16complex_geometry.wls \
    "$a/wolfram-geometry-report.json"
python scripts/check_dirac16complex_geometry.py --wolfram-report \
    "$a/wolfram-geometry-report.json"
python scripts/demo_grassmann_lagrangians.py
python scripts/check_dirac16complex_algebra.py --wolfram-report \
    "$a/wolfram-algebra-report.json"
python -m unittest discover -s tests -p "test_d16c_[ag]*.py" -v
python -m unittest discover -s tests -p "test_publication_tooling.py" -v
python scripts/build_stage1_summary.py --output "$a/stage1-summary.json"
python scripts/build_provenance_pdf.py provenance/DIRAC16COMPLEX_ARBITRARY_FIELD.md

```

Each verifier prints `check_count` and `failed_check_count` and exits nonzero if a check fails. The first algebra step runs without the Wolfram comparison (`--wolfram-report=` with an empty value), because the Wolfram report on disk may predate the run; the second one compares with the report just written. The PDF step builds this document twice (two builder runs, three pdflatex passes each), requires warning-free logs and byte-identical PDFs, and compares the result with the registered edition `dirac16complex-arbitrary-field` in `provenance/pdf-specifications.json`. After an edit of this document the edition is registered again with

```

python scripts/build_provenance_pdf.py --register \
    provenance/DIRAC16COMPLEX_ARBITRARY_FIELD.md

```

(in PowerShell the line continuation is a backtick instead of the backslash), and the sha256 pins in `tests/test_d16c_arbitrary_field_publication.py` are updated.

13 Limitations

1. The general statements of Sections 5 to 9 are proved by the arguments given here and checked exactly at test points: three points of one generic non-diagonal vielbein (G1), the primordial family (G2, symbolic in Python) and three times of a diagonal frame (G3). The checks are not a symbolic verification for every vielbein.
2. The metric variation of the action is carried out explicitly only for diagonal metrics. The off-diagonal components of $T_{\mu\nu}$ are tested through symmetry, Hermiticity and on-shell conservation in G1, and through the homogeneous forms of T_{ij} and T_{4i} .
3. Conservation of $T_{\mu\nu}$ is checked with on-shell jets at points, not as an off-shell Noether identity.
4. The energy-momentum, Lichnerowicz and invariance checks evaluate bilinears with commuting exact numbers, which is exact because $\Psi^\dagger$ is always on the left and Ψ on the right; the field equations, the canonical momentum and $\text{Lg}[]$ are checked in genuine Grassmann algebras.

5. The Dirac-bracket computation, the Hamiltonian density, the Heisenberg equations, the mode expansion, the Fock space and the expectation-value rule of Section 10 are derivations; only their algebraic ingredients (Results 10.1 to 10.6) are machine-checked.
6. The quantization is formal: the state space is a Krein space, the extra-time sector has complex frequencies, and no interacting theory, regularization or renormalization is constructed.
7. The homogeneous-sector formulas treat bilinears as classical mean-field quantities; the quantum expectation values of Section 10.11 are one-particle statements.
8. The signature (4,4) is not observed spacetime, and nothing here is compared with data. The connection to dark energy or dark matter, and the stability of solutions, are not decided in this document.
9. The cross-checks against dirac-main's published fixtures need the git-ignored dirac-main folder; in a fresh clone these sub-checks record "not-run" (measurement referenceFilesPresent in wolfram-algebra-report.json).