Provenance of the README of Pre-gravity6plane

- Every equation of the README is a restatement of a result computed in 6planes.nb (Wolfram Language 15.0.1, run 2026-10-30, 34 input cells, 0 messages) or verified by verify_dirac_matrices.wls (31 exact checks, all True): $T^A T^B + T^B T^A = 2\eta^{AB} I_{16}$; $\Gamma = \mathrm{diag}(-I_8, I_8)$; $\mathcal{L}^* = \mathcal{L}$; the Euler-Lagrange equations $T^A \partial_A \psi = (m + \lambda\bar{\psi}\psi)\psi$ and $\gamma^\mu D_\mu \psi = (m + \lambda\bar{\psi}\psi)\psi$; $K = -i\sigma_{16} T^4$ with eigenvalues ± 1 (8 each); the coupling blocks; $(6p+q)(6p-q) = m^2/H^2$; rank 2 and 28 parameterizations of M^6; $\mathcal{L}_m[\Gamma\psi^*] = \mathcal{L}_{-m}[\psi]$; $T^0 h_{KS} T^0 = -h_{KS}$; the zero energy of pair creation; the bridge data and $R \approx -\frac{31}{216 H^2 x_0^4}$; the occupation $1/(e^{\epsilon/T_H} + 1)$.

- The matrices T^A, σ_{16}, Γ, $P_{L,R}$ are those of Pre-gravity6plane.nb (P. L. Nash, (C) 2022, GPL), compared entry by entry with its stored output cells 494, 310, 312, 379, 381.

- The primordial vielbein is the one displayed in Pre-gravity6plane.nb (text cells 901, 913) with $a_4(Hx_4) = Hx_4$; the modern and future fields and the interpretation of "Universe" are definitions of this work, listed in Section 6 of the README.

- Literature: A. Einstein and N. Rosen, Phys. Rev. 48, 73 (1935); W. Kohn and L. J. Sham, Phys. Rev. 140, A1133 (1965); N. D. Mermin, Phys. Rev. 137, A1441 (1965); W. G. Unruh, Phys. Rev. D 14, 870 (1976); J. B. Hartle and S. W. Hawking, Phys. Rev. D 13, 2188 (1976); P. L. Nash, J. Math. Phys. 51, 042501 (2010).

- Every equation of Section 5 of the README is a result computed in deflation.nb (Wolfram Language 15.0.1, run 2026-10-30, 17 input cells, 0 messages): the line element with $g_{jj} = b_{jj} e^{-a x_4}$; the proper sizes $\sqrt{b_{jj}} e^{-a x_4/2}$; the x_0-only dependence of the Ricci scalar, Einstein tensor and Kretschmann scalar; the identity $-dx_4^2 - \sum_j g_{jj} dx_j^2 = -\frac{4}{a^2}(dw^2 + dy_5^2 + dy_6^2 + dy_7^2)/w^2$ with $w = \frac{2}{a} e^{a x_4/2}$, $y_j = \sqrt{b_{jj}} x_j$; $R = 3a^2$ and Kretschmann scalar $3a^4/2$ of the pure deflation field; the turning point $x_4^* = \frac{1}{a} \ln(b_{55}/p_5^2)$; the field equation of dirac16complex; the dispersion relation for ω^2 and $a_c = 2H$; $M^\dagger K + K M = 0$; and the -mass fraction $1.2 \times 10^{-31} \to 0.49999995$ of the numerical example ($a = b_{55} = k_1 = m = 1$, $q_5 = 1/2$, $-8 \le x_4 \le 5$).

- The deflation request and its clarifications are the user's (2026-10-30): $g_{jj} = b_{jj} e^{-a x_4}$ with $a > 0$, $b_{jj} > 0$, entering the line element as $-g_{jj} dx_j^2$, with the primordial field for all other components.