Symbols used in the README of Pre-gravity6plane

symbol	definition
$x_0..x_7$	coordinates: hidden, space (1-3), time (4), superluminal time (5-7)
η	$\mathrm{diag}(1,1,1,1,-1,-1,-1,-1)$
τ_A, $\bar{\tau}_A$	real 8x8 matrices of Pre-gravity6plane.nb, $\bar{\tau}_A = \sigma\tau_A^T\sigma$, $\sigma = \begin{pmatrix} 0 & I_4 \\ I_4 & 0 \end{pmatrix}$
$T^A = (T16^A)[A]$	$\begin{pmatrix} 0 & \bar{\tau}_A \\ \tau_A & 0 \end{pmatrix}$, real Dirac matrices
S^{AB}	$\frac{1}{4}[T^A, T^B]$
Γ, $P_{L,R}$, σ_{16}	$T^0\cdots T^7$; $\frac{1}{2}(I_{16} \mp \Gamma)$; $T^0T^1T^2T^3$
ψ, $\bar{\psi}$	dirac16complex (complex Grassmann 16-spinor), $\bar{\psi} = \psi^\dagger\sigma_{16}$
$\mathcal{L}$, m, λ	Lagrangian, mass, quartic coupling
K	$-i\sigma_{16}T^4$
$\psi_\pm$, $S_\pm$, $\mathcal{L}_D$	$K\psi_\pm = \pm\psi_\pm$; $S_\pm = \psi_\pm^\dagger(-iT^4)\psi_\pm$; $\mathcal{L}_D[\chi] = \chi^\dagger(-iT^4)(T^A\partial_A - m)\chi$
M, H	mass and Hubble parameters of Pre-gravity6plane.nb; $m = -2M$
z, t, p, q	$z = 6Hx_0$, $t = Hx_4$, mode $e^{pz+qt}u$
${F_A}^a$, ${F_a}^A$, V_1, V_2	triality frames and vector forms: $V_1 = \eta(\psi_1^T\sigma{F_A}^a)$, $V_2 = {F_a}^A\psi_2$
n_k, Z, M^6	$n_k = V_k[\mathrm{Re}\,\psi]$; $Z \in \mathbb{R}^8$; $M^6 = \{Z : n_1\cdot Z = n_2\cdot Z = 0\}$
f, r_s, H_Λ	$1 - r_s/x_0$ (modern), $1 - r_s/x_0 - H_\Lambda^2 x_0^2$ (future)
γ^μ, D_μ	${E_A}^\mu T^A$; $\partial_\mu + \frac{1}{4}{\omega_\mu}^A{}_B T_A T^B$
ρ_s, h_{KS}, α, β, p, B	Kohn-Sham density, $\alpha p + BM\beta$, $-T^4T^0$, $-iT^4$, $-i\,d/d\rho$, lapse e_4
$\Omega(T)$	Kohn-Sham grand potential at temperature T
R_1	$x_1 \mapsto -x_1$
R, κ, T_H, ϵ	scalar curvature, surface gravity, $\kappa/2\pi$, Killing energy
I_n	$n \times n$ identity matrix
g_{jj}, b_{jj}, a	deflating metric functions $g_{jj} = b_{jj}e^{-ax_4}$ $(j = 5,6,7)$, $a > 0$, $b_{jj} > 0$
w, y_j	coordinates of the deflating block: $w = \frac{2}{a}e^{ax_4/2}$, $y_j = \sqrt{b_{jj}}\,x_j$
$\mathbb{R}^4$	flat 4-space of x_0, x_1, x_2, x_3
ℓ, p_5, x_4^*	affine parameter; $p_5 = -g_{55}\,dx_5/d\ell$; turning point $\frac{1}{a}\ln(b_{55}/p_5^2)$
ω, $k_0..k_3$, q_5, q_6, q_7	frequency and wave numbers of a universe $e^{i(k_0x_0+\cdots+q_7x_7-\omega x_4)}u$ (the indexed q_j are wave numbers, not the exponent q); u a constant spinor
a_c	critical deflation rate $2H$
χ, M, c_A	mode spinor with $\chi' = d\chi/dx_4 = M\chi$; $M = T^4(i\sum_{A\neq 4} c_A T^A - m)$; real coefficients c_A
$P_\pm$	$\chi^\dagger\frac{1}{2}(I_{16} \pm K)\chi$, weights of the +mass and -mass sectors