Dirac matrices, Lagrangians and field equations on the deflating 4+4 and 7+1 spacetimes

With mass term and self-interaction. PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C: definitions, derivations, exact checks, and how to repeat them

Patrick Lee Nash, Ph.D.

November 13, 2026

0. What this document contains

This document defines the 16-component Dirac field on two eight-dimensional spacetimes whose coordinates $x5$, $x6$, $x7$ deflate exponentially, and derives its equations. It gives, for each of the two spacetimes,

1. the metric, its curvature and its symmetries (Part 1);
2. the eight real 16×16 Dirac matrices, printed in full, and the spin connection (Part 2);
3. the Lagrangian with its three parts: the kinetic term, the **mass term** and the **self-interaction**, which is quartic in the field (Part 3);
4. the field equation that follows from it, the change of normalization that removes the deflation from the equation of fields that do not depend on the deflating coordinates, and the conserved current (Part 4);
5. on the 4+4 spacetime, the two forms that the self-interaction can take once the theory is quantized: **form C**, the covariant form, which is the definition of the 4+4 model, and **form Q**, the quantized-sector form, which is the alternative; both are written out completely, with the exact relation between them (Part 5);
6. the algebra of the self-interaction: of its nine couplings only four are independent, and uniform matter and a pair of particles feel fixed combinations of them, which on 4+4 depend on the form; the consequences of the two forms for matter and for the vacuum, in numbers; and a discussion of the two forms (Part 6);
7. the solutions of the equation without self-interaction, mode by mode (Part 7), and the symmetries that survive the deflation (Part 8).

It is written for a reader who knows the Dirac equation of four-dimensional flat spacetime and the elements of general relativity, and nothing of this project. Every symbol is defined where it first appears, and Appendix B lists every symbol of the project with its meaning. Every equation is derived here or verified here; no other document is needed to read this one. Statements marked with an identifier such as `71d-C3` are verified by a program, exactly (in integer or rational arithmetic) unless the word “numerical” is used; Appendix A prints the complete output of these programs, and Section 0.3 says how to run them.

0.1 The hypotheses

The eight coordinates are written $x0, x1, \ldots, x7$, with the digit as part of the name. $x1, x2, x3$ are the coordinates of ordinary space and $x4$ is the ordinary time. $x0$ is a fourth, hidden space-like coordinate. The remaining three, $x5, x6, x7$, are extra coordinates, and the two models differ in their character:

	4+4 spacetime	7+1 spacetime
space-like coordinates	$x0, x1, x2, x3$	$x0, x1, x2, x3, x5, x6, x7$
time-like coordinates	$x4, x5, x6, x7$	$x4$
the Dirac field	ψ_{1644d}	ψ_{1671d}

Deflation. The author's hypothesis is that each of the three extra coordinates deflates exponentially in the ordinary time: in place of flat spacetime there is a pseudo-Riemannian space with a diagonal metric whose components along $x5, x6, x7$ are

$$\text{4+4:}\quad g_{tt}^{(4+4)} = -\,\text{const}_t^{\,2}\, e^{-2\,\text{rate}_t\, x4}, \qquad \text{7+1:}\quad g_{tt}^{(7+1)} = +\,\text{const}_t^{\,2}\, e^{-2\,\text{rate}_t\, x4}, \qquad t = 5, 6, 7. \tag{0.1}$$

Here $\text{rate}_5 = \text{g55}$, $\text{rate}_6 = \text{g66}$ and $\text{rate}_7 = \text{g77}$ are three positive constants, the **rates of deflation** (the names g55, g66, g77 are the author's; they are written upright and never carry a subscript), and const_5, const_6, const_7 are three positive constants that may differ from one another and between the two spacetimes. The superscript $(4+4)$ or $(7+1)$ marks a component of the metric, so that the component $g_{55}^{(4+4)}$ cannot be taken for the rate g55. The other five diagonal components are those of flat spacetime.

The field. The field has sixteen complex components. Its Lagrangian contains a mass term, bilinear in the field, and a self-interaction, quartic in the field; a theory of a scalar field ϕ would call them the "ϕ^2" and the "ϕ^4" term.

0.2 Conventions

- Units: $\hbar = c = 1$.
- **Sign of the metric:** space-like components are positive, time-like components are negative, as in Eq. (0.1).
- Greek indices $\mu, \nu, \rho, \sigma, \lambda$ label the coordinates and take the values $0, \ldots, 7$. Capital indices A, B, C label the directions of an orthonormal frame (Section 1.1) and take the same values. The index j takes the values $0, 1, 2, 3$ (the flat directions), and the indices t, s take the values $5, 6, 7$ (the deflating directions).
- Repeated Greek or capital indices, one up and one down, are summed. An index t or s is summed only where a sum sign says so.
- $\partial_\mu = \partial/\partial x\mu$; a prime is not used for derivatives.
- 1 denotes a unit matrix of the size that the context requires, 1_8 the 8×8 unit matrix; M^T is the transpose of a matrix M, $M^\dagger$ its Hermitian conjugate, $[M, N] = MN - NM$.
- Where a statement holds on both spacetimes with different signs, the sign is written with the constants c_γ and gs_{tt} of the following table.

constant	meaning	4+4	7+1
gs_{AA}	sign of the component AA of the flat metric	$+1$ for $A = 0, 1, 2, 3$; -1 for $A = 4, 5, 6, 7$	-1 for $A = 4$; $+1$ otherwise
gs_{tt}	that sign for the deflating directions	-1	$+1$
c_γ	sign in the anticommutator of the Dirac matrices, Eq. (2.1)	$+1$	-1

On both spacetimes $c_\gamma\, \text{gs}_{tt} = -1$ and $\text{gs}_{44} = -1$; several results below are the same on both for this reason.

0.3 How to repeat every result of this document

All commands are run from the folder `Revision_Deflation_CJT_and_Kohn-Sham_definition_C` of the repository `https://github.com/once-ere/spacetimes`. They need a Rust toolchain (`cargo`) and Python 3 with the packages numpy, scipy, sympy, mpmath and matplotlib; the last two lines also need pandoc and XeLaTeX.

```
# 1. compile the program and run its unit tests
cargo test  --release --manifest-path code/deflation-cjt-ks-c-rs/Cargo.toml
cargo build --release --manifest-path code/deflation-cjt-ks-c-rs/Cargo.toml
B=code/deflation-cjt-ks-c-rs/target/release/deflation_cjt_ks_c

# 2. geometry, matrices, spin connection, Dirac operators (Parts 1, 2, 4)
$B matrices
# 3. the self-interaction (Parts 3, 5, 6); the two forms on 4+4 in numbers (Section 6.6)
$B selfcoupling
$B forms
# 4. the solutions mode by mode (Part 7) and the symmetries (Part 8)
$B modes
$B symmetry
# 4a. the stability of the solution of Section 1.4
$B gravity

# 5. independent verification with computer algebra and other numerical tools
python3 code/scripts/verify_deflating_independent.py
python3 code/scripts/verify_cjt_ks_independent.py

# 6. the figures, and this document (Markdown, LaTeX, PDF)
python3 code/scripts/make_figures.py
python3 code/scripts/write_symbol_registry.py
python3 code/scripts/write_doc_tables.py
python3 code/scripts/assemble_docs.py
bash code/scripts/build_docs.sh "dirac matrices deflating.md"
```

What each step does. Step 1 compiles the program `deflation_cjt_ks_c` (written in Rust) and runs its unit tests, which must end with `test result: ok`; the third line abbreviates the path of the program. Each command of steps 2 to 4 computes, writes its data files into the folder `data`, and ends with a line that counts its checks, for example `forms: 9 of 9 checks passed`; the complete reports are the files `data/deflating_checks.txt`, `data/flat_matrix_checks.txt`, `data/selfcoupling_checks.txt`, `data/forms_checks.txt`, `data/mode_checks.txt`, `data/symmetry_checks.txt` and `data/gravity_checks.txt`, printed in Appendix A. (The command `forms` takes about ten minutes; the others take seconds to about a minute.) Step 5 repeats the central results with other tools (computer algebra for the geometry and for the algebra of the self-interaction, a general-purpose integrator and another library of special functions for the modes). Step 6 draws the figures from the data files and builds this document; the tables, the printed matrices and the program outputs in it are inserted by the scripts, not typed.

The source files in which each calculation is written: `metric.rs`, `geometry.rs`, `curved.rs` (geometry, spin connection, Dirac operators); `selfint.rs`, `cmd_selfcoupling.rs` (self-interaction); `halves.rs`, `cmd_forms.rs`, `cmd_forms_b.rs` (the two forms on 4+4: matter of two halves, vacuum); `modes.rs` (solutions); `symmetry_defl.rs`, `cmd_symmetry.rs` (symmetries); all in `code/deflation-cjt-ks-c-rs/src/`. Each file begins with a description of what it computes and how.

Part 1. Geometry of the deflating spacetimes

1.1 Metric, orthonormal frame, volume

The flat metrics of the two spacetimes are, in the coordinates $x0, \dots, x7$,

$$\begin{aligned}\eta^{(4+4)} &= \mathrm{diag}(+1,+1,+1,+1,-1,-1,-1,-1),\\ \eta^{(7+1)} &= \mathrm{diag}(+1,+1,+1,+1,-1,+1,+1,+1).\end{aligned} \tag{1.1}$$

Their diagonal entries are the signs gs_{AA} of Section 0.2. Define three **scale factors**, one for each deflating coordinate,

$$b_t(x4) = \mathrm{const}_t\, e^{-\mathrm{rate}_t\, x4}, \qquad t = 5,6,7. \tag{1.2}$$

The metric of the deflating spacetime is diagonal:

$$\begin{aligned}g_{\mu\nu}^{(4+4)} &= \mathrm{diag}(\,+1,+1,+1,+1,-1,-b_5^2,-b_6^2,-b_7^2),\\ g_{\mu\nu}^{(7+1)} &= \mathrm{diag}(\,+1,+1,+1,+1,-1,+b_5^2,+b_6^2,+b_7^2).\end{aligned} \tag{1.3}$$

Its last three components are those of the hypothesis (0.1) (checks `44d-G1`, `71d-G1`). In one formula for both spacetimes, $g_{tt} = \mathrm{gs}_{tt}\, b_t^2$. A coordinate interval dxt has the proper extent $b_t\,|dxt|$, which shrinks by the factor e in the time $1/\mathrm{rate}_t$: this is the meaning of "the coordinate xt deflates". On the 7+1 spacetime, where $x5, x6, x7$ are space-like, what shrinks is a length; on the 4+4 spacetime, where they are time-like, it is a duration measured along an additional time. The inverse metric is written $g^{\mu\nu}_{(4+4)}$, $g^{\mu\nu}_{(7+1)}$; its components along the deflating coordinates are gs_{tt}/b_t^2.

Frame. At each point choose eight vector fields $e_A = e_A{}^\mu\,\partial_\mu$ that are orthonormal, $g_{\mu\nu}\, e_A{}^\mu e_B{}^\nu = \eta_{AB}$. We take them along the coordinates:

$$e_A{}^\mu = \mathrm{diag}\Big(1,1,1,1,1,\frac{1}{b_5},\frac{1}{b_6},\frac{1}{b_7}\Big), \qquad e^A{}_\mu = \mathrm{diag}(1,1,1,1,1,b_5,b_6,b_7), \tag{1.4}$$

where $e^A{}_\mu$ is the inverse matrix. Frame indices are raised and lowered with the flat metric (1.1), coordinate indices with the metric (1.3).

Volume. The invariant volume element is $V_d\, d^8x$ with

$$V_d(x4) = \sqrt{|\det g|} = b_5\, b_6\, b_7 \;\propto\; e^{-\mathrm{gsum}\, x4}, \qquad \mathrm{gsum} = \mathrm{g55} + \mathrm{g66} + \mathrm{g77}. \tag{1.5}$$

V_d is the proper volume of a unit coordinate cell of the three deflating coordinates, and the **total rate** gsum is the rate at which it shrinks. Two further combinations of the rates occur:

$$\mathrm{gsq} = \mathrm{g55}^2 + \mathrm{g66}^2 + \mathrm{g77}^2, \qquad \mathrm{gprod} = \mathrm{g55}\,\mathrm{g66} + \mathrm{g55}\,\mathrm{g77} + \mathrm{g66}\,\mathrm{g77}, \qquad \mathrm{gsum}^2 = \mathrm{gsq} + 2\,\mathrm{gprod}. \tag{1.6}$$

1.2 Christoffel symbols

The Christoffel symbols are $\Gamma^\rho{}_{\mu\nu} = \frac{1}{2}\, g^{\rho\sigma}(\partial_\mu g_{\sigma\nu} + \partial_\nu g_{\sigma\mu} - \partial_\sigma g_{\mu\nu})$. The only components of the metric that depend on a coordinate are the three g_{tt}, and they depend on $x4$ alone, with $\partial_4 g_{tt} = -2\,\mathrm{rate}_t\, g_{tt}$. A Christoffel symbol therefore differs from zero only if it contains the index pair tt and the index 4. With the components g_{tt} and $g^{tt} = 1/g_{tt}$ of the metric (1.3) of either spacetime,

$$\Gamma^t{}_{4t} = \Gamma^t{}_{t4} = \tfrac{1}{2}\, g^{tt}\,\partial_4 g_{tt} = -\,\mathrm{rate}_t, \qquad \Gamma^4{}_{tt} = -\tfrac{1}{2}\, g^{44}\,\partial_4 g_{tt} = -\,\mathrm{gs}_{tt}\,\mathrm{rate}_t\, b_t^2, \tag{1.7}$$

without summation over t, where $g^{44} = -1$ was used. Thus $\Gamma^4{}_{tt} = +\mathrm{rate}_t\, b_t^2$ on the 4+4 spacetime and $-\mathrm{rate}_t\, b_t^2$ on the 7+1 spacetime; all other symbols vanish (checks `44d-G2`, `71d-G2`).

1.3 Curvature

Riemann tensor. With the convention $R^{\rho}{}_{\sigma\mu\nu} = \partial_\mu \Gamma^{\rho}{}_{\nu\sigma} - \partial_\nu \Gamma^{\rho}{}_{\mu\sigma} + \Gamma^{\rho}{}_{\mu\lambda}\Gamma^{\lambda}{}_{\nu\sigma} - \Gamma^{\rho}{}_{\nu\lambda}\Gamma^{\lambda}{}_{\mu\sigma}$ the components in the orthonormal frame that do not vanish are

$$R^{t}{}_{4t4} = -\,\mathrm{rate}_t^{\,2}, \qquad R^{4}{}_{t4t} = \mathrm{gs}_{tt}\,\mathrm{rate}_t^{\,2}, \qquad R^{t}{}_{sts} = \mathrm{gs}_{ss}\,\mathrm{rate}_t\,\mathrm{rate}_s \quad (t \neq s), \tag{1.8}$$

and those that follow from the antisymmetry in the last two indices (checks `44d-G4`, `71d-G4`). No component carries an index $0, 1, 2, 3$: *the four coordinates $x0, x1, x2, x3$ span a flat space, and the curvature lives in the four coordinates $x4, x5, x6, x7$.* The program obtains the curvature in two independent ways, from the Christoffel symbols and from Cartan's structure equations for the frame (checks `44d-G3`, `71d-G3`).

Sectional curvatures. For a pair of frame directions $A \neq B$ the number $K_{AB} = K_{BA} = R^{AB}{}_{AB}$ (first index raised with the flat metric, no summation) is the sectional curvature of the plane that the two directions span. From Eq. (1.8), on both spacetimes,

$$K_{4t} = \mathrm{rate}_t^{\,2}, \qquad K_{ts} = \mathrm{rate}_t\,\mathrm{rate}_s \quad (t \neq s), \qquad K_{AB} = 0 \text{ if } A \text{ or } B \in \{0,1,2,3\}. \tag{1.9}$$

(For example $K_{4t} = R^{4t}{}_{4t} = \mathrm{gs}_{tt}\,R^{4}{}_{t4t} = \mathrm{gs}_{tt}^2\,\mathrm{rate}_t^{\,2}$.) The Riemann tensor of this metric has no other components: it is "diagonal on pairs of directions".

Ricci tensor, curvature scalar, Einstein tensor. For such a Riemann tensor the mixed components of the Ricci tensor are $R^{A}{}_{A} = \sum_{B\neq A} K_{AB}$, the curvature scalar is $R = \sum_{A\neq B} K_{AB}$ (each pair counted twice), and the Einstein tensor $\mathcal{G}^{\mu}{}_{\nu} = R^{\mu}{}_{\nu} - \frac{1}{2} R\,\delta^{\mu}_{\nu}$ is diagonal with $\mathcal{G}^{A}{}_{A} = -\sum K_{BC}$, the sum running over the pairs $\{B, C\}$ of directions that do not contain A. With Eq. (1.9):

$$R^{4}{}_{4} = \mathrm{gsq}, \qquad R^{t}{}_{t} = \mathrm{rate}_t\,\mathrm{gsum}, \qquad R^{j}{}_{j} = 0, \qquad R = \mathrm{gsq} + \mathrm{gsum}^2, \tag{1.10}$$

$$\mathcal{G}^{4}{}_{4} = -\,\mathrm{gprod}, \qquad \mathcal{G}^{j}{}_{j} = -\tfrac{1}{2}(\mathrm{gsq} + \mathrm{gsum}^2), \qquad \mathcal{G}^{t}{}_{t} = \mathrm{rate}_t\,\mathrm{gsum} - \tfrac{1}{2}(\mathrm{gsq} + \mathrm{gsum}^2), \tag{1.11}$$

the same on both spacetimes in the sign convention of this document (checks `44d-G5`, `71d-G5`). All are constants: the curvature does not depend on the time.

1.4 The deflating spacetimes are not solutions of Einstein's equations without matter

The Einstein tensor (1.11) does not vanish, and it is not a multiple of the unit matrix unless all rates vanish: *the deflating spacetime is not a solution of Einstein's equations without matter, with or without a cosmological constant.* Something must sustain the deflation.

In more than four dimensions the general field equations of second order for the metric are those of Lovelock. Their first term beyond Einstein's is the Lanczos tensor $\mathcal{H}^{\mu}{}_{\nu}$ of the Gauss–Bonnet term, and the equations without matter read

$$\mathcal{G}^{\mu}{}_{\nu} + \Lambda_8\,\delta^{\mu}_{\nu} + \alpha_{\mathrm{GB}}\,\mathcal{H}^{\mu}{}_{\nu} = 0 \tag{1.12}$$

with a cosmological constant Λ_8 and a coupling α_{GB}. For a Riemann tensor that is diagonal on pairs of directions, $\mathcal{H}^{A}{}_{A} = -4\sum K_{BC}\,K_{B'C'}$, where the sum runs over the unordered couples of two pairs $\{B, C\}$, $\{B', C'\}$ that have no direction in common and do not contain A. A couple contributes only if its four directions are $4, 5, 6, 7$, so that

$$\mathcal{H}^{j}{}_{j} = -4\left(K_{45}K_{67} + K_{46}K_{57} + K_{47}K_{56}\right) = -4\,\mathrm{g55}\,\mathrm{g66}\,\mathrm{g77}\,\mathrm{gsum}, \qquad \mathcal{H}^{4}{}_{4} = \mathcal{H}^{t}{}_{t} = 0. \quad (1.13)$$

The components of Eq. (1.12) along $x4, \dots, x7$ contain no Gauss–Bonnet term and require $\mathcal{G}^{4}{}_{4} = \mathcal{G}^{5}{}_{5} = \mathcal{G}^{6}{}_{6} = \mathcal{G}^{7}{}_{7}$; since $\mathcal{G}^{5}{}_{5} - \mathcal{G}^{6}{}_{6} = (\mathrm{g55} - \mathrm{g66})\,\mathrm{gsum}$, this holds if and only if the three rates are equal. The components along $x0, \dots, x3$ then fix the coupling:

$$\mathrm{g55} = \mathrm{g66} = \mathrm{g77}: \qquad \Lambda_8 = 3\,\mathrm{g55}^2, \qquad \alpha_{\mathrm{GB}} = -\frac{1}{4\,\mathrm{g55}^2}. \quad (1.14)$$

With three equal rates the deflating spacetime is an exact solution of Einstein–Gauss–Bonnet gravity without matter; with unequal rates it is not, for any Λ_8 *and* α_{GB} (checks `44d-G10`, `71d-G10`; an independent script obtains the same Lanczos tensor from its definition by computer algebra). This solution is not stable. When the common rate of expansion of the four flat directions and the common rate of the three deflating directions are allowed to vary, the constraint among the initial data admits, near the solution, no deflation slower than g55; and for every perturbation that it does admit, with or without matter, the deflation departs from the exponential law and becomes faster and faster (check `gr-G5` of the command `gravity`, numerical; Appendix A.7 prints the report). What sustains the deflation is therefore an open question of the hypothesis, not answered by this document.

1.5 Symmetries

A **Killing vector** is a vector field along which the metric does not change. Flat spacetime in eight dimensions has 36: eight translations and 28 rotations and boosts, one for each pair of coordinates. The deflating spacetime has, for any rates, 14:

- the seven translations along all coordinates but $x4$;
- the six rotations of $x0, x1, x2, x3$;
- the **scaling**
$$\mathcal{X}_{\mathrm{scale}} = \partial_4 + \mathrm{g55}\,x5\,\partial_5 + \mathrm{g66}\,x6\,\partial_6 + \mathrm{g77}\,x7\,\partial_7, \quad (1.15)$$
which shifts the time and stretches the deflating coordinates so as to undo the shrinking.

Two equal rates add one Killing vector, the rotation of the two coordinates concerned (for $\mathrm{g55} = \mathrm{g66}$ it is $\mathrm{const}_6^2\,x6\,\partial_5 - \mathrm{const}_5^2\,x5\,\partial_6$, which contains the two constants). Three equal rates add the three rotations of $x5, x6, x7$ and three further vectors (Section 1.6), 20 in all (checks `44d-G6`, `71d-G6`). The following table counts, for three sets of rates, how many of the 28 rotations and boosts remain symmetries, and how many of six particular rotations, those of the four axes $x0, x5, x6, x7$, which Part 8 uses.

spacetime	rotations and boosts	isometries, rates (1, 2, 3)	rates (2, 2, 5)	rates (1, 1, 1)	internal generators	of these isometries: (1, 2, 3)	(2, 2, 5)	(1, 1, 1)
7+1	28	6	7	9	6	0	1	3
4+4	28	6	7	9	6	0	1	3

Three consequences. The spacetime is homogeneous: no place and no moment is distinguished. The constants const_t are not observable by themselves, since a rescaling of xt changes them. And there is no conserved energy, because a translation in time alone is not a symmetry. Nothing that involves the time survives among the rotations and boosts, in particular not the boosts of special relativity: the deflation defines a preferred state of rest, as the expansion of the universe does.




Figure 1: Symmetries of the deflating spacetimes. (a) The 28 rotations and boosts of flat spacetime, one square for each pair of coordinates: those that remain symmetries for any rates, those that remain symmetries only if the two coordinates deflate equally fast, and those that are never symmetries. Circles mark the six pairs taken from the axes $x0, x5, x6, x7$. (b) The number of Killing vectors.

1.6 Equal rates: a space of constant curvature

Let $\mathrm{g55} = \mathrm{g66} = \mathrm{g77}$. Then all sectional curvatures (1.9) of the four directions $4, 5, 6, 7$ are equal to $\mathrm{g55}^2$: the four coordinates $x4, \dots, x7$ span a space of constant curvature. With the new coordinates $\bar{x}4 = e^{\mathrm{g55}\,x4}/\mathrm{g55}$ and $\bar{x}t = \mathrm{const}_t\, xt$ the part of the line element that contains them becomes

$$-dx4^2 + \mathrm{gs}_{tt}\, e^{-2\,\mathrm{g55}\,x4} \sum_t d\bar{x}t^2 = \frac{-d\bar{x}4^2 + \mathrm{gs}_{tt} \sum_t d\bar{x}t^2}{(\mathrm{g55}\,\bar{x}4)^2}. \tag{1.16}$$

On the 7+1 spacetime ($\mathrm{gs}_{tt} = +1$) this is de Sitter space of the radius $1/\mathrm{g55}$ in the half of its flat slicing in which space contracts. On the 4+4 spacetime ($\mathrm{gs}_{tt} = -1$) it is minus the line element of four-dimensional hyperbolic space of that radius (checks `44d-G7`, `71d-G7`). The line element (1.16) is a multiple of a flat one, and the special conformal transformations of that flat metric along the three axes $\bar{x}s$ ($s = 5, 6, 7$) are the three further Killing vectors:

$$\mathcal{X}_{(s)} = 2\,\bar{x}s \left[\bar{x}4\,\frac{\partial}{\partial \bar{x}4} + \sum_t \bar{x}t\,\frac{\partial}{\partial \bar{x}t}\right] - \Big(\sum_t \bar{x}t^2 - \mathrm{gs}_{tt}\,\bar{x}4^2\Big)\frac{\partial}{\partial \bar{x}s}, \tag{1.17}$$

with $\bar{x}4\,\partial/\partial\bar{x}4 = \mathrm{g55}^{-1}\,\partial_4$.

1.7 Geodesics: what a classical particle does

The seven translations give seven conserved momenta P_μ ($\mu \neq 4$) of a classical particle of mass m. Let $E_\perp = (m^2 + \sum_j P_j^2)^{1/2}$ be the energy that the particle would have without momentum along the deflating coordinates. The conservation of the norm of the momentum, $g^{\mu\nu}P_\mu P_\nu = -m^2$, gives the velocity along the time, with an affine parameter ς along the path:

$$\Big(\frac{dx4}{d\varsigma}\Big)^2 = E_\perp^2 + \mathrm{gs}_{tt} \sum_t \frac{P_t^2}{b_t(x4)^2}. \tag{1.18}$$

The last term grows as $e^{2\,\mathrm{rate}_t\,x4}$: the momentum P_t belongs to a fixed coordinate wavelength, and the proper wavelength shrinks.

7+1 ($\mathrm{gs}_{tt} = +1$). The energy of a particle that moves along a deflating coordinate grows without bound, the reverse of the redshift in an expanding universe. For momentum along $x5$ alone the proper time that elapses between $x4$ and $x4 = \infty$ is finite, $\dfrac{m}{\mathrm{g55}\,E_\perp}\,\mathrm{arsinh}\Big(\dfrac{E_\perp\,\mathrm{const}_5\,e^{-\mathrm{g55}\,x4}}{|P_5|}\Big)$: the coordinates of Eq. (1.3) do not cover the whole future of such a particle (check `71d-G8`, numerical).

4+4 ($\mathrm{gs}_{tt} = -1$). The right-hand side of Eq. (1.18) vanishes at a **turning value** of the time; for momentum along $x5$ alone,

$$x4_{\mathrm{turn}} = \frac{1}{\mathrm{g55}}\,\ln\frac{E_\perp\,\mathrm{const}_5}{|P_5|}. \tag{1.19}$$

The path reaches this value and returns to smaller $x4$ (check `44d-G9`, numerical). A classical particle with momentum along a deflating time cannot pass the turning value of the ordinary time. Part 7 shows the quantum form of both results.




Figure 2: A classical particle of the mass $m = 1$ with the momenta $P_1 = 0.5$ along $x1$ and $P_5 = 0.3$ along the deflating coordinate $x5$, for $\mathrm{g55} = 1$ and $\mathrm{const}_5 = 0.8$, started at $x4 = -2$. (a) On the 7+1 spacetime the time $x4$ grows without bound within the finite proper time 3.386. (b) On the 4+4 spacetime the path reaches the turning value $x4 = 1.0924$ and returns to smaller $x4$.

Part 2. Dirac matrices and spin connection

2.1 The Dirac matrices of the orthonormal frame

At each point of a curved spacetime the tangent space is flat. The Dirac matrices of the orthonormal frame are therefore eight *constant* matrices γ^A, $A = 0, \ldots, 7$, the same at every point. For each of the two spacetimes they are real 16×16 matrices with entries 0 and ±1; each is a signed permutation matrix. They are written γ^A_{1644} and γ^A_{1671} where the two sets must be distinguished, and γ^A where a statement holds for either set. They satisfy

$$\gamma^A\gamma^B + \gamma^B\gamma^A = 2\,c_\gamma\,\eta^{AB}\,1, \qquad c_\gamma = +1\ (4{+}4), \qquad c_\gamma = -1\ (7{+}1), \tag{2.1}$$

with the flat metric (1.1) of the spacetime concerned. Written out: different matrices anticommute, and the squares are as follows ($j = 0, 1, 2, 3$ and $t = 5, 6, 7$).

set	$(\gamma^j)^2$	$(\gamma^4)^2$	$(\gamma^t)^2$	symmetric	antisymmetric
γ^A_{1644}	$+1$	-1	-1	$\gamma^0, \dots, \gamma^3$	$\gamma^4, \dots, \gamma^7$
γ^A_{1671}	-1	$+1$	-1	γ^4	the seven others

The matrices of the 4+4 spacetime square to the sign of the metric. Those of the 7+1 spacetime square to *minus* the sign of the metric: the matrix of the time squares to $+1$, as in the familiar Dirac theory with the metric $(+,-,-,-)$, while this document writes the metric with space-like components positive. The constant $c_\gamma = -1$ records this, and it accompanies the 7+1 matrices through all formulas (checks `44-R1`, `44-R2`, `44-C1`, `44-C2` and `71-R1`, `71-R2`, `71-C1`, `71-C2` of the report on the matrices of the frame, Appendix A.2, and `44d-C1`, `71d-C1`). In the reports of the programs the diagonal matrices of these squares are called `eta1644` and `eta1671`, in formulas η_{1644} and η_{1671}; they are related to the flat metrics (1.1) by $\eta_{1644} = \eta^{(4+4)}$ and $\eta_{1671} = -\eta^{(7+1)}$.

The eight matrices of the 4+4 spacetime

In the following displays $+$ stands for $+1$, $-$ for -1 and a dot for 0.

$$\gamma^0_{1644} = \begin{pmatrix}
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot\\
+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+\\
\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot
\end{pmatrix}
\qquad
\gamma^1_{1644} = \begin{pmatrix}
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&-&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot
\end{pmatrix}$$

$$\gamma^2_{1644} = \begin{pmatrix}
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&-&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&+&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
+&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot
\end{pmatrix}
\qquad
\gamma^3_{1644} = \begin{pmatrix}
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&-\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&+&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&-&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot
\end{pmatrix}$$

$$\gamma^4_{1644} = \begin{pmatrix}
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&+\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot
\end{pmatrix}
\qquad
\gamma^5_{1644} = \begin{pmatrix}
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&-&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&-&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&+&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&-&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&+&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&+&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&-&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
-&\cdot&-&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot
\end{pmatrix}$$

$$\gamma^6_{1644} = \begin{pmatrix}
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&+&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&+&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&+&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&+&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&+&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&-&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&-&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&-&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&-&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&-&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&-&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
-&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot
\end{pmatrix}
\qquad
\gamma^7_{1644} = \begin{pmatrix}
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
\cdot&+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot\\
+&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot&\cdot
\end{pmatrix}$$

$$\omega_{1644} = \begin{pmatrix} - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + \end{pmatrix} \qquad C^{-}_{1644} = \begin{pmatrix} \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot \end{pmatrix}$$

$$C^{+}_{1644} = \begin{pmatrix} \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot \end{pmatrix}$$

Matrices built from them (all verified in exact integer arithmetic):

matrix	definition	properties
ω_{1644}	$\gamma^0\gamma^1 \dots \gamma^7$	real, diagonal, diag$(-1_8, +1_8)$; anticommutes with every γ^A; it is the chirality matrix χ of this spacetime
P^L, P^R	$\frac{1}{2}(1-\chi)$, $\frac{1}{2}(1+\chi)$	chiral projectors; they select the upper and the lower eight components
C^{-}_{1644}	$\gamma^0\gamma^1\gamma^2\gamma^3$	real, symmetric, square 1; $C^{-}\gamma^A(C^{-})^{-1} = -(\gamma^A)^T$; defines the covariant adjoint $\tilde{\psi} = \psi^\dagger C^{-}$
C^{+}_{1644}	$C^{-}\omega_{1644}$	$C^{+}\gamma^A(C^{+})^{-1} = +(\gamma^A)^T$
β	$-i\gamma^4$	Hermitian, square 1
α^j	$-\gamma^4\gamma^j$ $(j = 0, 1, 2, 3)$	Hermitian, square 1; anticommute with one another and with β
W_{1644}	$i\,C^{-}\gamma^4$	Hermitian, square 1, trace 0; commutes with β and with every α^j; $C^{-} = -W_{1644}\,\beta$
τ^A, $\bar{\tau}^A$	the lower left and the upper right 8×8 block of γ^A	real; $\gamma^A = \begin{pmatrix} 0 & \bar{\tau}^A \\ \tau^A & 0 \end{pmatrix}$: every γ^A exchanges the upper and the lower eight components

The eight matrices of the 7+1 spacetime

$$\gamma^0_{1671} = \begin{pmatrix} \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - \end{pmatrix} \qquad \gamma^1_{1671} = \begin{pmatrix} \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}$$

$$\gamma^2_{1671} = \begin{pmatrix}
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & + & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
- & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & - & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot
\end{pmatrix}
\qquad
\gamma^3_{1671} = \begin{pmatrix}
\cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
+ & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & + & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & + \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot
\end{pmatrix}$$

$$\gamma^4_{1671} = \begin{pmatrix}
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot \\
+ & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot \\
\cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot
\end{pmatrix}
\qquad
\gamma^5_{1671} = \begin{pmatrix}
\cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
- & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & + \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot
\end{pmatrix}$$

$$\gamma^6_{1671} = \begin{pmatrix}
\cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
- & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & + \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot
\end{pmatrix}
\qquad
\gamma^7_{1671} = \begin{pmatrix}
\cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
- & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & - & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & - & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot
\end{pmatrix}$$

$$\omega_{1671} = \begin{pmatrix}
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot \\
+ & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot \\
\cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot
\end{pmatrix}
\qquad
C^-_{1671} = \begin{pmatrix}
+ & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & - & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & -
\end{pmatrix}$$

$$C^+_{1671} = \begin{pmatrix}
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot \\
+ & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot \\
\cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\
\cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & + & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot
\end{pmatrix}$$

matrix	definition	properties
ω_{1671}	$\gamma^0\gamma^1\cdots\gamma^7$	real, square -1; anticommutes with every γ^A

matrix	definition	properties
χ	$i\,\omega_{1671}$	the chirality matrix: Hermitian, square 1, anticommutes with every γ^A
P^L, P^R	$\frac{1}{2}(1-\chi)$, $\frac{1}{2}(1+\chi)$	chiral projectors; complex
β	γ^4	real, symmetric, square 1; defines the adjoint $\bar\psi = \psi^\dagger\beta$
α^A	$\gamma^4\gamma^A$ $(A\neq 4)$	real, symmetric, square 1; anticommute with one another and with β
C^-_{1671}	$\gamma^4\omega_{1671}$	$C^-\gamma^A(C^-)^{-1} = -(\gamma^A)^T$; enters the pair operators $\psi^T C^-\gamma^A\psi$
C^+_{1671}	γ^4	$C^+\gamma^A(C^+)^{-1} = +(\gamma^A)^T$
E^L, E^R	$\frac{1}{\sqrt{2}}\begin{pmatrix}1_8\\ -i\,1_8\end{pmatrix}$, $\frac{1}{\sqrt{2}}\begin{pmatrix}-i\,1_8\\ 1_8\end{pmatrix}$	complex 16×8 matrices whose columns span the ranges of P^L and P^R; $E^L E^{L\dagger} + E^R E^{R\dagger} = 1$
τ^A_{1671} $(A\neq 4)$	$i\,E^{L\dagger}\alpha^A E^L$	real 8×8 matrices; $E^{R\dagger}\alpha^A E^R = +i\,\tau^A_{1671}$, $E^{L\dagger}\alpha^A E^R = 0$, $E^{L\dagger}\beta E^R = 1_8$

2.2 The matrices in coordinate components

The Dirac matrices with a coordinate index are $\gamma^\mu = e_A{}^\mu\gamma^A$: five of them are the constant matrices, and the three along the deflating coordinates are the constant matrices divided by the scale factors,

$$\gamma^\mu = \gamma^{A=\mu} \quad (\mu = 0,\dots,4), \qquad \gamma^{\mu=t} = \frac{\gamma^{A=t}}{b_t(x4)} \quad (t = 5,6,7). \tag{2.2}$$

They satisfy $\gamma^\mu\gamma^\nu + \gamma^\nu\gamma^\mu = 2\,c_\gamma\,g^{\mu\nu}\,1$ with the inverse metric of Eq. (1.3) (checks `44d-C1`, `71d-C1`), and they depend on the time: $\partial_4\gamma^{\mu=t} = +\text{rate}_t\,\gamma^{\mu=t}$. From here on a matrix γ with a Greek index is a coordinate matrix (2.2) and a matrix with a capital index or with a digit is a constant matrix of the frame.

2.3 The spin connection

A spinor is defined relative to a frame. When the frame turns from point to point, the derivative of a spinor must account for the turning; the covariant derivative is

$$\nabla_\mu\psi = (\partial_\mu + \Omega_\mu)\,\psi, \qquad \Omega_\mu = \frac{c_\gamma}{4}\,\omega_{\mu AB}\,\gamma^A\gamma^B, \tag{2.3}$$

where the **spin connection** Ω_μ is built from the coefficients

$$\omega_\mu{}^A{}_B = e^A{}_\nu(\partial_\mu e_B{}^\nu + \Gamma^\nu{}_{\mu\lambda}\,e_B{}^\lambda), \qquad \omega_{\mu AB} = \eta_{AC}\,\omega_\mu{}^C{}_B = -\omega_{\mu BA}, \tag{2.4}$$

which express the covariant derivative of the frame vector e_B in the frame. The factor c_γ in Eq. (2.3) is that of Eq. (2.1): the matrices $\frac{c_\gamma}{4}[\gamma^A,\gamma^B]$ generate the Lorentz transformations of the spinors on either spacetime.

Evaluation. The frame (1.4) depends on $x4$ only, through $e_t{}^t = 1/b_t$, and the Christoffel symbols are those of Eq. (1.7). For $\mu = t$, $A = t$, $B = 4$ the first term of Eq. (2.4) vanishes ($e_4{}^\nu$ is constant) and the second gives

$$\omega_t{}^t{}_4 = e^t{}_t\,\Gamma^t{}_{t4}\,e_4{}^4 = -\,\mathrm{rate}_t\,b_t, \qquad \text{hence} \qquad \omega_{t\,t4} = -\,\mathrm{gs}_{tt}\,\mathrm{rate}_t\,b_t = -\,\omega_{t\,4t}. \tag{2.5}$$

In the same way $\omega_t{}^4{}_t = e^4{}_4\,\Gamma^4{}_{tt}\,e_t{}^t = -\mathrm{gs}_{tt}\,\mathrm{rate}_t\,b_t$, which confirms the antisymmetry after the first index is lowered with $\mathrm{gs}_{44} = -1$. For $\mu = 4$ the two terms of Eq. (2.4) cancel ($\partial_4 e_t{}^t = +\mathrm{rate}_t/b_t$ and $\Gamma^t{}_{4t}\,e_t{}^t = -\mathrm{rate}_t/b_t$), and for $\mu = 0,1,2,3$ both vanish. The coefficients (2.5) are thus the only ones that differ from zero: $+\mathrm{rate}_t\,b_t$ on the 4+4 spacetime and $-\mathrm{rate}_t\,b_t$ on the 7+1 spacetime (checks `44d-C2`, `71d-C2`). Inserted into Eq. (2.3), with $c_\gamma\,\mathrm{gs}_{tt} = -1$ and the anticommutation of γ^t and γ^4:

$$\Omega_t = \tfrac{1}{2}\,\mathrm{rate}_t\,b_t\,\gamma^t\gamma^4 \quad (t = 5,6,7), \qquad \Omega_0 = \Omega_1 = \Omega_2 = \Omega_3 = \Omega_4 = 0, \tag{2.6}$$

with the constant matrices of the frame on the right, *the same formula on both spacetimes* (checks `44d-C3`, `71d-C3`). Each Ω_t generates the Lorentz transformation in the plane of xt and $x4$.

Test. The spin connection is correct if and only if the coordinate matrices are covariantly constant,

$$\partial_\mu\gamma^\nu + \Gamma^\nu{}_{\mu\lambda}\,\gamma^\lambda + [\Omega_\mu, \gamma^\nu] = 0, \tag{2.7}$$

which holds for all 64 pairs of indices (checks `44d-C4`, `71d-C4`).

The combination that enters the Dirac equation. With Eqs. (2.2) and (2.6) and $(\gamma^t)^2 = -1$ on both spacetimes,

$$\gamma^\mu\Omega_\mu = \sum_t \frac{\gamma^t}{b_t}\cdot\tfrac{1}{2}\,\mathrm{rate}_t\,b_t\,\gamma^t\gamma^4 = -\tfrac{1}{2}\,\mathrm{gsum}\,\gamma^4. \tag{2.8}$$

The scale factors cancel, and only the total rate remains (checks `44d-C5`, `71d-C5`).

2.4 The curvature of the spinor derivative

The commutator of two covariant derivatives of a spinor is the Riemann tensor,

$$[\nabla_\mu, \nabla_\nu]\,\psi = (\partial_\mu\Omega_\nu - \partial_\nu\Omega_\mu + [\Omega_\mu, \Omega_\nu])\psi = \frac{c_\gamma}{4}\,R_{AB\mu\nu}\,\gamma^A\gamma^B\,\psi, \tag{2.9}$$

for all 64 pairs of indices (checks `44d-C6`, `71d-C6`); this is a third route to the curvature (1.8). Contracted with two Dirac matrices,

$$\tfrac{1}{2}\,\gamma^\mu\gamma^\nu\,[\nabla_\mu, \nabla_\nu] = -\frac{c_\gamma\,R}{4}\,1, \qquad R = \mathrm{gsq} + \mathrm{gsum}^2 \tag{2.10}$$

(checks `44d-C7`, `71d-C7`): the square of the Dirac operator $\gamma^\mu\nabla_\mu$ differs from c_γ times the wave operator of the spinor derivative by this constant.

Part 3. The Lagrangians

3.1 The three parts

The Lagrangian of the field is

$$\mathcal{L} = V_d \left[\mathcal{L}_{\text{kin}} + \mathcal{L}_{\text{mass}} + \mathcal{L}_{\text{self}}\right], \tag{3.1}$$

where the factor $V_d = \sqrt{|\det g|}$ of Eq. (1.5) makes $\mathcal{L}\, d^8x$ invariant. $\mathcal{L}_{\text{kin}}$ contains the derivative of the field and is bilinear in it; $\mathcal{L}_{\text{mass}}$ is bilinear without derivative; $\mathcal{L}_{\text{self}}$ is quartic without derivative. The components of the field anticommute with one another (they are Grassmann-valued, as the quantization of a fermion field requires). A product of two components is therefore an even element of the Grassmann algebra and commutes with everything; this is used repeatedly below.

3.2 The 7+1 spacetime

Adjoint. $\bar\psi = \psi^\dagger\gamma^4 = \psi^\dagger\beta$.

Kinetic term and mass term.

$$\mathcal{L}_{\text{kin}} + \mathcal{L}_{\text{mass}} = \bar\psi\,(i\,\gamma^\mu\nabla_\mu - m)\,\psi, \tag{3.2}$$

with the coordinate matrices (2.2), the covariant derivative (2.3) and a real constant m, the mass. The mass term $-m\,\bar\psi\psi$ is real because γ^4 is symmetric, and it is a scalar under the Lorentz transformations of the frame.

Self-interaction. The most general interaction without derivatives that is quartic in the field and invariant under the Lorentz transformations of the frame is built from the products of the Dirac matrices. Let

$$K = (A_1 < A_2 < \cdots < A_r) \tag{3.3}$$

be a strictly increasing list of r indices out of $0, \dots, 7$. The number $r = |K|$ is the **rank** of the list. There are $2^8 = 256$ lists: $1, 8, 28, 56, 70, 56, 28, 8, 1$ of the ranks $0, \dots, 8$. Define, for each list,

$$\Theta^K = \zeta_r\,\gamma^{A_1}\gamma^{A_2}\dots\gamma^{A_r}, \qquad \zeta_r = \begin{cases} i & \text{for } r = 2,3,6,7, \\ 1 & \text{for } r = 0,1,4,5,8, \end{cases} \qquad \Theta_K = (\Theta^K)^{-1} = \text{sgn}_K\,\Theta^K, \tag{3.4}$$

with $\Theta^{()} = 1$ for the empty list. The sign $\text{sgn}_K = \pm 1$ is the product of the squares of the matrices in K, that is, (-1) to the number of indices in K other than 4. The phase ζ_r is chosen so that $\gamma^4\Theta^K$ is Hermitian; every bilinear $\bar\psi\,\Theta^K\psi$ is then real (check `71s-S2`). The **invariant of rank** r and the self-interaction are

$$I_r = \sum_{|K|=r} (\bar\psi\,\Theta_K\,\psi)(\bar\psi\,\Theta^K\,\psi), \qquad \mathcal{L}_{\text{self}} = -\frac{1}{2}\sum_{r=0}^{8} g_r\, I_r. \tag{3.5}$$

The nine real constants $g_0, \dots, g_8$ are the **couplings** of the self-interaction. A coupling carries one digit as its subscript; it must not be confused with a rate (g55) or with a component of the metric ($g^{(7+1)}_{55}$). In a gravitational field each I_r is written with the constant matrices of the frame exactly as in flat spacetime, because it is invariant under the Lorentz transformations of the frame (check `71s-S4`); the deflation enters $\mathcal{L}_{\text{self}}$ only through the factor V_d of Eq. (3.1).

The four invariants of lowest and highest rank have familiar names. With $\gamma_A = (\gamma^A)^{-1}$ and the chirality matrix $\chi = i\,\omega_{1671}$:

invariant	written out	name	traditional coupling
I_0	$(\bar\psi\psi)^2$	square of the scalar density	$g_S = g_0$
I_1	$\sum_A(\bar\psi\gamma_A\psi)(\bar\psi\gamma^A\psi)$	square of the current	$g_V = g_1$
I_7	$-\sum_A(\bar\psi\gamma_A\chi\psi)(\bar\psi\gamma^A\chi\psi)$	minus the square of the axial current	$g_A = -g_7$
I_8	$-(\bar\psi\, i\chi\,\psi)^2$	minus the square of the pseudoscalar density	$g_P = -g_8$

The invariants of the ranks 2 to 6 are the squares of the antisymmetric tensors of those ranks. Part 6 shows that for anticommuting fields they are not independent of the four in the table.

3.3 The 4+4 spacetime: the covariant Lagrangian

Adjoint. $\tilde\psi = \psi^\dagger C^-$ with $C^- = C^-_{1644} = \gamma^0\gamma^1\gamma^2\gamma^3$. (The matrix γ^4 cannot serve: on this spacetime four matrices are time-like, and the adjoint that makes $\tilde\psi\psi$ invariant under all transformations of the frame is built from the product of the four space-like matrices.)

Kinetic term and mass term.

$$\mathcal{L}_{\text{kin}} + \mathcal{L}_{\text{mass}} = \tilde\psi\,(\gamma^\mu\nabla_\mu - m)\,\psi. \tag{3.6}$$

It contains no factor i: with the real matrices of 4+4, whose squares are the signs of the metric, the operator $\gamma^\mu\nabla_\mu - m$ already gives the mass shell of a particle of mass m (Section 4.4). The mass term $-m\,\tilde\psi\psi$ is real because C^- is symmetric.

Self-interaction. With the lists K of Eq. (3.3),

$$\Theta^K = \zeta_r\,\gamma^{A_1}\gamma^{A_2}\ldots\gamma^{A_r}, \qquad \zeta_r = \begin{cases} i & \text{for } r = 1,2,5,6,\\ 1 & \text{for } r = 0,3,4,7,8,\end{cases} \qquad \Theta_K = (\Theta^K)^{-1} = \text{sgn}_K\,\Theta^K, \tag{3.7}$$

where now sgn_K is $(-1)^r$ times the product of the squares of the matrices in K. The phases differ from those of Eq. (3.4) because the adjoint is built with C^- and not with a single Dirac matrix: they are chosen so that $C^-\Theta^K$ is Hermitian and every bilinear $\tilde\psi\,\Theta^K\psi$ is real (check `44c-S2`). The invariants and the self-interaction are

$$I_r = \sum_{|K|=r}(\tilde\psi\,\Theta_K\,\psi)(\tilde\psi\,\Theta^K\,\psi), \qquad \mathcal{L}_{\text{self}} = -\frac{1}{2}\sum_{r=0}^{8} g_r\,I_r. \tag{3.8}$$

Each I_r is invariant under the 28 transformations of $SO(4,4)$ of the frame (check `44c-S4`). Equations (3.6) and (3.8) are the **covariant Lagrangian** of the 4+4 model. Part 5 shows that its kinetic term is indefinite, describes what that means, and defines the model that is quantized: the invariants (3.8), which are called **form C** of the self-interaction and are the definition of the 4+4 model, together with a positive kinetic term. Part 5 also defines the alternative, **form Q**.

3.4 Why the bilinears are real, and why the invariants are invariant

Reality. For any matrix M, $(\psi^\dagger M\psi)^\dagger = \psi^\dagger M^\dagger\psi$. A bilinear $\psi^\dagger\,\mathrm{Adj}\,\Theta^K\psi$, with Adj $=\gamma^4$ on 7+1 and Adj $= C^-$ on 4+4, is therefore real if and only if Adj Θ^K is Hermitian. Reversing the order of the r matrices of a product costs the sign $(-1)^{r(r-1)/2}$, and moving the adjoint matrix through them costs a further sign that depends on r; the phase ζ_r is i exactly for the ranks at which the total sign is -1. The program verifies the Hermiticity for all 256 lists on each spacetime.

Invariance. An infinitesimal Lorentz transformation of the frame acts on the field as $\delta\psi = \frac{1}{2}\epsilon_{AB}\,\Sigma^{AB}\psi$ with $\Sigma^{AB} = \frac{1}{2}\gamma^A\gamma^B$ $(A < B)$ and 28 real parameters ϵ_{AB}. The scalar $\psi^\dagger\,\mathrm{Adj}\,\psi$ is invariant because $\Sigma^{AB\dagger}\,\mathrm{Adj} + \mathrm{Adj}\,\Sigma^{AB} = 0$. The commutator of Θ^K with $\gamma^A\gamma^B$ is either zero or a multiple of the single matrix $\Theta^{K'}$ of the same rank whose list K' is K with one of the two indices A, B exchanged for the other; the contributions of K and of K' to the variation of I_r cancel in pairs. The program verifies both statements for all 28 generators and all 256 lists.

3.5 Dictionary to the notation of the author's document on the 7+1 problem

The author's document that defines and solves the 7+1 problem with self-interaction uses the conventions of the first column. The definitions of Section 3.2 are his, in the notation of this project.

in the author's document	in this document	remark
time x_0; ordinary space x_1, x_2, x_3; auxiliary space $x_4, \dots, x_7$	time $x4$; ordinary space $x1, x2, x3$; the other space-like coordinates $x0, x5, x6, x7$	the index values 0 and 4 are exchanged
$\eta = \mathrm{diag}(+1, -1, \dots, -1)$ for the squares of the Dirac matrices and for the metric	squares of γ^A_{1671}; the metric is $\eta^{(7+1)}$ with space-like components positive	hence $c_\gamma = -1$ in Eq. (2.1)
Dirac matrices Γ_A	γ^A_{1671}	the same real matrices
Θ^K, Θ_K, I_r, g_r, $\mathcal{L}_{\mathrm{self}} = -\frac{1}{2}\sum_r g_r I_r$	the same symbols and the same definitions	
sign η_K; phase i^{σ_r}	sgn_K; ζ_r	renamed because η and σ have other meanings in the project
Fierz matrix F_{rs}	$\mathsf{F}_{rr'}$	sans-serif type, to distinguish it from amplitudes F

The example of that document, $g_0 = -10^5$ and $g_1 = +10^4$ with all other couplings zero, is evaluated in Section 6.3.

Part 4. The field equations

4.1 Variation of the action

The action is $\int \mathcal{L}\, d^8x$ with the Lagrangian (3.1). The field equation follows by varying the adjoint field, $\bar\psi$ on 7+1 and $\tilde\psi$ on 4+4, with ψ held fixed. The kinetic and the mass term are linear in the adjoint field and contribute the operator in parentheses of Eqs. (3.2) and (3.6), applied to ψ. In the self-interaction, a bilinear is an even element of the Grassmann algebra and commutes with everything, so that a product of two bilinears is varied as a product of ordinary numbers:

$$\delta\Big[(\bar\psi\Theta_K\psi)(\bar\psi\Theta^K\psi)\Big] = (\delta\bar\psi\,\Theta_K\psi)(\bar\psi\Theta^K\psi) + (\bar\psi\Theta_K\psi)(\delta\bar\psi\,\Theta^K\psi) = 2\,(\bar\psi\Theta_K\psi)(\delta\bar\psi\,\Theta^K\psi),$$

where the last step uses $\Theta_K = \mathrm{sgn}_K\,\Theta^K$. With the factor $-\frac{1}{2}g_r$ of Eqs. (3.5) and (3.8) the result is

$$\text{7+1:} \qquad (i\,\gamma^{\mu}\nabla_{\mu} - m)\,\psi_{1671d} = \sum_{r=0}^{8} g_r \sum_{|K|=r} (\bar{\psi}\,\Theta_K\,\psi)\,\Theta^K\psi, \tag{4.1}$$

$$\text{4+4 (covariant Lagrangian):} \qquad (\gamma^{\mu}\nabla_{\mu} - m)\,\psi_{1644d} = \sum_{r=0}^{8} g_r \sum_{|K|=r} (\tilde{\psi}\,\Theta_K\,\psi)\,\Theta^K\psi. \tag{4.2}$$

The left-hand side is the Dirac operator with the mass term. The right-hand side is the self-interaction, cubic in the field; without it the equation is linear. That the cubic term is the derivative of the quartic one is verified exactly, for each of the nine ranks, in the form in which it is used later: the mean-field potential is the derivative of the mean-field energy (checks `71s-S8`, `44q-S8`, `44c-S8`). The variation with respect to ψ gives the Hermitian conjugate of the same equation; this requires that the spin connection satisfy $\Omega_\mu^T\gamma^4 + \gamma^4\Omega_\mu = 0$ on 7+1 and $\Omega_\mu^T C^- + C^-\Omega_\mu = 0$ on 4+4, which holds (checks `71d-C8`, `44d-C8`).

4.2 The equations written out

With Eq. (2.8) for the spin connection and Eq. (2.2) for the coordinate matrices, Eq. (4.1) becomes

$$i\,\gamma^4\Big(\partial_4 - \frac{\mathrm{gsum}}{2}\Big)\psi + i\sum_{j=0}^{3}\gamma^j\,\partial_j\psi + i\sum_{t=5}^{7}\frac{\gamma^t}{b_t}\,\partial_t\psi - m\,\psi = \sum_{r=0}^{8} g_r \sum_{|K|=r} (\bar{\psi}\,\Theta_K\,\psi)\,\Theta^K\psi \qquad \text{(7+1)}, \tag{4.3}$$

and Eq. (4.2) becomes

$$\gamma^4\Big(\partial_4 - \frac{\mathrm{gsum}}{2}\Big)\psi + \sum_{j=0}^{3}\gamma^j\,\partial_j\psi + \sum_{t=5}^{7}\frac{\gamma^t}{b_t}\,\partial_t\psi - m\,\psi = \sum_{r=0}^{8} g_r \sum_{|K|=r} (\tilde{\psi}\,\Theta_K\,\psi)\,\Theta^K\psi \qquad \text{(4+4)}, \tag{4.4}$$

with the constant matrices of the frame throughout. Compared with flat spacetime, the derivative along the time is shifted by $\mathrm{gsum}/2$, and each derivative along a deflating coordinate is divided by its scale factor.

4.3 The weighted field: the deflation strengthens the self-interaction

The shift is removed by a change of normalization. Define the **weighted field**

$$\hat{\psi} = \sqrt{V_d}\,\psi. \tag{4.5}$$

Because $\partial_4\sqrt{V_d} = -\frac{1}{2}\,\mathrm{gsum}\,\sqrt{V_d}$, one has $(\partial_4 - \frac{1}{2}\mathrm{gsum})\,\psi = V_d^{-1/2}\,\partial_4\hat{\psi}$: the left-hand side of Eq. (4.3) or (4.4) applied to ψ equals $V_d^{-1/2}$ times the same expression *without* the term $\mathrm{gsum}/2$ applied to $\hat{\psi}$. The right-hand side is cubic in the field and equals $V_d^{-3/2}$ times the same expression in $\hat{\psi}$. Multiplying by $V_d^{1/2}$:

> **The weighted field obeys Eq. (4.3) or (4.4) without the term** $\mathrm{gsum}/2$ **and with every coupling** g_r **replaced by**
>
> $$\frac{g_r}{V_d(x4)} \;\propto\; g_r\,e^{\mathrm{gsum}\,x4}. \tag{4.6}$$
>
> **The mass** m **is not changed.**

The same statement in terms of the Lagrangian: V_d times the mass term and the self-interaction, written with the weighted field, is $-m\,\bar{\hat{\psi}}\hat{\psi} - \frac{1}{2}\sum_r (g_r/V_d)\,\hat{I}_r$, where $\hat{I}_r$ is the invariant (3.5) of the weighted field (check `def-S5`, exact). The quantity $\psi^\dagger\psi$ is a density per unit of proper volume and grows as the deflating coordinates shrink; $\hat{\psi}^\dagger\hat{\psi}$ is the density per unit of coordinate volume, from which this trivial growth is removed.

Consequence. *For a field that does not depend on $x5, x6, x7$ the equation of the weighted field is exactly the equation of flat spacetime, with a self-interaction whose couplings all grow as $e^{\mathrm{gsum}\,x4}$.* A bilinear term, such as the mass term, is unaffected; a quartic term is strengthened by the shrinking of the volume, because the same number of particles is confined to less and less proper volume. Every result that holds on flat spacetime for such fields holds for the weighted field at each moment, with the couplings of that moment.

4.4 The equations as evolution in the ordinary time

7+1. Multiply the equation of the weighted field by $\gamma^4 = \beta$ (its square is 1) and use $\alpha^A = \gamma^4\gamma^A$:

$$i\,\partial_4\hat{\psi} = \mathsf{h}\,\hat{\psi} + \frac{1}{V_d}\sum_{r=0}^{8} g_r \sum_{|K|=r} (\bar{\hat{\psi}}\,\Theta_K\,\hat{\psi})\,\beta\,\Theta^K\hat{\psi}, \qquad \mathsf{h} = \beta\,m + \sum_{j=0}^{3}\alpha^j\,(-i\partial_j) + \sum_{t=5}^{7}\frac{\alpha^t}{b_t}\,(-i\partial_t), \tag{4.7}$$

with $\bar{\hat{\psi}} = \hat{\psi}^\dagger\beta$. The matrix-valued operator h (sans-serif) is the Hamiltonian of a single particle.

4+4. Multiply the equation of the weighted field by $(\gamma^4)^{-1} = -\gamma^4$:

$$\partial_4\hat{\psi} = \Big[-m\,\gamma^4 + \sum_{j=0}^{3}\gamma^4\gamma^j\,\partial_j + \sum_{t=5}^{7}\frac{\gamma^4\gamma^t}{b_t}\,\partial_t\Big]\hat{\psi} - \frac{1}{V_d}\sum_{r=0}^{8} g_r \sum_{|K|=r} (\tilde{\bar{\hat{\psi}}}\,\Theta_K\,\hat{\psi})\,\gamma^4\,\Theta^K\hat{\psi}, \tag{4.8}$$

with $\tilde{\bar{\hat{\psi}}} = \hat{\psi}^\dagger C^-$.

Modes of the linear equation. Without the self-interaction the coefficients depend on $x4$ only, and a solution can be taken proportional to $\exp\left(i\sum_j p_j\,xj + i\sum_t k_t\,xt\right)$ with seven constant wavenumbers: four along the flat coordinates, written p_j on 7+1 and k_j on 4+4, and three along the deflating ones, k_t. Such a solution is a **mode**. Two quantities characterize it:

$$k_d(x4) = \Big(\sum_t \frac{k_t^2}{b_t(x4)^2}\Big)^{1/2}, \qquad E_\perp = \Big(m^2 + \sum_j p_j^2\Big)^{1/2}\ \text{(7+1)}, \quad E_\perp = \Big(m^2 + \sum_j k_j^2\Big)^{1/2}\ \text{(4+4)}. \tag{4.9}$$

k_d is the **physical wavenumber** along the deflating coordinates, which grows as $e^{\mathrm{rate}_t\,x4}$, and $E_\perp$ is the energy that the mode would have with $k_5 = k_6 = k_7 = 0$. For a mode, Eqs. (4.7) and (4.8) are ordinary differential equations:

$$\begin{aligned} \text{7+1:}\quad & i\,\partial_4\hat{\psi} = \mathsf{h}(x4)\,\hat{\psi}, \qquad \mathsf{h} = \mathsf{h}_\perp + k_d\,\alpha^d, \\ & \mathsf{h}_\perp = m\,\beta + \sum_j p_j\,\alpha^j, \qquad \alpha^d = \frac{1}{k_d}\sum_t \frac{k_t}{b_t}\,\alpha^t; \end{aligned} \tag{4.10}$$

$$\text{4+4:}\quad \partial_4\hat{\psi} = K(x4)\,\hat{\psi}, \qquad K = K_\perp + i\,k_d\,\gamma^4\gamma^d,$$
$$K_\perp = -m\,\gamma^4 + i\sum_j k_j\,\gamma^4\gamma^j, \qquad \gamma^d = \frac{1}{k_d}\sum_t \frac{k_t}{b_t}\,\gamma^t. \tag{4.11}$$

(The letter K without index, or with the index $\perp$, is this matrix; an index list K occurs only as a sub- or superscript of Θ and under a sum; a sectional curvature K_{AB} carries two indices.) Their properties decide everything that follows:

	7+1: h	4+4: K
square	$\mathsf{h}^2 = (E_\perp^2 + k_d^2)\,1$	$K^2 = (k_d^2 - E_\perp^2)\,1$
character	real and symmetric for all wavenumbers and all times	$K_\perp$ is anti-Hermitian; $i\gamma^4\gamma^d$ is Hermitian
evolution	unitary; $\hat{\psi}^\dagger\hat{\psi}$ is conserved	unitary, with $\hat{\psi}^\dagger\hat{\psi}$ conserved, if and only if $k_5 = k_6 = k_7 = 0$
what the deflation does	$\partial_4\mathsf{h} = \sum_t \text{rate}_t\,(k_t/b_t)\,\alpha^t$ anticommutes with β: the energy grows, and pairs are created	K^2 changes sign when k_d reaches $E_\perp$: the mode oscillates before, and grows or decays exponentially after
checks	`71d-D1`, `71d-D2`, `71d-D3`, `71d-D7`	`44d-D1`, `44d-D2`, `44d-D3`, `44d-D5`, `44d-D7`

The instantaneous mass shell is the same statement in other words: on 7+1 the frequency k_4 of a mode satisfies $k_4^2 = E_\perp^2 + k_d^2$; on 4+4, $k_4^2 = E_\perp^2 - k_d^2$, which is negative once the physical wavenumber along the additional times exceeds $E_\perp$. Part 7 solves both equations.

4.5 The conserved current, and charge conjugation

All three parts of the Lagrangian are invariant under a change of the phase of the field, $\psi \to e^{i\theta}\psi$. The corresponding conserved current is

$$j^\mu = \bar{\psi}\,\gamma^\mu\,\psi \;\text{ (7+1)}, \qquad j^\mu = i\,\tilde{\psi}\,\gamma^\mu\,\psi \;\text{ (4+4)}, \qquad \nabla_\mu j^\mu = 0 \iff \partial_\mu(V_d\,j^\mu) = 0, \tag{4.12}$$

with or without the self-interaction, which contains no derivative and does not change the current. (The factor i on 4+4 makes the current real: $C^-\gamma^\mu$ is antisymmetric.) The conserved charge is the integral of $V_d\,j^4$ over the seven coordinates other than $x4$, and

$$V_d\,j^4 = \hat{\psi}^\dagger\hat{\psi} \;\text{ (7+1)}, \qquad V_d\,j^4 = \hat{\psi}^\dagger\,W_{1644}\,\hat{\psi} \;\text{ (4+4)}, \qquad W_{1644} = i\,C^-\gamma^4. \tag{4.13}$$

On 7+1 the charge density is positive; after quantization the charge is the number of particles minus the number of antiparticles. On 4+4 the matrix W_{1644} has the eigenvalues $+1$ and -1, eight times each: *the conserved charge of the covariant 4+4 theory is not positive.* For a mode, $W_{1644}K + K^\dagger W_{1644} = 0$ for all wavenumbers (check `44d-D4`). Part 5 takes up the consequences.

Charge conjugation on 7+1. The matrix ω_{1671} commutes with every α^A and anticommutes with β; since the scale factors are real, the field $\omega_{1671}\,\psi^*$ solves the linear equation whenever ψ does (check `71d-D5`).

4.6 The two coupled 8-component equations

Each set of Dirac matrices exchanges the two chiralities, and the linear equation of the weighted field is a pair of equations for two 8-component fields.

4+4. With the blocks τ^A, $\bar\tau^A$ of Section 2.1 and $\hat\psi = (\hat\psi^L, \hat\psi^R)$, the upper and the lower eight components,

$$\bar\tau^A\, e_A{}^\mu \partial_\mu\, \hat\psi^R = m\, \hat\psi^L, \qquad \tau^A\, e_A{}^\mu \partial_\mu\, \hat\psi^L = m\, \hat\psi^R, \qquad e_A{}^\mu \partial_\mu = \Big(\partial_0, \dots, \partial_4, \frac{\partial_5}{b_5}, \frac{\partial_6}{b_6}, \frac{\partial_7}{b_7}\Big), \tag{4.14}$$

two equations for real 8-component fields if the initial data are real (check `44d-D6`).

7+1. With the matrices E^L, E^R and τ^A_{1671} of Section 2.1 and $\hat\varphi^L = E^{L\dagger}\hat\psi$, $\hat\varphi^R = E^{R\dagger}\hat\psi$,

$$\Big(i\,\partial_4 + \sum_{A\neq 4} \tau^A_{1671}\, e_A{}^\mu \partial_\mu\Big)\hat\varphi^L = m\,\hat\varphi^R, \qquad \Big(i\,\partial_4 - \sum_{A\neq 4} \tau^A_{1671}\, e_A{}^\mu \partial_\mu\Big)\hat\varphi^R = m\,\hat\varphi^L \tag{4.15}$$

(check `71d-D6`). In both cases the mass couples the two chiralities, and the self-interaction adds cubic terms to the right-hand sides.

Part 5. The 4+4 spacetime: the quantized sector and the two forms of the self-interaction

5.1 The covariant Lagrangian is indefinite

On the 4+4 spacetime only fields that do not depend on the three additional times $x5, x6, x7$ evolve unitarily in the ordinary time (Section 4.4). They form the **quantized sector**. For such fields the weighted field obeys the flat equations (Section 4.3), so that the flat form of the Lagrangian can be used. Write $\mathcal{L}_{\text{cov}}$ for the kinetic and mass terms (3.6) of the covariant Lagrangian in that form. With the matrices $\beta = -i\gamma^4$, $\alpha^j = -\gamma^4\gamma^j$ and $W_{1644} = i\,C^-\gamma^4$ of Section 2.1 one finds $C^-\gamma^4 = -i\,W_{1644}$, $C^-\gamma^j = -i\,W_{1644}\alpha^j$ and $C^- = -W_{1644}\beta$, and therefore

$$\mathcal{L}_{\text{cov}} = \tilde\psi\,(\gamma^A\partial_A - m)\,\psi = -\,\psi^\dagger\, W_{1644}\,(i\,\partial_4 - \mathsf{h})\,\psi, \qquad \mathsf{h} = \sum_{j=0}^{3} \alpha^j\,(-i\partial_j) + \beta\, m. \tag{5.1}$$

The Hermitian matrix W_{1644} has the eigenvalues $+1$ and -1, eight times each, and commutes with h. The sixteen components of the field therefore fall into two sets of eight, the two **halves** of the field,

$$\psi = \psi_+ + \psi_-, \qquad W_{1644}\,\psi_\pm = \pm\,\psi_\pm, \tag{5.2}$$

and h acts inside each half. At a given momentum four of the eight states of positive energy lie in each half: a particle has eight internal states, four in each half. Equation (5.1) says: *the covariant Lagrangian is the ordinary Lagrangian $\psi^\dagger(i\partial_4 - \mathsf{h})\psi$ of a Dirac field on the half $W_{1644} = -1$, and minus that Lagrangian on the half $W_{1644} = +1$.* The conserved charge (4.13) has the same structure.

For a free field the sign of a Lagrangian does not change the equation of motion: both halves obey $i\partial_4\psi = \mathsf{h}\psi$. The quantum theory of the free field is defined by the positive Lagrangian for all sixteen components,

$$\mathcal{L}_{\text{kin}} + \mathcal{L}_{\text{mass}} = \psi^\dagger(i\,\partial_4 - \mathsf{h})\psi = \bar\psi\,(i\,\beta\,\partial_4 - \beta\,\mathsf{h})\,\psi, \qquad \bar\psi = \psi^\dagger\beta, \tag{5.3}$$

in which every state of positive energy has positive norm. Its mass term is $-m\,\bar\psi\psi$, the same mass for both halves. The mass term of the covariant Lagrangian is different: because $C^- = -W_{1644}\beta$,

$$-m\,\tilde\psi\psi = +m\,\psi^\dagger\,W_{1644}\,\beta\,\psi = m\,(\psi_+^\dagger\beta\,\psi_+ - \psi_-^\dagger\beta\,\psi_-), \tag{5.4}$$

a mass term of opposite sign in the two halves. For the free field this difference is invisible, because the spectrum of h does not depend on the sign of m. With a self-interaction the two halves are no longer equivalent, and the quantized theory can be completed in two ways. They are called **form C** and **form Q**. Both keep the positive Lagrangian (5.3); they differ in the adjoint with which the nine invariants of the quartic term are built.

5.2 Form C, the covariant form: the definition of the 4+4 model

The quartic term is that of the covariant Lagrangian: the invariants (3.8), built with the adjoint $\tilde\psi = \psi^\dagger C^-$ and the matrices Θ^K of Eq. (3.7). They are written $I_r^{\mathrm C}$ where the two forms are compared:

$$\boxed{\mathcal{L}_{\mathrm{kin}} + \mathcal{L}_{\mathrm{mass}} + \mathcal{L}_{\mathrm{self}}^{\mathrm C} = \psi^\dagger(i\,\partial_4 - \mathsf{h})\psi - \frac{1}{2}\sum_{r=0}^{8} g_r\,I_r^{\mathrm C}, \qquad I_r^{\mathrm C} = \sum_{|K|=r} \mathrm{sgn}_K\,(\tilde\psi\,\Theta^K\psi)^2} \tag{5.5}$$

This is the definition of the 4+4 model in this project (instruction of the author of November 13, 2026). The quartic term is invariant under all 28 rotations and boosts of the 4+4 frame (check `44c-S4`), as the hypothesis requires of the invariants of the spacetime. The kinetic and mass terms (5.3) are invariant under the 13 of them that commute with W_{1644}: the ten rotations and boosts of $x0, \dots, x4$ and the three rotations of $x5, x6, x7$. The other fifteen mix a space-like coordinate or the ordinary time with an additional time; they do not preserve the quantized sector (they turn a field that does not depend on $x5, x6, x7$ into one that does).

The field equation of the quantized sector, for the weighted field on the deflating spacetime, follows by varying $\psi^\dagger$:

$$i\,\partial_4\hat\psi = \mathsf{h}\,\hat\psi + \frac{1}{V_d}\sum_{r=0}^{8} g_r \sum_{|K|=r} (\tilde{\hat\psi}\,\Theta_K\,\hat\psi)\,C^-\Theta^K\hat\psi. \tag{5.6}$$

It conserves $\int \hat\psi^\dagger\hat\psi$, the number of particles of both halves together.

5.3 Form Q, the quantized-sector form: the alternative

The bilinears are formed with the adjoint $\bar\psi = \psi^\dagger\beta$ of the positive Lagrangian (5.3) and with the eight matrices

$$\gamma_{44q}^A = -i\,\gamma_{1644}^A \quad (A = 0, \dots, 4), \qquad \gamma_{44q}^A = \gamma_{1644}^A \quad (A = 5, 6, 7). \tag{5.7}$$

They anticommute with one another, $\gamma_{44q}^4 = \beta$, and their squares are those of the Dirac matrices of a spacetime with one time: $+1$ for $A = 4$ and -1 otherwise (check `44q-S1`). In terms of them the positive Lagrangian (5.3) reads $\bar\psi\,(i\,\gamma_{44q}^A\partial_A - m)\,\psi$ for fields of the quantized sector, which has the form of the 7+1 Lagrangian (3.2). The matrices of the quartic term are built from them as in Eq. (3.4),

$$\Theta^K_Q = \zeta^Q_r\,\gamma^{A_1}_{44q}\cdots\gamma^{A_r}_{44q}, \qquad \zeta^Q_r = \begin{cases} i & \text{for } r = 2,3,6,7, \\ 1 & \text{for } r = 0,1,4,5,8, \end{cases}$$
$$(\Theta^K_Q)^2 = \mathrm{sgn}^Q_K\, 1, \qquad \mathrm{sgn}^Q_K = \prod_{A\in K} (\gamma^A_{44q})^2, \tag{5.8}$$

so that $\beta\,\Theta^K_Q$ is Hermitian (check `44q-S2`), and

$$I^{\mathrm{Q}}_r = \sum_{|K|=r} \mathrm{sgn}^Q_K\,(\bar\psi\,\Theta^K_Q\psi)^2, \qquad \mathcal{L}^{\mathrm{Q}}_{\mathrm{self}} = -\frac{1}{2}\sum_{r=0}^{8} g_r\, I^{\mathrm{Q}}_r. \tag{5.9}$$

The field equation of the quantized sector is

$$i\,\partial_4\hat\psi = \mathsf{h}\,\hat\psi + \frac{1}{V_d}\sum_{r=0}^{8} g_r \sum_{|K|=r} \mathrm{sgn}^Q_K\,(\bar{\hat\psi}\,\Theta^K_Q\,\hat\psi)\,\beta\,\Theta^K_Q\hat\psi, \tag{5.10}$$

term by term the equation (4.7) of the 7+1 spacetime for fields that do not depend on $x5, x6, x7$: the algebra of the matrices (5.7) is that of the 7+1 Dirac matrices. Every result that is obtained for such fields on 7+1 holds for form Q without change.

The invariants (5.9) are invariant under the group generated by the 28 products $\frac{1}{2}\gamma^A_{44q}\gamma^B_{44q}$. Of these, the 13 that commute with W_{1644} are the 13 transformations of the coordinates named in Section 5.2. The other 15 mix the two halves of the field and are not transformations of the coordinates of the 4+4 spacetime (check `44q-S4`).

5.4 The exact relation between the two forms

For every one of the 256 lists K (check `44c-S11`, exact),

$$C^-\Theta^K = z_K\, W_{1644}\,\beta\,\Theta^K_Q, \qquad z_K \in \{+1,-1,+i,-i\}, \tag{5.11}$$

where $z_K = \pm 1$ if K contains an even number of the indices $5,6,7$ (128 lists) and $z_K = \pm i$ if it contains an odd number (128 lists). Consequently

$$I^{\mathrm{C}}_r = \sum_{|K|=r} \mathrm{sgn}^Q_K\,(\psi^\dagger\, W_{1644}\,\beta\,\Theta^K_Q\,\psi)^2, \qquad \text{to be compared with} \qquad I^{\mathrm{Q}}_r = \sum_{|K|=r} \mathrm{sgn}^Q_K\,(\psi^\dagger\,\beta\,\Theta^K_Q\,\psi)^2. \tag{5.12}$$

Form C is form Q with the matrix W_{1644} inserted in every bilinear, and nothing else. What the insertion does is seen by decomposing the field into its halves (5.2). The matrix $\beta\,\Theta^K_Q$ is Hermitian, and with it each list K has four bilinears of the halves,

$$\mathcal{J}^{++}_K = \psi^\dagger_+\,\beta\,\Theta^K_Q\,\psi_+, \quad \mathcal{J}^{--}_K = \psi^\dagger_-\,\beta\,\Theta^K_Q\,\psi_-, \quad \mathcal{J}^{+-}_K = \psi^\dagger_+\,\beta\,\Theta^K_Q\,\psi_-, \quad \mathcal{J}^{-+}_K = \psi^\dagger_-\,\beta\,\Theta^K_Q\,\psi_+ = (\mathcal{J}^{+-}_K)^\dagger.$$

Lists with an even number of the indices 5, 6, 7. $\beta\,\Theta^K_Q$ commutes with W_{1644} and acts inside each half, so that $\mathcal{J}^{+-}_K = \mathcal{J}^{-+}_K = 0$. Form Q contains $(\mathcal{J}^{++}_K + \mathcal{J}^{--}_K)^2$ and form C contains $(\mathcal{J}^{++}_K - \mathcal{J}^{--}_K)^2$. The squares $(\mathcal{J}^{++}_K)^2$ and $(\mathcal{J}^{--}_K)^2$, which act between two particles of the same half, are equal in the

two forms. The product $2\,\mathcal{J}^{++}_K\mathcal{J}^{--}_K$, which acts between a particle of one half and a particle of the other, has opposite signs.

Lists with an odd number. $\beta\,\Theta^K_Q$ anticommutes with W_{1644} and connects the two halves, so that $\mathcal{J}^{++}_K = \mathcal{J}^{--}_K = 0$. Form Q contains $(\mathcal{J}^{+-}_K + \mathcal{J}^{-+}_K)^2$ and form C contains $(\mathcal{J}^{+-}_K - \mathcal{J}^{-+}_K)^2$. The squares $(\mathcal{J}^{+-}_K)^2$ and $(\mathcal{J}^{-+}_K)^2$ move two particles from one half to the other; they are equal in the two forms. The products $\mathcal{J}^{+-}_K\mathcal{J}^{-+}_K + \mathcal{J}^{-+}_K\mathcal{J}^{+-}_K$ exchange a particle of one half with a particle of the other; they have opposite signs.

> **Two particles of the same half interact identically in the two forms. A particle of the half $W_{1644} = +1$ and a particle of the half $W_{1644} = -1$ interact with opposite signs.**

Two further consequences are used below. (i) In both forms the interaction can move *two* particles together from one half to the other. The number of particles of each half is therefore not conserved separately; only the total number is, and the difference of the two numbers modulo 2. (ii) Both forms are symmetric under the exchange of the two halves. For fields in ordinary space this exchange is the transformation $\psi \to \gamma^0\gamma^5\psi$: the matrix $\gamma^0\gamma^5$ is Hermitian with square 1, it commutes with $\alpha^1, \alpha^2, \alpha^3$ and β, it anticommutes with W_{1644}, and it changes every bilinear of either form by a sign at most. For fields that depend on the hidden coordinate too, the same transformation is combined with the reflection $x0 \to -x0$.

5.5 What the covariant field equation itself says

If the quartic term of form C is combined with the *covariant* kinetic and mass terms (5.1) in place of the positive ones, the field equation is Eq. (4.2). Restricted to the quantized sector and multiplied by C^- it reads, with Eq. (5.1),

$$i\,\partial_4\psi = \mathsf{h}\,\psi + W_{1644}\,\frac{\partial\mathcal{L}^{\mathrm{C}}_{\mathrm{self}}}{\partial\psi^\dagger}, \tag{5.13}$$

whereas form C, Eq. (5.6), has $-\partial\mathcal{L}^{\mathrm{C}}_{\mathrm{self}}/\partial\psi^\dagger$ in place of $+W_{1644}\,\partial\mathcal{L}^{\mathrm{C}}_{\mathrm{self}}/\partial\psi^\dagger$: the interaction term of the covariant equation carries the factor $-W_{1644}$. On the half $W_{1644} = -1$ the two equations agree. On the half $W_{1644} = +1$ the covariant equation has the opposite sign of the whole interaction, the same change of sign as in Eq. (5.1). It conserves $\psi^\dagger W_{1644}\psi$ and not $\psi^\dagger\psi$, and it is not the equation of a Hamiltonian with states of positive norm when both halves are occupied. It is a consistent classical field equation, the one that the covariant Lagrangian implies literally; no quantum theory of particles is built on it in this project.

5.6 The three options side by side

option	kinetic and mass term	bilinears of the quartic term	status
form C	positive, Eq. (5.3)	$\tilde\psi\,\Theta^K\psi$ with $\tilde\psi = \psi^\dagger C^-$, Eq. (5.5)	**the definition of the 4+4 model**
form Q	positive, Eq. (5.3)	$\bar\psi\,\Theta^K_Q\psi$ with $\bar\psi = \psi^\dagger\beta$, Eq. (5.9)	the alternative; its results are reported beside those of the definition
covariant classical equation	indefinite, Eq. (5.1)	as in form C	described in Section 5.5; not quantized

The hypotheses of Section 0.1 fix the nine invariants as the invariants of the spacetime. On the 7+1 spacetime this determines the model. On the 4+4 spacetime it does not, because the adjoint that makes the invariants covariant (C^-) is not the adjoint that makes the energy positive (β): the

two differ by the matrix W_{1644}. The author has decided for form C. Part 6 derives what each form implies, and Section 6.7 discusses the two options.

Part 6. The algebra of the self-interaction, and what the two forms imply

This part analyzes the quartic term by exact algebra (Sections 6.1 to 6.5), gives the consequences of the two forms of the 4+4 model for matter and for the vacuum in numbers (Section 6.6), and discusses the two forms (Section 6.7). A statement that names no spacetime and no form holds for the 7+1 spacetime and for both forms of the 4+4 spacetime. The bilinears are $\psi^\dagger \,\mathrm{Adj}\,\Theta^K \psi$ with the adjoint matrix and the matrices of the model concerned: Adj $= \beta$ on 7+1; Adj $= C^-$ in form C; Adj $= \beta$ with the matrices Θ^K_Q and the signs sgn^Q_K of Eq. (5.8) in form Q.

6.1 The Fierz matrix

The 256 matrices Θ^K are a basis of all complex 16×16 matrices, with $\mathrm{Tr}(\Theta_K \Theta^J) = 16\,\delta_{KJ}$ for two lists K, J (checks `71s-S3`, `44q-S3`, `44c-S3`). Hence $\sum_K (\Theta_K)_{cd} (\Theta^K)_{ab} = 16\,\delta_{ad}\,\delta_{bc}$, the completeness relation, in which $a, b, c, d = 1, \ldots, 16$ number the components. With it a product of two matrices of the basis can be rearranged:

$$\sum_{|K|=r} (\Theta_K)_{ab}\,(\Theta^K)_{cd} = \sum_{r'=0}^{8} \mathsf{F}_{rr'} \sum_{|J|=r'} (\Theta_J)_{ad}\,(\Theta^J)_{cb},$$
$$\mathsf{F}_{rr'} = \frac{1}{16} \sum_l (-1)^{rr'-l} \binom{r'}{l} \binom{8-r'}{r-l}. \tag{6.1}$$

F (sans-serif) is the **Fierz matrix**; r, r' are ranks and l is a summation index. The derivation takes three lines. Expanding the left-hand side in the basis gives, for each list J of rank r', the coefficient $\frac{1}{256} \sum_{|K|=r} \mathrm{Tr}(\Theta_K \Theta_J \Theta^K \Theta^J)$. Two products of Dirac matrices either commute or anticommute: $\Theta_K \Theta_J \Theta^K = (-1)^{rr'-l}\,\Theta_J$ if the lists K and J have l indices in common. And $\binom{r'}{l}\binom{8-r'}{r-l}$ lists K of rank r share exactly l indices with a given J of rank r'. The integers $16\,\mathsf{F}_{rr'}$ are (rows r, columns r'):

r	0	1	2	3	4	5	6	7	8
0	1	1	1	1	1	1	1	1	1
1	8	-6	4	-2	0	2	-4	6	-8
2	28	14	4	-2	-4	-2	4	14	28
3	56	-14	-4	6	0	-6	4	14	-56
4	70	0	-10	0	6	0	-10	0	70
5	56	14	-4	-6	0	6	4	-14	-56
6	28	-14	4	2	-4	2	4	-14	28
7	8	6	4	2	0	-2	-4	-6	-8
8	1	-1	1	-1	1	-1	1	-1	1

The matrix satisfies $\mathsf{F}^2 = 1$; it has the eigenvalue $+1$ five times and -1 four times (checks `71s-S5`, `44q-S5`, `44c-S5`, which verify the identity (6.1) for all 9×65536 combinations of the rank and of the four indices). It is the same matrix on 7+1 and in both forms of 4+4.

6.2 Anticommuting fields: only four of the nine invariants are independent

Contract Eq. (6.1) with two adjoint fields (indices a and c) and two fields (indices b and d). The left-hand side is the invariant I_r. The right-hand side contains the two fields ψ in the opposite order, and the components of ψ anticommute, which costs a sign:

$$I_r = -\sum_{r'} \mathsf{F}_{rr'}\, I_{r'}, \qquad \text{that is} \qquad (1+\mathsf{F})\, I = 0. \tag{6.2}$$

Because F has the eigenvalue $+1$ five times, these are five independent linear relations among the nine invariants: **only four invariants are independent.** Solved for the invariants of the ranks 2 to 6:

$$\begin{aligned} &I_2 = -3I_0 - I_1 - I_7 - I_8, \qquad && I_3 = -4I_0 - I_7 + 4I_8, \qquad && I_4 = -5I_0 - 5I_8, \\ &I_5 = -4I_0 - I_1 + 4I_8, && I_6 = -I_0 + I_1 + I_7 - 3I_8. \end{aligned} \tag{6.3}$$

Inserted into the self-interaction,

$$\mathcal{L}_{\text{self}} = -\tfrac{1}{2}(g_0^{\text{eff}}\, I_0 + g_1^{\text{eff}}\, I_1 + g_7^{\text{eff}}\, I_7 + g_8^{\text{eff}}\, I_8), \tag{6.4}$$

with the four **effective couplings**

$$\begin{aligned} &g_0^{\text{eff}} = g_0 - 3g_2 - 4g_3 - 5g_4 - 4g_5 - g_6, \qquad && g_1^{\text{eff}} = g_1 - g_2 - g_5 + g_6, \\ &g_7^{\text{eff}} = g_7 - g_2 - g_3 + g_6, && g_8^{\text{eff}} = g_8 - g_2 + 4g_3 - 5g_4 + 4g_5 - 3g_6. \end{aligned} \tag{6.5}$$

A tensor coupling $g_2, \dots, g_6$ is not a new parameter of the theory: it shifts the scalar, vector, axial-vector and pseudoscalar couplings. Every physical quantity depends on the nine g_r only through the four combinations (6.5). The relations (6.3) and the combinations (6.5) are the same on 7+1 and in both forms of 4+4 (with I_r^{C} or I_r^{Q} for I_r): a model is specified by the four numbers $g_0^{\text{eff}}, g_1^{\text{eff}}, g_7^{\text{eff}}, g_8^{\text{eff}}$, and on 4+4 the choice of the form decides what these four numbers mean. The theorem (6.2) is verified in two independent ways: by exact integer arithmetic on the 16^4 coefficients of each quartic form, antisymmetrized as anticommuting fields require (checks `71s-S6`, `44q-S6`, `44c-S6`: the nine antisymmetrized forms span a space of dimension four), and, in the independent script, by evaluating the nine invariants on Grassmann-valued fields with an implementation of the Grassmann algebra.

6.3 Uniform matter: the mean-field couplings

In uniform matter at rest the expectation value of a product of two fields is the **density matrix** $\mathsf{n}_{ab} = \langle \psi_b^\dagger \psi_a \rangle$. By Wick's theorem the expectation value of a normal-ordered product of two bilinears is a direct (Hartree) term minus an exchange (Fock) term,

$$\big\langle (\psi^\dagger\, \text{Adj}\, \Theta^K \psi)^2 \big\rangle = \big[\text{Tr}(\text{Adj}\, \Theta^K \mathsf{n})\big]^2 - \text{Tr}(\text{Adj}\, \Theta^K \mathsf{n}\, \text{Adj}\, \Theta^K \mathsf{n}).$$

The 7+1 spacetime: two couplings. For a filled Fermi sea of particles the density matrix has the form $\mathsf{n} = (n\,1 + n_S\,\beta)/16$, where $n = \langle \psi^\dagger \psi \rangle$ is the number density and $n_S = \langle \bar\psi \psi \rangle$ the scalar density. The energy density of the self-interaction, $-\langle \mathcal{L}_{\text{self}} \rangle$, becomes

$$\varepsilon_{\text{int}} = \tfrac{1}{2}\, G_V^{\text{HF}}\, n^2 + \tfrac{1}{2}\, G_S^{\text{HF}}\, n_S^2, \tag{6.6}$$

$$G_S^{\text{HF}} = \tfrac{1}{16}(15\, g_0^{\text{eff}} - 8\, g_1^{\text{eff}} - 8\, g_7^{\text{eff}} - g_8^{\text{eff}}), \qquad G_V^{\text{HF}} = \tfrac{1}{16}(-\, g_0^{\text{eff}} + 22\, g_1^{\text{eff}} - 6\, g_7^{\text{eff}} + g_8^{\text{eff}}). \tag{6.7}$$

These are the **Hartree–Fock couplings**. In terms of the nine couplings, $G_S^{\rm HF} = \frac{1}{16}(15g_0 - 8g_1 - 28g_2 - 56g_3 - 70g_4 - 56g_5 - 28g_6 - 8g_7 - g_8)$ and $G_V^{\rm HF} = \frac{1}{16}(-g_0 + 22g_1 - 14g_2 + 14g_3 - 14g_5 + 14g_6 - 6g_7 + g_8)$ (check `71s-S7`). The direct terms alone would give the scalar coupling g_0 and the vector coupling g_1; the exchange terms reduce the scalar coupling by one sixteenth and add to the vector coupling. Uniform matter on 7+1 is thus described by two numbers, whatever the nine couplings are.

Example. For $g_0 = -10^5$, $g_1 = +10^4$ and all other couplings zero, the example of the author's document on the 7+1 problem (in units in which the mass m is one), Eq. (6.7) gives $G_S^{\rm HF} = -98750$ and $G_V^{\rm HF} = +20000$ (check `self-X1`).

The 4+4 spacetime: four couplings, whose roles depend on the form. When the two halves (5.2) of the field are filled with particles up to the Fermi momenta k_{F+} and k_{F-}, and no coherence between the halves is present, the density matrix has four terms,

$$\mathsf{n} = \tfrac{1}{16}(n\,1 + n_S\,\beta + n_W\,W_{1644} + n_{SW}\,W_{1644}\,\beta), \tag{6.8}$$

$$n = n_+ + n_-, \qquad n_W = n_+ - n_-, \qquad n_S = n_{S+} + n_{S-}, \qquad n_{SW} = n_{S+} - n_{S-},$$

where $n_\pm = \langle \psi_\pm^\dagger \psi_\pm \rangle$ are the number densities of the two halves and $n_{S\pm} = \langle \psi_\pm^\dagger \beta\, \psi_\pm \rangle$ their scalar densities. In both forms the energy density of the self-interaction is a sum of four squares (checks `44c-S7`, `44q-S7`, exact):

$$\varepsilon_{\rm int} = \tfrac{1}{2}(G_V^{\rm tot}\, n^2 + G_S^{\rm tot}\, n_S^2 + G_V^{\rm dif}\, n_W^2 + G_S^{\rm dif}\, n_{SW}^2). \tag{6.9}$$

$G_V^{\rm tot}, G_S^{\rm tot}$ are the couplings of the squares of the **total** densities, $G_V^{\rm dif}, G_S^{\rm dif}$ those of the squares of the **differences** between the halves. They are the two Hartree–Fock couplings (6.7) and two further combinations,

$$G_S^{\rm c} = \tfrac{1}{16}(-g_0^{\rm eff} - 2\,g_1^{\rm eff} + 2\,g_7^{\rm eff} + g_8^{\rm eff}), \qquad G_V^{\rm c} = -\tfrac{1}{16}(g_0^{\rm eff} + g_8^{\rm eff}), \tag{6.10}$$

the same four in both forms, with the roles of the totals and of the differences exchanged (check `44-S13`, exact):

coupling	multiplies	form C (the definition)	form Q (the alternative)
$G_V^{\rm tot}$	n^2	$G_V^{\rm c}$	$G_V^{\rm HF}$
$G_S^{\rm tot}$	n_S^2	$G_S^{\rm c}$	$G_S^{\rm HF}$
$G_V^{\rm dif}$	n_W^2	$G_V^{\rm HF}$	$G_V^{\rm c}$
$G_S^{\rm dif}$	n_{SW}^2	$G_S^{\rm HF}$	$G_S^{\rm c}$

In terms of the nine couplings: $G_S^{\rm c} = \frac{1}{16}(-g_0 - 2g_1 + 2g_2 + 6g_3 + 10g_5 - 2g_6 + 2g_7 + g_8)$ and $G_V^{\rm c} = \frac{1}{16}(-g_0 + 4g_2 + 10g_4 + 4g_6 - g_8)$. An equivalent way to write Eq. (6.9) shows what happens between particles:

$$\varepsilon_{\rm int} = \tfrac{1}{2}\,G_V^{\rm same}\,(n_+^2 + n_-^2) + \tfrac{1}{2}\,G_S^{\rm same}\,(n_{S+}^2 + n_{S-}^2) + G_V^{\rm opp}\,n_+ n_- + G_S^{\rm opp}\,n_{S+} n_{S-}, \tag{6.11}$$

with $G^{\rm same} = G^{\rm tot} + G^{\rm dif}$ and $G^{\rm opp} = G^{\rm tot} - G^{\rm dif}$, for the scalar and for the vector couplings alike:

$$
\begin{aligned}
G_S^{\text{same}} &= G_S^{\text{HF}} + G_S^{\text{c}} = \tfrac{1}{8}(7\,g_0^{\text{eff}} - 5\,g_1^{\text{eff}} - 3\,g_7^{\text{eff}}),\\
G_V^{\text{same}} &= G_V^{\text{HF}} + G_V^{\text{c}} = \tfrac{1}{8}(-\,g_0^{\text{eff}} + 11\,g_1^{\text{eff}} - 3\,g_7^{\text{eff}}),\\
G_S^{\text{opp}} &= \pm(G_S^{\text{HF}} - G_S^{\text{c}}) = \pm\tfrac{1}{8}(8\,g_0^{\text{eff}} - 3\,g_1^{\text{eff}} - 5\,g_7^{\text{eff}} - g_8^{\text{eff}}),\\
G_V^{\text{opp}} &= \pm(G_V^{\text{HF}} - G_V^{\text{c}}) = \pm\tfrac{1}{8}(11\,g_1^{\text{eff}} - 3\,g_7^{\text{eff}} + g_8^{\text{eff}}),
\end{aligned}
\tag{6.12}
$$

with the upper sign in form Q and the lower sign in form C. $G_S^{\text{same}}, G_V^{\text{same}}$ are the couplings between two particles of the same half; they are the same in both forms. $G_S^{\text{opp}}, G_V^{\text{opp}}$ are the couplings between a particle of one half and a particle of the other; they have opposite signs in the two forms. This is the statement of Section 5.4 for the mean field.

What this means for an attractive scalar coupling ($g_0 < 0$, the other couplings zero). In form Q the attraction stands in $G_S^{\text{tot}} = \frac{15}{16} g_0$: it lowers the energy of any scalar density, and matter with both halves equally filled feels it, as on 7+1. In form C it stands in $G_S^{\text{dif}} = \frac{15}{16} g_0$: it lowers the energy only if the scalar densities of the two halves differ, and the total scalar density is weakly repelled ($G_S^{\text{tot}} = -\frac{1}{16} g_0 > 0$). Between two particles of the same half the coupling attracts in both forms ($G_S^{\text{same}} = \frac{7}{8} g_0$); between particles of opposite halves it attracts in form Q ($G_S^{\text{opp}} = g_0$) and repels in form C ($G_S^{\text{opp}} = -g_0$). Section 6.6 gives the consequences in numbers.

6.4 Two particles: the pair couplings

A particle at rest has eight internal states (the eigenvalue $+1$ of β), and two particles at rest can be in $8 \cdot 7/2 = 28$ antisymmetric states. The self-interaction acts on a pair state Φ, an antisymmetric 16×16 matrix, as

$$
\Phi \;\mapsto\; \sum_r g_r \sum_{|K|=r} \text{sgn}_K \left(\text{Adj}\,\Theta^K\right) \Phi \left(\text{Adj}\,\Theta^K\right)^T,
\tag{6.13}
$$

projected on the states of positive energy with $\frac{1}{2}(1+\beta)$ on either side (in form Q with Θ_Q^K and sgn_K^Q). Its eigenvalue is the **pair coupling** G_{pair}, the interaction energy of the pair times the volume; a negative value means attraction.

7+1, and form Q of 4+4. The map has two eigenvalues (checks `71s-S9`, `44q-S9`):

$$
G_{\text{pair}} = \begin{cases} g_0^{\text{eff}} + g_1^{\text{eff}} + 5\,g_7^{\text{eff}} & \text{for 7 pairs,} \\ g_0^{\text{eff}} + g_1^{\text{eff}} - 3\,g_7^{\text{eff}} & \text{for 21 pairs.} \end{cases}
\tag{6.14}
$$

The seven pairs are those created by the **vector pair operators** $\psi^T C^- \gamma^A \psi$ along the seven axes other than the time ($A \neq 4$); among them is the pair along the hidden axis $x0$, the spin-singlet pair of the superfluid that the project studies. A scalar attraction ($g_0^{\text{eff}} < 0$) binds all 28 pairs equally; a vector repulsion ($g_1^{\text{eff}} > 0$) opposes all of them equally; the axial coupling distinguishes the seven from the twenty-one; the pseudoscalar coupling does not act between particles at rest.

Form C of 4+4. The state that a vector pair operator creates has a definite behavior under W_{1644} (check `44c-S9`):

$$
\begin{aligned}
W_{1644}\,\Phi\,W_{1644}^T &= -\Phi \quad \text{for the pairs along } x0, x1, x2, x3,\\
W_{1644}\,\Phi\,W_{1644}^T &= +\Phi \quad \text{for the pairs along } x5, x6, x7.
\end{aligned}
\tag{6.15}
$$

A pair along one of the four space-like axes consists of **one particle of each half**; a pair along one of the three additional time-like axes consists of **two particles of the same half**. By Section 5.4

the two forms agree on the pairs of two particles of the same half and have opposite signs on the pairs of one particle of each half. For all 28 pairs, in exact arithmetic (checks `44c-S9`, `44q-S9`):

form	number of pairs	the two particles are in	G_{pair}
C	4	different halves	$-g_0^{\text{eff}} - g_1^{\text{eff}} - 5\,g_7^{\text{eff}}$
C	12	different halves	$-g_0^{\text{eff}} - g_1^{\text{eff}} + 3\,g_7^{\text{eff}}$
C	9	same half	$g_0^{\text{eff}} + g_1^{\text{eff}} - 3\,g_7^{\text{eff}}$
C	3	same half	$g_0^{\text{eff}} + g_1^{\text{eff}} + 5\,g_7^{\text{eff}}$
Q	12	different halves	$g_0^{\text{eff}} + g_1^{\text{eff}} - 3\,g_7^{\text{eff}}$
Q	4	different halves	$g_0^{\text{eff}} + g_1^{\text{eff}} + 5\,g_7^{\text{eff}}$
Q	9	same half	$g_0^{\text{eff}} + g_1^{\text{eff}} - 3\,g_7^{\text{eff}}$
Q	3	same half	$g_0^{\text{eff}} + g_1^{\text{eff}} + 5\,g_7^{\text{eff}}$

In form Q the seven vector pairs are degenerate. In form C they split into the four pairs along the space-like axes, with $-(g_0^{\text{eff}} + g_1^{\text{eff}} + 5\,g_7^{\text{eff}})$, and the three pairs along the additional axes, with $+(g_0^{\text{eff}} + g_1^{\text{eff}} + 5\,g_7^{\text{eff}})$. *A scalar coupling that attracts in form Q, and on 7+1, repels in the hidden-axis pair of form C.*

6.5 Symmetries of the self-interaction

Chiral symmetry. The transformation $\psi \to e^{i\epsilon\chi}\psi$ with the chirality matrix χ of Section 2.1 and a real parameter ϵ leaves the self-interaction invariant if and only if

$$g_0^{\text{eff}} + g_8^{\text{eff}} = 0 \quad \text{(7+1, and form Q of 4+4)}, \qquad g_1^{\text{eff}} - g_7^{\text{eff}} = 0 \quad \text{(form C of 4+4)} \tag{6.16}$$

(checks `71s-S10`, `44q-S10`, `44c-S10`). On 7+1 and in form Q the transformation leaves I_1 and I_7 unchanged and rotates I_0 and I_8 into each other. In form C the roles of the even and of the odd ranks are exchanged, because the bilinears of the two forms differ by the matrix W_{1644}, which anticommutes with χ. A scalar coupling alone is chirally invariant in form C and not in form Q; a vector coupling alone is chirally invariant in form Q and not in form C. The mass term breaks this symmetry in any case.

The internal symmetry of fields in ordinary space. A field that depends on $x1, x2, x3, x4$ only has a kinetic term that contains $\alpha^1, \alpha^2, \alpha^3$ and β. Eight Hermitian matrices commute with these four (Part 8): the six generators proportional to $\gamma^a\gamma^b$ ($a < b$ in $\{0, 5, 6, 7\}$) of the rotations of the four extra axes, which form $SO(4)$; the unit matrix (the fermion number, with the phase symmetry $U(1)_F$); and the matrix P proportional to $\gamma^0\gamma^5\gamma^6\gamma^7$, which generates a second phase symmetry, $U(1)_E$. On 4+4 the three rotations of $x5, x6, x7$ among themselves commute with W_{1644}; the three generators $\gamma^0\gamma^t$ ($t = 5, 6, 7$) and P anticommute with W_{1644} and connect the two halves. Under which couplings is the self-interaction invariant under the unitary transformations that each of them generates (the field is multiplied by the exponential of i times an angle times the matrix)? In exact arithmetic (checks `71s-S12`, `44q-S12`, `44c-S12`; confirmed at a finite angle by the independent script, check `Icks-F2`):

generators	number	$7+1$	$4+4$, form Q	$4+4$, form C
phase of the field (fermion number)	1	all couplings	all couplings	all couplings
rotations of $x5, x6, x7$ among themselves	3	all couplings	all couplings	all couplings
rotations that mix $x0$ with $x5$, $x6$ or $x7$	3	all couplings	all couplings	no coupling ($g_0^{\text{eff}} = g_1^{\text{eff}} = g_7^{\text{eff}} = g_8^{\text{eff}} = 0$)

generators	number	$7+1$	$4+4$, form Q	$4+4$, form C
P, the generator of $U(1)_E$	1	$g_1^{\text{eff}} = g_7^{\text{eff}} = 0$	$g_1^{\text{eff}} = g_7^{\text{eff}} = 0$	no coupling ($g_0^{\text{eff}} = g_1^{\text{eff}} = g_7^{\text{eff}} = g_8^{\text{eff}} = 0$)

In form C every self-interaction that does not vanish breaks the three transformations that mix the hidden axis with $x5, x6, x7$, and breaks $U(1)_E$. The internal symmetry of fields in ordinary space is reduced from $SO(4)\times U(1)_F \times U(1)_E$ to $SO(3)_T \times U(1)_F$: the rotations $SO(3)_T$ of the three additional time-like coordinates among themselves, and the fermion number.

On 7+1 and in form Q the self-interaction keeps the six rotations and $U(1)_F$ for all couplings, and keeps $U(1)_E$ if and only if $g_1^{\text{eff}} = g_7^{\text{eff}} = 0$: a vector or an axial-vector coupling breaks $U(1)_E$, a scalar and a pseudoscalar coupling keep it. The four conditions of the table for form C, $5g_0^{\text{eff}} - 16g_7^{\text{eff}} + g_8^{\text{eff}} = 0$, $5g_1^{\text{eff}} - 4g_7^{\text{eff}} - g_8^{\text{eff}} = 0$, $4g_7^{\text{eff}} + g_8^{\text{eff}} = 0$ and $g_7^{\text{eff}} - g_8^{\text{eff}} = 0$, have only the solution in which all four effective couplings vanish.

Why. In the frame group of the 4+4 spacetime $\gamma^0\gamma^t$ generates a **boost** (it mixes a space-like with a time-like coordinate), which is not a unitary transformation of the field. The *unitary* transformation generated by the same matrix is a different transformation: it is a symmetry of the kinetic term of fields in ordinary space by accident, and it is not an element of the frame group. The quartic term of form C is invariant under the boost and has no reason to be invariant under the unitary transformation. In form Q, as on the 7+1 spacetime, where all four extra axes are space-like, the same matrices generate rotations, which are unitary, and the quartic term is invariant under them. For the pairs of Section 6.4 this means: in form Q the pair along $x0$ can be rotated into the pairs along $x5, x6, x7$, so that they are degenerate; in form C it cannot, and they are not.

6.6 What the two forms imply for matter and for the vacuum, in numbers

The numbers of this section are computed by the command `forms` of the program (checks `fm-F1` to `fm-F9`, Appendix A.4) for two sets of couplings of the Lagrangian, each with one strength $g > 0$:

- **set A**: a scalar attraction alone, $g_0 = -g$, all other couplings zero;
- **set B**: $g_0 = -g$ and $g_1 = +0.1\,g$ (the ratio of the author's example), all other couplings zero.

On the deflating spacetime every coupling of the weighted field grows as $1/V_d \propto e^{\text{gsum}\,x4}$ (Section 4.3): a strength g is reached at a definite time, and a ratio of two strengths is a difference of times, $\Delta x4 = \ln(\text{ratio})/\text{gsum}$.

The mean-field equations of two halves. In the mean field (6.9) the particles of the half $W_{1644} = \pm 1$ move with a mass $m_\pm^*$ and an energy shift $V_{0\pm}$,

$$m_\pm^* = m + G_S^{\text{tot}}\, n_S \pm G_S^{\text{dif}}\, n_{SW}, \qquad V_{0\pm} = G_V^{\text{tot}}\, n \pm G_V^{\text{dif}}\, n_W, \tag{6.17}$$

and fill the states up to the Fermi momenta $k_{F\pm}$. In D space dimensions ($D = 4$: the quantized sector, momenta along $x0, x1, x2, x3$; $D = 3$: fields that do not depend on the hidden coordinate $x0$ either), with the energy $E_\pm^*(k) = \sqrt{k^2 + m_\pm^{*2}}$ and four internal states per half,

$$n_\pm = 4\int_{|\vec{k}|<k_{F\pm}} \frac{d^Dk}{(2\pi)^D}, \qquad n_{S\pm} = 4\int_{|\vec{k}|<k_{F\pm}} \frac{d^Dk}{(2\pi)^D}\,\frac{m_\pm^*}{E_\pm^*}, \tag{6.18}$$

$$\varepsilon = \sum_\pm 4\int_{|\vec{k}|<k_{F\pm}} \frac{d^Dk}{(2\pi)^D}\,E_\pm^* + \tfrac{1}{2}(G_V^{\text{tot}}\, n^2 + G_V^{\text{dif}}\, n_W^2) - \tfrac{1}{2}(G_S^{\text{tot}}\, n_S^2 + G_S^{\text{dif}}\, n_{SW}^2). \tag{6.19}$$

ε is the energy density of matter. These equations count the particles only, not the filled states of negative energy (the **no-sea approximation**; the equations are those of the Kohn–Sham method for this interaction). Because the interaction can move two particles from one half to the other (Section 5.4), the state at a given total density n is the one of lowest energy with respect to the **polarization** $\mathrm{pol}_W = (n_+ - n_-)/n$.

Form C polarizes matter; form Q does not. For the same couplings of the Lagrangian, in 48 cases (two dimensions, two sets, three strengths, four densities), the state of lowest energy is polarized in 48 cases in form C and in 0 cases in form Q (check `fm-F3`). In form C a particle is attracted by the particles of its own half ($G_S^{\mathrm{same}} < 0$) and repelled by those of the other half ($G_S^{\mathrm{opp}} > 0$), so that the particles gather in one half; in form Q it is attracted by both.

The strength at which matter first binds. Matter is *self-bound* when its energy per particle, minimized over the density and the polarization, lies below the mass m. The smallest strength at which this happens is (units $m = 1$; "state": the state of lowest energy just above that strength; $k_{F,\mathrm{tot}}$: the Fermi momentum that the total density would have with equal halves; $m_\pm^*$: the masses of the particles of the two halves, for an empty half the mass that a particle of that half would have; Λ_{max} is explained below):

D	set	form	$g\,m^{D-1}$	state	$k_{F,\mathrm{tot}}/m$	m_+^*/m	m_-^*/m	Λ_{max}/m
4	A	C	224.7386	one half	0.6140	0.5239	1.5441	0.520
4	A	Q	82.1116	equal halves	0.8054	0.4946	0.4946	0.727
4	B	C	332.2917	one half	0.5211	0.5625	1.4842	0.449
4	B	Q	121.0858	equal halves	0.6792	0.5424	0.5424	0.628
3	A	C	24.5717	one half	0.5374	0.6454	1.4053	0.463
3	A	Q	9.8090	equal halves	0.7356	0.6276	0.6276	0.733
3	B	C	31.1674	one half	0.4635	0.6712	1.3638	0.400
3	B	Q	12.3819	equal halves	0.6351	0.6575	0.6575	0.635

In form C the bound state has one half filled, in form Q equal halves, in all eight cases (check `fm-F2`). Matter of form C needs 2.74 times (set A) and 2.74 times (set B) the strength of form Q for $D = 4$, and 2.51 and 2.52 times for $D = 3$: in units of $1/\mathrm{gsum}$ it binds later by 1.01 and 1.01 ($D = 4$), 0.92 and 0.92 ($D = 3$). Only four of the eight internal states take part, and the attraction inside one half, $G_S^{\mathrm{same}} = \frac{7}{8} g_0$, is weaker than the attraction $G_S^{\mathrm{tot}} = \frac{15}{16} g_0$ that matter of form Q with equal halves feels.

The vacuum. When the states of negative energy are filled up to a cutoff Λ of the momentum, the vacuum itself has scalar densities $n_{S\pm}^{\mathrm{vac}} = -4 \int_{|\vec{k}|<\Lambda} d^D k/(2\pi)^D \; m_\pm^*/E_\pm^*$, and the masses of the two halves in vacuum obey

$$\tfrac{1}{2}(m_+^* + m_-^*) - m = G_S^{\mathrm{tot}} \left(n_{S+}^{\mathrm{vac}} + n_{S-}^{\mathrm{vac}}\right), \qquad \tfrac{1}{2}(m_+^* - m_-^*) = G_S^{\mathrm{dif}} \left(n_{S+}^{\mathrm{vac}} - n_{S-}^{\mathrm{vac}}\right). \tag{6.20}$$

For a field without mass term ($m = 0$) the solution $m_+^* = m_-^* = 0$ always exists. A second solution appears when the more attractive of the two scalar couplings exceeds the **critical coupling** $G_c = -(2\pi)^D (D-1)/(8\, A_D\, \Lambda^{D-1})$, with A_D the area of the unit sphere in D dimensions ($A_4 = 2\pi^2$, $A_3 = 4\pi$). For the sets A and B that coupling is $G_S^{\mathrm{HF}} < 0$ in both forms, so that the critical strength g_{crit} is the same for both. The table gives it, and the masses of the two halves at a strength 2 per cent above it, in units of the cutoff (check `fm-F7`):

D	set	$G_c \Lambda^{D-1}$	$g_{\mathrm{crit}} \Lambda^{D-1}$	form Q: $(m_+^*, m_-^*)/\Lambda$	form C: $(m_+^*, m_-^*)/\Lambda$
4	A	-29.6088	31.5827	$(0.1243,\ 0.1243)$	$(0.1243,\ -0.1243)$
4	B	-29.6088	29.9836	$(0.1243,\ 0.1243)$	$(0.1243,\ -0.1243)$

D	set	$G_c\Lambda^{D-1}$	$g_{\text{crit}}\Lambda^{D-1}$	form Q: $(m_+^*, m_-^*)/\Lambda$	form C: $(m_+^*, m_-^*)/\Lambda$
3	A	-4.9348	5.2638	$(0.0860,\ 0.0860)$	$(0.0860,\ -0.0860)$
3	B	-4.9348	4.9973	$(0.0860,\ 0.0860)$	$(0.0860,\ -0.0860)$

In form Q the two halves acquire the same mass: a mass term proportional to $\bar{\psi}\psi$, the mass term of the positive Lagrangian (5.3). In form C they acquire masses of opposite sign: by Eq. (5.4) a mass term proportional to $\tilde{\psi}\psi$, the mass term of the covariant Lagrangian.

Each form generates dynamically the mass term that belongs to its own adjoint; the magnitude of the mass is the same. (The solutions are degenerate by the exchange of the halves and, without mass term, by the reversal of both signs; the table lists the representative with $m_+^* > 0$.)

Where the no-sea approximation applies. The matter above was computed without the filled negative-energy states. That is consistent only while the vacuum is stable, that is, while the most attractive scalar coupling is weaker than G_c for the cutoff of the model. The column Λ_{max} of the table of thresholds gives the largest cutoff, in units of m, for which this holds at the binding threshold. It is below m in all eight cases, and smaller in form C than in form Q: *in both forms, and more so in form C, self-bound matter by contact couplings alone lies outside the range in which the no-sea approximation is controlled.*

6.7 Discussion of the two forms

What each form is. Both forms keep what the quantum theory needs: the positive kinetic term, a real (Hermitian) interaction, and the conservation of the number of particles. Both use the same nine couplings and the same four effective couplings. They assign to these couplings different meanings. Form C takes the hypothesis literally: the quartic term is built from the invariants of the 4+4 spacetime. Form Q takes the *particle content* literally: the eight internal states of a particle are treated alike, as on the 7+1 spacetime.

Arguments for form C.

1. *It is the quartic term of the hypothesis.* The invariants I_r^{C} are the ones that are invariant under all rotations and boosts of the 4+4 frame. They can be written on the full deflating spacetime, for fields that depend on all eight coordinates, with the spin connection (Eq. (3.8)); the quantized sector is a restriction of that covariant theory. Form Q is defined in the quantized sector only: its invariance group is not a symmetry group of the 4+4 spacetime, and it has no covariant extension to fields that depend on the additional times.
2. *Its dynamics respects the distinction that the metric makes.* On the 4+4 spacetime $x0$ is space-like and $x5, x6, x7$ are time-like. Form C treats them differently (Sections 6.4 and 6.5); form Q has an accidental symmetry that rotates them into one another.
3. *Each structure appears with its own adjoint.* The mass that the interaction of form C generates in vacuum is the covariant mass term (Section 6.6).

Arguments against form C, or for form Q.

1. *The two adjoints are mixed.* Form C combines the quartic term of the covariant Lagrangian with the kinetic and mass terms of the positive Lagrangian. The Lagrangian as a whole is invariant only under the thirteen generators that both parts share, the same thirteen as for form Q. The covariance of the quartic term is not a symmetry of the model.
2. *The bare mass term and the dynamical mass have different structures* in form C: the bare mass term is $-m\,\bar{\psi}\psi$ and the mass that the interaction generates is proportional to $\tilde{\psi}\psi$. In form Q the two have the same structure.
3. *The results of the 7+1 spacetime do not carry over.* In form Q every result for fields that do not depend on $x5, x6, x7$ is the same on the two spacetimes; in form C the 4+4 model is a different model, with polarized matter, later binding, and pairs that are not degenerate.

4. *The symmetry pattern of the author's Higgs hypothesis is lost.* That hypothesis identifies the condensate of pairs along the hidden axis with the Higgs field: the six generators of the $SO(4)$ of Section 6.5 would play the role of the electroweak group, and the condensate along $x0$ would break it to the rotations of $x5, x6, x7$. In form Q this pattern is present for any couplings. In form C the self-interaction itself breaks $SO(4)$ to $SO(3)_T$ explicitly: the three generators that the condensate would have to break spontaneously are not symmetries of the model, and a scalar coupling of the sign that binds matter repels in the pair along the hidden axis (Section 6.4).
5. *A scalar coupling of the sign that binds matter on 7+1 binds less.* Matter needs two to three times the strength, which moves it further outside the range in which the no-sea approximation is controlled (Section 6.6).

What does not distinguish the forms. The reduction to four effective couplings and the Fierz matrix (Sections 6.1 and 6.2); the interaction between particles of the same half, hence everything about matter that occupies one half only; the pairs along $x5, x6, x7$; every result that does not involve the self-interaction; the critical strength at which the vacuum of a massless field becomes unstable, and the magnitude of the mass that it acquires.

What could decide. Three kinds of consideration could decide between the forms; none of them is settled by the calculations of this project.

1. *A principle.* If the model is required to be the restriction of a covariant theory on the full 4+4 spacetime, form C follows. If it is required to be the theory of eight equivalent internal states, form Q follows.
2. *A consistent quantization of the covariant Lagrangian itself.* If a quantization of Eq. (5.1) with its indefinite sign existed (with an indefinite metric on the space of states and a rule for the physical states), it would dictate the form of the interaction, and it would give neither form C nor form Q in general (Section 5.5). No such quantization is constructed in this project.
3. *An observable consequence.* The two forms differ in the internal states of self-bound matter (four or eight), in the pair that the self-interaction favors, and in the symmetry of the order parameter. A hypothesis that ties one of these to an observation would select a form.

The author's decision of November 13, 2026 is form C. Form Q remains recorded completely, in this document and in the program, as the alternative.

Part 7. Solutions of the linear equation, mode by mode

This part solves Eqs. (4.10) and (4.11), the equations of the weighted field without the self-interaction. They describe a field so dilute that the cubic term is negligible, and they supply the single-particle states from which a treatment of the interacting field starts.

7.1 Three kinds of modes

A mode with $k_5 = k_6 = k_7 = 0$ obeys the flat equation; its solutions are the plane waves of flat spacetime, with the energies $\pm E_\perp$, eight of each sign. A mode with some $k_t \neq 0$ behaves in the same way as long as its physical wavenumber is negligible, $k_d \ll E_\perp$. But k_d grows without bound, and every such mode sooner or later reaches $k_d \gg E_\perp$. The moment at which $k_d = E_\perp$ is its **crossing**; on 4+4 it is the turning value (1.19) of the classical particle, with the wavenumbers in place of the momenta.

The solution is exact when **one rate governs**: when only one of the k_t differs from zero, or when the three rates are equal. In both cases the matrix α^d or γ^d does not depend on the time, and $k_d \propto e^{\text{rate}_t\, x4}$ with the rate that governs. The natural variable is then

$$u_d = \frac{k_d(x4)}{\text{rate}_t}, \qquad \frac{d}{dx4} = \text{rate}_t\, u_d\, \frac{d}{du_d}. \tag{7.1}$$

The crossing is at $u_d = E_\perp/\text{rate}_t$, and this ratio of the energy to the rate is the one parameter on which the results depend.

7.2 The 7+1 spacetime: the deflation creates particles

Exact solution. In $\mathsf{h} = \mathsf{h}_\perp + k_d\,\alpha^d$ the two constant matrices satisfy $\mathsf{h}_\perp^2 = E_\perp^2$ and $(\alpha^d)^2 = 1$, and they anticommute. Let w be one of the eight eigenvectors of $\mathsf{h}_\perp$ with the eigenvalue $+E_\perp$; then $\alpha^d w$ is an eigenvector with the eigenvalue $-E_\perp$. Let Z_+ and Z_- denote two cylinder functions of the same kind (Bessel functions of the first or of the second kind, or Hankel functions) of the orders $\frac{1}{2} + iE_\perp/\text{rate}_t$ and $-\frac{1}{2} + iE_\perp/\text{rate}_t$. With the recurrence relations of the cylinder functions one verifies by substitution that

$$\hat{\psi} = \sqrt{u_d}\left[Z_+(u_d)\,w + i\,Z_-(u_d)\,\alpha^d w\right] \tag{7.2}$$

solves Eq. (4.10) for each of the eight w. Two independent kinds of cylinder function give all sixteen solutions. The program verifies Eq. (7.2) with the Bessel functions of an arbitrary-precision library, to a relative residual of 10^{-15}.

Creation. At early times, $u_d \to 0$, the solution that has positive energy, $\propto e^{-iE_\perp x4}\,w$, is Eq. (7.2) with Z_+ the Bessel function of the first kind of the order $-\frac{1}{2} - iE_\perp/\text{rate}_t$ and Z_- minus that of the order $+\frac{1}{2} - iE_\perp/\text{rate}_t$, a pair that obeys the same recurrence relations. At late times, $u_d \to \infty$, the solution of positive energy is Eq. (7.2) with the Hankel functions of the second kind, which behave as e^{-iu_d}, and the solution of negative energy is the one with the Hankel functions of the first kind. The connection formula of the Bessel functions expresses the first solution through the other two, and the weight of the negative-energy part divided by that of the positive-energy part is $e^{-2\pi E_\perp/\text{rate}_t}$. A mode that starts empty therefore ends with

$$n_{\text{cr}} = \frac{1}{e^{2\pi E_\perp/\text{rate}_t} + 1} \tag{7.3}$$

particles in each of its eight internal states, and with as many antiparticles. This is the Fermi–Dirac distribution in the energy $E_\perp$ at the temperature

$$T_{\text{defl}} = \frac{\text{rate}_t}{2\pi}. \tag{7.4}$$

For equal rates it is the temperature of de Sitter space found by Gibbons and Hawking; the creation of particles by a time-dependent metric is the phenomenon discovered by Parker (references at the end).

Numerical confirmation (checks `71d-M1` to `71d-M3`). The numerical solution of Eq. (4.10) for all sixteen components confirms the law. For $E_\perp/\text{g55}$ between 0.25 and 3 the created number agrees with Eq. (7.3) to a relative deviation of at most 3.2×10^{-6}, while the numbers themselves range from 0.17 to 6.5×10^{-9}; the eight internal states are created equally. With three equal rates the result depends neither on the direction of the wavenumber nor on the constants const_t. With unequal rates no exact solution is known. If the second physical wavenumber is still below $e^{-3}E_\perp$ when the first reaches $E_\perp$, the mean created number is Eq. (7.3) with the rate of the coordinate that becomes energetic first, to within 1.2 per cent. If two directions become energetic together, the eight internal states are created unequally, and the result is known only numerically.




Figure 3: Creation of particles on the deflating 7+1 spacetime. (a) Created quanta per state, multiplied by $e^{2\pi\,\mathrm{ratio}}$, against the ratio of the energy to the rate. Fermions follow the Fermi–Dirac law (7.3). For comparison the panel also shows the quanta of a scalar field on the same spacetime, which follow the Bose–Einstein law, Eq. (7.3) with -1 in place of $+1$ and with the energy of the scalar mode, at the same temperature. Lines: the exact laws; symbols: computed. (b) History of two modes, with $E_\perp/\mathrm{g55} = 1$ and 0.5: the number of particles in the mode against the physical wavenumber $k_d/E_\perp$, which grows with the time. After the crossing at $k_d = E_\perp$ the number settles at the value of Eq. (7.3) (dotted). (c) Two deflating directions, $\mathrm{g55} = 1$ and $\mathrm{g66} = 2$, 1.2 or 0.5: mean number of created particles per state against the delay between the two crossings; the dotted lines are Eq. (7.3) with each of the two rates.

Meaning. Each mode with $k_t \neq 0$ is populated once, at its crossing. Its particles then gain energy without bound, as $(E_\perp^2 + k_d^2)^{1/2}$, like the classical particle of Section 1.7. Modes with $k_t = 0$ stay empty. The stress of the created particles must eventually change the metric; that reaction has not been computed, and the results hold for as long as it can be neglected. The instantaneous energy of a quantum in a mode with $k_t \neq 0$ is at least $|k_t|/b_t$, which grows as $e^{\mathrm{rate}_t\, x4}$: if the deflating coordinates are periodic, the smallest wavenumber that differs from zero is fixed, and the sector of fields that do not depend on $x5, x6, x7$ is separated from the rest by a gap that grows exponentially.

7.3 The 4+4 spacetime: the deflation extinguishes the modes that depend on the additional times

Exact solution. In $K = K_\perp + i\,k_d\,\gamma^4\gamma^d$ the first matrix is anti-Hermitian with $K_\perp^2 = -E_\perp^2$, the matrix $i\gamma^4\gamma^d$ is Hermitian with the square one, and the two anticommute. Let w be one of the eight eigenvectors of $K_\perp$ with the eigenvalue $+iE_\perp$; then $i\gamma^4\gamma^d w$ is an eigenvector with the eigenvalue $-iE_\perp$ and is orthogonal to w. The equation is solved by the modified Bessel functions that decay at large argument, of the orders $\frac{1}{2} \mp iE_\perp/\mathrm{rate}_t$. Such a function is written here with its order as a subscript and with its argument; the letter K without subscript and argument is the matrix of Eq. (4.11). With the recurrence relations of these functions one verifies by substitution that

$$\hat{\psi} = \sqrt{u_d}\left[K_{\frac{1}{2}-iE_\perp/\mathrm{rate}_t}(u_d)\,w - K_{\frac{1}{2}+iE_\perp/\mathrm{rate}_t}(u_d)\,i\gamma^4\gamma^d w\right] \tag{7.5}$$

solves Eq. (4.11) for each of the eight w; the relative residual, computed with the Bessel functions of an arbitrary-precision library, is 2×10^{-15}. The other eight solutions are built in the same way from the modified Bessel functions that *grow* as e^{+u_d}.

Standing waves and their extinction. Since u_d itself grows exponentially with the time, a solution of the second kind grows as the exponential of an exponential. Such a solution is not admissible; it is the counterpart of the growing exponential under a potential barrier. Admissible are the eight

solutions (7.5), which stay bounded. Because w and $i\gamma^4\gamma^d w$ are orthogonal and the two functions in Eq. (7.5) are complex conjugates of each other,

$$|\hat{\psi}|^2 = 2\,u_d\,\left|K_{\frac{1}{2}+iE_\perp/\mathrm{rate}_t}(u_d)\right|^2\,|w|^2. \tag{7.6}$$

At early times, $u_d \to 0$, $|\hat{\psi}|^2 \to \pi\,|w|^2/\cosh(\pi E_\perp/\mathrm{rate}_t)$, a constant: the solution is a combination of $e^{+iE_\perp x4}\,w$ and $e^{-iE_\perp x4}\,i\gamma^4\gamma^d w$ with coefficients of *equal* magnitude. A bounded solution is a **standing wave** in the ordinary time, not a traveling wave, and its conserved charge $\hat{\psi}^\dagger W_{1644}\hat{\psi}$ vanishes. At late times, $u_d \to \infty$, $|\hat{\psi}|^2 \to \pi\,|w|^2\,e^{-2u_d}$: the mode dies out as the exponential of an exponential of the time.

The law of the extinction. In the semiclassical (WKB) approximation the bounded solution falls beyond the turning value, at which $k_d = E_\perp$, as $e^{-S_{\mathrm{WKB}}}$ with

$$S_{\mathrm{WKB}} = \int_{x4_{\mathrm{turn}}}^{x4} \sqrt{k_d^2 - E_\perp^2}\;dx4' = \frac{1}{\mathrm{rate}_t}\left[\sqrt{k_d^2 - E_\perp^2} - E_\perp \arccos\frac{E_\perp}{k_d}\right], \tag{7.7}$$

which approaches $(k_d - \frac{\pi}{2}E_\perp)/\mathrm{rate}_t$ for $k_d \gg E_\perp$. The exact solution fixes the constant that the approximation leaves open:

$$\ln\frac{|\hat{\psi}|_{\mathrm{early}}}{|\hat{\psi}|_{\mathrm{late}}} - S_{\mathrm{WKB}} \;\longrightarrow\; \tfrac{1}{2}\ln 2 - \tfrac{1}{2}\ln\left(1 + e^{-2\pi E_\perp/\mathrm{rate}_t}\right). \tag{7.8}$$

Numerical confirmation (checks `44d-M1` to `44d-M4`). Beyond the turning value exactly eight of the sixteen solutions stay bounded; followed backwards to early times they contain parts of positive and of negative frequency of equal weight, to 2×10^{-14}. Where one rate governs, the computed magnitude equals Eq. (7.6) over the range $0.02 < u_d < 12$ to a relative deviation of 3.6×10^{-12}, and Eq. (7.8) holds to 3.6×10^{-4} at $S_{\mathrm{WKB}} = 30$. With three unequal rates no exact solution is known; the bounded solutions fall as $e^{-S_{\mathrm{WKB}}}$ times a factor that depends on the internal state, with a spread of 0.40 in its logarithm at $S_{\mathrm{WKB}} = 30$.




Figure 4: Extinction on the deflating 4+4 spacetime. (a) A bounded solution: $|\hat{\psi}|^2$ against the physical wavenumber $k_d/E_\perp$, which grows with the time; computed (line) and the exact form (7.6) (symbols). It is constant long before the crossing at $k_d = E_\perp$ and falls after it. (b) Five cases, with one, two or three wavenumbers k_t different from zero and with equal and unequal rates, on a logarithmic scale. (c) The bounded modes of a scalar field on the same spacetime, which are extinguished in the same way; lines: exact, symbols: computed.

Meaning. On the flat 4+4 spacetime a field that depends on the additional times evolves non-unitarily in the ordinary time; this is the principal difficulty of four time-like dimensions, and on flat spacetime it can only be met by imposing that the fields do not depend on those times. On the deflating 4+4 spacetime every admissible mode that depends on a deflating time is a standing wave that is extinguished after its crossing. What remains at late times is the sector of fields that do not depend on $x5, x6, x7$, in which the evolution is unitary: the quantized sector of Part 5. *The deflation of the three additional times removes, dynamically, the sector that causes the difficulty.* The limits of this conclusion: the eight growing solutions are discarded because they are unbounded, which is a condition of admissibility and is not derived from a principle; and the theory still has no positive conserved charge for modes that depend on the deflating times.

7.4 The numerical method

The equations of this part are linear, with coefficients that grow exponentially, and an integrator with a fixed step fails on them. The program uses a commutator-free Magnus method of the fourth order: over each step the solution is advanced by the product of two matrix exponentials of linear combinations of the coefficient matrix at two Gauss points. Because the square of h or of K is a multiple of the unit matrix at any one time, each exponential is evaluated in closed form; on 7+1 every step is exactly unitary. The length of a step is chosen so that neither the phase accumulated in it nor the relative change of the coefficients exceeds a fixed bound. On 7+1 the number of particles is read off by projection on the eigenstates of the Hamiltonian corrected to the first order in the rate at which its direction changes, which removes the leading spurious oscillation of the particle number at finite time. The independent script, which uses an explicit Runge–Kutta method of the eighth order and the Bessel functions of another library, reproduces Eq. (7.3) to 2.2×10^{-5} and confirms Eq. (7.5) and, to the accuracy of the semiclassical form at the exponent that it reaches, Eq. (7.8).

Part 8. Which symmetries survive the deflation

Rotations and boosts. The flat spacetimes have 28 rotations and boosts, one for each pair of coordinates. On the deflating spacetime (checks `44d-S1`, `71d-S1`; table of Section 1.5):

- the six rotations of $x0, x1, x2, x3$ survive for any rates;
- the rotation of two deflating coordinates survives if and only if their rates are equal;
- nothing that involves $x4$ survives; on 7+1 these are the seven boosts, among them the three boosts of special relativity along $x1, x2, x3$;
- nothing that mixes a flat coordinate with a deflating coordinate survives.

For fields that do not depend on $x5, x6, x7$ the flat equations hold for the weighted field (Section 4.3), and their solutions are related by the ordinary boosts as on flat spacetime. What singles out the preferred state of rest for such fields is only the dependence of the couplings (4.6) on the time.

The internal symmetry of fields in ordinary space. For a field that depends on $x1, x2, x3, x4$ only, the linear equation of the weighted field contains four matrices: $\alpha^1, \alpha^2, \alpha^3$ and β. The matrices that commute with all four generate transformations of the sixteen components that leave the equation unchanged: an **internal symmetry**. Its algebra has the dimension 8 and the structure $su(2)_A \oplus su(2)_B \oplus u(1)_F \oplus u(1)_E$, on both spacetimes (check `def-S3`):

- six generators are the products $\gamma^a\gamma^b$ of the six pairs $a < b$ taken from the four axes $x0, x5, x6, x7$; they form the two $su(2)$;
- $u(1)_F$ is the phase symmetry of Section 4.5 (fermion number);
- $u(1)_E$ is a second phase symmetry, generated by a traceless Hermitian matrix P with $P^2 = 1$, proportional to $\gamma^0\gamma^5\gamma^6\gamma^7$; it distinguishes the states that are doublets of $su(2)_A$ from those that are doublets of $su(2)_B$: at a given momentum the eight states of positive energy are four

with $P = +1$, which are doublets of one $su(2)$ and do not transform under the other, and four with $P = -1$, for which the two algebras exchange their roles.

Because the weighted field of this sector obeys the flat equations, this symmetry is not changed by the deflation. But on flat spacetime the six products $\gamma^a\gamma^b$ are rotations of the four extra axes, symmetries of the spacetime; on the deflating spacetime 0, 1 or 3 of them are still symmetries of the spacetime, for three different, two equal or three equal rates (checks `44d-S2`, `71d-S2`; last three columns of the table of Section 1.5). The others remain exact symmetries of the sector *without* being symmetries of the spacetime. On 7+1 the self-interaction respects the six rotations and $u(1)_F$ for all couplings, and $u(1)_E$ only for $g_1^{\text{eff}} = g_7^{\text{eff}} = 0$. On 4+4, in form C (the definition), every self-interaction that does not vanish breaks the three generators that mix $x0$ with $x5, x6, x7$ and breaks $u(1)_E$; it keeps the three rotations of $x5, x6, x7$ among themselves and $u(1)_F$. In form Q the 4+4 model behaves as the 7+1 model (Section 6.5).

Discrete symmetries. The scale factors are real and depend on $x4$ only. Reflections of any coordinate other than $x4$, and charge conjugation on 7+1 (Section 4.5), survive. The reversal of the ordinary time does not: it turns deflation into inflation.

Part 9. What has been established, and what has not

Established (each item with the checks named in the text; exact unless marked numerical there):

1. The geometry of the two deflating spacetimes: Christoffel symbols, curvature, Einstein tensor, the 14 to 20 Killing vectors; the spacetimes are not vacuum solutions of Einstein's equations; with equal rates they solve Einstein–Gauss–Bonnet gravity for $\Lambda_8 = 3\,\mathrm{g55}^2$, $\alpha_{\text{GB}} = -1/(4\,\mathrm{g55}^2)$, and that solution is unstable (numerical).
2. The spin connection $\Omega_t = \frac{1}{2}\mathrm{rate}_t\, b_t\, \gamma^t\gamma^4$, the same formula on both spacetimes, and $\gamma^\mu\Omega_\mu = -\frac{1}{2}\mathrm{gsum}\,\gamma^4$.
3. The Lagrangians with kinetic term, mass term and self-interaction, Eqs. (3.2), (3.5), (3.6), (3.8), and on 4+4 the positive Lagrangian (5.3) with the two forms (5.5) and (5.9) of the self-interaction; every bilinear in them is real, and every invariant I_r is invariant under the transformations of the frame.
4. The field equations (4.1) and (4.2), and their forms (4.3), (4.4), (4.7), (4.8), (5.6), (5.10), (5.13).
5. The weighted field obeys the flat equations with every quartic coupling divided by V_d and with the mass unchanged.
6. Of the nine couplings only four combinations are independent, Eq. (6.5). Uniform matter on 7+1 feels the two Hartree–Fock couplings (6.7); on 4+4 it feels four couplings (6.9), the same four combinations in both forms with the roles of the total densities and of the differences between the halves exchanged. The pair couplings of the 28 pairs at rest, Eq. (6.14) and the table of Section 6.4. The conditions for chiral symmetry (6.16) and for the internal symmetry of fields in ordinary space; in particular, every self-interaction of form C breaks $SO(4)\times U(1)_E$ to $SO(3)_T$.
7. On 4+4 the covariant Lagrangian is indefinite, Eq. (5.1); form C of the self-interaction is form Q with W_{1644} inserted in every bilinear, Eqs. (5.11) and (5.12): two particles of the same half of the field interact identically in the two forms, a particle of each half with opposite signs.
8. Numerically, in the mean-field approximation of Section 6.6 and for the sets A and B of couplings: matter of form C is polarized and binds with one half of the field filled, at two to three times the strength at which matter of form Q binds with both halves equally filled; a massless field acquires in vacuum masses of opposite sign in the two halves in form C and of equal sign in form Q, at the same critical strength.
9. The exact solutions (7.2) and (7.5) of the linear equations when one rate governs; the law of creation (7.3) on 7+1; the extinction (7.6) to (7.8) on 4+4 (confirmed numerically).

Not established:

1. *What sustains the deflation.* The metric is a hypothesis. It is not a solution of Einstein's equations without matter, and the one theory in which it is a vacuum solution does not hold it stable.
2. *Which form of the self-interaction is the right one for the 4+4 model* (Sections 5.6 and 6.7). Form C is the definition, by the author's decision; form Q gives different interactions between the two halves of the field, and nothing in the calculations decides between them. 2a. *States with coherence between the two halves, and other couplings.* The density matrix (6.8) is diagonal in W_{1644}. Because the interaction connects the halves, states with $\langle \psi_+^\dagger \cdots \psi_- \rangle \neq 0$ are possible in principle in both forms; whether one of them has a lower energy than the states of Section 6.6 has not been examined. The numbers of Section 6.6 are for the sets A and B; the exact formulas hold for all couplings.
3. *The values of the couplings g_r and of the mass m.* Nothing in this document fixes them.
4. *Solutions of the nonlinear equations.* Part 7 solves the linear equations. With the self-interaction the field equations are solved in the project only in mean-field approximations for uniform matter and for pairs; an exact solution is not known.
5. *Modes with unequal rates and several $k_t \neq 0$* are known only numerically.
6. *The reaction of the created particles on the metric* (7+1), and *a principle that excludes the growing solutions* (4+4).

Appendix A. Complete output of the programs

The reports are printed as the programs wrote them (a line that is too long to be typeset is continued on the following lines, broken at spaces). In them the rates are called `g55`, `g66`, `g77`; `rate_t`, `b_t`, `const5` and so on are the quantities of the same names in the text; `gamma1644[A]` and `gamma1671[A]` are the matrices γ^A_{1644} and γ^A_{1671}; `Cm1644` is C^-_{1644}; `psihat` is the weighted field $\widehat{\psi}$; `gs` is the list of the signs gs_{AA} and `c` the constant c_γ of Section 0.2; `G_0` and `G_1` are G_S^{HF} and G_V^{HF}, `G_S0` and `G_V0` (also written `G_Sc`, `G_Vc`) are G_S^{c} and G_V^{c}; `G_Stot`, `G_Vtot`, `G_Sdif`, `G_Vdif`, `G_Ssame`, `G_Vsame`, `G_Sopp`, `G_Vopp` are the couplings of Eqs. (6.9) and (6.11); `eps_K` is sgn_K; `F` is the Fierz matrix `F`; `rho` and `rho_S` (in the report of the command `forms`: `n` and `n_S`) are the densities n and n_S, `s_+-` the scalar densities $n_{S\pm}$; `M`, `M_+`, `M_-` are the masses m^*, m^*_+, m^*_-; `W` is W_{1644}; "the covariant form" is form C and "the quantized-sector form" is form Q. "The main project" means the part of PROJECT spacetimes that treats the flat 4+4 and 7+1 spacetimes: its program is used as a library, the matrices of the frame are its matrices, and several checks compare with its data files. "The author's document" means his document on Kohn–Sham matter and gravity on the 7+1 spacetime.

A.1 Geometry, coordinate matrices, spin connection, Dirac operators (`deflation_cjt_ks_c matrices`)

```
PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C: checks of the geometry, of
 ↪  the Dirac matrices, of the spin connection and of the Dirac operators of the deflating
 ↪  spacetimes (44d-..: 4+4 with deflating x5, x6, x7; 71d-..: 7+1 with deflating x5, x6, x7)
generated by: deflation_cjt_ks_c (Rust crate spacetimes-deflation-cjt-ks-c, version 0.1.0;
 ↪  PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C); command:
 ↪  deflation_cjt_ks_c matrices
47 of 47 checks passed
```

[PASS] 44d-G1 (exact) metric of the deflating 4+4 spacetime in the author's sign convention (space-like +, time-like -): g = diag(gs_00, .., gs_44, gs_55 b_5^2, gs_66 b_6^2, gs_77 b_7^2) with the metric signs gs = diag[1, 1, 1, 1, -1, -1, -1, -1] and the scale factors b_5 = const5 exp(-g55 x4), b_6 = const6 exp(-g66 x4), b_7 = const7 exp(-g77 x4), that is g_55 = -const5^2 exp(-2 g55 x4) and likewise g_66, g_77 (the author's formulas for this spacetime, on which x5, x6, x7 are time-like; g55, g66, g77 > 0 are the three rates, and in statements with a general index t = 5, 6, 7 the rate of xt is written rate_t, gsum = g55 + g66 + g77). Verified: g_tt = -b_t^2 and d g_tt/dx4 = -2 rate_t g_tt; the frame e^A = (dx0, .., dx4, b_5 dx5, b_6 dx6, b_7 dx7) is orthonormal. All exact geometry checks use five rational parameter sets: (g55, g66, g77) = (1, 1, 1) at a point where (b_5, b_6, b_7) = (1, 1, 1); (g55, g66, g77) = (1, 2, 3) at a point where (b_5, b_6, b_7) = (1/2, 1/3, 1/5); (g55, g66, g77) = (1/2, 3/4, 5/3) at a point where (b_5, b_6, b_7) = (2, 1/3, 5/7); (g55, g66, g77) = (2, 2, 5) at a point where (b_5, b_6, b_7) = (3/2, 3/2, 1/4); (g55, g66, g77) = (7/3, 7/3, 7/3) at a point where (b_5, b_6, b_7) = (1/2, 3, 5/4)

[PASS] 44d-G2 (exact) the Christoffel symbols computed from their definition are Gamma^t_(4t) = Gamma^t_(t4) = -rate_t and Gamma^4_(tt) = -gs_tt rate_t b_t^2 = + rate_t b_t^2 (t = 5, 6, 7); all others vanish

[PASS] 44d-G3 (exact) Cartan's structure equations hold with the spin-connection coefficients of the tetrad postulate: the torsion d e^A + omega^A_B ^ e^B vanishes, and d omega^A_B + omega^A_C ^ omega^C_B equals the Riemann tensor computed from the Christoffel symbols (two independent routes to the curvature)

[PASS] 44d-G4 (exact) Riemann tensor in the orthonormal frame: R^t_(4t4) = -rate_t^2, R^4_(t4t) = gs_tt rate_t^2 = -rate_t^2, R^t_(sts) = gs_ss rate_t rate_s = -rate_t rate_s (t != s in {5, 6, 7}), and the components that follow from antisymmetry; all other components vanish. No component involves x0, x1, x2 or x3: the four space coordinates x0, x1, x2, x3 stay flat

[PASS] 44d-G5 (exact) with gsq = g55^2 + g66^2 + g77^2 and gsum = g55 + g66 + g77, in the author's sign convention of the metric: Ricci tensor R^4_4 = gsq, R^t_t = rate_t gsum, R^a_a = 0 (a = 0..3); curvature scalar R = gsq + gsum^2; Einstein tensor G^4_4 = -(g55 g66 + g55 g77 + g66 g77), G^t_t = rate_t gsum - (gsq + gsum^2)/2, G^a_a = -(gsq + gsum^2)/2; all are constants (they do not depend on x4), the mixed components are the same on the 4+4 and on the 7+1 spacetime, and the covariant divergence of the Einstein tensor vanishes (contracted Bianchi identity)

[PASS] 44d-G6 (exact) Killing vectors (the Lie derivative of the metric along each vanishes identically): 7 translations (x0, x1, x2, x3, x5, x6, x7), 6 rotations of (x0, x1, x2, x3), and the scaling d/dx4 + sum_t rate_t xt d/dxt, 14 in all; one more for every pair of equal rates (a rotation of the two coordinates), and three more if all three rates are equal (20 in all). Numbers found for the five parameter sets: [20, 14, 14, 15, 20]

[PASS] 44d-G7 (exact) if the three rates are equal, g55 = g66 = g77, the part (x4, x5, x6, x7) of the metric has constant curvature, R_(rho sigma mu nu) = +g55^2 (g_(rho mu) g_(sigma nu) - g_(rho nu) g_(sigma mu)): the four time-like coordinates carry minus the metric of hyperbolic space H^4 of radius 1/g55, written in horospherical coordinates (the upper-half-space form)

[PASS] 44d-G10 (exact) Lovelock gravity, in the author's sign convention of the metric (space-like components positive), with the action R - 2 Lambda_8 + alpha_GB (R^2 - 4 R_(mu nu) R^(mu nu) + R_(mu nu rho sigma) R^(mu nu rho sigma)) and the field equations G^mu_nu + Lambda_8 delta^mu_nu + alpha_GB Lanczos^mu_nu = 0: the curvature acts diagonally on pairs of frame directions; the Lanczos tensor of the Gauss-Bonnet term is Lanczos^a_a = -4 g55 g66 g77 gsum (a = 0..3), Lanczos^4_4 = Lanczos^t_t = 0; the Lovelock tensor of third order vanishes identically. For three equal rates (2 of the five parameter sets) the deflating spacetime solves the vacuum field equations with Lambda_8 = 3 g55^2 and alpha_GB = -1/(4 g55^2), that is 4 alpha_GB Lambda_8 = -3: the deflation needs no matter as its source in Einstein-Gauss-Bonnet gravity. For rates that are not all equal no choice of Lambda_8 and alpha_GB satisfies the components 4, 5, 6, 7

[PASS] 44d-G9 (numerical) geodesic with m = 1, momentum P_1 = 0.5 along x1 and P_5 = 0.3 along the deflating time x5 (g55 = 1, const5 = 0.8), started at x4 = -2 with increasing x4: P_1, P_5 and the norm are conserved to 1.7e-14; the path reaches the largest value x4 = 1.09240103 (predicted turning value ln(E_perp const5/|P_5|)/g55 with E_perp = sqrt(m^2 + P_1^2) = 1.09240103) after the affine parameter 3.38543615 (closed form 3.38543615) and then returns to smaller x4: a particle with momentum along a deflating time cannot pass the turning value

[PASS] 44d-C1 (exact) the coordinate matrices gamma^mu = e_A^mu gamma1644[A] of the deflating 4+4 spacetime (gamma^t = gamma1644[t]/b_t for t = 5, 6, 7; the other five unchanged) are real and satisfy gamma^mu.gamma^nu + gamma^nu.gamma^mu = 2 c g^(mu nu) 1 for all 64 pairs, with c = +1 and the inverse metric g^(mu nu) of the author's sign convention (space-like +, time-like -; g^(tt) = -1/b_t^2). The squares of the matrices are therefore equal to the signs of the metric. Verified for five rational parameter sets: (g55, g66, g77) = (1, 1, 1) at a point where (b_5, b_6, b_7) = (1, 1, 1); (g55, g66, g77) = (1, 2, 3) at a point where (b_5, b_6, b_7) = (1/2, 1/3, 1/5); (g55, g66, g77) = (1/2, 3/4, 5/3) at a point where (b_5, b_6, b_7) = (2, 1/3, 5/7); (g55, g66, g77) = (2, 2, 5) at a point where (b_5, b_6, b_7) = (3/2, 3/2, 1/4); (g55, g66, g77) = (7/3, 7/3, 7/3) at a point where (b_5, b_6, b_7) = (1/2, 3, 5/4)

[PASS] 44d-C2 (exact) the spin-connection coefficients computed from the tetrad postulate omega_mu^A_B = e^A_nu (d_mu e_B^nu + Gamma^nu_(mu lam) e_B^lam), with the Christoffel symbols obtained from the metric, and with the first frame index lowered by the metric signs gs, omega_(mu A B) = gs_AA omega_mu^A_B, are antisymmetric, omega_(mu A B) = -omega_(mu B A), and the only non-zero ones are omega_(t,t4) = -omega_(t,4t) = -gs_tt rate_t b_t = + rate_t b_t (t = 5, 6, 7); same five parameter sets

[PASS] 44d-C3 (exact) Omega_mu := (c/4) omega_(mu A B) gamma1644[A].gamma1644[B] with c = +1 equals +(1/2) rate_t b_t gamma1644[t].gamma1644[4] for mu = t in {5, 6, 7} and vanishes for mu = 0..4 (the same formula on both spacetimes, because c gs_tt = -1 on both): the spin connection is an element of the Lie algebra spanned by the scaled commutators Sigma^(t4)

[PASS] 44d-C4 (exact) d_mu gamma^nu + Gamma^nu_(mu lam) gamma^lam + [Omega_mu, gamma^nu] = 0 for all 64 pairs (mu, nu), with d_4 gamma^t = +rate_t gamma^t: the coordinate matrices are covariantly constant

[PASS] 44d-C5 (exact) sum_mu gamma^mu.Omega_mu = -(1/2) (g55 + g66 + g77) gamma1644[4]: in the Dirac operator the spin connection amounts to the replacement d_4 -> d_4 - gsum/2, which the weighted field psihat = sqrt(b_5 b_6 b_7) psi removes

[PASS] 44d-C6 (exact) d_mu Omega_nu - d_nu Omega_mu + [Omega_mu, Omega_nu] = (c/4) R_(A B mu nu) gamma1644[A].gamma1644[B] for all 64 pairs (mu, nu), with c = +1 and with the Riemann tensor that metric.rs computes from the metric, its first index lowered by the metric (d_4 Omega_t = -rate_t Omega_t): the curvature of the spinor derivative is the Riemann tensor

[PASS] 44d-C7 (exact) (1/2) sum_(mu nu) gamma^mu.gamma^nu.(d_mu Omega_nu - d_nu Omega_mu + [Omega_mu, Omega_nu]) = -(c R/4) 1 = - (R/4) 1 with the curvature scalar R = g55^2 + g66^2 + g77^2 + gsum^2 of the author's sign convention: the square of the Dirac operator differs from c times the wave operator of the spinor derivative by this term

[PASS] 44d-C8 (exact) transpose(Omega_mu).Cm1644 + Cm1644.Omega_mu = 0 (Cm1644 = gamma1644[0].gamma1644[1].gamma1644[2].gamma1644[3]): the adjoint field psi^dagger Cm1644 is differentiated with -Omega_mu on its right, as the conserved current requires

[PASS] 44d-D1 (exact) in momentum space (d_mu -> i k_mu) the Dirac operator of the weighted field at fixed x4, D = i gamma^mu k_mu with the coordinate matrices, satisfies (D + m)(D - m) = -(g^(mu nu) k_mu k_nu + m^2) 1 with g^(mu nu) k_mu k_nu = k_0^2 + k_1^2 + k_2^2 + k_3^2 - k_4^2 - sum_t k_t^2/b_t^2: the instantaneous mass shell is k_4^2 + sum_t k_t^2/b_t(x4)^2 = |k|^2 + m^2. Verified for five rational parameter sets and four rational momenta each

[PASS] 44d-D2 (exact) the generator of the evolution of the weighted field in x4, K(x4) = -m gamma1644[4] + i sum_a k_a gamma1644[4].gamma1644[a] + i sum_t (k_t/b_t(x4)) gamma1644[4].gamma1644[t], satisfies K^2 = (sum_t k_t^2/b_t^2 - m^2 - |k|^2) 1 and K + K^dagger = 2 i sum_t (k_t/b_t) gamma1644[4].gamma1644[t]

[PASS] 44d-D3 (exact) dK/dx4 = i sum_t rate_t (k_t/b_t) gamma1644[4].gamma1644[t] is a Hermitian matrix whose square is sum_t (rate_t k_t/b_t)^2 times the unit matrix; the weighted field obeys the second-order equation d_4^2 psihat = (K^2 + dK/dx4) psihat, a Klein-Gordon equation with a term that depends on the spin

[PASS] 44d-D4 (exact) W = i Cm1644.gamma1644[4] is Hermitian with W^2 = 1 and trace 0, and W.K + K^dagger.W = 0 for every momentum and all k_5, k_6, k_7: the covariant charge psihat^dagger W psihat of a mode is conserved on the deflating spacetime, and it is not positive

[PASS] 44d-D5 (exact) K is anti-Hermitian (the evolution in x4 is unitary and psihat^dagger psihat is conserved) if and only if k_5 = k_6 = k_7 = 0, that is, if the field does not depend on the deflating times

[PASS] 44d-D6 (exact) with gamma1644[A] = [[0, taubar1644[A]], [tau1644[A], 0]] the operator gamma^mu d_mu of the weighted field has the blocks [[0, sum_A taubar1644[A] e_A^mu d_mu], [sum_A tau1644[A] e_A^mu d_mu, 0]], e_A^mu d_mu = (d_0, .., d_4, d_5/b_5, d_6/b_6, d_7/b_7): the equation is the pair taubar^A e_A^mu d_mu psihat^R = m psihat^L, tau^A e_A^mu d_mu psihat^L = m psihat^R of real 8-component equations

```
[PASS] 44d-D7  (exact) worked example: m = 3, k = 0, (k_5, k_6, k_7) = (1, 0, 0) at the value of
↪  x4 where b_5 = 1/5 (so k_5/b_5 = 5 > m): K^2 = +16 1, and (4 + K)/8 and (4 - K)/8 are
↪  complementary projectors of trace 8: at this x4 eight independent solutions grow as exp(+4 x4)
↪  and eight decay as exp(-4 x4) to first order. A mode that depends on a deflating time is
↪  oscillatory only while k_t/b_t(x4) < sqrt(m^2 + |k|^2)
[PASS] 71d-G1  (exact) metric of the deflating 7+1 spacetime in the author's sign convention
↪  (space-like +, time-like -): g = diag(gs_00, .., gs_44, gs_55 b_5^2, gs_66 b_6^2, gs_77 b_7^2)
↪  with the metric signs gs = diag[1, 1, 1, 1, -1, 1, 1, 1] and the scale factors b_5 = const5
↪  exp(-g55 x4), b_6 = const6 exp(-g66 x4), b_7 = const7 exp(-g77 x4), that is g_55 = +const5^2
↪  exp(-2 g55 x4) and likewise g_66, g_77 (the author's formulas for this spacetime, on which x5,
↪  x6, x7 are space-like; g55, g66, g77 > 0 are the three rates, and in statements with a general
↪  index t = 5, 6, 7 the rate of xt is written rate_t, gsum = g55 + g66 + g77). Verified: g_tt =
↪  +b_t^2 and d g_tt/dx4 = -2 rate_t g_tt; the frame e^A = (dx0, .., dx4, b_5 dx5, b_6 dx6, b_7
↪  dx7) is orthonormal. All exact geometry checks use five rational parameter sets: (g55, g66,
↪  g77) = (1, 1, 1) at a point where (b_5, b_6, b_7) = (1, 1, 1); (g55, g66, g77) = (1, 2, 3) at
↪  a point where (b_5, b_6, b_7) = (1/2, 1/3, 1/5); (g55, g66, g77) = (1/2, 3/4, 5/3) at a point
↪  where (b_5, b_6, b_7) = (2, 1/3, 5/7); (g55, g66, g77) = (2, 2, 5) at a point where (b_5, b_6,
↪  b_7) = (3/2, 3/2, 1/4); (g55, g66, g77) = (7/3, 7/3, 7/3) at a point where (b_5, b_6, b_7) =
↪  (1/2, 3, 5/4)
[PASS] 71d-G2  (exact) the Christoffel symbols computed from their definition are Gamma^t_(4t) =
↪  Gamma^t_(t4) = -rate_t and Gamma^4_(tt) = -gs_tt rate_t b_t^2 = - rate_t b_t^2 (t = 5, 6, 7);
↪  all others vanish
[PASS] 71d-G3  (exact) Cartan's structure equations hold with the spin-connection coefficients of
↪  the tetrad postulate: the torsion d e^A + omega^A_B ^ e^B vanishes, and d omega^A_B +
↪  omega^A_C ^ omega^C_B equals the Riemann tensor computed from the Christoffel symbols (two
↪  independent routes to the curvature)
[PASS] 71d-G4  (exact) Riemann tensor in the orthonormal frame: R^t_(4t4) = -rate_t^2, R^4_(t4t) =
↪  gs_tt rate_t^2 = +rate_t^2, R^t_(sts) = gs_ss rate_t rate_s = +rate_t rate_s (t != s in {5, 6,
↪  7}), and the components that follow from antisymmetry; all other components vanish. No
↪  component involves x0, x1, x2 or x3: the four space coordinates x0, x1, x2, x3 stay flat
[PASS] 71d-G5  (exact) with gsq = g55^2 + g66^2 + g77^2 and gsum = g55 + g66 + g77, in the
↪  author's sign convention of the metric: Ricci tensor R^4_4 = gsq, R^t_t = rate_t gsum, R^a_a =
↪  0 (a = 0..3); curvature scalar R = gsq + gsum^2; Einstein tensor G^4_4 = -(g55 g66 + g55 g77 +
↪  g66 g77), G^t_t = rate_t gsum - (gsq + gsum^2)/2, G^a_a = -(gsq + gsum^2)/2; all are constants
↪  (they do not depend on x4), the mixed components are the same on the 4+4 and on the 7+1
↪  spacetime, and the covariant divergence of the Einstein tensor vanishes (contracted Bianchi
↪  identity)
[PASS] 71d-G6  (exact) Killing vectors (the Lie derivative of the metric along each vanishes
↪  identically): 7 translations (x0, x1, x2, x3, x5, x6, x7), 6 rotations of (x0, x1, x2, x3),
↪  and the scaling d/dx4 + sum_t rate_t xt d/dxt, 14 in all; one more for every pair of equal
↪  rates (a rotation of the two coordinates), and three more if all three rates are equal (20 in
↪  all). Numbers found for the five parameter sets: [20, 14, 14, 15, 20]
[PASS] 71d-G7  (exact) if the three rates are equal, g55 = g66 = g77, the part (x4, x5, x6, x7) of
↪  the metric has constant curvature, R_(rho sigma mu nu) = +g55^2 (g_(rho mu) g_(sigma nu) -
↪  g_(rho nu) g_(sigma mu)) in the author's sign convention: it is de Sitter space of radius
↪  1/g55 in the contracting half of its flat slicing
[PASS] 71d-G10  (exact) Lovelock gravity, in the author's sign convention of the metric
↪  (space-like components positive), with the action R - 2 Lambda_8 + alpha_GB (R^2 - 4 R_(mu nu)
↪  R^(mu nu) + R_(mu nu rho sigma) R^(mu nu rho sigma)) and the field equations G^mu_nu +
↪  Lambda_8 delta^mu_nu + alpha_GB Lanczos^mu_nu = 0: the curvature acts diagonally on pairs of
↪  frame directions; the Lanczos tensor of the Gauss-Bonnet term is Lanczos^a_a = -4 g55 g66 g77
↪  gsum (a = 0..3), Lanczos^4_4 = Lanczos^t_t = 0; the Lovelock tensor of third order vanishes
↪  identically. For three equal rates (2 of the five parameter sets) the deflating spacetime
↪  solves the vacuum field equations with Lambda_8 = 3 g55^2 and alpha_GB = -1/(4 g55^2), that is
↪  4 alpha_GB Lambda_8 = -3: the deflation needs no matter as its source in Einstein-Gauss-Bonnet
↪  gravity. For rates that are not all equal no choice of Lambda_8 and alpha_GB satisfies the
↪  components 4, 5, 6, 7
```

[PASS] 71d-G8 (numerical) geodesic of a particle of mass m = 1 with momentum P_1 = 0.5 along x1 and P_5 = 0.3 along the deflating coordinate x5 (g55 = 1, const5 = 0.8), integrated from x4 = -2 to x4 = 4 by the Runge-Kutta method: P_1, P_5 and the norm are conserved to 2.0e-10; the affine parameter (the proper time, since m = 1) that elapses is 3.3375401839, the closed form gives 3.3375401839; the proper time to x4 = infinity is finite, 3.386358: a particle that moves along a deflating coordinate leaves the coordinate patch after a finite proper time, with unbounded energy
[PASS] 71d-C1 (exact) the coordinate matrices gamma^mu = e_A^mu gamma1671[A] of the deflating 7+1 spacetime (gamma^t = gamma1671[t]/b_t for t = 5, 6, 7; the other five unchanged) are real and satisfy gamma^mu.gamma^nu + gamma^nu.gamma^mu = 2 c g^(mu nu) 1 for all 64 pairs, with c = -1 and the inverse metric g^(mu nu) of the author's sign convention (space-like +, time-like -; g^(tt) = +1/b_t^2). The squares of the matrices are therefore opposite to the signs of the metric. Verified for five rational parameter sets: (g55, g66, g77) = (1, 1, 1) at a point where (b_5, b_6, b_7) = (1, 1, 1); (g55, g66, g77) = (1, 2, 3) at a point where (b_5, b_6, b_7) = (1/2, 1/3, 1/5); (g55, g66, g77) = (1/2, 3/4, 5/3) at a point where (b_5, b_6, b_7) = (2, 1/3, 5/7); (g55, g66, g77) = (2, 2, 5) at a point where (b_5, b_6, b_7) = (3/2, 3/2, 1/4); (g55, g66, g77) = (7/3, 7/3, 7/3) at a point where (b_5, b_6, b_7) = (1/2, 3, 5/4)
[PASS] 71d-C2 (exact) the spin-connection coefficients computed from the tetrad postulate omega_mu^A_B = e^A_nu (d_mu e_B^nu + Gamma^nu_(mu lam) e_B^lam), with the Christoffel symbols obtained from the metric, and with the first frame index lowered by the metric signs gs, omega_(mu A B) = gs_AA omega_mu^A_B, are antisymmetric, omega_(mu A B) = -omega_(mu B A), and the only non-zero ones are omega_(t,t4) = -omega_(t,4t) = -gs_tt rate_t b_t = - rate_t b_t (t = 5, 6, 7); same five parameter sets
[PASS] 71d-C3 (exact) Omega_mu := (c/4) omega_(mu A B) gamma1671[A].gamma1671[B] with c = -1 equals +(1/2) rate_t b_t gamma1671[t].gamma1671[4] for mu = t in {5, 6, 7} and vanishes for mu = 0..4 (the same formula on both spacetimes, because c gs_tt = -1 on both): the spin connection is an element of the Lie algebra spanned by the scaled commutators Sigma^(t4)
[PASS] 71d-C4 (exact) d_mu gamma^nu + Gamma^nu_(mu lam) gamma^lam + [Omega_mu, gamma^nu] = 0 for all 64 pairs (mu, nu), with d_4 gamma^t = +rate_t gamma^t: the coordinate matrices are covariantly constant
[PASS] 71d-C5 (exact) sum_mu gamma^mu.Omega_mu = -(1/2) (g55 + g66 + g77) gamma1671[4]: in the Dirac operator the spin connection amounts to the replacement d_4 -> d_4 - gsum/2, which the weighted field psihat = sqrt(b_5 b_6 b_7) psi removes
[PASS] 71d-C6 (exact) d_mu Omega_nu - d_nu Omega_mu + [Omega_mu, Omega_nu] = (c/4) R_(A B mu nu) gamma1671[A].gamma1671[B] for all 64 pairs (mu, nu), with c = -1 and with the Riemann tensor that metric.rs computes from the metric, its first index lowered by the metric (d_4 Omega_t = -rate_t Omega_t): the curvature of the spinor derivative is the Riemann tensor
[PASS] 71d-C7 (exact) (1/2) sum_(mu nu) gamma^mu.gamma^nu.(d_mu Omega_nu - d_nu Omega_mu + [Omega_mu, Omega_nu]) = -(c R/4) 1 = + (R/4) 1 with the curvature scalar R = g55^2 + g66^2 + g77^2 + gsum^2 of the author's sign convention: the square of the Dirac operator differs from c times the wave operator of the spinor derivative by this term
[PASS] 71d-C8 (exact) transpose(Omega_mu).gamma1671[4] + gamma1671[4].Omega_mu = 0: the adjoint field psi^dagger gamma1671[4] is differentiated with -Omega_mu on its right, as the conserved current requires
[PASS] 71d-D1 (exact) in momentum space (d_mu -> -i k_mu) the Dirac operator of the weighted field at fixed x4 is gamma^mu k_mu - m with the coordinate matrices, and (gamma^mu k_mu + m)(gamma^nu k_nu - m) = -(g^(mu nu) k_mu k_nu + m^2) 1 with g^(mu nu) k_mu k_nu = -k_4^2 + k_0^2 + k_1^2 + k_2^2 + k_3^2 + sum_t k_t^2/b_t^2 (author's sign convention of the metric): the instantaneous mass shell is k_4^2 = |p|^2 + sum_t k_t^2/b_t(x4)^2 + m^2. Verified for five rational parameter sets and four rational momenta each
[PASS] 71d-D2 (exact) the Hamiltonian of the weighted field, h(x4) = m beta1671 + sum_j p_j alpha1671^j + sum_t (k_t/b_t(x4)) alpha1671^t (beta1671 = gamma1671[4], alpha1671^A = gamma1671[4].gamma1671[A]), is real and symmetric for every momentum and every x4, and h^2 = (m^2 + |p|^2 + sum_t k_t^2/b_t^2) 1: the evolution in x4 is unitary and psihat^dagger psihat is conserved, but the energy of a mode with k_t != 0 grows as exp(rate_t x4)
[PASS] 71d-D3 (exact) dh/dx4 = sum_t rate_t (k_t/b_t) alpha1671^t is real symmetric and anticommutes with beta1671: the Hamiltonian changes with x4 only through the momenta along the deflating coordinates, which is what creates pairs
[PASS] 71d-D5 (exact) omega1671 commutes with every alpha1671^A and anticommutes with beta1671; hence the charge-conjugate field omega1671 psi^* solves the Dirac equation on the deflating spacetime whenever psi does (the scale factors are real)

```
[PASS] 71d-D6  (exact) with the chiral 16x8 matrices E^L = (1_8, -i 1_8)^T/sqrt(2), E^R = (-i 1_8,
 ↪  1_8)^T/sqrt(2) (columns of 8 + 8 components) and the real 8x8 matrices tau1671^j of the
 ↪  author's notebook, (E^L)^dagger h E^L = -i sum_j (p_j/e_j) tau1671^j, (E^R)^dagger h E^R = +i
 ↪  sum_j (p_j/e_j) tau1671^j and (E^L)^dagger h E^R = (E^R)^dagger h E^L = m, where p_j/e_j =
 ↪  (p_0, p_1, p_2, p_3, k_5/b_5, k_6/b_6, k_7/b_7): the equation of the weighted field is the
 ↪  pair (i d_4 + tau^j e_j^mu d_mu) phihat^L = m phihat^R, (i d_4 - tau^j e_j^mu d_mu) phihat^R =
 ↪  m phihat^L of 8-component equations
[PASS] 71d-D7  (exact) worked example: m = 3, p = 0, (k_5, k_6, k_7) = (2, 0, 0) at the value of
 ↪  x4 where b_5 = 1/2 (so k_5/b_5 = 4): h^2 = 25 1, and (5 + h)/10, (5 - h)/10 are complementary
 ↪  projectors of trace 8: eight states of energy +5 and eight of energy -5 at this x4; one
 ↪  e-folding of the deflation later the energy is sqrt(9 + 16 e^2) = 11.28
[PASS] ALL  (exact) every check above passed
```

A.2 The matrices of the frame (`deflation_cjt_ks_c matrices`)

```
PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C: the checks of the real
 ↪  16x16 Dirac matrices of the main project, executed again (these matrices are the matrices of
 ↪  the orthonormal frame of the deflating spacetimes); identifiers 44-.. refer to gamma1644,
 ↪  71-.. to gamma1671
generated by: deflation_cjt_ks_c (Rust crate spacetimes-deflation-cjt-ks-c, version 0.1.0;
 ↪  PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C); command:
 ↪  deflation_cjt_ks_c matrices
106 of 106 checks passed

[PASS] 44-R1  (exact) every entry of gamma1644[A], A = 0..7, is one of the integers -1, 0, +1;
 ↪  hence the eight matrices are real
[PASS] 44-R2  (exact) every gamma1644[A] is a signed permutation matrix (one entry +1 or -1 in
 ↪  each row and in each column)
[PASS] 44-C1  (exact) gamma1644[A].gamma1644[B] + gamma1644[B].gamma1644[A] = 2 eta1644^(AB) 1
 ↪  for all 64 pairs (A, B), with eta1644 = diag[1, 1, 1, 1, -1, -1, -1, -1]
[PASS] 44-C2  (exact) transpose(gamma1644[A]) = eta1644^(AA) gamma1644[A]: matrices with square
 ↪  +1 are symmetric, matrices with square -1 are antisymmetric; hence transpose(gamma1644[A]) =
 ↪  inverse(gamma1644[A])
[PASS] 44-C3  (exact) the 256 ordered products gamma1644[A1]...gamma1644[Ak] (A1 < ... < Ak, k =
 ↪  0..8) are linearly independent: they span all real 16x16 matrices, so the representation is
 ↪  irreducible over the real numbers
[PASS] 44-C4  (exact) every one of the 255 non-empty products has trace 0, and the square of every
 ↪  product is +1 or -1
[PASS] 44-L1  (exact) Sigma1644^(AB) := (1/4)[gamma1644[A], gamma1644[B]] equals (1/2)
 ↪  gamma1644[A].gamma1644[B] for A != B, and Sigma1644^(AB) = -Sigma1644^(BA)
[PASS] 44-L2  (exact) the 28 matrices Sigma1644^(AB), A < B, are linearly independent
[PASS] 44-L3  (exact) [Sigma1644^(AB), gamma1644[C]] = eta1644^(BC) gamma1644[A] - eta1644^(AC)
 ↪  gamma1644[B] for all A, B, C: the Sigma1644^(AB) rotate the eight matrices as the components
 ↪  of a vector
[PASS] 44-L4  (exact) [Sigma1644^(AB), Sigma1644^(CD)] = eta1644^(BC) Sigma1644^(AD) -
 ↪  eta1644^(AC) Sigma1644^(BD) - eta1644^(BD) Sigma1644^(AC) + eta1644^(AD) Sigma1644^(BC) for
 ↪  all 4096 index combinations: the commutation relations of the Lie algebra so(p,q) of the
 ↪  metric eta1644
[PASS] 44-L5  (exact) the structure constants are integers and the Killing form is diagonal in the
 ↪  basis Sigma1644^(AB): B(Sigma1644^(AB), Sigma1644^(AB)) = -12 eta1644^(AA) eta1644^(BB); it
 ↪  has 16 positive and 12 negative eigenvalues (no zero eigenvalue: the algebra is semisimple)
[PASS] 44-L6  (exact) 12 generators are antisymmetric matrices (compact: they generate rotations)
 ↪  and 16 are symmetric (non-compact: they generate boosts); these are the numbers of the real
 ↪  form so(4,4), which has 12 compact and 16 non-compact generators
[PASS] 44-G1  (exact) each gamma1644[A] is invertible with inverse eta1644^(AA) gamma1644[A], and
 ↪  -gamma1644[A].gamma1644[B].inverse(gamma1644[A]) = gamma1644[B] for B != A, = -gamma1644[A]
 ↪  for B = A: gamma1644[A] acts on the vector index as the reflection that reverses the axis xA
[PASS] 44-G2  (numerical) exp(theta Sigma1644^(AB)) = cos(theta/2) 1 + sin(theta/2)
 ↪  gamma1644[A].gamma1644[B] when (gamma1644[A].gamma1644[B])^2 = -1 (rotation), and =
 ↪  cosh(theta/2) 1 + sinh(theta/2) gamma1644[A].gamma1644[B] when (gamma1644[A].gamma1644[B])^2
 ↪  = +1 (boost), for the 28 planes and random theta (largest deviation 2.3e-14)
```

```
[PASS] 44-G3  (numerical) for 20 random elements S = exp(sum_(A<B) theta_AB Sigma1644^(AB))
↪  (theta_AB random real numbers):  S.gamma1644[C].inverse(S) = sum_D Lambda^C_D gamma1644[D]
↪  with a real 8x8 matrix Lambda that preserves eta1644 (Lambda.eta1644.transpose(Lambda) =
↪  eta1644) and has determinant +1 (largest deviations 2.0e-13, 2.2e-15, 2.7e-15)
[PASS] 44-N1  (exact) the eight matrices rebuilt from the definitions equal, entry by entry, the
↪  eight matrices T16[0..7] printed by Mathematica in 4+4.nb (cell 314, Out[211])
[PASS] 44-N2  (exact) the rebuilt volume element equals T16[8] printed in 4+4.nb (cell 136,
↪  Out[99])
[PASS] 44-N3  (exact) the metric printed in 4+4.nb (cell 13, Out[9]) is eta4488 =
↪  diag(1,1,1,1,-1,-1,-1,-1) (PROJECT name: eta1644)
[PASS] 44-B1  (exact) the 8x8 matrix sigma (= sigma88) of 4+4.nb, PROJECT name sigma88: sigma88 =
↪  tau[1].tau[2].tau[3] = tau[4].tau[5].tau[6].tau[7], sigma88.sigma88 = 1,
↪  tau[1].tau[2]...tau[7] = 1, and tau[7] = diag(-1_4, +1_4) (4+4.nb Out[72], Out[89], Out[87],
↪  Out[63])
[PASS] 44-B2  (exact) tau[A].taubar[B] + tau[B].taubar[A] = 2 eta1644^(AB) 1 and taubar[A].tau[B]
↪  + taubar[B].tau[A] = 2 eta1644^(AB) 1 (8x8), A, B = 0..7 (4+4.nb Out[77])
[PASS] 44-B3  (exact) sigma88.taubar[A] = transpose(sigma88.tau[A]) for A = 0..7 (4+4.nb Out[76])
[PASS] 44-B4  (exact) T16[A] = [[0, taubar[A]], [tau[A], 0]] in 8x8 blocks, with tau[0] =
↪  taubar[0] = 1.  PROJECT names of these objects of 4+4.nb: gamma1644[A] := T16[A], tau1644[A]
↪  := tau[A], taubar1644[A] := taubar[A]
[PASS] 44-V1  (exact) omega1644 := gamma1644[0]...gamma1644[7] (= T16[8]) = diag(-1_8, +1_8):
↪  real, diagonal, symmetric, with square +1
[PASS] 44-V2  (exact) omega1644 anticommutes with every gamma1644[A] and commutes with every
↪  Sigma1644^(AB)
[PASS] 44-P1  (exact) PL1644 := (1 - omega1644)/2 = diag(1_8, 0_8) and PR1644 := (1 + omega1644)/2
↪  = diag(0_8, 1_8) are real matrices with integer entries
[PASS] 44-P2  (exact) PL1644 + PR1644 = 1, PL1644.PL1644 = PL1644, PR1644.PR1644 = PR1644,
↪  PL1644.PR1644 = PR1644.PL1644 = 0, trace PL1644 = trace PR1644 = 8 (4+4.nb Out[114])
[PASS] 44-P3  (exact) the rebuilt projectors PL1644 and PR1644 equal the matrices P_L and P_R
↪  printed in 4+4.nb (cells 163 and 165, Out[112] and Out[113])
[PASS] 44-P4  (exact) gamma1644[A].PL1644 = PR1644.gamma1644[A] and gamma1644[A].PR1644 =
↪  PL1644.gamma1644[A]: every Dirac matrix maps left-handed spinors to right-handed ones and
↪  conversely
[PASS] 44-P5  (exact) PL1644 and PR1644 commute with every Sigma1644^(AB): chirality is invariant
↪  under Spin(4,4)
[PASS] 44-S1  (exact) Cm1644 := gamma1644[0].gamma1644[1].gamma1644[2].gamma1644[3] (the matrix
↪  sigma16 of 4+4.nb) = diag(-sigma88, sigma88), symmetric, Cm1644.Cm1644 = 1 (4+4.nb Out[106])
[PASS] 44-S2  (exact) Cp1644 := gamma1644[4].gamma1644[5].gamma1644[6].gamma1644[7] (the matrix
↪  S44 of 7+1.nb) = diag(sigma88, sigma88), symmetric, Cp1644.Cp1644 = 1, and omega1644 =
↪  Cm1644.Cp1644
[PASS] 44-S3  (exact) Cm1644.gamma1644[A].Cm1644 = -transpose(gamma1644[A]) and
↪  Cp1644.gamma1644[A].Cp1644 = +transpose(gamma1644[A]) for A = 0..7
[PASS] 44-S4  (exact) Cm1644.gamma1644[A] is antisymmetric for A = 0..7 (4+4.nb Out[105]) and
↪  Cp1644.gamma1644[A] is symmetric
[PASS] 44-S5  (exact) gamma1644[4] (the matrix JT4 = T16[4] of 4+4.nb) is a complex structure:
↪  gamma1644[4].gamma1644[4] = -1, and transpose(gamma1644[4]).Cm1644.gamma1644[4] = Cm1644
[PASS] 71-R1  (exact) every entry of gamma1671[A], A = 0..7, is one of the integers -1, 0, +1;
↪  hence the eight matrices are real
[PASS] 71-R2  (exact) every gamma1671[A] is a signed permutation matrix (one entry +1 or -1 in
↪  each row and in each column)
[PASS] 71-C1  (exact) gamma1671[A].gamma1671[B] + gamma1671[B].gamma1671[A] = 2 eta1671^(AB) 1
↪  for all 64 pairs (A, B), with eta1671 = diag[-1, -1, -1, -1, 1, -1, -1, -1]
[PASS] 71-C2  (exact) transpose(gamma1671[A]) = eta1671^(AA) gamma1671[A]: matrices with square
↪  +1 are symmetric, matrices with square -1 are antisymmetric; hence transpose(gamma1671[A]) =
↪  inverse(gamma1671[A])
[PASS] 71-C3  (exact) the 256 ordered products gamma1671[A1]...gamma1671[Ak] (A1 < ... < Ak, k =
↪  0..8) are linearly independent: they span all real 16x16 matrices, so the representation is
↪  irreducible over the real numbers
[PASS] 71-C4  (exact) every one of the 255 non-empty products has trace 0, and the square of every
↪  product is +1 or -1
[PASS] 71-L1  (exact) Sigma1671^(AB) := (1/4)[gamma1671[A], gamma1671[B]] equals (1/2)
↪  gamma1671[A].gamma1671[B] for A != B, and Sigma1671^(AB) = -Sigma1671^(BA)
```

```
[PASS] 71-L2  (exact) the 28 matrices Sigma1671^(AB), A < B, are linearly independent
[PASS] 71-L3  (exact) [Sigma1671^(AB), gamma1671[C]] = eta1671^(BC) gamma1671[A] - eta1671^(AC)
↪  gamma1671[B] for all A, B, C: the Sigma1671^(AB) rotate the eight matrices as the components
↪  of a vector
[PASS] 71-L4  (exact) [Sigma1671^(AB), Sigma1671^(CD)] = eta1671^(BC) Sigma1671^(AD) -
↪  eta1671^(AC) Sigma1671^(BD) - eta1671^(BD) Sigma1671^(AC) + eta1671^(AD) Sigma1671^(BC) for
↪  all 4096 index combinations: the commutation relations of the Lie algebra so(p,q) of the
↪  metric eta1671
[PASS] 71-L5  (exact) the structure constants are integers and the Killing form is diagonal in the
↪  basis Sigma1671^(AB): B(Sigma1671^(AB), Sigma1671^(AB)) = -12 eta1671^(AA) eta1671^(BB); it
↪  has 7 positive and 21 negative eigenvalues (no zero eigenvalue: the algebra is semisimple)
[PASS] 71-L6  (exact) 21 generators are antisymmetric matrices (compact: they generate rotations)
↪  and 7 are symmetric (non-compact: they generate boosts); these are the numbers of the real
↪  form so(7,1), which has 21 compact and 7 non-compact generators
[PASS] 71-G1  (exact) each gamma1671[A] is invertible with inverse eta1671^(AA) gamma1671[A], and
↪  -gamma1671[A].gamma1671[B].inverse(gamma1671[A]) = gamma1671[B] for B != A, = -gamma1671[A]
↪  for B = A: gamma1671[A] acts on the vector index as the reflection that reverses the axis xA
[PASS] 71-G2  (numerical) exp(theta Sigma1671^(AB)) = cos(theta/2) 1 + sin(theta/2)
↪  gamma1671[A].gamma1671[B] when (gamma1671[A].gamma1671[B])^2 = -1 (rotation), and =
↪  cosh(theta/2) 1 + sinh(theta/2) gamma1671[A].gamma1671[B] when (gamma1671[A].gamma1671[B])^2
↪  = +1 (boost), for the 28 planes and random theta (largest deviation 7.1e-15)
[PASS] 71-G3  (numerical) for 20 random elements S = exp(sum_(A<B) theta_AB Sigma1671^(AB))
↪  (theta_AB random real numbers):  S.gamma1671[C].inverse(S) = sum_D Lambda^C_D gamma1671[D]
↪  with a real 8x8 matrix Lambda that preserves eta1671 (Lambda.eta1671.transpose(Lambda) =
↪  eta1671) and has determinant +1 (largest deviations 5.0e-14, 2.0e-15, 2.0e-15)
[PASS] 71-N1  (exact) the eight matrices rebuilt from the definitions equal, entry by entry, the
↪  eight matrices Gamma17[0..7] printed by Mathematica in 7+1.nb (cell 204, Out[178])
[PASS] 71-N2  (exact) PROJECT matrices: gamma1671[A] = Gamma17up[swap04(A)] with swap04 = (0 4)
↪  and Gamma17up[A] = eta17^(AA) Gamma17[A] (7+1.nb section 5.3); explicitly gamma1671[4] =
↪  Gamma17[0] (time-like), gamma1671[0] = -Gamma17[4], gamma1671[j] = -Gamma17[j] for j = 1, 2,
↪  3, 5, 6, 7.  (gamma1671 is NOT the complex set Gamma71 of 7+1.nb section 4.3, and eta1671 is
↪  NOT the matrix eta71 of 7+1.nb)
[PASS] 71-N3  (exact) in the labelling of 7+1.nb: Gamma17[A].Gamma17[B] + Gamma17[B].Gamma17[A] =
↪  2 eta17^(AB) 1 with eta17 = diag(+1,-1,-1,-1,-1,-1,-1,-1) (7+1.nb Out[173])
[PASS] 71-B1  (exact) tauO[A] = tau[A] (A = 1, 2, 3), tauO[A] = tau[A].sigma (A = 4..7): real,
↪  antisymmetric, tauO[A].tauO[A] = -1, mutually anticommuting, tauO[1]...tauO[7] = 1 (7+1.nb
↪  section 3.1)
[PASS] 71-B2  (exact) block form: Gamma17[0] = [[0, 1], [1, 0]] and Gamma17[i] = [[tauO[i], 0],
↪  [0, tauObar[i]]] with tauObar[i] = transpose(tauO[i]) = -tauO[i], i = 1..7 (7+1.nb Out[175])
[PASS] 71-B3  (exact) built from the 4+4 matrices only: Gamma17[0] = T16[0], Gamma17[i] =
↪  T16[0].T16[i] (i = 1, 2, 3), Gamma17[k] = T16[0].T16[k].S44 (k = 4..7) (7+1.nb Out[176])
[PASS] 71-B4  (exact) in the labels of this project: gamma1671[4] = gamma1644[0]; gamma1671[j] =
↪  -gamma1644[0].gamma1644[j] (j = 1, 2, 3); gamma1671[0] =
↪  +gamma1644[0].gamma1644[5].gamma1644[6].gamma1644[7]; gamma1671[5] =
↪  -gamma1644[0].gamma1644[4].gamma1644[6].gamma1644[7]; gamma1671[6] =
↪  +gamma1644[0].gamma1644[4].gamma1644[5].gamma1644[7]; gamma1671[7] =
↪  -gamma1644[0].gamma1644[4].gamma1644[5].gamma1644[6].  The time-like matrix of 7+1 IS the
↪  matrix of the hidden axis of 4+4; the other seven are products of two or of four of the 4+4
↪  matrices and therefore not linear combinations of them
[PASS] 71-V1  (exact) omega1671 := gamma1671[0]...gamma1671[7] = Gamma17[8] = [[0, -1], [1, 0]] in
↪  8x8 blocks (the omega17 of 7+1.nb: the seven sign changes of Gamma17up and the odd permutation
↪  swap04 compensate): real, antisymmetric, with square -1
[PASS] 71-V2  (exact) omega1671 anticommutes with every gamma1671[A] and commutes with every
↪  Sigma1671^(AB)
[PASS] 71-V3  (exact) the linear system X.gamma1671[A] + gamma1671[A].X = 0 (A = 0..7) for a real
↪  16x16 matrix X has rank 255, and omega1671 solves it (71-V2): the real matrices that
↪  anticommute with all eight gamma1671[A] are exactly the multiples of omega1671; since
↪  omega1671.omega1671 = -1, no real matrix with square +1 anticommutes with all of them: there
↪  is NO real chirality matrix in 7+1
[PASS] 71-P1  (exact) chi1671 := i omega1671 = i Gamma17[8] (= chi17 of 7+1.nb) is Hermitian,
↪  chi1671.chi1671 = 1, anticommutes with every gamma1671[A]
```

```
[PASS] 71-P2  (exact) PL1671 := (1 - chi1671)/2 and PR1671 := (1 + chi1671)/2: PL1671 + PR1671 =
↪  1, PL1671.PL1671 = PL1671, PR1671.PR1671 = PR1671, PL1671.PR1671 = 0, Hermitian, trace 8 each;
↪  they are complex (not real) matrices
[PASS] 71-P3  (exact) gamma1671[A].PL1671 = PR1671.gamma1671[A] for A = 0..7, and PL1671, PR1671
↪  commute with every Sigma1671^(AB)
[PASS] 71-P4  (exact) in 8x8 blocks chi1671 = [[0, -i], [i, 0]], 2 PL1671 = [[1, i], [-i, 1]] and
↪  2 PR1671 = [[1, -i], [i, 1]]; these are chi17, PL17 and PR17 of 7+1.nb (sections 4.6 and 7.10)
[PASS] 71-P5  (exact) with s := 1 - i Gamma17[0] (s.s^dagger = 2, so Uc1671 := s/sqrt(2) is
↪  unitary): s.chi1671 = -T16[8].s, i.e. Uc1671.chi1671.inverse(Uc1671) = -omega1644.  Hence
↪  Uc1671.PL1671.inverse(Uc1671) = PR1644 and Uc1671.PR1671.inverse(Uc1671) = PL1644: the
↪  complex chiral projectors of 7+1 are unitarily equivalent to the real projectors of 4+4.nb,
↪  with the names L and R exchanged.  (7+1.nb defines chi17 = +i Gamma17[8] for its real matrices
↪  and chi71 = -i Gamma71[8] for its complex matrices Gamma71; the PROJECT matrix chi1671 equals
↪  chi17, NOT chi71)
[PASS] 71-S1  (exact) Dirac adjoint: transpose(gamma1671[A]) =
↪  gamma1671[4].gamma1671[A].gamma1671[4] for A = 0..7 (gamma1671[4] is the time-like matrix; it
↪  is symmetric and squares to +1)
[PASS] 44-K1  (exact) 4+4: the matrices Cm1644.gamma1644[A1]...gamma1644[Ar] (A1 < ... < Ar) of
↪  rank r = 0..8 are all symmetric (+) or all antisymmetric (-): signs [1, -1, -1, 1, 1, -1, -1,
↪  1, 1]; the matrices Cp1644.gamma1644[A1]...gamma1644[Ar]: signs [1, 1, -1, -1, 1, 1, -1, -1,
↪  1]; in both cases exactly 120 of the 256 matrices are antisymmetric
[PASS] 44-K2  (exact) 4+4: the two Lorentz-scalar pair matrices Cm1644 (rank 0) and
↪  Cm1644.omega1644 (rank 8; equal to Cp1644 up to sign) are SYMMETRIC: the Lorentz-scalar
↪  bilinears dirac1644^T Cm1644 dirac1644 and dirac1644^T Cp1644 dirac1644 of one anticommuting
↪  16-component field vanish identically
[PASS] 71-K0  (exact) 7+1: Cp1671 := gamma1671[4] satisfies Cp1671.gamma1671[A] =
↪  +transpose(gamma1671[A]).Cp1671 and Cm1671 := gamma1671[4].omega1671 satisfies
↪  Cm1671.gamma1671[A] = -transpose(gamma1671[A]).Cm1671; both are symmetric matrices
[PASS] 71-K1  (exact) 7+1: signs (+ symmetric, - antisymmetric) of the matrices
↪  Cm1671.gamma1671[A1]...gamma1671[Ar] (A1 < ... < Ar) by rank r = 0..8: [1, -1, -1, 1, 1, -1,
↪  -1, 1, 1]; of Cp1671.gamma1671[A1]...gamma1671[Ar]: [1, 1, -1, -1, 1, 1, -1, -1, 1]; exactly
↪  120 of the 256 matrices are antisymmetric in both cases
[PASS] 71-K2  (exact) 7+1: the two Lorentz-scalar pair matrices Cm1671 (rank 0) and
↪  Cm1671.omega1671 = -Cp1671 (rank 8) are SYMMETRIC: the Lorentz-scalar bilinears dirac1671^T
↪  Cm1671 dirac1671 and dirac1671^T Cp1671 dirac1671 of one anticommuting 16-component field
↪  vanish identically
[PASS] 44-K3  (exact) 4+4: for all A != B in 1..7, transpose(Sigma1644^(AB)).Cm1644.gamma1644[0] +
↪  Cm1644.gamma1644[0].Sigma1644^(AB) = 0, while the same expression with A = 0 is not zero: the
↪  bilinear DeltaV1644^0 = dirac1644^T Cm1644 gamma1644^0 dirac1644 is INVARIANT under the 21
↪  generators of the pseudo-rotations of the seven coordinates x1..x7 (the group Spin(3,4), which
↪  contains the Lorentz group of x1 x2 x3 x4), and only the 7 generators that involve the hidden
↪  axis x0 change it
[PASS] 71-K3  (exact) 7+1: for all A != B in 1..7, transpose(Sigma1671^(AB)).Cm1671.gamma1671[0] +
↪  Cm1671.gamma1671[0].Sigma1671^(AB) = 0, while the same expression with A = 0 is not zero: the
↪  bilinear DeltaV1671^0 = dirac1671^T Cm1671 gamma1671^0 dirac1671 is INVARIANT under the 21
↪  generators of the Lorentz group of the seven coordinates x1..x7 (the group Spin(6,1), which
↪  contains the Lorentz group of x1 x2 x3 x4), and only the 7 generators that involve the hidden
↪  axis x0 change it
[PASS] 44-D1  (exact) (gamma1644^A k_A)^2 = (eta1644^(AB) k_A k_B) 1 for 40 random integer vectors
↪  k: the Dirac operator gamma1644^A d_A squares to the wave operator eta1644^(AB) d_A d_B = d0^2
↪  + d1^2 + d2^2 + d3^2 - d4^2 - d5^2 - d6^2 - d7^2
[PASS] 44-D2  (exact) (i gamma1644.k + m1644)(i gamma1644.k - m1644) = -(k.k + m1644^2) 1 for 40
↪  random integer (k, m1644): every component of a solution of (gamma1644^A d_A - m1644)
↪  dirac1644 = 0 obeys the Klein-Gordon equation (eta1644^(AB) d_A d_B - m1644^2) dirac1644 = 0
[PASS] 44-D3  (exact) worked example k = (0,2,2,4; 5,0,0,0), m1644 = 1 (on the mass shell:
↪  k4^2+k5^2+k6^2+k7^2 = k0^2+k1^2+k2^2+k3^2 + m1644^2): Pi1644 := (i gamma1644.k + m1644)/(2
↪  m1644) satisfies Pi1644.Pi1644 = Pi1644, trace Pi1644 = 8 and (i gamma1644.k - m1644).Pi1644 =
↪  0, so the plane-wave solutions u exp(i k.x) form an 8-dimensional space, u = Pi1644 w
```

```
[PASS] 44-D4  (exact) PL1644.gamma1644[A].PR1644 = [[0, taubar1644[A]], [0, 0]],
↪  PR1644.gamma1644[A].PL1644 = [[0, 0], [tau1644[A], 0]], PL1644.gamma1644[A].PL1644 =
↪  PR1644.gamma1644[A].PR1644 = 0: with dirac1644 = (psiL1644, psiR1644) the 16-component
↪  equation is the pair  taubar1644^A d_A psiR1644 = m1644 psiL1644,  tau1644^A d_A psiL1644 =
↪  m1644 psiR1644  of 8-component equations
[PASS] 44-D5  (exact) (tau1644^A k_A)(taubar1644^B k_B) = (taubar1644^A k_A)(tau1644^B k_B) =
↪  (eta1644^(AB) k_A k_B) 1_8 for 40 random integer k: each 8-component field obeys the
↪  Klein-Gordon equation
[PASS] 44-D6  (exact) gamma1644[4].gamma1644[a] (a = 0..3) are real symmetric with square +1 (and
↪  so are alpha1644^a := -gamma1644[4].gamma1644[a], the matrices of the Hamiltonian h1644 =
↪  alpha1644^a k_a + beta1644 m1644, beta1644 := -i gamma1644[4]); gamma1644[4].gamma1644[t] (t
↪  = 5, 6, 7) are real antisymmetric with square -1; gamma1644[4] is real antisymmetric with
↪  square -1
[PASS] 44-D7  (exact) x4-evolution d_4 dirac1644 = K1644 dirac1644 with K1644 = -m1644 gamma1644^4
↪  + i k_a gamma1644^4 gamma1644^a + i kappa_t gamma1644^4 gamma1644^t (a = 0..3 space, t = 5, 6,
↪  7 extra times): K1644.K1644 = -(m1644^2 + |k|^2 - |kappa|^2) 1 for 40 random integer (k,
↪  kappa, m1644): the frequencies in x4 are +-sqrt(m1644^2 + |k|^2 - |kappa|^2)
[PASS] 44-D8  (exact) K1644 + K1644^dagger = 2 i kappa_t gamma1644^4 gamma1644^t: K1644 is
↪  anti-Hermitian (so that h1644 := i K1644 is Hermitian and the x4-evolution preserves
↪  dirac1644^dagger dirac1644) if and only if kappa = 0, i.e. if the field does not depend on the
↪  extra times x5, x6, x7
[PASS] 44-D9  (exact) example of the instability: m1644 = 3, k = 0, kappa = (5, 0, 0) gives
↪  K1644.K1644 = +16, (K1644 - 4)(K1644 + 4) = 0 with trace K1644 = 0: K1644 has the real
↪  eigenvalues +4 and -4, eight times each, so eight solutions grow like exp(+4 x4)
[PASS] 44-D10  (exact) W1644 := i Cm1644.gamma1644[4] is Hermitian with W1644.W1644 = 1 and trace
↪  0 (eigenvalues +1 and -1, eight times each); W1644 commutes with gamma1644[4] and with
↪  gamma1644[4].gamma1644[a] (a = 0..3), hence with K1644 at kappa = 0, and anticommutes with
↪  gamma1644[4].gamma1644[t] (t = 5, 6, 7)
[PASS] 44-D11  (exact) the density of the conserved current of the Spin(4,4)-invariant Lagrangian,
↪  j1644^4 = dirac1644^dagger (i Cm1644.gamma1644[4]) dirac1644 = dirac1644^dagger W1644
↪  dirac1644, is NOT positive: W1644 has eight eigenvalues -1; the positive density
↪  dirac1644^dagger dirac1644 is conserved only for kappa = 0 (44-D8)
[PASS] 71-D1  (exact) (gamma1671^A k_A)^2 = (eta1671^(AB) k_A k_B) 1 = (k4^2 - k0^2 - k1^2 - k2^2
↪  - k3^2 - k5^2 - k6^2 - k7^2) 1 for 40 random integer vectors k
[PASS] 71-D2  (exact) (gamma1671.k + m1671)(gamma1671.k - m1671) = (k.k - m1671^2) 1 for 40 random
↪  integer (k, m1671): with dirac1671 = u exp(-i k_A x^A) the operator i gamma1671^A d_A - m1671
↪  becomes gamma1671.k - m1671, and every component of a solution obeys the Klein-Gordon equation
↪  (d4^2 - d0^2 - d1^2 - d2^2 - d3^2 - d5^2 - d6^2 - d7^2 + m1671^2) dirac1671 = 0
[PASS] 71-D3  (exact) worked example E = 5, momentum p = (p1, p2, p3) = (2, 2, 4), m1671 = 1 (on
↪  the mass shell, 25 = 24 + 1): Pi1671 := (gamma1671.k + m1671)/(2 m1671) satisfies
↪  Pi1671.Pi1671 = Pi1671, trace Pi1671 = 8, (gamma1671.k - m1671).Pi1671 = 0: eight independent
↪  positive-energy plane waves
[PASS] 71-D4  (exact) beta1671 := gamma1671[4] and alpha1671^j := gamma1671[4].gamma1671[j] (j =
↪  0, 1, 2, 3, 5, 6, 7) are real symmetric; alpha1671^i.alpha1671^j + alpha1671^j.alpha1671^i = 2
↪  delta^(ij), alpha1671^j.beta1671 + beta1671.alpha1671^j = 0, beta1671.beta1671 = 1: the
↪  Hamiltonian h1671 = -i alpha1671^j d_j + beta1671 m1671 is Hermitian
[PASS] 71-D5  (exact) h1671(p) = p^j alpha1671^j + m1671 beta1671 is a real symmetric matrix with
↪  h1671(p).h1671(p) = (|p|^2 + m1671^2) 1 for 40 random integer (p, m1671): the energies are +E
↪  and -E, E = sqrt(|p|^2 + m1671^2), eight times each
[PASS] 71-D6  (exact) tau1671^j := tauO[swap04(j)] (swap04 = (0 4)): tau1671^4 = 1_8; for the seven
↪  space indices tau1671^j is real, antisymmetric, tau1671^i.tau1671^j + tau1671^j.tau1671^i =
↪  -2 delta^(ij); in 8x8 blocks beta1671 = [[0, 1], [1, 0]] and alpha1671^j = [[0, tau1671^j],
↪  [-tau1671^j, 0]] (beta1671 = betaD and alpha1671^j = alphaD[swap04(j)] of 7+1.nb section 7.1)
[PASS] 71-D7  (exact) chiral form: with EL1671 = (1; -i)/sqrt(2), ER1671 = (-i; 1)/sqrt(2) (16x8,
↪  in 8x8 blocks; chi1671.EL1671 = -EL1671, chi1671.ER1671 = +ER1671, orthonormal):
↪  EL1671^dagger.h1671(p).EL1671 = -i p^j tau1671^j, ER1671^dagger.h1671(p).ER1671 = +i p^j
↪  tau1671^j, EL1671^dagger.h1671(p).ER1671 = ER1671^dagger.h1671(p).EL1671 = m1671 (20 random
↪  integer (p, m1671)); hence (i d_4 + tau1671^j d_j) phiL1671 = m1671 phiR1671 and (i d_4 -
↪  tau1671^j d_j) phiR1671 = m1671 phiL1671 for phiL1671 = EL1671^dagger dirac1671, phiR1671 =
↪  ER1671^dagger dirac1671 (EL1671, ER1671, phiL1671, phiR1671 are the objects EL, ER, phiL, phiR
↪  of 7+1.nb sections 7.10 and 7.11, in the PROJECT labelling of the coordinates)
```

```
[PASS] 71-D8  (exact) worked example E = 5, p = (2, 2, 4), m1671 = 1: with M1671 = m1671 + p^j
↪  tau1671^j (M1671.transpose(M1671) = E^2 1_8), the columns of Upos1671 := (E 1;
↪  transpose(M1671)) satisfy h1671(p).Upos1671 = +E Upos1671 and those of Vneg1671 := (E 1;
↪  -M1671) satisfy h1671(-p).Vneg1671 = -E Vneg1671; transpose(Upos1671).Upos1671 =
↪  transpose(Vneg1671).Vneg1671 = 2 E^2 1_8: the spinors u_s = Upos1671 e_s/sqrt(E), v_s =
↪  Vneg1671 e_s/sqrt(E) are REAL and normalized to 2E
[PASS] 71-D9  (exact) charge conjugation dirac1671_c := omega1671 conj(dirac1671): omega1671 is
↪  real and omega1671.gamma1671[A] = -gamma1671[A].omega1671, so (i gamma1671^A d_A - m1671)
↪  dirac1671_c = omega1671 conj((i gamma1671^A d_A - m1671) dirac1671); omega1671.omega1671 = -1
↪  gives (dirac1671_c)_c = -dirac1671: a 16-component Majorana condition dirac1671_c = dirac1671
↪  has only the solution dirac1671 = 0
[PASS] 71-Q1  (numerical) 7+1, one particle at rest: the 8 spin states form the spinor of SO(7)
↪  (Casimir 21/4) and split under SO(3) x SU(2)_A x SU(2)_B (ordinary rotations x rotations of
↪  the extra space x0 x5 x6 x7) as (2; 2, 1) + (2; 1, 2); [A_a, B_b] = 0 to 0e0, residual 6e-16
[PASS] 71-Q2  (numerical) 7+1, two identical particles in an s wave (28 states allowed by the
↪  Pauli principle): under SO(7) they form 7 + 21 (Casimir 6 and 10); there is NO SO(7) singlet.
↪  Under SO(3) x SU(2)_A x SU(2)_B:  7 = (1; 2, 2) + (3; 1, 1),  21 = (1; 3, 1) + (1; 1, 3) + (3;
↪  1, 1) + (3; 2, 2).  The pairs of ordinary spin 0 are the 10 states (1; 2, 2) + (1; 3, 1) + (1;
↪  1, 3) (residual 3e-13)
[PASS] 71-Q3  (exact) 7+1: for space indices i != j, gamma1671[i].gamma1671[j] = diag(tau1671^i
↪  tau1671^j, tau1671^i tau1671^j) in 8x8 blocks, so on the particles at rest, u = (e;
↪  e)/sqrt(2), the rotation generator Sigma1671^(ij) acts as the real antisymmetric 8x8 matrix
↪  (1/2) tau1671^i tau1671^j
[PASS] 71-Q4  (exact) 7+1: the 7 matrices tau1671^j and the 21 matrices tau1671^i tau1671^j (i <
↪  j; space indices) are antisymmetric and linearly independent: together they span all 28
↪  antisymmetric 8x8 matrices, i.e. all s-wave pair amplitudes
[PASS] 71-Q5  (exact) 7+1: the quadratic Casimir operator of SO(7), Cas7(Phi) := -sum_(i<j)
↪  [Sigma1671^(ij), [Sigma1671^(ij), Phi]], gives Cas7(Phi) = 6 Phi for every Phi = tau1671^j
↪  (the vector, 7) and Cas7(Phi) = 10 Phi for every Phi = tau1671^i tau1671^j (the adjoint, 21)
[PASS] 71-Q6  (exact) 7+1: the linear system [tau1671^i tau1671^j, X] = 0 (all 21 planes) for an
↪  8x8 matrix X has rank 63 and is solved by X = 1: the only rotation-invariant pair matrix is
↪  the unit matrix, which is SYMMETRIC and therefore excluded by the Pauli principle.  Two
↪  identical particles of the field cannot form an s-wave pair that is a singlet under all
↪  rotations of the seven space dimensions
[PASS] 71-Q7  (exact) 7+1: the ten antisymmetric matrices tau1671^0, tau1671^5, tau1671^6,
↪  tau1671^7 and tau1671^a tau1671^b (a < b in {0, 5, 6, 7}) are linearly independent and commute
↪  with the generators tau1671^2 tau1671^3, tau1671^3 tau1671^1, tau1671^1 tau1671^2 of ordinary
↪  rotations; the system [tau1671^i tau1671^j, Phi] = 0 (ordinary rotations) on antisymmetric Phi
↪  has rank 18 = 28 - 10: these ten matrices are ALL the s-wave pairs of ordinary spin 0
[PASS] 71-Q8  (exact) 7+1: each of tau1671^j (any space index j) and tau1671^i tau1671^j (i != j)
↪  is an orthogonal matrix: Phi.transpose(Phi) = 1.  A condensate whose pair matrix is
↪  proportional to tau1671^j or to tau1671^i tau1671^j therefore gives the SAME gap to all eight
↪  spin states (a 'unitary' state)
[PASS] 71-Q9  (exact) 7+1: two-particle content of the field bilinears, u_s^T X u_r for particles
↪  at rest: X = Cm1671.gamma1671[j] gives -tau1671^j (space index j) and X = Cm1671.gamma1671[4]
↪  gives 0; X = Cp1671.gamma1671[i].gamma1671[j] gives tau1671^i tau1671^j.  Hence the vector
↪  pair field is DeltaV1671^j := dirac1671^T Cm1671 gamma1671^j dirac1671 (its time component
↪  contains no pair of two particles) and the adjoint pair field is DeltaT1671^(ij) :=
↪  dirac1671^T Cp1671 gamma1671^i gamma1671^j dirac1671
[PASS] 71-Q10  (exact) 7+1: the system [tau1671^i tau1671^j, Phi] = 0 for the six rotation planes
↪  (1,2), (1,3), (2,3) of ordinary space and (5,6), (5,7), (6,7) of the extra space, on
↪  antisymmetric Phi, has rank 27 = 28 - 1, and Phi = tau1671^0 solves it: EXACTLY ONE s-wave
↪  pair is a singlet both of the ordinary rotations and of the rotations of x5 x6 x7.  Its pair
↪  matrix is tau1671^0, the two-particle content of DeltaV1671^0 = dirac1671^T Cm1671
↪  gamma1671^0 dirac1671, the component of the vector pair field along the hidden space dimension
↪  x0 (the same result as in 4+4, check 44-Q4)
[PASS] 44-Q1  (numerical) 4+4, fields independent of x5 x6 x7, one particle at rest (gamma1644[4]
↪  u = i u, 8 states; deviation 0e0): under SU(2)_X x SU(2)_Y (the two factors of the rotation
↪  group SO(4) of the four space coordinates x0 x1 x2 x3; they are NOT the factors SU(2)_A x
↪  SU(2)_B of the internal symmetry of the zero-mode sector, which do not act on ordinary space)
↪  x SU(2)_T (rotations of the extra times) the states are (2, 1; 2) with one sign of W1644 and
↪  (1, 2; 2) with the other; every state has ordinary spin 1/2 (residual 2e-16)
```

```
[PASS] 44-Q2  (numerical) 4+4, fields independent of x5 x6 x7, two identical particles in an s
 ↪  wave (28 states): (1, 1; 3) twice, (3, 1; 1), (1, 3; 1), (2, 2; 1), (2, 2; 3) under SU(2)_X x
 ↪  SU(2)_Y x SU(2)_T (X, Y as in 44-Q1).  The pairs that are singlets under ALL rotations of the
 ↪  four space dimensions are the 6 states (1, 1; 3) + (1, 1; 3): two triplets of the extra-time
 ↪  rotations, none of them a singlet of SU(2)_T.  10 states have ordinary spin 0 (residual 1e-14)
[PASS] 44-Q3  (numerical) 4+4: among the 120 Pauli-allowed bilinears dirac1644^T Cm1644
 ↪  gamma1644[A1]...gamma1644[Ar] dirac1644 (A1 < ... < Ar), those whose two-particle content
 ↪  (particles at rest) lies entirely in the SO(4)-singlet pairs have the index sets {A1,...,Ar} =
 ↪  {5} {0,1,2,3,5} {4,5} {0,1,2,3,4,5} {6} {0,1,2,3,6} {4,6} {0,1,2,3,4,6} {7} {0,1,2,3,7} {4,7}
 ↪  {0,1,2,3,4,7}; they include the components A = 5, 6, 7 (the extra-time components) of the
 ↪  vector pair field DeltaV1644^A := dirac1644^T Cm1644 gamma1644^A dirac1644
[PASS] 44-Q4  (numerical) 4+4: exactly 1 of the 28 pair states is a singlet both of the ordinary
 ↪  rotations (x1 x2 x3) and of the extra-time rotations (x5 x6 x7); it belongs to the space
 ↪  vector (2, 2; 1), is made of one particle with W1644 = +1 and one with W1644 = -1, and is the
 ↪  two-particle content of the bilinears with the index sets {A1,...,Ar} = {0} {0,4}: in
 ↪  particular of DeltaV1644^0 = dirac1644^T Cm1644 gamma1644^0 dirac1644, the component of the
 ↪  vector pair field along the hidden space dimension x0
[PASS] ALL  (exact) every check above passed
```

A.3 The self-interaction (`deflation_cjt_ks_c selfcoupling`)

```
PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C: exact checks of the
 ↪  self-interaction of the Dirac field (71s-..: 7+1; 44q-..: 4+4, quantized-sector form; 44c-..:
 ↪  4+4, covariant form; self-..: all)
generated by: deflation_cjt_ks_c (Rust crate spacetimes-deflation-cjt-ks-c, version 0.1.0;
 ↪  PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C); command:
 ↪  deflation_cjt_ks_c selfcoupling
36 of 36 checks passed

[PASS] 71s-S1  (exact) 7+1: the matrices are Gamma^A = gamma1671[A] (real); the adjoint is psibar
 ↪  = psi^dagger gamma1671[4]. The eight matrices anticommute in pairs and their squares are
 ↪  diag[-1, -1, -1, -1, 1, -1, -1, -1] times the unit matrix; the adjoint matrix is Hermitian
 ↪  with square 1
[PASS] 71s-S2  (exact) the 256 matrices Theta^K = phase_K Gamma^(A_1)..Gamma^(A_r) (A_1 < .. <
 ↪  A_r; 1, 8, 28, 56, 70, 56, 28, 8, 1 index sets for r = 0..8): with phase_K = i for the ranks r
 ↪  in [2, 3, 6, 7] and phase_K = 1 for the other ranks, every matrix M_K = Adj.Theta^K is
 ↪  Hermitian, so that every bilinear psibar Theta^K psi is real; (Theta^K)^2 = eps_K 1 with eps_K
 ↪  = eta_K, the product of the squares of the matrices with index in K, so that Theta_K = eta_K
 ↪  Theta^K (the author's definition); Theta_K := (Theta^K)^(-1) = eps_K Theta^K
[PASS] 71s-S3  (exact) Tr(Theta_K Theta^L) = 16 if K = L and 0 otherwise, for all 65536 pairs: the
 ↪  256 matrices Theta^K are a basis of the complex 16x16 matrices, and Theta_K is the dual basis
[PASS] 71s-S4  (exact) invariance: for the 28 generators (1/2) Gamma^A Gamma^B (A < B), (i)
 ↪  Sigma^dagger.Adj + Adj.Sigma = 0, so that psibar psi is invariant, and (ii) the commutator
 ↪  [Theta^K, Gamma^A Gamma^B] is zero or a multiple c_K of the single matrix Theta^K' of the same
 ↪  rank with K' = K with A, B exchanged in or out, and eps_K c_K + eps_K' c_K' = 0: each of the
 ↪  nine invariants I_r = sum_K (psibar Theta_K psi)(psibar Theta^K psi) is invariant. The group
 ↪  is Spin(7,1)
[PASS] 71s-S5  (exact) Fierz rearrangement. Every Theta^K either commutes or anticommutes with
 ↪  every Theta^L; the sum of these signs over the index sets K of rank r is the same for all L of
 ↪  rank s and equals 16 F_rs with the closed form F_rs = (1/16) sum_l (-1)^(rs - l) C(s, l) C(8 -
 ↪  s, r - l); the identity sum_(K of rank r) (Theta_K)_(ab) (Theta^K)_(cd) = sum_s F_rs sum_(L of
 ↪  rank s) (Theta_L)_(ad) (Theta^L)_(cb) holds for all 9 x 65536 combinations of r and of the
 ↪  spinor indices a, b, c, d (verified with the adjoint matrix multiplied in, that is for the
 ↪  matrices M_K); F^2 = 1, Tr F = 16/16, rank(1 + F) = 5 and rank(1 - F) = 4: F has the
 ↪  eigenvalue +1 five times and -1 four times
```

[PASS] 71s-S6 (exact) THEOREM (anticommuting fields). Write I_r = sum T^(r)_(abcd) psi_a^* psi_b psi_c^* psi_d. Two tensors give the same polynomial in Grassmann variables exactly if their parts antisymmetric in b and d agree, A_(abcd) = T_(abcd) - T_(adcb). The nine tensors T^(r) are linearly independent (rank 9), but their antisymmetric parts satisfy sum_s (delta_rs + F_rs) A^(s) = 0 for every r and span a space of dimension 4: the nine invariants obey (1 + F) I = 0, five independent linear relations, and four invariants are independent. In terms of I_0, I_1, I_7, I_8: I_2 = -3 I_0 - I_1 - I_7 - I_8, I_3 = -4 I_0 - I_7 + 4 I_8, I_4 = -5 I_0 - 5 I_8, I_5 = -4 I_0 - I_1 + 4 I_8, I_6 = -I_0 + I_1 + I_7 - 3 I_8 (each verified on all 65536 components). Therefore sum_r g_r I_r = g_0eff I_0 + g_1eff I_1 + g_7eff I_7 + g_8eff I_8 with g_0eff = g_0 - 3 g_2 - 4 g_3 - 5 g_4 - 4 g_5 - g_6, g_1eff = g_1 - g_2 - g_5 + g_6, g_7eff = g_7 - g_2 - g_3 + g_6, g_8eff = g_8 - g_2 + 4 g_3 - 5 g_4 + 4 g_5 - 3 g_6

[PASS] 71s-S7 (exact) Hartree-Fock. With the one-body density matrix n_ab = <psi_b^dagger psi_a> of a state without pairing, <:(psi^dagger M psi)^2:> = Tr(M n)^2 - Tr(M n M n) (Wick's theorem), and the mean-field potential sum_r g_r sum_K eps_K [Tr(M_K n) M_K - M_K n M_K] is again a combination of the matrices in n; for the density matrix of uniform matter n = (rho 1 + rho_S beta)/16 (rho = <psi^dagger psi>, rho_S = <psibar psi>) the Hartree-Fock energy is E_HF = (1/2)[G_1 rho^2 + G_0 rho_S^2] with G_1 = (-g_0 + 22 g_1 - 14 g_2 + 14 g_3 - 14 g_5 + 14 g_6 - 6 g_7 + g_8)/16 and G_0 = (15 g_0 - 8 g_1 - 28 g_2 - 56 g_3 - 70 g_4 - 56 g_5 - 28 g_6 - 8 g_7 - g_8)/16, that is G = (1 - F^T) g: the exchange (Fock) terms are the Fierz transforms of the direct (Hartree) terms g_1 rho^2 and g_0 rho_S^2; the mean-field potential is U = G_1 rho 1 + G_0 rho_S beta, so that the mass of the fermions in matter is M = m + G_0 rho_S and every single-particle energy is shifted by G_1 rho. In the four effective couplings: G_1 = (-g_0eff + 22 g_1eff - 6 g_7eff + g_8eff)/16, G_0 = (15 g_0eff - 8 g_1eff - 8 g_7eff - g_8eff)/16.

[PASS] 71s-S8 (exact) the mean-field potential U(n) = sum_K eps_K [Tr(M_K n) M_K - M_K n M_K] is the derivative of the Hartree-Fock energy E(n) = (1/2) sum_K eps_K [Tr(M_K n)^2 - Tr(M_K n M_K n)]: d E(n + t d)/dt at t = 0 equals Tr(U(n) d), verified exactly for each of the nine ranks with four pairs of Hermitian integer matrices n and d. The same derivative with respect to psibar gives the nonlinear term of the Dirac equation, sum_r g_r sum_K (psibar Theta_K psi) Theta^K psi

[PASS] 71s-S9 (exact) pairs. With the pair density kappa_ab = <psi_a psi_b> the pairing energy of the interaction is (1/2) sum_r g_r Tr(kappa^dagger P_r(kappa)), P_r(kappa) = sum_K eps_K M_K kappa M_K^T. Of the 256 real matrices X_J = Cm.gamma^J, 120 are antisymmetric (pair operators psi^T X_J psi that do not vanish); each is an eigenvector of all nine maps P_r with an integer eigenvalue. Projected on two particles at rest, kappa = P_+ X_J P_+^T with P_+ = (1 + beta)/2, 56 of them are non-zero, and each is an eigenvector of the projected maps P_+ P_r(kappa) P_+^T = q_r kappa, with the pair coupling G_pair = sum_r g_r q_r (attractive if negative). X_J and X_J.gamma^4 project on the same pair, so that there are 28 different pairs, as many as there are antisymmetric states of two particles with eight internal states each (8 x 7/2 = 28): 7 pairs with G_pair = g_0 + g_1 - 9 g_2 - 9 g_3 - 5 g_4 - 5 g_5 + 5 g_6 + 5 g_7; 21 pairs with G_pair = g_0 + g_1 - g_2 - g_3 - 5 g_4 - 5 g_5 - 3 g_6 - 3 g_7. For the hidden-axis pair psi^T Cm.gamma^0 psi, G_pair = g_0 + g_1 - 9 g_2 - 9 g_3 - 5 g_4 - 5 g_5 + 5 g_6 + 5 g_7; in the four effective couplings this is g_0eff + g_1eff + 5 g_7eff.

[PASS] 71s-S10 (exact) chiral phase rotation psi -> exp(i theta chi) psi with the Hermitian matrix chi = i omega1671 (omega1671 the product of the eight matrices) (chi^2 = 1): the invariants of the ranks [1, 3, 5, 7] are invariant one by one; the interaction sum_r g_r I_r is invariant exactly if g_0 - 4 g_2 - 10 g_4 - 4 g_6 + g_8 = 0 (1 condition(s) on the nine couplings; determined from the antisymmetrized variation of the quartic tensors). In the four effective couplings the condition reads g_0eff + g_8eff = 0: three of the four independent couplings are chirally invariant (vector, axial vector, and the combination I_0 - I_8 of scalar and pseudoscalar), one is not. A mass term m psibar psi is not invariant either.

[PASS] 44c-S1 (exact) 4+4, covariant form: the matrices are Gamma^A = gamma1644[A] (real); the adjoint is psitilde = psi^dagger Cm1644 with Cm1644 = gamma1644[0].gamma1644[1].gamma1644[2].gamma1644[3]. The eight matrices anticommute in pairs and their squares are diag[1, 1, 1, 1, -1, -1, -1, -1] times the unit matrix; the adjoint matrix is Hermitian with square 1; W = i Cm1644.gamma1644[4] is Hermitian with W^2 = 1, commutes with beta1644, and Cm1644 = -W.beta1644

[PASS] 44c-S2 (exact) the 256 matrices Theta^K = phase_K Gamma^(A_1)..Gamma^(A_r) (A_1 < .. < A_r; 1, 8, 28, 56, 70, 56, 28, 8, 1 index sets for r = 0..8): with phase_K = i for the ranks r in [1, 2, 5, 6] and phase_K = 1 for the other ranks, every matrix M_K = Adj.Theta^K is Hermitian, so that every bilinear psibar Theta^K psi is real; (Theta^K)^2 = eps_K 1 with eps_K = (-1)^r eta_K with eta_K the product of the squares of the matrices with index in K (the inverse matrix differs from eta_K Theta^K by the sign (-1)^r, because the adjoint is Cm1644 and not a single Dirac matrix); Theta_K := (Theta^K)^(-1) = eps_K Theta^K

[PASS] 44c-S3 (exact) Tr(Theta_K Theta^L) = 16 if K = L and 0 otherwise, for all 65536 pairs: the 256 matrices Theta^K are a basis of the complex 16x16 matrices, and Theta_K is the dual basis

[PASS] 44c-S4 (exact) invariance: for the 28 generators (1/2) Gamma^A Gamma^B (A < B), (i) Sigma^dagger.Adj + Adj.Sigma = 0, so that psibar psi is invariant, and (ii) the commutator [Theta^K, Gamma^A Gamma^B] is zero or a multiple c_K of the single matrix Theta^K' of the same rank with K' = K with A, B exchanged in or out, and eps_K c_K + eps_K' c_K' = 0: each of the nine invariants I_r = sum_K (psibar Theta_K psi)(psibar Theta^K psi) is invariant. The group is Spin(4,4); 13 of the 28 generators commute with W (the rotations and boosts of x0..x4 and the rotations of x5, x6, x7), and only these are symmetries of the positive Lagrangian of the quantized sector

[PASS] 44c-S5 (exact) Fierz rearrangement. Every Theta^K either commutes or anticommutes with every Theta^L; the sum of these signs over the index sets K of rank r is the same for all L of rank s and equals 16 F_rs with the closed form F_rs = (1/16) sum_l (-1)^(rs - l) C(s, l) C(8 - s, r - l); the identity sum_(K of rank r) (Theta_K)_(ab) (Theta^K)_(cd) = sum_s F_rs sum_(L of rank s) (Theta_L)_(ad) (Theta^L)_(cb) holds for all 9 x 65536 combinations of r and of the spinor indices a, b, c, d (verified with the adjoint matrix multiplied in, that is for the matrices M_K); F^2 = 1, Tr F = 16/16, rank(1 + F) = 5 and rank(1 - F) = 4: F has the eigenvalue +1 five times and -1 four times

[PASS] 44c-S6 (exact) THEOREM (anticommuting fields). Write I_r = sum T^(r)_(abcd) psi_a^* psi_b psi_c^* psi_d. Two tensors give the same polynomial in Grassmann variables exactly if their parts antisymmetric in b and d agree, A_(abcd) = T_(abcd) - T_(adcb). The nine tensors T^(r) are linearly independent (rank 9), but their antisymmetric parts satisfy sum_s (delta_rs + F_rs) A^(s) = 0 for every r and span a space of dimension 4: the nine invariants obey (1 + F) I = 0, five independent linear relations, and four invariants are independent. In terms of I_0, I_1, I_7, I_8: I_2 = -3 I_0 - I_1 - I_7 - I_8, I_3 = -4 I_0 - I_7 + 4 I_8, I_4 = -5 I_0 - 5 I_8, I_5 = -4 I_0 - I_1 + 4 I_8, I_6 = -I_0 + I_1 + I_7 - 3 I_8 (each verified on all 65536 components). Therefore sum_r g_r I_r = g_0eff I_0 + g_1eff I_1 + g_7eff I_7 + g_8eff I_8 with g_0eff = g_0 - 3 g_2 - 4 g_3 - 5 g_4 - 4 g_5 - g_6, g_1eff = g_1 - g_2 - g_5 + g_6, g_7eff = g_7 - g_2 - g_3 + g_6, g_8eff = g_8 - g_2 + 4 g_3 - 5 g_4 + 4 g_5 - 3 g_6

[PASS] 44c-S7 (exact) Hartree-Fock. With the one-body density matrix n_ab = <psi_b^dagger psi_a> of a state without pairing, <:(psi^dagger M psi)^2:> = Tr(M n)^2 - Tr(M n M n) (Wick's theorem), and the mean-field potential sum_r g_r sum_K eps_K [Tr(M_K n) M_K - M_K n M_K] is again a combination of the matrices in n; for the density matrix n = (rho 1 + rho_S beta + rho_W W + rho_SW W.beta)/16 (rho = <psi^dagger psi>, rho_S = <psi^dagger beta psi>, rho_W = <psi^dagger W psi> the covariant charge density, rho_SW = <psi^dagger W.beta psi> = -<psitilde psi>) the Hartree-Fock energy is diagonal, E_HF = (1/2)[G_V0 rho^2 + G_S0 rho_S^2 + G_1 rho_W^2 + G_0 rho_SW^2], with G_V0 = (-g_0 + 4 g_2 + 10 g_4 + 4 g_6 - g_8)/16, G_S0 = (-g_0 - 2 g_1 + 2 g_2 + 6 g_3 + 10 g_5 - 2 g_6 + 2 g_7 + g_8)/16, G_1 = (-g_0 + 22 g_1 - 14 g_2 + 14 g_3 - 14 g_5 + 14 g_6 - 6 g_7 + g_8)/16, G_0 = (15 g_0 - 8 g_1 - 28 g_2 - 56 g_3 - 70 g_4 - 56 g_5 - 28 g_6 - 8 g_7 - g_8)/16; the mean-field potential is U = G_V0 rho 1 + G_S0 rho_S beta + G_1 rho_W W + G_0 rho_SW W.beta: the two halves W = +1 and W = -1 of the field have the masses m + G_S0 rho_S +- G_0 rho_SW and the energy shifts G_V0 rho +- G_1 rho_W. In the four effective couplings: G_V0 = (-g_0eff - g_8eff)/16, G_S0 = (-g_0eff - 2 g_1eff + 2 g_7eff + g_8eff)/16, G_1 = (-g_0eff + 22 g_1eff - 6 g_7eff + g_8eff)/16, G_0 = (15 g_0eff - 8 g_1eff - 8 g_7eff - g_8eff)/16.

[PASS] 44c-S8 (exact) the mean-field potential U(n) = sum_K eps_K [Tr(M_K n) M_K - M_K n M_K] is the derivative of the Hartree-Fock energy E(n) = (1/2) sum_K eps_K [Tr(M_K n)^2 - Tr(M_K n M_K n)]: d E(n + t d)/dt at t = 0 equals Tr(U(n) d), verified exactly for each of the nine ranks with four pairs of Hermitian integer matrices n and d. The same derivative with respect to psibar gives the nonlinear term of the Dirac equation, sum_r g_r sum_K (psibar Theta_K psi) Theta^K psi

[PASS] 44c-S9 (exact) pairs. With the pair density kappa_ab = <psi_a psi_b> the pairing energy of the interaction is (1/2) sum_r g_r Tr(kappa^dagger P_r(kappa)), P_r(kappa) = sum_K eps_K M_K kappa M_K^T. Of the 256 real matrices X_J = Cm.gamma^J, 120 are antisymmetric (pair operators psi^T X_J psi that do not vanish); each is an eigenvector of all nine maps P_r with an integer eigenvalue. Projected on two particles at rest, kappa = P_+ X_J P_+^T with P_+ = (1 + beta)/2, 56 of them are non-zero, and each is an eigenvector of the projected maps P_+ P_r(kappa) P_+^T = q_r kappa, with the pair coupling G_pair = sum_r g_r q_r (attractive if negative). X_J and X_J.gamma^4 project on the same pair, so that there are 28 different pairs, as many as there are antisymmetric states of two particles with eight internal states each (8 x 7/2 = 28): 4 pairs (one particle in each of the halves W = +1 and W = -1) with G_pair = -g_0 - g_1 + 9 g_2 + 9 g_3 + 5 g_4 + 5 g_5 - 5 g_6 - 5 g_7; 3 pairs (both particles in the same half W = +1 or W = -1) with G_pair = g_0 + g_1 - 9 g_2 - 9 g_3 - 5 g_4 - 5 g_5 + 5 g_6 + 5 g_7; 9 pairs (both particles in the same half W = +1 or W = -1) with G_pair = g_0 + g_1 - g_2 - g_3 - 5 g_4 - 5 g_5 - 3 g_6 - 3 g_7; 12 pairs (one particle in each of the halves W = +1 and W = -1) with G_pair = -g_0 - g_1 + g_2 + g_3 + 5 g_4 + 5 g_5 + 3 g_6 + 3 g_7. For the hidden-axis pair psi^T Cm.gamma^0 psi, G_pair = -g_0 - g_1 + 9 g_2 + 9 g_3 + 5 g_4 + 5 g_5 - 5 g_6 - 5 g_7; in the four effective couplings this is -g_0eff - g_1eff - 5 g_7eff. In the quantized sector the hidden-axis pair satisfies W kappa W^T = - kappa: it joins a particle of the half W = +1 with a particle of the half W = -1.

[PASS] 44c-S10 (exact) chiral phase rotation psi -> exp(i theta chi) psi with the Hermitian matrix chi = omega1644 (the product of the eight matrices gamma1644, real) (chi^2 = 1): the invariants of the ranks [0, 2, 4, 6, 8] are invariant one by one; the interaction sum_r g_r I_r is invariant exactly if g_1 + g_3 - g_5 - g_7 = 0 (1 condition(s) on the nine couplings; determined from the antisymmetrized variation of the quartic tensors). In the four effective couplings the condition reads g_1eff - g_7eff = 0. (For the covariant form this rotation is the symmetry of the kinetic term of the quantized sector; the covariant kinetic term itself is not invariant under it, and the covariant mass term is. The roles of the even and of the odd ranks are exchanged with respect to the forms with a Dirac adjoint, because the matrix Cm1644 commutes with omega1644.)

[PASS] 44q-S1 (exact) 4+4, quantized-sector form: the matrices are Gamma_eff^A = -i gamma1644[A] for A = 0..4 and Gamma_eff^t = gamma1644[t] for t = 5, 6, 7; the adjoint is psibar = psi^dagger beta1644 with beta1644 = -i gamma1644[4] = Gamma_eff^4. The eight matrices anticommute in pairs and their squares are diag[-1, -1, -1, -1, 1, -1, -1, -1] times the unit matrix; the adjoint matrix is Hermitian with square 1; W = i Cm1644.gamma1644[4] is Hermitian with W^2 = 1, commutes with beta1644, and Cm1644 = -W.beta1644

[PASS] 44q-S2 (exact) the 256 matrices Theta^K = phase_K Gamma^(A_1)..Gamma^(A_r) (A_1 < .. < A_r; 1, 8, 28, 56, 70, 56, 28, 8, 1 index sets for r = 0..8): with phase_K = i for the ranks r in [2, 3, 6, 7] and phase_K = 1 for the other ranks, every matrix M_K = Adj.Theta^K is Hermitian, so that every bilinear psibar Theta^K psi is real; (Theta^K)^2 = eps_K 1 with eps_K = eta_K, the product of the squares of the matrices with index in K, so that Theta_K = eta_K Theta^K (the author's definition); Theta_K := (Theta^K)^(-1) = eps_K Theta^K

[PASS] 44q-S3 (exact) Tr(Theta_K Theta^L) = 16 if K = L and 0 otherwise, for all 65536 pairs: the 256 matrices Theta^K are a basis of the complex 16x16 matrices, and Theta_K is the dual basis

[PASS] 44q-S4 (exact) invariance: for the 28 generators (1/2) Gamma^A Gamma^B (A < B), (i) Sigma^dagger.Adj + Adj.Sigma = 0, so that psibar psi is invariant, and (ii) the commutator [Theta^K, Gamma^A Gamma^B] is zero or a multiple c_K of the single matrix Theta^K' of the same rank with K' = K with A, B exchanged in or out, and eps_K c_K + eps_K' c_K' = 0: each of the nine invariants I_r = sum_K (psibar Theta_K psi)(psibar Theta^K psi) is invariant. The group is the group generated by (1/2) Gamma_eff^A Gamma_eff^B, of which the 13 generators that commute with W act on the coordinates (rotations and boosts of x0..x4, rotations of x5, x6, x7: Spin(4,1) x Spin(3)); the other 15 mix a coordinate x0..x4 with one of x5, x6, x7 and are not symmetries of the kinetic term

[PASS] 44q-S5 (exact) Fierz rearrangement. Every Theta^K either commutes or anticommutes with every Theta^L; the sum of these signs over the index sets K of rank r is the same for all L of rank s and equals 16 F_rs with the closed form F_rs = (1/16) sum_l (-1)^(rs - l) C(s, l) C(8 - s, r - l); the identity sum_(K of rank r) (Theta_K)_(ab) (Theta^K)_(cd) = sum_s F_rs sum_(L of rank s) (Theta_L)_(ad) (Theta^L)_(cb) holds for all 9 x 65536 combinations of r and of the spinor indices a, b, c, d (verified with the adjoint matrix multiplied in, that is for the matrices M_K); F^2 = 1, Tr F = 16/16, rank(1 + F) = 5 and rank(1 - F) = 4: F has the eigenvalue +1 five times and -1 four times

[PASS] 44q-S6 (exact) THEOREM (anticommuting fields). Write I_r = sum T^(r)_(abcd) psi_a^* psi_b psi_c^* psi_d. Two tensors give the same polynomial in Grassmann variables exactly if their parts antisymmetric in b and d agree, A_(abcd) = T_(abcd) - T_(adcb). The nine tensors T^(r) are linearly independent (rank 9), but their antisymmetric parts satisfy sum_s (delta_rs + F_rs) A^(s) = 0 for every r and span a space of dimension 4: the nine invariants obey (1 + F) I = 0, five independent linear relations, and four invariants are independent. In terms of I_0, I_1, I_7, I_8: I_2 = -3 I_0 - I_1 - I_7 - I_8, I_3 = -4 I_0 - I_7 + 4 I_8, I_4 = -5 I_0 - 5 I_8, I_5 = -4 I_0 - I_1 + 4 I_8, I_6 = -I_0 + I_1 + I_7 - 3 I_8 (each verified on all 65536 components). Therefore sum_r g_r I_r = g_0eff I_0 + g_1eff I_1 + g_7eff I_7 + g_8eff I_8 with g_0eff = g_0 - 3 g_2 - 4 g_3 - 5 g_4 - 4 g_5 - g_6, g_1eff = g_1 - g_2 - g_5 + g_6, g_7eff = g_7 - g_2 - g_3 + g_6, g_8eff = g_8 - g_2 + 4 g_3 - 5 g_4 + 4 g_5 - 3 g_6

[PASS] 44q-S7 (exact) Hartree-Fock. With the one-body density matrix n_ab = <psi_b^dagger psi_a> of a state without pairing, <:(psi^dagger M psi)^2:> = Tr(M n)^2 - Tr(M n M n) (Wick's theorem), and the mean-field potential sum_r g_r sum_K eps_K [Tr(M_K n) M_K - M_K n M_K] is again a combination of the matrices in n; for the density matrix n = (rho 1 + rho_S beta + rho_W W + rho_SW W.beta)/16 (rho = <psi^dagger psi>, rho_S = <psibar psi>, rho_W = <psi^dagger W psi>, rho_SW = <psi^dagger W.beta psi>: the two halves W = +1 and W = -1 of the field filled differently) the Hartree-Fock energy is diagonal, E_HF = (1/2)[G_1 rho^2 + G_0 rho_S^2 + G_V0 rho_W^2 + G_S0 rho_SW^2], with G_1 = (-g_0 + 22 g_1 - 14 g_2 + 14 g_3 - 14 g_5 + 14 g_6 - 6 g_7 + g_8)/16 and G_0 = (15 g_0 - 8 g_1 - 28 g_2 - 56 g_3 - 70 g_4 - 56 g_5 - 28 g_6 - 8 g_7 - g_8)/16, that is G = (1 - F^T) g (the exchange terms are the Fierz transforms of the direct terms g_1 rho^2 and g_0 rho_S^2), and G_V0 = (-g_0 + 4 g_2 + 10 g_4 + 4 g_6 - g_8)/16, G_S0 = (-g_0 - 2 g_1 + 2 g_2 + 6 g_3 + 10 g_5 - 2 g_6 + 2 g_7 + g_8)/16; the mean-field potential is U = G_1 rho 1 + G_0 rho_S beta + G_V0 rho_W W + G_S0 rho_SW W.beta. For matter with equal halves (rho_W = rho_SW = 0) the mass of the fermions is M = m + G_0 rho_S and every single-particle energy is shifted by G_1 rho. In the four effective couplings: G_1 = (-g_0eff + 22 g_1eff - 6 g_7eff + g_8eff)/16, G_0 = (15 g_0eff - 8 g_1eff - 8 g_7eff - g_8eff)/16, G_V0 = (-g_0eff - g_8eff)/16, G_S0 = (-g_0eff - 2 g_1eff + 2 g_7eff + g_8eff)/16.

[PASS] 44q-S8 (exact) the mean-field potential U(n) = sum_K eps_K [Tr(M_K n) M_K - M_K n M_K] is the derivative of the Hartree-Fock energy E(n) = (1/2) sum_K eps_K [Tr(M_K n)^2 - Tr(M_K n M_K n)]: d E(n + t d)/dt at t = 0 equals Tr(U(n) d), verified exactly for each of the nine ranks with four pairs of Hermitian integer matrices n and d. The same derivative with respect to psibar gives the nonlinear term of the Dirac equation, sum_r g_r sum_K (psibar Theta_K psi) Theta^K psi

[PASS] 44q-S9 (exact) pairs. With the pair density kappa_ab = <psi_a psi_b> the pairing energy of the interaction is (1/2) sum_r g_r Tr(kappa^dagger P_r(kappa)), P_r(kappa) = sum_K eps_K M_K kappa M_K^T. Of the 256 real matrices X_J = Cm.gamma^J, 120 are antisymmetric (pair operators psi^T X_J psi that do not vanish); each is an eigenvector of all nine maps P_r with an integer eigenvalue. Projected on two particles at rest, kappa = P_+ X_J P_+^T with P_+ = (1 + beta)/2, 56 of them are non-zero, and each is an eigenvector of the projected maps P_+ P_r(kappa) P_+^T = q_r kappa, with the pair coupling G_pair = sum_r g_r q_r (attractive if negative). X_J and X_J.gamma^4 project on the same pair, so that there are 28 different pairs, as many as there are antisymmetric states of two particles with eight internal states each (8 x 7/2 = 28): 4 pairs (one particle in each of the halves W = +1 and W = -1) with G_pair = g_0 + g_1 - 9 g_2 - 9 g_3 - 5 g_4 - 5 g_5 + 5 g_6 + 5 g_7; 3 pairs (both particles in the same half W = +1 or W = -1) with G_pair = g_0 + g_1 - 9 g_2 - 9 g_3 - 5 g_4 - 5 g_5 + 5 g_6 + 5 g_7; 9 pairs (both particles in the same half W = +1 or W = -1) with G_pair = g_0 + g_1 - g_2 - g_3 - 5 g_4 - 5 g_5 - 3 g_6 - 3 g_7; 12 pairs (one particle in each of the halves W = +1 and W = -1) with G_pair = g_0 + g_1 - g_2 - g_3 - 5 g_4 - 5 g_5 - 3 g_6 - 3 g_7. For the hidden-axis pair psi^T Cm.gamma^0 psi, G_pair = g_0 + g_1 - 9 g_2 - 9 g_3 - 5 g_4 - 5 g_5 + 5 g_6 + 5 g_7; in the four effective couplings this is g_0eff + g_1eff + 5 g_7eff. In the quantized sector the hidden-axis pair satisfies W kappa W^T = - kappa: it joins a particle of the half W = +1 with a particle of the half W = -1.

[PASS] 44q-S10 (exact) chiral phase rotation psi -> exp(i theta chi) psi with the Hermitian matrix chi = omega1644 (the product of the eight matrices gamma1644, real) (chi^2 = 1): the invariants of the ranks [1, 3, 5, 7] are invariant one by one; the interaction sum_r g_r I_r is invariant exactly if g_0 - 4 g_2 - 10 g_4 - 4 g_6 + g_8 = 0 (1 condition(s) on the nine couplings; determined from the antisymmetrized variation of the quartic tensors). In the four effective couplings the condition reads g_0eff + g_8eff = 0: three of the four independent couplings are chirally invariant (vector, axial vector, and the combination I_0 - I_8 of scalar and pseudoscalar), one is not. A mass term m psibar psi is not invariant either.

[PASS] 44c-S11 (exact) relation between the two forms of the 4+4 interaction: for every index set K the covariant matrix is M_K(covariant) = z_K W.M_K(quantized sector) with z_K = +-1 if K contains an even number of the indices 5, 6, 7 (128 index sets; W commutes with M_K) and z_K = +-i if it contains an odd number (128 index sets; W anticommutes with M_K, and the bilinear connects the two halves W = +1 and W = -1 of the field). With eps_K and M_K of the quantized-sector form, I_r(covariant) = sum_K eps_K (psi^dagger W.M_K psi)^2 for all 256 index sets: the covariant form is the quantized-sector form with the matrix W inserted in every bilinear, and nothing else. Consequence: between two particles of the same half the two forms give the same interaction; between a particle of the half W = +1 and a particle of the half W = -1 they give interactions of opposite sign

[PASS] 44-S13 (exact) the two forms of the 4+4 interaction in uniform matter whose density matrix is diagonal in W. With n_+-, s_+- the number and scalar densities of the halves W = +-1, rho = n_+ + n_-, rho_W = n_+ - n_-, rho_S = s_+ + s_-, rho_SW = s_+ - s_-, the Hartree-Fock energies are E(quantized-sector form) = (1/2)[G_1 rho^2 + G_0 rho_S^2 + G_V0 rho_W^2 + G_S0 rho_SW^2] and E(covariant form) = (1/2)[G_V0 rho^2 + G_S0 rho_S^2 + G_1 rho_W^2 + G_0 rho_SW^2] with THE SAME four combinations of the nine couplings (compared coefficient by coefficient): G_0 = (15 g_0 - 8 g_1 - 28 g_2 - 56 g_3 - 70 g_4 - 56 g_5 - 28 g_6 - 8 g_7 - g_8)/16 = (15 g_0eff - 8 g_1eff - 8 g_7eff - g_8eff)/16, G_1 = (-g_0 + 22 g_1 - 14 g_2 + 14 g_3 - 14 g_5 + 14 g_6 - 6 g_7 + g_8)/16 = (-g_0eff + 22 g_1eff - 6 g_7eff + g_8eff)/16, G_S0 = (-g_0 - 2 g_1 + 2 g_2 + 6 g_3 + 10 g_5 - 2 g_6 + 2 g_7 + g_8)/16 = (-g_0eff - 2 g_1eff + 2 g_7eff + g_8eff)/16, G_V0 = (-g_0 + 4 g_2 + 10 g_4 + 4 g_6 - g_8)/16 = (-g_0eff - g_8eff)/16. Equivalently E = (1/2) G_Vsame (n_+^2 + n_-^2) + (1/2) G_Ssame (s_+^2 + s_-^2) +- [G_Vopp n_+ n_- + G_Sopp s_+ s_-], upper sign for the quantized-sector form and lower sign for the covariant form, with the couplings between particles of the same half G_Ssame = G_0 + G_S0 = (7 g_0eff - 5 g_1eff - 3 g_7eff)/8, G_Vsame = G_1 + G_V0 = (-g_0eff + 11 g_1eff - 3 g_7eff)/8 and the couplings between particles of different halves G_Sopp = G_0 - G_S0 = (8 g_0eff - 3 g_1eff - 5 g_7eff - g_8eff)/8, G_Vopp = G_1 - G_V0 = (11 g_1eff - 3 g_7eff + g_8eff)/8: the two forms agree inside each half and have opposite signs between the halves

[PASS] 71s-S12 (exact) internal symmetry of the fields that do not depend on the extra coordinates x0, x5, x6, x7 (their kinetic term contains alpha^1, alpha^2, alpha^3 and beta only). The eight Hermitian generators of the symmetry of the model with the Vorton-Scalar are the six rotations of the extra axes, (phase) gamma^a gamma^b with a < b in {0, 5, 6, 7}, the unit matrix (fermion number), and the generator P = (phase) gamma^0 gamma^5 gamma^6 gamma^7 of U(1)_E, with P^2 = 1 and trace 0; all commute with the four matrices of the kinetic term, and P commutes with the chirality matrix. The self-interaction is invariant under the six rotations and the phase for all values of the nine couplings (0 conditions found), because the rotations are among the 28 generators of check S4. Under psi -> exp(i theta P) psi the invariants of the ranks [0, 4, 8] are invariant one by one, and the interaction sum_r g_r I_r is invariant exactly if g_1 + g_3 - g_5 - g_7 (that is g_1eff - g_7eff) = 0 and g_2 + g_3 - g_6 - g_7 (that is -g_7eff) = 0 (2 condition(s))

[PASS] 44c-S12 (exact) internal symmetry of the fields that do not depend on the extra coordinates x0, x5, x6, x7 (their kinetic term contains alpha^1, alpha^2, alpha^3 and beta only). The eight Hermitian generators of the symmetry of the model with the Vorton-Scalar are the six rotations of the extra axes, (phase) gamma^a gamma^b with a < b in {0, 5, 6, 7}, the unit matrix (fermion number), and the generator P = (phase) gamma^0 gamma^5 gamma^6 gamma^7 of U(1)_E, with P^2 = 1 and trace 0; all commute with the four matrices of the kinetic term, and P commutes with the chirality matrix. The rotations of x5, x6, x7 among themselves commute with W (they act inside each half of the field); the three generators gamma^0 gamma^t and the generator P of U(1)_E anticommute with W (they connect the half W = +1 with the half W = -1). Under the three rotations of x5, x6, x7 among themselves, which are unitary elements of Spin(4,4), the self-interaction is invariant for all values of the couplings. Under each of the three unitary transformations psi -> exp(i theta gamma^0 gamma^t) psi (t = 5, 6, 7), which are NOT elements of Spin(4,4) (there gamma^0 gamma^t generates a boost, which is not unitary), the interaction sum_r g_r I_r is invariant exactly if (5 g_0 - 30 g_4 - 16 g_5 - 24 g_6 - 16 g_7 + g_8)/5 (that is (5 g_0eff - 16 g_7eff + g_8eff)/5) = 0 and (5 g_1 + 5 g_4 - 9 g_5 + 4 g_6 - 4 g_7 - g_8)/5 (that is (5 g_1eff - 4 g_7eff - g_8eff)/5) = 0 and (5 g_2 + 5 g_4 - 4 g_5 - g_6 - 4 g_7 - g_8)/5 (that is (-4 g_7eff - g_8eff)/5) = 0 and (5 g_3 - 5 g_4 + 4 g_5 - 4 g_6 - g_7 + g_8)/5 (that is (-g_7eff + g_8eff)/5) = 0 (4 condition(s) on the nine couplings, the same for the three) Under psi -> exp(i theta P) psi the invariants of the ranks [] are invariant one by one, and the

```
interaction sum_r g_r I_r is invariant exactly if (5 g_0 - 30 g_4 - 16 g_5 - 24 g_6 - 16 g_7 +
 ↪  g_8)/5 (that is (5 g_0eff - 16 g_7eff + g_8eff)/5) = 0 and (5 g_1 + 5 g_4 - 9 g_5 + 4 g_6 - 4
 ↪  g_7 - g_8)/5 (that is (5 g_1eff - 4 g_7eff - g_8eff)/5) = 0 and (5 g_2 + 5 g_4 - 4 g_5 - g_6 -
 ↪  4 g_7 - g_8)/5 (that is (-4 g_7eff - g_8eff)/5) = 0 and (5 g_3 - 5 g_4 + 4 g_5 - 4 g_6 - g_7 +
 ↪  g_8)/5 (that is (-g_7eff + g_8eff)/5) = 0 (4 condition(s))
[PASS] 44q-S12  (exact) internal symmetry of the fields that do not depend on the extra
 ↪  coordinates x0, x5, x6, x7 (their kinetic term contains alpha^1, alpha^2, alpha^3 and beta
 ↪  only). The eight Hermitian generators of the symmetry of the model with the Vorton-Scalar are
 ↪  the six rotations of the extra axes, (phase) gamma^a gamma^b with a < b in {0, 5, 6, 7}, the
 ↪  unit matrix (fermion number), and the generator P = (phase) gamma^0 gamma^5 gamma^6 gamma^7 of
 ↪  U(1)_E, with P^2 = 1 and trace 0; all commute with the four matrices of the kinetic term, and
 ↪  P commutes with the chirality matrix. The rotations of x5, x6, x7 among themselves commute
 ↪  with W (they act inside each half of the field); the three generators gamma^0 gamma^t and the
 ↪  generator P of U(1)_E anticommute with W (they connect the half W = +1 with the half W = -1).
 ↪  The self-interaction is invariant under the six rotations and the phase for all values of the
 ↪  nine couplings (0 conditions found), because the rotations are among the 28 generators of
 ↪  check S4. Under psi -> exp(i theta P) psi the invariants of the ranks [0, 4, 8] are invariant
 ↪  one by one, and the interaction sum_r g_r I_r is invariant exactly if g_1 + g_3 - g_5 - g_7
 ↪  (that is g_1eff - g_7eff) = 0 and g_2 + g_3 - g_6 - g_7 (that is -g_7eff) = 0 (2 condition(s))
[PASS] self-X1  (numerical) the example of the author's document, g_0 = g_S = -1e5 and g_1 = g_V =
 ↪  +1e4 with all other couplings zero, gives the Hartree-Fock couplings G_0 = -98750 and G_1 =
 ↪  +20000 (7+1 and the quantized-sector form of 4+4), the values stated in that document
```

A.4 The two forms on 4+4: matter of two halves and the vacuum (`deflation_cjt_ks_c forms`)

```
PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C: checks of the calculations
 ↪  for the two forms of the self-interaction on the 4+4 spacetime, C = covariant form (the
 ↪  definition) and Q = quantized-sector form (the alternative): uniform matter, effective
 ↪  potential, vacuum, Vorton-Scalar (fm-..)
generated by: deflation_cjt_ks_c (Rust crate spacetimes-deflation-cjt-ks-c, version 0.1.0;
 ↪  PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C); command:
 ↪  deflation_cjt_ks_c forms
9 of 9 checks passed

[PASS] fm-F1  (numerical) the mean-field couplings and the solver of two halves. (i) The four
 ↪  couplings G_S^HF = (15 g_0eff - 8 g_1eff - 8 g_7eff - g_8eff)/16, G_V^HF = (-g_0eff + 22
 ↪  g_1eff - 6 g_7eff + g_8eff)/16, G_Sc = (-g_0eff - 2 g_1eff + 2 g_7eff + g_8eff)/16, G_Vc =
 ↪  -(g_0eff + g_8eff)/16 computed from these closed forms equal those of the exact coefficient
 ↪  tables (checks 44c-S7, 44q-S7) for the sets of couplings used here: set A (scalar attraction
 ↪  alone: g_0 = -g, all other couplings zero): G_S^HF = -0.9375 g, G_V^HF = 0.0625 g, G_Sc =
 ↪  0.0625 g, G_Vc = 0.0625 g; set B (the ratio of the author's example: g_0 = -g, g_1 = +0.1 g,
 ↪  all other couplings zero): G_S^HF = -0.9875 g, G_V^HF = 0.2 g, G_Sc = 0.05 g, G_Vc = 0.0625 g.
 ↪  Covariant form C: G_Vtot = G_Vc, G_Stot = G_Sc (total densities), G_Vdif = G_V^HF, G_Sdif =
 ↪  G_S^HF (differences); quantized-sector form Q: the two pairs exchanged. (ii) The solver of the
 ↪  coupled gap equations (closed forms of the momentum integrals; all solutions) against the
 ↪  Kohn-Sham routines for matter of one kind of particle (Gauss-Legendre integrals), for 48 sets
 ↪  of form, dimension, couplings and Fermi momentum: with equal halves the symmetric solution is
 ↪  matter of eight states with the couplings G_Stot, G_Vtot (mass, energy density, pressure agree
 ↪  to 2.0e-15); with one half filled it is matter of four states with the couplings G_Ssame =
 ↪  G_Stot + G_Sdif, G_Vsame = G_Vtot + G_Vdif (2.2e-15). (iii) With unequal halves (k_- = 0.8
 ↪  k_+) the chemical potentials mu_+- = sqrt(k_+-^2 + M_+-^2) + V_+- are the derivatives of the
 ↪  energy density with respect to the densities of the halves (central differences: 2.7e-10), and
 ↪  the pressure from the momentum integrals equals mu_+ n_+ + mu_- n_- - eps (6.7e-16)
```

[PASS] fm-F2 (numerical) the binding thresholds of the two forms for the same couplings of the Lagrangian (fields that do not depend on x5, x6, x7; no-sea approximation; state of lowest energy with respect to the polarization): the smallest strength g = |g_0| at which matter is self-bound, verified in each case (bound 0.1 per cent above, not bound 0.1 per cent below): D = 4, set A, form C: g m^3 = 224.7386 (one half; k_F,total = 0.6140, M_+ = 0.5239, M_- = 1.5441); D = 4, set A, form Q: g m^3 = 82.1116 (equal halves; k_F,total = 0.8054, M_+ = 0.4946, M_- = 0.4946); D = 4, set B, form C: g m^3 = 332.2917 (one half; k_F,total = 0.5211, M_+ = 0.5625, M_- = 1.4842); D = 4, set B, form Q: g m^3 = 121.0858 (equal halves; k_F,total = 0.6792, M_+ = 0.5424, M_- = 0.5424); D = 3, set A, form C: g m^2 = 24.5717 (one half; k_F,total = 0.5374, M_+ = 0.6454, M_- = 1.4053); D = 3, set A, form Q: g m^2 = 9.8090 (equal halves; k_F,total = 0.7356, M_+ = 0.6276, M_- = 0.6276); D = 3, set B, form C: g m^2 = 31.1674 (one half; k_F,total = 0.4635, M_+ = 0.6712, M_- = 1.3638); D = 3, set B, form Q: g m^2 = 12.3819 (equal halves; k_F,total = 0.6351, M_+ = 0.6575, M_- = 0.6575). Where the bound state has equal halves or one half filled, the threshold equals the one computed for matter of one kind of particle by the routines of module kohn_sham (largest relative difference 5.7e-14). Ratio of the threshold of the covariant form to that of the quantized-sector form: D = 4, set A: 2.737; D = 4, set B: 2.744; D = 3, set A: 2.505; D = 3, set B: 2.517

[PASS] fm-F3 (numerical) the polarization of the two halves. For the same couplings of the Lagrangian (sets A and B, attractive scalar coupling g_0 < 0), 1.5, 3 and 6 times the strength at which the quantized-sector form first binds matter, and total densities of the Fermi momenta 0.4, 0.6, 0.8, 1.0 (48 cases): in the covariant form a particle is attracted by the particles of its own half and repelled by those of the other, and the state of lowest energy is polarized in 48 of the 48 cases; in the quantized-sector form it is polarized in 0 cases. Polarization (n_+ - n_-)/n of the state of lowest energy in the covariant form (four densities in the order given): D = 4, set A, 1.5 times: 1.00, 1.00, 1.00, 1.00; D = 4, set A, 3 times: 1.00, 1.00, 1.00, 1.00; D = 4, set A, 6 times: 1.00, 1.00, 1.00, 0.78; D = 4, set B, 1.5 times: 1.00, 1.00, 1.00, 0.35; D = 4, set B, 3 times: 1.00, 1.00, 0.65, 0.22; D = 4, set B, 6 times: 1.00, 1.00, 0.39, 0.19; D = 3, set A, 1.5 times: 1.00, 1.00, 1.00, 1.00; D = 3, set A, 3 times: 1.00, 1.00, 1.00, 1.00; D = 3, set A, 6 times: 1.00, 1.00, 1.00, 0.89; D = 3, set B, 1.5 times: 1.00, 1.00, 1.00, 0.45; D = 3, set B, 3 times: 1.00, 1.00, 0.81, 0.31; D = 3, set B, 6 times: 1.00, 1.00, 0.50, 0.22. Example: D = 4, set B, three times the threshold of the quantized-sector form, total density of the Fermi momentum 0.8 m: energy per particle minus m = +0.3287 m in the covariant form (polarization 0.65) and +0.1323 m in the quantized-sector form (polarization 0.00). Existence of the states: the energy was computed at 41 polarizations in each case and form (3936 states). In the covariant form every one of these states exists. In the quantized-sector form 108 states with unequal halves do not exist (D = 4, set A, form Q, 6 times, k_F,total = 1 m: 39 of 41; D = 4, set B, form Q, 3 times, k_F,total = 1 m: 2 of 41; D = 4, set B, form Q, 6 times, k_F,total = 0.8 m: 4 of 41; D = 4, set B, form Q, 6 times, k_F,total = 1 m: 31 of 41; D = 3, set A, form Q, 6 times, k_F,total = 1 m: 15 of 41; D = 3, set B, form Q, 6 times, k_F,total = 1 m: 17 of 41): the gap equations have no solution with non-negative masses, because the attraction by the more populated half (coupling G_Sopp between the halves, stronger than the coupling G_Ssame inside a half) would make the mass of the particles of the less populated half negative; this happens only at 3 or 6 times the threshold and k_F,total >= 0.8 m, and the states with equal halves and with one half filled exist in all cases

[PASS] fm-F4 (numerical) self-bound matter along the deflation clock in the two forms (clock counted from the moment at which matter of the respective form first binds; 101 clock readings from 0 to 5 in each of the 8 cases, the couplings growing to 148 times their value at the threshold): the energy per particle of the saturated state falls monotonically, its pressure vanishes (largest |p|/eps after the first two readings: 2.8e-8), and the kind of the state does not change: one half filled in the covariant form, equal halves in the quantized-sector form. D = 4, set A, form C (one half): E/N - m = -0.1092 m at tau_d = 1 and -0.2913 m at tau_d = 3 (mass of the particles of the filled half or halves 0.351 m and 0.183 m); D = 4, set A, form Q (equal halves): E/N - m = -0.1284 m at tau_d = 1 and -0.3438 m at tau_d = 3 (mass of the particles of the filled half or halves 0.319 m and 0.156 m); D = 4, set B, form C (one half): E/N - m = -0.0829 m at tau_d = 1 and -0.2209 m at tau_d = 3 (mass of the particles of the filled half or halves 0.392 m and 0.215 m); D = 4, set B, form Q (equal halves): E/N - m = -0.0968 m at tau_d = 1 and -0.2580 m at tau_d = 3 (mass of the particles of the filled half or halves 0.371 m and 0.199 m); D = 3, set A, form C (one half): E/N - m = -0.1226 m at tau_d = 1 and -0.3365 m at tau_d = 3 (mass of the particles of the filled half or halves 0.373 m and 0.157 m); D = 3, set A, form Q (equal halves): E/N - m = -0.1424 m at tau_d = 1 and -0.3945 m at tau_d = 3 (mass of the particles of the filled half or halves 0.347 m and 0.137 m); D = 3, set B, form C (one half): E/N - m = -0.0940 m at tau_d = 1 and -0.2567 m at tau_d = 3 (mass of the particles of the filled half or halves 0.407 m and 0.183 m); D = 3, set B, form Q (equal halves): E/N - m = -0.1093 m
at tau_d = 1 and -0.2990 m at tau_d = 3 (mass of the particles of the filled half or halves 0.389 m and 0.170 m). Regularity: the largest eigenvalue of the response matrix of the two gap equations stays below 1 at all readings (largest value 0.881), so that each state is the regular continuation of the dilute gas. Beyond the table the saturated state of the quantized-sector form, which has equal halves, reaches the eigenvalue 1 (2 G_Sdif ds/dM = 1: the two masses can split) at: D = 4, set A: tau_d = 6.416; D = 4, set B: not before tau_d = 9; D = 3, set A: tau_d = 5.296; D = 3, set B: tau_d = 7.323; the saturated state of the covariant form has one half filled and a negative eigenvalue throughout

[PASS] fm-F5 (numerical) matter of fixed density while the couplings grow with the deflation clock (clock counted from the binding threshold of the respective form; total densities of the Fermi momenta 0.3 m and 0.6 m; D = 4 and D = 3; sets A and B). In the covariant form the matter polarizes at a definite clock reading before the binding threshold: the two halves become unequal, and within at most 0.35 of the clock one half is empty; where the pressure along the coordinates of the motion becomes negative, it does so after the polarization, and in 3 of the 8 cases (the higher density) the pressure stays positive for -3 <= tau_d <= 5, because the repulsion G_Vsame between the particles of the filled half grows with the couplings while the attraction saturates as the mass falls. In the quantized-sector form the halves stay equal and the pressure becomes negative in all 8 cases. D = 4, set A, form C, k_F,total = 0.3 m: halves unequal from tau_d = -0.52, one half empty from -0.17; pressure negative from tau_d = 0.31; D = 4, set A, form C, k_F,total = 0.6 m: halves unequal from tau_d = -1.77, one half empty from -1.42; pressure negative from tau_d = -0.43; D = 4, set A, form Q, k_F,total = 0.3 m: halves equal throughout; pressure negative from tau_d = 0.78; D = 4, set A, form Q, k_F,total = 0.6 m: halves equal throughout; pressure negative from tau_d = -0.36; D = 4, set B, form C, k_F,total = 0.3 m: halves unequal from tau_d = -0.82, one half empty from -0.47; pressure negative from tau_d = 0.06; D = 4, set B, form C, k_F,total = 0.6 m: halves unequal from tau_d = -1.98, one half empty from -1.62; pressure positive throughout; D = 4, set B, form Q, k_F,total = 0.3 m: halves equal throughout; pressure negative from tau_d = 0.51; D = 4, set B, form Q, k_F,total = 0.6 m:
halves equal throughout; pressure negative from tau_d = -0.49; D = 3, set A, form C, k_F,total = 0.3 m: halves unequal from tau_d = -0.92, one half empty from -0.72; pressure negative from tau_d = -0.03; D = 3, set A, form C, k_F,total = 0.6 m: halves unequal from tau_d = -1.48, one half empty from -1.27; pressure positive throughout; D = 3, set A, form Q, k_F,total = 0.3 m: halves equal throughout; pressure negative from tau_d = 0.21; D = 3, set A, form Q, k_F,total = 0.6 m: halves equal throughout; pressure negative from tau_d = -0.21; D = 3, set B, form C, k_F,total = 0.3 m: halves unequal from tau_d = -1.02, one half empty from -0.88; pressure negative from tau_d = -0.12; D = 3, set B, form C, k_F,total = 0.6 m: halves unequal from tau_d = -1.52, one half empty from -1.32; pressure positive throughout; D = 3, set B, form Q, k_F,total = 0.3 m: halves equal throughout; pressure negative from tau_d = 0.10; D = 3, set B, form Q, k_F,total = 0.6 m: halves equal throughout; pressure negative from tau_d = -0.14

[PASS] fm-F6 (numerical) the Kohn-Sham method and the CJT effective potential agree for the two halves, in both forms: for 80 states (two dimensions, two sets, two forms, strengths 1.5 and 3 times the respective binding threshold, total densities of the Fermi momenta 0.3 to 1.1 m; 40 with equal halves, 17 with unequal halves, 23 with one half filled) the stationary point of Omega(M_+, M_-, V_+, V_-; mu) at the chemical potential of the Kohn-Sham state, found by Newton's method from displaced start values, has the same two masses, the same density and energy density, and -Omega equals the Kohn-Sham pressure (sum of the deviations at most 4.9e-11; residuals of the four stationarity conditions at most 1.4e-12). In the states with unequal halves the two chemical potentials agree to 6.7e-16 (relative): the polarization is the equilibrium one. In all 23 states with one half filled the lowest state of the empty half lies above the chemical potential

[PASS] fm-F7 (numerical) the vacuum of the two forms with the filled states of negative energy (cutoff Lambda). The gap equations of the vacuum are (M_+ + M_-)/2 - m = G_Stot (s_+ + s_-) and (M_+ - M_-)/2 = G_Sdif (s_+ - s_-) with s_+- = -(NU/2) Int^Lambda M_+-/w. (a) Without mass term (m = 0) and for an attractive scalar coupling (sets A and B: G_S^HF < 0 < G_Sc) a field acquires a mass when |G_S^HF| exceeds the critical coupling |G_c| (D = 4: G_c Lambda^3 = -29.608813, D = 3: G_c Lambda^2 = -4.934802), in both forms at the same strength of the couplings: no mass 2 per cent below it, a mass 2 per cent above it. In the quantized-sector form the two halves acquire the SAME mass (M_+ = M_-: the mass term of the positive Lagrangian); in the covariant form they acquire masses of OPPOSITE sign (M_+ = -M_-: the mass term -m psitilde psi of the covariant Lagrangian), verified to 1e-7. (b) With the bare mass 0.05 Lambda the quantized-sector form passes smoothly through the critical strength (the two masses stay equal at all 101 clock readings), whereas the covariant form has a sharp transition at which the two masses separate: D = 4, set A: tau_d = 0.0091 (linear criterion 0.0091); D = 4, set B: tau_d = 0.0094 (linear criterion 0.0094); D = 3, set A: tau_d = 0.0194 (linear criterion 0.0194); D = 3, set B: tau_d = 0.0199 (linear criterion 0.0199); the strength of the transition agrees with the linear criterion 1 + G_Sdif NU Int^Lambda d^Dk/(2 pi)^D k^2/w^3 = 0 at the common mass to 2.1e-10

[PASS] fm-F8 (numerical) bare Dirac mass, self-interaction and Vorton-Scalar together, in the two forms (D = 3, no-sea approximation): the smallest C at which matter is self-bound, for f_D = 0, 0.25, 0.5 and the contact couplings g_0 = -kappa_0 C, g_1 = +kappa_1 C. Without self-interaction both forms give the threshold of the model with the Vorton-Scalar alone, C = 3.70110 and 3.70110 against 3 pi^2/8 = 3.70110. In 18 of the 18 cases the threshold is that of matter of one kind (the smaller of the thresholds of matter with equal halves and of matter with one half filled, functional of module kohn_sham), confirmed by the functional of two halves minimized over density, field and polarization: at 1.001 times the threshold its lowest state is bound (E/N - m <= -1.6e-4 m) and is of the expected kind, at 0.999 times the threshold no state is bound (E/N - m >= 1.6e-4 m); in 0 cases the threshold was found by bisection with the functional of two halves (a bound state with unequal halves). With self-interaction: f_D = 0, kappa_0 = 1, kappa_1 = 0, form C: C = 3.9399 (shift +0.063; equal halves, x = 0.000); f_D = 0, kappa_0 = 1, kappa_1 = 0, form Q: C = 3.7156 (shift +0.004; equal halves, x = 0.614); f_D = 0, kappa_0 = 1, kappa_1 = 0.1, form C: C = 3.9399 (shift +0.063; equal halves, x = 0.000); f_D = 0, kappa_0 = 1, kappa_1 = 0.1, form Q: C = 4.1527 (shift +0.115; equal halves, x = 0.652); f_D = 0.25, kappa_0 = 1, kappa_1 = 0, form C: C = 2.6336 (shift -0.340; equal halves, x = 0.000); f_D = 0.25, kappa_0 = 1, kappa_1 = 0, form Q: C = 2.4758 (shift -0.402; equal halves, x = 0.000); f_D = 0.25, kappa_0 = 1, kappa_1 = 0.1, form C: C = 2.6307 (shift -0.341; equal halves, x = 0.000); f_D = 0.25, kappa_0 = 1, kappa_1 = 0.1, form Q: C = 2.6992 (shift -0.316; equal halves, x = 0.000);
f_D = 0.5, kappa_0 = 1, kappa_1 = 0, form C: C = 2.1617 (shift -0.538; equal halves, x = 0.000); f_D = 0.5, kappa_0 = 1, kappa_1 = 0, form Q: C = 1.8252 (shift -0.707; equal halves, x = 0.000); f_D = 0.5, kappa_0 = 1, kappa_1 = 0.1, form C: C = 2.1559 (shift -0.540; equal halves, x = 0.000); f_D = 0.5, kappa_0 = 1, kappa_1 = 0.1, form Q: C = 1.9335 (shift -0.649; equal halves, x = 0.000)

```
[PASS] fm-F9  (numerical) a scalar coupling of the opposite sign, g_0 > 0 alone. In the covariant
 ↪  form particles of different halves attract each other; the coupling of the square of the total
 ↪  density is G_Vtot = G_Vc = -g_0/16 < 0, and the energy per particle of uniform matter falls
 ↪  without bound as the density grows: the contact model then has no saturated state without a
 ↪  cutoff. In the quantized-sector form the same coupling repels (G_Stot = 15 g_0/16 > 0) and the
 ↪  energy per particle rises with the density. D = 4, g_0 = +100 m^-3, form C: E/N - m = +0.089,
 ↪  -1.518, -35.819, -641.710 m at k_F,total = 1, 2, 4, 8 m; D = 4, g_0 = +100 m^-3, form Q: E/N -
 ↪  m = +1.095, +15.250, +243.184, +3890.737 m at k_F,total = 1, 2, 4, 8 m; D = 3, g_0 = +15 m^-2,
 ↪  form C: E/N - m = +0.158, -0.054, -5.220, -58.160 m at k_F,total = 1, 2, 4, 8 m; D = 3, g_0 =
 ↪  +15 m^-2, form Q: E/N - m = +0.929, +6.205, +48.701, +389.107 m at k_F,total = 1, 2, 4, 8 m
```

A.5 The solutions mode by mode (`deflation_cjt_ks_c modes`)

```
PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C: checks of the solutions of
 ↪  the Dirac equations on the deflating spacetimes (71d-M..: pair creation on 7+1; 44d-M..:
 ↪  bounded solutions on 4+4)
generated by: deflation_cjt_ks_c (Rust crate spacetimes-deflation-cjt-ks-c, version 0.1.0;
 ↪  PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C); command:
 ↪  deflation_cjt_ks_c modes
7 of 7 checks passed

[PASS] 71d-M1  (numerical) pair creation by the deflation, one direction: for E_perp/g55 = 0.25,
 ↪  0.5, .., 3.0 the number of particles created per spin state in a mode that starts empty agrees
 ↪  with 1/(exp(2 pi E_perp/g55) + 1) to a relative deviation of at most 3.2e-6 (the numbers range
 ↪  from 0.17 to 6.5e-9); the eight spin states are created equally; the evolved states stay
 ↪  orthonormal to 2.0e-13
[PASS] 71d-M2  (numerical) equal rates g55 = g66 = g77 = 1.3 (constants 1, 0.7, 2), mode with m =
 ↪  1, p = (0, 0.1, 0.2, 0.3), (k_5, k_6, k_7) = (0.5, -0.8, 0.3): the created number per spin
 ↪  state is 5.70627486e-3 for all eight states, the law 1/(exp(2 pi E_perp/g55) + 1) gives
 ↪  5.70627482e-3 (relative deviation 6.7e-9): with equal rates the result depends neither on the
 ↪  direction of k_t nor on the constants
[PASS] 71d-M3  (numerical) unequal rates (g55 = 1 with g66 = 2, 1.2 or 0.5), mode with wavenumbers
 ↪  along x5 and x6: whenever the second physical wavenumber is still below exp(-3) E_perp at the
 ↪  moment at which the first reaches E_perp, the mean created number is the Fermi-Dirac law with
 ↪  the rate of the coordinate that becomes energetic first, to within 1.2 per cent; when the two
 ↪  become energetic together the eight spin states are created unequally (largest ratio of the
 ↪  largest to the smallest eigenvalue in the scan: 110) and no closed form is known
 ↪  (data/creation_anisotropic.csv)
[PASS] 44d-M1  (numerical) modes of the 4+4 field that depend on the deflating times (five cases:
 ↪  one, two or three non-zero k_t; equal and unequal rates and constants): beyond the turning
 ↪  value of x4 exactly eight of the sixteen solutions stay bounded; integrated backwards to early
 ↪  times they contain positive- and negative-energy parts of equal weight (relative difference of
 ↪  the two Gram matrices at most 2.0e-14): they are standing waves in x4, not travelling particles
[PASS] 44d-M2  (numerical) the covariant charge psihat^dagger W psihat of every bounded solution
 ↪  vanishes (ratio to psihat^dagger psihat at most 1.0e-14), as it must for a conserved quantity
 ↪  of a solution that dies out
[PASS] 44d-M3  (numerical) between the turning value and the late x4 every bounded solution falls
 ↪  by exp(-S) times a factor of order one, S = Int sqrt(k_d^2 - E_perp^2) dx4 = 30 being the WKB
 ↪  exponent (largest difference between the logarithm of the factor and S in the five cases:
 ↪  0.53). Where one rate governs (cases 1 to 4) the eight solutions fall by the same factor
 ↪  (spread of the logarithms at most 1.1e-14) and its logarithm is S + (1/2) ln 2 - (1/2) ln(1 +
 ↪  exp(-2 pi E_perp/rate_t)), as the exact solution requires (largest deviation 3.6e-4); with
 ↪  three unequal rates (case 5) the factor depends on the spin state (spread 0.40). The
 ↪  dependence on a deflating time dies out as exp(-S), and S grows as exp(rate_t x4)
[PASS] 44d-M4  (numerical) where one rate governs (one non-zero k_t, or equal rates) the computed
 ↪  |psihat|^2 equals the exact form 2 u_d |K_(1/2 + i E_perp/rate_t)(u_d)|^2 (modified Bessel
 ↪  function of the order 1/2 + i E_perp/rate_t, u_d = k_d/rate_t) over the range 0.02 < u_d < 12
 ↪  with a largest relative deviation of 3.6e-12
[PASS] ALL  (numerical) every check above passed
```

A.6 Symmetries (`deflation_cjt_ks_c symmetry`)

```
PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C: checks of the symmetries
 ↪  on the deflating spacetimes (isometries, internal symmetry of the fields that do not depend on
 ↪  the extra coordinates, zero modes on a vortex line, reduction of the couplings)
generated by: deflation_cjt_ks_c (Rust crate spacetimes-deflation-cjt-ks-c, version 0.1.0;
 ↪  PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C); command:
 ↪  deflation_cjt_ks_c symmetry
19 of 19 checks passed

[PASS] 44d-S1  (exact) of the 28 rotations and boosts of the flat 4+4 spacetime (one for each pair
 ↪  of coordinates) exactly 6 remain isometries of the deflating spacetime for three different
 ↪  rates (the rotations of x0, x1, x2, x3), 7 if two rates are equal (in addition the rotation of
 ↪  the two coordinates that deflate at the same rate) and 9 if the three rates are equal (in
 ↪  addition the three rotations of x5, x6, x7); no transformation that involves x4 survives, and
 ↪  none that mixes x0 with a deflating coordinate. Together with the 7 translations, the scaling
 ↪  and, for equal rates, the three special vectors of metric.rs this gives 14, 15 and 20 Killing
 ↪  vectors (Lie derivative of the metric zero at two points, exact arithmetic)
[PASS] 44d-S2  (exact) the products gamma^A gamma^B that commute with the five matrices alpha^1,
 ↪  alpha^2, alpha^3, beta, i beta chi (and so act as internal symmetries on fields that do not
 ↪  depend on the extra coordinates) are exactly those of the six pairs taken from x0, x5, x6, x7.
 ↪  Of these six, the deflating spacetime keeps as isometries 0 (three different rates), 1 (two
 ↪  equal rates) and 3 (three equal rates): the remaining 6, 5 and 3 are symmetries of the sector
 ↪  of such fields that have no counterpart among the isometries of the spacetime (hidden
 ↪  symmetries)
[PASS] 71d-S1  (exact) of the 28 rotations and boosts of the flat 7+1 spacetime (one for each pair
 ↪  of coordinates) exactly 6 remain isometries of the deflating spacetime for three different
 ↪  rates (the rotations of x0, x1, x2, x3), 7 if two rates are equal (in addition the rotation of
 ↪  the two coordinates that deflate at the same rate) and 9 if the three rates are equal (in
 ↪  addition the three rotations of x5, x6, x7); no transformation that involves x4 survives, and
 ↪  none that mixes x0 with a deflating coordinate. Together with the 7 translations, the scaling
 ↪  and, for equal rates, the three special vectors of metric.rs this gives 14, 15 and 20 Killing
 ↪  vectors (Lie derivative of the metric zero at two points, exact arithmetic)
[PASS] 71d-S2  (exact) the products gamma^A gamma^B that commute with the five matrices alpha^1,
 ↪  alpha^2, alpha^3, beta, i beta chi (and so act as internal symmetries on fields that do not
 ↪  depend on the extra coordinates) are exactly those of the six pairs taken from x0, x5, x6, x7.
 ↪  Of these six, the deflating spacetime keeps as isometries 0 (three different rates), 1 (two
 ↪  equal rates) and 3 (three equal rates): the remaining 6, 5 and 3 are symmetries of the sector
 ↪  of such fields that have no counterpart among the isometries of the spacetime (hidden
 ↪  symmetries)
[PASS] def-S3  (numerical) the internal symmetry of the fields that do not depend on the extra
 ↪  coordinates, computed again for both spacetimes (24 items of data/reduced_symmetry.csv, all
 ↪  as expected): the commutant of the five matrices has dimension 8 and is u(2) + u(2) = su(2)_A
 ↪  + su(2)_B + u(1)_F + u(1)_E; the eight one-particle states are (2,1) + (1,2); the four vector
 ↪  pair operators are the (2,2) with fermion number 2, and their invariant quadratic form is the
 ↪  frame metric of the axes x0, x5, x6, x7. The deflation does not change this symmetry, because
 ↪  the weighted field of this sector obeys the flat equations (checks 44d-D.., 71d-D.. of the
 ↪  command matrices)
[PASS] 44d-V1  (exact) alpha^1, alpha^2, alpha^3, beta and i beta chi are Hermitian, have square
 ↪  one and anticommute with one another
[PASS] 44d-V2  (exact) i beta alpha^1 and alpha^2 beta chi are commuting Hermitian matrices with
 ↪  square one
[PASS] 44d-V3  (exact) each of the four joint eigenspaces of the two matrices has dimension 4: a
 ↪  vortex line of unit winding carries four zero modes of the sixteen-component field
[PASS] 44d-V4  (exact) Pc = i alpha^1 alpha^2 alpha^3 chi equals the product of the five matrices,
 ↪  commutes with each of them, is Hermitian and traceless and has square one: it is, up to sign,
 ↪  the generator P of U(1)_E
[PASS] 44d-V5  (exact) winding +1: every w_z with (i beta alpha^1) w_z = -w_z, (alpha^2 beta chi)
 ↪  w_z = -w_z solves the zero-energy equations (i alpha^1 + beta) w_z = 0, (i alpha^2 + i beta
 ↪  chi) w_z = 0; on these four spinors alpha^3 = Pc, and its trace vanishes: two zero modes have
 ↪  alpha^3 = +1 (energy +k_3) and two alpha^3 = -1 (energy -k_3)
```

```
[PASS] 44d-V6  (exact) winding -1: the spinors w_z with (i beta alpha^1) w_z = -w_z, (alpha^2 beta
 ↪  chi) w_z = +w_z solve (i alpha^1 + beta) w_z = 0, (i alpha^2 - i beta chi) w_z = 0, and on
 ↪  them alpha^3 = -Pc: the directions of motion are exchanged
[PASS] 71d-V1  (exact) alpha^1, alpha^2, alpha^3, beta and i beta chi are Hermitian, have square
 ↪  one and anticommute with one another
[PASS] 71d-V2  (exact) i beta alpha^1 and alpha^2 beta chi are commuting Hermitian matrices with
 ↪  square one
[PASS] 71d-V3  (exact) each of the four joint eigenspaces of the two matrices has dimension 4: a
 ↪  vortex line of unit winding carries four zero modes of the sixteen-component field
[PASS] 71d-V4  (exact) Pc = i alpha^1 alpha^2 alpha^3 chi equals the product of the five matrices,
 ↪  commutes with each of them, is Hermitian and traceless and has square one: it is, up to sign,
 ↪  the generator P of U(1)_E
[PASS] 71d-V5  (exact) winding +1: every w_z with (i beta alpha^1) w_z = -w_z, (alpha^2 beta chi)
 ↪  w_z = -w_z solves the zero-energy equations (i alpha^1 + beta) w_z = 0, (i alpha^2 + i beta
 ↪  chi) w_z = 0; on these four spinors alpha^3 = Pc, and its trace vanishes: two zero modes have
 ↪  alpha^3 = +1 (energy +k_3) and two alpha^3 = -1 (energy -k_3)
[PASS] 71d-V6  (exact) winding -1: the spinors w_z with (i beta alpha^1) w_z = -w_z, (alpha^2 beta
 ↪  chi) w_z = +w_z solve (i alpha^1 + beta) w_z = 0, (i alpha^2 - i beta chi) w_z = 0, and on
 ↪  them alpha^3 = -Pc: the directions of motion are exchanged
[PASS] def-S4  (exact) reduction to fields that do not depend on x5, x6, x7: sqrt|g| times the
 ↪  Yukawa term and the potential of the Vorton-Scalar, written with the weighted fields psihat =
 ↪  sqrt(Vd) psi and Phihat = sqrt(Vd) Phi_VS, are the same expressions with the couplings
 ↪  y_VS/sqrt(Vd), lambda_VS/Vd and the vacuum value v_VS sqrt(Vd); the products y_VS v_VS and
 ↪  lambda_VS v_VS^2, hence the fermion mass and the mass of the radial mode, do not change
 ↪  (rational sample values, exact). The exponents of data/coupling_exponents.csv follow, with Vd
 ↪  proportional to exp(-gsum x4)
[PASS] def-S5  (exact) reduction to fields that do not depend on x5, x6, x7, mass term and
 ↪  self-interaction of the fermion field: sqrt|g| times [-m_D psibar psi - (1/2) g_r (psibar
 ↪  Theta_K psi)(psibar Theta^K psi)], written with the weighted field psihat = sqrt(Vd) psi, is
 ↪  the same expression with the unchanged bare Dirac mass m_D and with the coupling g_r/Vd; every
 ↪  coupling of the self-interaction therefore grows as exp(tau_d), as g_VS^2 does, and the ratio
 ↪  of the contact couplings to g_VS^2 is constant (rational sample values, three sets)
[PASS] ALL  (exact) every check above passed
```

A.7 The deflating spacetime as a solution of equations of gravity (`deflation_cjt_ks_c gravity`)

```
PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C: checks of the
 ↪  Einstein-Lovelock equations with Kohn-Sham matter (gr-..)
generated by: deflation_cjt_ks_c (Rust crate spacetimes-deflation-cjt-ks-c, version 0.1.0;
 ↪  PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C); command:
 ↪  deflation_cjt_ks_c gravity
6 of 6 checks passed

[PASS] gr-G1  (numerical) the Einstein-Lovelock tensors of a spacetime with two scale factors,
 ↪  computed from the pair curvatures K_(4i) = H_i' + H_i^2, K_(ij) = H_i H_j by E^(k)m_m =
 ↪  (-1)^(k+1) k! 2^(k-1) times the sum over the sets of 2k directions other than m of the sum
 ↪  over their divisions into pairs of the product of the pair curvatures: for 3 + 4 directions
 ↪  (the cosmological spacetime of the author's document KOHN_SHAM_GRAVITY) the nine components
 ↪  E^(1), E^(2), E^(3) (time, ordinary space, auxiliary space), each with its coefficients of
 ↪  H_a' and H_b', equal the 21 polynomials printed in that document at four pairs of rates
 ↪  (largest deviation 2.6e-15)
[PASS] gr-G2  (numerical) the self-consistent evolution of the author's document is reproduced:
 ↪  with his data (kappa = 1000, LambdaBar = 0, H_a(0) = 0.05, a(0) = b(0) = 1, Kohn-Sham matter
 ↪  of the saturation density) the constraint gives H_b(0), and the integration to x4 = 20 gives
 ↪  a, b, H_a, H_b; for Einstein gravity and for Einstein-Lovelock gravity with alpha_2 = 20,
 ↪  alpha_3 = 500 the ten numbers agree with the printed ones to 4.8e-8 (relative); the constraint
 ↪  is preserved to 1.3e-14. The conservation law eps' + 3 H_a (eps + p_a) + 4 H_b (eps + p_b) = 0
 ↪  holds at the test point of the document (a = 1 + 0.3 x + 0.05 x^2, b = 1/(1 + 0.2 x), x = 0.7)
 ↪  with the relative residual -9.2e-10
```

```
[PASS] gr-G3  (numerical) the deflating spacetime in the author's Einstein-Lovelock gravity (four
 ↪  flat coordinates x0..x3 with H_a = 0, three deflating coordinates with equal rates, H_b =
 ↪  -g55): (i) it solves the three field equations without matter for LambdaBar = 3 g55^2 and
 ↪  alpha_2 = 1/(4 g55^2) (residuals 5.4e-16 g55^2; alpha_3 is arbitrary because E^(3) vanishes),
 ↪  in agreement with the exact checks 44d-G10 and 71d-G10; (ii) at this solution the two
 ↪  equations for the accelerations are -6 H_a' + .. and +2 H_b' + .. (deviation of the matrix of
 ↪  coefficients from diag(-6, 2): 2.2e-15): the evolution is well defined; (iii) in Einstein
 ↪  gravity without the Gauss-Bonnet term and without LambdaBar the same spacetime needs a source
 ↪  with kappa eps = 3.0 g55^2, kappa p_a = -6.0 g55^2, kappa p_b = -3.0 g55^2, that is p_a = -2
 ↪  eps along the flat coordinates and p_b = -eps along the deflating ones. Kohn-Sham matter
 ↪  cannot be this source: its pressure along a coordinate on which the fields do not depend is Pi
 ↪  = (1/2) G_V rho^2 + (M - m)^2/(2 G_S), and Pi = -eps would require e_kin + G_V rho^2 = 0
[PASS] gr-G6  (numerical) the matter of the two forms of the 4+4 model as a source of gravity
 ↪  (fields that do not depend on x5, x6, x7, D = 4; couplings of the Lagrangian g_0 = -g, g_1 =
 ↪  +0.1 g; state of lowest energy per particle over all densities and polarizations, module
 ↪  halves). covariant form (the definition): threshold g m^3 = 332.2917; at three times that
 ↪  strength the saturated state has one half filled (Fermi momentum of the filled states 0.52859
 ↪  m, mass 0.37920 m, E/N - m = -0.09072 m, eps = 8.990698e-4 m^5, interaction pressure Pi =
 ↪  -7.826926e-5 m^5); as matter of one kind: 4 states, G_S = -934.5703, G_V = 261.6797.
 ↪  quantized-sector form (the alternative): threshold g m^3 = 121.0858; at three times that
 ↪  strength the saturated state has equal halves (Fermi momentum of the filled states 0.58171 m,
 ↪  mass 0.35844 m, E/N - m = -0.10601 m, eps = 2.592903e-3 m^5, interaction pressure Pi =
 ↪  -2.681363e-4 m^5); as matter of one kind: 8 states, G_S = -358.7168, G_V = 72.6515. The same
 ↪  states computed by the Kohn-Sham routines for matter of one kind of particle (module
 ↪  kohn_sham, other integrals) agree in mass, energy density, pressure and interaction pressure
 ↪  to 1.3e-15. In an evolution without collisions every particle keeps its state, so that a half
 ↪  that is empty stays empty: the source is then matter of one kind of particle throughout
[PASS] gr-G4  (numerical) the deflating spacetime with Kohn-Sham matter as a source (rate g55 =
 ↪  0.01 m; initial data a = b = 1, H_a = 0, H_b from the constraint; loading = kappa eps(0)/(3
 ↪  g55^2)): matter makes the three coordinates deflate FASTER than exponentially and makes the
 ↪  four flat coordinates EXPAND; in 8 of the 16 evolutions the rates exceed 50 g55 before the
 ↪  clock of the undisturbed deflation reads 6 (they grow without bound in a finite time). The
 ↪  constraint is preserved along every evolution to 6.7e-10 (relative). Results: [7+1, all
 ↪  momenta; tuned Einstein-Lovelock; loading 1e-3] H_b(0) = -1.00050 g55; at tau_ref = 6.00: a =
 ↪  1.2508, b = 4.908e-2, H_a = +1.4052 g55, H_b = -7.9908 g55 | [7+1, all momenta; tuned
 ↪  Einstein-Lovelock; loading 1e-1] H_b(0) = -1.04881 g55; the rates run away; at tau_ref = 3.58:
 ↪  a = 1.5647, b = 2.320e-2, H_a = +8.9453 g55, H_b = -50.0355 g55 | [7+1, all momenta; Einstein;
 ↪  loading 1e-3] H_b(0) = -0.03162 g55; at tau_ref = 6.00: a = 1.0011, b = 9.335e-1, H_a =
 ↪  +0.0012 g55, H_b = -0.0375 g55 | [7+1, all momenta; Einstein; loading 1e-1] H_b(0) = -0.31623
 ↪  g55; the rates run away; at tau_ref = 4.17: a = 1.5278, b = 9.199e-2, H_a = +11.2603 g55, H_b
 ↪  = -50.0924 g55 | [sector without momenta along x5, x6, x7, mean-field couplings; tuned
 ↪  Einstein-Lovelock; loading 1e-3] H_b(0) = -1.00050 g55; at tau_ref = 6.00: a = 1.2333, b =
 ↪  5.683e-2, H_a = +1.1823 g55, H_b = -6.6163 g55 | [sector without momenta along x5, x6, x7,
 ↪  mean-field couplings; tuned Einstein-Lovelock; loading 1e-1] H_b(0) = -1.04881 g55; the rates
 ↪  run away; at tau_ref = 3.74: a = 1.6348, b = 2.202e-2, H_a = +9.1382 g55, H_b = -50.0730 g55 |
 ↪  [sector without momenta along x5, x6, x7, mean-field couplings; Einstein; loading 1e-3] H_b(0)
 ↪  = -0.03162 g55; at tau_ref = 6.00: a = 1.0016, b =
```

```
9.331e-1, H_a = +0.0017 g55, H_b = -0.0380 g55 | [sector without momenta along x5, x6, x7,
 ↪  mean-field couplings; Einstein; loading 1e-1] H_b(0) = -0.31623 g55; the rates run away; at
 ↪  tau_ref = 4.05: a = 1.7118, b = 8.801e-2, H_a = +13.1968 g55, H_b = -50.2782 g55 | [4+4
 ↪  covariant form; tuned Einstein-Lovelock; loading 1e-3] H_b(0) = -1.00050 g55; at tau_ref =
 ↪  6.00: a = 1.5963, b = 1.521e-2, H_a = +6.2500 g55, H_b = -34.1042 g55 | [4+4 covariant form;
 ↪  tuned Einstein-Lovelock; loading 1e-1] H_b(0) = -1.04881 g55; the rates run away; at tau_ref =
 ↪  3.51: a = 1.6307, b = 2.463e-2, H_a = +9.1436 g55, H_b = -50.0904 g55 | [4+4 covariant form;
 ↪  Einstein; loading 1e-3] H_b(0) = -0.03162 g55; at tau_ref = 6.00: a = 1.0014, b = 9.332e-1,
 ↪  H_a = +0.0015 g55, H_b = -0.0378 g55 | [4+4 covariant form; Einstein; loading 1e-1] H_b(0) =
 ↪  -0.31623 g55; the rates run away; at tau_ref = 4.01: a = 1.5871, b = 9.687e-2, H_a = +11.7156
 ↪  g55, H_b = -50.0303 g55 | [4+4 quantized-sector form; tuned Einstein-Lovelock; loading 1e-3]
 ↪  H_b(0) = -1.00050 g55; at tau_ref = 6.00: a = 1.4087, b = 2.928e-2, H_a = +2.8489 g55, H_b =
 ↪  -15.6064 g55 | [4+4 quantized-sector form; tuned Einstein-Lovelock; loading 1e-1] H_b(0) =
 ↪  -1.04881 g55; the rates run away; at tau_ref = 3.57: a = 1.6321, b = 2.386e-2, H_a = +9.1359
 ↪  g55, H_b = -50.0502 g55 | [4+4 quantized-sector form; Einstein; loading 1e-3] H_b(0) =
 ↪  -0.03162 g55; at tau_ref = 6.00: a = 1.0015, b = 9.332e-1, H_a = +0.0016 g55, H_b = -0.0379
 ↪  g55 | [4+4 quantized-sector form; Einstein; loading 1e-1] H_b(0) = -0.31623 g55; the rates run
 ↪  away; at tau_ref = 4.02: a = 1.6299, b = 9.367e-2, H_a = +12.2541 g55, H_b = -50.1237 g55
[PASS] gr-G5  (numerical) the deflating vacuum solution of the tuned Einstein-Lovelock gravity,
 ↪  perturbed without matter. Near the solution the constraint reads -6 g55 dH_b - 48 H_a^2 =
 ↪  kappa eps (to lowest order; dH_b = H_b + g55): (a) for a deflation that is SLOWER than g55
 ↪  there are no initial data without matter near the solution (no root H_a between -g55 and +g55
 ↪  for H_b = -0.999 g55; the constraint is a polynomial of degree four in H_a whose roots for H_b
 ↪  = -g55 are 0, 0 and (6 +- sqrt(20)) g55, the two last ones far from the deflating solution);
 ↪  (b) for H_b = -1.001 g55 the constraint has the two roots H_a = +-g55 sqrt(0.001/8) =
 ↪  +-0.01118 g55 to lowest order (found to 4 per cent), and in both cases the deflation does not
 ↪  return to the rate g55 but becomes faster and faster: H_a(0) = +0.01147 g55: at tau_ref =
 ↪  6.00, H_a = +1.73339 g55, H_b = -6.42224 g55; H_a(0) = -0.01088 g55: the rates run away; at
 ↪  tau_ref = 5.20, H_a = -0.00498 g55, H_b = -50.15518 g55. The exactly exponential deflation is
 ↪  therefore an UNSTABLE solution of the gravity in which it is a vacuum solution: it lies on the
 ↪  edge of the allowed initial data, and every allowed perturbation, with or without matter,
 ↪  grows. Constraint preserved to 8.3e-9
```

A.8 The independent verifications (`verify_deflating_independent.py`, `verify_cjt_ks_independent.py`)

```
PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C: independent verification
 ↪  of the deflating spacetimes and of the model without contact terms (computer algebra and
 ↪  general-purpose numerical tools)
generated by: code/scripts/verify_deflating_independent.py (sympy 1.14.0, scipy, numpy 2.5.3,
 ↪  mpmath 1.3.0)
43 of 43 checks passed

[PASS] I44d-G1  4+4, arbitrary rates and constants (computer algebra), metric in the author's sign
 ↪  convention with g_tt = -b_t^2: the Christoffel symbols obtained from the metric by
 ↪  differentiation are Gamma^t_(4t) = -rate_t and Gamma^4_(tt) = -gs_tt rate_t b_t^2 = + rate_t
 ↪  b_t^2; all others vanish
[PASS] I44d-G2  4+4: Ricci tensor R^4_4 = gsq, R^t_t = rate_t gsum, zero along x0..x3; curvature
 ↪  scalar gsq + gsum^2; Einstein tensor -gprod (component 44), rate_t gsum - (gsq + gsum^2)/2
 ↪  (components tt), -(gsq + gsum^2)/2 (x0..x3), with gsq = g55^2 + g66^2 + g77^2, gsum = g55 +
 ↪  g66 + g77, gprod = g55 g66 + g55 g77 + g66 g77: identities in the six symbols g55, g66, g77,
 ↪  const5, const6, const7 (mixed components; the same on both spacetimes in the author's sign
 ↪  convention)
[PASS] I44d-G3  4+4: Riemann tensor in the orthonormal frame, R^t_(4t4) = -rate_t^2 and R^t_(sts)
 ↪  = gs_ss rate_t rate_s = -rate_t rate_s; no component carries an index 0, 1, 2 or 3
[PASS] I44d-G4  4+4: the Lie derivative of the metric vanishes identically along 14 vector fields
 ↪  (7 translations, 6 rotations of x0..x3, the scaling d_4 + sum rate_t xt d_t); it does not
 ↪  vanish for the rotation of x0 and x5 nor for the boost of x4 and x1; for g55 = g66 = g77 it
 ↪  vanishes in addition along the three rotations of x5, x6, x7 and three special vector fields
 ↪  (20 in all)
```

```
[PASS] I44d-G5  4+4: the list of Killing vectors is complete. The matrices of the Lorentz algebra
↪  of the tangent space that leave the Riemann tensor unchanged form a space whose dimension,
↪  plus 8, bounds the number of Killing vectors from above; exact ranks for (g55, g66, g77) = (1,
↪  2, 3): 6 + 8 = 14; (1, 2, 4): 6 + 8 = 14; (3, 5, 7): 6 + 8 = 14; (2, 2, 5): 7 + 8 = 15; (5, 2,
↪  2): 7 + 8 = 15; (1, 1, 1): 12 + 8 = 20; (7, 7, 7): 12 + 8 = 20: the bounds 14 (three different
↪  rates), 15 (two equal) and 20 (three equal) equal the numbers of Killing vectors exhibited
[PASS] I44d-L1  4+4, Einstein-Gauss-Bonnet gravity (metric in the author's sign convention;
↪  standard formula of the Lanczos tensor, computer algebra): Lanczos^a_a = -4 g55 g66 g77 gsum
↪  for a = 0..3 and Lanczos^4_4 = Lanczos^t_t = 0, identically in the rates and constants; for
↪  g55 = g66 = g77 the deflating spacetime solves G^mu_nu + Lambda_8 delta^mu_nu + alpha_GB
↪  Lanczos^mu_nu = 0 with Lambda_8 = 3 g55^2 and alpha_GB = -1/(4 g55^2); for unequal rates it
↪  does not, because G^5_5 - G^6_6 = +-(g55 - g66) gsum while the Lanczos tensor vanishes there
[PASS] I44d-C1  4+4: the coordinate matrices gamma^mu = e_A^mu gamma^A satisfy gamma^mu gamma^nu +
↪  gamma^nu gamma^mu = 2 c g^(mu nu) with c = +1 and the inverse metric of the author's sign
↪  convention, for all 64 pairs, identically in b_5, b_6, b_7
[PASS] I44d-C2  4+4: the spin connection Omega_mu = (c/4) omega_(mu A B) gamma^A gamma^B computed
↪  from the tetrad postulate (indices lowered with the metric) is antisymmetric in its frame
↪  indices and equals Omega_t = +(1/2) rate_t b_t gamma^t gamma^4 (frame matrices) for t = 5, 6,
↪  7, zero otherwise
[PASS] I44d-C3  4+4: d_mu gamma^nu + Gamma^nu_(mu lam) gamma^lam + [Omega_mu, gamma^nu] = 0 for
↪  all 64 pairs
[PASS] I44d-C4  4+4: sum_mu gamma^mu Omega_mu = -(gsum/2) gamma^4: the Dirac operator is gamma^4
↪  (d_4 - gsum/2) + sum_a gamma^a d_a + sum_t gamma^t d_t / b_t
[PASS] I44d-C5  4+4: (1/2) gamma^mu gamma^nu (d_mu Omega_nu - d_nu Omega_mu + [Omega_mu,
↪  Omega_nu]) = -(c R/4) 1 with c = +1 and R = gsq + gsum^2
[PASS] I44d-D1  4+4: the generator K = -m gamma^4 + i k_a gamma^4 gamma^a + i (k_t/b_t) gamma^4
↪  gamma^t of the evolution of the weighted field satisfies K^2 = (k_d^2 - E_perp^2) 1
↪  identically in the eight symbols m, k_0..k_3, k_5, k_6, k_7 and in b_5, b_6, b_7
[PASS] I71d-G1  7+1, arbitrary rates and constants (computer algebra), metric in the author's sign
↪  convention with g_tt = +b_t^2: the Christoffel symbols obtained from the metric by
↪  differentiation are Gamma^t_(4t) = -rate_t and Gamma^4_(tt) = -gs_tt rate_t b_t^2 = - rate_t
↪  b_t^2; all others vanish
[PASS] I71d-G2  7+1: Ricci tensor R^4_4 = gsq, R^t_t = rate_t gsum, zero along x0..x3; curvature
↪  scalar gsq + gsum^2; Einstein tensor -gprod (component 44), rate_t gsum - (gsq + gsum^2)/2
↪  (components tt), -(gsq + gsum^2)/2 (x0..x3), with gsq = g55^2 + g66^2 + g77^2, gsum = g55 +
↪  g66 + g77, gprod = g55 g66 + g55 g77 + g66 g77: identities in the six symbols g55, g66, g77,
↪  const5, const6, const7 (mixed components; the same on both spacetimes in the author's sign
↪  convention)
[PASS] I71d-G3  7+1: Riemann tensor in the orthonormal frame, R^t_(4t4) = -rate_t^2 and R^t_(sts)
↪  = gs_ss rate_t rate_s = +rate_t rate_s; no component carries an index 0, 1, 2 or 3
[PASS] I71d-G4  7+1: the Lie derivative of the metric vanishes identically along 14 vector fields
↪  (7 translations, 6 rotations of x0..x3, the scaling d_4 + sum rate_t xt d_t); it does not
↪  vanish for the rotation of x0 and x5 nor for the boost of x4 and x1; for g55 = g66 = g77 it
↪  vanishes in addition along the three rotations of x5, x6, x7 and three special vector fields
↪  (20 in all)
[PASS] I71d-G5  7+1: the list of Killing vectors is complete. The matrices of the Lorentz algebra
↪  of the tangent space that leave the Riemann tensor unchanged form a space whose dimension,
↪  plus 8, bounds the number of Killing vectors from above; exact ranks for (g55, g66, g77) = (1,
↪  2, 3): 6 + 8 = 14; (1, 2, 4): 6 + 8 = 14; (3, 5, 7): 6 + 8 = 14; (2, 2, 5): 7 + 8 = 15; (5, 2,
↪  2): 7 + 8 = 15; (1, 1, 1): 12 + 8 = 20; (7, 7, 7): 12 + 8 = 20: the bounds 14 (three different
↪  rates), 15 (two equal) and 20 (three equal) equal the numbers of Killing vectors exhibited
[PASS] I71d-L1  7+1, Einstein-Gauss-Bonnet gravity (metric in the author's sign convention;
↪  standard formula of the Lanczos tensor, computer algebra): Lanczos^a_a = -4 g55 g66 g77 gsum
↪  for a = 0..3 and Lanczos^4_4 = Lanczos^t_t = 0, identically in the rates and constants; for
↪  g55 = g66 = g77 the deflating spacetime solves G^mu_nu + Lambda_8 delta^mu_nu + alpha_GB
↪  Lanczos^mu_nu = 0 with Lambda_8 = 3 g55^2 and alpha_GB = -1/(4 g55^2); for unequal rates it
↪  does not, because G^5_5 - G^6_6 = +-(g55 - g66) gsum while the Lanczos tensor vanishes there
[PASS] I71d-C1  7+1: the coordinate matrices gamma^mu = e_A^mu gamma^A satisfy gamma^mu gamma^nu +
↪  gamma^nu gamma^mu = 2 c g^(mu nu) with c = -1 and the inverse metric of the author's sign
↪  convention, for all 64 pairs, identically in b_5, b_6, b_7
```

```
[PASS] I71d-C2  7+1: the spin connection Omega_mu = (c/4) omega_(mu A B) gamma^A gamma^B computed
↪  from the tetrad postulate (indices lowered with the metric) is antisymmetric in its frame
↪  indices and equals Omega_t = +(1/2) rate_t b_t gamma^t gamma^4 (frame matrices) for t = 5, 6,
↪  7, zero otherwise
[PASS] I71d-C3  7+1: d_mu gamma^nu + Gamma^nu_(mu lam) gamma^lam + [Omega_mu, gamma^nu] = 0 for
↪  all 64 pairs
[PASS] I71d-C4  7+1: sum_mu gamma^mu Omega_mu = -(gsum/2) gamma^4: the Dirac operator is gamma^4
↪  (d_4 - gsum/2) + sum_a gamma^a d_a + sum_t gamma^t d_t / b_t
[PASS] I71d-C5  7+1: (1/2) gamma^mu gamma^nu (d_mu Omega_nu - d_nu Omega_mu + [Omega_mu,
↪  Omega_nu]) = -(c R/4) 1 with c = -1 and R = gsq + gsum^2
[PASS] I71d-D1  7+1: the Hamiltonian h = m beta + p_j alpha^j + (k_t/b_t) alpha^t of the weighted
↪  field is symmetric and satisfies h^2 = (E_perp^2 + k_d^2) 1 identically in the eight symbols
↪  m, p_0..p_3, k_5, k_6, k_7 and in b_5, b_6, b_7
[PASS] I71d-M1  7+1: psihat = sqrt(u_d) [J_(1/2 + i E_perp/rate_t)(u_d) w + i J_(-1/2 + i
↪  E_perp/rate_t)(u_d) alpha^d w], with h_perp w = E_perp w and u_d = k_d/rate_t, satisfies i d
↪  psihat/dx4 = h psihat (Bessel functions of mpmath; relative residual 1.1e-15 at four values of
↪  x4; mode m = 1, p = (0, 0.3, -0.4, 0.5), k_5 = 0.7, g55 = 2)
[PASS] I71d-M2  7+1: the number of particles created in a mode that starts empty, integrated with
↪  the Runge-Kutta method DOP853 of scipy and projected on the negative-energy states at k_d = 80
↪  E_perp, equals 1/(exp(2 pi E_perp/rate_t) + 1) for all eight internal states in three modes
↪  (largest relative deviation 2.2e-05)
[PASS] I71d-M3  7+1, equal rates 1.3 and a mode that depends on all three deflating coordinates
↪  (constants 1, 0.7, 2): the same law holds (relative deviation 1.4e-05)
[PASS] I44d-M1  4+4: psihat = sqrt(u_d) [K_(1/2 - i E_perp/rate_t)(u_d) w - K_(1/2 + i
↪  E_perp/rate_t)(u_d) i gamma^4 gamma^d w], with K_perp w = i E_perp w and u_d = k_d/rate_t,
↪  satisfies d psihat/dx4 = K psihat (modified Bessel functions of mpmath; relative residual
↪  2.1e-15); the eigenspace of K_perp with the eigenvalue i E_perp has dimension 8, so that there
↪  are eight such solutions; they fall as exp(-u_d)
[PASS] I44d-M2  4+4: between early x4 and the value at which the WKB exponent is S_wkb = 7.63 the
↪  bounded solution falls by exp(-S_wkb) times a constant: ln(amplitude early/late) - S_wkb =
↪  0.33534, the exact value (1/2) ln 2 - (1/2) ln(1 + exp(-2 pi E_perp/rate_t)) = 0.33654 (the
↪  difference, 1.2e-03, is the correction of relative order 1/u_d to the WKB form)
[PASS] Idef-B1  7+1: a scalar mode that depends on one deflating coordinate and starts empty ends
↪  with 1/(exp(2 pi varpi_b/rate_t) - 1) quanta (DOP853 of scipy, three modes, largest relative
↪  deviation 4.6e-05)
[PASS] I44d-V1  4+4: the two zero-energy equations of a vortex line of unit winding have a
↪  4-dimensional space of constant spinor solutions (singular values, numpy); on it alpha^3 has
↪  the eigenvalues -1 -1 +1 +1 and equals the central matrix P_c = i alpha^1 alpha^2 alpha^3 chi:
↪  four zero modes, two for each direction along the line
[PASS] I71d-V1  7+1: the two zero-energy equations of a vortex line of unit winding have a
↪  4-dimensional space of constant spinor solutions (singular values, numpy); on it alpha^3 has
↪  the eigenvalues -1 -1 +1 +1 and equals the central matrix P_c = i alpha^1 alpha^2 alpha^3 chi:
↪  four zero modes, two for each direction along the line
[PASS] Idef-C1  cosmology: an independent integration (scipy) of the expansion with three equal
↪  rates gives, at fixed 100 theta_* = 1.03958 and with last scattering held at z_* = 1089.92,
↪  the Hubble constant for gsum = 0.01 H_0: 66.743 km/s/Mpc (Rust program: 66.743); for gsum =
↪  0.1 H_0: 60.773 km/s/Mpc (Rust program: 60.773)
[PASS] Idef-C2  recombination on the deflating background, independently: Peebles' three-level
↪  atom (scipy; not the recombination program of the main project) gives z_* = 1090.84 without
↪  deflation (Rust program: 1089.92) and 100 theta_* = 1.03905 (1.03965). For gsum = 0.03 H_0:
↪  last scattering moves by -0.551 (Rust: -0.550), the helium fraction by -0.00243 (-0.00243);
↪  Hubble constant 65.365 km/s/Mpc (Rust: 65.365), and 65.483 (65.483) if last scattering and
↪  helium are held fixed; for gsum = 0.1 H_0: last scattering moves by -1.882 (Rust: -1.877), the
↪  helium fraction by -0.00844 (-0.00844); Hubble constant 60.367 km/s/Mpc (Rust: 60.368), and
↪  60.773 (60.773) if last scattering and helium are held fixed. The shifts of last scattering
↪  agree to 0.3 per cent, the changes of the Hubble constant at the fixed angle to 0.01 per cent
↪  (full) and 0.00 per cent (last scattering and helium held fixed): the larger fall of the
↪  Hubble constant in the full calculation is confirmed by a calculation that shares no
↪  recombination code with it
```

```
[PASS] I44d-P1  4+4: Einstein tensor of ordinary space to first order in a perturbation (plane
 ↪  wave along x3, synchronous gauge, conformal time, scale factors of x5, x6, x7 unperturbed;
 ↪  arbitrary rates; sympy). With Hc = a'/a, He = Hc - a gsum/2 and the signs chosen so that the
 ↪  time-time component is -(8 pi G) rho: background 3 Hc^2 - 3 a gsum Hc + a^2 gprod (time-time)
 ↪  and -(2 Hc' + Hc^2 - 2 a gsum Hc + a^2 (gsq + gprod)) (space-space); G^4_4: (2 k^2 eta - He
 ↪  h') cos(k x3); G^4_3: 2 k eta' sin(k x3); trace of the space-space part: (2 k^2 eta - 2 He h'
 ↪  - h'') cos(k x3); G^3_3 - G^1_1: (1/2)[(h'' + 6 eta'') + 2 He (h' + 6 eta') - 2 k^2 eta] cos(k
 ↪  x3); all divided by a^2. These are the equations (E), (M), (T), (S) of assumption A3: those of
 ↪  Ma and Bertschinger with Hc replaced by He in three of them. The components G^0_0 and G^5_5
 ↪  along the extra coordinates are perturbed as well: an agent must hold those coordinates
[PASS] I71d-P1  7+1: Einstein tensor of ordinary space to first order in a perturbation (plane
 ↪  wave along x3, synchronous gauge, conformal time, scale factors of x5, x6, x7 unperturbed;
 ↪  arbitrary rates; sympy). With Hc = a'/a, He = Hc - a gsum/2 and the signs chosen so that the
 ↪  time-time component is -(8 pi G) rho: background 3 Hc^2 - 3 a gsum Hc + a^2 gprod (time-time)
 ↪  and -(2 Hc' + Hc^2 - 2 a gsum Hc + a^2 (gsq + gprod)) (space-space); G^4_4: (2 k^2 eta - He
 ↪  h') cos(k x3); G^4_3: 2 k eta' sin(k x3); trace of the space-space part: (2 k^2 eta - 2 He h'
 ↪  - h'') cos(k x3); G^3_3 - G^1_1: (1/2)[(h'' + 6 eta'') + 2 He (h' + 6 eta') - 2 k^2 eta] cos(k
 ↪  x3); all divided by a^2. These are the equations (E), (M), (T), (S) of assumption A3: those of
 ↪  Ma and Bertschinger with Hc replaced by He in three of them. The components G^0_0 and G^5_5
 ↪  along the extra coordinates are perturbed as well: an agent must hold those coordinates
[PASS] Idef-P2  background on the deflating spacetime (sympy; four species: pressureless matter,
 ↪  radiation, a fluid with a variable equation of state, cosmological term): the expression of
 ↪  the program for the derivative of the conformal Hubble rate is the derivative of the equation
 ↪  of the expansion Hc^2 - a gsum Hc + a^2 gprod/3 = (sum of 8 pi G_N a^2 rho)/3 with (8 pi G_N
 ↪  a^2 rho)' = (a gsum - (1 + 3 w) Hc) 8 pi G_N a^2 rho; the quantity B = He' - He^2 + 4 pi G_N
 ↪  a^2 (rho + P) equals a gsum [ (1/6) sum of 8 pi G_N a^2 rho (1 - 3 w) + (a gsum)^2/4 - a^2
 ↪  gprod/3 ] / (2 He), which vanishes without deflation; and the expression of the program for B'
 ↪  is its derivative
[PASS] Idef-P3  the two identities between the gaps (sympy; pressureless matter, radiation with
 ↪  anisotropic stress, a fluid with variable equation of state and its own sound speed,
 ↪  cosmological term; arbitrary rates). With D, T, DP, Pi the sums of 8 pi G_N a^2 times the
 ↪  density, momentum, pressure and shear perturbations, eta eliminated by the energy constraint
 ↪  (E), and the equations of motion of matter of Ma and Bertschinger: momentum gap = 2 k^2 eta' -
 ↪  T, trace gap = h'' + 2 He h' - 2 k^2 eta + 3 DP, shear gap = alpha' + 2 He alpha - eta + (3/2)
 ↪  Pi/k^2 with alpha = (h' + 6 eta')/(2 k^2) obey  momentum gap = He (trace gap) + h' B - (a
 ↪  gsum/2)(D - 3 DP)  and  2 k^2 (shear gap) = (trace gap) + (3/k^2)[(momentum gap)' + 2 He
 ↪  (momentum gap)].  Without deflation B = 0 and a gsum = 0: the three gaps then vanish together,
 ↪  as in general relativity; with deflation they cannot all vanish
[PASS] Idef-P4  growth of pressureless matter (sympy). From (E) with the momentum constraint:
 ↪  delta'' + [(2 He' + g)/(2 He)] delta' = [(Hc - a gsum)/(2 He)] g delta (closure "Momentum");
 ↪  from (E) with the trace equation: delta'' + He delta' = g delta/2, that is, in the time x4,
 ↪  delta'' + (2 H - gsum/2) delta' = 4 pi G_N rho_m delta (closure "Pressure"); g = 8 pi G_N a^2
 ↪  rho_m, primes in conformal time. Both hold for every wavenumber. For matter alone and three
 ↪  equal rates, to the first order in r = gsum/H the growth rate is d ln delta/d ln a = 1 + c r
 ↪  with c = -9/16 ("Pressure"), -13/16 ("Momentum"), and -11/16 for the four-dimensional equation
 ↪  delta'' + 2 H delta' = 4 pi G_N rho_m delta, which is neither
[PASS] Idef-P5  growth of pressureless matter from a = 0.01 to today at a fixed Hubble constant,
 ↪  relative to no deflation: an independent integration (scipy) of the three growth equations
 ↪  agrees with the Rust program (data/growth_deflation.csv) to 6.3e-07 for the rates up to 0.1
 ↪  H_0; at gsum = 0.1 H_0: closure "Momentum" 0.95525 (Rust 0.95525); closure "Pressure" 0.97830
 ↪  (Rust 0.97830); four-dimensional equation 0.96740 (Rust 0.96740)
[PASS] Idef-P6  the three closures as systems of equations for the density contrast of
 ↪  pressureless matter and the metric (matter and cosmological term; implicit integrator of
 ↪  scipy; gsum = 0.1 H_0, Hubble constant fixed; growth from a = 0.01 to today relative to no
 ↪  deflation). Closure "Momentum": 0.955197; closure "Pressure": 0.978268; neither depends on
 ↪  the wavenumber (to 7e-14 for k = 0.1, 0.03, 0.01, 0.003/Mpc). Closure "Shear" (energy
 ↪  constraint and equation of the shear; its equations are stiff, because the momentum density of
 ↪  the agent relaxes at the rate k^2/(3 Hc)): 0.978266 at k = 0.1/Mpc, 0.978251 at k = 0.03/Mpc,
 ↪  0.978119 at k = 0.01/Mpc, 0.976788 at k = 0.003/Mpc; it differs from the closure "Pressure" by
 ↪  1.5e-06, 1.7e-05, 1.5e-04, 1.5e-03, that is by 0.31, 0.30, 0.30, 0.27 times (H_0/k)^2: far
 ↪  inside the horizon the closure "Shear" is the closure "Pressure"
```

```
[PASS] Idef-P7  the Einstein-Boltzmann program of the revision against the growth equations: for
 ↪  the five rates of data/cmb_deflation_growth.csv and both closures, the growth of the density
 ↪  contrast of matter at k = 0.1/Mpc from a = 0.01 to today in the full calculation (photons,
 ↪  neutrinos, baryons, cold dark matter), relative to no deflation and at a fixed Hubble
 ↪  constant, agrees with the independent integration (scipy) of the growth equation of the
 ↪  closure to 1.3e-04
[PASS] Idef-P8  the agent in the unperturbed universe (sympy; arbitrary rates): if the time-time
 ↪  Einstein equation holds with matter alone (assumption A2: the agent that keeps the rates
 ↪  constant has no energy density), the space-space equation and the conservation of matter
 ↪  require the agent to have the pressure p_agent in ordinary space with 8 pi G_N p_agent =
 ↪  -(gsum/H) dH/dx4 - 2 gsum H + 2 gprod - gsum gprod/(3 H). To the first order in the rates this
 ↪  is -gsum H/2 in an era of matter: a tension, of the relative size gsum/H

PROJECT spacetimes, Revision_Deflation_CJT_and_Kohn-Sham_definition_C: independent verification
 ↪  of the self-interaction, of the Kohn-Sham and CJT calculations, of the pairing and of the
 ↪  gravity with Kohn-Sham matter (Grassmann algebra, computer algebra and general-purpose
 ↪  numerical tools)
generated by: code/scripts/verify_cjt_ks_independent.py (python3 with numpy, scipy, sympy)
45 of 45 checks passed

[PASS] Icks-S1  the Fierz matrix F_rs = (1/16) sum_l (-1)^(rs-l) C(s,l) C(8-s,r-l) (exact rational
 ↪  arithmetic of sympy): F^2 = 1, eigenvalue +1 five times and -1 four times, so that (1 + F) I =
 ↪  0 are five independent relations among the nine invariants and four invariants remain; solved
 ↪  for I_2 .. I_6 the relations are I_2 = -3 I_0 - I_1 - I_7 - I_8, I_3 = -4 I_0 - I_7 + 4 I_8,
 ↪  I_4 = -5 I_0 - 5 I_8, I_5 = -4 I_0 - I_1 + 4 I_8, I_6 = -I_0 + I_1 + I_7 - 3 I_8, the table
 ↪  selfcoupling_reduction.csv of the Rust program
[PASS] Icks-S2-71s  7+1: each of the 256 matrices Theta^K (products of the Dirac matrices with the
 ↪  phases of the model) squares to eps_K = +-1, with eps_K = eta_K (eta_K: product of the squares
 ↪  of the matrices in K) as in the table of the Rust program; every vertex matrix M_K = A Theta^K
 ↪  is Hermitian, so that every bilinear psibar Theta^K psi is real (largest deviation 0.0e+00);
 ↪  Tr(Theta_K Theta^L) = 16 delta_KL (0.0e+00): the 256 matrices are a basis of all 16 x 16
 ↪  matrices
[PASS] Icks-S3-71s  7+1: the Fierz matrix computed from the matrices, F_rs = (1/256) sum_(|K| = r)
 ↪  Tr(Theta_K Theta_L Theta^K Theta^L) for an L of rank s (the completeness relation of the
 ↪  basis; the same for every L of the rank, checked for six of each rank: 0.0e+00), equals the
 ↪  closed form (largest deviation 0.0e+00)
[PASS] Icks-S4-71s  7+1: the nine invariants evaluated on Grassmann-valued fields (psi and psi*
 ↪  independent random complex combinations of 6 anticommuting generators; each I_r is then an
 ↪  element of degree four with 15 coefficients; three random fields) obey (1 + F) I = 0 (largest
 ↪  residual 6.3e-16 of the largest coefficient) and the five relations of Icks-S1 (4.9e-16); the
 ↪  rank of the nine invariants over all samples is 4 (singular values 4.28e-01 and 2.68e-16 on
 ↪  either side of the gap): exactly four invariants are independent. (The products of the
 ↪  Grassmann-valued components, done one by one with the Grassmann class, give the same
 ↪  bilinears: 7.2e-15.)
[PASS] Icks-S5-71s  7+1: Hartree-Fock couplings by Wick's theorem, <:(psi^dagger M psi)^2:> = Tr(M
 ↪  n)^2 - Tr(M n M n). The density matrix n of a Fermi sea, summed from the eigenvectors of
 ↪  positive energy of h(k) = alpha.k + beta M for 40 random pairs k, -k, has the form (rho +
 ↪  rho_S beta)/16 (deviation 1.1e-17) with rho_S = sum M/w. The energy of each of the nine
 ↪  channels, E_HF = (1/2)(G_1 rho^2 + G_0 rho_S^2), reproduces the coefficients of the table
 ↪  selfcoupling_hartree_fock.csv of the Rust program (largest deviation 2.1e-13)
[PASS] Icks-S6-71s  7+1: the contact interaction among the 28 pairs of two particles at rest (the
 ↪  eight states with beta = +1), diagonalized numerically for nine random couplings: seven pairs
 ↪  (among them the pair along the hidden axis) have the coupling g_0eff + g_1eff + 5 g_7eff and
 ↪  twenty-one have g_0eff + g_1eff - 3 g_7eff (largest deviation 6.4e-14)
[PASS] Icks-S7-71s  7+1: under the finite chiral rotation psi -> exp(0.37 i chi) psi (chi
 ↪  Hermitian, chi^2 = 1) of Grassmann-valued fields the interaction sum_r g_r I_r is unchanged
 ↪  when g_0eff + g_8eff = g_0 - 4 g_2 - 10 g_4 - 4 g_6 + g_8 = 0 (relative change 5.8e-16) and
 ↪  changes for generic couplings (1.43)
```

```
[PASS] Icks-S2-44q  4+4, quantized-sector form: each of the 256 matrices Theta^K (products of the
↪  Dirac matrices with the phases of the model) squares to eps_K = +-1, with eps_K = eta_K
↪  (eta_K: product of the squares of the matrices in K) as in the table of the Rust program;
↪  every vertex matrix M_K = A Theta^K is Hermitian, so that every bilinear psibar Theta^K psi is
↪  real (largest deviation 0.0e+00); Tr(Theta_K Theta^L) = 16 delta_KL (0.0e+00): the 256
↪  matrices are a basis of all 16 x 16 matrices
[PASS] Icks-S3-44q  4+4, quantized-sector form: the Fierz matrix computed from the matrices, F_rs
↪  = (1/256) sum_(|K| = r) Tr(Theta_K Theta_L Theta^K Theta^L) for an L of rank s (the
↪  completeness relation of the basis; the same for every L of the rank, checked for six of each
↪  rank: 0.0e+00), equals the closed form (largest deviation 0.0e+00)
[PASS] Icks-S4-44q  4+4, quantized-sector form: the nine invariants evaluated on Grassmann-valued
↪  fields (psi and psi* independent random complex combinations of 6 anticommuting generators;
↪  each I_r is then an element of degree four with 15 coefficients; three random fields) obey (1
↪  + F) I = 0 (largest residual 9.4e-16 of the largest coefficient) and the five relations of
↪  Icks-S1 (7.6e-16); the rank of the nine invariants over all samples is 4 (singular values
↪  5.67e-01 and 2.73e-16 on either side of the gap): exactly four invariants are independent.
↪  (The products of the Grassmann-valued components, done one by one with the Grassmann class,
↪  give the same bilinears: 7.9e-15.)
[PASS] Icks-S5-44q  4+4, quantized-sector form: Hartree-Fock couplings by Wick's theorem,
↪  <:(psi^dagger M psi)^2:> = Tr(M n)^2 - Tr(M n M n). The density matrix n of a Fermi sea, summed
↪  from the eigenvectors of positive energy of h(k) = alpha.k + beta M for 40 random pairs k, -k,
↪  has the form (rho + rho_S beta)/16 (deviation 2.2e-17) with rho_S = sum M/w. The energy of
↪  each of the nine channels, E_HF = (1/2)(G_1 rho^2 + G_0 rho_S^2), reproduces the coefficients
↪  of the table selfcoupling_hartree_fock.csv of the Rust program (largest deviation 8.0e-15)
[PASS] Icks-S6-44q  4+4, quantized-sector form: the contact interaction among the 28 pairs of two
↪  particles at rest (the eight states with beta = +1), diagonalized numerically for nine random
↪  couplings: seven pairs (among them the pair along the hidden axis) have the coupling g_0eff +
↪  g_1eff + 5 g_7eff and twenty-one have g_0eff + g_1eff - 3 g_7eff (largest deviation 7.9e-14)
[PASS] Icks-S7-44q  4+4, quantized-sector form: under the finite chiral rotation psi -> exp(0.37 i
↪  chi) psi (chi Hermitian, chi^2 = 1) of Grassmann-valued fields the interaction sum_r g_r I_r
↪  is unchanged when g_0eff + g_8eff = g_0 - 4 g_2 - 10 g_4 - 4 g_6 + g_8 = 0 (relative change
↪  8.6e-16) and changes for generic couplings (0.59)
[PASS] Icks-S2-44c  4+4, covariant form: each of the 256 matrices Theta^K (products of the Dirac
↪  matrices with the phases of the model) squares to eps_K = +-1, with eps_K = (-1)^r eta_K
↪  (eta_K: product of the squares of the matrices in K) as in the table of the Rust program;
↪  every vertex matrix M_K = A Theta^K is Hermitian, so that every bilinear psibar Theta^K psi is
↪  real (largest deviation 0.0e+00); Tr(Theta_K Theta^L) = 16 delta_KL (0.0e+00): the 256
↪  matrices are a basis of all 16 x 16 matrices
[PASS] Icks-S3-44c  4+4, covariant form: the Fierz matrix computed from the matrices, F_rs =
↪  (1/256) sum_(|K| = r) Tr(Theta_K Theta_L Theta^K Theta^L) for an L of rank s (the completeness
↪  relation of the basis; the same for every L of the rank, checked for six of each rank:
↪  0.0e+00), equals the closed form (largest deviation 0.0e+00)
[PASS] Icks-S4-44c  4+4, covariant form: the nine invariants evaluated on Grassmann-valued fields
↪  (psi and psi* independent random complex combinations of 6 anticommuting generators; each I_r
↪  is then an element of degree four with 15 coefficients; three random fields) obey (1 + F) I =
↪  0 (largest residual 6.9e-16 of the largest coefficient) and the five relations of Icks-S1
↪  (4.3e-16); the rank of the nine invariants over all samples is 4 (singular values 4.26e-01 and
↪  2.47e-16 on either side of the gap): exactly four invariants are independent. (The products of
↪  the Grassmann-valued components, done one by one with the Grassmann class, give the same
↪  bilinears: 7.2e-15.)
[PASS] Icks-S5-44c  4+4, covariant form: Hartree-Fock couplings by Wick's theorem, <:(psi^dagger M
↪  psi)^2:> = Tr(M n)^2 - Tr(M n M n). The density matrix n of a Fermi sea, summed from the
↪  eigenvectors of positive energy of h(k) = alpha.k + beta M for 40 random pairs k, -k, has the
↪  form (rho + rho_S beta)/16 (deviation 2.2e-17) with rho_S = sum M/w. The energy of each of the
↪  nine channels, E_HF = (1/2)(G_V0 rho^2 + G_S0 rho_S^2 + G_1 rho_W^2 + G_0 rho_SW^2), W = i C^-
↪  gamma^4, without cross terms, reproduces the coefficients of the table
↪  selfcoupling_hartree_fock.csv of the Rust program (largest deviation 2.2e-16)
```

```
[PASS] Icks-S6-44c  4+4, covariant form: the contact interaction among the 28 pairs of two
 ↪  particles at rest (the eight states with beta = +1), diagonalized numerically for nine random
 ↪  couplings: the 12 pairs of two particles in the same half W = +-1 have the couplings of the
 ↪  quantized-sector form (3 with g_0eff + g_1eff + 5 g_7eff, 9 with g_0eff + g_1eff - 3 g_7eff),
 ↪  and the 16 pairs of particles in different halves have the OPPOSITE sign (4 with -(g_0eff +
 ↪  g_1eff + 5 g_7eff), 12 with -(g_0eff + g_1eff - 3 g_7eff)) (largest deviation 1.5e-14)
[PASS] Icks-S7-44c  4+4, covariant form: under the finite chiral rotation psi -> exp(0.37 i chi)
 ↪  psi (chi Hermitian, chi^2 = 1) of Grassmann-valued fields the interaction sum_r g_r I_r is
 ↪  unchanged when g_1eff - g_7eff = g_1 + g_3 - g_5 - g_7 = 0 (relative change 6.0e-16) and
 ↪  changes for generic couplings (0.27)
[PASS] Icks-S8  the two forms of the self-interaction on 4+4: with W = i C^- gamma^4 (Hermitian,
 ↪  W^2 = 1, commuting with beta) the invariants of the covariant form are those of the
 ↪  quantized-sector form with W inserted in every bilinear, I_r(covariant) = sum_K eps_K
 ↪  (psi^dagger W M_K psi)^2, on Grassmann-valued fields (largest relative deviation 3.0e-16):
 ↪  the two forms agree for particles in the same half W = +-1 and differ in sign for particles in
 ↪  different halves
[PASS] Icks-K1  the Fermi integrals rho_S = <M/w>, e_kin = <w> and <k^2/(D w)> over a ball of
 ↪  momenta in D = 7, 4, 3 space dimensions, three ways: Gauss-Legendre nodes of numpy (used
 ↪  below), the adaptive quadrature of scipy, and the closed forms from the recurrence I_n =
 ↪  [k_F^(n-1) E_F - (n-1) M^2 I_(n-2)]/n with I_0 = asinh(k_F/M), I_1 = E_F - M (40 digits):
 ↪  largest relative difference 1.2e-14
[PASS] Icks-K2  flat 7+1 spacetime, the example of the author's document KOHN_SHAM_GRAVITY (m = 1,
 ↪  G_S = -1e5, G_V = +1e4, k_F = 0.756636150701493): M = 0.348097340146, rho = 1.388100213e-05,
 ↪  eps = 1.348787233e-05, E/A - m = -0.02832142796 here, against the printed 0.348097340146,
 ↪  1.388100213e-5, 1.348787233e-5, -0.02832142796 (largest relative difference 1.7e-10); the
 ↪  minimum of the energy per particle, located by the vanishing of the pressure, is at k_F =
 ↪  0.756636150701 (printed: 0.756636150701493; pressure there 1.3e-21) and the chemical potential
 ↪  equals the energy per particle (difference 8.9e-15); p = mu rho - eps along the equation of
 ↪  state to 9.2e-15. With G_S = -98750, G_V = 20000, the Hartree-Fock couplings that g_0 = -1e5,
 ↪  g_1 = +1e4 give (Icks-S5; the same numbers are printed in section 4 of the author's document),
 ↪  the energy per particle stays above m for all Fermi momenta from 0.02 to 1.5 (lowest value
 ↪  +0.000156): uniform matter is not self-bound for that pair of couplings. The Kohn-Sham numbers
 ↪  of the document belong to G_S = -1e5, G_V = +1e4 taken as the mean-field couplings themselves
[PASS] Icks-K3  Kohn-Sham matter of 7+1 with momenta along all seven space coordinates while x5,
 ↪  x6, x7 deflate (collisionless): the integral over the ellipsoid of occupied states with the
 ↪  four flat momenta integrated in closed form reproduces the table ks_deflating_71.csv of the
 ↪  Rust program at the clock readings -2, -1, 0, 1, 2, 4 for equal rates and for rates 1 : 2 : 3
 ↪  (12 states; M, rho, rho_S, eps and the five pressures: largest deviation 1.4e-08); at the
 ↪  reference time it reproduces the ball (4.4e-15); the energy density obeys d eps/dx4 = sum_t
 ↪  rate_t (eps + p_t) (numerical derivative, four cases: 1.9e-08)
[PASS] Icks-K4  fields that do not depend on x5, x6, x7 (D = 4) nor on x0 (D = 3): the smallest
 ↪  scalar attraction |G_S| m^(D-1) at which uniform matter is self-bound, found with scipy
 ↪  (lowest energy per particle over the Fermi momentum equal to m): D = 4, G_V/|G_S| = 0.0000:
 ↪  62.0767; D = 4, G_V/|G_S| = 0.1000: 85.6504; D = 4, G_V/|G_S| = 0.2025: 119.5723; D = 3,
 ↪  G_V/|G_S| = 0.0000: 8.0671; D = 3, G_V/|G_S| = 0.1000: 9.8364; D = 3, G_V/|G_S| = 0.2025:
 ↪  12.2272; the table ks_reduced_thresholds.csv of the Rust program agrees to 1.0e-09
 ↪  (relative). The third ratio, 0.202532, is the ratio G_V/|G_S| of the Hartree-Fock couplings of
 ↪  g_1 = -0.1 g_0 (Icks-S5)
[PASS] Icks-K5  covariant form on 4+4, the two halves W = +-1 as two kinds of particles with four
 ↪  states each (g_0 = -363.26, g_1 = -0.1 g_0, total density of Fermi momentum 0.8; Hartree-Fock
 ↪  couplings of Icks-S5): the two gap equations M_+- = m + G_S0 rho_S +- G_0 rho_SW, solved with
 ↪  the hybrid Powell method of scipy from 81 start values that include negative masses, have
 ↪  exactly one solution at each of the polarizations 0, 0.5 and 1, with positive masses, and it
 ↪  is the state of the table ks_44c_polarization.csv of the Rust program (masses and energy:
 ↪  largest difference 1.5e-09): E/A - m = +0.3799 for equal halves, +0.3348 for the polarization
 ↪  0.5 and +0.3731 with all particles in one half; the same density in the quantized-sector form
 ↪  has E/A - m = +0.1323 (difference 1.1e-09). The state with equal halves is not the state of
 ↪  lowest energy. (A remark on the no-sea approximation: the Hartree-Fock energy as a function of
 ↪  unrestricted trial masses has no lowest value, because trial masses of opposite sign and large
 ↪  magnitude put particles into states of negative rest energy; for equal halves it falls to E/A
 ↪  - m = -2.74. The Kohn-Sham state is the solution of the gap equations, not the lowest value of
 ↪  that function.)
```

```
[PASS] Icks-J1  the CJT effective potential Omega(M, V_0; mu) = Omega_1(M, mu - V_0) - (M -
 ↪  m)^2/(2 G_S) - V_0^2/(2 G_V) evaluated in Python: for the 30 states of the table cjt_vs_ks.csv
 ↪  (D = 7, 4, 3) its stationary point at the chemical potential of the Kohn-Sham state has the
 ↪  mass, the density and the energy density Omega + mu rho of the Kohn-Sham state, and -Omega is
 ↪  the Kohn-Sham pressure, both of the Rust program and of the Kohn-Sham routine of this script
 ↪  (largest relative deviation 2.0e-09)
[PASS] Icks-J2  with the filled states of negative energy below a cutoff Lambda, a massless field
 ↪  acquires a mass in vacuum when the scalar attraction exceeds G_c = -(2 pi)^D (D - 1)/(nu A_D
 ↪  Lambda^(D-1)); computer algebra gives D = 7: G_c = -90*pi**4/Lambda**6; D = 4: G_c =
 ↪  -3*pi**2/Lambda**3; D = 3: G_c = -pi**2/(2*Lambda**2); the numerical minimum of the vacuum
 ↪  potential is at M = 0 for G_S = 0.98 G_c and at M > 0 for G_S = 1.02 G_c in all three cases.
 ↪  The coupling G_S = -1e5 of the author's example equals G_c for Lambda = (90 pi^4/1e5)^(1/6) =
 ↪  0.6665 m
[PASS] Icks-J3  the author's example with the filled states of negative energy below the cutoff
 ↪  included (vacuum mass kept at 1 by the choice of the bare mass m = M_vac + G_S nu Int^Lambda
 ↪  M_vac/w; energy of matter relative to the vacuum, minimized over the mass and the Fermi
 ↪  momentum with scipy): Lambda = 0.30: bare mass 0.9979, saturation at k_F = 0.7566 with E/A -
 ↪  M_vac = -0.02837; Lambda = 0.45: bare mass 0.9660, saturation at k_F = 0.7568 with E/A - M_vac
 ↪  = -0.02984; Lambda = 0.60: bare mass 0.7578, saturation at k_F = 0.7565 with E/A - M_vac =
 ↪  -0.04483; the values quoted by the Rust program (check cjt-J3) agree to 5e-05
[PASS] Icks-U1  the functional with bare Dirac mass, contact couplings and Vorton-Scalar in
 ↪  ordinary space (D = 3), minimized directly with scipy: 12 states of the table
 ↪  unified_states.csv of the Rust program are reproduced (eps, M and x: largest deviation
 ↪  1.7e-08); without bare mass and contact couplings matter is first self-bound at C = g_VS^2
 ↪  m^2/m_s^2 = 3.701102, the exact value 3 pi^2/8 = 3.701102 of a gas of massless fermions with
 ↪  the energy density m^2 m_s^2/(8 g_VS^2) of the scalar at phi = 0. Phase mode with a bare mass:
 ↪  the second derivative of the energy of the filled states with respect to the phase of the
 ↪  scalar equals (|<psibar psi>| - rho_S) m_D m_Y/m (cutoff 2 m, f_D = 0.3, k_F = m; adaptive
 ↪  quadrature; agreement with the formula and with the Rust table to 2.7e-07)
[PASS] Icks-P1  pairing amplitudes in the hidden-axis channel (pair matrix X = C^- gamma^0,
 ↪  antisymmetric; momenta p, -p -> p', -p' in ordinary space) from projectors that are summed
 ↪  from numerical eigenvectors of the single-particle Hamiltonian, twelve random cases on each of
 ↪  7+1 and 4+4 (quantized-sector form): the amplitudes of the ranks 0, 1, 7, 8 equal the closed
 ↪  forms F_0 = (E E' + M^2 - p.p')/(2 E E'), F_1 = (3 E E' - 2 M^2 - p.p')/(E E'), F_7 = (3 E E'
 ↪  + 2 M^2 - p.p')/(E E'), F_8 = -(E E' - M^2 - p.p')/(2 E E') (largest deviation 1.8e-15); the
 ↪  two spacetimes give the same amplitudes (3.7e-15); at rest the nine amplitudes are (1, 1, -9,
 ↪  -9, -5, -5, 5, 5, 0) (0.0e+00); the relations of Icks-S1 hold for the part of the amplitudes
 ↪  that is even under p' -> -p' (4.4e-15) and not for the odd part (deviation up to 6.55)
[PASS] Icks-P2  the gap equation of the hidden-axis pair solved again in Python (another grid of
 ↪  momenta: geometric panels around the Fermi momentum and up to the same upper limit 2e4 m as in
 ↪  the Rust program, ten Gauss-Legendre nodes each; plain iteration to a relative change below
 ↪  1e-12; the gap at the Fermi momentum from the equation itself). The closed angular average of
 ↪  the kernel agrees with the adaptive quadrature of scipy (1.2e-14). Gap in units of the Fermi
 ↪  kinetic energy at the clock reading 2.04, k_F = 0.5 m, for the exchange of the Vorton-Scalar
 ↪  alone (variant 0), with the scalar contact attraction g_0eff = -g_VS^2/m^2 (variant 1) and
 ↪  with the contact couplings in the ratio of the author's example (variant 2): m_s = 1.0,
 ↪  variant 0: 0.0247 (Rust: 0.0247); m_s = 1.0, variant 1: 0.2975 (Rust: 0.2975); m_s = 1.0,
 ↪  variant 2: 0.1963 (Rust: 0.1963); m_s = 0.3, variant 0: 0.4118 (Rust: 0.4118); m_s = 0.3,
 ↪  variant 1: 0.9035 (Rust: 0.9035); m_s = 0.3, variant 2: 0.7407 (Rust: 0.7407); largest
 ↪  relative difference 8.2e-07
[PASS] Icks-G1  spacetimes ds^2 = -dt^2 + a(t)^2 (dx_a)^2 + b(t)^2 (dy_b)^2 with 3 + 4 and with 4
 ↪  + 3 space coordinates, arbitrary functions a(t), b(t) (computer algebra, from the metric:
 ↪  Christoffel symbols, Riemann tensor): the Riemann tensor is diagonal in the pairs of
 ↪  directions, R^(ij)_(ij) = K_ij with K_(t i) = a_i''/a_i and K_(ij) = H_i H_j, and has no other
 ↪  components; the Einstein tensor is G^m_m = -sum of K over the pairs that do not contain m, and
 ↪  the Lanczos tensor 2 (R R^m_n - 2 R^m_a R^a_n - 2 R^(m a)_(n b) R^b_a + R^(m a)_(b c) R^(b
 ↪  c)_(n a)) - (1/2) delta^m_n (R^2 - 4 Ric^2 + Riem^2) is -4 times the sum over the sets of four
 ↪  directions without m of the sum over their three divisions into pairs of the products K K: the
 ↪  formula E^(k)m_m = (-1)^(k+1) k! 2^(k-1) [pair sum] of the Rust program for k = 1 (E^(1) = -G)
 ↪  and k = 2 (E^(2) = Lanczos tensor), time component and both space components
```

[PASS] Icks-G2 the Lovelock tensors of orders k = 1, 2, 3 derived from the reduced action S_k = Int dt N a^(n_a) b^(n_b) L_k, with the lapse N, the Lovelock density L_k = k! 2^k [sum over the sets of 2k directions and their divisions into pairs of the products of K] (L_1 = curvature scalar, L_2 = Gauss-Bonnet density, L_3 = cubic Lovelock density) and K_(t i) = (N a_i)^(-1) d/dt (a_i'/N): the Euler-Lagrange expressions for N, a and b at N = 1 (computer algebra, with second derivatives in the Lagrangian) reproduce E^(k)m_m = (-1)^(k+1) k! 2^(k-1) [pair sum without m] exactly, for the time component and both space components, for 3 + 4 and for 4 + 3 space coordinates (18 identities). All terms with third and fourth derivatives cancel: the equations are of second order

[PASS] Icks-G3 the self-consistent evolution of the author's document KOHN_SHAM_GRAVITY (three ordinary and four auxiliary space coordinates, kappa = 1000, LambdaBar = 0, H_a(0) = 0.05, a(0) = b(0) = 1, Kohn-Sham matter of the saturation density with G_S = -1e5, G_V = +1e4) integrated again with the Runge-Kutta method of order 8 of scipy, the field equations built from the pair sums with sympy and the matter from an independent quadrature: H_b(0), a(20), b(20), H_a(20), H_b(20) agree with the ten numbers printed in the document, for Einstein gravity and for Einstein-Lovelock gravity with alpha_2 = 20, alpha_3 = 500, to 4.8e-08 (relative); the constraint is preserved to 9.7e-14

[PASS] Icks-G4 the deflating spacetime (four flat coordinates, H_a = 0; three coordinates deflating at the same rate, H_b = -g55) in the author's Einstein-Lovelock gravity, by computer algebra in the symbol g55: it solves the three field equations without matter for LambdaBar = 3 g55^2 and alpha_2 = 1/(4 g55^2); E^(3) vanishes identically there (alpha_3 is arbitrary); the matrix of the coefficients of (H_a', H_b') in the two space equations is diag(-6, 2); in Einstein gravity alone the same spacetime needs kappa eps = 3 g55^2, kappa p_a = -6 g55^2, kappa p_b = -3 g55^2; near the solution the constraint is -6 g55 dH_b - 48 H_a^2 + (higher orders) = kappa eps, with no term linear in H_a; for H_b = -g55 its four roots in H_a are 0, 0 and (6 +- sqrt(20)) g55

[PASS] Icks-G6 the Einstein-Lovelock polynomials that the documents print for four directions with the rate H_a and three with the rate H_b (E^(1) and E^(2): time component and both space components; E^(3): time component) are the sums over pair curvatures, term by term (computer algebra, seven identities); every term of the three components of E^(3) contains at least two factors of H_a or of its derivative, so that E^(3) and its first variation vanish at H_a = 0

[PASS] Icks-G5 the deflating spacetime with Kohn-Sham matter of 7+1 (momenta along all seven space coordinates) as a source, g55 = 0.01 m, loading kappa eps(0)/(3 g55^2) = 1e-3, integrated with scipy to the clock reading 6 of the undisturbed deflation: tuned Einstein-Lovelock: H_b(0) = -1.00050 g55; at tau_ref = 6: a = 1.2508, b = 4.9085e-02, H_a = +1.4052 g55, H_b = -7.9908 g55; Einstein: H_b(0) = -0.03162 g55; at tau_ref = 6: a = 1.0011, b = 9.3353e-01, H_a = +0.0012 g55, H_b = -0.0375 g55; the summary of the Rust program (six decimals) agrees to 6.4e-06 (relative to |value| + 0.05; constraint preserved to 1.3e-11). Without matter, from H_b(0) = -1.001 g55 and the positive root H_a(0) = 0.01147 g55 of the constraint: H_a = +1.73339 g55, H_b = -6.42224 g55 at the same clock reading (Rust: +1.73339, -6.42224; relative difference 6.7e-10): the deflation does not return to the rate g55

[PASS] Icks-F1 the two forms of the 4+4 interaction in uniform matter whose two halves W = +1 and W = -1 are filled differently (numerical matrices; Wick's theorem with the density matrix (rho 1 + rho_S beta + rho_W W + rho_SW W.beta)/16): the Hartree-Fock energy is diagonal in the four densities for each of the nine ranks (largest off-diagonal element 0.0e+00); for random couplings g_0, g_1, g_7, g_8 the four coefficients are, in the covariant form, (G_Vc, G_Sc, G_V^HF, G_S^HF) for (rho^2, rho_S^2, rho_W^2, rho_SW^2) and, in the quantized-sector form, (G_V^HF, G_S^HF, G_Vc, G_Sc), with G_S^HF = (15 g_0 - 8 g_1 - 8 g_7 - g_8)/16, G_V^HF = (-g_0 + 22 g_1 - 6 g_7 + g_8)/16, G_Sc = (-g_0 - 2 g_1 + 2 g_7 + g_8)/16, G_Vc = -(g_0 + g_8)/16 (largest deviation 0.0e+00); for every one of the nine ranks the coefficients of the two forms are the same numbers with the sums and the differences of the densities exchanged (0.0e+00)

[PASS] Icks-F2 the internal symmetry of the fields in ordinary space and the self-interaction, tested at a finite angle (psi -> exp(0.37 i T) psi, numerical matrices, the nine antisymmetrized quartic tensors transformed index by index): number of independent conditions on the nine couplings for invariance, covariant form: 0 for a rotation of two of x5, x6, x7, 4 for the unitary transformation generated by gamma^0 gamma^5, 4 for the generator P of U(1)_E; quantized-sector form: 0, 0, 2. In the covariant form the couplings that are invariant under gamma^0 gamma^5 or under P are exactly those whose four effective couplings vanish (the five combinations that are zero for anticommuting fields): every self-interaction that does not vanish breaks these symmetries

```
[PASS] Icks-F3  the pairing amplitudes of the contact channels in the two forms (projectors on the
↪  states of positive energy from numerical eigenvectors; random masses and momenta in ordinary
↪  space): the pair along the hidden axis satisfies W kappa W^T = -kappa (one particle of each
↪  half of the field), the pairs along x5, x6, x7 satisfy W kappa W^T = +kappa (two particles of
↪  the same half); in the quantized-sector form the amplitudes of the ranks 0, 1, 7, 8 are the
↪  closed forms F_0, F_1, F_7, F_8 in both kinds of pairs, in the covariant form they are the
↪  closed forms in the pairs along x5, x6, x7 and MINUS the closed forms in the pair along the
↪  hidden axis (largest deviation 1.8e-15)
[PASS] Icks-F4  uniform matter of the two halves in both forms, recomputed with an independent
↪  solver of the two coupled gap equations (damped fixed-point iteration from several start
↪  values; Gauss-Legendre integrals), where the program uses closed forms of the integrals and a
↪  search for all solutions: (i) 36 states of the table of the energy against the polarization
↪  (both forms, D = 4 and 3, sets A and B): masses and energy per particle agree to 7.8e-11, and
↪  the iteration finds no solution of lower energy; (ii) the binding thresholds g m^(D-1) of the
↪  program (D = 4, set A, form C: 224.7386; D = 4, set A, form Q: 82.1116; D = 4, set B, form C:
↪  332.2917; D = 4, set B, form Q: 121.0858; D = 3, set A, form C: 24.5717; D = 3, set A, form Q:
↪  9.8090; D = 3, set B, form C: 31.1674; D = 3, set B, form Q: 12.3819): at 1.01 times each
↪  threshold the state that the program reports is bound, and at 0.99 times the threshold a scan
↪  of 36 densities and 5 polarizations finds no bound state; (iii) the saturated states at the
↪  clock readings 1 and 3 (16 states): energy per particle to 4.0e-12, pressure over energy
↪  density at most 1.1e-08
[PASS] Icks-F5  the effective potential of two halves, Omega(M_+, M_-, V_+, V_-; mu) = sum over the
↪  halves of the grand potential of free particles - muS^2/(2 G_Stot) - dS^2/(2 G_Sdif) - vT^2/(2
↪  G_Vtot) - vD^2/(2 G_Vdif), evaluated with adaptive quadrature at 16 of the Kohn-Sham states of
↪  the program (both forms; states with equal halves, one half, unequal halves): its four partial
↪  derivatives vanish (central differences; largest value 1.5e-09 of the density), -Omega equals
↪  the pressure and Omega + mu n the energy density of the Kohn-Sham state (4.7e-11)
[PASS] Icks-F6  the vacuum of the two forms with the filled states of negative energy (adaptive
↪  quadrature), at the clock readings -0.5, 0.2 and 0.5 counted from the critical strength, D = 4
↪  and 3, sets A and B, without bare mass and with the bare mass 0.05 Lambda: the masses that the
↪  program reports satisfy the two gap equations of the vacuum (largest residual 2.5e-11);
↪  without bare mass they vanish below the critical strength, and above it their magnitude is the
↪  solution of M = -G_S^HF NU Int^Lambda M/w (1.1e-11), with EQUAL signs in the quantized-sector
↪  form and OPPOSITE signs in the covariant form
[PASS] Icks-F7  the gap of the hidden-axis pair in the covariant form, where the contact couplings
↪  enter the pairing potential with the opposite sign: independent solution of the gap equation
↪  (another grid of momenta, the same upper limit 2e4 m, plain iteration) for the exchange of the
↪  Vorton-Scalar with a scalar contact attraction and with a vector contact repulsion (m_s = m,
↪  k_F = 0.5 m, clock reading 2.04): with the vector coupling the gap agrees with the program to
↪  7.8e-05 (relative) and exceeds the gap of the exchange alone; with the scalar coupling the
↪  program reports no gap and the independent iteration finds none above 1e-6 E_F
[PASS] Icks-F8  the ultraviolet criterion of the gap equation. (i) The integrals Int_0^inf dt
↪  (1/2) Int_-1^1 dx F(x)/(1 + t^2 - 2 t x) by nested adaptive quadrature: 0.733700550 for F = (1
↪  - x)/2 (closed form pi^2/8 - 1/2 = 0.733700550) and 6.402203301 for F = 3 - x (closed form 3
↪  pi^2/4 - 1 = 6.402203301). (ii) The kernel of this script at the momentum 3e4 m, integrated
↪  over the ratio of the momenta, equals U = g_VS^2 (pi^2/4 - 1) - m_ct^2 [(g_0eff -
↪  g_8eff)(pi^2/8 - 1/2) + (g_1eff + g_7eff)(3 pi^2/4 - 1)] for four sets of couplings (largest
↪  relative deviation 1.3e-05). (iii) The hidden-axis pair of the covariant form with the vector
↪  contact coupling g_1eff = +0.3 g_VS^2/m^2 (m_s = m, k_F = 0.5 m), solved with the upper limits
↪  2e3 m and 2e4 m: clock reading 2.04, U/(4 pi^2) = 0.659: gap/E_F = 0.22841 and 0.22848; clock
↪  reading 2.77, U/(4 pi^2) = 1.373: gap/E_F = 994.78 and 10958 (program, with its own grid:
↪  1044.3 and 11991); below the criterion the gap does not depend on the limit, and the two
↪  values and the values of U/(4 pi^2) agree with the program to 7.8e-05; above the criterion the
↪  gap grows with the limit in both programs by more than a factor 5 per decade (its value then
↪  depends on the limit and on the grid of momenta, and is not a property of the model)
```

Appendix B. The symbols

The following tables list every symbol of the project with its meaning, the names by which the programs and the data files call it, and its origin. The symbols of this document are among them; a

symbol has one meaning in every document of the project.

1 Coordinates and indices

symbol	names in programs and data files	meaning	origin
$X = \{x0, \dots, x7\}$	`x0` `x1` `x2` `x3` `x4` `x5` `x6` `x7`	the eight coordinates of both spacetimes, never renamed: $x0$ hidden space-like; $x1, x2, x3$ ordinary space; $x4$ ordinary time; $x5, x6, x7$ additional coordinates (time-like in 4+4, space-like in 7+1)	N01 (text at its head) and N04 `X` (section DEFCHART): the same eight names with the same meaning. N02 uses the same eight names with `x0` as the time: related by swap04
swap_{04}	`swap04`	the exchange of the index values 0 and 4; it carries the labels of `7+1.nb` (time at index 0) into those of the project (time at index 4)	new
A, B, C, D	—	index with the values $0, \dots, 7$; always labels the coordinate xA	new
a, b	—	indices whose range is stated at each place of use: the four extra axes, values 0, 5, 6, 7 (in everything that concerns pairs and the order parameter), the four flat coordinates other than the time, values 0, 1, 2, 3 (in the statements of the machine checks), or the sixteen components of the field, values 1 to 16 (in the density matrix n_{ab}). The letters `a`, `b` in sans-serif type are the scale factors of the Einstein-Lovelock evolution; a_{sp}, a_{sc}, $a(T)$, b_t and b have their own entries	new
i, j	—	index of space-like coordinates; values 1, 2, 3 (ordinary space) unless the range is stated: $j = 0, 1, 2, 3$ for the four flat coordinates other than the time where the deflating coordinates occur as well, and all seven space-like directions in the Hamiltonian of 7+1. In the Einstein-Lovelock equations i, j label the seven space directions	new
h (index)	—	index with the values 1, 2, 3 that labels the 4×4 matrices s_h, t_h	N01, N02, N05 `h` in `s4by4[h]`, `t4by4[h]`

2 Matrices of the 4+4 spacetime (tag 1644)

symbol	names in programs and data files	meaning	origin
η_{1644}	`eta1644`	metric of the flat 4+4 spacetime and matrix of the squares of its Dirac matrices, $\text{diag}(+1, +1, +1, +1, -1, -1, -1, -1)$: space-like $= +1$. It equals the flat metric $\eta^{(4+4)}$ of the author's sign convention for the deflating spacetimes	N01, N02, N05, N10–N12 `η4488`; in N01 also `η8` and `g_AB`; D08 G_{AB} with the index value 8 in the place of the value 0
s_h, t_h	—	the three self-dual and the three anti-self-dual real antisymmetric 4×4 matrices	N01, N02, N05, N10–N12 `s4by4[h]`, `t4by4[h]`; the letters are those of D06 (s^h self-dual, t^h anti-self-dual)

symbol	names in programs and data files	meaning	origin
τ^A_{1644}, $\bar\tau^A_{1644}$	`tau1644` `taubar1644`	the real 8×8 blocks of the 4+4 Dirac matrices (reduced generators of the split octonion algebra)	N01, N10–N12 `τ[A]` and `τ` with an overbar; N02 `τ[A]`, `τbar[A]`; matrices of this kind are the τ^A, $\bar\tau^A$ of D07 and D08, whose index runs from 1 to 8 (the value 8 there is the value 0 here)
σ_{88}	`sigma88`	the 8×8 matrix with the 4×4 blocks $(0, 1_4;\ 1_4, 0)$	N01 `σ`, `σ88`; N02 `σ`; N05 `σ88`
γ^A_{1644}	`gamma1644`	the eight real 16×16 Dirac matrices of the 4+4 spacetime	N01, N10–N12 `T16^A[A]`; N02 `T16[A]`; the t^A of D08 are built in the same way
Σ^{AB}_{1644}	`Sigma1644`	$\frac{1}{4}[\gamma^A_{1644}, \gamma^B_{1644}]$, the scaled commutators; they span $\mathfrak{so}(4,4)$	new name; the same matrices are `SAB` and `sAB[A][B]` of N10–N12; NOT `sAB` of N02, which is a sign
ω_{1644}	`omega1644`	the product $\gamma^0_{1644}\cdots\gamma^7_{1644}$; real chirality matrix of 4+4, $\mathrm{diag}(-1_8, +1_8)$	N01 `T16^A[8]`; N02 `T16[8]`
P^L_{1644}, P^R_{1644}	`PL1644` `PR1644`	the real chiral projectors $\frac{1}{2}(1 \mp \omega_{1644})$	N01 `P_L`, `P_R`; N02 `PL44`, `PR44`
C^-_{1644}, C^+_{1644}	`Cm1644` `Cp1644`	the products $\gamma^0\gamma^1\gamma^2\gamma^3$ and $\gamma^4\gamma^5\gamma^6\gamma^7$ of the 4+4 matrices; $C^\mp\gamma^A(C^\mp)^{-1} = \mp(\gamma^A)^T$	N01, N02, N10–N12 `σ16` (C^-); N02 `S44` (C^+)
W_{1644}	`W1644`	$i\,C^-_{1644}\gamma^4_{1644}$, the grading: Hermitian, $W^2 = 1$; density of the covariant current	new
K_{1644}	`K1644`	generator of the evolution of dirac1644 in $x4$: $\partial_4\psi = K\psi$	new
β_{1644}, α^j_{1644}	`beta1644` `alpha1644`	the Hermitian matrices $-i\gamma^4_{1644}$ and $-\gamma^4_{1644}\gamma^j_{1644}$ ($j = 0, 1, 2, 3$) of the quantized 4+4 theory	new
h_{1644}	`h1644`	Hamiltonian matrix of the quantized 4+4 theory, iK_{1644} for fields independent of $x5, x6, x7$	new
Π_{1644}	`Pi1644`	the projector $(i\gamma^A_{1644}k_A + m_{1644})/2m_{1644}$ on the plane-wave solutions of the worked example	new

3 Matrices of the 7+1 spacetime (tag 1671)

symbol	names in programs and data files	meaning	origin
η_{1671}	`eta1671`	the matrix of the squares of the Dirac matrices of 7+1, $\gamma^A_{1671}\gamma^B_{1671} + \gamma^B_{1671}\gamma^A_{1671} = 2\,\eta^{AB}_{1671}$: $\mathrm{diag}(-1,-1,-1,-1,+1,-1,-1,-1)$, time at index 4. The flat-space part of the project uses it as the metric (time-like $= +1$). It is minus the flat metric $\eta^{(7+1)}$ of the author's sign convention for the deflating spacetimes (space-like $= +1$)	N02 `η17` with the index values 0 and 4 exchanged; NOT `η71`
τ^A_{1671}	`tau1671`	the real 8×8 matrices of the 7+1 construction: $\tau^4_{1671} = 1_8$, the other seven represent the imaginary octonion units	N02 `τ0[swap04(A)]`

symbol	names in programs and data files	meaning	origin
γ^A_{1671}	`gamma1671`	the eight real 16×16 Dirac matrices of the 7+1 spacetime (contravariant index)	N02 `Γ17up[swap04(A)]`; NOT `Γ71[A]`
Σ^{AB}_{1671}	`Sigma1671`	$\frac{1}{4}[\gamma^A_{1671}, \gamma^B_{1671}]$, the scaled commutators; they span $\mathfrak{so}(7,1)$	N02 `Σ17[A,B]`, built from the `Γ17` with lower index: $\Sigma^{AB}_{1671} = \eta^{AA}_{1671}\eta^{BB}_{1671}$ `Σ17[A', B']` with $A' = \mathrm{swap}_{04}(A)$, $B' = \mathrm{swap}_{04}(B)$
ω_{1671}	`omega1671`	the product $\gamma^0_{1671} \cdots \gamma^7_{1671}$; real, $\omega^2_{1671} = -1$	N02 `Γ17[8]`, `ω17`
χ_{1671}	`chi1671`	$i\,\omega_{1671}$, the (complex) chirality matrix of 7+1	N02 `χ17`; NOT `χ71`
P^L_{1671}, P^R_{1671}	`PL1671` `PR1671`	the complex chiral projectors $\frac{1}{2}(1 \mp \chi_{1671})$	N02 `PL17`, `PR17`; NOT `PL71`, `PR71`
U^c_{1671}	`Uc1671`	the unitary matrix $(1 - i\gamma^4_{1671})/\sqrt{2}$ that carries the complex projectors of 7+1 into the real projectors of 4+4	new
C^+_{1671}, C^-_{1671}	`Cp1671` `Cm1671`	γ^4_{1671} and $\gamma^4_{1671}\omega_{1671}$; $C^\pm \gamma^A (C^\pm)^{-1} = \pm(\gamma^A)^T$	new
β_{1671}, α^j_{1671}	`beta1671` `alpha1671`	the real symmetric matrices γ^4_{1671} and $\gamma^4_{1671}\gamma^j_{1671}$ (seven space indices j) of the Hamiltonian	N02 `βD`, `αD[swap04(j)]`
h_{1671}	`h1671`	Hamiltonian matrix of the 7+1 theory, $\alpha^j_{1671} p^j + \beta_{1671} m_{1671}$	N02 `h`
Π_{1671}	`Pi1671`	the projector $(\gamma^A_{1671} k_A + m_{1671})/2m_{1671}$ of the worked example	new
E^L_{1671}, E^R_{1671}	`EL1671` `ER1671`	the chiral frames: 16×8 matrices whose columns span the two eigenspaces of χ_{1671}	N02 `EL`, `ER`
M_{1671}, U^+_{1671}, V^-_{1671}	`M1671` `Upos1671` `Vneg1671`	the 8×8 matrix $m_{1671} + p^j \tau^j_{1671}$ and the matrices of positive- and negative-energy spinors of the worked example (check `71-D8`)	N02 `M` of its Sec. 7.4 (in N02 the letter `M` also denotes other matrices and a mass); the two spinor matrices are new names

4 Fields and their equations

symbol	names in programs and data files	meaning	origin
ψ_{1644}	`dirac1644`	the field dirac1644: 16 complex Grassmann components on the 4+4 spacetime; $(\gamma^A_{1644}\partial_A - m_{1644})\psi_{1644} = 0$	new; `Ψ16` of N01 and N10–N12 is a column of 16 real quantities (in N10–N12 functions of `x0`, `x4` called the wave function of the universe), not this field
ψ^L_{1644}, ψ^R_{1644}	`psiL1644` `psiR1644`	the two 8-component chiral parts of dirac1644 (upper and lower eight components)	new
m_{1644}	`m1644`	mass of the field dirac1644	new
$\tilde{\psi}$	—	the covariant adjoint $\psi^\dagger C^-_{1644}$ of the 4+4 theory (invariant under $\mathrm{Spin}_0(4,4)$)	new
j^A_{1644}	`j1644`	the conserved current $i\,\tilde{\psi}\gamma^A_{1644}\psi$ of the covariant 4+4 theory (fermion number)	new

symbol	names in programs and data files	meaning	origin
ψ_{1671}	`dirac1671`	the field dirac1671: 16 complex Grassmann components on the 7+1 spacetime; $(i\gamma^A_{1671}\partial_A - m_{1671})\psi_{1671} = 0$	N02 `dirac16complex`
ϕ^L_{1671}, ϕ^R_{1671}	`phiL1671` `phiR1671`	the two 8-component chiral parts of dirac1671	N02 `φL`, `φR`
m_{1671}	`m1671`	mass of the field dirac1671	N02 `m`, `mD`
$\bar{\psi}$	—	the adjoint $\psi^\dagger\beta$ of the quantized theories ($\beta = \beta_{1671}$ or β_{1644}); in 4+4 it differs from $\tilde{\psi}$	new
β, α^j, χ, h, m	—	in the sections common to both spacetimes: the matrices and the mass of the spacetime at hand, i.e. the tagged symbols with the tag dropped; $\chi \equiv \chi_{1671}$ in 7+1 and $\chi \equiv \omega_{1644}$ in 4+4	new
$U(\vec{p})$	—	the 16×8 matrix of orthonormal positive-energy spinors at momentum $\vec{p}$	new

5 Symmetries, pairs and order parameter

symbol	names in programs and data files	meaning	origin
$U(1)_F$	—	the phase symmetry $\psi \to e^{i\epsilon}\psi$: fermion number	new
$U(1)_{VS}$	`UVS`	the chiral symmetry $\psi \to e^{i\epsilon\chi}\psi$, $\Phi_{VS} \to e^{-2i\epsilon}\Phi_{VS}$, broken by the vacuum value of the Vorton-Scalar	new
$SU(2)_A$, $SU(2)_B$	—	the two factors of the internal symmetry of the fields that do not depend on the four extra coordinates (zero-mode sector), in both spacetimes; $SO(4) = [SU(2)_A \times SU(2)_B]/Z_2$	new
$SU(2)_X$, $SU(2)_Y$	—	the two factors of the rotation group of the four space coordinates $x0, \dots, x3$ of the 4+4 spacetime (NOT the internal $SU(2)_A \times SU(2)_B$)	new
$SO(3)_T$, $SU(2)_T$	—	the rotations of the three additional times $x5, x6, x7$ of 4+4, and its covering group; the diagonal subgroup of $SU(2)_A \times SU(2)_B$	new
$U(1)_E$, P (generator)	—	the second central phase symmetry of the zero-mode sector and its traceless generator, $P^2 = 1$: $P = +1$ on the doublets of $SU(2)_A$, -1 on those of $SU(2)_B$	new
Δ^A_{1644}, Δ^A_{1671}	`DeltaV1644` `DeltaV1671`	the vector pair operators $\psi^T C^- \gamma^A \psi$ of each spacetime	new
Δ^{AB}_{1671}	`DeltaT1671`	the two-form pair operators $\psi^T C^+ \gamma^A \gamma^B \psi$ of the 7+1 spacetime	new
Δ^a	—	order parameter: expectation value of the vector pair operator along the extra axis $a \in \{0, 5, 6, 7\}$; $\lvert\Delta\rvert^2 \equiv \sum_a \lvert\Delta^a\rvert^2$	new
$\Phi_X(\vec{p})$	—	two-particle content $U(\vec{p})^T X U(-\vec{p})$ of the pair operator $\psi^T X \psi$	new
M	`M_gap`	the 8×8 gap matrix on the internal states of a pair (sans-serif; `M_gap` in the statements of the programs)	new
$d_\pm$	—	the two fourfold gaps of a general state, $d^2_\pm = \lvert\Delta\rvert^2 \pm \sqrt{\lvert\Delta\rvert^4 - \lvert\eta_{ab}\Delta^a\Delta^b\rvert^2}$	new
n^a	—	real unit vector: direction of the order parameter in the space of the four extra axes	new

6 The Vorton-Scalar (suffix VS)

symbol	names in programs and data files	meaning	origin
Φ_{VS}	`PhiVS`	the Vorton-Scalar: a complex scalar field	new (the name is that of the author's project NCSS, where the scalar is real)
v_{VS}	`vVS`	vacuum value: $\langle\Phi_{VS}\rangle = v_{VS}/\sqrt{2}$	new
λ_{VS}	`lambdaVS lamVS`	self-coupling of the Vorton-Scalar	new
y_{VS}	`yVS`	Yukawa coupling of the Vorton-Scalar to the fermion field	new
s_{VS}	`sVS`	the radial mode of the Vorton-Scalar (real field)	new
m_s	`msVS`	mass of the radial mode, $\sqrt{2\lambda_{VS}}\,v_{VS}$	new
ϑ_{VS}	`thetaVS thVS`	the phase mode of the Vorton-Scalar (real field, massless)	new
g_{VS}	`gVS`	coupling of both modes of the Vorton-Scalar to the fermions, $g_{VS} = y_{VS}/\sqrt{2} = m_Y/v_{VS}$, where m_Y is the part of the fermion mass that the Vorton-Scalar gives (equal to m/v_{VS} when there is no bare Dirac mass)	new
α_{VS}	`alphaVS aVS`	$g_{VS}^2/4\pi$	new
D_{VS}	`DVS`	$\alpha_{VS}\,m/m_s$: two fermions bind if and only if $D_{VS} > D_c$	new
J^A_{VS}	`JVS`	the neutral current of the Vorton-Scalar (bilinear in Φ_{VS})	new
J^A_5	`J5`	the axial current $\bar\psi\gamma^A\chi\psi$ of the fermion field	new
G_{VS}	`GVS`	constant of the Fermi type of the neutral-current interaction, $G_{VS}/\sqrt{2} = 1/(2v_{VS}^2)$	new
$\mathcal{L}^{VS}_{1644}$, $\mathcal{L}^{VS}_{1671}$	`LVS`	the Lagrangians that define the Vorton-Scalar in the two spacetimes	new
C (coupling)	—	$g_{VS}^2 m^2/m_s^2 = g_{VS}^4/2\lambda_{VS}$, the one number on which the fermion mass in matter depends	new

7 Pairing, gap equation and CJT effective action

symbol	names in programs and data files	meaning	origin
ν	—	the number of internal states of a particle of given momentum, $\nu = 8$	new
F_S, F_P	`F_scalar F_pseudoscalar`	exchange amplitudes of the radial and of the phase mode in a pair channel	new
$V_{\rm NC}$, $V_s(r)$	—	contact interaction of the neutral current in the hidden-axis channel; Yukawa potential of the radial mode	new
σ_X	—	the sign $+1$ (vector pairs along the extra axes) or -1 (their two-form pairs) of the phase-mode term of the kernel	new
$W(k,k')$	—	kernel of the gap equation: the s-wave average of the pairing attraction, $W(k,k') = \frac{1}{2}\int_{-1}^{1} d\cos\theta\,\big[-V_{VS} - V_{\rm ct}\big]$ for the momenta k, k' of the pair before and after the scattering (θ: the angle between them); without contact couplings the exchange attraction alone. Always written with its two arguments (W_{1644} is a matrix)	new
$\Delta(k)$, Δ_F	—	gap function; its value at the Fermi momentum	new
ξ_k, E_k, $\kappa(k)$	—	$\sqrt{m^2+k^2} - \mu$; quasi-particle energy $\sqrt{\xi_k^2 + \Delta(k)^2}$; pair correlation $\langle b(-\vec k)b(\vec k)\rangle$	new

symbol	names in programs and data files	meaning	origin
k_F, E_F, μ	`kF` `mu`	Fermi momentum; Fermi kinetic energy $\sqrt{m^2+k_F^2}-m$; chemical potential. The number density is $n=\nu V_D k_F^D/(2\pi)^D$ in D space dimensions ($n=\nu k_F^3/6\pi^2$ for $D=3$)	new
$N(0)$	—	density of states at the Fermi surface per internal state, $k_F\sqrt{m^2+k_F^2}/2\pi^2$	new
λ, λ_S, λ_P	—	dimensionless pairing coupling $N(0)W(k_F,k_F)$ and its parts from the radial and the phase mode	new
T_c	—	critical temperature of the paired state	new
D_c	—	critical value of D_{VS} for a bound pair in the Yukawa potential	new
$a_{\rm sc}$, a_{BB}	—	scattering length of two fermions; scattering length of two bound pairs	new
$\Gamma[\phi,G]$, Γ_2	—	the effective action for composite operators of Cornwall, Jackiw and Tomboulis; its two-particle-irreducible part	new (method of the cited paper)
$\mathcal{S}$, D_s, D_ϑ	—	propagator of the doubled fermion field; propagators of the radial and of the phase mode	new
$V[\Delta]$	—	the CJT effective potential of the paired state, measured from the unpaired state	new
$\eta(T)$	—	largest eigenvalue of the linearized gap equation, $\eta(T_c)=1$	new
$A_2(T)$, A_4	—	coefficients of the expansion of the CJT effective potential V along a gap function of fixed shape	new
$a(T)$, b, $K_{\rm GL}$	—	Ginzburg-Landau coefficients of the weak-coupling free energy (quadratic, quartic and gradient term)	new
c_s	—	speed of first sound of the superfluid, $c_s^2=dP/d\rho$	new

8 Matter, temperature and vacuum

symbol	names in programs and data files	meaning	origin
ϕ	—	expectation value of the Vorton-Scalar, $\langle\Phi_{VS}\rangle=\phi/\sqrt{2}$; the fermion mass is $g_{VS}\phi$	new
m^*, x	`M` `M_ks` `M_cjt` `M_cjt_global` `M_plus` `M_minus` `M_relaxed` `x` `Mp_ks` `Mp_cjt` `Mm_ks` `Mm_cjt` `Mp_below` `Mm_below` `Mp_above` `Mm_above` `M_large` `M_small`	m^*: the mass of the fermions in matter; $x=\phi/v_{VS}$: the expectation value of the Vorton-Scalar in units of its vacuum value. With a contact interaction $m^*=m+G_S\,n_S$ (gap equation of the Kohn-Sham and CJT calculations); with the Vorton-Scalar and a bare Dirac mass $m^*=m_D+m_Y\,x+G_S\,n_S$, so that $x=m^*/m$ if the whole mass comes from the Vorton-Scalar and no contact coupling acts. m_+^*, m_-^*: the masses of the particles of the two halves $W_{1644}=\pm1$ of the 4+4 field, $m_\pm^*=m+G_S^{\rm tot}\,n_S\pm G_S^{\rm dif}\,n_{SW}$ in both forms of the self-interaction (with the filled states of negative energy: the masses of the two halves in vacuum). The data files and the statements of the programs write m^* as `M` and $m_\pm^*$ as `M_plus`, `M_minus` (`Mp_..`, `Mm_..` with a suffix that names the method or the strength; `M_large`, `M_small`: the larger and the smaller of the two); the author's document KOHN_SHAM_GRAVITY writes M	new
$\varepsilon(k_F,x)$, $V_{\rm sea}(x)$	—	energy density of matter as a function of the Fermi momentum and of $x=m^*/m$ (in units of m^4 in ordinary space); renormalized energy of the filled negative-energy states. ε without arguments is the energy density of matter in the units of the calculation at hand	new

symbol	names in programs and data files	meaning	origin
B	—	energy density of the state without expectation value of the Vorton-Scalar (bag constant)	new
E_x	—	exchange (Fock) energy of the two modes	new
M_s, M_ϑ, a_s, a_ϑ	—	thermal masses of the two boson modes in the Hartree approximation, and the curvatures U''_{eff}, U'_{eff}/ϕ	new
$P(M)$, $Q(M)$	—	thermal part of the boson loop and of $\frac{1}{2}\mathrm{Tr}\ln$	new
T_{high}, T_N, $T_{1/2}$, T_0	—	two closed estimates of the temperature at which the fermion mass vanishes; temperatures at which the expectation value has fallen to half and to zero	new
$\sigma(p)$	—	self-energy that a pair condensate in the vacuum gives to the fermions: mass splitting $m \pm \sigma$	new
α_c, Λ (cutoff)	`Lambda`	critical value of α_{VS} for a pair condensate in the vacuum; Λ: a momentum cutoff (of the ladder equation in the vacuum, and of the filled states of negative energy in the calculations with contact interaction). Λ_8 and $\bar{\Lambda}$ are cosmological terms and have their own entries	new

9 Cosmology

symbol	names in programs and data files	meaning	origin
N	—	density of the dark radiation today in units of the density of one massless neutrino species	new
z_t, T_d	—	transition redshift of the stepped dark fluid, $m = T_{d0}(1+z_t)$; temperature of the dark sector	new
r_{step}	—	factor by which the density of the dark radiation rises when the fermions annihilate, $15^{1/3}$ for this model	new
ω_b, ω_c, h, H_0, A_s, n_s, τ	`tau_reio`	the six parameters of the standard cosmological model: densities of baryons and of cold dark matter (Ωh^2), Hubble constant $H_0 = 100\,h$ km s^{-1} Mpc^{-1}, amplitude and spectral index of the primordial fluctuations, optical depth of reionization (`tau_reio` in the data files; the letter τ with the index d is the deflation clock)	standard notation
$\Delta\chi^2$	—	worsening of the best fit to the mock data relative to the standard model	new
θ_s, r_s, D_A	—	angular size of the sound horizon; sound horizon; distance to last scattering	standard notation
c_{s0}	—	sound speed of superfluid dark matter today at the mean cosmic density	new
ρ_0, R_{TF}	—	constant of the equation of state of a condensate of bound pairs; Thomas-Fermi radius of a core supported by its pressure	new
m_{deg}, $m_{B,\text{max}}$	—	largest fermion mass for degeneracy, $m_{\text{deg}} = [6\pi^2\hbar^3\rho/(\nu\sigma_v^3)]^{1/4}$, and largest mass m_B of a boson for condensation, $m_{B,\text{max}} = [(2\pi)^{3/2}\hbar^3\rho/(2.612\,\sigma_v^3)]^{1/4}$, in a halo of the mass density ρ and the velocity dispersion σ_v	new
σ_v; κ_{SI}	—	the velocity dispersion of the dark matter of a halo (with the index v; σ_8 is the amplitude of clustering and $\sigma(p)$ a self-energy); the bound on the cross-section per mass of dark matter from colliding clusters, about 1 cm^2/g (with the index SI; κ_8 is the gravitational coupling)	new

D1 Deflating spacetimes: hypotheses and geometry

symbol	names in programs and data files	meaning	origin
g55, g66, g77	`g55` `g66` `g77`	the three rates of deflation of $x5$, $x6$, $x7$ (positive constants, dimension 1/length). These are the author's names; they are written upright and never with a subscript. They are *not* components of the metric: the components are g_{55}, g_{66}, g_{77}	author
rate_t	`rate_t` `rate`	the rate of the coordinate xt in a statement that holds for $t = 5, 6, 7$ alike: $\mathrm{rate}_5 = \mathrm{g55}$, $\mathrm{rate}_6 = \mathrm{g66}$, $\mathrm{rate}_7 = \mathrm{g77}$	new
gsum	`gsum` `gam`	$\mathrm{g55} + \mathrm{g66} + \mathrm{g77}$, the total rate: the proper volume of the three deflating coordinates falls as $e^{-\mathrm{gsum}\, x4}$ (`gam`: gsum in units of 100 km/s/Mpc)	new
gsq, gprod	`gsq` `gprod`	$\mathrm{g55}^2 + \mathrm{g66}^2 + \mathrm{g77}^2$ and $\mathrm{g55}\,\mathrm{g66} + \mathrm{g55}\,\mathrm{g77} + \mathrm{g66}\,\mathrm{g77}$; $\mathrm{gsum}^2 = \mathrm{gsq} + 2\,\mathrm{gprod}$	new
const5, const6, const7; const_t	`const5` `const6` `const7` `const_t` `cst`	the three positive constants of the author's metric (his word: const): the values of the scale factors at $x4 = 0$	author
b_5, b_6, b_7; b_t	`b_5` `b_6` `b_7` `b_t`	the scale factors of the deflating coordinates, $b_t(x4) = \mathrm{const}_t\, e^{-\mathrm{rate}_t\, x4}$: a coordinate distance dxt is the proper length $b_t\,\lvert dxt\rvert$. Always written with the index 5, 6, 7 or t (the letter b without index is a Ginzburg-Landau coefficient, and b in sans-serif type is the common scale factor of the three deflating coordinates in the Einstein-Lovelock evolution)	new
V_d	`Vd`	$b_5 b_6 b_7 = \sqrt{\lvert\det g\rvert}$, the proper volume of a unit coordinate cell of the deflating coordinates; $V_d \propto e^{-\mathrm{gsum}\, x4}$	new
μ, ν, ρ, σ (indices)	—	coordinate indices with the values $0, \ldots, 7$ on the deflating spacetimes (only as sub- or superscripts; as stand-alone letters μ is the chemical potential, ν the number of internal states and λ the pairing coupling). The capital indices A, B, C, D, also with a prime, label the orthonormal frame: the frame vector A points along the coordinate xA (as a stand-alone letter C is the coupling $g_{VS}^2 m^2/m_s^2$, and D the number of space dimensions with momentum). In the statements of the programs the indices are spelled out inside tensor expressions, as in `g^(mu nu)`	new
t, s (indices)	—	indices of the three additional coordinates $x5, x6, x7$, values 5, 6, 7; s is a second such index where two are needed (the matrices s_h carry the index h)	new
${e_A}^{\mu}$, ${e^A}_{\mu}$	`e_A^mu`	the orthonormal frame and its inverse: ${e^A}_{\mu} = \mathrm{diag}(1, 1, 1, 1, 1, b_5, b_6, b_7)$	new
${\Gamma^{\rho}}_{\mu\nu}$	`Gamma` `christoffel`	the Christoffel symbols of the deflating metric (three indices; $\Gamma[\phi, G]$ with square brackets is the CJT effective action). The author's document KOHN_SHAM_GRAVITY writes the Christoffel symbols with braces and uses Γ^A for the Dirac matrices, which are γ^A here	new
${R^{\rho}}_{\sigma\mu\nu}$, ${R^{\mu}}_{\nu}$, R	`riemann` `ricci` `Ric44` `Ric55` `Ric66` `Ric77` `Rscalar`	Riemann tensor, Ricci tensor and curvature scalar of the deflating metric	new
${\mathcal{G}^{\mu}}_{\nu}$	`Ein00` `Ein44` `Ein55` `Ein66` `Ein77` `einstein_mixed`	the Einstein tensor ${R^{\mu}}_{\nu} - \frac{1}{2} R\, {\delta^{\mu}}_{\nu}$ (script letter, so that it cannot be taken for Newton's constant or for a constant of the Fermi type)	new
$\mathcal{X}$; $\mathcal{X}_{\mathrm{scale}}$, $\mathcal{X}_{(s)}$	`killing_vectors` `killing_count`	a Killing vector field of the deflating metric (script letter; the plain letter X is the set of the eight coordinates); the scaling vector $\partial_4 + \sum_t \mathrm{rate}_t\, xt\, \partial_t$; the three special Killing vectors of the case of equal rates, $s = 5, 6, 7$	new
$\bar{x}4$; $\bar{x}5$, $\bar{x}6$, $\bar{x}7$	—	coordinates of the case of equal rates in which the line element of $x4, \ldots, x7$ is a multiple of a flat one: $\bar{x}4 = e^{\mathrm{g55}\, x4}/\mathrm{g55}$, $\bar{x}t = \mathrm{const}_t\, xt$	new

symbol	names in programs and data files	meaning	origin
$\mathcal{H}^{\mu}{}_{\nu}$; α_{GB}, Λ_8	`Lanczos` `alpha_GB` `Lambda_8`	the Lanczos tensor of the Gauss-Bonnet term, $\mathcal{H}^{\mu}{}_{\nu} = 2(R\,R^{\mu}{}_{\nu} - 2R^{\mu}{}_{\alpha}R^{\alpha}{}_{\nu} - 2R^{\mu\alpha}{}_{\nu\beta}R^{\beta}{}_{\alpha} + R^{\mu\alpha}{}_{\beta\gamma}R^{\beta\gamma}{}_{\nu\alpha}) - \frac{1}{2}\delta^{\mu}_{\nu}(R^2 - 4R^{\alpha}{}_{\beta}R^{\beta}{}_{\alpha} + R^{\alpha\beta}{}_{\gamma\delta}R^{\gamma\delta}{}_{\alpha\beta})$ (script letter; H is the Hubble rate); the coupling of the Gauss-Bonnet term and the cosmological constant of Einstein-Gauss-Bonnet gravity in eight dimensions, whose equations without matter are $\mathcal{G}^{\mu}{}_{\nu} + \Lambda_8\,\delta^{\mu}_{\nu} + \alpha_{GB}\,\mathcal{H}^{\mu}{}_{\nu} = 0$ (metric with positive space-like components). In the author's Einstein-Lovelock notation $E^{(1)} = -\mathcal{G}$, $E^{(2)} = \mathcal{H}$, $\alpha_2 = -\alpha_{GB}$, $\bar{\Lambda} = \Lambda_8$	new
K_{AB}	`ksec`	the sectional curvature of the plane of the frame directions A and B (metric with positive space-like components): on the deflating spacetimes $K_{4t} = \text{rate}_t^2$, $K_{ts} = \text{rate}_t\,\text{rate}_s$ for $t \neq s$, zero if one direction is flat; for scale factors that depend on $x4$ in any way, $K_{4i} = H_i' + H_i^2$ and $K_{ij} = H_iH_j$ with the rate H_i of the direction i (the pair curvatures of the author's Einstein-Lovelock equations). Always with two subscripts: the letter K without subscript is the generator of the evolution of the 4+4 field in $x4$, K as an index is an index set of the self-interaction, and K with an order as subscript and with an argument is a modified Bessel function	new
ς	`affine`	affine parameter along a geodesic; for a massive particle on the 7+1 spacetime, the proper time divided by the mass	new
P_{μ}	`P_1` `P_5`	the conserved covariant momenta of a geodesic (with an index; P without index is the generator of $U(1)_E$)	new
$x4_{\text{turn}}$	`x4_turn`	the value of the ordinary time at which a classical particle with momentum along a deflating time turns back, and beyond which a mode that depends on a deflating time is extinguished (4+4): there $k_d = E_{\perp}$	new

D2 Deflating spacetimes: matrices, fields and their equations

symbol	names in programs and data files	meaning	origin
γ^{μ}_{1644d}, γ^{μ}_{1671d}	`gamma_1644d` `gamma_1671d`	the Dirac matrices of the deflating spacetimes in coordinate components, $\gamma^{\mu} = e_A{}^{\mu}\gamma^A$: $\gamma^t = \gamma^{A=t}/b_t$ for $t = 5, 6, 7$, the other five are the constant matrices γ^A_{1644}, γ^A_{1671} of the frame	new
$\omega_{\mu AB}$	`omega_(mu A B)`	the coefficients of the spin connection, $\omega_{\mu AB} = \eta_{AC}\,e^C{}_{\nu}(\partial_{\mu}e_B{}^{\nu} + \Gamma^{\nu}{}_{\mu\lambda}e_B{}^{\lambda})$ with the flat metric $\eta = \eta^{(4+4)}$ or $\eta^{(7+1)}$ (three indices; ω_{1644}, ω_{1671} with a tag are the products of all eight Dirac matrices)	new
Ω_{μ}	`Omega_mu` `omega`	the spin connection as a matrix, $\Omega_{\mu} = \frac{1}{4}\,c_{\gamma}\,\omega_{\mu AB}\gamma^A\gamma^B$; the covariant derivative of a spinor is $\nabla_{\mu}\psi = (\partial_{\mu} + \Omega_{\mu})\psi$. On both deflating spacetimes $\Omega_t = \frac{1}{2}\,\text{rate}_t\,b_t\,\gamma^t\gamma^4$ (frame matrices) for $t = 5, 6, 7$ and $\Omega_{\mu} = 0$ otherwise. (Ω_{CJT} is an effective potential)	new
ψ_{1644d}, ψ_{1671d}	`dirac1644d` `dirac1671d`	the fields dirac1644d and dirac1671d: the 16-component fields on the deflating spacetimes. With mass term and self-interaction they obey $(\gamma^{\mu}\nabla_{\mu} - m)\psi_{1644d} = \sum_r g_r \sum_{\lvert K\rvert=r}(\tilde{\psi}\Theta_K\psi)\Theta^K\psi$ (covariant form of 4+4) and $(i\gamma^{\mu}\nabla_{\mu} - m)\psi_{1671d} = \sum_r g_r \sum_{\lvert K\rvert=r}(\bar{\psi}\Theta_K\psi)\Theta^K\psi$ (7+1)	new

symbol	names in programs and data files	meaning	origin
$\hat{\psi}$	psihat	the weighted field $\sqrt{V_d}\,\psi$, which obeys the equation without spin connection	new
$K(x4)$, $\mathsf{h}(x4)$	K h	the generator K_{1644} and the Hamiltonian matrix h_{1671} with the wavenumbers along the deflating coordinates divided by the scale factors: without self-interaction $\partial_4\hat{\psi} = K\hat{\psi}$ (4+4), $i\partial_4\hat{\psi} = \mathsf{h}\hat{\psi}$ (7+1)	new

D3 Deflating spacetimes: solutions mode by mode

symbol	names in programs and data files	meaning	origin
k_5, k_6, k_7; k_t	k_5 k_6 k_7 k_t	the conserved wavenumbers of a mode along the deflating coordinates (covariant components of the wave vector k_A)	new
$E_\perp$	E_perp	the energy of a mode, or of a classical particle, without its motion along the deflating coordinates: $\sqrt{m^2 + k_0^2 + k_1^2 + k_2^2 + k_3^2}$ (with the wavenumbers or momenta along $x0, \dots, x3$)	new
k_d	k_d	the physical wavenumber along the deflating coordinates, $k_d(x4) = \left[\sum_t k_t^2/b_t(x4)^2\right]^{1/2}$; it grows as the coordinates deflate	new
$\mathsf{h}_\perp$, $K_\perp$	h_perp K_perp	the parts of h and of K that do not contain the scale factors: the operators of the modes that do not depend on the deflating coordinates, $\mathsf{h}_\perp = m\beta + \sum_a p_a\alpha^a$ with $\mathsf{h}_\perp^2 = E_\perp^2$, and $K_\perp = -m\gamma^4 + i\sum_a k_a\gamma^4\gamma^a$ with $K_\perp^2 = -E_\perp^2$	new
γ^d, α^d	—	the frame matrix along the direction of the wave vector in the deflating coordinates, $\gamma^d = k_d^{-1}\sum_t (k_t/b_t)\gamma^t$, and $\alpha^d = \gamma^4\gamma^d$; constant in time when one rate governs	new
w; f_+, f_-; Z_+, Z_-	—	auxiliary objects of the exact solutions of the equations without self-interaction: a constant eigenvector of $\mathsf{h}_\perp$ or of $K_\perp$; the two coefficient functions of the ansatz; two cylinder functions whose orders differ by one	new
u_d	u_d	k_d/rate_t for a mode that depends on one deflating coordinate: the argument of the Bessel functions of the exact solutions	new
n_{cr}	n_cr n_min n_mean n_max n_exact fd5 fd6	the number of particles per internal state that the deflation creates in a mode that starts empty (7+1)	new
T_{defl}	—	the temperature of the thermal spectrum of the created quanta, $\mathrm{rate}_t/2\pi$ in units with $\hbar = k_B = 1$ (7+1)	new
S_{WKB}	S_wkb	$\int \sqrt{k_d^2 - E_\perp^2}\, dx4$ from the turning value on: exponent of the extinction of a bounded solution (4+4)	new
φ_{bos}, $\hat{\varphi}_{\mathrm{bos}}$, m_{bos}	phi_bos Shat Shat2 m_bos	a mode of the Vorton-Scalar (the radial mode, $m_{\mathrm{bos}} = m_s$, or the phase mode, $m_{\mathrm{bos}} = 0$), its weighted form $\sqrt{V_d}\,\varphi_{\mathrm{bos}}$, and its mass	new
ϖ_b	varpi_b	$\sqrt{k^2 + m_{\mathrm{bos}}^2 - \mathrm{gsum}^2/4}$: the frequency of the weighted boson mode that does not depend on the deflating coordinates	new

D4 The deflation clock and the couplings

symbol	names in programs and data files	meaning	origin
τ_d	`tau_d`	the deflation clock, $\tau_d = \mathrm{gsum}\,(x4 - x4_{\mathrm{ref}})$: the number of e-foldings by which the volume V_d has shrunk since the reference time; in an evolution in which the rates change, $\tau_d = -\ln(V_d/V_d^{\mathrm{ref}})$ (the letter τ without the index d is the optical depth of reionization)	new
τ_{pair}	`tau_pair`	the clock measured from the moment at which two fermions become bound: $\ln(D_{VS}/D_c)$	new
$\hat{\Phi}_{VS}$	`Phihat`	the weighted Vorton-Scalar $\sqrt{V_d}\,\Phi_{VS}$	new

D5 Dynamics of the paired state

symbol	names in programs and data files	meaning	origin
Ξ_d	`Xi_d`	$\mathrm{gsum}/(\lambda\,\Delta_{\mathrm{eq}})$: decides whether the paired state can follow the deflation ($\Xi_d \ll 1$: yes)	new
n_k	—	occupation of the pair of states of the single-particle energy ξ_k in the shell model of the paired state (with the index k; the letter n without index is a density)	new
Δ_{eq}	`gap_eq`	the equilibrium gap for the pairing strength of the moment	new
E_{sh}	`E_sh`	half-width in energy of the shell of states around the Fermi surface in the model of the pair dynamics; unit of energy and of inverse time there	new
n_{ex}	`n_ex` `n_ex_final`	density of broken pairs (in units of E_{sh} times the density of states)	new

D6 Cosmology and bounds on the deflating spacetimes

symbol	names in programs and data files	meaning	origin
G_N, G_8	`G_N` `GN_early` `GN_star` `G_8`	Newton's constant of ordinary space, and the gravitational coupling of the eight-dimensional Einstein theory (assumption A2): $G_N = G_8/(\text{proper volume of the extra coordinates}) \propto e^{\mathrm{gsum}\,x4}$	new
a_{sp}, H	—	scale factor of ordinary space (assumption A1) and its rate $H = (da_{\mathrm{sp}}/dx4)/a_{\mathrm{sp}}$; H_0 is its value today. (The rates of the Einstein-Lovelock evolution are H_{a}, H_{b}; the author's Einstein-Lovelock notebooks use the letter H for the rate in the exponent of their metric, which corresponds here to g55, g66, g77)	new
t_0	`age` `age_Gyr`	the age of the universe in the model at hand	new
θ_*, z_*	`theta_star` `z_star`	the angle under which the sound horizon at last scattering is seen, $r_s(z_*)/D_A(z_*)$ with the comoving distance D_A, and the redshift z_* of last scattering at which the optical depth without reionization is 1 (the $100\,\theta_*$ of the Planck papers). The angle θ_s is the same ratio at the maximum of the visibility function; for the model without deflation $100\,\theta_* = 1.03965$ and $100\,\theta_s = 1.040358$	new
δ_m, f_g, σ_8	`growth` `f_growth` `growth_ratio` `f_growth_ratio` `f_sigma8_ratio`	density contrast of matter in ordinary space; its growth rate $d\ln\delta_m/d\ln a_{\mathrm{sp}}$ today; the amplitude of clustering (only ratios of σ_8 are used)	new

symbol	names in programs and data files	meaning	origin
v_{EW}, n_{EW}	`v_EW` `n_EW`	the electroweak scale (the vacuum value of the Higgs field of the Standard Model, 246 GeV), and the exponent with which it would follow the coupling under the hypothesis H2: $n_{\mathrm{EW}} = d\ln v_{\mathrm{EW}}/d\ln\alpha_{VS}$	new
ϱ_c	`rho_c`	abbreviation in the formula for n_{EW} of the scale-invariant ladder equation: $\varrho_c = \frac{1}{2}\sqrt{\alpha_{VS}/\alpha_c - 1}$ (the letter ρ without index is a density)	new
m_p/m_e	`mu_pe`	the ratio of the proton mass to the electron mass (always written as this ratio in the documents; `mu_pe` in the statements of the programs)	new
G_{Fermi}, α_{em}	`G_Fermi`	the Fermi constant of the weak interaction and the fine-structure constant (they enter only in the discussion of bounds)	new

D7 Symmetries and vortex lines on the deflating spacetimes

symbol	names in programs and data files	meaning	origin
P_c	`Pc`	$i\alpha^1\alpha^2\alpha^3\chi$, the product of the five matrices of the sector that does not depend on the extra coordinates; equal, up to sign, to the generator P of $U(1)_E$	new
θ_{12}	—	the angle around a vortex line that lies along $x3$, in the plane of $x1$ and $x2$ (with the index 12; θ_* is the angle of the sound horizon and $\theta(\cdot)$ the step function)	new
w_z	`w_z`	a constant spinor of a fermion zero mode on a vortex line	new

D8 Tags and identifiers of the deflating spacetimes

symbol	names in programs and data files	meaning	origin
1644d, 1671d	`1644d` `1671d`	tags of the deflating 4+4 and 7+1 spacetimes and of every object defined on them (the tags 1644, 1671 of the flat spacetimes with the letter d)	new
A1, A2, A3, H2, HV	—	A1, A2: the two assumptions of the cosmological part (ordinary space expands; gravity is the eight-dimensional Einstein theory). A3: the assumption about the perturbations of ordinary space (the scale factors of the extra coordinates are not perturbed, and the agent that holds them has no energy density), with its closures. H2: the author's hypothesis that the superfluid state is the Higgs field. HV: the different assumption that the vacuum value of the Vorton-Scalar sets the electroweak scale	new
"Momentum", "Pressure", "Shear" (closures)	`Momentum` `Pressure` `Shear` `closure`	the three closures of the perturbation equations under assumption A3, named after the Einstein equation of ordinary space that holds with matter and radiation alone besides the energy constraint: the momentum constraint, the trace (pressure) equation, or the equation of the shear. In data files the suffixes `_M` and `_P` mark the first two; `_N4` marks the four-dimensional growth equation, which is none of them	new

D9 Perturbations of ordinary space and the microwave background

symbol	names in programs and data files	meaning	origin
$\tau_{\rm cf}$	`tau_cf` `tau0_Mpc`	the conformal time of ordinary space, $d\tau_{\rm cf} = dx4/a_{\rm sp}$, in Mpc (`tau0_Mpc`: its value today). In the equations of the perturbations a prime is the derivative with respect to $\tau_{\rm cf}$	new
$H_{\rm cf}$, H_e	`Hc` `He` `hub` `hub_e` `a_gsum_over_Hc`	the conformal Hubble rate $H_{\rm cf} = a_{\rm sp}H$, and $H_e = H_{\rm cf} - a_{\rm sp}\,$gsum$/2$, which takes its place in three of the Einstein equations of ordinary space when the extra coordinates deflate	new
k (perturbations), ℓ, D_ℓ^{TT}, D_ℓ^{TE}, D_ℓ^{EE}	—	the comoving wavenumber of a perturbation of ordinary space (in Mpc^{-1}; the physical wavenumber is $k/a_{\rm sp}$); the multipole; and the spectra of the temperature, of the polarization and their cross spectrum, $D_\ell = \ell(\ell+1)C_\ell/2\pi$ in μK^2	new
$h_{\rm sy}$, $\eta_{\rm sy}$, $\alpha_{\rm sy}$	—	the two functions that describe a scalar perturbation of the metric of ordinary space in the synchronous gauge (the h and η of Ma and Bertschinger; here with the index sy, because h is the Hubble parameter and η the flat metric), and the combination $\alpha_{\rm sy} = (h'_{\rm sy} + 6\eta'_{\rm sy})/2k^2$	new
$\mathcal{U}$	`potential`	$\eta_{\rm sy} + \alpha'_{\rm sy}$, the sum of the two gravitational potentials of the conformal Newtonian gauge (script letter): the potential that light feels	new
$\mathcal{D}$, $\mathcal{T}$, $\mathcal{P}$, $\mathcal{A}$	`d_sum` `three_dpres`	$8\pi G_N a_{\rm sp}^2$ times the perturbation of the density, of the divergence of the momentum density, of the pressure, and of the anisotropic stress of matter and radiation together (script letters; in the notation of Ma and Bertschinger: $\delta\rho$, $(\rho+P)\theta$, δP, $(\rho+P)\sigma$)	new
$\mathcal{E}$, $\mathcal{E}_P$, $\mathcal{E}_m$, $\mathcal{E}_r$, $\mathcal{B}$	`grho_sum` `half_rho_plus_p` `b_trace` `n_bg`	$8\pi G_N a_{\rm sp}^2$ times the density and the pressure of the unperturbed universe, summed over the species ($\mathcal{E}_m$, $\mathcal{E}_r$: the parts of $\mathcal{E}$ from pressureless matter and from radiation); and $\mathcal{B} = H'_e - H_e^2 + \frac{1}{2}(\mathcal{E} + \mathcal{E}_P)$, which vanishes without deflation (script letters)	new
$p_{\rm agent}$	—	the pressure in ordinary space of the agent, that is, of whatever keeps the three rates of deflation constant, in the unperturbed universe (under assumption A2 the agent has no energy density)	new
$\mathcal{Y}_M$, $\mathcal{Y}_T$, $\mathcal{Y}_S$; $\mathcal{Z}$	`momentum_gap` `trace_gap` `shear_gap` `x_gap` `shear_check` `trace_check`	the gaps: what the momentum constraint, the trace equation and the equation of the shear lack when only matter and radiation stand on their right-hand sides; the agent that holds the extra scale factors supplies them. $\mathcal{Z} = h'_{\rm sy}\,\mathcal{B} - \frac{1}{2}a_{\rm sp}\,$gsum$\,(\mathcal{D} - 3\mathcal{P})$ (script letters)	new
$Y_{\rm He}$	—	the mass fraction of helium after nucleosynthesis (with the index He; `Y_He` in the data files)	new

10 The metric of the deflating spacetimes in the author's sign convention

symbol	names in programs and data files	meaning	origin
$g^{(4+4)}_{\mu\nu}$, $g^{(7+1)}_{\mu\nu}$	`gs`	the metrics of the deflating spacetimes in the author's sign convention of November 13, 2026 (space-like components positive, time-like components negative): $g^{(4+4)} = \mathrm{diag}(+1,+1,+1,+1,-1,-b_5^2,-b_6^2,-b_7^2)$ and $g^{(7+1)} = \mathrm{diag}(+1,+1,+1,+1,-1,+b_5^2,+b_6^2,+b_7^2)$, that is $g^{(4+4)}_{tt} = -\mathrm{const}_t^2\, e^{-2\,\mathrm{rate}_t\, x4}$ and $g^{(7+1)}_{tt} = +\mathrm{const}_t^2\, e^{-2\,\mathrm{rate}_t\, x4}$ for $t = 5, 6, 7$. A component of the metric carries the superscript $(4+4)$ or $(7+1)$ and two coordinate indices; it is not a rate (g55) and not a coupling (g_r, g_{VS}). The first revision on the deflating spacetimes wrote $g_{1644d} = g^{(4+4)}$ and $g_{1671d} = -g^{(7+1)}$ (time-like component positive on 7+1); these two symbols are retired	author
$\eta^{(4+4)}$, $\eta^{(7+1)}$	—	the flat metrics of the author's sign convention, with which frame indices are raised and lowered on the deflating spacetimes: $\eta^{(4+4)} = \eta_{1644} = \mathrm{diag}(+1,+1,+1,+1,-1,-1,-1,-1)$ and $\eta^{(7+1)} = -\eta_{1671} = \mathrm{diag}(+1,+1,+1,+1,-1,+1,+1,+1)$; $g_{\mu\nu}\, e_A{}^{\mu} e_B{}^{\nu} = \eta_{AB}$	author
c_γ	`clifford_sign`	the sign that relates the Dirac matrices to the metric of the author's sign convention, $\gamma^A\gamma^B + \gamma^B\gamma^A = 2\, c_\gamma\, \eta^{AB}$ with $\eta = \eta^{(4+4)}$ or $\eta^{(7+1)}$: $c_\gamma = +1$ on 4+4 and $c_\gamma = -1$ on 7+1. The Dirac matrices themselves are those of the flat-space part of the project, unchanged	new
b_t^{ref}	`s5` `s6` `s7`	the value of the scale factor b_t at the reference time $x4_{\mathrm{ref}}$. A particle that keeps its wavenumber k_t has the proper momentum k_t/b_t, larger than at the reference time by the stretch factor $b_t^{\mathrm{ref}}/b_t = e^{\mathrm{rate}_t (x4 - x4_{\mathrm{ref}})}$ (columns `s5`, `s6`, `s7`)	new
τ_d^0	`tau_ref` `tau_ref_end`	the reading that the deflation clock would have if the rates had kept their initial value, $\tau_d^0 = 3\,\mathrm{g55}\, x4$ in the evolutions with gravity (equal rates), where the clock actually reached is $\tau_d = -3 \ln \mathrm{b}$	new

11 Mass term and self-interaction of the fermion field

symbol	names in programs and data files	meaning	origin
$\mathcal{L}_{\mathrm{self}}$; $\mathcal{L}^{\mathrm{C}}_{\mathrm{self}}$, $\mathcal{L}^{\mathrm{Q}}_{\mathrm{self}}$	—	the self-interaction of the fermion field (the author's definition, document KOHN_SHAM_GRAVITY): $\mathcal{L}_{\mathrm{self}} = -\frac{1}{2}\sum_{r=0}^{8} g_r\, I_r$, a sum of nine quartic invariants. On the 4+4 spacetime the invariants can be formed in two ways that are not equivalent: $\mathcal{L}^{\mathrm{C}}_{\mathrm{self}} = -\frac{1}{2}\sum_r g_r\, I^{\mathrm{C}}_r$ (form C, the definition of the 4+4 model) and $\mathcal{L}^{\mathrm{Q}}_{\mathrm{self}} = -\frac{1}{2}\sum_r g_r\, I^{\mathrm{Q}}_r$ (form Q, the alternative)	K1; the two forms on 4+4 are new
g_r; $g_0, \ldots, g_8$; g_S, g_V, g_A, g_P	`g_0` `g_1` `g_2` `g_3` `g_4` `g_5` `g_6` `g_7` `g_8`	the nine couplings of the self-interaction, one for each rank r (the author's symbols; dimension length^{D-1} in D space dimensions); $g_0 = g_S$ (scalar), $g_1 = g_V$ (vector), $g_7 = -g_A$ (axial vector), $g_8 = -g_P$ (pseudoscalar). Such a coupling carries one digit as subscript; it is not a rate of deflation (g55: two digits, upright, no subscript), not a component of the metric (superscript $(4+4)$ or $(7+1)$, two indices) and not the coupling g_{VS} of the Vorton-Scalar	K1

symbol	names in programs and data files	meaning	origin
K, J (index sets); r, r' (ranks)	`mask` `rank` `J`	K, J: a strictly increasing list $A_1 < \cdots < A_r$ of indices out of $0, \ldots, 7$ (there are 256); $r = \lvert K\rvert$ is its rank, $0 \le r \le 8$; r' is a second rank and l the summation index in the closed form of the Fierz matrix. In the data files an index set is given by its bit mask (`mask`: the sum of 2^A over its indices)	K1
Θ^K, Θ_K; ζ_r; sgn_K	`eps_K` `phase` `eta_K`	the 256 matrices of the self-interaction (the author's symbols Θ^K, Θ_K): $\Theta^K = \zeta_r\, \gamma^{A_1} \cdots \gamma^{A_r}$, with the phase $\zeta_r = 1$ or i chosen so that the bilinear formed with Θ^K is real, and $\Theta_K = (\Theta^K)^{-1} = \mathrm{sgn}_K\, \Theta^K$ with the sign $\mathrm{sgn}_K = \pm 1$, so that $(\Theta^K)^2 = \mathrm{sgn}_K$. On 7+1: $\zeta_r = i$ for $r = 2, 3, 6, 7$ (the author's i^{σ_r} with $\sigma_r = r(r-1)/2 \bmod 2$), and sgn_K is the product of the squares of the Dirac matrices in K (the author's η_K; column `eta_K`). On 4+4 (form C, the definition): $\zeta_r = i$ for $r = 1, 2, 5, 6$, and sgn_K is $(-1)^r$ times that product. (The sign is written with the letters sgn, because η and ε have other meanings; columns `eps_K`, `phase`.)	K1; form C of 4+4 is new
γ^A_{44q}; Θ^K_Q, ζ^Q_r, sgn^Q_K	`gamma44q`	the matrices with which form Q (the quantized-sector form) of the self-interaction on 4+4 is built: $\gamma^A_{44q} = -i\,\gamma^A_{1644}$ for $A = 0, \ldots, 4$ and $\gamma^A_{44q} = \gamma^A_{1644}$ for $A = 5, 6, 7$; they have the squares of the 7+1 matrices, and $\gamma^4_{44q} = \beta_{1644}$. $\Theta^K_Q = \zeta^Q_r\, \gamma^{A_1}_{44q} \cdots \gamma^{A_r}_{44q}$ with $\zeta^Q_r = i$ for $r = 2, 3, 6, 7$ and $\zeta^Q_r = 1$ for the other ranks; $(\Theta^K_Q)^2 = \mathrm{sgn}^Q_K$, the product of the squares of the matrices γ^A_{44q} in K	new
I_r; I^{C}_r, I^{Q}_r	—	the invariant of rank r (the author's symbol): $I_r = \sum_{\lvert K\rvert=r} (\psi^\dagger\, \mathrm{Adj}\, \Theta_K \psi)(\psi^\dagger\, \mathrm{Adj}\, \Theta^K \psi)$ with the adjoint matrix Adj of the model. On 7+1: $I_r = \sum_{\lvert K\rvert=r} \mathrm{sgn}_K (\bar\psi\, \Theta^K \psi)^2$. On 4+4 there are two forms: $I^{\mathrm{C}}_r = \sum_{\lvert K\rvert=r} \mathrm{sgn}_K (\tilde\psi\, \Theta^K \psi)^2$ (form C, the definition) and $I^{\mathrm{Q}}_r = \sum_{\lvert K\rvert=r} \mathrm{sgn}^Q_K (\bar\psi\, \Theta^K_Q \psi)^2$ (form Q, the alternative); exactly, $I^{\mathrm{C}}_r = \sum_{\lvert K\rvert=r} \mathrm{sgn}^Q_K (\psi^\dagger W_{1644}\, \beta\, \Theta^K_Q \psi)^2$. I_r without superscript is the invariant of the model at hand; in a statement that holds in both forms of 4+4, of either form. I_0 is the square of the scalar density, I_1 the square of the current	K1; the two forms on 4+4 are new
F, $\mathsf{F}_{rr'}$	`fierz` `F_r0` `F_r1` `F_r2` `F_r3` `F_r4` `F_r5` `F_r6` `F_r7` `F_r8`	the Fierz matrix (9×9): $\mathsf{F}_{rr'} = \frac{1}{16} \sum_l (-1)^{rr'-l} \binom{r'}{l} \binom{8-r'}{r-l}$, with $\mathsf{F}^2 = 1$. For anticommuting fields the nine invariants obey $(1 + \mathsf{F})\, I = 0$. (Sans-serif type; F_S, F_P, F_0, F_1, F_7, F_8 in italic type are pairing amplitudes. The author's document writes F_{rs}.)	K1
g^{eff}_0, g^{eff}_1, g^{eff}_7, g^{eff}_8	`g0eff` `g1eff` `g7eff` `g8eff` `g_0eff` `g_1eff` `g_7eff` `g_8eff`	the four independent couplings that remain for anticommuting fields, when $I_2, \ldots, I_6$ are expressed by I_0, I_1, I_7, I_8: $g^{\mathrm{eff}}_0 = g_0 - 3g_2 - 4g_3 - 5g_4 - 4g_5 - g_6$, $g^{\mathrm{eff}}_1 = g_1 - g_2 - g_5 + g_6$, $g^{\mathrm{eff}}_7 = g_7 - g_2 - g_3 + g_6$, $g^{\mathrm{eff}}_8 = g_8 - g_2 + 4g_3 - 5g_4 + 4g_5 - 3g_6$; then $\mathcal{L}_{\mathrm{self}} = -\frac{1}{2}(g^{\mathrm{eff}}_0 I_0 + g^{\mathrm{eff}}_1 I_1 + g^{\mathrm{eff}}_7 I_7 + g^{\mathrm{eff}}_8 I_8)$	new
n	—	the one-body density matrix of matter (16×16), $\mathsf{n}_{ab} = \langle \psi^\dagger_b \psi_a \rangle$, with $\mathrm{Tr}\, \mathsf{n} = n$ and $\mathrm{Tr}(\beta\, \mathsf{n}) = n_S$; for uniform matter at rest $\mathsf{n} = (n + n_S\, \beta)/16$. (The author's document writes R for the matrix $\mathsf{n}\, \beta$.)	new

symbol	names in programs and data files	meaning	origin
G_S^{HF}, G_V^{HF}; G_S^{c}, G_V^{c}	`G_0` `G_1` `G_S0` `G_V0` `hf_rho` `hf_rhoS` `hf_rhoW` `hf_rhoSW`	the four combinations of the couplings that the Hartree-Fock approximation gives for uniform matter: $G_S^{\mathrm{HF}} = (15g_0^{\mathrm{eff}} - 8g_1^{\mathrm{eff}} - 8g_7^{\mathrm{eff}} - g_8^{\mathrm{eff}})/16$, $G_V^{\mathrm{HF}} = (-g_0^{\mathrm{eff}} + 22g_1^{\mathrm{eff}} - 6g_7^{\mathrm{eff}} + g_8^{\mathrm{eff}})/16$, $G_S^{\mathrm{c}} = (-g_0^{\mathrm{eff}} - 2g_1^{\mathrm{eff}} + 2g_7^{\mathrm{eff}} + g_8^{\mathrm{eff}})/16$ and $G_V^{\mathrm{c}} = -(g_0^{\mathrm{eff}} + g_8^{\mathrm{eff}})/16$. On 7+1 the interaction energy density is $\frac{1}{2}(G_V^{\mathrm{HF}} n^2 + G_S^{\mathrm{HF}} n_S^2)$: there G_S^{HF}, G_V^{HF} are the mean-field couplings G_S, G_V. On 4+4 it is $\frac{1}{2}(G_V^{\mathrm{tot}} n^2 + G_S^{\mathrm{tot}} n_S^2 + G_V^{\mathrm{dif}} n_W^2 + G_S^{\mathrm{dif}} n_{SW}^2)$ with the same four combinations: $G^{\mathrm{tot}} = G^{\mathrm{c}}$, $G^{\mathrm{dif}} = G^{\mathrm{HF}}$ in form C and $G^{\mathrm{tot}} = G^{\mathrm{HF}}$, $G^{\mathrm{dif}} = G^{\mathrm{c}}$ in form Q. (The superscript c is upright; G_c with the subscript c is the critical coupling of the vacuum. The author's document and the data files write G_0, G_1 for G_S^{HF}, G_V^{HF}; the data files write `G_S0`, `G_V0` for G_S^{c}, G_V^{c}.)	K1; the combinations of 4+4 are new
G_{pair}	`G_pair`	the contact coupling of a pair of two particles at rest: the interaction energy of the pair times the volume; attractive if negative. For the seven vector pairs on 7+1 and in form Q of 4+4: $G_{\mathrm{pair}} = g_0^{\mathrm{eff}} + g_1^{\mathrm{eff}} + 5\, g_7^{\mathrm{eff}}$. In form C of 4+4: $-(g_0^{\mathrm{eff}} + g_1^{\mathrm{eff}} + 5\, g_7^{\mathrm{eff}})$ for the four vector pairs along $x0, x1, x2, x3$ (one particle of each half of the field) and $+(g_0^{\mathrm{eff}} + g_1^{\mathrm{eff}} + 5\, g_7^{\mathrm{eff}})$ for the three vector pairs along $x5, x6, x7$ (two particles of the same half)	new
n_+, n_-, n_{S+}, n_{S-}; n_W, n_{SW}; pol_W	`rho_W` `rho_SW` `pol_W` `pol`	number densities $n_\pm = \langle \psi_\pm^\dagger \psi_\pm \rangle$ and scalar densities $n_{S\pm} = \langle \psi_\pm^\dagger \beta\, \psi_\pm \rangle$ of the particles in the two halves $W_{1644} = \pm 1$ of the 4+4 field; $n = n_+ + n_-$, $n_S = n_{S+} + n_{S-}$, $n_W = n_+ - n_-$, $n_{SW} = n_{S+} - n_{S-}$; the polarization $\mathrm{pol}_W = (n_+ - n_-)/n$ (columns `pol_W`, `pol`)	new
Adj	—	the adjoint matrix of a form of the self-interaction: its bilinears are $\psi^\dagger \mathrm{Adj}\, \Theta^K \psi$. $\mathrm{Adj} = \beta$ on 7+1 and in form Q of 4+4, where $\psi^\dagger \mathrm{Adj} = \bar{\psi}$; $\mathrm{Adj} = C^-_{1644}$ in form C of 4+4, where $\psi^\dagger \mathrm{Adj} = \tilde{\psi}$. (Upright letters; used where a statement holds for every form.)	new
ψ_+, ψ_-	—	the two halves of the 4+4 field: $\psi = \psi_+ + \psi_-$ with $W_{1644}\, \psi_\pm = \pm \psi_\pm$, eight components each. The Hamiltonian of the quantized sector acts inside each half, and a particle has four internal states in each	new
$\mathcal{L}_{\mathrm{cov}}$	—	the covariant Lagrangian of the free field on 4+4, $\mathcal{L}_{\mathrm{cov}} = \tilde{\psi}\, (\gamma^A \partial_A - m)\, \psi$. For fields of the quantized sector $\mathcal{L}_{\mathrm{cov}} = -\psi^\dagger W_{1644}\, (i\partial_4 - \mathsf{h})\, \psi$: it is $\mathcal{L}_{\mathrm{kin}} + \mathcal{L}_{\mathrm{mass}}$ on the half $W_{1644} = -1$ and minus that on the half $W_{1644} = +1$	new
z_K	—	the phase in the exact relation between the matrices of the two forms on 4+4, $C^-_{1644}\, \Theta^K = z_K\, W_{1644}\, \beta\, \Theta^K_Q$: $z_K = \pm 1$ if the index set K contains an even number of the indices 5, 6, 7, and $z_K = \pm i$ if it contains an odd number	new
$\mathcal{J}_K^{++}$, $\mathcal{J}_K^{--}$, $\mathcal{J}_K^{+-}$, $\mathcal{J}_K^{-+}$	—	the bilinears of the two halves of the 4+4 field for the index set K: $\mathcal{J}_K^{++} = \psi_+^\dagger \beta\, \Theta^K_Q \psi_+$, $\mathcal{J}_K^{--} = \psi_-^\dagger \beta\, \Theta^K_Q \psi_-$, $\mathcal{J}_K^{+-} = \psi_+^\dagger \beta\, \Theta^K_Q \psi_-$ and $\mathcal{J}_K^{-+} = \psi_-^\dagger \beta\, \Theta^K_Q \psi_+ = (\mathcal{J}_K^{+-})^\dagger$. If K contains an even number of the indices 5, 6, 7 only the first two differ from zero, otherwise only the last two. (Script letter with an index set and two signs; J^A_{VS}, J^A_5 are currents and J is an index set.)	new

symbol	names in programs and data files	meaning	origin
$G_S^{\rm tot}$, $G_V^{\rm tot}$; $G_S^{\rm dif}$, $G_V^{\rm dif}$	`G_Stot` `G_Vtot` `G_Sdif` `G_Vdif`	the mean-field couplings of the two halves of the 4+4 field: the interaction energy density of uniform matter is $\frac{1}{2}(G_V^{\rm tot}n^2 + G_S^{\rm tot}n_S^2 + G_V^{\rm dif}n_W^2 + G_S^{\rm dif}n_{SW}^2)$. $G^{\rm tot}$ multiply the squares of the total densities, $G^{\rm dif}$ the squares of the differences between the densities of the two halves. Form C: $G_S^{\rm tot} = G_S^{\rm c}$, $G_V^{\rm tot} = G_V^{\rm c}$, $G_S^{\rm dif} = G_S^{\rm HF}$, $G_V^{\rm dif} = G_V^{\rm HF}$. Form Q: $G_S^{\rm tot} = G_S^{\rm HF}$, $G_V^{\rm tot} = G_V^{\rm HF}$, $G_S^{\rm dif} = G_S^{\rm c}$, $G_V^{\rm dif} = G_V^{\rm c}$. (In the data files and in the statements of the programs: `G_Stot`, `G_Vtot`, `G_Sdif`, `G_Vdif`.)	new
$G_S^{\rm same}$, $G_V^{\rm same}$; $G_S^{\rm opp}$, $G_V^{\rm opp}$	`G_Ssame` `G_Vsame` `G_Sopp` `G_Vopp`	the mean-field couplings between two particles of the same half and between particles of opposite halves of the 4+4 field: $G^{\rm same} = G^{\rm tot} + G^{\rm dif}$ and $G^{\rm opp} = G^{\rm tot} - G^{\rm dif}$, so that the interaction energy density is $\frac{1}{2}G_V^{\rm same}(n_+^2 + n_-^2) + G_V^{\rm opp}\, n_+ n_- + \frac{1}{2}G_S^{\rm same}(n_{S+}^2 + n_{S-}^2) + G_S^{\rm opp}\, n_{S+}n_{S-}$. $G^{\rm same} = G^{\rm HF} + G^{\rm c}$ is the same in both forms: $G_S^{\rm same} = (7g_0^{\rm eff} - 5g_1^{\rm eff} - 3g_7^{\rm eff})/8$ and $G_V^{\rm same} = (-g_0^{\rm eff} + 11g_1^{\rm eff} - 3g_7^{\rm eff})/8$. $G^{\rm opp}$ has opposite signs in the two forms: $G^{\rm opp} = G^{\rm HF} - G^{\rm c}$ in form Q and $G^{\rm opp} = -(G^{\rm HF} - G^{\rm c})$ in form C, with $G_S^{\rm HF} - G_S^{\rm c} = (8g_0^{\rm eff} - 3g_1^{\rm eff} - 5g_7^{\rm eff} - g_8^{\rm eff})/8$ and $G_V^{\rm HF} - G_V^{\rm c} = (11g_1^{\rm eff} - 3g_7^{\rm eff} + g_8^{\rm eff})/8$. (In the data files and in the statements of the programs: `G_Ssame`, `G_Vsame`, `G_Sopp`, `G_Vopp`.)	new
g (strength)	—	the common strength of the couplings of the Lagrangian in the sets A and B of the numerical examples on 4+4: $g = \|g_0\| > 0$, of dimension length^{D-1} (column `g` of the files `forms_..`, in units $m = 1$; in the files of the gaps the column `g` is the coupling g_{VS}). On the deflating spacetime g grows as e^{τ_d}	new
$\mathcal{L}_{\rm kin}$, $\mathcal{L}_{\rm mass}$	—	the kinetic term and the mass term of the Lagrangian of the fermion field, $\mathcal{L} = V_d\,[\mathcal{L}_{\rm kin} + \mathcal{L}_{\rm mass} + \mathcal{L}_{\rm self}]$. On 4+4, for fields of the quantized sector, $\mathcal{L}_{\rm kin} + \mathcal{L}_{\rm mass} = \psi^\dagger(i\partial_4 - \mathsf{h})\psi$ (the positive Lagrangian, with the mass term $-m\,\bar\psi\psi$)	new

12 Kohn-Sham and CJT calculations with contact interaction

symbol	names in programs and data files	meaning	origin
G_S, G_V	`G_S` `G_V` `GS_abs`	the scalar and the vector coupling of the Kohn-Sham (mean-field) description of uniform matter (the author's symbols): interaction energy density $\frac{1}{2}G_V n^2 + \frac{1}{2}G_S n_S^2$; attraction for $G_S < 0$, repulsion for $G_V > 0$; dimension length^{D-1}. The author's example: $G_S = -10^5$, $G_V = +10^4$ in units $m = 1$, $D = 7$	K1
n, n_S	`rho` `rho_S` `rho_ks` `rho_cjt` `n_ks` `n_cjt`	number density $\langle\psi^\dagger\psi\rangle$ and scalar density $\langle\bar\psi\psi\rangle$ of the fermions of matter (the filled states of negative energy not counted). The data files and the author's document write ρ (`rho`) and ρ_S (`rho_S`); the files of the two halves of the 4+4 field write `n`	K1 (there ρ, ρ_S)
D	`D`	the number of space dimensions in which the particles of matter have momentum: 7 (all space coordinates of 7+1), 4 (fields that do not depend on $x5, x6, x7$) or 3 (fields that depend on ordinary space only). $V_D = \pi^{D/2}/\Gamma_{\rm E}(D/2+1)$ is the volume and $A_D = D\,V_D$ the area of the unit ball in D dimensions ($\Gamma_{\rm E}$: Euler's gamma function)	new

symbol	names in programs and data files	meaning	origin
$E^*(k)$; $\langle f\rangle$	—	$E^*(k)=\sqrt{k^2+m^{*2}}$, the energy of a particle of matter without the vector term; $\langle f\rangle=\nu\int d^Dk\, f/(2\pi)^D$ over the occupied states: $n=\langle 1\rangle$, $n_S=\langle m^*/E^*\rangle$. For the two halves of the 4+4 field: $E^*_\pm(k)=\sqrt{k^2+m^{*2}_\pm}$	K1 (there ω for E^*)
$\varepsilon_{\rm kin}$; p_j; $p_{\rm int}$	`e_kin` `eps` `eps_ks` `eps_cjt` `eps_rel` `p` `p_ks` `p_flat` `p5` `p6` `p7` `p_a` `p_b` `pa_rel` `pb_rel` `Pi`	$\varepsilon_{\rm kin}=\langle E^*\rangle$: the kinetic part of the energy density $\varepsilon=\varepsilon_{\rm kin}+\frac12 G_V n^2-(m^*-m)^2/(2G_S)$; p_j: the pressure of matter along the coordinate xj, $\langle k_j^2/E^*\rangle+p_{\rm int}$; $p_{\rm int}=\frac12 G_V n^2+(m^*-m)^2/(2G_S)$: the interaction part of the pressure, which is the whole pressure along a coordinate on which the fields do not depend (`Pi` in the data files, Π in the author's document)	K1
V_0	`V_0`	the vector mean field $V_0=G_V\,n$ (the author's symbol); it shifts the chemical potential, $\mu\to\mu-V_0$. For the two halves of the 4+4 field: $V_{0\pm}=G_V^{\rm tot}\,n\pm G_V^{\rm dif}\,n_W$	K1
$\Omega_{\rm CJT}$; $\Omega_{\rm half}$	`minus_Omega`	the CJT effective potential of uniform matter with contact interaction per unit volume, a function of the mass m^* and of the vector field V_0 at the chemical potential μ: $\Omega_{\rm CJT}=-\nu\int\frac{d^Dk}{(2\pi)^D}\,(\mu-V_0-E^*)\,\theta(\mu-V_0-E^*)-\frac{(m^*-m)^2}{2G_S}-\frac{V_0^2}{2G_V}$; at its stationary point $-\Omega_{\rm CJT}$ is the pressure. For the two halves of the 4+4 field it is a function of two masses and two vector fields: $\Omega_{\rm CJT}=\Omega_{\rm half}(m^*_+,\mu-V_{0+})+\Omega_{\rm half}(m^*_-,\mu-V_{0-})$ $-[\frac12(m^*_++m^*_-)-m]^2/(2G_S^{\rm tot})-[\frac12(m^*_+-m^*_-)]^2/(2G_S^{\rm dif})$ $-[\frac12(V_{0+}+V_{0-})]^2/(2G_V^{\rm tot})-[\frac12(V_{0+}-V_{0-})]^2/(2G_V^{\rm dif})$, with the grand potential of free particles of one half (four internal states), $\Omega_{\rm half}(m^*,\mu)=-4\int\frac{d^Dk}{(2\pi)^D}\,(\mu-E^*)\,\theta(\mu-E^*)$	new
G_c	`Gc`	the critical coupling of the vacuum: with the filled states of negative energy below the cutoff Λ a massless field acquires a mass when $G_S<G_c=-(2\pi)^D(D-1)/(\nu A_D\Lambda^{D-1})$; $G_c\Lambda^6=-90\pi^4$ for $D=7$	K2 ($D=7$)
$m_{\rm vac}$, $m_{\rm bare}$; $\langle\bar\psi\psi\rangle_{\rm vac}$	`M_vac` `m_bare` `condensate`	with the filled states of negative energy: the mass of the fermions in vacuum, the mass parameter of the Lagrangian (bare mass) that produces it, and the scalar density of the filled states, $\langle\bar\psi\psi\rangle_{\rm vac}=-\nu\int^\Lambda\frac{d^Dk}{(2\pi)^D}\frac{m_{\rm vac}}{\sqrt{k^2+m_{\rm vac}^2}}$ (negative); $m_{\rm vac}=m_{\rm bare}+G_S\langle\bar\psi\psi\rangle_{\rm vac}$	new
$n^{\rm vac}_{S+}$, $n^{\rm vac}_{S-}$; $\Omega_{\rm vac}$	`Omega`	for the two halves of the 4+4 field with the filled states of negative energy below the cutoff Λ: the scalar densities of the filled states, $n^{\rm vac}_{S\pm}=-4\int_{\vec k<\Lambda}\frac{d^Dk}{(2\pi)^D}\,\frac{m^*_\pm}{E^*_\pm}$, which enter the gap equations of the vacuum $\frac12(m^*_++m^*_-)-m=G_S^{\rm tot}(n^{\rm vac}_{S+}+n^{\rm vac}_{S-})$, $\frac12(m^*_+-m^*_-)=G_S^{\rm dif}(n^{\rm vac}_{S+}-n^{\rm vac}_{S-})$; and the energy density of the vacuum relative to the state in which both halves have the mass m of the Lagrangian, $\Omega_{\rm vac}=-4\sum_\pm\int_{\vec k<\Lambda}\frac{d^Dk}{(2\pi)^D}\,[E^*_\pm-\sqrt{k^2+m^2}]$ $-[\frac12(m^*_++m^*_-)-m]^2/(2G_S^{\rm tot})-[\frac12(m^*_+-m^*_-)]^2/(2G_S^{\rm dif})$; among several solutions of the gap equations the one of lowest $\Omega_{\rm vac}$ is the vacuum (column `Omega`)	new
$g_{\rm crit}$; $\Lambda_{\rm max}$	`g_crit` `cutoff_max`	$g_{\rm crit}$: the critical strength of a set of couplings, the strength g at which the more attractive of the scalar couplings $G_S^{\rm tot}$, $G_S^{\rm dif}$ reaches the critical coupling G_c of the vacuum of a massless field. $\Lambda_{\rm max}$: at a given strength, the largest cutoff Λ for which that coupling is still weaker than G_c, so that the vacuum of the massless field is stable (column `cutoff_max`, in units of m)	new

symbol	names in programs and data files	meaning	origin
k_{F+}, k_{F-}, $k_{F,\mathrm{tot}}$; μ_+, μ_-	`kF_total` `mu_plus` `mu_minus`	for the two halves of the 4+4 field: the Fermi momenta of the two halves, $n_\pm = 4V_D k_{F\pm}^D/(2\pi)^D$ (four internal states each); the Fermi momentum $k_{F,\mathrm{tot}}$ that the total density would have with equal halves, $n = 8V_D k_{F,\mathrm{tot}}^D/(2\pi)^D$; and the chemical potentials $\mu_\pm = \sqrt{k_{F\pm}^2 + m_\pm^{*2}} + V_{0\pm}$ of the two halves (for an empty half: the energy of its lowest state)	new
$\mathsf{R}_{\mathrm{gap}}$	`response`	the response matrix of the two gap equations $m_\pm^* = m + G_S^{\mathrm{same}} n_{S\pm} + G_S^{\mathrm{opp}} n_{S\mp}$ of the halves of the 4+4 field: the 2×2 matrix with the elements $(\mathsf{R}_{\mathrm{gap}})_{++} = G_S^{\mathrm{same}}\, \partial n_{S+}/\partial m_+^*$, $(\mathsf{R}_{\mathrm{gap}})_{+-} = G_S^{\mathrm{opp}}\, \partial n_{S-}/\partial m_-^*$, $(\mathsf{R}_{\mathrm{gap}})_{-+} = G_S^{\mathrm{opp}}\, \partial n_{S+}/\partial m_+^*$, $(\mathsf{R}_{\mathrm{gap}})_{--} = G_S^{\mathrm{same}}\, \partial n_{S-}/\partial m_-^*$ (derivatives at fixed Fermi momenta). While its eigenvalues are below 1 the solution is the regular continuation of the dilute gas; at the eigenvalue 1 solutions with other masses branch off. Column `response`: the largest eigenvalue. (Sans-serif with the suffix gap; R in italic type is a curvature.)	new
$-g_0/C$, g_1/C	`kappa_0` `kappa_1`	the couplings g_0 and g_1 of the Lagrangian in units of the coupling C of the Vorton-Scalar (units $m = 1$), in the calculations with bare mass, self-interaction and Vorton-Scalar together on 4+4; they are written as these ratios, without a letter of their own (columns `kappa_0`, `kappa_1`)	new
m_D, m_Y, f_D	`f_D`	when a bare Dirac mass and the Vorton-Scalar are both present: the bare Dirac mass m_D, the part $m_Y = g_{VS} v_{VS}$ of the mass that the Vorton-Scalar gives, and the fraction $f_D = m_D/m$ of the vacuum mass $m = m_D + m_Y$	new
m_ϑ	`mph2_over_C_ms2`	the mass of the phase mode ϑ_{VS} of the Vorton-Scalar: zero while the chiral symmetry $U(1)_{VS}$ is exact; a bare Dirac mass breaks the symmetry and gives $m_\vartheta^2 v_{VS}^2 = (\lvert\langle\bar\psi\psi\rangle_{\mathrm{vac}}\rvert - n_S) m_D m_Y/m$	new
$-G_S/C$, G_V/C	`kappa_S` `kappa_V`	the contact couplings in units of the coupling $C = g_{VS}^2 m^2/m_s^2$ of the Vorton-Scalar (units $m = 1$); they are written as these ratios, without a letter of their own (columns `kappa_S`, `kappa_V`)	new
h_{KS}	—	the Hamiltonian matrix of a particle of matter in the Kohn-Sham description, $\mathsf{h}_{\mathrm{KS}} = \sum_j \alpha^j k_j + \beta\, m^* + V_0$	K1
$\Gamma[S]$; S, S_0	—	the CJT effective action as a functional of the propagator of the fermions, $\Gamma[S] = -i\,\mathrm{Tr}\ln S^{-1} - i\,\mathrm{Tr}(S_0^{-1}S - 1) + \Gamma_2[S]$, with the full propagator S and the propagator S_0 of free particles of the mass m (the argument in brackets tells it from the action $\Gamma[\phi, G]$ of the model with the Vorton-Scalar)	new
n_{ref}	—	the number density of matter at the reference time $x4_{\mathrm{ref}}$; matter that keeps its particles has $n = n_{\mathrm{ref}}\, e^{\tau_d}$	new

13 Pairing with contact couplings

symbol	names in programs and data files	meaning	origin
F_0, F_1, F_7, F_8	`F0` `F1` `F7` `F8` `F0c` `F1c` `F7c` `F8c`	the pairing amplitudes of the four independent contact channels in the pair channel along the hidden axis, for particles of mass m^* that scatter from the momenta $\vec{p}, -\vec{p}$ to $\vec{p}\,', -\vec{p}\,'$ in ordinary space ($E = \sqrt{m^{*2} + p^2}$, E' likewise): $F_0 = (EE' + m^{*2} - \vec{p}\cdot\vec{p}\,')/(2EE')$, $F_1 = (3EE' - 2m^{*2} - \vec{p}\cdot\vec{p}\,')/(EE')$, $F_7 = (3EE' + 2m^{*2} - \vec{p}\cdot\vec{p}\,')/(EE')$, $F_8 = -(EE' - m^{*2} - \vec{p}\cdot\vec{p}\,')/(2EE')$. $F_0 = F_S$ and $F_8 = -F_P$. The same closed forms hold in the pair channels along $x5, x6, x7$; the sign with which they enter the pairing potential is $s_{\rm ct}$	new
$m_{\rm ct}$	`m_ct`	the range parameter of the contact couplings in the gap equation: each effective coupling $g_r^{\rm eff}$ is replaced by $g_r^{\rm eff}\, m_{\rm ct}^2/(q^2 + m_{\rm ct}^2)$, the form that the exchange of a boson of mass $m_{\rm ct}$ gives (q: momentum transfer)	new
$V_{\rm ct}$, V_{VS}	—	the pairing potentials in the pair channel along the hidden axis and in those along $x5, x6, x7$: of the contact interaction, $V_{\rm ct} = s_{\rm ct}\sum_{r=0,1,7,8} g_r^{\rm eff} F_r\, m_{\rm ct}^2/(q^2 + m_{\rm ct}^2)$, and of the exchange of the two modes of the Vorton-Scalar, $V_{VS} = -g_{VS}^2[F_S/(q^2 + m_s^2) + F_P/(q^2 + m_\vartheta^2)]$; attractive where negative	new
$s_{\rm ct}$	`ct_sign`	the sign with which the contact couplings enter the pairing potential $V_{\rm ct}$ of a vector pair whose axis is perpendicular to ordinary space: $s_{\rm ct} = +1$ on 7+1, in form Q of 4+4 (the pair along the hidden axis $x0$ and the pairs along $x5, x6, x7$) and in form C for the pairs along $x5, x6, x7$; $s_{\rm ct} = -1$ in form C for the pair along the hidden axis (column `ct_sign`)	new
$k_{\rm max}$	`K_MAX` `K_MAX_EARLIER`	the upper limit of the momentum integral of the gap equation. In the calculations of this folder $k_{\rm max} = 2\times 10^4 \max(m, k_F)$ (constant `K_MAX` of the program); the earlier calculations of the project used $60\max(m, k_F)$ (`K_MAX_EARLIER`)	new
$w_\infty(t)$; s_Δ; $U_{\rm uv}$; u_S, u_V	`uv` `UV_I_S` `UV_I_V` `UV_CRITICAL`	the quantities of the ultraviolet criterion of the gap equation. $w_\infty(t)$: the kernel at momenta far above all masses, where $W(k,k') \to w_\infty(k'/k)/k^2$. s_Δ: the exponent of a power-law solution $\Delta(k) \propto k^{-s_\Delta}$ at large momenta, a root of $\int_0^\infty dt\, t^{1-s_\Delta}\, w_\infty(t) = 4\pi^2$. $U_{\rm uv} = \int_0^\infty dt\, w_\infty(t)$: the ultraviolet strength of the kernel; the gap equation has a solution that does not depend on $k_{\rm max}$ if and only if $U_{\rm uv} < 4\pi^2$. With both modes of the Vorton-Scalar, $U_{\rm uv} = 2u_S\, g_{VS}^2 - s_{\rm ct}\, m_{\rm ct}^2[(g_0^{\rm eff} - g_8^{\rm eff})\, u_S + (g_1^{\rm eff} + g_7^{\rm eff})\, u_V]$, with the two integrals $u_S = \pi^2/8 - \frac{1}{2}$ and $u_V = 3\pi^2/4 - 1$ (constants `UV_I_S`, `UV_I_V`; `UV_CRITICAL` is $4\pi^2$). Column `uv`: $U_{\rm uv}/4\pi^2$	new
Φ, $\Phi(\vec{p})$	—	a state of a pair of two particles, an antisymmetric 16×16 matrix; $\Phi(\vec{p})$: the state of the pair along the hidden axis with the momenta $\vec{p}, -\vec{p}$, the matrix $(C^-\gamma^0)^*$ between the projectors on the states of positive energy. On 4+4: $W_{1644}\,\Phi\, W_{1644}^T = -\Phi$ for a pair of one particle of each half of the field and $+\Phi$ for a pair of two particles of the same half. (Without suffix; the Vorton-Scalar always carries the suffix VS.)	new

14 Einstein-Lovelock gravity with Kohn-Sham matter

symbol	names in programs and data files	meaning	origin
$E^{(k)\mu}{}_{\nu}$	—	the Lovelock tensor of order $k = 1, 2, 3$ in the author's notation and sign: $E^{(1)} = -\mathcal{G}$ (minus the Einstein tensor), $E^{(2)} = \mathcal{H}$ (the Lanczos tensor). For a metric whose scale factors depend only on $x4$: $E^{(k)m}{}_{m} = (-1)^{k+1} k!\, 2^{k-1}$ times the sum, over the sets of $2k$ frame directions other than m and over their divisions into pairs, of the products of the sectional curvatures K_{AB} of the pairs. (With the order in parentheses as a superscript; E_F, E_k, $E_\perp$, E^* are energies.)	K1
α_1, α_2, α_3; $\bar{\Lambda}$; κ_8	`LambdaBar` `kappa` `kappa_eps`	the couplings of the Lovelock terms ($\alpha_1 = 1$), the cosmological term and the gravitational coupling of the author's Einstein-Lovelock equations: $\sum_k \alpha_k E^{(k)4}{}_{4} - \bar{\Lambda} = \kappa_8\, \varepsilon$ for the time component and $\sum_k \alpha_k E^{(k)m}{}_{m} - \bar{\Lambda} = -\kappa_8\, p_m$ for the space direction m. $\alpha_2 = -\alpha_{GB}$ and $\bar{\Lambda} = \Lambda_8$. The author's document writes κ for $\kappa_8 = 8\pi G_8$ (the letter κ with an argument, $\kappa(k)$, is a pair correlation). A Lovelock coupling carries a digit as subscript; α^j with a superscript are Dirac matrices	K1
a, b; H_{a}, H_{b}; n_{a}, n_{b}	`a` `b` `H_a` `H_b` `Ha_over_g` `Hb_over_g` `Hb0_over_g` `a_end` `b_end` `Ha_end_over_g` `Hb_end_over_g`	the two scale factors of a spacetime whose space coordinates form two groups of n_{a} and n_{b} coordinates, and their rates $H_{\mathsf{a}} = \mathsf{a}'/\mathsf{a}$, $H_{\mathsf{b}} = \mathsf{b}'/\mathsf{b}$ (a prime: derivative with respect to the time $x4$). In the author's document: a for ordinary space ($n_{\mathsf{a}} = 3$) and b for the four auxiliary coordinates. On the deflating spacetimes with equal rates: a for $x0, \ldots, x3$ ($n_{\mathsf{a}} = 4$) and b $= b_5 = b_6 = b_7$ ($n_{\mathsf{b}} = 3$); the undisturbed deflation is $H_{\mathsf{a}} = 0$, $H_{\mathsf{b}} = -\mathsf{g55}$	K1 (there a, b, H_a, H_b)
loading	`loading`	$\kappa_8\, \varepsilon(0)/(3\, \mathsf{g55}^2)$: the energy density of the matter at the start of an evolution, in units of the energy density that would make three coordinates deflate at the rate g55 in Einstein gravity	new
$T^{\mu}{}_{\nu}$; p_{a}, p_{b}; δH_{b}	—	the energy-momentum tensor of uniform matter in mixed components, $T^4{}_4 = \varepsilon$, $T^m{}_m = -p_m$; its pressures along the coordinates that carry the scale factors a and b; the deviation $H_{\mathsf{b}} + \mathsf{g55}$ of the rate of the deflating coordinates from the undisturbed deflation	new

15 Tags and identifiers of the calculations with mass term and self-interaction

symbol	names in programs and data files	meaning	origin
71s, 44q, 44c	`71s` `44q` `44c`	tags of the three models of the self-interaction in the programs and the data files: 71s on 7+1; 44c form C on 4+4, the definition (bilinears with $\tilde{\psi} = \psi^\dagger C^-_{1644}$); 44q form Q on 4+4, the alternative (bilinears with $\bar{\psi} = \psi^\dagger \beta_{1644}$ and the matrices γ^A_{44q})	new

symbol	names in programs and data files	meaning	origin
form C, form Q	—	the two forms of the self-interaction on the 4+4 spacetime. Form C, the covariant form, is the definition of the 4+4 model (instruction of the author of November 13, 2026): the invariants I_r^{C}, built with the adjoint $\tilde{\psi}$, which are invariant under all rotations and boosts of the 4+4 frame, together with the positive kinetic and mass terms. Form Q, the quantized-sector form, is the alternative: the invariants I_r^{Q}, built with the adjoint $\bar{\psi}$ of the positive kinetic term. In the data files the column `form` holds C or Q, or 0 for C and 1 for Q. (Upright capitals; C in italic type is the coupling of the Vorton-Scalar, and $C^{\pm}$ with a sign are charge-conjugation matrices.)	new
set A, set B	—	the two sets of couplings of the Lagrangian for which the numbers of the two forms on 4+4 are given, each with one strength g: set A, a scalar attraction alone, $g_0 = -g$; set B, $g_0 = -g$ and $g_1 = +0.1\,g$ (the ratio of the author's example); all other couplings zero. In the data files of the two forms the column `set` holds A or B, or 0 for A and 1 for B (in the files of the gaps the column `set` numbers the pairs of values of m_s and k_F)	new
identifiers of the machine checks	—	`44d-..`, `71d-..`, `def-..`: checks on the deflating 4+4 spacetime, on the deflating 7+1 spacetime, and of results that hold on both; `44-..`, `71-..`: checks of the flat spacetimes, executed again; `71s-S..`, `44q-S..`, `44c-S..`, `self-X..`: the algebra of the self-interaction; `ks-K..`: Kohn-Sham calculations; `cjt-J..`: CJT effective potential with contact interaction; `uni-U..`: contact interaction, bare mass and Vorton-Scalar together; `pc-P..`: pairing with contact couplings; `gr-G..`: gravity with Kohn-Sham matter; `fm-F..`: the two forms of the self-interaction on 4+4 (matter of two halves, vacuum, thresholds). A leading capital I (`Icks-..`, `Idef-..`, `I44d-..`, `I71d-..`) marks a check of the independent verification in Python	new

Key to the column "origin":

code	source
N01	the author's notebook `4+4.nb` (Dirac matrices of 4+4; split octonions)
N02	the author's notebook `7+1.nb` (octonions; Dirac matrices and Dirac equation of 7+1)
N04	the author's notebook `Dirac_Einstein-Lovelock_4+4_Nash.nb` (meaning of the coordinates)
N05	the author's notebook `Dirac_Nash_Identity.nb`
N10	the author's notebook `Pair_Creation_of_Universes_WaveFunctionOfUniverse-4+4-Einstein-Lovelock-Nash.nb`
N12	the author's notebook `WaveFunctionUniverse-4+4-Einstein-Lovelock.nb`
D06	P. L. Nash, "Identities satisfied by the generators of the Dirac algebra", J. Math. Phys. 25, 204 (1984)
D07	P. L. Nash, "On the structure of the split octonion algebra", Nuovo Cimento B 105, 31 (1990)
D08	P. L. Nash, "Second gravity", J. Math. Phys. 51, 042501 (2010)
K1	the author's document `KOHN_SHAM_GRAVITY` (folder 7+1; sections 15 and 16 of the notebook `7+1.nb`): nine-channel self-coupling, Fierz matrix, Kohn-Sham theory, Einstein-Lovelock evolution

code	source
K2	the author's document `DIRAC16COMPLEX` (folder 7+1): the critical coupling of the vacuum in seven space dimensions
author	a name that an instruction of the author fixes (the rates g55, g66, g77; the constants; the signs of the metric)
new	introduced by this project

Appendix C. Files

file	content
`code/deflation-cjt-ks-c-rs/src/metric.rs`, `geometry.rs`, `curved.rs`	metric, Christoffel symbols, curvature, Killing vectors, coordinate matrices, spin connection, Dirac operators (reports A.1, A.2)
`code/deflation-cjt-ks-c-rs/src/selfint.rs`, `cmd_selfcoupling.rs`	the 256 matrices Θ^K, the Fierz matrix, the theorem on the four invariants, Hartree–Fock and pair couplings, symmetries, the forms of 4+4 (report A.3)
`code/deflation-cjt-ks-c-rs/src/halves.rs`, `cmd_forms.rs`, `cmd_forms_b.rs`	the two forms on 4+4: mean-field couplings and matter of two halves, polarization, binding thresholds, the vacuum (report A.4)
`code/deflation-cjt-ks-c-rs/src/modes.rs`	integration of the modes; creation and extinction (report A.5)
`code/deflation-cjt-ks-c-rs/src/symmetry_defl.rs`, `cmd_symmetry.rs`	isometries, internal symmetry, the weighted Lagrangian (report A.6)
`code/deflation-cjt-ks-c-rs/src/gravity.rs`, `cmd_gravity.rs`	Einstein–Lovelock equations with two scale factors (report A.7)
`code/scripts/verify_deflating_independent.py`, `verify_cjt_ks_independent.py`	the independent verifications (report A.8)
`data/forms_couplings.csv`, `data/forms_thresholds.csv`, `data/forms_energy_vs_polarization.csv`, `data/forms_vacuum_critical.csv`, `data/selfcoupling_internal_symmetry.csv`	the mean-field couplings of the sets A and B, the binding thresholds, the energy against the polarization, the vacuum at the critical strength, and the conditions of invariance under the internal symmetry, in both forms (Sections 6.5, 6.6)
`data/selfcoupling_fierz.csv`, `data/selfcoupling_reduction.csv`, `data/selfcoupling_hartree_fock.csv`, `data/selfcoupling_pairs.csv`, `data/selfcoupling_channels.csv`, `data/selfcoupling_examples.csv`	the Fierz matrix, the reduction to four invariants, the Hartree–Fock couplings, the pair couplings, the channels, and the example, as tables
`data/isometries.csv`, `data/geodesic_1644d.csv`, `data/geodesic_1671d.csv`	data of the figures of Part 1
`data/creation_single.csv`, `data/creation_history.csv`, `data/creation_anisotropic.csv`, `data/boson_creation.csv`, `data/extinction_profile.csv`, `data/extinction_checks.csv`	data of the figures of Part 7
`figures/fig_isometries.png`, `fig_geodesics.png`, `fig_creation.png`, `fig_extinction.png`	the four figures, drawn by `code/scripts/make_figures.py`
`docs_src/dirac matrices deflating.src.md`	the source of this document

References

1. D. Lovelock, "The Einstein tensor and its generalizations", J. Math. Phys. **12**, 498 (1971).
2. D. G. Boulware and S. Deser, "String-generated gravity models", Phys. Rev. Lett. **55**, 2656 (1985).
3. D. R. Brill and J. A. Wheeler, "Interaction of neutrinos and gravitational fields", Rev. Mod. Phys. **29**, 465 (1957).
4. M. Fierz, "Zur Fermischen Theorie des β-Zerfalls", Z. Phys. **104**, 553 (1937).

5. L. Parker, “Quantized fields and particle creation in expanding universes. I”, Phys. Rev. **183**, 1057 (1969).
6. G. W. Gibbons and S. W. Hawking, “Cosmological event horizons, thermodynamics, and particle creation”, Phys. Rev. D **15**, 2738 (1977).
7. N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, 1982).