

Applied Data Analysis (CS401)



Lecture 5
Regression
analysis
19 Oct 2022

EPFL

Robert West



Announcements

- Homework H1 due end of next week
 - Normally Fri, but grace period until Sat 29 Oct 23:59 (due to slightly delayed release date)
- Project milestone P1 feedback to be released next week
- Friday's lab session:
 - Quiz 4
 - Exercise on regression analysis
 - Homework office hours
 - Questions? Ask them on Ed under "[FAQ](#)"!
- Indicative course feedback is being collected (until Sun 23 Oct)

Feedback


Give us feedback on this lecture here:

<https://go.epfl.ch/ada2022-lec5-feedback>

- What did you (not) like about this lecture?
- What was (not) well explained?
- On what would you like more (fewer) details?
- ...

Course eval (“indicative feedback”) open until 23 Oct

Go to <https://isa.epfl.ch> now!



The screenshot displays the ISA EPFL website interface. The top navigation bar includes links for Home, Courses, Courses booklets, Results, Exams, Language Centre, Personal Details, Delegates, Tutoring, Exchange application, Projects, and Internships and projects in enterprise. The main content area is titled 'Horaires' and features a date selection tool set to 'Week starting 15.11.2021 to 21.11.2021'. Below this is a timetable table showing courses for Monday through Saturday. The right sidebar contains a list of links, with 'Indicative feedback (BA/MA)' highlighted by a large green arrow. A red box at the bottom right contains a message about the deadline for submitting grades.

Courses

- Biological modeling of neural networks
- Computational linear algebra
- Data in context: Critical Data Studies II
- Deep learning
- Distributed intelligent systems
- Image processing II
- Introduction to database systems
- Optimization for machine learning
- Software security
- Systems for data science

Data Science, 2021-2022, Master semester 3

- Distributed information systems
- Foundations of Data Science
- Information security and privacy
- Numerical integration of dynamical systems
- Stochastic simulation

List of students on the course

Horaires

Refresh Export iCalendar Export Excel Export PDF List

Choose the date: <<from 15.11.2021 to 21.11.2021>> One week

Week starting 15.11.2021 to 21.11.2021

	Mo	Tu	We	Th	Fr	Sa
8h - 9h		Numerical integration of dynamical systems MAA110 MATH-452 Lecture		Stochastic simulation MAB1486 MATH-414 Exercises	Foundations of Data Science INM200 COM-406 Lecture	
9h - 10h		Numerical integration of dynamical systems MAA110 MATH-452 Lecture		Stochastic simulation MAB1486 MATH-414 Exercises	Foundations of Data Science INM200 COM-406 Lecture	
10h -			Foundations of Data Science INM200 COM-406 Lecture		Foundations of Data Science	

Administration messages

Délai de rendu des notes pour les enseignants

Vos notes ne seront pas forcément toutes visibles dans votre portail au terme du délai de rendu de notes. Si c'est le cas, merci de patienter quelques jours ou de contacter votre enseignant.



Linear
regression

Credits

- Much of the material in this lecture is based on Andrew Gelman and Jennifer Hill's great book "Data Analysis Using Regression and Multilevel/Hierarchical Models", available for free [here](#)
- For a neat and gentle written intro to linear regression, especially check out chapters 3 and 4

What you should already know about linear regression



POLLING TIME

- “How familiar are you with linear regression?”
- Scan QR code or go to <https://web.speakup.info/room/join/66626>



Linear regression as you know it

- **Given:** n data points (X_i, y_i) , where X_i is k -dimensional vector of predictors (a.k.a. features) of i -th data point, and y_i is scalar outcome
- **Goal:** find the optimal coefficient vector $\beta = (\beta_1, \dots, \beta_k)$ for approximating the y_i 's as a linear function of the X_i 's:

$$y_i = X_i \beta + \epsilon_i$$

Scalar product (a.k.a. dot product) of 2 vectors


$$= \beta_1 X_{i1} + \dots + \beta_k X_{ik} + \epsilon_i, \quad \text{for } i = 1, \dots, n$$

where ϵ_i are error terms that should be as small as possible

- X_{i1} usually the constant 1 (by def) $\Rightarrow \beta_1$ a constant intercept

Example with one predictor

$$y \approx \beta_1 + \beta_2 X$$



Linear regression as you know it

- **Given:** n data points (X_i, y_i) , where X_i is k -dimensional vector of predictors (a.k.a. features), and y_i is scalar outcome, of i -th data point

- **Goal:** find the optimal coefficient vector $\beta = (\beta_1, \dots, \beta_k)$ for approximating the y 's as a linear function of the X 's:

$$y_i = X_i \beta + \epsilon_i$$

$$= \beta_1 X_{i1} + \dots + \beta_k X_{ik} + \epsilon_i, \quad \text{for } i = 1, \dots, n$$

where ϵ_i are error terms that should be as small as possible


- X_{i1} usually the constant 1 $\rightarrow \beta_1$ a constant intercept

Optimality criterion: least squares

$$y_i = X_i\beta + \epsilon_i \quad \text{for } i = 1, \dots, n$$

- Intuitively, want errors ϵ_i to be as small as possible
- Technically, want sum of squared errors as small as possible
 \Leftrightarrow find $\hat{\beta}$ such that we minimize

$$\sum_{i=1}^n (y_i - X_i\hat{\beta})^2$$



Use cases of regression



- **Prediction:** use fitted model to estimate outcome y for a new X not seen during model fitting (if you've seen regression before, then probably in the context of prediction)
- **Descriptive data analysis:** compare average outcomes across subgroups of data (today!)
- **Causal modeling:** understand how outcome y changes when you manipulate predictors X (next lecture is about causality, although not primarily using regression)

Regression as comparison of average outcomes


Example with one binary predictor X_i

No Yes

- $X_i = \text{mom_hs} = \text{"Did mother finish high school?"} \in \{0, 1\}$
- $y_i = \text{kid_score} = \text{child's score on cognitive test} \in [0, 140]$

$$y_i = \beta_1 + \beta_2 X_i + \epsilon_i$$

$$\text{kid_score} = 78 + 12 \cdot \text{mom_hs} + \text{error}$$



One binary predictor X_i :

Interpretation of fitted parameters β


$$y_i = \beta_1 + \beta_2 X_i + \epsilon_i$$

- **Intercept** β_1 : mean outcome for data points i with $X_i = 0$
- **Slope** β_2 : difference in mean outcomes between data points with $X_i = 1$ and data points with $X_i = 0$

- Reason: means minimize least-squares criterion:

$\sum_{i=1}^n (y_i - m)^2$ is minimized w.r.t. m when

$-2 \sum_{i=1}^n (y_i - m) = 0$, i.e., when $m = (1/n) \sum_{i=1}^n y_i$



One binary predictor X_i :

Interpretation of fitted parameters β

$$y_i = \beta_1 + \beta_2 X_i + \epsilon_i$$



- **Intercept** β_1 : mean outcome for data points i with $X_i = 0$
- **Slope** β_2 : difference in mean outcomes between data points with $X_i = 1$ and data points with $X_i = 0$
- Reason: means minimize least-squares criterion:

$$\sum_{i=1}^n (y_i - \beta_1 - \beta_2 X_i)^2$$

So why not just compute the two means separately and then compare them?

when

$$(1/n)$$



What a mean monkey!


Example with one continuous predictor X_i

- $X_i = \text{mom_iq} = \text{mother's IQ score} \in [70, 140]$
- $y_i = \text{kid_score} = \text{child's score on cognitive test} \in [0, 140]$

$$y_i = \beta_1 + \beta_2 X_i + \epsilon_i$$

$$\text{kid_score} = 26 + 0.6 \cdot \text{mom_iq} + \text{error}$$

estimated
(hypothetical) mean
kid_score for moms
with IQ = 0: 26




estimated mean
kid_score for moms
with IQ = 100: $26 +$
 $0.6 \cdot 100 = 86$



One continuous predictor X_i :

Interpretation of fitted parameters β

$$y_i = \beta_1 + \beta_2 X_i + \epsilon_i$$


- **Intercept** β_1 : estimated mean outcome for data points i with $X_i = 0$
- **Slope** β_2 : difference in estimated mean outcomes between data points whose X_i 's differ by 1
- Why “estimated”? → e.g., A scatter plot showing a cloud of blue data points. A horizontal red line is drawn across the plot, representing a regression line with a slope of zero. The axes are represented by black arrows.
- NB: for binary predictor, we got “exact” instead of “estimated”

Example with multiple predictors

- ($X_{i1} = 1 = \text{constant}$)
- $X_{i2} = \text{mom_hs} = \text{“Did mother finish high school?”} \in \{0, 1\}$ No Yes
- $X_{i3} = \text{mom_iq} = \text{mother’s IQ score} \in [70, 140]$
- $y_i = \text{kid_score} = \text{child’s score on cognitive test} \in [0, 140]$


$$y_i = \beta_1 + \beta_2 X_{i2} + \beta_3 X_{i3} + \epsilon_i$$

$$\text{kid_score} = 26 + 6 \cdot \text{mom_hs} + 0.6 \cdot \text{mom_iq} + \text{error}$$



Example with multiple predictors

$$\text{kid_score} = 26 + 6 \cdot \text{mom_hs} + 0.6 \cdot \text{mom_iq} + \text{error}$$



Example with **interaction** of predictors


- $X_{i2} = \text{mom_hs} = \text{"Did mother finish high school?"}$ $\overset{\text{No Yes}}{\in \{0, 1\}}$
- $X_{i3} = \text{mom_iq} = \text{mother's IQ score} \in [70, 140]$
- $y_i = \text{kid_score} = \text{child's score on cognitive test} \in [0, 140]$

$$y_i = \beta_1 + \beta_2 X_{i2} + \beta_3 X_{i3} + \beta_4 \text{mom_hs} \cdot \text{mom_iq} + \epsilon_i$$



$$\text{kid_score} = -11 + 51 \cdot \text{mom_hs} + 1.1 \cdot \text{mom_iq} - 0.5 \cdot \text{mom_hs} \cdot \text{mom_iq} + \text{error}$$

Example with **interaction** of predictors



$$\text{kid_score} = -11 + 51 \cdot \text{mom_hs} + 1.1 \cdot \text{mom_iq} - 0.5 \cdot \text{mom_hs} \cdot \text{mom_iq} + \text{error}$$



So why not just compute
the two means separately
and then compare them?



So why not just compute the two means separately and then compare them?



Mom drives Mercedes Mom doesn't drive Mercedes

Mom finished high school

avg kid_score

90

avg kid_score

90

Mom didn't finish high school

avg kid_score

78

avg kid_score

78

Mom drives Mercedes Mom doesn't drive Mercedes

Mom finished high school

990
women

10
women

Mom didn't finish high school

10
women

990
women


	Mom drives Mercedes	Mom doesn't drive Mercedes		Mom drives Mercedes	Mom doesn't drive Mercedes
Mom finished high school	avg kid_score 90	avg kid_score 90	Mom finished high school	990 women	10 women
Mom didn't finish high school	avg kid_score 78	avg kid_score 78	Mom didn't finish high school	10 women	990 women

THINK FOR A MINUTE:

**What is the mean outcome for Mercedes-driving moms vs. for non-Mercedes-driving moms?
Compare the two means! What does the comparison tell you about the link between Mercedes-driving and kid_score?**

(Feel free to discuss with your neighbor.)

- Mean kid_score for Mercedes drivers: $0.99 \cdot 90 + 0.01 \cdot 78 \approx 90$
- Mean kid_score for non-Mercedes drivers: $0.01 \cdot 90 + 0.99 \cdot 78 \approx 78$
- But really driving Mercedes makes no difference (for fixed high-school predictor)!
- Root of evil: **correlation** between finishing high school and driving Mercedes
- **Regression** to the rescue: $\text{kid_score} = 78 + 12 \cdot \text{mom_hs} + 0 \cdot \text{mercedes} + \text{error}$

	Mercedes	No Mercedes		Mercedes	No Mercedes
Mom finished high school	mean kid_score 90	mean kid_score 90	<div>Aha!</div> 	990 women	10 women
Mom didn't finish high school	mean kid_score 78	mean kid_score 78		10 women	990 women

Quantifying uncertainty

Quantifying uncertainty

- Statistical software gives you more than just coefficients β :

Aha!

Residuals:

Min	1Q	Median	3Q	Max
-52.873	-12.663	2.404	11.356	49.545

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	25.73154	5.87521	4.380	1.49e-05	***
mom.hs	5.95012	2.21181	2.690	0.00742	**
mom.iq	0.56391	0.06057	9.309	< 2e-16	***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 18.14 on 431 degrees of freedom
Multiple R-Squared: 0.2141, Adjusted R-squared: 0.2105
F-statistic: 58.72 on 2 and 431 DF, p-value: < 2.2e-16

p-value: probability of estimating such an extreme coefficient if the true coefficient were zero (= null hypothesis)

Residuals and R^2

- **Residual** for data point i : estimation error on data point i :

$$r_i = y_i - X_i\hat{\beta}$$

- Mean of residuals = 0
(total overestimation = total underestimation)

- Variance of residuals


= avg squared distance of predicted value from observed value

= “unexplained variance”

- Fraction of variance explained by the model:

$$R^2 = 1 - \hat{\sigma}^2 / s_y^2$$


Variance of
outcomes y



Residuals and R^2

- **Residual** for data point i : estimation error on data point i :

$$r_i = y_i - X_i \hat{\beta}$$



Aha!

Residuals = 0

(Estimation = total underestimation)


Residuals

avg squared distance of predicted value from observed value
= “unexplained variance”

Fraction of variance explained by the model:

$$R^2 = 1 - \hat{\sigma}^2 / s_y^2$$


Variance of
outcomes y



Coefficient of determination: R^2

$$R^2 = 1 - \hat{\sigma}^2 / s_y^2$$


Fitted Line Plot
Output1 = 44.53 + 2.024 Input



S	283.042
R-Sq	14.7%
R-Sq(adj)	14.3%

$$R^2 = 0.147$$





Fitted Line Plot
Output2 = 44.86 + 2.134 Input







S	48.8860
R-Sq	86.5%
R-Sq(adj)	86.5%

$$R^2 = 0.865$$

Coefficient of determination: R^2



Coefficient of determination: R^2



$R^2 = 0.67$ everywhere!

Course eval (“indicative feedback”) open until 23 Oct

Go to <https://isa.epfl.ch> now!

The screenshot displays the ISA EPFL website interface. At the top, a navigation bar includes links for Home, Courses, Courses booklets, Results, Exams, Language Centre, Personal Details, Delegates, Tutoring, Exchange application, Projects, and Internships and projects in enterprise. The main content area is divided into three sections:

- Courses:** A list of courses including Biological modeling of neural networks, Computational linear algebra, Data in context: Critical Data Studies II, Deep learning, Distributed intelligent systems, Image processing II, Introduction to database systems, Optimization for machine learning, Software security, and Systems for data science. Below this, a section for "Data Science, 2021-2022, Master semester 3" lists Distributed information systems, Foundations of Data Science, Information security and privacy, Numerical integration of dynamical systems, and Stochastic simulation.
- Horaires:** A timetable section for the week starting 15.11.2021 to 21.11.2021. It includes a date picker set to "One week" and a table of courses.
- Administration messages:** A section titled "Délai de rendu des notes pour les enseignants" (Note submission deadline for teachers) with a message in French.

A large green arrow points to the "Administration messages" section.

	Mo	Tu	We	Th	Fr	Sa
8h - 9h		Numerical integration of dynamical systems MAA110 MATH-452 Lecture		Stochastic simulation MAB1486 MATH-414 Exercises	Foundations of Data Science INM200 COM-406 Lecture	
9h - 10h		Numerical integration of dynamical systems MAA110 MATH-452 Lecture		Stochastic simulation MAB1486 MATH-414 Exercises	Foundations of Data Science INM200 COM-406 Lecture	
10h -			Foundations of Data Science INM200 COM-406 Lecture		Foundations of Data Science	

Délai de rendu des notes pour les enseignants

Vos notes ne seront pas forcément toutes visibles dans votre portail au terme du délai de rendu de notes. Si c'est le cas, merci de patienter quelques jours ou de contacter votre enseignant.

Assumptions made in regression modeling

Assumptions for regression modeling

1. Validity:
 - a. Outcome measure should accurately reflect the phenomenon of interest
 - b. Model should include all relevant predictors
 - c. Model should generalize to cases to which it will be applied

Assumptions for regression modeling (2)

2. Additivity and linearity:

$$\begin{aligned}y_i &= X_i\beta + \epsilon_i \\ &= \beta_1 X_{i1} + \cdots + \beta_k X_{ik} + \epsilon_i, \quad \text{for } i = 1, \dots, n\end{aligned}$$

But very flexible: we require linearity in predictors (not necessarily in raw inputs); predictors can be arbitrary functions of raw inputs, e.g.,

- logarithms, polynomials, reciprocals, ...
- interactions (i.e., products) of multiple inputs
- discretization of raw inputs, coded as indicator variables

Assumptions for regression modeling (3)

- 3. Independence of errors: no interaction between data points
 - 4. Equal variance of errors
 - 5. Normality (Gaussianity) of errors
- } less important
in practice

Transformations of predictors and outcomes

Transformations of predictors

- When we apply linear transformations to predictors, the model remains “equally good”:
 - The fitted coefficients may change, but predicted outcomes and model fit (R^2) won't change
- For instance,

$$\text{earnings} = -61000 + 51 \cdot \text{height (in millimeters)} + \text{error}$$

$$\text{earnings} = -61000 + 81000000 \cdot \text{height (in miles)} + \text{error}.$$


Mean-centering of predictors

- Compute the mean value of a predictor over all data points, and subtract it from each value of that predictor:


$$X_{ik} \leftarrow X_{ik} - \text{mean}(X_{1k}, \dots, X_{nk})$$

- \Rightarrow the predictor X_{ik} now has mean 0

(hypothetical) mean
kid_score for moms
with IQ = 0: 26



mean kid_score for
moms with mean IQ: 86



After mean-centering of predictors, ...

... you have a convenient interpretation of coefficients β_k of main predictors (i.e., non-interaction predictors):

- $k = 1$ (i.e., intercept):
 - Estimated mean outcome when each predictor has its mean value
- $k > 1$:
 - Model w/o interactions: estimated mean increase in outcome y for each unit increase in X_{ik}
 - Model with interactions: estimated mean increase in outcome y for each unit increase in X_{ik} **when each other predictor has its mean value**

Standardization via z-scores


- First **mean-center** all predictors, then **divide them by their standard deviations**:

$$X_{ik} \leftarrow [X_{ik} - \text{mean}(X_{1k}, \dots, X_{nk})] / \text{sd}(X_{1k}, \dots, X_{nk})$$

- Resulting values are called “**z-scores**”
- All predictors now have the same units:
distance (in terms of standard deviations) from the mean
- This lets us compare coefficients for predictors with previously incomparable units of measurement, e.g., IQ score vs. earnings in Swiss francs vs. height in centimeters

Logarithmic outcomes

- **Practical:** makes sense if the outcome y follows a heavy-tailed distribution
- Only works for non-negative outcomes
- **Theoretical:** turns an additive model into a **multiplicative model**:



$$\log y_i = b_0 + b_1 X_{i1} + b_2 X_{i2} + \cdots + \epsilon_i$$

Exponentiating both sides yields

$$\begin{aligned} y_i &= e^{b_0 + b_1 X_{i1} + b_2 X_{i2} + \cdots + \epsilon_i} \\ &= B_0 \cdot B_1^{X_{i1}} \cdot B_2^{X_{i2}} \cdots E_i \end{aligned}$$

Logarithmic outcomes: Interpreting coefficients

$$\begin{aligned}y_i &= e^{b_0 + b_1 X_{i1} + b_2 X_{i2} + \dots + \epsilon_i} \\ &= B_0 \cdot B_1^{X_{i1}} \cdot B_2^{X_{i2}} \dots E_i\end{aligned}$$

- An **additive** increase of 1 in predictor $X_{.1}$ is associated with a **multiplicative** increase of $B_1 = \exp(b_1)$ in the outcome
- If $b_1 \approx 0$, we can immediately interpret b_1 (without needing to exponentiate it first to get B_1 !) as the **relative increase** in outcomes, since $\exp(b_1) \approx 1 + b_1$
- E.g., $b_1 = 0.05 \Rightarrow B_1 = \exp(b_1) \approx 1.05$
 \Rightarrow “+1 in predictor $X_{.1}$ ” is associated with “+5% in outcome”


Going beyond linear regression for comparing means

Beyond linear regression: generalized linear models

- Logistic regression: binary outcomes
- Poisson regression: non-negative integer outcomes (e.g., counts)

Beyond comparing means; or, A taste of causality: “Difference in differences”

- Two groups: P , S
- At time 2, group P receives a **treatment**, group S doesn't
- Question: Did the treatment have an **effect**? If so, how large was it?
- P and S don't start out the same at time 1
- There is a temporal “baseline effect”




Beyond comparing means; or, A taste of causality: “Difference in differences” (2)

- Elegant linear model with binary predictors:

$$y_{it} = a + b \cdot \text{treated}_i + c \cdot \text{time2}_t + d \cdot (\text{treated}_i \cdot \text{time2}_t) + \text{error}$$

- d = treatment effect
- All of this with one single regression!
- You get quantification of uncertainty (significance) for free!




Beyond comparing means; or, A taste of causality: “Difference in differences” (2)

What a treat!

- Elegant predictors.


$$y_{it} = a + b \cdot \text{treated}_i + c \cdot \text{time2}_t + d \cdot (\text{treated}_i \cdot \text{time2}_t) + \text{error}$$

- d = treatment effect
- All of this with one single regression!
- You get quantification of uncertainty (significance) for free!



A bonanza of causality: Next lecture!

It!, my banana is
non-linear...




Summary

- Linear regression as a tool for comparing means across subgroups of data
- How? Read group means off from fitted coefficients
- Advantages over plain comparison of means “by hand”:
 - Accounting for correlations among predictors
 - Quantification of uncertainty (significance) “for free”
 - Additive or multiplicative model: all it takes is a log
- Caveat emptor:
 - Model must be appropriately specified, else nonsense results → stay critical, run diagnostics (e.g., R^2)

Course eval (“indicative feedback”) open until 23 Oct

Go to <https://isa.epfl.ch> now!



The screenshot displays the ISA EPFL website interface. The top navigation bar includes links for Home, Courses, Courses booklets, Results, Exams, Language Centre, Personal Details, Delegates, Tutoring, Exchange application, Projects, and Internships and projects in enterprise. The main content area is titled 'Horaires' and features a date selection tool set to 'Week starting 15.11.2021 to 21.11.2021'. Below this is a timetable table showing courses for Monday through Saturday. The left sidebar lists various courses, including 'Data Science, 2021-2022, Master semester 3'. The right sidebar contains a list of links, with 'Indicative feedback (BA/MA)' highlighted by a large green arrow. Below the links is a section for 'Administration messages' with a message about the deadline for submitting grades.

Courses

- Biological modeling of neural networks
- Computational linear algebra
- Data in context: Critical Data Studies II
- Deep learning
- Distributed intelligent systems
- Image processing II
- Introduction to database systems
- Optimization for machine learning
- Software security
- Systems for data science

Data Science, 2021-2022, Master semester 3

- Distributed information systems
- Foundations of Data Science
- Information security and privacy
- Numerical integration of dynamical systems
- Stochastic simulation

List of students on the course

Horaires

Refresh Export iCalendar Export Excel Export PDF List

Choose the date: <<from 15.11.2021 to 21.11.2021>>

One week

Week starting 15.11.2021 to 21.11.2021

	Mo	Tu	We	Th	Fr	Sa
8h - 9h		Numerical integration of dynamical systems MAA110 MATH-452 Lecture		Stochastic simulation MAB1486 MATH-414 Exercises	Foundations of Data Science INM200 COM-406 Lecture	
9h - 10h		Numerical integration of dynamical systems MAA110 MATH-452 Lecture		Stochastic simulation MAB1486 MATH-414 Exercises	Foundations of Data Science INM200 COM-406 Lecture	
10h -			Foundations of Data Science INM200 COM-406 Lecture		Foundations of Data Science	

Administration messages

Délai de rendu des notes pour les enseignants

Vos notes ne seront pas forcément toutes visibles dans votre portail au terme du délai de rendu de notes. Si c'est le cas, merci de patienter quelques jours ou de contacter votre enseignant.

Feedback

Give us feedback on this lecture here:

<https://go.epfl.ch/ada2022-lec5-feedback>

- What did you (not) like about this lecture?
- What was (not) well explained?
- On what would you like more (fewer) details?
- ...

Credits

- Much of the material in this lecture is based on Andrew Gelman and Jennifer Hill's great book "Data Analysis Using Regression and Multilevel/Hierarchical Models", available for free [here](#)
- For a neat and gentle written intro to linear regression, especially check out chapters 3 and 4

Bonus: Logarithmic outcomes and predictors

Interpretation of coefficient of logarithmic predictor:

- **Multiplicative** increase by 1% in predictor X_1 is associated with a **multiplicative** increase by $b_1\%$ in the outcome
- Why?
 - $\log(y) = a + b \log(X) \Rightarrow y = \exp(a) * X^b$
 - Multiplying X by a factor c multiplies y by a factor of c^b
 - $c^b \approx 1 + b*(c-1)$ for $c \approx 1$ (hint: Taylor approximation!)
 - Example when using $c = 1.01$ (i.e., increase by 1%):
 $b = 2 \Rightarrow$ increasing X by 1% increases y by 2%