

MASM22/FMSN30: Linear and Logistic Regression, 7.5 hp

FMSN40: ... with Data Gathering, 9 hp

Lecture 1, spring 2023

Linear regression: assumptions and estimates

<https://canvas.education.lu.se/courses/23000>

Mathematical Statistics / Centre for Mathematical Sciences
Lund University


20/3-23

Simple Linear Regression

We measure two variables, x and Y . How does the value of Y depend on the value of x ? Is there a linear relationship? How can we estimate this relationship using observed data?

Example: Ice cream

An ice cream manufacturer suspects that storing ice cream at low temperatures leads to weight loss.




Multiple linear regression

We measure several variables, x_1, \dots, x_p , and Y . How does the value of Y depend on the values of x_1, \dots, x_p ?

Example: Iris

The petal length depends on sepal length, sepal width and species.



Basic assumptions

- ▶ Y : **continuous** dependent variable, “response” or “outcome”, assumed **random**.
- ▶ x_1, \dots, x_p : explanatory variables, “covariates”; assumed **non-random**.
- ▶ We *hypothesize* that Y has a linear relationship with x_1, \dots, x_p , on average, and follows the **linear model**:

$$E(Y) = \beta_0 + \beta_1 x_1 + \dots + \beta_p x_p$$

- ▶ $\beta_0, \beta_1, \dots, \beta_p$: unknown parameters, assumed **non-random**.
- ▶ β_0 = intercept; $E(Y)$ when $x_1 = \dots = x_p = 0$,
- ▶ β_1, \dots, β_p = slopes in the corresponding x -directions; the additive change in $E(Y)$ when the corresponding x -variable is increased by 1 unit and the others are held fixed.


Model: multiple linear regression

- ▶ We denote with Y_i , where $i = 1, \dots, n$, the i th observation from a set of n measurements of Y .
- ▶ We denote with x_{ij} , where $i = 1, \dots, n$ and $j = 1, \dots, p$, the corresponding i th observation of the j th x -variable.
- ▶ The model for a generic observation Y_i is

$$Y_i = \beta_0 + \beta_1 x_{i1} + \dots + \beta_p x_{ip} + \epsilon_i, \quad i = 1, \dots, n$$


where ϵ_i is the “measurement error” which contains all the random variation not explained by the linear model.

Ice cream: weight loss by time in storage



data and fitted line

Iris: petal length as function of sepal length and width, $p = 2$



Assumptions for the measurement error

Besides linearity, we also assume for all $i = 1, \dots, n$

$$E(\epsilon_i) = 0$$

$$E(Y_i) = \beta_0 + \beta_1 x_{i1} + \dots + \beta_p x_{ip} = \mu_i$$

$$V(\epsilon_i) = \sigma^2$$

$$V(Y_i) = \sigma^2$$

$$\epsilon_i \sim N(0, \sigma^2)$$

$$Y_i \sim N(\mu_i, \sigma^2)$$

ϵ_i are pairwise independent

Y_i are pairwise independent

A note on notation

Strictly speaking, we assume properties of Y_i *conditional* on $X_1 = x_{i1}$:

$$E(Y_i \mid X_1 = x_{i1}, \dots, X_p = x_{ip}) = \beta_0 + \beta_1 x_{i1} + \dots + \beta_p x_{ip} = \mu_i,$$



$$V(Y_i \mid X_1 = x_{i1}, \dots, X_p = x_{ip}) = \sigma^2,$$

$$Y_i \mid X_1 = x_{i1}, \dots, X_p = x_{ip} \sim N(\mu_i, \sigma^2)$$

We consider this notation as implicit and always write $E(Y_i)$, etc., in place of $E(Y_i \mid X_1 = x_{i1}, \dots)$... except on the next slide...

Warning!

Our assumptions imply that the conditional distribution of $Y_i | X = x_i$ is Normal but we don't know the marginal distribution of Y_i ignoring X !



The marginal distribution of Y is clearly not Normal but this is just due to the strange distribution of the x -values.

To assess normality you should instead inspect **residuals** (introduced later), which means you cannot always see that your model will be wrong (or right!) until after you have fitted it!

Multiple linear regression with matrices

$$\mathbf{Y} = \begin{pmatrix} Y_1 \\ Y_2 \\ \vdots \\ Y_n \end{pmatrix}, \mathbf{X} = \begin{pmatrix} 1 & x_{11} & \cdots & x_{1p} \\ 1 & x_{21} & \cdots & x_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & x_{n1} & \cdots & x_{np} \end{pmatrix}, \boldsymbol{\epsilon} = \begin{pmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_n \end{pmatrix}, \boldsymbol{\beta} = \begin{pmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_p \end{pmatrix}$$

$$\mathbf{Y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon} = \begin{pmatrix} 1 \cdot \beta_0 + x_{11} \cdot \beta_1 + \dots + x_{1p} \cdot \beta_p \\ 1 \cdot \beta_0 + x_{21} \cdot \beta_1 + \dots + x_{2p} \cdot \beta_p \\ \vdots \\ 1 \cdot \beta_0 + x_{n1} \cdot \beta_1 + \dots + x_{np} \cdot \beta_p \end{pmatrix} + \begin{pmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_n \end{pmatrix}$$
$$\sim N_n(\mathbf{X}\boldsymbol{\beta}, \sigma^2 \mathbf{I})$$

(n -dimensional multivariate normal distribution)

We assume $\epsilon \sim N_n(\mathbf{0}, \sigma^2 \mathbf{I})$ where $E(\epsilon) = \mathbf{0} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$ and

the covariance matrix $\text{Var}(\epsilon)$ is given by

$$\begin{aligned} \text{Var}(\epsilon) &= \begin{pmatrix} V(\epsilon_1) & C(\epsilon_1, \epsilon_2) & \dots & C(\epsilon_1, \epsilon_n) \\ C(\epsilon_2, \epsilon_1) & V(\epsilon_2) & \dots & C(\epsilon_2, \epsilon_n) \\ \vdots & \vdots & \ddots & \vdots \\ C(\epsilon_n, \epsilon_1) & C(\epsilon_n, \epsilon_2) & \dots & V(\epsilon_n) \end{pmatrix} \\ &= \begin{pmatrix} \sigma^2 & 0 & \dots & 0 \\ 0 & \sigma^2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \sigma^2 \end{pmatrix} = \sigma^2 \cdot \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix} = \sigma^2 \mathbf{I} \end{aligned}$$

Least squares estimates: simple linear without matrices

Find β_0, β_1 that minimize the loss function

$$Q(\beta_0, \beta_1) = \sum_{i=1}^n (Y_i - E(Y_i))^2 = \sum_{i=1}^n (Y_i - (\beta_0 + \beta_1 x_i))^2.$$

Partial derivatives

Find the minimum by solving the linear equation system

$$\frac{\partial Q}{\partial \beta_0} = -2 \sum_{i=1}^n (Y_i - \beta_0 - \beta_1 x_i) = 0 \Leftrightarrow n\beta_0 + \beta_1 \sum_{i=1}^n x_i = \sum_{i=1}^n Y_i$$

$$\frac{\partial Q}{\partial \beta_1} = -2 \sum_{i=1}^n x_i (Y_i - \beta_0 - \beta_1 x_i) = 0 \Leftrightarrow \beta_0 \sum_{i=1}^n x_i + \beta_1 \sum_{i=1}^n x_i^2 = \sum_{i=1}^n x_i Y_i$$

$$\hat{\beta}_0 = \bar{Y} - \hat{\beta}_1 \bar{x},$$

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(Y_i - \bar{Y})}{\sum_{i=1}^n (x_i - \bar{x})^2}.$$

Least squares estimates: with matrices

Find β that minimizes $Q(\beta) = (\mathbf{Y} - \mathbf{X}\beta)'(\mathbf{Y} - \mathbf{X}\beta)$.

Partial derivatives

Expand $Q(\beta)$ and use the fact that all the terms are scalar

$$\begin{aligned} Q(\beta) &= \mathbf{Y}'\mathbf{Y} - \mathbf{Y}'\mathbf{X}\beta - \beta'\mathbf{X}'\mathbf{Y} + \beta'\mathbf{X}'\mathbf{X}\beta \\ &= \mathbf{Y}'\mathbf{Y} - 2\beta'\mathbf{X}'\mathbf{Y} + \beta'\mathbf{X}'\mathbf{X}\beta \\ \frac{\partial Q(\beta)}{\partial \beta'} &= -2\mathbf{X}'\mathbf{Y} + 2\mathbf{X}'\mathbf{X}\beta = \mathbf{0} \end{aligned}$$

The solution satisfies the normal equations: $\mathbf{X}'\mathbf{X}\beta = \mathbf{X}'\mathbf{Y}$.

$$\begin{pmatrix} n & \sum_{i=1}^n x_{i1} & \cdots & \sum_{i=1}^n x_{ip} \\ \sum_{i=1}^n x_{i1} & \sum_{i=1}^n x_{i1}^2 & \cdots & \sum_{i=1}^n x_{i1}x_{ip} \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{i=1}^n x_{ip} & \sum_{i=1}^n x_{i1}x_{ip} & \cdots & \sum_{i=1}^n x_{ip}^2 \end{pmatrix} \begin{pmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_p \end{pmatrix} = \begin{pmatrix} \sum_{i=1}^n Y_i \\ \sum_{i=1}^n x_{i1}Y_i \\ \vdots \\ \sum_{i=1}^n x_{ip}Y_i \end{pmatrix}$$

- **Parameter estimates:** $\hat{\beta}$ is the solution to the normal equations:

$$\hat{\beta} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{Y}$$

- **Predicted values** = fitted line (plane):

$$\hat{Y}_i = \hat{E}(Y_i) = \hat{\beta}_0 + \hat{\beta}_1 x_{i1} + \dots + \hat{\beta}_p x_{ip} \quad \hat{\mathbf{Y}} = \mathbf{X}\hat{\beta}$$

- **Residuals:** the difference between observations and predictions:

$$e_i = Y_i - \hat{Y}_i \quad \mathbf{e} = \mathbf{Y} - \hat{\mathbf{Y}} = \mathbf{Y} - \mathbf{X}\hat{\beta}$$

- **Residual variance:** s^2 is an estimate of the variance of the error, a measure of the “residual variability” unexplained by the model.

$$\hat{\sigma}^2 = s^2 = \frac{Q(\hat{\beta})}{n - (p + 1)} = \frac{\sum_{i=1}^n (Y_i - \hat{Y}_i)^2}{n - (p + 1)} = \frac{\mathbf{e}'\mathbf{e}}{n - (p + 1)}$$

Note: $p + 1$ is the total number of β -parameters in the model.

Properties of parameter estimates

Note: For a constant matrix \mathbf{A} and a random matrix \mathbf{Y} we have

$$E(\mathbf{AY}) = \mathbf{A}E(\mathbf{Y}) \text{ and } \text{Var}(\mathbf{AY}) = \mathbf{A}\text{Var}(\mathbf{Y})\mathbf{A}'.$$

$$E(\hat{\beta}) = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}' \underbrace{E(\mathbf{Y})}_{\mathbf{X}\beta} = \underbrace{(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{X}}_{\mathbf{I}}\beta = \beta$$

$$\begin{aligned} \text{Var}(\hat{\beta}) &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\text{Var}(\mathbf{Y})((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}')' \\ &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}' \cdot \sigma^2\mathbf{I} \cdot \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1} = \sigma^2(\mathbf{X}'\mathbf{X})^{-1} \\ &= \begin{pmatrix} V(\hat{\beta}_0) & \cdots & C(\hat{\beta}_0, \hat{\beta}_p) \\ \vdots & \ddots & \vdots \\ C(\hat{\beta}_p, \hat{\beta}_0) & \cdots & V(\hat{\beta}_p) \end{pmatrix} \end{aligned}$$

When $p = 1$:

$$\begin{aligned} V(\hat{\beta}_0) &= \sigma^2 \left(\frac{1}{n} + \frac{\bar{x}^2}{\sum_{i=1}^n (x_i - \bar{x})^2} \right), \quad V(\hat{\beta}_1) = \frac{\sigma^2}{\sum_{i=1}^n (x_i - \bar{x})^2} \\ C(\hat{\beta}_0, \hat{\beta}_1) &= C(\hat{\beta}_1, \hat{\beta}_0) = -\bar{x}V(\hat{\beta}_1) \end{aligned}$$

For a specific set of x -values, $\mathbf{x}_0 = (1 \ x_{01} \ \dots \ x_{0p})$, we have

- on average (the fitted line) $\hat{Y}_0 = \mathbf{x}_0\hat{\beta}$:

$$E(\hat{Y}_0) = \mathbf{x}_0 E(\hat{\beta}) = \mathbf{x}_0 \beta = \beta_0 + \beta_1 x_{01} + \dots + \beta_p x_{0p} = \mu_0,$$

$$V(\hat{Y}_0) = \mathbf{x}_0 \text{Var}(\hat{\beta}) \mathbf{x}_0' = \sigma^2 \mathbf{x}_0 (\mathbf{X}'\mathbf{X})^{-1} \mathbf{x}_0'$$

$$[p = 1] = \sigma^2 \left(\frac{1}{n} + \frac{(x_0 - \bar{x})^2}{\sum_{i=1}^n (x_i - \bar{x})^2} \right)$$

- for a new observation $\hat{Y}_{\text{pred}_0} = \mathbf{x}_0\hat{\beta} + \epsilon_0$:

$$E(\hat{Y}_{\text{pred}_0}) = \mathbf{x}_0 E(\hat{\beta}) + E(\epsilon_0) = \mathbf{x}_0 \beta = \mu_0,$$

$$V(\hat{Y}_{\text{pred}_0}) = \mathbf{x}_0 \text{Var}(\hat{\beta}) \mathbf{x}_0' + V(\epsilon_0) = \sigma^2 (1 + \mathbf{x}_0 (\mathbf{X}'\mathbf{X})^{-1} \mathbf{x}_0')$$

$$[p = 1] = \sigma^2 \left(1 + \frac{1}{n} + \frac{(x_0 - \bar{x})^2}{\sum_{i=1}^n (x_i - \bar{x})^2} \right)$$

Standard error

The **standard error** (s.e. = medelfel), $d(\hat{\theta}) = \sqrt{\hat{V}(\hat{\theta})}$, of an estimate $\hat{\theta}$, replaces any unknown parameters in $V(\hat{\theta})$ by their estimates. Here, replace σ by $\hat{\sigma} = s$.

Comments

- ▶ $\hat{\beta}$ exists only when $\mathbf{X}'\mathbf{X}$ is non-singular. Near-singularity makes all β -estimates very uncertain.
- ▶ Never calculate $(\mathbf{X}'\mathbf{X})^{-1}$ “by hand”. Take a course in Numerical analysis (or rely on R).
- ▶ The uncertainty of the β -estimates is mostly due to the sample size, n , and the structure of the x -variables, as expressed in $(\mathbf{X}'\mathbf{X})^{-1}$.
- ▶ The uncertainty of the predictions \hat{Y}_0 is **also** due to how far \mathbf{x}_0 is from the midpoint of the x -variables.
- ▶ $V(\hat{Y}_{\text{pred}_0}) \rightarrow \sigma^2 > 0$ when $V(\hat{Y}_0) \rightarrow 0$.

Ice cream: estimates

Y = weight loss (g), x = storage time (weeks).

Model $Y = \beta_0 + \beta_1 x + \epsilon$

Variable	parameter	estimate	s.e.	unit
intercept (time = 0)	β_0	-5.7	0.81	g
storage time	β_1	1.33	0.03	g/week
resid.std.dev	σ	0.80		g

Fitted line: $\hat{Y} = -5.7 + 1.33x$.

Predictions

If we store the ice cream for $x_0 = 34$ weeks, how much weight loss can we expect on average? in a single package?

	estimate	s.e.	unit
on average	$\hat{Y}_0 = -5.7 + 1.33 \cdot 34 = 39.7$	0.15	g
single package	$\hat{Y}_{\text{pred}_0} = 39.7 + \epsilon_0$	0.82	g

Note: $0.82 = \sqrt{0.80^2 + 0.15^2}$.